Contraction-induced endocardial *id2b* plays a dual role in regulating myocardial contractility

and valve formation

Shuo Chen[1, 2, 3#], Jinxiu Liang[1, 2, 3#], Jie Yin[1, 2, 3], Weijia Zhang[1, 2, 3], Peijun Jiang[1, 2, 3], Wenyuan Wang[4, 5],

Xiaoying Chen[6, 7], Yuanhong Zhou[8], Peng Xia[8], Fan Yang[6, 7], Ying Gu[1,2, 3], Ruilin Zhang[4, 5], Peidong

Han[1, 2, 3*]

[1]Department of Cardiology, Center for Genetic Medicine, The Fourth Affiliated Hospital of School of

Medicine, and International School of Medicine, International Institutes of Medicine, Zhejiang

University, Yiwu, China, 322000.

[2]Division of Medical Genetics and Genomics, The Children's Hospital, Zhejiang University School of

Medicine, National Clinical Research Center for Child Health, Hangzhou, Zhejiang 310058, China.

[3]Institute of Genetics, Zhejiang University School of Medicine, Hangzhou, Zhejiang, China.

[4]TaiKang Medical School (School of Basic Medical Sciences), Wuhan University, Wuhan, China

[5]Hubei Provincial Key Laboratory of Developmentally Originated Disease, Wuhan, China

[6]Department of Biophysics, and Kidney Disease Center of the First Affiliated Hospital, Zhejiang University

School of Medicine, Hangzhou, China.

[7]Liangzhu Laboratory, Zhejiang University Medical Center, Hangzhou, China.

[8]Zhejiang Provincial Key Laboratory for Cancer Molecular Cell Biology, Life Sciences Institute, Zhejiang

University, Hangzhou, Zhejiang 310058, China.

#Shuo Chen and Jinxiu Liang contributed equally to this work.

*To whom correspondence should be addressed.

hanpd@zju.edu.cn

+86-571-88206360

Abstract

Biomechanical cues play an essential role in sculpting organ formation. Comprehending how cardiac

cells perceive and respond to biomechanical forces is a biological process with significant medical

implications that remains poorly understood. Here we show that biomechanical forces activate

endocardial *id2b* (inhibitor of DNA binding 2b) expression, thereby promoting cardiac contractility and

valve formation. Taking advantage of the unique strengths of zebrafish, particularly the viability of

embryos lacking heartbeats, we systematically compared the transcriptomes of hearts with impaired

contractility to those of control hearts. This comparison identified *id2b* as a gene sensitive to blood

flow. By generating a knockin reporter line, our results unveiled the presence of *id2b* in the

endocardium, and its expression is sensitive to both pharmacological and genetic perturbations of

contraction. Furthermore, *id2b* loss-of-function resulted in progressive heart malformation and early

lethality. Combining RNA-seq analysis, electrophysiology, calcium imaging, and echocardiography,

we discovered profound impairment in atrioventricular (AV) valve formation and defective excitation-

contraction coupling in *id2b* mutants. Mechanistically, deletion of *id2b* reduced AV endocardial cell

proliferation and led to a progressive increase in retrograde blood flow. In the myocardium, *id2b*

directly interacted with the bHLH component *tcf3b* (transcription factor 3b) to restrict its activity.

Inactivating *id2b* unleashed its inhibition on *tcf3b*, resulted in enhanced repressor activity of *tcf3b*,

which subsequently suppressed the expression of *nrg1* (neuregulin 1), an essential mitogen for heart

development. Overall, our findings identify *id2b* as an endocardial cell-specific, biomechanical

signaling-sensitive gene, which mediates intercellular communications between endocardium and

myocardium to sculpt heart morphogenesis and function.

Keywords: Zebrafish, Heart, *id2b*, Valve, Contraction

Introduction

The heart develops with continuous contraction, and biomechanical cues play an essential role in cardiac morphogenesis[1, 2]. Blood flow is directly sensed by the surrounding endocardium, which undergoes multiscale remodeling during zebrafish heart development. In the atrioventricular canal (AVC) endocardium, oscillatory flow promotes valvulogenesis through transient receptor potential (TRP) channel-mediated expression of Krüppel-like factor 2a (*klf2a*)[3, 4, 5]. Meanwhile, mechanical forces trigger ATP-dependent activation of purinergic receptors, inducing expression of nuclear factor of activated T cells 1 (*nfatc1*) and subsequent valve formation[6]. In the chamber endocardium, blood flow induces endocardial cells to adopt chamber- and region-specific cell morphology during cardiac ballooning[7]. A recent study further emphasized that blood flow is essential for endocardial cell accrual in assembling the outflow tract[8]. Beyond their role in endocardial cells, proper biomechanical cues are indispensable for shaping the myocardium. For instance, in contraction compromised *tnnt2a*[9] and *myh6*[10] mutants, trabeculation is markedly reduced. Moreover, apart from the tissue-scale regulatory effect, the shape changes[11] and myofibril content[12] at the single-cardiomyocyte level are also sculpted by the interplay of contractility and blood flow in the developing heart.

In ventricular myocardium morphogenesis, biomechanical forces coordinate intra-organ communication between endocardial and myocardial cells by regulating BMP, Nrg/Erbb, and Notch signaling. The Nrg-Erbb axis stands as one of the most extensively studied signaling pathway mediating cell-cell communications in the heart[13, 14, 15, 16]. In particular, endocardial Notch activity induced by cardiac contraction promotes the expression of Nrg, which then secretes into the extracellular space, binding to Erbb2/4 receptor tyrosine kinases on cardiomyocyte and promoting their delamination[17, 18, 19]. In agreement with the pivotal role of this signaling pathway in heart development, genetic mutations in zebrafish *nrg2a* and *erbb2* result in severely compromised trabeculae formation[20, 21, 22].

The development of cardiomyocytes encompasses the specification of subcellular structure, metabolic state, gene expression profile, and functionality[23, 24]. The rhythmic contraction of cardiomyocytes relies on precisely regulated excitation-contraction coupling (E-C coupling), transducing electrical activity into contractile forces. This intricate signaling cascade involves

membrane action potential, calcium signaling, and sarcomeric structure[25, 26]. Specifically, membrane

depolarization triggers the opening of L-type calcium channel (LTCC), allowing calcium influx. The

calcium signaling then activates the ryanodine receptor on the sarcoplasmic reticulum (SR) membrane,

releasing additional calcium[25]. E-C coupling is essential for heart development, as evidenced by the

complete silence of the ventricle and reduced cardiomyocyte number in *cacna1c* (LTCC α1 subunit in

zebrafish) mutant[27]. Beyond its role in modulating cardiac structure formation, previous studies

indicate that Nrg-Erbb2 signaling is also necessary for cardiac function, as *erbb2* mutants exhibit

severely compromised fractional shortening and an immature conduction pattern[20, 28].

Id (inhibitor of DNA binding) proteins belong to the helix-loop-helix (HLH) family and function

as transcriptional repressors[29]. Notably, Id2 lacks a DNA-binding domain and forms heterodimer with

other bHLH proteins, acting in a dominant-negative manner[30]. Id2 plays a crucial role in heart

development, and its genetic deletion results in severe cardiac defects in mice[31, 32, 33]. In zebrafish, *id2a*

and *id2b* are homologs of the mammalian *Id2* gene. However, their expression pattern and function in

the zebrafish heart remains largely unknown. In the present study, we identified that *id2b* is specifically

expressed in endocardial cells of the developing heart, and its expression is regulated by cardiac

contraction and blood flow. Genetic deletion of *id2b* led to impaired AV valve formation and reduced

cardiac contractility. Therefore, *id2b* serves as a crucial mediator linking biomechanical cues to heart

morphogenesis.

Results

Transcriptome analysis identifies *id2b* as a blood flow sensitive gene

Blood circulation is dispensable for early embryonic development in zebrafish, presenting an ideal

model to investigate biomechanical cues influencing heart morphogenesis. To identify genes affected

by cardiac contraction or blood flow, we treated *myl7:mCherry* zebrafish embryos with tricaine to

inhibit cardiac contractility from 72 hours post-fertilization (hpf) to 96 hpf. Hearts from control and

tricaine-treated zebrafish embryos were manually collected under a fluorescence stereoscope as

previously reported[34]. Subsequently, approximately 1,000 hearts from each group were subjected to

RNA sequencing (Figure 1A). A total of 4,530 genes with differential expression were identified,

comprising 2,013 up-regulated and 2,517 down-regulated genes. With a specific focus on identifying

key transcription factors (TFs) affected by perturbing biomechanical forces, differentially expressed

genes (DEGs) encoding TFs were enriched into signaling pathways through KEGG analysis.

Interestingly, our analysis identified several pathways known to be involved in heart development,

including the transforming growth factor beta (TGFβ) signaling and Notch signaling pathways (Figure

1B). In particular, the scaled expression levels of the top 6 DEGs ($|\log_2 FC| \geq 0.585$), exhibiting up- or

down-regulation, were listed (Figure 1C). Intriguingly, *Id2* has been shown to regulate murine AV

valve formation, a process notably influenced by alterations in blood flow directionality. Moreover,

loss of *Id2* leads to malformation of both the arterial and venous poles of the heart and disrupts AV

valve morphogenesis[31, 33]. Therefore, we interrogated the expression and function of *id2b* in developing

embryos.

Quantitative real-time PCR (qRT-PCR) analysis of purified embryonic hearts revealed a

significant reduction in *id2b* mRNA levels and an increase in *id2a* levels following tricaine treatment

from 72 to 96 hpf compared to controls (Figure 1D). Furthermore, in situ hybridization was performed

to visualize *id2b* expression under tricaine or 10 µM blebbistatin (an inhibitor of sarcomeric function

and cardiac contractility) treatment from 48 to 72 or from 72 to 96 hpf as previously described[5].

Consistently, our results showed a reduction in *id2b* signal in contraction-deficient hearts compared to

the control (Figure 1E). In cardiomyocytes, *tnnt2a* encodes a key sarcomeric protein essential for

contractility. Similarly, injection of a previously characterized *tnnt2a* morpholino[35] at the 1-cell stage

also led to compromised contraction and diminished expression of *id2b* (Figure 1F). Taken together,

these results indicate that biomechanical cues are essential for activating *id2b* in embryonic hearts.

Visualization of the spatiotemporal expression of *id2b* in developing embryos

Due to technical challenges in visualizing the cell-type-specific expression of *id2b* in the developing

heart using whole-mount in situ hybridization, we employed an intron targeting-mediated approach[36] to

generate a knockin *id2b:eGFP* reporter line. This method allowed us to achieve specific labeling

without perturbing the integrity and function of the endogenous gene[36] (Figure 2A). Comparison of

id2b:eGFP fluorescence with in situ hybridization at 24, 48, and 72 hpf revealed that the reporter signal

closely recapitulates the endogenous *id2b* expression pattern. The fluorescence was notably enriched in

the heart, brain, retina, and notochord (Figure 2B), mirroring observations from a previously reported

id2b transgenic line generated through BAC-mediated recombination[37].

To further elucidate the spatiotemporal expression of *id2b* in developing hearts, we crossed

id2b:eGFP with *myl7:mCherry* or *kdrl:mCherry*, labelling cardiomyocytes or endocardial cells,

respectively. Confocal images revealed minimal, if any, presence of *id2b:eGFP* in *myl7:mCherry*+

cardiomyocytes (Figure 2C). In sharp contrast, clear co-localization between *id2b:eGFP* and

kdrl:mCherry was evident at 48, 72, and 96 hpf (Figure 2D). Endocardial localization of *id2b* was

further confirmed by RNAscope analysis (Figure 2E). In adult hearts, *id2b:eGFP* fluorescence was

enriched in the chamber endocardium and the endothelium lining AVC, outflow tract (OFT), and

bulbus arteriosus (Figure 2F). Interestingly, there was an absence of *id2b:eGFP* signal in

kdrl:mCherry+ endothelial cells in trunk blood vessel and brain vasculature (Figure 2G). Collectively,

these results indicate that *id2b* is expressed in endocardial cells across different developmental stages.

BMP signaling and cardiac contraction regulate *id2b* expression

Taking advantage of live imaging on developing embryos, we explored the *in vivo* dynamics of *id2b* in

response to biomechanical force at single-cell resolution. When embryos were treated with tricaine or

blebbistatin, the intensity of *id2b:eGFP* in atrial and ventricular endocardium was significantly reduced

(Figure 3A, B). Similarly, injection of *tnnt2a* morpholino also markedly suppressed *id2b:eGFP* signal

(Figure 3A, B), in agreement with the results obtained from in situ hybridization. Strikingly, the

reduction in fluorescence intensity was particularly pronounced in AVC endothelial cells (Figure 3A, B,

asterisks).

We then explored how cardiac contraction modulated *id2b* expression. Given that endocardial

cells can sense blood flow through primary cilia[38, 39], we used a characterized morpholino[38] to

knockdown *ift88*, an intraflagellar transporter essential for primary cilia formation. Previously work

demonstrated a complete loss of primary cilia in endocardial cells upon *ift88* knockdown[38]. As

expected, a significant decrease in *id2b:eGFP* intensity was observed in the chamber and AVC

endocardium of *ift88* morphants compared to control (Figure 3C, D), suggesting that biomechanical

forces promote the expression of *id2b* via primary cilia. In the developing heart, a central hub for

mediating biomechanical cues is the Klf2 gene, which includes the *klf2a* and *klf2b* paralogues in

zebrafish[3, 4, 5, 38, 40]. Previous studies in mammals and zebrafish have highlighted the essential role of

Klf2 transcription factor activity in cardiac valve and myocardial wall formation[3, 40]. As a flow-

responsive gene, *klf2a* expression has been observed throughout the entire endocardium, evidenced by

mRNA expression and transgenic studies[3, 4]. Interestingly, in situ hybridization on 48 and 72 hpf *klf2a*[-/-]

and *klf2b*[-/-] embryos unveiled a drastic decrease in *id2b* expression compared with wild-type zebrafish

(Figure 3E), supporting the notion that *klf2*-mediated biomechanical signaling is essential for activating

id2b expression.

Given that *id2b* has been reported as a target gene of bone morphogenetic protein (BMP)

signaling, we explored whether BMP also played a role in regulating *id2b* expression. To this end, we

knocked down *bmp2b*, *bmp4*, and *bmp7a* in 1-cell stage embryos. Live imaging at 24 hpf revealed a

significant reduction in *id2b:eGFP* fluorescence signal in morpholino-injected hearts compared to

controls (Figure 3-figure supplement 1A, B), suggesting that *id2b* is a target gene of BMP signaling

during early embryonic development. Similarly, treatment with the BMP inhibitor Dorsomorphin from

10 to 24 hpf resulted in a marked decrease in *id2b:eGFP* signal (Figure 3-figure supplement 1C, D).

Considering that heartbeats in zebrafish commence at approximately 22 hpf, we treated embryos with

Dorsomorphin from 24 to 48 hpf or from 36 to 60 hpf. While the number of endocardial cells was

slightly reduced upon Dorsomorphin exposure as previously reported[7], surprisingly, quantification of

the average *id2b:eGFP* fluorescence intensity in individual endocardial cells revealed no significant

differences between Dorsomorphin and DMSO-treated controls (Figure 3-figure supplement 1C, D).

We further visualized BMP activity using the *BRE:d2GFP* reporter line. Confocal images

revealed strong fluorescence in the myocardium at 72 hpf, with minimal signal present in the

endocardium except for the AVC endothelium (Figure 3-figure supplement 1E). Moreover, after

tricaine treatment, endocardial *BRE:d2GFP* slightly increased (Figure 3-figure supplement 1E), as

opposed to the reduced *id2b:eGFP* signal (Figure 3A, B). Likewise, endocardial *BRE:d2GFP* intensity

was barely affected after completely blocking contraction with *tnnt2a* MO injection (Figure 3-figure

supplement 1E). These observations align with previous work using pSmad-1/5/8 as a readout of BMP

activity, indicating that endocardial BMP signaling is independent of blood flow[7]. Collectively, these

results suggest that *id2b* expression is regulated by both BMP and biomechanical signaling, with the

relative contribution of each pathway varying across developmental stages.

Compromised AV valve formation in *id2b* mutants

To investigate the role of the contractility-*id2b* axis in zebrafish heart development, we generated a

loss-of-function mutant line using CRISPR/Cas9. A pair of sgRNAs designed to target exon 1 was

injected with zCas9 protein into 1-cell-stage embryos. Consequently, we identified a mutant allele with

a 157 bp truncation, leading to the generation of a premature stop codon (Figure 4A, left). In *id2b*

mutants (*id2b$^{-/-}$*), the expression levels of *id2b* were significantly decreased, while *id2a* expression

levels were increased compared to *id2b$^{+/+}$* siblings (Figure 4A, right). The overall morphology of *id2b$^{-/-}$*

remained unaltered at 72 and 96 hpf (Figure 4B). However, *id2b$^{-/-}$* zebrafish experienced early lethality

starting around 31 weeks post-fertilization (Figure 4C). Strikingly, pericardial edema was observed in

20% (9/45) of adult *id2b$^{-/-}$* zebrafish (Figure 4D, top). Upon dissecting hearts from these *id2b$^{-/-}$*

zebrafish, a prominent enlargement in the atrium with a smaller ventricle was detected (Figure 4D,

bottom), which has been characterized as cardiomyopathy in zebrafish[41, 42]. Histological analysis

further revealed malformation in the AV valves of these *id2b$^{-/-}$* mutants compared to controls (Figure

4E, right). Specifically, we noted that the superior and inferior leaflets were significantly thinner,

comprising only 1-2 layer of cells in *id2b$^{-/-}$* zebrafish with an enlarged atrium. This was in sharp

contrast to *id2b$^{+/+}$* zebrafish, which exhibited multilayers of cells (Figure 4E, left). Subsequent

examination of the remaining 80% of *id2b$^{-/-}$* zebrafish (36/45) that did not display prominent pericardial

edema also revealed AV valve malformation, albeit to a lesser extent (Figure 4E, middle).

To further interrogate the effect of *id2b* inactivation on AV valve formation and function, we

analyzed the number of AVC endothelial cells using *kdrl:nucGFP*. At 96 hpf, a reduced number of

kdrl:nucGFP^{+} cells was detected in the AVC region of *id2b$^{-/-}$* embryos compared with *id2b$^{+/+}$* (Figure

4-figure supplement 1A, B). In contrast, the number of atrial and ventricular endocardial cells did not

differ between *id2b$^{-/-}$* and *id2b$^{+/+}$* (Figure 4-figure supplement 1C, D). Subsequently, we assessed

hemodynamic flow by conducting time-lapse imaging of red blood cells labelled by *gata1:dsred*.

Surprisingly, the pattern of hemodynamics was largely preserved in $id2b^{-/-}$ embryos compared to

$id2b^{+/+}$ siblings at 96 hpf (Figure 4-figure supplement 1E, Video 1, 2), suggesting that the reduced

number of endocardial cells in the AVC region was not sufficient to induce functional defects.

Additionally, we performed echocardiography to analyze blood flow in adult zebrafish as previously

described[43]. In $id2b^{-/-}$ hearts, prominent retrograde blood flow was detected in the AVC region (8/13)

(Figure 4G), while unidirectional blood flow was observed in $id2b^{+/+}$ (10/10) (Figure 4F).

Quantification analysis showed ~32% retrograde blood flow in $id2b^{-/-}$, compared to 0% in $id2b^{+/+}$

zebrafish (Figure 4H). Consistently, the superior and inferior leaflets were much thinner in $id2b^{-/-}$

exhibiting retrograde flow compared with control fish (Figure 4-figure supplement 1F). Overall, these

histological and functional analyses indicate that $id2b$ deletion leads to progressive defects in AV valve

morphology and hemodynamic flow.

***id2b* deletion perturbs calcium signaling and contractile function in the myocardium**

Although similar defects in AV valve formation have been reported in both *klf2a* and *nfatc1* mutants,

they do not display noticeable pericardial edema at the adult stage, nor do they experience early

lethality[3, 38, 40, 43, 44]. Therefore, we sought to investigate whether other cardiac properties have also been

affected by $id2b$ loss-of-function. To this end, we employed RNA-seq analysis on purified embryonic

$id2b^{-/-}$ and $id2b^{+/+}$ hearts (Figure 5-figure supplement 1). As expected, enrichment analysis of DEGs

demonstrated that the top-ranked anatomical structures affected by $id2b$ deletion included the heart

valve, the compact layer of ventricle, and the atrioventricular canal (Figure 5-figure supplement 1A).

Interestingly, $id2b$ inactivation also impacted phenotypes such as cardiac muscle contraction and heart

contraction (Figure 5-figure supplement 1B). Therefore, we investigated cardiac contractile function

through time-lapse imaging on the *myl7:mCherry* background. At 72 and 120 hpf, a significant

decrease in cardiac function was observed in $id2b^{-/-}$ compared with $id2b^{+/+}$ (Figure 5A-C, Figure 5-

figure supplement 2A-C). Similarly, echocardiography analysis showed that the contractile function in

adult $id2b^{-/-}$ heart was dramatically reduced compared with age-matched $id2b^{+/+}$ (Figure 5D, F). These

functional defects in $id2b$-deleted hearts could not be attributed to differences in cardiomyocyte

number, as we counted cardiomyocytes using the *myl7:H2A-mCherry* line and found no apparent

changes between *id2b*[-/-] and *id2b*[+/+] embryos at 72 and 120 hpf (Figure 5-figure supplement 2D, E).

Similarly, *id2b*[-/-] also developed regular trabecular structures (Figure 5-figure supplement 2F). Through

α-actinin immunostaining, we observed similar sarcomeric structures in *id2b*[-/-] and control

cardiomyocytes at 72 hpf and adult stages (115 dpf) (Figure 5-figure supplement 2G), corroborating

that the reduced contractility in *id2b*-depleted heart was independent of structural defects.

The key functional unit that transmits electrical activity to contractile function is E-C coupling.

Because *id2b*[-/-] displayed reduced cardiac function, we visualized calcium signaling in the developing

heart using *actb2:GCaMP6s* zebrafish (Figure 5G). Compared to *id2b*[+/+] controls, *id2b*[-/-] embryos

exhibited markedly decreased calcium transient amplitude (Figure 5H), consistent with compromised

calcium handling observed in other zebrafish cardiomyopathy models[42, 45]. In cardiomyocyte, the entry

of extracellular calcium is mainly mediated through the LTCC. As previously reported, a defect in

zebrafish LTCC pore forming α1 subunit *cacna1c* leads to compromised cardiac function[27]. We

collected hearts from 72 hpf and 5 months post-fertilization (mpf) zebrafish and detected

downregulated *cacna1c* in *id2b*[-/-] compared to *id2b*[+/+] (Figure 5I). In addition, we measured cardiac

action potential using intracellular recording[46]. Compared to *id2b*[+/+] zebrafish, the duration of the

action potential in *id2b*[-/-] was significantly shorter (Figure 5J, K), consistent with the decreased

expression level of *cacna1c*. Together, these data indicate that *id2b* loss-of-function leads to

compromised calcium signaling and cardiac contractile function.

Reduced expression of *nrg1* mediates the compromised contractility in *id2b*[-/-]

Because the deficiency of *id2b* in the endocardium disrupted the function of myocardium, we

speculated that the crosstalk between these two types of cells was affected in *id2b*[-/-]. Interestingly,

comparing the differentially expressed genes in embryonic *id2b*[-/-] and *id2b*[+/+] hearts identified a

significant reduction in the expression level of Nrg1, a key mitogen regulating the intra-organ

communications between endocardial cells and cardiomyocytes (Figure 6A). Remarkably, analysis of a

zebrafish single-cell database[47] revealed enriched expression of *nrg1* in endocardial cells (Figure 6-

figure supplement 1). However, attempts to detect *nrg1* expression through in situ hybridization were

unsuccessful, likely due to its low abundance in the heart. Alternatively, qRT-PCR analysis of purified

120 hpf embryonic hearts validated decreased *nrg1* levels in *id2b^{-/-}* compared to control (Figure 6B).

Previous studies have demonstrated that perturbations in Nrg-Erbb2 signaling, as seen in zebrafish

erbb2 mutants, result in dysfunctional cardiac contractility[20]. Consistently, a decrease in heart rate was

observed in embryos treated with the *erbb2* inhibitor AG1478 (Figure 6C).

Remarkably, injecting *nrg1* mRNA at the 1-cell stage not only rescued the reduced expression of

cacna1c in *id2b^{-/-}* hearts (Figure 6D) but also restored the diminished heart rate (Figure 6E). This is

consistent with prior study showing that Nrg1 administration can restore LTCC expression and calcium

current in failing mammalian cardiomyocytes[48]. Overall, our data suggest that endocardial *id2b*

promotes Nrg1 synthesis, thereby enhancing cardiomyocyte contractile function.

Id2b interacts with Tcf3b to limit its repressor activity on *nrg1* expression

We further interrogated how *id2b* promotes the expression of *nrg1*. As a HLH factor lacking a DNA

binding motif, Id2b has been reported to form a heterodimer with Tcf3b to limits its function as a

potent transcriptional repressor[49]. Notably, we detected expression of *tcf3b* in endocardial cells by

analyzing a zebrafish single-cell database[47] (Figure 7-figure supplement 1). To determine if zebrafish

Id2b and Tcf3b interact *in vitro*, Flag-*id2b* and HA-*tcf3b* were co-expressed in HEK293 cells. Co-

immunoprecipitation analysis confirmed their interaction (Figure 7A), although whether they interact *in*

vivo remains to be further investigated. Subsequently, qRT-PCR analysis on purified 120 hpf

embryonic hearts revealed a significant increase in the expression of *socs3b* and *socs1a*, target genes of

tcf3b, in *id2b^{-/-}* compared to *id2b^{+/+}* (Figure 7B). This suggests an elevation in *tcf3b* activity

with the loss of *id2b* function. Notably, the expression levels of *tcf3a* and *tcf3b* remained consistent

between *id2b^{-/-}* and *id2b^{+/+}* hearts (Figure 7B).

To understand how the altered interaction between *id2b* and *tcf3b* influences *nrg1* expression, we

analyzed the promoter region of zebrafish *nrg1* using JASPAR and identified two potential *tcf3b*

binding sites (Figure 7C). Subsequently, a DNA fragment containing the zebrafish *nrg1* promoter

region was subcloned into a vector carrying the luciferase reporter gene. Co-injection of this construct

with *tcf3b* mRNA into 1-cell stage embryos resulted in a significant decrease in luciferase signal.

Conversely, co-injection with a previously characterized *tcf3b* morpholino led to enhanced luciferase

intensity (Figure 7D). These results suggest a possible mechanism by which Tcf3b represses *nrg1*

expression in zebrafish.

Lastly, injecting *tcf3b* morpholino into *id2b*$^{-/-}$ embryos was performed to assess whether

attenuating the overactive *tcf3b* in *id2b*$^{-/-}$ could restore the expression level of *nrg1*. qRT-PCR analysis

of purified 72 hpf hearts revealed a partial restoration of the diminished *nrg1* expression in *id2b*$^{-/-}$ upon

tcf3b inhibition (Figure 7E). Taken together, our results indicate that biomechanical cues activate

endocardial Id2b expression, leading to its interaction with Tcf3b to alleviate repression on the

nrg1 promoter. Consequently, the depletion of *id2b* unleashes Tcf3b's repressor activity, leading to a

reduction in *nrg1* expression, which further acts through *erbb2* to regulate cardiomyocyte function

(Figure 7F).

Discussion

Biomechanical forces play an essential role in regulating the patterning and function of the heart. At

AVC, oscillatory flow promotes the expression of *klf2a* and *nfatc1* to modulate valve morphogenesis.

In chamber endocardium, blood flow induces endocardial cells to acquire distinct cell morphology.

However, it still lacks a systematic analysis of the transcriptome underlying compromised heartbeats.

In the present study, we analyzed embryonic zebrafish hearts without contractility and identified genes

that are regulated by biomechanical forces. Specifically, our results unveiled the endocardial-specific

expression of *id2b*, which was tightly regulated by flow-sensitive primary cilia-*klf2* axis. Genetic

deletion of *id2b* resulted in compromised valve formation and progressive atrium enlargement. In

addition, a reduction in heart rate and contractile force was observed in *id2b*$^{-/-}$, owing to decreased

expression of LTCC α1 subunit *cacna1c*. Mechanistically, *id2b* interacts with bHLH transcription

factor *tcf3*b to limit its repressor activity. Hence genetic deletion of *id2b* unleashes *tcf3*b activity, which

further represses endocardial *nrg1* expression. As a result, Injection of *nrg1* mRNA partially rescues

the phenotype of *id2b* deletion. Overall, our findings identify *id2b* as a novel mediator that regulate the

interplay between endocardium and myocardium during heart development.

In mammals, the deletion of *Id2* leads to malformations in the arterial and venous poles of the

heart, as well as affects AV valve morphogenesis[31, 33]. Interestingly, approximately 20% perinatal

lethality is reported in *Id2* knockout mice, exhibiting AV septal defects and membranous ventricular

septal defects[33]. Remarkably, pericardial edema is evident in 20% of adult *id2b*[-/-] zebrafish, with a

prominent enlargement of the atrium. The superior and inferior leaflets of AV valves in *id2b*[-/-] mutants

are significantly thinner compared to the control. Therefore, our results suggest that *id2b* may play a

similar role in regulating AV valve formation in zebrafish as its mammalian orthologue *Id2*. It is

proposed that the loss of *Id2* in mice results in compromised endocardial proliferation and aberrant

endothelial-to mesenchymal transformation (EMT), collectively leading to defective valve

morphogenesis[33]. Nevertheless, the mechanism by which *id2b* loss-of-function causes a reduction in

leaflet thickness in zebrafish remains to be determined in future studies.

id2b has been recognized as a target gene of the BMP signaling pathway. As expected,

knockdown of *bmp2b*, *bmp4*, and *bmp7a* at 1-cell stage confirms that endocardial *id2b* expression is

controlled by BMP activity during early embryonic development. Surprisingly, treatment with the BMP

inhibitor Dorsomorphin at 24 and 36 hpf, when cardiac contractions have already initiated, fails to alter

id2b expression in the endocardium, suggesting that BMP is dispensable for *id2b* activation at these

stages. Instead, endocardial *id2b* expression is reduced upon loss-of-function of *klf2a*, *klf2b*, and *ift88*,

suggesting an essential role of the primary cilia-*klf2* axis in mediating *id2b* activation. In endocardial

cells, Trp, Piezo, and ATP-dependent P2X/P2Y channels[4, 6, 50, 51] are well-established sensors for

biomechanical stimulation. The activation of these channels further promotes the activities of Klf2 and

Nfatc1 to drive heart development and valvulogenesis. However, whether these channels are also

required for the activation of *id2b* warrants further investigation.

The Nrg-Erbb signaling plays an essential role in regulating heart morphogenesis and function. In

the mammalian heart, the genetic deletion of Nrg1 or Erbb2 results in severely perturbed cardiac

trabeculae formation[13, 14, 15]. Zebrafish *erbb2* mutants exhibit a similar defect in cardiomyocyte

proliferation and trabeculation[20]. Interestingly, *nrg1* mutant zebrafish display grossly normal cardiac

structure during early embryonic devolpment[21, 52]. Nevertheless, zebrafish *nrg2a* loss-of-function leads

to defective trabeculae formation, suggesting that *nrg2a* is the predominant ligand secreted from

endocardium, promoting ventricular morphogenesis[21]. In the adult stage, perivascular cells[53] or

regulatory T cells[54]-derived *nrg1* promotes cardiomyocyte proliferation during heart regeneration.

Hence, the specific ligand/receptor and the spatiotemporal regulation of the Nrg-Erbb axis appear to be

more complicated in both embryonic and adult zebrafish. Interestingly, the *nrg1* mutant heart exhibits a

defect in cardiac nerve expansion and heart maturation at the juvenile stage despite normal cardiac

structure formation[52], suggesting its potential role in regulating cardiac function. Our findings

demonstrate that the expression of *nrg1* in embryonic endocardial cells is influenced by biomechanical

cues and *id2b* activity. This signaling axis is essential for coordinating endocardium-myocardium

interaction and establishing proper cardiac function.

Materials and methods

Zebrafish handling and lines

All animal procedures were approved by the Animal Care and Use Committee of the Zhejiang

University School of Medicine (application no.29296). Embryonic and adult fish were raised and

maintained under standard conditions at 28 ℃ on a 14/10 hour day/night cycle. The following

zebrafish lines were used in this study: *Tg(myl7:mCherry)*[sd7] [55], *Tg(myl7:H2A-mCherry)*[sd12] [56],

Tg(kdrl:mCherry)[S896] [57], *Tg(kdrl:nucGFP)*[y7] [58], *Tg(BRE:d2GFP)*[mw30] [59], and *Tg(actb2:Gcamp6s)*. To

generate the *id2b* mutant, two short guide RNAs (sgRNAs) targeting exon 1 were generated using the

MAXIscript T7 transcription kit (ambion, AM1314). The sgRNAs were as follows: sgRNA1: 5' -

GAAGGCAGTCAGTCCGGTG - 3'; sgRNA2: 5' - GAACCGGAGCGTGAGTAAGA - 3'. The two

sgRNAs, along with zCas9 protein, were co-injected into one-cell stage embryos. Embryos were raised

to adulthood and crossed to wild-type zebrafish to obtain F_1 progenies. Through PCR analysis, a

mutant line with a 157 bp truncation was identified.

The knock-in *id2b:eGFP* line was generated using a previously reported method[36]. Briefly, three

sgRNAs were designed to target the intron of *id2b* (sgRNA1: 5' - GAGACAAATATCTACTAGTG - 3';

sgRNA2: 5' - GTTGAACACATGACGATATT - 3'; sgRNA3: 5' - GCACAACTTAGATTTCAAGT -

3'). Co-injection of each individual sgRNA with zCas9 protein into one-cell stage zebrafish embryos

yielded varying cleavage efficiency. Since sgRNA2 displayed the highest gene editing efficiency, it

was selected for subsequent studies. Next, a donor plasmid containing the sgRNA targeting sequence of

the intron, exon 2 of *id2b*, and P2A-eGFP was generated. Co-injection of sgRNA, donor plasmid, and

zCas9 protein into one-cell stage embryos led to concurrent cleavage of the sgRNA targeting sites in

both the zebrafish genome and the donor plasmid (Figure 2A). Accordingly, eGFP fluorescence was

observed in injected 24 hpf zebrafish embryos, indicating the incorporation of the donor. The insertion

of the *id2b*-p2A-eGFP donor into the genome was confirmed by PCR analysis with primers

recognizing target site or donor sequences, respectively (Figure 2A). Embryos with mosaic eGFP

expression were raised to adulthood and crossed with wild-type zebrafish to obtain F_1 progenies.

Overall, two founders were identified. The junction region of the F_1 embryos was sequenced to

determine the integration sites. Although the two founders had slightly different integration sites in the

intron, the expression pattern and fluorescence intensity of eGFP were indistinguishable between the

two lines.

Morpholinos

All morpholinos (Gene Tools) used in this study have been previously characterized. *tnnt2a* MO (5' -

CATGTTTGCTCTGATCTGACACGCA - 3')[35]; *ift88* MO (5' -

CTGGGACAAGATGCACATTCTCCAT - 3')[38]; *bmp2b* MO (5' -

ACCACGGCGACCATGATCAGTCAGT - 3')[60]; *bmp4* MO (5' -

AACAGTCCATGTTTGTCGAGAGGTG - 3')[61]; *bmp7a* MO (5' -

GCACTGGAAACATTTTTAGAGTCAT - 3')[60]; *tcf3b* MO (5' -

CGCCTCCGTTAAGCTGCGGCATGTT - 3')[62]. For each morpholino, a 1 nL solution was injected

into one-cell stage embryos at the specified concentrations: 0.5 µg/µL *tnnt2a* MO, 2 µg/µL *ift88* MO,

0.5 µg/µL *bmp2b* MO, 2 µg/µL *bmp4* MO, 4 µg/µL *bmp7a* MO, and 1 µg/µL *tcf3b* MO.

Small molecules treatment

To inhibit cardiac contraction, embryos were incubated in 1 mg/mL tricaine (Sigma, A5040) or 10 µM

blebbistatin (MedChemExpress, HY13441) PTU-added egg water for 12-24 hours. In order to inhibit

erbb2 signaling pathway, 10 µM AG1478 (Sigma, 658552) were used to treat 4 dpf larvae. To inhibit

BMP signaling pathway, 10 µM Dorsomorphin (Sigma, P5499) was used to treat 10, 24, and 36 hpf

embryos.

***In situ* hybridization and RNAscope**

Whole mount *in situ* hybridization was performed as previously described[46]. The probes were

synthesized using the DIG RNA labeling kit (Roche). The primers used for obtaining the *id2b* probe

template were as follows: Forward 5' - ATGAAGGCAGTCAGTCCGGTGAGGT - 3'; Reverse 5' -

TCAACGAGACAGGGCTATGAGGTCA - 3'. RNAscope analysis was performed using the probe Dr-

id2b (Advanced Cell Diagnostics, 517541) and the Multiplex Fluorescent Detection Kit version 2

(Advanced Cell Diagnostics, 323100) as previously described[19].

Embryonic heart isolation and RNA-seq analysis

Hearts were isolated from embryos carrying the *Tg*(*myl7:mCherry)* transgene following an established

protocol[34]. A minimum of 1,000 hearts for each experimental group were manually collected under a

Leica M165FC fluorescence stereomicroscope and transferred into ice-cold PBS buffer. After

centrifugation at 12, 000 g for 2 min at 4°C, the supernatant was removed, and hearts were lysed in

cold Trizol buffer (Ambion, 15596). Total RNA was extracted for subsequent quantitative real-time

PCR or RNA-seq analysis.

Duplicate samples from control and Tricaine-treated embryonic hearts underwent RNA-seq. Raw

sequencing reads were preprocessed to remove adapters and filter low-quality reads. Clean sequencing

reads were then mapped to the zebrafish reference[63] using STAR with default parameters[64].

Subsequently, gene quantification was carried out with RSEM[65]. The gene expected count was applied

to identify differential expression genes (DEGs), retaining only genes with counts per million (CPM) of

10 in at least two samples. DESeq2[66] was employed for differential expression analysis, and *P*-values

were adjusted using BH correction. DEGs were defined as those with $|\log_2 \text{fold change}| \geq 0.585$ and an

adjusted *P*-value < 0.1. The primary focus was on genes related to transcription regulation, and gene

enrichment analysis was conducted using ClusterProfiler[67]. To analyze DEGs in *id2b*[-/-] and control

embryonic hearts, we performed enrichment analysis with the R package EnrichR, dissecting the

potential anatomy expression pattern and underlying phenotypes. We mainly focused on genes with the

heart related phenotypes, including cardiac muscle tissue development, cardiomyocyte differentiation,

heart morphogenesis and cardiac chamber development. All the analysis on identifying DEGs were

batch corrected.

Quantitative real-time PCR analysis

After extraction from isolated embryonic hearts, 50 ng-1 ug of mRNA was reverse transcribed to

cDNA using the PrimeScript RT Master Mix kit (Takara, RR036A). Real-time PCR was performed

using the TB Green Premix Ex Taq kit (Takara, RR420A) on the Roche LightCycler 480. Expression

levels of the target genes were normalized to *actb1* as an internal control. All experiments were

repeated three times. The following primer sets were used: *id2b* Forward 5' -

ACCTTCAGATCGCACTGGAC - 3', Reverse 5' - CTCCACGACCGAAACACCATT - 3'; *nrg1*

Forward 5' - CTGCATCATGGCTGAGGTGA - 3', Reverse 5' - TTAACTTCGGTTCCGCTTGC - 3';

cacnα1c Forward 5' - GCCCTTATTGTAGTGGGTAGTG - 3', Reverse 5' -

AGTGTTTTGGAGGCCCATTG - 3'; *tcf3a* Forward 5' - CCTCCGGTCATGAGCAACTT - 3',

Reverse 5' - TTTCCCATGATGCCTTCGCT - 3'; *tcf3b* Forward 5' - CCTTTAATGCGCCGTGCTTC -

3', Reverse 5' - GCGTTCTTCCATTCCTGTACCA - 3'; *socs1a* Forward 5' -

TCAGCCTGACAGGAAGCAAG - 3', Reverse 5' - GTTGCACAGGGATGCAGTCG - 3'; *socs3b*

Forward 5' - GGGACAGTGAGTTCCTCCAA - 3', Reverse 5' - ATGGGAGCATCGTACTCCTG - 3';

actb1 Forward 5' - ACCACGGCCGAAAGAGAAAT - 3', Reverse 5' -

GCAAGATTCCATACCCAGGA - 3'.

Co-immunoprecipitation (Co-IP) and western blot

Zebrafish *tcf3b* and *id2b* were overexpressed in 293T cell for 48 hours. The transfected cells were then

collected and lysed using IP lysis buffer (Sangon Biotech, C500035) containing protease and

phosphatase inhibitors (Sangon Biotech, C510009, C500017). For the IP experiment, anti-Flag

antibody (Cell Signaling Technology, 14793, 1:100) and IgG antibody (ABclonal, AC005, 1:100) was

incubated with cell lysates overnight at 4°C. Pretreated magnetic beads were bound with the antigen-

antibody complex for 4 hours at 4°C, followed by washing with IP lysis buffer three times. For western

blot, samples were denatured at 95°C for 10 min, separated on a 5%-12% gradient gel. Proteins were

then transferred to a PVDF membrane (Sigma, ISEQ00010). The membrane was blocked for 1 hour

with 5% nonfat milk or 5% BSA (Sangon Biotech) dissolved in TBST and then incubated with primary

antibodies (anti-FLAG, Cell Signaling Technology, 14793, 1:1000; anti-HA, Sigma, H3663, 1:1000)

overnight at 4°C, followed by three times 10-min TBST washes. HRP-conjugated secondary antibodies

(Invitrogen, 31430, 31460) were incubated for 1 hour at room temperature, followed by three times 10-

min TBST washes. The detection of immunoreactive bands was performed with a chemiluminescent

substrate (Thermo Scientific, 34577) and imaged using the Azure Biosystems 400.

Immunofluorescence

For immunofluorescence on adult zebrafish hearts, we fixed the hearts overnight in 4%

paraformaldehyde at 4 □, followed by equilibration through 15% and 30% sucrose in PBS solution.

The hearts were embedded and frozen in O.C.T. compound (Epredia, 6502), and 10 μm thick

cryosections were prepared using a CryoStar NX50 cryostat. Immunofluorescence experiments were

performed as previously described[68]. For immunofluorescence on embryonic hearts, embryos were

fixed overnight in 4% paraformaldehyde at 4□, washed twice quickly in 100% methanol, and then

dehydrated overnight at -20□ in 100% methanol. Subsequently, rehydration was performed through a

methanol gradient (100%, 75%, 50%, 25%, 10 min each), followed by three times washes in PBST (1%

PBS/0.1% Triton-X100, 10 min each). The embryos were treated with 10 μg/mL proteinase K diluted

in PBST for 20 min at room temperature, re-fixed in 4% paraformaldehyde for 20 min, washed in

PBST, and immersed in blocking solution (PBST/1% BSA/2% goat serum) for 1 hour at room

temperature. Following this, the embryos were incubated in the primary antibody diluted in blocking

solution overnight at 4□. After washing in PBST, they were incubated in the secondary antibody

(1:200) for 2 hours at room temperature. The primary antibody used was: Anti-GFP antibody (Santa

Cruz Biotechnology, sc9996, 1:200), anti-α-actinin antibody (Sigma, A7811, 1:200). The secondary

antibody used was: Anti-mouse IgG-Alexa 488 (Invitrogen, A11011, 1:400). DAPI was used to stain

cell nuclei.

Cardiac function analysis

To assess cardiac function in embryonic hearts, embryos were incubated in 0.16 mg/mL tricaine

(Sigma, A5040) and embedded in 1% low melting agarose. Heart contractions were recorded for 1 min

using a Nikon Ti2 microscope at a rate of 25 frames per second. Fractional shortening and heart rate

were measured as described previously[46]. For cardiac contractile functions in adulthood, zebrafish were

fixed on a sponge soaked with system water with the belly facing up and echocardiography was

performed[69]. Videos and images in color Doppler mode and B-mode were obtained using the

Vevo1100 imaging system at a frequency of 50 MHz. Nikon NIS elements AR analysis and Image J

software were employed for data extraction. To evaluate AV valve function, the ratio of inflow and

outflow area in the same frame was quantified[43].

Calcium imaging

At 120 hpf, embryos were treated with 10 mM 2,3-butanedione monoxime (BDM, Sigma, B0753) and

mounted in 1% low melting agarose. Time-lapse images were acquired using a Nikon Ti2 microscope

at a rate of 50 frames per second. Data were analyzed using Nikon NIS elements AR analysis software.

Intracellular action potential recording

Electrophysiology study was performed on adult zebrafish ventricles as previously described[46]. Briefly,

hearts were mounted in a chamber containing Tyrode's solution: NaCl 150 mM, KCl 5.4 mM, MgSO$_4$

1.5 mM, NaH$_2$PO$_4$ 0.4 mM, CaCl$_2$ 2 mM, Glucose 10 mM, HEPES 10 mM, pH was adjusted to 7.4.

Glass pipettes with tip resistance 30-40 MΩ were filled with 3M KCl solution. Intracellular action

potentials were recorded using an HEKA amplifier and pClamp10.3 software (Molecular Devices).

Histology and HE staining

Adult hearts were dissected and fixed overnight at 4℃ in 4% PFA, followed by three times PBS

washes. Dehydration involved an ethanol gradient (70%, 80%, 95%, 100%, 100%, 30 min each),

followed by three soaks in dimethylbenzene at 65℃, before embedding in paraffin. Sections of 5 µm

thickness were prepared using the Leica RM2235 manual rotary microtome for hematoxylin and eosin

(HE) staining.

Injection of mRNA

533 The embryonic zebrafish cDNA library was used as a template to amplify the *nrg1* and *tcf3b* fragment,

534 which was then subcloned into the pCS2 vector. The vector was linearized using Not I restriction

535 endonuclease, and mRNA was transcribed in vitro using the mRNA transcription kit (Ambion,

536 AM1340). 100 pg of purified mRNA was injected into one-cell stage embryos.

Luciferase assay

The LCR (luciferase reporter) plasmid was generated by subcloning the 5′ UTR of *nrg1* into the

upstream region of renilla luciferase on the psiCheck2 plasmid. Following construction, 25 pg of the

LCR plasmid was co-injected with either 100 pg of *tcf3b* mRNA or 1 ng of *tcf3b* MO into 1-cell stage

zebrafish embryos. At 48 hpf, twenty embryos were gathered into one group and fully lysed.

Subsequently, firefly and renilla luciferase activities were sequentially measured using a microplate

reader with the dual luciferase reporter gene assay kit (Yeasen, 11402ES60), according to the

manufacturer′s instructions. The experiment was independently replicated three times. The relative

renilla luciferase activity, normalized by firefly luciferase activity, served as an indicator of *nrg1*

expression level under the influence of *tcf3b* overexpression or reduction.

Image processing and statistical analysis

Whole mount *in situ* hybridization images were captured with a Leica M165FC stereomicroscope. Live

imaging of zebrafish embryos involved mounting anesthetized embryos in 1% low melting agarose

(Sangon Biotech, A600015) and manually oriented them for optimal visual access to the heart.

Confocal images were obtained with a Nikon Ti2 confocal microscope. Fluorescence intensity and cell

number counting were processed using Nikon NIS Elements AR analysis and Image J software.

Statistical analysis was performed using Graphpad prism 8 software. No statistical methods were used

to predetermine sample size. Unpaired two-tailed Student's t-tests was used to determine statistical

significance. Data are presented as mean ± s.e.m, $^{*}P < 0.05$ was considered to be statistically significant.

Acknowledgements: We thank Dr. Pengfei Xu for providing morpholinos. We also thank Dr. Jia Li for

the support in generating the *id2b:eGFP* line.

Author Contributions: Shuo Chen and Peidong Han contributed to the study design. Shuo Chen,

Jinxiu Liang, Weijia Zhang, Peijun Jiang, Wenyuan Wang, Xiaoying Chen and Yuanhong Zhou

conducted experiments. Jie Yin analyzed RNA-sequencing data. Peng Xia, Fan Yang, Ying Gu, and

Ruilin Zhang provided key reagents and analytic tool. Shuo Chen, Jinxiu Liang, Jie Yin, and Peidong

Han prepared the manuscript.

Source of Funding: This work was supported by the National Natural Science Foundation of China

(32170823, 92468104, 31871462), and the National Key R&D Program of China (2023YFA1800600).

Disclosures: The authors declare that they have no conflict of interest.

Data availability: The authors declare that all data supporting the findings in the paper are available in

the article and the supplementary files. RNA-seq data have been deposited in GEO under accession

number GSE295737 and GSE295738.

References

1.   Duchemin AL, Vignes H, Vermot J, Chow R. Mechanotransduction in cardiovascular
579   morphogenesis and tissue engineering. *Curr Opin Genet Dev* **57**, 106-116 (2019).

581   2.   Sidhwani P, Yelon D. Fluid forces shape the embryonic heart: Insights from zebrafish.
582   *Curr Top Dev Biol* **132**, 395-416 (2019).

584   3.   Vermot J, *et al.* Reversing blood flows act through klf2a to ensure normal
585   valvulogenesis in the developing heart. *PLoS Biol* **7**, e1000246 (2009).

587   4.   Heckel E, *et al.* Oscillatory Flow Modulates Mechanosensitive klf2a Expression
588   through trpv4 and trpp2 during Heart Valve Development. *Curr Biol* **25**, 1354-1361
589   (2015).

5.   Galvez-Santisteban M, *et al.* Hemodynamic-mediated endocardial signaling controls
in vivo myocardial reprogramming. *Elife* **8**, (2019).

6.   Fukui H, *et al.* Bioelectric signaling and the control of cardiac cell identity in response
to mechanical forces. *Science* **374**, 351-354 (2021).

7.   Dietrich AC, Lombardo VA, Veerkamp J, Priller F, Abdelilah-Seyfried S. Blood flow and
Bmp signaling control endocardial chamber morphogenesis. *Dev Cell* **30**, 367-377
(2014).

8.   Sidhwani P, *et al.* Cardiac function modulates endocardial cell dynamics to shape the
cardiac outflow tract. *Development* **147**, (2020).

9.   Staudt DW, Liu J, Thorn KS, Stuurman N, Liebling M, Stainier DY. High-resolution
imaging of cardiomyocyte behavior reveals two distinct steps in ventricular
trabeculation. *Development* **141**, 585-593 (2014).

10.   Peshkovsky C, Totong R, Yelon D. Dependence of cardiac trabeculation on neuregulin
signaling and blood flow in zebrafish. *Dev Dyn* **240**, 446-456 (2011).

11.   Auman HJ, Coleman H, Riley HE, Olale F, Tsai HJ, Yelon D. Functional modulation of
cardiac form through regionally confined cell shape changes. *PLoS Biol* **5**, e53 (2007).

12.   Lin YF, Swinburne I, Yelon D. Multiple influences of blood flow on cardiomyocyte
hypertrophy in the embryonic zebrafish heart. *Dev Biol* **362**, 242-253 (2012).

13.   Gassmann M, *et al.* Aberrant neural and cardiac development in mice lacking the
ErbB4 neuregulin receptor. *Nature* **378**, 390-394 (1995).

14.   Lee KF, Simon H, Chen H, Bates B, Hung MC, Hauser C. Requirement for neuregulin
receptor erbB2 in neural and cardiac development. *Nature* **378**, 394-398 (1995).

15.   Meyer D, Birchmeier C. Multiple essential functions of neuregulin in development.
Nature **378**, 386-390 (1995).

16.   Zhang W, Liang J, Han P. Cardiac cell type-specific responses to injury and
627   contributions to heart regeneration. *Cell Regen* **10**, 4 (2021).

628
629   17.   Lai D*, et al.* Neuregulin 1 sustains the gene regulatory network in both trabecular
630   and nontrabecular myocardium. *Circ Res* **107**, 715-727 (2010).

631
632   18.   Rentschler S*, et al.* Neuregulin-1 promotes formation of the murine cardiac
633   conduction system. *Proc Natl Acad Sci U S A* **99**, 10464-10469 (2002).

634
635   19.   Liang J*, et al.* Genetically encoded tension heterogeneity sculpts cardiac
636   trabeculation. *Sci Adv* **11**, eads2998 (2025).

637
638   20.   Liu J*, et al.* A dual role for ErbB2 signaling in cardiac trabeculation. *Development* **137**,
639   3867-3875 (2010).

640
641   21.   Rasouli SJ, Stainier DYR. Regulation of cardiomyocyte behavior in zebrafish
642   trabeculation by Neuregulin 2a signaling. *Nat Commun* **8**, 15281 (2017).

643
644   22.   Han P*, et al.* Coordinating cardiomyocyte interactions to direct ventricular chamber
645   morphogenesis. *Nature* **534**, 700-704 (2016).

646
647   23.   Guo Y, Pu WT. Cardiomyocyte Maturation: New Phase in Development. *Circ Res* **126**,
648   1086-1106 (2020).

649
650   24.   Alvarez-Dominguez JR, Melton DA. Cell maturation: Hallmarks, triggers, and
651   manipulation. *Cell* **185**, 235-249 (2022).

652
653   25.   Bers DM. Cardiac excitation-contraction coupling. *Nature* **415**, 198-205 (2002).

654
655   26.   Xu C*, et al.* Ptpn23 Controls Cardiac T-Tubule Patterning by Promoting the Assembly
656   of Dystrophin-Glycoprotein Complex. *Circulation* **149**, 1375-1390 (2024).

657
658   27.   Rottbauer W, Baker K, Wo ZG, Mohideen MA, Cantiello HF, Fishman MC. Growth and
659   function of the embryonic heart depend upon the cardiac-specific L-type calcium
660   channel alpha1 subunit. *Dev Cell* **1**, 265-275 (2001).

662   28.   Samsa LA, Givens C, Tzima E, Stainier DY, Qian L, Liu J. Cardiac contraction activates
663   endocardial Notch signaling to modulate chamber maturation in zebrafish.
664   *Development* **142**, 4080-4091 (2015).

665
666   29.   Benezra R, Davis RL, Lockshon D, Turner DL, Weintraub H. The protein Id: a negative
667   regulator of helix-loop-helix DNA binding proteins. *Cell* **61**, 49-59 (1990).

668
669   30.   Wong MV, Jiang S, Palasingam P, Kolatkar PR. A divalent ion is crucial in the structure
670   and dominant-negative function of ID proteins, a class of helix-loop-helix
671   transcription regulators. *PLoS One* **7**, e48591 (2012).

672
673   31.   Jongbloed MR, *et al.* Expression of Id2 in the second heart field and cardiac defects
674   in Id2 knock-out mice. *Dev Dyn* **240**, 2561-2577 (2011).

675
676   32.   Moskowitz IP, *et al.* A molecular pathway including Id2, Tbx5, and Nkx2-5 required
677   for cardiac conduction system development. *Cell* **129**, 1365-1376 (2007).

678
679   33.   Moskowitz IP, *et al.* Transcription factor genes Smad4 and Gata4 cooperatively
680   regulate cardiac valve development. [corrected]. *Proc Natl Acad Sci U S A* **108**, 4006-
681   4011 (2011).

682
683   34.   Burns CG, MacRae CA. Purification of hearts from zebrafish embryos. *Biotechniques*
684   **40**, 274, 276, 278 passim (2006).

685
686   35.   Sehnert AJ, Huq A, Weinstein BM, Walker C, Fishman M, Stainier DY. Cardiac
687   troponin T is essential in sarcomere assembly and cardiac contractility. *Nat Genet* **31**,
688   106-110 (2002).

689
690   36.   Li J, Zhang BB, Ren YG, Gu SY, Xiang YH, Du JL. Intron targeting-mediated and
691   endogenous gene integrity-maintaining knockin in zebrafish using the CRISPR/Cas9
692   system. *Cell Res* **25**, 634-637 (2015).

693
694   37.   Forster D, *et al.* Genetic targeting and anatomical registration of neuronal
695   populations in the zebrafish brain with a new set of BAC transgenic tools. *Sci Rep* **7**,
696   5230 (2017).

697
698   38.   Li X, *et al.* Primary cilia mediate Klf2-dependant Notch activation in regenerating
699   heart. *Protein Cell* **11**, 433-445 (2020).

39.   Nauli SM, Kawanabe Y, Kaminski JJ, Pearce WJ, Ingber DE, Zhou J. Endothelial cilia
are fluid shear sensors that regulate calcium signaling and nitric oxide production
through polycystin-1. *Circulation* **117**, 1161-1171 (2008).

40.   Rasouli SJ*, et al.* The flow responsive transcription factor Klf2 is required for
myocardial wall integrity by modulating Fgf signaling. *Elife* **7**, (2018).

41.   Weeks O*, et al.* Embryonic alcohol exposure in zebrafish predisposes adults to
cardiomyopathy and diastolic dysfunction. *Cardiovasc Res* **120**, 1607-1621 (2024).

42.   Kamel SM, Koopman CD, Kruse F, Willekers S, Chocron S, Bakkers J. A Heterozygous
Mutation in Cardiac Troponin T Promotes Ca(2+) Dysregulation and Adult
Cardiomyopathy in Zebrafish. *J Cardiovasc Dev Dis* **8**, (2021).

43.   Gunawan F, Gentile A, Gauvrit S, Stainier DYR, Bensimon-Brito A. Nfatc1 Promotes
Interstitial Cell Formation During Cardiac Valve Development in Zebrafish. *Circ Res*
126, 968-984 (2020).

44.   Novodvorsky P*, et al.* klf2ash317 Mutant Zebrafish Do Not Recapitulate Morpholino-
Induced Vascular and Haematopoietic Phenotypes. *PLoS One* **10**, e0141611 (2015).

45.   Kamel SM*, et al.* Istaroxime treatment ameliorates calcium dysregulation in a
zebrafish model of phospholamban R14del cardiomyopathy. *Nat Commun* **12**, 7151
(2021).

46.   Zhang R*, et al.* In vivo cardiac reprogramming contributes to zebrafish heart
regeneration. *Nature* **498**, 497-501 (2013).

47.   Jiang M*, et al.* Characterization of the Zebrafish Cell Landscape at Single-Cell
Resolution. *Front Cell Dev Biol* **9**, 743421 (2021).

48.   Wang X, Zhuo X, Gao J, Liu H, Lin F, Ma A. Neuregulin-1beta Partially Improves
Cardiac Function in Volume-Overload Heart Failure Through Regulation of Abnormal
Calcium Handling. *Front Pharmacol* **10**, 616 (2019).

49.   Slattery C, Ryan MP, McMorrow T. E2A proteins: regulators of cell phenotype in
normal physiology and disease. *Int J Biochem Cell Biol* **40**, 1431-1436 (2008).

50. Li J, *et al.* Piezo1 integration of vascular architecture with physiological force. *Nature*
515, 279-282 (2014).

51. Nonomura K, *et al.* Mechanically activated ion channel PIEZO1 is required for
lymphatic valve formation. *Proc Natl Acad Sci U S A* **115**, 12817-12822 (2018).

52. Brown D, *et al.* Neuregulin-1 is essential for nerve plexus formation during cardiac
maturation. *J Cell Mol Med* **22**, 2007-2017 (2018).

53. Gemberling M, Karra R, Dickson AL, Poss KD. Nrg1 is an injury-induced
cardiomyocyte mitogen for the endogenous heart regeneration program in zebrafish.
Elife **4**, (2015).

54. Hui SP, *et al.* Zebrafish Regulatory T Cells Mediate Organ-Specific Regenerative
Programs. *Dev Cell* **43**, 659-672 e655 (2017).

55. Palencia-Desai S, Kohli V, Kang J, Chi NC, Black BL, Sumanas S. Vascular endothelial
and endocardial progenitors differentiate as cardiomyocytes in the absence of
Etsrp/Etv2 function. *Development* **138**, 4721-4732 (2011).

56. Schumacher JA, Bloomekatz J, Garavito-Aguilar ZV, Yelon D. tal1 Regulates the
formation of intercellular junctions and the maintenance of identity in the
endocardium. *Dev Biol* **383**, 214-226 (2013).

57. Chi NC, *et al.* Foxn4 directly regulates tbx2b expression and atrioventricular canal
formation. *Genes Dev* **22**, 734-739 (2008).

58. Roman BL, *et al.* Disruption of acvrl1 increases endothelial cell number in zebrafish
cranial vessels. *Development* **129**, 3009-3019 (2002).

59. Collery RF, Link BA. Dynamic smad-mediated BMP signaling revealed through
transgenic zebrafish. *Dev Dyn* **240**, 712-722 (2011).

60. Lele Z, Bakkers J, Hammerschmidt M. Morpholino phenocopies of the swirl,
snailhouse, somitabun, minifin, silberblick, and pipetail mutations. *Genesis* **30**, 190-
194 (2001).

61.   Weber S, *et al.* SIX2 and BMP4 mutations associate with anomalous kidney

777   development. *J Am Soc Nephrol* **19**, 891-903 (2008).

779   62.   Dorsky RI, Itoh M, Moon RT, Chitnis A. Two tcf3 genes cooperate to pattern the

780   zebrafish brain. *Development* **130**, 1937-1947 (2003).

782   63.   Weinberger M, Simões FC, Patient R, Sauka-Spengler T, Riley PR. Functional

783   Heterogeneity within the Developing Zebrafish Epicardium. *Dev Cell* **52**, 574-

784   590.e576 (2020).

786   64.   Dobin A, *et al.* STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* **29**, 15-21

787   (2013).

789   65.   Li B, Dewey CN. RSEM: accurate transcript quantification from RNA-Seq data with or

790   without a reference genome. *BMC Bioinformatics* **12**, 323 (2011).

792   66.   Love MI, Huber W, Anders S. Moderated estimation of fold change and dispersion

793   for RNA-seq data with DESeq2. *Genome Biology* **15**, 550 (2014).

795   67.   Wu T, *et al.* clusterProfiler 4.0: A universal enrichment tool for interpreting omics

796   data. *The Innovation* **2**, 100141 (2021).

798   68.   Han P, *et al.* Hydrogen peroxide primes heart regeneration with a derepression

799   mechanism. *Cell Res* **24**, 1091-1107 (2014).

801   69.   Wang LW, *et al.* Standardized echocardiographic assessment of cardiac function in

802   normal adult zebrafish and heart disease models. *Dis Model Mech* **10**, 63-76 (2017).

Figure legends

Figure 1. Identification of *id2b* as a blood flow sensitive gene.

(**A**) Schematic showing the experimental procedures, including treatment, heart collection, and RNA-

sequencing of zebrafish embryonic hearts. (**B**) KEGG enrichment analysis depicting differentially

expressed genes encoding transcription factors and transcriptional regulators between control (ctrl) and

tricaine-treated embryonic hearts. Red and blue rectangles represent up-regulated and down-regulated

gene sets, respectively. |log$_2$fold change|≥0.585, adjusted *p*-value<0.1. Each replicate contains

approximately 1,000 hearts. (**C**) Heat map exhibiting representative genes from KEGG pathways

mentioned in (**B**). (**D**) qRT-PCR analysis of *id2b* and *id2a* mRNA in control and tricaine-treated

embryonic hearts. Data were normalized to the expression of *actb1*. Each sample contains ~1,000

embryonic hearts. N=three biological replicates. (**E**) *In situ* hybridization of *id2b* in 72 hpf and 96 hpf

ctrl, tricaine (1 mg/mL), and blebbistatin (10 µM)-treated embryos. Numbers at the bottom of each

panel indicate the ratio of representative staining. (**F**) *In situ* hybridization showing reduced *id2b*

expression in *tnnt2a* morpholino-injected embryos at 72 hpf compared to control. [***]*p*<0.001. Scale

bars, 50 µm.

Figure 2. The spatiotemporal expression of *id2b*.

(**A**) Schematic of the intron targeting-mediated *eGFP* knockin at the *id2b* locus using the CRISPR-

Cas9 system. The sgRNA targeting sequence and the protospacer adjacent motif (PAM) sequence are

shown in orange and blue, respectively. The donor plasmid comprises left and right arm sequences and

a *linker-FLAG-P2A-eGFP* cassette denoted by black lines with double arrows and green box,

respectively. The *linker-FLAG-P2A-eGFP* cassette was integrated into the *id2b* locus upon co-injection

of the donor plasmid with sgRNA and zCas9 protein, enabling detection by PCR using two pairs of

primers (F1, R1 and F2, R2)-the former length yielding a length of about 2.2 kb and the latter about 2.7

kb. (**B**) Zebrafish *id2b* expression pattern, as indicated by in situ hybridization of embryos at

designated time points, was consistent with the fluorescence localization of *id2b:eGFP*, revealing

expression in the heart, brain, retina, notochord, pronephric duct, and other tissues. (**C**) Maximum

intensity projections (top) and confocal sections (bottom) of *id2b:eGFP; Tg(myl7:mCherry)* hearts at

designated time points. (**D**) Maximum intensity projections (top) and confocal sections (bottom) of

id2b:eGFP; Tg(kdrl:mCherry) embryos at specific time points. Magenta, *id2b:eGFP*; yellow,

kdrl:mCherry. (**E**) RNAscope analysis of *id2b* in 96 hpf embryonic heart. White dashed line outlines

the heart. OFT, outflow tract. (**F**) Immunofluorescence of adult *id2b:eGFP; Tg(myl7:mCherry)* heart

section (left panel). Enlarged views of boxed areas are shown in the right panel. Green, eGFP; red,

mCherry; blue, DAPI. BA, bulbus arteriosus; V, ventricle; A, atrium. AVC, atrioventricular canal. (**G**)

Confocal z-stack maximum intensity projection of *id2b:eGFP;Tg(kdrl:mCherry)* embryos at 96 hpf

showing the whole body (lateral view) and the head (top view). Scale bars, 500 μm (**B**, **F**, left and **G**),

50 μm (**C** and **D**), 25 μm (**E**), 100 μm (**G**, right).

Figure 3. Cardiac contraction promotes endocardial *id2b* expression through primary cilia.

(**A**) Representative confocal z-stack (maximal intensity projection) of *id2b:eGFP* embryos under

different conditions: control, tricaine-treated, blebbistatin-treated, and *tnnt2a* morpholino-injected.

Images were captured using the same magnification. (**B**) Quantification of mean fluorescence intensity

(MFI) of *id2b:eGFP* in the working myocardium (atrium and ventricle, A+V) and atrioventricular

canal (AVC) in (**A**). Data normalized to the mean fluorescence intensity of control hearts. n=(11, 15)

(ctrl versus tricaine); n=(5, 6) (ctrl versus blebbistatin); n=(10,11) (ctrl versus *tnnt2a* MO). (**C**)

Representative confocal z-stack (maximal intensity projection) of control and *ift88* morpholino-injected

id2b:eGFP embryos. (**D**) Normalized MFI of *id2b:eGFP* in the working myocardium (A+V) and AVC

in (**C**). n=(17, 9). (**E**) Whole-mount *in situ* hybridization showing *id2b* expression in control, *klf2a*[-/-],

and *klf2b*[-/-] embryos at 48 hpf and 72 hpf. Numbers at the bottom of each panel indicate the ratio of

representative staining. Data are presented as mean ± sem. [**]*p*<0.01, [***]*p*<0.001, [****]*p*<0.0001. Scale

bars, 50 μm.

Figure 4. *id2b*[-/-] adults exhibit thinner atrioventricular valve leaflets and prominent retrograde

blood flow.

(**A**) Two sgRNAs, represented by short vertical lines, were designed to create *id2b*[-/-] mutants. Co-

injection of the two sgRNAs with zCas9 protein induces a 157 bp truncation in the exon 1 of *id2b*,

which can be detected by genotyping primers marked with arrows. This genetic modification leads to

the formation of a premature stop codon and the subsequent loss of the HLH domain. Right, qRT-PCR

analysis of *id2b* and *id2a* mRNA levels in *id2b*[+/+] and *id2b*[-/-] adult hearts. (**B**) No discernible

morphological differences were observed between *id2b*[+/+] and *id2b*[-/-] larvae at both 72 hpf and 96 hpf.

(**C**) Kaplan-Meier survival curve analysis and logrank test of *id2b*[+/+] (n=50) and *id2b*[-/-] (n=46). Wpf,

weeks post-fertilization. (**D**) Pericardial edema and an enlarged atrium are evident in a subset of $id2b^{-/-}$

adults. V, ventricle; A, atrium. (**E**) $id2b^{-/-}$ adults developed thinner atrioventricular valve leaflets

(denoted by arrowheads) compared to $id2b^{+/+}$. Enlarged views of boxed areas are shown in the bottom

panels. (**F**, **G**) Echocardiograms of adult $id2b^{+/+}$ (**F**) and $id2b^{-/-}$ (**G**) hearts. Unidirectional blood flow

was observed in the $id2b^{+/+}$ heart, while retrograde blood flow was evident in the $id2b^{-/-}$ heart. (**H**) Ratio

of retrograde flow area over inflow area shows a significant increase in retrograde flow in $id2b^{-/-}$ (n=13)

compared to $id2b^{+/+}$ (n=10). Data are presented as mean ± sem. $^*p<0.05$, $^{**}p<0.01$. Scale bars, 500 µm

(**B** and **D**, bottom), 2 mm (**D**, top), 200 µm (**E**).

Figure 5. Reduced cardiac contractile function and compromised calcium handling in $id2b^{-/-}$

mutants.

(**A**) Time-lapse imaging (from T1 to T8) illustrates the cardiac contraction-relaxation cycle of 120 hpf

$id2b^{+/+}$ and $id2b^{-/-}$ hearts carrying *myl7:mCherry*. (**B** and **C**) $id2b^{-/-}$ larvae (n=20) display a significant

decrease in heart rate and fractional area change compared to $id2b^{+/+}$ (n=20). (**D**) Echocardiograms of

adult $id2b^{+/+}$ and $id2b^{-/-}$ hearts. (**E** and **F**) $id2b^{-/-}$ fish (n=12) exhibit reduced cardiac contractile function

with preserved heart rate compared to $id2b^{+/+}$ (n=14). (**G**) Time-lapse imaging illustrates the calcium

dynamics of 120 hpf $id2b^{+/+}$ and $id2b^{-/-}$ hearts carrying *actb2:GCaMP6s*. (**H**) Ratio of maximal

fluorescence intensity (F) over basal fluorescence intensity (F0) of GCaMP6s signal. n=(26, 17). (**I**)

qRT-PCR analysis of *cacnα1c* mRNA in $id2b^{+/+}$ and $id2b^{-/-}$ hearts at 120 hpf (N=three biological

replicates, with each sample containing 500-1,000 embryonic hearts) and adult stage (N=five biological

replicates). Data were normalized to the expression of *actb1*. (**J**) The action potential of ventricular

cardiomyocyte in adult $id2b^{+/+}$ (n=9) and $id2b^{-/-}$ (n=9) hearts. (**K**) Statistical data showed a notable

difference between $id2b^{+/+}$ and $id2b^{-/-}$ accordingly. Data are presented as mean ± sem. $^{***}p<0.001$, ****

$p<0.0001$, ns, not significant. Scale bars, 50 µm.

Figure 6. Nrg1 serves as a pivotal mitogen mediating the function of $id2b$.

(**A**) Identification of genes (*fgf8a*, *nkx2.5*, *myh6*, *nrg1*, and *nkx2.7*) associated with four distinct heart

development processes: cardiac muscle tissue development, cardiomyocyte differentiation, heart

morphogenesis, and cardiac chamber development. The heatmap illustrates scaled-normalized

expression values for the mentioned genes. (**B**) qRT-PCR analysis of *nrg1* mRNA in 120 hpf *id2b*$^{+/+}$

and *id2b*$^{-/-}$ embryonic hearts. Data were normalized to the expression of *actb1*. N=four biological

replicates, with each sample containing 500-1,000 embryonic hearts. (**C**) Heart rate in 120 hpf larvae

treated with AG1478 (n=20) and DMSO (n=20). (**D**) *id2b*$^{+/+}$ and *id2b*$^{-/-}$ larvae were injected with *nrg1*

mRNA at the 1-cell stage, followed by qRT-PCR analysis of *cacnα1c* mRNA at 72 hpf. Data were

normalized to the expression of *actb1*. N=three biological replicates, with each sample containing 100-

200 embryonic hearts. (**E**) The heart rate of 72 hpf *id2b*$^{+/+}$ and *id2b*$^{-/-}$ larvae injected with *nrg1* mRNA

at 1-cell stage. n=(16, 12, 14, 14). Data are presented as mean ± sem. $^{*}p<0.05$, $^{**}p<0.01$, $^{***}p<0.001$,

$^{****}p<0.0001$. ns, not significant.

Figure 7. Id2b interacts with Tcf3b to restrict its inhibition on *nrg1* expression.

(**A**) Immunoprecipitation (IP) assays of Flag-*id2b* and HA-*tcf3b* co-transfected 293T cells. (**B**) qRT-

PCR analysis of *tcf3a*, *tcf3b*, *socs1a*, and *socs3b* mRNA in 120 hpf *id2b*$^{+/+}$ and *id2b*$^{-/-}$ embryonic hearts.

Data were normalized to the expression of *actb1*. N=four biological replicates, with each sample

containing 500-1,000 embryonic hearts. (**C**) Two potential Tcf3b binding sites, with sequences

corresponding to the human TCF3 (left) and mouse Tcf3 (right) binding motifs, were predicted in the

2,000bp DNA sequence upstream of the zebrafish *nrg1* transcription start site using JASPAR. (**D**)

Luciferase assay showing the expression of *nrg1* in embryos with *tcf3b* overexpression (*tcf3b* OE) and

morpholino-mediated *tcf3b* knockdown (*tcf3b* MO). N=three biological replicates. (**E**) qRT-PCR

analysis of *nrg1* mRNA in 72 hpf *id2b*$^{+/+}$ and *id2b*$^{-/-}$ embryonic hearts injected with control and *tcf3b*

morpholino. Data were normalized to the expression of *actb1*. N=four biological replicates, with each

sample containing 100-200 embryonic hearts. (**F**) Schematic model for *id2b*-mediated regulation of

myocardium function. During heart development, blood flow operates through primary cilia, initiating

endocardial *id2b* expression. Subsequently, the interaction between Id2b and Tcf3b restricts the activity

of Tcf3b, ensuring proper *nrg1* expression, which in turn promotes LTCC expression (left). However,

in the absence of Id2b, Tcf3b inhibits *nrg1* expression. The reduced Nrg1 hinders LTCC expression in

cardiomyocytes, resulting in decreased extracellular calcium entry and disruption of myocardial

function. Data are presented as mean ± sem. $^*p<0.05$, $^{**}p<0.01$, $^{***}p<0.001$. ns, not significant.

Figure 3-figure supplement 1. Blood flow and BMP signaling independently activates *id2b*

expression.

(**A**) Representative confocal maximal intensity projection of control (non-injected), *bmp2b*, *bmp4*, and

bmp7a morpholino-injected *id2b:eGFP*; *Tg*(*myl7:mCherry*) hearts at 24 hpf. White circles outline

eGFP signal. (**B**) Quantification of mean fluorescence intensity of *id2b:eGFP* in (**A**). Data normalized

to the mean fluorescence intensity of control hearts. n=(7, 9, 8, 10). (**C**) Representative confocal

maximal intensity projection of DMSO and Dorsomorphin (DM)-treated *id2b:eGFP* hearts at 24, 48

and 60 hpf. Embryos were treated from 10 to 24 hpf, from 24 to 48 hpf, or from 36 to 60 hpf. White

circles outline eGFP signal. (**D**) Quantification of mean fluorescence intensity of *id2b:eGFP* in (**C**).

Data normalized to the mean fluorescence intensity of DMSO-treated hearts. n=(6, 10) (24 hpf); n=(14,

16) (48 hpf); n=(9, 9) (60 hpf). (**E**) Confocal optical sections of control, tricaine-treated, and *tnnt2a*

morpholino-injected 72 hpf *Tg*(*BRE:d2GFP*); *Tg*(*kdrl:mCherry*) hearts. Yellow arrowheads,

endocardial cells. Numbers at the top of each panel indicate the ratio of representative images. (**F**)

Schematic diagram of blood flow and BMP signaling-mediated *id2b* expression. Data are presented as

mean ± sem. $^{***}p<0.001$, $^{****}p<0.0001$. ns, not significant. Scale bars, 50 µm.

Figure 4-figure supplement 1. *id2b$^{-/-}$* larvae exhibit a decreased number of valve endocardial cells

while maintaining normal atrioventricular valve function.

(**A**) Representative confocal images of valve endocardial cells in 96 hpf *id2b$^{+/+}$* and *id2b$^{-/-}$* hearts

carrying *Tg*(*kdrl:nucGFP*). (**B**) Quantification of the number of valve endocardial cells in *id2b$^{+/+}$* and

id2b$^{-/-}$ hearts. VECs, valve endocardial cells. n=(10, 10). (**C**) Representative confocal images of 72 and

120 hpf *id2b$^{+/+}$* and *id2b$^{-/-}$* hearts in *Tg*(*kdrl:nucGFP*); *Tg*(*myl7:mCherry*) transgenic background. (**D**)

Statistical analysis of the number of endocardial cells in the ventricle (V), atrium (A), and combined

(A+V) in *id2b$^{+/+}$* and *id2b$^{-/-}$* hearts. n=(13, 13) (72 hpf); n=(12, 15) (120 hpf). (**E**) Quantification of

blood flow patterns in 96 hpf *id2b$^{+/+}$* (n=16) and *id2b$^{-/-}$* (n=15) hearts. (**F**) H&E staining of adult *id2b$^{-/-}$*

and $id2b^{+/+}$ hearts after echocardiographic analysis in **Figure 4F** and **4G**. Enlarged views of boxed

areas are shown in the bottom panels. Data are presented as mean ± sem. [****]$p<0.0001$. ns, not

significant. Scale bars, 20 µm (**A**), 50 µm (**C**), 200 µm (**F**).

Figure 5-figure supplement 1. *id2b* loss-of-function impacts both valve formation and cardiac

contraction.

(**A**) The top 20 predicted tissue expression patterns based on the differential expressed genes from

$id2b^{+/+}$ and $id2b^{-/-}$ hearts at 120 hpf were displayed. (**B**) The top 20 predicted phenotypes related to the

differential expressed genes from $id2b^{+/+}$ and $id2b^{-/-}$ hearts at 120 hpf were illustrated.

Figure 5-figure supplement 2. *id2b^{-/-}* hearts develop normal trabeculae and sarcomeres.

(**A**) Time-lapse imaging (from T1 to T8) illustrates the cardiac contraction-relaxation cycle of 72 hpf

$id2b^{+/+}$ and $id2b^{-/-}$ hearts carrying *myl7:mCherry*. (**B** and **C**) Heart rate and fractional area change in

$id2b^{-/-}$ (n=21) and $id2b^{+/+}$ (n=21) at 72 hpf. (**D**) Representative confocal z-stack of 72 and 120 hpf

$id2b^{+/+}$ and $id2b^{-/-}$ hearts with *Tg(myl7:H2A-mCherry)* transgene. (**E**) Quantification of the number of

cardiomyocytes in the ventricle (V), atrium (A), and combined (A+V) in (**D**). n=(8, 10) (72 hpf); n=(10,

10) (120 hpf). (**F**) Representative confocal images of 72 and 120 hpf $id2b^{+/+}$ and $id2b^{-/-}$ hearts carrying

Tg(myl7:mCherry). (**G**) Confocal immunofluorescence images of α-actinin (green) in embryonic (72

hpf) and adult (115 dpf) hearts (left). Right: fluorescence intensity profiles for α-actinin. Data are

presented as mean ± sem. ns, not significant. Scale bars, 50 µm (**D** and **F**), 5 µm (**G**).

Figure 6-figure supplement 1. *nrg1* is expressed in the endocardial cells.

(**A**) Analysis of *nrg1* expression using the zebrafish single-cell landscape database. The red rectangle

highlights endocardial cells. (**B**) Identification of the endocardial cell population in the zebrafish cell

landscape. Purple dots represent endocardial cells, and the red oval denotes cell populations from the

heart. Images were generated using the Zebrafish Cell Landscape (ZCL) at http://bis.zju.edu.cn/ZCL/.

Figure 7-figure supplement 1. Expression landscape of zebrafish *tcf3b*.

(A) The major cell types expressing *tcf3b* were displayed according to the zebrafish single-cell

landscape. **(B)** Identification of the endocardial cell population in the zebrafish cell landscape. Purple

dots represent endocardial cells, and the red oval denotes cell populations from the heart. Images were

generated using the Zebrafish Cell Landscape (ZCL) at http://bis.zju.edu.cn/ZCL/.

Video 1. 96 hpf *id2b*$^{+/+}$ larvae displayed unidirectional blood flow in the AV canal. Scale bar, 50

μm.

Video 2. 96 hpf *id2b*$^{-/-}$ larvae displayed unidirectional blood flow in the AV canal. Scale bar, 50 μm.

A

sgRNA target   PAM
TCTGTTGAACACATGACGATATTTGGAAG

Endogenous *id2b* locus   Exon1   Exon2   stop codon
F1   Left arm   Right arm   R2

Left arm   Right arm
F2   Exon2   *linker-Flag-P2A-eGFP*   R1

id2b-linker-Flag-P2A-EGFP donor

+ *sgRNA*
+*zCas9 protein*

Targeted *id2b* locus
2.2kb   2.7kb
Exon1   Exon2   *linker-Flag-P2A-eGFP*   Exon2
F1   R1   F2   R2
plasmid fragment

B   24 hpf   48 hpf   72 hpf

C   48 hpf   72 hpf   96 hpf
id2b:eGFP; *myl7:mCherry*

D   *id2b:eGFP*; *kdrl:mCherry*

E   *id2b*   96 hpf
OFT
Ventricle

F   BA   AVC   A
V
DAPI
id2b:eGFP
myl7:mCherry

G   *id2b:eGFP*; *kdrl:mCherry*   96 hpf
id2b:eGFP
kdrl:mCherry

Figure 2