Contraction-induced endocardial *id2b* plays a dual role in regulating myocardial contractility and valve formation

Shuo Chen[1, 2, 3], Jie Yin[1, 2, 3], Jinxiu Liang[1, 2, 3], Weijia Zhang[1, 2, 3], Peijun Jiang[1, 2, 3], Wenyuan Wang[4, 5], Xiaoying Chen[6, 7], Yuanhong Zhou[8], Peng Xia[8], Fan Yang[6, 7], Ying Gu[1,2, 3], Ruilin Zhang[4, 5], Peidong Han[1, 2, 3*]

[1]*Center for Genetic Medicine, The Fourth Affiliated Hospital, Zhejiang University School of Medicine, Hangzhou, Zhejiang, China.*

[2]*Institute of Genetics, Zhejiang University International School of Medicine, Hangzhou, Zhejiang, China.*

[3]*Division of Medical Genetics and Genomics, The Children's Hospital, Zhejiang University School of Medicine, National Clinical Research Center for Child Health, Hangzhou, Zhejiang 310058, China.*

[4]*TaiKang Medical School (School of Basic Medical Sciences), Wuhan University, Wuhan, China*

[5]*Hubei Provincial Key Laboratory of Developmentally Originated Disease, Wuhan, China*

[6]*Department of Biophysics, and Kidney Disease Center of the First Affiliated Hospital, Zhejiang University School of Medicine, Hangzhou, China.*

[7]*Liangzhu Laboratory, Zhejiang University Medical Center, Hangzhou, China.*

[8]*Zhejiang Provincial Key Laboratory for Cancer Molecular Cell Biology, Life Sciences Institute, Zhejiang University, Hangzhou, Zhejiang 310058, China.*

*To whom correspondence should be addressed.

hanpd@zju.edu.cn

+86-571-88206360

Abstract

Biomechanical cues play an essential role in sculpting organ formation. Comprehending how cardiac

cells perceive and respond to biomechanical forces is a biological process with significant medical

implications that remains poorly understood. Here we show that biomechanical forces activate

endocardial *id2b* (inhibitor of DNA binding 2b) expression, thereby promoting cardiac contractility and

valve formation. Taking advantage of the unique strengths of zebrafish, particularly the viability of

embryos lacking heartbeats, we systematically compared the transcriptomes of hearts with impaired

contractility to those of control hearts. This comparison identified *id2b* as a gene sensitive to blood

flow. By generating a knockin reporter line, our results unveiled the presence of *id2b* in the

endocardium, and its expression is sensitive to both pharmacological and genetic perturbations of

contraction. Furthermore, *id2b* loss-of-function resulted in progressive heart malformation and early

lethality. Combining RNA-seq analysis, electrophysiology, calcium imaging, and echocardiography,

we discovered profound impairment in atrioventricular (AV) valve formation and defective excitation-

contraction coupling in *id2b* mutants. Mechanistically, deletion of *id2b* reduced AV endocardial cell

proliferation and led to a progressive increase in retrograde blood flow. In the myocardium, *id2b*

directly interacted with the bHLH component *tcf3b* (transcription factor 3b) to restrict its activity.

Inactivating *id2b* unleashed its inhibition on *tcf3b*, resulted in enhanced repressor activity of *tcf3b*,

which subsequently suppressed the expression of *nrg1* (neuregulin 1), an essential mitogen for heart

development. Overall, our findings identify *id2b* as an endocardial cell-specific, biomechanical

signaling-sensitive gene, which mediates intercellular communications between endocardium and

myocardium to sculpt heart morphogenesis and function.

Keywords: Zebrafish, Heart, *id2b*, Valve, Contraction

Introduction

The heart develops with continuous contraction, and biomechanical cues play an essential role in cardiac morphogenesis[1,2]. Blood flow is directly sensed by the surrounding endocardium, which undergoes multiscale remodeling during zebrafish heart development. In the atrioventricular canal (AVC) endocardium, oscillatory flow promotes valvulogenesis through transient receptor potential (TRP) channel-mediated expression of Krüppel-like factor 2a (*klf2a*)[3-5]. Meanwhile, mechanical forces trigger ATP-dependent activation of purinergic receptors, inducing expression of nuclear factor of activated T cells 1 (*nfatc1*) and subsequent valve formation[6]. In the chamber endocardium, blood flow induces endocardial cells to adopt chamber- and region-specific cell morphology during cardiac ballooning[7]. A recent study further emphasizes that blood flow is essential for endocardial cell accrual in assembling the outflow tract[8]. Beyond their role in endocardial cells, proper biomechanical cues are indispensable for shaping the myocardium. For instance, in contraction compromised *tnnt2a*[9] and *myh6*[10] mutants, trabeculation is markedly reduced. Moreover, apart from the tissue-scale regulatory effect, the shape changes[11] and myofibril content[12] at the single-cardiomyocyte level are also sculpted by the interplay of contractility and blood flow in the developing heart.

In ventricular myocardium morphogenesis, biomechanical forces coordinate intra-organ communication between endocardial and myocardial cells by regulating BMP, Nrg/Erbb, and Notch signaling. The Nrg-Erbb axis stands as one of the most extensively studies signaling pathway mediating cell-cell communications in the heart[13-15]. In particular, endocardial Notch activity induced by cardiac contraction promotes the expression of Nrg, which then secretes into the extracellular space, binding to Erbb2/4 receptor tyrosine kinases on cardiomyocyte and promoting their delamination[16,17]. In agreement with the pivotal role of this signaling pathway in heart development, genetic mutations in zebrafish *nrg2a* and *erbb2* result in severely compromised trabeculae formation[18-20].

The development of cardiomyocytes encompasses the specification of subcellular structure, metabolic state, gene expression profile, and functionality[21,22]. The rhythmic contraction of cardiomyocytes relies on precisely regulated excitation-contraction coupling (E-C coupling), transducing electrical activity into contractile forces. This intricate signaling cascade involves membrane action potential, calcium signaling, and sarcomeric structure[23]. Specifically, membrane

depolarization triggers the opening of L-type calcium channel (LTCC), allowing calcium influx. The

calcium signaling then activates the ryanodine receptor on the sarcoplasmic reticulum (SR) membrane,

releasing additional calcium[23]. E-C coupling is essential for heart development, as evidenced by the

complete silence of the ventricle and reduced cardiomyocyte number in *cacna1c* (LTCC α1 subunit in

zebrafish) mutant[24]. Beyond its role in modulating cardiac structure formation, previous studies

indicate that Nrg-Erbb2 signaling is also necessary for cardiac function, as *erbb2* mutants exhibit

severely compromised fractional shortening and an immature conduction pattern[18,25]. However, the

precise mechanism through which the blood flow-Nrg-Erbb2 axis controls cardiomyocyte functionality

remains poorly understood.

Results

Transcriptome analysis identifies *id2b* as a blood flow sensitive gene

Blood circulation is dispensable for early embryonic development in zebrafish, presenting an ideal

model to investigate biomechanical cues influencing heart morphogenesis. To identify genes affected

by cardiac contraction or blood flow, we treated *myl7:mCherry* zebrafish embryos with tricaine to

inhibit cardiac contractility from 72 hours post-fertilization (hpf) to 96 hpf. Hearts from control and

tricaine-treated zebrafish embryos were manually collected under a fluorescence stereoscope as

previously reported[26]. Subsequently, approximately 1,000 hearts from each group were subjected to

RNA sequencing (Figure 1A). A total of 4,530 genes with differential expression were identified,

comprising 2,013 up-regulated and 2,517 down-regulated genes. With a specific focus on identifying

key transcription factors (TFs) affected by perturbing biomechanical forces, differentially expressed

genes (DEGs) encoding TFs were enriched into signaling pathways through KEGG analysis.

Interestingly, our analysis identified several pathways known to be involved in heart development,

including the transforming growth factor beta (TGFβ) signaling and Notch signaling pathways (Figure

1B). In particular, the scaled expression levels of the top 6 DEGs ($|\log_2 FC| \geqslant 0.585$), exhibiting up- or

down-regulation, were listed (Figure. 1C). Notably, as a component of the helix-loop-helix (HLH)

family, *id2b* is the zebrafish homolog of the mammalian *Id2* gene, functioning as a transcriptional

repressor. Intriguingly, *Id2* has been shown to regulate the formation of the murine AV valve, a process

notably influenced by alterations in blood flow directionality. Consequently, we interrogated the

expression and function of *id2b* in developing embryos.

Through quantitative real-time PCR (qRT-PCR) analysis on purified embryonic hearts, we

observed a ~50% reduction in *id2b* mRNA level following tricaine treatment from 72 to 96 hpf

compared to the control (Figure 1D). Furthermore, in situ hybridization was performed to visualize

id2b expression under tricaine or 10 uM blebbistatin (an inhibitor of sarcomeric function and cardiac

contractility) treatment from 48 to 72 or from 72 to 96 hpf. Consistently, our results showed a reduction

in *id2b* signal in contraction-deficient hearts compared to the control (Figure 1E). Similarly, injection

of a previously characterized *tnnt2* morpholino at the 1-cell stage also led to compromised contraction

and diminished expression of *id2b* (Figure 1F). Taken together, these results indicate that

biomechanical cues are essential for activating *id2b* in embryonic hearts.

Visualization of the spatiotemporal expression of *id2b* in developing embryos

Due to technical challenges in visualizing the cell-type-specific expression of *id2b* in the developing

heart using whole-mount in situ hybridization, we employed an intron targeting-mediated approach[27] to

generate a knockin *id2b:eGFP* line. This method allowed us to achieve specific labeling without

perturbing the integrity and function of the endogenous gene[27] (Figure 2A). Comparison of *id2b:eGFP*

fluorescence with in situ hybridization at 24, 48, and 72 hpf revealed a substantial recapitulation of the

eGFP signal with that of endogenous *id2b* expression. The fluorescence was notably enriched in the

heart, brain, retina, and notochord (Figure 2B), mirroring observations from a previously reported *id2b*

transgenic line generated through BAC-mediated recombination[28].

To further elucidate the spatiotemporal expression of *id2b* in developing hearts, we crossed

id2b:eGFP with *myl7:mCherry* or *kdrl:mCherry*, labelling cardiomyocytes or endocardial cells,

respectively. Confocal images revealed minimal, if any, presence of *id2b:eGFP* in *myl7:mCherry*[+]

cardiomyocytes (Figure 2C). In sharp contrast, clear co-localization between *id2b:eGFP* and

kdrl:mCherry was evident at 48, 72, and 96 hpf (Figure 2D). Interestingly, there was an absence of

id2b:eGFP signal in *kdrl:mCherry*[+] endothelial cells in trunk blood vessel and brain vasculature

(Figure 2E). In adult hearts, *id2b:eGFP* fluorescence was enriched in the chamber endocardium and the

endothelium lining AVC, outflow tract (OFT), and bulbus arteriosus (Figure 2F).

Cardiac contraction promotes endocardial *id2b* expression through primary cilia but not BMP

Taking advantage of live imaging on developing embryos, we explored the *in vivo* dynamics of *id2b* in

response to biomechanical force at single-cell resolution. When embryos were treated with tricaine or

blebbistatin, the intensity of *id2b:eGFP* in atrial and ventricular endocardium was significantly reduced

(Figure 3A, B). Similarly, injection of *tnnt2a* morpholino also markedly suppressed *id2b:eGFP* signal

(Figure 3A, B), in agreement with the results obtained from in situ hybridization. Strikingly, the

reduction in fluorescence intensity was particularly pronounced in AVC endothelial cells (Figure 3A, B,

asterisks).

We then explored how cardiac contraction modulated *id2b* expression. Given that endocardial

cells can sense blood flow through primary cilia[29,30], we used a characterized morpholino[29] to

knockdown *ift88*, an intraflagellar transporter essential for primary cilia formation. Previously work

demonstrated a complete loss of primary cilia in endocardial cells upon *ift88* knockdown[29]. As

expected, a significant decrease in *id2b:eGFP* intensity was observed in the chamber and AVC

endocardium of *ift88* morphants compared to control (Figure 3C, D), suggesting that biomechanical

forces promote the expression of *id2b* via primary cilia. In the developing heart, a central hub for

mediating biomechanical cues is the Klf2 gene, which includes the *klf2a* and *klf2b* paralogues in

zebrafish[3-5,29,31]. Previous studies in mammals and zebrafish have highlighted the essential role of Klf2

transcription factor activity in cardiac valve and myocardial wall formation[3,31]. As a flow-responsive

gene, *klf2a* expression has been observed throughout the entire endocardium, evidenced by mRNA

expression and transgenic studies[3,4]. Interestingly, in situ hybridization on 48 and 72 hpf *klf2a*$^{-/-}$ and

klf2b$^{-/-}$ embryos unveiled a drastic decrease in *id2b* expression compared with wild-type zebrafish

(Figure 3E), supporting the notion that *klf2*-mediated biomechanical signaling is essential for activating

id2b expression.

Because *id2b* has been reported to be a target gene of bone morphogenetic protein (BMP)

signaling, we explored whether BMP also played a role in regulating *id2b* when blood flow was

perturbed. To this end, we knocked down *bmp2b*, *bmp4*, and *bmp7a* in 1-cell stage *id2b:eGFP* embryos

and found a significant reduction in fluorescence signal in morpholino-injected hearts compared to

control (Figure 3-figure supplement 1A, B), suggesting that *id2b* is a target gene of BMP signaling

during early embryonic development. Considering that heartbeats in zebrafish commence at

approximately 22 hpf, we treated embryos with the BMP inhibitor Dorsomorphin from 24 to 48 hpf or

from 36 to 60 hpf. While the number of endocardial cells was slightly reduced upon Dorsomorphin

exposure, as previously reported[7], surprisingly, quantification of the average *id2b:eGFP* fluorescence

intensity in individual endocardial cells revealed no significant differences between Dorsomorphin and

DMSO controls (Figure 3-figure supplement 1C, D).

We further visualized BMP activity using the *Bre:dGFP* reporter line. Confocal images revealed

strong fluorescence in the myocardium at 72 hpf, with minimal signal present in the endocardium

except for the AVC endothelium (Figure 3-figure supplement 1E). Moreover, after tricaine treatment,

endocardial *Bre:dGFP* slightly increased (Figure 3-figure supplement 1E), as opposed to the reduced

id2b:eGFP signal (Figure 3A, B). Likewise, endocardial *Bre:dGFP* intensity was barely affected after

completely blocking contraction with *tnnt2a* MO injection (Figure 3-figure supplement 1E). These

observations align with previous work using pSmad-1/5/8 as a readout of BMP activity, indicating that

endocardial BMP signaling is independent of blood flow[7]. Collectively, these results suggest that BMP

signaling and blood flow modulate *id2b* expression in a developmental-stage-dependent manner.

Compromised AV valve formation in *id2b* mutants

To investigate the role of the contractility-*id2b* axis in zebrafish heart development, we generated a

loss-of-function mutant line using CRISPR/Cas9. A pair of sgRNAs designed to target exon 1 was

injected with zCas9 protein into 1-cell-stage embryos. Consequently, we identified a mutant allele with

a 157 bp truncation, leading to the generation of a premature stop codon (Figure 4A). The overall

morphology of *id2b* mutants (*id2b$^{-/-}$*) remained unaltered compared to *id2b$^{+/+}$* siblings at 72 and 96 hpf

(Figure 4B). However, *id2b$^{-/-}$* zebrafish were subjected to early lethality starting around 31 weeks post-

fertilization (Figure 4C). Strikingly, pericardial edema was observed in 20% (9/45) of adult *id2b$^{-/-}$*

zebrafish (Figure 4D, top). Upon dissecting hearts from these *id2b$^{-/-}$* zebrafish, a prominent enlargement

in the atrium was detected (Figure 4D, bottom). Histological analysis further revealed malformation in

the AV valves of these $id2b^{-/-}$ mutants compared to the control (Figure 4E). Upon closer inspection of

AV valve anatomy, we noted that the superior and inferior leaflets were significantly thinner,

comprising only 1-2 layer of cells in $id2b^{-/-}$ zebrafish with an enlarged atrium. This was in sharp

contrast to $id2b^{+/+}$ zebrafish, which exhibited multilayers of cells (Figure 4E). Subsequent examination

of the remaining 80% of $id2b^{-/-}$ zebrafish (36/45) that did not display prominent pericardial edema also

revealed an enlarged atrium and AV valve malformation, albeit to a lesser extent (data not shown). In

contrast, $id2b^{+/+}$ controls (48/48) displayed normal heart morphology with intact AV valve (Figure 4E,

left).

To further interrogate the effect of $id2b$ inactivation on AV valve formation and function, we

analyzed the number of AVC endothelial cells using $kdrl:nucGFP$. At 96 hpf, a reduced number of

$kdrl:nucGFP^{+}$ cells was detected in the AVC region of $id2b^{-/-}$ embryos compared with $id2b^{+/+}$ (Figure

4-figure supplement 1A, B). In contrast, the number of atrial and ventricular endocardial cells did not

differ between $id2b^{-/-}$ and $id2b^{+/+}$ (Figure 4-figure supplement 1C, D). Subsequently, we assessed

hemodynamic flow by conducting time-lapse imaging of red blood cells labelled by $gata1:dsred$.

Surprisingly, the pattern of hemodynamics was largely preserved in $id2b^{-/-}$ compared with $id2b^{+/+}$

siblings at 96 hpf (Figure 4-figure supplement 1E, Video 1, 2). Additionally, we performed

echocardiography to analyze blood flow in adult zebrafish as previously described[32]. In $id2b^{-/-}$ hearts,

prominent retrograde blood flow was detected in the AVC region (8/13) (Figure 4G), while

unidirectional blood flow was observed in $id2b^{+/+}$ (10/10) (Figure 4F). Quantification analysis showed

~32% retrograde blood flow in $id2b^{-/-}$, compared to 0% in $id2b^{+/+}$ zebrafish (Figure 4H). Consistently,

the superior and inferior leaflets were much thinner in $id2b^{-/-}$ exhibiting retrograde flow compared with

control fish (data not shown). Overall, these histological and functional analyses indicate that $id2b$

deletion leads to progressive defects in AV valve morphology and hemodynamic flow.

$id2b$ deletion perturbs calcium signaling and contractile function in the myocardium

Although similar defects in AV valve formation have been reported in both $klf2a$ and $nfatc1$ mutants,

they do not display noticeable pericardial edema at the adult stage, nor do they experience early

lethality[3,29,31-33]. Therefore, we sought to investigate whether other cardiac properties have also been

affected by *id2b* loss-of-function. To this end, we employed RNA-seq analysis on purified embryonic

id2b$^{-/-}$ and *id2b*$^{+/+}$ hearts (Figure 5-figure supplement 1). As expected, enrichment analysis of DEGs

demonstrated that the top-ranked anatomical structures affected by *id2b* deletion included the heart

valve, the compact layer of ventricle, and the atrioventricular canal (Figure 5-figure supplement 1A).

Interestingly, *id2b* inactivation also impacted phenotypes such as cardiac muscle contraction and heart

contraction (Figure 5-figure supplement 1B). Therefore, we investigated cardiac contractile function

through time-lapse imaging on the *myl7:mCherry* background. At 120 hpf, a significant decrease in

cardiac contractility was observed in *id2b*$^{-/-}$, as evidenced by reduced fractional area change and heart

rate, compared with *id2b*$^{+/+}$ (Figure 5A-C). Similarly, echocardiography analysis showed that the

contractile function in adult *id2b*$^{-/-}$ heart was dramatically reduced compared with age-matched *id2b*$^{+/+}$

(Figure 5D, E). These functional defects in *id2b*-deleted hearts could not be attributed to differences in

cardiomyocyte number, as we counted cardiomyocytes using the *myl7:H2A-mCherry* line and found no

apparent changes between *id2b*$^{-/-}$ and *id2b*$^{+/+}$ embryos at 72 and 120 hpf (Figure 5-figure supplement

2A, B). Similarly, *id2b*$^{-/-}$ also developed regular trabecular structures at embryonic (Figure 5-figure

supplement 2C) and adult stages (Figure 4E). Through α-actinin immunostaining, we observed similar

sarcomeric structures in 72 hpf and adult (115 dpf) *id2b*$^{-/-}$ and control cardiomyocytes (Figure 5-figure

supplement 2D), corroborating that the reduced contractility in *id2b*-depleted heart was independent of

structural defects.

The key functional unit that transmits electrical activity to contractile function is E-C coupling.

Because *id2b*$^{-/-}$ displayed reduced cardiac function without conspicuous myocardial structural defects,

we visualized calcium signaling in the developing heart using *actb2:GCaMP6s* zebrafish (Figure 5G).

Compared to *id2b*$^{+/+}$ control, *id2b*$^{-/-}$ exhibited markedly decreased calcium amplitude at 120 hpf (Figure

5H), which is consistent with the reduced fractional area change. In cardiomyocyte, the entry of

extracellular calcium is mainly mediated through the LTCC. As previously reported, a defect in

zebrafish LTCC pore forming α1 subunit *cacna1c* leads to compromised cardiac function[24]. We

collected hearts from 72 hpf and 5 months post-fertilization (mpf) zebrafish and detected

downregulated *cacna1c* in *id2b*$^{-/-}$ compared to *id2b*$^{+/+}$ (Figure 5I). In addition, we measured cardiac

action potential using intracellular recording[34]. Compared to $id2b^{+/+}$ zebrafish, the duration of the

action potential in $id2b^{-/-}$ was significantly shorter (Figure 5J, K), consistent with the decreased

expression level of *cacna1c*. Together, these data indicate that *id2b* loss-of-function leads to

compromised calcium signaling and cardiac contractile function.

Reduced expression of *nrg1* mediates the compromised contractility in *id2b*$^{-/-}$

Because the deficiency of *id2b* in the endocardium disrupted the function of myocardium, we

speculated that the crosstalk between these two types of cells was affected in $id2b^{-/-}$. Interestingly,

comparing the differentially expressed genes in embryonic $id2b^{-/-}$ and $id2b^{+/+}$ hearts identified a

significant reduction in the expression level of *nrg1*, a key mitogen regulating the intra-organ

communications between endocardial cells and cardiomyocytes (Figure 6A). Remarkably, analysis of a

zebrafish single-cell database[35] revealed enriched expression of *nrg1* in endocardial cells (Figure 6-

figure supplement 1). However, attempts to detect *nrg1* expression through in situ hybridization were

unsuccessful, likely due to its low abundance in the heart. Alternatively, qRT-PCR analysis of purified

72 hpf embryonic hearts validated decreased *nrg1* levels in $id2b^{-/-}$ compared to control (Figure 6B).

Previous studies have demonstrated that perturbations in Nrg-Erbb2 signaling, as seen in zebrafish

erbb2 mutants, result in dysfunctional cardiac contractility[18]. Consistently, a decrease in heart rate was

observed in embryos treated with the *erbb2* inhibitor AG1478 (Figure 6E).

Remarkably, injecting *nrg1* mRNA at the 1-cell stage not only rescued the reduced expression of

cacna1c in $id2b^{-/-}$ hearts (Figure 6C) but also restored the diminished heart rate (Figure 6D). This is

consistent with prior study showing that Nrg1 administration can restore LTCC expression and calcium

current in failing mammalian cardiomyocytes[36]. Overall, our data suggest that endocardial *id2b*

promotes *nrg1* synthesis, thereby enhancing cardiomyocyte contractile function.

***id2b* interacts with *tcf3b* to limit its repressor activity on *nrg1* expression**

We further interrogated how *id2b* promotes the expression of *nrg1*. As a HLH factor lacking a DNA

binding motif, *id2b* has been reported to form a heterodimer with *tcf3b* and inhibit its function. Notably,

we detected expression of *tcf3b* in endocardial cells by analyzing a zebrafish single-cell database[35]

277 (Figure 7-figure supplement 1). To determine whether this interaction occurs in zebrafish, Flag-*id2b*

278 and HA-*tcf3b* were co-expressed in HEK293 cells. Co-immunoprecipitation analysis confirmed the

279 direct binding of *id2b* to *tcf3b* protein (Figure 7A). qRT-PCR analysis on purified 72 hpf embryonic

280 hearts revealed a significant increase in the expression of *socs3b* and *socs1a*, target genes of *tcf3b*, in

281 *id2b$^{-/-}$* compared to *id2b$^{+/+}$* (Figure 7B). This suggests an elevation in *tcf3b* activity associated with the

282 loss of *id2b* function. Notably, the expression levels of *tcf3a* and *tcf3b* remained consistent between

283 *id2b$^{-/-}$* and *id2b$^{+/+}$* hearts (Figure 7B).

284  To understand how the altered interaction between *id2b* and *tcf3b* influences *nrg1* expression, we

285 analyzed the promoter region of zebrafish *nrg1* using JASPAR and identified two potential *tcf3b*

286 binding sites (Figure 7C). Subsequently, a DNA fragment containing the zebrafish *nrg1* promoter

287 region was subcloned into a vector carrying the luciferase reporter gene. Co-injection of this construct

288 with *tcf3b* mRNA into 1-cell stage embryos resulted in a significant decrease in luciferase signal.

289 Conversely, co-injection with a previously characterized *tcf3b* morpholino led to enhanced luciferase

290 intensity (Figure 7D). These results suggest a possible mechanism by which *tcf3b* represses *nrg1*

291 expression in zebrafish.

292  Lastly, injecting *tcf3b* morpholino into *id2b$^{-/-}$* embryos was performed to assess whether

293 attenuating the overactive *tcf3b* in *id2b$^{-/-}$* could restore the expression level of *nrg1*. qRT-PCR analysis

294 of purified 72 hpf hearts revealed a partial restoration of the diminished *nrg1* expression in *id2b$^{-/-}$* upon

295 *tcf3b* inhibition (Figure 7E). Taken together, our results indicate that biomechanical cues activate

296 endocardial *id2b* expression, leading to its interaction with *tcf3b* to alleviate repression on the

297 *nrg1* promoter. Consequently, the depletion of *id2b* unleashes *tcf3b*'s repressor activity, leading to a

298 reduction in *nrg1* expression, which further acts through *erbb2* to regulate cardiomyocyte function

299 (Figure 7F).

301 **Discussion**

302 Biomechanical forces play an essential role in regulating the patterning and function of the heart. At

303 AVC, oscillatory flow promotes the expression of *klf2a* and *nfatc1* to modulate valve morphogenesis.

304 In chamber endocardium, blood flow induces endocardial cells to acquire distinct cell morphology.

However, it still lacks a systematic analysis of the transcriptome underlying compromised heartbeats.

In the present study, we analyzed embryonic zebrafish hearts without contractility and identified genes

that are regulated by biomechanical forces. Specifically, our results unveiled the endocardial-specific

expression of *id2b*, which was tightly regulated by flow-sensitive primary cilia-*klf2* axis. Genetic

deletion of *id2b* resulted in compromised valve formation and progressive atrium enlargement. In

addition, a reduction in heart rate and contractile force was observed in *id2b*[-/-], owing to decreased

expression of LTCC α1 subunit *cacna1c*. Mechanistically, *id2b* interacts with bHLH transcription

factor *tcf3*b to limit its repressor activity. Hence genetic deletion of *id2b* unleashes *tcf3*b activity, which

further represses endocardial *nrg1* expression. As a result, Injection of *nrg1* mRNA partially rescues

the phenotype of *id2b* deletion. Overall, our findings identify *id2b* as a novel mediator that regulate the

interplay between endocardium and myocardium during heart development.

In mammals, the deletion of *Id2* leads to malformations in the arterial and venous poles of the

heart, as well as affects AV valve morphogenesis[37,38]. Interestingly, approximately 20% perinatal

lethality is reported in *Id2* knockout mice, exhibiting AV septal defects and membranous ventricular

septal defects[37]. Remarkably, pericardial edema is evident in 20% of adult *id2b*[-/-] zebrafish, with a

prominent enlargement of the atrium. The superior and inferior leaflets of AV valves in *id2b*[-/-] mutants

are significantly thinner compared to the control. Therefore, our results suggest that *id2b* may play a

similar role in regulating AV valve formation in zebrafish as its mammalian orthologue *Id2*. It is

proposed that the loss of *Id2* in mice results in compromised endocardial proliferation and aberrant

endothelial-to mesenchymal transformation (EMT), collectively leading to defective valve

morphogenesis[37]. Nevertheless, the mechanism by which *id2b* loss-of-function causes a reduction in

leaflet thickness in zebrafish remains to be determined in future studies.

id2b has been recognized as a target gene of the BMP signaling pathway. As expected,

knockdown of *bmp2b*, *bmp4*, and *bmp7a* at 1-cell stage confirms that endocardial *id2b* expression is

controlled by BMP activity during early embryonic development. Surprisingly, treatment with the BMP

inhibitor Dorsomorphin at 24 and 36 hpf, when cardiac contractions have already initiated, fails to alter

id2b expression in the endocardium, suggesting that BMP is dispensable for *id2b* activation at these

stages. Instead, endocardial *id2b* expression is reduced upon loss-of-function of *klf2a*, *klf2b*, and *ift88*,

suggesting an essential role of the primary cilia-*klf2* axis in mediating *id2b* activation. In endocardial

cells, Trp, Piezo, and ATP-dependent P2X/P2Y channels[4,6,39,40] are well-established sensors for

biomechanical stimulation. The activation of these channels further promotes the activities of Klf2 and

Nfatc1 to drive heart development and valvulogenesis. However, whether these channels are also

required for the activation of *id2b* warrants further investigation.

The Nrg-Erbb signaling plays an essential role in regulating heart morphogenesis and function. In

the mammalian heart, the genetic deletion of Nrg1 or Erbb2 results in severely perturbed cardiac

trabeculae formation[13-15]. Zebrafish *erbb2* mutants exhibit a similar defect in cardiomyocyte

proliferation and trabeculation[18]. Interestingly, *nrg1* mutant zebrafish display grossly normal cardiac

structure during early embryonic devolpment[19,41]. Nevertheless, zebrafish *nrg2a* loss-of-function leads

to defective trabeculae formation, suggesting that *nrg2a* is the predominant ligand secreted from

endocardium, promoting ventricular morphogenesis[19]. In the adult stage, perivascular cells[42] or

regulatory T cells[43]-derived *nrg1* promotes cardiomyocyte proliferation during heart regeneration.

Hence, the specific ligand/receptor and the spatiotemporal regulation of the Nrg-Erbb axis appear to be

more complicated in both embryonic and adult zebrafish. Interestingly, the *nrg1* mutant heart exhibits a

defect in cardiac nerve expansion and heart maturation at the juvenile stage despite normal cardiac

structure formation[41], suggesting its potential role in regulating cardiac function. Our findings

demonstrate that the expression of *nrg1* in embryonic endocardial cells is influenced by biomechanical

cues and *id2b* activity. This signaling axis is essential for coordinating endocardium-myocardium

interaction and establishing proper cardiac function.

Materials and methods

Zebrafish handling and lines

All animal procedures were approved by the Animal Care and Use Committee of the Zhejiang

University School of Medicine. Embryonic and adult fish were raised and maintained under standard

conditions at 28 °C on a 14/10 hour day/night cycle. The following zebrafish lines were used in this

study: *Tg(myl7:mCherry)*[sd7 44], *Tg(myl7:H2A-mCherry)*[sd12 45], *Tg(kdrl:mCherry)*[S896 46]*, Tg(kdrl:*

nucGFP)[y7 47], *Tg(Bre:d2GFP)*[mw30 48], and *Tg(actb2:Gcamp6s).* To generate the *id2b* mutant, two short

guide RNAs (sgRNAs) targeting exon 1 were generated using the MAXIscript T7 transcription kit

(ambion, AM1314). The sgRNAs were as follows: sgRNA1: 5' - GAAGGCAGTCAGTCCGGTG - 3';

sgRNA2: 5' - GAACCGGAGCGTGAGTAAGA - 3'. The two sgRNAs, along with zCas9 protein,

were co-injected into one-cell stage embryos. Embryos were raised to adulthood and crossed to wild-

type zebrafish to obtain F_1 progenies. Through PCR analysis, a mutant line with a 157 bp truncation

was identified.

The knock-in *id2b:eGFP* line was generated using a previously reported method[27]. Briefly, three

sgRNAs were designed to target the intron of *id2b* (sgRNA1: 5' - GAGACAAATATCTACTAGTG -

3'; sgRNA2: 5' - GTTGAACACATGACGATATT - 3'; sgRNA3: 5' -

GCACAACTTAGATTTCAAGT - 3'). Co-injection of each individual sgRNA with zCas9 protein into

one-cell stage zebrafish embryos yielded varying cleavage efficiency. Since sgRNA2 displayed the

highest gene editing efficiency, it was selected for subsequent studies. Next, a donor plasmid

containing the sgRNA targeting sequence of the intron, exon 2 of *id2b*, and P2A-eGFP was generated.

Co-injection of sgRNA, donor plasmid, and zCas9 protein into one-cell stage embryos led to

concurrent cleavage of the sgRNA targeting sites in both the zebrafish genome and the donor plasmid

(Figure 2A). Accordingly, eGFP fluorescence was observed in injected 24 hpf zebrafish embryos,

indicating the incorporation of the donor. The insertion of the *id2b*-p2A-eGFP donor into the genome

was confirmed by PCR analysis with primers recognizing target site or donor sequences, respectively

(Figure 2A). Embryos with mosaic eGFP expression were raised to adulthood and crossed with wild-

type zebrafish to obtain F_1 progenies. Overall, two founders were identified. The junction region of the

F_1 embryos was sequenced to determine the integration sites. Although the two founders had slightly

different integration sites in the intron, the expression pattern and fluorescence intensity of eGFP were

indistinguishable between the two lines.

Morpholinos

All morpholinos (Gene Tools) used in this study have been previously characterized. *tnnt2a* MO (5' -

CATGTTTGCTCTGATCTGACACGCA - 3')[49]; *ift88* MO (5' -

CTGGGACAAGATGCACATTCTCCAT - 3')[29]; *bmp2b* MO (5' -

ACCACGGCGACCATGATCAGTCAGT - 3')[50]; *bmp4* MO (5' -

AACAGTCCATGTTTGTCGAGAGGTG - 3')[51]; *bmp7a* MO (5' -

GCACTGGAAACATTTTTAGAGTCAT - 3')[50]; *tcf3b* MO (5' -

CGCCTCCGTTAAGCTGCGGCATGTT - 3')[52]. For each morpholino, a 1 nl solution was injected

into one-cell stage embryos at the specified concentrations: 0.5 ng *tnnt2a* MO, 2 ng *ift88* MO, 0.5 ng

bmp2b MO, 2 ng *bmp4* MO, 4 ng *bmp7a* MO, and 1 ng *tcf3b* MO.

Small molecules treatment

To inhibit cardiac contraction, embryos were incubated in 1 mg/mL tricaine (Sigma, A5040) or 10 uM

blebbistatin (MedChemExpress, HY13441) PTU-added egg water for 12-24 hours. In order to inhibit

erbb2 signaling pathway, 10 uM AG1478 (Sigma, 658552) were used to treat 4 dpf larvae. To inhibit

BMP signaling pathway, 10 uM Dorsomorphin (Sigma, P5499) was used to treat 24 and 36 hpf

embryos.

In situ **hybridization**

Whole mount *in situ* hybridization was performed as previously described[34]. The probes were

synthesized using the DIG RNA labeling kit (Roche). The primers used for obtaining the *id2b* probe

template were as follows: Forward 5' - ATGAAGGCAGTCAGTCCGGTGAGGT - 3'; Reverse 5' -

TCAACGAGACAGGGCTATGAGGTCA - 3'.

Embryonic heart isolation and RNA-seq analysis

Hearts were isolated from embryos carrying the *Tg*(*myl7: mCherry*) transgene following an established

protocol[26]. A minimum of 1,000 hearts for each experimental group were manually collected under a

Leica M165FC fluorescence stereomicroscope and transferred into ice-cold PBS buffer. After

centrifugation at 12, 000 g for 2 min at 4°C, the supernatant was removed, and hearts were lysed in

cold Trizol buffer (Ambion, 15596). Total RNA was extracted for subsequent quantitative real-time

PCR or RNA-seq analysis.

Duplicate samples from control and Tricaine-treated embryonic hearts underwent RNA-seq. Raw

sequencing reads were preprocessed to remove adapters and filter low-quality reads. Clean sequencing

reads were then mapped to the zebrafish reference[53] using STAR with default parameters[54].

Subsequently, gene quantification was carried out with RSEM[55]. The gene expected count was applied

to identify differential expression genes (DEGs), retaining only genes with counts per million (CPM) of

10 in at least two samples. DESeq2[56] was employed for differential expression analysis, and *P*-values

were adjusted using BH correction. DEGs were defined as those with $|\log_2 \text{fold change}| \geq 0.585$ and an

adjusted *P*-value < 0.1. The primary focus was on genes related to transcription regulation, and gene

enrichment analysis was conducted using ClusterProfiler[57]. To analyze DEGs in *id2b*[-/-] and control

embryonic hearts, we performed enrichment analysis with the R package EnrichR, dissecting the

potential anatomy expression pattern and underlying phenotypes. We mainly focused on genes with the

heart related phenotypes, including cardiac muscle tissue development, cardiomyocyte differentiation,

heart morphogenesis and cardiac chamber development. All the analysis on identifying DEGs were

batch corrected.

Quantitative real-time PCR analysis

After extraction from isolated embryonic hearts, 50 ng-1 ug of mRNA was reverse transcribed to

cDNA using the PrimeScript RT Master Mix kit (Takara, RR036A). Real-time PCR was performed

using the TB Green Premix Ex Taq kit (Takara, RR420A) on the Roche LightCycler 480. Expression

levels of the target genes were normalized to *actb1* as an internal control. All experiments were

repeated three times. The following primer sets were used: *id2b* Forward 5' -

ACCTTCAGATCGCACTGGAC - 3', Reverse 5' - CTCCACGACCGAAACACCATT - 3'; *nrg1*

Forward 5' - CTGCATCATGGCTGAGGTGA - 3', Reverse 5' - TTAACTTCGGTTCCGCTTGC - 3';

cacnα1c Forward 5' - GCCCTTATTGTAGTGGGTAGTG - 3', Reverse 5' -

AGTGTTTTGGAGGCCCATTG - 3'; *tcf3a* Forward 5' - CCTCCGGTCATGAGCAACTT - 3',

Reverse 5' - TTTCCCATGATGCCTTCGCT - 3'; *tcf3b* Forward 5' - CCTTTAATGCGCCGTGCTTC

- 3', Reverse 5' - GCGTTCTTCCATTCCTGTACCA - 3'; *socs1a* Forward 5' -

TCAGCCTGACAGGAAGCAAG - 3', Reverse 5' - GTTGCACAGGGATGCAGTCG - 3'; *socs3b*

Forward 5' - GGGACAGTGAGTTCCTCCAA - 3', Reverse 5' - ATGGGAGCATCGTACTCCTG -

3'; *actb1* Forward 5' - ACCACGGCCGAAAGAGAAAT - 3', Reverse 5' -

GCAAGATTCCATACCCAGGA - 3'.

Co-immunoprecipitation (Co-IP) and western blot

Zebrafish *tcf3b* and *id2b* were overexpressed in 293T cell for 48 hours. The transfected cells were then

collected and lysed using IP lysis buffer (Sangon Biotech, C500035) containing protease and

phosphatase inhibitors (Sangon Biotech, C510009, C500017). For the IP experiment, anti-Flag

antibody (Cell Signaling Technology, 14793, 1:100) was incubated with cell lysates overnight at 4°C.

Pretreated magnetic beads were bound with the antigen-antibody complex for 4 hours at 4°C, followed

by washing with IP lysis buffer three times. For western blot, samples were denatured at 95°C for 10

min, separated on a 5%-12% gradient gel. Proteins were then transferred to a PVDF membrane (Sigma,

ISEQ00010). The membrane was blocked for 1 hour with 5% nonfat milk or 5% BSA (Sangon Biotech)

dissolved in TBST and then incubated with primary antibodies (anti-FLAG, Cell Signaling Technology,

14793, 1:1000; anti-HA, Sigma, H3663, 1:1000) overnight at 4°C, followed by three 10-min TBST

washes. HRP-conjugated secondary antibodies (Invitrogen, 31430, 31460) were incubated for 1 hour at

room temperature, followed by three 10-min TBST washes. The detection of immunoreactive bands

was performed with a chemiluminescent substrate (Thermo Scientific, 34577) and imaged using the

Azure Biosystems 400.

Immunofluorescence

For immunofluorescence on adult zebrafish hearts, we fixed the hearts overnight in 4%

paraformaldehyde at 4 °C, followed by equilibration through 15% and 30% sucrose in PBS solution.

The hearts were embedded and frozen in O.C.T. compound (Epredia, 6502), and 10 μm thick

cryosections were prepared using a CryoStar NX50 cryostat. Immunofluorescence experiments were

performed as previously described[58]. For immunofluorescence on embryonic hearts, embryos were

fixed overnight in 4% paraformaldehyde at 4°C, washed twice quickly in 100% methanol, and then

dehydrated overnight at -20°C in 100% methanol. Subsequently, rehydration was performed through a

methanol gradient (100%, 75%, 50%, 25%, 10 min each), followed by three washes in PBST (1%

PBS/0.1% Triton-X100, 10 min each). The embryos were treated with 10 μg/mL proteinase K diluted

in PBST for 20 min at room temperature, re-fixed in 4% paraformaldehyde for 20 min, washed in

PBST, and immersed in blocking solution (PBST/1% BSA/2% goat serum) for 1 hour at room

temperature. Following this, the embryos were incubated in the primary antibody diluted in blocking

solution overnight at 4°C. After washing in PBST, they were incubated in the secondary antibody

(1:200) for 2 hours at room temperature. The primary antibody used was: Anti-GFP antibody (Santa

Cruz Biotechnology, sc9996, 1: 200), anti-α-actinin antibody (Sigma, A7811, 1: 200). The secondary

antibody used was: Anti-mouse IgG-Alexa 488 (Invitrogen, A11011, 1: 400). DAPI was used to stain

cell nuclei.

Cardiac function analysis

To assess cardiac function in embryonic hearts, embryos were embedded in 1% low melting agarose,

and heart contractions were recorded for 1 min using a Nikon Ti2 microscope at a rate of 25 frames per

second. Fractional shortening and heart rate were measured as described previously[34]. For cardiac

contractile functions in adulthood, zebrafish were fixed on a sponge soaked with system water with the

belly facing up and echocardiography was performed[59]. Videos and images in color Doppler mode and

B-mode were obtained using the Vevo1100 imaging system at a frequency of 50 MHz. Nikon NIS

elements AR analysis and Image J software were employed for data extraction. To evaluate AV valve

function, the ratio of inflow and outflow area in the same frame was quantified[32].

Calcium imaging

At 120 hpf, embryos were treated with 10 mM 2,3-butanedione monoxime (BDM, Sigma, B0753) and

mounted in 1% low melting agarose. Time-lapse images were acquired using a Nikon Ti2 microscope

at a rate of 50 frames per second. Data were analyzed using Nikon NIS elements AR analysis software.

Intracellular action potential recording

Electrophysiology study was performed on adult zebrafish ventricles as previously described[34]. Briefly,

hearts were mounted in a chamber containing Tyrode's solution: NaCl 150 mM, KCl 5.4 mM, $MgSO_4$

1.5 mM, NaH_2PO_4 0.4 mM, $CaCl_2$ 2 mM, Glucose 10 mM, HEPES 10 mM, pH was adjusted to 7.4.

Glass pipettes with tip resistance 30-40 MΩ were filled with 3M KCl solution. Intracellular action

potentials were recorded using an HEKA amplifier and pClamp10.3 software (Molecular Devices).

Histology and HE staining

Adult hearts were dissected and fixed overnight at 4°C in 4% PFA, followed by three PBS washes.

Dehydration involved an ethanol gradient (70%, 80%, 95%, 100%, 100%, 30 min each), followed by

three soaks in dimethylbenzene at 65°C, before embedding in paraffin. Sections of 5 um thickness were

prepared using the Leica RM2235 manual rotary microtome for hematoxylin and eosin (HE) staining.

Injection of mRNA

The embryonic zebrafish cDNA library was used as a template to amplify the *nrg1* and *tcf3b* fragment,

which was then subcloned into the pCS2 vector. The vector was linearized using Not I restriction

endonuclease, and mRNA was transcribed in vitro using the mRNA transcription kit (Ambion,

AM1340). 100 pg of purified mRNA was injected into one-cell stage embryos.

Luciferase assay

The LCR (luciferase reporter) plasmid was generated by subcloning the 5′ UTR of *nrg1* into the

upstream region of renilla luciferase on the psiCheck2 plasmid. Following construction, 25 pg of the

LCR plasmid was co-injected with either 100 pg of *tcf3b* mRNA or 1 ng of *tcf3b* MO into 1-cell stage

zebrafish embryos. At 48 hpf, twenty embryos were gathered into one group and fully lysed.

Subsequently, firefly and renilla luciferase activities were sequentially measured using a microplate

reader with the dual luciferase reporter gene assay kit (Yeasen, 11402ES60), according to the

manufacturer′s instructions. The experiment was independently replicated three times. The relative

renilla luciferase activity, normalized by firefly luciferase activity, served as an indicator of *nrg1*

expression level under the influence of *tcf3b* overexpression or reduction.

Image processing and statistical analysis

Whole mount *in situ* hybridization images were captured with a Leica M165FC stereomicroscope. Live

imaging of zebrafish embryos involved mounting anesthetized embryos in 1% low melting agarose

(Sangon Biotech, A600015) and manually oriented them for optimal visual access to the heart.

Confocal images were obtained with a Nikon Ti2 confocal microscope. Fluorescence intensity and cell

number counting were processed using Nikon NIS Elements AR analysis and Image J software.

Statistical analysis was performed using Graphpad prism 8 software. No statistical methods were used

to predetermine sample size. Unpaired two-tailed Student's t-tests was used to determine statistical

significance. Data are presented as mean $\pm$ s.e.m, $*P < 0.05$ was considered to be statistically

significant.

Acknowledgements: We thank Dr. Pengfei Xu for providing morpholinos. We also thank Dr. Jia Li for

the support in generating the *id2b:eGFP* line.

Author Contributions: Shuo Chen and Peidong Han contributed to the study design. Shuo Chen,

Jinxiu Liang, Weijia Zhang, Peijun Jiang, Wenyuan Wang, Xiaoying Chen and Yuanhong Zhou

conducted experiments. Jie Yin analyzed RNA-sequencing data. Peng Xia, Fan Yang, Ying Gu, and

Ruilin Zhang provided key reagents and analytic tool. Shuo Chen, Jie Yin, and Peidong Han prepared

the manuscript.

Source of Funding: This work was supported by the National Key R&D Program of China

(2018YFA0800102, 2023YFA1800602, 2018YFA0800501), and the National Natural Science

Foundation of China (32170823, 31871462).

Disclosures: The authors declare that they have no conflict of interest.

555   **References**

556   1   Duchemin, A. L., Vignes, H., Vermot, J. & Chow, R. Mechanotransduction in
557   cardiovascular morphogenesis and tissue engineering. *Curr Opin Genet Dev* **57**, 106-
558   116 (2019).
559   2   Sidhwani, P. & Yelon, D. Fluid forces shape the embryonic heart: Insights from
560   zebrafish. *Curr Top Dev Biol* **132**, 395-416 (2019).
561   3   Vermot, J. *et al.* Reversing blood flows act through klf2a to ensure normal
562   valvulogenesis in the developing heart. *PLoS Biol* **7**, e1000246 (2009).
563   4   Heckel, E. *et al.* Oscillatory Flow Modulates Mechanosensitive klf2a Expression
564   through trpv4 and trpp2 during Heart Valve Development. *Curr Biol* **25**, 1354-1361
565   (2015).
566   5   Galvez-Santisteban, M. *et al.* Hemodynamic-mediated endocardial signaling controls
567   in vivo myocardial reprogramming. *Elife* **8** (2019).
568   6   Fukui, H. *et al.* Bioelectric signaling and the control of cardiac cell identity in
569   response to mechanical forces. *Science* **374**, 351-354 (2021).
570   7   Dietrich, A. C., Lombardo, V. A., Veerkamp, J., Priller, F. & Abdelilah-Seyfried, S.
571   Blood flow and Bmp signaling control endocardial chamber morphogenesis. *Dev Cell*
572   **30**, 367-377 (2014).
573   8   Sidhwani, P. *et al.* Cardiac function modulates endocardial cell dynamics to shape
574   the cardiac outflow tract. *Development* **147** (2020).
575   9   Staudt, D. W. *et al.* High-resolution imaging of cardiomyocyte behavior reveals two
576   distinct steps in ventricular trabeculation. *Development* **141**, 585-593 (2014).
577   10   Peshkovsky, C., Totong, R. & Yelon, D. Dependence of cardiac trabeculation on
578   neuregulin signaling and blood flow in zebrafish. *Dev Dyn* **240**, 446-456 (2011).
579   11   Auman, H. J. *et al.* Functional modulation of cardiac form through regionally
580   confined cell shape changes. *PLoS Biol* **5**, e53 (2007).
581   12   Lin, Y. F., Swinburne, I. & Yelon, D. Multiple influences of blood flow on
582   cardiomyocyte hypertrophy in the embryonic zebrafish heart. *Dev Biol* **362**, 242-253
583   (2012).
584   13   Gassmann, M. *et al.* Aberrant neural and cardiac development in mice lacking the
585   ErbB4 neuregulin receptor. *Nature* **378**, 390-394 (1995).
586   14   Lee, K. F. *et al.* Requirement for neuregulin receptor erbB2 in neural and cardiac
587   development. *Nature* **378**, 394-398 (1995).
588   15   Meyer, D. & Birchmeier, C. Multiple essential functions of neuregulin in
589   development. *Nature* **378**, 386-390 (1995).
590   16   Lai, D. *et al.* Neuregulin 1 sustains the gene regulatory network in both trabecular
591   and nontrabecular myocardium. *Circ Res* **107**, 715-727 (2010).
592   17   Rentschler, S. *et al.* Neuregulin-1 promotes formation of the murine cardiac
593   conduction system. *Proc Natl Acad Sci U S A* **99**, 10464-10469 (2002).
594   18   Liu, J. *et al.* A dual role for ErbB2 signaling in cardiac trabeculation. *Development* **137**,
595   3867-3875 (2010).

19 Rasouli, S. J. & Stainier, D. Y. R. Regulation of cardiomyocyte behavior in zebrafish
trabeculation by Neuregulin 2a signaling. *Nat Commun* **8**, 15281 (2017).

20 Han, P. *et al.* Coordinating cardiomyocyte interactions to direct ventricular chamber
morphogenesis. *Nature* **534**, 700-704 (2016).

21 Guo, Y. & Pu, W. T. Cardiomyocyte Maturation: New Phase in Development. *Circ Res*
126, 1086-1106 (2020).

22 Alvarez-Dominguez, J. R. & Melton, D. A. Cell maturation: Hallmarks, triggers, and
manipulation. *Cell* **185**, 235-249 (2022).

23 Bers, D. M. Cardiac excitation-contraction coupling. *Nature* **415**, 198-205 (2002).

24 Rottbauer, W. *et al.* Growth and function of the embryonic heart depend upon the
cardiac-specific L-type calcium channel alpha1 subunit. *Dev Cell* **1**, 265-275 (2001).

25 Samsa, L. A. *et al.* Cardiac contraction activates endocardial Notch signaling to
modulate chamber maturation in zebrafish. *Development* **142**, 4080-4091 (2015).

26 Burns, C. G. & MacRae, C. A. Purification of hearts from zebrafish embryos.
Biotechniques **40**, 274, 276, 278 passim (2006).

27 Li, J. *et al.* Intron targeting-mediated and endogenous gene integrity-maintaining
knockin in zebrafish using the CRISPR/Cas9 system. *Cell Res* **25**, 634-637 (2015).

28 Forster, D. *et al.* Genetic targeting and anatomical registration of neuronal
populations in the zebrafish brain with a new set of BAC transgenic tools. *Sci Rep* **7**,
5230 (2017).

29 Li, X. *et al.* Primary cilia mediate Klf2-dependant Notch activation in regenerating
heart. *Protein Cell* **11**, 433-445 (2020).

30 Nauli, S. M. *et al.* Endothelial cilia are fluid shear sensors that regulate calcium
signaling and nitric oxide production through polycystin-1. *Circulation* **117**, 1161-
1171 (2008).

31 Rasouli, S. J. *et al.* The flow responsive transcription factor Klf2 is required for
myocardial wall integrity by modulating Fgf signaling. *Elife* **7** (2018).

32 Gunawan, F., Gentile, A., Gauvrit, S., Stainier, D. Y. R. & Bensimon-Brito, A. Nfatc1
Promotes Interstitial Cell Formation During Cardiac Valve Development in Zebrafish.
Circ Res **126**, 968-984 (2020).

33 Novodvorsky, P. *et al.* klf2ash317 Mutant Zebrafish Do Not Recapitulate Morpholino-
Induced Vascular and Haematopoietic Phenotypes. *PLoS One* **10**, e0141611 (2015).

34 Zhang, R. *et al.* In vivo cardiac reprogramming contributes to zebrafish heart
regeneration. *Nature* **498**, 497-501 (2013).

35 Jiang, M. *et al.* Characterization of the Zebrafish Cell Landscape at Single-Cell
Resolution. *Front Cell Dev Biol* **9**, 743421 (2021).

36 Wang, X. *et al.* Neuregulin-1beta Partially Improves Cardiac Function in Volume-
Overload Heart Failure Through Regulation of Abnormal Calcium Handling. *Front
Pharmacol* **10**, 616 (2019).

37 Moskowitz, I. P. *et al.* Transcription factor genes Smad4 and Gata4 cooperatively
regulate cardiac valve development. [corrected]. *Proc Natl Acad Sci U S A* **108**, 4006-
4011 (2011).

38 Jongbloed, M. R. *et al.* Expression of Id2 in the second heart field and cardiac defects
in Id2 knock-out mice. *Dev Dyn* **240**, 2561-2577 (2011).

39 Li, J. *et al.* Piezo1 integration of vascular architecture with physiological force.
Nature **515**, 279-282 (2014).

40 Nonomura, K. *et al.* Mechanically activated ion channel PIEZO1 is required for
lymphatic valve formation. *Proc Natl Acad Sci U S A* **115**, 12817-12822 (2018).

41 Brown, D. *et al.* Neuregulin-1 is essential for nerve plexus formation during cardiac
maturation. *J Cell Mol Med* **22**, 2007-2017 (2018).

42 Gemberling, M., Karra, R., Dickson, A. L. & Poss, K. D. Nrg1 is an injury-induced
cardiomyocyte mitogen for the endogenous heart regeneration program in zebrafish.
Elife **4** (2015).

43 Hui, S. P. *et al.* Zebrafish Regulatory T Cells Mediate Organ-Specific Regenerative
Programs. *Dev Cell* **43**, 659-672 e655 (2017).

44 Palencia-Desai, S. *et al.* Vascular endothelial and endocardial progenitors
differentiate as cardiomyocytes in the absence of Etsrp/Etv2 function. *Development*
138, 4721-4732 (2011).

45 Schumacher, J. A., Bloomekatz, J., Garavito-Aguilar, Z. V. & Yelon, D. tal1 Regulates
the formation of intercellular junctions and the maintenance of identity in the
endocardium. *Dev Biol* **383**, 214-226 (2013).

46 Chi, N. C. *et al.* Foxn4 directly regulates tbx2b expression and atrioventricular canal
formation. *Genes Dev* **22**, 734-739 (2008).

47 Roman, B. L. *et al.* Disruption of acvrl1 increases endothelial cell number in zebrafish
cranial vessels. *Development* **129**, 3009-3019 (2002).

48 Collery, R. F. & Link, B. A. Dynamic smad-mediated BMP signaling revealed through
transgenic zebrafish. *Dev Dyn* **240**, 712-722 (2011).

49 Sehnert, A. J. *et al.* Cardiac troponin T is essential in sarcomere assembly and cardiac
contractility. *Nat Genet* **31**, 106-110 (2002).

50 Lele, Z., Bakkers, J. & Hammerschmidt, M. Morpholino phenocopies of the swirl,
snailhouse, somitabun, minifin, silberblick, and pipetail mutations. *Genesis* **30**, 190-
194 (2001).

51 Weber, S. *et al.* SIX2 and BMP4 mutations associate with anomalous kidney
development. *J Am Soc Nephrol* **19**, 891-903 (2008).

52 Dorsky, R. I., Itoh, M., Moon, R. T. & Chitnis, A. Two tcf3 genes cooperate to pattern
the zebrafish brain. *Development* **130**, 1937-1947 (2003).

53 Weinberger, M., Simões, F. C., Patient, R., Sauka-Spengler, T. & Riley, P. R. Functional
Heterogeneity within the Developing Zebrafish Epicardium. *Dev Cell* **52**, 574-
590.e576 (2020).

54 Dobin, A. *et al.* STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* **29**, 15-21
(2013).

55 Li, B. & Dewey, C. N. RSEM: accurate transcript quantification from RNA-Seq data
with or without a reference genome. *BMC Bioinformatics* **12**, 323 (2011).

56 Love, M. I., Huber, W. & Anders, S. Moderated estimation of fold change and
dispersion for RNA-seq data with DESeq2. *Genome Biology* **15**, 550 (2014).

57   Wu, T. *et al.* clusterProfiler 4.0: A universal enrichment tool for interpreting omics

data. *The Innovation* **2**, 100141 (2021).

58   Han, P. *et al.* Hydrogen peroxide primes heart regeneration with a derepression

mechanism. *Cell Res* **24**, 1091-1107 (2014).

59   Wang, L. W. *et al.* Standardized echocardiographic assessment of cardiac function in

normal adult zebrafish and heart disease models. *Dis Model Mech* **10**, 63-76 (2017).

Figure legends

Figure 1. Identification of *id2b* as a blood flow sensitive gene.

(**A**) Schematic showing the experimental procedures, including treatment, heart collection, and RNA-

sequencing of zebrafish embryonic hearts. (**B**) KEGG enrichment analysis depicting differentially

expressed genes encoding transcription factors and transcriptional regulators between control (ctrl) and

tricaine-treated embryonic hearts. Red and blue rectangles represent up-regulated and down-regulated

gene sets, respectively. |$\log_2$fold change|≥0.585, adjusted *p*-value<0.1. Each replicate contains

approximately 1,000 hearts. (**C**) Heat map exhibiting representative genes from KEGG pathways

mentioned in (**B**). (**D**) qRT-PCR analysis of *id2b* mRNA in control and tricaine-treated embryonic

hearts. Data were normalized to the expression of *actb1*. Each sample contains ~1,000 embryonic

hearts. (**E**) *In situ* hybridization of *id2b* in 72 hpf and 96 hpf ctrl, tricaine (1 mg/mL), and blebbistatin

(10 μM)-treated embryos. Numbers at the bottom of each panel indicate the ratio of representative

staining. (**F**) *In situ* hybridization showing reduced *id2b* expression in *tnnt2a* morpholino-injected

embryos at 72 hpf compared to control. ∗∗∗ *p*<0.001. Scale bars: 50 μm.

Figure 2. The spatiotemporal expression of *id2b*.

(**A**) Schematic of the intron targeting-mediated *eGFP* knockin at the *id2b* locus using the CRISPR-

Cas9 system. The sgRNA targeting sequence and the protospacer adjacent motif (PAM) sequence are

shown in orange and blue, respectively. The donor plasmid comprises left and right arm sequences and

a *linker-FLAG-P2A-eGFP* cassette denoted by black lines with double arrows and green box,

respectively. The *linker-FLAG-P2A-eGFP* cassette was integrated into the *id2b* locus upon co-injection

of the donor plasmid with sgRNA and zCas9 protein, enabling detection by PCR using two pairs of

primers (F1, R1 and F2, R2)-the former length yielding a length of about 2.2 kb and the latter about

2.7 kb. (**B**) Zebrafish *id2b* expression pattern, as indicated by in situ hybridization of embryos at

designated time points, was consistent with the fluorescence localization of *id2b:eGFP*, revealing

expression in the heart, brain, retina, notochord, pronephric duct, and other tissues. (**C**) Three

dimensional (3D) reconstructions (top) and confocal sections (bottom) of *id2b:eGFP;*

Tg(myl7:mCherry) hearts at designated time points. (**D**) 3D reconstructions (top) and confocal sections

(bottom) of *id2b:eGFP; Tg(kdrl:mCherry)* embryos at specific time points. Magenta: *id2b:eGFP*;

yellow: *kdrl:mCherry*. (**E**) Confocal z-stack maximal intensity projection of

id2b:eGFP;Tg(kdrl:mCherry) embryos at 96 hpf showing the whole body (lateral view) and the head

(top view). (**F**) Immunofluorescence of adult *id2b:eGFP; Tg(kdrl:mCherry)* heart section (middle

panel). Enlarged views of boxed areas are shown in the left and right panels. Green: eGFP; red:

mCherry; blue, DAPI. Scale bars: 500 μm (**B**, **E**, left and **F**); 100 μm (**E**, right); 50 μm (**C** and **D**).

Figure 3. Cardiac contraction promotes endocardial *id2b* expression through primary cilia.

(**A**) Representative confocal z-stack (maximal intensity projection) of *id2b:eGFP* embryos under

different conditions: control, tricaine-treated, blebbistatin-treated, and *tnnt2a* morpholino-injected. (**B**)

Quantification of mean fluorescence intensity (MFI) of *id2b:eGFP* in the working myocardium (atrium

and ventricle, A+V) and atrioventricular canal (AVC) in (**A**). Data normalized to the mean fluorescence

intensity of control hearts. (**C**) Representative confocal z-stack (maximal intensity projection) of

control and *ift88* morpholino-injected *id2b:eGFP* embryos. (**D**) Normalized MFI of *id2b:eGFP* in the

working myocardium (A+V) and AVC in (**C**). (**E**) Whole-mount *in situ* hybridization showing *id2b*

expression in control, *klf2a*[-/-]*,* and *klf2b*[-/-] embryos at 48 hpf and 72 hpf. Numbers at the bottom of each

panel indicate the ratio of representative staining. Data are presented as mean ± sem. ** $p<0.01$, ***

$p<0.001$, **** $p<0.0001$. Scale bars: 50 μm.

Figure 4. *id2b*[-/-] adults exhibit thinner atrioventricular valve leaflets and prominent retrograde

blood flow.

(**A**) Two sgRNAs, represented by short vertical lines, were designed to create *id2b*[-/-] mutants. Co-

injection of the two sgRNAs with zCas9 protein induces a 157 bp truncation in the exon 1 of *id2b*,

which can be detected by genotyping primers marked with arrows. This genetic modification leads to

the formation of a premature stop codon and the subsequent loss of the HLH domain. (**B**) No

discernible morphological differences were observed between $id2b^{+/+}$ and $id2b^{-/-}$ larvae at both 72 hpf

and 96 hpf. (**C**) Kaplan-Meier survival curve analysis and logrank test of $id2b^{+/+}$ (n=50) and $id2b^{-/-}$

(n=46). Wpf: weeks post-fertilization. (**D**) Pericardial edema and an enlarged atrium are evident in a

subset of $id2b^{-/-}$ adults. V, ventricle; A, atrium. (**E**) $id2b^{-/-}$ adults developed thinner atrioventricular

valve leaflets (denoted by arrowheads) compared to $id2b^{+/+}$. Enlarged views of boxed areas are shown

in the right panels. (**F**, **G**) Echocardiograms of adult $id2b^{+/+}$ (**F**) and $id2b^{-/-}$ (**G**) hearts. Unidirectional

blood flow was observed in the $id2b^{+/+}$ heart, while retrograde blood flow was evident in the $id2b^{-/-}$

heart. (**H**) Ratio of retrograde flow area over inflow area shows a significant increase in retrograde flow

in $id2b^{-/-}$ (n=13) compared to $id2b^{+/+}$ (n=10). Data are presented as mean ± sem. ** $p<0.01$. Scale bars:

200 μm (**E**); 500 μm (**B** and **D**, bottom); 2 mm (**D**, top).

Figure 5. Reduced cardiac contractile function and compromised calcium handling in $id2b^{-/-}$

mutants.

(**A**) Time-lapse imaging (from T1 to T8) illustrates the cardiac contraction-relaxation cycle of 120 hpf

$id2b^{+/+}$ and $id2b^{-/-}$ hearts carrying *myl7:mCherry*. (**B** and **C**) $id2b^{-/-}$ larvae display a significant decrease

in heart rate and fractional area change compared to $id2b^{+/+}$. (**D**) Echocardiograms of adult $id2b^{+/+}$ and

$id2b^{-/-}$ hearts. (**E** and **F**) $id2b^{-/-}$ fish exhibit reduced cardiac contractile function with preserved heart

rate compared to $id2b^{+/+}$. (**G**) Time-lapse imaging illustrates the calcium dynamics of 120 hpf $id2b^{+/+}$

and $id2b^{-/-}$ hearts carrying *actb2:GCaMP6s*. (**H**) Ratio of maximal fluorescence intensity (F) over basal

fluorescence intensity (F0) of GCaMP6s signal. (**I**) qRT-PCR analysis of *cacnα1c* mRNA in $id2b^{+/+}$

and $id2b^{-/-}$ hearts at 120 hpf (N=three replicates, with each sample containing 500-1,000 embryonic

hearts) and adult stage (N=five replicates). Data were normalized to the expression of *actb1*. (**J**) The

action potential of ventricular cardiomyocyte in adult $id2b^{+/+}$ and $id2b^{-/-}$ hearts. (**K**) Statistical data

showed a notable difference between $id2b^{+/+}$ and $id2b^{-/-}$ accordingly. Data are presented as mean ± sem.

*** $p<0.001$, **** $p<0.0001$, ns, not significant. Scale bar: 50 μm.

Figure 6. *nrg1* serves as a pivotal mitogen mediating the function of *id2b*.

(**A**) Identification of genes (*fgf8a*, *nkx2.5*, *myh6*, *nrg1*, and *nkx2.7*) associated with four distinct heart

development processes: cardiac muscle tissue development, cardiomyocyte differentiation, heart

morphogenesis, and cardiac chamber development. The heatmap illustrates scaled-normalized

expression values for the mentioned genes. (**B**) qRT-PCR analysis of *nrg1* mRNA in 120 hpf *id2b*$^{+/+}$

and *id2b*$^{-/-}$ embryonic hearts. Data were normalized to the expression of *actb1*. N=four replicates, with

each sample containing 500-1,000 embryonic hearts. (**C**) *id2b*$^{+/+}$ and *id2b*$^{-/-}$ larvae were injected with

nrg1 mRNA at the 1-cell stage, followed by qRT-PCR analysis of *cacnα1c* mRNA at 72 hpf. Data

were normalized to the expression of *actb1*. N=three replicates, with each samples containing 100-200

embryonic hearts. (**D**) The heart rate of 72 hpf *id2b*$^{+/+}$ and *id2b*$^{-/-}$ larvae injected with *nrg1* mRNA at 1-

cell stage. (**E**) Heart rate in 120 hpf larvae treated with AG1478 and DMSO. Data are presented as

mean ± sem. * $p<0.05$, ** $p<0.01$, *** $p<0.001$, **** $p<0.0001$. ns, not significant.

Figure 7. *id2b* interacts with *tcf3b* to restrict its inhibition on *nrg1* expression.

(**A**) Immunoprecipitation (IP) assays of FLAG-*id2b* and HA-*tcf3b* co-transfected 293T cells. (**B**) qRT-

PCR analysis of *tcf3a*, *tcf3b*, *socs1a*, and *socs3b* mRNA in 120 hpf *id2b*$^{+/+}$ and *id2b*$^{-/-}$ embryonic hearts.

Data were normalized to the expression of *actb1*. N=three replicates, with each samples containing

500-1,000 embryonic hearts. (**C**) Two potential *tcf3*b binding sites, with sequences corresponding to

the human tcf3 (left) and mouse tcf3 (right) binding motifs, were predicted in the 2,000bp DNA

sequence upstream of the zebrafish *nrg1* transcription start site using JASPAR. (**D**) Luciferase assay

showing the expression of *nrg1* in embryos with *tcf3b* overexpression (*tcf3b* OE) and morpholino-

mediated *tcf3b* knockdown (*tcf3b* MO). N=three replicates. (**E**) qRT-PCR analysis of *nrg1* mRNA in

72 hpf *id2b*$^{+/+}$ and *id2b*$^{-/-}$ embryonic hearts injected with control and *tcf3b* morpholino. Data were

normalized to the expression of *actb1*. N=three replicates, with each samples containing 100-200

embryonic hearts. (**F**) Schematic model for *id2b*-mediated regulation of myocardium function. During

heart development, blood flow operates through primary cilia, initiating endocardial *id2b* expression.

Subsequently, the interaction between *id2b* and *tcf3b* restricts the activity of *tcf3b*, ensuring proper

nrg1 expression, which in turn promotes LTCC expression (left). However, in the absence of *id2b*,

tcf3b inhibits *nrg1* expression. The reduced *nrg1* hinders LTCC expression in cardiomyocytes,

resulting in decreased extracellular calcium entry and disruption of myocardial function. Data are

presented as mean ± sem. * $p<0.05$, ** $p<0.01$, *** $p<0.001$, **** $p<0.0001$. ns, not significant.

Figure 3-figure supplement 1. Blood flow and BMP signaling independently activates *id2b*

expression.

(**a**) Representative confocal maximal intensity projection of control (non-injected), *bmp2b*, *bmp4*, and

bmp7a morpholino-injected *id2b:eGFP*; *Tg(myl7:mCherry)* hearts at 72 hpf. (**b**) Quantification of

mean fluorescence intensity of *id2b:eGFP* in (**a**). Data normalized to the mean fluorescence intensity

of control hearts. (**c**) Representative confocal maximal intensity projection of DMSO and

Dorsomorphin (DM)-treated *id2b:eGFP* hearts at 48 and 60 hpf. Embryos were treated from 24 to 48

hpf, or from 36 to 60 hpf. (**d**) Quantification of mean fluorescence intensity of *id2b:eGFP* in (**c**). Data

normalized to the mean fluorescence intensity of DMSO-treated hearts. (**e**) Confocal optical sections of

control, tricaine-treated, and *tnnt2a* morpholino-injected 72 hpf *Tg(Bre:d2GFP)*; *Tg(kdrl:mCherry)*

hearts. Yellow arrowheads: endocardial cells. Numbers at the top of each panel indicate the ratio of

representative images. (**f**) Schematic diagram of blood flow and BMP signaling-mediated *id2b*

expression. Data are presented as mean ± sem. *** $p<0.001$, **** $p<0.0001$. ns, not significant.

Figure 4-figure supplement 1. *id2b^{-/-}* larvae exhibit a decreased number of valve endocardial cells

while maintaining normal atrioventricular valve function.

(**A**) Representative confocal images of valve endocardial cells in 96 hpf *id2b^{+/+}* and *id2b^{-/-}* hearts

carrying *Tg(kdrl:nucGFP)*. (**B**) Quantification of the number of valve endocardial cells in *id2b^{+/+}* and

id2b^{-/-} hearts. VECs: valve endocardial cells. (**C**) Representative confocal images of 72 and 120 hpf

id2b^{+/+} and *id2b^{-/-}* hearts in *Tg(kdrl:nucGFP)*; *Tg(myl7:mCherry)* transgenic background. (**D**) Statistical

analysis of the number of endocardial cells in the ventricle (V), atrium (A), and combined (A+V) in

id2b^{+/+} and *id2b^{-/-}* hearts. (**E**) Quantification of blood flow patterns in 96 hpf *id2b^{+/+}* and *id2b^{-/-}* hearts.

Data are presented as mean ± sem. **** $p<0.0001$. ns, not significant. Scale bars: 20 μm (**A**); 50 μm

(**B**).

Figure 5-figure supplement 1. *id2b* **loss-of-function impacts both valve formation and cardiac**

contraction.

(**A**) The top 20 predicted tissue expression patterns based on the differential expressed genes from

id2b$^{+/+}$ and *id2b*$^{-/-}$ hearts at 120 hpf were displayed. (**B**) The top 20 predicted phenotypes related to the

differential expressed genes from *id2b*$^{+/+}$ and *id2b*$^{-/-}$ hearts at 120 hpf were illustrated.

Figure 5-figure supplement 2. *id2b*$^{-/-}$ **hearts develop normal trabeculae and sarcomeres.**

(**A**) Representative confocal z-stack of 72 and 120 hpf *id2b*$^{+/+}$ and *id2b*$^{-/-}$ hearts with *Tg(myl7:H2A-*

mCherry) transgene. (**B**) Quantification of the number of cardiomyocytes in the ventricle (V), atrium

(A), and combined (A+V) in (**A**). (**C**) Representative confocal images of 72 and 120 hpf *id2b*$^{+/+}$ and

id2b$^{-/-}$ hearts carrying *Tg(myl7:H2A-mCherry)*. (**D**) Confocal immunofluorescence images of α-actinin

(green) in embryonic (72 hpf) and adult (115 dpf) hearts (left). Right: fluorescence intensity profiles for

α-actinin. Data are presented as mean ± sem. ns, not significant. Scale bars: 50 μm (**A** and **C**); 5 μm

(**D**).

Figure 6-figure supplement 1. *nrg1* **is expressed in the endocardial cells.**

(**A**) Analysis of *nrg1* expression using the zebrafish single-cell landscape database. The red rectangle

highlights endocardial cells. (**B**) Identification of the endocardial cell population in the zebrafish cell

landscape. Purple dots represent endocardial cells, and the red oval denotes cell populations from the

heart. Images were generated using the Zebrafish Cell Landscape (ZCL) at http://bis.zju.edu.cn/ZCL/.

Figure 7-figure supplement 1. Expression landscape of zebrafish *tcf3b***.**

The major cell types expressing *tcf3b* were displayed according to the zebrafish single-cell landscape.

Images were generated using the Zebrafish Cell Landscape (ZCL) at http://bis.zju.edu.cn/ZCL/.

Video 1. 96 hpf *id2b*$^{+/+}$ **larvae displayed unidirectional blood flow in the AV canal.** Scale bar: 50

μm.

853   **Video 2. 96 hpf *id2b⁻ᐟ⁻* larvae displayed unidirectional blood flow in the AV canal.** Scale bar: 50 μm.

Figure 1

A

sgRNA target   PAM
TCTGTTGAACACATGACGATATTTGGAAG

Endogenous *id2b* locus   stop codon
Exon1 ━━ Exon2
F1   Left arm   Right arm   R2

Left arm   Right arm
Exon2 *linker-Flag-P2A-eGFP*
F2   R1

id2b-linker-Flag-P2A-EGFP donor

+ *sgRNA*
+*zCas9 protein*

Targeted *id2b* locus
2.2kb   2.7kb
Exon1   Exon2 *linker-Flag-P2A-eGFP*   Exon2
F1   R1   F2   R2
plasmid fragment

B   24 hpf   48 hpf   72 hpf

C   48 hpf   72 hpf   96 hpf
id2b:eGFP; myl7: mCherry

D   *id2b:eGFP; kdrl: mCherry*

E   *id2b:eGFP*; *kdrl: mCherry*   96 hpf
id2b:eGFP
kdrl: mCherry

F   DAPI
id2b:eGFP
myl7:mCherry

Figure 2

Figure 6