Evolutionary Insights from the Mitochondrial Genome of *Oikopleura dioica*: Sequencing Challenges, RNA Editing, Gene Transfers to the Nucleus, and tRNA Loss

Yael Klirs[1], Maria Novosolov[1,2], Carmela Gissi[3,4,5], Rade Garic[6], Tal Pupko[7], Thomas Stach[8], Dorothée Huchon[1,9]

[1]School of Zoology, George S. Wise Faculty of Life Sciences, Tel Aviv University, Tel Aviv 6997801, Israel

[2]GeoGenetics Centre, GLOBE Institute, Faculty of Health and Medical Sciences, University of Copenhagen, Copenhagen, Denmark.

[3]Department of Biosciences, Biotechnology and Environment, University of Bari Aldo Moro, 70126 Bari, Italy

[4]Institute of Biomembranes, Bioenergetics and Molecular Biotechnologies, Consiglio Nazionale delle Ricerche, 70126 Bari, Italy

[5]CoNISMa, Consorzio Nazionale Interuniversitario per le Scienze del Mare, 00196 Roma, Italy

[6]Institute for Marine and Coastal Research, University of Dubrovnik, Kneza D. Jude 12, 20000 Dubrovnik, Croatia

[7]The Shmunis School of Biomedicine and Cancer Research, George S. Wise Faculty of Life Sciences, Tel Aviv University, Tel Aviv, 6997801 Israel

[8]Department of Molecular Parasitology, Institute of Biology, Humboldt-Universität zu Berlin, Berlin, Germany

[9]The Steinhardt Museum of Natural History and National Research Center, Tel Aviv University, Tel Aviv 6997801, Israel

ORCID IDs: 0009-0006-2572-0612 (Yael Klirs), 0000-0002-4034-3441 (Maria Novosolov),

0000-0002-2269-079X (Carmela Gissi), 0000-0001-6477-5559 (Rade Garic), 0000-0001-

9463-2575 (Tal Pupko), 0000-0001-5461-9069 (Thomas Stach), 0000-0002-5047-244X

(Dorothée Huchon).

* Author for Correspondence: Dorothée Huchon, School of Zoology, George S. Wise

Faculty of Life Sciences, Tel Aviv University, Tel Aviv 6997801, Israel, +972 36409817,

huchond@tauex.tau.ac.il

Abstract

Sequencing the mitochondrial genome of the tunicate *Oikopleura dioica* is a challenging task due to the presence of long poly-A/T homopolymer stretches, which impair sequencing and assembly. Here, we report on the sequencing and annotation of the majority of the mitochondrial genome of *O. dioica* by means of combining several DNA and amplicon reads obtained by Illumina and MinIon Oxford Nanopore Technologies (ONT) with public RNA sequences. We document extensive RNA editing, since all homopolymer stretches present in the mitochondrial DNA correspond to 6U-regions in the mitochondrial RNA. Out of the 13 canonical protein-coding genes, we were able to detect eight, plus an unassigned ORF that lacked sequence similarity to canonical mitochondrial protein-coding genes. We show that the *nad3* gene has been transferred to the nucleus and acquired a mitochondria-targeting signal. In addition to two very short rRNAs, we could only identify a single tRNA (tRNA-Met), suggesting multiple losses of tRNA genes, supported by a corresponding loss of mitochondrial aminoacyl-tRNA synthetases in the nuclear genome. Based on the eight canonical protein-coding genes identified, we reconstructed maximum likelihood and Bayesian phylogenetic trees and inferred an extreme evolutionary rate of this mitochondrial genome. The phylogenetic position of appendicularians among tunicates, however, could not be accurately determined.

Keywords

mtDNA, phylogenomics, Oxford Nanopore Technologies, poly-A/T homopolymers, Tunicata, Appendicularia.

Significance

Sequencing and annotating the mitochondrial genome of fast-evolving organisms is difficult because they often present unusual characteristics. The tunicate *Oikopleura dioica* is a model species for understanding tunicate and chordate genome evolution. However, no complete annotated mitochondrial genome for this species has been published to date. Here, we determined the major part of the mitochondrial genome of *O. dioica*. Our results indicate the presence of highly modified rRNA genes and the absence of all tRNAs except tRNA-Met. Moreover, we show that the mitochondrial genome undergoes editing at the RNA level. Our study demonstrates that utilizing a combination of public RNA data and DNA from long-and short-read sequencing platforms significantly improves our ability to study mitochondrial genomes with atypical characteristics.

Introduction

Most mitochondrial genomes of the Bilateria are circular DNA molecules of approximately 14-19 kb in length, characterized by the absence of introns and the presence of only short intergenic regions (Lavrov and Pett 2016). These genomes typically contain 37 genes: 13 protein-coding genes: ATP synthase subunits 6 and 8 (*atp6* and *atp8*), cytochrome oxidase subunits (*cox1, cox2, cox3*), cytochrome b (*cytb*), and dehydrogenase subunits (*nad1, nad2, nad3, nad4, nad5, nad6*, and *nad4L*); two ribosomal genes (rRNA): small-subunit (*rns*) and large-subunit (*rnl*) and ~22 tRNAs (Boore 1999; Gissi et al. 2008; Rubinstein et al. 2013). However, bilaterian mitochondrial genomes are substantially diverse in their structure, organization, evolutionary rate, and genetic code (Lavrov 2007).

Tunicate mitochondrial genomes are fast evolving, both in terms of substitutions (Singh et al. 2009) and genome rearrangements (Gissi et al. 2008; Stach et al. 2010). As a result of this high rearrangement rate, there is no standard tunicate mitochondrial gene order. In addition, the ascidian mitochondrial genetic code differs from the vertebrate one. Of note, the "ascidian" genetic code, which was initially determined based on the mitochondrial sequences of *Halocynthia roretzi* and *Pyura stolonifera* (Durrheim et al. 1993; Yokobori et al. 1993), seems to be valid for all other tunicate classes (Pichon et al. 2020).

The subphylum Tunicata is divided into three classes: Ascidiacea, Thaliacea, and Appendicularia. Appendicularia is a key class for understanding tunicate and chordate genome evolution, as it seems to be the sister-clade to all other tunicates. *Oikopleura dioica* is a frequent model tunicate for the class Appendicularia in developmental biology, evolutionary biology, and comparative genomics research (Delsuc et al. 2006; Nishida 2008). Despite the importance of this clade, no complete annotated appendicularian mitochondrial genome is currently available in public databases. One non-annotated mitochondrial genome of *O.*

dioica from Okinawa Japan is available (accession OU015571; 9,225 bp; (Bliznina, et al.

2021)).

The mitochondrial genome of *O. dioica* from Norway has been partially reconstructed

by Denoeud et al. (2010) from genomic DNA and EST data. However, the obtained

mitochondrial contigs of this genome are not publicly available. This genome has been

inferred to harbor long poly-A/T stretches interrupting the open reading frames (ORFs) and it

has been suggested that these interrupted ORFs are corrected by RNA editing (Denoeud et al.

2010). Such homopolymer stretches cannot be sequenced using Sanger sequencing or

Illumina short-read sequencing, since they are error hot spots for the DNA polymerases used

in these sequencing techniques. Thus, together with the difficulty in isolation and

identification of appendicularian species, this peculiarity explains the paucity of publicly

available mitochondrial data on *O. dioica*. Annotation of the resulting genome is also

expected to be challenging, due to RNA editing. RNA editing refers to modifications such as

insertions, deletions, and base substitutions of RNA transcripts. As a result, the RNA and

computationally inferred protein sequences differ from the DNA template (Gott and Emeson

2000). To complicate matters, in *O. dioica* RNA editing seems to be prevalent in those

regions that are hard to sequence, i.e., the poly-A/T stretches. These stretches appear as 6U-

sites in mitochondrial RNA sequences, regardless of the number of T/A at the DNA level

(Denoeud et al. 2010).

The phylogenetic analyses performed by Denoeud et al. (2010) concluded that *O.*

dioica mitochondrial protein-coding genes are very fast evolving, even when compared to the

already high rate of other tunicate mitochondrial genomes. Using a combination of PCR and

EST mapping approaches, they determined the order of nine out of the 13 mitochondrial

protein-coding genes: *cox3, nad1, cox1, cytb,* a putative *nad2*, *atp6, cox2, nad5*, along a large

mitochondrial contig, and *nad4* on a separate contig. They did not identify any tRNA or

rRNA genes. Unfortunately, the mitochondrial contigs obtained by Denoeud et al. (2010)

have not been deposited in public databases and only the raw ESTs data are available in the

Oikobase database (Danks et al. 2012).

A draft mitochondrial genome scaffold of *O. dioica* from Okinawa (Japan) has been

published by Bliznina et al. (2021) (accession OU015571; 9,225 bp), with the annotation

reported only in the publication (Figure 7 of their article). The annotation obtained is,

however, preliminary as several genes contain stop codons and frameshifts. Unlike in

Denoeud et al. (2010), the mitochondrial genome is presented as a circular molecule that

encompasses seven mitochondrial genes (Bliznina et al. 2021). The mitochondrial gene order

resembles that from Figure S4 from Denoeud et al. (2010). However, some differences exist.

First, the *nad2* and *nad5* genes are missing from the annotation of Bliznina et al. (2021).

Second, the *rnl* and four tRNAs: tRNA-Met, tRNA-Trp, tRNA-Ala, and tRNA-His are

annotated. Third, they identified a LAGLIDADG homing endonuclease in the middle of the

cox1 gene (Figure 7 from Bliznina et al. 2021). It should be noted that *O. dioica* from

Okinawa most likely represents a different species from *O. dioica* from Norway (Masunaga et

al. 2022; Plessy et al. 2024).

Here, we determined the major part of the mitochondrial genome sequence of *O.*

dioica from Norway, its gene content, and its order. We observed a substantial gene loss

compared to other metazoan mitochondrial genomes and provide evidence for extensive RNA

editing.

Results

Sequencing the Mitochondrial Genome

DNA extracted from *O. dioica* individuals from Norway was sent for Illumina sequencing. Since it was impossible to assemble these reads due to the prevalence of homopolymers (poly-A or poly-T stretches), we next performed long-read sequencing using Oxford Nanopore Technology (ONT) sequencing. However, in this case, the flow-cell pores became blocked within an hour (Minion sequencing was performed twice by different companies), resulting in too little data for a reliable assembly. Therefore, we PCR-amplified the mitochondrial genome in 12 fragments (supplementary Fig. S1, Supplementary Material online) and sequenced all barcoded fragments using a single ONT flow cell. Primers were designed based on the *O. dioica* EST and RNA data available in public databases (see Materials and Methods and supplementary table S1, Supplementary Material online). We successfully assembled each PCR fragment (except one) and, using the Illumina data, we obtained a single contig. Unfortunately, we could not circularize the genome because we were unable to assemble one fragment completely. The length of the *O. dioica* mitochondrial contig we assembled is 18,390 bp (Figure 1). This genome size is larger than the average of other tunicates, 14,808 ± 821 bp, and compatible with the PCR prediction of a ~19,000 bp genome (supplementary table S1, Supplementary Material online).

Multiple instances of poly-A/T stretches were found in the assembled contig (Figure 1, supplementary Fig. S2, Supplementary Material online). The exact number of either A/T repeats within poly-A/T regions could not be accurately determined. These stretches are too long to be fully contained in the short Illumina sequences (Ari and Arikan 2016), while ONT basecallers do not accurately determine the length of homopolymers (Wick et al. 2019). Moreover, the length of homopolymers has been correctly resolved up to a length of 11 bp in an R10.4 flow cell (Sereika et al. 2022), while in an R9.4.1 flow cell, a good accuracy (70%

errorless) was observed only in homopolymers up to 5 bp (Delahaye and Nicolas 2021).

However, in our case, the length of the 65 poly-A/T stretches found in the mitochondrial

sequence ranged from 2 to 509 bp, with an average of 143 ± 103.5 bp (Materials and

Methods, supplementary table S2, Supplementary Material online), which is considerably

larger. Moreover, it has been observed that regions harboring inverted repeats are poorly

sequenced using ONT (Spealman et al. 2020). When in a PCR-fragment a poly-A region is

followed by a poly-T region, these two homopolymers can form structures similar to inverted

repeats, thus disrupting ONT sequencing. In addition, Taq polymerase slippage errors can

further contribute to uncertainty associated with the length of long homopolymers. All these

factors contribute to heterogeneity among ONT reads that map to the same mitochondrial

locus harboring a poly-A/T region. One such example is shown in Figure 2. Note that these

stretches are sometimes imperfect, i.e., there may be a few G residues within a poly-A stretch

or a few C residues with a poly-T stretch (Figure 2). These G and C residues are unlikely to

reflect ONT sequencing errors, as they could sometimes be verified using Illumina reads. In

total, these poly-A/T stretches span ~9,200 bp. This suggests that while the total

mitochondrial genome size is longer than in other tunicates, the mitochondrial genome

without the poly-A/T stretches is, in fact, shorter.

Evidence of rRNA Editing

A total of 65 poly-A/T sites were found in the assembled *O. dioica* mitochondrial genome. By

mapping (1) the mitochondrial transcripts identified from the EST from Norway, and (2) the

RNA contigs assembled from publicly available RNA reads from mainland Japan (Wang et

al. 2015) and the North Sea (Raghavan et al. 2023) onto the DNA sequence assembled from

the ONT reads, we were able to confirm the editing of 59 of these 65 poly-A/T sites to 6U-

sites (see an example in Figure 2 and supplementary table S2, Supplementary Material

online). Of note, we performed a barcoding analysis that supports the hypothesis that our

DNA sample, the EST data, and the North Sea RNA reads originated from the same species,

while the RNA reads from mainland Japan likely originated from a different species (see

Materials and Methods, Masunaga et al. 2022; Plessy et al. 2024). For the remaining six sites,

no direct RNA evidence for editing was obtained as we could not detect RNA reads or ESTs

that mapped to the corresponding region. These poly-A/T stretches are present both in coding

and noncoding mitochondrial DNA regions. Specifically, we observed that exactly six

consecutive U residues (6U-sites) in the RNA corresponded to a poly-T stretch in the DNA

sequence. These results suggest that the longer U stretches in the pre-mRNA are edited,

resulting in shorter stretches of 6U. The exact nature of this RNA editing is unknown. It is,

however, remarkable that protein-coding genes as well as the two rRNAs and the tRNA-Met

only include poly-T regions and not poly-A regions. When poly-A regions are found on the

direct strand, this indicates that the genes are encoded on the reverse strand. We detected a

few instances of EST reads that include a poly-U stretch, i.e., before the excision of poly-U to

6U. These observations suggest that the shortening of the poly-U stretch, which is necessary

to recover the reading frame, occurs at the RNA level co-transcriptionally or very soon after

the synthesis of the precursor transcript.

Interestingly we noticed that while poly-T regions are present within all genes, the

genes often end with a poly-A. The poly-A ending is part of the TAA stop codon in most

protein-coding genes, except for *cox1*, which has a TAG stop codon (Table 1). The contigs

assembled from RNA reads support the view that these poly-A stretches are reduced to 6A in

the mRNA. Similarly, we noticed that a poly-A stretch is often present in the DNA before the

start of a protein-coding gene (Figure 1). No conserved sequence motif could be identified

around the poly-A/T stretches. The only major departure we could detect around the poly-A/T

stretches was the absence of cytosine just after the 6T site. No other signal could be identified

(Supplementary Fig. S3, Supplementary Material online).

Oikopleura dioica **Mitochondrial Genome Sequence and Structure**

Figure 1 presents the assembled mitochondrial genome of *O. dioica*. We could not determine

the sequence in the region labeled as "Gap" between the *nad5* and *rnl* genes, probably due to

the presence of secondary structures that impaired sequencing. Running the PCR product of

fragment 12 on a gel indicated that the length of the "Gap" region is approximately 400 bp

(supplementary Fig. S1, supplementary table S1, Supplementary Material online). However,

the fact that we could amplify this missing region using primers from both sides of the gap

suggests that the genome is circular. Notably, although we could not assemble the entire

fragment 12, the ONT-contigs we obtained allowed us to determine the end of the *nad5* gene

and the *nad4-rnl* region, supporting the view that we had amplified the correct region of the

mitochondrial genome. The circularity of the mitochondrial genome is in line with Bliznina et

al. (2021).

225
Gene Content and Gene Loss in *O. dioica* Mitochondrial Genome

Most tunicate mitochondrial genomes encompass 13 protein-coding genes, two rRNA genes,

and 24 tRNA genes that are encoded on the same strand (Shenkar et al. 2016; Singh et al.

2009). Surprisingly, in the mitochondrial genome of *O. dioica* we could only identify eight

canonical protein-coding genes: ATP synthase subunit 6, subunits 1-3 of cytochrome c

oxidase (*cox*), cytochrome *b* (*cytb*), and NAD dehydrogenase (*nad*) 1, 4, and 5. In addition to

the two rRNA genes encoding the small and large subunits, we were able to identify a single

tRNA gene, tRNA-Met (Figure 1, Table 1). The genes *cox1, cox2, cox3, cytb, nad1, nad5*, and

tRNA-Met (UAU) are encoded on the forward strand, while *atp6, nad4*, *rns*, *rnl,* and an

unidentified protein-coding gene named *ODI_mt_ORF196* (see below), are encoded on the

reverse strand (Table 1). No incomplete stop codons were found (Table 1). Moreover, we did

not find any LAGLIDADG homing endonuclease ORF encoded within the *cox1* gene. These

findings suggest the loss of five protein-coding genes: *atp8, nad 2, 3, 4L*, and *6* and, all tRNA

genes except tRNA-Met.

An additional mitochondrial protein-coding gene was identified following a Blastn

search of the obtained mitochondrial sequence against the EST data available in Oikobase

(Danks et al. 2012). Notably, programs such as "Find ORFs" (implemented in Geneious

Prime v.2023.2.1) are inapplicable for this mitochondrial genome, due to the extremely fast

rate of evolution and the numerous poly-A/T stretches that disrupt the reading frames of

protein-coding genes. This additional ORF is 1,426 bp long and does not show any significant

sequence similarity to any known protein-coding gene present in the NCBI public database

(i.e., based on Blastp searches). We thus termed it gene "*ODI_mt_ORF196*" based on its

amino-acid length (Figure 1, Table 1). This protein was inferred to have six transmembrane

domains and, based on its position, it corresponds to the putative *nad2* gene identified by

Denoeud et al. (2010). An Alpha fold-based prediction of this gene (supplementary Fig. S4,

Supplementary Material online) was used as query against multiple protein structure

databases: AlphaFold/UniProt50 v4, AlphaFold/Swiss-Prot v4, AlphaFold/Proteome v4,

CATH50 4.3.0, MGnify-ESM30 v1, PDB100 2201222, GMGCL 2204 from Foldseek (van

Kempen et al. 2024). No significant structural hits were obtained.

We identified the two rRNA genes, *rns* and *rnl,* by searching for highly transcribed

regions along unannotated regions of the assembled genome. Two non-protein-coding regions

were found to be highly expressed based on Illumina RNA read mapping (supplementary Fig.

S5, Supplementary Material online). Further support for the transcription of these regions lies

in their presence in the EST data. Specifically, among these two highly expressed regions, the

longer one, which resides between the "gap region" and the *nad4* gene, was assumed to be the

rnl gene. This annotation is supported by RNA structure similarities between the putative *rnl*

and the structure model profile of this gene, as implemented in Mitos2 (Donath et al. 2019).

The second one, located between the *atp6* and the *cox2* genes, was annotated as *rns*, although

no structural similarity was found between this sequence and the *rns* structure profile in

Mitos2. Both the *rnl* and *rns* genes were found to be extremely short: 291 and 232 bp,

respectively, compared to 1,129 and 660 bp in the model tunicate *Ciona intestinalis* (see also

Table 1).

268   We were able to identify a single tRNA gene, tRNA-Met (UAU). This tRNA presents

a typical secondary cloverleaf structure (supplementary Fig. S6, Supplementary Material

online), and UAU anti-codon, corresponding to a methionine codon according to the ascidian

mitochondrial genetic code. Unlike in Bliznina et al. (2021), no other tRNAs were identified.

Aminoacyl-tRNA Synthetase (aaRS) Profiling

We used hidden Markov models (HMMs) to identify nuclear-encoded aaRSs in published

sequences of *O. dioica* and a few other chordate representatives. We found the presence of all

cytosolic aaRSs in published proteomes of *O. dioica* (accessions: GCA_000209535, Norway

and GCA_907165135, Okinawa) (Figure 3). However, in contrast to the other chordates

investigated, *O. dioica* is inferred to have lost most of its mitochondrial aaRSs except for

aaRS-Gly (G), aaRS-Phe (F), and aaRS-Met (M). Similarly, the mitochondrial aaRS-Val (V)

is absent in the tunicates *C. intestinalis* (GCA_000224145) and *Styela clava*

(GCF_000001405), and the mitochondrial aaRS-Lys (K) and aaRS-Thr (T) are absent in all

the species included in this comparison. Furthermore, *O. dioica* encodes two copies of aaRS-

Gly (one cytosolic and the other mitochondrial), while the other chordates encode only a

single copy that functions both in the cytosol and the mitochondrion, suggesting a duplication

event in the lineage leading to appendicularians.

Gene Transfer to the Mitochondrial Genome

HMMs were used to search for the missing canonical mitochondrially-encoded genes (i.e., *nad2*, *nad3*, *nad4L*, *nad6,* and *atp8*) in the nuclear genome. Using these models, we found a protein sequence with a significant similarity (E-value < 5e-07) to the NAD3 model in the two published proteomes of *O. dioica* (proteins CAG5094752 and CBY12996 for the Norwegian and Okinawan genomes, respectively). Domain searches, using Cdsearch (Wang et al. 2022), confirmed that the protein sequences identified include a NAD-ubiquinone/plastoquinone oxidoreductase, chain 3 domain. These searches were conducted using the CAG5094752 and CBY12996 protein sequences as queries against the CDD v3.21 database and yielded E-values of 4.94e-03 and 3.80e-03, respectively. This protein was predicted to include two putative transmembrane domains, in support of its role in the electron transport chain (supplementary Fig. S7, Supplementary Material online). This protein-coding gene was found to be encoded on the X specific region of the X chromosome of *O. dioica* and to harbor three introns. Conversely, no nuclear-encoded protein showing significant similarity to the NAD3 model was found in the proteomes of *C. intestinalis*, *S. clava*, and *H. sapiens*.

To confirm that the identified NAD3 proteins of the Norwegian and Japanese *O. dioica* are expressed in the mitochondria we searched for the presence of mitochondrial target peptides. A significant target peptide signal was found in these putative NAD3 proteins, with a likelihood above 0.89. We could not confirm the transfer of other missing genes (i.e., *nad2*, *nad4L*, *nad6,* and *atp8*) to the nuclear genome, possibly due to the fast evolutionary rate of *O. dioica* and the short length of these proteins (150-152 amino-acid), or to the incompleteness of proteomic data for *O. dioica*.

Phragenetic Reconstructions

Phylogenetic Reconstructions

Phylogenetic reconstructions were performed using two datasets. In the first, *Doliolum*

nationalis was the sole representative of the tunicate class Thaliacea. The second dataset

differed from the first in including two additional thaliacean sequences: the fast-evolving

sequences of *Salpa thompsoni* and *Salpa fusiformis* (supplementary table S4, Supplementary

Material online). Phylogenetic trees were reconstructed under the maximum likelihood (ML)

criterion using various models of sequence evolution implemented in IQtree (Minh et al.

2020; Wang et al. 2017), including complex models (i.e., MtZoa+F+G+I,

MtZoa+EHO+F+G+I, Poisson+C10+G, Poisson+C20+G), and under the Bayesian CAT

model implemented in Phylobayes (Lartillot and Philippe 2004). In all trees, the monophyly

of tunicates was supported with maximal support (bootstrap percentage (BP) = 100, posterior

probability (PP) = 1.0). However, the position of *O. dioica* was highly variable, depending on

the dataset and the evolutionary model chosen (Figure 4, supplementary Fig. S8,

Supplementary Material online). When the tree was reconstructed without members of the

Salpa genus, *O. dioica* was the first tunicate to diverge when using the MtZoa+EHO+F+G+I

or the mtZOA+F+I+G4 models (see Figure 4 and supplementary Fig. S8A, Supplementary

Material online, respectively). Remarkably, the branch support values for a sister-clade

position of *O. dioica* to all other tunicates were low (BP < 50%). Conversely, support for the

monophyly of accepted tunicate clades (i.e., Stolidobranchia and

Thaliacea+Phlebobranchia+Aplousobranchia) was moderate to high (BP = 60-93%). When

using more complex models such as the Poisson+C10+G or the Poisson+C20+G

(supplementary Fig. S8B-C, Supplementary Material online) or the Bayesian CAT model

(supplementary Fig. S8H, Supplementary Material online), *O. dioica* branched within the

Stolidobranchia. This positioning and the resulting relationships among Stolidobranchia

clades were not supported (BP < 50%; PP < 0.5). When members of the genus *Salpa* were

included in the phylogenetic analyses, *O. dioica* systematically grouped with the *Salpa*

members except when using the most complex Poisson+C20+G ML model (supplementary

Fig. S8G, Supplementary Material online). Interestingly, support for this relationship

decreased with the complexity of the ML model used (BP = 85%, MtZOA+G+F; BP = 77%,

mtZOA+EHO+F+I+G4, BP = 51%, Poisson+C10+G) (supplementary Fig. S8D-F,

Supplementary Material online). In the tree reconstructed with the Poisson+20+G model, the

Thaliacea monophyly was recovered (BP = 59%), but *O. dioica* branched within the

Stolidobranchia and, the relationships among tunicate classes were not resolved (BP < 50%).

When using the Bayesian CAT model, the phylogenetic position of *O. dioica* could not be

resolved when *Salpa* was included. Specifically, convergence parameters suggested that

convergence was achieved. However, different topologies were obtained depending on the

chains considered. With chains 1, 2, and 3 the Thaliacea monophyly was recovered but not

supported (PP = 0.53). When chains 1, 2, and 4 were considered, *Salpa* and *O. dioica* were

found to be sister clades (PP = 0.57) (supplementary Fig. S8I-J, Supplementary Material

online). The different branching positions of *O. dioica* among tunicates are summarized in

supplementary Fig. S8K (Supplementary Material online).

Discussion

The mitochondrial genome of *O. dioica* is extremely fast-evolving and characterized by the

presence of numerous poly-A/T stretches undergoing RNA editing. Several canonical

metazoan mitochondrial protein-coding genes and almost all tRNA genes are absent from the

genome; there is a substantial reduction in the size of the rRNA genes; and there is evidence

of a protein-coding gene transfer to the nuclear genome. Such high mitochondrial genome

plasticity characterizes fast-evolving animal clades (Pett et al. 2011; Yahalomi et al. 2017).

Protein-Coding Genes

Five canonical protein-coding genes are absent (*atp8, nad2, nad3, nad4L,* and *nad6*) from the mitochondrial genome of *O. dioica*, of which only *nad3* was identified in the nuclear genome. The four absent genes may be encoded in either the mitochondrial or the nuclear genome, but we were unable to identify them. Indeed, the *O. dioica* mitochondrial and nuclear genomes are extremely fast evolving and, consequently, these genes may have been missed by algorithms for remote homology detection, including HMM searches. Notably, these missing genes (*atp8, nad2, nad4L,* and *nad6*) are among the fastest evolving mitochondrial genes (Castellana et al. 2011; Pesole et al. 1999). The unassigned gene *ODI_mt_ORF196* is predicted to encode a protein harboring four alpha helices (supplementary figure S4, supplementary material online), unlike all other missing proteins in tunicates: the NDH2 protein of most metazoans harbors six alpha helices, while the other canonical mitochondrial proteins are predicted to harbor three or fewer alpha helices (The UniProt The UniProt Consortium 2022). Moreover, none of the methods used (Blastp, HMMer, and Cdsearch) detected a significant similarity with any other canonical proteins. We thus refrained from annotating this gene as one of the four missing genes, although Denoeud et al. (2010) annotated this gene as *nad2*.

After removing the poly-A/T regions, the largest non-coding region identified was 206 bp (between *ODI_mt_ORF196* and *cytb*) and it is possible that at least one among the *atp8* and *nad4L* genes, which are shorter than 300 bp in most tunicates, are encoded within this region. The *atp8* gene, in particular, has been overlooked in the past from the original annotation of several mitochondrial genomes (e.g. Helfenbein et al. 2004; Hoffmann et al. 1992; Le et al. 2002; Okimoto et al. 1992), including tunicates (Yokobori et al. 1999; Yokobori et al. 2003) and was then identified in ascidians following mitochondrial EST analyses (Gissi et al. 2004; Gissi and Pesole 2003).

Our results suggest that *nad3* has been relocated to the nuclear genome. While this is

the first evidence of relocation of this mitochondrial gene to the nucleus in animals, a case of

nuclear *nad3* has been reported in the green alga *Chlamydomonas reinhardtii* together with

the transfer of *nad4L* (Cardol et al. 2006). Similar transfers to the nuclear genome have been

described for other animal mitochondrial genes, for example, the *atp6* and *atp9* genes of the

ctenophore *Mnemiopsis leidyi* (Pett et al. 2011) and the mitochondrial gene *atp9* in the

demosponge *Amphimedon queenslandica* (Erpenbeck et al. 2006). Short ORFs are more

difficult to identify than long ones, and because *atp8* and *nad4L* are relatively short it is

possible that this factor, in addition to their high rate of evolution, has impaired their

identification. The fact that *nad6* was detected neither in the mitochondrial genome nor in the

nuclear one, is more surprising. This gene is long in other tunicates (450.9 ± 20.2 bp), and

except for the gene ODI-mt-*ORF196,* no other putative ORFs longer than 300 bp were

detected. The possible incompleteness of the two public nuclear genome assemblies of *O.*

dioica and their inferred proteomes, or the specific criteria and bioinformatic pipeline used to

annotate the protein genes in these nuclear assemblies, could also explain the failure to

identify *nad6*, *atp8*, and *nad4L* in the available nuclear assemblies.

A third explanation could be that the genes that we could not identify have become

fused with other mitochondrial genes. For example, the ORF we annotated as *atp6* could

include regions corresponding to both *atp6* and *atp8*. An example of ORF fusion is found in

the *cox*1/2 gene of amoebas (e.g., Lonergan and Gray 1996). However, this explanation is

unlikely in our case since all *O. dioica* ORFs are shorter than their tunicate counterparts

(Table 1), while in the case of a fusion, we would expect these ORFs to be longer.

rRNA Genes

Tunicate rRNA genes are known to be poorly conserved. As a case in point, the boundaries of

tunicate mitochondrial rRNA genes have been mostly inferred from the start and end of the

flanking genes, indicating that the length estimates of rRNA genes are error-prone

(Rubinstein et al. 2013; Singh et al. 2009). Only for the ascidians *Halocynthia roretzi* and *C.*

intestinalis, could the rRNA boundaries be better defined, based on EST transcript data (Gissi

and Pesole 2003). Although the two rRNA genes are retained in all tunicate mitochondrial

genomes, they exhibit a great range of sizes. Tunicates present the shortest mitochondrial

rRNA genes among chordates, with lengths varying from 596 to 984 bp for *rns* and from 921

to 1,299 bp for *rnl* (Table 1). Both rRNAs are considerably shorter in *O. dioica,* with the *rns*

and *rnl* being 242 bp and 215 bp, respectively, following removal of the poly-T stretches

(Table 1). Additionally, they have lost most structural similarity with other metazoan rRNA

genes (HMM searches could only identify a part of the *rnl* gene). Interestingly, extreme

mitochondrial rRNA reduction and lack of structural similarity are characteristics shared

between appendicularians and ctenophores (Arafat et al. 2018), another lineage with an

extreme rate of mitochondrial genome evolution. We annotated the two rRNA regions as

different genes because most mitochondrial genomes contain these two genes. However,

given the absence of sequence similarity, we cannot exclude the possibility that the *rns* region

may instead be a fragment of the *rnl*. Indeed, in calcareous sponges and Placozoa, rRNA

genes are fragmented and encoded on different regions of the mitochondrial genome

(reviewed in Lavrov and Pett 2016). Of note, *trans*-splicing is unknown in animals.

tRNA Genes

Tunicate tRNA gene content is variable and ranges from 22-27 tRNAs (Iannelli et al. 2007;

Rubinstein et al. 2013). Specifically, the loss and acquisition of additional tRNA genes seem

to be tolerated and the number of tRNA genes has been found to vary in a species-specific

manner (Gissi et al. 2008; Rubinstein et al. 2013). In *O. dioica* the situation is again extreme,

as all tRNAs are absent except for tRNA-Met (UAU).

Nuclear-encoded cytoplasmic tRNAs import to the mitochondria has evolved many

times over the course of mitochondrial evolution (Adams and Palmer 2003). We thus assume

that the missing mitochondrial tRNAs in *O. dioica*'s mitochondrial genome are encoded in the

nuclear genome and imported to the mitochondria.

The retention of a single tRNA-Met in the mitochondrial genome, despite the loss of

all or most tRNAs, is not novel to appendicularians and has been observed in cnidarians,

certain chaetognaths, and Keratosa sponges (Lavrov et al. 2023). The conventional

explanation is based on the fact that mitochondrial protein synthesis starts with N-

formylmethionine, as in prokaryotes, rather than methionine, the initiating amino acid in

eukaryotic cytosolic protein synthesis. The addition of a formyl group to methionine is

catalyzed by the enzyme methionyl-tRNA formyltransferase (Bianchetti et al. 1977). As

found in prokaryotes, and in accordance with the prokaryotic endosymbiotic origin of

mitochondria, this modification occurs not on free methionine but only after methionine is

loaded onto a specific tRNA: namely, the initiator tRNA-fmet. Hence, it is usually assumed

that the retained tRNA-Met in mitochondria is the initiator tRNA-fmet (e.g., Beagley et al.

1998).

Interestingly, animal mitochondrial genomes usually retain tRNAs that are specific to

their mitochondrial genetic codes. For example, most cnidarians retain the tRNA-Trp gene

necessary for the translation of the UGA codon, which is a stop codon in the nuclear genetic

code (Beagley et al. 1998). In the mitochondrial genetic code of ascidians, four codons have a

different assignment compared with the nuclear genetic code: methionine (AUA), tryptophan

(UGA), and glycine (AGA and AGG). We thus expected the presence of a tRNA-Trp (for the

translation of the UGA codon) and a tRNA-Gly (to translate the codons AGA and AGG) in

the mitochondrial genome of *O. dioica*. Therefore, we computed the codon frequency across

all protein-coding genes of *O. dioica*, thus verifying the usage of all these AUA, UGA, AGA,

and AGG codons in the mitochondrial genome (supplementary table S6, Supplementary

Material online). Indeed, it has been found that translation of the AGA and AGG codons to

Gly improves the similarity between *O. dioica* and tunicate mitochondrial proteins,

supporting the use of the ascidian mitochondrial genetic code in appendicularians (Pichon et

al. 2020). The presence of a cytosolic and a mitochondrial aaRS-Gly, observed while

searching for mitochondrial aaRS in the *O. dioica* nuclear genome, further supports the view

that a specific mitochondrial tRNA recognizes the glycine AGA and AGG codons. Regarding

protein-coding genes, two explanations are possible for the lack of mitochondrial Gly and Trp

tRNAs: either we failed to identify these tRNA genes in the sequenced mitochondrial genome

due to their very unusual secondary structure or because they are interrupted by poly-A/T

stretches; or these tRNAs are nuclear-encoded and imported from the cytosol. In support of

the latter scenario, it was found that among animals, both ctenophores and myxozoans have

lost all their mitochondrial tRNA but still use the UGA codon as tryptophan (Arafat et al.

2018; Pett et al. 2011; Yahalomi et al. 2017).

Genome Editing of Poly-A/T Stretches

The sequencing of PCR fragments using ONT long-reads allowed us to assemble a large part

of the mitochondrial genome of *O. dioica*. It confirmed the presence of numerous poly-A/T

stretches in the DNA sequence that are reduced to 6U-sites in the RNA transcript. Because

accurate sequencing of long homopolymers is currently unfeasible using Sanger, Illumina

(Ari and Arikan 2016), or Nanopore sequencing (Delahaye and Nicolas 2021; Sereika et al.

2022), assessing the length of T/A stretches in our assembly is unreliable. The average size of

these regions is likely slightly higher than that reported in the assembly because sequencing

methods tend to underestimate the length of homopolymer stretches. Another source of

uncertainty regarding the length of the homopolymer stretches stems from the fact that we

sequenced amplicons obtained from DNA extracted from a pool of polymorphic individuals.

Intra-species variation in the length of these stretches likely exists, further contributing to

inference uncertainty.

What is the molecular mechanism underlying the editing to 6U-sites and, more

specifically, how are these sites recognized? We could not identify a conserved sequence

motif surrounding these sites (Supplementary Fig. S3, Supplementary Material online). Our

analysis revealed the following general motif for the poly-A/T stretch (see Materials and

Methods): (1) the 5'-end of this strand starts with a perfect 6T homopolymer; (2) downstream

of this region, there is a poly-pyrimidine track whose length varies from 2 to 509 bases; (3) it

ends when a purine is first encountered. Only the poly-pyrimidine track is excised. We

speculate that the perfect 6T motif in the upstream region may be the beginning of the

recognition editing signal and that a purine marks its end. In between these motifs, the

polypyrimidine track is excised. Notably, the enzymes involved in this excision mechanism

are still unknown.

Another kind of editing was observed within the cytochrome *b* gene: rather than a

perfect 6T motif, it starts with "TTTTTCTT", followed by an irregular excised region,

"AACTT", as it contains two purines (supplementary Fig. S9, Supplementary Material

online).

Mitochondrial genome editing is rare in animals and is usually limited to the insertion

of a few bases (Lavrov et al. 2016). Here, rather, the deletion of a large polypyrimidine region

is inferred. The removal of such a long stretch at the RNA level is a characteristic of introns

(Chorev and Carmel 2012). Spliceosomal introns are characterized by short signal sequences

such as GT/AG and AT/AC (Berná and Alvarez-Valin 2014; Rogozin et al. 2012). The RNA

editing, we observed here, seems to differ greatly from such a splicing mechanism, as we

could not find similar motifs. Similarly, the poly-A/T stretches bore no sequence similarity to

self-splicing introns, which have well-defined secondary structures (Waring and Davies

1984).

Trypanosomatid protists present a cleavage-ligation pathway through which multiple

uridines can be deleted or inserted by a U-specific exonuclease and a terminal uridil

transferase, respectively (Cruz-Reyes and Sollner-Webb 1996). This particular editing

mechanism involves guide RNAs, which determine the number and positions of the U

nucleotides deleted or inserted within the mitochondrial genome of these protists, e.g.,

Leishmania tarentolae (Connell et al. 1997). Future studies are necessary to determine

whether some components of the editing mechanisms in these protists and in *O. dioica* are

evolutionarily related.

Appendicularians have classically been divided into three families: Fritillariidae,

Kowaleskiidae, and Oikopleuridae (Garić 2024). However, preliminary results from our

group suggest that the Kowaleskiidae are actually members of the Fritillariidae (Garić et al.

2018). The presence of edited homopolymer stretches has been noted for the COI gene of *O.

vanhoeffeni* (Latour 2021). Interestingly, we observed that large poly-A/T inserts are also

present in members of the Fritillariidae, suggesting that this mitochondrial editing mechanism

was present in the ancestor of Appendicularia (Garić et al. 2018). Conversely, while

examining published data from Naville et al. (2019), we identified a clade within the

Oikopleuridae that includes the species *Bathochordaeus stygius*, *Mesochordaeus*

erythrocephalus, and *Oikopleura longicauda*. These three species do not harbor large poly-

A/T stretches (Garić et al. 2018), suggesting that this RNA editing mechanism may have been

lost in Oikopleuridae. Additional genomic and transcriptomic data from a larger taxon

sampling of appendicularians are needed to better understand the distribution and

evolutionary dynamics of this editing mechanism.

536   EST and transcriptomic RNAseq data are available for *O. dioica* from Norway and

mainland Japan, respectively. For the Norway sample, the genomic data obtained in this work

indicated the presence of RNA editing. To evaluate whether the poly-A/T stretches are

conserved among *O. dioica* lineages, we compared the location of 6U-regions in the DNA and

transcriptomic data from Norway and mainland Japan, respectively (Supplementary Fig. S2,

Supplementary Material online). Our results suggest that while many of the 6U-loci are

conserved, differences do exist. For example, in the *cox1* gene, a 6U-site in the Norwegian

sequence and a UUCUUU site in the Japanese sequence can be reliably aligned. These

differences suggest that the poly-A/T stretches loci may not be conserved in the mitochondrial

genomes of different populations or species. Of note, it was recently suggested that *O. dioica*

from Japan and Norway represent different species (Masunaga et al. 2022; Plessy et al. 2024).

The birth and death dynamics of these poly-A/T stretches will be better clarified when

genomic and transcriptomic data from a larger appendicularian taxonomic sampling become

available.

550   That we were able to assemble large contigs encompassing several genes from RNA

sequences indicates that the pre-mRNA transcripts in *O. dioica* are polycistronic. In

vertebrates, it has been shown that the cleavage of tRNAs that flank protein-coding genes

constitutes the mechanism (termed tRNA punctuation) that releases monocistronic or

bicistronic mature mRNAs of mitochondrial coding genes from the large RNA precursor

(Ojala et al. 1981). In ascidians, the transcription and RNA maturation process of the

mitochondrial genome has been reconstructed through EST analyses, revealing that the main

features of mitochondrial transcription are conserved between vertebrates and ascidians (*i.e.,*

the presence of polycistronic and immature transcripts; the creation of stop codons by

polyadenylation; tRNA signal punctuation; and rRNA transcript termination signals) (Gissi

and Pesole 2003). Since most tRNAs have been lost in *O. dioica,* a mechanism differing from

the tRNA signal punctuation is likely employed to release mature mRNAs.

The Phylogenetic Position of *O. dioica*

Based on both molecular (Delsuc et al. 2018) and morphological (Braun et al. 2020) analyses,

O. dioica was placed as a sister-clade to all other tunicates. However, it was recognized that

this placement, in molecular studies, may stem from a long-branch-attraction (LBA) artifact

(Delsuc et al. 2018). In our analysis, we found that the phylogenetic position of *O. dioica* was

highly variable (supplementary Fig. S8K, Supplementary Material online). The grouping of

O. dioica and members of the genus *Salpa* is most likely an LBA artifact (Philippe et al.

2005). However, even when the two *Salpa* representatives were removed from the

phylogenetic analyses, the phylogenetic position of *O. dioica* was never well supported and

depended on assumed probabilistic model used. The lack of a phylogenetic signal in the

mitochondrial dataset, which precludes the reconstruction of phylogenetic relationships at the

phylum level, is a feature of fast-evolving species (Arafat et al. 2018; Pett et al. 2011;

Yahalomi et al. 2017).

An *O. dioica* Mitochondrial Genome from Okinawa, Japan

After completing this manuscript, a description of the complete mitochondrial genome of *O.*

dioica from Okinawa (Japan) (accession number PP146516) was reported in a preprint

(Dierckxsens et al. 2024). While numerous substitutions differentiate the two genomic

sequences (113 bp difference in the barcoding region corresponding to a K2P distance of

20.3%), the gene order was identical. One difference is that of the presence of a region

containing several tandem repeats between the *cox1* and the gene we have labeled

ODI_mt_ORF196. Remarkably, Dierckxsens et al. (2024) succeeded in assembling a circular

molecule. The region in the Okinawa sample that is homologous to the part that we failed to

sequence, is a large non-coding region of ~28,000 bp composed of numerous repetitive

elements. A similar structure may be present in the Norway sequence analyzed here.

Conclusions

We have demonstrated that a combination of DNA and RNA data from long- and short-read

sequencing could significantly improve our understanding of the structure and gene content of

the *O. dioica* mitochondrial genome.

The mitochondrial genome of *O. dioica* shows a unique gene content and an extreme

rate of sequence evolution. A total of 28 genes are missing, including *atp8*, *nad2*, *nad3*,

nad4L, *nad6*, and all tRNA genes except tRNA-Met (UAU). The mitochondrial gene *nad3*

was transferred to the nucleus and harbors a mitochondrial target peptide. Further work is

however needed in order to decipher the complete sequence of the mitochondrial genome, and

in particular to unravel the accurate sequence and length of the poly-A/T regions.

Unfortunately, no sequencing techniques currently unable us to perfectly sequence these

regions, as the sequencing methods Illumina, ONT, and Sanger approaches failed to correctly

sequence long polynucleotide regions. However, information obtained from the sequencing of

PCR fragments helped us to assemble a draft mitochondrial genome.

Because of the fast-evolutionary rate, we were unable to accurately determine the

phylogenetic position of *O. dioica*. LBA artefacts are expected due to such a high

evolutionary rate, and the inferred tree often varied depending on the set of species included

in the analysis. Thus, mitochondrial sequences are not the markers of choice when the goal is

the accurate inference of phylogenetic relationships within appendicularians. We expect that

the availability of nuclear genomes whose rate of evolution is lower than that of the

mitochondrial genome, will enable more accurate reconstruction of the phylogenetic

relationships among appendicularian species. Of note, comparative genomic studies of *O.*

dioica and additional members of the appendicularians are expected to provide us with a

better understanding of the evolution of RNA editing in this group.

Materials and Methods

DNA Extraction, Sequencing, and Assembly

Living specimens of *O. dioica* were provided by the Sars International Centre for Marine Molecular Biology (SICMMB) (Bergen, Norway) in March 2019 and cultured at Humboldt-University zu Berlin (HU) following the methods detailed in Bouquet et al. (2009). The animals sequenced in this study originated from a cross between new animals from the field collected at Rosslandspollen, north of Bergen (60°33'54.6"N 5°2'24.5"E), and an existing culture (generation 29). These animals were grown for ~20 generations in the lab. Ethanol-preserved samples were shipped to Tel-Aviv University. Genomic DNA was extracted from ~100 whole individuals following the protocol of Fulton (2012). About 100 ng were sent to the Technion sequencing center (Haifa) for Illumina sequencing (November 2019), the libraries were prepared using the TruSeq DNA Nano library prep kit (Illumina), and were sequenced on an Illumina Hiseq 2500 platform. Similarly, about 1 μg of total DNA was sent to Hy Laboratories Ltd (Hylabs, Rehovot) for ONT sequencing using a MinION Mk 1 R9.4 flow cell with a MinION Mk1B device and 1D sequencing kits (May 2019). However, the ONT sequencing only provided 962 reads (mean length 750 bp). ONT sequencing was thus performed again using a different DNA extract and a different sequencing service (the Technion sequencing center) (on July 2019 and December 2021). The flow cell and device setting and sequencing kits were the same as previously. The first sequencing (July 2019) provided ~2 million reads (mean length 2,330 bp) but we could not assemble the mitochondrial genome using these data. *A posteriori,* we found that only ~600 reads mapped to the mitochondrial genome (mean length 300 bp) and that these reads only mapped to a few regions of the mitochondrial genome (data not shown). For this second ONT sequencing, fragments around 3,000 bp were selected. Because we realized that mitochondrial DNA might be fragmented, we performed the last ONT sequencing (from December 2021) without

fragment size selection. This ONT sequencing also provided only 3,144 reads (mean length

1,070 bp).

The Illumina 100 bp paired-ended reads were cleaned from adaptors using Cutadapt

v.2.8 (Martin 2011) and assembled using IDBA-UD v.1.1.3 (Zhu et al. 2010) with default

parameters. Efforts to assemble the mitochondrial genome from these reads were unsuccessful

because contigs were broken at the long poly-A/T regions.

Neither Illumina nor ONT sequencing enabled us to assemble a complete

mitochondrial sequence. We thus decided to amplify the mitochondrial genome from several

PCR fragments. To do so, we needed to design primers in regions that did not include

homopolymers. We used published RNA reads and EST sequences to identify mitochondrial

genes and design primers.

Identification of Canonical Mitochondrial Protein Transcripts in Published RNA and

EST Sequences

Illumina RNA reads from *O. dioica* specimens from Japan and the North Sea were

downloaded from the National Center for Biotechnology Information (NCBI) (accessions:

SRR1693762, SRR1693765, SRR1693766, and SRR1693767) (Wang et al. 2015), and the

European Nucleotide Archive (ENA) (accessions: SRR20015061) (Raghavan et al. 2023),

respectively. These reads were assembled using IDBA-UD v.1.1.3 (Zhu et al. 2010) with

default parameters. tBlastn searches were conducted against the two transcriptome assemblies

obtained using the amino acid sequences of the protein-coding genes of the mitochondrial

genome of *Ciona robusta* (AJ517314) as query. The reads from Japan could be assembled in

four contigs and will be referred to as "Japan" (supplementary Fig. S10A, Supplementary

Material online). The genes from the North Sea could be assembled in five contigs and will be

referred to as "North Sea" (supplementary Fig. S10B, Supplementary Material online). All

contigs were annotated with the webserver MITOS V2 (Donath et al. 2019).

665   Mapping Illumina DNA reads and the few ONT reads to these contigs confirmed the

presence of homopolymers and also indicated that the Japanese *O. dioica* presented many

substitutions when compared to our sample, which originated from Norway. EST sequences

from *O. dioica* samples from Norway were available in the OikoBase database (Danks et al.

2012). These data originated from natural populations of *O. dioica* caught in the fjords near

Bergen, Norway (Denoeud et al. 2010). For each mitochondrial ORF identified in the

Illumina North Sea dataset, we mapped ~500 EST sequences to the reference. We next

filtered the flanking regions of the extracted EST sequences against the UniVec Vector

database using Geneious Prime v.2023.2.1. The consensus sequences were extracted for each

gene, which helped us determine the exact gene boundaries. We verified that all coding

transcripts we had identified could be successfully translated to proteins using the ascidian or

the invertebrate mitochondrial genetic codes. We found that the ascidian genetic code yielded

proteins with a higher sequence similarity to mitochondrial proteins in other animals than

when using the invertebrate genetic code, in agreement with Pichon et al. (2020).

PCR Amplification, Sequencing, and Assembly

Primers of ~ 30 bp with a TM ~ 60 C° were designed based on the ORFs assembled from the

EST, accounting for the gene order inferred from the contigs (supplementary table S1,

Supplementary Material online). Long PCRs were performed with several primer

combinations (one from each pair of contigs and in various orientations), to determine gene

arrangements along the mitochondrial genome. Because of the high AT content of the

genome, we were unable to design primers that would result in overlapping PCR fragments.

Instead, we used the same region for forward and reverse primers, leading to non-overlapping

fragments. The sequence of the primer region was confirmed using Illumina data. PCRs were

performed in a 25µl volume with 2u of TaKaRa Ex-Taq polymerase (Cat.# RR001), 1X of

Ex-Taq buffer, 1M of Betain, 0.2mM of dNTPs (TaKaRa mixture), 0.5µM of each primer and

1µl *O. dioica* DNA extract. The PCR program used: (a) a long denaturation step at 95°C for 5

min; (b) 35 cycles of a short denaturation step at 95°C for 45 sec followed by an annealing

plus elongation step at 68°C for 10 min; and (c) a long elongation step at 72°C for 10 min.

Reamplifications were performed to improve the PCR yield of some fragments.

Reamplifications were performed using 1 µl of PCR products in place of genomic DNA.

Twelve PCR reactions were successful and sent for ONT sequencing (supplementary Fig. S1,

Supplementary Material online). Each PCR product was barcoded and the sequencing was

performed on May 2023 by the Technion Sequencing Center on an Oxford Nanopore

MinION device (MinKNOW software v.22.12.7), using one R9.4.1 flow cell (Oxford

Nanopore, cat no. FLO-MIN106D). The ONT reads obtained were filtered from adaptors

using Porechop v.0.2.4 (Wick et al. 2017) with the parameter -middle_threshold 85. Next, we

assembled the trimmed ONT reads with Flye v.2.9.1 (Kolmogorov et al. 2020) as

implemented in Geneious Prime v.2023.2.1 with two rounds of polishing. The same

parameters could not be used for all PCR fragments, as in some cases the assembly program

returned an error message or failed to terminate. In these cases, other parameters were tested:

e.g., if the default parameter "genome size" of 1,000 bp resulted in a failed run, we tried

running the program with increasing values of genome size from 2,000 to 5,000 bp. If this

also failed, we added the option "metagenome" or the option "reduced contig = 30".

However, even with these modifications the assembly of the PCR fragment #12 was

unsuccessful.

Many of the assembled contigs were palindromic, with the middle of the contig

corresponding to the PCR primer. In this case, the consensus sequence was split into two

halves, reads were mapped to each half (using Geneious Prime v.2023.2.1 with default

options), and the region with the highest number of mapped reads was kept as the consensus

sequence.

The entire mitochondrial sequence was determined by connecting the obtained contigs

in their overlapping primer regions, i.e., the same primer region was the forward primer of

one PCR fragment and the reverse primer of another fragment. The exact sequences of the

primer regions were verified by mapping the Illumina reads to the assembled genome

sequence.

Annotation of the Mitochondrial Genome

Protein-coding genes

Gene annotation was based on Blast searches against the NCBI database and was confirmed

using MITOS v.2 (Donath et al. 2019). The end positions of protein-coding genes were

verified using EST data (the presence of a poly-A region after the stop codon). The start

codon was determined by searching for the first start codon upstream of the region that

corresponded to the EST data.

rRNAs

The mapping of Illumina RNA reads from Japan (SRR1693762, SRR1693765, SRR1693766,

SRR1693767) and the North Sea (SRR20015061) to the mitochondrial genome of *O. dioica*

revealed two regions with higher coverage that we inferred to be *rns* and *rnl* (supplementary

Fig. S5, Supplementary Material online). Read mapping was done using the Geneious

Mapper, Geneious Prime v.2023.2.1 with the following parameters: medium-low sensitivity,

randomly map multiple best matches, only map paired reads which both map, and trim paired

read overhangs. Average coverage was computed with Geneious Prime v.2023.2.1 (i.e., panel

statistics). The region that we annotated as *rnl* was also identified as such using gene

prediction with MITOS v.2 (Donath et al. 2019). The *rnl* and *rns* sequences were also present

in the EST data, which helped determine the boundaries of these ribosomal genes.

tRNAs

Putative tRNAs were searched using MITOS v.2 (Donath et al. 2019) and the ARWEN v.1.2

webserver (Laslett and Canbäck 2008). Both programs identify a single tRNA gene

(supplementary Fig. S6, Supplementary Material online).

Aminoacyl-tRNA Synthetase Profiling

To corroborate the absence of tRNA genes in the mitochondrial genome, we searched for

aaRS genes in the nuclear proteome of *O. dioica* (i.e., the amino acid sequences of *O. dioica*

in the NCBI "protein" database). The aaRS search was done using HMMER v.3.3.2 software

(Eddy 2011) and was based on profiles developed by Pett and Lavrov (2015). These profiles

were used to search against proteome datasets downloaded from NCBI on April 2023 of *O.*

dioica from Norway (GCA_000209535), *O. dioica* from Okinawa (GCA_907165135), *H.*

sapiens (GCF_000001405.40 Human Build 38 patch release 14), *C. intestinalis*

(GCF_000224145.3), and *S. clava* (GCF_013122585.1).

Nuclear Gene Transfer

We used HMM searches against the proteome of *O. dioica* from Norway GCA_000209535

and *O. dioica* from Okinawa GCA_907165135 to detect canonical mitochondrial-encoded

genes that could have been transferred to the nuclear genomes. The searches were performed

with HMMER v3.4 (Eddy 2011) using the *atp8*, *nad2*, *nad3*, *nad4L*, and *nad6* gene models

developed by Gabriel Alves Vieira and available in the mitos2_wrapper source code

repository (Vieira 2023). As negative control we also searched the proteomes of *H. sapiens*

(GCF_000001405.40 Human Build 38 patch release 14), *C. intestinalis* (GCF_000224145.3),

and *S. clava* (GCF_013122585.1). Putative mitochondrial proteins that were found to be

encoded in the nuclear genome were searched for mitochondrial targeting signal using the

TargetP-2.0 webserver (Armenteros et al. 2019).

Codon Usage and Amino-Acid Composition

Codon usage and amino acid composition were computed using Geneious Prime v.2023.2.1.

Characterization of ODI_mt_ORF196

The number of transmembrane domains of ODI_mt_ORF196 was predicted using TMHMM

v.1.0.2 in Geneious Prime v.2023.2.1 (Viklund and Elofsson 2004). The protein structure was

predicted using Alphafold (Jumper et al. 2021). The predicted structure was used as query

against a large database of structures using Foldseek (van Kempen et al. 2024).

Characterization of the Poly-A/T Stretches

We defined poly-T stretches by manually examining 59 cases of edited poly-T sites, i.e., these

editing cases were validated by comparing RNA and DNA sequences of these loci. A classic

poly-T stretch is defined as follows: (1) the 5'-end of this stretch starts with a perfect 6T

homopolymer. This region is not excised; (2) downstream of this region, there is a C/T stretch

(i.e., a Polypyrimidine track). This region is excised; (3) the stretch ends when either an A or

a G nucleotide is encountered. This A or G nucleotide is not part of the poly-T stretch and is

not excised. In six validated cases this pattern was violated. In five of these cases there was a

single A nucleotide within the long C/T stretch and in one case a G. We thus include in the

definition also cases in which the C/T stretch comprises a single purine, as long as it is

followed by at least 10 more C/T nucleotides. Searching the genome with this definition, we

found six additional poly-T stretches (four classic without purine in the C/T stretch and two

non-classic with one purine in the C/T stretch) for which we lack RNA support. In total, we

annotated 65 poly-T stretches in the mitochondrial genome of *O. dioica*. As we searched for

the above motif or its reverse complement to account for both strands, we termed these

stretches poly-A/T stretches.

For each homopolymer site in the genome we extracted 50 bp before the site, the 6 T/A in the

site itself, and 50 bp after the site using our own python v.3.11.5 script on Jupiter Notebook

v.7.1.2. Nucleotide frequencies before and after the homopolymer sites were plotted

(Supplementary Fig. S3, Supplementary Material online) using WebLogo v.2.8.2 with

parameter "Frequency Plot" selected in the section "Advanced Logo Options". (Crooks et al.

2004). To determine the conservation of the poly-A/T stretches, we aligned the mitochondrial

gene regions extracted from the Illumina RNA data from Japan to our own assembly.

Pairwise alignments were computed using MAFFT v.7.490 under the L-INS-i algorithm.

Barcoding Analysis to Confirm Species Identity

We used a barcoding approach to confirm that our DNA samples and the EST deposited in

Oikobase belong to the same *Oikopleura* species. We computed the evolutionary Kimura two

parameters (K2P) distance using MEGA v.11.0.13 (Tamura et al. 2021) between the

barcoding region of the mitochondrial sequence we had assembled (i.e., the CDS region

without the poly-A/T stretches) and the consensus barcoding region we had assembled from

the EST sequence. We extracted the extended tunicate barcoding *cox1* region of Salonna et al.

(2021) from the mitochondrial DNA sequence. We next searched for sequence similarity

against the EST data using Blastn. Out of the 100 best hits, a consensus sequence of the EST

cox1 region was computed. Specifically, the K2P distance between the two sequences (i.e.,

the EST consensus and the mitochondrial genome) was 0.0093 substitutions per site (6 bp

differences, 0.9%). Because the number of bp differences among the EST sequences ranged

from 0 to 20 (average ~8bp), we inferred that our sample and the EST data had originated

from the same species. Moreover, the barcoding region from the North Sea RNA contigs was

also found to be identical to the EST consensus sequence, suggesting that these RNA reads

had originated from the same species as well.

In contrast, the barcoding region of the Japanese RNA contigs presented 93 bp

differences from the EST and the mitochondrial genome from Norway assembled in this work

(K2P distance of above 0.163 substitutions per site). This suggests that these mainland

Japanese sequences originated from a different species than the *Oikopleura* species whose

mitochondrial genome was determined in this work (Masunaga et al. 2022; Plessy et al.

2024).

The sequence PP146516 from Okinawa (Dierckxsens et al. 2024) not only greatly

differed from the Norway sequences, (above 111 bp difference in the barcoding region and

K2P distance of above 0.198 substitutions per site) but also greatly differed from the Japanese

contig we had assembled from mainland Japan (94 bp difference in the barcoding region and

K2P distance of 0.163 substitutions per site), supporting the view that PP146516 belongs to a

third *Oikopleura* lineage (Masunaga et al. 2022; Plessy et al. 2024).

Mitochondrial Genome Length among Tunicates

The average mitochondrial genome size of tunicates and the average gene lengths were

calculated based on 34 complete mitochondrial genomes of tunicates downloaded from the

NCBI database (supplementary table S4, Supplementary Material online).

Phylogenetic Analysis of Mitochondrial Protein Sequences

The phylogenetic analyses encompassed 34 tunicate species with a complete mitochondrial

genome that were deposited in NCBI (last accessed: June 2023) and 14 outgroup species

(supplementary table S4, Supplementary Material online). The outgroup species were chosen

following Singh et al. (2009). Eight protein-coding gene (*atp6, cox1, cox2, cox3, cytb, nad1,*

nad4, nad5) were extracted from the Genbank flatfile of each species using Geneious Prime

v.2023.2.1. We used the Guidance2 server version 2.02 (Sela et al. 2015) with default

parameters to identify unreliable regions of the protein alignment, which was generated using

MAFFT version 7.407, also with default parameters. We excluded these regions as well as

alignment columns that included more than 50% missing data or gaps. The different protein-

coding alignments were then concatenated to form a super-matrix with 2,027 columns.

Because members of the tunicate genus *Salpa* are extremely fast evolving, we prepared a

second dataset, repeating the above steps, but excluding *S. thompsoni* and *S, fusiformi.* The

resulting super-matrix of 46 species included 2,176 columns.

The phylogenetic trees were reconstructed under the maximum likelihood (ML)

criterion using IQ-TREE 2.1.2 (Minh et al. 2020). First, we identified the best-fitting model

by using the ModelFinder-Plus (MFP) option of IQ-TREE 2.1.2 and computed the ML tree

using this model. Using IQtree, the model that best fitted both datasets (with and without the

genus *Salpa*) was mtZOA+F+G+I. We also reconstructed the evolutionary relationships using

complex models that are not implemented in MFP: mtZOA+EHO+F+G+I, Poisson+C10+G,

and Poisson+C20+G. The robustness of each internal branch of the ML trees obtained was

evaluated with 100 bootstrap replications (Felsenstein 1985).

Bayesian tree reconstructions were performed using the site-heterogeneous CAT

model implemented in Phylobayes MPI v1.5 (Lartillot et al. 2013), which is expected to lower

the impact of long-branch attraction on the inferred phylogenetic tree (Lartillot et al. 2007).

For the first and second datasets, four independent chains were run for 71,880 and 188,000

cycles, respectively. The first 5,000 and 10,000 trees from each chain, for the first and second

datasets, respectively, were discarded as burn-in and trees were sampled every 10 trees (i.e.,

the sample size was 3,160, and 17,800 trees per chain for the first and second dataset

respectively). Chain convergence was assessed using the bpcomp and tracecomp scripts,

which are part of Phylobayes. Specifically, the bpcomp maxdiff values were 0.23 and the

tracecomp effsize and rel_diff values were >800 and below 0.15, supporting a proper

convergence for the first dataset. For the second dataset, the bpcomp maxdiff value was 0.40

when the four chains were considered together, which indicated insufficient convergence.

However, when considering only chains 1, 2, and 3 or chains 1, 2, and 4 this value dropped to

0.18 and 0.27, indicating proper convergence. In both cases, rel_diff values were >150 and

below 0.16, supporting convergence.

Acknowledgments

We want to thank Nitzan Fourier from the Technion Genome Center for her help with Nanopore sequencing. We thank Yair Shimony for his help with the structure predictions.

Author Contributions

D.H., C.G., and T.S. conceived the study and designed the research. T.S. provided the biological samples. M.N. and Y.K. performed the bench work. D.H. and Y.K. assembled and annotated the mt genome. D.H., M.N., T.P., and Y.K. performed the various evolutionary analyses and drew the figures. All authors contributed to interpretation of the results. D.H., T.P., and Y.K. wrote the first draft of the manuscript. All authors assisted in revising the manuscript and approved the final version of the text.

Funding

This work was supported by a German-Israeli Foundation for Scientific Research and Development grant to DH and TS (Grant No. I-1454-203.13/2018).

Data Availability

The Illumina read data have been deposited in the European Nucleotide Archive under study ID number PRJEB72855. The mtDNA sequence was deposited under accession number OZ035851. The ONT reads obtained for each of the PCR fragments, the phylogenetic trees,

as well as the sequences assembled from published RNA and EST data are available on

GitHub (https://github.com/dorohuchon/Oikopleura_dioica_mitochondrial_genome).

Literature Cited

Adams KL, Palmer JD 2003. Evolution of mitochondrial gene content: gene loss and transfer

to the nucleus. Mol Phylogenet Evol. 29: 380-395. https://doi.org/10.1016/S1055-

7903(03)00194-5

Arafat H, Alamaru A, Gissi C, Huchon D 2018. Extensive mitochondrial gene rearrangements

in Ctenophora: insights from benthic Platyctenida. BMC Evol Biol. 18: 65.

https://doi.org/10.1186/s12862-018-1186-1

Ari Ş, Arikan M. 2016. Next-generation sequencing: advantages, disadvantages, and future.

In: Hakeem KR, Tombuloğlu H, Tombuloğlu G, editors. Plant Omics: Trends and

Applications. Cham: Springer International Publishing. p. 109-135,

https://doi.org/10.1007/978-3-319-31703-8_5

Armenteros JJA, et al. 2019. Detecting sequence signals in targeting peptides using deep

learning. Life Sci Alliance. 2: e201900429. https://doi.org/10.26508/lsa.201900429

Beagley CT, Okimoto R, Wolstenholme DR 1998. The mitochondrial genome of the sea

anemone *Metridium senile* (Cnidaria): introns, a paucity of tRNA genes, and a near-

standard genetic code. Genetics. 148: 1091-1108.

https://doi.org/10.1093/genetics/148.3.1091

Berná L, Alvarez-Valin F 2014. Evolutionary genomics of fast evolving tunicates. Genome

Biol Evol 6: 1724-1738. https://doi.org/10.1093/gbe/evu122

Bianchetti R, Lucchini G, Crosti P, Tortora P 1977. Dependence of mitochondrial protein

synthesis initiation on formylation of the initiator methionyl-tRNAf. J Biol Chem.

252: 2519-2523. https://doi.org/10.1016/S0021-9258(17)40489-3

Bliznina A, et al. 2021. Telomere-to-telomere assembly of the genome of an individual

Oikopleura dioica from Okinawa using Nanopore-based sequencing. BMC Genom.

22: 222. https://doi.org/10.1186/s12864-021-07512-6

Boore JL 1999. Animal mitochondrial genomes. Nucleic Acids Res. 27: 1767-1780.

https://doi.org/10.1093/nar/27.8.1767

Bouquet J-M, et al. 2009. Culture optimization for the emergent zooplanktonic model

organism *Oikopleura dioica*. J Plankton Res. 31: 359-370.

https://doi.org/10.1093/plankt/fbn132

Braun K, Leubner F, Stach T 2020. Phylogenetic analysis of phenotypic characters of

Tunicata supports basal Appendicularia and monophyletic Ascidiacea. Cladistics. 36:

259-300. https://doi.org/10.1111/cla.12405

Cardol P, et al. 2006. ND3 and ND4L subunits of mitochondrial complex I, both nucleus

encoded in *Chlamydomonas reinhardtii*, are required for activity and assembly of the

enzyme. Eukaryot Cell. 5: 1460-1467. https://doi.org/10.1128/ec.00118-06

Castellana S, Vicario S, Saccone C 2011. Evolutionary patterns of the mitochondrial genome

in Metazoa: exploring the role of mutation and selection in mitochondrial protein–

coding genes. Genome Biol Evol. 3: 1067-1079. https://doi.org/10.1093/gbe/evr040

Chorev M, Carmel L 2012. The function of introns. Front Genet. 3: 55.

https://doi.org/10.3389/fgene.2012.00055

Connell GJ, Byrne EM, Simpson L 1997. Guide RNA-independent and guide RNA-

dependent uridine insertion into *Cytochrome b* mRNA in a mitochondrial lysate from

Leishmania tarentolae: role of RNA secondary structure. Biol Chem. 272: 4212-4218.

https://doi.org/10.1074/jbc.272.7.4212

Crooks GE, Hon G, Chandonia JM, Brenner SE 2004. WebLogo: a sequence logo generator.

Genome Res. 14: 1188-1190. https://doi.org/10.1101/gr.849004

Cruz-Reyes J, Sollner-Webb B 1996. Trypanosome U-deletional RNA editing involves guide

RNA-directed endonuclease cleavage, terminal U exonuclease, and RNA ligase

activities. Proc Natl Acad Sci USA. 93: 8901-8906.

https://doi.org/10.1073/pnas.93.17.8901

Danks G, et al. 2012. OikoBase: a genomics and developmental transcriptomics resource for

the urochordate *Oikopleura dioica*. Nucleic Acids Res. 41: D845-D853.

https://doi.org/10.1093/nar/gks1159

Delahaye C, Nicolas J 2021. Sequencing DNA with nanopores: troubles and biases. PLOS

ONE. 16: e0257521. https://doi.org/10.1371/journal.pone.0257521

Delsuc F, Brinkmann H, Chourrout D, Philippe H 2006. Tunicates and not cephalochordates

are the closest living relatives of vertebrates. Nature. 439: 965-968.

https://doi.org/10.1038/nature04336

Delsuc F, et al. 2018. A phylogenomic framework and timescale for comparative studies of

tunicates. BMC Biol. 16: 1-14. https://doi.org/10.1186/s12915-018-0499-2

Denoeud F, et al. 2010. Plasticity of animal genome architecture unmasked by rapid evolution

of a pelagic tunicate. Science. 330: 1381-1385.

https://doi.org/10.1126/science.1194167

Dierckxsens N, et al. 2024. Tracing homopolymers in *Oikopleura dioica's*

mitogenome. bioRxiv. 2024.2005.2016.594446. https://doi.org/10.1101/2024.05.16.594446

Donath A, et al. 2019. Improved annotation of protein-coding genes boundaries in metazoan

mitochondrial genomes. Nucleic Acids Res. 47: 10543-10552.

https://doi.org/10.1093/nar/gkz833

Durrheim GA, Corfield VA, Harley EH, Ricketts MH 1993. Nucleotide sequence of

cytochrome oxidase (subunit III) from the mitochondrion of the tunicate *Pyura*

stolonifera: evidence that AGR encodes glycine. Nucleic Acids Res. 21: 3587-3588.

https://doi.org/10.1093/nar/21.15.3587

Eddy SR 2011. Accelerated profile HMM searches. PLoS Comput Biol. 7: e1002195. .

https://doi.org/10.1371/journal.pcbi.1002195

Erpenbeck D, et al. 2006. Mitochondrial diversity of early-branching Metazoa is revealed by

the complete mt genome of a haplosclerid demosponge. Mol Biol Evol 24: 19-22.

https://doi.org/10.1093/molbev/msl154

Felsenstein J 1985. Confidence limits on phylogenies: an approach using the bootstrap.

Evolution. 39: 783-791. https://doi.org/10.2307/2408678

Fulton TL, Wagner SM, Shapiro B. 2012. Case study: recovery of ancient nuclear DNA from

toe pads of the extinct passenger pigeon. In: Shapiro B, Hofreiter M, editors. Ancient

DNA: methods   and   protocols.   Totowa,   NJ: Humana   Press.   p.   29-35,

https://doi.org/10.1007/978-1-61779-516-9_4

Garić R 2024. World List of Appendicularia. Appendicularia. Accessed through: World

Register   of   Marine   Species.

https://www.marinespecies.org/aphia.php?p=taxdetails&id=146421

Garić R, et al. 2018. Challenges in appendicularian integrative taxonomy and barcoding.

ICES Annual Science Conference 2018.   https://www.imp-du.com/datoteke/Ices-

2018-rade-garic.pdf

Gissi C, Iannelli F, Pesole G 2004. Complete mtDNA of *Ciona intestinalis* reveals extensive

gene rearrangement and the presence of an *atp8* and an extra *trnM* gene in ascidians. J

Mol Evol. 58: 376-389. https://doi.org/10.1007/s00239-003-2559-6

Gissi C, Iannelli F, Pesole G 2008. Evolution of the mitochondrial genome of Metazoa as

exemplified by comparison of congeneric species. Heredity. 101: 301-320.

https://doi.org/10.1038/hdy.2008.62

Gissi C, Pesole G 2003. Transcript mapping and genome annotation of ascidian mtDNA using

EST data. Genome Res. 13: 2203-2212. https://doi.org/10.1101/gr.1227803

Gott JM, Emeson RB 2000. Functions and mechanisms of RNA editing. Annu Rev Genet. 34:

499-531. https://doi.org/10.1146/annurev.genet.34.1.499

Helfenbein KG, Fourcade HM, Vanjani RG, Boore JL 2004. The mitochondrial genome of

Paraspadella goto is highly reduced and reveals that chaetognaths are a sister group to

protostomes.   Proc   Natl   Acad   Sci   USA.   101:   10639-10643.

https://doi.org/10.1073/pnas.0400941101

Hoffmann RJ, Boore JL, Brown WM 1992. A novel mitochondrial genome organization for

the   blue   mussel,   *Mytilus   edulis*.   Genetics.   131:   397-412.

https://doi.org/10.1093/genetics/131.2.397

Iannelli F, Griggio F, Pesole G, Gissi C 2007. The mitochondrial genome of *Phallusia*

mammillata and *Phallusia fumigata* (Tunicata, Ascidiacea): high genome plasticity at

intra-genus level. BMC Evol Biol. 7: 155. https://doi.org/10.1186/1471-2148-7-155

Jumper J, et al. 2021. Highly accurate protein structure prediction with AlphaFold. Nature.

596: 583-589. https://doi.org/10.1038/s41586-021-03819-2

Kolmogorov M, et al. 2020. metaFlye: scalable long-read metagenome assembly using repeat

graphs. Nat Methods. 17: 1103-1110. https://doi.org/10.1038/s41592-020-00971-x

Lartillot N, Brinkmann H, Philippe H 2007. Suppression of long-branch attraction artefacts in

the animal phylogeny using a site-heterogeneous model. BMC Evol Biol. 7: S4.

https://doi.org/10.1186/1471-2148-7-S1-S4

Lartillot N, Philippe H 2004. A Bayesian mixture model for across-site heterogeneities in the

amino-acid   replacement   process.   Mol   Biol   Evol.   21:   1095-1109.

https://doi.org/10.1093/molbev/msh112

Lartillot N, Rodrigue N, Stubbs D, Richer J 2013. PhyloBayes MPI: Phylogenetic

reconstruction with infinite mixtures of profiles in a parallel environment. Syst Biol.

62: 611-615. https://doi.org/10.1093/sysbio/syt022

Laslett D, Canbäck B 2008. ARWEN: a program to detect tRNA genes in metazoan

mitochondrial nucleotide sequences. Bioinformatics. 24: 172-175.

https://doi.org/10.1093/bioinformatics/btm573

Latour AR 2021. Molecular divergence and gene flow in cold ocean tunicates: insights from

cytochrome oxidase 1. Diss Memorial University of Newfoundland. 73-87.

https://doi.org/10.48336/hkd1-9x29

Lavrov DV 2007. Key transitions in animal evolution: a mitochondrial DNA perspective.

Integr Comp Biol. 47: 734-743. https://doi.org/10.1093/icb/icm045

Lavrov DV, Adamski M, Chevaldonné P, Adamska M 2016. Extensive mitochondrial mRNA

editing and unusual mitochondrial genome organization in calcaronean sponges. Curr

Biol. 26: 86-92. https://doi.org/10.1016/j.cub.2015.11.043

Lavrov DV, et al. 2023. Phylomitogenomics bolsters the high-level classification of

Demospongiae (phylum Porifera). PLOS ONE. 18: e0287281.

https://doi.org/10.1371/journal.pone.0287281

Lavrov DV, Pett W 2016. Animal mitochondrial DNA as we do not know it: mt-genome

organization and evolution in nonbilaterian lineages. Genome Biol Evol. 8: 2896-

2913. https://doi.org/10.1093/gbe/evw195

Le TH, Blair D, McManus DP 2002. Mitochondrial genomes of parasitic flatworms. Trends

Parasitol. 18: 206-213. https://doi.org/10.1016/S1471-4922(02)02252-3

Lonergan KM, Gray MW 1996. Expression of a continuous open reading frame encoding

subunits 1 and 2 of cytochrome *c* oxidase in the mitochondrial DNA of *Acanthamoeba*

castellanii. J Mol Biol. 257: 1019-1030. https://doi.org/10.1006/jmbi.1996.0220

Martin M 2011. Cutadapt removes adapter sequences from high-throughput sequencing reads.

EMBnet j. 17: 3. https://doi.org/10.14806/ej.17.1.200

1052 Masunaga A, et al. 2022. The cosmopolitan appendicularian *Oikopleura dioica* reveals hidden

1053    genetic diversity around the globe. Mar Biol. 169: 157.

1054    https://doi.org/10.1007/s00227-022-04145-5

1055 Minh BQ, et al. 2020. IQ-TREE 2: New models and efficient methods for phylogenetic

1056    inference in the genomic era. Mol Biol Evol. 37: 1530-1534.

1057    https://doi.org/10.1093/molbev/msaa015

1058 Naville M, et al. 2019. Massive changes of genome size driven by expansions of non-

1059    autonomous transposable elements. Curr Biol. 29: 1161-1168.e1166.

1060    https://doi.org/10.1016/j.cub.2019.01.080

1061 Nishida H 2008. Development of the appendicularian *Oikopleura dioica*: culture, genome,

1062    and cell lineages. Dev Growth Differ. 50: S239-S256. https://doi.org/10.1111/j.1440-

1063    169X.2008.01035.x

1064 Ojala D, Montoya J, Attardi G 1981. tRNA punctuation model of RNA processing in human

1065    mitochondria. Nature. 290: 470-474. https://doi.org/10.1038/290470a0

1066 Okimoto R, Macfarlane JL, Clary DO, Wolstenholme DR 1992. The mitochondrial genomes

1067    of two nematodes, *Caenorhabditis elegans* and *Ascaris suum*. Genetics. 130: 471-498.

1068    https://doi.org/10.1093/genetics/130.3.471

1069 Pesole G, Gissi C, De Chirico A, Saccone C 1999. Nucleotide substitution rate of mammalian

1070    mitochondrial genomes. J Mol Evol. 48: 427-434.

1071    https://doi.org/10.1007/PL00006487

1072 Pett W, Lavrov DV 2015. Cytonuclear interactions in the evolution of animal mitochondrial

1073    tRNA metabolism. Genome Biol Evol. 7: 2089-2101.

1074    https://doi.org/10.1093/gbe/evv124

Pett W, et al. 2011. Extreme mitochondrial evolution in the ctenophore *Mnemiopsis leidyi*:

Insight from mtDNA and the nuclear genome. Mitochondrial DNA. 22: 130-142.

https://doi.org/10.3109/19401736.2011.624611

Philippe H, Zhou Y, Brinkmann H, Rodrigue N, Delsuc F 2005. Heterotachy and long-branch

attraction in phylogenetics. BMC Evol Biol. 5: 50.   https://doi.org/10.1186/1471-

2148-5-50

Pichon J, Luscombe NM, Plessy C 2020. Widespread use of the "ascidian" mitochondrial

genetic   code   in   tunicates.   F1000Research.   8:   2072-2072.

https://doi.org/10.12688/f1000research.21551.2

Plessy C, et al. 2024. Extreme genome scrambling in marine planktonic *Oikopleura dioica*

cryptic species. Genome Res. 34: 426-440.   https://doi.org/10.1101%2Fgr.278295.123

Raghavan V, Eichele G, Larink O, Karin EL, Söding J 2023. RNA sequencing indicates

widespread conservation of circadian clocks in marine zooplankton. NAR Genom

Bioinform. 5.   https://doi.org/10.1093/nargab/lqad007

Rogozin IB, Carmel L, Csuros M, Koonin EV 2012. Origin and evolution of spliceosomal

introns. Biol Direct. 7: 11.   https://doi.org/10.1186/1745-6150-7-11

Rubinstein ND, et al. 2013. Deep sequencing of mixed total DNA without barcodes allows

efficient assembly of highly plastic ascidian mitochondrial genomes. Genome Biol

Evol. 5: 1185-1199.   https://doi.org/10.1093/gbe/evt081

Salonna M, et al. 2021. An elongated COI fragment to discriminate botryllid species and as an

improved ascidian DNA barcode. Sci Rep. 11: 4078.   https://doi.org/10.1038/s41598-

021-83127-x

Sela I, Ashkenazy H, Katoh K, Pupko T 2015. GUIDANCE2: Accurate detection of

unreliable alignment regions accounting for the uncertainty of multiple parameters.

Nucleic Acids Res. 43: W7-W14.   https://doi.org/10.1093/nar/gkv318

Sereika M, et al. 2022. Oxford Nanopore R10.4 long-read sequencing enables the generation

of near-finished bacterial genomes from pure cultures and metagenomes without

short-read or reference polishing. Nat Methods. 19: 823-826.

https://doi.org/10.1038/s41592-022-01539-7

Shenkar N, Koplovitz G, Dray L, Gissi C, Huchon D 2016. Back to solitude: Solving the

phylogenetic position of the diazonidae using molecular and developmental

characters. Mol Phylogenetics Evol. 100: 51-56.

https://doi.org/10.1016/j.ympev.2016.04.001

Singh TR, et al. 2009. Tunicate mitogenomics and phylogenetics: peculiarities of the

Herdmania momus mitochondrial genome and support for the new chordate

phylogeny. BMC Genom. 10: 534. https://doi.org/10.1186/1471-2164-10-534

Spealman P, Burrell J, Gresham D 2020. Inverted duplicate DNA sequences increase

translocation rates through sequencing nanopores resulting in reduced base calling

accuracy. Nucleic Acids Res. 48: 4940-4945. https://doi.org/10.1093/nar/gkaa206

Stach T, Braband A, Podsiadlowski L 2010. Erosion of phylogenetic signal in tunicate

mitochondrial genomes on different levels of analysis. Mol Phylogenetics Evol. 55:

860-870. https://doi.org/10.1016/j.ympev.2010.03.011

Tamura K, Stecher G, Kumar S 2021. Mega11: molecular evolutionary genetics analysis

version 11. Mol Biol Evol. 38: 3022-3027. https://doi.org/10.1093/molbev/msab120

The UniProt Consortium 2022. UniProt: the universal protein knowledgebase in 2023.

Nucleic Acids Res. 51: D523-D531. https://doi.org/10.1093/nar/gkac1052

van Kempen M, et al. 2024. Fast and accurate protein structure search with Foldseek. Nature

Biotechnology. 42: 243-246. https://doi.org/10.1038/s41587-023-01773-0

[source code repository] Vieira GA 2023. mitos2_wrapper. GitHub.

https://github.com/gavieira/mitos2_wrapper/tree/master/refseq63m/featureProtHMM

Viklund H, Elofsson A 2004. Best alpha-helical transmembrane protein topology predictions

are achieved using hidden Markov models and evolutionary information. Prot Sci. 13:

1908-1917. https://doi.org/10.1110/ps.04625404

Wang H-C, Minh BQ, Susko E, Roger AJ 2017. Modeling site heterogeneity with posterior

mean site frequency profiles accelerates accurate phylogenomic estimation. Syst Biol.

67: 216-235. https://doi.org/10.1093/sysbio/syx068

Wang J, et al. 2022. The conserved domain database in 2023. Nucleic Acids Res. 51: D384-

D388. https://doi.org/10.1093/nar/gkac1096

Wang K, Omotezako T, Kishi K, Nishida H, Onuma TA 2015. Maternal and zygotic

transcriptomes in the appendicularian, *Oikopleura dioica*: novel protein-encoding

genes, intra-species sequence variations, and trans-spliced RNA leader. Dev Genes

Evol. 225: 149-159. https://doi.org/10.1007/s00427-015-0502-7

Waring RB, Davies RW 1984. Assessment of a model for intron RNA secondary structure

relevant to RNA self-splicing - a review. Gene. 28: 277-291.

https://doi.org/10.1016/0378-1119(84)90145-8

Wick RR, Judd LM, Gorrie CL, Holt KE 2017. Completing bacterial genome assemblies with

multiplex MinION sequencing. Microb Genom. 3: e000132.

https://doi.org/10.1099/mgen.0.000132

Wick RR, Judd LM, Holt KE 2019. Performance of neural network basecalling tools for

Oxford Nanopore sequencing. Genome Biol. 20: 129. https://doi.org/10.1186/s13059-

019-1727-y

Yahalomi D, et al. 2017. The multipartite mitochondrial genome of *Enteromyxum leei*

(Myxozoa): eight fast-evolving megacircles. Mol Biol Evol. 34: 1551–1556.

https://doi.org/10.1093/molbev/msx072

Yokobori S-I, et al. 1999. Complete DNA sequence of the mitochondrial genome of the

ascidian *Halocynthia roretzi* (Chordata, Urochordata). Genetics. 153: 1851-1862.

https://doi.org/10.1093/genetics/153.4.1851

Yokobori S-I, Ueda T, Watanabe K 1993. Codons AGA and AGG are read as glycine in

ascidian mitochondria. J Mol Evol. 36: 1-8. https://doi.org/10.1007/BF02407301

Yokobori S-I, Watanabe Y, Oshima T 2003. Mitochondrial genome of *Ciona savignyi*

(Urochordata, Ascidiacea, Enterogona): comparison of gene arrangement and tRNA

genes with *Halocynthia roretzi* mitochondrial genome. J Mol Evol. 57: 574-587.

https://doi.org/10.1007/s00239-003-2511-9

Zhu W, Lomsadze A, Borodovsky M 2010. *Ab initio* gene identification in metagenomic

sequences. Nucleic Acids Res. 38: e132-e132. https://doi.org/10.1093/nar/gkq275

1160

1161

1162 **Figure legend**

1163

1164 **Figure 1. *Oikopleura dioica* mitochondrial genome map.**

1165 Protein-coding genes are in green, rRNA genes (*rns*, *rnl*) in dark blue, tRNA-Met (UAU) in

1166 pink, coding sequences (CDS) in yellow, and poly-A/T stretches in light blue. The genes are

1167 pointing in the direction of their transcriptional orientation. A non-sequenced region,

1168 indicated as "gap", is in gray.

1169

Figure 2. A typical poly-A/T region in the *O. doica cox1* gene.

The reference sequence represents the assembled mitochondrial genome. The sequence of part

of the *cox1* gene is indicated in dark green. Above it are the nucleotide and amino-acid

sequences of the reference sequence. The poly-A/T region is in light blue. Reads from five

different sequencing sources mapped to this region are shown below the reference sequence:

our ONT data, our Illumina reads, public EST data from Norway, public RNA reads from

Japan, and public RNA reads from the North Sea (see Supplementary table S3A,

Supplementary Material online for accession numbers). Each line represents a different read.

Note that the different ONT reads do not have the same length, exemplifying the fact that the

length of the poly-T region could not be accurately determined. The EST and RNA reads lack

a long poly-U region. Rather, the poly-T region corresponds to 6U-bases. This difference

(long poly-T stretch in DNA *versus* 6U in the EST and RNA reads) provides evidence of

editing at the RNA level. Illumina reads can map to each side of the poly-T stretch, but

because poly-T stretches can be longer than the read length, we were unable to assemble the

entire region using Illumina only or to determine the length of the poly-T region.

1185

Figure 3. Presence-absence of aaRS orthologs in different species.

Single eukaryotic (Euk): single copy aaRSs that function both in the cytosol and

mitochondria. Other aaRSs are either expressed only in the cytosol (cytosolic) or in the

mitochondrion (mitochondrial). aaRSs presence is shown in pink, aaRSs absence in white,

and aaRS-Gly in blue. Protein accession numbers are provided in Supplementary table S5,

Supplementary Material online.

1192

Figure 4. The position of *O. dioica* among chordates.

The phylogenetic tree was reconstructed based on mitochondrial protein sequences under the

mtZOA+EHO+F+I+G4. Branch supports are only indicated for those cases in which the

bootstrap support values were lower than 100%. The tunicate classes: Stolidobranchia in blue,

Thaliacea in red, Phlebobranchia in green, Aplousobranchia in yellow, and Appendicularia in

purple

1199

1200 .

1201 **Table**

1202

1203 **Table 1.** *Oikopleura dioica* **genes, position, length, direction (F- forward, R- reverse),**

1204 **start and stop codons, and comparison to the minimal, maximal, and average length of**

1205 **orthologous genes in 38 tunicate sequences (supplementary table S4, Supplementary**

1206 **Material online).**

1207

| Gene | Type | Position | | Length [bp] | | Direction | Codon | | Length range in tunicates [bp] | | Average length in tunicates [bp] |
		start	end	gene	CDS		start	stop	min	max	
rnl	rRNA	13	303	291	242	R			921	1,299	1,105
nad4	gene	934	2,847	1,914	876	R	ATC	TAA	1,209	1,401	1,321
cox3	gene	2,906	4,574	1,669	870	F	ATT	TAA	768	795	786
tRNA-Met	tRNA	4,671	4,763	93	69	F			61	76	66
nad1	gene	4,805	5,800	996	615	F	TTG	TAA	853	1,035	914
cox1	gene	5,836	7,834	1,999	1,512	F	ATT	TAG	1,482	1,575	1,546
ODI_mt_ ORF196	gene	8,462	9,887	1,426	591	R	TTG	TAA			
cytb	gene	10,115	12,085	1,971	993	F	GTG	TAA	1,035	1,122	1,093
atp6	gene	12,312	13,063	752	516	R	ATT	TAA	492	642	619
rns	rRNA	13,064	13,295	232	215	R			596	984	696
cox2	gene	13,305	14,301	997	582	F	ATT	TAA	651	942	689
nad5	gene	14,456	18,371	3,916	1,248	F	ATA	TAA	1,458	1,788	1,695

1208

1209

1210

Oikopleura dioica mitochondrial genome 18,390 bp

	AA	Oikopleura dioica Norway	Oikopleura dioica Osaka	Ciona intestinalis	Styela clava	Homo sapiens
Single Euk.	A	■	■	■	■	■
	G			◆	◆	◆
	H	■	■	■	■	■
	Q	■	■	■	■	■
Cytosolic	G	◆	◆			
	C	■	■	■	■	■
	D	■	■	■	■	■
	E/P	■	■	■	■	■
	Fa	■	■	■	■	■
	Fb	■	■	■	■	■
	I	■	■	■	■	■
	K	■	■	■	■	■
	L	■	■	■	■	■
	M	■	■	■	■	■
	N	■	■	■	■	■
	R	■	■	■	■	■
	S	■	■	■	■	■
	T	■	■	■	■	■
	V	■	■	■	■	■
	W	■	■	■	■	■
	Y	■	■	■	■	■
Mitochondrial	G	◆	◆			
	C			■	■	■
	D			■	■	■
	E			■	■	■
	F	■	■	■	■	■
	I			■	■	■
	K					
	L			■	■	■
	M	■	■	■	■	■
	N			■	■	■
	P			■	■	■
	R			■	■	■
	S			■	■	■
	T					
	V					■
	W			■	■	■
	Y			■	■	■