Deciphering the mitochondrial genome of *Oikopleura dioica* (Chordata: Tunicata:

Appendicularia)

Yael Klirs[1], Maria Novosolov[1,2], Carmela Gissi[3,4,5], Rade Garić[6], Tal Pupko[7], Thomas Stach[8],

Dorothée Huchon[1,9]

[1]School of Zoology, George S. Wise Faculty of Life Sciences, Tel Aviv University, Tel Aviv

6997801, Israel

[2]GeoGenetics Centre, GLOBE Institute, Faculty of Health and Medical Sciences, University

of Copenhagen, Copenhagen, Denmark.

[3]Department of Biosciences, Biotechnology and Environment, University of Bari Aldo Moro,

70126 Bari, Italy

[4]Institute of Biomembranes, Bioenergetics and Molecular Biotechnologies, Consiglio

Nazionale delle Ricerche, 70126 Bari, Italy

[5]CoNISMa, Consorzio Nazionale Interuniversitario per le Scienze del Mare, 00196 Roma,

Italy

[6]Institute for Marine and Coastal Research, University of Dubrovnik, Kneza D. Jude 12,

20000 Dubrovnik, Croatia

[7]The Shmunis School of Biomedicine and Cancer Research, George S. Wise Faculty of Life

Sciences, Tel Aviv University, Tel Aviv, 6997801 Israel

[8]Department of Molecular Parasitology, Institute of Biology, Humboldt-Universität zu Berlin,

Berlin, Germany

[9]The Steinhardt Museum of Natural History and National Research Center, Tel Aviv

University, Tel Aviv 6997801, Israel

ORCID IDs: 0009-0006-2572-0612 (Yael Klirs) 0000-0002-4034-3441 (Maria Novosolov),

0000-0002-2269-079X (Carmela Gissi), 0000-0001-6477-5559 (Rade Garić), 0000-0001-

9463-2575 (Tal Pupko), 0000-0001-5461-9069 (Thomas Stach), 0000-0002-5047-244X

(Dorothée Huchon).

* Author for Correspondence: Dorothée Huchon, School of Zoology, George S. Wise

Faculty of Life Sciences, Tel Aviv University, Tel Aviv 6997801, Israel, +972 36409817,

huchond@tauex.tau.ac.il

Abstract

Sequencing the mitochondrial genome of the tunicate *Oikopleura dioica* is a challenging task because of the presence of long poly-T/A homopolymer stretches, which impair sequencing and assembly. Here, we report the sequencing and annotation of the majority of the mitochondrial genome of *O. dioica* by combining several DNA and amplicon reads obtained by Illumina and MinIon Oxford Nanopore Technologies (ONT) with public RNA sequences. We document extensive RNA editing, since all homopolymer stretches present in the mitochondrial DNA correspond to 6U-regions in the mitochondrial RNA. Out of the 13 canonical protein-coding genes, we were able to detect eight plus an unassigned ORF, which lacked sequence similarity to canonical mitochondrial protein-coding genes. We showed that the *nad3* gene has been transferred to the nucleus and acquired a mitochondria-targeting signal. In addition to two very short rRNAs, we could only identify a single tRNA (tRNA-Met), suggesting multiple losses of tRNA genes, supported by a corresponding loss of mitochondrial aminoacyl-tRNA synthetases in the nuclear genome.

Based on the eight canonical protein-coding genes identified, we reconstructed maximum likelihood and Bayesian phylogenetic trees and inferred an extreme evolutionary rate of this mitochondrial genome. However, the phylogenetic position of appendicularians among tunicates could not be accurately determined.

Keywords

mtDNA, phylogenomics, Oxford Nanopore Technologies, RNA editing, tRNA loss, Mitochondrial-nuclear gene transfer

Significance

Sequencing and annotating the mitochondrial genome of fast-evolving organisms is difficult because they often present unusual characteristics. The tunicate *O. dioica* is a model species for understanding tunicate and chordate genome evolution. However, no complete annotated mitochondrial genome for this species has been published to date. Here, we determined the major part of the mitochondrial genome of *O. dioica*. Our results indicate the presence of highly modified rRNA genes and the absence of all tRNAs except tRNA-Met. Moreover, we show that the mitochondrial genome undergoes editing at the RNA level. Our study demonstrates that a combination of public RNA data and DNA from long-and short-read sequencing platforms significantly improves our ability to study mitochondrial genomes with atypical characteristics.

Introduction

Most mitochondrial genomes of Bilateria are circular DNA molecules of approximately 14-19 kb in length, characterized by the absence of introns and the presence of only short intergenic regions (Lavrov and Pett 2016). These genomes typically contain 37 genes: 13 protein-coding genes: ATP synthase subunits 6 and 8 (*atp6* and *atp8*), cytochrome oxidase subunits (*cox1, cox2, cox3*), cytochrome b (*cytb*), and dehydrogenase subunits (*nad1, nad2, nad3, nad4, nad5, nad6*, and *nad4L*); two ribosomal genes (rRNA): small-subunit (*rns*) and large-subunit (*rnl*) and ~22 tRNAs (Boore 1999; Gissi, et al. 2008; Rubinstein, et al. 2013). However, bilaterian mitochondrial genomes are substantially diverse in their structure, organization, evolutionary rate, and genetic code (Lavrov 2007).

Tunicate mitochondrial genomes are fast evolving, both in terms of substitutions (Singh, et al. 2009) and genome rearrangements (Gissi, et al. 2008; Stach, et al. 2010). As a result of this high rearrangement rate, there is no standard tunicate mitochondrial gene order. In addition, the ascidian mitochondrial genetic code differs from the vertebrate one. Of note, the "ascidian" genetic code, which was initially determined based on the mitochondrial sequences of *Halocynthia roretzi* and *Pyura stolonifera* (Durrheim, et al. 1993; Yokobori, et al. 1993), seems to be valid for all other tunicate classes (Pichon, et al. 2019).

Tunicata are divided into three classes: Ascidiacea, Thaliacea, and Appendicularia. Appendicularia is a key class for understanding tunicate and chordate genome evolution, as it is an early diverging clade within tunicates. *Oikopleura dioica* is a frequent model tunicate for the Appendicularia class in developmental biology, evolutionary biology, and comparative genomics research (Delsuc, et al. 2006; Nishida 2008). Despite the importance of this clade, no complete annotated appendicularian mitochondrial genome is currently available in public databases. One non-annotated mitochondrial genome of *O. dioica* from Okinawa Japan is available (accession OU015571; 9,225 bp; (Bliznina, et al. 2021)).

The mitochondrial genome of *O. dioica* from Norway has been partially reconstructed

by Denoeud *et al.* (2010) from genomic DNA and EST data. However, the obtained

mitochondrial contigs of this genome are not publicly available. This genome has been

inferred to harbor long poly-T/A stretches interrupting the open reading frames (ORFs) and it

has been proposed that these interrupted ORFs are corrected by RNA editing (Denoeud, et al.

2010). Such homopolymer stretches can neither be sequenced using Sanger sequencing nor

using Illumina short-read sequencing, since they are error hot spots for the DNA polymerases

used in these sequencing techniques. Thus, together with the difficulty in isolation and

identification of appendicularian species, this peculiarity explains the paucity of publicly

available mitochondrial data for *O. dioica*. Annotation of the resulting genome is also

expected to be challenging, due to RNA editing. RNA editing describes modifications such as

insertions, deletions, and base substitutions of RNA transcripts. As a result, the RNA and

computationally inferred protein sequences differ from the DNA template (Gott and Emeson

2000). To complicate matters, in *O. dioica* RNA editing seems to be prevalent in the same

regions that are hard to sequence, i.e., the poly-T/A stretches. These stretches appear as 6U-

sites in mitochondrial RNA sequences, regardless of the number of T/A at the DNA level

(Denoeud, et al. 2010).

The phylogenetic analyses performed by Denoeud *et al.* (2010) concluded that *O.*

dioica mitochondrial protein-coding genes are very fast evolving, even when compared to the

already high rate of other tunicate mitochondrial genomes. Using a combination of PCR and

EST mapping approaches they determined the order of nine out of the 13 mitochondrial

protein-coding genes: *cox3, nad1, cox1, cytb,* a putative *nad2*, *atp6, cox2, nad5,* along a large

mitochondrial contig and *nad4* on a separate contig. They did not identify any tRNA or rRNA

genes. Unfortunately, the mitochondrial contigs obtained by Denoeud *et al*. (2010) have not

been deposited in public databases and only the raw ESTs data are available in the Oikobase

database (Danks, et al. 2012).

A draft mitochondrial genome scaffold of *O. dioica* from Okinawa (Japan) has been

published by Bliznina, et al. (2021) (accession OU015571; 9,225 bp), with the annotation

reported only in the publication (Figure 7 of their article). The annotation obtained is,

however, preliminary as several genes contain stop codons and frameshifts. Unlike Denoeud,

et al. (2010), the mitochondrial genome is presented as a circular molecule that encompasses

seven mitochondrial genes (Bliznina, et al. 2021). The mitochondrial gene order resembles

the order from Figure S4 of Denoeud, et al. (2010). Yet, some differences exist. First, the

nad2 and *nad5* genes are missing from the annotation of Bliznina, et al. (2021). Second, the

rnl and four tRNAs: tRNA-Met, tRNA-Trp, tRNA-Ala, and tRNA-His are annotated. Third,

they identified a LAGLIDADG homing endonuclease in the middle of the *cox1* gene (Figure

7 of Bliznina et al. 2021). It should be noted that *O. dioica* from Okinawa most likely

represents a different species from *O. dioica* from Norway (Masunaga, et al. 2022).

Here, we determine the major part of the mitochondrial genome sequence of *O. dioica*

from Norway, its gene content and its order. We observe substantial gene loss compared to

other metazoan mitochondrial genomes and provide evidence for extensive RNA editing.

Results

Sequencing the Mitochondrial Genome

DNA extracted from *O. dioica* individuals from Norway was sent to Illumina sequencing.

Since, it was impossible to assemble these reads because of the prevalence of homopolymers

(poly-T or poly-A stretches), we next performed long-read sequencing using Oxford

Nanopore Technology (ONT) sequencing. However, in this case, the flow-cell pores became

blocked within an hour (Minion sequencing was performed twice by different companies),

resulting in too little data for reliable assembly. Therefore, we PCR-amplified the

mitochondrial genome in 12 fragments (supplementary Fig. S1, Supplementary Material

online) and sequenced all barcoded fragments using a single ONT flow cell. Primers were

designed based on the *O. dioica* EST and RNA data available in public databases (see

Methods and supplementary table S1, Supplementary Material online). We successfully

assembled each PCR fragment (except one), and using the Illumina data, we obtained a single

contig. Unfortunately, we could not circularize the genome because we could not entirely

assemble one fragment. The length of *O. dioica* mitochondrial contig assembled is 18,390 bp

(Figure 1). This genome size is larger than the average of other tunicates, 14,808 ± 821 bp,

and compatible with the PCR prediction of a ~19,000 bp genome (supplementary table S1,

Supplementary Material online).

Multiple instances of poly-T/A stretches were found in the assembled contig (Figure

1, supplementary Fig. S2, Supplementary Material online). The exact number of either A/T

repeats within poly-T/A regions could not be accurately determined. These stretches are too

long to be fully contained in the short Illumina sequences (Ari and Arikan 2016), while ONT

basecallers do not accurately determine the length of homopolymers (Wick, et al. 2019).

Moreover, the length of homopolymers is correctly resolved up to a length of 11 bp in R10.4

flow cell (Sereika, et al. 2022), while in the R9.4.1 flow cell, a good accuracy (70% errorless)

is observed only in homopolymers up to 5 bp (Delahaye and Nicolas 2021). However, in our

case, the length of the 65 poly-T/A stretches found in the mitochondrial sequence ranged from

2 to 509 bp, with an average of 143 ± 103.5 bp (See Methods, supplementary table S2,

Supplementary Material online), which is considerably larger. Moreover, it has been observed

that regions harboring inverted repeats are poorly sequenced using ONT (Spealman, et al.

2020). When in a PCR-fragment a poly-A region is followed by a poly-T region, these two

homopolymers can form structures similar to inverted repeats, thus disrupting ONT

sequencing. In addition, Taq polymerase slippage errors can further contribute to uncertainty

associated with the length of long homopolymers. All these factors contribute to

heterogeneity among ONT reads that map to the same mitochondrial locus harboring a poly-

T/A region. One such example is shown in Figure 2. Note, that these stretches are sometimes

imperfect, i.e., there may be a few G residues within a poly-A stretch or a few C residues with

a poly-T stretch (Figure 2). These G and C residues are unlikely to reflect ONT sequencing

errors, as they could sometimes be verified using Illumina reads. In total, these poly-T/A

stretches span ~9,200 bp. This suggests that while the total mitochondrial genome size is

longer than in other tunicates, the mitochondrial genome without the poly-T/A stretches is, in

fact, shorter.

Evidence for rRNA Editing

A total of 65 poly-T/A sites were found in the assembled *O. dioica* mitochondrial genome. By

mapping (1) the mitochondrial transcripts identified from the EST from Norway, and (2) the

RNA contigs assembled from publicly available RNA reads from mainland Japan (Wang, et

al. 2015) and the North Sea (Raghavan, et al. 2023) onto the DNA sequence assembled from

the ONT reads, we were able to confirm the editing of 59 of these 65 poly-T/A sites to 6U-

sites (see an example in Figure 2 and supplementary table S2, Supplementary Material

online). For the remaining 6 sites, no direct RNA evidence for editing exists as we could not

detect RNA reads or ESTs that mapped to the corresponding region. These poly-T/A stretches

are present both in coding and noncoding mitochondrial DNA regions. Specifically, we

observed that exactly six consecutive U residues (6U-sites) in the RNA corresponded to a

poly-T stretch in the DNA sequence. These results suggest that the longer U stretches in the

pre-mRNA are edited, resulting in only shorter stretches of 6U. The exact nature of this RNA

editing is unknown. It is, however, remarkable that protein-coding genes as well as the two

rRNAs and the tRNA-Met only include poly-T regions and not poly-A regions. When poly-A

regions are found on the direct strand, it indicates that the genes are encoded on the reverse

strand. We detected a few instances of EST reads that include a poly-U stretch, i.e., before the

excision of poly-U to 6U. These observations suggest that the shortening of the poly-U

stretch, which is necessary to recover the reading frame, occurs at the RNA level co-

transcriptionally or very fast after the synthesis of the precursor transcript.

Interestingly we noticed that while poly-T regions are present within all genes, the

genes often end with a poly-A. The poly-A ending is part of the TAA stop codon in most

protein-coding genes, except for *cox1*, which has a TAG stop codon (Table 1). The contigs

assembled from RNA reads support the view that these poly-A stretches are reduced to 6A in

the mRNA. Similarly, we noticed that a poly-A stretch is often present in the DNA before the

start of a protein-coding gene (Figure 1). No conserved sequence motif could be identified

around the poly-T/A stretches. The only major departure we could detect around the poly-T/A

stretches is the absence of cytosine just after the 6T site. No other signal could be identified

(Supplementary Fig. S3, Supplementary Material online).

***Oikopleura dioica* Mitochondrial Genome Sequence and Structure**

Figure 1 presents the assembled mitochondrial genome of *O. dioica*. We could not determine

the sequence in the region labeled as "Gap" between the *nad5* and *rnl* genes, probably

because of the presence of secondary structures that impaired sequencing. Running the PCR

product of fragment 12 on a gel suggests that the length of the "Gap" region is approximately

400 bp (supplementary Fig S1, Supplementary Material online). However, the fact that we

could amplify this missing region using primers from both sides of the gap suggests that the

genome is circular. Notably, although we could not assemble the entire fragment 12, the

ONT-contigs we obtained allowed us to determine the end of the *nad5* gene and the *nad4-rnl*

region, supporting the view that we amplified the correct region of the mitochondrial genome.

The circularity of the mitochondrial genome is in line with Bliznina, et al. (2021).

Gene Content and Gene Loss in *O. dioica* Mitochondrial Genome

Most tunicate mitochondrial genomes encompass 13 protein-coding genes, two rRNA genes,

and 24 tRNA genes that are encoded on the same strand (Shenkar, et al. 2016; Singh, et al.

2009). Surprisingly, in the mitochondrial genome of *O. dioica* we could only identify eight

canonical protein-coding genes: ATP synthase subunit 6, subunits 1-3 of cytochrome c

oxidase (*cox*), cytochrome *b* (*cytb*), and NAD dehydrogenase (*nad*) 1, 4, and 5. In addition to

the two rRNA genes encoding the small and large subunits, we could identify a single tRNA

gene, tRNA-Met (Figure 1, Table 1)*.* The genes *cox1, cox2, cox3, cytb, nad1, nad5*, and

tRNA-Met (UAU) are encoded on the forward strand, while *atp6, nad4*, *rns*, *rnl,* and an

unidentified protein-coding gene named *ODI_mt_ORF196* (see below) are encoded on the

reverse strand (Table

Table 1). No incomplete stop codons were found (Table 1). Moreover, no LAGLIDADG

ORF was found to be encoded within the *cox1* gene. These results suggest the loss of five

protein-coding genes: *atp8, nad 2, 3, 4L*, and *6* and all tRNA genes except for tRNA-Met.

An additional mitochondrial protein-coding gene was identified following a Blastn

search of the obtained mitochondrial sequence against the EST data available in Oikobase

(Danks, et al. 2012). Notably, programs such as "Find ORFs" (implanted in Geneious Prime

v.2023.2.1.) are inapplicable for this mitochondrial genome, because of the extreme fast rate

of evolution and the numerous poly-T/A stretches that disrupt the reading frames of protein-

coding genes. This additional ORF is 1,426 bp long and does not show any significant

sequence similarity to any known protein-coding gene present in NCBI public database (i.e.,

based on Blastp searches). We thus dubbed it gene "*ODI_mt_ORF196*" based on its amino-

acid length (Figure 1, Table 1). This protein was inferred to have six transmembrane domains

and, based on its position, it corresponds to the putative *nad2* gene identified by Denoeud, et

al. (2010). An Alpha fold-based prediction of this gene (supplementary Fig. S4,

Supplementary Material online) was used as query against multiple protein structure

databases: AlphaFold/UniProt50 v4, AlphaFold/Swiss-Prot v4, AlphaFold/Proteome v4,

CATH50 4.3.0, MGnify-ESM30 v1, PDB100 2201222, GMGCL 2204 from Foldseek (van

Kempen, et al. 2024). No significant structural hits were obtained.

We identified the two rRNA genes, *rns* and *rnl,* by searching for highly transcribed

regions along unannotated regions of the assembled genome. Two noncoding regions were

found to be highly expressed based on Illumina RNA read mapping (supplementary Fig. S5,

Supplementary Material online). Further support for the transcription of these regions is their

presence in the EST data. Specifically, the longer expressed non-coding region that resides

between the "gap region" and the *nad4* gene was assumed to be the *rnl* gene. This annotation

is supported by RNA structure similarity between the putative *rnl* and structure model profile

of *rnl* genes, as implemented in Mitos2 (Donath, et al. 2019). The second, shorter expressed

noncoding region, located between the *atp6* and the *cox2* genes, was annotated as *rns*,

although no structural similarity was found between this sequence and *rns* structure profile in

Mitos2. Both the *rnl* and *rns* genes were found to be extremely short: 291 and 232 bp,

respectively, compared to 1,129 and 660 bp in the model tunicate *Ciona intestinalis* (see also

Table 1).

We could identify a single tRNA gene, tRNA-Met (UAU). This tRNA presents a

typical secondary cloverleaf structure (supplementary Fig. S6, Supplementary Material

online), and UAU anti-codon, corresponding to a methionine codon according to the ascidian

mitochondrial code. Unlike in Bliznina, et al. (2021) no other tRNAs were identified.

Aminoacyl-tRNA Synthetase (aaRS) Profiling

We used hidden Markov models (HMMs) to identify nuclear-encoded aaRSs in published

sequences of *O. dioica* and few other chordate representatives (Figure 3, supplementary table

S4, Supplementary Material online). We found the presence of all cytosolic aaRSs in

published proteomes of *O. dioica* (accessions: GCA_000209535, Norway and

GCA_907165135, Okinawa) (Figure 3). However, in contrast to the other chordates

investigated, *O. dioica* is inferred to have lost most of the mitochondrial aaRSs except for

aaRS-Gly (G), aaRS-Phe (F), and aaRS-Met (M). Likewise, the mitochondrial aaRS-Val (V)

is absent in the tunicates *C. intestinalis* (GCA_000224145) and *Styela clava*

(GCF_000001405), and the mitochondrial aaRS-Lys (K) and aaRS-Thr (T) are absent in all

species included in this comparison. Furthermore, *O. dioica* encodes two copies of aaRS-Gly

(one cytosolic and the other mitochondrial) while the other chordates encode only a single

copy that functions both in the cytosol and the mitochondrion, suggesting a duplication event

in the lineage leading to appendicularians.

Gene Transfer to the Mitochondrial Genome

HMMs were used to search for the missing canonical mitochondrially-encoded genes (i.e.,

nad2, nad3, *nad4L*, *nad6,* and *atp8*) in the nuclear genome. Using these models, we found a

protein sequence with a significant similarity (E-value < 5e-07) to the NAD3 model in the

two published proteomes of *O. dioica* (proteins CAG5094752 and CBY12996 for the

Norwegian and Okinawan genomes, respectively). A domain search confirmed that the

protein sequence identified includes a NAD-ubiquinone/plastoquinone oxidoreductase, chain

3 domain (Cdsearch (Wang, et al. 2022) search using the CAG5094752 and CBY12996

protein sequences as queries against the CDD v3.21 database yielded E-values of 4.94e-03

and 3.80e-03, respectively). This protein was predicted to include two putative

transmembrane domains, in support of its role in the electron transport chain (supplementary

Fig. S7, Supplementary Material online). This protein-coding gene was found to be encoded

on the X specific region of the X chromosome of *O. dioica* and to harbor three introns.

Conversely, no nuclear-encoded protein showing significant similarity to the NAD3 model

was found in the proteomes of *C. intestinalis*, *S. clava* and *H. sapiens*.

To confirm that the identified NAD3 proteins of the Norwegian and Japanese *O.*

dioica are expressed in the mitochondria we search for the presence of mitochondrial target

peptides. A significant target peptide signal was found in these putative NAD3 proteins with a

likelihood above 0.89. We could not confirm the transfer of other missing genes (i.e., *nad2*,

nad4L, *nad6,* and *atp8*) to the nuclear genome: this can be due to the fast-evolutionary rate of

O. dioica and the short length of these proteins (150-152 amino-acid), or to the

incompleteness of proteomic data for *O. dioica*.

Phylogenetic Reconstructions

Phylogenetic reconstructions were performed using two datasets. In the first, *Doliolum*

nationalis was the sole representative of the tunicate class Thaliacea. The second dataset

differed from the first by including two additional thaliacean sequences: the fast-evolving

sequences of *Salpa thompsoni* and *Salpa fusiformis* (supplementary table S4, Supplementary

Material online). Phylogenetic trees were reconstructed under the maximum likelihood (ML)

criterion using various models of sequence evolution implemented in IQtree (Minh, et al.

2020; Wang, et al. 2017), including complex models (i.e., MtZoa+F+G+I,

MtZoa+EHO+F+G+I, Poisson+C10+G, Poisson+C20+G) and under the Bayesian CAT

model implemented in Phylobayes (Lartillot and Philippe 2004). In all trees the monophyly of

tunicates was supported with maximal support (bootstrap percentage (BP) = 100, posterior

probability (PP) = 1.0), however, the position of *O. dioica* was highly variable depending on

the dataset and the evolutionary model chosen (Figure 4, supplementary Fig. S8 A-J,

Supplementary Material online). When the tree was reconstructed without members of the

Salpa genus, *O. dioica* was the first tunicate to diverge when using the MtZoa+F+G+I or the

MtZoa+EHO+F+G+I models (see Figure 4 and supplementary Fig. S8 A, Supplementary

Material online). Remarkably, the branch support values for the early divergence of *O. dioica*

were low (BP < 50%). Conversely, support for the monophyly of accepted tunicate clades

(i.e., Stolidobranchia and Thaliacea+Phlebobranchia+Aplousobranchia) was moderate to high

(BP = 60-93%). When using more complex models such as the Poisson+C20+G

(supplementary Fig. S8 C, Supplementary Material online) or the Bayesian CAT model

(supplementary Fig. S8 H-J, Supplementary Material online), *O. dioica* branched within

Stolidobranchia. This positioning and the resulting relationships among Stolidibranchia clades

were not supported (BP < 50%; PP < 0.5). When members of the genus *Salpa* were included

in the phylogenetic analyses, *O. dioica* systematically grouped with the *Salpa* members

except when using the most complex Poisson+C20+G ML model (supplementary Fig. S8 G,

Supplementary Material online). Interestingly, support for this relationship decreased with the

complexity of the ML model used (BP = 85%, MtZOA+G+F; BP = 77%, MtZoa+EHO+G+F,

BP = 51%, Poisson+C10+G) (supplementary Fig. S7 D-F, Supplementary Material online). In

the tree reconstructed with the Poisson+20+G model, the Thaliacea monophyly was recovered

(BP = 59%) but *O. dioica* branched within Stolidobranchia, and basal relationships within

tunicates were not supported (BP < 50%). When using the Bayesian CAT model, the

phylogenetic position of *O. dioica* could not be resolved when *Salpa* was included.

Specifically, convergence parameters suggested that convergence was achieved, however,

different topologies were obtained depending on the chains considered. With chains 1, 2 and 3

the Thaliacea monophyly was recovered but not supported (PP = 0.53). When chains 1, 2, and

4 were considered, *Salpa* and *O. dioica* were found to be sister clade (PP = 0.57)

(supplementary Fig. S8 I-J, Supplementary Material online).

Discussion

The mitochondrial genome of *O. dioica* appears extremely divergent and fast-evolving. It is

characterized by the presence of numerous poly-T/A stretches undergoing RNA editing. A

few canonical metazoan mitochondrial protein genes and almost all tRNA genes are absent

from the genome, there is a substantial reduction in the size of the rRNA genes, and there is

evidence of a protein-coding gene transfer to the nuclear genome. Such high mitochondrial

genome plasticity characterizes fast-evolving animal clades (Pett, et al. 2011; Yahalomi, et al.

2017).

Protein-Coding Genes

Five canonical protein-coding genes are absent (*atp8, nad2, nad3, nad4L,* and *nad6*) in the

mitochondrial genome of *O. dioica*, four of which were not identified in the nuclear genome

(i.e., all excluding *nad3*). These four genes may be encoded in either the mitochondrial or the

nuclear genome, but we were unable to identify them. Indeed, the *O. dioica* mitochondrial and

nuclear genomes are extremely fast evolving, and consequently, these genes may have been

missed by algorithms for remote homology detection, including HMM searches. Notably,

these missing genes (*atp8, nad2, nad4L,* and *nad6*) are among the fastest evolving

mitochondrial genes (Castellana, et al. 2011; Pesole, et al. 1999). The unassigned gene

ODI_mt_ORF196 is predicted to encode a protein harboring four alpha helices

(supplementary figure S4, supplementary material online), unlike all other missing proteins in

tunicates: the NDH2 protein of most metazoans harbors six alpha helices and the other

canonical mitochondrial proteins are predicted to harbor three or fewer alpha helices (The

UniProt The UniProt Consortium 2022). Moreover, none of the methods used (Blastp,

HMMer, and Cdsearch) detected a significant similarity with any other canonical proteins.

We thus refrained from annotating this gene as one of the four missing gene, although

Denoeud *et al.* (2010) annotated this gene as *nad2*.

After removing poly-T/A regions, the largest non-coding region identified is 206 bp

(between *ODI_mt_ORF196* and *cytb*) and it is possible that at least one among the *atp8* and

nad4L genes, which are shorter than 300 bp in most tunicates, are encoded within this region.

The *atp8* gene, in particular, has been overlooked in the past from the original annotation of

several mitochondrial genomes (e.g. Helfenbein, et al. 2004; Hoffmann, et al. 1992; Le, et al.

2002; Okimoto, et al. 1992), including tunicates (Yokobori, et al. 1999; Yokobori, et al. 2003)

and was then identified in ascidians following mitochondrial EST analyses (Gissi, et al. 2004;

Gissi and Pesole 2003).

Our results suggest that *nad3* has been relocated to the nuclear genome. While this is

the first evidence for relocation of this mitochondrial gene to the nucleus in animals, a case of

nuclear *nad3* has been reported in the green alga *Chlamydomonas reinhardtii* together with

the transfer of *nad4L* (Cardol, et al. 2006). Similar transfers to the nuclear genome have been

described for other animal mitochondrial genes, for example, the *atp6* and *atp9* genes of the

ctenophore *Mnemiopsis leidyi* (Pett, et al. 2011), and the mitochondrial gene *atp9* in the

demosponge *Amphimedon queenslandica* (Erpenbeck, et al. 2006). Short ORFs are more

difficult to identify than long ones, and because *atp8* and *nad4L* are relatively short, it is

possibly that this factor, in additional to their high rate of evolution, impaired their

identification. The fact that *nad6* was detected neither in the mitochondrial genome nor in the

nuclear one, is more surprising. This gene is longer (450.9 bp ± 20.2 bp in other tunicates),

and except for the gene ODI-mt-*ORF196,* no other putative ORFs longer than 300 bp were

detected. The possible incompleteness of the two public *O. dioica* assemblies and the inferred

proteomes, or the specific criteria and bioinformatic pipeline used to annotate the protein

genes in these nuclear assemblies, could also explain the failure to identify *nad6*, *atp8* and

nad4L in the available nuclear assemblies.

A third explanation could be that the genes that we could not identify have been fused

with other mitochondrial genes. For example, the ORF we annotated as *atp6* could include

regions corresponding to both *atp6* and *atp8*. An example of ORF fusion is found in the

*cox*1/2 gene of amoebas (e.g., Lonergan and Gray 1996). However, this explanation is

unlikely in our case since all *O. dioica* ORFs are shorter than their tunicate counterparts

(Table 1), while in the case of a fusion, we would expect these ORFs to be longer.

rRNA Genes

Tunicate rRNA genes are known to be poorly conserved. As a case in point, the boundaries of tunicate mitochondrial rRNA genes are mostly inferred from the end and start of the flanking genes, indicating that the length estimates of rRNA genes are error-prone (Rubinstein, et al. 2013; Singh, et al. 2009). Only for the ascidians *Halocynthia roretzi* and *C. intestinalis*, some rRNA boundaries were tentatively better defined based on EST transcript data (Gissi and Pesole 2003). Although the two rRNA genes are retained in all tunicate mitochondrial genomes, they exhibit a great range of sizes. Tunicates present the shortest mitochondrial rRNA genes among chordates with lengths varying from 596 to 984 bp for *rns* and from 921 to 1,299 bp for *rnl* (Table 1). Both rRNAs are considerably shorter in *O. dioica,* with the *rns* and *rnl* being 242 bp and 215 bp, respectively, after the removal of the poly-T stretches (Table 1). Additionally, they have lost most structural similarity with other metazoan rRNA genes (HMM searches could only identify a part of the *rnl* gene). Interestingly, extreme mitochondrial rRNA reduction and lack of structural similarity are characteristics shared between appendicularians and ctenophores (Arafat, et al. 2018), another lineage with an extreme rate of mitochondrial genome evolution. We annotated the two rRNA regions as different genes because most mitochondrial genomes contain these two genes. However, given the absence of sequence similarity we cannot exclude the possibility that the *rns* region may be, instead, a fragment of the *rnl*. Indeed, in calcareous sponges and Placozoa, rRNA genes are fragmented and encoded on different regions of the mitochondrial genome (reviewed in Lavrov and Pett 2016). Of note, *trans*-splicing is unknown in animals.

tRNA Genes

Though most tunicates encode 24 tRNAs, their tRNA gene content is variable and ranges from 22-27 tRNAs (Iannelli, et al. 2007; Rubinstein, et al. 2013). Specifically, the loss and

acquisition of additional tRNA genes seem to be tolerated and the number of tRNA genes has

been found to vary in a species-specific manner (Gissi, et al. 2008; Rubinstein, et al. 2013). In

O. dioica the situation is again extreme as all tRNAs are absent except for the tRNA-Met

(UAU).

Nuclear-encoded cytoplasmic tRNAs import to the mitochondria has evolved many

times over the course of mitochondrial evolution (Adams and Palmer 2003). We thus assume

that the missing mitochondrial tRNAs in *O. dioica* mitochondrial genome are now encoded in

the nuclear genome and are imported to the mitochondria.

The retention of a single tRNA-Met in the mitochondrial genome, despite the loss of

all other, or most, tRNAs, is not novel to appendicularians and has been observed in

cnidarians, certain chaetognaths, and Keratosa sponges (Lavrov, et al. 2023). The

conventional explanation is based on the fact that mitochondrial protein synthesis starts with

N-formylmethionine, as in prokaryotes, instead of methionine, the initiating amino acid in

eukaryotic cytosolic protein synthesis. The addition of a formyl group to methionine is

catalyzed by the enzyme methionyl-tRNA formyltransferase (Bianchetti, et al. 1977). As

found in prokaryotes, and in accordance with the prokaryotic endosymbiotic origin of

mitochondria, this modification occurs not on free methionine but after methionine is loaded

onto a specific tRNA, namely the initiator tRNA-fmet. Hence, it is usually assumed that the

retained tRNA-Met in mitochondria is the initiator tRNA-fmet (e.g., Beagley, et al. 1998).

Interestingly, animal mitochondrial genomes usually keep tRNAs that are specific to

their mitochondrial genetic codes. For example, most cnidarians retain the tRNA-Trp gene

necessary for the translation of the UGA codon, which is a stop codon in the nuclear genetic

code (Beagley, et al. 1998). In the mitochondrial genetic code of ascidians, four codons have

a different assignment compared with the nuclear genetic code: methionine (AUA),

tryptophan (UGA), and glycine (AGA and AGG). We thus expected the presence of a tRNA-

Trp (for the translation of the UGA codon) and a tRNA-Gly (to translate the codons AGA and

AGG) in the mitochondrial genome of *O. dioica*. Therefore, we computed the codon

frequency across all protein-coding genes of *O. dioica*, thus verifying the usage of all these

AUA, UGA, AGA and AGG codons in the mitochondrial genome (supplementary table S5,

Supplementary Material online). Indeed, it has been found that translation of the AGA and

AGG codons to Gly improves the similarity between *O. dioica* and tunicate mitochondrial

proteins, supporting the use of the ascidian mitochondrial genetic code in appendicularians

(Pichon, et al. 2020). The presence of a cytosolic and a mitochondrial aaRS-Gly, observed

while searching for mitochondrial aaRS in *O. dioica* nuclear genome, further supports the

view that a specific mitochondrial tRNA recognizes the glycine AGA and AGG codons. As

for the case of protein-coding genes, two explanations are possible for the lack of

mitochondrial Gly and Trp tRNAs: either we failed to identify these tRNA genes in the

sequenced mitochondrial genome, because of their very unusual secondary structure or

because they are interrupted by poly T/A stretches, or these tRNAs are nuclear-encoded and

imported from the cytosol. In support of the import scenario, it was found that among

animals, both ctenophores and Myxozoa have lost all their mitochondrial tRNA but still use

the UGA codon as tryptophan (Arafat, et al. 2018; Pett, et al. 2011; Yahalomi, et al. 2017).

Genome Editing of Poly-T/A Stretches

The sequencing of PCR fragments using ONT long-read allowed us to assemble a large part

of the mitochondrial genome of *O. dioica*. It confirms the presence of numerous poly-T/A

stretches in the DNA sequence that are reduced to 6U-sites in the RNA transcript. Because

accurate sequencing of long homopolymers is currently infeasible with Sanger, Illumina (Ari

and Arikan 2016) or Nanopore sequencing (Delahaye and Nicolas 2021; Sereika, et al. 2022),

the length of T/A stretches in our assembly is unreliable. The average size of these regions is

likely slightly higher than that reported in the assembly because sequencing methods tend to

underestimate the length of homopolymer stretches. Another source of uncertainty regarding

the length of the homopolymer stretches stems from the fact that we sequenced amplicons

obtained from DNA extracted from a pool of polymorphic individuals. Intra-species variation

in the length of these stretches likely exists, further contributing to inference uncertainty.

What is the molecular mechanism underlying the editing to 6U-sites, and more

specifically, how are these sites recognized? We could not identify a conserved sequence

motif surrounding these sites (Supplementary Fig. S3, Supplementary Material online). Our

analysis reveals the following general motif for the poly- T/A stretch (see Methods): (1) the

5'-end of this strand starts with a perfect 6T homopolymer; (2) downstream of this region,

there is a poly-pyrimidine track whose length varies from 2 to 509 bases; (3) it ends when a

purine. Only the poly-pyrimidine track is excised. We speculate that the perfect 6T motif in

the upstream region may be the beginning of the recognition editing signal and that a purine

marks its end. In between these motifs, the polypyrimidine track is excised. Notably, the

enzymes involved in this excision mechanism are still unknown.

Another kind of editing was observed within the cytochrome *b* gene: instead of a

perfect 6T motif, it starts with "TTTTTCTT", then there is the excised region, "AACTT",

which is irregular as it contains two purines (supplementary Fig. S9, Supplementary Material

online).

Mitochondrial genome editing is rare in animals, and it is usually limited to the

insertion of a few bases (Lavrov, et al. 2016). Here we instead have a deletion of a large

polypyrimidine region. The removal of such a long stretch at the RNA level is a characteristic

of introns (Chorev and Carmel 2012). Spliceosomal introns are characterized by short signal

sequences such as GT/AG and AT/AC (Berná and Alvarez-Valin 2014; Rogozin, et al. 2012).

It seems that the RNA editing observed here is very different from such a splicing mechanism

as we could not find such motifs. Similarly, the poly-T/A stretches have no sequence

similarity with self-splicing introns that have well-defined secondary structures (Waring and

Davies 1984).

Trypanosomatid protists present a cleavage-ligation pathway through which multiple

uridines can be deleted or inserted by a U-specific exonuclease and a terminal uridil

transferase, respectively (Cruz-Reyes and Sollner-Webb 1996). This particular editing

mechanism involves guide RNAs, which determine the number and positions of the U

nucleotides deleted or inserted within the mitochondrial genome of these protists, e.g.,

Leishmania tarentolae (Connell, et al. 1997). Future studies are necessary to determine if

some components of the editing mechanisms in these protists and in *O. dioica* are

evolutionary related.

Appendicularians have classically been divided into three families Fritillariidae,

Kowaleskiidae, and Oikopleuridae (Garić 2024). However preliminary results from our group

suggest that Kowaleskiidae are members of the Fritillariidae (Garić, et al. 2018). The presence

of edited homopolymer stretches has been mentioned for the COI gene of *O. vanhoeffeni*

(Latour 2021). Interestingly, we have observed that large poly-T inserts are also present in

members of the Fritillariidae, suggesting that this mitochondrial editing mechanism was

present in the ancestor of Appendicularia (Garić, et al. 2018). Conversely, while examining

published data from Naville et al. (Naville, et al. 2019), we identified a clade within

Oikopleuridae, which includes the species *Bathochordaeus stygius*, *Mesochordaeus*

erythrocephalus, and *Oikopleura longicauda* that do not harbor large poly-T/A stretches

(Garić, et al. 2018), suggesting that this RNA editing mechanism may have been lost in

Oikopleuridae. Additional genomic and transcriptomic data from a larger taxon sampling of

appendicularians are needed to better understand the distribution and evolutionary dynamics

of this editing mechanism.

EST and transcriptomic RNAseq data are available for *O. dioica* from Norway and

mainland Japan, respectively. For the Norway sample, genomic data were obtained in this

work, indicating the presence of RNA editing. To evaluate if the Poly-T stretches are

conserved among *O. dioica* lineages, we compared the location of 6U-regions in the DNA and

transcriptomic data from Norway and mainland Japan, respectively (Supplementary Fig. S2,

Supplementary Material online). Our results suggest that while many of the 6U-loci are

conserved, differences exist. For example, in the *cox1* gene, a 6U-site in the Norwegian

sequence and a UUCUUU site in the Japanese sequence can be reliably aligned. These

differences suggest that the poly-T/A stretches loci may not be conserved in the mitochondrial

genomes of different populations or species. Of note, it was recently suggested that *O. dioica*

from Japan and Norway represent different species (Masunaga, et al. 2022). The birth and

death dynamics of these poly-T/A stretches will be better clarified when genomic and

transcriptomic data from a larger appendicularian taxonomic sampling becomes available.

The fact that we could assemble large contigs encompassing several genes from RNA

sequences indicated that the pre-mRNA transcripts in *O. dioica* are polycistronic. In

vertebrates, it has been shown that the cleavage of tRNAs which flank protein-coding genes

(named tRNA punctuation) is the mechanism that releases monocistronic or bicistronic

mature mRNAs of mitochondrial coding genes from the large RNA precursor (Ojala, et al.

1981). In ascidians, the transcription and RNA maturation process of the mitochondrial

genome has been reconstructed through EST analyses, showing that the main features of

mitochondrial transcription are conserved between vertebrates and ascidians (*i.e.,* presence of

polycistronic and immature transcripts, creation of stop codons by polyadenylation, tRNA

signal punctuation, and rRNA transcript termination signals) (Gissi and Pesole 2003). Since

most tRNAs are lost in *O. dioica,* a mechanism different from the tRNA signal punctuation is

likely employed to release mature mRNAs.

The Phylogenetic Position of *O. dioica*

Based on both molecular (Delsuc, et al. 2018) and morphological (Braun, et al. 2020) analyses *O. dioica* was placed as the first diverging branch of the tunicates tree, however it was recognized that this placement, in molecular studies, may be a long-branch-attraction (LBA) artifact (Delsuc, et al. 2018). In our analysis, we found that the phylogenetic position of *O. dioica* was highly variable. The grouping of *O. dioica* and members of the genus *Salpa* is most likely an LBA artifact (Philippe, et al. 2005), however, even when the two *Salpa* representatives were removed from the phylogenetic analyses, the phylogenetic position of *O. dioica* was never well supported and depended on the model used. This suggests the fast-evolutionary rate of this species impedes the use of molecular markers to resolve its phylogenetic position. Again, the lack of phylogenetic signal in the mitochondrial dataset, which precludes the reconstruction of phylogenetic relationships at the phylum level, is a feature of fast-evolving species (Arafat, et al. 2018; Pett, et al. 2011; Yahalomi, et al. 2017).

Conclusions

We demonstrated that a combination of DNA and RNA data from long- and short-read

sequencing could significantly improve our understanding of the structure and gene content of

O. dioica mitochondrial genome.

The mitochondrial genome of *O. dioica* shows unique gene content and an extreme

rate of sequence evolution. A total of 28 genes are missing, including *atp8*, *nad2*, *nad3*

nad4L, *nad6*, and all tRNA genes except for tRNA-Met (UAU). The mitochondrial gene *nad3*

was transferred to the nucleus, and harbors a mitochondrial target peptide. Further work is,

however, needed to decipher the complete sequence of the mitochondrial genome and in

particular to unravel the accurate sequence and length of the poly-T regions. Unfortunately,

currently, no sequencing techniques allow us to perfectly sequence these regions, as the

sequencing methods Illumina, ONT, and Sanger approaches failed to correctly sequence long

polynucleotide regions. However, information obtained from the sequencing of PCR

fragments helped us assembled a draft mitochondrial genome.

Because of the fast-evolutionary rate, we were unable to accurately determine the

phylogenetic position of *O. dioica*. Of note, comparative genomic studies of *O. dioica* and

additional members of the appendicularians are expected to provide us with a better

understanding of the evolution of RNA editing in this group.

Materials and Methods

DNA Extraction, Sequencing and Assembly

Living specimens of *O. dioica* animals were provided by Sars International Centre for Marine Molecular Biology (SICMMB) (Bergen, Norway) in March 2019 and cultured at Humboldt-University zu Berlin (HU) following the methods detailed in Bouquet, et al. (2009). Ethanol-preserved samples were shipped to Tel-Aviv University. Genomic DNA was extracted from ~100 whole individuals following the protocol of Fulton (2012). About 100 ng were sent to the Technion sequencing center (Haifa) for Illumina sequencing (November 2019), the libraries were prepared using the TruSeq DNA Nano library prep kit (Illumina) and were sequenced on an Illumina Hiseq 2500 platform. Similarly, about 1μg of total DNA was sent to Hy Laboratories Ltd (Hylabs, Rehovot) for ONT sequencing using a MinION Mk 1 R9.4 flow cell with a MinION Mk1B device and 1D sequencing kits (May 2019). However, the ONT sequencing only provided 962 reads (mean length 750 bp). ONT sequencing was thus again performed using a different DNA extract and a different sequencing service (the Technion sequencing center) (on July 2019 and December 2021). The flow cell and device setting and sequencing kits were the same as previously. The first sequencing (July 2019) provided ~2 million reads (mean length 2,330 bp) but we could not assemble the mitochondrial genome using these data. *A posteriori,* we found that only ~600 reads mapped to the mitochondrial genome (mean length 300 bp) and that these reads only mapped to a few regions of the mitochondrial genome (data not shown). For this second ONT sequencing fragments around 3,000 bp were selected. Because we realized that mitochondrial DNA may be fragmented, we performed the last ONT sequencing (from December 2021) without fragment size selection. This ONT sequencing also provided only 3,144 reads (mean length 1,070 bp).

The Illumina 100 bp paired-ended reads were cleaned from adaptors using Cutadapt

v.2.8 (Martin 2011) and assembled using IDBA-UD v.1.1.3 (Zhu, et al. 2010) using default

parameters. Efforts to assemble the mitochondrial genome from these reads were unsuccessful

because contigs were broken in the long poly-T regions.

Because whole genome sequencing using Illumina or ONT did not allow us to

assemble a complete mitochondrial sequence we decided to amplify the mitochondrial

genome into several PCR fragments. To do so we needed to design primers in conserved

regions that did not include homopolymers. We used published RNA reads and EST

sequences to identify mitochondrial genes and design primers.

Identification of Canonical Mitochondrial Protein Transcripts in Published RNA and

EST Sequences

Illumina RNA reads from *O. dioica* specimens from Japan and the North Sea were

downloaded from the National Center for Biotechnology Information (NCBI) (accessions:

SRR1693762, SRR1693765, SRR1693766, and SRR1693767) (Wang, et al. 2015), and the

European Nucleotide Archive (ENA) (accessions: SRR20015061) (Raghavan, et al. 2023),

respectively. The reads were then assembled with IDBA-UD v.1.1.3 (Zhu, et al. 2010) using

default parameters. tBlastn searches were conducted against the transcriptome assembly

obtained using the amino acid sequences of the protein-coding genes of the mitochondrial

genome of *Ciona robusta* (AJ517314) as query. The reads from Japan could be assembled in

four contigs and will be referred to as "Japan" (supplementary Fig. S10 A, Supplementary

Material online). These contigs were annotated with the webserver MITOS V2 (Donath, et al.

2019) The genes from the North Sea could be assembled in five contigs and will be referred

to as "North Sea" (supplementary Fig. S10 B, Supplementary Material online). All contigs

were annotated with the webserver MITOS V2 (Donath, et al. 2019).

Mapping Illumina DNA reads and the few ONT reads to these contigs confirmed the

presence of homopolymers and also indicated that the Japanese *O. dioica* presented many

substitutions when compared to our sample which originated from Norway. EST sequences

from *O. dioica* samples from Norway were available in the OikoBase database (Danks, et al.

2012). For each mitochondrial ORF identified in the Illumina North Sea dataset, we mapped

~500 EST sequences to the reference. We next filtered the flanking regions of the extracted

EST sequences against the UniVec Vector using Geneious Prime v.2023.2.1. The consensus

sequences were extracted for each gene, which helped us determine the exact gene

boundaries. We verified that all coding transcripts we identified could be successfully

translated to protein using the ascidian or the invertebrate mitochondrial codes. We found that

the ascidian genetic code yields proteins that have higher sequence similarity to mitochondrial

proteins in other animals than when using the invertebrate genetic code, in agreement with

Pichon, et al. (2020).

PCR Amplification, Sequencing and Assembly

Primers of ~ 30 bp with a TM ~ 60 C° were designed based on ORF assembled from the EST,

accounting for the gene order inferred from the contigs (supplementary table S1,

Supplementary Material online). Long PCRs were performed, with several primer

combinations (one from each pair of contigs and in various orientations), to determine gene

arrangements along the mitochondrial genome. Because of the high AT content of the

genome, we were unable to design primers that would result in overlapping PCR fragments.

Instead, we used the same region for forward and reverse primers, leading to non-overlapping

fragments. The sequence of the primer region was confirmed using Illumina data. PCRs were

performed in a 25µl volume with 2u of TaKaRa Ex-Taq polymerase (Cat.# RR001), 1X of

Ex-Taq buffer, 1M of Betain, 0.2mM of dNTPs TaKaRa mixture, 0.5µM of each primer and

1μl *O. dioica* DNA extraction. The PCR program used: (a) long denaturation step at 95°c for

5 min; (b) 35 cycles of a short denaturation step at 95°c for 45 sec followed by an annealing

plus elongation step at 68°c for 10 min; (c) a long elongation step at 72°c for 10 min.

Reamplifications was performed to improve the PCR yield of some fragments.

Reamplifications were performed using 1 μl of PCR products in place of genomic DNA. 12

PCR reactions were successful and sent to ONT sequencing (supplementary Fig. S1,

Supplementary Material online). Each PCR product was barcoded and the sequencing was

performed in May 2023 by the Technion Sequencing Center on an Oxford Nanopore MinION

device (MinKNOW software v.22.12.7), using one R9.4.1 flow cell (Oxford Nanopore, cat

no. FLO-MIN106D). The ONT reads obtained were filtered from adaptors using Porechop

v.0.2.4 (Wick, et al. 2017) using the parameters -middle_threshold 85. Next, we assembled

the trimmed ONT reads with Flye v.2.9.1 (Kolmogorov, et al. 2020) as implemented in

Geneious Prime v.2023.2.1 with two rounds of polishing. The same parameters could not be

used for all PCR fragments as in some cases, the assembly program returned an error message

or failed to terminate. In these cases, other parameters were tested, e.g., if the default

parameter "genome size" of 1,000 bp resulted in a failed run, we tried running the program

with increasing values of genome size from 2,000 to 5,000 bp. If this also failed, we added the

option "metagenome" or the option "reduced contig = 30". Even with these modifications, the

assembly of PCR fragment 12 was unsuccessful.

Many of the assembled contigs were palindromic, with the middle of the contig

corresponding to the PCR primer. In this case, the consensus sequence was split into two

halves, reads were mapped to each half (using Geneious Prime v.2023.2.1 with default

options) and the region with the highest number of mapped reads was kept as the consensus

sequence.

The entire mitochondrial sequence was determined by connecting the obtained contigs

in their overlapping primer regions, i.e., the same primer region was the forward primer of

one PCR fragment and the reverse primer on another fragment. The exact sequences of the

primer regions were verified by mapping the Illumina reads to the assembled genome

sequence.

Annotation of the Mitochondrial Genome

Protein-coding genes

Gene annotation was based on Blast search against the NCBI databases and was confirmed

using MITOS v.2 (Donath, et al. 2019). The end positions of protein-coding genes were

verified using EST data (the presence of a poly-A region after the stop codon). The start

codon was determined by searching for the first start codon upstream of the region that

corresponded to the EST data.

rRNAs

The mapping of Illumina RNA reads from Japan (SRR1693762, SRR1693765, SRR1693766,

SRR1693767) and North Sea (SRR20015061) to the mitochondrial genome of *O. dioica*

revealed two regions with higher coverage that we inferred to be *rns* and *rnl* (supplementary

Fig. S5, Supplementary Material online). Read mapping was done using the Geneious

Mapper, Geneious Prime v.2023.2.1 with the parameters: medium-low sensitivity, randomly

map multiple best matches, only map paired reads which both map, and trim paired reads

overhangs. Average coverage was computed with Geneious Prime v.2023.2.1 statistics. The

region that we annotated as *rnl* was also identified as such using gene prediction with MITOS

v.2 (Donath, et al. 2019). The *rnl* and *rns* sequences were also present in the EST data, which

helped determine the boundaries of these ribosomal genes.

 tRNAs

Putative tRNAs were searched using MITOS v.2 (Donath, et al. 2019) and the ARWEN v.1.2

webserver (Laslett and Canbäck 2008). Both programs identify a single tRNA gene

(supplementary Fig. S6, Supplementary Material online).

Aminoacyl-tRNA Synthetase Profiling

To corroborate the absence of tRNA genes in the mitochondrial genome, we searched for

aaRS genes in the nuclear proteome (i.e., the amino acid sequences of *O. dioica* in the NCBI

"protein" database). The aaRS search was done using HMMER v.3.3.2 software (Eddy 2011)

and was based on profiles developed by Pett and Lavrov (2015). These profiles were used to

search against proteome datasets downloaded from NCBI on April 2023 of *O. dioica* from

Norway (GCA_000209535), *O. dioica* from Okinawa (GCA_907165135), *H. sapiens*

(GCF_000001405.40 Human Build 38 patch release 14), *C. intestinalis* (GCF_000224145.3),

and *S. clava* (GCF_013122585.1).

Nuclear Gene Transfer

We used HMM searches against the proteome of *O. dioica* from Norway GCA_000209535

and *O. dioica* from Okinawa GCA_907165135 to detect canonical mitochondrial-encoded

genes that could have been transferred to the nuclear genomes. The searches were performed

using HMMER v3.4 (Eddy 2011) using the *atp8*, *nad2*, *nad3*, *nad4L*, and *nad6* gene models

developed by Gabriel Alves Vieira and available in the mitos2_wrapper source code

repository (Vieira 2023). As negative control we also searched the proteomes of *H. sapiens*

GCF_000001405.40 Human Build 38 patch release 14, *C. intestinalis* GCF_000224145.3, and

S. clava GCF_013122585.1. Putative mitochondrial proteins that were found to be encoded in

the nuclear genome were searched for mitochondrial targeting signal using the TargetP-2.0

webserver (Armenteros, et al. 2019).

Codon Usage and Amino-Acid Composition

Codon usage and amino acid composition were computed using Geneious Prime v.2023.2.1.

Characterization of ODI_mt_ORF196

The number of transmembrane domains of ODI_mt_ORF196 was predicted using TMHMM

v.1.0.2 in Geneious Prime v.2023.2.1 (Viklund and Elofsson 2004). The protein structure was

predicted using Alphafold (Jumper, et al. 2021). The predicted structure was used as query

against a large database of structures using Foldseek (van Kempen, et al. 2024).

Characterization of the Poly-T/A Stretches

We define poly-T stretches, by manually examining 59 cases of edited poly-T sites, i.e., these

editing cases were validated by comparing RNA and DNA sequences of these loci. A classic

poly-T stretch is defined as follows: (1) the 5'-end of this stretch starts with a perfect 6T

homopolymer. This region is not excised; (2) downstream of this region, there is a C/T stretch

(i.e., a Polypyrimidine track). This region is excised; (3) the stretch ends when either an A or

a G nucleotide is encountered. This A or G nucleotide is not part of the poly-T stretch and not

excised. In six validated cases, this pattern was violated. In five of these cases there was a

single A nucleotide within the long C/T stretch, and in one case a G. We thus include in the

definition cases in which the C/T stretch includes a single purine, as long as it is followed by

at least 10 more C/T nucleotides. Searching the genome with this definition, we found six

additional poly-T stretches (four classic and two non-classic) for which we lack RNA support.

In total we annotated 65 poly-T stretches in these genomes. As we search for the above motif

or its reverse complement to account for both strands, we term these stretches poly-T/A

stretches.

For each homopolymer site in the genome we extracted 50 bp before the site, the 6 T/A in the

site itself, and 50 bp after the site using our own python v.3.11.5 script on Jupiter Notebook

v.7.1.2. Nucleotide frequencies before and after the homopolymer sites were plotted

(Supplementary Fig. S3, Supplementary Material online) by WebLogo with parameters of

Advanced Logo Options: Frequency Plot.v.2.8.2 (Crooks, et al. 2004). To determine the

conservation of the poly T/A stretches, we aligned the mitochondrial gene regions extracted

from the Illumina RNA data from Japan and to our own assembly. Alignments were

computed using MAFFT v.7.490 under the L-INS-i algorithm.

Mitochondrial Genome Length among Tunicates

The average mitochondrial genome size of tunicates was calculated from the length of the

whole genome sequences of 34 tunicate species (supplementary table S3, Supplementary

Material online) downloaded from the NCBI database.

Phylogenetic Analysis of Mitochondrial Protein Sequences

The phylogenetic analyses included 34 tunicate species with a complete mitochondrial

genome that were deposited in NCBI (last accessed: June 2023) and 14 outgroup species

(supplementary table S3, Supplementary Material online). The outgroup species were chosen

following Bourlat, et al. (2006); Bourlat, et al. (2008); Shenkar, et al. (2016); Singh, et al.

(2009). Eight protein-coding gene (*atp6, cox1, cox2, cox3, cytb, nad1, nad4, nad5*) were first

extracted from each species and aligned separately. We used the Guidance2 server (Sela, et al.

2015) to identify unreliable regions of the alignment. We excluded these regions as well as

alignment columns that included more than 50% missing data or gaps. The different protein-

coding alignments were then concatenated to form a super-matrix with 2,027 columns.

Because members of the tunicate genus *Salpa* are extremely fast evolving, we prepared a

second dataset, repeating the above steps, but excluding *S. thompsoni* and *S, fusiformi*. The

resulting super-matrix of 46 species had 2,176 columns.

The phylogenetic trees were reconstructed under the maximum likelihood (ML)

criterion using IQ-TREE 2.1.2 (Minh, et al. 2020). First, we identified the best-fitting model

by using the ModelFinder-Plus (MFP) option of IQ-TREE 2.1.2 and computed the ML tree

using this model. The model that best fitted both datasets (with and without the genus *Salpa*)

was mtZOA+F+G+I. We reconstructed the tree using complex models that are not

implemented in MFP: mtZOA+EHO+F+G+I, Poisson+C10+G, and Poisson+C20+G. The

Robustness of each internal branch of the ML trees were evaluated with 100 bootstrap

replications (Felsenstein 1985).

Bayesian tree reconstructions were performed using the site-heterogeneous CAT

model implemented in Phylobayes MPI v1.5 (Lartillot, et al. 2013), which is expected to

lower the impact of long-branch attraction on the phylogenetic tree inferred (Lartillot, et al.

2007). For the first and second datasets, four independent chains were run for 71,880 and

188,000 cycles, respectively. The first 5,000 and 10,000 trees from each chain, for the first

and second data sets, respectively, were discarded as burn-in and trees were sampled every 10

trees (i.e., the sample size was 3,160, and 17,800 trees per chain for the first and second data

set respectively). Chain convergence was assessed using the bpcomp and tracecomp scripts,

which are part of Phylobayes. Specifically, the bpcomp maxdiff values were 0.23 and the

tracecomp effsize and rel_diff values were >800 and below 0.15, supporting a proper

convergence for the first dataset. For the second dataset, the bpcomp maxdiff value was 0.40

when the 4 chains were considered together, which indicated insufficient convergence.

However, when considering only chains 1, 2, and 3 or chains 1, 2 and 4 this value dropped to

0.18 and 0.27 indicating proper convergence. In both cases, rel_diff values were >150 and

below 0.16 supporting convergence. We thus considered these two results.

Supplementary Material

Supplementary files Tables S1-S6. Figures S1-S10 are available at Genome Biology and

Evolution online.

Acknowledgments

This work was supported by the German-Israeli Foundation for Scientific Research and

Development to DH and TS (Grant No. I-1454-203.13/2018). We want to thank Nitzan

Fourier from the Technion Genome Center for her help with Nanopore sequencing. We thank

Yair Shimony for his help with the structure predictions.

Data Availability

The Illumina read data have been deposited in the European Nucleotide Archive under study

ID number PRJEB72855. The mtDNA sequence was deposited under accession number

OZ035851. The ONT reads obtained for each of the PCR fragments, as well as the sequences

assembled from published RNA and EST data are available on GitHub

(https://github.com/dorohuchon/Oikopleura_dioica_mitochondrial_genome).

Literature Cited

Adams KL, Palmer JD 2003. Evolution of mitochondrial gene content: gene loss and transfer

to the nucleus. Mol Phylogenetics Evol. 29:380-395. https://doi.org/10.1016/S1055-

7903(03)00194-5

Arafat H, Alamaru A, Gissi C, Huchon D 2018. Extensive mitochondrial gene rearrangements

in Ctenophora: insights from benthic Platyctenida. BMC Evol Biol. 18:65.

https://doi.org/10.1186/s12862-018-1186-1

Ari Ş, Arikan M. 2016. Next-generation sequencing: Advantages, disadvantages, and future.

In: Hakeem KR, Tombuloğlu H, Tombuloğlu G, editors. Plant Omics: Trends and

Applications. Springer International Publishing. 109-135. https://doi.org/10.1007/978-

3-319-31703-8_5

Armenteros JJA, et al. 2019. Detecting sequence signals in targeting peptides using deep

learning. Life Sci Alliance. 2:e201900429. https://doi.org/10.26508/lsa.201900429

Beagley CT, Okimoto R, Wolstenholme DR 1998. The mitochondrial genome of the sea

anemone *Metridium senile* (Cnidaria): introns, a paucity of tRNA genes, and a near-

standard genetic code. Genetics. 148:1091-1108.

https://doi.org/10.1093/genetics/148.3.1091

Berná L, Alvarez-Valin F 2014. Evolutionary genomics of fast evolving tunicates. Genome

Biol Evol 6:1724-1738. https://doi.org/10.1093/gbe/evu122

Bianchetti R, Lucchini G, Crosti P, Tortora P 1977. Dependence of mitochondrial protein

synthesis initiation on formylation of the initiator methionyl-tRNAf. J Biol Chem.

252:2519-2523. https://doi.org/10.1016/S0021-9258(17)40489-3

Bliznina A, et al. 2021. Telomere-to-telomere assembly of the genome of an individual

Oikopleura dioica from Okinawa using Nanopore-based sequencing. BMC Genom.

22:222. https://doi.org/10.1186/s12864-021-07512-6

Boore JL 1999. Animal mitochondrial genomes. Nucleic Acids Res. 27:1767-1780.
https://doi.org/10.1093/nar/27.8.1767

Bouquet J-M, et al. 2009. Culture optimization for the emergent zooplanktonic model
organism *Oikopleura dioica*. J Plankton Res. 31:359-370.
https://doi.org/10.1093/plankt/fbn132

Bourlat SJ, et al. 2006. Deuterostome phylogeny reveals monophyletic chordates and the new
phylum Xenoturbellida. Nature. 444:85-88. https://doi.org/10.1038/nature05241

Bourlat SJ, Nielsen C, Economou AD, Telford MJ 2008. Testing the new animal phylogeny: a
phylum level molecular analysis of the animal kingdom. Mol Phylogenet Evol. 49:23-
31. https://doi.org/10.1016/j.ympev.2008.07.008

Braun K, Leubner F, Stach T 2020. Phylogenetic analysis of phenotypic characters of
Tunicata supports basal Appendicularia and monophyletic Ascidiacea. Cladistics.
36:259-300. https://doi.org/10.1111/cla.12405

Cardol P, et al. 2006. ND3 and ND4L subunits of mitochondrial complex I, both nucleus
encoded in *Chlamydomonas reinhardtii*, are required for activity and assembly of the
enzyme. Eukaryot Cell. 5:1460-1467. https://doi.org/10.1128/ec.00118-06

Castellana S, Vicario S, Saccone C 2011. Evolutionary patterns of the mitochondrial genome
in Mtazoa: exploring the role of mutation and selection in mitochondrial protein–
coding genes. Genome Biol Evol. 3:1067-1079. https://doi.org/10.1093/gbe/evr040

Chorev M, Carmel L 2012. The function of introns. Front Genet. 3:55.
https://doi.org/10.3389/fgene.2012.00055

Connell GJ, Byrne EM, Simpson L 1997. Guide RNA-independent and guide RNA-
dependent uridine insertion into *Cytochrome b* mRNA in a mitochondrial lysate from
Leishmania tarentolae: role of RNA secondary structure. Biol Chem. 272:4212-4218.
https://doi.org/10.1074/jbc.272.7.4212

Crooks GE, Hon G, Chandonia JM, Brenner SE 2004. WebLogo: a sequence logo generator.

Genome Res. 14:1188-1190. https://doi.org/10.1101/gr.849004

Cruz-Reyes J, Sollner-Webb B 1996. Trypanosome U-deletional RNA editing involves guide

RNA-directed endonuclease cleavage, terminal U exonuclease, and RNA ligase

activities. Proc Natl Acad Sci USA. 93:8901-8906.

https://doi.org/10.1073/pnas.93.17.8901

Danks G, et al. 2012. OikoBase: a genomics and developmental transcriptomics resource for

the urochordate *Oikopleura dioica*. Nucleic Acids Res. 41:D845-D853.

https://doi.org/10.1093/nar/gks1159

Delahaye C, Nicolas J 2021. Sequencing DNA with nanopores: troubles and biases. PLOS

ONE. 16:e0257521. https://doi.org/10.1371/journal.pone.0257521

Delsuc F, Brinkmann H, Chourrout D, Philippe H 2006. Tunicates and not cephalochordates

are the closest living relatives of vertebrates. Nature. 439:965-968.

https://doi.org/10.1038/nature04336

Delsuc F, et al. 2018. A phylogenomic framework and timescale for comparative studies of

tunicates. BMC Biol. 16. https://doi.org/10.1186/s12915-018-0499-2

Denoeud F, et al. 2010. Plasticity of animal genome architecture unmasked by rapid evolution

of a pelagic tunicate. Science. 330:1381-1385.

https://doi.org/10.1126/science.1194167

Donath A, et al. 2019. Improved annotation of protein-coding genes boundaries in metazoan

mitochondrial genomes. Nucleic Acids Res. 47:10543-10552.

https://doi.org/10.1093/nar/gkz833

Durrheim GA, Corfield VA, Harley EH, Ricketts MH 1993. Nucleotide sequence of

cytochrome oxidase (subunit III) from the mitochondrion of the tunicate *Pyura*

stolonifera: evidence that AGR encodes glycine. Nucleic Acids Res. 21:3587-3588.

https://doi.org/10.1093/nar/21.15.3587

Eddy SR 2011. Accelerated profile HMM searches. PLoS Comput Biol. 7:e1002195. .

https://doi.org/10.1371/journal.pcbi.1002195

Erpenbeck D, et al. 2006. Mitochondrial diversity of early-branching metazoa is revealed by

the complete mt genome of a halosclerid demosponge. Mol Biol Evol 24:19-22.

https://doi.org/10.1093/molbev/msl154

Felsenstein J 1985. Confidence limits on phylogenies: an approach using the bootstrap.

Evolution. 39:783-791. https://doi.org/10.2307/2408678

Fulton TL, Wagner SM, Shapiro B. 2012. Case study: recovery of ancient nuclear DNA from

toe pads of the extinct passenger pigeon. In: Shapiro B, Hofreiter M, editors. Ancient

DNA: methods and protocols. Humana Press. 29-35. https://doi.org/10.1007/978-1-

61779-516-9_4

Garić R 2024. World List of Appendicularia. Appendicularia. Accessed through: World

Register   of   Marine   Species.

https://www.marinespecies.org/aphia.php?p=taxdetails&id=146421

Garić R, et al. 2018. Challenges in appendicularian integrative taxonomy and barcoding //

ICES Annual Science Conference 2018. https://www.imp-du.com/datoteke/Ices-2018-

rade-garic.pdf

Gissi C, Iannelli F, Pesole G 2004. Complete mtDNA of *Ciona intestinalis* reveals extensive

gene rearrangement and the presence of an *atp8* and an extra *trnM* gene in ascidians. J

Mol Evol. 58:376-389. https://doi.org/10.1007/s00239-003-2559-6

Gissi C, Iannelli F, Pesole G 2008. Evolution of the mitochondrial genome of Metazoa as

exemplified   by   comparison   of   congeneric   species.   Heredity.   101:301-320.

https://doi.org/10.1038/hdy.2008.62

Gissi C, Pesole G 2003. Transcript mapping and genome annotation of ascidian mtDNA using

EST data. Genome Res. 13:2203-2212. https://doi.org/10.1101/gr.1227803

Gott JM, Emeson RB 2000. Functions and mechanisms of RNA editing. Annu Rev Genet.

34:499-531. https://doi.org/10.1146/annurev.genet.34.1.499

Helfenbein KG, Fourcade HM, Vanjani RG, Boore JL 2004. The mitochondrial genome of

Paraspadella goto is highly reduced and reveals that chaetognaths are a sister group to

protostomes. Proc Natl Acad Sci USA. 101:10639-10643.

https://doi.org/10.1073/pnas.0400941101

Hoffmann RJ, Boore JL, Brown WM 1992. A novel mitochondrial genome organization for

the blue mussel, *Mytilus edulis*. Genetics. 131:397-412.

https://doi.org/10.1093/genetics/131.2.397

Iannelli F, Griggio F, Pesole G, Gissi C 2007. The mitochondrial genome of *Phallusia*

mammillata and *Phallusia fumigata* (Tunicata, Ascidiacea): high genome plasticity at

intra-genus level. BMC Evol Biol. 7:155. https://doi.org/10.1186/1471-2148-7-155

Jumper J, et al. 2021. Highly accurate protein structure prediction with AlphaFold. Nature.

596:583-589. https://doi.org/10.1038/s41586-021-03819-2

Kolmogorov M, et al. 2020. metaFlye: scalable long-read metagenome assembly using repeat

graphs. Nat Methods. 17:1103-1110. https://doi.org/10.1038/s41592-020-00971-x

Lartillot N, Brinkmann H, Philippe H 2007. Suppression of long-branch attraction artefacts in

the animal phylogeny using a site-heterogeneous model. BMC Evol Biol. 7:S4.

https://doi.org/10.1186/1471-2148-7-S1-S4

Lartillot N, Philippe H 2004. A bayesian mixture model for across-site heterogeneities in the

amino-acid replacement process. Mol Biol Evol. 21:1095-1109.

https://doi.org/10.1093/molbev/msh112

Lartillot N, Rodrigue N, Stubbs D, Richer J 2013. PhyloBayes MPI: Phylogenetic
reconstruction with infinite mixtures of profiles in a parallel environment. Syst Biol.
62:611-615. https://doi.org/10.1093/sysbio/syt022

Laslett D, Canbäck B 2008. ARWEN: a program to detect tRNA genes in metazoan
mitochondrial nucleotide sequences. Bioinformatics. 24:172-175.
https://doi.org/10.1093/bioinformatics/btm573

Latour AR 2021. Molecular divergence and gene flow in cold ocean tunicates: insights from
cytochrome oxidase 1. Diss Memorial University of Newfoundland.73-87.
https://doi.org/10.48336/hkd1-9x29

Lavrov DV 2007. Key transitions in animal evolution: a mitochondrial DNA perspective.
Integr Comp Biol. 47:734-743. https://doi.org/10.1093/icb/icm045

Lavrov DV, Adamski M, Chevaldonné P, Adamska M 2016. Extensive mitochondrial mRNA
editing and unusual mitochondrial genome organization in calcaronean sponges. Curr
Biol. 26:86-92. https://doi.org/10.1016/j.cub.2015.11.043

Lavrov DV, et al. 2023. Phylomitogenomics bolsters the high-level classification of
Demospongiae (phylum Porifera). PLOS ONE. 18:e0287281.
https://doi.org/10.1371/journal.pone.0287281

Lavrov DV, Pett W 2016. Animal mitochondrial DNA as we do not know it: mt-genome
organization and evolution in nonbilaterian lineages. Genome Biol Evol. 8:2896-2913.
https://doi.org/10.1093/gbe/evw195

Le TH, Blair D, McManus DP 2002. Mitochondrial genomes of parasitic flatworms. Trends
Parasitol. 18:206-213. https://doi.org/10.1016/S1471-4922(02)02252-3

Lonergan KM, Gray MW 1996. Expression of a continuous open reading frame encoding
subunits 1 and 2 of cytochrome *c* oxidase in the mitochondrial DNA of *Acanthamoeba*
castellanii. J Mol Biol. 257:1019-1030. https://doi.org/10.1006/jmbi.1996.0220

Martin M 2011. Cutadapt removes adapter sequences from high-throughput sequencing reads.

EMBnet j. 17:3. https://doi.org/10.14806/ej.17.1.200

Masunaga A, et al. 2022. The cosmopolitan appendicularian *Oikopleura dioica* reveals hidden

genetic diversity around the globe. Mar Biol. 169:157. https://doi.org/10.1007/s00227-

022-04145-5

Minh BQ, et al. 2020. IQ-TREE 2: New models and efficient methods for phylogenetic

inference in the genomic era. Mol Biol Evol. 37:1530-1534.

https://doi.org/10.1093/molbev/msaa015

Naville M, et al. 2019. Massive changes of genome size driven by expansions of non-

autonomous transposable elements. Curr Biol. 29:1161-1168.e1166.

https://doi.org/10.1016/j.cub.2019.01.080

Nishida H 2008. Development of the appendicularian *Oikopleura dioica*: culture, genome,

and cell lineages. Dev Growth Differ. 50:S239-S256. https://doi.org/10.1111/j.1440-

169X.2008.01035.x

Ojala D, Montoya J, Attardi G 1981. tRNA punctuation model of RNA processing in human

mitochondria. Nature. 290:470-474. https://doi.org/10.1038/290470a0

Okimoto R, Macfarlane JL, Clary DO, Wolstenholme DR 1992. The mitochondrial genomes

of two nematodes, *Caenorhabditis elegans* and *Ascaris suum*. Genetics. 130:471-498.

https://doi.org/10.1093/genetics/130.3.471

Pesole G, Gissi C, De Chirico A, Saccone C 1999. Nucleotide substitution rate of mammalian

mitochondrial genomes. J Mol Evol. 48:427-434. https://doi.org/10.1007/PL00006487

Pett W, Lavrov DV 2015. Cytonuclear interactions in the evolution of animal mitochondrial

tRNA metabolism. Genome Biol Evol. 7:2089-2101.

https://doi.org/10.1093/gbe/evv124

Pett W, et al. 2011. Extreme mitochondrial evolution in the ctenophore *Mnemiopsis leidyi*:

Insight from mtDNA and the nuclear genome. Mitochondrial DNA. 22:130-142.

https://doi.org/10.3109/19401736.2011.624611

Philippe H, Zhou Y, Brinkmann H, Rodrigue N, Delsuc F 2005. Heterotachy and long-branch

attraction in phylogenetics. BMC Evol Biol. 5:50. https://doi.org/10.1186/1471-2148-

5-50

Pichon J, Luscombe NM, Plessy C 2020. Widespread use of the "ascidian" mitochondrial

genetic   code   in   tunicates.   F1000Research.   8:2072-2072.

10.12688/f1000research.21551.2

Pichon J, Luscombe NM, Plessy C 2019. Widespread use of the "ascidian" mitochondrial

genetic   code   in   tunicates.   F1000Research.   8:2072-2072.

https://doi.org/10.12688%2Ff1000research.21551.2

Raghavan V, Eichele G, Larink O, Karin EL, Söding J 2023. RNA sequencing indicates

widespread conservation of circadian clocks in marine zooplankton. NAR Genom

Bioinform. 5. https://doi.org/10.1093/nargab/lqad007

Rogozin IB, Carmel L, Csuros M, Koonin EV 2012. Origin and evolution of spliceosomal

introns. Biol Direct. 7:11. https://doi.org/10.1186/1745-6150-7-11

Rubinstein ND, et al. 2013. Deep sequencing of mixed total DNA without barcodes allows

efficient assembly of highly plastic ascidian mitochondrial genomes. Genome Biol

Evol. 5:1185-1199. https://doi.org/10.1093/gbe/evt081

Sela I, Ashkenazy H, Katoh K, Pupko T 2015. GUIDANCE2: Accurate detection of

unreliable alignment regions accounting for the uncertainty of multiple parameters.

Nucleic Acids Res. 43:W7-W14. https://doi.org/10.1093/nar/gkv318

Sereika M, et al. 2022. Oxford Nanopore R10.4 long-read sequencing enables the generation

of near-finished bacterial genomes from pure cultures and metagenomes without

short-read or reference polishing. Nat Methods. 19:823-826.

https://doi.org/10.1038/s41592-022-01539-7

Shenkar N, Koplovitz G, Dray L, Gissi C, Huchon D 2016. Back to solitude: Solving the

phylogenetic position of the diazonidae using molecular and developmental

characters. Mol Phylogenetics Evol. 100:51-56.

https://doi.org/10.1016/j.ympev.2016.04.001

Singh TR, et al. 2009. Tunicate mitogenomics and phylogenetics: peculiarities of the

Herdmania momus mitochondrial genome and support for the new chordate

phylogeny. BMC Genom. 10:534. https://doi.org/10.1186/1471-2164-10-534

Spealman P, Burrell J, Gresham D 2020. Inverted duplicate DNA sequences increase

translocation rates through sequencing nanopores resulting in reduced base calling

accuracy. Nucleic Acids Res. 48:4940-4945. https://doi.org/10.1093/nar/gkaa206

Stach T, Braband A, Podsiadlowski L 2010. Erosion of phylogenetic signal in tunicate

mitochondrial genomes on different levels of analysis. Mol Phylogenetics Evol.

55:860-870. https://doi.org/10.1016/j.ympev.2010.03.011

The UniProt Consortium 2022. UniProt: the universal protein knowledgebase in 2023.

Nucleic Acids Res. 51:D523-D531. https://doi.org/10.1093/nar/gkac1052

van Kempen M, et al. 2024. Fast and accurate protein structure search with Foldseek. Nature

Biotechnology. 42:243-246. https://doi.org/10.1038/s41587-023-01773-0

[source code repository] Vieira GA. 2023. mitos2_wrapper. GitHub.

https://github.com/gavieira/mitos2_wrapper/tree/master/refseq63m/featureProtHMM.

Viklund H, Elofsson A 2004. Best alpha-helical transmembrane protein topology predictions

are achieved using hidden Markov models and evolutionary information. Prot Sci.

13:1908-1917. https://doi.org/10.1110/ps.04625404

Wang H-C, Minh BQ, Susko E, Roger AJ 2017. Modeling site heterogeneity with posterior

mean site frequency profiles accelerates accurate phylogenomic estimation. Syst Biol.

67:216-235. https://doi.org/10.1093/sysbio/syx068

Wang J, et al. 2022. The conserved domain database in 2023. Nucleic Acids Res. 51:D384-

D388. https://doi.org/10.1093/nar/gkac1096

Wang K, Omotezako T, Kishi K, Nishida H, Onuma TA 2015. Maternal and zygotic

transcriptomes in the appendicularian, *Oikopleura dioica*: novel protein-encoding

genes, intra-species sequence variations, and trans-spliced RNA leader. Dev Genes

Evol. 225:149-159. https://doi.org/10.1007/s00427-015-0502-7

Waring RB, Davies RW 1984. Assessment of a model for intron RNA secondary structure

relevant   to   RNA   self-splicing   -   a   review.   Gene.   28:277-291.

https://doi.org/10.1016/0378-1119(84)90145-8

Wick RR, Judd LM, Gorrie CL, Holt KE 2017. Completing bacterial genome assemblies with

multiplex   MinION   sequencing.   Microb   Genom.   3:e000132.

https://doi.org/10.1099/mgen.0.000132

Wick RR, Judd LM, Holt KE 2019. Performance of neural network basecalling tools for

Oxford Nanopore sequencing. Genome Biol. 20:129. https://doi.org/10.1186/s13059-

019-1727-y

Yahalomi D, et al. 2017. The multipartite mitochondrial genome of *Enteromyxum leei*

(Myxozoa):   eight   fast-evolving   megacircles.   Mol   Biol   Evol.   34:1551–1556.

https://doi.org/10.1093/molbev/msx072

Yokobori S-i, et al. 1999. Complete DNA sequence of the mitochondrial genome of the

ascidian *Halocynthia roretzi* (Chordata, Urochordata). Genetics. 153:1851-1862.

https://doi.org/10.1093/genetics/153.4.1851

Yokobori S-i, Ueda T, Watanabe K 1993. Codons AGA and AGG are read as glycine in

ascidian mitochondria. J Mol Evol. 36:1-8. https://doi.org/10.1007/BF02407301

Yokobori S-i, Watanabe Y, Oshima T 2003. Mitochondrial genome of *Ciona savignyi*

(Urochordata, Ascidiacea, Enterogona): comparison of gene arrangement and tRNA

genes with *Halocynthia roretzi* mitochondrial genome. J Mol Evol. 57:574-587.

https://doi.org/10.1007/s00239-003-2511-9

Zhu W, Lomsadze A, Borodovsky M 2010. *Ab initio* gene identification in metagenomic

sequences. Nucleic Acids Res. 38:e132-e132. https://doi.org/10.1093/nar/gkq275

1082

1083

1084

1085   **Figure legend**

1086

1087   **Figure 1. *Oikopleura dioica* mitochondrial genome map.**

1088   Protein-coding genes are in green, rRNA genes (*rns*, *rnl*) are in dark blue, tRNA-Met (UAU)

1089   is in pink, coding sequences (CDS) are in yellow, and the poly-T/A stretches are in light blue.

1090   The genes are pointing in the direction of their transcriptional orientation. A non-sequenced

1091   region is indicated as "gap" is in gray.

1092

Figure 2. A typical poly-T/A region in *O. doica cox1* gene.

The reference sequence represents the assembled mitochondrial genome. The sequence of part

of the *cox1* gene is indicated in dark green. Above it are the nucleotide and amino-acid

sequences of the reference sequence. In light blue is the poly-T/A region. Reads from five

different sequencing sources mapped to this region are shown below the reference sequence:

our ONT data, our Illumina reads, public EST data from Norway, public RNA reads from

Japan, and public RNA reads from the North Sea (see Supplementary table S3,

Supplementary Material online A for accession numbers). Each line represents a different

read. Note that the different ONT reads do not have the same length, exemplifying the fact

that the length of the poly-T region could not be accurately determined. The EST and RNA

reads lack a long poly-T region. Rather, the poly-T region corresponds to 6U-bases. This

difference (long poly-T stretch in DNA *versus* 6U in the EST and RNA reads) provides

evidence for editing at the RNA level. Illumina reads can map to each side of the poly-T

stretch, but because poly-T stretches can be longer than the read length, we are unable to

assemble the entire region using Illumina only or determine the length of the poly-T region.

1108

1109

Figure 3. Presence-absence of aaRS orthologs in different species.

1110

1111 Single eukaryotic (Euk): single copy aaRSs that function both in the cytosol and

1112 mitochondria. Other aaRSs are either expressed only in the cytosol (cytosolic) or in the

1113 mitochondrion (mitochondrial). In pink aaRSs presence, in white absence, and in blue the

1114 aaRS-Gly. Protein accession numbers are provided in Supplementary table S3, Supplementary

1115 Material online.

1116

1117

Figure 4. The position of *O. dioica* among chordates.

1118

1119 The phylogenetic tree was reconstructed based on mitochondrial protein sequences. Branch

1120 supports are only indicated for those cases in which the bootstrap support values were lower

1121 than 100%. The tunicates classes: Stolidobranchia in blue, Thaliacea in red, Phlebobranchia in

1122 green, Aplousobranchia in yellow, and Appendicularia in purple.

1123

1124

1125 **Table**

1126 **Table 1.** *Oikopleura dioica* **genes, position, length, direction (F- forward, R- reverse),**

1127 **start and stop codons, and comparison to the minimal, maximal, and average length of**

1128 **orthologous genes in 38 tunicate sequences (supplementary table S4, Supplementary**

1129 **Material online).**

Gene	Type	Position start	Position end	Length [bp] gene	Length [bp] CDS	Direction	Codon start	Codon stop	Length range in tunicates [bp] min	Length range in tunicates [bp] max	Average length in tunicates [bp]
srnl	rRNA	13	303	291	242	R			921	1,299	1,105
nad4	gene	934	2,847	1,914	876	R	ATC	TAA	1,209	1,401	1,321
cox3	gene	2,906	4,574	1,669	870	F	ATT	TAA	768	795	786
tRNA-M	tRNA	4,671	4,763	93	69	F			61	76	66
nad1	gene	4,805	5,800	996	615	F	TTG	TAA	853	1,035	914
cox1	gene	5,836	7,834	1,999	1,512	F	ATT	TAG	1,482	1,575	1,546
ODI_mt_ ORF196	gene	8,462	9,887	1,426	591	R	TTG	TAA			
cytb	gene	10,115	12,085	1,971	993	F	GTG	TAA	1,035	1,122	1,093
atp6	gene	12,312	13,063	752	516	R	ATT	TAA	492	642	619
rns	rRNA	13,064	13,295	232	215	R			596	984	696
cox2	gene	13,305	14,301	997	582	F	ATT	TAA	651	942	689
nad5	gene	14,456	18,371	3,916	1,248	F	ATA	TAA	1,458	1,788	1,695

1130

1131

Oikopleura dioica
mitochondrial genome
18,390 bp

	AA	Oikopleura dioica Norway	Oikopleura dioica Osaka	Ciona intestinalis	Styela clava	Homo sapiens
Single Euk.	A	■	■	■	■	■
	G			▧	▧	▧
	H	■	■	■	■	■
	Q	■	■	■	■	■
Cytosolic	G	▧	▧			
	C	■	■	■	■	■
	D	■	■	■	■	■
	E/P	■	■	■	■	■
	Fa	■	■	■	■	■
	Fb	■	■	■	■	■
	I	■	■	■	■	■
	K	■	■	■	■	■
	L	■	■	■	■	■
	M	■	■	■	■	■
	N	■	■	■	■	■
	R	■	■	■	■	■
	S	■	■	■	■	■
	T	■	■	■	■	■
	V	■	■	■	■	■
	W	■	■	■	■	■
	Y	■	■	■	■	■
Mitochondrial	G	▧	▧			
	C			■	■	■
	D			■	■	■
	E			■	■	■
	F	■	■	■	■	■
	I			■	■	■
	K					
	L			■	■	■
	M	■	■			
	N			■	■	■
	P			■	■	■
	R			■	■	■
	S			■	■	■
	T					
	V					■
	W			■	■	■
	Y			■	■	■