Silybin B, resveratrol and epigallocatechin-3 gallate (EGCG) bind to troponin to restore the loss of lusitropy caused by cardiomyopathy mutations *in vitro, in vivo*, and *in silico*

Zeyu Yang − Institute of Chemical Biology, Molecular Sciences Research Hub and Department of Chemistry, Molecular Sciences Research Hub, Imperial College London, London W12 0BZ, U.K. Email: zeyu.yang14@imperial.ac.uk

Alice Sheehan   NHLI, Imperial College London, London, W12 ONN, United Kingdom, Email: sheehan.ecila@gmail.com

Andrew Messer   NHLI, Imperial College London, London, , W12 ONN, United Kingdom, Email: a.messer@outlook.com

Sharmane Tsui   NHLI, Imperial College London, London, , W12 ONN, United Kingdom, sharmanetsui98@gmail.com

Alexander Sparrow Division of Cardiovascular Medicine, Radcliffe Department of Medicine, University of Oxford, Oxford, United Kingdom   and British Heart Foundation Centre of Research Excellence, University of Oxford, Oxford, United Kingdom; https://orcid.org/0000-0002-3354-3314 Email: alexander.sparrow@cardiov.ox.ac.uk

Charles Redwood Division of Cardiovascular Medicine, Radcliffe Department of Medicine, University of Oxford, Oxford, United Kingdom and British Heart Foundation Centre of Research Excellence, University of Oxford, Oxford, United Kingdom; https://orcid.org/0000-0002-8673-7542 Email: charles.redwood@cardiov.ox.ac.uk

Vladimir Kren: Laboratory of Biotransformation, Institute of Microbiology of theCzech Academy of Sciences, Prague, Czechia https://orcid.org/0000-0002-1091-4020. Email: kren@biomed.cas.cz

Ian R. Gould − Department of Chemistry, Molecular Sciences Research Hub and Institute of Chemical Biology, Molecular Sciences Research Hub, Imperial College London, London W12 0BZ, U.K.; orcid.org/0000-0003-3559-0234; Email: i.gould@imperial.ac.uk

Steven B. Marston − National Heart & Lung Institute, Imperial College London, London W12 0NN, U.K. orcid.org/0000-0001-6054-6070; Email: s.marston@ imperial.ac.uk

ABSTRACT

Adrenergic activation of protein kinase A (PKA) targets the sarcolemma, sarcoplasmic reticulum, and contractile apparatus to increase contractile force and heart rate. In the thin filaments of the contractile apparatus, cTroponinI Ser22 and Ser23 are the targets for PKA phosphorylation. The effect of phosphorylation is a 2-3 fold decrease of affinity of cTn for Ca^{2+}, associated with a higher rate of Ca^{2+} dissociation from cTnC leading to a faster relaxation rate of the cardiac muscle (lusitropy). This modulation of Ca^{2+}-sensitivity is often suppressed by mutations that cause cardiomyopathy (uncoupling) and this could be sufficient to induce cardiomyopathy. Therefore, a drug that could restore the phosphorylation-dependent modulation of Ca^{2+}-sensitivity could have potential for treatment of these pathologies. We have found that in single filament assays that a number of small molecules including silybin B, resveratrol and EGCG can restore coupling.

We performed molecular dynamics simulations of the unphosphorylated and phosphorylated cardiac Troponin core with the TNNC1 G159D DCM mutation. We found that silybin B, EGCG and resveratrol restored the phosphoryation-induced change of most metrics to wild-type values, whilst silybin A, an inactive isomer of silybin B, and ECG did not. We analysed the atomic-level changes induced by ligand binding to explain recoupling.

In parallel, we have extended our studies to intact TNNT2 R92Q-transfected cardiomyocytes. The mutation blunts the increased relaxation speed response to β1 adrenergic stimulation and we found that resveratrol, EGCG and silybin B could restore the β1 adrenergic response whilst silybin A did not.

INTRODUCTION

In cardiac muscle, contractility is controlled by changes in the concentration of intracellular Ca^{2+} ion that binds to troponin, a component of the thin filaments, to activate the contractile interaction of thick and thin filaments. In addition to this Ca^{2+} switch, cardiac muscle possesses a unique modulatory mechanism that allows the heart to meet increased oxygen demand during exercise.

Release of adrenaline and noradrenaline activates *β*-1 receptors in cardiac myocytes and leads to activation of adenylyl cyclase *via* stimulatory G-protein (Gs). The

resulting increase of the cytosolic cyclic adenosine monophosphate (cAMP) levels leads to activation of protein kinase A (PKA), which phosphorylates a number of targets in the sarcolemma, sarcoplasmic reticulum, and contractile apparatus to increase contractile force and heart rate. In the thin filaments of the contractile apparatus, troponinI (cTnI) is the target for PKA phosphorylation and is phosphorylated at residues Ser22 and Ser23 in the cardiac-specific N-terminal peptide (NcTnI: residues 1 to 32). The effect of phosphorylation is a 2-3 fold decrease of affinity of cTn for Ca^{2+} due to altered cTnC-cTnI interactions, linked to a higher rate of Ca^{2+} dissociation from cTnC [1-4]. The consequent faster relaxation rate of the cardiac muscle due to TnI phosphorylation (lusitropy) is essential for shortening the cardiac muscle contraction-relaxation cycle allowing for efficient contraction at a faster heart rate.

It is well established that familial hypertrophic cardiomyopathy (HCM) and familial dilated cardiomyopathy (DCM) are often caused by single mutations in one of the contractile protein genes. These mutations primarily affect Ca^{2+} regulation or the PKA-dependent modulatory system, either mimicking the Ca^{2+}-sensitivity shifts caused by phosphorylation or interfering with them such that Ca^{2+}-sensitivity becomes independent of phosphorylation level (uncoupling) [5-7]. Examples of well-studied mutations that cause uncoupling include the mutation in cardiac troponin C, cTnC G159D, that causes familial dilated cardiomyopathy [8-11] and the HCM-linked mutations in troponinT (TnT) R92Q and cardiac actin E99K [12-17]. Uncoupling results in blunting of the lusitropic response to β-1 receptor stimulation, thus abnormalities of myofilament Ca^{2+}-sensitivity and its physiological modulation by phosphorylation could be sufficient to induce the disease phenotype [7, 18, 19].

A drug that could restore the phosphorylation-dependent modulation of Ca^{2+}-sensitivity could have potential for treatment of these pathologies. We have investigated small molecules that can specifically reverse these abnormalities *in vitro*. Based on our lead compound, epigallocatechin-3-gallate (EGCG) [20, 21] that is both a desensitiser and a recoupler, we examined 40 compounds and found 23 compounds that reversed the uncoupling. Many of these are pure recouplers whilst 3 of the compounds desensitized but did not recouple and the rest had no effect [22]. Importantly, recoupling was complete and independent of the causative mutation and the nature of the compound. Thus, small molecules that can act as desensitisers or

recouplers have the potential as treatments for HCM or DCM. However we do not know the molecular mechanism of the recoupling phenomenon, nor do we know whether the recoupling compounds we investigated *in vitro* are effective in intact muscle. In this paper, we aim to answer these questions *in vitro, in silico, and in vivo.*

Understanding how phosphorylation modulates troponin function is not straightforward since troponin contains several disordered regions that are crucial to regulation and are not accessible to methodologies such as X-ray diffraction or cryoelectron microscopy that only resolve static structures. We recently used molecular dynamics (MD) simulation to determine the atomistic changes that occur when Ser22, 23 of cTnI is phosphorylated in the regulatory core of troponin. These studies showed that phosphorylation induced no significant stable structural changes but rather phosphorylation changed the dynamics of the transitions between a small number of states [23, 24]. We established metrics characteristic of the ser22/23 phosphorylation transition: changes in the distribution of the angle between helices A and B of troponin C and the angle between the NcTnC domain and the ITC domain. Changes in these metrics were consistent with phosphorylation destabilising the open state of TnC. Importantly, the introduction of the TnC G159D mutation profoundly changed these metrics. Unphosphorylated G159D tended to have values equivalent to phosphorylated wild-type and upon phosphorylation the values for G159D moved in the opposite direction to WT [24].

The analysis of MD simulations of G159D troponin clearly shows that the mutation abrogates the effects of phosphorylation on structure and dynamics and thus explains its uncoupling activity. We may also use these metrics to determine the effects of small molecules on the system and to test whether recoupling *in vitro* corresponds to the restoration of WT dynamics *in silico*.

We analysed molecular dynamics simulations of unphosphorylated and phosphorylated cardiac Troponin core with the G159D mutation in the presence of two pure recouplers, silybin B (SB) and resveratrol, and epigallocatechin-3 gallate (EGCG), a mixed recoupler and desensitiser. These were compared with silybin A, the stereoisomer of silybin B and epicatechin gallate (ECG), closely related to EGCG, that have no measurable recoupling activity *in vitro* as controls. We found that silybin B, EGCG, and resveratrol restored most key metrics towards wild-type

values, whilst silybin A (SA) and ECG did not. EGCG gave a complex response consistent with its dual function as recoupler and desensitiser.

In parallel, we have extended our studies to intact cardiomyocytes. Mutations that uncouple *in vitro* blunt the response to β1 adrenergic stimulation in intact cardiomyocytes and cardiac tissue [19]. We have therefore tested the effects of small molecules, that recouple *in vitro,* on the response to a β1 adrenergic agonist. We found that resveratrol, EGCG and silybin B could restore the β1 adrenergic response whilst silybin A did not; in the case of EGCG the response was partly confounded by off-target effects.

2. RESULTS

2.1 *In vitro effects of small molecules*

In previous studies, we demonstrated that small molecules, including EGCG, silybin B, and resveratrol were able to restore the lack of response to troponin I phosphorylation caused by mutations in thin filament proteins, whereas the stereoisomer silybin A and ECG did not [21, 22]. Figure 1A summarises the effects of these compounds on wild-type thin filaments and upon thin filaments containing the uncoupling HCM mutation TPM1 E180G. Figure 1B shows the effects of EGCG, silybin B and silybin A on the Ca^{2+}-regulation of thin filaments containing the TNNC1 (cardiac TnC) G159D mutation. None of the compounds affect the phosphorylation-dependent Ca^{2+}-sensitivity shift in wild-type thin filaments. The Ca^{2+}-sensitivity of mutant thin filaments is not modulated by phosphorylation (uncoupled), however, coupling is restored to both TPM1 E180G and TNNC G159D by 100µM EGCG, resveratrol or silybin B but not by silybin A or ECG (supplement Figure 4C). In addition, EGCG, ECG and silybin A are desensitisers, reducing Ca^{2+}-sensitivity in wild-type and mutant by about 1.5-fold independently of the recoupling property, thus silybin B and resveratrol are 'pure' recouplers whilst EGCG is a mixed desensitiser and recoupler. These results are in accord with the results of previous studies showing that the recoupling effect of small molecules, silybin B, EGCG, and resveratrol is independent of the causative mutation and that a full restoration of the effects of troponin I phosphorylation is achieved by all recouplers [21, 22] see supplement table 4.

Figure 1

 Small molecules that can restore the response to phosphorylation in mutant thin filaments.

 Measurement made by in vitro motility assay in reconstituted human thin filaments.

A- Fixed [Ca²⁺] assay (~EC50) using WT or TPM1 E180G HCM mutant . The increased motility due to dephosphorylation is plotted. Small molecules do not affect wild-type response (red and pink bars) but EGCG, silybin B and resveratrol restore the dephosphorylation effect to E180G whilst silybin A does not (black and green bars).

B- The same effect of the small molecules is shown by the measurements of the Ca²⁺-activation curve for phosphorylated and unphosphorylated DCM mutant (TNNC1 G159D) thin filaments. Purple G159D thin filaments, green G159D + small molecule. Solid line phosphorylated, dotted line unphosphorylated. EGCG and silybin B restore the Ca²⁺ shift due to phosphorylation of troponin I. Note that with Silybin A, Phosphorylated and unphosphorylated fits are superimposed whilst with Silybin B, the unphosphorylated fit is superimposed on apo (purple) fits. EC50 values are given in Supp Table 1.

2.2 Contractility of isolated cardiomyocytes

In order to assess the effects of EGCG, silybin A and silybin B on uncoupling in intact cells, we isolated cardiomyocytes from wild-type and two well-characterised models of hypertrophic cardiomyopathy that are uncoupled *in vitro* due to mutations in a thin filament protein. The models were ACTC E99K transgenic mouse myocytes

and guinea-pig cardiomyocytes with the TNNT2 (TnT) R92Q mutation replacing the wild-type protein, introduced by transfection [12, 15, 16].

The contractility of wild-type and mutant myocytes was measured. We studied three main parameters: the contractile amplitude (% peak height), the time of contraction (time to 90% peak, ttp_{90}), and the time of relaxation (time to 90% baseline, ttb_{90}) , see Figure 2A and Supp Table 2. Whilst the contraction amplitude of wild-type mouse and guinea-pig myocytes was similar, ttp_{90} and ttb_{90} in guinea-pig were about double that of mouse, indicating slower kinetics.

Cardiomyopathic mutations affected the baseline contractility of cardiomyocytes as previously noted. Both ttp_{90} and ttb_{90} were greater in ACTC E99K mouse cardiac cells compared to WT cells by 41% and 46% respectively, indicating slower contraction and relaxation whilst contractile amplitude was not different. In the Guinea-pig TNNT2 R92Q cells, ttp_{90} was the same but ttb_{90} was 35% greater than the wild type indicating slower relaxation due to the mutation.

The normal response to β1 receptor activation is increased amplitude of shortening and faster contraction and relaxation (lusitropy). In contrast, mutations that are observed to be uncoupled *in vitro* have a blunted response to β1 adrenergic stimulation in intact cardiomyocytes and cardiac tissue [19]. We confirmed this finding in our preparations (Figure 2A).

We treated cells with the β1-selective agonist dobutamine in the presence of the highly selective β2 receptor antagonist, ICI-118,551. In mouse cells, dobutamine increased the amplitude of shortening by 47% and decreased ttb_{90} by 21%, thus indicating both inotropic and lusitropic effects. In guinea-pig cells dobutamine did not affect amplitude but ttb_{90} was decreased by 21% indicating normal lusitropy.

In the mutant cells, we did not observe any significant lusitropy, indicating that β1 adrenergic stimulation and the consequent troponin I phosphorylation was uncoupled from its effect on the rate of relaxation (Supp Table 2, Figure 2A). We then tested the ability of small molecules to restore ttb_{90} to WT levels in the presence of dobutamine.

Figure 2A

Effects of dobutamine on cardiac myocyte contractility.

Amplitude, ttp_{90} and ttb_{90} parameters are shown. Blue- WT and E99K mouse compared. Purple- WT and R92Q guinea-pig compared. Full data and statistical analysis in Supplement 2

Figure 2B

Lusitropy and the effect of small molecules measured in cardiomyocytes

Lusitropy is defined as the fractional change of ttb_{90} due to dobutamine [1-(ttb_{90} +Dob/ ttb_{90} -Dob)]. Positive lusitropy (increase in relaxation speed) corresponds to a decrease in this parameter. Blue, mouse experiments, Purple, Guinea-pig experiments. Original data and statistical analysis in Supplement Table 3

EGCG, silybin A, silybin B, and resveratrol were examined as potential re-coupling compounds. Mouse E99K or guinea-pig R92Q myocytes were incubated in the presence of the small molecules and contractility was measured before and after adding 0.4µM dobutamine. Preliminary titrations indicated that maximal effects of these molecules were achieved with micromolar concentrations, much lower than the

concentrations needed for the *in vitro* motility assays. We quantified lusitropy as the fractional change of ttb_{90} due to dobutamine [1- (ttb_{90} +Dob/ttb_{90} -dob)]. The baseline lusitropy in wild-type mouse or guinea-pig was a 20-24% *decrease* whilst in the uncoupled mutant myocytes dobutamine induced a non-significant *increase* in ttb_{90} (12% in mouse, 9% in guinea-pig) (Figure 2).

In the presence of the known recoupling molecules, there was a clear restoration of the lusitropic effect. In the presence of silybin B, dobutamine reduced ttb_{90} by 25% in E99K mouse and 13% in R92Q guinea pig. In the presence of resveratrol, dobutamine reduced ttb_{90} by 41% in E99K mouse and 17% in R92Q guinea-pig. With EGTA, dobutamine reduced ttb_{90} by 22% in E99K mouse and reduced ttb_{90} by 17% in R92Q guinea-pig. In contrast, silybin A, which does not recouple *in vitro*, did not restore lusitropy; we observed a non-significant 4% *increase* in ttb_{90} in E99K mouse and an 18% *increase* in R92Q guinea pig myocytes which is similar to the effect of dobutamine in the absence of small molecules (9% increase), see Figure 2B and Supplement table 3.

Thus, the small molecules that were shown to be recouplers *in vitro* also restored the blunted lusitropic response to β1 adrenergic stimulation in intact mutant myocytes. The specificity of this effect is confirmed by the inability of silybin A to restore lusitropy.

2.3 Effects of small molecules on troponin phosphorylation determined by Molecular Dynamics Simulations

To understand how the small molecules modulated troponin behaviour at the atomic level we performed molecular dynamics simulations (5×1500ns) on phosphorylated and unphosphorylated wild-type and TnC G159D DCM mutant troponin in the presence of silybin A, silybin B, EGCG, ECG and resveratrol.

2.3.1 Recoupling ligands can restore the phosphorylation-dependent changes in molecular dynamics metrics that are suppressed in TnC G159D mutant

In previous molecular dynamics simulations, we demonstrated that phosphorylation of troponin I does not significantly alter the structure of the troponin core. However,

phosphorylation does change the dynamics of troponin and the uncoupling mutant TnC G159D. We have derived several metrics that are characteristic of the phosphorylated and unphosphorylated state. In particular, the distribution of the angle between cTnC helices A and B and the angle begween quasi-rigid troponin domains (NcTnC and the ITC domain) may be used since the effect of TnI phosphorylation on the metrics of the uncoupling TnC G159D mutation are strikingly different from wild-type with phosphorylation-dependent changes in G159D being mostly in the opposite direction to those in wild-type [24]. Figure 3A shows the angle distributions comparing WT and G159D troponin whilst Figure 3B shows the effect of small molecules and phosphorylation on the angle distributions.

2.3.2   Angle between troponin C helices A and B

The A/B helix angle may be used as a metric for the opening and closing of the hydrophobic patch (Table 1A). In WT, the effect of phosphorylation is a decrease of the mean A/B angle, from 101.8° to 96.1° (Δ= - 5.6°). For the G159D mutant we observe the complete inverse behaviour: upon phosphorylation, the A/B angle increases from 97.1° to 99.0° (Δ= +1.9°). Comparing the WT and G159D, unphosphorylated A/B angles we see that the WT angle is significantly higher than G159D.

Figure 3 The distribution of a helix A/B and Interdomain angles.

A   The effects of the G159D mutation and phosphorylation on the distribution of angles

B   The effects of phosphorylation and small molecules on the distribution of A/B angle and interdomain angle. For G159D. Table 1 shows the quantification of the distributions. Distribution plots for ECG are shown in supplement Figure 4D.

Figure 3

We determined the effect of the small molecule ligands on the interhelix angle distribution in G159D and wild-type troponin (Figure 3, Table 1B). In wild-type the ligands tended to have no effect or to increase A/B angle in both unphosphorylated and phosphorylated states, however, the decrease in mean angle upon phosphorylation, associated with a normal response to phosphorylation was preserved. In G159D the effects of the small molecules were different, the three recouplers restored the decrease in A/B angle on phosphorylation (SB, -3.1°, EGCG, -13.3°, RVL, -11.1° compared with -5.6° in the wild-type) whereas with silybin A, which has no recoupling activity *in vitro,* the change resembled that of G159D (SA, +4.88°, ECG, -0.2° compared with ApoG159D +1.9°). This pattern of results was also clear when considering the percentage of the area under the curve for the A/B angle above 110°. With G159D this parameter increased from 1 to 7% on phosphorylation. It increased from 11 to 24% in the presence of silybin A and from 4 to 6% with ECG but with the three recoupling molecules the change was in the opposite direction (SB, 14>9%, EGCG, 11>3%, RVL 25>1% compared with 18 > 1% in wild-type troponin). Thus the three small molecules that are physiological recouplers restore the phosphorylation-dependent changes in the disease-causing, uncoupled, G159D mutation at the atomic level and silybin A and ECG that are not recouplers, do not restore the A/B angle change.

Table 1A

A/B helix angle metrics

troponin	ligand		uP mean (std)	Δ	P		uP	Δ	P		uP		P	
			Mean angle				**% > 110**				**FWHM**			
WT	apo		101.81 (9.15)	-5.65	96.16 (5.73)		18	-17	1		17.11	-4.95	12.16	
G159D	apo		97.06 (5.48)	1.91	98.97 (6.73)		1	6	7		12.48	1.16	13.64	
G159D	EGCG		102.25 (6.07)	-13.25	89.0 (18.07)	✅	11	-8	3	✅	12.77	1.36	14.13	
G159D	RVL		105.48 (9.16)	-11.14	94.34 (8.72)	✅	25	-24	1	✅	14.81	5.58	20.39	
G159D	SilybinA		98.55 (10.98)	4.88	103.43 (10.94)	❎	11	13	24	❎	17.39	-3.28	14.11	
G159D	SilybinB		103.61 (5.81)	-3.09	100.52 (7.45)	✅	14	-5	9	✅	14.24	0.81	15.05	
G159D	ECG		98.1 (6.4)	-0.2	97.9 (6.9)	→	4.0	2	6.0	❎	16.1	-2.6	13.5	*

Table 1B

troponin	ligand	Mean hinge angle (std)				FWHM			
		uP		P		uP	Δ	P	
WT	apo	121.61 (7.24)	1.65	123.26 (5.95)		21.97	-5.54	16.43	
G159D	apo	121.51 (4.82)	-5.31	116.2 (6.04)		11.49	0.82	12.31	
G159D	EGCG	114.62 (9.18)	-7.03	107.59 (14.53)	?	23.99	-1.51	22.48	✅
G159D	RVL	112.61 (7.64)	9.28	121.89 (6.99)	✅	18.34	-4.41	13.93	✅
G159D	SilybinA	126.65 (5.77)	-6.35	120.3 (8.47)	❌	14.13	6.64	20.77	
G159D	SilybinB	117.74 (5.6)	3.58	121.32 (4.98)	✅	11.53	0.02	11.55	✅
G159D	ECG	118.9 (9.0)	-8.3	110.6 (9.0)	❌	11.2	0.5	11.7	

1A   *Quantification of the effects of phosphorylation and small molecules on A/B angle metrics*.

1B   *Quantification of the effect of phosphorylation and small molecules on interdomain hinge angle metrics.* Ticks indicate where the wild-type changes on phosphorylation are restored by the small molecules. Crosses indicate no restoration.

2.3.3   Effect of small molecules on hinge motion.

The principal motion of troponin is a hinge-like motion between the two quasi-rigid domains of troponin, NcTnC and the ITC domain. The distribution of hinge angles in wild-type troponin shows an increase in the mean angle upon phosphorylation and a lengthening of the TnC peptide link between the domains. In contrast, phosphorylation of G159D shifts the inter-domain angle distribution towards a lower mean angle. Thus the wild-type response to phosphorylation is disrupted by the G159D mutation at the atomic level (Figure 3B, Table 1B, Table 2A).

When the recoupling molecules SB and RVL were added to the G159D simulation, the increase in mean angle upon phosphorylation was restored toward wild-type (SB, +3.58°, RVL, +9.28° compared with WT, +1.65°). In contrast, adding silybin A or ECG to G159D did not reverse the hinge angle change and resembled untreated G159D (SA, -6.4°, ECG -8.3° compared with apoG159D. -5.3°). The effect of EGCG on G159D hinge angle was anomalous: the mean angle decreased by 7.0° despite EGCG being a recoupling molecule. However, The EGCG angle distribution is bimodal or tri-modal suggesting multiple actions. If just the prominent peak centred on 103° is considered, EGCG may behave like the other recouplers. EGCG is both a recoupler and a desensitiser of actomyosin and this second property, not shared with SB or RVL, may be responsible for the occupancy below 80°. This is supported by the observation that the distribution plot for ECG which is a pure desensitizer is simpler and lacks the 80° component (Supplerment Figure 4D).

It is also notable that phosphorylation and mutations affect the broadness of the distribution of hinge angles which is reduced on phosphorylation of wild-type but not with G159D. G159D is substantially more rigid than WT and it was proposed by Yang et al. that the strong bond between TnC R83 and D159 across the hinge played a part here. It is thus noteworthy that this interaction is substantially weakened by all the ligands binding even though there is no evidence for any ligand binding close to this site, implying an allosteric effect (Table 2A).

Table 2A

		Interdomain distance metrics								TnC G/D159 – R83 ionic bond	
		Mean distance				FWHM					
troponin	ligand	uP mean (std)	Δ	P		uP	Δ	P		uP	P
WT	apo	30.18 (1.34)	-0.28	29.9 (0.9)		2.99	-0.61	2.38		0	0
G159D	apo	29.22 (0.98)	0.35	29.59 (1.18)		1.57	-0.06	1.51		30062	28544
G159D	EGCG	31.17 (1.48)	0.81	31.98 (1.59)	?	2.38	-0.57	1.81	✅	8326	19890
G159D	RVL	32.13 (1.69)	-0.74	31.39 (1.51)	✅	4.94	-2.82	2.12	✅	18945	17269
G159D	SilybinA	30.09 (1.5)	0.75	30.84 (1.04)		4.4	-1.97	2.43	?	3101	15131
G159D	SilybinB	31.83 (1.83)	-1.11	30.72 (1.13)	✅	1.9	0.74	2.64	?	6554	8506
G159D	ECG	30.2 (1.2)	0	30.2 (1.4)	<>	1.8	0.7	2.5		30750	32250

Table 2B

		MMPBSA metrics							Fraction of time attached	
		MMPBSA Switch				MMPBSA Ligand binding				
troponin	ligand	uP mean (std)	Δ	P		uP	Δ	P	uP	P
WT	apo	-108.43 (24.05)	-3.46	-111.89 (13.91)						
G159D	apo	-113.47 (13.75)	-3.53	-117.0 (15.01)						
G159D	EGCG	-109.26 (13.73)	-13.6	-122.87 (13.56)		-26.12 (11.19)	-0.71	-25.41 (10.14)	0.93	0.88
G159D	RVL	-104.91 (15.69)	-2.02	-106.93 (15.23)		-13.71 (8.46)	-1.36	-12.35 (7.56)	0.49	0.33
G159D	SilybinA	-113.65 (15.03)	1.8	-111.85 (16.08)		-25.48 (9.21)	-6.36	-19.12 (9.26)	0.95	0.82
G159D	SilybinB	-106.72 (17.92)	0.07	-106.79 (13.52)		-26.16 (10.04)	0.47	-26.63 (10.69)	0.95	0.88
G159D	ECG	-108.1 (15.2)	-4.9	-113.0 (16.6)		-25.0	0.9	-25.9	0.92	

2A   Quantification of the effect of phosphorylation and small molecules on interdomain distance and on the interdomain ionic bond between TnC G/D159 and R83. Ticks indicate where the wild-type changes on phosphorylation are restored by the small molecules.

2B   Quantification of the effect of phosphorylation and small molecules on switch peptide of TnI binding to troponin C and ligand binding to troponin. Ligand binding parameters

2.3.4   Binding of small molecules to troponin

The molecular dynamics simulations allowed us to determine how the small molecules interact with G159D troponin and how this modifies the response to phosphorylation.

Figure 4   Parameterised solution configurations

silybin A (magenta) silybin B (green), EGCG (yellow) and resveratrol (blue). A and D rings of silybin A and B and A and C rings of EGCG are aligned. See Figure 1 and Supplement Figure 1 for identification of the atoms and rings.

Figure 4 illustrates the preferred solution conformation of the ligands considered in this paper. silybin B and A are approximately flat; they are stereoisomers with the C ring in opposite orientations. The A, C and D rings of EGCG has considerable homology with silybin but the orientation of the B ring gives the molecule a propellor-like shape. Resveratrol likewise has some structural homology with silybin and EGCG.

2.3.4.1 Binding energy

MMPBSA was used to calculate ΔG for ligand binding (Table 2B). ΔG was in the range 25.5-26.2 kJ/mole for EGCG, ECG, silybin A and silybin B. This corresponds to a dissociation constant of around 6nM. ΔG for resveratrol appeared to be lower, consistent with its radically different binding footprint on troponin compared with the other recouplers. For binding to G159D, ΔG did not change much with

phosphorylation except for silybin A, which does not recouple, where binding became significantly weaker on phosphorylation.

2.3.4.2 Ligand binding hotspots on troponin

We identified the most probable contacts on troponin using the CPTraj procedures. The plot in Supp Figure 2A shows the probability of interactions between the ligand and each amino acid of the troponin core. Although it is quantitative, the plot is one-dimensional and gives no idea of the stereochemistry of the interactions. In every case, discrete contact patterns were observed in unphosphorylated G159D that were different for each of the four small molecules and that changed upon phosphorylation as would be expected if they influence the phosphorylation-dependent structural and dynamic changes we have demonstrated.

One way of visualizing the interaction hotspots is to plot all contacts above a threshold (in this case interactions present >10% of the time) on representative models of the troponin core structure, see Supp Figure 2B and movies, SUPP 2C. Taken together these plots show that ligand binding is phosphorylation dependent. In general, hotspots tended to be more dispersed after phosphorylation indicating that ligands explore a wider range of interactions compared with unphosphorylated although overall ΔG was not changed. Many of the phosphorylation-dependent hotspots are near the NcTnI-NctnC interface or the interdomain interface. It is also apparent that ligand interaction can change the conformation of the troponin core, consistent with the angle changes discussed above. Apparent contacts with the tail of troponin (around the TnI helix 1-helix 2 turns and the linker before TnT helix1) are likely to be artefactual since these parts of troponin are likely to be in contact with the rest of the thin filament *in vivo* [25] [26].

In these plots there appear to be to be multiple discrete hotspots; examples of this is silybin B binding to unphosphorylated G159D where the hotspot appears to extend over a large area of the NcTnC surface which would not be possible for a single site given the size of silybin B and, in EGCG binding to unphosphorylated G159D, there are two hotspots on opposite sides of the troponin again not possible to be occupied by EGCG simultaneously. This is most likely due to the time-averaged nature of the CPTraj calculations whereas in reality, ligand binding is a dynamic process. The

ligand may occupy a number of locations at different times all of which appear in the model.

In order to understand ligand binding it is necessary to study it in 4 dimensions- this can only be done by directly observing the binding trajectories (Figure 7 and supplement Figure 3). Observation confirms the stochastic nature of ligand binding. A ligand can bind at several discrete locations as was suggested for silybin B and EGCG. In the five 1500ns runs, the initial locations can be different. Often the ligand goes to one location and stays there for prolonged periods, however, the ligand can move during a run, either by 'creeping' across the troponin or more often by dissociation and rebinding. In some cases the ligand is free for a substantial part of the simulation, most notably with reseveratol (See Table 2B).

We searched for a pattern of contacts that was associated with recoupling. The supplementary Figure 3 shows the visual analysis of selected whole trajectories and highlights those positions where the ligand is often found. Observation suggests for silybin B there are two common positions: at the end of Tn, intercalated between the NcTnI and NcTnC A and B helices and at the 'top' of the hinge, interacting with CcTnT and inhibitory peptide. SB is bound nearly all the time to unphosphorylated G159D and tends to go to one site or the other. Therefore the time average in Supplement Figure 2 is a overwhelmingly a mixture of two separate hotspots. Figure 5 shows a frame from one run where SB occupies site A for most of the 1500ns run. When phosphorylated (SEP), silybin B is almost never at site A. It is often in parts of the tail and CcTnC and sometimes also at top of hinge like uP SB site B.

Silybin A also binds primarily near the N terminus of TnI and the helix A-B loop of EF I (Figure5 and Supp Figure3). Unlike Silybin B Silybin A does not appear to be intercalated between NcTnI and NcTnC but sits on the surface with only the A ring engaged . A secondary site is at the 'top' of the interdomain hinge region. All of these contacts are lost on phosphorylation.

Silybin A   Resveratrol

Figure 7

Silybin A   Silybin B

EGCG   Resveratrol

ECG

Figure 5

Ligands bound to troponin in their most probable state.

Top- ball and stick represenstaion of the ligand interacting with TnC(light grey) and TnI(blue). TnT is light brown. Ligand is red The molecule is orientated to give the best view of the interaction.

Bottom- Whole troponin surface representations. TnI is blue and TnC is cyan (light brown for resveratrol). Ligand is red.

 Full trajectories are illustrated in Supplement Figure 3.

____

In unphosphorylated G159D, EGCG occupies the site intercalated between NcTnI and NcTnC (site A) identified in silybin B part of the time, although it does not use the same set of contacts and there are additional hydrogen bonding to the switch peptide, perhaps associated with its desensitising property (Figure 5 and Supp 3). None of these sites are occupied when troponin is phosphorylated.

In contrast ECG primarily occupies a site on the opposite side of NcTnC where it sits on the surface approximately flat with the B ring extending away from troponin.

Resveratrol binds less frequently to troponin than the others (it was free for 45% of the MD runs see Table 2B) and most often makes contacts with a completely different region of troponin, inserted into the C-terminal domain of troponin C close to the interdomain interface (Figure 5 and Supp Figures 3 and 4E).

3 DISCUSSION

3.1 Silybin B, EGCG and resveratrol can restore lusitropy to mutant myocytes that have impaired lusitropy.

Lusitropy, the increase in the rate of cardiac muscle relaxation, is an essential component of the heart's response to adrenergic activation [19]. It is predominantly mediated by PKA phosphorylation of cardiac troponin I [27, 28]. However in genetic cardiomyopathies due to mutations in thin filament proteins, lusitropy is much reduced and *in vitro* PKA phosphorylation does not alter Ca^{2+}-sensitivity, the usual measure of the effect of phosphorylation [7, 19]. Basic and animal studies indicate that specific suppression of lusitropy can induce symptoms of heart failure under stress and clinical studies have shown that lack of response to adrenergic activation is associated with increased adverse cardiac events [29] [19]. Thus suppression of lusitropy by mutations may be a significant disease mechanism for cardiomyopathy.

Restoration of the lusitropic response is therefore a suitable target for therapy. We have investigated a series of small molecules, acting on troponin, that can restore the phosphorylation-dependent modulation of Ca^{2+}-sensitivity (recoupling) at the single myofilament level as potential treatments [22] [21] . These molecules appeared to be promising since they all gave total restoration *in vitro* that was independent of the mutation that caused the dysfunction in lusitropy (Summarised in Supplement Table 6). However, it was not established these small molecules had any effect on intact muscle and the molecular mechanism by which many different small molecules could cause a single outcome, recoupling, was unknown.

In this study, we have addressed these questions for five small molecules with a range of functions and structures. Silybin B and resveratrol are structurally distinct yet both have a pure recoupling activity *in vitro*, whilst EGCG is bifunctional, being both a recoupler and a desensitiser. As negative controls, we studied silybin A, the stereoisomer of silybin B, and ECG, a close analogue of EGCG, that have no recoupling activity i*n vitro*.

We tested these small molecules for their ability to restore lusitropy in two previously studied mutant systems. Myocytes from ACTC E99K mice and guinea-pig myocytes transfected with the TNNT2 R92Q mutation were challenged with the β1 specific adrenergic agent dobutamine [12, 16]. The results correlated with *in vitro* results: the

mutant myocytes had suppressed lusitropy but wild-type lusitropy could be restored by silybin B, EGCG and Resveratrol but not by silybin A. This result confirms that recouplers studied *in vitro* do restore lusitropy, meaning that these small molecules have therapeutic potential.

3.2 Silybin B, EGCG and resveratrol can restore the dynamics of the G159D mutant troponin core to wild-type values.

To understand the molecular mechanism by which silybinB, EGCG and resveratrol restore lusitropy we extended our molecular dynamics studies of the structure and dynamics of the troponin core that have defined how phosphorylation at cTnI Ser22 and 23 modulates the dynamics of troponin and how the DCM-associated mutation, G159D interferes with this modulation [24, 30] . Several phosphorylation-dependent metrics may be derived from the MD simulations; the most useful for this study are the change in the distribution of the angle between the A and B helices of cTnC (a measure of the opening and closing of the hydrophobic patch) and changes in the hinge angle between the two quasi rigid domains of troponin (NcTnC and cTnI 1-33 vs CcTnC / TnT / TnI helices H1 and H2 see [31]).

Phosphorylation, in wild-type, causes an increase in the mean A/B helix angle and a reduction in the range of motion (FWHM) ; phosphorylation decreases the mean hinge angle whilst also reducing the FWHM parameter.

In contrast, phosphorylation of G159D troponin has an opposite effect on most of the metrics compared with wild-type. The G159D mutant troponin has an intrinsically lower A/B helix angle and FWHM than wild-type and phosphorylation causes an increased mean helix A/B angle whilst there is a decrease in interdomain hinge angle when G159D is phosphorylated. We have proposed that these abnormalities seen in G159D dynamics are causative of the uncoupling effect of the mutation and therefore could be used to test for recoupling.

In the presence of the three recoupling compounds the parameters for A/B helix angle and the interdomain hinge angle of G159D troponin are largely restored to wild-type values (See Figure 6 for a summary), thus recoupling involves restoration of wild-type dynamic behaviour in troponin at the atomic level. It is perhaps relevant that the strong ionic and H bonding between the mutant amino acid cTnC D159 and

T83 across the interdomain interface, thought to be responsible for the reduced range of hinge motion in the mutant, is much reduced by ligand binding. It has been noted that ligands may occupy multiple sites only one of which may be relevant to recoupling and also that ligand is not always bound, especially with resveratrol (Table 2B). These factors may obscure the full effects of the ligands on the MD metrics.

Importantly, silybin A and ECG that do not recouple do not alter the phosphorylation-dependent metrics of G159D significantly, confirming the specific effect of the recouplers. It is notable that silybin B, EGCG and resveratrol affect the parameters for unphosphorylated G159D with little effect on phosphorylated G159D in parallel with the i*n vitro* motility measurements that also show that G159D behaves like phosphorylated wild-type independent of phosphorylation and the small molecules primarily alter the properties of unphosphorylated troponin (Figure 1).

3.3   Ligand binding and recoupling

Understanding how the recouplers function cannot be precise because of the disordered nature of troponin in the regions where the ligands contact and the weak, stochastic binding of the ligands. Nevertheless, the ligands do satisfy the basic criterion of allosteric effectors since they bind differently to unphosphorylated and phosphorylated G159D troponin. There is no significant change in binding energetics of the ligands to unphosphorylated and phosphorylated troponin, instead we observe that ligands are concentrated on a few 'hotspots' in unphosphorylated troponin, that are not occupied in phosphorylated troponin and that the 'hotspots' are more dispersed in phosohorylated troponin. We have concentrated our analysis on the effects of the ligands on unphosphorylated G159D troponin. The most common poses of bound silybin B and EGCG are found to be intercalated between the disordered N terminus of cTnI, adjacent to the phosphorylatable serines 22 and 23, and the helix A and B region of TnC. Since these poses are only seen in unphosphorylated troponin it is feasible that their occupancy is the cause of recoupling. Compatible with this hypothesis is the observation that silybin A binds to roughly the same region of NcTnI but is not intercalated between TnI and TnC,

presumably because the stereoisomer does not fit the space. Similarly, ECG binds predominanatly to the surface of cTnC at a site not occupied by EGCG.

Our study included resveratrol since it was functionally identical to silybin B in single filament and myocyte assays but had a radically different chemical structure. In 45% of the MD runs resveratrol is unbound (Table 2B) and its contacts with unphosphorylated troponin are also very different from silybin B and EGCG. In unphosphorylated troponin it appears that resveratrol is mostly located in the C terminal domain of cTnC near the hinge interface, possibly intercalated between the two domains. As with the other recouplers, binding is reduced and dispersed on phosphorylation. Therefore, we cannot propose a single mechanism for recoupling; a study of a wider range of recouplers would be needed to attempt to resolve the question.

Figure 6

Comparison of the effect of phosphorylation of cardiac troponin on its biochemical, physiological and molecular dynamics parameters, its suppression by the TnC G159D DCM related mutation and its restoration by small molecules.

Parameters compared are lusitropy in intact myocytes (from Fig 3), Fixed [Ca2+] screen for coupling in thin filaments measured by IVMA from Sheehan et al., [22] and effect of phosphorylation on EC_{50}, $(EC_{50}P/EC_{50}uP)$ in thin filaments measured by IVMA (from Figure 1, Supp Table1), Change on A/B angle on phosphorylation and change in hinge angle upon phosphorylation from Molecular dynamics simulations (from Figure 4, Tables 1 and 2). The recouplers are highlighted in blue.

3.4   CONCLUSIONS

A primary aim of this study was to demonstrate that the uncoupling phenomenon caused by cardiomyopathy-related mutations and the ability of small molecules to restore coupling, first found with pure thin filaments *in vitro*, is also relevant at the cellular and atomistic levels. Figure 6 summarises our results that support this proposition. Uncoupling is manifested at the myocyte level as an insensitivity of relaxation rate to β adrenergic stimulation and at the atomistic level as the failure of phosphorylation to increase interdomain angle or decrease A/B angle. It is also evident that silybin B, resveratrol and EGCG restore the native parameters to the mutant troponin whilst silybin A and ECG do not, as predicted.

We have shown Silybin B plays a significant role in safeguarding the heart against cardiomyopathy and Silybin A does not compromise this activity [22]. Racemic Silybin is a readily available and safe nutraceutical, as is resveratrol [32]. Further research on these compounds as treatments to restore lusitropy is thus indicated.

The compounds studied here are all polyphenols that may have multiple pharmacological activities that render them unsuitable as specific treatments [33-35]. However, at least 25 small molecules have been shown to be recouplers. Further work on these *in silico* could define the specific properties needed for recoupling and allow for the prediction and design of new molecules that recouple without additional activities that could be used in the treatment of genetic cardiomyopathies.

METHODS

 4.1   The movement of synthetic thin filaments containing G159D troponinC over immobilised myosin was measured by *in vitro* motility assay as described by [15].

4.2   Mouse cardiomyocyte isolation protocol and measurement of cell shortening was based on previously described methods using the Ionoptix system (Ionoptix, Milton, MA). Cell contraction was measured using a charge-coupled device video camera (MyoCam-S, IonOptix) connected to a personal computer running IonWizard software [13]. Guinea pig myocyte contractility was measured in the Cytocypher instrument as described by [36] using transfected guinea-pig myocytes prepared as described by [16]. Incubation medium was Krebs-Hensleit buffer with added 1mM CaCl$_2$. Reagents were obtained from Sigma-Aldrich, except for Silybin A and Silybin B which were prepared by Prof Vladimir Kren and Dr David Biedermann [37].

4.3   Molecular Dynamics studies

4.3.1 MD simulations of troponin were performed as described by Yang et al [24]. Parameterisation of the ligands in this study were based on a general AMBER force field (GAFF) which is suitable for small organic molecules [38]. Ligand partial charges were derived from the ground state structures which were found by following conformational search, geometry optimisation, and thermal correction procedures, as detailed below: An ensemble of conformers were generated for each ligand using the ETKDG algorithm as implemented by RDKit. [39]

Each conformer was optimised to a local minima. Each optimised structure was subjected to thermal correction with frequency to calculate the Gibbs free energy. [40-42] . The structure with the lowest ground state energy was picked as the ground state structure

From the ground state structures, restrained electrostatic potential (RESP) calculations were used to obtain the charge distribution on the molecule [43].

Gaussian16 [44] was used for both geometry optimisation and RESP calculation. Geometry optimisation was done with B3LYP functional with cc-pVDZ basis set and the polarisable continuum model (PCM) in water as implicit solvent [45-48].   RESP

calculation was done with Hartree-Fock method and 6-31G* basis set in vacuum. [49]. AmberTools was then used to generate the charge parameters for each atom.

4.3.2   Analysis   Representative structures, binding energy estimation by MMBPSA, interhelical angle distribution and hinge angle analysis were performed as described by [24]. CPPTraj was used to determine the contact/interactions. Pytraj [50], a Python package binding to cppraj program [51], was used for the distance measurements in this work. The ligand was deemed to be in contact with a residue when the minimal distance between any atom of the ligand and a residue was lower than 2.5 Å. The same was applied to ligand atom interactions where a ligand atom was deemed to be in contact with the protein when the minimal distance between atoms of the ligand and any part of the protein was lower than 2.5 Å.

ACKNOWLEDGEMENT

This study was supported by British Heart Foundation programme grant (RG/11/20/29266).

AS has a Ph.D. studentship funded by the BHF Centre of ResearchExcellence (RE/13/4/30184).

ZY is supported by an awarded Ph.D. studentship from the Wellcome Trust grant number 108908/Z/15/Z A 4 year PhD in theoretical systems biology and bioinformatics.

VK is supported by the Czech Science Foundation, grant number GA21-01799S

CR and AS are supported by British Heart Foundation Grant PG/18/68/33883.

REFERENCES

[1] W.J. Dong, J.J. Jayasundar, J. An, J. Xing, H.C. Cheung, Effects of PKA phosphorylation of cardiac troponin I and strong crossbridge on conformational transitions of the N-domain of cardiac troponin C in regulated thin filaments, Biochemistry 46 (2007) 9752-61.

[2] S.P. Robertson, J.D. Johnson, M.J. Holroyde, E.G. Kranias, J.D. Potter, R.J. Solaro, The effect of troponin I phosphorylation on the Ca2+-binding properties of the Ca2+-regulatory site of bovine cardiac troponin, J Biol Chem 257 (1982) 260-3.

[3] O.K. Baryshnikova, M.X. Li, B.D. Sykes, Modulation of cardiac troponin C function by the cardiac-specific N-terminus of troponin I: influence of PKA phosphorylation and involvement in cardiomyopathies., J Mol Biol 375(3) (2008) 735-751.

[4] S.C. Little, B.J. Biesiadecki, A. Kilic, R.S.D. Higgins, P.M.L. Janssen, J.P. Davis, The rates of Ca2+ dissociation and cross-bridge detachment from ventricular myofibrils as reported by a fluorescent cardiac Troponin C., J Biol Chem 287 (2012) 27930-27940.

[5] S.B. Marston, How do mutations in contractile proteins cause the primary familial cardiomyopathies?, J Cardiovasc Transl Res 4(3) (2011) 245-55.

[6] J.A. Spudich, Hypertrophic and Dilated Cardiomyopathy: Four Decades of Basic Research on Muscle Lead to Potential Therapeutic Approaches to These Devastating Genetic Diseases, Biophys J 106(6) (2014) 1236-1249.

[7] A. Messer, S. Marston, Investigating the role of uncoupling of Troponin I phosphorylation from changes in myofibrillar Ca2+-sensitivity in the pathogenesis of Cardiomyopathy, Front Physiol 5 (2014) 315.

[8] E. Dyer, A. Jacques, A. Hoskins, D. Ward, C. Gallon, A. Messer, J. Kaski, M. Burch, J. Kentish, S. Marston, Functional Analysis of a Unique Troponin C Mutation, Gly159Asp that Causes Familial Dilated Cardiomyopathy, Studied in Explanted Heart Muscle, Circ Heart Fail 2 (2009) 456-464.

[9] B.J. Biesiadecki, T. Kobayashi, J.S. Walker, R. John Solaro, P.P. de Tombe, The troponin C G159D mutation blunts myofilament desensitization induced by troponin I Ser23/24 phosphorylation, Circ Res 100(10) (2007) 1486-93.

[10] W. Dong, J. Xing, Y. Ouyang, J. An, H.C. Cheung, Structural kinetics of Cardiac troponin C mutants linked to familial hypertrophic and dilated cardiomyopathy in troponin complexes, J Biol Chem 283(6) (2008) 3424-3432.

[11] J.P. Kaski, M. Burch, P.M. Elliott, Mutations in the cardiac Troponin C gene are a cause of idiopathic dilated cardiomyopathy in childhood, Cardiol Young 17(6) (2007) 675-7.

[12] W. Song, E. Dyer, D. Stuckey, O. Copeland, M. Leung, C. Bayliss, A.E. Messer, R. Wilkinson, J. Tremoleda, M. Schneider, S. Harding, C. Redwood, K. Clarke, K. Nowak, L. Monserrat, D. Wells, S. Marston, Molecular mechanism of the Glu99lys mutation in cardiac actin (ACTC gene) that causes apical hypertrophy in man and mouse, J Biol Chem 286(31) (2011) 27582-27593.

[13] C. Rowlands, T. Owen, S. Lawal, S. Cao, S. Pandey, H.Y. Yang, W. Song, R. Wilkinson, A. Alvarez-Laviada, K. Gehmlich, S. Marston, K.T. MacLeod, Age and strain related aberrant Ca(2+) release is associated with sudden cardiac death in the ACTC E99K mouse model of hypertrophic cardiomyopathy, American journal of physiology. Heart and circulatory physiology 313 (2017) H1213-H1226.

[14] J.G.W. Smith, T. Owen, J.R. Bhagwan, D. Mosqueira, E. Scott, I. Mannhardt, A. Patel, R. Barriales-Villa, L. Monserrat, A. Hansen, T. Eschenhagen, S.E. Harding, S. Marston, C. Denning, Isogenic Pairs of hiPSC-CMs with Hypertrophic Cardiomyopathy/LVNC-Associated

ACTC1 E99K Mutation Unveil Differential Functional Deficits, Stem Cell Reports 11 (2018) 1226-1242.

[15] A.E. Messer, C.R. Bayliss, M. El-Mezgueldi, C.S. Redwood, D.G. Ward, M. Papadaki, C.D. Remedios, S.B. Marston, Mutations in troponin T associated with Hypertrophic Cardiomyopathy increase Ca2+-sensitivity and suppress the modulation of Ca2+-sensitivity by troponin I phosphorylation, Arch biochem biophys 601 (2016) 113 - 120.

[16] P. Robinson, X. Liu, A. Sparrow, S. Patel, Y.H. Zhang, B. Casadei, H. Watkins, C. Redwood, Hypertrophic cardiomyopathy mutations increase myofilament Ca2 buffering, alter intracellular Ca2 handling, and stimulate Ca2-dependent signaling, J Biol Chem 293(27) (2018) 10487-10499.

[17] F. Baudenbacher, T. Schober, J.R. Pinto, V.Y. Sidorov, F. Hilliard, R.J. Solaro, J.D. Potter, B.C. Knollmann, Myofilament Ca2+ sensitization causes susceptibility to cardiac arrhythmia in mice, J Clin Invest 118(12) (2008) 3893-903.

[18] R. Wilkinson, W. Song, N. Smoktunowicz, S. Marston, A dilated cardiomyopathy mutation blunts adrenergic response and induces contractile dysfunction under chronic angiotensin II stress, Am J Physiol H&C 309(11) (2015) H1936 - 46.

[19] S. Marston, J.R. Pinto, Suppression of lusitropy as a disease mechanism in cardiomyopathies, Front Cardiovasc Med 9 (2023) 1080965.

[20] N. Tadano, C. Du, F. Yumoto, S. Morimoto, M. Ohta, M. Xie, K. Nagata, D. Zhan, Q. Lu, Y. Miwa, F. Takahashi-Yanaga, M. Tanokura, I. Ohtsuki, T. Sasaguri, Biological actions of green tea catechins on cardiac troponin C, Brit J Pharmacol 161(5) (2010) 1034-1043.

[21] M. Papadaki, P.G. Vikhorev, S.B. Marston, A.E. Messer, Uncoupling of myofilament Ca2+ sensitivity from troponin I phosphorylation by mutations can be reversed by epigallocatechin-3-gallate, Cardiovasc Res 108(1) (2015) 99 - 110.

[22] A. Sheehan, A.E. Messer, M. Papadaki, A. Choudhry, V. Kren, D. Biedermann, B. Blagg, A. Khandelwal, S.B. Marston, Molecular Defects in Cardiac Myofilament Ca2+-Regulation Due to Cardiomyopathy-Linked Mutations Can Be Reversed by Small Molecules Binding to Troponin, Front Physiol 9(243) (2018) 300-12.

[23] J.E. Zamora, M. Papadaki, A.E. Messer, S.B. Marston, I.R. Gould, Troponin structure: its modulation by Ca(2+) and phosphorylation studied by molecular dynamics simulations., Phys Chem Chem Phys 18(30) (2016) 20691-20707.

[24] Z. Yang, S.B. Marston, I.R. Gould, Modulation of Structure and Dynamics of Cardiac Troponin by Phosphorylation and Mutations Revealed by Molecular Dynamics Simulations, J Phys Chem B 127(41) (2023) 8736-8748.

[25] Y. Yamada, K. Namba, T. Fujii, Cardiac muscle thin filament structures reveal calcium regulatory mechanism, Nature Comm 11(1) (2020) 153.

[26] C.M. Risi, B. Belknap, J. Atherton, I.L. Coscarella, H.D. White, P.B. Chase, J.R. Pinto, V.E. Galkin, Troponin Structural Dynamics in the Native Cardiac Thin Filament Revealed by Cryo Electron Microscopy, J Mol Biol 436(6) (2024) 168498.

[27] J.C. Kentish, D.T. McCloskey, J. Layland, S. Palmer, J.M. Leiden, A.F. Martin, R.J. Solaro, Phosphorylation of troponin I by protein kinase A accelerates relaxation and crossbridge cycle kinetics in mouse ventricular muscle, Circ Res 88(10) (2001) 1059-65.

[28] J. Layland, R.J. Solaro, A.M. Shah, Regulation of cardiac contractile function by troponin I phosphorylation, Cardiovasc Res 66(1) (2005) 12-21.

[29] P.H. Waddingham, S. Bhattacharyya, J.V. Zalen, G. Lloyd, Contractile reserve as a predictor of prognosis in patients with non-ischaemic systolic heart failure and dilated cardiomyopathy: a systematic review and meta-analysis, Echo Res Pract 5(1) (2018) 1-9.

[30] J.E. Zamora, PhD Thesis "Investigating cardiomyopathies with atomistic simulations of cardiac Troponin", Institute of Chemical Biology, Imperial College London, London, 2019.

[31] S. Marston, J.E. Zamora, Troponin structure and function: a view of recent progress, Journal of muscle research and cell motility 41(1) (2020) 71-89.

[32] V. Křen, K. Valentová, V. Kren, Silybin and its congeners: from traditional medicine to molecular effects, Nat Prod Rep 39(6) (2022) 1264-1281.

[33] H.I. Ingólfsson, P. Thakur, K.F. Herold, E.A. Hobart, N.B. Ramsey, X. Periole, D.H. de Jong, M. Zwama, D. Yilmaz, K. Hall, T. Maretzky, H.C. Hemmings, C. Blobel, S.J. Marrink, A. Koçer, J.T. Sack, O.S. Andersen, Phytochemicals perturb membranes and promiscuously alter protein function., ACS Chem Biol 9(8) (2014) 1788-1798.

[34] V. Křen, K. Valentová, Silybin and its congeners: from traditional medicine to molecular effects, Nat Prod Rep 39(6) (2022) 1264-1281.

[35] J. Baell, M.A. Walters, Chemistry: Chemical con artists foil drug discovery., Nature 513(7519) (2014) 481-483.

[36] P.T. Wright, S.F. Tsui, A.J. Francis, K.T. MacLeod, S.B. Marston, Approaches to High-Throughput Analysis of Cardiomyocyte Contractility, Frontiers in Physiology 11 (2020) 612-612.

[37] D. Monti, R. Gažák, P. Marhol, D. Biedermann, K.i. Purchartová, M. Fedrigo, S. Riva, V. Křen, Enzymatic Kinetic Resolution of Silybin Diastereoisomers, J Nat Prod 73(4) (2010) 613-619.

[38] J. Wang, R. Wolf, J. Caldwell, P. Kollman, D.A. Case, Development and testing of a general amber force field, Journal of computational chemistry 25(9) (2004) 1157-74.

[39] S. Riniker, G.A. Landrum, Better Informed Distance Geometry: Using What We Know To Improve Conformation Generation, JCTC 55(12) (2015) 2562-2574.

[40] G. Luchini, J.V. Alegre-Requena, I. Funes-Ardoiz, R.S. Paton, GoodVibes: automated thermochemistry for heterogeneous computational chemistry data, F1000research 9(Chem Inf Sci) (2020) 291.

[41] Y.-P. Li, J. Gomes, S.M. Sharada, A.T. Bell, M. Head-Gordon, Improved Force-Field Parameters for QM/MM Simulations of the Energies of Adsorption for Molecules in Zeolites and a Free Rotor Correction to the Rigid Rotor Harmonic Oscillator Model for Adsorption Enthalpies, J Phys Chem C 119(4) (2015) 1840-1850.

[42] S. Grimme, Supramolecular Binding Thermodynamics by Dispersion-Corrected Density Functional Theory, Chem European J 18(32) (2012) 9955-9964.

[43] C.I. Bayly, P. Cieplak, W. Cornell, P.A. Kollman, A well-behaved electrostatic potential based method using charge restraints for deriving atomic charges: the RESP model, J Phys Chem 97(40) (1993) 10269-10280.

[44] M.J. Frisch, e. al., Gaussian~16 Revision C.01, (2016).

[45] A.D. Becke, Density-functional thermochemistry. III. The role of exact exchange, J Chem Phys 98(7) (1993) 5648-5652.

[46] C. Lee, W. Yang, R.G. Parr, Development of the Colle-Salvetti correlation-energy formula into a functional of the electron density, Phys Rev B 37(2) (1988) 785-789.

[47] T.H. Dunning, Gaussian basis sets for use in correlated molecular calculations. I. The atoms boron through neon and hydrogen, J Chem Phys 90(2) (1989) 1007-1023.

[48] S. Miertuš, E. Scrocco, J. Tomasi, Electrostatic interaction of a solute with a continuum. A direct utilizaion of AB initio molecular potentials for the prevision of solvent effects, Chem Phys 55(1) (1981) 117-129.

[49] R. Ditchfield, W.J. Hehre, J.A. Pople, Self-Consistent Molecular-Orbital Methods. IX. An

Extended Gaussian-Type Basis for Molecular-Orbital Studies of Organic Molecules, J Chem

Phys 54(2) (1971) 724-728.

[50] N. H, R. DR, S. J, D.A. Case, PYTRAJ v1.0.0.dev1: Interactive data analysis for molecular
dynamics simulations, (2016).

[51] D.R. Roe, T.E. Cheatham, 3rd, PTRAJ and CPPTRAJ: Software for Processing and Analysis
of Molecular Dynamics Trajectory Data, J Chem Theory Comput 9(7) (2013) 3084-95.