Title: **A conserved chronobiological complex times *C. elegans* development**

Rebecca K. Spangler[1]*, Kathrin Braun[2]*, Guinevere E. Ashley[3]* Marit van der Does[2,4],

Daniel Wruck[1], Andrea Ramos Coronado[1], James Matthew Ragle[3], Vytautas

Iesmantavicius[2], Lucas J. Morales Moya[2], Keya Jonnalagadda[3], Carrie L. Partch[1,5,6],

Helge Großhans[2,4‡], Jordan D. Ward[3‡]

[1]Department of Chemistry and Biochemistry, University of California-Santa Cruz, Santa

Cruz, CA 95064, USA.

[2]Friedrich Miescher Institute for Biomedical Research, 4056 Basel, Switzerland

[3]Department of Molecular, Cell, and Developmental Biology, University of California-

Santa Cruz, Santa Cruz, CA 95064, USA.

[4]University of Basel, 4002 Basel, Switzerland

[5]Center for Circadian Biology, University of California-San Diego, La Jolla, CA 92093,

USA

[6]Howard Hughes Medical Institute, University of California-Santa Cruz, Santa Cruz

95064, USA

‡Author for correspondence.

*These authors contributed equally to this work

Abstract

The mammalian PAS-domain protein PERIOD (PER) and its *C. elegans* orthologue LIN-42 have been proposed to constitute an evolutionary link between two distinct, circadian and developmental, timing systems. However, while the function of PER in animal circadian rhythms is well understood molecularly and mechanistically, this is not true for LIN-42's function in timing rhythmic development. Here, using targeted deletions, we find that the LIN-42 PAS domains are dispensable for the protein's function in timing molts. Instead, we observe arrhythmic molts upon deletion of a distinct sequence element, conserved with PER. We show that this element, designated CK1δ-binding domain (CK1BD), mediates stable binding to KIN-20, the *C. elegans* CK1δ/ε orthologue. We demonstrate that CK1δ phosphorylates LIN-42 and define two conserved helical motifs in the CK1BD, CK1BD-A and CK1BD-B, that have distinct roles in controlling CK1δ-binding and kinase activity *in vitro*. KIN-20 and the LIN-42 CK1BD are required for proper molting timing *in vivo*, and loss of LIN-42 binding changes KIN-20 subcellular localization. The interactions mirror the central role of a stable circadian PER–CK1 complex in setting a robust ~24-hour period. Hence, our results establish LIN-42/PER – KIN-20/CK1δ/ε as a functionally conserved signaling module of two distinct chronobiological systems.

Introduction

Chronobiology is the study of biological rhythms. Circadian rhythms which enable organisms to anticipate daily cycles of light, temperature, and other environmental variables are one of the best studied examples. [1] PERIOD (PER) proteins are central to this timekeeping mechanism. Their cellular abundance, stability, and nuclear localization changes rhythmically over a ~24-hour cycle. In mammals, the paralogs PERIOD1 and PERIOD2 (PER1 and PER2) associate with Casein kinase 1 δ/ε (CK1δ/ε) as well as the CRYPTOCHROME proteins, CRY1 and CRY2, to form nuclear transcriptional complexes which repress the transcriptional activators circadian locomotor output cycles protein kaput (CLOCK) and brain and muscle ARNT-like 1 (BMAL1)(Fig. S1A).[2–5] Since the targets of the CLOCK-BMAL1 heterodimer include PER and CRY themselves, PER and CRY thus eventually repress their own production, permitting subsequent re-expression of CLOCK-BMAL1 to start a new cycle.[2,6]

To maintain the 24-hour period of circadian rhythms, a delay between the activation and

the repression of CLOCK-BMAL1 transcription is required. Central to this delay is the

post-translational modification of PER by one of the two closely related CK1δ and CK1ε

kinases. For convenience and because of the redundancy in their function, we will

henceforth refer to the two isoforms generically as CK1. Anchored through a CK1-Binding

Domain (CK1BD)[7], CK1 is associated with PER for its entire existence in the cell, even

translocating from the cytoplasm to the nucleus together (Fig. S1A)[4,8], allowing PER to

deliver CK1 to other targets at clock-controlled promoters.[8,9] CK1-mediated

phosphorylation of a PER degron licenses PER ubiquitylation and subsequent

degradation, thereby limiting PER abundance and its period of activity (Fig. S1B).[10–12] In

addition to the degron, CK1 phosphorylates multiple serines in another PER sequence

element known as FASP ("Familial Advanced Sleep Phase")(Fig. S1B).[13,14]

Phosphorylation of the FASP site turns PER into an inhibitor of CK1 kinase activity.[15,16]

In humans, a single residue mutation in this region that blocks all FASP phosphorylation

results in a short circadian period that manifests as the eponymous Familial Advanced

Sleep Phase, where affected individuals wake up very early in the morning.[17,18]

In contrast to circadian timing, the chronobiology of development is mechanistically less

well understood. In *C. elegans*, a single PER orthologue, LIN-42, appears to diverge

substantially in structure and function from the mammalian and fly PER proteins. At 598

amino acids for its longest isoform, LIN-42 is substantially shorter than mouse PER2 at

1225 amino acids.[19] LIN-42 lacks a CRY-binding domain, consistent with a lack of a CRY

orthologue in *C. elegans*, and in its tandem PER-ARNT-SIM (PAS) domains, only PAS-B

is well conserved, adopts a canonical fold, and mediates dimerization in a mode identical

to mammalian PER.[20,21] Finally, two stretches of sequence, previously termed SYQ and

LT according to their first amino acids[22], bear sequence homology to the two PER CK1BD

subdomains, CK1BD-A and CK1BD-B, but their function has remained unexplored. *lin-42*

further differs from canonical PERs in its expression dynamics and functions: rather than

exhibiting a ~24-hour, temperature-invariant period, *lin-42* expression cycles exhibit a

temperature-dependent length which can be as short as ~7 hours at 25˚C.[19,23] Indeed,

lin-42 was identified as a developmental gene whose mutation causes heterochronic

phenotypes, i.e., defects in temporal cell fate specification where cells adopt adult cell

fates precociously, in larvae.[24,25] Hence, it was proposed that *lin-42/PER* constitute an

evolutionary link between two distinct, circadian and developmental, timing systems.[19]

The extent of functional similarity between these timing systems generally, and LIN-42

and PER function specifically, has remained uncertain, despite the realization that

additional orthologues of mammalian clock genes exist and cause heterochronic

phenotypes when mutated.[19,21,23,26–28] Conceptually, whereas mutations of circadian

clock genes such as *PER* change the tempo and/or robustness of circadian rhythms,

heterochronic mutations are defined by their ability to alter the sequence of

developmental events, such that certain events are skipped.[29,30] Accordingly, and despite

its rhythmic expression, the heterochronic function of LIN-42 does not appear to involve

recurring activity but rather the stage-specific repression of the *let-7* miRNA prior to the

third larval stage.[31–33]

LIN-42 is also required for the rhythmic occurrence of molts. Wild-type animals molt, i.e.,

regenerate a new collagenous apical extracellular matrix (cuticle), at the end of each of

the four larval stages.[34] Under constant environmental conditions, individually grown

animals enter and exit molts with great temporal uniformity, revealing robust temporal

control. *lin-42(ok2385)* mutant animals were shown to cause a slow-down of development

as well as an arrhythmic molting phenotype where this uniformity is lost such that

individual animals molt at different times.[35,36] It is unknown how LIN-42 contributes to

rhythmic molting mechanistically. In addition to *CRY*, the *C. elegans* genome lacks

obvious orthologues of *BMAL1* and *CLOCK*, arguing for a mechanism that differs from

that of the circadian clock at least in the identity of several core components.[27] In a yeast

two-hybrid assay, LIN-42 was shown to be capable of binding to numerous transcription

factors.[37] Although the functional relevance of such binding for molting has remained

unexplored, LIN-42 binding to the REV-ERB orthologue NHR-85 appears important for

robust periodic transcription of the heterochronic *lin-4* miRNA. Finally, loss of KIN-20, the

C. elegans orthologue of CK1δ/ε, replicates some heterochronic phenotypes of *lin-42*

mutation and slows development[26], but its heterochronic functions were argued not to
involve LIN-42[38], and it is unknown whether KIN-20 is required for rhythmic molting.

Here, we set out to further characterize the molecular and developmental functions of
LIN-42. Using targeted mutations, we find that its PAS domains are largely dispensable
both for heterochronic pathway activity and rhythmic molting. By contrast, the SYQ/LT
regions are specifically required for rhythmic molting. We demonstrate that these domains
function as a CK1BD, mediating stable KIN-20/CK1 binding to LIN-42 *in vivo* and *in vitro*.
CK1 phosphorylates LIN-42 *in vitro* and this activity is distinctly modulated through the
two CK1BD subdomains: CK1BD-B is particularly important for stable CK1 binding,
whereas the CK1BD-A element inhibits CK1 enzymatic activity. Moreover, loss of LIN-42
binding impairs nuclear localization of KIN-20. Our results identify LIN-42/PER–KIN-
20/CK1 as a conserved chronobiological complex utilized by two distinct biological
oscillators. Although CK1 activity in the circadian clock has previously been viewed mostly
through the lens of its effects on PER, our findings align well with the growing notion that
PER-mediated regulation of CK1 may be an additional important mechanism to support
robust circadian rhythms.

Results

The LIN-42 CK1BD mediates a subset of developmental functions of LIN-42

Previous analysis of the two partial deletion alleles of *lin-42*, *n1089* and *ok2385* (Fig. 1A),
have suggested the possibility that different developmental functions could be genetically
separable. Although both deletions caused precocious heterochronic phenotypes, only
ok2385 showed evidence of slow and arrhythmic molting.[22,35,39] To confirm these findings,
we quantified molt, intermolt, and larval stage duration for mutant and wild-type animals
at high temporal resolution, using a luciferase assay that monitors entry and exit from the
lethargus state associated with molting on animals grown in isolation.[36,40] Indeed, we
found that *lin-42(ok2385)* mutant larvae developed more slowly than wild-type animals
and became increasingly arrhythmic (Fig. 1B, S2A-C). Moreover, and as observed
previously[36], most mutant animals underwent only three molts (33/39 animals) within the
duration of this assay, suggesting either a precocious exit from the molting cycle, or a

very delayed or abnormal fourth molt (Fig. 1B, S2D). By contrast, essentially all *lin-*

42(n1089) animals completed a normal number of molts (42/42 animals) and maintained

robust synchrony, although they developed significantly slower than wild-type animals

(Fig. 1B). At the same time, both alleles caused robust heterochronic phenotypes,

illustrated by precocious formation of alae (Fig. 1C) in L3 to early L4 larvae. Alae are a

cuticular structure normally secreted by the terminally differentiated epidermal seam cells

at the L4-adult transition.

Figure 1. *lin-42ΔPAS* mutations cause mild heterochronic phenotypes while *lin-42ΔCK1BD* mutations cause asynchronous molting (A) Schematic depiction of the *lin-42* genomic locus. The gene structure of the *lin-42 a, b,* and *c* isoforms are shown. The location of sequences encoding putative PAS-A, PAS-B, SYQ, and LT sequence motifs are indicated. Published *lin-42* deletion alleles *n1089* and *ok2385* are indicated in dark and light grey, respectively, newly generated targeted *ΔPAS* and *ΔCK1BD* deletions in navy and teal, respectively. **(B)** Heatmaps showing trend-corrected luminescence traces from the indicated genotype. Each horizontal line represents one animal. Traces are sorted by entry into the first molt. Darker color indicates low luminescence signal and corresponds to the molts. **(C)** Bar plot quantifying the percentage of animals of the indicated genotype with precocious, complete, or partial alae at the L3-L4 molt. **(D)** Bar plot quantifying the percentage of animals of the indicated genotype that exhibited egg-laying defects as determined by the presence of hatched larvae in the animal. **(E)** Bar plot depicting the average number of live progeny from hermaphrodites of the indicated genotype. **(F)** Bar plot quantifying the percentage of animals of the indicated genotype that arrested or died as larvae. (C-F) Statistical significance was determined using an ordinary one-way ANOVA. p<0.05 was considered statistically significant. **** indicates < 0.0001. Error bars in C-F represent standard deviation.

The complex nature of the alleles, each with at least one breakpoint in a noncoding region,

precludes a straightforward attribution of the phenotypes to specific features of the LIN-

42 protein. Therefore, we decided to generate precise deletions of the two conserved

regions, PAS and SYQ/LT (Fig. 1A). Henceforth, and based on the functional

characterization which we describe below, we will refer to SYQ/LT as CK1BD for CK1-

Binding Domain. To our surprise, *lin-42(wrd67[ΔPAS])* mutant animals exhibited

essentially wild-type molting patterns in the luciferase assay, resembling neither *lin-*

42(n1089) nor *lin-42(ok2385)* mutant animals. By contrast, *lin-42(wrd63[ΔCK1BD])*

mutant animals exhibited increasing loss of molting synchrony over time (Fig. 1B).

However, unlike *lin-42(ok2385)*, *lin-42(ΔCK1BD)* mutant animals executed four

detectable molts during the assay (Fig. 1B, S2D).

To get a better understanding of the developmental relevance of each domain, we

examined additional phenotypes (Fig. 1D-F; Table S1). We found that all four alleles

affected brood sizes, albeit to varying degrees, in the order

ok2385>>ΔCK1BD≈n1089>ΔPAS of decreasing severity (Fig. 1E, Table S1). Only

ok2385 and *n1089* caused highly penetrant egg-laying (Egl) and precocious alae

defects[22,35,39], whereas ΔPAS and ΔCK1BD mutants had completely wild-type egg laying

and only modest precocious alae (Fig. 1C and D, Table S1). Finally, both *ok2385* and, to

a lesser extent, ΔCK1BD caused larval arrest and lethality, a phenotype interestingly not

observed in the luciferase assay, suggesting an environmental dependency (Fig. 1F,

Table S1). Taken together, these results indicate surprisingly mild phenotypes upon loss

of the LIN-42 PAS region and highlight an important function of the CK1BD for rhythmic

molting.

LIN-42 and KIN-20 interact *in vivo*

Since the genetic results suggested separable functions of LIN-42, we hypothesized that

these could depend on specific interaction partners. To identify interaction partners we

performed anti-FLAG immunoprecipitations on endogenously tagged *lin-*

42(xe321[3xFLAG::lin-42b+c]) mixed-stage animal lysates followed by mass-

spectrometry. As a control for non-specific binding, we also performed an anti-FLAG

immunoprecipitation in an unrelated strain (*sart-3::GFP::3xFLAG*).[41] We observed a

single highly enriched interaction partner for LIN-42: KIN-20 – the orthologue of

mammalian CK1δ/ε (Fig. 2A).

Figure 2 LIN-42 and KIN-20 interact *in vivo* (A) Volcano blot comparing protein enrichments in 3xFLAG::LIN-42b+c and 3xFLAG::SART-3 (control) immunoprecipitations, determined by mass spectrometry. **(B)** mRNA expression profile for *lin-42* (dark blue) and *kin-20* (light blue) mRNAs throughout larval development. Data Source: Meeuse et al., 2020. **(C)** Western blot with extracts from *3xflag::lin-42b+c; 3xflag-ha::kin-20* (HW3479) larvae collected hourly for 11 hours, starting at 13 h after plating synchronized L1 stage animals on food at 25°C. Top part probed with anti-FLAG-HRP (1:1,000), lower part with anti-actin-1 (1:7,500). Arrows indicate bands for LIN-42b, KIN-20 and ACT-1 (see Fig. S5A). **(D)** Anti-HA pulldown from *3xflag::lin-42b+c; 3xflag-ha::kin-20* animals. Blot probed with anti-FLAG (1:1,000).

The mRNA levels of both *lin-42* and *kin-20* oscillate.[40,42] Further analysis of the published

sequencing data revealed that the peak phases of the two proteins differ, but that there

are periods where both transcripts accumulate (Fig. 2B). This is also true for the proteins:

we observed by Western blotting that the levels of endogenously tagged KIN-20 protein

were largely invariant over time, whereas those of endogenously tagged LIN-42b protein

levels changed rhythmically (Fig. 2C). Finally, we confirmed a physical interaction with a
reciprocal anti-HA pulldown from *lin-42(xe321[3xFLAG::lin-42b+c]); kin-*
20(xe328[3xFLAG-HA::kin-20]) animal lysates, again using endogenously tagged
proteins (Fig. 2D). Taken together, we conclude that LIN-42 and KIN-20 form a complex
in vivo.

The LIN-42 SYQ-LT regions form a functional CK1BD capable of binding
mammalian CK1

Mammalian CK1 binds PER through its kinase domain. This region is highly conserved,
with CK1δ and KIN-20 exhibiting 79% and 100% sequence identity within the kinase
domain and active sites, respectively (Fig. S3).[43,44] By contrast, the CK1BD is less well
conserved in LIN-42 with only ~30% sequence identity and a different spacing of the
relative LIN-42 CK1BD-A and -B sites (Fig. 3A). Hence, it was unclear whether this region
could function as a CK1BD in LIN-42.

We produced recombinant proteins to test this possibility. We were unable to express
soluble KIN-20 from a bacterial or insect system, so we used recombinant human CK1δ
kinase domain (hereafter referred to as CK1), similar to previous studies of KIN-20.[45] *In*
vitro pulldown assays with biotinylated LIN-42 protein containing the SYQ/CK1BD-A and
LT/CK1BD-B domains as well as C-terminal tail (residues 402-589) revealed that CK1
bound to wild-type LIN-42 protein. Deletion of either the SYQ/CK1BD-A or LT/CK1BD-B
motifs reduced this interaction (Fig. 3B).

To test whether the CK1BD is sufficient to bind CK1, we purified a truncated fragment of
LIN-42 lacking the C-terminal tail (CK1BD) and performed bio-layer interferometry (BLI)
assays with biotinylated CK1 and titrating LIN-42 fragments. This optical technique
measures macromolecular interactions in real-time, providing k_{on} and k_{off} rates from which
affinity constants (K_D) can be calculated. The smaller size of the CK1BD compared to the
CK1BD+Tail (residues 402-589; same construct used in Fig. 3B) resulted in a lower
response signal, however, both proteins bound to biotinylated CK1 with comparable,
nanomolar affinities (348 ± 123 nM for CK1BD and 209 ± 78 nM for CK1BD + Tail; Fig.

3C-E). These results demonstrate that the SYQ and LT motifs encompass a minimal

CK1BD.

Figure 3 The LIN-42 SYQ and LT domains constitute a functional CK1-binding domain (A) Schematic representing PER2 and LIN-42 protein domains. Protein constructs used in this study, CK1BD + Tail and CK1BD, are indicated. CK1BD, Casein Kinase 1-Binding Domain; CBD, CRY-binding domain. **(B)** Representative pulldown assay of human CK1 and biotinylated LIN-42 CK1BD + Tail proteins using the indicated protein variants. **(C)** Values for K_D from kinetic analysis of BLI data (D-H) based on a 2:1 heterogeneous ligand binding model and global analysis (Octet). Mean ± SEM, ordinary one-way ANOVA. p<0.05 was considered statistically significant. **** indicates p< 0.0001. **(D-H)** Bio-layer interferometry (BLI) data for indicated LIN-42 protein binding to immobilized, biotinylated CK1. Inset values represent the concentrations of LIN-42 for individual binding reactions. Model fit to association and dissociation over time is represented by black lines. Data shown from one representative experiment of n ≥ 3 assays.

We next performed BLI experiments to quantitatively explore the significance of the

CK1BD-A and -B motifs in binding to CK1. LIN-42-binding to CK1 was not detectable

when CK1BD-B was deleted, either singly (*CK1BD-ΔB*) or in combination with CK1BD-A

(*CK1BD-ΔA/B*) even at higher concentrations (Fig. 3C,F,G). In contrast, deletion of the

CK1BD-A domain reduced the affinity of LIN-42 for CK1 approximately 10-fold (K_D = ~3

µM)(Fig. 3C,H), necessitating higher concentrations of the CK1BD-ΔA +Tail protein to

determine an accurate K_D (Fig. 3H). These data indicate that CK1BD-A and B both

contribute to high-affinity binding of CK1 *in vitro*, while revealing a more important
contribution of the CK1BD-B motif, consistent with recent work that demonstrated an
essential role of the human PER2 CK1BD-B motif for stable association with CK1 in a
mouse model and mammalian cells.[46] Collectively, our data establish that the LIN-42
CK1BD is functionally conserved and mediates stable binding to the CK1 kinase domain.
249
Phosphorylation of the LIN-42 C-terminus by CK1 exhibits a conserved mode of
feedback inhibition Mammalian CK1 regulates PER abundance by controlling its
degradation post-translationally. Mutations on CK1 or its PER phosphorylation sites can
induce changes in circadian period as great as ~4-hours.[16,18,47,48] To test whether LIN-42
is a CK1 substrate, we performed *in vitro* ^{32}P-ATP kinase assays using the CK1BD alone
or the CK1BD+Tail as substrates. CK1BD protein contains 4 serine and 6 threonine
residues while the CK1BD + Tail construct contains an additional 22 serine and 15
threonine residues. We observed phosphorylation for both constructs, yet more robustly
on LIN-42 CK1BD+Tail than on CK1BD alone (Fig. 4A-C). Hence, CK1 appears capable
of phosphorylating both CK1BD and the extended C-terminal tail. Deletion of CK1BD-B
reduced phosphorylation relative to the wild-type protein, whereas phosphorylation of the
CK1BD-ΔA + Tail mutant protein remained largely unaltered (Fig. 4A-C). Thus, in the two
assays, loss of CK1BD-B impacts both binding and phosphorylation by CK1 more strongly
than loss of CK1BD-A (Fig. 3C, 4A-C).
264
To explore the kinetics of CK1-mediated phosphorylation of the LIN-42 extended C-
terminus, we performed substrate titration experiments using an ADP-Glo enzymatic
assay. Here, we observed a decrease in kinase activity with high levels of wild-type LIN-
42 CK1BD+Tail (Fig. 4D). This result mirrors the ability of phosphorylated PER as well as
the apoptosis substrate p63, to inhibit the activity of CK1 (DBT in Drosophila) in both
mammalian and *Drosophila* systems through product inhibition.[14]
271
Closer examination revealed that deletion of the LIN-42 CK1BD-A motif relieved feedback
inhibition (Fig. 4D), explaining why the ^{32}P-ATP data had shown little decrease in the
overall phosphorylation level (Fig. 4A-C) despite having reduced affinity for the kinase

(Fig. 3C). Conversely, deletion of the B motif either in isolation or with the A motif, caused

a significant reduction in overall phosphorylation of CK1BD+Tail. No serine and only 4

threonine residues are removed in this deletion, so this effect was likely not due to loss

of phosphosites. The catalytic efficiency (k_{cat}/K_m) of CK1 for each LIN-42 substrate

decreased significantly with deletion of the CK1BD motifs relative to wild-type, with activity

most severely impacted upon loss of the CK1BD-B motif (Fig. 4D,E). Taken together,

these results suggest that while both motifs are involved in binding to CK1, deletion of the

CK1BD-A region predominantly relieves feedback inhibition without fully disrupting the

anchoring interaction seen with the deletion of the CK1BD-B domain. Moreover, and

similar to the mammalian PER–CK1 complex[49], loss of the anchoring interaction

compromises, but does not fully abrogate phosphorylation.

Identification of potential LIN-42 phosphosites

To determine potential CK1-dependent LIN-42 phosphorylation sites, we performed *in*

vitro kinase reactions followed by phosphoenrichment and mass spectrometry to identify

phosphopeptides.There are a total of 26 serine and 21 threonine residues in the CK1BD

+ Tail construct. Of these, 12 serine and 3 threonine sites in the C-terminus were

phosphorylated upon incubation of LIN-42 CK1BD + Tail with CK1 at least once from

three replicates *in vitro* (Fig. 4F). Although we detected low levels of phosphorylation on

LIN-42 constructs lacking the tail in the [32]P-ATP assay (Fig. 4A), no serine and only 1

threonine phosphorylation within the CK1BD was detected *in vitro,* reflecting possible

limitations in the assays.

To test whether LIN-42 was phosphorylated *in vivo*, we immunoprecipitated

endogenously tagged 3xFLAG::LIN-42b+c at various time points from a population of

synchronized L2 stage larvae and subjected precipitates to mass spectrometry. We

identified 15 serine residues, 1 threonine residue, and 1 tyrosine residue that were

phosphorylated on LIN-42. All but four of these phosphosites (all serine residues) are

located on the CK1BD + Tail (Fig. 4F). Among these, 6 serine residues overlap with the

13 phosphoserines identified after the *in vitro* reaction while no threonine or tyrosine

residues were phosphorylated in both the *in vitro* and *in vivo* datasets (Fig. 4F). The

Figure 4 The extended C-terminus of LIN-42 is phosphorylated by CK1 (A) Representative ^{32}P-radiolabeled ATP enzymatic assay of CK1 activity on LIN-42 CK1BD + Tail and CK1BD wild-type and indicated domain deletion mutants. **(B)** Quantification of band intensity depicted in (A) and replicates. Model fit to Michaelis- Menten, n = 3. **(C)** Fold-change quantification of band intensity from panels A & B compared to CK1 autophosphorylation at 120 minutes. Ordinary one-way ANOVA. $p<0.05$ was considered statistically significant. ****$p < 0.0001$, n ≥ 3, mean ± SEM. (D) Representative ADP-Glo enzymatic assay comparing activity of CK1 on LIN-42 CK1BD + Tail wild-type (maroon), CK1BD-ΔA (orange), CK1BD-ΔB (yellow), and CK1BD-ΔA/B (green). **(E)** Values of k_{cat}/K_m calculated from (D). n=3, mean ± SEM, Ordinary one-way ANOVA. $p<0.05$ was considered statistically significant. * indicates $p<0.03$ and ** indicates $p<0.002$. **(F)** LIN-42 phosphorylation sites identified via mass spectrometry from *in vivo* samples from whole worm lysate collected hourly from synchronized L1 animals from 14 hrs (early L2) to 25 hrs (late L3) after plating at 25°C (top arrows) and *in vitro* kinase reactions of CK1 activity on LIN-42 CK1BD + Tail (bottom arrows). Serine, threonine, and tyrosine residues are indicated via vertical blue, purple, and pink ticks, respectively, along the CK1BD + Tail construct schematic.

overlap of *in vitro* and *in vivo* results is consistent with the possibility that LIN-42 is a CK1

substrate *in vivo*. Whether incomplete overlap reflects differences in kinase activities in

the activity of additional kinases and/or specificity factors *in vivo*, or technical differences

in the experimental approaches remains to be determined.

LIN-42 C-terminal tail deletion causes a heterochronic phenotype but leaves

molting timing largely unaffected

Given the extensive phosphorylation of the LIN-42 tail *in vitro* and *in vivo*, we sought to

explore its functional relevance. Unexpectedly, *lin-42(ΔTail)* mutant animals did not

recapitulate the *lin-42(ΔCK1DB)* arrhythmic molting phenotype but instead resembled *lin-*

42(n1089) animals phenotypically (Fig. 1C-F; Fig. 5). Specifically, *lin-42(ΔTail)* mutants

exhibited regular timing of the first three molts, yet an extended and irregular fourth molt

(Fig. 5A, S4A-C, S5). The severity of this extended 4[th] molt was variable and could reflect

unaccounted environmental differences during assay runs. The mutant animals also

exhibited a modest reduction in brood size, egg-laying defects and precocious alae but

no larval arrest (Fig. 5B-E). These data indicate that the LIN-42 C-terminus is largely

dispensable for timing of larval molting and suggest instead that it plays a previously

unrecognized role in the heterochronic functions of LIN-42.

***kin-20* null and catalytically dead mutants exhibit arrhythmic molting**

Deletion of a major phosphorylation target, the LIN-42 tail, did not recapitulate *lin-*

42(ΔCK1BD) mutant molting timing defects which prompted the question whether KIN-20

was at all required for rhythmic molting. To address this question, we monitored molt

timing for *kin-20(ok505)* null mutant animals (*kin-20(0)*). *kin-20* loss recapitulated both the

slow development and arrhythmic molt phenotypes of *lin-42(ok2835)* and *lin-42(ΔCK1DB)*

mutations (Fig. 1B; Fig. 5A, S4A-C). Like *lin-42(ok2835)* and *lin-42(ΔCK1DB)* mutations,

the *kin-20(0)* mutation also caused larval arrest when animals were grown on plate but

not in the liquid culture luciferase assay (Fig. 1F; Fig. 5E). Additionally, we found that *kin-*

20(0) mutant animals resembled *lin-42(ok2835)* mutants in their severe egg-laying

defects and brood size reduction, phenotypes which are weaker or even absent from *lin-*

42(ΔCK1DB) animals (Fig. 1D,E; Fig. 5B,C). Finally, like *lin-42(ΔCK1DB)* animals but

unlike *lin-42(ok2385)* mutants, *kin-20* null mutants did not develop precocious alae (Fig.

1C; Fig. 5D), consistent with earlier work.[26]

To assess the relevance of KIN-20's enzymatic activity for rhythmic molting, we

engineered a D310A point mutation into the endogenous *kin-20* locus to abrogate

catalytic activity. This mutation disrupts the HRD motif in the catalytic loop that

coordinates phosphorylated residues (Fig. S3).[50,51] The resulting *kin-20(xe355[D310A])*

animals recapitulated the *kin-20(0)* mutant arrhythmic molting phenotype, although KIN-

20 levels were not decreased (Fig. S6A-D). Indeed, we observed a trend towards

increased accumulation of the mutant KIN-20 protein, possibly indicating autoregulation

but not further pursued by us (Fig. 5A; S4A-C; S6B). We conclude that KIN-20 and

especially its enzymatic activity are required for rhythmic molting.

Figure 5 *kin-20* null and catalytically dead alleles exhibit larval arrest and asynchronous molting similar to *lin-42(ΔCK1BD)* mutants (A) Heatmaps showing trend-corrected luminescence traces from the indicated genotype. Each horizontal line represents one animal. Traces are sorted by entry into the first molt. Darker color indicates low luminescence signal and corresponds to the molts. **(B)** Bar plot depicting the average number of live progeny from hermaphrodites of the indicated genotype. **(C)** Bar plot quantifying the percentage of animals of the indicated genotype that exhibited egg-laying defects as determined by the presence of hatched larvae in the animal. **(D)** Bar plot quantifying the percentage of animals of the indicated genotype with precocious complete or partial alae at the L3-L4 molt. **(E)** Bar plot quantifying the percentage of animals of the indicated genotype that arrested or died as larvae. **(B-E)** Statistical significance was determined using ordinary one-way ANOVA. p<0.05 was considered statistically significant. ** indicates p<0.0001****, indicates p<0.0001. Error bars in B-E represent standard deviation.

KIN-20 exhibits dynamic changes in subcellular localization

Although KIN-20 kinase activity and its binding to the LIN-42 CK1BD are required for

rhythmic molting, LIN-42 tail phosphorylation appeared dispensable. Moreover, *kin-20*

deletion or inactivation appeared to cause more pronounced defects than deletion of the

LIN-42-CK1BD. Hence, we wondered whether the significance of the interaction between

the two proteins could lie in regulation of KIN-20 by LIN-42, rather than vice versa. We

therefore examined KIN-20 levels and localization. Using GFP::KIN-20 (with a split-GFP

system to enhance the signal; Methods), we observed both nuclear and cytoplasmic

signal in the epidermis. Strikingly, the relative distribution between these compartments

varied with developmental stage. Microfluidics-based observation of early larvae[52]

revealed substantial cytoplasmic GFP::KIN-20 signal during molts and at the beginning

of larval stages but a predominantly nuclear signal in the middle of the larval stage (Fig.

S7). We recapitulated these dynamics in L4 stage animals grown on plates (Fig. 6A),

Figure 6. The LIN-42 CKBD is required for KIN-20 nuclear localization. (A) Confocal images of *splitgfp::kin-20* animals staged according to vulva morphology (Mok et al., 2015) from late L3 to adult. Arrows indicate seam cell nuclear (white) and cytoplasmic (blue) localization; white arrowheads indicate nuclear hyp7 localization. Scale bar=20 μm. **(B)** Confocal images of *gfp::lin-42b+c; wrmscarlet::kin-20* staged according to vulva morphology. An arrow indicates seam cell, an arrowhead hyp7 nuclear localization. Scale bar=20 μm. **(C)** Confocal images of *splitgfp::kin-20* in wild type and *lin-42ΔCK1BD* mutant animals during early - mid L4 stage based on the vulvae morphology. Arrows indicate seam cell nuclear (white) and cytoplasmic (blue) localization, white arrowheads indicate hyp7 nuclear localization. Scale bar=20 μm. **(D)** Quantification of GFP::KIN-20 signal in hyp7 (nucleus), seam cells (nucleus) and hyp7 (cytoplasm) of wild-type (blue) and *lin-42ΔCK1BD* (orange) animals. Each condition with n=9 mid-L4 stage worms.

364 which additionally revealed an apparent membrane-bound pool of KIN-20 in seam cells
365 (Fig. 6A).
366

LIN-42 binding promotes nuclear localization of KIN-20
The timing of increased nuclear KIN-20 in the middle of a larval stage coincides with peak
accumulation of LIN-42, especially LIN-42b (Fig. 2C). Moreover, using endogenously
tagged LIN-42::GFP and wrmScarlet::KIN-20, we found that the two proteins co-localized
in the nuclei of epidermal seam and hyp7 cells (Fig. 6B). Hence, we wondered if LIN-42
contributed to the change in KIN-20 localization. To test this notion, we examined how
lack of LIN-42 binding affected KIN-20 localization dynamics. Specifically, we visualized
GFP::KIN-20 in *lin-42(ΔCK1BD)* mutant animals during mid-L4 stage. We observed
reduced nuclear GFP signal relative to wild-type animals (Fig. 6C,D) in both hypodermal
and seam cell nuclei with no significant change in cytoplasmic levels. Together, these
data indicate that LIN-42 promotes nuclear accumulation of KIN-20.
378

Discussion
PER and its nematode orthologue LIN-42 control two distinct types of biological timing:
mammalian and fly circadian rhythms (PER) and *C. elegans* developmental progression
(LIN-42). The identification of additional orthologous pairs functioning in these respective
pathways, ROR/NHR-23, CK1/DBT/KIN-20, Timeless/TIM-1, and REV-ERB/NHR-85,
suggests that there is a conserved set of biological timing genes, whose function can be
exploited in different contexts. The wiring among these components, in the few cases
where this has been explored, appeared strikingly different, making it unclear why and
how these genes would be evolutionarily selected for distinct timing functions. Our work
shows the first example of a conserved module in these clocks: the LIN-42/PER – KIN-
20/CK1 interaction.
390

The PAS domains are largely dispensable for LIN-42 developmental timing function
The PAS domains are the most notable feature of LIN-42 sequence conservation and
were thought to be an interaction platform for a developmental stage-specific binding
partner.[19] Previous work characterizing the *lin-42(ok2385)* and *n1089* alleles suggested

that the PAS domains were necessary for the heterochronic functions of LIN-42.[22,35] Yet,
our precise deletion of the LIN-42 PAS domains caused only a mild precocious alae
phenotype with significantly less penetrance than in *lin-42(n1089)* mutants, suggesting
that this model is incorrect. While *lin-42(n1089)* removes the sequence of the PAS
domains, it also results in a truncation of *lin-42b* and removal of part of the *lin-42a*
promoter sequence.[22,35] The *n1089* phenotype could reflect loss of the PAS domains
combined with potential reduction in expression of other elements such as the C-terminal
tail or an NHR-85 interaction site.[37]

The weak phenotypes seen with our precise deletion are particularly striking when
considering that the PAS domains, and specifically the highly conserved PAS-B domain,
mediate LIN-42 dimerization[20] - a function that is important for PER activity in circadian
rhythms[53]. Although it remains to be determined whether the PAS domains are necessary
for LIN-42 dimerization *in vivo*, it seems possible that LIN-42's molting timing function
does not require dimerization.

LIN-42 contains a CK1BD that is important for rhythmic molting
Contrasting with the PAS domain, we found the LIN-42 SYQ/LT regions to be required
for rhythmic molting, and we could demonstrate *in vitro* that they constitute a functional
CK1BD that anchors KIN-20/CK1, promotes LIN-42 phosphorylation, and mediates
kinase inhibition. Moreover, the patterns of phosphorylation that CK1 generates on LIN-
42 *in vitro* overlap with those that we observed on LIN-42 *in vivo*. We note that it remains
to be demonstrated to which extent the *in vivo* phosphorylation events depend on KIN-
20. Our attempts to address this question by examining LIN-42 phosphorylation in strains
lacking active KIN-20 were thwarted by the slow growth, sickness, and arrhythmic molting
observed in these strains.

The two CK1BD subdomains have separable functions *in vitro*. The CK1BD-B plays a
more prominent role in kinase binding, mirroring the anchoring function of its mammalian
and *Drosophila* equivalents.[14,46,49] Anchoring allows CK1 to target lower affinity, non-
consensus motifs, which are key regulatory sites in fly/mammalian circadian

clocks.[14,46,49,54,55] However, without the PER CK1BD-B not all capacity for
phosphorylation is lost, which suggests that high-affinity binding is not essential for
phosphorylation but promotes it.[49] For LIN-42-ΔCK1BD-B, we also see low levels of
phosphorylation by CK1 *in vitro* (Fig. 4), suggesting that it may play a similar role in
controlling KIN-20 activity.

The LIN-42 CK1BD-A plays a less prominent role in binding but rather is involved in CK1
inhibition. CK1δ-dependent phosphorylation of both mammalian and *Drosophila* PER at
sites near the CK1BD leads to feedback inhibition of CK1 through conserved anion
binding sites.[14] Our *in vitro* LIN-42 phosphorylation data reveal a decrease in enzyme
catalytic activity as substrate concentration increases, dependent on the CK1BD-A,
consistent with feedback product inhibition of KIN-20 (Fig. 4). Alternatively, this result
could reflect substrate inhibition, where increasing amounts of LIN-42 CK1BD+Tail result
in a non-productive enzyme-substrate complex. Such substrate inhibition has also been
reported in several recent studies of CK1 showing phosphorylated substrates binding the
enzyme and regulating its activity.[43,56–58] Further enzymatic and structural studies will be
required to distinguish between these modes of regulation.

Function of the LIN-42-CK1 interaction
Many models of circadian clock function highlight phosphorylation of PER by CK1 as the
central consequence of their interaction, controlling PER stability and thereby setting the
period length. Intriguingly, although CK1 heavily phosphorylates the LIN-42 C-terminal
tail, including on serine and threonine residues that are conserved among *Caenorhabditis*
nematodes, but not on more distantly related nematodes (Fig. S8), deletion of this
sequence does not affect molting rhythmicity. These data suggest either phosphosite
redundancy, or that regulated phosphorylation of LIN-42 is not the key function of this
complex for controlling molting timing.

For PER–CK1, several lines of evidence show that reciprocal to CK1-mediated regulation
of PER, PER also regulates CK1 activity, both on itself and potentially other targets such
as CLOCK. Indeed, PER and CRY function as a bridging complex to provide CK1 access

to CLOCK[8,9], and mutation of the CK1BD causes defects in PER and CK1 nuclear

localization as well as CK1-mediated phosphorylation of CLOCK.[9,59] Product inhibition

through phosphorylated PER may thus limit phosphorylation not only of PER itself, but

potentially also of other substrates, including CLOCK. Strikingly, KIN-20 also exhibits

dynamic subcellular localization, with nuclear accumulation depending on LIN-42 binding.

Whether this nuclear accumulation reflects co-transport, nuclear retention, or stabilization

of the nuclear KIN-20 pool is not known. However, together with the fact that the LIN-42

tail, despite being heavily phosphorylated, is dispensable for rhythmic molting, this

localization data indeed supports a model where a key function of the LIN-42–KIN-20

complex in the context of molting could be regulation of KIN-20 rather than LIN-42. Such

a scenario may also explain the previous observation that forced expression of the short

LIN-42a isoform, largely comprising the CK1BD + Tail, caused extended molt durations[35],

as this might sequester KIN-20 and/or inhibit its kinase activity.

Distinct functions of LIN-42 and KIN-20 in molting versus heterochronic timing?

Our data not only support the existence of LIN-42 and KIN-20 in a stable complex that is

important for molting timing, but also suggest that LIN-42 and KIN-20 can function

independently of one another in some contexts. Neither *kin-20(lf)* nor *lin-42(ΔCK1DB)*

mutations cause precocious alae, a phenotype observed with several other *lin-42* mutant

alleles. These findings agree with previous genetic data that suggested independent

functions of *lin-42* and *kin-20* in regulating the expression of the heterochronic *let-7*

microRNA.[38] A recent study on LIN-42's function in regulating another heterochronic

miRNA, *lin-4*, indicated that physical interaction with the NHR-85 transcription factor may

support LIN-42's activity in this process, and perhaps the heterochronic pathway more

generally.[37] Relatively mild effects upon *nhr-85* deletion, and LIN-42's ability to interact

with additional transcription factors in a yeast two-hybrid assay suggest that additional

functionally relevant interactions partners remain to be identified. The apparent

differences in wiring among components of the heterochronic pathway and the molting

timer may reflect their different modes of timekeeping where the heterochronic pathway

is chiefly concerned with the order of events and the molting timer - similar to the circadian

clock - with the tempo and/or robustness of their execution.

Indeed, although LIN-42/PER have previously been suggested to provide an evolutionary

link between circadian (PER) and heterochronic (LIN-42) timing systems[19], our data

suggest that the link primarily derives from functional conservation between circadian and

molting timing. Hence, we may speculate that the heterochronic function of LIN-42, which

appears to involve neither rhythmic activity nor conserved interactions with orthologues

of other circadian clock [19,21,23,26–28] arose secondarily to, or independent from, a rhythmic

timing function. Finally, we note with interest that the phosphorylation of PER by CK1 is

an important component of temperature compensation in the circadian clock[60], which

allows this clock to maintain its period despite variation in temperature. By contrast, the

period of *C. elegans* molting and oscillatory genes expression changes with temperature.

Our data that the LIN-42 tail, where the bulk of CK1 mediated phosphorylation occurs, is

dispensable for rhythmic molting may thus reflect the lack of robust temperature

compensation in the *C. elegans* developmental clock.

Materials and Methods

***C. elegans* culture**

C. elegans were grown and maintained at 20°C unless indicated otherwise and cultured

as described[61] on MYOB[62] or NGM 2% agar plates with *Escherichia coli* OP50 bacteria.

Full genotypes of strains used in this work and the respective growth conditions are listed

in Table S2. All mutant strains were backcrossed to N2 at least 2X. The Bristol N2 isolate

was used as the wildtype. Below, we describe how we generated novel strains for this

study. In Table S3, we list oligo repair templates, plasmids and genotyping primers. In

Table S4 we list crRNAs used.

Genome editing

Novel *lin-42* mutant alleles (*(wrd67[ΔPAS])*, (*wrd63[ΔCK1BD)* and (*wrd107[ΔTail])*) using

CRISPR-Cas9 were generated by injection of Cas9 or Cas12 ribonucleoprotein

complexes as described.[63] Specified deletions were made in the endogenous *lin-42* locus

in N2 animals. Repair templates were used at 100 ng/µl along with Cas9 or Cas12 at 250

ng/µl, crRNA oligos at 60 ng/µl, and 10 ng/µl co-injection marker (pCFJ90).[64] Strains used

in the luciferase assays were generated by crossing the mutant strains with a previously

described luciferase reporter strain (HW1993)[40] (Table S2).

Strains containing endogenously tagged *lin-42* and *kin-20* were obtained using

CRISPR/Cas9 as described previously.[64] In short: 5 µg of of Alt-R S.p. Cas9 Nuclease

V3 (IDT, Cat # 1081058), 2 µg of Alt-R® CRISPR-Cas9 tracrRNA (IDT, Cat # 1072532)

and 1.5 µg of crRNA were incubated at 37°C for 15 min to form the RNP complex. 500

ng repair template and co-injection markers pIK127 (10 ng/µl) and pRF4 (*rol-6(su1006)*)

(40 ng/µl) were added to a final volume of 20 µl in water. Mix was injected into N2 animals.

For *lin-42*, a *gfp::3xflag* (amplified from plasmid pIK384 with primers KK41/KK42) or

3xflag tag (Ultramer DNA Oligo, KK90) was inserted at the N-terminus of LIN-42b+c

isoforms.

For kin-20, a *3x::flag-ha* (Ultramer DNA oligo, KK107), *4xgfp11::linker:.flag* (amplified

from plasmid pIK401 with primers KK145/KK146) or *wrmScarlet* (amplified from plasmid

pIK385 with primers KK102/KK103) was inserted at the first common shared exon of all

KIN-20 isoforms with following flanking sequence: 5′ tatattcaaattttcagCGGAGATG-

[insert]-GAACTTCGTGTCGGCAATCGTTTCC 3′).

To reconstitute GFP with the split-GFP system, we crossed the obtained

4xgfp11::linker::flag::kin-20 line into a MosSCI strain containing the *gfp1-10* fragment

expressed from an *eft-3* promoter. For this strain the *gfp1-10* sequence from plasmid

pCZGY2254[65] for *C. elegans* was codon optimized. After adding two synthetic introns the

codon-optimized sequence was ordered as a gBlock from IDT. The sequence was cloned

into a MosSCI-compatible backbone together with *eft-3*p promoter and *tbb-2* 3'UTR

sequences to create plasmid pIK407 by Gibson assembly. pIK407 (*eft-3p::gfp1-*

10(codon-optimized)::tbb-2 3'UTR) was inserted into the ttTi5605 site on chromosome II

by MosSCI.[66]

To generate a catalytic dead KIN-20, the D310A mutation was introduced by CRISPR-
Cas9 into N2 and *3xflag-ha::kin-20* animals, respectively. Injection was performed as
described previously for the *lin-42* and *kin-20* tagged strains. The ultramer DNA oligo
KK137 containing the D310A mutation and two silent mutations (to abrogate crRNA
recognition) was used as a repair template.

Immunoprecipitation

For the 3xFLAG-LIN-42 IP, mixed stage animals were obtained by plating eggs after a
standard hypochlorite treatment and growth at different temperatures (15°C, 20°C and
25°C). 24 hrs after plating, worms were collected by pooling 30,000 worms from each
temperature condition per strain and replicate (triplicates for each strain). For the HA-KIN-
20 IP, synchronized worms were collected after 32 hrs at 25°C. For the mapping of LIN-
42 phosphorylation sites, synchronized worms were grown at 25°C and collected hourly
from 14 hrs (L2 larval stage) to 25 hrs (L4 larval stage). Worm extracts followed by
immunoprecipitation were done as previously described.[67] In brief, extracts were made in
lysis buffer (50mM Tris-HCl, pH 7.5, 150 mM NaCl, 1% TRITON X-100, 1 mM EDTA)
supplemented with 1 mM PMSF and 1 tablet of cOmplete Protease inhibitor (Roche, cat.
No. 11 873 580 001) per 50 ml using an MP Biomedical Fast-Prep-24 5G bead beater.
For the FLAG IP, cleared worm extract was incubated with 50 µl of 50% bead suspension
of anti-FLAG M2 Magnetic Beads (Sigma. Catalog Number M8823) for 2 hours at 4°C.
For the HA-IP worm extract was incubated with 50 µl of bead suspension of Pierce™
Anti-HA Magnetic Beads (Thermo Scientific, Cat. No. 88837).

Proteolytic digest

The beads were incubated for 4 hrs at RT with 800 rpm shaking in 5 µl of digestion buffer
(3 M guanidine hydrochloride, 20 mM EPPS (4-(2-Hydroxyethyl)-1-
piperazinepropanesulfonic acid) pH 8.5, 10 mM CAA (2-Chloroacetamide), 5 mM TCEP
(Tris(2-carboxyethyl)phosphine hydrochloride)) and 1 µl of Lys-C (0.2 µg/µl in 50 mM[68],
pH 8.5). 1 µl of trypsin (0.2 µg/µl) was added and incubated at 37°C overnight. After 12
hrs another 1 µl of trypsin was added and incubation continued for 4 more hours at 37°C

Mass spectrometry to identify LIN-42 interacting factors

The generated peptides were acidified with 0.8% TFA (final concentration) and analyzed by LC–MS/MS on an EASY-nLC 1000 [Thermo Scientific] with a two-column set-up. The peptides were applied onto a peptide 75 µm x 2 cm PepMap Trap trapping column [Thermo Scientific] in 0.1% formic acid, 2% acetonitrile in H2O at a constant pressure of 800 bar. Using a flow rate of 150 nl/min, peptides were separated on a 50 µm x 15 cm PepMap C18, 2 µm, 100 A [Thermo Scientific] at 45 °C with a linear gradient of 2%–6% buffer B in buffer A in 3 min followed by a linear increase from 6 to 22% in 40 min, 22%–28% in 9 min, 28%–36% in 8 min, 36%–80% in 1 min, and the column was finally washed for 14 min at 80% buffer B in buffer A (buffer A: 0.1% formic acid; buffer B: 0.1% formic acid in acetonitrile). The separation column was mounted on an EASY-Spray™ source [Thermo Scientific] connected to an Orbitrap Fusion LUMOS [Thermo Scientific]. The data were acquired using 120,000 resolution for the peptide measurements in the Orbitrap and a top T (3 s) method with HCD fragmentation for each precursor and fragment measurement in the ion trap according the recommendation of the manufacturer (Thermo Scientific). For the analysis protein identification and relative quantification of the proteins was performed with MaxQuant v.1.5.3.8 using Andromeda as search engine, and label-free quantification (LFQ). The *C. elegans* subset of the UniProt v.2021_05 combined with the contaminant database from MaxQuant was searched and the protein and peptide FDR were set to 0.01. The LFQ intensities estimated by MaxQuant were analyzed with the einprot R package (https://github.com/fmicompbio/einprot) v0.5.4. Features classified by MaxQuant as potential contaminants or reverse (decoy) hits or identified only by site, as well as features identified based on a single peptide or with a score below 10, were filtered out. The LFQ intensities were log2 transformed and missing values were imputed using the 'MinProb' method from the imputeLCMD R package v2.0 with default settings. Pairwise comparisons were performed using limma v3.50.0, considering only features with at least 2 non-imputed values across all the samples in the comparison. Estimated log2-fold changes and P-values (moderated t-test) from limma were used to construct volcano plots. Data will be available via ProteomeXchange with identifier PXD058601 upon publication.

Mass spectrometry to map *in vivo* LIN-42 phosphorylation sites

The peptides from each sample were labeled with TMTpro reagents and pooled. The TMT labeled peptide mixture was subjected to off-line high pH fractionation on a YMC Triart C18 0.5 × 250 mm column (YMC Europe GmbH) using the Agilent 1100 system (Agilent Technologies). A total of 96 fractions was collected for each experiment and concatenated into 48 fractions as previously described. For each LC-MS analysis, all available peptides were loaded onto a PepMap Neo trap (Thermo Fisher) using the Vanquish Neo UHPLC system (Thermo Fisher). On-line peptide separation was performed on a 15-cm EASY-Spray™ C18 column (ES75150PN, Thermo Fisher) by applying a linear gradient of increasing ACN concentration at a flowrate of 200 nL/min. Orbitrap Fusion Lumos Tribrid (Thermo Fisher) mass spectrometer was operated in the data-dependent mode. The ions for the survey scan were collected for a maximum of 50 ms to reach the standard AGC target value and the scan recorded using an Orbitrap detector at a resolution of 120,000. The topmost intense precursor ions from the Orbitrap survey scan recorded every 3 sec were selected for stepped higher-energy C-trap dissociation (HCD) at 29%, 32% and 35% normalized collision energy scan. To reach an AGC value of 100,000 ions, the maximum ion accumulation time for the MS2 scan was set to 500 ms. The TMT reporter ions were quantified using an MS2 scan recorded using the Orbitrap analyzer at a resolution of 50,000. Thermo RAW files were processed using Proteome Discoverer 2.4 software (Thermo Fisher) as described in the manufacturer's instructions. Briefly, the Sequest search engine was used to search the MS2 spectra against the *C. elegans* UniProt database (downloaded on 05/2023) supplemented with common contaminating proteins. For peptide identification, cysteine carbamidomethylation and TMTpro tags on lysine and peptide N-termini were set as static modifications, whereas oxidation of methionine residues and acetylation protein N-termini were set as variable modifications. The assignments of the MS2 scans were filtered to allow 1% FDR. For reporter quantification, the S/N values were corrected for isotopic impurities of the TMTpro reagent using the values provided by the manufacturer. The sums across all TMTpro reporter channels were normalized assuming equal total protein content in each sample. Data will be available via ProteomeXchange with identifier PXD058598 upon publication.

Western Blot

Synchronized animals were collected hourly from 13 hrs - 23 hrs after plating at 25°C. Extracts were made by boiling at 95°C for 5 minutes in lysis buffer (63 mM Tris-HCl (pH 6.8), 5 mM DTT, 2% SDS, 5% sucrose) followed by sonication with a BioRupter Plus (Diagnode) with the following settings: 13 cycles, 30 s on/off at 4°C. Samples were cleared by centrifugation, before separating proteins by SDS-PAGE (loading: 50 µg protein extract per well) and transferring them to PVDF membranes by semi-dry blotting. The following antibodies were used: Monoclonal mouse anti-FLAG M2-Peroxidase (HRP)(Sigma-Aldrich; A8592, dilution: 1:1000), monoclonal mouse anti-Actin clone C4 (Millipore; MAB1501, dilution 1:7500) and anti-mouse HRP conjugated antibody (GE Healthcare #NXA931, dilution 1:2000). The membrane was incubated with ECL™ Prime Western-Blot-Reagent (Cytiva, #RPN2236) and bands detected with an Amersham Imager 680 (GE Healthcare).

Phenotypic analysis

For phenotypic analyses, gravid adults were bleached, and embryos were hatched at low density onto MYOB plates seeded with OP50, or gravid adults were picked onto seeded plates and removed after 1-2 hours. Animals were maintained at 20°C and observed daily or twice daily. Bag-of-worms phenotype was determined when live progeny were observed inside adult animals. For brood counts, animals were individually transferred to fresh wells daily for five days after L4 and live progeny were counted and averaged.

To score premature alae, synchronized animals were collected from MYOB plates by washing off plates. 1000 µl of M9 + 2% gelatin was added to the plate or well, agitated to suspend animals in M9+gelatin, and then transferred to a 1.5 ml tube. Animals were spun at 700xg for 1 min. The media was then aspirated off and animals were resuspended in 500µl M9 + 2% gelatin with 5 mM levamisole. 12 µl of animals in M9 +gel with levamisole solution were placed on slides with a 2% agarose pad and secured with a coverslip. Images were acquired using a Plan-Apochromat 40x/1.3 Oil DIC lens or a Plan-Apochromat 63x/1.4 Oil DIC lens on an AxioImager M2 microscope (Carl Zeiss

Microscopy, LLC) equipped with a Colibri 7 LED light source and an Axiocam 506 mono

camera. Acquired images were processed through Affinity photo software (version:

1.9.2.1035). For the confocal microscopy, L3/L4 stage-animals were picked from OP50

plates grown at 25°C. Worms were mounted on a glass slide with a 2 % agarose patch

immobilized with 5 µl of 10 mM levamisole (Fluca Analytical, #31742). Images were

acquired in channels for red (561 nm laser), green (488 nm laser) and DIC with a 40x/1.3

immersion objective on a Zeiss LSM700 confocal microscope. Acquired images were

processed with FiJi.[68]

Quantification of GFP levels

For quantification, mid-L4 stage worms (defined by vulval morphology) were selected.

Images acquired in the GFP fluorescence channel were imported into Napari for

analysis.[69] The following regions of interest were manually annotated for quantification:

Nuclei of hyp7 and seam cells. For the cytoplasm of hyp7, the layer with the maximum

intensity of nuclear GFP signal in hyp7 was selected and a region between the annotated

hyp7 nuclei was chosen. To define the background, a region in the gonad was used

(where no GFP signal was detected). For the quantification, average GFP levels were

computed per region and worm. Background levels were subtracted to the GFP levels of

all other regions.

Microfluidics

Worms were prepared and loaded into the device as described in Berger et al. (2021).

Briefly, gravid adult hermaphrodites were harvested and treated with bleach. The

resulting embryos were collected by centrifugation at 1,600 g for 1 minute and washed

three times with S-Basal buffer. For synchronization, embryos were incubated overnight

in S-Basal, allowing larvae to arrest. Arrested larvae were then passed through a 10 µm

filter (pluriStrainer Mini 10 µm, PluriSelect), and 6,000 worms were transferred to 3 NGM

plates, where they were incubated for 12 hours at 25°C. Once the worms reached the

late L1 stage, they were harvested using M9 buffer, washed twice to remove debris, and

loaded into the experimental device. During the experiment, bacterial food was supplied

at a constant rate of 1 µl/h, with periodic increases to 100 µl/h for 5 seconds every 30

minutes to clear debris. Images were acquired every 10 minutes over 24 hours using a
spinning disk confocal scanning microscope (Yokogawa CSU W1 with Dual T2).
Brightfield and fluorescent signals (488 nm laser) were recorded simultaneously using
two sCMOS Photometrics Prime 95B cameras with a 40 x oil immersion lens (NA = 1.3).
Imaging was conducted with a 25 ms exposure time and a motorized z-drive, acquiring
z-stacks with a 1 μm step size and 20 images per stack. Analysis was performed using
Fiji/ImageJ software.
711

Luciferase assays
Assays were performed and analyzed as described.[40] Briefly, gravid adults were
bleached, and eggs were immediately singled into wells containing 90 μl OP50/S-
Basal/D-Luciferin solution per well and left in the luminometer machine and measured
every 10 minutes for 0.5 seconds. The assays lasted 96 hrs or 130 hrs. For statistical
analysis, we performed a Wilcoxon-Mann-Whitney test (implemented in the python
package SciPy version 1.4.1 as the function Mann-Whitney-U).
719

Expression and purification of recombinant proteins
All proteins were expressed from a pET22-based vector in *Escherichia coli* Rosetta (DE3)
cells based on the Parallel vector series.[70] All LIN-42 constructs (CK1BD + Tail, residues
402-598; CK1BD, residues 402-475; CK1BD-ΔA, -ΔB, and ΔAB mutants) were expressed
downstream of an N-terminal TEV-cleavable His-NusA tag. Human CK1δ catalytic
domains (CK1δ ΔC, residues 1–317) were all expressed with a TEV-cleavable His-GST
tag. All proteins expressed from Parallel vectors have an additional N-terminal vector
artifact of 'GAMDPEF' remaining after TEV cleavage. Cells were grown in LB media at
37°C until the O.D.600 reached ~0.8; expression was induced with 0.5 mM IPTG, and
cultures were grown for approximately 16–20 hr more at 18°C. Cells were centrifuged at
3,200 x g, resuspended in 50 mM Tris, pH 7.5, 300 mM NaCl, 20 mM imidazole, 5%
(vol/vol) glycerol, 1 mM tris(2-carboxyethyl)phosphine (TCEP), and 0.05% Tween-20. For
purification of recombinant protein, cells were lysed with a microfluidizer followed by
sonication and then the lysate was clarified via centrifugation at 140,500 x g for 1 hour at
4°C. Ni-NTA affinity chromatography was used to extract his-tagged proteins from the

lysate and then the affinity/solubility tags were cleaved using His6-TEV (GST-TEV for
CK1δ ΔC) protease overnight at 4ºC. The cleaved protein was then separated from
solubility tag and TEV by a second Ni-NTA (GST for CK1δ ΔC) affinity column and further
purified using size exclusion chromatography (SEC) in 50 mM Tris, pH 7.5, 200 mM NaCl,
1 mM EDTA, 5% (vol/vol) glycerol, 1 mM TCEP, and 0.05% Tween-20. Small aliquots of
protein were frozen in liquid nitrogen and stored at -70ºC for long-term storage.
741

In vitro biotinylation, pull-down assays, and bio-layer interferometry
LIN-42 CK1BD + Tail constructs and CK1δ ΔC were biotinylated via Sortase A-mediated
reactions between a Sortase A recognition motif peptide (biotin-LPETGG) and our LIN-
42 CK1BD + Tail/CK1δ ΔC (N-terminal G from 'GAMDPEF' artifact). Reactions were
carried out in 50 mM Tris pH 7.5 and 150 mM NaCl using 5 μM His6-Sortase A, 300 μM
biotin-LPETGG, and 50 μM protein. Ni-NTA affinity chromatography followed by SEC was
used to purify labeled protein from His6-Sortase A and excess biotin, respectively. Pull
down assays were performed using magnetic streptavidin beads to bind biotinylated LIN-
42 CK1BD + Tail WT and mutants (final concentration 5 μM) in the presence and absence
of CK1δ ΔC (final concentration 5 μM). All BLI assays were performed in SEC buffer
supplemented with 7.5mM BSA as previously described[71,72] using an eight-channel Octet-
RED96e.
754

ADP-Glo kinase assays
Substrate titration kinase reactions were performed on the indicated recombinant LIN-42
proteins (CK1BD + Tail WT or mutants) using the ADP-Glo kinase assay kit (Promega)
according to the manufacturer's instructions. All reactions were performed in 30 μL
volumes in duplicate (n=3 independent experiments) using 1x kinase buffer (25mM Tris
pH 7.5, 100 mM NaCl, 10 mM $MgCl_2$, and 2 mM TCEP) supplemented with 100μM ATP,
0.2μM recombinant CK1, and indicated LIN-42 proteins. Reactions were held at room
temperature for 1 hour and then 5 μL aliquots were taken and quenched with ADP-Glo
reagents. Luminescent measurements were measured at room temperature with a
SYNERGY2 microplate reader in 384-well microplates. Data analysis was performed
using Excel (Microsoft) and Prism (GraphPad).

^{32}P-ATP kinase assays

1 μM CK1δ ΔC was incubated with 10 μM LIN-42 (CK1BD + Tail/CK1BD WT and mutant proteins) in 1x kinase buffer (25 mM Tris pH 7.5, 100 mM NaCl, 10 mM MgCl2, 2 mM TCEP). Reactions were started by the addition of ^{32}P-ATP (final concentration 2 mM) and samples were collected at indicated time points and quenched in an equivalent volume of 2x SDS-PAGE loading buffer. Proteins labeled with ^{32}P were separated and analyzed via SDS-PAGE and the gels were dried at 80ºC for 2 hours before overnight exposure in a phosphor screen (Amersham Biosciences). A Typhoon Trio (Amersham Biosciences) phosphorimager was used to visualize exposed gels and ^{32}P-labeled protein bands were quantified via densitometry using ImageJ (NIH), Excel (Microsoft) and Prism (GraphPad).

In vitro **Mass Spectrometry**

Sample preparation:

Kinase reactions were performed in 1x kinase buffer (25 mM Tris pH 7.5, 100 mM NaCl, 10 mM MgCl2, 2 mM TCEP) for 60 minutes and then quenched with 20mM EDTA. Samples were denatured, reduced, alkylated, and digested according to the In-solution Protein Digestion (Promega) protocol. In brief, phosphorylated samples were denatured/reduced in 8M urea/50mM ammonium bicarbonate/5mM Dithiothreitol (DTT) for 1hr at 37°C. Samples were incubated for 30 minutes in the dark with 15mM iodoacetamide and then digested with X Grade Modified Trypsin according to the Trypsin Digestion Protocol (Promega). Digested Samples were phospho-enriched using the High-Select Fe-NTA Phosphopeptide Enrichment Kit (Thermo Scientific) and sent for analysis at the University of California, Davis Proteomic Core Facility.

LC-MS:

For each sample, equal volumes were loaded onto a disposable Evotip C18 trap column (Evosep Biosytems, Denmark) as per the manufacturer's instructions. Briefly, Evotips were wetted with 2-propanol, equilibrated with 0.1% formic acid, and then loaded using centrifugal force at 1200g. Evotips were subsequently washed with 0.1% formic acid, and then 200 µL of 0.1% formic acid was added to each tip to prevent drying. The tipped samples were subjected to nanoLC on a Evosep One instrument (Evosep Biosystems).

Tips were eluted directly onto a PepSep analytical column, dimensions: 15 cm x 75 um

C18 column (PepSep, Denmark) with 1.5 µm particle size (100 Å pores) (Bruker

Daltonics), and a ZDV spray emmiter (Bruker Daltronics). Mobile phases A and B were

water with 0.1% formic acid (v/v) and 80/20/0.1% ACN/water/formic acid (v/v/vol),

respectively. The standard pre-set method of 40 samples-per-day whisper method was

used, which is a 31-minute run.

Mass Spectrometry – Performed on a hybrid trapped ion mobility spectrometry-

quadrupole time of flight mass spectrometer (timsTOF Pro, (Bruker Daltonics, Bremen,

Germany) with a modified nano-electrospray ion source (CaptiveSpray, Bruker

Daltonics). In the experiments described here, the mass spectrometer was operated in

diaPASEF mode. Desolvated ions entered the vacuum region through the glass capillary

and deflected into the TIMS tunnel which is electrically separated into two parts (dual

TIMS). Here, the first region is operated as an ion accumulation trap that primarily stores

all ions entering the mass spectrometer, while the second part performs trapped ion

mobility analysis.

DIA PASEF:

The dual TIMS analyzer was operated at a fixed duty cycle close to 100% using equal

accumulation and ramp times of 85 ms each.

Data-independent analysis (DIA) scheme consisted of one MS scan followed by MSMS

scans taken with 19 precursor windows at width of 50Th per 0.57s cycle over the mass

range 300-1200 Dalton. The TIMS scans layer the doubly and triply charged peptides

over a ion mobility -1/k0- range of 0.7-1.3 V*sec/cm2. The collision energy was ramped

linearly as a function of the mobility from 59 eV at 1/K0=1.4 to 20 eV at 1/K0=0.6.

Data Analysis:

DIA

LCMS files were processed with Spectronaut version 16.1 (Biognosys, Zurich,

Switzerland) using DirectDIA analysis mode. Mass tolerance/accuracy for precursor and

fragment identification was set to default settings. The unreviewed FASTA for *C. elegans*

was downloaded from Uniprot and a database of common laboratory contaminants were

used.[73] A maximum of two missing cleavages were allowed, the required minimum

peptide sequence length was 7 amino acids, and the peptide mass was limited to a

maximum of 4600 Da. Carbamidomethylation of cysteine residues was set as a fixed

modification, and methionine oxidation and acetylation of protein N termini as variable

modifications. A decoy false discovery rate (FDR) at less than 1% for peptide spectrum

matches and protein group identifications was used for spectra filtering (Spectronaut

default). Decoy database hits, proteins identified as potential contaminants, and proteins

identified exclusively by one site modification were excluded from further analysis.

Data availability:

Data is available from the massive online repository https://massive.ucsd.edu/ ID number

MSV000096641 and proteome Exchange ID number PXD058784

Author contributions

Rebecca K. Spangler: Conceptualization; formal analysis; investigation; writing –

original draft. **Guinevere E. Ashley:** Conceptualization; formal analysis; investigation;

writing – original draft. **Kathrin Braun:** Conceptualization; Formal analysis; investigation;

writing – review and editing. **Marit van der Does:** Formal analysis; investigation; writing

– review and editing. **Daniel Wruck:** Formal analysis; investigation; writing – review and

editing. **Andrea Ramos-Coronado:** Formal analysis; investigation; writing – review and

editing. **James Matthew Ragle:** Formal analysis; investigation; writing – review and

editing. **Vytautas Iesmantavicius:** Formal analysis; investigation; writing – review and

editing. **Lucas J Morales Moya:** Formal analysis; writing – review and editing. **Keya**

Jonnalagadda: Formal analysis; writing – review and editing. **Carrie L. Partch:**

Conceptualization; supervision; funding acquisition; writing – review and editing; project

administration. **Helge Großhans:** Conceptualization; supervision; funding acquisition;

writing – original draft; project administration. **Jordan D. Ward:** Conceptualization;

supervision; funding acquisition; writing – original draft; project administration.

Acknowledgements

Some strains were provided by the *Caenorhabditis* Genetics Center, which is funded by

the NIH Office of Research Infrastructure Programs [P40 OD010440]. We thank Iskra

Katic for worm injections and technical support, Daniel Hess and Jan Seebacher for mass

spectrometry analysis, and Dimos Gaidatzis for help with proteomics data analysis. We

are thankful to Laurent Gelman and Laure Plantard for help with confocal imaging. Some

analysis for this project was performed in the Proteomics Core Facility of the Genome

Center, University of California, Davis with instrument funding provided by the NIH

(S10OD026918-01A1). We thank Michelle Salemi (UC Davis) for performing the

proteomics sample prep, the LC-MS/MS method writing and running and data analysis.

The Bruker tims-TOF HT MS/Evosep LC system was supported by the Howard Hughes

Medical Institute Investigator Award for Dr. Neal Hunter, UC Davis. Biolayer

Interferometry data were collected with instrument funding provided by the NIH

(S10OD027012).

Funding

This work was funded by the National Institutes of Health (NIH) National Institute of

General Medical Sciences (NIGMS) to J.D.W (R01GM138701) and C.L.P.

(R35GM141849), and the Howard Hughes Medical Institute to C.L.P. This work is part of

a project that has received funding from the European Research Council (ERC) under the

European Union's Horizon 2020 research and innovation program (Grant agreement No.

741269, to H.G.) and from the Swiss National Science Foundation (#310030_207470, to

H.G.). The FMI is core-funded by the Novartis Research Foundation.

Data availability

All relevant data can be found within the article and its supplementary information.

Figure legends

Figure 1. *lin-42ΔPAS* mutations cause mild heterochronic phenotypes while *lin-

***42ΔCK1BD* mutations cause asynchronous molting (A)** Schematic depiction of the

lin-42 genomic locus. The gene structure of the *lin-42 a, b*, and *c* isoforms are shown.

The location of sequences encoding putative PAS-A, PAS-B, SYQ, and LT sequence

motifs are indicated. Published *lin-42* deletion alleles *n1089* and *ok2385* are indicated in

dark and light grey, respectively, newly generated targeted ΔPAS and ΔCK1BD deletions

in navy and teal, respectively. **(B)** Heatmaps showing trend-corrected luminescence

traces from the indicated genotype. Each horizontal line represents one animal. Traces

are sorted by entry into the first molt. Darker color indicates low luminescence signal and

corresponds to the molts. **(C)** Bar plot quantifying the percentage of animals of the

indicated genotype with precocious complete or partial alae at the L3-L4 molt. **(D)** Bar

plot quantifying the percentage of animals of the indicated genotype that exhibited egg-

laying defects as determined by the presence of hatched larvae in the animal. **(E)** Bar plot

depicting the average number of live progeny from hermaphrodites of the indicated

genotype. **(F)** Bar plot quantifying the percentage of animals of the indicated genotype

that arrested or died as larvae. (C-F) Statistical significance was determined using an

ordinary one-way ANOVA. $p < 0.05$ was considered statistically significant. **** indicates

< 0.0001. Error bars in C-F represent standard deviation.

Figure 2 LIN-42 and KIN-20 interact *in vivo* (A) Volcano blot comparing protein

enrichments   in   3xFLAG::LIN-42b+c   and   3xFLAG::SART-3   (control)

immunoprecipitations, determined by mass spectrometry. **(B)** mRNA expression profile

for *lin-42* (dark blue) and *kin-20* (light blue) mRNAs throughout larval development. Data

Source: Meeuse et al., 2020. **(C)** Western blot with extracts from *3xflag::lin-42b+c; 3xflag-*

ha::kin-20 (HW3479) larvae collected hourly for 11 hours, starting at 13 h after plating

synchronized L1 stage animals on food at 25°C. Top part probed with anti-FLAG-HRP

(1:1,000), lower part with anti-actin-1 (1:7,500). Arrows indicate bands for LIN-42b, KIN-

20 and ACT-1 (see Fig. S5A). **(D)** Anti-HA pulldown from *3xflag::lin-42b+c; 3xflag-ha::kin-*

20 animals. Blot probed with anti-FLAG (1:1,000).

Figure 3 The LIN-42 SYQ and LT domains constitute a functional CK1-binding

domain (A) Schematic representing PER2 and LIN-42 protein domains. Protein

constructs used in this study, CK1BD + Tail and CK1BD, are indicated i. CK1BD, Casein

Kinase 1-Binding Domain; CBD, CRY-binding domain. **(B)** Representative pulldown

assay of human CK1 and biotinylated LIN-42 CK1BD + Tail proteins using the indicated

protein variants. **(C)** Values for K_D from kinetic analysis of BLI data (D-H) based on a 2:1

heterogeneous ligand binding model and global analysis (Octet). Mean ± SEM, ordinary

one-way ANOVA. $p < 0.05$ was considered statistically significant. **** indicates $p < 0.0001$.

(D-H) Bio-layer interferometry (BLI) data for indicated LIN-42 protein binding to

immobilized, biotinylated CK1. Inset values represent the concentrations of LIN-42 for

individual binding reactions. Model fit to association and dissociation over time is

represented by black lines. Data shown from one representative experiment of n ≥ 3

assays.

Figure 4 The extended C-terminus of LIN-42 is phosphorylated by CK1 (A)

Representative [32]P-radiolabeled ATP enzymatic assay of CK1 activity on LIN-42 CK1BD

+ Tail and CK1BD wild-type and indicated domain deletion mutants. **(B)** Quantification of

band intensity depicted in (A) and replicates. Model fit to Michaelis- Menten, n = 3. **(C)**

Fold-change quantification of band intensity from panels A & B compared to CK1

autophosphorylation at 120 minutes. Ordinary one-way ANOVA. p<0.05 was considered

statistically significant. ****p < 0.0001, n ≥ 3, mean ± SEM. (D) Representative ADP-Glo

enzymatic assay comparing activity of CK1 on LIN-42 CK1BD + Tail wild-type (maroon),

CK1BD-ΔA (orange), CK1BD-ΔB (yellow), and CK1BD-ΔA/B (green). **(E)** Values of k_{cat}/K$_m$

calculated from (D). n=3, mean ± SEM, Ordinary one-way ANOVA. p<0.05 was

considered statistically significant. * indicates p<0.03 and ** indicates p<0.002. **(F)** LIN-

42 phosphorylation sites identified via mass spectrometry from *in vivo* samples from

whole worm lysate collected hourly from synchronized L1 animals from 14 hrs (early L2)

to 25 hrs (late L3) after plating at 25°C (top arrows) and *in vitro* kinase reactions of CK1

activity on LIN-42 CK1BD + Tail (bottom arrows). Serine, threonine, and tyrosine residues

are indicated via vertical blue, purple, and pink ticks, respectively, along the CK1BD +

Tail construct schematic.

Figure 5 *kin-20* null and catalytically dead alleles exhibit larval arrest and

asynchronous molting similar to *lin-42(ΔCK1BD)* mutants (A) Heatmaps showing

trend-corrected luminescence traces from the indicated genotype. Each horizontal line

represents one animal. Traces are sorted by entry into the first molt. Darker color indicates

low luminescence signal and corresponds to the molts. **(B)** Bar plot depicting the average

number of live progeny from hermaphrodites of the indicated genotype. **(C)** Bar plot

quantifying the percentage of animals of the indicated genotype that exhibited egg-laying

defects as determined by the presence of hatched larvae in the animal. **(D)** Bar plot

quantifying the percentage of animals of the indicated genotype with precocious complete

or partial alae at the L3-L4 molt. **(E)** Bar plot quantifying the percentage of animals of the

indicated genotype that arrested or died as larvae. **(B-E)** Statistical significance was

determined using ordinary one-way ANOVA. p<0.05 was considered statistically

significant. ** indicates p<0.0001****, indicates p<0.0001. Error bars in B-E represent

standard deviation.

Figure 6. The LIN-42 CKBD is required for KIN-20 nuclear localization. (A) Confocal

images of *splitgfp::kin-20* animals staged according to vulva morphology (Mok et al.,

2015) from late L3 to adult. Arrows indicate seam cell nuclear (white) and cytoplasmic

(blue) localization; white arrowheads indicate nuclear hyp7 localization. Scale bar=20 µm.

(B) Confocal images of *gfp::lin-42b+c; wrmscarlet::kin-20* staged according to vulva

morphology. An arrow indicates seam cell, an arrowhead hyp7 nuclear localization. Scale

bar=20 µm. **(C)** Confocal images of *splitgfp::kin-20* in wild type and *lin-42ΔCK1BD* mutant

animals during early - mid L4 stage based on the vulvae morphology. Arrows indicate

seam cell nuclear (white) and cytoplasmic (blue) localization, white arrowheads indicate

hyp7 nuclear localization. Scale bar=20 µm. **(D)** Quantification of GFP::KIN-20 signal in

hyp7 (nucleus), seam cells (nucleus) and hyp7 (cytoplasm) of wild-type (blue) and *lin-*

42ΔCK1BD (orange) animals. Each condition with n=9 mid-L4 stage worms.

Figure S1 PER proteins have a crucial role in mammalian circadian rhythms. (A)

Cartoon schematic of the primary transcription-translation feedback loop which generates

~24-hour rhythms in mammals. CCGs, Clock-controlled genes. **(B)** Cartoon schematic

illustrating the mammalian phosphoswitch that dictates CK1-dependent regulation of PER

stability. Crystal structure of human CK1δ (gray) bound to phosphorylated PER2 FASP

(orange, 4pFASP) peptide, PDB: 8d7o. pD, phosphodegron; FASP, Familial Advanced

Sleep Phase.

Figure S2 Larval stage durations for *lin-42* mutant animals. Boxplots showing

durations (in hours) for **(A)** larval stages **(B)** molts and **(C)** intermolts from luciferase

assay. Wild type in white, *lin-42(n1089)* in dark grey, *lin-42(ok2385)* in light grey, *lin-*

42ΔPAS in blue and *lin-42ΔCK1BD* in green. **(D)** Bar plot showing the number of molts

from the luciferase assay of the indicated genotypes.

Figure S3 Conservation and structural prediction of *C. elegans* KIN-20. (A)

Sequence alignment of *C. elegans* KIN-20 and KIN-19, *H. sapiens* CK1δ, *M. musculus*

CK1δ, *A. thalia* CK1, *S. cerevisiae* HRR25 (HRR), and *D. melanogaster* Doubletime

(DBT). Important enzymatic sequence features are boxed; full kinase domain (blue box -

79% identical between KIN-20 and human CK1δ), catalytic lysine (purple - conserved in

KIN-20), catalytic loop (pink - 100% conserved between KIN-20 and human CK1δ),

Magnesium (Mg)-binding loop (orange - 12 out of 13 residues conserved between KIN-

20 and human CK1), anion coordination sites 1 (yellow), 2 (green), and 3 (blue) (all

conserved in KIN-20). **(B)** Crystal structure of *H. sapiens* CK1δ (PDB 6pxo). Dark blue

indicates residue is conserved in KIN-20; residues that diverge are highlighted in pale

blue (similar amino acid) or not conserved (white). **(C)** AlphaFold structural prediction of

KIN-20 (https://alphafold.ebi.ac.uk/entry/A8X4B3) colored by the model confidence. **(D)**

KIN-20 AlphaFold plot of predicted aligned error.

Figure S4 Larval stage durations for *kin-20 and lin-42* mutant animals. Boxplots

showing durations (in hours) for **(A)** Larval stages **(B)** Molts **(C)** Intermolts from luciferase

assay. Wild type (*Wt*) in white, *lin-42(ΔTail)* in blue, *kin-20(0)* in light grey, *kin-20 D310A*

in dark grey. Statistics were done using the Mann-Whitney U-test. Stars indicate the

significance of difference between the *Wt* strain and the different *lin-42* and *kin-20* mutant

animals: * $p<0.05$, ** $p<0.01$, *** $p<0.001$, **** $p<0.0001$. **(D)** Bar plot showing the number

of molts detected in the assay in percentage of animals.

1007

1008   **Figure S5 Luciferase replicate of *lin-42(ΔTail).*** Heatmaps showing trend-corrected

1009   luminescence traces from the indicated genotype. Each horizontal line represents one

1010   animal. Traces are sorted to the entry of the first molt. Darker color indicates low

1011   luminescence signal and corresponds to the molts.

1012

Figure S6 *kin-20(D310A)* affects molt timing. (A) Western Blot with extracts from

3xflag::lin-42b+c, 3xflag::kin-20 and *3xflag::lin-42b+c; 3xflag::kin-20* larvae (mid L4

stage). Blot probed with anti-FLAG-HRP (1:1000). Arrows indicate bands for LIN-42b and

KIN-20. **(B)** Western Blot with extracts from *3xflag::kin-20* wild-type, *(xe401[D310A])* and

(xe400[D310A]) mutant animals. Top panel probed with anti-FLAG-HRP (1:1000). Lower

panel probed with anti-actin-1 (1:7500). Arrows indicate bands for KIN-20. **(C)** Heatmaps

showing trend-corrected luminescence traces from the indicated genotype. Each

horizontal line represents one animal. Traces are sorted by entry into the first molt. Darker

color indicates low luminescence signal and corresponds to the molts. **(D)** Boxplots

showing the duration (in hours) for the larval stage from the luciferase assay.

1023

1024 **Figure S7 KIN-20 dynamics during L2 - L3 stage.** Microscopy images of one

1025 *splitgfp::kin-20* larva from a microfluidics experiment at indicated timepoints. Time

1026 indicated in minutes after Molt 1 (M1) or Molt 2 (M2) exit. Arrows indicate seam cell

1027 nuclear (white) and cytoplasmic (blue) localization; white arrowheads indicate nuclear

1028 hyp7 localization. Scale bar=50 μm.

1029

Figure S8 Alignment of nematode LIN-42 protein sequences. LIN-42 homologs from

the indicated nematode species were aligned using Clustal Omega. The length in amino

acids of each homolog follows the species and homolog name. To the left and right of the

alignment are amino acid positions of the end residues for each protein. Blue shading

indicates conserved sequences and the histogram at the bottom depicts the degree of

conservation with a consensus sequence listed below. The positions of the *C. elegans*

CK1BD-A and CK1BD-B motifs are indicated. The location of the phosphosites found in

our *in vivo*, *in vitro* and both datasets are indicated.

1038

1039 **TABLES**

1040 **Table S1 *lin-42* mutant phenotypes**

1041 All animals grew from eggs hatched onto seeded plates at low density after sodium

1042 hypochlorite treatment. n ≥ 20 for all analyses. [a]Percentage of animals that failed to reach

1043 adulthood by day 8 after hatching or died as young larva. [b]Percentage of animals with

alae formation in early L4 (L4.0-L4.2). [c]Percentage of animals that exhibited bag-of-worms phenotype. [d]Average number of progeny from fertile adults; arrested and bagged animals were not included in the calculation. *Broods were not counted in animals with high instance of BOW phenotype.

Table S2 strains

C. elegans strains used in this work.

Table S3 oligos and plasmids

Oligos and plasmids used in this work

Table S4 crRNAs

CRISPR RNAs used in this work

References

1. Takahashi, J.S. (2017). Transcriptional architecture of the mammalian circadian clock. Nat. Rev. Genet. *18*, 164–179.

2. Gallego, M., and Virshup, D.M. (2007). Post-translational modifications regulate the ticking of the circadian clock. Nat. Rev. Mol. Cell Biol. *8*, 139–148.

3. Gekakis, N., Staknis, D., Nguyen, H.B., Davis, F.C., Wilsbacher, L.D., King, D.P., Takahashi, J.S., and Weitz, C.J. (1998). Role of the CLOCK protein in the mammalian circadian mechanism. Science *280*, 1564–1569.

4. Lee, C., Etchegaray, J.P., Cagampang, F.R., Loudon, A.S., and Reppert, S.M. (2001). Posttranslational mechanisms regulate the mammalian circadian clock. Cell *107*, 855–867.

5. Lowrey, P.L., and Takahashi, J.S. (2011). Genetics of circadian rhythms in Mammalian model organisms. Adv. Genet. *74*, 175–230.

6. Preußner, M., and Heyd, F. (2016). Post-transcriptional control of the mammalian circadian clock: implications for health and disease. Pflugers Arch. *468*, 983–991.

7. Lee, C., Weaver, D.R., and Reppert, S.M. (2004). Direct Association between Mouse PERIOD and CKIε Is Critical for a Functioning Circadian Clock. Mol Cell Biol *24*, 584–594. https://doi.org/10.1128/MCB.24.2.584-594.2004.

8. Aryal, R.P., Kwak, P.B., Tamayo, A.G., Gebert, M., Chiu, P.-L., Walz, T., and Weitz, C.J. (2017). Macromolecular Assemblies of the Mammalian Circadian Clock. Mol Cell *67*, 770-782.e6. https://doi.org/10.1016/j.molcel.2017.07.017.

9.   Cao, X., Yang, Y., Selby, C.P., Liu, Z., and Sancar, A. (2021). Molecular mechanism of the
repressive phase of the mammalian circadian clock. Proc Natl Acad Sci U S A *118*,
e2021174118. https://doi.org/10.1073/pnas.2021174118.

10. Akashi, M., Tsuchiya, Y., Yoshino, T., and Nishida, E. (2002). Control of Intracellular
Dynamics of Mammalian Period Proteins by Casein Kinase I ε (CKIε) and CKIδ in Cultured
Cells. Mol Cell Biol *22*, 1693–1703. https://doi.org/10.1128/MCB.22.6.1693-1703.2002.

11. Eide, E.J., Woolf, M.F., Kang, H., Woolf, P., Hurst, W., Camacho, F., Vielhaber, E.L.,
Giovanni, A., and Virshup, D.M. (2005). Control of mammalian circadian rhythm by
CKIepsilon-regulated proteasome-mediated PER2 degradation. Mol. Cell. Biol. *25*, 2795–
2807.

12. Shirogane, T., Jin, J., Ang, X.L., and Harper, J.W. (2005). SCFbeta-TRCP controls clock-
dependent transcription via casein kinase 1-dependent degradation of the mammalian
period-1 (Per1) protein. J Biol Chem *280*, 26863–26872.
https://doi.org/10.1074/jbc.M502862200.

13. Narasimamurthy, R., Hunt, S.R., Lu, Y., Fustin, J.-M., Okamura, H., Partch, C.L., Forger,
D.B., Kim, J.K., and Virshup, D.M. (2018). CK1δ/ε protein kinase primes the PER2 circadian
phosphoswitch. Proc Natl Acad Sci U S A *115*, 5986–5991.
https://doi.org/10.1073/pnas.1721076115.

14. Philpott, J.M., Freeberg, A.M., Park, J., Lee, K., Ricci, C.G., Hunt, S.R., Narasimamurthy,
R., Segal, D.H., Robles, R., Cai, Y., et al. (2023). PERIOD phosphorylation leads to
feedback inhibition of CK1 activity to control circadian period. Molecular Cell *83*, 1677-
1692.e8. https://doi.org/10.1016/j.molcel.2023.04.019.

15. Philpott, J.M., Narasimamurthy, R., Ricci, C.G., Freeberg, A.M., Hunt, S.R., Yee, L.E.,
Pelofsky, R.S., Tripathi, S., Virshup, D.M., and Partch, C.L. (2020). Casein kinase 1
dynamics underlie substrate selectivity and the PER2 circadian phosphoswitch. eLife *9*,
e52343. https://doi.org/10.7554/eLife.52343.

16. Xu, Y., Toh, K.L., Jones, C.R., Shin, J.-Y., Fu, Y.-H., and Ptácek, L.J. (2007). Modeling of a
human circadian mutation yields insights into clock regulation by PER2. Cell *128*, 59–70.
https://doi.org/10.1016/j.cell.2006.11.043.

17. Jones, C.R., Campbell, S.S., Zone, S.E., Cooper, F., DeSano, A., Murphy, P.J., Jones, B.,
Czajkowski, L., and Ptček, L.J. (1999). Familial advanced sleep-phase syndrome: A short-
period circadian rhythm variant in humans. Nat Med *5*, 1062–1065.
https://doi.org/10.1038/12502.

18. Toh, K.L., Jones, C.R., He, Y., Eide, E.J., Hinz, W.A., Virshup, D.M., Ptácek, L.J., and Fu,
Y.H. (2001). An hPer2 phosphorylation site mutation in familial advanced sleep phase
syndrome. Science *291*, 1040–1043. https://doi.org/10.1126/science.1057499.

19. Jeon, M., Gardner, H.F., Miller, E.A., Deshler, J., and Rougvie, A.E. (1999). Similarity of the
C. elegans Developmental Timing Protein LIN-42 to Circadian Rhythm Proteins. Science
286, 1141–1146. https://doi.org/10.1126/science.286.5442.1141.

20. Lamberti, M.L., Spangler, R.K., Cerdeira, V., Ares, M., Rivollet, L., Ashley, G.E., Coronado,
A.R., Tripathi, S., Spiousas, I., Ward, J.D., et al. (2024). Clock gene homologs *lin-42* and
kin-20 regulate circadian rhythms in *C. elegans*. Sci Rep *14*, 12936.
https://doi.org/10.1038/s41598-024-62303-9.

21. Migliori, M.L., Goya, M.E., Lamberti, M.L., Silva, F., Rota, R., Bénard, C., and Golombek,
D.A. (2023). *Caenorhabditis elegans* as a Promising Model Organism in Chronobiology. J
Biol Rhythms, 07487304221143483. https://doi.org/10.1177/07487304221143483.

22. Tennessen, J.M., Gardner, H.F., Volk, M.L., and Rougvie, A.E. (2006). Novel heterochronic
functions of the *Caenorhabditis elegans* period-related protein LIN-42. Dev Biol *289*, 30–43.
https://doi.org/10.1016/j.ydbio.2005.09.044.

23. Gissendanner, C.R., Crossgrove, K., Kraus, K.A., Maina, C.V., and Sluder, A.E. (2004).
Expression and function of conserved nuclear receptor genes in *Caenorhabditis elegans*.
Dev Biol *266*, 399–416. https://doi.org/10.1016/j.ydbio.2003.10.014.

24. Abrahante, J.E., Miller, E.A., and Rougvie, A.E. (1998). Identification of Heterochronic
Mutants in *Caenorhabditis elegans*: Temporal Misexpression of a Collagen::Green
Fluorescent Protein Fusion Gene. Genetics *149*, 1335–1351.
https://doi.org/10.1093/genetics/149.3.1335.

25. Liu, Z. (1990). Genetic Control of Stage-specific Developmental Events in *C. elegans*.

26. Banerjee, D., Kwok, A., Lin, S.-Y., and Slack, F.J. (2005). Developmental timing in *C.
elegans* is regulated by *kin-20* and *tim-1*, homologs of core circadian clock genes. Dev Cell
8, 287–295. https://doi.org/10.1016/j.devcel.2004.12.006.

27. Hasegawa, K., Saigusa, T., and Tamai, Y. (2005). *Caenorhabditis elegans* opens up new
insights into circadian clock mechanisms. Chronobiol. Int. *22*, 1–19.

28. Kostrouchova, M., Krause, M., Kostrouch, Z., and Rall, J.E. (1998). CHR3: a *Caenorhabditis
elegans* orphan nuclear hormone receptor required for proper epidermal development and
molting. Development (Cambridge, England) *125*, 1617–1626.

29. Ambros, V. (1989). A hierarchy of regulatory genes controls a larva-to-adult developmental
switch in *C. elegans*. Cell *57*, 49–57.

30. Ambros, V., and Horvitz, H.R. (1984). Heterochronic mutants of the nematode
Caenorhabditis elegans. Science *226*, 409–416.

31. McCulloch, K.A., and Rougvie, A.E. (2014). *Caenorhabditis elegans period* homolog *lin-42*
regulates the timing of heterochronic miRNA expression. Proceedings of the National
Academy of Sciences *111*, 15450–15455. https://doi.org/10.1073/pnas.1414856111.

32. Perales, R., King, D.M., Aguirre-Chen, C., and Hammell, C.M. (2014). LIN-42, the
Caenorhabditis elegans PERIOD homolog, Negatively Regulates MicroRNA Transcription.
PLoS Genet *10*, e1004486. https://doi.org/10.1371/journal.pgen.1004486.s008.

33. Van Wynsberghe, P.M., Finnegan, E.F., Stark, T., Angelus, E.P., Homan, K.E., Yeo, G.W.,
and Pasquinelli, A.E. (2014). The Period protein homolog LIN-42 negatively regulates

microRNA biogenesis in *C. elegans*. Dev Biol *390*, 126–135.
https://doi.org/10.1016/j.ydbio.2014.03.017.

34. Lažetić, V., and Fay, D.S. (2017). Molting in *C. elegans*. Worm *6*, 1–20.
https://doi.org/10.1080/21624054.2017.1330246.

35. Monsalve, G.C., Van Buskirk, C., and Frand, A.R. (2011). LIN-42/PERIOD controls cyclical
and developmental progression of *C. elegans* molts. Curr Biol *21*, 2033–2045.
https://doi.org/10.1016/j.cub.2011.10.054.

36. Olmedo, M., Geibel, M., Artal-Sanz, M., and Merrow, M. (2015). A High-Throughput Method
for the Analysis of Larval Developmental Phenotypes in *Caenorhabditis elegans*. Genetics
201, 443. https://doi.org/10.1534/genetics.115.179242.

37. Kinney, B., Sahu, S., Stec, N., Hills-Muckey, K., Adams, D.W., Wang, J., Jaremko, M.,
Joshua-Tor, L., Keil, W., and Hammell, C.M. (2023). A circadian-like gene network
programs the timing and dosage of heterochronic miRNA transcription during *C. elegans*
development. Dev Cell *58*, 2563-2579.e8. https://doi.org/10.1016/j.devcel.2023.08.006.

38. Rhodehouse, K., Cascino, K., Aseltine, L., Padula, A., Weinstein, R., Spina, J.S., Olivero,
C.E., and Van Wynsberghe, P.M. (2018). The Doubletime Homolog KIN-20 Mainly
Regulates *let-7* Independently of Its Effects on the Period Homolog LIN-42 in
Caenorhabditis elegans. G3 (Bethesda) *8*, 2617–2629.
https://doi.org/10.1534/g3.118.200392.

39. Edelman, T.L.B., McCulloch, K.A., Barr, A., Frøkjær-Jensen, C., Jorgensen, E.M., and
Rougvie, A.E. (2016). Analysis of a *lin-42/period* Null Allele Implicates All Three Isoforms in
Regulation of *Caenorhabditis elegans* Molting and Developmental Timing. G3 (Bethesda) *6*,
4077–4086. https://doi.org/10.1534/g3.116.034165.

40. Meeuse, M.W., Hauser, Y.P., Morales Moya, L.J., Hendriks, G., Eglinger, J., Bogaarts, G.,
Tsiairis, C., and Großhans, H. (2020). Developmental function and state transitions of a
gene expression oscillator in *Caenorhabditis elegans*. Mol Syst Biol *16*, e9975.
https://doi.org/10.15252/msb.20209498.

41. Rüegger, S., Miki, T.S., Hess, D., and Großhans, H. (2015). The ribonucleotidyl transferase
USIP-1 acts with SART3 to promote U6 snRNA recycling. Nucleic Acids Res *43*, 3344–
3357. https://doi.org/10.1093/nar/gkv196.

42. Hendriks, G.-J., Gaidatzis, D., Aeschimann, F., and Großhans, H. (2014). Extensive
oscillatory gene expression during *C. elegans* larval development. Mol Cell *53*, 380–392.
https://doi.org/10.1016/j.molcel.2013.12.013.

43. Crosby, P., Goularte, N.F., Sharma, D., Chen, E., Parico, G.C.G., Philpott, J.M., Harold, R.,
Gustafson, C.L., and Partch, C.L. (2023). CHRONO participates in multi-modal repression
of circadian transcriptional complexes. Preprint at bioRxiv,
https://doi.org/10.1101/2022.10.04.510902 https://doi.org/10.1101/2022.10.04.510902.

44. Dahlberg, C.L., Nguyen, E.Z., Goodlett, D., and Kimelman, D. (2009). Interactions between
Casein Kinase Iε (CKIε) and Two Substrates from Disparate Signaling Pathways Reveal

Mechanisms for Substrate-Kinase Specificity. PLOS ONE *4*, e4766.
https://doi.org/10.1371/journal.pone.0004766.

45. LaBella, M.L., Hujber, E.J., Moore, K.A., Rawson, R.L., Merrill, S.A., Allaire, P.D., Ailion, M.,
Hollien, J., Bastiani, M.J., and Jorgensen, E.M. (2020). Casein Kinase 1δ Stabilizes Mature
Axons by Inhibiting Transcription Termination of Ankyrin. Dev Cell *52*, 88-103.e6.
https://doi.org/10.1016/j.devcel.2019.12.005.

46. An, Y., Yuan, B., Xie, P., Gu, Y., Liu, Z., Wang, T., Li, Z., Xu, Y., and Liu, Y. (2022).
Decoupling PER phosphorylation, stability and rhythmic expression from circadian clock
function by abolishing PER-CK1 interaction. Nat Commun *13*, 3991.
https://doi.org/10.1038/s41467-022-31715-4.

47. Lowrey, P.L., Shimomura, K., Antoch, M.P., Yamazaki, S., Zemenides, P.D., Ralph, M.R.,
Menaker, M., and Takahashi, J.S. (2000). Positional syntenic cloning and functional
characterization of the mammalian circadian mutation tau. Science *288*, 483–492.
https://doi.org/10.1126/science.288.5465.483.

48. Xu, Y., Padiath, Q.S., Shapiro, R.E., Jones, C.R., Wu, S.C., Saigoh, N., Saigoh, K., Ptácek,
L.J., and Fu, Y.-H. (2005). Functional consequences of a CKIdelta mutation causing familial
advanced sleep phase syndrome. Nature *434*, 640–644.
https://doi.org/10.1038/nature03453.

49. Marzoll, D., Serrano, F.E., Shostak, A., Schunke, C., Diernfellner, A.C.R., and Brunner, M.
(2022). Casein kinase 1 and disordered clock proteins form functionally equivalent,
phospho-based circadian modules in fungi and mammals. Proc. Natl. Acad. Sci. U. S. A.
119.

50. Johnson, L.N., Noble, M.E., and Owen, D.J. (1996). Active and inactive protein kinases:
structural basis for regulation. Cell *85*, 149–158. https://doi.org/10.1016/s0092-
8674(00)81092-2.

51. Nolen, B., Taylor, S., and Ghosh, G. (2004). Regulation of Protein Kinases: Controlling
Activity through Activation Segment Conformation. Molecular Cell *15*, 661–675.
https://doi.org/10.1016/j.molcel.2004.08.024.

52. Berger, S., Spiri, S., deMello, A., and Hajnal, A. (2021). Microfluidic-based imaging of
complete *Caenorhabditis elegans* larval development. Development *148*, dev199674.
https://doi.org/10.1242/dev.199674.

53. Zheng, B., Larkin, D.W., Albrecht, U., Sun, Z.S., Sage, M., Eichele, G., Lee, C.C., and
Bradley, A. (1999). The *mPer2* gene encodes a functional component of the mammalian
circadian clock. Nature *400*, 169–173. https://doi.org/10.1038/22118.

54. Kim, E.Y., Ko, H.W., Yu, W., Hardin, P.E., and Edery, I. (2007). A DOUBLETIME kinase
binding domain on the *Drosophila* PERIOD protein is essential for its hyperphosphorylation,
transcriptional repression, and circadian clock function. Mol Cell Biol *27*, 5014–5028.
https://doi.org/10.1128/MCB.02339-06.

55. Nawathean, P., Stoleru, D., and Rosbash, M. (2007). A small conserved domain of
Drosophila PERIOD is important for circadian phosphorylation, nuclear localization, and

transcriptional repressor activity. Mol Cell Biol *27*, 5002–5013.
https://doi.org/10.1128/MCB.02338-06.

56. Cullati, S.N., Chaikuad, A., Chen, J.-S., Gebel, J., Tesmer, L., Zhubi, R., Navarrete-Perea,
J., Guillen, R.X., Gygi, S.P., Hummer, G., et al. (2022). Kinase domain autophosphorylation
rewires the activity and substrate specificity of CK1 enzymes. Mol Cell *82*, 2006-2020.e8.
https://doi.org/10.1016/j.molcel.2022.03.005.

57. Gebel, J., Tuppi, M., Chaikuad, A., Hötte, K., Schröder, M., Schulz, L., Löhr, F., Gutfreund,
N., Finke, F., Henrich, E., et al. (2020). p63 uses a switch-like mechanism to set the
threshold for induction of apoptosis. Nat Chem Biol *16*, 1078–1086.
https://doi.org/10.1038/s41589-020-0600-3.

58. Harold, R.L., Tulsian, N.K., Narasimamurthy, R., Yaitanes, N., Ayala Hernandez, M.G., Lee,
H.-W., Crosby, P., Tripathi, S.M., Virshup, D.M., and Partch, C.L. (2024). Isoform-specific C-
terminal phosphorylation drives autoinhibition of Casein kinase 1. Proc Natl Acad Sci U S A
121, e2415567121. https://doi.org/10.1073/pnas.2415567121.

59. Chiou, Y.-Y., Yang, Y., Rashid, N., Ye, R., Selby, C.P., and Sancar, A. (2016). Mammalian
Period represses and de-represses transcription by displacing CLOCK-BMAL1 from
promoters in a Cryptochrome-dependent manner. Proc Natl Acad Sci U S A *113*, E6072–
E6079. https://doi.org/10.1073/pnas.1612917113.

60. Narasimamurthy, R., and Virshup, D.M. (2021). The phosphorylation switch that regulates
ticking of the circadian clock. Mol Cell *81*, 1133–1146.
https://doi.org/10.1016/j.molcel.2021.01.006.

61. Brenner, S. (1974). The genetics of *Caenorhabditis elegans*. Genetics *77*, 71–94.

62. Church, D.L., Guan, K.L., and Lambie, E.J. (1995). Three genes of the MAP kinase
cascade, *mek-2*, *mpk-1/sur-1* and *let-60 ras*, are required for meiotic cell cycle progression
in *Caenorhabditis elegans*. Development *121*, 2525–2535.

63. Ragle, J.M., Levenson, M.T., Clancy, J.C., Vo, A.A., Pham, V., and Ward, J.D. (2022). The
conserved, secreted protease inhibitor MLT-11 is necessary for *C. elegans* molting and
embryogenesis. Preprint at bioRxiv, https://doi.org/10.1101/2022.06.29.498124
https://doi.org/10.1101/2022.06.29.498124.

64. Frøkjaer-Jensen, C., Davis, M.W., Hopkins, C.E., Newman, B.J., Thummel, J.M., Olesen,
S.-P., Grunnet, M., and Jorgensen, E.M. (2008). Single-copy insertion of transgenes in
Caenorhabditis elegans. Nat Genet *40*, 1375–1383. https://doi.org/10.1038/ng.248.

65. Noma, K., Goncharov, A., Ellisman, M.H., and Jin, Y. (2017). Microtubule-dependent
ribosome localization in *C. elegans neurons*. Elife *6*, e26376.
https://doi.org/10.7554/eLife.26376.

66. Frøkjær-Jensen, C., Davis, M.W., Ailion, M., and Jorgensen, E.M. (2012). Improved Mos1-
mediated transgenesis in *C. elegans*. Nat Meth *9*, 117–118.
https://doi.org/10.1038/nmeth.1865.

67. Gudipati, R.K., Braun, K., Gypas, F., Hess, D., Schreier, J., Carl, S.H., Ketting, R.F., and
Großhans, H. (2021). Protease-mediated processing of Argonaute proteins controls small
RNA association. Mol Cell *81*, 2388-2402.e8. https://doi.org/10.1016/j.molcel.2021.03.029.

68. Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T.,
Preibisch, S., Rueden, C., Saalfeld, S., Schmid, B., et al. (2012). Fiji: an open-source
platform for biological-image analysis. Nat Methods *9*, 676–682.
https://doi.org/10.1038/nmeth.2019.

69. Chiu, C.-L., Clack, N., and Community, T.N. (2022). napari: a Python Multi-Dimensional
Image Viewer Platform for the Research Community. Microscopy and Microanalysis *28*,
1576–1577. https://doi.org/10.1017/S1431927622006328.

70. Sheffield, P., Garrard, S., and Derewenda, Z. (1999). Overcoming expression and
purification problems of RhoGDI using a family of "parallel" expression vectors. Protein Expr
Purif *15*, 34–39. https://doi.org/10.1006/prep.1998.1003.

71. Fribourgh, J.L., Srivastava, A., Sandate, C.R., Michael, A.K., Hsu, P.L., Rakers, C., Nguyen,
L.T., Torgrimson, M.R., Parico, G.C.G., Tripathi, S., et al. (2020). Dynamics at the serine
loop underlie differential affinity of cryptochromes for CLOCK:BMAL1 to control circadian
timing. Elife *9*, e55275. https://doi.org/10.7554/eLife.55275.

72. Parico, G.C.G., Perez, I., Fribourgh, J.L., Hernandez, B.N., Lee, H.-W., and Partch, C.L.
(2020). The human CRY1 tail controls circadian timing by regulating its association with
CLOCK:BMAL1. Proc Natl Acad Sci U S A *117*, 27971–27979.
https://doi.org/10.1073/pnas.1920653117.

73. Frankenfield, A.M., Ni, J., Ahmed, M., and Hao, L. (2022). Protein Contaminants Matter:
Building Universal Protein Contaminant Libraries for DDA and DIA Proteomics. J Proteome
Res *21*, 2104–2113. https://doi.org/10.1021/acs.jproteome.2c00145.

1296