Title

Evolutionary and functional analyses reveal a role for the RHIM in tuning RIPK3 activity across vertebrates

Authors

Elizabeth J. Fay[1], Kolya Isterabadi[1], Charles M. Rezanka[1], Jessica Le[1], Matthew D. Daugherty[1]*

Affiliations

[1]Department of Molecular Biology, School of Biological Sciences, University of California, San Diego, La Jolla, CA, 92093

* To whom correspondence should be addressed: Matthew D. Daugherty (mddaugherty@ucsd.edu)

Abstract

Receptor interacting protein kinases (RIPK) RIPK1 and RIPK3 play important roles in diverse innate immune pathways. Despite this, some RIPK1/3-associated proteins are absent in specific vertebrate lineages, suggesting that some RIPK1/3 functions are conserved while others are more evolutionarily labile. Here, we perform comparative evolutionary analyses of RIPK1-5 and associated proteins in vertebrates to identify lineage-specific rapid evolution of RIPK3 and RIPK1 and recurrent loss of RIPK3-associated proteins. Despite this, diverse vertebrate RIPK3 proteins are able to activate NF-κB and cell death in human cells. Additional analyses revealed a striking conservation of the RIP homotypic interaction motif (RHIM) in RIPK3, as well as other human RHIM-containing proteins. Interestingly, diversity in the RIPK3 RHIM can tune activation of NF-κB while retaining the ability to activate cell death. Altogether, these data suggest that NF-κB activation is a core, conserved function of RIPK3, and the RHIM can tailor RIPK3 function to specific needs within and between species.

Introduction

Receptor interacting protein kinase 3 (RIPK3), and the closely related kinase RIPK1, have critical roles in mediating cell death and inflammatory signaling [1]. Downstream of signals from innate immune receptors like Z-DNA binding protein 1 (ZBP1) and tumor necrosis factor receptor 1 (TNFR1), RIPK3 and RIPK1 interact via their shared RIP homotypic interaction motif (RHIM), leading to their autophosphorylation and activation of additional effector proteins to carry out their innate immune functions. For example, RIPK3 can phosphorylate MLKL to activate highly inflammatory programmed necrotic cell death (necroptosis) [2], RIPK1/3 can engage caspase-8 (CASP8) to activate apoptotic caspases [3, 4], or RIPK1/3 can activate the pro-survival and pro-inflammatory transcription factor NF-κB [5, 6]. Several factors are known to impact the outcome of RIPK1/3 activation, such as the activating signal and caspase activity [1, 7]. Altogether, RIPK3 and RIPK1 are central to determining cell fate downstream of various innate immune stimuli.

Like many innate immune proteins, RIPK3 is evolutionarily dynamic in vertebrates. Phylogenomic analyses of a small number of species from different vertebrate clades suggest that RIPK3, and the associated proteins ZBP1 and MLKL, have been lost in some vertebrate lineages including birds, carnivores, and marsupials [8, 9]. In addition, RIPK3, RIPK1, MLKL, and ZBP1 are known to interact with viral proteins [10-15] and have been shown to be evolving under recurrent positive selection in both primates and bats [16, 17], a hallmark of host proteins that directly interact with viral proteins [18]. Despite the known evolutionary divergence of RIPK3 in vertebrates, most of what is known about their activation and regulation is from studies conducted in humans and mice. Even between humans and mice, there are known differences in the roles of specific genes in regulation of NF-κB, suggesting this pathway can be tailored to the distinct environments of different species [19]. These differences likely extend beyond humans and mice, and likely include RIPK1/RIPK3 mediated activation of NF-κB. Furthermore, RIPK1 and RIPK3 are part of a larger family of RIPKs, including RIPK2, RIPK4, and ANKK1/RIPK5, which, with the exception of RIPK5, are known to activate NF-κB in various cellular contexts [20-22]. The

degree to which these other RIP kinases and their interacting partners have diversified
across vertebrates is unknown.

Here, we use evolutionary and functional approaches to characterize
diversification of RIPK3-mediated activation of NF-κB. We identified RIPK1- and RHIM-
mediated activation of NF-κB by diverse vertebrate RIPK3 proteins, as well as distinct
mechanisms of RIPK3-mediated activation of NF-κB. Phylogenetic and phylogenomic
analyses of RIPK1-5 and several key associated proteins revealed the dynamic evolution
of RIPK3 and RIPK1. In addition to observing the loss of RIPK3 and necroptosis-
associated proteins in some vertebrate lineages, we identify changes in regulatory
features—including catalytic sites and CASP cleavage sites—in RIPK3 and RIPK1 in
specific mammalian lineages that may affect their function. Intriguingly, we found that the
RHIM domain of RIPK3 and RIPK1, as well as other RHIM-containing proteins, is highly
conserved in vertebrates and in a non-vertebrate RIPK1 protein. Consistent with this
strong conservation of the RHIM domain, we found that RIPK1 proteins from diverse
vertebrates, and even a non-vertebrate RIPK1, are able to activate NF-κB, suggesting
that activation of this inflammatory response is a conserved function of RIPK3, RIPK1,
and potentially other RHIM-containing proteins in vertebrates. Altogether, these data
suggest that activation of NF-κB is the ancestral and conserved feature of RIPK3 and
RIPK1, while other functions such as necroptosis are more evolutionarily labile and
rapidly evolving, likely as a response to evolutionary pressure from pathogens.

Results

Comparative evolutionary analysis of RIPK1-5

To understand the evolution and diversification of RIPK function, we first wished to
compare the functions of human RIPKs. There are canonically four RIPKs in humans,
RIPK1-4. ANKK1 is a close paralog of RIPK4 and has therefore been denoted as RIPK5.
Additional proteins have also been described as RIPK proteins, including DSTYK,
LRRK1, and LRRK2, but these proteins are phylogenetically distinct and will therefore not
be considered here [22]. The shared domain architecture of RIPK1-5 includes a
conserved N-terminal kinase domain, an extended disordered intermediate domain, and
distinct protein interaction domains at the C-terminus that can include a death domain

(DD), RIP homotypic interaction motif (RHIM), caspase activation and recruitment domain

(CARD), or ankyrin repeats (ANK) (Figure 1A). Human RIPK1-4 activate NF-κB, albeit

through different mechanisms, including differential dependence on endogenous RIPK1

and kinase activity (Supplementary Figure 1-3). Interestingly, despite its close homology

to RIPK4, RIPK5 did not activate NF-κB. Altogether, these data highlight NF-κB as a

shared function of RIPK1-4.

We next wished to perform comparative evolutionary analyses of RIPK1-5 in

vertebrates to determine which RIPKs might be undergoing functional divergence. We

first analyzed RIPK1-5 in four distinct mammalian clades to determine whether there was

evidence of recurrent positive selection, which can be a hallmark of host-pathogen

interactions and can drive functional diversification. Notably, we only found evidence for

positive selection acting on RIPK1 and RIPK3 (Fig 1B, Supplementary files 1-2).

Interestingly, RIPK1 and RIPK3 are not uniformly evolving under positive selection in all

mammalian clades analyzed. For instance, in primates, RIPK3 is evolving under strong

positive selection, where RIPK1 is not, consistent with previous data [16, 17]. In contrast,

ungulate RIPK3 does not show evidence for positive selection whereas ungulate RIPK1

does. These data indicate that among RIPK1-5, only RIPK1 and RIPK3 genes have

undergone recurrent positive selection, and that RIPK1 and RIPK3 may have faced

different selective pressures across distinct mammalian lineages.

In addition to evolving under positive selection in primates and bats [16], RIPK3,

ZBP1, and MLKL are known to be lost in some vertebrate lineages [8]. However, it is

unknown if other RIPK and RIPK-associated proteins are similarly lost. We therefore

performed phylogenomic analysis on 489 vertebrate species for the presence or absence

of RIPK1-5 and several key RIPK-associated proteins (Figure 1C, Supplementary Files 4

and 5). Due to the difficulty in discerning whether a gene absence in an individual species

is due to true gene loss or incomplete genome assembly, we focused on multi-species

patterns of gene loss within and between vertebrate lineages. Given that NF-κB is broadly

present in many vertebrate and non-vertebrate species, including horseshoe crab [23],

Drosophila melanogaster [24], and cnidarians and bivalves (reviewed in [25]), we

analyzed proteins associated with other RIPK functions. We first confirmed the loss of

necroptosis associated proteins that has been previously reported [8]: RIPK3 and ZBP1

in birds; RIPK3, ZBP1, and MLKL in marsupials; MLKL in carnivores. In addition, we found

that ZBP1 is absent in all fish species, suggesting that this protein, and ZBP1-mediated

necroptosis, arose only during tetrapod evolution.

We then analyzed RIPK1 and its associated proteins. RIPK1 is found nearly every

vertebrate genome queried, as well as clear homologs in several non-vertebrate species,

including zebra mussel (*Dreissena polymorpha*). Additional RIPK-like proteins have been

found in other non-vertebrate species, including a RIPK1/2-like protein in *Drosophila* and

other protostomes [8], although these proteins were not considered here due to minimal

sequence similarity to the human RIP kinase domains and our focus on vertebrate RIP

kinases. Likewise, caspase-8 (CASP8), a known regulator of RIPK1, is found in most

tetrapod genomes, although it is absent in fish and sea lamprey. This emergence of

CASP8 in tetrapods coincides with the emergence of ZBP1 in tetrapods. Further

supporting the link between CASP8 and ZBP1, CASP8 is lost in some reptiles that have

also lost ZBP1 (Supplementary Table 1). CASP8 is a known negative regulator of

necroptosis and is important for mitigating ZBP1-induced inflammation [26]. These data

suggest that loss of ZBP1 is sufficient to allow for loss of CASP8. In contrast to CASP8,

its close paralog, CASP10 is present in fish and sea lamprey but displays patterns of

lineage specific loss. For example, we confirmed the loss of CASP10 in some rodents

[27], revealing that suborder Myomorpha ("mouse-like") and Castorimorpha ("beaver-

like") rodents have lost CASP10, whereas Sciuromorpha ("squirrel-like") and

Hystricomorpha ("porcupine-like") rodents have retained CASP10. We also found loss of

CASP10 in all marsupials and several ungulates. Finally, we found that FADD and TRIF

are well conserved in all vertebrates.

We also analyzed the phylogenomic distribution of RIPK2, 4, and 5 and their

associated proteins. RIPK2 and its associated protein NOD1 are found in most

vertebrates [20, 21]. Intriguingly, we observed a second paralog of RIPK2, which we call

RIPK2B, in many vertebrates but lost in therian (live-bearing) mammals (Supplementary

Figure 4). NOD2 is present in lamprey and fish, but absent in birds, reptiles, and some

amphibians, including some of the same species that contain RIPK2B. While RIPK2B

retains both the kinase domain and CARD, there may be distinct regulatory mechanisms,

specifically those related to NOD2, compared to RIPK2. RIPK4 is present in jawed

vertebrates. IRF6, the only known phosphorylation target of RIPK4 [20, 21], emerged prior

to RIPK4 and has been lost in many vertebrate species, particularly in carnivores,

cetaceans, and ungulates. RIPK5, which has no known function and no known interacting

proteins, has been lost in several lineages, including monotremes and lagomorphs.

Overall, our analyses revealed that RIPK1 and RIPK3 are evolutionarily distinct from other

RIP kinases, and the recurrent loss of some RIPK1/3-associated necroptosis proteins

(ZBP1, MLKL, CASP8) and RIPK3 suggests that these proteins may have functions

tailored to the needs of specific vertebrate lineages.

Figure 1. Comparative evolutionary analysis of RIPK1-5. (A) Domain structures of human RIP kinases. RHIM = RIP homotypic interaction motif, DD = death domain, CARD = caspase activation and recruitment domain, ANK = ankyrin repeats. (B) Positive selection analysis of RIPK1-5 in the indicated mammalian order. Input sequences and PAML p-values can be found in Supplementary Files 1 and 2. Images of model species generated using BioRender. (C) Heat amp showing the percentage of species within a clade that have the indicated protein. The clades and the number of species within each clade are indicated on the left. Complete lists of proteins and species in each group can be found in Supplementary Files 4 and 5.

Diverse vertebrate RIPK3 proteins activate NF-κB

Our analyses revealed dynamic evolution of RIPK3 in vertebrates, including

signatures of positive selection in multiple mammalian clades and recurrent loss of RIPK3

and its associated proteins. We hypothesized that this rapid RIPK3 evolution may have

resulted in different functions of RIPK3 in different vertebrate lineages. We therefore

tested the ability of RIPK3 proteins from diverse vertebrates, ranging from 23-63%

similarity to humans (Fig 2A, Supplementary Figure 5), to activate NF-κB in human cells.

To our surprise, among all vertebrates tested, including mammals, reptiles, and sea

lamprey, only pig and sea lamprey RIPK3 failed to activate NF-κB in human cells (Fig

2B). This striking conservation of function indicates that NF-κB activation has remained

intact despite lineage specific RIPK3 divergence and loss of interacting partners. We then

analyzed whether, like human RIPK3 (Supplementary Figure 1C) [28], other vertebrate

RIPK3 proteins require RIPK1 to activate NF-κB by repeating experiments in RIPK1 KO

cells. Interestingly, while mammalian RIPK3 activation is dependent on RIPK1, both lizard

and turtle RIPK3 activate NF-κB independently of RIPK1 (Fig 2B). These results indicate

that the requirements for NF-κB activation by RIPK3 are species-specific.

Finally, we tested whether another function of RIPK3, activation of ZBP1- and

MLKL-dependent cell death (Supplementary Figure 6), is also conserved. Most

mammalian RIPK3 proteins were able to activate cell death using human ZBP1 and MLKL

(Fig 2C). This included pig RIPK3, which is unable to activate NF-κB, suggesting that

activation of cell death and NF-κB by RIPK3 proteins have been separated in a species-

specific manner. Likewise, we observed that cat RIPK3 can activate NF-κB but fails to

activate cell death. Based on this observation, we analyzed the sequences of carnivore

RIPK3 proteins and found that feline RIPK3 have mutated kinase catalytic residue

(Supplementary Figure 7), explaining the loss of cell death activation by cat RIPK3.

Outside of mammals, only lizard RIPK3 was able to activate cell death, despite the ability

of both lizard and turtle RIPK3 to activate NF-κB, further suggesting that these functions

of RIPK3 have distinct, species-specific requirements.

A

| 100 | 59 | 63 | 61 | 24 | 29 | 23 | % identical to human RIPK3 |
| Human | Mouse | Pig | Cat | Lizard | Turtle | Lamprey | |

B

C

Figure 2. Diverse vertebrate RIPK3 proteins activate NF-κB. (A) Percent similarity of RIPK3 from the indicated species compared to humans. (B) RIPK3 proteins were transfected into WT or RIPK1 KO HEK293T cells, along with NF-κB firefly luciferase and control renilla luciferase reporter plasmids (see Materials and Methods), and NF-κB activity was measured at 18h post-transfection. (C) RIPK3 proteins were transfected into HEK293T cells with and without human ZBP1 and MLKL. At 18h post-transfection, cells were stained using the ReadyProbe Cell Viability kit and fluorescence was measured using a plate reader. Species shown are mouse (*Mus musculus*), cat (*Felis catus*), pig (*Sus scrofa*), lizard (*Anolis carolinensis*), turtle (*Chelonia mydas*), and lamprey (*Petromyzon marinus*). Data are representative of 3-5 independent experiments with n=3-6 replicates per group. Data were analyzed using two-way ANOVA with Šidák's multiple comparisons test. ns = not significant, **** = $p < 0.0001$.

Conservation of the RHIM sequence determines RIPK3 activation of NF-κB

The conservation of NF-κB activation by RIPK3 proteins that share as little as 24%
sequence identity and are up to >300 million years diverged [29] suggests some regions
of the protein are likely highly conserved. To identify such regions, we returned to our
positive selection analyses. While we identified rapidly evolving sites throughout the
primate RIPK3, there was a cluster of sites in and around the C-terminal RHIM (Fig 3A
Supplementary File 5), which is a known region of interaction between RIPK1 and RIPK3
and RIPK3 homooligomerization [30, 31]. Intriguingly, despite multiple rapidly evolving
sites within the RHIM, a core tetrad (VQVG) is not rapidly evolving. Similarly, the core
tetrads of bat and carnivore RIPK3 are not rapidly evolving (Fig 3B). Furthermore, the
RIPK3 core RHIM tetrad is highly conserved across vertebrates (Fig 3C). This is despite
loss of other functional motifs in specific lineages, including loss of the RIPK3 catalytic
site in felines (Supplementary Figure 7).

We therefore tested whether this conserved RHIM tetrad was responsible for
activation of NF-kB by diverse vertebrate RIPK3s. We mutated the RIPK3 RHIM core
tetrad (V/I-Q-V/I-G to AAAA) in our panel of vertebrate RIPK3 proteins and tested the
ability of these proteins to activate NF-kB in human cells. Mutation of human, mouse, and
cat RIPK3 RHIMs reduced activation of NF-κB as expected (Fig 3D, Supplementary
Figure 5). Surprisingly, despite activating independently of RIPK1 (Fig 2B), activation of
NF-κB by lizard and turtle RIPK3 remains dependent on the RHIM (Fig 3E, left), and
independent of RIPK3 kinase activity (Supplementary Figure 8-9). These data support
that reptile RIPK3 activates NF-κB through a distinct mechanism from mammalian RIPK3,
but one that is still highly dependent on the RHIM core tetrad. Interestingly, while
analyzing RHIM domains of non-human RIPK3, we observed that the sea lamprey RHIM
diverges from the canonical V/I-Q-V/I-G tetrad and instead contains TQIG (Fig 3C). To
determine if this was responsible for the limited activation of NF-κB by sea lamprey
RIPK3, we mutated this motif to IQIG and observed an increase in NF-κB activation (Fig
3E, right). Altogether, these data indicate that NF-κB activation is a conserved function of
RIPK3 that requires conservation of the core RHIM tetrad, but allows for substantial
sequence divergence elsewhere in the protein.

Figure 3. Conservation of the RHIM sequence determines RIPK3 NF-κB activation. A) Residues evolving under positive selection in primate RIPK3 and the primate RIPK3 RHIM domain mapped on the human sequence. (B) Residues evolving under positive selection in bat and carnivore RIPK3, mapped on to the *Sturnira hondurensis* and *Felis catus* sequences respectively. (C) Alignments of the RIPK3 RHIM across diverse vertebrates. Residue numbers refer to the human sequence. (D-E) Mammalian (D) and non-mammalian (E) RIPK3 proteins were transfected into WT HEK293T cells along with NF-κB firefly luciferase and control renilla luciferase reporter plasmids (see Materials and Methods). NF-κB activity was measured at 18h post-transfection. Species shown are mouse (*Mus musculus*), cat (*Felis catus*), pig (*Sus scrofa*), lizard (*Anolis carolinensis*), turtle (*Chelonia mydas*), and lamprey (*Petromyzon marinus*). Data are representative of 3-5 independent experiments with n=3-6 replicates per group. Data were analyzed using two-way ANOVA with Šidák's multiple comparisons test. ns = not significant, **** = p<0.0001.

NF-κB activation is a shared function of RHIM-containing proteins and can be tuned by
the RHIM

Given that activation of NF-κB by RIPK3 is dependent the RHIM, we hypothesized
that the core RHIM tetrad and activation of NF-κB would be broadly conserved across all
RHIM-containing proteins, including RIPK1, ZBP1 and TRIF. We found that the RIPK1
RHIM is highly conserved in vertebrates and in RIPK1-like protein found in zebra mussel
(Fig 4A). Human RIPK1 is known to activate NF-κB independently of its RHIM through
polyubiquitination of the intermediate domain [32, 33]. However, the strong conservation
of the RHIM domain tetrad sequence (V/I-Q-V/I-G) of RIPK1 across species led us to
hypothesize that RHIM-mediated activation of NF-κB by RIPK1 is also conserved. We
therefore tested a diverse panel of metazoan RIPK1 proteins in our NF-κB assay. To
avoid the effects of both kinase activity and kinase-mediated interactions [34], we
generated RIPK1 C-terminus (RIPK1^{CT}) proteins from diverse species (Supplementary
Figure 10). Both WT and RHIM mutant human RIPK1^{CT} activate NF-κB to the same extent
as full length RIPK1 (Supplementary Figure 11), validating our use of this system to
characterize RIPK1 functions independent from the kinase domain. Diverse vertebrate
RIPK1^{CT} proteins activate NF-κB independent of endogenous RIPK1 (Supplementary
Figure 12), consistent with our hypothesis that NF-κB activation is an ancestral and
conserved function of these proteins. However, like RIPK3, the specific sequence
requirements for RIPK1-mediated NF-κB activation varied by species. Specifically,
mutation of the RHIM domain does not affect activation of NF-κB by mammalian proteins,
including human, mouse, cat, and pig (Figure 4B, left). In contrast, mutation of the RHIM
domain attenuated or completely prevented the ability of non-mammalian (lizard, sea
lamprey, and zebra mussel) RIPK1^{CT} to activate NF-κB (Figure 4B, right).

Other than RIPK1 and RIPK3, only two other human proteins contain a RHIM
domain, TRIF and ZBP1. The TRIF RHIM and the first RHIM of ZBP1 are also highly
conserved in mammals and some other tetrapods, whereas the second RHIM of ZBP1
diverges even within mammals (Fig 4C, Supplementary Figure 13A). We then tested for
RHIM-dependence of NF-κB activation by human TRIF and ZBP1. As described
previously [35], ZBP1 activates NF-κB in a RHIM- and RIPK1-dependent manner (Fig

4D). Consistent with our evolutionary prediction, only the first RHIM tetrad is required for

NF-κB activation by ZBP1 (Fig 4D). Conversely, mutation of the TRIF RHIM domain does

not affect activation of NF-κB, and activation is only moderately reduced in RIPK1 KO

cells (Supplementary Figure 13B-C), likely due to the TIR domain in TRIF that is known

mediate its innate immune signaling [36].

Finally, due to the central role of the core tetrad of the RHIM domain, we tested

the plasticity of RHIM function with regard to sequence divergence. In our phylogenetic

analyses of RIPK3, we identified several species in which the RHIM core tetrad has

diverged, particularly outside of tetrapods, including tetrads that have diverged from the

conserved motif (V/I-Q-V/I-G) at only a single residue (Supplementary Table 2). To test

for functional differences, we inserted these naturally-occurring RHIM tetrads into human

RIPK3 and characterized the ability to activate NF-κB and cell death. We included tetrads

found in rodents, bats, eulipotyphlans, reptiles, and fish (VQFG), amphibians and fish

(LQIG, VQSG), fish (CQIG), and sea lamprey (TQIG). Interestingly, while most tested

variants activate NF-κB compared to the inactive RIPK3 RHIM mutant (AAAA tetrad),

albeit lower than the WT (VQVG) tetrad, TQIG and VQSG did not activate NF-κB (Fig 4E,

Supplementary Figure 14-15). These data suggest that, while there is some plasticity in

the RHIM tetrad, not all residues are functional. All tetrad variants were able to activate

ZBP1-dependent cell death similar to WT RIPK3 (Fig 4F), further revealing the separation

of RIPK3-mediated NF-κB activation from cell death activation. Altogether, these data

suggest that diversity in the RIPK3 RHIM domain may tune activation of NF-κB and tailor

RIPK3 function to the specific needs of species.

Figure 4. NF-κB activation is a shared function of RHIM-containing proteins and can be tuned by
the RHIM. (A) Alignment of RIPK1 RHIM across diverse vertebrates. Residue numbers refer to
the human sequence. (B) Diverse vertebrate RIPK1^CT proteins. were transfected into HEK293T cells
along with NF-κB firefly luciferase and control renilla luciferase reporter plasmids (see Materials and
Methods), and NF-κB activity was measured at 18h post-transfection. (C) Alignment of ZBP1 RHIMs
across diverse vertebrates. Residue numbers refer to the human sequence. (D) Activation of NF-κB by
WT and RHIM mutant ZBP1 proteins. (E) NF-κB activation by human RIPK3 with the indicated RHIM
tetrad variant. (F) Human RIPK3 proteins with the indicated RHIM tetrad variants were transfected into
HEK293T cells with MLKL (gray circles) or MLKL and ZBP1 (orange squares) and viability was
measured at 18h post-transfection. Data are representative of 2-5 independent experiments with n=3-
6 replicates per group. Data were analyzed using two-way ANOVA with Šidák's multiple comparisons
test (A, D) one-way ANOVA with Tukey's multiple comparisons test (E), or two-way ANOVA with
Tukey's multiple comparisons test (F). ns = not significant, **** = p<0.0001.

Discussion

Activation of NF-κB is critical to both the innate and adaptive immune response in

humans and mice [19], and is a deeply conserved pathway in many metazoans [23-25].

Despite this deep conservation of NF-κB signaling, differences in the function and

regulation of NF-κB-associated proteins between mice and humans have been identified

[19], suggesting that this functionally well-conserved pathway can be adapted to different

species. These data indicate that broader phylogenetic analysis of the genes and

functions that are associated with NF-κB signaling are needed to understand immune

responses across species, including those that harbor pathogens that pose a zoonotic

threat to humans.

Here, we apply broad phylogenetic and functional sampling to the RIP kinase family

of NF-κB activators. Using a phylogenomic approach, we identified widespread

conservation of RIPK1, RIPK2, and RIPK4 and their associated proteins within

vertebrates. We also show conservation of activation of NF-κB signaling by RIPK1 and

RIPK3 across diverse vertebrate species that is largely reliant on the highly conserved

core tetrad of the RHIM domains found in both proteins. This striking conservation of

RIPK1- and RIPK3-mediated activation of NF-κB, from species that span >500 million

years of vertebrate evolution, underscores the core functionality of RIP kinases as NF-κB

activating proteins. Despite this conservation of NF-κB activation, we also observed

lineage specific changes in RIPK presence or mechanisms of NF-κB activation. For

instance, we observed recurrent gene loss of RIPK3 and RIPK5 throughout the vertebrate

phylogeny, and discovered a second paralog of RIPK2 in many non-mammalian

vertebrate species (RIPK2B). In addition, although RIPK3-mediated activation of NF-κB

is highly dependent on RIPK1 across diverse mammals, RIPK3 activation of NF-κB in

reptiles is independent of RIPK1. Activation of NF-κB is part of the antiviral immune

response and is known to be antagonized by viruses across a range of species, including

vertebrates and non-vertebrates [37-39]. This antagonism could result in adaptation of

NF-κB-associated proteins, including RIP kinases, leading to lineage-specific

mechanisms of activation that are tailored to the specific contexts within and between

species.

The most striking differences in RIPK-associated genes and functions are those
involved in necroptosis. Necroptosis is highly inflammatory and the associated proteins
must therefore be tightly regulated, as it could be a cause of immunopathology during
infection. Despite this, necroptosis has been shown to be important for the response to
several viruses [40-42]. It is hypothesized that necroptosis, defined by RIPK3-mediated
activation of MLKL, arose in vertebrates [8, 43]. Interestingly, we find that two critical
regulators of necroptosis, CASP8 and ZBP1, are only found in tetrapods, suggestive of
an additional innovation within the necroptosis pathway during the divergence of
tetrapods. Moreover, consistent with previous observations [8, 9], we observe a large
number of necroptosis-associated gene losses, while also discovering important
functional motifs in necroptosis proteins in distinct tetrapod lineages. These data support
a model where necroptosis has been constructed as an important facet of the innate
immune response in some tetrapods, but is not universal and has been deconstructed in
species where the immunopathologic effects outweigh the benefits.

There have been several hypotheses to explain the loss of necroptosis associated
proteins in specific vertebrate lineages, such as diet (*e.g.,* carnivores) or behavior (*e.g.,*
torpor in reptiles) [8]. While the cause of a specific instance of gene loss is difficult to
identify, it is clear that gene loss can provide an evolutionary tool for adaptation [44].For
example, caspases, which are key regulators of both immunologically silent apoptosis
and inflammatory pyroptosis, differ greatly across mammals, with lineage-specific
caspase repertoires including both loss and gain of novel caspases [27]. Even within a
single mammalian order, homologous caspases have been found to serve different
functions in response to pathogens [45]. This type of lineage-specific innovation likely
exists for necroptosis, and gene loss could be a cause or a result of adaptations. Indeed,
differences in MLKL activation by RIPK3 across species have been identified [46], which
may be the result of additional diverse regulatory mechanisms for necroptosis even
amongst species with an intact pathway. Strikingly, despite the lack of a complete
necroptosis pathway in some lineages (cat, turtle, lamprey), our functional data indicate
that RIPK3 retains the ability to activate NF-κB. Activation of NF-κB could therefore
represent the ancestral function of RIPK3, and it has been co-opted to activate
necroptosis in vertebrates.

The RHIM core tetrad was previously known to be conserved between humans and
mice [47]. Here, we expand this analysis and identify striking conservation of the RIPK3,
RIPK1, ZBP1, and TRIF RHIM domains across most vertebrates. While activation of NF-
κB by all tested non-human RIPK3 proteins was highly dependent on the RHIM domain,
only non-mammalian RIPK1^{CT} proteins required the RHIM for maximal NF-κB activation.
These species (lizard, lamprey, and zebra mussel) also lack ZBP1, with lamprey and
zebra mussel predating the emergence of ZBP1 in tetrapods. Necroptosis may have
driven evolution of RHIM-independent activation of NF-κB by RIPK1. This may have also
shaped RHIM-dependent functions of RIPK3 across vertebrates, as the RHIM tetrad is
much less conserved in fish compared to tetrapods, and insertion of these tetrad variants
into human RIPK3 alters its function. Virus antagonism has also likely played a role in the
evolution of RIPK3 and RIPK1 function. For example, vaccinia virus E3 prevents Z-DNA
sensing by ZBP1 [40], human cytomegalovirus UL36 promotes MLKL degradation, and
poxviruses encode an MLKL homolog to prevent RIPK3-mediated activation of host MLKL
[48]. Conservation of the RHIM domain may be a way to retain some RIPK3/RIPK1 innate
immune functions in the face of such antagonism. However, several viruses have taken
advantage of this conservation and encode RHIM-containing proteins to prevent RHIM-
dependent functions [43, 47]. Intriguingly, several of these mechanisms are known to be
species-specific and differ between humans and mice [13, 14]. These data support a
model where viruses and host necroptosis are engaged in a molecular arms race that has
shaped the function of RIPK3 and RIPK1 across vertebrate species.

Altogether, our combined phylogenetic and functional approaches reveal both
conservation and a striking lack of conservation in vertebrate RIPKs and their associated
proteins and functions. Such characteristics are common hallmarks of pathways that are
required for the innate immune response to pathogens, but are under constant
evolutionary pressure to innovate to avoid pathogen antagonism. The strong conservation
of NF-κB activation by RIPK1 across species is consistent with the model in which RIPK1
function is central to determining cell fate downstream of TNF and TLR signaling [49].
Moreover, because some pathogens have evolved strategies to inhibit CASP8, RIPK3-
and RIPK1-mediated necroptosis is activated as a secondary cell death strategy when
CASP8 is inhibited [50]. This places both RIPK3 and RIPK1, critical regulators of

necroptosis, at the center of determining cell fate during pathogen infection. The
divergence of RIPK3 and RIPK1 across vertebrates identified here, as well as the
previous finding that these proteins are evolving under positive selection [9, 16, 17],
reveals that pathogens are likely the driving force behind the lineage-specific regulatory
mechanisms or functions of these proteins across species that we observe here. Thus,
the functional divergence identified in our work highlights the evolutionary innovation that
can arise in innate signaling pathways across diverse species, and underscores the
importance of considering natural diversity when characterizing the innate immune
response to pathogens.

Materials & Methods

Positive selection analysis

Positive selection analysis was performed using three independent methods as we have
done previously [51, 52]. Nucleotide sequences from the indicated mammalian clade that
aligned to full-length human RIPK1-5 were downloaded from NCBI and aligned using
ClustalOmega [53]. Maximum likelihood (ML) tests were performed with codeml in the
PAML software suite [54]. For PAML, aligned sequences were subjected to ML tests
using NS sites models disallowing (M7) or allowing (M8) positive selection. The reported
p-values were calculated using a chi-squared test on twice the difference of the log
likelihood (lnL) values between the two models using 2 degrees of freedom. We confirmed
convergence of lnL values by performing each analysis using two starting omega (dN/dS)
values (0.4 and 1.5). Codons evolving under positive selection from PAML analyses have
a posterior probability greater than 0.90 using Native Empirical Bayes (NEB) and Bayes
Empirical Bayes (BEB) analysis and the F61 codon frequency model. The same
nucleotide alignments were used as input for FUBAR [55] and MEME [56] using the
DataMonkey server [57]. In both cases, default parameters were used and codons with a
signature of positive selection with a p-value of <0.01 are reported. Accession numbers
for input sequences, PAML p-values, and sites identified can be found in Supplementary
Files 1-3.

Gene loss analysis

Using the aligned and filtered vertebrate RIPK sequences (4,550 sequences in

total, see 'Phylogenetic analysis' section), a tree was built using FastTree [58, 59]

implemented using Geneious. Using this tree, we were able to identify and extract *bona*

fide RIP kinases and remove similar, non-RIP kinases. The extracted sequences were

again aligned using Clustal Omega and filtered using the criteria above (See

'Phylogenetic analysis' section). A second tree was generated using FastTree. From this

tree, we extracted individual sequence lists for RIPK1-5 to determine which vertebrate

species did or did not have each RIPK. This method was repeated for CASP2 and

CASP8-10, ZBP1, MLKL, TRIF, FADD, NOD 1 and 2, and IRF6. The list of species and

lists of accession numbers for each protein can be found in Supplementary Files 4 and 5.

Plasmids and constructs

Coding sequences for human RIPK1 (Addgene #78834), human RIPK2

(ORFeome ID #4886), human RIPK3 (Addgene #78804), mouse RIPK1 (Addgene

#115341), and mouse RIPK3 (Addgene #78805) were cloned into apcDNA5/FRT/TO

backbone (Invitrogen, Carlsbad, CA) with an N-terminal V5 tag and linker using Gibson

Assembly (New England Biolabs, Ipswich, MA). Human RIPK4 (NP_065690.2) and

RIPK5 (NP_848605.1), and non-human RIPK1 C-termini (cat, XP_023109490.2; pig,

XP_003128209.1; chicken, NP_989733.3; lizard, XP_003224434.1; lamprey,

XP_032813622.1; zebra mussel, XP_052232845.1) and RIPK3 (cat, XP_003987615.3;

pig, XP_001927459.3; lizard, XP_003223896.2; turtle, XP_037771912.1; lamprey,

XP_032816533.1) homologues were ordered from Twist Biosciences (San Francisco,

CA, USA) and were cloned along with RHIM mutants targeting the core tetrad into the

pcDNA5/FRT/TO backbone (Invitrogen, Carlsbad, CA) with an N-terminal V5 tag and

linker using Gibson Assembly (New England Biolabs, Ipswich, MA). Kinase mutants for

human proteins RIPK1 (D138N), RIPK2 (D146N), RIPK3 (D142N), RIPK4 (D143N), and

RIPK5 (D145N), and RHIM mutants for RIPK1 (IQIG542AAAA) and RIPK3

(VQVG461AAAA) were generated using Gibson Assembly.

Cell culture and transient transfections

HEK293T cells (obtained from ATCC) and the generated HEK293T RIPK1 KO cells
were maintained at a low passage number to maintain less than one year since
purchase, acquisition, or generation. Both cell lines were grown in complete media
containing DMEM (Gibco, Carlsbad, CA), 10% FBS (Peak Serum, Wellington, CO), and
1% penicillin/streptomycin (Gibco, Carlsbad, CA). A day prior to transfection, HEK293T
and RIPK1 KO cells were seeded into 24-well plates with 500 µL complete media or 96-
well plates with 80 µL complete media. Cells were transiently transfected with 500 ng
total DNA and 1.5 µL of TransIT-X2 (Mirus Bio, Madison, WI) in 100 µL Opti-MEM
(Gibco, Carlsbad, CA) for a 24-well plate or 100 ng DNA and 0.3 µL TransIT-X2 in 10
µL Opti-MEM for a 96-well plate. DNA and TransIT-X2 were incubated at room
temperature for 25-30 minutes, then added dropwise to the appropriate well. Cells were
harvested or analyzed at 18-22 hours post-transfection.
488
Generation of knockout cell lines
RIPK1 knockout HEK293T cells were generated using CRISPR/Cas9 as
previously described [52, 60]. Briefly, plasmids were generated order to produce
lentivirus-like particles containing the CRISPR/Cas9 machinery and guide RNA targeting
exon 5 (ENSE00003586162) of RIPK1. The protocol for the molecular cloning of this
plasmid was adapted from Feng Zhang (Sanjana et al., 2014) using the transfer plasmid
pLB-Cas9 (gift from Feng Zhang, Addgene plasmid # 52962). We designed the gRNA
target sequence using the web tool CHOPCHOP (Labun et al., 2016), available at
https://chopchop.cbu.uib.no/, and synthesized oligonucleotides from Integrated DNA
Technologies (San Diego, CA). The synthesized oligonucleotide pair was phosphorylated
and annealed using T4 Polynucleotide Kinase (NEB M0201S) with T4 Ligation Buffer
(NEB). Duplexed oligonucleotides were ligated into dephosphorylated and BsmBI-
digested pLB-Cas9 using the Quick Ligase kit (NEB M2200S) to generate transfer
plasmid with RIPK1 guide sequence. To generate lentivirus-like particles, this transfer
plasmid was transfected alongside two packaging plasmids, pMD2.G (gift from Didier
Trono, Addgene plasmid # 12259) and psPAX2 (gift from Didier Trono, Addgene plasmid
12260), into HEK293T cells with a 1:1:1 stoichiometry. Forty-eight hours post-
transfection supernatant was harvested and syringe-filtered (0.45µm). Supernatant

containing sgRNA-encoding lentivirus-like particles was then used to transduce HEK293T

cells. Transduced cells were cultured in growth media for 48 hours, then cultured in

growth media supplemented with 1 µg/ml puromycin for 72 hours. Using limiting dilution

in 96-well plates, the monoclonal cell line was then obtained. RIPK1 knockout was

confirmed by Sanger sequencing and by western blot using an α-RIPK1 antibody.

NF-κB and IFN luciferase reporter assays

To quantify NF-κB and IFN activation, we used the Dual-Glo Luciferase Assay

System E2920 (Promega, USA). WT or RIPK1 KO HEK293T cells were seeded in a white

96-well plate and transfected with firefly luciferase fused to either the NF-κB response

element (pGL4.32, Promega) or the human IFN-beta promoter (IFN-Beta-pGL3, Addgene

#102597), renilla luciferase fused to herpes simplex virus thymidine kinase promoter

(pTK-Renilla, Thermo Scientific), and the indicated RIPK. Firefly luciferase was used as

the primary reporter with renilla luciferase being a normalization control. Eighteen to 24

hours after transfection, Dual-Glo Luciferase Assay Reagent was added to each

transfected well as well as 3-6 untransfected wells to serve as a negative control.

Following a 10-minute incubation, the firefly luciferase signal was measured using a

BioTek Cytation imaging reader with Gen5 software (Agilent Technologies, San Diego,

CA). Then, Dual-Glo Stop & Glo reagent was added to each well. Following a 10-minute

incubation, the renilla luciferase signal was measured. The background luminescence

signal from the buffer-treated untransfected conditions was subtracted from other

conditions, firefly values were normalized to renilla values, and all samples were

normalized to the empty vector control condition. Values are reported as NF-κB activity

(firefly/renilla).

Cell death assay

Cell viability was measured using the ReadyProbes Cell Viability Imaging Kit, Blue/Green

(ThermoFisher Scientific). HEK293T cells were seeded in 80 µL of DMEM+/+ (See "Cell

culture and transient transfections") in a clear 96-well plate and transfected with the

indicated plasmids or a vector control. Eighteen hours post-transfection, cells were

stained using NucBlue Live reagent, which stains all nuclei, and NucGreen Dead reagent,

which stains only dead cells. A 2x concentrated mix (4 drops/mL of each dye) was made
and added to wells to 1x concentration. Cells were incubated for 20 minutes at room
temperature. Fluorescence for NucBlue (DAPI, excitation 377/20, emission 447/20) and
NucGreen (GFP, excitation 469/20, emission 525/20) were read using a BioTek Cytation
imaging reader with Gen5 software (Agilent Technologies, San Diego, CA). Unstained
wells were read and used as controls for background fluorescence. GFP values were
normalized to DAPI values, and all samples were normalized to the vector control
condition. Values are reported as Cytotoxicity (GFP/DAPI).
546
Immunoblotting and antibodies
Eighteen to 22 hours post-transfection, cells were washed with 1X PBS and lysed with
boiling 1x NuPAGE LDS sample buffer containing 5% β-mercaptoethanol at room
temperature for 5 minutes and then at 98°C for 7-10 minutes. Lysates were separated
by SDS-PAGE (4-15% Bis-tris gel; Life Technologies, San Diego, CA) with 1X MOPS
buffer (Life Technologies, San Diego, CA). Proteins were transferred onto a
nitrocellulose membrane (Life Technologies, San Diego, CA) and blocked with PBS–T
containing 5% bovine serum albumin (BSA) (Spectrum, New Brunswick, NJ).
Membranes were incubated with the indicated rabbit primary antibodies diluted with 5%
BSA and PBS-T at 1:1000 overnight at 4°C (α-V5 clone D3H8Q, α-GAPDH clone
14C10, α-RIPK1 clone E8S7U XP, α-RIPK2 clone D10B11, α-RIPK3 clone E1Z1D, α-
RIPK4 #12636; Cell Signaling Technology, Danvers, MA). Membranes were rinsed
three times with PBS-T, incubated with HRP-conjugated rabbit secondary antibody
diluted at 1:10,000 with 5% BSA and PBS-T for 30 minutes at room temperature, and
developed with SuperSignal West Pico PLUS Chemiluminescent Substrate (Thermo
Fisher Scientific, Carlsbad, CA).
563
564 *Statistical analysis*
565 Statistical analyses were completed using GraphPad Prism 10 software. Tests were
566 performed as indicated. Error bars were calculated using SEM.
567
568 **Acknowledgements**

This work was supported by National Institutes of Health grant R35 GM133633 (to MDD); Pew Biomedical Scholars Program (to MDD); Burroughs Wellcome Investigators in the Pathogenesis of Infectious Disease Program (to MDD); and BrightSpinnaker Fellowship (to CMR). We thank all members of the Daugherty lab for helpful discussions and Scott Biering, Ryan Langlois, and Patrick Mitchell for their critical reading of this manuscript.

Conflict of Interest

The authors declare that they have no conflict of interest.

References

1. Newton, K., *RIPK1 and RIPK3: critical regulators of inflammation and cell death.* Trends Cell Biol, 2015. **25**(6): p. 347-53.
2. Sun, L., et al., *Mixed lineage kinase domain-like protein mediates necrosis signaling downstream of RIP3 kinase.* Cell, 2012. **148**(1-2): p. 213-27.
3. Wang, L., F. Du, and X. Wang, *TNF-alpha induces two distinct caspase-8 activation pathways.* Cell, 2008. **133**(4): p. 693-703.
4. Mandal, P., et al., *RIP3 induces apoptosis independent of pronecrotic kinase activity.* Mol Cell, 2014. **56**(4): p. 481-95.
5. Meylan, E., et al., *RIP1 is an essential mediator of Toll-like receptor 3-induced NF-kappa B activation.* Nat Immunol, 2004. **5**(5): p. 503-7.
6. Moriwaki, K., et al., *The necroptosis adaptor RIPK3 promotes injury-induced cytokine expression and tissue repair.* Immunity, 2014. **41**(4): p. 567-78.
7. Orozco, S. and A. Oberst, *RIPK3 in cell death and inflammation: the good, the bad, and the ugly.* Immunol Rev, 2017. **277**(1): p. 102-112.
8. Dondelinger, Y., et al., *An evolutionary perspective on the necroptotic pathway.* Trends Cell Biol, 2016. **26**(10): p. 721-732.
9. Agueda-Pinto, A., et al., *Convergent Loss of the Necroptosis Pathway in Disparate Mammalian Lineages Shapes Viruses Countermeasures.* Front Immunol, 2021. **12**: p. 747737.
10. Harris, K.G., et al., *RIP3 Regulates Autophagy and Promotes Coxsackievirus B3 Infection of Intestinal Epithelial Cells.* Cell Host Microbe, 2015. **18**(2): p. 221-32.
11. Croft, S.N., E.J. Walker, and R. Ghildyal, *Human Rhinovirus 3C protease cleaves RIPK1, concurrent with caspase 8 activation.* Sci Rep, 2018. **8**(1): p. 1569.
12. Wagner, R.N., J.C. Reed, and S.K. Chanda, *HIV-1 protease cleaves the serine-threonine kinases RIPK1 and RIPK2.* Retrovirology, 2015. **12**: p. 74.
13. Muscolino, E., et al., *Species-Specific Inhibition of Necroptosis by HCMV UL36.* Viruses, 2021. **13**(11).
14. Huang, Z., et al., *RIP1/RIP3 binding to HSV-1 ICP6 initiates necroptosis to restrict virus propagation in mice.* Cell Host Microbe, 2015. **17**(2): p. 229-42.

15. Liu, X., et al., *Epstein-Barr virus encoded latent membrane protein 1 suppresses necroptosis through targeting RIPK1/3 ubiquitination.* Cell Death Dis, 2018. **9**(2): p. 53.

16. Palmer, S., et al., *Evolutionary profile for (host and viral) MLKL indicates its activities as a battlefront for extensive counteradaptation.* Mol Biol Evol, 2021. **38**(12): p. 5405-522.

17. Cariou, M., et al., *Distinct evolutionary trajectories of SARS-CoV-2-interacting proteins in bats and primates identify important host determinants of COVID-19.* Proc Natl Acad Sci U S A, 2022. **119**(35): p. e2206610119.

18. Daugherty, M.D. and H.S. Malik, *Rules of engagement: molecular insights from host-virus arms races.* Annu Rev Genet, 2012. **46**: p. 677-700.

19. Zhang, Q., M.J. Lenardo, and D. Baltimore, *30 Years of NF-kappaB: A Blossoming of Relevance to Human Pathobiology.* Cell, 2017. **168**(1-2): p. 37-57.

20. Cuny, G.D. and A. Degterev, *RIPK protein kinase family: Atypical lives of typical kinases.* Semin Cell Dev Biol, 2021. **109**: p. 96-105.

21. Eng, V.V., M.A. Wemyss, and J.S. Pearson, *The diverse roles of RIP kinases in host-pathogen interactions.* Semin Cell Dev Biol, 2021. **109**: p. 125-143.

22. Lv, S., et al., *Comparative and evolutionary analysis of RIP kinases in immune responses.* Front Genet, 2022. **13**: p. 796291.

23. Wang, X.W., et al., *Evidence for the ancient origin of the NF-kappaB/IkappaB cascade: its archaic role in pathogen infection and immunity.* Proc Natl Acad Sci U S A, 2006. **103**(11): p. 4204-9.

24. Hetru, C. and J.A. Hoffmann, *NF-kappaB in the immune response of Drosophila.* Cold Spring Harb Perspect Biol, 2009. **1**(6): p. a000232.

25. Williams, L.M. and T.D. Gilmore, *Looking Down on NF-kappaB.* Mol Cell Biol, 2020. **40**(15).

26. Rodriguez, D.A., et al., *Caspase-8 and FADD prevent spontaneous ZBP1 expression and necroptosis.* Proc Natl Acad Sci U S A, 2022. **119**(41): p. e2207240119.

27. Eckhart, L., et al., *Identification of novel mammalian caspases reveals an important role of gene loss in shaping the human caspase repertoire.* Mol Biol Evol, 2008. **25**(5): p. 831-41.

28. Yatim, N., et al., *RIPK1 and NF-kappaB signaling in dying cells determines cross-priming of CD8(+) T cells.* Science, 2015. **350**(6258): p. 328-34.

29. Benton, M.J. and P.C. Donoghue, *Paleontological evidence to date the tree of life.* Mol Biol Evol, 2007. **24**(1): p. 26-53.

30. Wu, X., et al., *The structure of a minimum amyloid fibril core formed by necroptosis-mediating RHIM of human RIPK3.* Proc Natl Acad Sci U S A, 2021. **118**(14).

31. Sun, X., et al., *Identification of a novel homotypic interaction motif required for the phosphorylation of receptor-interacting protein (RIP) by RIP3.* J Biol Chem, 2002. **277**(11): p. 9505-11.

32. Li, H., et al., *Ubiquitination of RIP is required for tumor necrosis factor alpha-induced NF-kappaB activation.* J Biol Chem, 2006. **281**(19): p. 13636-13643.

33. Ea, C.K., et al., *Activation of IKK by TNFalpha requires site-specific ubiquitination of RIP1 and polyubiquitin binding by NEMO.* Mol Cell, 2006. **22**(2): p. 245-57.

654   34.   Delanghe, T., Y. Dondelinger, and M.J.M. Bertrand, *RIPK1 Kinase-Dependent*
655   *Death: A Symphony of Phosphorylation Events.* Trends Cell Biol, 2020. **30**(3): p.
656   189-200.
657   35.   Rebsamen, M., et al., *DAI/ZBP1 recruits RIP1 and RIP3 through RIP homotypic*
658   *interaction motifs to activate NF-kappaB.* EMBO Rep, 2009. **10**(8): p. 916-22.
659   36.   Nanson, J.D., B. Kobe, and T. Ve, *Death, TIR, and RHIM: Self-assembling*
660   *domains involved in innate immunity and cell-death signaling.* J Leukoc Biol, 2019.
661   **105**(2): p. 363-375.
662   37.   Albarnaz, J.D., et al., *Molecular mimicry of NF-kappaB by vaccinia virus protein*
663   *enables selective inhibition of antiviral responses.* Nat Microbiol, 2022. **7**(1): p.
664   154-168.
665   38.   Gonzalez Lopez Ledesma, M.M., et al., *Dengue virus NS5 degrades ERC1 during*
666   *infection to antagonize NF-kB activation.* Proc Natl Acad Sci U S A, 2023. **120**(23):
667   p. e2220005120.
668   39.   Palmer, W.H., et al., *Induction and Suppression of NF-kappaB Signalling by a DNA*
669   *Virus of Drosophila.* J Virol, 2019. **93**(3).
670   40.   Koehler, H., et al., *Vaccinia virus E3 prevents sensing of Z-RNA to block ZBP1-*
671   *dependent necroptosis.* Cell Host Microbe, 2021. **29**(8): p. 1266-1276 e5.
672   41.   Balachandran, S. and G.F. Rall, *Benefits and Perils of Necroptosis in Influenza*
673   *Virus Infection.* J Virol, 2020. **94**(9).
674   42.   Shubina, M., et al., *Necroptosis restricts influenza A virus as a stand-alone cell*
675   *death mechanism.* J Exp Med, 2020. **217**(11).
676   43.   Tummers, B. and D.R. Green, *The evolution of regulated cell death pathways in*
677   *animals and their evasion by pathogens.* Physiol Rev, 2022. **102**(1): p. 411-454.
678   44.   Albalat, R. and C. Canestro, *Evolution by gene loss.* Nat Rev Genet, 2016. **17**(7):
679   p. 379-91.
680   45.   Devant, P., A. Cao, and J.C. Kagan, *Evolution-inspired redesign of the LPS*
681   *receptor caspase-4 into an interleukin-1beta converting enzyme.* Sci Immunol,
682   2021. **6**(62).
683   46.   Tanzer, M.C., et al., *Evolutionary divergence of the necroptosis effector MLKL.* Cell
684   Death Differ, 2016. **23**(7): p. 1185-97.
685   47.   Riebeling, T., U. Kunzendorf, and S. Krautwald, *The role of RHIM in necroptosis.*
686   Biochem Soc Trans, 2022. **50**(4): p. 1197-1205.
687   48.   Petrie, E.J., et al., *Viral MLKL Homologs Subvert Necroptotic Cell Death by*
688   *Sequestering Cellular RIPK3.* Cell Rep, 2019. **28**(13): p. 3309-3319 e5.
689   49.   Clucas, J. and P. Meier, *Roles of RIPK1 as a stress sentinel coordinating cell*
690   *survival and immunogenic cell death.* Nat Rev Mol Cell Biol, 2023. **24**(11): p. 835-
691   852.
692   50.   Remick, B.C., M.M. Gaidt, and R.E. Vance, *Effector-Triggered Immunity.* Annu
693   Rev Immunol, 2023. **41**: p. 453-481.
694   51.   Tsu, B.V., et al., *Host-specific sensing of coronaviruses and picornaviruses by the*
695   *CARD8 inflammasome.* PLoS Biol, 2023. **21**(6): p. e3002144.
696   52.   Stevens, D.A., et al., *Antiviral function and viral antagonism of the rapidly evolving*
697   *dynein activating adaptor NINL.* Elife, 2022. **11**.
698   53.   Sievers, F., et al., *Fast, scalable generation of high-quality protein multiple*
699   *sequence alignments using Clustal Omega.* Mol Syst Biol, 2011. **7**: p. 539.

700 54. Yang, Z., *PAML 4: phylogenetic analysis by maximum likelihood.* Mol Biol Evol,
701   2007. **24**(8): p. 1586-91.
702 55. Murrell, B., et al., *FUBAR: a fast, unconstrained bayesian approximation for*
703   *inferring selection.* Mol Biol Evol, 2013. **30**(5): p. 1196-205.
704 56. Murrell, B., et al., *Detecting individual sites subject to episodic diversifying*
705   *selection.* PLoS Genet, 2012. **8**(7): p. e1002764.
706 57. Weaver, S., et al., *Datamonkey 2.0: A Modern Web Application for Characterizing*
707   *Selective and Other Evolutionary Processes.* Mol Biol Evol, 2018. **35**(3): p. 773-
708   777.
709 58. Price, M.N., P.S. Dehal, and A.P. Arkin, *FastTree: computing large minimum*
710   *evolution trees with profiles instead of a distance matrix.* Mol Biol Evol, 2009. **26**(7):
711   p. 1641-50.
712 59. Price, M.N., P.S. Dehal, and A.P. Arkin, *FastTree 2--approximately maximum-*
713   *likelihood trees for large alignments.* PLoS One, 2010. **5**(3): p. e9490.
714 60. Tsu, B.V., et al., *Diverse viral proteases activate the NLRP1 inflammasome.* Elife,
715   2021. **10**.
716