Title

Evolutionary and functional analyses reveal conserved and non-conserved functions of vertebrate RIP kinases

Authors

Elizabeth J. Fay[1], Kolya Isterabadi[1], Charles M. Rezanka[1], Jessica Le[1], Matthew D. Daugherty[1]*

Affiliations

[1]Department of Molecular Biology, School of Biological Sciences, University of California, San Diego, La Jolla, CA, 92093

* To whom correspondence should be addressed: Matthew D. Daugherty (mddaugherty@ucsd.edu)

Abstract

Receptor interacting protein kinases (RIPKs) play important roles in diverse innate immune signaling pathways. Despite this, some RIPK-associated proteins are absent in specific vertebrate lineages, suggesting that some aspects of RIPK protein function are well conserved while other functions are more evolutionarily labile. Here, we perform phylogenomic analyses of RIPK1-5 and associated proteins in vertebrates to identify RIPK functions that are well conserved, such as NF-κB signaling, and functions that have undergone pervasive gene loss, such as necroptosis. We further use functional approaches to characterize RIPK1-5 in humans and RIPK1 and RIPK3 across diverse species. Consistent with our phylogenomic inferences, we identify NF-κB as a core function of RIP kinases in humans, albeit through diverse mechanisms. We additionally identified conservation of the RIP homotypic interaction motif in RIPK1 and RIPK3, despite loss of key mediators of RIPK1/3 function in vertebrates. The preservation of this motif has also led to the conservation of NF-κB activation by RIPK1 and RIPK3, with some variation in regulatory mechanisms, from diverse vertebrate and non-vertebrate species. Altogether, these data suggest that NF-κB activation is a core, conserved

32 function of RIP kinases, while additional RIPK functions have been tailored to specific
33 needs within and between species.
34
Introduction
Receptor-interacting protein kinases 1-5 (RIPK1-5) are a family of intracellular
signaling proteins with diverse and overlapping roles in innate immunity [1, 2]. These
proteins share a similar domain structure marked by a homologous N-terminal kinase
domain, an intermediate domain, and a C-terminal interaction domain [3]. RIPK1-3 have
all been shown to play a role in the response to pathogens through activation of the pro-
inflammatory and pro-survival transcription factor NF-κB, inflammatory and non-
inflammatory cell death, and type I interferon (IFN-I) signaling [2]. RIPK4 can activate NF-
κB [4], but has no defined role in the response to pathogens and is a regulator of
keratinocyte growth and homeostasis [5, 6]. RIPK5, which is most similar to RIPK4, has
no known role in innate immunity.
RIPK1 and RIPK3 are perhaps the best described members of the RIPK family
and have critical roles in mediating cell death and inflammatory signaling [7]. Downstream
of signals from innate immune receptors like Z-DNA binding protein 1 (ZBP1) and tumor
necrosis factor receptor 1 (TNFR1), RIPK1 and/or RIPK3 are activated by
autophosphorylation and proceed to activate additional effector proteins to carry out their
innate immune functions. For example, RIPK3 can phosphorylate MLKL to activate highly
inflammatory programmed necrotic cell death (necroptosis) [8], RIPK1/3 can engage
caspase-8 (CASP8) to activate apoptotic caspases [9, 10], or RIPK1/3 can activate the
pro-survival and pro-inflammatory transcription factor NF-κB [11, 12]. Several factors are
known to impact the outcome of RIPK1/3 activation, such as the activating signal and
caspase activity [7, 13]. Altogether, RIPK1 and RIPK3 are central to determining cell fate
downstream of various innate immune stimuli.
Like many innate immune proteins, RIPK1 and RIPK3 are evolutionarily dynamic
in vertebrates. Phylogenomic analyses of a small number of species from different
vertebrate clades suggest that RIPK3, and the associated proteins ZBP1 and MLKL, have
been lost in some vertebrate lineages including birds, carnivores, and marsupials [14,
15]. In addition, RIPK1, RIPK3, MLKL, and ZBP1 are known to interact with viral proteins

[16-21] and have been shown to be evolving under recurrent positive selection in both

primates and bats [22], a hallmark of host proteins that directly interact with viral proteins

[23]. The degree to which other RIP kinases (RIPK2, 4, and 5) and their interacting

partners have diversified across vertebrates is unknown. Moreover, the vast majority of

functional data on RIPKs are from human or mouse homologs, despite evolutionary

studies that suggest functional diversification in these proteins may have occurred.

Here, we use evolutionary and functional approaches to characterize RIPK1-5 and

their associated proteins across vertebrate species. We identify unique patterns of

conservation of RIP kinases and their associated proteins across vertebrate lineages.

These distinct evolutionary histories have resulted in different mechanisms of activation

of NF-κB for human RIPK1-4. In addition to observing the loss of RIPK3 and necroptosis-

associated proteins in some vertebrate lineages, we identify changes in regulatory

features—including catalytic sites and CASP cleavage sites—in RIPK1 and RIPK3 in

specific mammalian lineages that may affect their function. Intriguingly, we found that the

RIP homotypic interaction motif (RHIM) domain of RIPK1 and RIPK3, as well as other

RHIM-containing proteins, is highly conserved in vertebrates and in a non-vertebrate

RIPK1 protein. Consistent with this strong conservation of the RHIM domain, we found

that RIPK1 and RIPK3 proteins from diverse vertebrates, and even a non-vertebrate

RIPK1, are able to activate NF-κB in a RHIM-dependent manner, suggesting that

activation of this inflammatory response is a conserved function of RIPK1 and RIPK3 in

vertebrates. Altogether, these data suggest that activation of NF-κB is the ancestral and

conserved feature of RIP kinases, while other RIPK-mediated functions such as

necroptosis are more evolutionarily labile.

Materials & Methods

Phylogenetic analysis

Phylogenetic trees were generated for RIPK1-5 using previously described

methods [24]. Briefly, the kinase domains from human RIPK1 (NP_001341859.1), RIPK2

(NP_003812.1), RIPK3 (NP_006862.2), RIPK4 (NP_065690.2), and RIPK5

(NP_848605.1) were used to query the RefSeq protein database using BLASTp [25].

While LRRK1 and LRRK2 share kinase homology with RIPK1-5 and are sometimes

known as RIPK6 and -7 [3], they have a distinct domain structure and function compared

to RIPK1-5 and will therefore not be considered as RIP kinases for the purposes of this

study. The five resulting sequence lists were consolidated to a single list of unique

sequences using Geneious software. These sequences were then aligned using Clustal

Omega [26] with two refinement iterations implemented in Geneious software. Following

alignment, we filtered sequences based on alignment of an N-terminal kinase domain to

human RIPK kinase domains and the presence of a C-terminal domain. Proteins with

additional N-terminal domains, proteins lacking homology to human kinase domains, and

proteins lacking C-terminal domains were removed. Nearly-identical sequences were

reduced to a single unique sequence using CD-HIT [24] with a 0.8 sequence identity

cutoff. The resulting 736 sequences were realigned using Clustal Omega with two

refinement iterations. Phylogenetic trees were generated for both full length protein

alignments and for alignments trimmed to include only the kinase domains using IQ-TREE

[24] with 1000 bootstrap and SH-aLRT support values. The best substitution model was

determined by ModelFinder [27]. Accession numbers with species names and the tree

file can be found Supplementary Files 1 and 2.

Gene loss analysis

Using the aligned and filtered vertebrate RIPK sequences (4,550 sequences in

total, see 'Phylogenetic analysis' section), a tree was built using FastTree [28, 29]

implemented using Geneious. Using this tree, we were able to identify and extract *bona*

fide RIP kinases and remove similar, non-RIP kinases. The extracted sequences were

again aligned using Clustal Omega and filtered using the criteria above (See

'Phylogenetic analysis' section). A second tree was generated using FastTree. From this

tree, we extracted individual sequence lists for RIPK1-5 to determine which vertebrate

species did or did not have each RIPK. This method was repeated for CASP2 and

CASP8-10, ZBP1, MLKL, TRIF, FADD, NOD 1 and 2, and IRF6. The list of species and

lists of accession numbers for each protein can be found in Supplementary Files 3 and 4.

Plasmids and constructs

Coding sequences for human RIPK1 (Addgene #78834), human RIPK2
(ORFeome ID #4886), human RIPK3 (Addgene #78804), mouse RIPK1 (Addgene
#115341), and mouse RIPK3 (Addgene #78805) were cloned into apcDNA5/FRT/TO
backbone (Invitrogen, Carlsbad, CA) with an N-terminal V5 tag and linker using Gibson
Assembly (New England Biolabs, Ipswich, MA). Human RIPK4 (NP_065690.2) and
RIPK5 (NP_848605.1), and non-human RIPK1 C-termini (cat, XP_023109490.2; pig,
XP_003128209.1;   chicken,   NP_989733.3;   lizard,   XP_003224434.1;   lamprey,
XP_032813622.1; zebra mussel, XP_052232845.1) and RIPK3 (cat, XP_003987615.3;
pig, XP_001927459.3; lizard, XP_003223896.2; turtle, XP_037771912.1; lamprey,
XP_032816533.1) homologues were ordered from Twist Biosciences (San Francisco,
CA, USA) and were cloned along with RHIM mutants targeting the core tetrad into the
pcDNA5/FRT/TO backbone (Invitrogen, Carlsbad, CA) with an N-terminal V5 tag and
linker using Gibson Assembly (New England Biolabs, Ipswich, MA). Kinase mutants for
human proteins RIPK1 (D138N), RIPK2 (D146N), RIPK3 (D142N), RIPK4 (D143N), and
RIPK5 (D145N), and RHIM mutants for RIPK1 (IQIG542AAAA) and RIPK3
(VQVG461AAAA) were generated using Gibson Assembly.
140
Cell culture and transient transfections
HEK293T cells (obtained from ATCC) and the generated HEK293T RIPK1 KO cells
were maintained at a low passage number to maintain less than one year since
purchase, acquisition, or generation. Both cell lines were grown in complete media
containing DMEM (Gibco, Carlsbad, CA), 10% FBS (Peak Serum, Wellington, CO), and
1% penicillin/streptomycin (Gibco, Carlsbad, CA). A day prior to transfection, HEK293T
and RIPK1 KO cells were seeded into 24-well plates with 500 µL complete media or 96-
well plates with 80 µL complete media. Cells were transiently transfected with 500 ng
total DNA and 1.5 µL of TransIT-X2 (Mirus Bio, Madison, WI) in 100 µL Opti-MEM
(Gibco, Carlsbad, CA) for a 24-well plate or 100 ng DNA and 0.3 µL TransIT-X2 in 10
µL Opti-MEM for a 96-well plate. DNA and TransIT-X2 were incubated at room
temperature for 25-30 minutes, then added dropwise to the appropriate well. Cells were
harvested or analyzed at 18-22 hours post-transfection.
154

Generation of knockout cell lines

RIPK1 knockout HEK293T cells were generated using CRISPR/Cas9 as
previously described [30, 31]. Briefly, plasmids were generated order to produce
lentivirus-like particles containing the CRISPR/Cas9 machinery and guide RNA targeting
exon 5 (ENSE00003586162) of RIPK1. The protocol for the molecular cloning of this
plasmid was adapted from Feng Zhang (Sanjana et al., 2014) using the transfer plasmid
pLB-Cas9 (gift from Feng Zhang, Addgene plasmid # 52962). We designed the gRNA
target sequence using the web tool CHOPCHOP (Labun et al., 2016), available at
https://chopchop.cbu.uib.no/, and synthesized oligonucleotides from Integrated DNA
Technologies (San Diego, CA). The synthesized oligonucleotide pair was phosphorylated
and annealed using T4 Polynucleotide Kinase (NEB M0201S) with T4 Ligation Buffer
(NEB). Duplexed oligonucleotides were ligated into dephosphorylated and BsmBI-
digested pLB-Cas9 using the Quick Ligase kit (NEB M2200S) to generate transfer
plasmid with RIPK1 guide sequence. To generate lentivirus-like particles, this transfer
plasmid was transfected alongside two packaging plasmids, pMD2.G (gift from Didier
Trono, Addgene plasmid # 12259) and psPAX2 (gift from Didier Trono, Addgene plasmid
12260), into HEK293T cells with a 1:1:1 stoichiometry. Forty-eight hours post-
transfection supernatant was harvested and syringe-filtered (0.45µm). Supernatant
containing sgRNA-encoding lentivirus-like particles was then used to transduce HEK293T
cells. Transduced cells were cultured in growth media for 48 hours, then cultured in
growth media supplemented with 1 µg/ml puromycin for 72 hours. Using limiting dilution
in 96-well plates, the monoclonal cell line was then obtained. RIPK1 knockout was
confirmed by Sanger sequencing and by western blot using an α-RIPK1 antibody.

NF-κB and IFN luciferase reporter assays

To quantify NF-κB and IFN activation, we used the Dual-Glo Luciferase Assay
System E2920 (Promega, USA). WT or RIPK1 KO HEK293T cells were seeded in a white
96-well plate and transfected with firefly luciferase fused to either the NF-κB response
element (pGL4.32, Promega) or the human IFN-beta promoter (IFN-Beta-pGL3, Addgene
#102597), renilla luciferase fused to herpes simplex virus thymidine kinase promoter
(pTK-Renilla, Thermo Scientific), and the indicated RIPK. Firefly luciferase was used as

the primary reporter with renilla luciferase being a normalization control. Eighteen to 24

hours after transfection, Dual-Glo Luciferase Assay Reagent was added to each

transfected well as well as 3-6 untransfected wells to serve as a negative control.

Following a 10-minute incubation, the firefly luciferase signal was measured using a

BioTek Cytation imaging reader with Gen5 software (Agilent Technologies, San Diego,

CA). Then, Dual-Glo Stop & Glo reagent was added to each well. Following a 10-minute

incubation, the renilla luciferase signal was measured. The background luminescence

signal from the buffer-treated untransfected conditions was subtracted from other

conditions, firefly values were normalized to renilla values, and all samples were

normalized to the empty vector control condition. Values are reported as NF-κB activity

(firefly/renilla).

Cell death assay

Cell viability was measured using the ReadyProbes Cell Viability Imaging Kit, Blue/Green

(ThermoFisher Scientific). HEK293T cells were seeded in 80 µL of DMEM+/+ (See "Cell

culture and transient transfections") in a clear 96-well plate and transfected with the

indicated plasmids or a vector control. Eighteen hours post-transfection, cells were

stained using NucBlue Live reagent, which stains all nuclei, and NucGreen Dead reagent,

which stains only dead cells. A 2x concentrated mix (4 drops/mL of each dye) was made

and added to wells to 1x concentration. Cells were incubated for 20 minutes at room

temperature. Fluorescence for NucBlue (DAPI, excitation 377/20, emission 447/20) and

NucGreen (GFP, excitation 469/20, emission 525/20) were read using a BioTek Cytation

imaging reader with Gen5 software (Agilent Technologies, San Diego, CA). Unstained

wells were read and used as controls for background fluorescence. GFP values were

normalized to DAPI values, and all samples were normalized to the vector control

condition. Values are reported as Cytotoxicity (GFP/DAPI).

Immunoblotting and antibodies

Eighteen to 22 hours post-transfection, cells were washed with 1X PBS and lysed with

boiling 1x NuPAGE LDS sample buffer containing 5% β-mercaptoethanol at room

temperature for 5 minutes and then at 98°C for 7-10 minutes. Lysates were separated

by SDS-PAGE (4-15% Bis-tris gel; Life Technologies, San Diego, CA) with 1X MOPS

buffer (Life Technologies, San Diego, CA). Proteins were transferred onto a

nitrocellulose membrane (Life Technologies, San Diego, CA) and blocked with PBS–T

containing 5% bovine serum albumin (BSA) (Spectrum, New Brunswick, NJ).

Membranes were incubated with the indicated rabbit primary antibodies diluted with 5%

BSA and PBS-T at 1:1000 overnight at 4°C (α-V5 clone D3H8Q, α-GAPDH clone

14C10, α-RIPK1 clone E8S7U XP, α-RIPK2 clone D10B11, α-RIPK3 clone E1Z1D, α-

RIPK4 #12636; Cell Signaling Technology, Danvers, MA). Membranes were rinsed

three times with PBS-T, incubated with HRP-conjugated rabbit secondary antibody

diluted at 1:10,000 with 5% BSA and PBS-T for 30 minutes at room temperature, and

developed with SuperSignal West Pico PLUS Chemiluminescent Substrate (Thermo

Fisher Scientific, Carlsbad, CA).

Statistical analysis

Statistical analyses were completed using GraphPad Prism 10 software. Tests were

performed as indicated. Error bars were calculated using SEM.

Results

Comparative evolutionary and functional analysis of RIPK1-5

To directly compare the evolution of all RIP kinases, we performed phylogenetic

analysis on RIPK1-5 in vertebrates. We analyzed both the full-length proteins (Figure 1A,

Supplementary Files 1 and 2) and the kinase domains (Supplementary Figure 1). RIPK1

and RIPK3 are closely related, as are RIPK4 and RIPK5, reflecting their similar C-terminal

domains (Figure 1B). Moreover, RIPK1, 3, and 5 have greater sequence diversity across

vertebrates when compared to more well conserved RIPK2 and RIPK4 (Figure 1C).

These data, which are consistent with prior data showing that RIPK1 and RIPK3 are

evolving under diversifying positive selection in two mammalian clades [22], indicate that

different selective pressures have shaped the evolution of different RIPK genes in

vertebrates.

Figure 1: Diversity of RIPK evolution in vertebrates A) Maximum likelihood phylogenetic tree of vertebrate RIP kinases. The tree is rooted on an outgroup of RIPK proteins from *Petromyzon marinus* and cartilaginous fish. Bootstrap branch support values are indicated. The complete phylogenetic tree and list of protein accession numbers can be found in Supplementary Files 1 and 2. B) Domain structures of human RIP kinases. RHIM = RIP homotypic interaction motif, DD = death domain, CARD = caspase activation and recruitment domain, ANK = ankyrin repeats. C) Percent identity of macaque (*Macaca mulatta*), mouse (*Mus musculus*), and zebrafish (*Danio rerio*) RIP kinases compared to human proteins. Images of model species generated using BioRender.

RIPK1-4 have been reported to activate NF-κB in various cellular contexts while RIPK5 has no reported function in innate immune signaling [2]. We used a luciferase reporter assay to quantify activation of NF-κB by RIPK1-5 in HEK293T cells. As expected, overexpression of RIPK1-4 led to robust activation NF-κB (Figure 2A). In contrast, we observed no activation of NF-κB by RIPK5, despite expression levels similar to RIPK1 (Supplementary Figure 2). Interestingly, NF-κB activation by RIPK1-3 is independent of their kinase activity, while mutation of the RIPK4 kinase domain prevented activation of NF-κB (Figure 2B).

Of RIPK1-4, only RIPK1 is highly expressed in HEK293T cells (Supplementary Figure 3A-D). To better isolate the functions of individual RIP kinases, we generated *RIPK1* knockout HEK293T cells (RIPK1 KO; Supplementary Figure 3A, E). While RIPK2 and RIPK4 retained their ability to activate NF-κB, activation of NF-κB by RIPK3 is lost in RIPK1 KO cells (Figure 2C). This was expected given the critical role of RIPK1 in

regulating RIPK3 function [32]. These data suggest that the unique evolutionary histories

of RIPK1-4 have led to distinct mechanisms for activation of NF-κB.

Figure 2: Differential mechanisms of NF-κB activation by human RIPK1-4. Human V5-RIPK1-5
proteins were transfected into WT (A-C) or RIPK1 KO (C) HEK293T cells, along with NF-κB firefly
luciferase and control renilla luciferase reporter plasmids (see Materials and Methods), and NF-
κB activity was measured at 18h post-transfection. A) Human RIPK1-5 activation of NF-κB in WT
293T cells. B) Activation of NF-κB by catalytically active versus kinase-mutant RIPK1-5 in WT
HEK293T cells. C) Activation of NF-κB by RIPK1-5 in WT versus RIPK1 KO HEK293T cells) Data
are representative of 3 independent experiments with n=3-6 replicates per group. Data were
analyzed using one-way ANOVA with Dunnett's multiple comparisons test (A) or two-way ANOVA
with Šidák's multiple comparisons test (B, C). ns = not significant, *** = p<0.001, **** = p<0.0001.

RIP kinases have also been reported to activate type I interferons (IFN-I) [33-35]

and multiple forms of cell death [7]. Overexpression of RIPK2 and RIPK4 lead to moderate

IFN-I activation (Supplementary Figure 4A), which has been previously reported for

RIPK2 but not RIPK4 [33], although the overall activation was <1% the level of activation

seen upon overexpression of MAVS (mitochondrial antiviral signaling protein). Only

RIPK1 was able to induce any cell death over background levels (Supplementary Figure

4B) indicative of its central role in regulating cell survival and death [36], although this is

also minimal compared to a known activator of cell death. In contrast, RIPK3, which on

its own does not induce cell death, potently induces cell death when co-expressed with

ZBP1 and MLKL (Supplementary Figure 4B-C) [8, 37]. Altogether, these data highlight

the diverse functions of human RIPK1-4, as well as the conserved function of NF-κB
activation.
295
Phylogenomic analysis of RIPK and associated proteins in vertebrates
RIPK3, ZBP1, and MLKL are known to be lost in some vertebrate lineages [14]. It
is unknown if other RIPK and RIPK-associated proteins are similarly lost. We therefore
performed phylogenomic analysis on 490 vertebrate species for the presence or absence
of RIPK1-5 and several key RIPK-associated proteins (Figure 3, Supplementary Files 3
and 4). Due to the difficulty in discerning whether a gene absence in an individual species
is due to true gene loss or incomplete genome assembly, we focused on multi-species
patterns of gene loss within and between vertebrate lineages. We first confirmed the loss
of necroptosis associated proteins that has been previously reported [14]: RIPK3 and
ZBP1 in birds; RIPK3, ZBP1, and MLKL in marsupials; MLKL in carnivores. ZBP1 is
absent in fish, suggesting that this protein, and ZBP1-mediated necroptosis, arose only
during tetrapod evolution. Analysis of other RIP kinases revealed that RIPK1, RIPK2, and
RIPK4 are found in most jawed vertebrate species, while RIPK5 is recurrently lost in
different vertebrate lineages similar to RIPK3. Moreover, RIPK1-3 are found in the most
distant vertebrate species, relative to humans, sea lamprey (*Petromyzon marinus*),
whereas RIPK4 and RIPK5 are absent in this deeply branched vertebrate species.
Intriguingly, we observed a second paralog of RIPK2, which we call RIPK2B, in many
vertebrates but lost in therian (live-bearing) mammals (Supplementary Figure 5).
Additionally, we found a RIPK1-like protein in several non-vertebrate species, including
the zebra mussel *Dreissena polymorpha*. Additional RIPK-like proteins have been found
in other non-vertebrate species, including a RIPK1/2-like protein in *Drosophila* and other
protostomes [14], although these proteins were not considered here due to minimal
sequence similarity to the human RIP kinase domains and our focus on vertebrate RIP
kinases. Together, these data indicate that RIP kinases have duplicated, diversified, and,
in the cases of RIPK3 and RIPK5, have been recurrently lost throughout vertebrate
evolution.

Figure 3: Loss of specific RIPK and associated proteins in some vertebrate lineages. Heat map showing the percentage of species within a clade that have the indicated protein. The clades and the number of species within each clade are indicated on the left. Complete lists of proteins and species in each group can be found in Supplementary Files 3 and 4.

We also analyzed the phylogenomic distribution of other RIPK-associated proteins. We analyzed caspase-8 (CASP8) and additional activator caspases CASP2, CASP9, and CASP10. We confirmed the loss of CASP10 in some rodents [38], revealing that suborder Myomorpha ("mouse-like") and Castorimorpha ("beaver-like") rodents have lost CASP10, whereas Sciuromorpha ("squirrel-like") and Hystricomorpha ("porcupine-like") rodents have retained CASP10. We found that CASP8 is absent in fish and sea lamprey. This coincides with the emergence of ZBP1 in tetrapods. Further supporting the link between CASP8 and ZBP1, CASP8 is lost in some reptiles that have also lost ZBP1 (Supplementary Table 1). CASP8 is a known negative regulator of necroptosis and is important for mitigating ZBP1-induced inflammation [39]. These data suggest that loss of

ZBP1 is sufficient to allow for loss of CASP8. In contrast to CASP8, FADD, which may
also contribute to regulation of ZBP1 activity [39], is found in all jawed vertebrates,
indicating that it emerged prior to ZBP1. Additional RIPK-associated proteins, including
TRIF (RIPK1-interacting), NOD 1 and NOD2 (RIPK2-interacting), and IRF6 (RIPK4-
interacting) were also analyzed [1, 2]. TRIF is broadly expressed across vertebrates.
NOD1 is only found in jawed vertebrates. NOD2 is present in lamprey and fish, but absent
in birds, reptiles, and some amphibians, including some of the same species that contain
RIPK2B, the paralog of RIPK2 (Supplementary Figure 5). While RIPK2B retains both the
kinase domain and CARD, there may be distinct regulatory mechanisms, specifically
those related to NOD2, compared to RIPK2. Altogether, our large-scale phylogenomic
analysis revealed patterns of conservation and evolutionary lability in RIPK and RIPK-
associated proteins in vertebrates.

Lineage-specific losses of RIPK functional motifs despite broad conservation of RHIM
domains across vertebrates

In addition to analyzing the presence or absence of a protein across vertebrates,
we characterized the presence of specific regulatory and functional sites within proteins,
including catalytic sites, phosphorylation sites, and caspase cleavage sites. For instance,
the known CASP8 cleavage site in RIPK1 is absent in many cetaceans (Figure 4A).
Cetaceans have been shown to have disrupted necroptosis through truncation of RIPK3
(identified here as a loss of RIPK3) [15], suggesting that CASP8 cleavage of RIPK1 in
these species is likely not necessary to regulate necroptosis. However, there could be
other CASP cleavage sites within these RIPK1 proteins that serves a similar regulatory
function. Interestingly, while RIPK3 is retained in carnivores, the key catalytic residue
(D142 in human RIPK3), is mutated to an asparagine (Figure 4B). This, combined with
the loss of MLKL in carnivores, suggests that the role of RIPK3 in this order is independent
of its kinase activity.

Interestingly, despite the loss of key functional domains of RIPK1 and RIPK3, the
core tetrad of the RHIM domain (V/I-Q-V/I-G), is conserved across vertebrates (Figure
4C-F). This motif has been shown to be important for RIPK1-RIPK3 interactions, and
facilitates known kinase-independent functions of RIPK3 [40]. Moreover, the first RHIM

domain of ZBP1 and the TRIF RHIM domain are also highly conserved in vertebrates

(Supplementary Figure 6A, left and 6B). Interestingly, the second RHIM domain of ZBP1

diverges outside of mammals (Supplementary Figure 6A, right). Altogether, the strong

conservation of the RHIM domain across RIPK1 and RIPK3, despite the loss of other

functional motifs, suggests the ancestral and conserved functions of these proteins are

RHIM-dependent.

A

Cetacean RIPK1 CASP8 cleavage site

```
                         321        ▼        328
Human........................L Q L D C V A V
Sperm whale...............L Q I D C V A I
Blue whale ..................L Q I D C V A I
Orca ...........................L Q I G C V A M
Bottlenose dolphin .........L Q I G C V A V
Pilot whale..................L Q I G C V A V
Vaquita ......................L Q I D C V A M
Narwhal ......................L Q I G C V A M
Beluga whale ...............L Q I G C V A M
```

B

Carnivore RIPK3 catalytic site

```
                         137        ⬇        147
Human.......................V L L H R D L K P S N
Dog............................V L L H R D L K P S N
Ferret.........................V L L H R D L K P S N
Bear...........................V L L H R D L K P S N
Lynx..........................V L L H R N L K P S N
Tiger..........................V L L H R N L K P S N
Lion............................V L L H R N L K P S N
Cat.............................V L L H R N L K P S N
Cheetah......................V L L H R N L K P S N
```

C

Cetacean RIPK1 RHIM domain

```
                              RHIM tetrad
                              ⏞
                         536              546
Human..............................S T G I Q I G A Y N Y
Sperm whale.....................S S G V Q I G K N N Y
Blue whale .........................S S G I Q I G N N N Y
Orca ..................................S F G V Q I G N N N F
Bottlenose dolphin ..............C S G V Q I G N N N F
Pilot whale........................C S G V Q I G N N N F
Vaquita .............................C S G I Q I G N N N Y
Narwhal .............................S F G V Q I G N N N Y
Beluga whale .....................S F G V Q I G N N N Y
```

D

Carnivore RIPK3 RHIM domain

```
                              RHIM tetrad
                              ⏞
                         455              465
Human...........................C S G V Q V G D N N Y
Dog..................................S R A V Q V G D Y N Y
Ferret...............................S H A V Q V G D N N Y
Bear.................................S H A V Q V G D N N Y
Lynx................................S H A V Q V G D N N Y
Tiger................................S H A V Q V G D N N H
Lion..................................S H A V Q V G D N N H
Cat...................................S H A V Q V G D N N Y
Cheetah............................S H A V Q V G D N N Y
```

E

Vertebrate RIPK1 RHIM domain

```
                         536              546
Human ..........................S T G I Q I G A Y N Y
Mouse ...........................S S G I Q I G N H N Y
Pig ..................................N S G V Q I G D H N Y
Lizard .............................S T G I Q I G N Y N H
Frog ................................S N S I Q I G N H N T
Fish .................................A S A I Q I G S Y N T
Lamprey ........................G E N I Q I G N N N V
Zebra mussel .................P P N I Q I G N N N K
```

F

Vertebrate RIPK3 RHIM domain

```
                         455              465
Human ..........................C S G V Q V G D N N Y
Mouse ...........................C S E V Q I G N Y N S
Pig ..................................N Q G V Q I G N S N Y
Lizard .............................V S G I Q I G N N N T
Frog ................................A T G V Q I G N N N F
Fish .................................V F G V Q I G N N N S
Lamprey ........................L K N T Q I G N N N T
```

Figure 4: Conservation of the RIPK1 and RIPK3 RHIM domain across vertebrates. RIPK1 sequences from cetaceans highlighting the CASP8 cleavage site (A) and the RHIM tetrad (C). RIPK3 sequences from carnivore species highlighting the catalytic site in the kinase domain (B) and the RHIM tetrad (D). The RHIM domain of RIPK1 (E) and RIPK3 (F) from diverse vertebrate species. Residue numbers refer to the human sequence. Cetacean species include sperm whale (*Physeter catodon*), blue whale (*Balaenoptera musculus*), orca (*Orcinus orca*), bottlenose dolphin (*Tursiops truncatus*), pilot whale (*Globicephala melas*), vaquita (*Phocoena sinus*), narwhal (*Monodon monoceros*), and beluga whale (*Delphinapterus leucas*). Carnivore species include dog (*Canis lupus familiaris*), ferret (*Mustela putorius furo*), bear (*Ursus americanus*), lynx (*Lynx canadensis*), tiger (*Panthera tigris*), lion (*Panthera leo*), cat (*Felis catus*), cheetah (*Acinonyx jubatus*). RIPK3 (B) and RIPK1 (C) sequences from vertebrate species highlighting the core tetrad of the RHIM domain. Residue numbers refer to the human sequence. Vertebrate species shown include mouse (*Mus musculus*), bat (*Pteropus alecto*), pig (*Sus scrofa*), cat (*Felis catus*), lizard (*Anolis carolinensis*), frog (*Xenopus laevis*), fish (*Danio rerio*), lamprey (*Petromyzon marinus*), and zebra mussel (*Dreissena polymorpha*).

RHIM-dependent NF-κB activation by RIPK1 and RIPK3 is conserved in vertebrates

Activation of NF-κB by RIPK3 is dependent on the presence of RIPK1 (Figure 2C). We hypothesized this is due to direct interaction with RIPK1 via the RHIM domain. We therefore tested for activation of NF-κB by a RIPK3 RHIM mutant. Activation of NF-κB by RIPK3 is completely ablated upon mutation of the RHIM core tetrad (Supplementary Figure 7A). Interestingly, while mutation of the RIPK1 RHIM domain reduces activation of NF-κB, there is still robust activation (Supplementary Figure 7A). This is likely due to death domain-mediated interactions. We also tested NF-κB activation of other RHIM-containing proteins. As described previously [41], ZBP1 activates NF-κB in a RHIM- and RIPK1-dependent manner (Supplementary Figure 7B-C), although only the first RHIM is required for activation. Conversely, mutation of the TRIF RHIM domain does not affect activation of NF-κB, and activation is only moderately reduced in RIPK1 KO cells (Supplementary Figure 7D-E), likely due to the TIR domain in TRIF that is known mediate its innate immune signaling [40].

The strong conservation of the RHIM domain tetrad sequence (V/I-Q-V/I-G) of RIPK1 and RIPK3 across species led us to hypothesize that activation of NF-κB by RIPK1 and RIPK3 is also conserved. We therefore decided to tested a diverse panel of metazoan RIPK1 proteins in our NF-κB assay. To avoid the effects of both kinase activity and kinase-mediated interactions [42], we generated RIPK1 C-terminus (RIPK1^{CT}) proteins from diverse species (Supplementary Figure 8A). Both WT and RHIM mutant RIPK1^{CT} activate NF-κB to the same extent as full length RIPK1 (Supplementary Figure 8B).

Mammalian and non-mammalian RIPK1^{CT} proteins were all able to activate NF-κB

independently of endogenous RIPK1 (Figure 5A, Supplementary Figure 9). Similar to

human RIPK1, mutation of the RHIM domain does not affect activation of NF-κB by other

mammalian proteins, including mouse, cat, and pig (Figure 5B, left). In contrast, mutation

of the RHIM domain attenuated the ability of lizard and zebra mussel RIPK1^{CT} to activate

NF-κB, and mutation of the lamprey RHIM domain completely prevented activation of NF-

κB (Figure 5B, right).

We then tested diverse RIPK3 proteins for activation of NF-κB, including species

that do (mouse, pig, turtle) and do not (cat, lizard, sea lamprey) have a complete

necroptosis pathway as defined by presence of ZBP1, MLKL, and catalytically active

RIPK3. Mouse and cat RIPK3 activated NF-κB in a RIPK1-dependent manner, while pig

RIPK3 did not activate NF-κB (Figure 5C, left, Supplementary Figure 10). We additionally

tested the ability of mammalian RIPK3 to activate cell death using human ZBP1 and MLKL

(Supplementary Figure 11). Interestingly, pig RIPK3 activated cell death despite being

unable to activate NF-κB. Conversely, cat RIPK3 failed to activate cell death

(Supplementary Figure 11), confirming its catalytic inactivity, but is able to activate NF-

κB (Figure 5C, left). These data suggest that activation of cell death and NF-κB are two

distinct functions of RIPK3. Both lizard and turtle RIPK3 were able to activate NF-κB

independent of RIPK1 (Figure 5C, right). Activation of NF-κB was dependent on the RHIM

domain for all RIPK3 proteins tested, regardless of their dependence on RIPK1 (Figure

5D). Sea lamprey RIPK3, which has a RHIM tetrad of TQIG, had limited activation of NF-

κB, but activation increased upon insertion of a IQIG tetrad (Figure 5D). This highlights

the importance of the RHIM domain, particularly the core tetrad, in driving activation of

NF-κB. Altogether, these data reveal the conservation of NF-κB activation by vertebrate

RIPK1 and RIPK3 via the RHIM domain, while also highlighting subtle differences

between mammalian and non-mammalian RIP kinases.

Figure 5: NF-κB activation is a conserved function of RIPK1 and RIPK3 across vertebrates. RIPK1^{CT} (A-B) or RIPK3 (C-D) proteins were transfected into WT or RIPK1 KO HEK293T cells, along with NF-κB firefly luciferase and control renilla luciferase reporter plasmids (see Materials and Methods), and NF-κB activity was measured at 18h post-transfection A) Activation of NF-κB by mammalian (left) and non-mammalian (right) RIPK1^{CT} in WT and RIPK1 KO HEK293T cells. B) Activation of NF-κB by WT and RHIM mutant mammalian (left) and non-mammalian (right RIPK1^{CT} in WT HEK293T cells. C) Activation of NF-κB by mammalian (left) and non-mammalian (right) RIPK3 in WT and RIPK1 KO HEK293T cells. D) Activation of NF-κB by WT and RHIM mutant mammalian (left) and non-mammalian (right) RIPK3 in HEK293T cells. Squares in (D) represent inactivating alanine mutations, triangles represent a gain of function TQIG>IQIG mutation. Species shown are mouse (*Mus musculus*), cat (*Felis catus*), pig (*Sus scrofa*), lizard (*Anolis carolinensis*), turtle (*Chelonia mydas*), lamprey (*Petromyzon marinus*), and mussel (*Dreissena polymorpha*). Data are representative of 3-5 independent experiments with n=3-6 replicates per group. Data were analyzed using two-way ANOVA with Šidák's multiple comparisons test. ns = not significant* = p<0.05, ** = p<0.01, *** = p<0.001, **** = p<0.0001.

Discussion

Activation of NF-κB is critical to both the innate and adaptive immune response in humans and mice [43]. NF-κB and analogous pathways have been found in many non-vertebrate species, including horseshoe crab [44], *Drosophila melanogaster* [45], and cnidarians and bivalves (reviewed in [46]). Despite this deep conservation of NF-κB signaling, differences in the function and regulation of NF-κB-associated proteins between mice and humans have been identified [43], suggesting that this functionally well-conserved pathway can be adapted to different species. These data indicate that broader phylogenetic analysis of the genes and functions that are associated with NF-κB signaling are needed to understand the pathogen response across species, including those that harbor viruses and bacteria that pose a zoonotic threat to humans.

Here, we apply broad phylogenetic and functional sampling to the RIP kinase family of NF-κB activators. Using a phylogenomic approach, we identified widespread conservation of RIPK1, RIPK2, and RIPK4 within vertebrates. We also show conservation of activation of NF-κB signaling by RIPK1 and RIPK3 across diverse vertebrate species that is reliant on the highly conserved core tetrad of the RHIM domains found in both proteins. This striking conservation of RIPK1- and RIPK3-mediated activation of NF-κB, from species that span >500 million years of vertebrate evolution, underscores the core functionality of RIP kinases as NF-κB activating proteins. Despite this conservation of NF-κB activation, we also observed lineage specific changes in RIPK presence or mechanisms of NF-κB activation. For instance, we observed recurrent gene loss of RIPK3

and RIPK5 throughout the vertebrate phylogeny, and discovered a second paralog of
RIPK2 in many non-mammalian vertebrate species (RIPK2B). In addition, although
RIPK3-mediated activation of NF-κB is highly dependent on RIPK1 across diverse
mammals, RIPK3 activation of NF-κB in reptiles is independent of RIPK1. Activation of
NF-κB is part of the antiviral immune response and is known to be antagonized by viruses
across a range of species, including vertebrates and non-vertebrates [47-49]. This
antagonism could result in adaptation of NF-κB-associated proteins, including RIP
kinases, leading to lineage-specific mechanisms of activation that are tailored to the
specific contexts within and between species.

The most striking differences in RIPK-associated genes and functions are those
involved in necroptosis. Necroptosis is highly inflammatory and the associated proteins
must therefore be tightly regulated, as it could be a cause of immunopathology during
infection. Despite this, necroptosis has been shown to be important for the response to
several viruses [50-52]. It is hypothesized that necroptosis, defined by RIPK3-mediated
activation of MLKL, arose in vertebrates [14, 53]. Interestingly, we find that two critical
regulators of necroptosis, CASP8 and ZBP1, are only found in tetrapods, suggestive of
an additional innovation within the necroptosis pathway during the divergence of
tetrapods. Moreover, consistent with previous observations [14, 15], we observe a large
number of necroptosis-associated gene losses, while also discovering important
functional motifs in necroptosis proteins in distinct tetrapod lineages. These data support
a model where necroptosis has been constructed as an important facet of the innate
immune response in some tetrapods, but is not universal and has been deconstructed in
species where the immunopathologic effects outweigh the benefits.

There have been several hypotheses to explain the loss of necroptosis associated
proteins in specific vertebrate lineages, such as diet (*e.g.,* carnivores) or behavior (*e.g.,*
torpor in reptiles) [14]. While the cause of a specific instance of gene loss is difficult to
identify, it is clear that gene loss can provide an evolutionary tool for adaptation [54].For
example, caspases, which are key regulators of both immunologically silent apoptosis
and inflammatory pyroptosis, differ greatly across mammals, with lineage-specific
caspase repertoires including both loss and gain of novel caspases [38]. Even within a
single mammalian order, homologous caspases have been found to serve different

functions in response to pathogens [55]. This type of lineage-specific innovation likely
exists for necroptosis, and gene loss could be a cause or a result of adaptations. Indeed,
differences in MLKL activation by RIPK3 across species have been identified [56], which
may be the result of additional diverse regulatory mechanisms for necroptosis even
amongst species with an intact pathway. Strikingly, despite the lack of a complete
necroptosis pathway in some lineages (cat, turtle, lamprey), our functional data indicate
that RIPK3 retains the ability to activate NF-κB. Activation of NF-κB could therefore
represent the ancestral function of RIPK3, and it has been co-opted to activate
necroptosis in vertebrates.

Altogether, our combined phylogenetic and functional approaches reveal both
conservation and a striking lack of conservation in vertebrate RIPKs and their associated
proteins and functions. The strong conservation of RIPK1 proteins and their ability to
activate NF-κB across species is consistent with the model in which RIPK1 function is
central to determining cell fate downstream of TNF and TLR signaling [36]. Indeed, RIPK1
has been described as a guardian of antimicrobial proinflammatory signaling, where
RIPK1 activates proinflammatory and pro-survival signals downstream of multiple innate
immune receptors as a way to preserve activation in the face of pathogen antagonism
[57]. For instance, because some pathogens have evolved strategies to inhibit CASP8,
RIPK1- and RIPK3-mediated necroptosis is activated as a secondary cell death strategy
when CASP8 is inhibited [57]. This places both RIPK1 and RIPK3, a critical regulator of
necroptosis, at the center of determining cell fate during pathogen infection. The
divergence of RIPK1 and RIPK3 across vertebrates identified here, as well as the
previous finding that these proteins are evolving under positive selection [15, 22], may
have also resulted in distinct regulatory mechanisms or functions of these proteins across
species, such as the loss of the CASP8 cleavage site in some cetacean RIPK1 proteins
identified here. Thus, the functional divergence identified in our work highlights the
evolutionary innovation that can arise in innate signaling pathways across diverse
species, and underscores the importance of considering natural diversity when
characterizing the innate immune response to pathogens.

References

1. Cuny, G.D. and A. Degterev, *RIPK protein kinase family: Atypical lives of typical kinases.* Semin Cell Dev Biol, 2021. **109**: p. 96-105.
2. Eng, V.V., M.A. Wemyss, and J.S. Pearson, *The diverse roles of RIP kinases in host-pathogen interactions.* Semin Cell Dev Biol, 2021. **109**: p. 125-143.
3. Lv, S., et al., *Comparative and evolutionary analysis of RIP kinases in immune responses.* Front Genet, 2022. **13**: p. 796291.
4. Meylan, E., et al., *RIP4 (DIK/PKK), a novel member of the RIP kinase family, activates NF-kappa B and is processed during apoptosis.* EMBO Rep, 2002. **3**(12): p. 1201-8.
5. Kwa, M.Q., G.M. Scholz, and E.C. Reynolds, *RIPK4 activates an IRF6-mediated proinflammatory cytokine response in keratinocytes.* Cytokine, 2016. **83**: p. 19-26.
6. Kwa, M.Q., et al., *Receptor-interacting protein kinase 4 and interferon regulatory factor 6 function as a signaling axis to regulate keratinocyte differentiation.* J Biol Chem, 2014. **289**(45): p. 31077-87.
7. Newton, K., *RIPK1 and RIPK3: critical regulators of inflammation and cell death.* Trends Cell Biol, 2015. **25**(6): p. 347-53.
8. Sun, L., et al., *Mixed lineage kinase domain-like protein mediates necrosis signaling downstream of RIP3 kinase.* Cell, 2012. **148**(1-2): p. 213-27.
9. Wang, L., F. Du, and X. Wang, *TNF-alpha induces two distinct caspase-8 activation pathways.* Cell, 2008. **133**(4): p. 693-703.
10. Mandal, P., et al., *RIP3 induces apoptosis independent of pronecrotic kinase activity.* Mol Cell, 2014. **56**(4): p. 481-95.
11. Meylan, E., et al., *RIP1 is an essential mediator of Toll-like receptor 3-induced NF-kappa B activation.* Nat Immunol, 2004. **5**(5): p. 503-7.
12. Moriwaki, K., et al., *The necroptosis adaptor RIPK3 promotes injury-induced cytokine expression and tissue repair.* Immunity, 2014. **41**(4): p. 567-78.
13. Orozco, S. and A. Oberst, *RIPK3 in cell death and inflammation: the good, the bad, and the ugly.* Immunol Rev, 2017. **277**(1): p. 102-112.
14. Dondelinger, Y., et al., *An evolutionary perspective on the necroptotic pathway.* Trends Cell Biol, 2016. **26**(10): p. 721-732.
15. Agueda-Pinto, A., et al., *Convergent Loss of the Necroptosis Pathway in Disparate Mammalian Lineages Shapes Viruses Countermeasures.* Front Immunol, 2021. **12**: p. 747737.
16. Harris, K.G., et al., *RIP3 Regulates Autophagy and Promotes Coxsackievirus B3 Infection of Intestinal Epithelial Cells.* Cell Host Microbe, 2015. **18**(2): p. 221-32.
17. Croft, S.N., E.J. Walker, and R. Ghildyal, *Human Rhinovirus 3C protease cleaves RIPK1, concurrent with caspase 8 activation.* Sci Rep, 2018. **8**(1): p. 1569.
18. Wagner, R.N., J.C. Reed, and S.K. Chanda, *HIV-1 protease cleaves the serine-threonine kinases RIPK1 and RIPK2.* Retrovirology, 2015. **12**: p. 74.
19. Muscolino, E., et al., *Species-Specific Inhibition of Necroptosis by HCMV UL36.* Viruses, 2021. **13**(11).
20. Huang, Z., et al., *RIP1/RIP3 binding to HSV-1 ICP6 initiates necroptosis to restrict virus propagation in mice.* Cell Host Microbe, 2015. **17**(2): p. 229-42.

583 21. Liu, X., et al., *Epstein-Barr virus encoded latent membrane protein 1 suppresses*
584   *necroptosis through targeting RIPK1/3 ubiquitination.* Cell Death Dis, 2018. **9**(2):
585   p. 53.
586 22. Palmer, S., et al., *Evolutionary profile for (host and viral) MLKL indicates its*
587   *activities as a battlefront for extensive counteradaptation.* Mol Biol Evol, 2021.
588   **38**(12): p. 5405-522.
589 23. Daugherty, M.D. and H.S. Malik, *Rules of engagement: molecular insights from*
590   *host-virus arms races.* Annu Rev Genet, 2012. **46**: p. 677-700.
591 24. Fu, L., et al., *CD-HIT: accelerated for clustering the next-generation sequencing*
592   *data.* Bioinformatics, 2012. **28**(23): p. 3150-2.
593 25. Altschul, S.F., et al., *Gapped BLAST and PSI-BLAST: a new generation of protein*
594   *database search programs.* Nucleic Acids Res, 1997. **25**(17): p. 3389-402.
595 26. Sievers, F., et al., *Fast, scalable generation of high-quality protein multiple*
596   *sequence alignments using Clustal Omega.* Mol Syst Biol, 2011. **7**: p. 539.
597 27. Kalyaanamoorthy, S., et al., *ModelFinder: fast model selection for accurate*
598   *phylogenetic estimates.* Nat Methods, 2017. **14**(6): p. 587-589.
599 28. Price, M.N., P.S. Dehal, and A.P. Arkin, *FastTree: computing large minimum*
600   *evolution trees with profiles instead of a distance matrix.* Mol Biol Evol, 2009. **26**(7):
601   p. 1641-50.
602 29. Price, M.N., P.S. Dehal, and A.P. Arkin, *FastTree 2--approximately maximum-*
603   *likelihood trees for large alignments.* PLoS One, 2010. **5**(3): p. e9490.
604 30. Tsu, B.V., et al., *Diverse viral proteases activate the NLRP1 inflammasome.* Elife,
605   2021. **10**.
606 31. Stevens, D.A., et al., *Antiviral function and viral antagonism of the rapidly evolving*
607   *dynein activating adaptor NINL.* Elife, 2022. **11**.
608 32. Liu, S., et al., *RIPK3 signaling and its role in the pathogenesis of cancers.* Cell Mol
609   Life Sci, 2021. **78**(23): p. 7199-7217.
610 33. Pandey, A.K., et al., *NOD2, RIP2 and IRF5 play a critical role in the type I interferon*
611   *response to Mycobacterium tuberculosis.* PLoS Pathog, 2009. **5**(7): p. e1000500.
612 34. Lei, Y., et al., *Cooperative sensing of mitochondrial DNA by ZBP1 and cGAS*
613   *promotes cardiotoxicity.* Cell, 2023. **186**(14): p. 3013-3032 e22.
614 35. Saleh, D., et al., *Kinase Activities of RIPK1 and RIPK3 Can Direct IFN-beta*
615   *Synthesis Induced by Lipopolysaccharide.* J Immunol, 2017. **198**(11): p. 4435-
616   4447.
617 36. Clucas, J. and P. Meier, *Roles of RIPK1 as a stress sentinel coordinating cell*
618   *survival and immunogenic cell death.* Nat Rev Mol Cell Biol, 2023. **24**(11): p. 835-
619   852.
620 37. Upton, J.W., W.J. Kaiser, and E.S. Mocarski, *DAI/ZBP1/DLM-1 complexes with*
621   *RIP3 to mediate virus-induced programmed necrosis that is targeted by murine*
622   *cytomegalovirus vIRA.* Cell Host Microbe, 2012. **11**(3): p. 290-7.
623 38. Eckhart, L., et al., *Identification of novel mammalian caspases reveals an important*
624   *role of gene loss in shaping the human caspase repertoire.* Mol Biol Evol, 2008.
625   **25**(5): p. 831-41.
626 39. Rodriguez, D.A., et al., *Caspase-8 and FADD prevent spontaneous ZBP1*
627   *expression and necroptosis.* Proc Natl Acad Sci U S A, 2022. **119**(41): p.
628   e2207240119.

40.   Nanson, J.D., B. Kobe, and T. Ve, *Death, TIR, and RHIM: Self-assembling*
domains involved in innate immunity and cell-death signaling. J Leukoc Biol, 2019.
105(2): p. 363-375.
41.   Rebsamen, M., et al., *DAI/ZBP1 recruits RIP1 and RIP3 through RIP homotypic*
interaction motifs to activate NF-kappaB. EMBO Rep, 2009. **10**(8): p. 916-22.
42.   Delanghe, T., Y. Dondelinger, and M.J.M. Bertrand, *RIPK1 Kinase-Dependent*
Death: A Symphony of Phosphorylation Events. Trends Cell Biol, 2020. **30**(3): p.
189-200.
43.   Zhang, Q., M.J. Lenardo, and D. Baltimore, *30 Years of NF-kappaB: A Blossoming*
of Relevance to Human Pathobiology. Cell, 2017. **168**(1-2): p. 37-57.
44.   Wang, X.W., et al., *Evidence for the ancient origin of the NF-kappaB/IkappaB*
cascade: its archaic role in pathogen infection and immunity. Proc Natl Acad Sci U
S A, 2006. **103**(11): p. 4204-9.
45.   Hetru, C. and J.A. Hoffmann, *NF-kappaB in the immune response of Drosophila.*
Cold Spring Harb Perspect Biol, 2009. **1**(6): p. a000232.
46.   Williams, L.M. and T.D. Gilmore, *Looking Down on NF-kappaB.* Mol Cell Biol,
2020. **40**(15).
47.   Albarnaz, J.D., et al., *Molecular mimicry of NF-kappaB by vaccinia virus protein*
enables selective inhibition of antiviral responses. Nat Microbiol, 2022. **7**(1): p.
154-168.
48.   Gonzalez Lopez Ledesma, M.M., et al., *Dengue virus NS5 degrades ERC1 during*
infection to antagonize NF-kB activation. Proc Natl Acad Sci U S A, 2023. **120**(23):
p. e2220005120.
49.   Palmer, W.H., et al., *Induction and Suppression of NF-kappaB Signalling by a DNA*
Virus of Drosophila. J Virol, 2019. **93**(3).
50.   Koehler, H., et al., *Vaccinia virus E3 prevents sensing of Z-RNA to block ZBP1-*
dependent necroptosis. Cell Host Microbe, 2021. **29**(8): p. 1266-1276 e5.
51.   Balachandran, S. and G.F. Rall, *Benefits and Perils of Necroptosis in Influenza*
Virus Infection. J Virol, 2020. **94**(9).
52.   Shubina, M., et al., *Necroptosis restricts influenza A virus as a stand-alone cell*
death mechanism. J Exp Med, 2020. **217**(11).
53.   Tummers, B. and D.R. Green, *The evolution of regulated cell death pathways in*
animals and their evasion by pathogens. Physiol Rev, 2022. **102**(1): p. 411-454.
54.   Albalat, R. and C. Canestro, *Evolution by gene loss.* Nat Rev Genet, 2016. **17**(7):
p. 379-91.
55.   Devant, P., A. Cao, and J.C. Kagan, *Evolution-inspired redesign of the LPS*
receptor caspase-4 into an interleukin-1beta converting enzyme. Sci Immunol,
2021. **6**(62).
56.   Tanzer, M.C., et al., *Evolutionary divergence of the necroptosis effector MLKL.* Cell
Death Differ, 2016. **23**(7): p. 1185-97.
57.   Remick, B.C., M.M. Gaidt, and R.E. Vance, *Effector-Triggered Immunity.* Annu
Rev Immunol, 2023. **41**: p. 453-481.
671