Cntnap2 loss drives striatal neuron hyperexcitability and behavioral inflexibility

Katherine R. Cording[1], Emilie M. Tu[2], Hongli Wang[2], Alexander H. C. W. Agopyan-Miu[2], and Helen S. Bateup[1,2,*]

[1]Helen Wills Neuroscience Institute, University of California, Berkeley, Berkeley, CA USA
[2]Department of Molecular and Cell Biology, University of California, Berkeley, Berkeley, CA USA

*Correspondence to: bateup@berkeley.edu

Abstract

Autism spectrum disorder (ASD) is a neurodevelopmental disorder characterized by two major diagnostic criteria - persistent deficits in social communication and interaction, and the presence of restricted, repetitive patterns of behavior (RRBs). Evidence from both human and animal model studies of ASD suggest that alteration of striatal circuits, which mediate motor learning, action selection, and habit formation, may contribute to the manifestation of RRBs. *CNTNAP2* is a syndromic ASD risk gene, and loss of function of *Cntnap2* in mice is associated with RRBs. How loss of *Cntnap2* impacts striatal neuron function is largely unknown. In this study, we utilized *Cntnap2*$^{-/-}$ mice to test whether altered striatal neuron activity contributes to aberrant motor behaviors relevant to ASD. We find that *Cntnap2*$^{-/-}$ mice exhibit increased cortical drive of striatal projection neurons (SPNs), with the most pronounced effects in direct pathway SPNs. This enhanced drive is likely due to increased intrinsic excitability of SPNs, which make them more responsive to cortical inputs. We also find that *Cntnap2*$^{-/-}$ mice exhibit spontaneous repetitive behaviors, increased motor routine learning, and cognitive inflexibility. Increased corticostriatal drive, in particular of the direct pathway, may contribute to the acquisition of repetitive, inflexible behaviors in *Cntnap2* mice.

Introduction

Autism spectrum disorder (ASD) is a neurodevelopmental disorder characterized by alterations in social communication and interaction, as well as the presentation of restricted, repetitive behaviors (APA, 2022). Given that ASD has high heritability (Sandin et al., 2017), much work has been done in the last 30 years to identify genes that confer risk of developing ASD (De Rubeis et al., 2014; Iossifov et al., 2014; Sanders et al., 2015). Through this, hundreds of high-confidence risk genes have been identified, varying greatly in the proteins for which they code (Satterstrom et al., 2020). These include transcriptional and translational regulators, ion channels, receptors, cell adhesion molecules, and others (De Rubeis et al., 2014; Delorme et al., 2013; Ebert & Greenberg, 2013). Recent work has focused on identifying brain regions and circuits that may be commonly affected by ASD-related mutations. The basal ganglia, in particular the striatum, represents one such commonly altered brain region and prior studies have demonstrated changes in striatal circuit function and striatum-associated behaviors in mice with mutations in ASD risk genes (Benthall et al., 2021; Fuccillo, 2016; Peca et al., 2011; Peixoto et al., 2016; Platt et al., 2017; Rothwell et al., 2014; Wang et al., 2016). However, whether basal ganglia circuits are convergently changed in ASD mouse models is an open question. Here we investigated whether loss of function of the syndromic ASD risk gene *Cntnap2* alters striatal physiology and basal ganglia-dependent behaviors in mice.

Cntnap2 codes for a neurexin-like cell adhesion molecule called Contactin-associated protein-like 2 (Caspr2) (Poliak et al., 1999; Poliak et al., 2003). In mice, Caspr2 is expressed in several cortical and subcortical regions, including the striatum, from embryonic day 14 (E14) onward into adulthood (Penagarikano et al., 2011). Caspr2 is primarily localized at the juxtaparanodes of myelinated axons and is involved in the clustering of potassium channels (Poliak et al., 2003; Scott et al., 2019). In mice, *in vitro* work suggests that Caspr2 may also play a role in AMPAR trafficking and cell morphology (Anderson et al., 2012; Gdalyahu et al., 2015; Varea et al., 2015), and *ex vivo* studies indicate that it can control cell excitability and circuit synchronicity (Martin-de-Saavedra et al., 2022). Caspr2 is important during neurodevelopment and has been implicated in neuronal migration (Penagarikano et al., 2011), the maturation and function of parvalbumin-positive GABAergic interneurons (Penagarikano et al., 2011; Scott et al., 2019; Vogt et al., 2018), and the timing of myelination (Scott et al., 2019). *CNTNAP2* mutations in people lead to a neurodevelopmental syndrome that can include language disorders, epilepsy, obsessive compulsive disorder, and ASD (Penagarikano & Geschwind, 2012; Rodenas-Cuadrado et al., 2014). A mouse model of this syndrome, *Cntnap2*[-/-], has been shown to exhibit good face validity for ASD-relevant social and motor behavior alterations (Brunner et al., 2015; Dawes et al., 2018; Penagarikano et al., 2011; Scott et al., 2019). However, the impact of *Cntnap2* loss on striatal synaptic physiology and excitability and corticostriatal-dependent behaviors has not been comprehensively assessed.

The striatum is primarily composed of GABAergic striatal projection neurons (SPNs), which make up two functionally distinct output pathways: the D1-receptor expressing cells of the direct pathway (dSPNs), which project to substantia nigra pars reticulata (SNr), and the D2-receptor expressing cells of the indirect pathway (iSPNs), which project to external globus pallidus (GPe) (Calabresi et al., 2014; Gerfen & Surmeier, 2011; Kravitz et al., 2010; Tai et al.,

2012). The two types of SPNs are intermixed throughout the striatum and receive excitatory glutamatergic inputs from cortex and thalamus, as well as dopaminergic input from the midbrain (Ding et al., 2008; Doig et al., 2010; Gerfen & Surmeier, 2011). Coordinated activity between the populations of SPNs in response to these inputs mediates action selection, motor learning and habit formation (Hawes et al., 2015; Santos et al., 2015; Yin et al., 2005, 2006). Although SPNs comprise upwards of 95% of the cells in striatum, there are a wide range of distinct interneurons, primarily GABAergic, that contribute significantly to the inhibitory circuitry of the striatum. Parvalbumin (PV) interneurons, which make up ~2% of the cells in striatum, provide the largest feedforward inhibition onto SPNs (Burke et al., 2017). Changes in the number and/or function of PV interneurons have been identified in several ASD mouse models, including *Cntnap2*[-/-] mice, implicating PV circuitry as a potential common alteration across ASD mouse models (Filice et al., 2020; Juarez & Martinez Cerdeno, 2022).

  To determine how loss of *Cntnap2* affects striatal function, we assessed the physiology of SPNs and striatal PV-interneurons in the dorsolateral striatum and utilized a range of assays to assess striatum-associated behaviors in *Cntnap2*[-/-] mice. We find that SPNs of both the direct and indirect pathways exhibit increased corticostriatal drive, despite unchanged excitatory cortical input onto these cells. Although decreased inhibitory function has been identified in other brain regions in *Cntnap2*[-/-] mice, we find no deficit in broad or PV-specific inhibitory input onto SPNs in the case of *Cntnap2* loss. Instead, we identify a significant increase in the intrinsic excitability of SPNs in *Cntnap2*[-/-] mice, in particular in dSPNs, which likely underlies the increased corticostriatal drive of these cells. Behaviorally, we find that *Cntnap2*[-/-] mice exhibit RRB-like behaviors including increased grooming, marble burying and nose poking in the holeboard assay. *Cntnap2*[-/-] mice also exhibit increased motor routine learning in the accelerating rotarod, and cognitive inflexibility in an odor-based reversal learning task. Taken together, these findings suggest that enhanced direct pathway drive may play a role in the spontaneous and learned repetitive behaviors exhibited by these mice.

Results

Cntnap2^{-/-} SPNs exhibit increased cortical drive

Emerging evidence suggests that corticostriatal synapses are a common site of alteration in mouse models of ASD (Li & Pozzo-Miller, 2020). To test whether mice with loss of *Cntnap2* exhibit changes in corticostriatal connectivity, we crossed *Cntnap2^{-/-}* mice to *Thy1*-ChR2-YFP mice, which express channelrhodopsin in a subset of layer V pyramidal neurons (Fig. 1A) (Arenkiel et al., 2007; Poliak et al., 1999). These mice were crossed to a D1-tdTomato reporter line to visually identify dSPNs (Ade et al., 2011). We recorded from SPNs in the dorsolateral striatum (DLS), as this sensorimotor striatal subregion is implicated in the acquisition of habitual and procedural behaviors (Packard & Knowlton, 2002). Changes in physiological function in this area may be connected to the acquisition of repetitive motor behaviors in ASD (APA, 2022; Fuccillo, 2016). To simulate a train of cortical inputs, we applied ten pulses of blue light over the recording site in DLS and measured the number of action potentials (APs) fired by SPNs in the absence of synaptic blockers (Fig. 1A). We altered the light intensity to vary the probability of eliciting subthreshold depolarizations or AP firing. dSPNs were identified using tdTomato fluorescence, and tdTomato negative neurons were designated putative iSPNs.

We quantified the number of evoked APs at different light intensities and found that both dSPNs and iSPNs in young adult *Cntnap2^{-/-}* mice exhibited significantly increased spike probability compared to wild-type (WT) SPNs (Fig. 1B-E). This suggests increased corticostriatal drive, consistent with findings in another mouse model with loss of function of the ASD-risk gene *Tsc1* (Benthall et al., 2021). To test whether the enhanced spiking probability was due to a change in the synaptic excitability of *Cntnap2^{-/-}* SPNs, we applied blue light of varying intensity over the recording site in DLS to evoke AMPAR-driven excitatory postsynaptic currents (EPSCs). We found that the average optically-evoked EPSC amplitude was not significantly different across a range of light intensities in *Cntnap2^{-/-}* dSPNs or iSPNs compared to WT (Fig. 1F-I).

To further assess synaptic inputs, we measured the number of dendritic spines in *Cntnap2^{-/-}* and WT SPNs, which are typically the sites of cortical synaptic innervation (Bouyer et al., 1984; Xu et al., 1989). To visualize spines, we injected neonate *Cntnap2;Drd1a*-tdTomato mice with AAV5-*Syn1*-GFP virus to sparsely label dSPNs and iSPNs in the DLS (Fig. S1) (Keaveney et al., 2018). We found that *Cntnap2^{-/-}* SPNs in adult mice had similar spine density as WT (Fig. S1), suggesting no overall change in synapse number. Together, these results show that dSPNs and iSPNs in *Cntnap2^{-/-}* mice exhibit enhanced cortically-driven spiking. However, this is not due to a change in corticostriatal synaptic strength or overall synapse density.

Increased corticostriatal drive in *Cntnap2^{-/-}* mice is not due to reduced inhibition

Previous work has indicated a reduction in the number and/or function of fast-spiking parvalbumin-expressing (PV) interneurons across multiple brain regions in *Cntnap2^{-/-}* mice (Ahmed et al., 2023; Antoine et al., 2019; Jurgensen & Castillo, 2015; Paterno et al., 2021; Penagarikano et al., 2011; Vogt et al., 2018). While inhibitory deficits have been identified in the

cortex and hippocampus (Antoine et al., 2019; Jurgensen & Castillo, 2015), and the number of PV interneurons has been reported to be decreased in striatum (Penagarikano et al., 2011), a comprehensive assessment of inhibitory synaptic function has yet to be completed in the striatum of *Cntnap2$^{-/-}$* mice. We first determined whether there were global deficits in inhibition onto SPNs in *Cntnap2$^{-/-}$* mice using intrastriatal electrical stimulation to evoke inhibitory postsynaptic currents (IPSCs) (Fig. 2A). In *Cntnap2$^{-/-}$* dSPNs, the average electrically-evoked IPSCs across a range of stimulation intensities were not different from WT dSPNs (Fig. 2B,C). However, in *Cntnap2$^{-/-}$* iSPNs, IPSCs were, on average, slightly larger than those in WT iSPNs, although we note that the responses were variable across cells (Fig. 2D,E).

There are many sources of inhibition in the striatum (Burke et al., 2017), which can all be activated with electrical stimulation. To assess whether inhibition from PV interneurons specifically is altered in *Cntnap2$^{-/-}$* mice, we crossed *Cntnap2$^{-/-}$*;D1-tdTomato mice to *Pvalb*-Cre; RCL-ChR2-H134R-EYFP (Ai32) mice to express channelrhodopsin in PV interneurons (Fig. 2F) (Hippenmeyer et al., 2005; Madisen et al., 2010). We applied a blue light pulse of varying intensity over the recording site to evoke PV interneuron-specific IPSCs in SPNs, in the presence of excitatory synaptic blockers (Fig. 2F). We found that the average optically-driven IPSCs did not differ significantly in amplitude in *Cntnap2$^{-/-}$* dSPNs or iSPNs compared to WT controls (Fig. 2G-J).

To directly measure PV neuron function, we assessed the intrinsic excitability of PV interneurons in *Cntnap2$^{-/-}$* mice. To visualize PV interneurons for recordings, we crossed *Cntnap2$^{-/-}$* mice to *Pvalb*-Cre;RCL-tdT (Ai9) mice (Fig. S2A). Plotting the number of APs fired as a function of current step size indicated that there were no significant differences in the intrinsic excitability of PV interneurons in *Cntnap2$^{-/-}$* mice compared to controls (Fig. S2B,C). There were also no changes in intrinsic cell properties such as rheobase, membrane resistance, capacitance, resting membrane potential, or AP shape in PV interneurons in *Cntnap2$^{-/-}$* mice (Fig. S2D-K).

Given prior reports of altered PV cell number in *Cntnap2$^{-/-}$* mice (Paterno et al., 2021; Penagarikano et al., 2011; Vogt et al., 2018), we counted PV-expressing cells in the striatum, using immunohistochemistry and fluorescent in situ hybridization. We found no significant difference in the number of PV-positive cells in the dorsal striatum of *Cntnap2$^{-/-}$* mice compared to WT (Fig. S3A-F and K-N). We also found no changes in the average single cell or total level of PV protein expression in the dorsal striatum (Fig. 3G-J). Overall, we did not observe significant changes in either PV interneuron number, PV expression, or PV interneuron-mediated inhibition in the adult *Cntnap2$^{-/-}$* striatum compared to WT controls.

SPN intrinsic excitability is increased in *Cntnap2$^{-/-}$* mice

Given that the increased cortical drive onto *Cntnap2$^{-/-}$* SPNs could not be explained by changes in excitatory or inhibitory synaptic function, we tested whether it could be due to a change in the intrinsic excitability of SPNs. To measure this, we recorded from dSPNs and iSPNs in *Cntnap2$^{-/-}$*;D1-tdTomato mice and injected current steps of increasing amplitude. We found that *Cntnap2$^{-/-}$* SPNs exhibited significantly increased intrinsic excitability compared to WT SPNs, which was most pronounced for dSPNs (Fig. 3A-L). *Cntnap2$^{-/-}$* dSPNs, but not iSPNs,

also had reduced rheobase current (Fig. 3C), the minimum current required to evoke an AP, as well as increased membrane resistance (Fig. 3D). Membrane capacitance (Fig. 3E,K), resting membrane potential (Fig. 3F,L), and AP shape (Fig. S4) were not significantly changed in *Cntnap2-/-* SPNs. Given the lack of synaptic changes observed in *Cntnap2$^{-/-}$* SPNs, the increase in SPN intrinsic excitability likely underlies their enhanced corticostriatal drive (see Fig. 1).

Cntnap2$^{-/-}$ mice exhibit increased repetitive behaviors

RRBs comprise one of the primary symptom domains of ASD (APA, 2022). Alterations in striatal circuits are thought to be involved in the manifestation of RRBs, given the striatum's role in action selection and motor control (Estes et al., 2011; Fuccillo, 2016; Hollander et al., 2005; Langen et al., 2014). To determine whether changes in motor behavior accompanied the altered striatal physiology in *Cntnap2$^{-/-}$* mice, we assessed locomotor activity and spontaneous repetitive behaviors using the open field, marble burying, and holeboard assays (Fig. 4A-D). In the open field, we found no significant difference in the total distance traveled, average speed, or number of rears in *Cntnap2$^{-/-}$* mice compared to WT controls (Fig. 4E-G). We did find that *Cntnap2$^{-/-}$* mice made significantly more entries into the center of the open field arena than WT mice, which may reflect a reduction in avoidance behavior in these mice (Fig. 4H). Manually scored grooming behavior revealed that *Cntnap2$^{-/-}$* mice initiated more grooming bouts than WT controls in the open field (Fig. 4I).

To further assess motor behaviors in *Cntnap2$^{-/-}$* mice, we utilized the marble burying assay (Fig. 4C), which takes advantage of a mouse's natural tendency to dig or bury. The number of marbles buried is used as a measure of persistent or repetitive behavior (Angoa-Perez et al., 2013). We found that *Cntnap2$^{-/-}$* mice buried significantly more marbles on average than WT controls (Fig. 4J). Another measure of repetitive behavior, which is based on the natural exploratory behavior of mice is the holeboard assay (Fig. 4D). In this task, the number of nose pokes made into unbaited holes is recorded. We found that *Cntnap2$^{-/-}$* mice made significantly more nose pokes within a 10-minute period than WT mice (Fig. 4K). This was largely due to increased poking during the last 5 minutes of the test (Fig. 4L-M), indicating persistent poking behavior in *Cntnap2$^{-/-}$* mice. Together with the increased grooming identified in the open field, increased marble burying and nose poking indicate an increase in RRBs in *Cntnap2$^{-/-}$* mice. A summary of behavior test results by genotype and sex is shown in Table S1.

To gain further insight into the spontaneous behavior profile of *Cntnap2$^{-/-}$* mice, we utilized a combination of DeepLabCut and Keypoint-MoSeq to perform unbiased, machine learning-based assessment of general locomotion and behavior in an additional cohort of *Cntnap2$^{-/-}$* mice (Fig. 4N-P, S5) (Mathis et al., 2018; Wiltschko et al., 2020). Again, we found that *Cntnap2$^{-/-}$* mice did not exhibit major changes in general locomotor activity compared to WT littermates (Fig. S5A). Analysis of movement "syllables" using Keypoint-MoSeq revealed that across the 25 most frequently performed syllables, two syllables associated with grooming were performed with significantly increased frequency in *Cntnap2$^{-/-}$* mice (Fig. 5N). *Cntnap2$^{-/-}$* mice also had an increase in the total number of grooming bouts (Fig. S5B), replicating the findings in the manually scored cohort (see Fig. 4I). While syllable usage was generally similar between WT and *Cntnap2$^{-/-}$* mice, transitions between syllables differed between the groups (Fig. S5C).

A measure of the entropy of syllable transitions revealed that *Cntnap2*[-/-] mice exhibited less entropy, suggesting less variability in the transition from one movement syllable to the next (Fig. 4O). This rigidity in motor sequence may be indicative of more restricted motor behavior overall. Finally, we tested whether a trained decoder could accurately distinguish WT and *Cntnap2*[-/-] mice using information about movement, syllable usage, or syllable transitions. The decoding models performed significantly better than chance at identifying WT and *Cntnap2*[-/-] mice based on their syllable usage and transitions, but not general locomotor activity (Fig. 4P). Together, this analysis demonstrates that while overall locomotor activity is not strongly affected in *Cnantp2*[-/-] mice, the behavior patterns of these mice are distinct from WT, reflecting enhanced presence of RRBs.

Cntnap2[-/-] mice exhibit enhanced motor learning

The accelerating rotarod is a striatal-dependent measure of motor coordination and learning that has been used across a range of ASD mouse models (Cording & Bateup, 2023). Changes in corticostriatal circuits have been identified in mouse models of ASD with altered performance in the task (Cording & Bateup, 2023). Given the altered corticostriatal drive in *Cntnap2*[-/-] mice, we tested whether motor coordination and learning were affected in these mice. In the rotarod test, mice learn to walk and then run to stay on a rotating rod as it increases in speed over the course of five minutes. Mice perform three trials a day for four days. In trials one through six, the rod increases in speed from five to 40 revolutions per minute (RPM), while in trials seven through 12 the rod increases from 10 to 80 RPM (Fig. 5A). Learning occurs over trials within a day, as well as across days, as the mouse develops and hones a stereotyped motor pattern to stay on the rod for increasing amounts of time (Rothwell et al., 2014; Yin et al., 2009). We found that *Cntnap2*[-/-] mice performed significantly better than WT mice in this task, in particular, in the later trials when the rod rotates at the faster 10 to 80 RPM speed (Fig. 5B). Initial performance (terminal velocity on trial one) was not significantly different between WT and *Cntnap2*[-/-] mice (Fig. 5C), but the learning rate from trial one to trial 12 was increased in *Cntnap2*[-/-] mice (Fig. 5D). These findings expand upon previous work indicating increased performance on both steady-state and accelerating rotarod tasks utilizing slower speeds in *Cntnap2*[-/-] mice (Dawes et al., 2018; Penagarikano et al., 2011). These results also align with the increased rotarod performance seen in other ASD mouse models exhibiting enhanced corticostriatal drive (Benthall et al., 2021; Cording & Bateup, 2023).

Cntnap2[-/-] mice exhibit cognitive inflexibility

RRBs include not just stereotyped movements, but also insistence on sameness and perseverative interests (APA, 2022). Cognitive inflexibility, a deficit in the ability to flexibly adapt to changes in the environment and update behavior, is a manifestation of ASD and some other psychiatric disorders, which are associated with striatal dysfunction (Fuccillo, 2016). Indeed, in individuals with ASD, the severity of RRBs is associated with measures of cognitive inflexibility, and evidence from imaging studies suggests that altered corticostriatal connectivity may be present in the case of both repetitive behaviors and cognitive inflexibility (Uddin, 2021). To assess cognitive flexibility in *Cntnap2*[-/-] mice, we utilized a four-choice odor-based reversal

learning assay (Johnson et al., 2016; Lin et al., 2022). Briefly, mice are trained to dig for a reward in one of four pots containing scented wood shavings (Fig. 6A). On the first day of the task (acquisition), the rewarded pot is scented with odor one (O1). Mice reach criterion when they have chosen O1 for at least eight of 10 consecutive trials. On day two, mice are given a recall test in which the rewarded pot is again scented with O1. After reaching criterion, the reversal trials begin, and the rewarded pot is scented with the previously unrewarded odor two (O2). To reach criterion, mice must learn the new association of O2 and reward and choose O2 for eight of 10 consecutive trials.

During acquisition, $Cntnap2^{-/-}$ mice performed similarly to WT controls, not differing in the average number of trials needed to reach criterion, the number of quadrant entries made before making a choice, or the latency to choose a pot (Fig. 6B-D). On day two, $Cntnap2^{-/-}$ mice performed similarly to controls during recall, demonstrating successful consolidation of the odor-reward pairing (Fig. 6E). However, we found that $Cntnap2^{-/-}$ mice exhibited a deficit in reversal learning, requiring significantly more trials on average than WTs to reach criterion once the odor-reward pairing was changed (Fig. 6F). Interestingly, during reversal, $Cntnap2^{-/-}$ mice made fewer quadrant entries before making a digging choice and exhibited significantly decreased latency to make a choice compared to controls (Fig. 6G,H). Even after the first correct choice of O2 during reversal, $Cntnap2^{-/-}$ mice took more trials to reach criterion than WTs (Fig. 6I). In terms of errors, $Cntnap2^{-/-}$ mice made more reversal errors than WT mice (Fig. 6J), in particular perseverative (continuing to choose O1) and regressive (choosing O1 after correctly choosing O2 once) errors (Fig. 6K). $Cntnap2^{-/-}$ mice did not differ from WT controls in choices of the novel (newly introduced during reversal) or irrelevant (never rewarded) odors, or in the number of omitted trials (timing out without making a choice) (Fig. 6K). Instead, the persistence in choosing O1, even after at least one correct choice of O2, drove the cognitive inflexibility in these mice (Fig. 6L). This persistence in choice may be reflective of the broader scope of RRBs in $Cntnap2^{-/-}$ mice.

Discussion

In this study we tested whether loss of the ASD risk gene *Cntnap2* results in altered striatal physiology or striatal-dependent behavior. We found that both direct and indirect pathway SPNs exhibited significantly increased cortical drive in *Cntnap2*$^{-/-}$ mice. This change was likely not due to a change in excitatory or inhibitory synapses, as excitatory cortical input onto SPNs was unchanged and there were no significant deficits in inhibition onto SPNs in these mice. Instead, loss of *Cntnap2* resulted in a significant increase in the intrinsic excitability of SPNs, in particular dSPNs. At the behavioral level, *Cntnap2*$^{-/-}$ mice exhibited repetitive behaviors including increased grooming, nose poking, and marble burying. These mice also had enhanced motor learning, performing significantly better than controls in the accelerating rotarod task. Finally, *Cntnap2*$^{-/-}$ mice exhibited cognitive inflexibility in the four-choice reversal learning task.

Cellular phenotypes of *Cntnap2* loss

The loss of Caspr2 has a variable impact on intrinsic excitability across brain regions and cell types. The increased intrinsic excitability that we identified in SPNs has also been observed in Purkinje cells of the cerebellum (Fernandez et al., 2021) and pyramidal cells of the cortex (Antoine et al., 2019; Cifuentes-Diaz et al., 2023). However, we note hypoactivity (Brumback et al., 2018) or unchanged excitability (Lazaro et al., 2019) of pyramidal cells in some cortical regions in *Cntnap2*$^{-/-}$ mice. Caspr2 is involved in the clustering of voltage-gated potassium channels, in particular at the juxtaparanodes of myelinated axons (Poliak et al., 1999; Poliak et al., 2003) and the axon initial segment (Inda et al., 2006). Indeed, there are profound deficits in the clustering of Kv1-family channels in *Cntnap2*$^{-/-}$ mice, particularly Kv1.2 channels (Scott et al., 2019). These channels play an important role in regulating the intrinsic excitability of SPNs (Nisenbaum et al., 1994), and when blocked, result in decreased rheobase and increased firing (Shen et al., 2004). The loss of Caspr2 in *Cntnap2*$^{-/-}$ mice may result in improper localization of these channels, which could contribute to the decreased rheobase and increased SPN excitability we observed. While Caspr2 has most often been discussed for this interaction with Kv1-family channels, recent work has shown that it can also play a role in calcium signaling. The extracellular domain of Caspr2, after undergoing cleavage through proteolytic processing, can interact with the calcium extrusion pump PMCA2, promoting calcium export and likely decreasing cell excitability (Martin-de-Saavedra et al., 2022). This process, which increases with neuronal activity, may be disrupted in *Cntnap2*$^{-/-}$ mice, which could also contribute to the increased intrinsic excitability that we found in SPNs.

Prior studies of *Cntnap2*$^{-/-}$ mice have identified changes in the number of PV-expressing interneurons in the cortex (Penagarikano et al., 2011; Vogt et al., 2018), hippocampus (Paterno et al., 2021; Penagarikano et al., 2011) and striatum (Penagarikano et al., 2011). However, this finding is inconsistent across studies, as others have reported no change in the number of PV interneurons in these regions (Ahmed et al., 2023; Lauber et al., 2018; Scott et al., 2019). One possible explanation for this disparity is altered PV protein expression in *Cntnap2*$^{-/-}$ mice such that immunoreactivity varies in cell counting assessments. This is supported by the finding that the number of Vicia Villosa Agglutinin-positive (VVA+) perineuronal nets that preferentially

surround PV cells is unchanged in *Cntnap2*[-/-] mice, even when PV immunoreactivity varies (Hartig et al., 1992; Haunso et al., 2000; Lauber et al., 2018). Parvalbumin, a Ca^{2+} buffer, plays an important role in the intrinsic fast-spiking properties of PV interneurons, such that a reduction in PV protein expression is known to change PV intrinsic function (Orduz et al., 2013). However, altered intrinsic properties of PV interneurons has also been variably reported across brain regions and studies of *Cntnap2*[-/-] mice, with subtle changes in PV firing properties reported in the developing striatum (Ahmed et al., 2023) and adult cortex (Vogt et al., 2018), but unchanged in the hippocampus (Paterno et al., 2021) and medial prefrontal cortex (Lazaro et al., 2019). In this study, we find no significant change in the number of PV interneurons or the striatal expression of PV protein in *Cntnap2*[-/-] mice. Following from this, we find no significant deficit in inhibition onto SPNs nor do we find any differences in PV interneuron intrinsic excitability in these mice. Interestingly, we find that broad inhibition measured through intrastriatal stimulation is enhanced specifically onto iSPNs in *Cntnap2*[-/-] mice. This change is likely driven by altered function or connectivity of another type of striatal interneuron, or other SPNs, which provide lateral inhibition onto iSPNs (Burke et al., 2017; Koos et al., 2004).

<u>Loss of *Cntnap2* alters striatal-dependent behaviors</u>

The striatum can be separated into functionally distinct subregions. We focused on the dorsal striatum in this study because of its role in controlling motor and cognitive functions (Voorn et al., 2004), which are relevant to ASD (Fuccillo, 2016; Subramanian et al., 2017). The dorsal striatum can be further subdivided into the dorsomedial striatum (DMS) and the dorsolateral striatum (DLS), with the former considered an associative region involved in goal-directed action-outcome learning and the latter implicated in the acquisition of habitual or procedural behaviors (Packard & Knowlton, 2002). We focused on cell function in DLS as stereotyped, perseverative or persistent behaviors likely recruit DLS circuitry (Evans et al., 2024; Fuccillo, 2016). In the accelerating rotarod assay, learning and performance in the task has been associated with changes in the DLS. Positive modulation of the firing rate of DLS neurons occurs during rotarod training, in particular in later trials of the task, and synaptic potentiation of DLS SPNs in late training is necessary for intact performance (Yin et al., 2009). In line with this, lesions of the DLS impair both early and late rotarod learning (Yin et al., 2009). We found that *Cntnap2*[-/-] mice had increased rotarod performance, most notably at the later stages when DLS function is strongly implicated. Functionally, we also found increased cortical drive of DLS SPNs in these mice, in particular of dSPNs, a change that was associated with increased rotarod performance in another mouse model with disruption of an ASD-risk gene (Benthall et al., 2021). Together, this supports a connection between the functional change observed in DLS SPN physiology and the increased motor routine learning in *Cntnap2*[-/-] mice.

In terms of restricted, repetitive behaviors, we replicated prior studies showing increased spontaneous grooming in *Cntnap2*[-/-] mice (Penagarikano et al., 2011). Early evidence implicates the striatum in the control of the syntax or sequence of movements in a rodent grooming bout, such that very small lesions of DLS are capable of disrupting grooming (Cromwell & Berridge, 1996). However, recent work has also outlined roles for cellular modulation in DMS and ventral striatal Islands of Calleja in the control of grooming behavior (Ramirez-Armenta et al., 2022;

Zhang et al., 2021). *Cntnap2*$^{-/-}$ mice also exhibited increased marble burying and nose poking. The precise neurobiological substrates of these behaviors are yet unclear, but evidence linking increases in these behaviors to changes in cortico-striatal and amygdala-striatal function supports the notion that these behaviors may fit into a broader basal ganglia-associated RRB-like domain (Albelda & Joel, 2012; Lee et al., 2024).

In the four-choice reversal learning task, *Cntnap2*$^{-/-}$ mice showed no differences during the acquisition phase, suggesting that there were no broad deficits in learning. However, in the reversal stage of the task, *Cntnap2*$^{-/-}$ mice took significantly more trials to learn a new odor-reward pairing, owing primarily to continued choice of the previously rewarded odor. The DMS and ventral striatum (nucleus accumbens) have been shown to play an important role in reversal learning (Izquierdo et al., 2017), and in the four choice task specifically (Delevich et al., 2022). Additionally, decreased dopamine release in the DLS is associated with deficits in reversal learning in this task (Kosillo et al., 2019; Lin et al., 2022). Together, the learning phenotypes seen in *Cntnap2*$^{-/-}$ mice in the accelerating rotarod and reversal learning assay share an underlying rigidity in behavioral choice. In both cases, changes in striatal circuits likely underlie the repetitive, stereotyped behaviors.

In summary, our results fit into a model whereby divergent cellular changes in the striatum driven by a functionally diverse set of ASD risk genes similarly enhance corticostriatal drive, in particular, of the direct pathway. This in turn may facilitate striatal-dependent motor routine learning and behavioral perseveration. We speculate that a shared gain-of-function in striatal circuits may play a role in the formation of perseverative or repetitive behaviors in a sub-set of ASDs more broadly.

Figures and Legends
Figure 1

Figure 1. *Cntnap2^{-/-}* SPNs exhibit increased cortical drive.

(A) Left: schematic of the corticostriatal connectivity experiments. For corticostriatal excitability, cortical terminals expressing ChR2 were stimulated with 10 pulses of blue light at 10 Hz and responses were recorded from dSPNs (red) and iSPNs (grey) in dorsolateral striatum. For corticostriatal synaptic strength, cortical terminals expressing ChR2 were stimulated with blue light at increasing intensity and synaptic currents were recorded from dSPNs (red) and iSPNs (grey) in dorsolateral striatum. Center: 10x confocal image of the striatum from a *Cntnap2^{+/+}*;D1-tdTomato;Thy1-ChR2 mouse. Right: 20x confocal image of dorsolateral striatum from a *Cntnap2^{+/+}*;D1-tdTomato;Thy1-ChR2 mouse. YFP (green) labels cell bodies and axons of a subset of layer V pyramidal neurons, tdTomato (red) labels dSPNs, and DAPI stained nuclei are in blue.

(B) Example single traces of action potentials (APs) in dSPNs evoked by cortical terminal stimulation at 20% light intensity for the indicated genotypes.

(C) Quantification (mean ± SEM) of the number of APs evoked in dSPNs at different light intensities. *Cntnap2^{+/+}* n=9 mice, 24 cells, *Cntnap2^{-/-}* n=10 mice, 22 cells. Two-way ANOVA p values are shown; geno $F_{(1, 221)} = 16.75$, stim $F_{(12, 221)} = 34.19$, g x s $F_{(12, 221)} = 0.7682$.

(D) Example single traces of action potentials (APs) in iSPNs evoked by cortical terminal stimulation at 20% light intensity for the indicated genotypes.

(E) Quantification (mean ± SEM) of the number of APs evoked in iSPNs at different light intensities. *Cntnap2^{+/+}* n=9 mice, 23 cells, *Cntnap2^{-/-}* n=10 mice, 21 cells. Two-way ANOVA p values are shown; geno $F_{(1, 234)} = 8.067$, stim $F_{(12, 234)} = 24.88$, g x s $F_{(12, 234)} = 0.2057$.

(F) Average EPSC traces from example dSPNs of each genotype induced by optogenetic cortical terminal stimulation at 14% light intensity.

(G) Quantification of EPSC amplitude evoked in dSPNs at different light intensities (line represents the mean, dots/squares represent average EPSC amplitude for each mouse). *Cntnap2^{+/+}* n=8 mice, 17 cells, *Cntnap2^{-/-}* n=5 mice, 13 cells. Two-way ANOVA p values are shown; geno $F_{(1, 33)} = 0.6527$, stim $F_{(2, 33)} = 11.80$, g x s $F_{(2, 33)} = 0.05117$.

(H) Average EPSC traces from example iSPNs of each genotype induced by optogenetic cortical terminal stimulation at 14% light intensity.

(I) Quantification of EPSC amplitude evoked in iSPNs at different light intensities (line represents mean, dots/squares represent average for mouse). *Cntnap2^{+/+}* n=6 mice, 13 cells, *Cntnap2^{-/-}* n=4 mice, 11 cells. Two-way ANOVA p values are shown; geno $F_{(1, 24)} = 2.770$, stim $F_{(2, 24)} = 3.466$, g x s $F_{(2, 24)} = 0.1408$.

Figure 2

Figure 2. Inhibition is not reduced in *Cntnap2⁻/⁻* SPNs.
(A) Top: schematic of the experiment. A bipolar stimulating electrode was placed roughly 200 µm from the recording site. A range of electrical stimulation intensities was applied to the tissue while IPSCs were recorded from dSPNs (red) and iSPNs (grey) in dorsolateral striatum. Bottom: 20x confocal image of dorsolateral striatum from a *Cntnap2⁺/⁺;*D1-tdTomato mouse. tdTomato (red) labels dSPNs, DAPI stained nuclei are in blue.

(B) Average IPSC traces from example dSPNs of each genotype evoked by electrical stimulation at 1.5 (x0.01 mA) intensity for the indicated genotypes.

(C) Quantification (mean ± SEM) of IPSC amplitude in dSPNs at different stimulation intensities. $Cntnap2^{+/+}$ n=17 cells from 9 mice, $Cntnap2^{-/-}$ n=16 cells from 9 mice. Two-way ANOVA p values are shown; geno F (1, 248) = 0.004612, stim F (7, 248) = 16.09, g x s F (7, 249) = 0.3164.

(D) Average IPSC traces from example iSPNs of each genotype evoked by electrical stimulation at 1.5 (x0.01 mA) intensity for the indicated genotypes.

(E) Quantification (mean ± SEM) of IPSC amplitude in iSPNs at different stimulation intensities. $Cntnap2^{+/+}$ n=16 cells from 9 mice, $Cntnap2^{-/-}$ n=16 cells from 10 mice. Two-way ANOVA p values are shown; geno F (1, 240) = 11.25, stim F (7, 240) = 10.17, g x s F (7, 240) = 0.1947.

(F) Top: schematic of the experiment. PV interneuron terminals expressing ChR2 were stimulated with blue light at a range of intensities and optically-evoked IPSCs were recorded from dSPNs (red) and iSPNs (grey) in dorsolateral striatum. Bottom: 20x confocal image of dorsolateral striatum from a $Cntnap2^{+/+}$;D1-tdTomato;PV-Cre;Ai32 mouse. YFP (green) labels PV interneurons, tdTomato (red) labels dSPNs, DAPI stained nuclei are in blue.

(G) Average IPSC traces from example dSPNs of each genotype evoked by optogenetic PV interneuron stimulation at 30% light intensity.

(H) Quantification (mean ± SEM) of IPSC amplitude in dSPNs at different light intensities. $Cntnap2^{+/+}$ n=29 cells from 15 mice, $Cntnap2^{-/-}$ n=23 cells from 11 mice. Two-way ANOVA p values are shown; geno F (1, 504) = 0.006524, stim F (7, 504) = 14.26, g x s F (7, 504) = 0.1124.

(I) Average IPSC traces from example iSPNs of each genotype evoked by optogenetic PV interneuron stimulation at 30% light intensity.

(J) Quantification (mean ± SEM) of IPSC amplitude in iSPNs at different light intensities. $Cntnap2^{+/+}$ n=24 cells from 14 mice, $Cntnap2^{-/-}$ n=27 cells from 13 mice. Two-way ANOVA p values are shown; geno F (1, 392) = 2.638, stim F (7, 392) = 13.59, g x s F (7, 392) = 0.5088.

Figure 3

Figure 3. Intrinsic excitability is increased in *Cntnap2*$^{-/-}$ SPNs.

(A) Example AP traces in dSPNs evoked by a 200 pA current step for the indicated genotypes.

(B) Quantification (mean ± SEM) of the number of APs evoked in dSPNs at different current step amplitudes. *Cntnap2*$^{+/+}$ n=22 cells from 8 mice, *Cntnap2*$^{-/-}$ n=23 cells from 8 mice. Two-way ANOVA p values are shown; geno F (1, 688) = 48.80, stim F (15, 688) = 107.5, g x s F (15, 688) = 1.737.

(C) Quantification (mean ± SEM) of the rheobase current in dSPNs. Dots/squares represent the rheobase current for each neuron. n is the same as in panel B. **p=0.0016, two-tailed unpaired t test.

(D-F) Quantification (mean ± SEM) of dSPN membrane resistance (D), *p=0.0328, Mann-Whitney test; membrane capacitance, p=0.2182, Mann-Whitney test; and resting membrane potential (F), p=0.9914, two-tailed unpaired t test. Dots/squares represent the average value for each neuron. n is the same as in panel B.

(G) Example AP traces in iSPNs evoked by a 200 pA current step for the indicated genotypes.

(H) Quantification (mean ± SEM) of the number of APs evoked in iSPNs at different current step amplitudes. *Cntnap2*$^{+/+}$ n=22 cells from 8 mice, *Cntnap2*$^{-/-}$ n=21 cells from 8 mice. Two-way ANOVA p values are shown; geno F (1, 656) = 4.186, stim F (15, 656) = 118.7, g x s F (15, 656) = 0.7780.

(I) Quantification (mean ± SEM) of the rheobase current in iSPNs. Dots/squares represent the rheobase current for each neuron. n is the same as in panel H. p=0.0923, two-tailed unpaired t test.

(J-K) Quantification (mean ± SEM) of iSPN membrane resistance (J), p=0.8193, Mann-Whitney test; membrane capacitance (K), p=0.6886, two-tailed unpaired t test; and resting membrane potential (L), p=0.4859, two-tailed unpaired t test. Dots/squares represent the average value for each neuron. n is the same as in panel H.

Figure 4

Figure 4. *Cntnap2$^{-/-}$* mice exhibit increased repetitive behaviors.

(A-D) Schematics of the behavioral assays used to measure repetitive behaviors in *Cntnap2$^{+/+}$* and *Cntnap2$^{-/-}$* mice, created with BioRender.com. Male and female mice were used for all tests.

(E-H) Quantification (mean ± SEM) of open field activity over 60 minutes. (E) total distance traveled, p= 0.3538, Mann-Whitney test; (F) average speed, p=0.3832, Mann-Whitney test; (G) number of rears, p=0.1892, two-tailed unpaired t test; (H) number of center entries, *p=0.0101, two-tailed unpaired t test. *Cntnap2$^{+/+}$* n=41 mice, *Cntnap2$^{-/-}$* n=34 mice.

(I) Quantification (mean ± SEM) of the number of manually scored grooming bouts in the first 20 minutes of the open field test, **p=0.0034, Mann-Whitney test, *Cntnap2$^{+/+}$* n=41 mice, *Cntnap2$^{-/-}$* n=34 mice.

(J) Quantification (mean ± SEM) of total marbles buried in the marble burying assay. *Cntnap2$^{+/+}$* n=33 mice and *Cntnap2$^{-/-}$* n=33 mice, *p=0.0396, two-tailed unpaired t test.

(K-M) Quantification (mean ± SEM) of performance in the holeboard assay. (K) Total number of nose pokes in 10 minutes, *p=0.0212, Mann-Whitney test; (L) nose pokes in the first five minutes, p=0.4811, two-tailed unpaired t test; and (M) nose pokes in the last five minutes, *p=0.0116, two-tailed unpaired t test. *Cntnap2$^{+/+}$* n=25 mice, *Cntnap2$^{-/-}$* n=22 mice.

(N) Quantification (mean ± SEM) of the frequency of movement syllables (top 25 most frequent syllables) in the open field assay defined by Keypoint-MoSeq. *Cntnap2$^{+/+}$* n=13 mice and *Cntnap2$^{-/-}$* n=11 mice, **p=0.0013 for grooming 6, **p=0.0013 for grooming 7, Kruskal-Wallis test with Dunn's correction for multiple comparisons.

(O) Quantification (mean ± SEM) of the entropy of syllable transitions in the open field assay. *Cntnap2$^{+/+}$* n=13 mice and *Cntnap2$^{-/-}$* n=11 mice, *p=0.0236, two-tailed unpaired t test.

(P) Accuracy of a Random Forest decoder trained on DeepLabCut basic locomotor data (Movement), Keypoint-MoSeq syllable usage data (Syllable usage), or Keypoint-MoSeq syllable transition data (Syllable transition) in distinguishing between *Cntnap2$^{+/+}$* and *Cntnap2$^{-/-}$* mice. Dotted line represents chance performance.

For panels E-M and O, dots/squares represent the value for each mouse.

Figure 5

Figure 5. *Cntnap2^-/-* mice exhibit enhanced motor learning.
(A) Schematic of the rotarod apparatus (top), and design of the task (bottom), created with BioRender.com. Mice walk to stay on the rotating rod for three 5-minute trials a day for two days at 5-40 RPM acceleration over 5 minutes, followed by three trials a day for two days at 10-80 RPM.
(B) Quantification (mean ± SEM) of accelerating rotarod performance across 12 trials for the indicated genotypes. *Cntnap2^+/+* n=30 mice, *Cntnap2^-/-* n=29 mice. Two-way ANOVA p values are shown; geno F (1, 683) = 92.17, trial F (11, 683) = 58.45, g x t F (11, 683) = 2.820.
(C) Quantification (mean ± SEM) of rotarod performance on trial 1 quantified as terminal speed. Dots/squares represent the performance of individual mice. n is same as in panel B, p=0.1518, Mann-Whitney test.
(D) Quantification (mean ± SEM) of learning rate (RPM/day) calculated as the slope of the line of performance from the first trial (1) to the last trial (12) for each mouse. Dots/squares represent the learning rate for individual mice. n is same as in panel B, ***p=0.0002, two-tailed unpaired t test.

Figure 6

Figure 6. *Cntnap2^{-/-}* mice exhibit cognitive inflexibility.

(A) Schematic of the four-choice odor-based reversal learning task, created with BioRender.com. Colored circles represent pots with different scented wood shavings. Yellow ring represents the food reward. Red circle in the Day 2 Reversal panel indicates a novel odor.

(B-D) Quantification of parameters during acquisition learning. Mean ± SEM number of trials to reach criterion (B, at least 8 out of last 10 trials correct), p=0.5397, Mann-Whitney test; number of quadrant entries before making a choice (C), p=0.9118, Mann-Whitney test; and latency to make a choice (D), p=0.7224, two-tailed unpaired t test. Dots/squares represent the value for each mouse.

(E) Quantification (mean ± SEM) of the number of trials to reach criterion (at least 8 out of last 10 trials correct) during the recall test on day 2, p=0.3737, Mann-Whitney test.

(F-I) Quantification of parameters during reversal learning. Mean ± SEM number of trials to reach criterion (F, at least 8 out of last 10 trials correct), **p=0.0048, two-tailed unpaired t test; number of quadrant entries before making a choice (G), *p=0.0158, two-tailed unpaired t test; latency to make a choice (H), **p=0.0013, two-tailed unpaired t test; and number of trials to reach criterion after the first correct choice (I), *p=0.0183, two-tailed unpaired t test.

(J) Quantification (mean ± SEM) of the total number of errors made during reversal learning, **p=0.0034, Mann-Whitney test.

(K) Quantification (mean ± SEM) of the different error types made during reversal learning. Perseverative errors, ***p=0.0005, regressive errors **p=0.0068, novel errors, p=0.9955, irrelevant errors p=0.9988, omissions, p=>0.9999, repeated measures two-way ANOVA with Šídák's multiple comparisons test; geno F (1, 18) = 10.91, error F (4, 72) = 53.49, g x e F (4, 72) = 4.292, animal F (18, 72) = 0.9638.

(L) Quantification (mean ± SEM) of the percent of choices made for each odor, binned across three trials, during reversal learning. Odor 1 was rewarded during acquisition learning and odor 2 was rewarded during reversal learning.

For panels B-L, n=10 *Cntnap2^{+/+}* mice and 10 *Cntnap2^{-/-}* mice.

Supplementary Figures and Table

Supplementary Figure 1

Supplementary Figure 1. *Cntnap2*$^{-/-}$ SPNs do not have altered spine density.
(A) Representative deconvoluted images of dendritic spines from dSPNs for the indicated genotypes.
(B) Quantification (mean ± SEM) of dendritic spine density per 10 µm of dendrite in *Cntnap2*$^{+/+}$ and *Cntnap2*$^{-/-}$ dSPNs. Dots/squares represent the average spine density per neuron. *Cntnap2*$^{+/+}$ n=9 neurons (15 dendrites) from 6 mice, *Cntnap2*$^{-/-}$ n=8 neurons (15 dendrites) from 6 mice. p=0.9964, two-tailed unpaired t test.
(C) Representative deconvoluted images of dendritic spines from iSPNs for the indicated genotypes.
(D) Quantification (mean ± SEM) of dendritic spine density per 10 µm of dendrite in *Cntnap2*$^{+/+}$ and *Cntnap2*$^{-/-}$ iSPNs. Dots/squares represent the average spine density per neuron. *Cntnap2*$^{+/+}$ n=7 neurons (15 dendrites) from 4 mice, *Cntnap2*$^{-/-}$ n=10 neurons (15 dendrites) from 7 mice. p=0.5362, Mann-Whitney test.

Supplementary Figure 2

Supplementary Figure 2. PV interneuron intrinsic excitability is unchanged in *Cntnap2*^{-/-} mice.

(A) Top: schematic of the experiment. tdTomato-expressing PV interneurons were recorded in the dorsolateral striatum and injected with current steps of increasing magnitude to evoke firing. Bottom: 20x confocal image of dorsolateral striatum from a *Cntnap2*$^{+/+}$;PV-Cre;Ai9 mouse. tdTomato (red) labels PV interneurons, DAPI stained nuclei are in blue.

(B) Example traces of APs in PV interneurons evoked by a 200 pA current step for the indicated genotypes.

(C) Quantification (mean $\pm$ SEM) of the number of APs evoked in PV interneurons at different current step amplitudes. *Cntnap2*$^{+/+}$ n=28 cells from 6 mice, *Cntnap2*$^{-/-}$ n=23 cells from 4 mice. Two-way ANOVA p values are shown; geno $F_{(1, 784)}$ = 2.728e-006, stim $F_{(15, 784)}$ = 67.25, g x s $F_{(15, 784)}$ = 0.1039.

(D) Quantification (mean $\pm$ SEM) of the rheobase current. Dots/squares represent the rheobase current for each neuron. n is the same as in panel C. p=0.8852, two-tailed unpaired t test.

(E-G) Quantification (mean $\pm$ SEM) of the membrane resistance (E), p=0.3422, Mann-Whitney test; membrane capacitance (F), p=0.9055, two-tailed unpaired t test; and resting membrane potential (G), p=0.3141, two-tailed unpaired t test. Dots/squares represent the average value for each neuron. n is the same as in panel C.

(H) Phase plane plot of a single AP evoked in PV interneurons from *Cntnap2*$^{+/+}$ (black) and *Cntnap2*$^{-/-}$ (red) mice.

(I-K) Quantification (mean $\pm$ SEM) of the AP threshold (I), p=0.6025, two-tailed unpaired t test; AP height (J), p=0.2850, two-tailed unpaired t test; and AP width at half max (K), p=0.2561, two-tailed unpaired t test. Dots/squares represent the value for the first spike evoked at rheobase for each neuron. n is the same as in panel C.

Supplementary Figure 3

Supplementary Figure 3. *Cntnap2⁻/⁻* **mice do not exhibit changes in Parvalbumin positive cell number or expression.**

(A) Representative confocal images of dorsolateral striatum labeled with an antibody against Parvalbumin (PV) protein for the indicated genotypes, mice aged P21.

(B) Quantification (mean ± SEM) of the number of PV positive cells per ROI at P21, p=0.2554, two-tailed unpaired t test. *Cntnap2⁺/⁺* n=8 sections from 4 mice (2 sections imaged per mouse), *Cntnap2⁻/⁻* n=8 sections from 4 mice (2 sections imaged per mouse).

(C) Representative confocal images of dorsolateral striatum labeled with an antibody against PV for the indicated genotypes, mice aged postnatal day (P) 40-50.

(D) Quantification (mean ± SEM) of the number of PV positive cells per ROI, p=0.5144, two-tailed unpaired t test. *Cntnap2⁺/⁺* n=24 sections from 12 mice (2 sections imaged per mouse), *Cntnap2⁻/⁻* n=24 sections from 12 mice (2 sections imaged per mouse).

(E) Representative confocal images of dorsal striatum labeled with an antibody against PV for the indicated genotypes, mice aged P40-50.

(F) Quantification (mean ± SEM) of the number of PV positive cells in the whole dorsal striatum at P40-50, p=0.4201, two-tailed unpaired t test. *Cntnap2⁺/⁺* n=8 sections from 4 mice (2 sections imaged per mouse), *Cntnap2⁻/⁻* n=8 sections from 4 mice (2 sections imaged per mouse).

(G) Quantification (mean ± SEM) of the average single cell fluorescence of PV antibody staining in dorsal striatum, p=0.9996, two-tailed unpaired t test. *Cntnap2⁺/⁺* n=8 sections from 4 mice (2 sections imaged per mouse), *Cntnap2⁻/⁻* n=8 sections from 4 mice (2 sections imaged per mouse).

(H) Quantification (mean ± SEM) of the average bulk fluorescence of PV antibody staining in dorsal striatum per section, p=0.4878, two-tailed unpaired t test. *Cntnap2⁺/⁺* n=8 sections from 4 mice (2 sections imaged per mouse), *Cntnap2⁻/⁻* n=8 sections from 4 mice (2 sections imaged per mouse).

(I) Representative western blots for CASPR2, PV, and GAPDH protein in dorsal striatal tissue punches. 10 independent samples per genotype are shown.

(J) Quantification (mean ± SEM) of PV protein expression normalized to GAPDH. Data are presented as a percentage of *Cntnap2⁺/⁺* levels, p=0.1485, two-tailed unpaired t test. *Cntnap2⁺/⁺* n=10 mice, *Cntnap2⁻/⁻* n=10 mice.

(K) Representative confocal images of in situ hybridization for *Pvalb* in dorsolateral striatum for the indicated genotypes, mice aged P40-50.

(L-N) Quantification (mean ± SEM) of the number of *Pvalb* positive cells in the dorsolateral striatum per section (J), p>0.9999, Mann-Whitney test; dorsomedial striatum (K), p=0.7429, Mann-Whitney test; and central striatum (L), p=0.6286, Mann-Whitney test. *Cntnap2⁺/⁺* n=4 sections from 2 mice (2 sections imaged per mouse), *Cntnap2⁻/⁻* n=4 sections from 2 mice (2 sections imaged per mouse).

Supplementary Figure 4

Supplementary Figure 4. Action potential properties are unchanged in *Cntnap2^{-/-}* SPNs.
(A) Phase plane plot of a single AP evoked in dSPNs from *Cntnap2^{+/+}* (black) and *Cntnap2^{-/-}* (orange) mice.
(B-D) Quantification (mean ± SEM) of dSPN AP threshold (B), p=0.0516, two-tailed unpaired t test; AP height (C), p=0.0833, two-tailed unpaired t test; and AP width at half max (D), p=0.3792, two-tailed unpaired t test. Dots/squares represent the value for the first spike evoked at rheobase for each neuron. *Cntnap2^{+/+}* n=22 cells from 8 mice, *Cntnap2^{-/-}* n=23 cells from 8 mice.
(E) Phase plane plot of a single AP evoked in iSPNs in *Cntnap2^{+/+}* (green) and *Cntnap2^{-/-}* (blue) mice.
(F-H) Quantification (mean ± SEM) of iSPN AP threshold (F), p=0.1495, two-tailed unpaired t test; AP height (G), p=0.1794, Mann-Whitney test; and AP width at half max (H), p=0.2409, two-tailed unpaired t test. Dots/squares represent the value for the first spike evoked at rheobase for each neuron. *Cntnap2^{+/+}* n=22 cells from 8 mice, *Cntnap2^{-/-}* n=21 cells from 8 mice.

Supplementary Figure 5

Supplementary Figure 5. DeepLabCut and Keypoint-MoSeq analysis of *Cntnap2*^{-/-} mice.
(A) Quantification (mean ± SEM) of the total distance traveled in the open field measured by
DeepLabCut, p=0.8201, Mann-Whitney test; *Cntnap2*$^{+/+}$ n=13 mice and *Cntnap2*$^{-/-}$ n=11 mice.
(B) Quantification (mean ± SEM) of the number of grooming bouts in the open field assessed by
Keypoint-MoSeq, calculated as the number of instances of all grooming syllables. *Cntnap2*$^{+/+}$
n=13 mice and *Cntnap2*$^{-/-}$ n =11 mice, *p = 0.0352, Mann-Whitney test.
(C) Map depicting the difference in transitions between syllables (top 25 most frequent syllables)
for *Cntnap2*$^{+/+}$ and *Cntnap2*$^{-/-}$ mice (*Cntnap2*$^{-/-}$ minus *Cntnap2*$^{+/+}$). A red circle around a syllable
number indicates increased usage of that syllable in *Cntnap2*$^{-/-}$ mice, a blue circle indicates
decreased usage of that syllable in *Cntnap2*$^{-/-}$ mice. A red line between syllable numbers
indicates increased usage of that syllable transition in *Cntnap2*$^{-/-}$ mice, a blue line indicates
decreased usage of that syllable transition in *Cntnap2*$^{-/-}$ mice. The thickness of the line scales
with the size of the difference and the size of the circle surrounding the syllable number scales
with usage of the syllable.

Supplementary Table 1. Summary of behavior data by sex and genotype

Group	N (mice)	Mean	Std. deviation	Std. error of the mean
Open field				
Cntnap2$^{+/+}$ male distance traveled	22	127.7 m	54.34	11.58
Cntnap2$^{-/-}$ male distance traveled	16	156 m	28.36	7.091
Cntnap2$^{+/+}$ female distance traveled	19	139.2 m	48.71	11.17
Cntnap2$^{-/-}$ female distance traveled	18	131.7 m	57.52	13.56
Cntnap2$^{+/+}$ male avg. speed	22	0.03536 m/s	0.01500	0.003198
Cntnap2$^{-/-}$ male avg. speed	16	0.04331 m/s	0.007964	0.001991
Cntnap2$^{+/+}$ female avg. speed	19	0.03868 m/s	0.01341	0.003076
Cntnap2$^{-/-}$ female avg. speed	18	0.03644 m/s	0.01590	0.003749
Cntnap2$^{+/+}$ male rears	22	557.6 rears	201.8	43.03
Cntnap2$^{-/-}$ male rears	16	462.8 rears	173.0	43.25
Cntnap2$^{+/+}$ female rears	19	361.2 rears	188.3	43.19
Cntnap2$^{-/-}$ female rears	18	355.1 rears	158.8	37.42
Cntnap2$^{+/+}$ male center entries	22	254.3 entries	123.3	26.29
Cntnap2$^{-/-}$ male center entries	16	350.8 entries	102.4	25.59
Cntnap2$^{+/+}$ female center entries	19	204.8 entries	71.51	16.41
Cntnap2$^{-/-}$ female center entries	18	247.6 entries	88.30	20.81
Cntnap2$^{+/+}$ male number of grooming bouts (20 min.)	22	14.91 bouts	8.602	1.834
Cntnap2$^{-/-}$ male number of grooming bouts (20 min.)	16	19.56 bouts	11.52	2.879
Cntnap2$^{+/+}$ female number of grooming bouts (20 min.)	19	13.37 bouts	6.353	1.457
Cntnap2$^{-/-}$ female number of grooming bouts (20 min.)	18	28.44 bouts	18.99	4.475

Marble burying				
Cntnap2$^{+/+}$ male marbles buried	16	8.125 marbles	6.206	1.552
Cntnap2$^{-/-}$ male marbles buried	17	10.82 marbles	5.187	1.258
Cntnap2$^{+/+}$ female marbles buried	17	8.235 marbles	5.641	1.368
Cntnap2$^{-/-}$ female marbles buried	16	11.25 marbles	5.310	1.328
Holeboard				
Cntnap2$^{+/+}$ male 10 min. nose pokes	11	156.5 pokes	21.18	6.385
Cntnap2$^{-/-}$ male 10 min. nose pokes	11	192.3 pokes	38.79	11.70
Cntnap2$^{+/+}$ female 10 min. nose pokes	14	175.9 pokes	37.51	10.03
Cntnap2$^{-/-}$ female 10 min. nose pokes	11	180.5 pokes	47.50	14.32
Cntnap2$^{+/+}$ male first 5 min. nose pokes	11	105.4 pokes	16.26	4.901
Cntnap2$^{-/-}$ male first 5 min. nose pokes	11	114.5 pokes	30.25	9.121
Cntnap2$^{+/+}$ female first 5 min. nose pokes	14	110.5 pokes	22.93	6.128
Cntnap2$^{-/-}$ female first 5 min. nose pokes	11	112.3 pokes	30.09	9.073
Cntnap2$^{+/+}$ male last 5 min. nose pokes	11	51.18 pokes	10.09	3.042
Cntnap2$^{-/-}$ male last 5 min. nose pokes	11	77.73 pokes	14.80	4.462
Cntnap2$^{+/+}$ female last 5 min. nose pokes	14	65.36 pokes	20.82	5.565
Cntnap2$^{-/-}$ female last 5 min. nose pokes	11	68.27 pokes	20.15	6.077
Rotarod				
Cntnap2$^{+/+}$ male learning rate	15	1.839 RPM/day	1.416	0.3655
Cntnap2$^{-/-}$ male learning rate	14	2.856 RPM/day	1.166	0.3115
Cntnap2$^{+/+}$ female learning rate	15	2.162 RPM/day	1.083	0.2798
Cntnap2$^{-/-}$ female learning rate	15	3.607 RPM/day	1.103	0.2849
Four choice reversal learning				
Cntnap2$^{+/+}$ male	5	20.80 trials	6.834	3.056

acquisition trials				
Cntnap2^{-/-} male acquisition trials	5	20 trials	7.106	3.178
Cntnap2^{+/+} female acquisition trials	5	17.20 trials	4.550	2.035
Cntnap2^{-/-} female acquisition trials	5	19.60 trials	2.302	1.030
Cntnap2^{+/+} male recall trials	5	11 trials	1.871	0.8367
Cntnap2^{-/-} male recall trials	5	11 trials	4.243	1.897
Cntnap2^{+/+} female recall trials	5	10.60 trials	1.140	0.5099
Cntnap2^{-/-} female recall trials	5	10.20 trials	1.095	0.4899
Cntnap2^{+/+} male reversal trials	5	18.40 trials	1.817	0.8124
Cntnap2^{-/-} male reversal trials	5	30.40 trials	4.037	1.806
Cntnap2^{+/+} female reversal trials	5	26.40 trials	9.737	4.354
Cntnap2^{-/-} female reversal trials	5	32.60 trials	4.827	2.159
Cntnap2^{+/+} male reversal errors	5	7.400 errors	0.8944	0.4000
Cntnap2^{-/-} male reversal errors	5	17.60 errors	2.608	1.166
Cntnap2^{+/+} female reversal errors	5	12.20 errors	6.380	2.853
Cntnap2^{-/-} female reversal errors	5	15.20 errors	1.789	0.8000
DeepLabCut Keypoint-MoSeq Open Field				
Cntnap2^{+/+} male distance traveled	7	272 m	6112	2310
Cntnap2^{-/-} male distance traveled	5	232 m	4876	2181
Cntnap2^{+/+} female distance traveled	6	202 m	1604	654.9
Cntnap2^{-/-} female distance traveled	6	227 m	4651	1899
Cntnap2^{+/+} male number of grooming bouts (60 min.)	7	324.7 bouts	166.1	62.77
Cntnap2^{-/-} male number of grooming bouts (60	5	411.6 bouts	131.6	58.84

min.)				
Cntnap2[+/+] female number of grooming bouts (60 min.)	6	361.0 bouts	92.18	37.63
Cntnap2[-/-] female number of grooming bouts (60 min.)	6	534.0 bouts	206.9	84.47

Experimental procedures

Mice

All animal procedures were conducted in accordance with protocols approved by the University of California, Berkeley Institutional Animal Care and Use Committee (IACUC) and Office of Laboratory Animal Care (OLAC) (AUP-2016-04-8684-2). *Cntnap2*$^{-/-}$ mice and littermate *Cntnap2*$^{+/+}$ controls with the following alleles were utilized for each experiment.

Table 1. Summary of mouse lines used.

Experiment	Mouse line	Allele 1 (reference; JAX strain #)	Allele 2 (reference; JAX strain #)	Allele 3 (reference; JAX strain #)	Allele 4 (reference; JAX strain #)
Corticostriatal transmission (Fig. 1)	*Cntnap2*;D1-tdTomato;Thy1-ChR2	*Cntnap2* (Poliak et al., 2003); #017482)	Drd1a-tdTomato (Ade et al., 2011); #016204)	Thy1-ChR2-YFP (Arenkiel et al., 2007); #007612)	
Broad inhibition (Fig. 2A-E)	*Cntnap2*;D1-tdTomato	*Cntnap2* (Poliak et al., 2003); #017482)	Drd1a-tdTomato (Ade et al., 2011); #016204)		
PV-specific inhibition (Fig. 2F-J)	*Cntnap2*;D1-tdTomato;PV-Cre;Ai32	*Cntnap2* (Poliak et al., 2003); #017482)	Drd1a-tdTomato (Ade et al., 2011); #016204)	*Pvalb*-Cre (Hippenmeyer et al., 2005); #017320)	Ai32 (Madisen et al., 2012); #012569)
SPN intrinsic excitability (Fig. 3, S4)	*Cntnap2*;D1-tdTomato	*Cntnap2* (Poliak et al., 2003); #017482)	Drd1a-tdTomato (Ade et al., 2011); #016204)		
Behavior experiments* (Fig. 4-6, S5)	*Cntnap2*;D1-tdTomato	*Cntnap2* (Poliak et al., 2003); #017482)	Drd1a-tdTomato (Ade et al., 2011); #016204)		
Morphology, PV cell counting (Fig. S1, S3A-H)	*Cntnap2*;D1-tdTomato	*Cntnap2* (Poliak et al., 2003); #017482)	Drd1a-tdTomato (Ade et al., 2011); #016204)		
PV intrinsic excitability (Fig. S2)	*Cntnap2*;PV-Cre;Ai9	*Cntnap2* (Poliak et al., 2003); #017482)	*Pvalb*-Cre (Hippenmeyer et al., 2005); #017320)	Ai9 (Madisen et al., 2010); #007909)	

PV in situ, western blot (Fig. S3I-N)	*Cntnap2*	*Cntnap2* (Poliak et al., 2003); #017482)			

*Littermate animals both positive and negative for D1-tdTomato were used in behavior experiments

Mice were group housed on a 12 h light/dark cycle (dark cycle 9:00 AM – 9:00 PM) and given ad libitum access to standard rodent chow and water. Both male and female animals were used for experimentation. The ages, sexes, and numbers of mice used for each experiment are indicated in the respective method details and figure legends. All mice used for experiments were heterozygous or hemizygous for the Drd1a-tdTomato, Thy1-ChR2-YFP, PV-Cre, Ai32, or Ai9 transgenes to avoid potential physiological or behavioral alterations.

Electrophysiology

Mice (P50-60) were briefly anesthetized with isoflurane and perfused transcardially with ice-cold ACSF (pH = 7.4) containing (in mM): 127 NaCl, 25 NaHCO3, 1.25 NaH2PO4, 2.5 KCl, 1 MgCl2, 2 CaCl2, and 25 glucose, bubbled continuously with carbogen (95% O2 and 5% CO2). Brains were rapidly removed and coronal slices (275 μm) were cut on a VT1000S vibratome (Leica) in oxygenated ice-cold choline-based external solution (pH = 7.8) containing (in mM): 110 choline chloride, 25 NaHCO3, 1.25 NaHPO4, 2.5 KCl, 7 MgCl2, 0.5 CaCl2, 25 glucose, 11.6 sodium ascorbate, and 3.1 sodium pyruvate. Slices were recovered in ACSF at 36°C for 15 min and then kept at room temperature (RT) before recording. Recordings were made with a MultiClamp 700B amplifier (Molecular Devices) at RT using 3-5 MOhm glass patch electrodes (Sutter, #BF150-86-7.5). Data were acquired using ScanImage software, written and maintained by Dr. Bernardo Sabatini (https://github. com/bernardosabatini/ SabalabAcq). Traces were analyzed in Igor Pro (Wavemetrics). Recordings with a series resistance > 25 MOhms or holding current above 200 pA were rejected.

Current-clamp recordings

Current clamp recordings were made using a potassium-based internal solution (pH = 7.4) containing (in mM): 135 KMeSO4, 5 KCl, 5 HEPES, 4 Mg-ATP, 0.3 Na-GTP, 10 phosphocreatine, and 1 EGTA. For corticostriatal excitability experiments, optogenetic stimulation consisted of a full-field pulse of blue light (470 nm, 0.5 ms pulse width, CoolLED) through a 63x objective (Olympus, LUMPLFLN60XW). Light power was linear over the range of intensities tested. No synaptic blockers were included. For intrinsic excitability experiments (both SPN and PV interneuron), NBQX (10 μM, Tocris, #1044), CPP (10 μM, Tocris, #0247) and picrotoxin (50 μM, Abcam, #120315) were added to the external solution to block synaptic transmission. One second depolarizing current steps were applied to induce action potentials. No holding current was applied to the membrane.

Voltage-clamp recordings

Voltage-clamp recordings were made using a cesium-based internal solution (pH = 7.4) containing (in mM): 120 CsMeSO4, 15 CsCl, 10 TEA-Cl, 8 NaCl, 10 HEPES, 1 EGTA, 5 QX-314, 4 Mg-ATP, and 0.3 Na-GTP. Recordings were acquired with the amplifier Bessel filter set at 3 kHz. Corticostriatal synaptic stimulation experiments to measure evoked AMPA-mediated EPSCs were performed in picrotoxin (50 µM), and optogenetic stimulation consisted of a full-field pulse of blue light (470 nm, 0.15 ms pulse width) through a 63x objective. Synaptic stimulation experiments to measure evoked IPSCs were performed in NBQX (10 µM) and CPP (10 µM). For electrically evoked IPSCs, a concentric bipolar stimulating electrode (FHC, #30214) was placed in dorsal striatum, roughly 200 µm medial to the recording site in dorsolateral striatum, and a 0.15 ms stimulus was applied. For PV-interneuron optically evoked IPSCs, a full-field pulse of blue light (470 nm, 0.15 ms pulse width) was applied through a 63x objective at the recording site.

Dendritic imaging and spine analysis

Neonatal (P1-3) *Cntnap2*$^{-/-}$;D1-tdT and *Cntnap2*$^{+/+}$;D1-tdT mice were cryoanesthetized and injected bilaterally with 200 nL AAV1.hSyn.eGFP.WPRE.bGH (Penn Vector Core, #p1696 (Keaveney et al., 2018)), diluted 1:75 in saline to achieve sparse transduction. Injections were targeted to the dorsal striatum, with coordinates approximately 1.3 mm lateral to midline, 2.0 mm posterior to bregma, and 1.5 mm ventral to the head surface. At P50-60, mice were anesthetized with isoflurane and transcardial perfusion was performed with 10 mL of 1x PBS followed by 10 mL of ice cold 4% PFA (EMS, #15710-S) in 1x PBS. Brains were post-fixed in 4% PFA in 1x PBS overnight at 4° C. 80 µm coronal sections were made using a freezing microtome (American Optical, AO 860) and stored in 1x PBS at 4° C. Sections were blocked for 1 hour at RT in BlockAid (ThermoFisher, #B10710) and incubated for 48 hours with gentle shaking at 4° C with antibodies against GFP (1:2500, Abcam, #13970) and RFP (1:1000, Rockland (VWR, #600-401-379) diluted in PBS-Tx (1x PBS with 0.25% Triton X-100 (Sigma, #T8787). Sections were washed 3 x 10 min in PBS-Tx and incubated with gentle shaking for 1 hour at RT with Alexa Fluor 488 and 546 secondary antibodies (1:500, Invitrogen, #A11039, #A11035). Sections were washed 3 x 10 min in 1x PBS and mounted onto SuperFrost slides (VWR, #48311- 703) using VECTASHIELD HardSet Antifade Mounting Medium (Vector Laboratories, #H-1400-10). Z stack images of individual dendrites were taken on a confocal microscope (Olympus FLUOVIEW FV3000) with a 60x oil immersion objective (Olympus #1-U2B832) at 2.5x zoom with a step size of 0.4 µm and deconvoluted using Olympus CellSens software. To quantify spine density, dendrites and spines were reconstructed using the FilamentTracer module in Imaris software (Oxford Instruments). The spine density of each dendrite was calculated using Imaris. Dendrites analyzed varied in total length, but excluded the most proximal and distal portions of the dendrite.

Brain sectioning and immunohistochemistry

Adult mice were perfused as above and brains were post-fixed with 4% paraformaldehyde overnight, then sectioned coronally at 30 µm. For immunohistochemistry,

individual wells of sections were washed for 3 x 5 min with 1x PBS, then blocked for 1 hour at RT with BlockAid blocking solution. Primary antibodies diluted in PBS-Tx were added and tissue was incubated for 48 hours with gentle shaking at 4° C. Sections were then washed 3 x 10 min with PBS-Tx. Secondary antibodies diluted 1:500 in PBS-Tx were added and incubated with gentle shaking for 1 hour at RT. Sections were washed 3 x 10 min in 1x PBS. Sections were mounted onto SuperFrost slides (VWR, #48311- 703) and coverslipped with VECTASHIELD HardSet with DAPI (Vector Laboratories, #H-1500-10) or VECTASHIELD HardSet Antifade Mounting Medium (Vector Laboratories, #H-1400-10). The following antibodies were used: mouse anti-PV (1:1000, Sigma, #P3088), rabbit anti-PV (1:1000, Abcam, #11427), anti-RFP (1:500, Rockland, #600-401-379), Alexa Fluor 405, 488 and 546 conjugated secondary antibodies (1:500, Invitrogen, #A-31553, #A-11001, #A-11003, and #A-11035).

PV cell counting

To count PV+ interneurons, Z-stack images of immunostained striatal sections were taken on a confocal microscope (Olympus FLUOVIEW FV3000) with a 10x or 20x objective (Olympus # 1-U2B824 or Olympus # 1-U2B825) and step size of 2 μm. For quantification, image stacks were Z-projected to maximum intensity using Fiji (ImageJ) and cropped to a 400 μm x 400 μm image in anatomically matched sections of the DLS. All PV-expressing cells within this region were counted using the ROI manager tool in ImageJ. Designation of a cell as PV positive was determined by the experimenter and consistently maintained across animals. Experimenter was blind to genotype and ROIs were made on the DAPI channel to avoid selecting regions based on PV expression. To quantify bulk PV fluorescence, ROIs were manually defined in ImageJ using the Freehand tool to cover as much of the DLS as possible, and mean fluorescence intensity was measured. To quantify individual cell PV fluorescence, ROIs were manually defined around every PV positive cell in the previosuly drawn DLS ROI using the Freehand tool, and mean fluorescence intensity was measured.

<u>Western Blot</u>

Adult mice (P48-55) were deeply anesthetized with isoflurane and decapitated. Brains were rapidly dissected and 1.5 mm dorsal striatum punches (Biopunch, Ted Pella, #15111-15) were collected from both hemispheres, flash-frozen in liquid nitrogen, and stored at –80° C. On the day of analysis, frozen samples were sonicated (QSonica Q55) until homogenized in 200 μl lysis buffer containing 1% SDS in 1x PBS with Halt phosphatase inhibitor cocktail (Thermo Fisher Scientific, #PI78420) and Complete mini EDTA-free protease inhibitor cocktail (Roche, #4693159001). Sample homogenates were then boiled on a heat block at 95° C for 5 min and allowed to cool to RT. Total protein content was determined using a BCA assay (Thermo Fisher Scientific, #23227). Following the BCA assay, protein homogenates were mixed with 4x Laemmli sample buffer (BioRad, #161-0747). 12.5μg of total protein per sample were then loaded onto 12% Criterion TGX gels (BioRad, #5671044) and run at 65 V. Proteins were transferred to a PVDF membrane (BioRad, #1620177) at 11 V for 14 hours at 4° C using the BioRad Criterion Blotter (BioRad, #1704070). Membranes (BioRad, #1620177) were briefly reactivated in methanol and rinsed in water 3x. After rinsing, membranes were blocked in 5%

milk in 1x TBS with 1% Tween (TBS-Tween) for 1 hour at RT before being incubated with primary antibodies diluted in 5% milk in TBS-Tween overnight at 4° C. The following day, after 3 x 10 min washes with TBS-Tween, membranes were incubated with secondary antibodies for 1 hour at RT. Following 6 × 10 min washes, membranes were incubated with chemiluminescence substrate (PerkinElmer #NEL105001EA) for 1 min and exposed to Amersham Hyperfilm ECL (VWR, #95017-661).

Bands were quantified by densitometry using ImageJ software. GAPDH was used to normalize protein content and data is expressed as a percentage of control within a given experiment. The following antibodies were used: anti-Caspr2 (1:5000, Abcam, #153856), anti-PV (1:2500, Abcam, #11427), anti-GAPDH (1:5000, Cell Signaling, #51745S), and anti-rabbit goat HRP conjugate (1:5000, BioRad, #1705046).

In situ hybridization

Fluorescent in situ hybridization was performed to quantify *Pvalb* mRNA expression in the striatum of *Cntnap2*$^{+/+}$ and *Cntnap2*$^{-/-}$ mice. Mice were briefly anesthetized with isoflurane and brains were harvested, flash-frozen in OCT mounting medium (Thermo Fisher Scientific, #23-730-571) on dry ice and stored at -80° C for up to 6 months. 16 µm sections were collected using a cryostat (Thermo Fisher Scientific, Microm HM 550), mounted directly onto Superfrost Plus glass slides (VWR, #48311-703) and stored at -80° C for up to 6 months. In situ hybridization was performed according to the protocols provided with the RNAscope Multiplex Fluorescent Reagent Kit (ACD, #323100). *Drd1a* mRNA was visualized with a probe in channel 2 (ACD, #406491-C2) and *Pvalb* mRNA in channel 3 (ACD, #421931-C3). After incubation, sections were secured on slides using ProLong Gold Antifade Mountant with DAPI (Invitrogen, P36935) and 60 x 24 mm rectangular glass coverslips (VWR, #16004-096). Sections were imaged on an Olympus FluoView 3000 confocal microscope using a 10x objective with 1.5x zoom and a step size of 2 µm. *Pvalb*-expressing cells were quantified across the entire striatum using the ROI manager tool in ImageJ. A cell was considered *Pvalb* positive if over 50% of the cell contained fluorescent puncta when compared to the DAPI channel. Experimenter was blind to genotype.

Behavioral analysis

All behavior studies were carried out in the dark phase of the light cycle under red lights (open field) or white lights (marble burying, holeboard, rotarod, and four choice reversal learning). Mice were habituated to the behavior testing room for at least 30 min prior to testing. Mice were given at least one day between different tests. All behavior equipment was cleaned between each trial and mouse with 70% ethanol and rinsed in diluted soap followed by water at the end of the day. If male and female mice were to be tested on the same day, male mice were run first then returned to the housing room, after which all equipment was thoroughly cleaned prior to bringing in female mice for habituation. Behavioral tests were performed with young adult male and female mice (7-11 weeks old). The experimenter was blind to genotype throughout the testing and scoring procedures.

Open field assay

        Exploratory behavior in a novel environment and general locomotor activity were assessed by a 60 min session in an open field chamber (40 cm L x 40 cm W x 34 cm H) made of transparent plexiglass. Horizontal infrared photobeams (Stoelting, 60001-02A) were positioned to detect rearing. The mouse was placed in the bottom right-hand corner of the arena and behavior was recorded using an overhead camera and analyzed using ANY-maze software (Stoelting). An observer manually scored self-grooming behavior during the first 20 minutes of the test. A grooming bout was defined as an unbroken series of grooming movements, including licking of body, paws, or tail, as well as licking of forepaws followed by rubbing of face with paws.

Open field assay with DeepLabCut Keypoint-MoSeq analysis

        Mice were placed in the open field arena and video recorded with a monochrome camera (FLIR Grasshopper 3, GS3-U3-41C6NIR-C) and a 16 mm wide angle lens (Kowa, LM16HC) placed above the arena from a height of 50 cm. To extract the body part (keypoint) coordinates from the video recordings, DeepLabCut (DLC) 2.3.4 (Mathis, et al. 2018; Nath, et al. 2019) was used. Fourteen body parts including nose, head, left ear, right ear, left forelimb, right forelimb, spine 1, spine 2, spine 3, left hindlimb, right hindlimb, tail 1, tail 2, and tail 3 were manually labeled on a small subset of the video frames. A DLC model was then trained using the annotated frames to label those 14 body parts for all videos recorded. The total distance traveled, and number of center entries were calculated using the coordinate of bodypart tail 1. Discrete behavior syllables were extracted using Keypoint-MoSeq 0.4.4 (Weinreb, et al. 2023). Syllable usage and transition data were obtained using built-in functions of the Keypoint-MoSeq package. Decoding and entropy analysis were performed using customized Python 3.9 script. Code available upon request in Github. Entropy was calculated using the following equation, where u_i denotes the frequency of the syllable i and $p_{i,j}$ denotes the transition probability from syllable i to syllable j.:

$$Entropy = -\sum_{i,j} u_i \times p_{i,j} \times \log_2 p_{i,j}$$

Marble burying assay

        The marble burying assay was used to test for repetitive behavior. 20 black marbles were arranged in an orderly 4 x 5 grid on top of 5 cm of clean corn cob bedding in a standard mouse cage. Overhead room lights were on and white noise was played to induce mild stress. Mice were placed in the cage with the marbles for 30 minutes. The number of unburied marbles (>50% exposed) was recorded after the session.

Holeboard assay

        The holeboard assay was used to measure exploratory and repetitive behavior. The holeboard apparatus consisted of a smooth, flat, opaque gray plastic platform, suspended 10 cm from the base by four plastic pegs in each corner. The board contained 16 evenly spaced 2 cm diameter holes and was surrounded by a 30 cm high clear plastic square encasing. During

testing, mice were placed into the center of the holeboard. Mice explored the board for 10 minutes while video was recorded from both an above and side-view camera. Videos were used post-hoc to manually count and map the number of nose pokes made during the task. Nose pokes were defined as the mouse's nose passing through the board barrier when viewed through the side-view camera.

Accelerating rotarod assay

The accelerating rotarod test was used to examine motor coordination and learning. Mice were trained on a rotarod apparatus (Ugo Basile, #47650) for four consecutive days. Three trials were completed per day with a 5 min break between trials. The rotarod was accelerated from 5-40 revolutions per minute (rpm) over 300 s for trials 1-6 (days 1 and 2), and from 10-80 rpm over 300 s for trials 7-12 (days 3 and 4). On the first testing day, mice were first acclimated to the apparatus by being placed on the rotarod rotating at a constant 5 rpm for 60 s and returned to their home cage for 5 min prior to starting trial 1. Latency to fall, or to rotate off the top of the rotarod barrel, was measured by the rotarod stop-trigger timer.

Four choice odor-based reversal learning test

The four-choice odor-based reversal learning test was used to assess learning and cognitive flexibility. Animals were food restricted for 6 days in total, with unrestricted access to drinking water, and maintained at 90-95% of ad lib feeding body weight. Food was given at the end of the day once testing was completed. Food restriction and introduction to Froot Loop cereal piece (Kellogg's, Battle Creek, MI) began 48 hours before pre-training. The four-choice test was performed in a custom-made square box (30.5 cm L × 30.5 cm W × 23 cm H) constructed of clear acrylic. Four internal walls 7.6 cm wide partially divided the arena into four quadrants. A 15.2 cm diameter removable cylinder fit in the center of the maze and was lowered between trials (after a digging response) to isolate the mouse from the rest of the maze. Odor stimuli were presented mixed with wood shavings in white ceramic pots measuring 7.3 cm in diameter and 4.5 cm deep. All pots were sham baited with a piece of Froot Loop cereal secured underneath a mesh screen at the bottom of the pot. This was to prevent mice from using the odor of the Froot Loop to guide their choice. The apparatus was cleaned with 2.5% acetic acid followed by water and the pots were cleaned with 70% ethanol followed by water between mice. The apparatus was cleaned with diluted soap and water at the end of each testing day.

On the first habituation day of pre-training (day 1), animals were allowed to freely explore the testing arena for 30 min and consume small pieces of Froot Loops placed inside empty pots positioned in each of the four corners. On the second shaping day of pre-training (day 2), mice learned to dig to find cereal pieces buried in unscented coarse pine wood shavings (Harts Mountain Corporation, Secaucus, NJ). A single pot was used and increasing amounts of unscented wood shavings were used to cover each subsequent cereal reward. The quadrant containing the pot was alternated on each trial and all quadrants were rewarded equally. Trials were untimed and consisted of (in order): two trials with no shavings, two trials with a dusting of shavings, two trials with the pot a quarter full, two trials with the pot half full,

and four trials with the cereal piece completely buried by shavings. The mouse was manually returned to the center cylinder between trials.

On the days for odor discrimination (day 3, acquisition) and reversal (day 4), wood shavings were freshly scented on the day of testing. Anise extract (McCormick, Hunt Valley, MD) was used undiluted at 0.02 ml/g of shavings. Clove, litsea, and eucalyptus oils (San Francisco Massage Supply Co., San Francisco, CA) were diluted 1:10 in mineral oil and mixed at 0.02 ml/g of shavings. Thymol (thyme; Alfa Aesar, A14563) was diluted 1:20 in 50% ethanol and mixed at 0.01 ml/g of shavings. During the discrimination phase (day 3), mice had to discriminate between four pots with four different odors and learn which one contained a buried food reward. Each trial began with the mouse confined to the start cylinder. Once the cylinder was lifted, timing began, and the mouse could freely explore the arena until it chose to dig in a pot. Digging was defined as purposefully moving the shavings with both front paws. A trial was terminated if no choice was made within 3 min and recorded as omission. Criterion was met when the animal completed eight out of ten consecutive trials correctly. The spatial location of the odors was shuffled on each trial. The rewarded odor during acquisition was anise.

The first four odor choices made during acquisition were analyzed to determine innate odor preference by the percentage of choices for a given odor: $Cntnap2^{+/+}$ mice: 60% thyme, 25% anise, 12.5% clove, and 2.5% litsea. $Cntnap2^{-/-}$ mice: 47.5% thyme, 45% anise, 7.5% clove, 0% litsea. We note that both $Cntnap2^{+/+}$ and $Cntnap2^{-/-}$ mice exhibited the strongest innate preference for thyme, an unrewarded odor. There were no significant differences in innate odor preference.

The reversal phase of the task was carried out on day 4. Mice first performed the task with the same rewarded odor as the discrimination day to ensure they learned and remembered the task. After reaching criterion on recall (eight out of ten consecutive trials correct), the rewarded odor was switched, and mice underwent a reversal learning test in which a previously unrewarded odor (clove) was rewarded. A novel odor (eucalyptus) was also introduced, which replaced thyme. Perseverative errors were choices to dig in the previously rewarded odor that was no longer rewarded. Regressive errors were choosing the previously rewarded odor after the first correct choice of the newly rewarded odor. Novel errors were choices to dig in the pot with the newly introduced odor (eucalyptus). Irrelevant errors were choices to dig in the pot that had never been rewarded (litsea). Omissions were trials in which the mouse failed to make a digging choice within 3 min from the start of the trial. Total errors were the sum of perseverative, regressive, irrelevant, novel, and omission errors. Criterion was met when the mouse completed eight out of ten consecutive trials correctly. The spatial location of the odors was shuffled on each trial.

Quantification and statistical analysis

Experiments were designed to compare the main effect of genotype. The sample sizes were based on prior studies and are indicated in the figure legend for each experiment. Whenever possible, quantification and analyses were performed blind to genotype. GraphPad Prism version 10 was used to perform statistical analyses. The statistical tests and outcomes for each experiment are indicated in the respective figure legend. Two-tailed unpaired t tests were

used for comparisons between two groups. For data that did not pass the D'Agostino & Pearson normality test, a Mann-Whitney test was used. Two-way ANOVAs were used to compare differences between groups for experiments with two independent variables. Statistical significance was defined in the figure panels as follows: $*p < 0.05$, $**p < 0.01$, $***p < 0.001$.

Acknowledgements
This work was supported by SFARI research grant #514428 to H.S.B. and NIH fellowship #F31NS124499 to K.R.C.

Conflicts of interest
The authors declare no competing interests.

References

Ade, K. K., Wan, Y., Chen, M., Gloss, B., & Calakos, N. (2011). An Improved BAC Transgenic Fluorescent Reporter Line for Sensitive and Specific Identification of Striatonigral Medium Spiny Neurons. *Front Syst Neurosci*, *5*, 32. https://doi.org/10.3389/fnsys.2011.00032

Ahmed, N. Y., Knowles, R., Liu, L., Yan, Y., Li, X., Schumann, U., Wang, Y., Sontani, Y., Reynolds, N., Natoli, R., Wen, J., Del Pino, I., Mi, D., & Dehorter, N. (2023). Developmental deficits of MGE-derived interneurons in the Cntnap2 knockout mouse model of autism spectrum disorder. *Front Cell Dev Biol*, *11*, 1112062. https://doi.org/10.3389/fcell.2023.1112062

Albelda, N., & Joel, D. (2012). Animal models of obsessive-compulsive disorder: exploring pharmacology and neural substrates. *Neurosci Biobehav Rev*, *36*(1), 47-63. https://doi.org/10.1016/j.neubiorev.2011.04.006

Anderson, G. R., Galfin, T., Xu, W., Aoto, J., Malenka, R. C., & Sudhof, T. C. (2012). Candidate autism gene screen identifies critical role for cell-adhesion molecule CASPR2 in dendritic arborization and spine development. *Proc Natl Acad Sci U S A*, *109*(44), 18120-18125. https://doi.org/10.1073/pnas.1216398109

Angoa-Perez, M., Kane, M. J., Briggs, D. I., Francescutti, D. M., & Kuhn, D. M. (2013). Marble burying and nestlet shredding as tests of repetitive, compulsive-like behaviors in mice. *J Vis Exp*(82), 50978. https://doi.org/10.3791/50978

Antoine, M. W., Langberg, T., Schnepel, P., & Feldman, D. E. (2019). Increased Excitation-Inhibition Ratio Stabilizes Synapse and Circuit Excitability in Four Autism Mouse Models. *Neuron*, *101*(4), 648-661 e644. https://doi.org/10.1016/j.neuron.2018.12.026

APA. (2022). *Diagnostic and Statistical Manual of Mental Disorders* (5th ed., text rev. ed.). https://doi.org/https://doi.org/10.1176/appi.books.9780890425787

Arenkiel, B. R., Peca, J., Davison, I. G., Feliciano, C., Deisseroth, K., Augustine, G. J., Ehlers, M. D., & Feng, G. (2007). In vivo light-induced activation of neural circuitry in transgenic mice expressing channelrhodopsin-2. *Neuron*, *54*(2), 205-218. https://doi.org/10.1016/j.neuron.2007.03.005

Benthall, K. N., Cording, K. R., Agopyan-Miu, A., Wong, C. D., Chen, E. Y., & Bateup, H. S. (2021). Loss of Tsc1 from striatal direct pathway neurons impairs endocannabinoid-LTD and enhances motor routine learning. *Cell Rep*, *36*(6), 109511. https://doi.org/10.1016/j.celrep.2021.109511

Bouyer, J. J., Park, D. H., Joh, T. H., & Pickel, V. M. (1984). Chemical and structural analysis of the relation between cortical inputs and tyrosine hydroxylase-containing terminals in rat neostriatum. *Brain Res*, *302*(2), 267-275. https://doi.org/10.1016/0006-8993(84)90239-7

Brumback, A. C., Ellwood, I. T., Kjaerby, C., Iafrati, J., Robinson, S., Lee, A. T., Patel, T., Nagaraj, S., Davatolhagh, F., & Sohal, V. S. (2018). Identifying specific prefrontal neurons that contribute to autism-associated abnormalities in physiology and social behavior. *Mol Psychiatry*, *23*(10), 2078-2089. https://doi.org/10.1038/mp.2017.213

Brunner, D., Kabitzke, P., He, D., Cox, K., Thiede, L., Hanania, T., Sabath, E., Alexandrov, V., Saxe, M., Peles, E., Mills, A., Spooren, W., Ghosh, A., Feliciano, P., Benedetti, M., Luo Clayton, A., & Biemans, B. (2015). Comprehensive Analysis of the 16p11.2 Deletion and Null Cntnap2 Mouse Models of Autism Spectrum Disorder. *PLoS One*, *10*(8), e0134572. https://doi.org/10.1371/journal.pone.0134572

Burke, D. A., Rotstein, H. G., & Alvarez, V. A. (2017). Striatal Local Circuitry: A New Framework for Lateral Inhibition. *Neuron*, *96*(2), 267-284. https://doi.org/10.1016/j.neuron.2017.09.019

Calabresi, P., Picconi, B., Tozzi, A., Ghiglieri, V., & Di Filippo, M. (2014). Direct and indirect pathways of basal ganglia: a critical reappraisal. *Nat Neurosci*, *17*(8), 1022-1030. https://doi.org/10.1038/nn.3743

Cifuentes-Diaz, C., Canali, G., Garcia, M., Druart, M., Manett, T., Savariradjane, M., Guillaume, C., Le Magueresse, C., & Goutebroze, L. (2023). Differential impacts of Cntnap2 heterozygosity and Cntnap2 null homozygosity on axon and myelinated fiber development in mouse. *Front Neurosci*, *17*, 1100121. https://doi.org/10.3389/fnins.2023.1100121

Cording, K. R., & Bateup, H. S. (2023). Altered motor learning and coordination in mouse models of autism spectrum disorder. *Front Cell Neurosci*, *17*, 1270489. https://doi.org/10.3389/fncel.2023.1270489

Cromwell, H. C., & Berridge, K. C. (1996). Implementation of action sequences by a neostriatal site: a lesion mapping study of grooming syntax. *J Neurosci*, *16*(10), 3444-3458. https://doi.org/10.1523/JNEUROSCI.16-10-03444.1996

Dawes, J. M., Weir, G. A., Middleton, S. J., Patel, R., Chisholm, K. I., Pettingill, P., Peck, L. J., Sheridan, J., Shakir, A., Jacobson, L., Gutierrez-Mecinas, M., Galino, J., Walcher, J., Kuhnemund, J., Kuehn, H., Sanna, M. D., Lang, B., Clark, A. J., Themistocleous, A. C., . . . Bennett, D. L. (2018). Immune or Genetic-Mediated Disruption of CASPR2 Causes Pain Hypersensitivity Due to Enhanced Primary Afferent Excitability. *Neuron*, *97*(4), 806-822 e810. https://doi.org/10.1016/j.neuron.2018.01.033

De Rubeis, S., He, X., Goldberg, A. P., Poultney, C. S., Samocha, K., Cicek, A. E., Kou, Y., Liu, L., Fromer, M., Walker, S., Singh, T., Klei, L., Kosmicki, J., Shih-Chen, F., Aleksic, B., Biscaldi, M., Bolton, P. F., Brownfeld, J. M., Cai, J., . . . Buxbaum, J. D. (2014). Synaptic, transcriptional and chromatin genes disrupted in autism. *Nature*, *515*(7526), 209-215. https://doi.org/10.1038/nature13772

Delevich, K., Hoshal, B., Zhou, L. Z., Zhang, Y., Vedula, S., Lin, W. C., Chase, J., Collins, A. G. E., & Wilbrecht, L. (2022). Activation, but not inhibition, of the indirect pathway disrupts choice rejection in a freely moving, multiple-choice foraging task. *Cell Rep*, *40*(4), 111129. https://doi.org/10.1016/j.celrep.2022.111129

Delorme, R., Ey, E., Toro, R., Leboyer, M., Gillberg, C., & Bourgeron, T. (2013). Progress toward treatments for synaptic defects in autism. *Nat Med*, *19*(6), 685-694. https://doi.org/10.1038/nm.3193

Ding, J., Peterson, J. D., & Surmeier, D. J. (2008). Corticostriatal and thalamostriatal synapses have distinctive properties. *J Neurosci*, *28*(25), 6483-6492. https://doi.org/10.1523/JNEUROSCI.0435-08.2008

Doig, N. M., Moss, J., & Bolam, J. P. (2010). Cortical and thalamic innervation of direct and indirect pathway medium-sized spiny neurons in mouse striatum. *J Neurosci*, *30*(44), 14610-14618. https://doi.org/10.1523/JNEUROSCI.1623-10.2010

Ebert, D. H., & Greenberg, M. E. (2013). Activity-dependent neuronal signalling and autism spectrum disorder. *Nature*, *493*(7432), 327-337. https://doi.org/10.1038/nature11860

Estes, A., Shaw, D. W., Sparks, B. F., Friedman, S., Giedd, J. N., Dawson, G., Bryan, M., & Dager, S. R. (2011). Basal ganglia morphometry and repetitive behavior in young children with autism spectrum disorder. *Autism Res*, *4*(3), 212-220. https://doi.org/10.1002/aur.193

Evans, M. M., Kim, J., Abel, T., Nickl-Jockschat, T., & Stevens, H. E. (2024). Developmental Disruptions of the Dorsal Striatum in Autism Spectrum Disorder. *Biol Psychiatry*, *95*(2), 102-111. https://doi.org/10.1016/j.biopsych.2023.08.015

Fernandez, M., Sanchez-Leon, C. A., Llorente, J., Sierra-Arregui, T., Knafo, S., Marquez-Ruiz, J., & Penagarikano, O. (2021). Altered Cerebellar Response to Somatosensory Stimuli in the Cntnap2 Mouse Model of Autism. *eNeuro*, *8*(5). https://doi.org/10.1523/ENEURO.0333-21.2021

Filice, F., Janickova, L., Henzi, T., Bilella, A., & Schwaller, B. (2020). The Parvalbumin Hypothesis of Autism Spectrum Disorder. *Front Cell Neurosci*, *14*, 577525. https://doi.org/10.3389/fncel.2020.577525

Fuccillo, M. V. (2016). Striatal Circuits as a Common Node for Autism Pathophysiology. *Front Neurosci*, *10*, 27. https://doi.org/10.3389/fnins.2016.00027

Gdalyahu, A., Lazaro, M., Penagarikano, O., Golshani, P., Trachtenberg, J. T., & Geschwind, D. H. (2015). Correction: The Autism Related Protein Contactin-Associated Protein-Like 2 (CNTNAP2) Stabilizes New Spines: An In Vivo Mouse Study. *PLoS One*, *10*(5), e0129638. https://doi.org/10.1371/journal.pone.0129638

Gerfen, C. R., & Surmeier, D. J. (2011). Modulation of striatal projection systems by dopamine. *Annu Rev Neurosci*, *34*, 441-466. https://doi.org/10.1146/annurev-neuro-061010-113641

Hartig, W., Brauer, K., & Bruckner, G. (1992). Wisteria floribunda agglutinin-labelled nets surround parvalbumin-containing neurons. *Neuroreport*, *3*(10), 869-872. https://doi.org/10.1097/00001756-199210000-00012

Haunso, A., Ibrahim, M., Bartsch, U., Letiembre, M., Celio, M. R., & Menoud, P. (2000). Morphology of perineuronal nets in tenascin-R and parvalbumin single and double knockout mice. *Brain Res*, *864*(1), 142-145. https://doi.org/10.1016/s0006-8993(00)02173-9

Hawes, S. L., Evans, R. C., Unruh, B. A., Benkert, E. E., Gillani, F., Dumas, T. C., & Blackwell, K. T. (2015). Multimodal Plasticity in Dorsal Striatum While Learning a Lateralized Navigation Task. *J Neurosci*, *35*(29), 10535-10549. https://doi.org/10.1523/JNEUROSCI.4415-14.2015

Hippenmeyer, S., Vrieseling, E., Sigrist, M., Portmann, T., Laengle, C., Ladle, D. R., & Arber, S. (2005). A developmental switch in the response of DRG neurons to ETS transcription factor signaling. *PLoS Biol*, *3*(5), e159. https://doi.org/10.1371/journal.pbio.0030159

Hollander, E., Anagnostou, E., Chaplin, W., Esposito, K., Haznedar, M. M., Licalzi, E., Wasserman, S., Soorya, L., & Buchsbaum, M. (2005). Striatal volume on magnetic resonance imaging and repetitive behaviors in autism. *Biol Psychiatry*, *58*(3), 226-232. https://doi.org/10.1016/j.biopsych.2005.03.040

Inda, M. C., DeFelipe, J., & Munoz, A. (2006). Voltage-gated ion channels in the axon initial segment of human cortical pyramidal cells and their relationship with chandelier cells. *Proc Natl Acad Sci U S A*, *103*(8), 2920-2925. https://doi.org/10.1073/pnas.0511197103

Iossifov, I., O'Roak, B. J., Sanders, S. J., Ronemus, M., Krumm, N., Levy, D., Stessman, H. A., Witherspoon, K. T., Vives, L., Patterson, K. E., Smith, J. D., Paeper, B., Nickerson, D. A., Dea, J., Dong, S., Gonzalez, L. E., Mandell, J. D., Mane, S. M., Murtha, M. T., . . . Wigler, M. (2014). The contribution of de novo coding mutations to autism spectrum disorder. *Nature*, *515*(7526), 216-221. https://doi.org/10.1038/nature13908

Izquierdo, A., Brigman, J. L., Radke, A. K., Rudebeck, P. H., & Holmes, A. (2017). The neural basis of reversal learning: An updated perspective. *Neuroscience*, *345*, 12-26. https://doi.org/10.1016/j.neuroscience.2016.03.021

Johnson, C. M., Peckler, H., Tai, L. H., & Wilbrecht, L. (2016). Rule learning enhances structural plasticity of long-range axons in frontal cortex. *Nat Commun*, *7*, 10785. https://doi.org/10.1038/ncomms10785

Juarez, P., & Martinez Cerdeno, V. (2022). Parvalbumin and parvalbumin chandelier interneurons in autism and other psychiatric disorders. *Front Psychiatry*, *13*, 913550. https://doi.org/10.3389/fpsyt.2022.913550

Jurgensen, S., & Castillo, P. E. (2015). Selective Dysregulation of Hippocampal Inhibition in the Mouse Lacking Autism Candidate Gene CNTNAP2. *J Neurosci*, *35*(43), 14681-14687. https://doi.org/10.1523/JNEUROSCI.1666-15.2015

Keaveney, M. K., Tseng, H. A., Ta, T. L., Gritton, H. J., Man, H. Y., & Han, X. (2018). A MicroRNA-Based Gene-Targeting Tool for Virally Labeling Interneurons in the Rodent Cortex. *Cell Rep*, *24*(2), 294-303. https://doi.org/10.1016/j.celrep.2018.06.049

Koos, T., Tepper, J. M., & Wilson, C. J. (2004). Comparison of IPSCs evoked by spiny and fast-spiking neurons in the neostriatum. *J Neurosci*, *24*(36), 7916-7922. https://doi.org/10.1523/JNEUROSCI.2163-04.2004

Kosillo, P., Doig, N. M., Ahmed, K. M., Agopyan-Miu, A., Wong, C. D., Conyers, L., Threlfell, S., Magill, P. J., & Bateup, H. S. (2019). Tsc1-mTORC1 signaling controls striatal dopamine release and cognitive flexibility. *Nat Commun*, *10*(1), 5426. https://doi.org/10.1038/s41467-019-13396-8

Kravitz, A. V., Freeze, B. S., Parker, P. R., Kay, K., Thwin, M. T., Deisseroth, K., & Kreitzer, A. C. (2010). Regulation of parkinsonian motor behaviours by optogenetic control of basal ganglia circuitry. *Nature*, *466*(7306), 622-626. https://doi.org/10.1038/nature09159

Langen, M., Bos, D., Noordermeer, S. D., Nederveen, H., van Engeland, H., & Durston, S. (2014). Changes in the development of striatum are involved in repetitive behavior in autism. *Biol Psychiatry*, *76*(5), 405-411. https://doi.org/10.1016/j.biopsych.2013.08.013

Lauber, E., Filice, F., & Schwaller, B. (2018). Dysregulation of Parvalbumin Expression in the Cntnap2-/- Mouse Model of Autism Spectrum Disorder. *Front Mol Neurosci*, *11*, 262. https://doi.org/10.3389/fnmol.2018.00262

Lazaro, M. T., Taxidis, J., Shuman, T., Bachmutsky, I., Ikrar, T., Santos, R., Marcello, G. M., Mylavarapu, A., Chandra, S., Foreman, A., Goli, R., Tran, D., Sharma, N., Azhdam, M., Dong, H., Choe, K. Y., Penagarikano, O., Masmanidis, S. C., Racz, B., . . . Golshani, P. (2019). Reduced Prefrontal Synaptic Connectivity and Disturbed Oscillatory Population Dynamics in the CNTNAP2 Model of Autism. *Cell Rep*, *27*(9), 2567-2578 e2566. https://doi.org/10.1016/j.celrep.2019.05.006

Lee, I. B., Lee, E., Han, N. E., Slavuj, M., Hwang, J. W., Lee, A., Sun, T., Jeong, Y., Baik, J. H., Park, J. Y., Choi, S. Y., Kwag, J., & Yoon, B. J. (2024). Persistent enhancement of basolateral amygdala-dorsomedial striatum synapses causes compulsive-like behaviors in mice. *Nat Commun*, *15*(1), 219. https://doi.org/10.1038/s41467-023-44322-8

Li, W., & Pozzo-Miller, L. (2020). Dysfunction of the corticostriatal pathway in autism spectrum disorders. *J Neurosci Res*, *98*(11), 2130-2147. https://doi.org/10.1002/jnr.24560

Lin, W. C., Liu, C., Kosillo, P., Tai, L. H., Galarce, E., Bateup, H. S., Lammel, S., & Wilbrecht, L. (2022). Transient food insecurity during the juvenile-adolescent period affects adult weight, cognitive flexibility, and dopamine neurobiology. *Curr Biol*, *32*(17), 3690-3703 e3695. https://doi.org/10.1016/j.cub.2022.06.089

Madisen, L., Mao, T., Koch, H., Zhuo, J. M., Berenyi, A., Fujisawa, S., Hsu, Y. W., Garcia, A. J., 3rd, Gu, X., Zanella, S., Kidney, J., Gu, H., Mao, Y., Hooks, B. M., Boyden, E. S., Buzsaki, G., Ramirez, J. M., Jones, A. R., Svoboda, K., . . . Zeng, H. (2012). A toolbox of Cre-dependent optogenetic transgenic mice for light-induced activation and silencing. *Nat Neurosci*, *15*(5), 793-802. https://doi.org/10.1038/nn.3078

Madisen, L., Zwingman, T. A., Sunkin, S. M., Oh, S. W., Zariwala, H. A., Gu, H., Ng, L. L., Palmiter, R. D., Hawrylycz, M. J., Jones, A. R., Lein, E. S., & Zeng, H. (2010). A robust and high-throughput Cre reporting and characterization system for the whole mouse brain. *Nat Neurosci*, *13*(1), 133-140. https://doi.org/10.1038/nn.2467

Martin-de-Saavedra, M. D., Dos Santos, M., Culotta, L., Varea, O., Spielman, B. P., Parnell, E., Forrest, M. P., Gao, R., Yoon, S., McCoig, E., Jalloul, H. A., Myczek, K., Khalatyan, N., Hall, E. A., Turk, L. S., Sanz-Clemente, A., Comoletti, D., Lichtenthaler, S. F., Burgdorf, J. S., . . . Penzes, P. (2022). Shed CNTNAP2 ectodomain is detectable in CSF and regulates Ca(2+) homeostasis and network synchrony via PMCA2/ATP2B2. *Neuron*, *110*(4), 627-643 e629. https://doi.org/10.1016/j.neuron.2021.11.025

Mathis, A., Mamidanna, P., Cury, K. M., Abe, T., Murthy, V. N., Mathis, M. W., & Bethge, M. (2018). DeepLabCut: markerless pose estimation of user-defined body parts with deep learning. *Nat Neurosci*, *21*(9), 1281-1289. https://doi.org/10.1038/s41593-018-0209-y

Nisenbaum, E. S., Xu, Z. C., & Wilson, C. J. (1994). Contribution of a slowly inactivating potassium current to the transition to firing of neostriatal spiny projection neurons. *J Neurophysiol*, *71*(3), 1174-1189. https://doi.org/10.1152/jn.1994.71.3.1174

Orduz, D., Bischop, D. P., Schwaller, B., Schiffmann, S. N., & Gall, D. (2013). Parvalbumin tunes spike-timing and efferent short-term plasticity in striatal fast spiking interneurons. *J Physiol*, *591*(13), 3215-3232. https://doi.org/10.1113/jphysiol.2012.250795

Packard, M. G., & Knowlton, B. J. (2002). Learning and memory functions of the Basal Ganglia. *Annu Rev Neurosci*, *25*, 563-593. https://doi.org/10.1146/annurev.neuro.25.112701.142937

Paterno, R., Marafiga, J. R., Ramsay, H., Li, T., Salvati, K. A., & Baraban, S. C. (2021). Hippocampal gamma and sharp-wave ripple oscillations are altered in a Cntnap2 mouse model of autism spectrum disorder. *Cell Rep*, *37*(6), 109970. https://doi.org/10.1016/j.celrep.2021.109970

Peca, J., Feliciano, C., Ting, J. T., Wang, W., Wells, M. F., Venkatraman, T. N., Lascola, C. D., Fu, Z., & Feng, G. (2011). Shank3 mutant mice display autistic-like behaviours and striatal dysfunction. *Nature*, *472*(7344), 437-442. https://doi.org/10.1038/nature09965

Peixoto, R. T., Wang, W., Croney, D. M., Kozorovitskiy, Y., & Sabatini, B. L. (2016). Early hyperactivity and precocious maturation of corticostriatal circuits in Shank3B(-/-) mice. *Nat Neurosci*, *19*(5), 716-724. https://doi.org/10.1038/nn.4260

Penagarikano, O., Abrahams, B. S., Herman, E. I., Winden, K. D., Gdalyahu, A., Dong, H., Sonnenblick, L. I., Gruver, R., Almajano, J., Bragin, A., Golshani, P., Trachtenberg, J. T., Peles, E., & Geschwind, D. H. (2011). Absence of CNTNAP2 leads to epilepsy, neuronal migration abnormalities, and core autism-related deficits. *Cell*, *147*(1), 235-246. https://doi.org/10.1016/j.cell.2011.08.040

Penagarikano, O., & Geschwind, D. H. (2012). What does CNTNAP2 reveal about autism spectrum disorder? *Trends Mol Med*, *18*(3), 156-163. https://doi.org/10.1016/j.molmed.2012.01.003

Platt, R. J., Zhou, Y., Slaymaker, I. M., Shetty, A. S., Weisbach, N. R., Kim, J. A., Sharma, J., Desai, M., Sood, S., Kempton, H. R., Crabtree, G. R., Feng, G., & Zhang, F. (2017). Chd8 Mutation Leads to Autistic-like Behaviors and Impaired Striatal Circuits. *Cell Rep*, *19*(2), 335-350. https://doi.org/10.1016/j.celrep.2017.03.052

Poliak, S., Gollan, L., Martinez, R., Custer, A., Einheber, S., Salzer, J. L., Trimmer, J. S., Shrager, P., & Peles, E. (1999). Caspr2, a new member of the neurexin superfamily, is localized at the juxtaparanodes of myelinated axons and associates with K+ channels. *Neuron*, *24*(4), 1037-1047. https://doi.org/10.1016/s0896-6273(00)81049-1

Poliak, S., Salomon, D., Elhanany, H., Sabanay, H., Kiernan, B., Pevny, L., Stewart, C. L., Xu, X., Chiu, S. Y., Shrager, P., Furley, A. J., & Peles, E. (2003). Juxtaparanodal clustering of Shaker-like K+ channels in myelinated axons depends on Caspr2 and TAG-1. *J Cell Biol*, *162*(6), 1149-1160. https://doi.org/10.1083/jcb.200305018

Ramirez-Armenta, K. I., Alatriste-Leon, H., Verma-Rodriguez, A. K., Llanos-Moreno, A., Ramirez-Jarquin, J. O., & Tecuapetla, F. (2022). Optogenetic inhibition of indirect pathway neurons in the dorsomedial striatum reduces excessive grooming in Sapap3-knockout mice. *Neuropsychopharmacology*, *47*(2), 477-487. https://doi.org/10.1038/s41386-021-01161-9

Rodenas-Cuadrado, P., Ho, J., & Vernes, S. C. (2014). Shining a light on CNTNAP2: complex functions to complex disorders. *Eur J Hum Genet*, *22*(2), 171-178. https://doi.org/10.1038/ejhg.2013.100

Rothwell, P. E., Fuccillo, M. V., Maxeiner, S., Hayton, S. J., Gokce, O., Lim, B. K., Fowler, S. C., Malenka, R. C., & Sudhof, T. C. (2014). Autism-associated neuroligin-3 mutations commonly impair striatal circuits to boost repetitive behaviors. *Cell*, *158*(1), 198-212. https://doi.org/10.1016/j.cell.2014.04.045

Sanders, S. J., He, X., Willsey, A. J., Ercan-Sencicek, A. G., Samocha, K. E., Cicek, A. E., Murtha, M. T., Bal, V. H., Bishop, S. L., Dong, S., Goldberg, A. P., Jinlu, C., Keaney, J. F., 3rd, Klei, L., Mandell, J. D., Moreno-De-Luca, D., Poultney, C. S., Robinson, E. B., Smith, L., . . . State, M. W. (2015). Insights into Autism Spectrum Disorder Genomic Architecture and Biology from 71 Risk Loci. *Neuron*, *87*(6), 1215-1233. https://doi.org/10.1016/j.neuron.2015.09.016

Sandin, S., Lichtenstein, P., Kuja-Halkola, R., Hultman, C., Larsson, H., & Reichenberg, A. (2017). The Heritability of Autism Spectrum Disorder. *JAMA*, *318*(12), 1182-1184. https://doi.org/10.1001/jama.2017.12141

Santos, F. J., Oliveira, R. F., Jin, X., & Costa, R. M. (2015). Corticostriatal dynamics encode the refinement of specific behavioral variability during skill learning. *Elife*, *4*, e09423. https://doi.org/10.7554/eLife.09423

Satterstrom, F. K., Kosmicki, J. A., Wang, J., Breen, M. S., De Rubeis, S., An, J.-Y., Peng, M., Collins, R., Grove, J., Klei, L., Stevens, C., Reichert, J., Mulhern, M. S., Artomov, M., Gerges, S., Sheppard, B., Xu, X., Bhaduri, A., Norman, U., . . . Walters, R. K. (2020). Large-Scale Exome Sequencing Study Implicates Both Developmental and Functional Changes in the Neurobiology of Autism. *Cell*, *180*(3), 568-584.e523. https://doi.org/10.1016/j.cell.2019.12.036

Scott, R., Sanchez-Aguilera, A., van Elst, K., Lim, L., Dehorter, N., Bae, S. E., Bartolini, G., Peles, E., Kas, M. J. H., Bruining, H., & Marin, O. (2019). Loss of Cntnap2 Causes Axonal Excitability Deficits, Developmental Delay in Cortical Myelination, and Abnormal Stereotyped Motor Behavior. *Cereb Cortex*, *29*(2), 586-597. https://doi.org/10.1093/cercor/bhx341

Shen, W., Hernandez-Lopez, S., Tkatch, T., Held, J. E., & Surmeier, D. J. (2004). Kv1.2-containing K+ channels regulate subthreshold excitability of striatal medium spiny neurons. *J Neurophysiol*, *91*(3), 1337-1349. https://doi.org/10.1152/jn.00414.2003

Subramanian, K., Brandenburg, C., Orsati, F., Soghomonian, J. J., Hussman, J. P., & Blatt, G. J. (2017). Basal ganglia and autism - a translational perspective. *Autism Res*, *10*(11), 1751-1775. https://doi.org/10.1002/aur.1837

Tai, L. H., Lee, A. M., Benavidez, N., Bonci, A., & Wilbrecht, L. (2012). Transient stimulation of distinct subpopulations of striatal neurons mimics changes in action value. *Nat Neurosci*, *15*(9), 1281-1289. https://doi.org/10.1038/nn.3188

Uddin, L. Q. (2021). Cognitive and behavioural flexibility: neural mechanisms and clinical considerations. *Nat Rev Neurosci*, *22*(3), 167-179. https://doi.org/10.1038/s41583-021-00428-w

Varea, O., Martin-de-Saavedra, M. D., Kopeikina, K. J., Schurmann, B., Fleming, H. J., Fawcett-Patel, J. M., Bach, A., Jang, S., Peles, E., Kim, E., & Penzes, P. (2015). Synaptic abnormalities and cytoplasmic glutamate receptor aggregates in contactin associated protein-like 2/Caspr2 knockout neurons. *Proc Natl Acad Sci U S A*, *112*(19), 6176-6181. https://doi.org/10.1073/pnas.1423205112

Vogt, D., Cho, K. K. A., Shelton, S. M., Paul, A., Huang, Z. J., Sohal, V. S., & Rubenstein, J. L. R. (2018). Mouse Cntnap2 and Human CNTNAP2 ASD Alleles Cell Autonomously Regulate PV+ Cortical Interneurons. *Cereb Cortex*, *28*(11), 3868-3879. https://doi.org/10.1093/cercor/bhx248

Voorn, P., Vanderschuren, L. J., Groenewegen, H. J., Robbins, T. W., & Pennartz, C. M. (2004). Putting a spin on the dorsal-ventral divide of the striatum. *Trends Neurosci*, *27*(8), 468-474. https://doi.org/10.1016/j.tins.2004.06.006

Wang, X., Bey, A. L., Katz, B. M., Badea, A., Kim, N., David, L. K., Duffney, L. J., Kumar, S., Mague, S. D., Hulbert, S. W., Dutta, N., Hayrapetyan, V., Yu, C., Gaidis, E., Zhao, S., Ding, J. D., Xu, Q., Chung, L., Rodriguiz, R. M., . . . Jiang, Y. H. (2016). Altered mGluR5-Homer scaffolds and corticostriatal connectivity in a Shank3 complete knockout model of autism. *Nat Commun*, *7*, 11459. https://doi.org/10.1038/ncomms11459

Wiltschko, A. B., Tsukahara, T., Zeine, A., Anyoha, R., Gillis, W. F., Markowitz, J. E., Peterson, R. E., Katon, J., Johnson, M. J., & Datta, S. R. (2020). Revealing the structure of pharmacobehavioral space through motion sequencing. *Nat Neurosci*, *23*(11), 1433-1443. https://doi.org/10.1038/s41593-020-00706-3

Xu, Z. C., Wilson, C. J., & Emson, P. C. (1989). Restoration of the corticostriatal projection in rat neostriatal grafts: electron microscopic analysis. *Neuroscience*, *29*(3), 539-550. https://doi.org/10.1016/0306-4522(89)90129-2

Yin, H. H., Knowlton, B. J., & Balleine, B. W. (2005). Blockade of NMDA receptors in the dorsomedial striatum prevents action-outcome learning in instrumental conditioning. *Eur J Neurosci*, *22*(2), 505-512. https://doi.org/10.1111/j.1460-9568.2005.04219.x

Yin, H. H., Knowlton, B. J., & Balleine, B. W. (2006). Inactivation of dorsolateral striatum enhances sensitivity to changes in the action-outcome contingency in instrumental conditioning. *Behav Brain Res*, *166*(2), 189-196. https://doi.org/10.1016/j.bbr.2005.07.012

Yin, H. H., Mulcare, S. P., Hilário, M. R. F., Clouse, E., Holloway, T., Davis, M. I., Hansson, A. C., Lovinger, D. M., & Costa, R. M. (2009). Dynamic reorganization of striatal circuits during the acquisition and consolidation of a skill. *Nature Neuroscience*, *12*(3), 333-341. https://doi.org/10.1038/nn.2261

Zhang, Y. F., Vargas Cifuentes, L., Wright, K. N., Bhattarai, J. P., Mohrhardt, J., Fleck, D., Janke, E., Jiang, C., Cranfill, S. L., Goldstein, N., Schreck, M., Moberly, A. H., Yu, Y., Arenkiel, B. R., Betley, J. N., Luo, W., Stegmaier, J., Wesson, D. W., Spehr, M., . . . Ma, M. (2021). Ventral striatal islands of Calleja neurons control grooming in mice. *Nat Neurosci*, *24*(12), 1699-1710. https://doi.org/10.1038/s41593-021-00952-z