An early, novel arginine methylation of KCa3.1 attenuates subsequent T cell exhaustion.

Authors:

Piyush Sharma[1], Ao Guo[1], Suresh Poudel[1], Emilio Boada-Romero[1], Katherine C. Verbist[1],

Gustavo Palacios[1], Kalyan Immadisetty[2], Mark J. Chen[1,3], Dalia Haydar[4], Ashutosh Mishra[5],

Junmin Peng[5,6], M. Madan Babu[2], Giedre Krenciute[4], Evan S. Glazer[7], Douglas R. Green[1].

[1]Department of Immunology, St. Jude Children's Research Hospital, Memphis, TN 38105, USA

[2]Center of Excellence in Data Driven Discovery, Department of Structural Biology, St. Jude

Children's Research Hospital, Memphis, TN 38105, USA

[3]Department of Laboratory Medicine, Cleveland Clinic, Cleveland, OH 44106, USA

[4]Department of Bone Marrow Transplantation and Cellular Therapy, St. Jude Children's Research

Hospital, Memphis, TN 38105, USA

[5]Center for Proteomics and Metabolomics, St. Jude Children's Research Hospital, Memphis, TN

38105, USA

[6]Department of Structural Biology, Department of Developmental Neurobiology, St. Jude

Children's Research Hospital, Memphis, TN 38105, USA

[7]Department of Surgery, University of Tennessee Health Science Center, 910 Madison Ave., Suite

300, Memphis, TN 38163, USA

Correspondence to piyush.sharma@stjude.org, douglas.green@stjude.org

ABSTRACT

T cell receptor (TCR) engagement initiates the activation process, and this signaling event is

regulated in multifaceted ways. Nutrient availability in the immediate niche is one such mode of

regulation[1-3]. Here, we investigated how the availability of an essential amino acid methionine

(Met) and TCR signaling might interplay in the earliest events of T cell activation to affect

subsequent T cell fate and function. We found that limiting Met during only the initial 30 minutes

of CD8$^+$ T cell activation increased Ca^{2+} influx, Ca^{2+}-mediated NFAT1 (*Nfatc2*) activation,

NFAT1 promoter occupancy, and T cell exhaustion. We identified changes in the protein arginine

methylome during the initial 30 min of TCR engagement and discovered a novel arginine

methylation of a Ca^{2+}-activated potassium transporter, KCa3.1, which regulates Ca^{2+}-mediated

NFAT1 signaling to ensure optimal activation. Ablation of arginine methylation in KCa3.1 led to

increased NFAT1 activation, rendering T cells dysfunctional in murine tumour and infection

models. Furthermore, acute Met supplementation at early stages reduced nuclear NFAT1 in

tumour-infiltrating T cells and augmented their anti-tumour activity. Our findings identify a

metabolic event occurring early after T cell activation that influences the subsequent fate of the

cell.

Methionine limitation promotes T cell exhaustion.

During TCR engagement, nutrient metabolism ensures effective and optimal T cell activation and

function[4]. To determine which amino acids were rapidly consumed upon T cell activation, we

activated OT-I transgenic CD8[+] T cells in complete medium and performed targeted mass

spectrometry. We observed a significant decrease in intracellular Met and other amino acids as

early as 10 min (Fig. 1A). Among the amino acids important for T cells, Met is the sole amino acid

responsible for cellular methylome maintenance via the S-adenosyl Met (SAM) pathway, and

alterations in the SAM pathway adversely affect T cell functions[5,6]. In complete media, no

concomitant changes were observed in SAM or S-adenosyl homocysteine (SAH) (Fig. 1B, upper

panel). T cells consumed their intracellular Met pool only in Met-deprived medium, correlating

with decreases in SAM concentration and corresponding increases in SAH (Fig.1B, bottom panel).

These observations indicate that extracellular Met availability is required to maintain the SAM

cycle during T cell activation.

Mice infected with chronic Lymphocytic choriomeningitis virus (LCMV) have ~65% less Met

than in serum as early as day 2 post infection[7] (Extended Data Fig. 1A), suggesting that T cells

can experience low Met conditions during activation. To determine whether Met limitation during

TCR engagement affects T cell proliferation, we lowered the available amount of Met to 0.03 mM

in the culture medium. Cell-trace violet (CTV) labelled OT-I CD8[+] T cells were activated in media

with control (0.1 mM) or 0.03 mM Met for 30 min to 6 hr, after which Met was restored to 0.1

mM, and we observed no difference in the proliferation of T cells activated in reduced Met for

different times (Fig. 1C). When then assessed their activity *in vivo* upon transferring these

activated T cells into mice bearing B16/MC38 tumours expressing ovalbumin (B16-OVA/ MC38-

OVA) 24 h after activation (Fig. 1D), we observed defective tumour control in T cells activated in

0.03 mM Met compared to 0.1 mM Met, across all time points, with the earliest being 30 min

(Rag1[-/-] mice: Fig. 1E-G-B16-OVA; B6 WT mice: Extended Data Fig. 1B, C-MC38-OVA). To

further assess this effect, we generated LCMV GP33-specific memory T cells and activated them

for 24 hrs with GP33 peptide in either 0.1 mM Met or 0.03 mM Met for 30 min, after which Met

was restored to 0.1 mM and then transferred them into B16-GP33 (B16 expressing GP33 peptide)

tumour-bearing Rag1[-/-] mice (Extended Data Fig. 1D). We observed defective tumour control

leading to poor survival in mice receiving T cells activated in 0.03 mM Met compared to 0.1 mM

Met (Fig. 1H and I).

To examine the effects of activation in reduced Met on the T cell epigenome, we performed

ATAC-Seq at 24 hrs post activation (Extended Data Fig. 1E) and observed increased chromatin

accessibility in T cells activated in 0.03 mM Met during the initial 30 min (Extended Data Fig. 2A

and B). Overrepresentation analysis showed enrichment of gene sets associated with an

exhaustion-like phenotype (Extended Data Fig. 2C). Further HOMER motif analysis showed that

transcription factor (TF) motifs known to be associated with exhausted T cells[8,9] were highly

accessible in 0.03 mM-activated T cells (Extended Data Fig. 2D).

We activated OT-I CD8[+] T cells T cells in either control or 0.03 mM Met for the initial 30 min

followed by 24 hours culture in 0.1 mM Met and injected them into B16-OVA bearing Rag1[-/-]

mice. TILs were collected at day 9 post T cell injection. Again, animals receiving T cells activated

initially in 0.03 mM Met showed defective tumour control (Extended Data Fig. 2E and F). The

CD8[+] TILs displayed reduced CD62L[+], CD62L[high] CD44[high] central memory cells (Tcm) with

increased TOX expression and reduced IFNγ production compared to TILs from T cells activated

in Met replete media (Extended Data Fig. 2G).

T cell dysfunction is known to contribute to poor prognoses in cancer[10]. This dysfunctional state

is at least in part driven by the transcription factor (TF) TOX[11] and is accompanied by markers

such as increased surface expression of PD1 and Tim-3 and reduced effector cytokine

production[12,13]. To further examine the effect of activation in limiting Met, we activated CD45.1[+]

or CD45.2[+] OT-I CD8[+] T cells in 0.1 (control) or 0.03 mM Met for 30 min, restored Met to 0.1

mM for 24 hrs, and mixed them at a ratio of 1:1 before transferring them into B16-OVA tumour-

bearing $Rag1^{-/-}$ mice (Fig. 2A). Even though CD8$^+$ TIL initially activated in 0.03 mM outnumbered

the 0.1 mM-activated CD8$^+$ TIL at day 12 (Fig. 2B and C), these cells exhibited significant

increases in exhaustion-associated expression of PD1 and Tim-3, reduced effector cytokines (Fig.

2D,E and Extended Data Fig. 2H), increased expression of exhaustion associated TF, TOX (Fig.

2F and G), and reduced expression of the stemness-associated TF, TCF-1 (Extended Data Fig. 2I).

Altogether, these results suggest that reduced Met during early TCR signaling promotes

acquisition of epigenomic changes associated with exhaustion and drives T cells towards a

dysfunctional, exhausted state.

Exhausted T cells have been further defined as precursors of exhaustion (TEXprog) and

TEXprogenitor-like cells, both of which produce a replicative burst and effector function upon

checkpoint blockade [14], and terminally exhausted cells (TEXterm), which fail to mount an effective

effector response [15,16]. Although TOX is associated with maintenance of exhausted T cells, it is

also expressed in TEXprog [17]. We therefore further dissected the TOX$^+$ CD8$^+$ T cells from the

experiment in Fig. 2F (Fig. 2H) and found decreased TEXprog (TCF1$^+$Tim-3$^-$) and increased

TEXterm (TCF1$^-$Tim-3$^+$) among T cells initially activated in 0.03 mM Met, suggesting a decreased

pool of stem-like TEXprog cells (Fig. 2I). Furthermore, evaluation of total CD8$^+$ TILs from mice

receiving T cells initially activated in 0.03 mM Met for 30 minutes revealed a lower frequency of

TCF1$^+$Tim-3$^-$ TEXprog cells and CX3CR1$^+$Ly108$^+$ TEXprogenitor-like cells (Extended Data Fig.

2J) and a corresponding increase in the frequency of TEXterm cells defined as TCF1$^-$Tim-3$^+$

(Extended Data Fig. 2K). We also observed a decrease in memory-associated markers such as

CD62L$^+$, CD62L$^+$CD44$^+$ (Extended Data Fig. 2L) and CD127$^+$CD27$^+$ (Extended Data Fig. 2M

and N) and an increase in KLRG1$^+$CD127$^-$ terminal effector-like T cells (Extended Data Fig. 2O

and P) when initially activated in 0.03 mM Met.

To examine the transcriptomic state, we performed global RNA-seq of day 9 TIL from T cells

originally activated for 30 min in 0.1 mM or 0.03 mM Met and then cultured in 0.1 mM Met for

24hrs prior to transfer into B16-OVA tumour-bearing $Rag1$-/- mice. We found several

transcriptomic changes, the majority of which represented increased expression of differentially

enriched (DE) genes in T cells initially activated in 0.03 mM Met (Extended Data Fig. 3A and B).

Over-representation analysis of the DE genes in T cells activated in 0.1 mM Met revealed
enrichment of GOBP pathways which are associated with T cell effector functions along with
increases in transcription factors associated with a stem-like T cell state and memory generation
[18-21] (Extended Data Fig. 3C). Cells initially activated in 0.03 mM Met showed enrichment of genes
which are associated with terminal differentiation or dysfunction of T cells and transcription
factors associated with hyperactivation and development and maintenance of T cell exhaustion [8,22-
24] (Extended Data Fig. 3D). GSEA analysis also showed an enriched exhaustion-associated gene
set [25,26] in CD8[+] TIL initially activated in 0.03 mM Met (Fig. 2J) and enrichment of hallmark gene
sets associated with T cell effector function in the control (0.1 mM Met) T cells (Extended Data
Fig. 3E). These results suggest that Met availability during early TCR engagement influences T
cell differentiation, with initial, reduced Met availability promoting hyperactivation and driving T
cells towards exhaustion.
137
Extracellular Met metabolism regulates Ca^{2+} flux and NFAT1 activation.
Among the earliest events in T cell activation is the influx of Ca^{2+} [27,28], which is accompanied by
the activation of calcineurin, which dephosphorylates Nuclear Factor of Activated T cells-1
(NFAT1), resulting in its nuclear translocation and activation[29,30]. Because low Met levels for only
the first 30 min of activation influenced the subsequent T cell phenotype, we investigated changes
in Ca^{2+} flux using Fluo-8. We observed increased Ca^{2+} influx upon stimulation of T cells with anti-
CD3/CD28 antibody in 0.03 mM compared to 0.1 mM Met (Fig. 3A), which was reduced by the
CRAC channel inhibitor YM-58483 (Extended Data Fig. 4A). NFAT1 cooperates with the AP1
complex to ensure optimal T cell activation and differentiation, but high levels of NFAT1 activity
have been associated with T cell dysfunction and exhaustion [29,31]. Confocal imaging revealed an
increase in activated NFAT1 (nuclear localization of NFAT1) in T cells activated for 30 min with
anti-CD3/28 beads in 0.03 mM Met compared to 0.1 mM Met (Fig. 3B, Extended Data Fig. 4B),
quantified as either a ratio of nuclear NFAT1 to total cell NFAT (Fig. 3C and Extended Data Fig.
4C(0.0 mM Met)) or as nuclear NFAT1 fluorescence intensity (Extended Data Fig. 4D), which
were abolished upon treatment with either the calcineurin inhibitor Cyclosporin A (CsA) (Fig. 3B
and C) or CRAC channel inhibitor YM-58483 (Extended Data Fig. 4E). Examination of NFAT2
at 30 min of activation revealed no difference in its activation in T cells activated in either 0.1 mM
or 0.03 mM Met (Extended Data Fig. 4F). Next, we activated CD8[+] T cells in complete medium

for 24hrs and rested them for 48hrs before secondary activation in 0.03 mM or 0.1 mM Met and

observed increased nuclear NFAT1 levels in 0.03 mM Met 30 min after activation (Extended Data

Fig. 4G), suggesting that the effect of limiting Met on NFAT1 activation is independent of previous

activation events.

NFAT1 is known to regulate several exhaustion-associated genes such as *Lag3, Pdcd1, Havcr2,*

Ctla4, Tnfrsf9 [31]. We performed NFAT1 CUT&RUN on T cells activated in 0.1 mM or 0.03 mM

Met for 30 min, followed by 0.1 mM Met for 24hrs and observed increased NFAT1 binding in the

low Met group (Fig. 3D). We observed that Met limitation for the initial 30 min resulted in

increased NFAT1 binding to *Pdcd1, Havcr2, Lag3, and Tnfrsf9* (Fig. 3E) as well as *Ctla4 and Tox*

gene loci (Fig. 3F). This further correlated with increased transcription of most of these genes at

24 hrs (Extended Data Fig. 4H). Altogether, our results suggest that reduced Met availability

during the initial stages of TCR activation directs a trajectory of T cells towards hyperactivation

and eventual exhaustion, and the maintenance of extracellular Met levels in the face of rapid Met

consumption upon TCR engagement dampens Ca^{2+}-mediated NFAT1 activity and T cell fate.

KCa3.1 methylation regulates NFAT1.

Methionine metabolism is the sole pathway for epigenetic and proteomic methylome maintenance,

which is known to affect T cell function [32]. When converted to SAH, SAM acts as the methyl

donor; thus, the ratio of SAM to SAH is known as the cellular methylation potential [33], an index

of a cell's capacity to maintain its methylome. We calculated the SAM/SAH ratio over time after

TCR engagement in the absence of extracellular Met and found this ratio to be significantly

reduced as early as 10 min post-activation (Extended Data Fig. 5A), suggesting alterations in the

methylome. Met metabolism is essential for DNA methylation [34]; however, we found no

significant differences in global 5-methylcytosine levels at either 30 min or 24 hrs in T cells

activated as above (Extended Data Fig. 5B). We similarly assessed H3me3K27 and H3me2K4 by

CUT&RUN at 24 hrs post activation, but again, no global changes were observed (Extended Data

Fig. 5C and D). Thus, Met restriction for only the first 30 min of T cell activation does not appear

to directly affect global DNA or histone methylation.

Therefore, we postulated that Met restriction influences early methylation events independently of
direct effects on DNA and histone methylation. Post-translational protein arginine methylation
mediated by protein arginine methyltransferases regulates protein function [35,36]. We found that
arginine methylation in T cells activated in 0.0 mM or 0.03 mM Met was significantly reduced 30
min post-activation when compared to T cells activated in 0.1 mM Met (Fig. 4A and B, Extended
Data Fig. 5E). To identify specific proteins differentially methylated upon TCR engagement in
limited Met, CD8[+] T cells activated by anti-CD3 antibody with IgG cross-linking for 30 min were
processed for methyl-arginine immunoprecipitation and quantification using multiplexed tandem
mass tag-mass spectrometry (TMT-MS). We identified common proteins which were
differentially methylated in 0.0 mM and 0.03 mM Met activation conditions compared to 0.1 mM
Met (Fig. 4C) and found the top candidate, KCa3.1 to be di-methylated at Arginine-350 (R350) in
0.1 mM Met, which was reduced in 0.0 mM and 0.03 mM Met (Fig. 4D, Supplementary Table. 1).
Furthermore, protein-BLAST analysis showed that KCa3.1 R350 is conserved across species,
including humans, which suggested that R350 may be important for KCa3.1 regulation (Extended
Data Fig. 5F).
201
KCa3.1 is a calcium-activated potassium transporter encoded by *Kcnn4* and is activated by
calmodulin binding upon TCR-mediated Ca^{2+} flux[37]. During T cell activation, the plasma
membrane is hyperpolarized to ensure sufficient Ca^{2+} flux for signaling[38], and this is maintained
by K^+ efflux through potassium transporters such as KCa3.1[39,40]. Based on our earlier observation
of increased Ca^{2+} signaling under low Met conditions, we hypothesized that R350 methylation of
KCa3.1 regulates its activity, and loss of this methyl group may result in hyperactivation, leading
to increased Ca^{2+} influx and NFAT1 activation. To understand how arginine methylation may
mechanistically modulate KCa3.1 activity, we performed MD simulations with the solved crystal
structure of human KCa3.1(hKCa3.1) bound to calmodulin (CaM) at R352 (corresponding to R350
in murine KCa3.1) (Extended Data Fig. 5G). The calmodulin-binding pocket encompasses D78-
T79-D80-S81-E82-E83-E84, which is predominantly negatively charged/polar, where E84 forms
a salt bridge with R352 of hKCa3.1. Our results showed that both symmetrical di-methylation
(SDMA) and asymmetrical di-methylation (ADMA) of R352 introduced a hydrophobic methyl
group into this negatively charged/polar CaM binding pocket, thereby weakening the hKCa3.1 and
CaM interaction (Fig. 4E). Furthermore, MD simulations showed that the salt bridge interaction

of R352 with E84 of the CaM binding pocket diminished over time, especially in the case of

SDMA where both guanidino nitrogens are methylated, compared to ADMA in which only one

nitrogen is methylated (Fig. 4F, Extended Data Fig. 5H), suggesting that disruption of arginine

methylation promoted stabilization of calmodulin binding, leading to increased activation of the

channel and Ca^{2+} flux.

To validate our *in-silico* simulation results, we used the chemical inhibitor TRAM-34 to reduce

KCa3.1 activity [41,42] during T cell activation and observed reduced Ca^{2+} flux upon TRAM-34

treatment only in 0.03 mM Met and not in 0.1 mM Met (Fig. 4G) at the concentration of TRAM-

34 we employed. Measurement of nuclear NFAT1 levels revealed that the treatment normalized

nuclear NFAT1 levels in T cells activated in 0.03 mM Met in a dose-dependent manner and had

no effect on T cells activated in 0.1 mM Met (Extended Data Fig. 5I). We then activated OT-I T

cells in 0.1 mM or 0.03 mM Met with either DMSO or TRAM-34 for 30 min, washed and then

cultured the cells for an additional 24 hrs in 0.1 mM Met before transferring into B16-OVA

tumour-bearing *Rag1*[-/-] mice (Extended Data Fig. 5J). Chemical inhibition of KCa3.1 for 30 min

significantly improved anti-tumour activity in T cells activated for 30 min in 0.03 mM Met (Fig.

4H), suggesting that increased KCa3.1 activity was responsible for the eventual decreased T cell

function under this activation condition.

Methylation-deficient KCa3.1 increases Ca^{2+}-mediated NFAT1 activation.

To specifically interrogate the role of R350 methylation of KCa3.1 in T cell activation, we cloned

KCa3.1 (Kca3.1[WT]) and a mutant with an alanine substitution (KCa.3.1[R350A]) into retroviral vectors

and expressed them in primary murine CD8[+] T cells in which endogenous KCa3.1 was deleted

using CRISPR/Cas9 (Extended Data Fig. 6A, B and C), and expression of the two was similar

(Extended Data Fig. 6D). Fluo-8 Ca^{2+} flux analysis showed significantly increased flux upon

activation in KCa3.1[R350A] T cells (Fig. 5A). Due to the expression of GFP by transduced T cells,

which might interfere with the Fluo-8 signal, we used another Ca^{2+}-sensitive dye, ICR-1AM, and

observed similar results (Extended Data Fig. 6E). Increased Ca^{2+} flux correlated with an increase

in nuclear NFAT1 30 min post activation (Fig. 5B, C and Extended Data Fig. 6F). This heightened

nuclear NFAT1 in cells expressing KCa3.1[R350A] was visible under resting conditions and increased

upon activation in a time-dependent manner (Extended Data Fig. 6G).

To investigate the epigenomic state of WT or mutant KCa3.1 T cells, we analyzed global

chromatin accessibility via ATAC-Seq after 24 hrs of *in vitro* activation. We found that the

activated KCa3.1[R350A] T cells showed distinct chromatin accessibility which was increased

compared to KCa3.1[WT] T cells (Fig. 5D, Extended Data Fig. 7A). Furthermore, GSEA analysis

revealed enrichment of GOBP terms associated with negative regulators of T cell-mediated

immunity and gene sets associated with T cell exhaustion [25,26] in T cells expressing mutant

KCa3.1[R350A] (Fig. 5E and F), suggesting that ablation of R350 methylation drives T cells towards

exhaustion early after activation. Upon comparing these ATAC-Seq results with ATAC-Seq of T

cells activated in 0.03 mM vs. 0.1 mM Met (Extended Data Fig. 2B), we found several common

differentially accessible regions (DAR) (Extended Data Fig. 7B) with enrichment of gene sets

associated with T cell exhaustion (Extended Data Fig. 7C). HOMER motif analysis of WT or

mutant T cells and common DARs was similarly enriched (Extended Data Fig. 2D, 7D and E).

Despite these results, KCa.3.1[R350A] did not fully phenocopy the chromatin effects of activation in

0.03 mM Met, suggesting that other differentially methylated proteins in addition to KCa.3.1 may

also impact chromatin accessibility following activation in the latter condition.

To investigate whether the observed exhaustion-specific chromatin state led to defective

functionality of mutant KCa3.1 T cells, we transferred T cells expressing KCa.3.1[WT] or

KCa.3.1[R350A] into B16-OVA tumour-bearing Rag1[-/-] mice. We found that T cells expressing

KCa.3.1[R350A] displayed impaired tumour control and survival (Fig. 5G, H and I), suggesting that

the absence of KCa3.1 methylation drives T cells towards an exhaustion trajectory, leading to

defective effector function and tumour control.

Ablation of KCa3.1 methylation promotes T cell exhaustion.

To determine whether ablation of R350 methylation drives T cell exhaustion in tumours, we

transferred T cells expressing KCa.3.1[WT] or KCa.3.1[R350A] into B16-OVA tumour-bearing Rag1[-/-]

mice. Tumours were isolated and measured at day 12 post-T cell injection, and consistently, much

larger tumours were extracted from animals treated with T cells expressing KCa.3.1[R350A]

(Extended Data Fig. 8A). KCa.3.1[R350A]-expressing TIL showed high surface expression of the

exhaustion markers PD1, Lag3, and Tim-3 (Extended Data Fig. 8B) that correlated with increased

TOX expression (Extended Data Fig. 8C). Among these cells we also found fewer TEXprog

(CD69^{lo}Ly108^{hi})[43] and IFNγ- and TNFα-producing T cells (Extended Data Fig. 8D and E). Next,

we performed a similar experiment in a competitive setting, wherein CD45.1$^+$ KCa.3.1^{WT} and

CD45.1$^+$CD45.2$^+$ Kca.3.1^{R350A}-expressing OT-I CD8$^+$ T cells were mixed at a 1:1 ratio and

transferred into B16-OVA tumour-bearing *Rag1$^{-/-}$* mice (Fig. 6A). TIL analysis at D12 showed

increased PD1$^+$Tim-3$^+$, Eomes$^+$PD1$^+$ (Fig. 6B, Extended Data Fig. 8F) and TOX expression in

Kca.3.1^{R350A} -expressing T cells, which correlated with reduced expression of the effector

cytokines IFNγ and TNFα (Fig. 6C and D). Further analysis revealed a decreased frequency of

TCF1$^+$Tim-3$^-$ TEXprog and increased TCF1$^-$Tim-3$^+$ TEXterm populations (Extended Data Fig. 8G

and Fig. 6E). We also observed a decrease in CD62L$^+$ and CD127$^+$CD27$^+$ memory T cells (Fig.

6F) with an increase in the KLRG1$^+$CD127$^-$ terminal effector-like phenotype in KCa.3.1^{R350A}-

expressing T cells, as compared to T cells expressing WT KCa3.1 (Extended Data Fig. 8H and I).

T cell exhaustion is a prominent feature of chronic viral infection [44]. We utilized the LCMV

infection model in WT (CD45.2, Thy 1.2) mice into which CD45.1$^+$CD45.2$^+$ KCa.3.1^{WT} and

CD45.2$^+$ KCa.3.1^{R350A}-expressing Thy1.1 (CD90.1$^+$) P14$^+$ T cells were injected at a 1:1 ratio into

mice infected with either an acute Armstrong or a chronic Clone-13 strain. Spleens were analyzed

on day 9 post-infection (Fig. 6G). The KCa.3.1^{R350A} population frequency was significantly

reduced in Clone-13 infection, but no significant change was observed in acute infection (Extended

Data Fig. 8J). KCa.3.1^{R350A}-expressing T cells presented with increased TOX, PD1 and Tim-3

expression in both acute and chronic infections (Fig. 6H-K, Extended Data Fig. 8K). We also

found that KCa.3.1^{R350A}-expressing T cells harbored reduced frequencies of CX3CR1$^+$PD1$^+$

TEXprog cells (Fig. 6L) in both acute and chronic infection and increased Tim-3$^+$Cx3CR1$^+$PD1$^+$

TEXeff cells (T exhausted effector-like cells) and TCF1$^-$Tim-3$^+$ TEXterm cells in chronic infection

(Extended Data Fig. 8L and Fig. 6M). KCa.3.1^{R350A}-expressing P14$^+$ T cells displayed significantly

increased KLRG1$^+$CD127$^-$ terminal effector-like cells and a concurrent decrease in KLRG1$^-$

CD127$^+$ memory precursor cells in chronic infection but not in acute infection (Extended Data Fig.

8M-O). Thus, ablation of R350 methylation drives T cells towards an exhaustion trajectory even

at the early stages of infection. Together, these experiments support the idea that R350 methylation

maintains optimal KCa3.1 function, ensuring optimal T cell activation and effector function in

both tumours and viral infections.

Acute Met supplementation promotes anti-tumour T cell function.

CD8[+] T cells are activated in both secondary lymphoid tissues and within tumours [45,46]. While T cells are known to be primarily activated in draining lymph nodes (dLNs), a recent study showed that activation of tumour infiltrating T cells in tumour microenvironment is necessary to acquire an effector-like phenotype[47]. Our results in Fig. 1H and Extended Data Fig. 4G support the idea that Met is required for optimal TCR activation, and this requirement is independent of the prior activation status of the T cell. Thus, we hypothesized that upon tumour infiltration, CD8[+] T cells experience antigen presentation in a Met-deficient microenvironment, leading to altered effector function. To evaluate Met levels in TILs, we performed mass spectrometric analysis of CD8[+] TIL and dLN CD8[+] T cells from B16 subcutaneous tumour-bearing mice at day 9. We observed a ~50% reduction in intracellular Met in CD8[+] TIL compared to that in dLN CD8[+] T cells (Extended Data Fig. 9A, Supplementary table 2). We also observed a similar decrease in intracellular Met in CD8[+] TIL isolated from primary human colorectal carcinomas compared with corresponding CD8[+] T cells from PBMC (Extended Data Fig. 9B). To determine whether CD8[+] TIL with reduced intracellular Met showed increased nuclear NFAT1, we sorted activated CD44[+] CD8[+] T cells from B16 tumours and respective dLN at day 9 post implantation and imaged for NFAT1. We observed increased levels of nuclear NFAT1 in CD44[+]CD8[+] TIL compared to counterparts from dLN (Fig. 7A), suggesting that reduced Met availability in the TME may have led to increased NFAT1 activation in those T cells.

Continuous Met supplementation in tumours has been shown to promote anti-tumour T cell activity, associated with global epigenetic maintenance [32]. Our findings, however, highlight an early role for Met in T cell fate, and we hypothesized that Met supplementation up to that of serum levels at the early stages of tumour development would increase T cell anti-tumour activity. We therefore injected 50 μl of 61μM Met peri-tumourally once per day for 5 days when tumour volumes were 70-100 mm^3 (Extended Data Fig. 9C) and found that treatment decreased nuclear NFAT1 levels in CD44[+] CD8[+] TIL as compared to TIL from HBSS-injected tumours (Fig. 7B). Indeed, nuclear NFAT1 in TIL was similar to or lower than that in CD8[+] T cells from dLN (Extended data Fig. 9D). Furthermore, acute Met supplementation delayed tumour growth in multiple murine tumour models, thereby improving overall survival (Fig. 7C, D and Extended Data Fig. 9E-G). Met supplementation resulted in decreased PD1 and Tim-3 surface expression in

CD8$^+$ TIL, along with increased CD62L^{hi}CD44^{hi} Tcm and reduced CD62L^{lo}CD44^{hi} Tem

(Extended Data Fig. 9H and I) compared to TIL in HBSS-treated tumours. In contrast to

immunocompetent WT mice, Met injected peri-tumourally in tumour-bearing NSG mice resulted

in no difference in tumour growth (Extended Data Fig. 9J), implying a lymphocyte-mediated

effect. Depletion with an anti-CD8 antibody (Extended Data Fig. 9K) in B16-OVA tumour-bearing

mice abrogated the effect of Met supplementation on tumour control (Fig. 7E and F), supporting

the conclusion that the observed Met supplementation-mediated tumour control is dependent on

CD8$^+$ T cells. We then employed a contralateral tumour model with B16-OVA implanted in one

shoulder and in the opposite abdominal flank and injected Met peri-tumourally to the latter tumour.

Met supplementation of the abdominal flank tumours impeded contralateral tumour growth

(Extended Data Fig. 10A and B), further suggesting a T cell response. These results are consistent

with the idea that acute Met supplementation promoted tumour control by CD8$^+$ T cells. We also

found that an 0.5% increase in dietary Met supplementation in B16 tumour-bearing mice showed

better tumour growth control (Extended Data Fig. 10C), which is concordant with a recent study

showing that Met supplementation in humans increased the effector function of CD8$^+$ T cells [32].

Met supplementation, therefore, enhances the anti-tumour function of CD8$^+$ T cells.

Met supplementation enhances CAR and ICB treatment.

CAR-T cell treatment has been clinically successful in the treatment of hematological

malignancies, but less so in solid tumours [48,49]. In a solid tumour model in *Rag1$^{-/-}$* mice, we

supplemented Met in the presence of transferred murine B7-H3 CAR-T cells into animals bearing

a B7-H3$^+$ F420 tumour[50] (Extended Data Fig. 10D). Again, Met supplementation improved tumour

control and animal survival in this model (Fig. 7G and H).

Immune checkpoint blockade (ICB) has been widely used either alone or in combination with

cancer treatments[51]. To query whether Met supplementation might cooperate with ICB, we

administered either IgG isotype or anti-PD1 antibody along with HBSS or Met supplementation

in subcutaneous MC38 tumour-bearing mice (Extended Data Fig. 10E) and found that

peritumoural Met supplementation with anti-PD1 treatment resulted in significantly reduced

tumour growth and increased survival of these animals compared to controls (Extended Data Fig.

10F and G). As local Met supplementation may not be feasible in the clinical setting, we performed

an ICB experiment with dietary supplementation of Met (Extended Data Fig. 10H). We observed

that a diet containing 1.5% Met (0.5% increase in dietary Met) enhanced the efficacy of anti-PD-

1, leading to significant control of tumour growth and survival of the mice, compared to that of

anti-PD1 with control diet (Fig. 7I-K, Supplementary Table. 3). These results suggest the potential

application of Met as a combinatorial treatment in the existing immunotherapies.

Discussion

T cell exhaustion is generally understood to be a consequence of persistent antigen stimulation

[12,44]; however, situations exist in which antigen persistence does not result in exhaustion, such as

in autoimmune T cells in Type I diabetes [21]. It is therefore possible that in addition to chronic

antigen exposure, nutrient limitation contributes to exhaustion [52,53]. A recent study suggested that

T cells can follow an exhaustion trajectory within hours of antigen recognition [54]. Here, we propose

that the state of T cell exhaustion is the result of a multifactorial pathway that involves the

assimilation of different input signals, including nutrients, and is engaged as early as the initial

recognition of cognate antigen.

Studies have shown that strong TCR signals result in hyperactivation and activation-induced cell

death, whereas weak TCR signals result in dampened responses [55]. NFAT family members are key

regulators of T cell activation and interact with key partners to induce activation-associated genes

including effector- and exhaustion-associated genes [28]. Continuous or partner-less NFAT

activation is associated with hyperactivation, anergy and exhaustion [29,31,56]; therefore, NFAT

activation must be tightly controlled for optimal T cell function, but little is known about how

niche-specific nutrients regulate NFAT. We found that during rapid tumour proliferation in the

early growth phase, $CD8^+$ TIL display increased NFAT1 activation, which is regulated

metabolically. Our data provide a novel mechanism of Met metabolism in the regulation of NFAT1

activation via post-translational modification of KCa3.1, without which $CD8^+$ T cells progress

towards exhaustion (Extended Data Fig. 11). We found that arginine methylation in KCa3.1 limits

Ca^{2+}-dependent NFAT1 signaling, ensuring optimal activation in tumours and infections. Met

supplementation to the TME preserves anti-tumour immunity [32] and limits exhaustion (this study),

a treatment that can be harnessed to improve patient prognosis in combination with ICB therapies.

Our results emphasize the importance of non-histone protein methylation in the functional

maintenance of T cells in diseases and describe a link between TCR signaling, metabolic

regulation, and transcription factor activation. These results suggest a change in the current dogma

of T cell exhaustion from a state driven only by persistent antigen exposure—acquired over a long

period of time—to a dynamic state controlled by multiple pathways over the course of T cell

activation and differentiation. Our study supports the idea that metabolism-dependent, rapid

proteomic modifications that occur upon TCR engagement influence subsequent T cell fate and

provides mechanistic insight into how niche-specific nutrient deprivation renders T cells

dysfunctional

Acknowledgements

We thank P. Fitzgerald, S. Sirasanagandla, M. Yang and L. Harris, for technical and administrative

assistance; L. Long and H. Chi for providing retroviral vectors and Cas9-OTI mice; C. Guy for

microscope imaging assistance; M. Morrison for organizing patient samples; the staff at the St.

Jude Immunology Flow Cytometry Core Facility for cell sorting. Statistical calculations were

performed using GraphPad Prism. We thank Dr. S. Pounds for the additional assistance with the

statistical analysis of the data in Fig. 4A. Illustrations were created using BioRender software. This

work was supported by National Institutes of Health grants AI123322 and CA231620 to D.R.G.,

American Lebanese Syrian Associated Charities (ALSAC) to G.K., and National Cancer Institute

grant K99CA256262 to D.H.

Author Contributions

P.S. conceived the project, designed, performed most experiments, interpreted results, and wrote

the manuscript. A.G. contributed to CUT&RUN, ATAC-Seq, and RNA-Seq experiments. K.V.

and E.B.R. helped in the development of the flow cytometry panel, experiments, and review of the

manuscript. G.P. contributed to and processed the LC-MS/MS experiments. D.H. and G.K.

assisted with CAR-T cell experiments. K.I. and M.B. performed MD simulations and analyses.

S.P. and M.J.C. analyzed the CUT&RUN, ATAC-seq, and RNA-seq data. A. M. and J.P.

performed and analyzed methyl-arginine immunoprecipitation and TMT-MS experiments. E.S.G

provided patient tumour samples. D.R.G. supervised the experimental designs, interpreted the

results, and wrote the manuscript.

References

1   Lim, A. R., Rathmell, W. K. & Rathmell, J. C. The tumor microenvironment as a
metabolic barrier to effector T cells and immunotherapy. *Elife* **9** (2020).
https://doi.org:10.7554/eLife.55185
2   Kedia-Mehta, N. & Finlay, D. K. Competition for nutrients and its role in controlling
immune responses. *Nat Commun* **10**, 2123 (2019).
https://doi.org:10.1038/s41467-019-10015-4
3   Wei, J., Raynor, J., Nguyen, T. L. & Chi, H. Nutrient and Metabolic Sensing in T
Cell Responses. *Front Immunol* **8**, 247 (2017).
https://doi.org:10.3389/fimmu.2017.00247
4   Cohen, S., Danzaki, K. & MacIver, N. J. Nutritional effects on T-cell
immunometabolism. *Eur J Immunol* **47**, 225-235 (2017).
https://doi.org:10.1002/eji.201646423
5   Klein Geltink, R. I. & Pearce, E. L. The importance of methionine metabolism. *Elife*
8 (2019). https://doi.org:10.7554/eLife.47221
6   Yang, X. *et al.* MAT2A-Mediated S-Adenosylmethionine Level in CD4(+) T Cells
Regulates HIV-1 Latent Infection. *Front Immunol* **12**, 745784 (2021).
https://doi.org:10.3389/fimmu.2021.745784
7   Lercher, A. *et al.* Type I Interferon Signaling Disrupts the Hepatic Urea Cycle and
Alters Systemic Metabolism to Suppress T Cell Function. *Immunity* **51**, 1074-1087
e1079 (2019). https://doi.org:10.1016/j.immuni.2019.10.014
8   Beltra, J. C. *et al.* Stat5 opposes the transcription factor Tox and rewires exhausted
CD8(+) T cells toward durable effector-like states during chronic antigen exposure.
Immunity **56**, 2699-2718 e2611 (2023).
https://doi.org:10.1016/j.immuni.2023.11.005
9   Belk, J. A. *et al.* Genome-wide CRISPR screens of T cell exhaustion identify
chromatin remodeling factors that limit T cell persistence. *Cancer Cell* **40**, 768-786
e767 (2022). https://doi.org:10.1016/j.ccell.2022.06.001
10 Thommen, D. S. & Schumacher, T. N. T Cell Dysfunction in Cancer. *Cancer Cell*
33, 547-562 (2018). https://doi.org:10.1016/j.ccell.2018.03.012
11 Khan, O. *et al.* TOX transcriptionally and epigenetically programs CD8(+) T cell
exhaustion. *Nature* **571**, 211-218 (2019). https://doi.org:10.1038/s41586-019-
1325-x
12 Wherry, E. J. & Kurachi, M. Molecular and cellular insights into T cell exhaustion.
Nat Rev Immunol **15**, 486-499 (2015). https://doi.org:10.1038/nri3862
13 Zhang, Z. *et al.* Pan-cancer landscape of T-cell exhaustion heterogeneity within
the tumor microenvironment revealed a progressive roadmap of hierarchical
dysfunction associated with prognosis and therapeutic efficacy. *EBioMedicine* **83**,
104207 (2022). https://doi.org:10.1016/j.ebiom.2022.104207
14 Kallies, A., Zehn, D. & Utzschneider, D. T. Precursor exhausted T cells: key to
successful immunotherapy? *Nat Rev Immunol* **20**, 128-136 (2020).
https://doi.org:10.1038/s41577-019-0223-7
15 Hudson, W. H. *et al.* Proliferating Transitory T Cells with an Effector-like
Transcriptional Signature Emerge from PD-1(+) Stem-like CD8(+) T Cells during
Chronic Infection. *Immunity* **51**, 1043-1058 e1044 (2019).
https://doi.org:10.1016/j.immuni.2019.11.002

16 Utzschneider, D. T. *et al.* Early precursor T cells establish and propagate T cell
exhaustion in chronic infection. *Nat Immunol* **21**, 1256-1266 (2020).
https://doi.org:10.1038/s41590-020-0760-z
17 Alfei, F. *et al.* TOX reinforces the phenotype and longevity of exhausted T cells in
chronic viral infection. *Nature* **571**, 265-269 (2019).
https://doi.org:10.1038/s41586-019-1326-9
18 Manolios, N., Papaemmanouil, J. & Adams, D. J. The role of ion channels in T cell
function and disease. *Front Immunol* **14**, 1238171 (2023).
https://doi.org:10.3389/fimmu.2023.1238171
19 Feng, Q. *et al.* Lactate increases stemness of CD8 + T cells to augment anti-tumor
immunity. *Nat Commun* **13**, 4981 (2022). https://doi.org:10.1038/s41467-022-
32521-8
20 Hart, G. T., Hogquist, K. A. & Jameson, S. C. Kruppel-like factors in lymphocyte
biology. *J Immunol* **188**, 521-526 (2012).
https://doi.org:10.4049/jimmunol.1101530
21 Abdelsamed, H. A. *et al.* Beta cell-specific CD8(+) T cells maintain stem cell
memory-associated epigenetic programs during type 1 diabetes. *Nat Immunol* **21**,
578-587 (2020). https://doi.org:10.1038/s41590-020-0633-5
22 Sun, Q. *et al.* STAT3 regulates CD8+ T cell differentiation and functions in cancer
and acute infection. *J Exp Med* **220** (2023). https://doi.org:10.1084/jem.20220686
23 Zheng, C. *et al.* Transcriptomic profiles of neoantigen-reactive T cells in human
gastrointestinal cancers. *Cancer Cell* **40**, 410-423 e417 (2022).
https://doi.org:10.1016/j.ccell.2022.03.005
24 Belk, J. A., Daniel, B. & Satpathy, A. T. Epigenetic regulation of T cell exhaustion.
Nat Immunol **23**, 848-860 (2022). https://doi.org:10.1038/s41590-022-01224-z
25 Wherry, E. J. *et al.* Molecular signature of CD8+ T cell exhaustion during chronic
viral infection. *Immunity* **27**, 670-684 (2007).
https://doi.org:10.1016/j.immuni.2007.09.006
26 Doering, T. A. *et al.* Network analysis reveals centrally connected genes and
pathways involved in CD8+ T cell exhaustion versus memory. *Immunity* **37**, 1130-
1144 (2012). https://doi.org:10.1016/j.immuni.2012.08.021
27 Trebak, M. & Kinet, J. P. Calcium signalling in T cells. *Nat Rev Immunol* **19**, 154-
169 (2019). https://doi.org:10.1038/s41577-018-0110-7
28 Quintana, A., Griesemer, D., Schwarz, E. C. & Hoth, M. Calcium-dependent
activation of T-lymphocytes. *Pflugers Arch* **450**, 1-12 (2005).
https://doi.org:10.1007/s00424-004-1364-4
29 Hogan, P. G. Calcium-NFAT transcriptional signalling in T cell activation and T cell
exhaustion. *Cell Calcium* **63**, 66-69 (2017).
https://doi.org:10.1016/j.ceca.2017.01.014
30 Hogan, P. G., Chen, L., Nardone, J. & Rao, A. Transcriptional regulation by
calcium, calcineurin, and NFAT. *Genes Dev* **17**, 2205-2232 (2003).
https://doi.org:10.1101/gad.1102703
31 Martinez, G. J. *et al.* The transcription factor NFAT promotes exhaustion of
activated CD8(+) T cells. *Immunity* **42**, 265-278 (2015).
https://doi.org:10.1016/j.immuni.2015.01.006

32   Bian, Y. *et al.* Cancer SLC43A2 alters T cell methionine metabolism and histone
methylation. *Nature* **585**, 277-282 (2020). https://doi.org/10.1038/s41586-020-
2682-1

33   Caudill, M. A. *et al.* Intracellular S-adenosylhomocysteine concentrations predict
global DNA hypomethylation in tissues of methyl-deficient cystathionine beta-
synthase heterozygous mice. *J Nutr* **131**, 2811-2818 (2001).
https://doi.org:10.1093/jn/131.11.2811

34   Zhang, N. Role of methionine on epigenetic modification of DNA methylation and
gene expression in animals. *Anim Nutr* **4**, 11-16 (2018).
https://doi.org:10.1016/j.aninu.2017.08.009

35   Hwang, J. W., Cho, Y., Bae, G. U., Kim, S. N. & Kim, Y. K. Protein arginine
methyltransferases: promising targets for cancer therapy. *Exp Mol Med* **53**, 788-
808 (2021). https://doi.org:10.1038/s12276-021-00613-y

36   Al-Hamashi, A. A., Diaz, K. & Huang, R. Non-Histone Arginine Methylation by
Protein Arginine Methyltransferases. *Curr Protein Pept Sci* **21**, 699-712 (2020).
https://doi.org:10.2174/1389203721666200507091952

37   Lam, J. & Wulff, H. The Lymphocyte Potassium Channels Kv1.3 and KCa3.1 as
Targets for Immunosuppression. *Drug Dev Res* **72**, 573-584 (2011).
https://doi.org:10.1002/ddr.20467

38   Rader, R. K. *et al.* T cell activation is regulated by voltage-dependent and calcium-
activated potassium channels. *J Immunol* **156**, 1425-1430 (1996).

39   Cahalan, M. D. & Chandy, K. G. The functional network of ion channels in T
lymphocytes. *Immunol Rev* **231**, 59-87 (2009). https://doi.org:10.1111/j.1600-
065X.2009.00816.x

40   Vodnala, S. K. *et al.* T cell stemness and dysfunction in tumors are triggered by a
common mechanism. *Science* **363** (2019).
https://doi.org:10.1126/science.aau0135

41   Bachmann, M. *et al.* Pharmacological targeting of the mitochondrial calcium-
dependent potassium channel KCa3.1 triggers cell death and reduces tumor
growth and metastasis in vivo. *Cell Death Dis* **13**, 1055 (2022).
https://doi.org:10.1038/s41419-022-05463-8

42   Wulff, H. *et al.* Design of a potent and selective inhibitor of the intermediate-
conductance Ca2+-activated K+ channel, IKCa1: a potential immunosuppressant.
Proc Natl Acad Sci U S A **97**, 8151-8156 (2000).
https://doi.org:10.1073/pnas.97.14.8151

43   Beltra, J. C. *et al.* Developmental Relationships of Four Exhausted CD8(+) T Cell
Subsets Reveals Underlying Transcriptional and Epigenetic Landscape Control
Mechanisms. *Immunity* **52**, 825-841 e828 (2020).
https://doi.org:10.1016/j.immuni.2020.04.014

44   McLane, L. M., Abdel-Hakeem, M. S. & Wherry, E. J. CD8 T Cell Exhaustion
During Chronic Viral Infection and Cancer. *Annu Rev Immunol* **37**, 457-495 (2019).
https://doi.org:10.1146/annurev-immunol-041015-055318

45   Thompson, E. D., Enriquez, H. L., Fu, Y. X. & Engelhard, V. H. Tumor masses
support naive T cell infiltration, activation, and differentiation into effectors. *J Exp*
Med **207**, 1791-1804 (2010). https://doi.org:10.1084/jem.20092454

46   Yu, P. *et al.* Priming of naive T cells inside tumors leads to eradication of
established   tumors.   *Nat   Immunol*   **5**,   141-149   (2004).
https://doi.org:10.1038/ni1029
47   Prokhnevska, N. *et al.* CD8(+) T cell activation in cancer comprises an initial
activation phase in lymph nodes followed by effector differentiation within the
tumor.   *Immunity*   **56**,   107-124   e105   (2023).
https://doi.org:10.1016/j.immuni.2022.12.002
48   Sterner, R. C. & Sterner, R. M. CAR-T cell therapy: current limitations and potential
strategies. *Blood Cancer J* **11**, 69 (2021). https://doi.org:10.1038/s41408-021-
00459-7
49   Marofi, F. *et al.* CAR T cells in solid tumors: challenges and opportunities. *Stem
Cell Res Ther* **12**, 81 (2021). https://doi.org:10.1186/s13287-020-02128-1
50   Du, H. *et al.* Antitumor Responses in the Absence of Toxicity in Solid Tumors by
Targeting B7-H3 via Chimeric Antigen Receptor T Cells. *Cancer Cell* **35**, 221-237
e228 (2019). https://doi.org:10.1016/j.ccell.2019.01.002
51   Sun, Q. *et al.* Immune checkpoint therapy for solid tumours: clinical dilemmas and
future   trends.   *Signal   Transduct   Target   Ther*   **8**,   320   (2023).
https://doi.org:10.1038/s41392-023-01522-4
52   Scharping, N. E. *et al.* The Tumor Microenvironment Represses T Cell
Mitochondrial Biogenesis to Drive Intratumoral T Cell Metabolic Insufficiency and
Dysfunction.   *Immunity*   **45**,   374-388   (2016).
https://doi.org:10.1016/j.immuni.2016.07.009
53   Franco, F., Jaccard, A., Romero, P., Yu, Y. R. & Ho, P. C. Metabolic and epigenetic
regulation   of   T-cell   exhaustion.   *Nat   Metab*   **2**,   1001-1012   (2020).
https://doi.org:10.1038/s42255-020-00280-9
54   Rudloff, M. W. *et al.* Hallmarks of CD8(+) T cell dysfunction are established within
hours of tumor antigen encounter before cell division. *Nat Immunol* **24**, 1527-1539
(2023). https://doi.org:10.1038/s41590-023-01578-y
55   Shakiba, M. *et al.* TCR signal strength defines distinct mechanisms of T cell
dysfunction   and   cancer   evasion.   *J   Exp   Med*   **219**   (2022).
https://doi.org:10.1084/jem.20201966
56   Abe, B. T., Shin, D. S., Mocholi, E. & Macian, F. NFAT1 supports tumor-induced
anergy   of   CD4(+)   T   cells.   *Cancer   Res*   **72**,   4642-4651   (2012).
https://doi.org:10.1158/0008-5472.CAN-11-3775

FIGURE 1

Figure 1: TCR-mediated, rapid methionine consumption governs T cell effector function. a-b, Quantification of intracellular amino acids at 10 min (a) and SAM and SAH up to 60 min (b) in OT-I T cells activated with 10 ng/ml SIINFEKL (n=3). **c,** T cell proliferation via cell-trace violet staining of OT-I CD8$^+$ T cells activated in either 0.1 mM Met or 0.03 mM Met for the indicated times before restoration to 0.1 mM Met in 0.03 mM Met conditions and then analyzed 72 hrs post

activation (n=3). Representative of 3 samples per group. **d,** Schematic design for OT-I T cell

activation initially in 0.1 or 0.03 mM Met, followed by restoration of Met to 0.1 mM for 24 hrs

before injection into B16-OVA tumour-bearing mice. **e,** Tumour growth of B16-OVA in *Rag1*$^{-/-}$

treated with OT-I CD8$^+$ T cells activated as described in (d) for 30 min-6hrs (n=5). **f-g,** Tumour

growth (f) and survival (g) of B16-OVA tumours in *Rag1*$^{-/-}$ mice after transfer of 24 hr-activated

OT-I T cells with first 30 min of stimulation being in 0.1 or 0.03 mM Met with 2.5ng/ml

SIINFEKL before restoration to 0.1 mM Met (n=5). **h-i,** Tumour growth (h) and survival (i) of

B16-OVA tumours treated with activated GP33$^+$-memory T cells as described in extended data

fig. 1d (n=5). Data are mean±s.e.m. Unpaired two-tailed Student's t-test (a, b), two-way ANOVA

(e, f, h), Mantel-Cox log rank test (g, i) and paired two-tailed Student's t-test (G-L) *P < 0.05, **P

< 0.01, ***P < 0.001, ****P < 0.0001.

FIGURE 2

Figure 2: Reduced methionine availability during TCR signaling promotes T cell exhaustion.

a, Schematic of experimental design. OT-I CD8$^+$ T cells with different congenic markers were initially activated in 0.1 mM or 0.03 mM Met for 30 min with replenishment of Met in 0.03 mM to 0.1 mM Met for 24 hrs, transferred into B16-OVA tumour-bearing *Rag1$^{-/-}$* mouse at a 1:1 ratio, and analyzed day 12 post T cell transfer. **b-c,** Frequencies (b) and absolute number (c) of

transferred T cells isolated from B16-OVA tumours at D12 post transfer (n=7). **d-e,** Frequency of

PD1$^+$ and IFNγ^+TNFα^+ OT-I CD8$^+$ TIL, initially activated in 0.1 mM and 0.03 mM Met and

assessed at D12 post injection (as in a) (n=7). **f-g,** Contour plot (f) and frequency of TOX$^+$ cells

(g) and of OT-I CD8$^+$ TIL as in d (n=7). **h-i,** Representative contour plot (h) and quantification (i)

of TCF1 and Tim-3 expressing cells from TOX$^+$ CD8$^+$ TIL from OT-I CD8$^+$ T cells isolated at

D12 post transfer as in (a) (n=7). **j,** T cell exhaustion gene sets in GSEA analysis of RNA-Seq of

OT-I CD8$^+$ TIL on d9 post T cell injection as in Fig. 1f (n=4). Data are mean±s.e.m. paired two-

tailed Student's t-test. *P < 0.05, **P < 0.01, ***P < 0.001.

FIGURE 3

 Figure 3: Extracellular methionine availability regulates Ca^{2+}-mediated NFAT1 activity.

 a, Fluo-8 AM analysis of Ca^{2+} flux in $CD8^+$ T cells activated with anti-CD3 and anti-CD28 by

 anti-hamster IgG crosslinking in either 0.1 mM or 0.03 mM Met containing Ca^{2+} Ringer solution

 with addition of 2 mM Ca^{2+} to measure Ca^{2+} influx. The area under the curve is measured from

 350-500 sec. Ionomycin then added as a positive control for maximal flux. Representative of 2

 experiments. **b,** Representative images of T cells, cultured with or without anti-CD3/28 dynabeads

 (dark grey masked), in 0.1 mM and 0.03 mM Met, with or without cyclosporin-A treatment for 30

 min, stained with NFAT1 (red, marked), Hoechst (blue) and Phalloidin (green). **c,** Quantification

 of NFAT1 intensity as nuclear to total cell ratio (c) (See Fig. S4A for unmasked figure). Scale

 bar=10μm. (Each circle represents one cell, n=40 cells). **d,** Histogram of NFAT1 binding signals

 (read count per million reads normalized to background) from NFAT1 CUT&RUN on T cells

 initially activated in 0.1 or 0.03 mM Met for 30 min and assessed 24 hrs post activation. **e-f,**

 NFAT1 CUT&RUN peaks at the known target genes (e) and quantification of known NFAT1-

 binding regions of *Lag3, Pdcd1, Havcr2, Ctla4* and *Tnfrsf9 and Tox* (f, normalized read count (see

 differential binding analysis)) in T cells initially activated in 0.1 or 0.03 mM Met for 30 min and

 assessed 24 hrs post activation (n=2). Data are mean±s.e.m. Unpaired two-tailed Student's t-test

 (C, D). *P < 0.05, **P < 0.01, ***P < 0.001, ****P < 0.0001.

FIGURE 4

Figure 4. Methionine mediated KCa3.1 methylation upon TCR activation regulates T cell effector function.

a-b, Representative images (a) and quantification (b) of methyl-arginine in CD8$^+$ T cells activated with anti-CD3/28 dynabeads (circled in red channel images) in 0.1, 0.0, and 0.03 mM Met for 30 min, stained with pan-meArg (red, with arrows), Hoechst (blue), and Phalloidin (green) (scale bar=10 μm) (each circle represents one cell. n= 25 cells). **c,** Venn diagram of enriched methyl-arginine proteins at 30 min post activation of T cells in 0.1 mM vs 0.0 mM or 0.1 mM vs 0.03 mM Met, as identified by TMT-MS. **d,** Heat-map (Log$_2$ fold change) of TMT-MS identified proteins ($p<0.05$) with enriched methyl-arginine in T cells activated in 0.1 mM vs 0.0mM or 0.03 mM Met by IgG crosslinking with anti-CD3 for 30 min (n=3). **e,** Predicted interaction of KCa3.1 monomer (blue cartoon) with calmodulin (yellow cartoon) as un-methylated (left), SDMA (center) and ADMA (right). The Calmodulin-binding pocket is represented as surface rendering with interacting amino acids as sticks (negative charges in red, polar residues in green) and R352 as ball and sticks; introduction of a hydrophobic methyl group (middle and bottom) weakens the interactions between calmodulin and the KCa3.1 monomer. **f,** MD simulation analysis: Analysis of variations of distance of salt bridge between calmodulin E84 and demethylated R352 WT (purple), SDMA R352 (green) and ADMA R352 (blue) over the course of simulation time (averaged across all four monomers, three MD trials). **g,** Fluo-8 AM analysis of Ca^{2+} flux in CD8$^+$ T cells activated with anti-CD3 and anti-CD28 by anti-hamster IgG crosslinking in Ca^{2+}-free Ringer solution supplemented with either 0.1 mM or 0.03 mM Met and treated with either with DMSO or 1μM TRAM-34. Area under the curve measured from 250-450 sec (representative of 2 experiments). **h,** Tumour growth of B16-OVA tumours in *Rag1$^{-/-}$* mouse after transfer of OT-I CD8$^+$ T cells activated in 0.1 mM or 0.03 mM Met with either DMSO or 1 μM TRAM-34 for 30 min and cultured for 24 hrs (n=5). Data are mean±s.e.m. Unpaired two-tailed Student's t-test (b), 2-way ANOVA (h), two-sample T-test (g) as described in the methods. *$P < 0.05$, **$P < 0.01$, ***$P < 0.001$.

FIGURE 5

Figure 5. Ablation of KCa3.1 R350 methylation increased Ca²⁺-mediated NFAT1 activity,

promoting T cell dysfunction.

a, Fluo-8 analysis of Ca²⁺ flux in KCa3.1^{WT} and KCa3.1^{R350A} T cells activated with anti-CD3 and

anti-CD28 by anti-hamster IgG crosslinking in Ca²⁺-free Ringer solution. Area under the curve

calculated from 200-350 sec. **b-c,** Representative image (c) and quantification of NFAT1intensity

as nuclear to total cell ratio (c) in KCa3.1^{WT} and KCa3.1^{R350A} OT-I T cells activated for 30 min

700 with anti-CD3/28 dynabeads (dark grey masked) (See Extended Data Fig. 6F for unmasked figure),

701 scale bar=10 μm (each circle represents one cell, n=40 cells). **d,** Chromatin accessibility heatmap

702 of T cells expressing activated KCa3.1^{WT} or KCa3.1^{R350A} with each row representing peaks

703 (p<0.05 and log2 fold-change >1.5) displayed over the span of 2kb window with peak as center

704 (grouped from least to max differential region), analyzed 24 hrs post activation with 2.5ng/ml

705 SIINFEKL (n=3). **e,** GOBP terms associated with negative regulation of T cells enriched in OT-I

706 T cells expressing KCa3.1^{R350A} vs KCa3.1^{WT} activated as in (b) (n=3). **f,** T cell exhaustion gene

707 sets in GSEA analysis of differential accessible promoter reads from ATAC-Seq of OT-I T cells

708 expressing KCa3.1^{WT} or KCa3.1^{R350A}, activated with 2.5 ng/ml SIINFEKL for 24 hrs (n=3). **g-h,**

709 Tumour growth (g) and survival (h) of B16-OVA tumour-bearing RAG$^{-/-}$ mice after transfer of

710 KCa3.1^{WT} and KCa3.1^{R350A} OT-I T cells (n=5). **i,** B16-OVA tumours harvested at D12 post

711 transfer of KCa3.1^{WT} and KCa3.1^{R350A} OT-I T cells (n=4). Data are mean±s.e.m. Two-sample T-

712 test (a) as described in the methods, unpaired two-tailed Student's t-test (c), 2-way ANOVA (g),

713 and Mantel-Cox log rank test (h). **P < 0.01, ****P < 0.0001.

FIGURE 6

Figure 6. Ablation of KCa3.1 R350 methylation promotes T cell exhaustion in tumours and infection models.

a, Schematic of experimental design to assess congenically distinct KCa3.1^{WT} and KCa3.1^{R350A} OT-I T cells that were mixed at a 1:1 ratio and transferred into B16-OVA tumour-bearing $Rag1^{-/-}$ mice and analyzed day 12 post T cell transfer. **b-d** Frequency of Tim-3$^+$PD1$^+$ (b), histogram and quantification of TOX MFI (c), and frequency of IFNγ$^+$TNFα$^+$ (d) among KCa3.1^{WT} and KCa3.1^{R350A} OT-I TIL at D12 post T-cell transfer as in (a) (n=7). **e,** Frequency of TCF1$^+$Tim-3$^-$ TEXprog and TCF1$^-$Tim-3$^+$ TEXterm in KCa3.1^{WT} and KCa3.1^{R350A} OT-I TIL at D12 post T-cell transfer as in (a) (n=7). **f,** Expression of memory markers CD62L and CD127$^+$CD27$^+$ on OT-I T cells expressing KCa3.1^{WT} or KCa3.1^{R350A} isolated at D12 post transfer into B16-OVA tumour-bearing mice as in (a) (n=7). **g,** Schematic of experimental design to assess congenically distinct KCa3.1^{WT} and KCa3.1^{R350A} Thy1.1 P14 T cells that were mixed at a 1:1 ratio and transferred into WT mice infected with either LCMV-Armstrong or LCMV-Clone-13. Spleens were analyzed at D9 post infection. **h-i,** Expression of TOX (h) and frequency of TOX$^+$ and PD1$^+$Tim-3$^+$ (i) in KCa3.1^{WT} and KCa3.1^{R350A} P14 CD8$^+$ T cells from spleens at D9 post infection with LCMV-Armstrong (n=4). **j-k,** Expression of TOX (j) and frequency of TOX$^+$ and PD1$^+$Tim-3$^+$ (k) in KCa3.1^{WT} and KCa3.1^{R350A} P14 CD8$^+$ T cells from spleens at D9 post infection with LCMV-Clone-13 (p) (n=4). **l-m,** Frequency of CX3CR1$^+$PD1$^+$ TEXprog (l) and TCF1$^-$Tim-3$^+$ TEXterm (m) in KCa3.1^{WT} and KCa3.1^{R350A} P14 CD8$^+$ T cells from spleens at D9 post infection as in (g) (n=4). Data are mean±s.e.m. Paired two-tailed Student's t-test. *P < 0.05, **P < 0.01, ***P < 0.001, ****P < 0.0001.

Figure 7. Acute methionine supplementation promotes CD8$^+$ T cell-mediated tumour control and enhances ICB treatment.

a, Nuclear NFAT1 quantification in CD44$^+$CD8$^+$ T cells isolated from B16 tumours and respective dLNs at D9 post implantation (each circle represents one cell, n=40 cells, n=5). **b,** Quantification of nuclear NFAT1 in CD44$^+$ CD8$^+$ T cells isolated from B16 tumours after 5 days of peri-tumoural supplementation of 50 μl HBSS or 61 μM Met per day (each circle represents one cell, n=40 cells, n=4). **c-d,** Tumour growth (c) and survival (d) of B16 tumour-bearing WT mice, supplemented either with HBSS or 61 μM Met peri-tumourally for five days as in (b) (n=10). **e-f,** B16 tumour growth (e) and survival (f) of WT mice, treated intraperitoneally with IgG isotype or anti-CD8 antibody at D1 and D3 and treated peri-tumourally with HBSS or Met from D7-D12 as in (b) (n=5, representative of two experiments). **g-h,** Tumour growth (g) and survival (h) of F420 tumour-bearing *Rag1*$^{-/-}$ mice injected with B7-H3 CAR T-cells at D0 and peri-tumourally treated with HBSS or Met from D1-D6 as in (b) (n=5). **i-k,** Tumour growth (i, j) and survival (k) of MC38 tumour-bearing WT mice, treated either with anti-PD1 or IgG-isotype (shown by arrows in i) and fed with either control (1% Met) or Met-rich (1.5% Met) diet as shown in Extended Data Fig. 10H (n=10). Data are mean±s.e.m. Unpaired two-tailed Student's t-test (a), 2-way ANOVA (c, e, g and j), and Mantel-Cox log rank test (d, f, h and k). Statistical analysis of (b) is described in methods. *$P < 0.05$, **$P < 0.01$, ***$P < 0.001$, ****$P < 0.0001$.

Extended Data Fig. 1

Extended Data Figure 1. Methionine availability during TCR activation determines T cell fate.

a, Log2FC in serum Met levels post LCMV-C13 chronic infection analyzed by Lercher et. al, *Immunity,* 2019 (ref. 7). **b-c,** Tumour growth (b) and survival (c) of B16-OVA tumour-bearing WT mice treated with or without OT-I CD8$^+$ T cells activated in 0.1 or 0.03 mM Met for 30 min before restoring Met to 0.1 mM in 0.03 mM Met for 24 hrs, as described in Fig. 1d (n=5). **d,** Schematic of experimental design for generation of LCMV-GP33 specific memory T cells, which were then activated as described in (Fig. 1d) before injection into B16-OVA tumour-bearing *Rag1$^{-/-}$* mice. **e,** Schematic design for ATAC-Seq on OT-I T cell activation initially in 0.1 or 0.03 mM

768   Met followed by restoration of Met to 0.1 mM for 24 hrs. Data are mean±s.e.m. 2-way ANOVA

769   (b), and Mantel-Cox log rank test (c). **$P < 0.01$, ***$P < 0.001$.

Extended Data Fig. 2

Extended Data Figure 2. Reduced methionine during TCR activation promotes T cell exhaustion.

a, PCA analysis of ATAC-Seq of OT-I T cells activated with 2.5ng/ml SIINFEKL in 0.1 mM and 0.03 mM Met and cultured in 0.1 mM Met for 24 hrs. n=3. **b,** Chromatin accessibility heatmap of T cells initially activated in 0.1 or 0.03 mM Met with each row representing peaks (p<0.05 and log2 fold-change >1.5) displayed over the span of 2kb window with peak as center (grouped from least to max differential region), analyzed from ATAC-Seq from (a). (n=3). **c,** Over-representation analysis of the DAR from (b) showing the top 5 genesets associated with exhaustion. **d,** Homer analysis of (a) showing top 20 motifs and their percent coverage. Colour corresponds to the common exhaustion-associated motifs also described in the indicated studies (blue and purple). **e-f,** Tumour growth (e) and weight (f) of B16-OVA tumour-bearing $Rag1^{-/-}$ mice injected with OT-I T cells as in (Fig. 1d) (n=5). **g,** Frequency of CD62L$^+$, Tcm (CD62L^{hi} CD44^{hi}), TOX expression (MFI) and IFN-γ^+ in OT-I CD8$^+$ TIL isolated from B16-OVA tumour-bearing $Rag1^{-/-}$ mice at D9 post-injection with OT-I T cells activated as in Fig. 1d (n=5). **h,** Frequency of PD1$^+$ Tim-3$^+$ on OT-I CD8$^+$ T cells at D12 post transfer of OT-I T cells as in Fig. 2a (n=7). **i,** Frequency of TCF1$^+$ on OT-I CD8$^+$ T cells at D12 post transfer of OT-I T cells as in Fig. 2a (n=7). **j-l,** Frequency of TCF1$^+$Tim-3$^-$ TEXprog, CX3CR1$^+$Ly108$^+$ TEXprogenitor-like (j) and, TCF1$^-$Tim-3$^+$ TEXterm (k) CD62L$^+$ T memory and CD62L$^+$CD44$^+$ T central memory cells (l) in OT-I CD8$^+$ T cells as in Fig. 2a (n=7, n=4). **m-p,** Representative contour plot and quantification of CD127$^+$CD27$^+$ memory cells (m, n) and KLRG1$^+$CD127$^-$ terminal effector cells (o, p) in OT-I CD8$^+$ TIL isolated at D12 post of cells activated as in Fig. 1d into B16-OVA tumour-bearing $Rag1^{-/-}$ mice, as in Fig. 2a (n=4). Data are mean±s.e.m. Statistical analyses performed using 2-way ANOVA (e), unpaired two-tailed Student's t-test (f, g) paired two-tailed Student's t-test (h-p). *P < 0.05, **P < 0.01, ***P < 0.001.

Extended Data Figure 3. RNA-Seq analysis shows enrichment of exhaustion signature in low Met.

a, Volcano plot and top ten differentially expressed genes (red) (0.1 mM vs 0.03 mM, p<0.05, FC>1.5) from OT-I CD8$^+$ TIL isolated from mice injected with OT-I T cells, initially activated in 0.1 mM Met or 0.03 mM Met (as in Fig. 1d), at day 9 post injection into B16-OVA tumour-bearing mice (n=4). **b,** Heatmap of DE genes, which shows increased expression in cells initially activated in either 0.03 mM Met or 0.1 mM Met from the RNAseq in (a) (n=4). **c,** Over-representation analysis of DE genes with higher expression in TIL initially activated in 0.1 mM Met, showing enrichment of biological processes and transcription factors associated with T cell effector function and regulation. **d,** Over-representation analysis of DE genes with higher expression in TIL initially activated in 0.03 mM Met, showing enrichment of gene sets and transcription factors associated with promotion and maintenance of the exhausted T cell state. **e,** GSEA analysis of hallmark genesets as analyzed from RNA-Seq of OT-I TIL initially activated in 0.1 or 0.03 mM Met, isolated from B16-OVA tumours at D9 post transfer (n=4). RNA-Seq statistics were applied as described in methods.

Extended Data Fig. 4

Extended Data Figure 4. Met limitation results in increased TCR-mediated NFAT1 activation.

a, Fluo-8 AM analysis of Ca^{2+} flux in $CD8^+$ T cells activated with anti-CD3 and anti-CD28 and anti-hamster IgG crosslinking in Ca^{2+}-free Ringer solution containing either 0.1 mM or 0.03 mM Met and treated for one hour with DMSO or 5μM YM58483. Area under the curve calculated from 150-300 sec (n=2). **b,** Un-edited image of Figure. 3b showing the unmasked, red anti-CD3/38 dynabeads which are masked grey in the main figure to highlight NFAT1 staining. **c,** NFAT1 intensity quantification (nuclear to total cell ratio) of $CD8^+$ T cells activated in either 0.1 mM, 0.00 mM or 0.03 mM Met for 30 min with anti CD3/28 dynabeads and stained for NFAT1 (each circle represents one cell, n=40 cells). **d,** Quantification of nuclear NFAT1 from the experiment shown in Figure 3b. **e,** NFAT1 intensity quantification (nuclear to total cell ratio) of $CD8^+$ T cells activated for 30 min by CD3/28 dynabeads in either 0.1 mM or 0.03 mM Met in the presence of DMSO or 5μM YM58483 (each circle represents one cell, n=40 cells). **f,** NFAT2 quantification (nuclear to total cell ratio) of $CD8^+$ T cells, activated in 0.1 mM or 0.03 mM Met for 30 min with anti-CD3/28 dynabeads (each circle represents one cell, n=40 cells). **g,** NFAT1 quantification (nuclear to total cell ratio) of previously activated OT-I T cells, treated with anti-CD3/28 dynabeads for 30 min. T cells were initially activated with 2.5ng/ml SIINFEKL for 24 hrs in control medium and rested with 10 ng/ml mIL-7 for 48 hrs (each circle represents one cell, n=40). **h,** Normalized RNA expression quantified by quantitative-PCR performed 24 hrs post activation of OT-I $CD8^+$ T cells activated with 2.5 ng/ml SIINFEKL in 0.1 mM or 0.03 mM Met for 30 min followed by 0.1 mM Met (n=6). Data are mean±s.e.m. Two-sample T-test (a) as described in the methods, unpaired two-tailed Student's t-test (c-g) and one-tailed Student's t-test (h). *P < 0.05, **P < 0.01, ***P < 0.001, ****P < 0.0001.

Extended Data Fig. 5

Extended Data Figure 5. Initial Met limitation doesn't affect DNA or histone methylation but alters KCa3.1 arginine methylation.

a, Methylation potential (SAM/SAH) of 10 ng/ml SIINFEKL activated OT-I T cells in 0.1 mM or 0.0 mM Met at 10, 30 and 60 min (n=3). **b,** % 5-methylcytosine (%5-mc) of OT-I CD8$^+$ T cells activated with 2.5 ng/ml SIINFEKL in 0.1 mM or 0.03 mM Met for 30 min, then 0.1 mM Met for 24 hr (n=3). **c-d,** Histograms (top) and heat-map (bottom) of H3K4me3 (c) and H3K27me3 (d) CUT&RUN reads (read count per million, normalized to background) in OT-I CD8$^+$ T cells activated initially in 0.1 mM or 0.03 mM Met with restoration to 0.1 mM after 30 min and processed at 24 hrs post activation (n=3). **e,** Un-edited image of Figure. 4a showing the unmasked, red CD3/38 dynabeads which are masked grey in the main figure to highlight me-Arg staining. **f,** Conserved arginine (highlighted red) in KCa3.1 as analyzed by protein-BLAST in multiple species and visualized using Jalview. **g,** Tetrameric assembly of human KCa3.1-Calmodulin complex within the membrane (top). KCa3.1 is represented as ribbons and calmodulin quartet as cartoon with interacting Ca^{2+} ions as purple spheres. K$^+$ ions are represented as cyan spheres. KCa3.1 monomer (blue) is shown in complex with calmodulin (yellow) along with demethylated arginine-352 (green) (bottom). **h,** MD simulations analysis: Illustration of total count of negatively charged residues within a 4Å radius of all four R352/SDMA/ADMA of huKCa3.1 the course of simulation time (averaged across all four monomers, three MD trials). **i,** NFAT1 quantification (nuclear to total cell ratio) in OT-I CD8$^+$ T cells activated for 30 min with anti CD3/28 dynabeads in 0.03 mM Met (left) or 0.1 mM Met (right) in the presence of 0.5, 1 or 2 μM TRAM-34 (each circle represents one cell, n=45). **j,** Schematic design for OT-I T cell activation initially in 0.1 or 0.03 mM Met, treated with DMSO or TRAM-34 for 30 min, followed by washing and culturing in 0.1 mM Met+SIINFEKL (2.5ng/ml) for 24 hrs before injection into B16-OVA tumour-bearing *Rag1$^{-/-}$* mice. Data are mean±s.e.m. Unpaired one-tailed Student's t-test (a) or unpaired two-tailed Student's t-test (b, i). *$P < 0.05$, **$P < 0.01$.

Extended Data Fig. 6

Extended Data Figure 6. KCa3.1 R350 methylation regulates Ca^{2+}- mediated NFAT1 activation.

a, Schematic of experimental design of endogenous KCa3.1 knockdown with expression of KCa3.1^{WT} and KCa3.1^{R350A} in Cas9-OT-I or P14$^+$ CD8$^+$ T cells. **b,** Sorting strategy for *in vitro*-generated KCa3.1^{WT} and KCa3.1^{R350A} T cells; gate shows the cells that express WT or R350A KCa3.1 with endogenous KCa3.1 knockdown. **c,** Immunoblot showing knockdown of KCa3.1 with sgRNA1 and sgRNA2, compared to control. **d,** Immunoblot of KCa3.1 expression in KCa3.1^{WT}- and KCa3.1^{R350A}-expressing CD8$^+$ T cells. **e,** ICR-1 AM analysis of Ca^{2+} flux in KCa3.1^{WT} and KCa3.1^{R350A} T cells activated with anti-CD3 and anti-CD28 by anti-hamster IgG crosslinking in Ca^{2+}-free Ringer solution. Area under the curve is calculated from 200-350 sec. **f,** Un-edited image of Figure. 5b showing the unmasked CD3/38 dynabeads which are masked grey in the main figure to highlight NFAT1 staining. **g,** Nuclear NFAT1quantification of KCa3.1^{WT} and KCa3.1^{R350A}, either control (no Dynabeads) or activated with anti-CD3/28 Dynabeads for 10 min (each circle represents one cell, n=45 cells). Data are mean±s.e.m. Two-sample T-test (a) as described in the methods and unpaired two-tailed Student's t-test. ****P < 0.0001.

Extended Data Fig. 7

Extended Data Figure 7. Ablation of KCa3.1 R350 methylation shows similar ATAC-Seq
886

signature as of T cells activated in reduced Met.
887

a, PCA analysis of ATAC-Seq of KCa3.1^WT- and KCa3.1^R350A- activated with 2.5 ng/ml
888

SIINFEKL for 24 hrs. n=3. **b,** Common differentially accessible regions (DAR) (p<0.05, fold
889

change>1.5) between ATAC-Seq of OT-I cells expressing KCa3.1^WT or KCa3.1^R350A, or OT-I cells
890

initially activated in 0.1 mM or 0.03 mM Met as in (a) and Extended Data Fig. 2b. **c,** Over-
891

representation analysis of the common DAR from (b) showing the top 8 gene sets associated with
892

exhaustion and activation. **d-e,** Top 10 motifs analyzed by HOMER in ATAC-Seq DAR's from
893

894   24hrs activated OT-I CD8+ T cells expressing KCa3.1^{R350A} vs KCa3.1^{WT} (d), and from the

895   common DAR's from (b) (e). Colour corresponds to the common exhaustion-associated motifs

896   also described in the indicated studies (blue and purple).

Extended Data Figure 8. Ablation of KCa3.1 R350 methylation promotes T cell exhaustion *in-vivo*.

a, Tumour weight of B16-OVA tumours isolated at D12 post transfer of OT-I T cells expressing KCa3.1^{WT} or KCa3.1^{R350A} (n=4). **b,** Surface expression of PD1$^+$, Lag3$^+$ and PD1$^+$Tim-3$^+$ in CD8$^+$ TIL isolated at D12 post transfer into B16-OVA tumour-bearing mice of OT-I T cells expressing KCa3.1^{WT} or KCa3.1^{R350A} (n=4). **c,** Representative histogram of TOX expression and quantification in CD8$^+$ TIL isolated at D12 post transfer of OT-I T cell expressing KCa3.1^{WT} or KCa3.1^{R350A} as (b) (n=4). **d,** Frequency of CD69^{lo}Ly108^{hi} TEXprog in KCa3.1^{WT} and KCa3.1^{R350A} OT-I T cells at D12 post transfer (n=4). **e,** Representative contour plot (left) and quantification of IFNγ$^+$TNFα$^+$ (right) in OT-I T cells expressing KCa3.1^{WT} or KCa3.1^{R350A} at D12 post transfer as (b) (n=4). **f-g,** Expression of EOMES$^+$PD1$^+$ (f) and representative contour plot of expression of TCF1 and Tim-3 (g) on congenically marked OT-I T cells expressing KCa3.1^{WT} or KCa3.1^{R350A}, mixed and transferred at a ratio of 1:1. TIL were isolated from B16-OVA tumours at D12 post T cell transfer (n=7). **h-i,** Representative contour plot (h) and quantification of KLRG1$^+$CD127$^-$ terminal effector cells (i) on congenically marked OT-I T cells expressing KCa3.1^{WT} or KCa3.1^{R350A} as in (f), isolated at D12 post transfer into B16-OVA tumour-bearing mice (n=7). **j,** Frequency of KCa3.1^{WT}- and KCa3.1^{R350A}-expressing P14 T cells in spleens, D9 post infection with either LCMV-Armstrong or LCMV-Clone-13 (n=4). **k,** Representative contour plot of PD1$^+$Tim3$^+$ KCa3.1^{WT}- and KCa3.1^{R350A}-expressing P14 T cells in spleens, D9 post infection with either LCMV-Armstrong or LCMV-Clone-13 (n=4). **j,** Frequency of Tim-3$^+$Cx3CR1$^+$PD1$^+$ TEXeff KCa3.1^{WT}- and KCa3.1^{R350A}-expressing P14 T cells in spleens, D9 post infection with either LCMV-Armstrong or LCMV-Clone-13 (n=4). **m-o,** Representative contour plot (m) and quantification of KLRG1$^+$CD127$^-$ terminal effector cells (n) and KLRG1$^-$CD127$^+$ memory precursor cells (o) in spleens of animals injected with congenically marked P14 T cells expressing KCa3.1^{WT} or KCa3.1^{R350A}, mixed 1:1, and infected with either LCMV-Armstrong or LCMV-Clone-13. Analysis on d9 post infection (n=4). Data are mean±s.e.m. Unpaired two-tailed Student's t-test (a-e) and paired two-tailed Student's t-test (f-n). *P < 0.05, **P < 0.01, ***P < 0.001, ****P < 0.0001.

Extended Data Fig. 9

Extended Data Figure 9. Acute Met supplementation promotes anti-tumour T cell differentiation and tumour control.

a, Quantification of essential amino acids (left) and peak area of Met (right) in LN CD8$^+$ T cells and CD8$^+$ TIL, isolated at day 12 post tumour implantation (n=3). **b,** Quantification of intracellular Met in CD8$^+$ PBMC and CD8$^+$ TIL isolated from patient primary colorectal tumours (n= 8). **c,** Schematic of experimental design of acute HBSS/Met peritumoural treatment in sub-cutaneous tumours. **d,** Nuclear NFAT1 quantification of CD44$^+$ CD8$^+$ T cells isolated from B16 subcutaneous tumours and dLN, 5 days post daily peri-tumoural supplementation of 50 ml 61μM Met (M=1 mouse, each circle represents one cell, n=40 cells). **e-g,** Tumour growth (left) and survival (right) of WT mice with sub-cutaneous MC38 colorectal tumours (e) (n=10), Lewis-lung carcinoma (LLC) tumours (f) (representative of two experiments of n=5), and F420 osteosarcoma tumours (g) (n=5), treated with either HBSS or 50 ml 61μM Met peri-tumourally daily for 5 days. **h-i,** Expression of PD1, PD1$^+$Tim-3$^+$ (h), Tcm (CD62L^{hi}Cd44^{hi}) and Tem (CD62L^{lo}CD44^{hi}) (i) on CD8$^+$ TIL isolated from B16 tumour-bearing mice 2 days after 5 daily peri-tumoural injection with HBSS or Met (n=5-4). **j,** Tumour growth of B16f10 and MC38 tumours in NSG mice treated peri-tumourally with 50 μl HBSS or 50 μl 61μM Met daily for 5 days (n=5). **k,** Representative flow cytometry plot of blood showing CD8$^+$ T cell depletion in anti-CD8 antibody treated mice compared to IgG-Isotype-treated mice (n=4). Data are mean±s.e.m. Unpaired (a, d, h and i) and paired (b) two-tailed Student's t-test, two-way ANOVA (e-g (left) and j), Mantel-Cox log rank test (e-f (right)). *P < 0.05, **P < 0.01, ***P < 0.001, ****P < 0.0001.

Extended Data Fig. 10

Extended Data Figure 10. Acute Met supplementation promote T cell mediated tumour
control and enhances tumour immunotherapies.
a, Schematic of experimental design of contralateral tumour experiment. **b,** Growth of B16f10
tumours implanted on left shoulder (dashed lines) and right flank (solid lines) following peri-
tumoural injection of flank tumour with 50 µl HBSS or 61µM Met daily for 5 days (n=5,
representative of two experiments). **c,** B16 tumour growth in WT mice fed with either 1 % Met
chow or 1.5 % Met chow for 7 days post tumour implantation (n=5). **d,** Schematic of experimental
design to assess the effect of HBSS or Met supplementation on CAR-T cell therapy in murine solid
tumour model. **e,** Schematic of design to assess tumour growth with anti-PD1 treatment and with
either peri-tumoural HBSS or Met supplementation. **f-g,** Tumour growth (f) and survival (g) of
MC38 tumour-bearing WT mice, treated either with anti-PD1 or IgG-isotype and supplemented
with HBSS or Met peri-tumourally for five days as in (e) (n=5, representative of two experiments).
h, Schematic of experimental design to assess tumour growth with anti-PD1 treatment on sub-
cutaneous tumour-bearing mouse, supplemented with either control diet or Met-rich diet. Data are
mean±s.e.m. 2-way ANOVA (b, c, and f), and Mantel-Cox log rank test (g). *$P < 0.05$, **$P < 0.01$,
***$P < 0.001$, ****$P < 0.0001$.
969

Extended Fig. 11

Extended Data Fig. 11: Methionine metabolism regulates TCR-mediated Ca²⁺ flux and T cell

fate. Diagram illustrating the impact of methionine metabolism on TCR-dependent methylation of

KCa3.1, leading to regulation of Ca²⁺ flux and subsequent activation of NFAT1. Low extracellular

Met levels lead to decreased methylation potential, reducing KCa3.1 R350 methylation. This

results in increased Ca²⁺ flux and downstream NFAT1 activation and consequent T cell

hyperactivation and exhaustion.

Materials and Methods:

Reagents

RPMI 1640 (cat. 11875093, Gibco) and methionine-free RPMI 1640 (cat. A1451701; Gibco) supplemented with 10% dialyzed FCS (Atlanta Biologicals), sodium pyruvate (cat. 11360070, Gibco), NEAA (11140050, Gibco), glutamine (cat. 25030081, Thermo-Fisher), and penicillin-streptomycin cocktail (cat. 15070063, Thermo-Fisher) and stored at 4°C. Methionine (cat. M5308, Sigma-Aldrich) was reconstituted in PBS (0.1 M) and stored at 4°C. SIINFEKL peptide (cat. 60193, AnaSpec) peptide was reconstituted in PBS at 0.5 mg/ml and stored at -20°C. TRAM-34 (cat. HY-13519, MedChem Express) was reconstituted at 1 M with DMSO. Cyclosporin A (CsA) (cat. 30024, Sigma-Aldrich) was reconstituted to a concentration of 50 mg/ml in DMSO. YM-58483 (cat. 3939, Tocris) was reconstituted at 20 mM stock in DMSO. All reconstituted reagents were stored at -20°C and aliquoted to avoid repeated thawing.

Mice

All mouse studies were conducted in accordance with protocols approved by the St. Jude Children's Research Hospital Committee on Care and Use of Animals, and in compliance with all relevant ethical guidelines. All mice were kept under a 12h-12h light-dark cycle in a specific-pathogen-free (SPF) facility at the institute's Animal Resource Center. Age-and sex-matched mice were randomly assigned to the experimental and control groups. Cas9-OT/P14^{+}-I (Cas9 Knock-in with OVA-specific transgenic TCR (OT-I) or LCMV-GP33 specific TCR(p14)), C57BL/6J (WT), OT-I, and $Rag1^{-/-}$ mice were bred in-house. The investigators were not blinded to the experiments and outcome assessments.

Cell lines

B16F10, B16F10-Ova (B16-Ova) melanoma cells were provided by Dr. Hongbo Chi. MC38 and MC38-Ova colon carcinoma cell lines were provided by Dr. Dario Vignali. The F420 sarcoma cell line was provided by Dr. Jason T Yustein. HEK293T and Lewis lung carcinoma (LLC) cells were purchased from the ATCC. Plat-e cells used for retroviral packaging were purchased from Cell Biolabs (cat. RV-101). All the cell lines used were tested and confirmed to be mycoplasma-negative but not independently authenticated.

Murine and Human CD8$^+$ T cell isolation and stimulation

Murine CD8$^+$ T cells were isolated from the spleen and lymph nodes (LN) by physical disruption through a 70μ strainer before treatment with erythrocyte-lysis buffer. Cells were subjected to CD8$^+$ T cell enrichment using the EasySep mouse CD8$^+$ T cell isolation kit (cat. 19853, StemCell), according to the manufacturer's protocol. Isolated T cells were then either activated with SIINFEKL peptide or mouse CD3/28 Dynabeads (cat. 11456D, Thermo-Fisher) in RPMI medium, as described above. Human CD8$^+$ T cells were isolated from apheresis rings from the St Jude Blood Donor Center. Blood obtained from an apheresis ring was mixed at a 1:1 ratio with PBS containing 2% FBS and carefully layered onto Lymphoprep (cat. 07801, StemCell) solution. The samples were then centrifuged at 800g for 30 min at room temperature without braking. The peripheral blood mononuclear cell (PBMC) layer was aspirated and washed twice in PBS with 2% FBS before enrichment of CD8$^+$ T cells using the Human CD8$^+$ T Cell Isolation Kit (cat. 17953, StemCell), according to the manufacturer's protocol. CD8$^+$ T cells were activated using human CD3/28 Dynabeads (cat. 11131D, Thermo Fisher Scientific) in RPMI medium. Apheresis rings were collected with written consent from the donors for research upon review and approval by the Institutional Review Board at St Jude Children's Research Hospital.

Ca^{2+} Flux measurement and analysis

Isolated T cells were labelled with 3 μM Fluo-8 AM (AAT Bioquest, cat. 21080) or 5 μM ICR-1AM (Ion Biosciences) with 0.04% pluronic F-127 (Thermo-Fisher, cat. P6867) for 2 hrs in HHBS at 37 °C and washed twice with Ca^{2+}-free Krebs-Ringer's solution (Thermo Fisher Scientific, cat. no. J67839-AP). The cells were then resuspended in Ca^{2+}-free Krebs-Ringer's solution and incubated with anti-CD3ε (4 μg/ml; 2C11, cat. BE0001, Bio-X-Cell) and anti-CD28 (4 μg/ml; 37.51, cat. BE0015, Bio-X-Cell) on ice for 10 min. 0.1 mM Met or 0.03 mM Met was added to the samples and the baseline was recorded before the addition of hamster-IgG for crosslinking. Ca^{2+} influx was measured after the addition of 2mM Ca^{2+}, 1uM ionomycin was added as a positive control for Ca^{2+} flux. All measurements were recorded on either Cytek Aurora (SpectroFlo) or BD-Fortessa flow cytometer.

To analyze temporal changes in fluorescence intensity, we binned the time variable into 10-second
intervals to calculate the mean log2-transformed fluorescence intensity within each 10-second bin
for each sample. We calculated the AUC across the entire time series, as well as within a focused
time window (x–y seconds) using the trapezoidal rule, which sums the area under each segment
formed by the binned time points. This approach captures cumulative fluorescence intensity over
time, providing a metric for comparing samples. To evaluate statistical significance between
groups, we performed two-sample t-tests on the log2-transformed fluorescence values within the
specified time window (x–y seconds).

Quantitative PCR

Isolated T cells were activated with 2.5ng/ml SIINFEKL in 0.1 mM Met or 0.03 mM Met for 30
min after which Met in 0.03 mM was restored to 0.1 mM. T cells were then further cultured for 24
hrs. Cells were washed twice with PBS and RNA was isolated using Direct-zol RNA miniprep Kit
(Zymo research, cat. no. R2050). RNA was quantified and processed for cDNA synthesis using
Reverse Transcriptase (Thermo Fisher Scientific, cat. no. 28025-013). cDNA was then used to
perform quantitative PCR with the specific primers (Table S2) using SYBR green PCR master mix
(Thermo Fisher Scientific, cat. no. 4309155). PCR reaction was run and quantified by QuantStudio
7 flex.

LCMV infection

LCMV Armstrong and Clone-13 was obtained from Dr. Shaabani, Scripps Research Institute. The
virus was propagated in BHK21(baby hamster kidney) cells as described[1] and stored at -80 degree
Celsius. For experiments, $2x10^6$ pfu of both Armstrong and Clone-13 was used at $2x10^6$
pfu/animal. The virus was injected intravenously, and the infected animals were maintained under
a 12h-12h light-dark cycle in a SPF, Biosafety Level-2 facility in the institute's Animal Resource
Center.

Cloning and virus production

Kcnn4 gRNA's were designed using the Broad Institute platform
(https://portals.broadinstitute.org/gpp/public/analysis-tools/sgrna-design) and cloned into the

ametrine-expressing retroviral vector LMPd-gRNA-mPGK-Ametrine

(sgRNA1:CACCGGGCAGGCTGTCAATGCCACG,

AAACCGTGGCATTGACAGCCTGCCC; sgRNA2:CACCGTGTGGGGCAAGATTGTCTGC,

AAACGCAGACAATCTTGCCCCACAC). KCa3.1^{WT} and KCa3.1^{R350A} cDNA were synthesized

by Genscript; the latter contained an alanine substitution at R350. Both KCa3.1^{WT} and KCa3.1^{R350A}

had substitutions at the PAM recognition sites of the gRNAs (Table S3). Each cDNA was

subcloned into the pMIGII. Plat-e cells were co-transfected with the cloned vectors and pCL-eco

(cat. 12371, Addgene) in Opti-MEM (cat. 31985070, Thermo-Fisher) with Trans-IT-293

transfection reagent (cat. MIR 2705, Mirus Bio) and cultured at 37°C for 12-15 hrs prior to

changing the medium and culturing for an additional 24 hrs at 37°C, followed by 24 hrs at 32°C,

5% CO_2. The viral supernatant was harvested and spun at 1800rpm at RT to remove the cell debris.

Viral supernatant was either used immediately or stored at -80°C till further use.

1081

Viral transduction

Naive Cas9-OT-I CD8$^+$ cells were isolated from the spleen and peripheral LNs of Cas9-OT-I mice

using a Magnisort naïve CD8$^+$ T cell isolation kit according to the manufacturer's instructions (cat.

8804-6825-74, Thermo-Fisher). Purified CD8$^+$ T cells were activated *in vitro* for 20 h with plate-

bound anti-CD3ε (5 μg/ml; 2C11, cat. BE0001, Bio-X-Cell) and anti-CD28 (5 μg/ml; 37.51, cat.

BE0015, Bio-X-Cell) antibodies. Viral transduction was performed by spinfection at 900 *g* at

25 °C for 3 h with 10 μg/ml polybrene (Sigma-Aldrich), followed by resting for 3 h at 37 °C in

5% CO_2. Cells were washed and cultured in medium supplemented with mouse rIL-2 (20 U/ml;

cat. 212-12, Peprotech) for 4 days. GFP-Ametrine double-positive cells were sorted using the

Reflection cell sorter (iCyt) or the MoFlo XDP cell sorter and rested for 24 hrs in medium

containing rIL-2 (10U/ml, cat. 212-12 Peprotech) before activation or adoptive transfer into the

animals.

Flow cytometry

Antibodies from BioLegend included Pacific Blue anti-mouse Ly108 (330-AJ, 134608), BV510

anti-mouse KLRG1 (2F1, 138421), BV570 anti-mouse CD62L (MEL-14, 104433), BV711 anti-

mouse Tim-3 (RMT3-23, 119727), BV785 anti-mouse CD127 (A7R34, 135037), PE-Cy5 anti-

mouse Granzyme-B (QA16A02, 372226), PE-Fire 700, anti-mouse CD4 (GK1.5, 100484),

APC/Cy7 anti-mouse TNFα (MP6-XT22, 506344), PE-Cy7 anti-mouse KLRG1 (2F1/KLRG1,

138416), PerCP-Cy 5.5 anti-mouse CD62L (MEL-14, 104432), BV605 anti-mouse CD127

(A7R34, 135025), Pacific blue anti-mouse CD69 (H1.2F3, 104524), Pacific blue anti-mouse

CD45.1 (110722), APC anti-mouse PD1 (RL388, 109111), BV711 anti-mouse PD1 (29F.1A12,

135231), PE anti-mouse IFNγ (XMG1.2, 505808), FITC anti-mouse TNFα (MP6-XT22, 506304),

Pacific Blue anti-human CD45 (HI30, 982306), APC/Cy7 anti-human CD8 (RPA-T8, 344713).

BUV563 anti-mouse LAG3 (C9B7W, 741350), BUV615 anti-mouse CD69 (H1.2F3, 751593),

BUV805 anti-mouse CXCR3 (173, 748700), BV421 anti-mouse EOMES (X4-83, 567166),

BV480 anti-mouse CD45.1 (A20, 746666), Alexa-Fluor 488 anti-mouse TCF1 (S33-966, 567018),

Alexa-Fluor 647 TOX (NAN448B, 568356), BUV 496 anti-mouse Ly-108 (13G3, 750046),

APC/Cy7 anti-mouse CD44 (IM7, 560568) and BUV805 anti-mouse CD8α (53-6.7, 612898) were

acquired from BD Biosciences. BUV395 anti-mouse CD44 (IM7, 363-0441-82), PerCP-Cy 5.5

anti-mouse IL-2 (JES6-5H4, 45-7021-82), PerCP-eF710 anti-mouse CD27 (O323, 46-0279-42),

and PE-Cy7 anti-mouse Tim-3 (RMT3-23, 12-5870-82, eBioscience) were obtained from Thermo

Fisher. APC anti-human/mouse Tox (REA473, 130-118-335) was obtained from Miltenyi Biotech.

For surface markers, the cells were stained for 30 min on ice in PBS containing 2% FCS and 0.1%

sodium azide (cat. S2002, Sigma-Aldrich). To examine intracellular cytokine production, the

isolated cells were stimulated with a cell activation cocktail (cat. 5476, TOCRIS) in the presence

of monensin (cat. 554724, BD Biosciences) for 4 hr, and stained for intracellular cytokines and

transcription factors using a fixation/permeabilization kit (cat. 00-5523-00, Ebiosciences). Fixable

viability dye (Thermo Fisher) was used to exclude dead cells. Samples were acquired on a Cytek

Aurora (SpectroFlo) or BD LSRII (FACSDiva) flow cytometer and analysed using FlowJo

software.

***In vivo* tumor analysis**

For all tumor experiments, 0.5×10^6 tumor cells were injected subcutaneously into the right flank

of naive mice (aged 6–10 weeks). Met was diluted in HBSS to a concentration of 61 μM per 50μl.

Mice were injected peri-tumorally for 5 days with either 50 μl Met or HBSS. Tumours were

measured with calipers every 2 days and tumor volumes were calculated using the formula:

(width^2 × length)/2. For adoptive transfer experiments, T cells (1×10^6) were transferred i.v. 7 -9

days after tumor injection. For contralateral tumor model, 0.5×10^6 B16 tumor cells were injected

subcutaneously into the right flank and left shoulder of the same mouse. Met or HBSS was injected

peri-tumorally to the right flank tumor (abdominal). For check-point blockade experiment,

0.5×10^6 MC38 tumor cells were injected subcutaneously into the right flank of WT mice. The

mice were given either anti-PD1 antibody (RMP1-14, Bio X Cell) or rat IgG2a (2A3, Bio X Cell),

intraperitoneally twice at a dose of 150 mg/kg in 100 μl PBS at day 9 and 12. Met or HBSS was

given peri-tumorally for 5 days from day 9 to 13. For the CAR-T tumor model, 0.5×10^6 F420

tumor cells (derived from singly floxed *p53+/F-Col2.3* transgenic mice[2]); were injected

subcutaneously into the right flank of *Rag1*[-/-] mice. 5×10^6 B7-H3 CAR-T cells (see below) were

transferred i.v. 8-10 days post implantation. Tumour size was measured every 2-4 days.

Generation of mouse CAR-T cells

Generation of mouse B7-H3-CAR constructs and CAR-T generation have been previously

described[3]. Briefly, HEK293T cells were transiently transfected with the CAR-encoding plasmid,

the Peg-Pam plasmid encoding MoMLV gag-pol, and a plasmid encoding the VSVG envelope.

Viral supernatants were collected at 48 h and filtered with a 0.45m filter. The virus was then used

to transduce the GPE86 producer cell line. B7-H3-CAR-expressing GPE86 cells were stained with

Alexa Fluor-647 anti-human IgG, F(ab′)2 fragment antibody (cat. 109-606-006, Jackson

ImmunoResearch) and sorted using the BD FACSAria III system.

Naive $CD8^+$T cells from 6–8 weeks old $CD45.1^+$ mice were activated with plate-bound anti-CD3ε

(1 μg/ml, 145-2C11, cat. BE0001, BioXcell), anti-CD28 (2 μg/ml, 37.51, cat. BE0015, BioXcell)

and 50 U ml^{-1} rhIL-2 (cat. 200-02, Peprotech). On day 2 post activation, activated $CD8^+$ T cells

were transduced with retrovirus expressing B7-H3-CAR on retronectin-coated (cat. T100B,

Takara) non-tissue-culture-treated plate in complete RPMI medium supplemented with

50 U/ml rhIL-2. At 48 h after transduction, CAR-T cells were collected and expanded in the

presence of 50 U/ml rhIL-2 for another 3 days before adoptive transfer into tumor-bearing mice.

Tumour-infiltrating lymphocyte isolation

B16 tumors were harvested at the indicated time points and digested using a digestion cocktail

(0.1% Collagenase IV and 0.01% DNaseI) for 5 min at 37 °C. Digested tumors were passed

through a 70m strainer and resuspended in PBS plus 2% dialyzed FCS and processed for analysis

and/or sorting by flow cytometry, as described above.

Patient samples were collected following surgery and were immediately processed. Tumour

samples were dissected into small pieces and digested at 37°C with digestion cocktail for 40 min.

Digested tumor samples were physically disrupted and passed through 70μ filter. Samples were

washed with PBS-2% dialyzed FCS twice and stained for CD45, CD3ε and CD8 and sorted for

CD8$^+$ T cells by flow cytometry. Purified T cells were washed once with 1% saline and flash

frozen in liquid nitrogen. Samples were collected upon informed consent and procedures approved

by the IRB of St. Jude Children's Research Hospital.

Immunofluorescence staining and imaging.

CD8$^+$ T cells were immobilized in 8-well poly L-lysine-coated IBIDI chambers and stimulated by

the addition of CD3/28 Dynabeads (cat. 11452D, Thermo Fisher) for 30 min, followed by 10 min

of fixation with 4% paraformaldehyde (PFA) (cat. 15710, Electron Microscopy Science) at 37°C.

The cells were washed once with TBS (50 mM Tris, 100 mM NaCl, pH 8.0) and permeabilized

with permeabilization buffer (50mM Tris, 100 mM NaCl, pH 8.0, 0.3% (v/v) Triton X-100) for

20 min at room temperature. Cells were washed again with TBS before blocking with

TBS containing 2% BSA (Cat. 001-000-182, Jackson ImmunoResearch) for 60 min at room

temperature. The cells were stained overnight at 4 °C with the following primary antibodies: anti-

NFAT1 (1:200, cat. 4389, Cell Signaling Technology), anti-NFAT2 (1:200, cat. D15F1, Cell

Signaling Technology) and anti-mono and dimethyl arginine (1:500, cat. ab412, Abcam). The

samples were washed twice with TBS and incubated for 1 h at room temperature with the following

antibodies: anti-rabbit Alexa Fluor plus 595 (1:1,000, cat. A-11012, Thermo Fisher), anti-mouse

CD8-APC (1:500; cat. 100712, BioLegend), Alexa Fluor 488 Phalloidin (1:1000, cat. A12379,

Thermo Fisher) and Hoechst 33258 (1:1000, cat. H3569, Thermo-Fisher). Samples were then

imaged by Marianas spinning-disc confocal microscopy (Intelligent Imaging Innovations), using

the Prime95B sCMOS camera, and the 405, 488, 561 and 640 nm laser lines. Sum fluorescence

intensities were analyzed using Slide Book 6 (Intelligent Imaging Innovations).

Immunoblotting

T cells were washed once with ice-cold PBS and immediately lysed with 4x laemmli buffer (Bio-

Rad, cat. 1610747). Cell lysates were separated by SDS-PAGE and blotted with anti-Kcnn4

antibody (cat. PA5-33875, Thermo Fisher) and HRP-conjugated anti-Actin (cat. sc4777-HRP,

Santa Cruz Biotechnology). Images were developed using Clarity Western ECL substrate (cat.

1705061, Bio-Rad) and acquired on Bio-Rad Chemi-Doc.

1189

Amino acid measurement

Subcutaneous tumors were isolated and processed as described above. Sorted T cells were washed

once with ice-cold saline, flash-frozen with liquid nitrogen, and stored at –80ºC before analysis.

Cell pellets were extracted using 750 µl of methanol/acetonitrile/water (5:3:2, v/v/v) and the

supernatant was dried by lyophilization. Aliquots of 20-50 µl from plasma and TIF were extracted

with at least a 15-fold excess volume of the methanol/acetonitrile/water solution, and the

supernatant was then collected and dried by lyophilization. Dried extracts containing the

hydrophilic metabolites were dissolved in 30 µl of water/acetonitrile (8:2, v/v), and 10 µl was used

to derivatize amino acids as described [4] with some modifications. The samples were placed in glass

autosampler vials, and 35 µl of sodium borate buffer (100 mM, pH 9.0) was added and mixed by

pipetting. 10 µl of the 6-aminoquinolyl-N-hydroxysuccinimidyl carbamate (AQC, 10 mM in

acetonitrile)-derivatizing reagent (Cayman Chemical) was added, the vial was sealed, mixed by

vortexing, and then incubated at 55°C for 15 min. The vial was cooled to room temperature and

then 1 µL was analyzed by liquid chromatography with tandem mass spectrometry (LC-MS/MS).

An ACQUITY Premier UPLC System (Waters Corp) was used for the LC separations, using a

non-linear gradient condition as follows: 0-0.4 min 3% B; 0.4-8 min 3 to 96% B (using Curve #8

of the inlet condition in MassLynx™); 8-12 min 96% B; 12-12.5 min 96 to 3% B; 12.5-14 min 3%

B. Mobile phase A was water supplemented with 0.15% acetic acid, and mobile phase B was

acetonitrile with 0.15% acetic acid. The column used was an Accucore C30 (50 × 2.1 mm, 2.6 µm)

(Thermo Fisher Scientific) operated at 50°C. The flow rate was 300 µl/min and the injection

volume used was 1 µl. All LC/MS solvents and reagents were the highest purity available (water,

acetonitrile, acetic acid, boric acid, sodium hydroxide) and were purchased from Thermo Fisher

Scientific. A Xevo TQ-XS Triple Quadrupole Mass Spectrometry (TQ-XS) (Waters Corp)

equipped with a multi-mode ESI/APCI/ESCi ion source was employed as detector. The TQ-XS

was operated in positive ion mode using a multiple reaction monitoring mass spec method (MRM).

The MRM conditions were set to a minimum of 15 points per peak with an automatic dwell time.

The operating conditions of the source were: Capillary Voltage 3.8 kV, Cone Voltage 40 V,

Desolvation Temp 550°C, Desolvation Gas Flow 1000 L/h, Cone Gas Flow 150 L/h, Nebuliser

7.0 Bar, Source Temp 150°C. Authentic amino acid standards were purchased from Sigma-Aldrich

and used to establish MRM conditions and calibration curves. The monitored parent/daughter ions,

fragmentation collision energy (CE), and retention time window for each amino acid are listed in

Supplementary Table 4. The acquired MRM data was processed using the software application

Skyline 21.2 (MacCoss Lab Software).

1223

***In-silico* model building of KCa3.1(SK4)-Calmodulin complex and simulation.**

To study the impact of SK4s arginine dimethylation on the binding of calmodulin, we simulated

the SK4 channel:Calmodulin complex in a membrane layer via all-atom molecular dynamics

(MD), which is considered a computational microscope[5]. The Cryo-EM structure of the human

SK4 channel:Calmodulin complex state (PDB:6CNN[6]) was used as the starting structure. Thisa

tetramer compleconsistsng of four SK4 monomers, four calmodulins (CAMs), three calcium ions

bound to each calmodulin (CAM) (total 1,2) and four K^+ ions bound to the channel selectivity

filter.

Initially, protein was prepared was using the molecular operating environment (MOE) software.

The missing residues in the cryo-EM structure were modeled based on the human AlphaFold

structure (AF-015554-F1). Protonate3D module in MOE was used to assign appropriate

protonation states at pH 7.4 and then energy minimized. Further, simulation system was prepared

using CHARMM-GUI web server[7-9]. Protein was placed in the POPE membrane bilayer and

protein orientation was obtained using the PPM web server[10]. The protein-membrane bilayer

complex was immersed in the TIP3P[11] water box of size $\approx$150 X 150 X 230 $Å^3$ with an edge

distance of 40 Å on all sides. KCl salt was added to balance the charges and maintain a

physiological salt concentration of 0.15M. Overall, each wildtype simulation system contains four

SK4 monomers, four CAMs, 12 Ca^{2+} ions, $\approx$355 K^+ ions, $\approx$367 Cl^- ions, $\approx$618 POPE lipids, and

$\approx$128,144 TIP3P waters. The total numbers of atoms in this system are $\approx$498,944. Two more

systems were generated via a protocol similar to that explained above, except that in one system,

R352 of the SK4 monomer was symmetrically dimethylated and in the second system, R352 was

asymmetrically dimethylated. Dimethylation was introduced using the CHARMM-GUI web

server. Note that arginine's at position 352 in all four SK4 monomers were dimethylated in both

systems.

Further, all three systems were simulated using AMBER20/22 MD software[12]. The MD simulation

was performed in three steps. In step-1, energy minimization was carried out to remove potential

steric clashes in two stages. In stage 1, each system was energy-minimized for 5000 steps, while

positional restraints were applied to the protein and membrane: the first 2500 steps with the

steepest-descent algorithm and the remaining 2500 steps using the conjugate gradient algorithm[13].

In stage 2, the second round of minimization was carried out with all restraints removed in 50,000

steps. The steepest-descent algorithm[14] was used for the first 25,000 steps, while the conjugate

gradient algorithm was used for the remaining 25,000 steps. In step-2, equilibration was carried in

six stages: (1) equilibrated for 125 ps with a time step of 1fs while applying positional restraints

on protein (K=10 kcal/mol.Å^2) and membrane (K=2.5 kcal/mol.Å^2), (2) equilibrated for 125 ps

with a time step of 1fs w while applying positional restraints on the protein (K=5.0 kcal/mol.Å^2)

and membrane (K=2.5 kcal/mol.Å^2), (3) equilibrated for 125 ps with a time step of 1fs while

applying positional restraints on the protein (K=2.5 kcal/mol.Å^2) and membrane (K=1.0

kcal/mol.Å^2), (4) equilibrated for 500 ps with a time step of 2 fs while applying positional restraints

on the protein (K=1.0 kcal/mol.Å^2) and membrane (K=0.5 kcal/mol.Å^2), (5) equilibrated for 500

ps with a time step of 2 fs while applying positional restraints on the protein (K=0.5 kcal/mol.Å^2)

and membrane (K=0.1 kcal/mol.Å^2) and (6) equilibrated for 500 ps with a time step of 2 fs while

applying weak positional restraints on the protein (K=0.5 kcal/mol.Å^2) and restraints on the rest

of the system were completely removed. The first two equilibration simulations were performed

under NVT conditions, whereas the last four equilibration simulations were performed under NPT

conditions. In step-3, production simulations were carried out under periodic boundary conditions

using the NPT ensemble, during which the temperature (310 K) and pressure (1 atm) were kept

constant. Three replicates of each system were simulated, and each replicate was simulated for ≈1

μs. Langevin integrator and Monte Carlo barostat were used[15,16] to conduct NPT simulations,

whereas the SHAKE algorithm was used to restrain bonds with hydrogens[17]. Particle-mesh Ewald

(PME) method was used to treat long-ranged electrostatic interactions[18] and the nonbonded cutoff

was 12Å (force switching = 10Å). CHARM36 all-atom additive force field[19,20] was used to treat

the entire system. Overall, nine MD simulations were conducted across three systems, with each

system comprising three replicates that each ran for approximately 1μs. Consequently, the total

MD simulation data accumulated for analysis amounts to approx. 9 µs, derived from the sum of

simulation times for all systems and replicates (3 systems X 3 replicates X ≈1µs/per replicate =

≈9µs). Two frames per nanosecond were used for the data analysis. Visual molecular dynamics

(VMD)[21] was used for visualization and graphic generation, whereas Tcl was used for data

analysis. Gnuplot was used to generate the time series plots.

1282

ATAC-seq

OT-I CD8$^+$ T cells were activated with 2.5ng/ml SIINFEKL in either 0.1 mM Met or 0.03 mM

Met with restoration of Met to 0.1 mM in 0.03 mM condition post 30 min of activation and

incubated for 24 hrs. KCa3.1^{WT} and KCa3.1^{R350A} cells were activated by 2.5ng/ml SIINFEKL for

24 hrs. Cells were processed for ATAC-Seq after 24 hrs of activation. Briefly, 50,000 cells were

washed with 1 ml cold PBS by centrifugation at 500 g for 5 min at 4°C. Cells were then

resuspended in 50 ml cold cell lysis buffer (10 mM Tris pH 7.4, 10 mM NaCl, 3 mM MgCl$_2$, and

0.1% NP-40). Isolated nuclei were pelleted at 1,000 × g for 10 min at 4°C, resuspended in 50 µl

transposition reaction mixture (25 µl TD buffer, 2.5 µl TDE1, 22.5 µl ddH$_2$O) (20034197,

Illumina), and incubated at 42°C for 45 min. DNA was purified using a MiniElute PCR purification

kit (cat. 28004, Qiagen). Samples were then processed for library barcoding and amplification with

Q5 High-Fidelity 2× Master Mix (cat. M0492S, NEB). Prepared libraries were sent for sequencing

after quantification using Qubit and size distribution as determined using an Agilent 4200

TapeStation.

Analysis of ATAC-seq data

Paired-end sequencing reads were trimmed using Trim Galore (version 0.5.0)

(https://github.com/FelixKrueger/TrimGalore) with default parameters. Reads were aligned to the

reference mouse mm10 assembly using Bowtie 2 (version 2.3.5.1)[22] with settings --end-to-end --

very-sensitive -X 2000. The resulting alignments, recorded in a BAM file, were sorted, indexed, and

marked for duplicates with Picard MarkDuplicates function (version 2.19.0). Afterward, the BAM

file was filtered with SAMtools (version 1.9)[23], BamTools (version 2.5.1)[24] and scripts of nf-

core/chipseq[25] to discard reads, mates that were unmapped, PCR/optical duplicates, not primary

alignments, mapped to multiple locations, or mapped to ENCODE blacklisted regions[26]; only reads

mapped in proper pair were kept (-F 1804 -f 2 -q 30). The alignments were shifted by deepTools

alignmentSieve[27] with --ATACshift and default parameters. Nucleosome-free reads (fragment

length < 109 bp) were separated from the BAM files with SAMtools (version 1.9). MACS (version

2.1.2)[28] was used to call peaks from the BAM files with narrowPeak setting, --extsize 200, and

recommended mappable genome size (default value for other parameters). Chromatin accessibility

signal was normalized by scaling to 1 million mapped reads using BEDTools (version 2.27.1)[29]

and bedGraphToBigWig (version 377)[30] and visualized as heatmaps using deepTools plotHeatmap

(version 3.2.1)[27].

Differentially accessibility analysis was performed using DiffBind[31] (v.3.4.7) (summit = 50,

normalized by trimmed mean of M values[32] (TMM) approach, library size estimated by reads

mapped to "background" regions, the genomic regions (binned into 15000 bp) overlapped with

peaks). Peaks were annotated to the nearest genes with annotatePeak function in R package

ChIPseeker[33] (v.1.30.3) at the gene level using the default parameters. Genes were ranked by fold

change (log2 transformed) of associated peaks located within +/- 1.5 kb of transcription start sites

(if a gene was associated with multiple peaks, the mean was used). Gene Set Enrichment Analysis[34]

(GSEA) was then performed against MSigDB collections with R package clusterProfiler[35]

(v.4.2.2). The common differentially accessible regions (DAR) between ATAC-Seq of KCa3.1WT

and KCa3.1R350A and T cells activated in 0.1 mM or 0.03 mM Met were obtained by the

overlapping of the DARs (at least 1 bp). Enrichment analysis was performed using the over-

representation approach against MSigDB using the R package clusterProfiler.

1326

CUT&RUN

CUT&RUN experiments were performed as previously described[36] with slight modifications.

Purified OT-I CD8$^+$ T cells were activated with 2.5ng/ml SIINFEKL in 0.1 mM and 0.03 mM Met

for 30 min. Met was restored to 0.1 mM post 30 min and cultured for 24 hrs. Cells were washed

with cold PBS and dead cells were removed using Easysep dead cell removal kit (cat. 17899,

StemCell). Cells (2 × 10^5) were washed twice with wash buffer (20 mM HEPES (H3375, Sigma-

Aldrich], 150 mM NaCl (cat. AM9760G, Invitrogen), 0.5 mM spermidine (cat. S0266, Sigma-

Aldrich), and protease inhibitor cocktail (cat. 5056489001, Sigma-Aldrich)), resuspended and

bound to concanavalin-A coated magnetic beads (cat. BP531, Bang Laboratories). The

resuspension buffer was removed using a magnetic stand and beads were resuspended in 100 µl
antibody buffer (20 mM HEPES, 150 mM NaCl, 0.5 mM spermidine, 0.01% digitonin (cat.
300410, Millipore), 2 mM EDTA (cat. AM9260G, Invitrogen), and protease inhibitor cocktail).
Primary antibodies (1:100) were added to the samples and incubated overnight at 4 °C. Next,
samples were washed twice with cold Dig-wash buffer (20 mM HEPES, 150 mM NaCl, 0.5 mM
spermidine, 0.01% digitonin, and protease inhibitor) and pAG-MNase (cat. 123461, Addgene) was
added and rotated at 4 °C for 1 h. Samples were washed twice and resuspended in 50 µl Dig-wash
buffer and $CaCl_2$ (2 µl of 100 mM, cat. 2115, Sigma-Aldrich) was added followed by brief
vortexing and incubation on ice for 30 min. Next, 50 µl of 2× STOP buffer (340 mM NaCl, 20 mM
EDTA, 4 mM EGTA (cat. E3889, Sigma-Aldrich) and 100 µg/ml RNase A (cat. EN0531, Thermo
Fisher), and 50 µg/ml GlycoBlue (cat. AM9515, Invitrogen)) was added and mixed by gentle
vortexing and incubated for 30 min at 37 °C to release CUT&RUN fragments. Fragmented DNA
was purified using the NEB Monarch PCR&DNA Cleanup Kit (cat. T1030S, New England
Biolabs). DNA libraries were prepared using the NEBNext Ultra II DNA Library Prep Kit (cat.
E7645S, New England Biolabs) and purified with AMPure SPRI beads (cat. B23318, Beckman-
Coulter). Prepared libraries were quantified using Qubit and size distribution was determined with
a Agilent 4200 TapeStation analysis before paired-end sequencing.

CUT&RUN data processing

Paired-end sequencing reads were trimmed using Trim Galore (v.0.5.0)
(https://github.com/FelixKrueger/TrimGalore) with default parameters. The reads were then
aligned to the reference mouse mm10 assembly using Bowtie 2 (v.2.3.5.1)[22] with settings --end-
to-end --very-sensitive --no-mixed --no-discordant -q --phred33 -I 10 -X 700. The resulting
alignments, recorded in the BAM file, were sorted, indexed, and marked for duplicates using the
Picard MarkDuplicates function (v.2.19.0) (Picard toolkit. *Broad Institute GitHub Repository,*
2019). Next, the BAM file was filtered with SAMtools (v.1.9)[23], BamTools (v.2.5.1)[24] and scripts
of nf-core/chipseq[25] to discard reads, including mates that were unmapped, PCR/optical duplicates,
non-primary alignments, reads that mapped to multiple locations or to ENCODE-blacklisted
regions[26], or with more than 4 mismatches (-F 0x004 -F 0x008 -F 0x0100 -F 0x0400 -f 0x001 -q
1). MACS (v.2.1.2)[28] was used to call peaks from the BAM file with IgG control and recommended
mappable genome size (the default values were used for the other parameters). NarrowPeak mode

was used for NFAT1, H3K27me3, and H3K4me3. Binding signal was normalized by scaling to 1

million mapped reads using BEDTools (version 2.27.1)[29] and bedGraphToBigWig (version 377)[30]

and visualized as heatmaps using deepTools plotHeatmap (version 3.2.1)[27]

In the CUT&RUN experiment with spike-in E. coli, two modifications were made: i) a hybrid

reference of mouse mm10 and E. coli ASM584v2, and ii) signals were normalized by scaling to

per million reads mapped to E. coli.

Differential binding analysis was performed using DiffBind (v.3.4.7) (summit = 75 (for NFAT1)

and 100 (for H3K27me3 and K3K4me3 modifications), normalized by the trimmed mean of M

values (TMM) approach library size estimated by reads mapped to "background" regions. Peaks

were annotated to the nearest genes with the annotatePeak function in R package ChIPseeker

(v.1.30.3) at the gene level using default parameters. Genes were ranked by fold-change (log2

transformed; if a gene was associated with multiple peaks, the mean was used). GSEA was then

performed against MSigDB collections with R package clusterProfiler (v.4.2.2).

1379

RNA-seq

The isolated CD8$^+$ TIL were washed once with cold PBS and pelleted at 1500 rpm at 4°C. Total

RNA was isolated using the Direct-zol RNA Microprep Kit (cat. R2061, Zymo Research)

according to the manufacturer's instructions and quantified using an Agilent 4200 TapeStation.

Libraries were prepared using the KAPA RNA HyperPrep Kit with RiboErase (HMR) (cat.

08098131702, Roche) and purified by AMPure SPRI beads (cat. B23318, Beckman-Coulter).

Libraries were quantified and size distribution was determined using a Agilent 4200 TapeStation

before paired-end sequencing was performed.

RNA-seq data processing

Paired-end sequencing reads were mapped by the pipeline of St. Jude Center for Applied

Bioinformatics. The reads were trimmed using Trim Galore (version 0.5.0)

(https://github.com/FelixKrueger/TrimGalore) with default parameters. Reads were aligned to the

reference mouse mm10 assembly plus ERCC spike-in sequences using STAR (version 2.7.5a)[37].

The resulting alignments, recorded in BAM file, were sorted, indexed, and marked for duplicates

with Picard MarkDuplicates function (version 2.19.0). Transcript quantification was calculated

using RSEM[38]. Differential gene expression analysis was carried out with DESeq2 with Wald test

(default parameters)[39]. To examine if global changes in gene expression were present, the RUVg

function in the RUVSeq package was used for normalization with ERCC spike-in followed by

DESeq2 according to the RUVSeq manual[40]. Gene set enrichment analysis (GSEA) was performed

against the MSigDB database[41] with R package clusterProfiler (version 4.2.2)[35] with the genes

ranked by the Wald statistic from DESeq2 analysis.

1401

Quantitative Proteomics by tandem mass tag mass spectrometry (TMT-MS)

$CD8^+$ T cells were isolated from WT mice and activated with anti-CD3 antibody by crosslinking

with IgG for 30 min. Activated and untreated cells were washed once with ice-cold PBS and flash-

frozen in liquid nitrogen. Proteomic profiling of the whole proteome and methylome was carried

out using a previously optimized protocol[42] with modifications. Briefly, the cells were lysed in 8M

Urea lysis buffer using pulse sonication, and approximately 100 µg of protein per sample was

digested with Lys-C (Wako, 1:100 w/w) at 21 °C for 2 h, followed by dilution to reduce urea to

2M, and further digestion with trypsin (Promega, 1:50 w/w) at 21 °C overnight. The protein digests

were acidified (trifluoroacetic acid to 1%), centrifuged at 21,000 x g for 10 min at 4°C to remove

any insoluble material, desalted with Sep-Pak C18 cartridge (Waters), and dried by Speedvac.

Each sample was resuspended in 50 mM HEPES (pH 8.5), TMT labeled, mixed equally, desalted,

and fractionated by offline HPLC (Agilent 1220) using basic-pH reverse phase LC (Waters

XBridge C18 column, 3.5 µm particle size, 4.6 mm x 25 cm, 180 min gradient, 80 fractions).

Fractions were aliquoted (5% v/v) for whole proteome and methylome (95% v/v) profiling. The

methylome fractions were further concatenated into 10 fractions for antibody-based sequential

mono-methyl-arginine and di-methyl-arginine peptide enrichment.

1418

Antibody-based Sequential Methylome Enrichment

Each concatenated fraction (approximately 1 mg) was resuspended in 400 µL ice-cold IAP buffer

(50mM MOPS, pH 7.2, 10 mM sodium phosphate, and 50 mM NaCl) and centrifuged at 21,000 ×

g for 10 min at 4°C to remove any insoluble material. The peptides were then incubated with

mono-methyl-Arginine antibody beads (mono-me-R, 12235S Cell Signaling Technology) at an

antibody-to-peptide ratio of 1:20 (w/w, optimized through a pilot experiment) for 2 hr at 4°C with

gentle end-over-end rotation. The antibody beads were then collected by a brief centrifugation,

washed 3 times with 1 mL ice-cold IAP buffer and twice with 1 mL ice-cold PBS, while the

supernatants were carefully removed and used for the sequential di-methyl-Arginine peptide

enrichment (di-me-R symmetric, cat.13563S Cell Signaling Technology). Peptides were eluted

from the beads twice at room temperature with 50 µL of 0.15% TFA, dried, and analyzed by LC-

MS/MS.

1431

LC-MS/MS Analysis

Each sample was dried, reconstituted, and analyzed by LC-MS/MS (a CoAnn 75 µm x 30 cm

column, packed with 1.9 µm C18 resin from Dr. Maisch GmbH) interfaced with the Orbitrap

Fusion MS (Thermo Fisher). LC settings included an 80 min gradient of 15%-40% buffer B (70%

ACN, 2.5% DMSO, and 0.1% FA) with buffer A (2.5% DMSO, and 0.1% FA) at a flow rate of

~0.25 µL /min flow rate. MS settings included data-dependent (3 s cycle) mode with a survey scan

in the Orbitrap (60,000 resolution, scan range 410–1600 m/z, 1×106 AGC target, and 50 ms

maximal ion time), followed by sequential isolation of abundant ions in a 3 s duty cycle, with

fragmentation by higher-energy collisional dissociation (HCD, 38 normalized collision energy),

and high-resolution detection of MS/MS ions in the Orbitrap (60,000 resolution, 1×105 AGC

target, 105 ms maximal ion time,1.0 m/z isolation window, and 20 s dynamic exclusion).

1443

Database Search and TMT Quantification

The MS/MS spectra were searched against the UniProt Mouse protein database (version

2020.04.22) using the COMET algorithm (v2018.013[43]) with the JUMP software suite[44]. Search

parameters included MS1 mass tolerance of 20 ppm and MS/MS of 0.02 Da, fully tryptic, static

mass shift for the TMT16 tags (+304.2071453) and carbamidomethyl modification of 57.02146

on cysteine, dynamic mass shift for Met oxidation (+15.99491), for mono-methyl-Arginine

(+14.01565), and for di-methyl-Arginine (+28.0313), maximal missed cleavage (n = 3), and

maximal dynamic modifications per peptide (n=5). All matched MS/MS spectra were filtered by

mass accuracy and matching scores to reduce false discovery rate (FDR) below 1% for either

proteins (whole proteome analysis) or peptides (methylome analysis), based on the target-decoy

strategy[45,46].

1455

TMT quantification analysis was performed as previously reported[47] with the following

modifications: (i) extracting reporter ion intensities from each PSM; (ii) correcting the intensities

according to the isotopic distribution of each TMT reagent; (iii) removing PSMs of very low

intensities (e.g., minimum value of 1,000 and median value of 5,000); (iv) normalizing sample

loading bias with the trimmed median intensity of all PSMs; (v) calculating the mean-centered

intensities across samples (e.g., relative intensities between each sample and the mean), (vi)

summarizing protein relative intensities by averaging related PSMs; (vii) finally deriving protein

absolute intensities by multiplying the relative intensities by the grand-mean of the three most

highly abundant PSMs. (vii) Protein fold change and p-values of different comparisons were

calculated based on the protein intensities, using a log transformation and moderated t-test with

the limma R package[48,49].

1467

Quantification and statistical analysis

Data were plotted and analysed using GraphPad Prism (GraphPad Software, v.9.2.0). Statistical

significance was calculated using unpaired one- or two-tailed Student's *t*-tests. Two-way ANOVA

was performed to compare tumor growth curves. The log-rank (Mantel–Cox) test was performed

to compare the mouse survival curves. For Fig. 7b, a linear mixed effects model with $\log_{10}(y)$ as

the response variable, group as a fixed-effect predictor variable, and mouse as a random-effect

predictor variable was fit to the data. This model represents data with cells as random samples

from each mouse. The results showed that the mean $\log_{10}(y)$ values of the two mouse groups

differed by -0.131 units (95% CI: -0.163, -0.0999; $p = 6.98 \cdot 10^{-15}$). Statistical significance was set

at $P < 0.05$. Data are presented as mean ± SEM. No statistical methods were used to determine

sample size.

References

1480
1   Welsh, R. M. & Seedhom, M. O. Lymphocytic choriomeningitis virus (LCMV):
propagation, quantitation, and storage. *Curr Protoc Microbiol* **Chapter 15**, Unit 15A
11 (2008). https://doi.org:10.1002/9780471729259.mc15a01s8

Zhao, S. *et al.* NKD2, a negative regulator of Wnt signaling, suppresses tumor growth and metastasis in osteosarcoma. *Oncogene* **34**, 5069-5079 (2015). https://doi.org/10.1038/onc.2014.429

Haydar, D. *et al.* Cell-surface antigen profiling of pediatric brain tumors: B7-H3 is consistently expressed and can be targeted via local or systemic CAR T-cell delivery. *Neuro Oncol* **23**, 999-1011 (2021). https://doi.org/10.1093/neuonc/noaa278

Sharma, G. *et al.* Analysis of 26 amino acids in human plasma by HPLC using AQC as derivatizing agent and its application in metabolic laboratory. *Amino Acids* **46**, 1253-1263 (2014). https://doi.org/10.1007/s00726-014-1682-6

Meszaros, B. *et al.* Recent breakthroughs in computational structural biology harnessing the power of sequences and structures. *Curr Opin Struct Biol* **80**, 102608 (2023). https://doi.org/10.1016/j.sbi.2023.102608

Lee, C. H. & MacKinnon, R. Activation mechanism of a human SK-calmodulin channel complex elucidated by cryo-EM structures. *Science* **360**, 508-513 (2018). https://doi.org/10.1126/science.aas9466

Jo, S., Kim, T., Iyer, V. G. & Im, W. CHARMM-GUI: a web-based graphical user interface for CHARMM. *J Comput Chem* **29**, 1859-1865 (2008). https://doi.org/10.1002/jcc.20945

Lee, J. *et al.* CHARMM-GUI Input Generator for NAMD, GROMACS, AMBER, OpenMM, and CHARMM/OpenMM Simulations Using the CHARMM36 Additive Force Field. *J Chem Theory Comput* **12**, 405-413 (2016). https://doi.org/10.1021/acs.jctc.5b00935

Wu, E. L. *et al.* CHARMM-GUI Membrane Builder toward realistic biological membrane simulations. *J Comput Chem* **35**, 1997-2004 (2014). https://doi.org/10.1002/jcc.23702

Lomize, M. A., Pogozheva, I. D., Joo, H., Mosberg, H. I. & Lomize, A. L. OPM database and PPM web server: resources for positioning of proteins in membranes. *Nucleic Acids Res* **40**, D370-376 (2012). https://doi.org/10.1093/nar/gkr703

Jorgensen, W. L., Chandrasekhar, J., Madura, J. D., Impey, R. W. & Klein, M. L. Comparison of simple potential functions for simulating liquid water. *The Journal of Chemical Physics* **79**, 926-935 (1983). https://doi.org/10.1063/1.445869

Case, D. A. *et al.* AmberTools. *J Chem Inf Model* **63**, 6183-6191 (2023). https://doi.org/10.1021/acs.jcim.3c01153

B. N, P. *Mathematics of Computation* **28**, 669-671 (1974). https://doi.org/10.2307/2005951

in *Oeuvres complètes: Series 1* Vol. 10 *Cambridge Library Collection - Mathematics* (ed Augustin-Louis Cauchy) 399-402 (Cambridge University Press, 2009).

Feller, S. E., Zhang, Y., Pastor, R. W. & Brooks, B. R. Constant pressure molecular dynamics simulation: The Langevin piston method. *The Journal of Chemical Physics* **103**, 4613-4621 (1995). https://doi.org/10.1063/1.470648

Åqvist, J., Wennerström, P., Nervall, M., Bjelic, S. & Brandsdal, B. O. Molecular dynamics simulations of water and biomolecules with a Monte Carlo constant

pressure algorithm. *Chemical Physics Letters* **384**, 288-294 (2004). https://doi.org:https://doi.org/10.1016/j.cplett.2003.12.039

Ryckaert, J.-P., Ciccotti, G. & Berendsen, H. J. C. Numerical integration of the cartesian equations of motion of a system with constraints: molecular dynamics of n-alkanes. *Journal of Computational Physics* **23**, 327-341 (1977). https://doi.org:https://doi.org/10.1016/0021-9991(77)90098-5

Darden, T., York, D. & Pedersen, L. Particle mesh Ewald: An N·log(N) method for Ewald sums in large systems. *The Journal of Chemical Physics* **98**, 10089-10092 (1993). https://doi.org:10.1063/1.464397

Huang, J. *et al.* CHARMM36m: an improved force field for folded and intrinsically disordered proteins. *Nat Methods* **14**, 71-73 (2017). https://doi.org:10.1038/nmeth.4067

Best, R. B. *et al.* Optimization of the additive CHARMM all-atom protein force field targeting improved sampling of the backbone phi, psi and side-chain chi(1) and chi(2) dihedral angles. *J Chem Theory Comput* **8**, 3257-3273 (2012). https://doi.org:10.1021/ct300400x

Humphrey, W., Dalke, A. & Schulten, K. VMD: visual molecular dynamics. *J Mol Graph* **14**, 33-38, 27-38 (1996). https://doi.org:10.1016/0263-7855(96)00018-5

Langmead, B. & Salzberg, S. L. Fast gapped-read alignment with Bowtie 2. *Nat Methods* **9**, 357-359 (2012). https://doi.org:10.1038/nmeth.1923

Li, H. *et al.* The Sequence Alignment/Map format and SAMtools. *Bioinformatics* **25**, 2078-2079 (2009). https://doi.org:10.1093/bioinformatics/btp352

Barnett, D. W., Garrison, E. K., Quinlan, A. R., Stromberg, M. P. & Marth, G. T. BamTools: a C++ API and toolkit for analyzing and managing BAM files. *Bioinformatics* **27**, 1691-1692 (2011). https://doi.org:10.1093/bioinformatics/btr174

Ewels, P. A. *et al.* The nf-core framework for community-curated bioinformatics pipelines. *Nat Biotechnol* **38**, 276-278 (2020). https://doi.org:10.1038/s41587-020-0439-x

Amemiya, H. M., Kundaje, A. & Boyle, A. P. The ENCODE Blacklist: Identification of Problematic Regions of the Genome. *Sci Rep* **9**, 9354 (2019). https://doi.org:10.1038/s41598-019-45839-z

Ramirez, F. *et al.* deepTools2: a next generation web server for deep-sequencing data analysis. *Nucleic Acids Res* **44**, W160-165 (2016). https://doi.org:10.1093/nar/gkw257

Zhang, Y. *et al.* Model-based analysis of ChIP-Seq (MACS). *Genome Biol* **9**, R137 (2008). https://doi.org:10.1186/gb-2008-9-9-r137

Quinlan, A. R. & Hall, I. M. BEDTools: a flexible suite of utilities for comparing genomic features. *Bioinformatics* **26**, 841-842 (2010). https://doi.org:10.1093/bioinformatics/btq033

Kent, W. J., Zweig, A. S., Barber, G., Hinrichs, A. S. & Karolchik, D. BigWig and BigBed: enabling browsing of large distributed datasets. *Bioinformatics* **26**, 2204-2207 (2010). https://doi.org:10.1093/bioinformatics/btq351

Ross-Innes, C. S. *et al.* Differential oestrogen receptor binding is associated with clinical outcome in breast cancer. *Nature* **481**, 389-393 (2012). https://doi.org:10.1038/nature10730

32   Chen, Y., Lun, A. T. & Smyth, G. K. From reads to genes to pathways: differential
expression analysis of RNA-Seq experiments using Rsubread and the edgeR
quasi-likelihood   pipeline.   *F1000Res*   **5**,   1438   (2016).
https://doi.org:10.12688/f1000research.8987.2
33   Yu, G., Wang, L. G. & He, Q. Y. ChIPseeker: an R/Bioconductor package for ChIP
peak annotation, comparison and visualization. *Bioinformatics* **31**, 2382-2383
(2015). https://doi.org:10.1093/bioinformatics/btv145
34   Subramanian, A. *et al.* Gene set enrichment analysis: a knowledge-based
approach for interpreting genome-wide expression profiles. *Proc Natl Acad Sci U*
S A **102**, 15545-15550 (2005). https://doi.org:10.1073/pnas.0506580102
35   Yu, G., Wang, L. G., Han, Y. & He, Q. Y. clusterProfiler: an R package for
comparing biological themes among gene clusters. *OMICS* **16**, 284-287 (2012).
https://doi.org:10.1089/omi.2011.0118
36   Meers, M. P., Bryson, T. D., Henikoff, J. G. & Henikoff, S. Improved CUT&RUN
chromatin profiling tools. *Elife* **8** (2019). https://doi.org:10.7554/eLife.46314
37   Dobin, A. & Gingeras, T. R. Mapping RNA-seq Reads with STAR. *Curr Protoc*
Bioinformatics   **51**,   11   14   11-11   14   19   (2015).
https://doi.org:10.1002/0471250953.bi1114s51
38   Li, B. & Dewey, C. N. RSEM: accurate transcript quantification from RNA-Seq data
with or without a reference genome. *BMC Bioinformatics* **12**, 323 (2011).
https://doi.org:10.1186/1471-2105-12-323
39   Love, M. I., Huber, W. & Anders, S. Moderated estimation of fold change and
dispersion for RNA-seq data with DESeq2. *Genome Biol* **15**, 550 (2014).
https://doi.org:10.1186/s13059-014-0550-8
40   Risso, D., Ngai, J., Speed, T. P. & Dudoit, S. Normalization of RNA-seq data using
factor analysis of control genes or samples. *Nat Biotechnol* **32**, 896-902 (2014).
https://doi.org:10.1038/nbt.2931
41   Liberzon, A. *et al.* The Molecular Signatures Database (MSigDB) hallmark gene
set   collection.   *Cell   Syst*   **1**,   417-425   (2015).
https://doi.org:10.1016/j.cels.2015.12.004
42   Yu, K. *et al.* High-Throughput Profiling of Proteome and Posttranslational
Modifications by 16-Plex TMT Labeling and Mass Spectrometry. *Methods Mol Biol*
2228, 205-224 (2021). https://doi.org:10.1007/978-1-0716-1024-4_15
43   Eng, J. K. *et al.* A deeper look into Comet--implementation and features. *J Am Soc*
Mass Spectrom **26**, 1865-1874 (2015). https://doi.org:10.1007/s13361-015-1179-
x
44   Wang, X. *et al.* JUMP: a tag-based database search tool for peptide identification
with high sensitivity and accuracy. *Mol Cell Proteomics* **13**, 3663-3673 (2014).
https://doi.org:10.1074/mcp.O114.039586
45   Peng, J., Elias, J. E., Thoreen, C. C., Licklider, L. J. & Gygi, S. P. Evaluation of
multidimensional chromatography coupled with tandem mass spectrometry
(LC/LC-MS/MS) for large-scale protein analysis: the yeast proteome. *J Proteome*
Res **2**, 43-50 (2003). https://doi.org:10.1021/pr025556v
46   Elias, J. E. & Gygi, S. P. Target-decoy search strategy for increased confidence in
large-scale protein identifications by mass spectrometry. *Nat Methods* **4**, 207-214
(2007). https://doi.org:10.1038/nmeth1019

47   Niu, M. *et al.* Extensive Peptide Fractionation and y(1) Ion-Based Interference
Detection Method for Enabling Accurate Quantification by Isobaric Labeling and
Mass   Spectrometry.   *Anal   Chem*   **89**,   2956-2963   (2017).
https://doi.org:10.1021/acs.analchem.6b04415
48   Bai, B. *et al.* Deep Multilayer Brain Proteomics Identifies Molecular Networks in
Alzheimer's   Disease   Progression.   *Neuron*   **105**,   975-991   e977   (2020).
https://doi.org:10.1016/j.neuron.2019.12.015
49   Smyth, G. K. Linear models and empirical bayes methods for assessing differential
expression in microarray experiments. *Stat Appl Genet Mol Biol* **3**, Article3 (2004).
https://doi.org:10.2202/1544-6115.1027
1630
1631