Rapid metabolic regulation of a novel arginine methylation of KCa3.1 attenuates T cell exhaustion.

Authors:

Piyush Sharma[1*], Ao Guo[1#], Suresh Poudel[1], Emilio Boada-Romero[1], Katherine C. Verbist[1],

Gustavo Palacios[1], Kalyan Immadisetty[2], Mark J. Chen[1,3], Dalia Haydar[4], Ashutosh Mishra[5],

Junmin Peng[5,6], M. Madan Babu[2], Giedre Krenciute[4], Evan S. Glazer[7], Douglas R. Green[1*].

[1]Department of Immunology, St. Jude Children's Research Hospital, Memphis, TN 38105, USA

[2]Center of Excellence in Data Driven Discovery, Department of Structural Biology, St. Jude

Children's Research Hospital, Memphis, TN 38105, USA

[3]Department of Laboratory Medicine, Cleveland Clinic, Cleveland, OH 44106, USA

[4]Department of Bone Marrow Transplantation and Cellular Therapy, St. Jude Children's Research

Hospital, Memphis, TN 38105, USA

[5]Center for Proteomics and Metabolomics, St. Jude Children's Research Hospital, Memphis, TN

38105, USA

[6]Department of Structural Biology, Department of Developmental Neurobiology, St. Jude

Children's Research Hospital, Memphis, TN 38105, USA

[7]Department of Surgery, University of Tennessee Health Science Center, 910 Madison Ave., Suite

300, Memphis, TN 38163, USA

[#]Current address: Key Laboratory of Immune Response and Immunotherapy, School of Basic

Medical Sciences, Division of Life Sciences and Medicine, University of Science and Technology

of China, Hefei 230027, Anhui, China

*Corresponding author: piyush.sharma@stjude.org, douglas.green@stjude.org

ABSTRACT

T cell exhaustion is linked to persistent antigen exposure and perturbed activation events,

correlating with poor disease prognosis. Tumor-mediated T cell exhaustion is well documented;

however, how the nutrient-deprived tumor niche affects T cell receptor (TCR) activation is largely

unclear. We show that methionine metabolism licenses optimal TCR signaling by regulating the

protein arginine methylome, and limiting methionine availability during early TCR signaling

promotes subsequent T cell exhaustion. We discovered a novel arginine methylation of a Ca^{2+}-

activated potassium transporter, KCa3.1, prevention of which results in increased Ca^{2+}-mediated

NFAT1 activation, NFAT1 promoter occupancy, and T cell exhaustion. Furthermore, methionine

supplementation reduces nuclear NFAT1 in tumor-infiltrating T cells and augments their anti-

tumor activity. These findings demonstrate metabolic regulation of T cell exhaustion determined

during TCR engagement.

Main Text

To provide an effective anti-tumor response, CD8[+] T cells are activated in draining lymph nodes

(dLN) and within the nutrient-deprived tumor microenvironment (TME) (*1, 2*). The TME is a

constraining niche that leads to defective cytotoxic function of CD8[+] T cells and tumor evasion, at

least in part by promoting T cell exhaustion (*3, 4*). T cell exhaustion is a differentiation state

characterized by a loss of proliferative capacity and a reduction in effector molecules(*5*). To date,

T cell exhaustion has been described as being the result of chronic antigen exposure and associated

changes in specific epigenetic signatures (*6*). Recent studies, however, have shown that T cells can

acquire an exhaustion signature early after activation (*7*), which raises the possibility that the

exhaustion program is driven by factors that occur early after activation in addition to chronic

antigen exposure.

T cell receptor (TCR) engagement is stringently regulated to ensure optimal activation. TCR

activation leads to a rapid increase in metabolic and proliferative capacities that are necessary for

T cell function and expansion. Metabolic flux during TCR engagement is crucial, and changes in

extracellular nutrient availability have been shown to modulate T cell activation (*8*). Activation of

T cells within the TME is vital for an effective anti-tumor response (*2*), an environment known for

its paucity of nutrients. It is therefore likely that nutrient metabolites play a role in optimal TCR

activation. One such metabolite, Methionine (Met), is known to be crucial for epigenetic and

proteomic maintenance and is among the amino acids depleted in the TME. Tumor cells act as a

Met sink, depriving T cells, and supplementation of Met enhances anti-tumor immunity (*9*). It is

unclear, however, how reduced levels of extracellular Met affect TCR-mediated T cell activation

and eventual fate.

Results

Early methionine limitation promotes T cell exhaustion.

To determine which amino acids were rapidly consumed upon T cell activation and whether Met was among them, we activated OT-I transgenic CD8$^+$ T cells in complete medium and performed targeted mass spectrometry. We observed a significant decrease in intracellular Met and other amino acids in only 10 min (Fig. 1A). Among the amino acids important for T cells, Met is the sole amino acid responsible for cellular methylome maintenance via the S-adenosyl Met (SAM) pathway, and alterations in the SAM pathway adversely affect T cell function (*10, 11*). In complete media, no significant changes were observed in SAM or S-adenosyl homocysteine (SAH) (Fig. 1B, upper panel). Instead, T cells consumed the intracellular Met pool only in Met-deprived medium, correlating with decreases in SAM concentrations and corresponding increases in SAH (Fig.1B, bottom panel). These observations indicate that extracellular Met is required to maintain the SAM cycle during T cell activation. By fueling their own rapid proliferation (*12*), tumor cells act as nutrient sinks, creating a nutrient-deprived niche for T cells. To determine whether tumor cells deplete Met in this T cell niche, we cultured CD8$^+$ T cells along with B16f10 (B16) melanoma cells in a trans-well system and analyzed the amino acid profile in a time-dependent fashion. We observed that tumor cells consume Met (and other amino acids), reducing availability for T cells and diminishing the intracellular pool (Fig. S1A, Table. 1). Furthermore, mass spectrometric analysis of tumor-infiltrating CD8$^+$ lymphocytes (CD8$^+$ TIL) and draining lymph node (dLN) CD8$^+$ T cells from B16 subcutaneous tumor-bearing mice at day 9 revealed a ~50% reduction in intracellular Met in CD8$^+$ TIL compared to that in dLN CD8$^+$ T cells (Fig. 1C and Fig. S1B). We also observed a similar decrease in intracellular Met in CD8$^+$ TIL isolated from primary human colorectal carcinomas compared with corresponding CD8$^+$ T cells from PBMC (Fig. 1D) and compared with that of the tumor cells themselves (Fig. S1C).

CD8$^+$ T cells are activated in both secondary lymphoid tissues and within tumors(*1, 13*). Thus, upon tumor infiltration, CD8$^+$ T cells will experience antigen presentation in a nutrient-deficient microenvironment. To determine whether Met limitation during early TCR engagement affects T cell proliferation, we lowered the available amount of Met to 0.03 mM in the culture medium, approximately half that of murine serum (~0.061mM). Cell-trace violet (CTV) labelled OT-I CD8$^+$ T cells were activated in media with control (0.1 mM) or 0.03 mM Met for 30 min to 6 hr, after

which Met was restored to 0.1 mM, and we observed no difference in the proliferation of T cells

activated in reduced Met for different times (Fig. S1D). Upon transferring these activated T cells

into $Rag1^{-/-}$ mice bearing B16 tumors expressing ovalbumin (B16-OVA) 24 h after activation (Fig.

S1E), we observed defective tumor control in T cells activated in 0.03mM Met compared to 0.1

mM Met, across all time points, with the earliest being 30 min (Fig. S1F). Next, we generated

LCMV-GP33 specific memory T cells (Fig. S1G) and activated them for 24 hrs with GP33 peptide

in either 0.1 mM Met or 0.03 mM Met for 30 min, after which Met was restored to 0.1 mM and

then transferred them into B16-GP33 (B16 expressing GP33 peptide) tumor-bearing $Rag1^{-/-}$ mice.

We observed defective tumor control leading to poor survival in mice receiving T cells activated

in 0.03 mM Met compared to 0.1 mM Met (Fig. S1H and I). We then activated OT-I CD8$^+$ T cells

in either 0.1 mM or 0.03 mM Met for 30 min as previously and transferred into B16-OVA tumor-

bearing $Rag1^{-/-}$ or C57BL6/J (WT) mice. We again observed that T cells activated in 0.03 mM Met

for the initial 30 min presented with defective tumor control and resultant poor survival of the

recipient animals (Fig. 1E and Fig. S1J). Further analysis of these same animals revealed increased

tumor burden at day 9, a reduction in CD62L$^+$, CD62L^{hi}CD44^{hi} central memory cells (Tcm),

increased TOX expression in T cells, and reduced IFNγ production (Fig. S1K, L and M).

T cell exhaustion is known to contribute to poor prognoses in cancer patients[14]. This

dysfunctional state is at least in part driven by the transcription factor TOX and is accompanied by

markers such as increased surface expression of PD1 and Tim-3 and reduced effector cytokine

production[5, 15]. To examine exhaustion here we activated CD45.1$^+$ or CD45.2$^+$ OT-I CD8$^+$ T

cells in 0.1 (control) or 0.03 mM Met for 30 min, restored Met to 0.1 mM for 24 hrs, and mixed

them at a ratio of 1:1 before transferring them into B16-OVA tumor-bearing $Rag1^{-/-}$ mice (Fig.

1F). Even though CD8$^+$ TIL initially activated in 0.03mM outnumbered the 0.1 mM-activated

CD8$^+$ TIL at day 12 (Fig. S2A), these cells exhibited significant increases in exhaustion-associated

expression of PD1 and Tim-3 (Fig. 1G) and TOX (Fig. 1H), reduced TCF-1 expression (Fig. S2B),

and reduced effector cytokines (Fig. 1I).

Exhausted T cells have been further defined as precursors of exhaustion (TEXprog) and

TEXprogenitor-like cells, which both produce a replicative burst and effector function upon

checkpoint blockade (16), and terminally exhausted cells (TEXterm), which fail to mount an

effective effector response ($17, 18$). Although TOX is associated with maintenance of exhausted
T cells, it is also expressed in TEXprog (19). We therefore further dissected the TOX$^+$ CD8$^+$ T
cells from Fig. 1H and found decreased TEXprog (TCF1$^+$Tim-3$^-$) and increased TEXterm (TCF1$^-$
Tim-3$^+$) among T cells initially activated in 0.03 mM Met, suggesting a decreased pool of stem-
like TEXprog cells (Fig. S2C and D). Furthermore, evaluation of total CD8$^+$ TILs from mice
receiving T cells activated in 0.03 mM Met revealed a lower frequency of TEXprog cells as
TCF1$^+$Tim-3$^-$ (Fig. 1J), Tim-3$^-$Ly108$^+$ (Fig. S1E), CX3CR1$^+$Ly108$^+$ TEXprogenitor-like cells
(Fig. 1K) and a corresponding increase in frequency of TEXterm cells defined by TCF1$^-$Tim-3$^-$
(Fig. 1L), Tim-3$^+$Ly108$^-$ and CXCR3$^-$Tim-3$^+$ (Fig. S2F). We also observed a decrease in memory-
associated markers such as CD62L$^+$, CD62L$^+$CD44$^+$ (Fig. S2G) and CD127$^+$CD27$^+$ (Fig. S1H and
I) and an increase in KLRG1$^+$CD127$^-$ terminal effector-like T cells when initially activated in 0.03
mM Met (Fig. S1J and K).

To examine the transcriptomic state, we performed global RNA-seq of day 9 TIL from T cells
originally activated for 30 min in 0.1 mM or 0.03 mM Met and then cultured in 0.1 mM Met for
24hrs prior to transfer into B16-OVA tumor-bearing mice. We found several transcriptomic
changes, the majority of which being increased expression of differentially enriched (DE) genes
in T cells initially activated in 0.03 mM Met (Fig. S3A and B). Over-representation analysis of the
DE genes in T cells activated in 0.1 mM Met revealed enrichment of GOBP pathways which are
associated with T cell effector functions along with increases in transcription factors associated
with a stem-like T cell state and memory generation(20-23) (Fig. S3C). Similar analysis of cells
initially activated in 0.03 mM Met showed enrichment of genes which are associated with terminal
differentiation or dysfunction of T cells and transcription factors associated with hyperactivation
and development and maintenance of T cell exhaustion(24-27) (Fig. S3D). GSEA analysis also
showed an enriched exhaustion-associated gene set ($28, 29$) in CD8$^+$ TIL initially activated in 0.03
mM Met (Fig. 1M) and enrichment of hallmark gene sets associated with T cell effector function
in control T cells (Fig. S3E). These results suggest that Met availability during initial TCR
engagement influences T cell differentiation, with initial, reduced Met availability promoting
hyperactivation and driving T cells towards exhaustion.

Methionine regulated Ca^{2+} flux induces NFAT1.

Among the earliest events in T cell activation is the influx of Ca^{2+} (*30*). This is accompanied by

the activation of calcineurin, which dephosphorylates nuclear factor of activated T cells 1

(NFAT1), resulting in its nuclear translocation and activation (*31*). Because low Met levels for

only the first 30 min of activation influenced the T cell phenotype, we investigated changes in Ca^{2+}

flux using Fluo-8. We observed increased Ca^{2+} influx upon stimulation of T cells with anti-

CD3/CD28 antibody in 0.03 mM compared to 0.1 mM Met (Fig. 2A). NFAT1 cooperates with the

AP1 complex to ensure optimal T cell activation and differentiation, but high levels of NFAT1

activity have been associated with T cell dysfunction and exhaustion (*31, 32*). Confocal imaging

revealed an increase in activated NFAT1(nuclear localization of NFAT1) in T cells 30 min

activated with anti-CD3/28 in 0.03 mM Met compared to 0.1 mM Met (Fig. 2B, Fig. S4A),

quantified as either a ratio of nuclear NFAT1 to whole cell NFAT (Fig. 2C and S4C(0.0 mM Met))

or as nuclear NFAT1 fluorescence intensity (Fig. S4B), which were abolished upon treatment with

either the calcineurin inhibitor Cyclosporin A (CsA) (Fig. 2B and C) or the CRAC channel

inhibitor YM-58483 (Fig. S4D and E). We then activated $CD8^+$ T cells in complete medium for

24hrs and rested them for 48hrs before secondary activation in 0.03 mM or 0.1 mM Met and

observed increased nuclear NFAT1 levels in 0.03 mM Met 30 min after activation (Fig. S4F),

suggesting that the effect of limiting Met on NFAT1 activation is independent of previous

activation events. We also examined NFAT2 at 30 min post activation and found no difference in

its activation in T cells activated in either 0.1 mM or 0.03 mM Met (Fig. S4G).

NFAT1 is known to regulate several exhaustion associated genes such as *Lag3, Pdcd1, Havcr2,*

Ctla4, Tnfrsf9 (32). We performed NFAT1 CUT&RUN on T cells activated in 0.1 mM or 0.03

mM Met for 30 min, followed by 0.1 mM Met for 24hrs and observed more reads in the low Met

group (Fig. 2D). Met limitation for the initial 30 min resulted in increased binding of NFAT1 to

Lag3, Pdcd1, Havcr2, Ctla4, Tnfrsf9, and Tox gene loci (Fig. 2E and S4H), which also correlated

with increased transcription of these genes (Fig. S4I). Altogether, our results suggest that reduced

Met availability during the initial stages of TCR activation directs the trajectory of T cells towards

hyperactivation and eventually exhaustion, and the maintenance of extracellular Met levels in the

face of rapid Met consumption upon TCR engagement regulates Ca^{2+}-mediated NFAT1 activity

and T cell fate.

KCa3.1 methylation regulates NFAT1.

Methionine metabolism is the sole pathway for epigenetic and proteomic methylome maintenance, which is known to affect T cell function(*9*). When converted to SAH, SAM acts as the methyl donor; thus, the ratio of SAM to SAH is known as the cellular methylation potential (*33*), an index of a cell's capacity to maintain its methylome. We calculated the SAM/SAH ratio over time after TCR engagement in the absence of extracellular Met and found this ratio to be significantly reduced as early as 10 min post-activation (Fig. S5A), suggesting alterations in the methylome. Met metabolism is essential for DNA methylation (*34*); however, we found no significant differences in global 5-methylcytosine levels at either 30 min or 24 hrs in T cells activated as above (Fig. S5B). We similarly assessed H3me3K27 and H3me2K4 by CUT&RUN at 24 hrs post activation, but again, no global changes were observed (Fig. S5C and D). Thus, Met restriction for only the first 30 min of T cell activation does not appear to directly affect global DNA or histone methylation.

Therefore, we postulated that Met restriction influences early methylation events independently of DNA and histone methylation. Post-translational protein arginine methylation mediated by protein arginine methyltransferases regulates protein function(*35, 36*). We found that arginine methylation in T cells activated in 0.0 mM or 0.03 mM Met was significantly reduced 30 min post-activation when compared to T cells activated in 0.1 mM Met (Fig. S5E and Fig. 3A). To identify specific proteins differentially methylated upon TCR engagement in limited Met, CD8[+] T cells activated for 30 min were processed for methyl-arginine immunoprecipitation and quantification using multiplexed tandem mass tag-mass spectrometry (TMT-MS). We identified common proteins which were differentially methylated in 0.0 mM and 0.03 mM Met activation conditions compared to 0.1 mM Met (Fig. S5F) and found the top candidate, KCa3.1, to be less methylated at Arginine-350 (R350) (Fig. S5G). Furthermore, protein-BLAST analysis showed that KCa3.1 R350 is conserved across species, including humans, which suggested that R350 may be important for KCa3.1 regulation (Fig. S5H).

KCa3.1 is a calcium-activated potassium transporter encoded by *Kcnn4* and is activated by calmodulin binding upon TCR-mediated Ca^{2+} flux (*37*) (Fig. S5I). During T cell activation, the plasma membrane is hyperpolarized to ensure sufficient Ca^{2+} flux for optimal signaling (*38*), and

this is maintained by K^+ efflux through potassium transporters such as KCa3.1 (*39, 40*). Based on

our earlier observation of increased Ca^{2+} signaling, we hypothesized that R350 methylation of

KCa3.1 limits its activity, and loss of this methyl group may result in hyperactivation, leading to

increased Ca^{2+} influx and NFAT1 activation. To test this, we used the chemical inhibitor TRAM-

34 to reduce KCa3.1 activity (*41, 42*) during T cell activation and observed reduced Ca^{2+} flux upon

TRAM-34 treatment only in 0.03 mM Met and not in 0.1 mM Met (Fig. 3B) at the concentration

of TRAM-34 we employed. Measurement of nuclear NFAT1 levels showed that the treatment

normalized nuclear NFAT1 levels in T cells activated in 0.03 mM Met in a dose-dependent manner

and had no effect on T cells activated in 0.1 mM Met (Fig. S6A and B). We then activated OT-I T

cells in 0.1 mM or 0.03 mM Met with either DMSO or TRAM-34 for 30 min and then cultured

the cells for an additional 24 hrs in 0.1 mM Met before transferring into B16-OVA tumor-bearing

Rag1[-/-] mice. Chemical inhibition of KCa3.1 for 30 min significantly improved anti-tumor activity

in T cells activated for 30 min in 0.03 mM Met (Fig. 3C), suggesting that KCa3.1 activity

influences T cell function.

Methylation-deficient KCa3.1 increases NFAT1 activation.

To specifically interrogate the role of R350 methylation of KCa3.1 in T cell activation, we cloned

KCa3.1 (Kca3.1[WT]) and a mutant with an alanine substitution (KCa.3.1[R350A]) into retroviral vectors

and expressed them in primary murine CD8[+] T cells in which endogenous KCa3.1 was deleted

using CRISPR/Cas9 (Fig. S6C, D and E), and expression of the two was similar (Fig. S6F). Ca^{2+}

flux analysis using ICR-1 AM showed increased flux upon activation in KCa3.1[R350A] T cells (Fig.

3D) with concordant increases in nuclear NFAT1 30 min post activation (Fig. 3E, F and Fig. S6G).

This heightened nuclear NFAT1 in cells expressing KCa3.1[R350A] was visible under basal

conditions increased upon activation in a time-dependent manner (Fig. S6H).

To better understand how arginine methylation mechanistically modulates KCa3.1 activity, we

performed MD simulations with the solved crystal structure of human KCa3.1(hKCa3.1) bound to

calmodulin (CaM) at R352, corresponding to R350 in murine KCa3.1 (Fig. S7A). The calmodulin

binding pocket encompasses D78-T79-D80-S81-E82-E83-E84, which is predominantly

negatively charged/polar, where E84 forms a salt bridge with R352 of hKCa3.1. Our results

showed that both symmetrical di-methylation (SDMA) and asymmetrical di-methylation (ADMA)

of R352 introduced a hydrophobic methyl group into this negatively charged/polar CaM binding

pocket, thereby weakening the hKCa3.1 and CaM interaction (Fig. S7B). Furthermore, MD

simulations showed that the salt bridge interaction of R352 with E84 of the CaM binding pocket

diminished over time, especially in the case of SDMA where both guanidino nitrogens are

methylated, compared to ADMA in which only one nitrogen is methylated (Fig. S7C), suggesting

that disruption of arginine methylation promoted stabilization of calmodulin binding, leading to

increased activation of the channel and Ca^{2+} flux, consistent with our experimental observations.

To investigate whether the genomic state of T cells activated in reduced Met is similar to those

with mutant KCa3.1, we analyzed global chromatin accessibility via ATAC-Seq after 24 hrs of *in-*

vitro activation of T cells initially activated in 0.1 mM or 0.03 mM Met and T cells expressing WT

or mutant KCa3.1. This analysis showed a separation of groups in principle component analysis

(PCA), suggesting distinct phenotypes (Fig. S8A and B) with significantly more accessible regions

in T cells initially activated in 0.03 mM Met, similar to KCa3.1^{R350A} T cells (Fig. S8C). Upon

comparing the ATAC-Seq results, we found several common differentially accessible regions

(DAR) (Fig. S8D) with enrichment of gene sets associated with T cell exhaustion (Fig. S8E).

HOMER motif analysis was similarly enriched in exhaustion-associated motifs (*24*) in both

ATAC-Seq and the common DAR (Fig. S8F). Furthermore, GSEA of ATAC-Seq of WT or mutant

KCa3.1-expressing T cells revealed enrichment of GOBP terms associated with negative

regulators of T cell-mediated immunity and gene sets associated with T cell exhaustion (*28, 29*) in

T cells expressing mutant KCa3.1^{R350A} (Fig. S8G and H), suggesting that ablation of R350

methylation drives T cells towards exhaustion early after activation. Despite these results,

KCa.3.1^{R350A} did not fully phenocopy the chromatin effects of activation in 0.03 mM Met,

suggesting that differentially methylated proteins in addition to KCa.3.1 may also impact

chromatin accessibility following activation in the latter condition.

Ablation of KCa3.1 methylation promotes T cell exhaustion.

To examine the consequences of ablation of R350 methylation *in vivo*, we transferred T cells

expressing KCa.3.1^{WT} or KCa.3.1^{R350A} into B16-OVA tumor-bearing Rag1$^{-/-}$ mice. We found that

T cells expressing KCa.3.1^{R350A} displayed impaired tumor control with less survival of recipient

animals (Fig. 3G and H). Tumors were isolated at day 12 post-T cell injection, and consistently,

much larger tumors were extracted from animals treated with T cells expressing KCa.3.1^{R350A} (Fig.

3I and Fig. S9A). KCa.3.1^{R350A}-expressing TIL showed high surface expression of the exhaustion

markers PD1, Lag3 and Tim-3 (Fig. S9B) that correlated with increased TOX expression (Fig.

S9C). Among these cells we also found fewer TEXprog (CD69^{lo}Ly108^{hi})([43]) and IFNγ- and

TNFα-producing T cells (Fig. S9D and E). Next, we performed a similar experiment in a

competitive setting, wherein CD45.1$^+$ KCa.3.1^{WT} and CD45.1$^+$CD45.2$^+$ Kca.3.1^{R350A}-expressing

OT-I CD8$^+$ T cells were mixed at a 1:1 ratio and transferred into B16-OVA tumor-bearing *Rag1$^{-/-}$*

mice (Fig. 3J). TIL analysis at D12 showed increased PD1$^+$Tim-3$^+$, Eomes$^+$PD1$^+$ and TOX

expression in Kca.3.1^{R350A} T cells, which correlated with reduced expression of the effector

cytokines IFNγ and TNFα (Fig. 3K, L and Fig. S9F). Further analysis revealed a decreased

frequency of TCF1$^+$Tim3$^-$ TEXprog and increased TCF1$^-$Tim-3$^+$ TEXterm populations (Fig. 3M

and Fig. S9G). We also observed a decrease in CD62L$^+$ and CD127$^+$CD27$^+$ memory T cells (Fig.

S9H) with an increase in the KLRG1$^+$CD127$^-$ terminal effector-like phenotype in KCa.3.1^{R350A}-

expressing T cells (Fig. S9I).

In addition to tumors, T cell exhaustion is also a prominent feature of chronic viral infection ([44]).

We utilized the Lymphocytic-Choriomeningitis Virus (LCMV) infection system and infected WT

mice containing CD45.1$^+$CD45.2$^+$ KCa.3.1^{WT} and CD45.2$^+$ KCa.3.1^{R350A}-expressing Thy1.1$^+$ P14$^+$

T cells at a 1:1 ratio into mice, which were infected with either an acute Armstrong or a chronic

Clone-13 strain, and their spleens were analyzed on day 9 post-infection (Fig. 3N). The

KCa.3.1^{R350A} population frequency was significantly reduced in Clone-13 infection, but no

significant change was observed in acute infection (Fig. S9J). In both acute and chronic infections,

KCa.3.1^{R350A}-expressing P14$^+$ T cells displayed increased KLRG1$^+$CD127$^-$ terminal effector-like

cells and a concurrent decrease in KLRG1$^-$CD127$^+$ memory precursor cells (Fig. S9K, L and M).

KCa.3.1^{R350A}-expressing T cells also presented with increased TOX, PD1 and Tim3 expression

(Fig. 3O and P). Furthermore, we found that KCa.3.1^{R350A}-expressing T cells harbored reduced

frequencies of CX3CR1$^+$PD1$^+$ TEXprog cells and increased Tim-3$^+$Cx3CR1$^+$PD1$^+$ TEXeff and

TCF1$^-$Tim3$^+$ TEXterm cells, especially in the setting of chronic infection (Fig. 3Q, R and S). Thus,

ablation of R350 methylation drives T cells towards exhaustion trajectory even at the early stages

of infection. Together, these experiments show that R350 methylation maintains optimal KCa3.1

function, ensuring optimal T cell activation and effector function in both tumors and viral
infections.
312
Acute Met supplementation promotes anti-tumor T cell function.
To determine whether CD8$^+$ TIL in the TME (where Met is limited) also showed increased nuclear
NFAT1, we sorted activated CD44$^+$ CD8$^+$ T cells from B16 tumors and respective dLN at day 9
post implantation and imaged for NFAT1. We observed increased levels of nuclear NFAT1 in
CD44$^+$CD8$^+$ TIL compared to counterparts from dLN (Fig. 4A), suggesting that reduced Met
availability in the TME may have led to increased NFAT1 activation in those T cells. Continuous
Met supplementation in tumors has been shown to promote anti-tumor T cell activity by promoting
global epigenetic maintenance (9). Our findings, however, highlight an early role for Met in T cell
fate, and we hypothesized that Met supplementation up to that of serum levels would increase T
cell anti-tumor activity at the early stages of tumor development. We therefore injected 50 μl of
61μM Met peri-tumorally once per day for 5 days when tumor volumes were 70-100 mm^3 (Fig.
S10A) and found that treatment decreased nuclear NFAT1 levels in CD44$^+$ CD8$^+$ TIL as compared
to TIL from HBSS-injected tumors (Fig. 4B). Indeed, nuclear NFAT1 in TIL was similar to or
lower than that in CD8$^+$ T cells from dLN (Fig. S10B). Furthermore, acute Met supplementation
delayed tumor growth in multiple murine tumor models, thereby improving overall survival (Fig.
3C, D; Fig. S10C, D and E). Met supplementation resulted in decreased PD1, Tim3, and CD69
surface expression (Fig. S10F) along with increased CD62L^{hi}CD44^{hi} Tcm and reduced
CD62L^{lo}CD44^{hi} Tem (Fig. S10G) compared to TIL in HBSS-treated tumors. In contrast to
immunocompetent WT mice, Met injected peri-tumorally in tumor-bearing NSG mice resulted in
no difference in tumor growth (Fig. S10H), implying a lymphocyte-mediated effect. Depletion
with an anti-CD8 antibody (Fig. S10I) in B16-OVA tumor-bearing mice abrogated the effect of
Met supplementation on tumor control (Fig. 3E). We then employed a contralateral tumor model
with B16-OVA implanted in one shoulder and in the opposite abdominal flank and injected Met
peri-tumorally to the latter tumor and Met supplementation of the abdominal tumors impeded
contralateral tumor growth (Fig. S10J and K), further indicating a specific T cell response. We
also found that dietary Met supplementation in B16 tumor-bearing mice showed better tumor
growth control (Fig. S10L), which is concordant with a recent study showing that Met

supplementation in humans increased the effector function of CD8$^+$ T cells(9). Met

supplementation, therefore, enhances the anti-tumor function of CD8$^+$ T cells.

Acute Met supplementation enhances ICB and CAR treatment.

Immune checkpoint blockade (ICB) has been widely used either alone or in combination with

cancer treatments (45). To query whether Met supplementation might cooperate with ICB, we

administered either IgG isotype or anti-PD1 antibody along with HBSS or Met supplementation

in subcutaneous MC38 tumor-bearing mice (Fig. 4F) and found that Met plus anti-PD1 treatment

resulted in significantly reduced tumor growth and increased survival of these animals compared

to controls (Fig. 3G and H). CAR-T cell treatment has been clinically successful in the treatment

of hematological malignincies, but less so in solid tumors ($46, 47$). In a solid tumor model (48) in

RAG1$^{-/-}$ mice, we supplemented Met in the presence of transferred murine B7-H3 CAR-T cells in

a B7-H3$^+$ F420 (Fig. 4I), and this too improved tumor control and animal survival (Fig. 4J and K).

Discussion

T cell exhaustion is generally understood to be a consequence of persistent antigen stimulation ($5,$

44); however, situations exist in which antigen persistence does not result in exhaustion, such as

in autoimmune T cells in Type I diabetes (23). It is therefore possible that in addition to chronic

antigen exposure, nutrient limitation drives exhaustion ($49, 50$). A recent study suggested that T

cells can follow an exhaustion trajectory within hours of antigen recognition (7). Here, we propose

that the state of T cell exhaustion is the result of a multifactorial pathway that involves the

assimilation of different input signals and is engaged as early as the initial cognate antigen

recognition.

Studies have shown that strong TCR signals result in hyperactivation and activation-induced cell

death, whereas weak TCR signals result in dampened responses (51). NFAT family members are

key regulators of T cell activation and interact with key partners to induce activation-associated

genes including effector- and exhaustion-associated genes (52). Continuous or partner-less NFAT

activation is associated with hyperactivation, anergy and exhaustion ($31, 32, 53$); therefore, NFAT

activation must be tightly controlled for optimal T cell function, but little is known about how

niche-specific nutrients regulate NFAT. We show that during rapid tumor proliferation in the early

growth phase, CD8[+] TIL display increased NFAT1 activation, which is regulated metabolically. Our data provide a novel mechanism of Met metabolism in the regulation of NFAT1 activation via post-translational modification of KCa3.1, without which CD8[+] T cells progresses towards exhaustion. We found that arginine methylation in KCa3.1 limits Ca^{2+}-dependent NFAT1 signaling, ensuring optimal activation in tumors and infections. Met supplementation to the TME preserves anti-tumor immunity (9) and limits exhaustion (this study), a treatment that can be harnessed to improve patient prognosis in combination with ICB therapies. Our results emphasize the importance of non-histone protein methylation in the functional maintenance of T cells in diseases and describe a link between TCR signaling, metabolic regulation, and transcription factor activation. These results suggest a change in the current dogma of T cell exhaustion from a state driven by persistent antigen exposure—acquired over a long period of time—to a dynamic state controlled by multiple pathways over the course of T cell activation and differentiation. Our study demonstrates that metabolism-dependent, rapid proteomic modifications that occur upon TCR engagement influence subsequent T cell fate and provides mechanistic insight into how niche-specific nutrient deprivation renders T cells dysfunctional.

Acknowledgements: We thank P. Fitzgerald, S. Sirasanagandla, M. Yang and L. Harris, for technical and administrative assistance; L. Long and H. Chi for providing retroviral vectors and Cas9-OTI mice; C. Guy for microscope imaging assistance; M. Morrison for organizing patient samples; the staff at the St. Jude Immunology Flow Cytometry Core Facility for cell sorting. We thank Dr. S. Pounds for additional assistance with the statistical analysis of the data in Fig. 4B. Statistical calculations were performed using GraphPad Prism. Illustrations were created using BioRender software.

Funding: This worked in supported by National Institutes of Health grants AI123322 and CA231620 (D.R.G.), American Lebanese Syrian Associated Charities (ALSAC) (G.K.) and National Cancer Institute grant K99CA256262 (D.H.)

Author Contributions

Conceptualization: PS
Methodology: PS, AG, GP, AM, JP, MMB, GK, ESG, DRG

Investigation: PS, AG, EBR, GP, KCV, KI, AM, DH

Visualization: PS, SP, MJC

Software: SP, MJC

Supervision: DRG

Funding acquisition: DRG, DH, GK

Writing – original draft: PS, DRG

Writing – review & editing: PS, KCV, DRG

Competing Interests: D.R.G. consulted for Venus Therapeutics, Boehringer-Ingelheim and Asha Pharmaceuticals. E.S.G consulted for Anviron. G.K. has patent applications in field of cell or gene therapy in cancer and has received honoraria from Cell Signaling and Kineticos.

References:

1. E. D. Thompson, H. L. Enriquez, Y. X. Fu, V. H. Engelhard, Tumor masses support naive T cell infiltration, activation, and differentiation into effectors. *J Exp Med* **207**, 1791-1804 (2010).
2. N. Prokhnevska *et al.*, CD8(+) T cell activation in cancer comprises an initial activation phase in lymph nodes followed by effector differentiation within the tumor. *Immunity* **56**, 107-124 e105 (2023).
3. A. R. Lim, W. K. Rathmell, J. C. Rathmell, The tumor microenvironment as a metabolic barrier to effector T cells and immunotherapy. *Elife* **9**, (2020).
4. J. Wei, J. Raynor, T. L. Nguyen, H. Chi, Nutrient and Metabolic Sensing in T Cell Responses. *Front Immunol* **8**, 247 (2017).
5. E. J. Wherry, M. Kurachi, Molecular and cellular insights into T cell exhaustion. *Nat Rev Immunol* **15**, 486-499 (2015).
6. X. Lan, C. C. Zebley, B. Youngblood, Cellular and molecular waypoints along the path of T cell exhaustion. *Sci Immunol* **8**, eadg3868 (2023).
7. M. W. Rudloff *et al.*, Hallmarks of CD8(+) T cell dysfunction are established within hours of tumor antigen encounter before cell division. *Nat Immunol* **24**, 1527-1539 (2023).
8. N. Kedia-Mehta, D. K. Finlay, Competition for nutrients and its role in controlling immune responses. *Nat Commun* **10**, 2123 (2019).
9. Y. Bian *et al.*, Cancer SLC43A2 alters T cell methionine metabolism and histone methylation. *Nature* **585**, 277-282 (2020).
10. R. I. Klein Geltink, E. L. Pearce, The importance of methionine metabolism. *Elife* **8**, (2019).
11. X. Yang *et al.*, MAT2A-Mediated S-Adenosylmethionine Level in CD4(+) T Cells Regulates HIV-1 Latent Infection. *Front Immunol* **12**, 745784 (2021).
12. V. Papalazarou, O. D. K. Maddocks, Supply and demand: Cellular nutrient uptake and exchange in cancer. *Mol Cell* **81**, 3731-3748 (2021).

13. P. Yu *et al.*, Priming of naive T cells inside tumors leads to eradication of established tumors. *Nat Immunol* **5**, 141-149 (2004).
14. D. S. Thommen, T. N. Schumacher, T Cell Dysfunction in Cancer. *Cancer Cell* **33**, 547-562 (2018).
15. Z. Zhang *et al.*, Pan-cancer landscape of T-cell exhaustion heterogeneity within the tumor microenvironment revealed a progressive roadmap of hierarchical dysfunction associated with prognosis and therapeutic efficacy. *EBioMedicine* **83**, 104207 (2022).
16. A. Kallies, D. Zehn, D. T. Utzschneider, Precursor exhausted T cells: key to successful immunotherapy? *Nat Rev Immunol* **20**, 128-136 (2020).
17. W. H. Hudson *et al.*, Proliferating Transitory T Cells with an Effector-like Transcriptional Signature Emerge from PD-1(+) Stem-like CD8(+) T Cells during Chronic Infection. *Immunity* **51**, 1043-1058 e1044 (2019).
18. D. T. Utzschneider *et al.*, Early precursor T cells establish and propagate T cell exhaustion in chronic infection. *Nat Immunol* **21**, 1256-1266 (2020).
19. F. Alfei *et al.*, TOX reinforces the phenotype and longevity of exhausted T cells in chronic viral infection. *Nature* **571**, 265-269 (2019).
20. N. Manolios, J. Papaemmanouil, D. J. Adams, The role of ion channels in T cell function and disease. *Front Immunol* **14**, 1238171 (2023).
21. Q. Feng *et al.*, Lactate increases stemness of CD8 + T cells to augment anti-tumor immunity. *Nat Commun* **13**, 4981 (2022).
22. G. T. Hart, K. A. Hogquist, S. C. Jameson, Kruppel-like factors in lymphocyte biology. *J Immunol* **188**, 521-526 (2012).
23. H. A. Abdelsamed *et al.*, Beta cell-specific CD8(+) T cells maintain stem cell memory-associated epigenetic programs during type 1 diabetes. *Nat Immunol* **21**, 578-587 (2020).
24. J. C. Beltra *et al.*, Stat5 opposes the transcription factor Tox and rewires exhausted CD8(+) T cells toward durable effector-like states during chronic antigen exposure. *Immunity* **56**, 2699-2718 e2611 (2023).
25. Q. Sun *et al.*, STAT3 regulates CD8+ T cell differentiation and functions in cancer and acute infection. *J Exp Med* **220**, (2023).
26. C. Zheng *et al.*, Transcriptomic profiles of neoantigen-reactive T cells in human gastrointestinal cancers. *Cancer Cell* **40**, 410-423 e417 (2022).
27. J. A. Belk, B. Daniel, A. T. Satpathy, Epigenetic regulation of T cell exhaustion. *Nat Immunol* **23**, 848-860 (2022).
28. E. J. Wherry *et al.*, Molecular signature of CD8+ T cell exhaustion during chronic viral infection. *Immunity* **27**, 670-684 (2007).
29. T. A. Doering *et al.*, Network analysis reveals centrally connected genes and pathways involved in CD8+ T cell exhaustion versus memory. *Immunity* **37**, 1130-1144 (2012).
30. M. Trebak, J. P. Kinet, Calcium signalling in T cells. *Nat Rev Immunol* **19**, 154-169 (2019).
31. P. G. Hogan, Calcium-NFAT transcriptional signalling in T cell activation and T cell exhaustion. *Cell Calcium* **63**, 66-69 (2017).
32. G. J. Martinez *et al.*, The transcription factor NFAT promotes exhaustion of activated CD8(+) T cells. *Immunity* **42**, 265-278 (2015).

33.   M. A. Caudill *et al.*, Intracellular S-adenosylhomocysteine concentrations predict
global DNA hypomethylation in tissues of methyl-deficient cystathionine beta-
synthase heterozygous mice. *J Nutr* **131**, 2811-2818 (2001).
34.   N. Zhang, Role of methionine on epigenetic modification of DNA methylation and
gene expression in animals. *Anim Nutr* **4**, 11-16 (2018).
35.   J. W. Hwang, Y. Cho, G. U. Bae, S. N. Kim, Y. K. Kim, Protein arginine
methyltransferases: promising targets for cancer therapy. *Exp Mol Med* **53**, 788-
808 (2021).
36.   A. A. Al-Hamashi, K. Diaz, R. Huang, Non-Histone Arginine Methylation by Protein
Arginine Methyltransferases. *Curr Protein Pept Sci* **21**, 699-712 (2020).
37.   J. Lam, H. Wulff, The Lymphocyte Potassium Channels Kv1.3 and KCa3.1 as
Targets for Immunosuppression. *Drug Dev Res* **72**, 573-584 (2011).
38.   R. K. Rader *et al.*, T cell activation is regulated by voltage-dependent and calcium-
activated potassium channels. *J Immunol* **156**, 1425-1430 (1996).
39.   M. D. Cahalan, K. G. Chandy, The functional network of ion channels in T
lymphocytes. *Immunol Rev* **231**, 59-87 (2009).
40.   S. K. Vodnala *et al.*, T cell stemness and dysfunction in tumors are triggered by a
common mechanism. *Science* **363**, (2019).
41.   M. Bachmann *et al.*, Pharmacological targeting of the mitochondrial calcium-
dependent potassium channel KCa3.1 triggers cell death and reduces tumor
growth and metastasis in vivo. *Cell Death Dis* **13**, 1055 (2022).
42.   H. Wulff *et al.*, Design of a potent and selective inhibitor of the intermediate-
conductance Ca2+-activated K+ channel, IKCa1: a potential immunosuppressant.
Proc Natl Acad Sci U S A **97**, 8151-8156 (2000).
43.   J. C. Beltra *et al.*, Developmental Relationships of Four Exhausted CD8(+) T Cell
Subsets Reveals Underlying Transcriptional and Epigenetic Landscape Control
Mechanisms. *Immunity* **52**, 825-841 e828 (2020).
44.   L. M. McLane, M. S. Abdel-Hakeem, E. J. Wherry, CD8 T Cell Exhaustion During
Chronic Viral Infection and Cancer. *Annu Rev Immunol* **37**, 457-495 (2019).
45.   Q. Sun *et al.*, Immune checkpoint therapy for solid tumours: clinical dilemmas and
future trends. *Signal Transduct Target Ther* **8**, 320 (2023).
46.   R. C. Sterner, R. M. Sterner, CAR-T cell therapy: current limitations and potential
strategies. *Blood Cancer J* **11**, 69 (2021).
47.   F. Marofi *et al.*, CAR T cells in solid tumors: challenges and opportunities. *Stem
Cell Res Ther* **12**, 81 (2021).
48.   H. Du *et al.*, Antitumor Responses in the Absence of Toxicity in Solid Tumors by
Targeting B7-H3 via Chimeric Antigen Receptor T Cells. *Cancer Cell* **35**, 221-237
e228 (2019).
49.   N. E. Scharping *et al.*, The Tumor Microenvironment Represses T Cell
Mitochondrial Biogenesis to Drive Intratumoral T Cell Metabolic Insufficiency and
Dysfunction. *Immunity* **45**, 374-388 (2016).
50.   F. Franco, A. Jaccard, P. Romero, Y. R. Yu, P. C. Ho, Metabolic and epigenetic
regulation of T-cell exhaustion. *Nat Metab* **2**, 1001-1012 (2020).
51.   M. Shakiba *et al.*, TCR signal strength defines distinct mechanisms of T cell
dysfunction and cancer evasion. *J Exp Med* **219**, (2022).

52. A. Quintana, D. Griesemer, E. C. Schwarz, M. Hoth, Calcium-dependent activation of T-lymphocytes. *Pflugers Arch* **450**, 1-12 (2005).
53. B. T. Abe, D. S. Shin, E. Mocholi, F. Macian, NFAT1 supports tumor-induced anergy of CD4(+) T cells. *Cancer Res* **72**, 4642-4651 (2012).

Supplementary Materials References.

1. S. Zhao *et al.*, NKD2, a negative regulator of Wnt signaling, suppresses tumor growth and metastasis in osteosarcoma. *Oncogene* **34**, 5069-5079 (2015).
2. D. Haydar *et al.*, Cell-surface antigen profiling of pediatric brain tumors: B7-H3 is consistently expressed and can be targeted via local or systemic CAR T-cell delivery. *Neuro Oncol* **23**, 999-1011 (2021).
3. G. Sharma *et al.*, Analysis of 26 amino acids in human plasma by HPLC using AQC as derivatizing agent and its application in metabolic laboratory. *Amino Acids* **46**, 1253-1263 (2014).
4. B. Meszaros *et al.*, Recent breakthroughs in computational structural biology harnessing the power of sequences and structures. *Curr Opin Struct Biol* **80**, 102608 (2023).
5. C. H. Lee, R. MacKinnon, Activation mechanism of a human SK-calmodulin channel complex elucidated by cryo-EM structures. *Science* **360**, 508-513 (2018).
6. S. Jo, T. Kim, V. G. Iyer, W. Im, CHARMM-GUI: a web-based graphical user interface for CHARMM. *J Comput Chem* **29**, 1859-1865 (2008).
7. J. Lee *et al.*, CHARMM-GUI Input Generator for NAMD, GROMACS, AMBER, OpenMM, and CHARMM/OpenMM Simulations Using the CHARMM36 Additive Force Field. *J Chem Theory Comput* **12**, 405-413 (2016).
8. E. L. Wu *et al.*, CHARMM-GUI Membrane Builder toward realistic biological membrane simulations. *J Comput Chem* **35**, 1997-2004 (2014).
9. M. A. Lomize, I. D. Pogozheva, H. Joo, H. I. Mosberg, A. L. Lomize, OPM database and PPM web server: resources for positioning of proteins in membranes. *Nucleic Acids Res* **40**, D370-376 (2012).
10. W. L. Jorgensen, J. Chandrasekhar, J. D. Madura, R. W. Impey, M. L. Klein, Comparison of simple potential functions for simulating liquid water. *The Journal of Chemical Physics* **79**, 926-935 (1983).
11. D. A. Case *et al.*, AmberTools. *J Chem Inf Model* **63**, 6183-6191 (2023).
12. P. B. N, *Mathematics of Computation* **28**, 669-671 (1974).
13. in *Oeuvres complètes: Series 1,* A.-L. Cauchy, Ed. (Cambridge University Press, Cambridge, 2009), vol. 10, pp. 399-402.
14. S. E. Feller, Y. Zhang, R. W. Pastor, B. R. Brooks, Constant pressure molecular dynamics simulation: The Langevin piston method. *The Journal of Chemical Physics* **103**, 4613-4621 (1995).
15. J. Åqvist, P. Wennerström, M. Nervall, S. Bjelic, B. O. Brandsdal, Molecular dynamics simulations of water and biomolecules with a Monte Carlo constant pressure algorithm. *Chemical Physics Letters* **384**, 288-294 (2004).
16. J.-P. Ryckaert, G. Ciccotti, H. J. C. Berendsen, Numerical integration of the cartesian equations of motion of a system with constraints: molecular dynamics of n-alkanes. *Journal of Computational Physics* **23**, 327-341 (1977).

17. T. Darden, D. York, L. Pedersen, Particle mesh Ewald: An N·log(N) method for Ewald sums in large systems. *The Journal of Chemical Physics* **98**, 10089-10092 (1993).

18. J. Huang *et al.*, CHARMM36m: an improved force field for folded and intrinsically disordered proteins. *Nat Methods* **14**, 71-73 (2017).

19. R. B. Best *et al.*, Optimization of the additive CHARMM all-atom protein force field targeting improved sampling of the backbone phi, psi and side-chain chi(1) and chi(2) dihedral angles. *J Chem Theory Comput* **8**, 3257-3273 (2012).

20. W. Humphrey, A. Dalke, K. Schulten, VMD: visual molecular dynamics. *J Mol Graph* **14**, 33-38, 27-38 (1996).

21. B. Langmead, S. L. Salzberg, Fast gapped-read alignment with Bowtie 2. *Nat Methods* **9**, 357-359 (2012).

22. H. Li *et al.*, The Sequence Alignment/Map format and SAMtools. *Bioinformatics* **25**, 2078-2079 (2009).

23. D. W. Barnett, E. K. Garrison, A. R. Quinlan, M. P. Stromberg, G. T. Marth, BamTools: a C++ API and toolkit for analyzing and managing BAM files. *Bioinformatics* **27**, 1691-1692 (2011).

24. P. A. Ewels *et al.*, The nf-core framework for community-curated bioinformatics pipelines. *Nat Biotechnol* **38**, 276-278 (2020).

25. H. M. Amemiya, A. Kundaje, A. P. Boyle, The ENCODE Blacklist: Identification of Problematic Regions of the Genome. *Sci Rep* **9**, 9354 (2019).

26. F. Ramirez *et al.*, deepTools2: a next generation web server for deep-sequencing data analysis. *Nucleic Acids Res* **44**, W160-165 (2016).

27. Y. Zhang *et al.*, Model-based analysis of ChIP-Seq (MACS). *Genome Biol* **9**, R137 (2008).

28. A. R. Quinlan, I. M. Hall, BEDTools: a flexible suite of utilities for comparing genomic features. *Bioinformatics* **26**, 841-842 (2010).

29. W. J. Kent, A. S. Zweig, G. Barber, A. S. Hinrichs, D. Karolchik, BigWig and BigBed: enabling browsing of large distributed datasets. *Bioinformatics* **26**, 2204-2207 (2010).

30. C. S. Ross-Innes *et al.*, Differential oestrogen receptor binding is associated with clinical outcome in breast cancer. *Nature* **481**, 389-393 (2012).

31. Y. Chen, A. T. Lun, G. K. Smyth, From reads to genes to pathways: differential expression analysis of RNA-Seq experiments using Rsubread and the edgeR quasi-likelihood pipeline. *F1000Res* **5**, 1438 (2016).

32. G. Yu, L. G. Wang, Q. Y. He, ChIPseeker: an R/Bioconductor package for ChIP peak annotation, comparison and visualization. *Bioinformatics* **31**, 2382-2383 (2015).

33. A. Subramanian *et al.*, Gene set enrichment analysis: a knowledge-based approach for interpreting genome-wide expression profiles. *Proc Natl Acad Sci U S A* **102**, 15545-15550 (2005).

34. G. Yu, L. G. Wang, Y. Han, Q. Y. He, clusterProfiler: an R package for comparing biological themes among gene clusters. *OMICS* **16**, 284-287 (2012).

35. M. P. Meers, T. D. Bryson, J. G. Henikoff, S. Henikoff, Improved CUT&RUN chromatin profiling tools. *Elife* **8**, (2019).

36. A. Dobin, T. R. Gingeras, Mapping RNA-seq Reads with STAR. *Curr Protoc Bioinformatics* **51**, 11 14 11-11 14 19 (2015).

37. B. Li, C. N. Dewey, RSEM: accurate transcript quantification from RNA-Seq data with or without a reference genome. *BMC Bioinformatics* **12**, 323 (2011).

38. M. I. Love, W. Huber, S. Anders, Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. *Genome Biol* **15**, 550 (2014).

39. D. Risso, J. Ngai, T. P. Speed, S. Dudoit, Normalization of RNA-seq data using factor analysis of control genes or samples. *Nat Biotechnol* **32**, 896-902 (2014).

40. A. Liberzon *et al.*, The Molecular Signatures Database (MSigDB) hallmark gene set collection. *Cell Syst* **1**, 417-425 (2015).

41. K. Yu *et al.*, High-Throughput Profiling of Proteome and Posttranslational Modifications by 16-Plex TMT Labeling and Mass Spectrometry. *Methods Mol Biol* **2228**, 205-224 (2021).

42. J. K. Eng *et al.*, A deeper look into Comet--implementation and features. *J Am Soc Mass Spectrom* **26**, 1865-1874 (2015).

43. X. Wang *et al.*, JUMP: a tag-based database search tool for peptide identification with high sensitivity and accuracy. *Mol Cell Proteomics* **13**, 3663-3673 (2014).

44. J. Peng, J. E. Elias, C. C. Thoreen, L. J. Licklider, S. P. Gygi, Evaluation of multidimensional chromatography coupled with tandem mass spectrometry (LC/LC-MS/MS) for large-scale protein analysis: the yeast proteome. *J Proteome Res* **2**, 43-50 (2003).

45. J. E. Elias, S. P. Gygi, Target-decoy search strategy for increased confidence in large-scale protein identifications by mass spectrometry. *Nat Methods* **4**, 207-214 (2007).

46. M. Niu *et al.*, Extensive Peptide Fractionation and y(1) Ion-Based Interference Detection Method for Enabling Accurate Quantification by Isobaric Labeling and Mass Spectrometry. *Anal Chem* **89**, 2956-2963 (2017).

47. B. Bai *et al.*, Deep Multilayer Brain Proteomics Identifies Molecular Networks in Alzheimer's Disease Progression. *Neuron* **105**, 975-991 e977 (2020).

48. G. K. Smyth, Linear models and empirical bayes methods for assessing differential expression in microarray experiments. *Stat Appl Genet Mol Biol* **3**, Article3 (2004).

Main Figure:

FIGURE 1

Figure 1: TCR-mediated, rapid methionine consumption governs anti-tumor effector function. (A,B) Quantification of intracellular amino acids 10 min (a), SAM and SAH up to 60 min (b) in OT-I T cells activated with 10 ng/ml SIINFEKL (n=3). **(C)** Quantification of essential amino acids (left) and peak area of Met (right) in LN CD8$^+$ T cells and CD8$^+$ TIL, isolated at day 12 post tumor implantation (n=3). **(D)** Quantification of intracellular Met in CD8$^+$ PBMC and CD8$^+$ TIL isolated from patient primary colorectal tumors (n= 8). **(E)** Tumor growth (left) and survival (right) of B16-OVA tumors in Rag1$^{-/-}$ mice after transfer of 24 hr activated OT-I T cells initially activated in 0.1 or 0.03 mM Met with 2.5ng/ml SIINFEKL for 30 min, followed by restoration of 0.03 mM to 0.1 mM Met (n=5). **(F)** Schematic of experimental design. OT-I CD8$^+$ T cells with different congenic markers were initially activated in 0.1 mM or 0.03 mM Met for 30 min with replenishment of Met in 0.03 mM to 0.1 mM Met for 24 hrs, transferred into B16-OVA tumor bearing Rag1$^{-/-}$ mouse at a 1:1 ratio and analyzed day 12 post T cell transfer. **(G)** Frequency of PD1$^+$ and PD1$^+$Tim-3$^+$ cells of OT-I CD8$^+$ TIL, initially activated in 0.1 mM and 0.03 mM Met and assessed at D12 post injection (n=7). **(H-I)** Contour plot and frequency of TOX$^+$ cells (h) and frequency of IFNγ$^+$TNFα$^+$ (i) of OT-I CD8$^+$ TIL as in G (n=7). **(J-L)** Frequency of (j) TCF1$^+$Tim-3$^-$ TEXprog, CX3CR1$^+$Ly108$^+$ TEXprogenitor-like (k) and, TCF1$^-$Tim-3$^+$ TEXterm (l) cells in OT-I CD8$^+$ T cells initially activated as in E (n=7, n=4). **(M)** T cell exhaustion gene sets in GSEA analysis of RNA-Seq of OT-I CD8$^+$ TIL as in E n=4. Data are mean±s.e.m. Unpaired two-tailed Student's t-test (A-D), two-way ANOVA (E-left), Mantel-Cox log rank test (E--right) and paired two-tailed Student's t-test (G-L) *P < 0.05, **P < 0.01, ***P < 0.001, ****P < 0.0001.

FIGURE 2

Figure 2: Extracellular methionine availability regulates NFAT1 activation and T cell exhaustion. (A) Fluo-8 AM analysis of Ca^{2+} flux in $CD8^+$ T cells activated with anti-CD3 and anti-CD28 by anti-hamster IgG crosslinking in either 0.1 mM or 0.03 mM Met containing Ca^{2+}

Ringer solution with addition of 2 mM Ca^{2+} to measure Ca^{2+} influx. Representative of 2

experiments. **(B, C)** Representative images of T cells, cultured with or without anti-CD3/28

dynabeads (dark grey masked), in 0.1 mM and 0.03 mM Met, with or without cyclosporin-A

treatment for 30 min, stained with NFAT1 (red, marked), Hoechst (blue) and Phalloidin (green)

(b). Quantification of NFAT1 intensity as nuclear to cell ratio (c) (See Fig. S4A for unmasked

figure). Scale bar=10μm. (Each circle represents one cell, n=40 cells). **(D, E)** Histogram of NFAT1

binding signals (d), read count per million reads normalized to background) and quantification of

known NFAT1-binding regions of *Lag3, Pdcd1, Havcr2, Ctla4* and *Tnfrsf9 and Tox* (e, normalized

read count (see differential binding analysis)) in T cells initially activated in 0.1 or 0.03 mM Met

for 30 min and assessed 24 hrs post activation (n=2). Data are mean±s.e.m. Unpaired two-tailed

Student's t-test (C, D). *$P < 0.05$, **$P < 0.01$, ***$P < 0.001$, ****$P < 0.0001$.

FIGURE 3

Figure 3. KCa3.1 arginine 350 methylation regulates NFAT1-activity upon T cell activation.
(A) Quantification of methyl-arginine in $CD8^+$ T cells activated with anti-CD3/28 dynabeads in 0.1, 0.0, or 0.03 mM Met for 30 min (each circle represents one cell. n= 25 cells). **(B)** Fluo-8 AM analysis of Ca^{2+} flux in $CD8^+$ T cells activated with anti-CD3 and anti-CD28 by anti-hamster IgG crosslinking in either 0.1 mM or 0.03 mM Met containing Ca^{2+}-free Ringer solution treated either with DMSO or 1μM TRAM-34. **(C)** Tumor growth of B16-OVA tumors in $Rag1^{-/-}$ mouse after transfer of OT-I $CD8^+$ T cells activated in 0.1 mM or 0.03 mM Met with either DMSO or 1 μM TRAM-34 for 30 min and cultured for 24 hrs (n=5). **(D)** ICR-1 AM analysis of Ca^{2+} flux in $KCa3.1^{WT}$ and $KCa3.1^{R350A}$ T cells activated with anti-CD3 and anti-CD28 by anti-hamster IgG crosslinking in Ca^{2+}-free Ringer solution. **(E, F)** Representative image (e) and quantification of NFAT1intensity as nuclear to cell ratio (f) in $KCa3.1^{WT}$ and $KCa3.1^{R350A}$ OT-I T cells activated for 30 min with anti-CD3/28 dynabeads (dark grey masked) (See Fig. S6C for unmasked figure), scale bar=10 μm (each circle represents one cell, n=40 cells). **(G, H)** Tumor growth (g) and survival (h) of B16-OVA tumor-bearing $RAG^{-/-}$ mice after transfer of $KCa3.1^{WT}$ and $KCa3.1^{R350A}$ OT-I T cells (n=5). **(I)** B16-OVA tumors harvested at D12 post transfer of $KCa3.1^{WT}$ and $KCa3.1^{R350A}$ OT-I T cells (n=4). **(J)** Schematic of experimental design to assess congenically distinct $KCa3.1^{WT}$ and $KCa3.1^{R350A}$ OT-I T cells, mixed at a 1:1 ratio and transferred into B16-OVA tumor bearing $Rag1^{-/-}$ mice and analyzed day 12 post T cell transfer. **(K-L)** Frequency of $Tim-3^+PD1^+$ (k), histogram and quantification of TOX MFI, and frequency of $IFNγ^+TNFα^+$ (l) in $KCa3.1^{WT}$ and $KCa3.1^{R350A}$ OT-I TIL at D12 post T-cell transfer as in J (n=7). **(M)** Frequency of $TCF1^+Tim-3^-$ TEXprog and $TCF1^-Tim-3^+$ TEXterm in $KCa3.1^{WT}$ and $KCa3.1^{R350A}$ OT-I TIL at D12 post T-cell transfer as in J (n=7). **(N)** Schematic of experimental design to assess congenically distinct $KCa3.1^{WT}$ and $KCa3.1^{R350A}$ Thy1.1 P14 T cells, mixed at a 1:1 ratio, transferred into WT mice, and infected with either LCMV-Armstrong or LCMV-Clone-13. Spleens were analyzed at D9 post infection. **(O, P)** Frequency of Tox^+ and $PD1^+Tim-3^+$ in $KCa3.1^{WT}$ and $KCa3.1^{R350A}$ P14 $CD8^+$ T cells from spleens at D9 post infection with either LCMV-Armstrong (o) or LCMV-Clone-13 (p) (n=4). **(Q-S)** Frequency of $CX3CR1^+PD1^+$ TEXprog (q), $Tim-3^+$ of $CX3CR1^+PD1^+$ TEXeffector-like (r) and $TCF1^-Tim-3^+$ TEXterm (s) in $KCa3.1^{WT}$ and $KCa3.1^{R350A}$ P14 $CD8^+$ T cells from spleens at D9 post infection as in N (n=4). Data are mean±s.e.m. Unpaired two-tailed Student's t-test (A, F), 2-way ANOVA (C, G), Mantel-Cox test (H) and paired two-tailed Student's t-test (K-M, O-S). *$P < 0.05$, **$P < 0.01$, ***$P < 0.001$, ****$P < 0.0001$.

FIGURE 4

Figure 4. Acute methionine supplementation promotes CD8$^+$ T cell-mediated tumor control.
(A) Nuclear NFAT1 quantification in CD44$^+$CD8$^+$ T cells isolated from B16 tumors and respective dLNs at D9 post implantation (each circle represents one cell, n=40 cells, n=5). **(B)** Quantification of nuclear NFAT1 in CD44$^+$ CD8$^+$ T cells isolated from B16 tumors, after 5 days of peri-tumoral supplementation per day of 50 µl HBSS or 61 µM Met (each circle represents one cell, n=40 cells, n=4). **(C, D)** Tumor growth (c) and survival (d) of B16 tumor-bearing WT mice, supplemented either with HBSS or 61 µM Met peri-tumorally for five days as in B (n=10). **(E)** B16 tumor growth (left) and survival (right) of WT mice, treated intraperitoneally with IgG isotype or anti-CD8 antibody at D-1 and D3 and treated peri-tumorally with HBSS or Met from D7-D12 as in B (n=5, representative of two experiments). **(F)** Schematic design to assess tumor growth with anti-PD1 treatment and with either HBSS or Met supplementation. **(G, H)** Tumour growth (g) and survival (h) of MC38 tumor-bearing WT mice, treated either with anti-PD1 or IgG-isotype and supplemented with HBSS or Met peri-tumorally for five days as in B (n=5, representative of two experiments). **(I)** Schematic of experimental design to assess the effect of HBSS or Met supplementation on CAR-T cell therapy in mouse solid tumors. **(J, K)** Tumor growth (j) and survival (k) of F420 tumor-bearing Rag1$^{-/-}$ mice injected with B7-H3 CAR T-cells at D0 and peri-tumorally treated with HBSS or Met from D1-D6 as in B (n=5). Data are mean±s.e.m. Unpaired two-tailed Student's t-test (A), 2-way ANOVA (C, E-left, G and J), and Mantel-Cox log rank test (D, E-right, H and K). Statistical analysis of (B) is described in methods. *P < 0.05, **P < 0.01, ***P < 0.001, ****P < 0.0001.s

SUPPLEMENTARY FIGURES

Supplementary Fig. 1

Supplementary Figure 1: Met availability during TCR engagement determines anti-tumor T cell function. (A) Quantification of intracellular Met in CD8$^+$ T cells and B16 cells at 0, 6, and 12 hrs of co-culture (n=3). **(B)** Quantification of intracellular non-essential amino acids from LN CD8$^+$ and CD8$^+$ TIL, isolated at D12 post implantation of B16 tumors (n=3). **(C)** Quantification of intracellular Met from tumor cells and CD8$^+$ TIL isolated from patient primary colorectal tumors (n=4). **(D)** T cell proliferation via cell-trace violet staining of OT-I CD8$^+$ T cells activated in either 0.1 mM Met or 0.03 mM Met for the indicated times before restoration to 0.1 mM Met in 0.03 mM Met conditions, analyzed 72 hrs post activation (n=3). Representative of 3 samples per group. **(E)** Schematic design for OT-I T cell activation initially in 0.1 or 0.03 mM Met, followed by restoration of Met to 0.1 mM for 24 hrs before injection into B16-OVA tumor bearing mice. **(F)** Tumor growth of B16-OVA in Rag1$^{-/-}$ treated with OT-I CD8$^+$ T cells activated as described in (E) for 30 min-6hrs (n=5). **(G)** Schematic experimental design for generation of LCMV-GP33 specific memory T cells, which were then activated as in described in (E) before injection into B16-OVA tumor-bearing Rag1$^{-/-}$ mice. **(H, I)** Tumor growth (h) and survival (i) of B16-OVA tumors treated as described in (G) (n=5). **(J)** Tumor growth and survival of B16-OVA tumor bearing WT mice treated with or without OT-I CD8$^+$ T cells activated in 0.1 or 0.03 mM Met for 30 min before restoring Met to 0.1 mM in 0.03 mM Met for 24 hrs, as described in (E) (n=5). **(K)** Tumor growth of B16-OVA tumor bearing Rag1$^{-/-}$ mice injected with OT-I T cells as in (E) (n=5). **(L, M)** Tumor weight (l) and frequency of CD62L$^+$, Tcm (CD62L^{hi} CD44^{hi}), IFN-γ^+, and TOX expression (MFI) (m) in OT-I CD8$^+$ TIL isolated from B16-OVA tumor bearing Rag1$^{-/-}$ mice at D9 post-injection with OT-I T cells activated as in (E) (n=5). Data are mean±s.e.m. Unpaired two-tailed Student's t-test (A, B, C, L and M), two-way ANOVA (F and H) and Mantel-Cox test (I and J-right). *P < 0.05, **P < 0.01, ****P < 0.0001

Supplementary Fig. 2

Supplementary Figure 2: Met limitation promotes T cell exhaustion and hampers memory generation. (A) Frequencies of T cells isolated from B16-OVA tumors at D12 post transfer of OT-

I T cells activated as in Fig. S1E (n=7). **(B)** Expression of TCF1$^+$ on OT-I CD8$^+$ T cells at D12 post

transfer of OT-I T cells, activated as in Fig. S1E (n=7). **(C, D)** Representative contour plot (c) and

quantification (d) of TCF1 and Tim-3 expressing cells from TOX$^+$ CD8$^+$ TIL from OT-I CD8$^+$ T cells

isolated at D12 post transfer of OT-I T cells activated as in Fig. S1E into B16-OVA tumor-bearing

Rag1$^{-/-}$ mice (n=7). **(E-G)** Frequency of Tim-3$^-$Ly108$^+$ TEXprog (e), Tim-3$^+$Ly108$^-$, CXCR3$^-$Tim-3$^+$

TEXterm (f), CD62L$^+$ T memory and CD62L$^+$CD44$^+$ T central memory cells (g) in OT-I TIL as in (A)

(n=7 (f), n=4 (g)). **(H-J)** Representative contour plot and quantification of CD127$^+$CD27$^+$ memory

cells (h, i) and KLRG1$^+$CD127$^-$ terminal effector cells (j) in OT-I CD8$^+$ TIL isolated at D12 post of

cells activated as in Fig. S1E into B16-OVA tumor-bearing Rag1$^{-/-}$ mice (n=4). Data are mean±s.e.m

with statistical analyses performed using paired two-tailed Student's t-test (*$P < 0.05$, **$P < 0.01$,

***$P < 0.001$).

Supplementary Fig. 3

Supplementary Figure 3: RNA-Seq analysis shows enrichment of exhaustion signature in low Met. (A) Volcano plot and top ten differentially expressed genes (red) (0.1 mM vs 0.03 mM, p<0.05, FC>1.5) from OT-I CD8[+] TIL isolated from mice injected with OT-I T cells, initially activated in 0.1 mM Met or 0.03 mM Met (as in Fig. S1E), at day 12 post injection into B16-OVA tumor-bearing mice (n=4). **(B)** Heatmap of DE genes, which shows either increased expression in cells initially activated in 0.03 mM Met or 0.1 mM Met, from the RNAseq in (A) (n=4). **(C)** Over-representation analysis of DE genes with higher expression in TIL initially activated in 0.1 mM Met, showing enrichment of biological processes and transcription factors associated with T cell effector function and regulation. **(D)** Over-representation analysis of DE genes with higher expression in TIL initially activated in 0.03 mM Met, showing enrichment of gene sets and transcription factors associated with promotion and maintenance of the exhausted T cell state. **(E)** GSEA analysis of hallmark genesets as analyzed from RNA-Seq of OT-I TIL initially activated in 0.1 or 0.03 mM Met, isolated from B16-OVA tumors at D12 post transfer (n=4). RNA-Seq statistics were applied as described in methods.

Supplementary Fig. 4

Supplementary Fig. 4: Met limitation results in increased TCR-mediated NFAT1 activation.

(A) Un-edited image of Figure. 2B showing the unmasked, Red CD3/38 dynabeads which are masked grey in the main figure to highlight NFAT1 staining. **(B)** Quantification of nuclear NFAT1 from the experiment shown in Figure 2B. **(C)** NFAT1 intensity quantification (nuclear to cell ratio) of CD8$^+$ T cells activated in either 0.1 mM, 0.00 mM or 0.03 mM Met for 30 min with anti CD3/28 dynabeads and stained for NFAT1 (each circle represents one cell, n=40 cells). **(D)** Fluo-8 AM analysis of Ca^{2+} flux in CD8$^+$ T cells activated with anti-CD3 and anti-CD28 and anti-hamster IgG crosslinking in either 0.1 mM or 0.03 mM Met containing Ca^{2+}-free Ringer solution treated with DMSO or 5μM YM58483. **(E)** NFAT1 intensity quantification (nuclear to cell ratio) of CD8$^+$ T cells activated for 30 min by CD3/28 dynabeads in either 0.1 mM or 0.03 mM Met in the presence of DMSO or 5μM YM58483 (n=40 cells). **(F)** NFAT1 quantification (nuclear to cell ratio) of previously activated OT-I T cells, treated with anti-CD3/28 dynabeads for 30 min. T cells were initially activated with 2.5ng/ml SIINFEKL for 24 hrs in control medium and rested with 10 ng/ml mIL-7 for 48 hrs (each circle represents one cell, n=40). **(G)** NFAT2 quantification (nuclear to cell ratio) of CD8$^+$ T cells, activated in 0.1 mM or 0.03 mM Met for 30 min with anti-CD3/28 dynabeads (each circle represents one cell, n=40 cells). **(H)** NFAT1 Cut&Run tracks at *Pdcd1, Havcr2, Lag3, and Tnfrsf9* locus. The gray highlights the peaks showing increased NFAT1 binding (n=2). **(I)** Normalized RNA expression quantified by quantitative-PCR performed 24 hrs post activation of OT-I CD8$^+$ T cells activated with 2.5 ng/ml SIINFEKL in 0.1 mM or 0.03 mM Met for 30 min followed by 0.1 mM (n=6). Data are mean±s.e.m. Unpaired two-tailed Student's t-test (B, C, E, F, G) and one-tailed Student's t-test (I). *P < 0.05, **P < 0.01, ***P < 0.001, ****P < 0.0001.

Supplementary Fig. 5

Supplementary Fig. 5: Initial Met limitation results in reduced arginine methylation but not DNA or histone methylation. (A) Methylation potential (SAM/SAH) of 10 ng/ml SIINFEKL activated OT-I T cells in 0.1 mM or 0.0 mM Met, at 10, 30 and 60 min (n=3). **(B)** % 5-methylytosine (%5-mc) of OT-I CD8$^+$ T cells activated with 2.5 ng/ml SIINFEKL in 0.1 mM or 0.03 mM Met for 30 min, then 0.1 mM Met for 24 hr (n=3). **(C, D)** Histograms (top) and heat-map (bottom) of H3K4me3 (c) and H3K27me3 (d) CUT&RUN reads (read count per million, normalized to background) in OT-I CD8$^+$ T cells activated initially in 0.1 mM or 0.03 mM Met, with restoration to 0.1 mM after 30 min and processed at 24 hrs post activation (n=3). **(E)** Representative images of CD8$^+$ T cells activated with anti-CD3/28 dynabeads (circled bright red) in 0.1, 0.0, and 0.03 mM Met for 30 min, stained with methyl-arginine (red, marked), Hoechst (blue), and Phalloidin (green) (scale bar=10 μm) **(F)** Venn diagram of enriched methyl-arginine proteins at 30 min post activation of T cells in 0.1 mM vs 0.0 mM or 0.1 mM vs 0.03 mM Met, as identified by TMT-MS. **(G)** Heat-map (Log$_2$ fold change) of TMT-MS identified proteins (p<0.05) with enriched methyl-arginine in T cells activated in 0.1 mM vs 0.0mM or 0.03 mM Met by IgG crosslinking with anti-CD3 for 30 min (n=3). **(H)** Conserved arginine (highlighted red) in KCa3.1 as analyzed by protein-BLAST in multiple species and visualized using Jalview. **(I)** Schematic pathway of TCR mediated KCa3.1 activation and function. Data are mean±s.e.m. Unpaired one-tailed Student's t-test (A) or unpaired two-tailed Student's t-test (B).

Supplementary Fig. 6

Supplementary Fig. 6: KCa3.1 R350 methylation regulates NFAT1 activation. (A, B) NFAT1 quantification (nuclear to cell ratio) in OT-I CD8$^+$ T cells activated for 30 min with anti CD3/28 dynabeads in 0.03 mM Met (a) or 0.1 mM Met (b) in the presence of 0.5, 1 or 2 μM TRAM-34 (each circle represents one cell, n=45). **(C)** Schematic of experimental design of endogenous KCa3.1 knockdown with expression of KCa3.1^{WT} and KCa3.1^{R350A} in Cas9-OT-I or P14$^+$ CD8$^+$ T cells. **(D)** Sorting strategy for *in vitro*-generated KCa3.1^{WT} and KCa3.1^{R350A} T cells; gate shows the cells that express WT or R350A KCa3.1 with endogenous KCa3.1 knockdown. **(E)** Immunoblot showing knockdown of KCa3.1 with sgRNA1 and sgRNA2, compared to control. **(F)** Immunoblot of KCa3.1 expression in KCa3.1^{WT}- and KCa3.1^{R350A}-expressing CD8$^+$ T cells. **(G)** Un-edited image of Figure. 3E showing the unmasked CD3/38 dynabeads which are masked grey in the main figure to highlight NFAT1 staining. F. Nuclear NFAT1quantification of KCa3.1^{WT} and KCa3.1^{R350A}, either control (no Dynabeads) or activated with anti-CD3/28 Dynabeads for 10 min (each circle represents one cell, n=45 cells). Data are mean±s.e.m. Unpaired two-tailed Student's t-test. *P < 0.05, **P < 0.01, ****P < 0.0001.

Supplementary Fig. 7

Supplementary Fig. 7: huKCa3.1 R352 methylation weakens KCa3.1-Calmodulin interaction *in-silico*. (A) Tetrameric assembly of human KCa3.1-Calmodulin complex within the membrane (left). KCa3.1 is represented as ribbons and calmodulin quartet as cartoon with interacting Ca^{2+}

ions as purple spheres. K$^+$ ions are represented as cyan spheres. KCa3.1 monomer (blue) is shown

in complex with calmodulin (yellow) along with demethylated arginine-352 in green (right). **(B)**

Interaction of KCa3.1 monomer (blue cartoon) with calmodulin (yellow cartoon) as un-

methylated (top), SDMA (middle) and ADMA (bottom). The Calmodulin-binding pocket is

represented as surface with interacting amino acids as sticks (negatively charged-red, polar

residue-green) and R352 as ball and sticks; introduction of hydrophobic methyl group (middle

and bottom) weakens the interactions between calmodulin and KCa3.1 monomer. **(C)** MD

simulations analysis: Analysis of variations of distance of salt bridge between calmodulin E84 and

demethylated R352 (WT), SDMA R352 and ADMA R352 over the course of simulation time (top).

Illustration of total count of negatively charged residues within a 4Å radius of all four

R352/SDMA/ADMA of huKCa3.1 the course of simulation time (bottom) (averaged across all four

monomers, three MD trials).

Supplementary Fig. 8

Supplementary Fig. 8: ATAC-Seq analysis of KCa3.1 R350A similar exhaustion T cell signature as

of initial Met limitation. (A,B) PCA analysis of ATAC-Seq of OT-I T cells, activated with 2.5ng/ml

SIINFEKL in 0.1 mM and 0.03 mM Met and cultured in 0.1 mM Met for 24 hrs (a), or KCa3.1^{WT}-

and KCa3.1^{R350A}- activated with 2.5 ng/ml SIINFEKL for 24 hrs (b). n=3. **(C)** Chromatin accessibility

heatmap of of T cells initially activated in 0.1 or 0.03 mM Met (left) or expressing activated

KCa3.1^{WT} or KCa3.1^{R350A} (right) with each row representing peaks (p<0.05 and log2 fold-change

>1.5) displayed over the span of 2kb window with peak as center (grouped from least to max

differential region), analyzed 24 hrs post activation with 2.5ng/ml SIINFEKL (n=3). **(D)** Common

differentially accessible regions (DAR) (p<0.05, fold change>1.5) between ATAC-Seq of OT-I cells

expressing KCa3.1^{WT} or KCa3.1^{R350A}, or OT-I cells initially activated in 0.1 mM or 0.03 mM Met as

in (A,B) **(E)** Over-representation analysis of the common DAR from (d) showing the top 8 gene

sets associated with exhaustion and activation. **(F)** Top 10 motifs analyzed by HOMER in ATAC-

Seq DAR's from OT-I CD8+ T cells initially activated in 0.03 mM Met vs 0.1 mM Met (left) or

expressing KCa3.1^{R350A} vs KCa3.1^{WT} and activated as (b) (middle), and from the common DAR's

from (d). **(G)** GOBP terms associated with negative regulation of T cells enriched in OT-I T cells

expressing KCa3.1^{R350A} vs KCa3.1^{WT} (activated at b) GSEA analysis of promoter reads from ATAC-

Seq post 24 hrs of activation (n=3). **(H)** T cell exhaustion gene sets in GSEA analysis of differential

accessible promoter reads from ATAC-Seq of OT-I T cells expressing KCa3.1^{WT} or KCa3.1^{R350A},

activated with 2.5 ng/ml SIINFEKL for 24 hrs (n=3).

Supplementary Fig. 9: Ablation of KCa3.1 R350 methylation promotes T cell exhaustion *in-vivo*.
(A) Tumor weight of B16-OVA tumors isolated at D12 post transfer of OT-I T cells expressing KCa3.1^{WT} or KCa3.1^{R350A} (n=4). **(B)** Surface expression of PD1$^+$, Lag3$^+$ and PD1$^+$Tim-3$^+$ in CD8$^+$ TIL isolated at D12 post transfer into B16-OVA tumor-bearing mice of OT-I T cells expressing KCa3.1^{WT} or KCa3.1^{R350A} (n=4). **(C)** Representative histogram of TOX expression and quantification in CD8$^+$ TIL isolated at D12 post transfer of OT-I T cell expressing KCa3.1^{WT} or KCa3.1^{R350A} as (b) (n=4). **(D)** Representative contour plot (left) and quantification of IFNγ^+TNFα^+ (right) in OT-I T cells expressing KCa3.1^{WT} or KCa3.1^{R350A} at D12 post transfer as (b) (n=4). **(F, G)** Expression of EOMES$^+$PD1$^+$ (f) and representative contour plot of expression of TCF1 and Tim-3 (g) on congenically marked OT-I T cells expressing KCa3.1^{WT} or KCa3.1^{R350A}, mixed and transferred at a ratio of 1:1. TIL were isolated from B16-OVA tumors at D12 post T cell transfer (n=7). **(H, I)** Expression of memory markers CD62L and CD127$^+$CD27$^+$ (h), representative contour plot and quantification of KLRG1$^+$CD127$^-$ terminal effector cells (i) on congenically marked OT-I T cells expressing KCa3.1^{WT} or KCa3.1^{R350A}, as in (F), isolated at D12 post transfer into B16-OVA tumor-bearing mice (n=7). **(J)** Frequency of KCa3.1^{WT}- and KCa3.1^{R350A} -expressing P14 T cells in spleens, D9 post infection with either LCMV-Armstrong or LCMV-Clone-13 (n=4). **(K-M)** Representative contour plot (k) and quantification of KLRG1$^+$CD127$^-$ terminal effector cells (l) and KLRG1$^-$CD127$^+$ memory precursor cells (m) in spleens of animals injected with congenically marked P14 T cells expressing KCa3.1^{WT} or KCa3.1^{R350A}, mixed 1:1, and infected with either LCMV-Armstrong or LCMV-Clone-13. Analysis on d9 post infection (n=4). Data are mean±s.e.m. Unpaired two-tailed Student's t-test (A-D) and paired two-tailed Student's t-test (F-M). *P < 0.05, **P < 0.01, ***P < 0.001, ****P < 0.0001.

Supplementary Fig. 10

Supplementary Fig. 10: Acute Met supplementation promotes anti-tumor T cell differentiation

and tumor control. (A) Schematic of experimental design of acute HBSS/Met treatment in sub-

cutaneous tumors. **(B)** Nuclear NFAT1 quantification of CD44$^+$ CD8$^+$ T cells isolated from B16

subcutaneous tumors and dLN, 5 days post daily peri-tumoral supplementation of 50 μl 61μM

Met (n=4 mice, each circle represents one cell, n=40 cells). **(C-E)** Tumor growth (left) and survival

(right) of WT mice with sub-cutaneous MC38 colorectal tumors (c, n=10), Lewis-lung carcinoma

(LLC) tumors (d, (representative of two experiments, n=5) and F420 osteosarcoma tumors (e,

n=5), treated with either HBSS or 50 μl 61μM Met peri-tumorally daily for 5 days (n=10). **(F, G)**

Expression of PD1, Tim-3, CD69$^+$ (f), Tcm (CD62L^{hi}Cd44^{hi}) and Tem (CD62L^{lo}CD44^{hi}) (g) on CD8$^+$ TIL

isolated from B16 tumor bearing mice 2 days after 5 daily peri-tumoral injection with HBSS or

Met (n=4). **(H)** Tumor growth of B16f10 and MC38 tumors in NSG mice treated peri-tumorally

with 50 μl HBSS or 50 μl 61μM Met daily for 5 days (n=5). **(I)** Representative flow cytometry plot

of blood showing CD8$^+$ T cell depletion in anti-CD8 antibody treated mice compared to IgG-

Isotype treated mice (n=4). **(J)** Schematic of experimental design of contralateral tumor

experiment. **(K)** Growth of B16f10 tumors implanted on left shoulder (dashed lines) and right

flank (solid lines) following peri-tumoral injection of flank tumor with 50 μl HBSS or 61μM Met

daily for 5 days (n=5, representative of two experiments). **(L)** B16 tumor growth in WT mice fed

with either 1 % Met chow or 1.5 % Met chow, 7 days post tumor implantation (n=5). Data are

mean±s.e.m. Two-way ANOVA (C-F, J and K), Mantel-Cox log rank test (C, D and E), unpaired two-

tailed Student's t-test (B, G). *P < 0.05, **P < 0.01, ***P < 0.001, ****P < 0.0001.

1018

1019

1020

1021

1022

Supplementary Fig. 11

1023

Supplementary Fig. 11: Methionine metabolism regulates TCR-mediated Ca^{2+} flux and T cell

fate. Diagram illustrating the impact of methionine metabolism on TCR-dependent methylation

of KCa3.1, leading to regulation of Ca2+ flux and subsequent activation of NFAT1. When

extracellular Met levels are low, the potential for methylation decreases, which in turn reduces

KCa3.1 methylation. This results in excessive channel activation and increased Ca2+ flux,

ultimately triggering NFAT1 activation. NFAT1 then moves to the nucleus and activates gene

signatures associated with activation and exhaustion, contributing to hyperactivation and

exhaustion.

1032

Materials and Methods:

Reagents

RPMI 1640 (cat. 11875093, Gibco) and methionine-free RPMI 1640 (cat. A1451701; Gibco)

supplemented with 10% dialyzed FCS (Atlanta Biologicals), sodium pyruvate (cat. 11360070,

Gibco), NEAA (11140050, Gibco), glutamine (cat. 25030081, Thermo-Fisher), and penicillin-

streptomycin cocktail (cat. 15070063, Thermo-Fisher) and stored at 4°C. Methionine (cat. M5308,

Sigma-Aldrich) was reconstituted in PBS (0.1 M) and stored at 4°C. SIINFEKL peptide (cat. 60193,

AnaSpec) peptide was reconstituted in PBS at 0.5 mg/ml and stored at -20°C. TRAM-34 (cat. HY-

13519, MedChem Express) was reconstituted at 1 M with DMSO. Cyclosporin A (CsA) (cat. 30024,

Sigma-Aldrich) was reconstituted to a concentration of 50 mg/ml in DMSO. YM-58483 (cat. 3939,

Tocris) was reconstituted at 20 mM stock in DMSO. All reconstituted reagents were stored at -

20°C and aliquoted to avoid repeated thawing.

Mice

All mouse studies were conducted in accordance with protocols approved by the St. Jude

Children's Research Hospital Committee on Care and Use of Animals, and in compliance with all

relevant ethical guidelines. All mice were kept under a 12h-12h light-dark cycle in a specific-

pathogen-free (SPF) facility at the institute's Animal Resource Center. Age-and sex-matched

mice were randomly assigned to the experimental and control groups. Cas9-OT/P14$^+$-I (Cas9

Knock-in with OVA-specific transgenic TCR (OT-I) or LCMV-GP33 specific TCR(p14)), C57BL/6J

(WT), OT-I, and Rag1$^{-/-}$ mice were bred in-house. The investigators were not blinded to the

experiments and outcome assessments.

Cell lines

B16F10, B16F10-Ova (B16-Ova) melanoma cells were provided by Dr. Hongbo Chi. MC38 and

MC38-Ova colon carcinoma cell lines were provided by Dr. Dario Vignali. The F420 sarcoma cell

line was provided by Dr. Jason T Yustein. HEK293T and Lewis lung carcinoma (LLC) cells were

purchased from the ATCC. Plat-e cells used for retroviral packaging were purchased from Cell

Biolabs (cat. RV-101). All the cell lines used were tested and confirmed to be mycoplasma-

negative but not independently authenticated.

Murine and Human CD8$^+$ T cell isolation and stimulation

Murine CD8$^+$ T cells were isolated from the spleen and lymph nodes (LN) by physical disruption

through a 70μ strainer before treatment with erythrocyte-lysis buffer. Cells were subjected to

CD8$^+$ T cell enrichment using the EasySep mouse CD8$^+$ T cell isolation kit (cat. 19853, StemCell),

according to the manufacturer's protocol. Isolated T cells were then either activated with

SIINFEKL peptide or mouse CD3/28 Dynabeads (cat. 11456D, Thermo-Fisher) in RPMI medium, as

described above. Human CD8$^+$ T cells were isolated from apheresis rings from the St Jude Blood

Donor Center. Blood obtained from an apheresis ring was mixed at a 1:1 ratio with PBS containing

2% FBS and carefully layered onto Lymphoprep (cat. 07801, StemCell) solution. The samples were

then centrifuged at 800g for 30 min at room temperature without braking. The peripheral blood

mononuclear cell (PBMC) layer was aspirated and washed twice in PBS with 2% FBS before

enrichment of CD8$^+$ T cells using the Human CD8$^+$ T Cell Isolation Kit (cat. 17953, StemCell),

according to the manufacturer's protocol. CD8$^+$ T cells were activated using human CD3/28

Dynabeads (cat. 11131D, Thermo Fisher Scientific) in RPMI medium. Apheresis rings were

collected with written consent from the donors for research upon review and approval by the

Institutional Review Board at St Jude Children's Research Hospital.

Ca^{2+} Flux measurement

Isolated T cells were labelled with 3 μM Fluo-8 AM (AAT Bioquest, cat. 21080) or 5 μM ICR-1AM

(Ion Biosciences) with 0.04% pluronic F-127 (Thermo-Fisher, cat. P6867) for 2 hrs in HHBS at 37 °C

and washed twice with Ca^{2+}-free Krebs-Ringer's solution (Thermo Fisher Scientific, cat. no.

J67839-AP). The cells were then resuspended in Ca^{2+}-free Krebs-Ringer's solution and incubated

with anti-CD3ε (4 μg/ml; 2C11, cat. BE0001, Bio-X-Cell) and anti-CD28 (4 μg/ml; 37.51, cat.

BE0015, Bio-X-Cell) on ice for 10 min. 0.1 mM Met or 0.03 mM Met was added to the samples

and the baseline was recorded before the addition of hamster-IgG for crosslinking. Ca^{2+} influx was

measured after the addition of 2mM Ca^{2+}, 1uM ionomycin was added as a positive control for

Ca^{2+} flux. All measurements were recorded on either Cytek Aurora (SpectroFlo) or BD-Fortessa

flow cytometer.

Quantitative PCR

Isolated T cells were activated with 2.5ng/ml SIINFEKL in 0.1 mM Met or 0.03 mM Met for 30 min

after which Met in 0.03 mM was restored to 0.1 mM. T cells were then further cultured for 24

hrs. Cells were washed twice with PBS and RNA was isolated using Direct-zol RNA miniprep Kit

(Zymo research, cat. no. R2050). RNA was quantified and processed for cDNA synthesis using

Reverse Transcriptase (Thermo Fisher Scientific, cat. no. 28025-013). cDNA was then used to

perform quantitative PCR with the specific primers (Table S2) using SYBR green PCR master mix

(Thermo Fisher Scientific, cat. no. 4309155). PCR reaction was run and quantified by QuantStudio

7 flex.

Cloning and virus production

Kcnn4 gRNA's were designed using the Broad Institute platform

(https://portals.broadinstitute.org/gpp/public/analysis-tools/sgrna-design) and cloned into the

ametrine-expressing retroviral vector LMPd-gRNA-mPGK-Ametrine

(sgRNA1:CACCGGGCAGGCTGTCAATGCCACG, AAACCGTGGCATTGACAGCCTGCCC;

sgRNA2:CACCGTGTGGGGCAAGATTGTCTGC, AAACGCAGACAATCTTGCCCCACAC). KCa3.1^{WT} and

KCa3.1^{R350A} cDNA were synthesized by Genscript; the latter contained an alanine substitution at

R350. Both KCa3.1^{WT} and KCa3.1^{R350A} had substitutions at the PAM recognition sites of the gRNAs

(Table S3). Each cDNA was subcloned into the pMIGII. Plat-e cells were co-transfected with the

cloned vectors and pCL-eco (cat. 12371, Addgene) in Opti-MEM (cat. 31985070, Thermo-Fisher)

with Trans-IT-293 transfection reagent (cat. MIR 2705, Mirus Bio) and cultured at 37°C for 12-15

hrs prior to changing the medium and culturing for an additional 24 hrs at 37°C, followed by 24

hrs at 32°C, 5% CO_2. The viral supernatant was harvested and spun at 1800rpm at RT to remove

the cell debris. Viral supernatant was either used immediately or stored at -80°C till further use.

1111

Viral transduction

Naive Cas9-OT-I CD8$^+$ cells were isolated from the spleen and peripheral LNs of Cas9-OT-I mice

using a Magnisort naïve CD8$^+$ T cell isolation kit according to the manufacturer's instructions (cat.

8804-6825-74, Thermo-Fisher). Purified CD8$^+$ T cells were activated *in vitro* for 20 h with plate-

bound anti-CD3ε (5 µg/ml; 2C11, cat. BE0001, Bio-X-Cell) and anti-CD28 (5 µg/ml; 37.51, cat.

BE0015, Bio-X-Cell) antibodies. Viral transduction was performed by spinfection at 900 g at 25 °C

for 3 h with 10 µg/ml polybrene (Sigma-Aldrich), followed by resting for 3 h at 37 °C in 5% CO_2.

Cells were washed and cultured in medium supplemented with mouse rIL-2 (20 U/ml; cat. 212-

12, Peprotech) for 4 days. GFP-Ametrine double-positive cells were sorted using the Reflection

cell sorter (iCyt) or the MoFlo XDP cell sorter and rested for 24 hrs in medium containing rIL-2

(10U/ml, cat. 212-12 Peprotech) before activation or adoptive transfer into the animals.

Flow cytometry

Antibodies from BioLegend included Pacific Blue anti-mouse Ly108 (330-AJ, 134608), BV510 anti-

mouse KLRG1 (2F1, 138421), BV570 anti-mouse CD62L (MEL-14, 104433), BV711 anti-mouse Tim-

3 (RMT3-23, 119727), BV785 anti-mouse CD127 (A7R34, 135037), PE-Cy5 anti-mouse Granzyme-

B (QA16A02, 372226), PE-Fire 700, anti-mouse CD4 (GK1.5, 100484), APC/Cy7 anti-mouse TNFα

(MP6-XT22, 506344), PE-Cy7 anti-mouse KLRG1 (2F1/KLRG1, 138416), PerCP-Cy 5.5 anti-mouse

CD62L (MEL-14, 104432), BV605 anti-mouse CD127 (A7R34, 135025), Pacific blue anti-mouse

CD69 (H1.2F3, 104524), Pacific blue anti-mouse CD45.1 (110722), APC anti-mouse PD1 (RL388,

109111), BV711 anti-mouse PD1 (29F.1A12, 135231), PE anti-mouse IFNγ (XMG1.2, 505808), FITC

anti-mouse TNFα (MP6-XT22, 506304), Pacific Blue anti-human CD45 (HI30, 982306), APC/Cy7

anti-human CD8 (RPA-T8, 344713). BUV563 anti-mouse LAG3 (C9B7W, 741350), BUV615 anti-

mouse CD69 (H1.2F3, 751593), BUV805 anti-mouse CXCR3 (173, 748700), BV421 anti-mouse

EOMES (X4-83, 567166), BV480 anti-mouse CD45.1 (A20, 746666), Alexa-Fluor 488 anti-mouse

TCF1 (S33-966, 567018), Alexa-Fluor 647 TOX (NAN448B, 568356), BUV 496 anti-mouse Ly-108

(13G3, 750046), APC/Cy7 anti-mouse CD44 (IM7, 560568) and BUV805 anti-mouse CD8α (53-6.7,

612898) were acquired from BD Biosciences. BUV395 anti-mouse CD44 (IM7, 363-0441-82),

PerCP-Cy 5.5 anti-mouse IL-2 (JES6-5H4, 45-7021-82), PerCP-eF710 anti-mouse CD27 (O323, 46-

0279-42), and PE-Cy7 anti-mouse Tim-3 (RMT3-23, 12-5870-82, eBioscience) were obtained from

Thermo Fisher. APC anti-human/mouse Tox (REA473, 130-118-335) was obtained from Miltenyi

Biotech. For surface markers, the cells were stained for 30 min on ice in PBS containing 2% FCS

and 0.1% sodium azide (cat. S2002, Sigma-Aldrich). To examine intracellular cytokine production,

the isolated cells were stimulated with a cell activation cocktail (cat. 5476, TOCRIS) in the

presence of monensin (cat. 554724, BD Biosciences) for 4 hr, and stained for intracellular

cytokines and transcription factors using a fixation/permeabilization kit (cat. 00-5523-00,

Ebiosciences). Fixable viability dye (Thermo Fisher) was used to exclude dead cells. Samples were

acquired on a Cytek Aurora (SpectroFlo) or BD LSRII (FACSDiva) flow cytometer and analysed

using FlowJo software.

***In vivo* tumor analysis**

For all tumor experiments, 0.5×10^6 tumor cells were injected subcutaneously into the right flank

of naive mice (aged 6–10 weeks). Met was diluted in HBSS to a concentration of 61 µM per 50µl.

Mice were injected peri-tumorally for 5 days with either 50 µl Met or HBSS. Tumours were

measured with calipers every 2 days and tumor volumes were calculated using the formula:

$(width^2 \times length)/2$. For adoptive transfer experiments, T cells (1×10^6) were transferred i.v. 7 -9

days after tumor injection. For contralateral tumor model, 0.5×10^6 B16 tumor cells were

injected subcutaneously into the right flank and left shoulder of the same mouse. Met or HBSS

was injected peri-tumorally to the right flank tumor (abdominal). For check-point blockade

experiment, 0.5×10^6 MC38 tumor cells were injected subcutaneously into the right flank of WT

mice. The mice were given either anti-PD1 antibody (RMP1-14, Bio X Cell) or rat IgG2a (2A3, Bio

X Cell), intraperitoneally twice at a dose of 150 mg/kg in 100 µl PBS at day 9 and 12. Met or HBSS

was given peri-tumorally for 5 days from day 9 to 13. For the CAR-T tumor model, 0.5×10^6 F420

tumor cells (derived from singly floxed *p53+/F-Col2.3* transgenic mice(*1*)); were injected

subcutaneously into the right flank of Rag1$^{-/-}$ mice. 5×10^6 B7-H3 CAR-T cells (see below) were

transferred i.v. 8-10 days post implantation. Tumour size was measured every 2-4 days.

Generation of mouse CAR-T cells

Generation of mouse B7-H3-CAR constructs and CAR-T generation have been previously

described(*2*). Briefly, HEK293T cells were transiently transfected with the CAR-encoding plasmid,

the Peg-Pam plasmid encoding MoMLV gag-pol, and a plasmid encoding the VSVG envelope. Viral

supernatants were collected at 48 h and filtered with a 0.45m filter. The virus was then used to

transduce the GPE86 producer cell line. B7-H3-CAR-expressing GPE86 cells were stained with

Alexa Fluor-647 anti-human IgG, F(ab')2 fragment antibody (cat. 109-606-006, Jackson

ImmunoResearch) and sorted using the BD FACSAria III system.

Naive CD8$^+$T cells from 6–8 weeks old CD45.1$^+$ mice were activated with plate-bound anti-CD3ε

(1 µg/ml, 145-2C11, cat. BE0001, BioXcell), anti-CD28 (2 µg/ml, 37.51, cat. BE0015, BioXcell) and

50 U ml^{-1} rhIL-2 (cat. 200-02, Peprotech). On day 2 post activation, activated CD8$^+$ T cells were

transduced with retrovirus expressing B7-H3-CAR on retronectin-coated (cat. T100B, Takara)

non-tissue-culture-treated plate in complete RPMI medium supplemented with 50 U/ml rhIL-2.

At 48 h after transduction, CAR-T cells were collected and expanded in the presence of

50 U/ml rhIL-2 for another 3 days before adoptive transfer into tumor-bearing mice.

Tumour-infiltrating lymphocyte isolation

B16 tumors were harvested at the indicated time points and digested using a digestion cocktail

(0.1% Collagenase IV and 0.01% DNaseI) for 5 min at 37 °C. Digested tumors were passed through

a 70m strainer and resuspended in PBS plus 2% dialyzed FCS and processed for analysis and/or

sorting by flow cytometry, as described above.

Patient samples were collected following surgery and were immediately processed. Tumour

samples were dissected into small pieces and digested at 37°C with digestion cocktail for 40 min.

Digested tumor samples were physically disrupted and passed through 70µ filter. Samples were

washed with PBS-2% dialyzed FCS twice and stained for CD45, CD3ε and CD8 and sorted for CD8$^+$

T cells by flow cytometry. Purified T cells were washed once with 1% saline and flash frozen in

liquid nitrogen. Samples were collected upon informed consent and procedures approved by the

IRB of St. Jude Children's Research Hospital.

Immunofluorescence staining and imaging.

CD8$^+$ T cells were immobilized in 8-well poly L-lysine-coated IBIDI chambers and stimulated by

the addition of CD3/28 Dynabeads (cat. 11452D, Thermo Fisher) for 30 min, followed by 10 min

of fixation with 4% paraformaldehyde (PFA) (cat. 15710, Electron Microscopy Science) at 37°C.

The cells were washed once with TBS (50 mM Tris, 100 mM NaCl, pH 8.0) and permeabilized with
permeabilization buffer (50mM Tris, 100 mM NaCl, pH 8.0, 0.3% (v/v) Triton X-100) for 20 min at
room temperature. Cells were washed again with TBS before blocking with TBS containing 2%
BSA (Cat. 001-000-182, Jackson ImmunoResearch) for 60 min at room temperature. The cells
were stained overnight at 4 °C with the following primary antibodies: anti-NFAT1 (1:200, cat.
4389, Cell Signaling Technology), anti-NFAT2 (1:200, cat. D15F1, Cell Signaling Technology) and
anti-mono and dimethyl arginine (1:500, cat. ab412, Abcam). The samples were washed twice
with TBS and incubated for 1 h at room temperature with the following antibodies: anti-rabbit
Alexa Fluor plus 595 (1:1,000, cat. A-11012, Thermo Fisher), anti-mouse CD8-APC (1:500; cat.
100712, BioLegend), Alexa Fluor 488 Phalloidin (1:1000, cat. A12379, Thermo Fisher) and Hoechst
33258 (1:1000, cat. H3569, Thermo-Fisher). Samples were then imaged by Marianas spinning-
disc confocal microscopy (Intelligent Imaging Innovations), using the Prime95B sCMOS camera,
and the 405, 488, 561 and 640 nm laser lines. Sum fluorescence intensities were analyzed using
Slide Book 6 (Intelligent Imaging Innovations).

Immunoblotting
T cells were washed once with ice-cold PBS and immediately lysed with 4x laemmli buffer (Bio-
Rad, cat. 1610747). Cell lysates were separated by SDS-PAGE and blotted with anti-Kcnn4
antibody (cat. PA5-33875, Thermo Fisher) and HRP-conjugated anti-Actin (cat. sc4777-HRP, Santa
Cruz Biotechnology). Images were developed using Clarity Western ECL substrate (cat. 1705061,
Bio-Rad) and acquired on Bio-Rad Chemi-Doc.
1217

Amino acid measurement
Subcutaneous tumors were isolated and processed as described above. Sorted T cells were
washed once with ice-cold saline, flash-frozen with liquid nitrogen, and stored at –80ºC before
analysis. Cell pellets were extracted using 750 µl of methanol/acetonitrile/water (5:3:2, v/v/v)
and the supernatant was dried by lyophilization. Aliquots of 20-50 µl from plasma and TIF were
extracted with at least a 15-fold excess volume of the methanol/acetonitrile/water solution, and

the supernatant was then collected and dried by lyophilization. Dried extracts containing the

hydrophilic metabolites were dissolved in 30 µl of water/acetonitrile (8:2, v/v), and 10 µl was

used to derivatize amino acids as described (*3*) with some modifications. The samples were

placed in glass autosampler vials, and 35 µl of sodium borate buffer (100 mM, pH 9.0) was added

and mixed by pipetting. 10 µl of the 6-aminoquinolyl-N-hydroxysuccinimidyl carbamate (AQC, 10

mM in acetonitrile)-derivatizing reagent (Cayman Chemical) was added, the vial was sealed,

mixed by vortexing, and then incubated at 55°C for 15 min. The vial was cooled to room

temperature and then 1 µL was analyzed by liquid chromatography with tandem mass

spectrometry (LC-MS/MS). An ACQUITY Premier UPLC System (Waters Corp) was used for the LC

separations, using a non-linear gradient condition as follows: 0-0.4 min 3% B; 0.4-8 min 3 to 96%

B (using Curve #8 of the inlet condition in MassLynx™); 8-12 min 96% B; 12-12.5 min 96 to 3% B;

12.5-14 min 3% B. Mobile phase A was water supplemented with 0.15% acetic acid, and mobile

phase B was acetonitrile with 0.15% acetic acid. The column used was an Accucore C30 (50 × 2.1

mm, 2.6 µm) (Thermo Fisher Scientific) operated at 50°C. The flow rate was 300 µl/min and the

injection volume used was 1 µl. All LC/MS solvents and reagents were the highest purity available

(water, acetonitrile, acetic acid, boric acid, sodium hydroxide) and were purchased from Thermo

Fisher Scientific. A Xevo TQ-XS Triple Quadrupole Mass Spectrometry (TQ-XS) (Waters Corp)

equipped with a multi-mode ESI/APCI/ESCi ion source was employed as detector. The TQ-XS was

operated in positive ion mode using a multiple reaction monitoring mass spec method (MRM).

The MRM conditions were set to a minimum of 15 points per peak with an automatic dwell time.

The operating conditions of the source were: Capillary Voltage 3.8 kV, Cone Voltage 40 V,

Desolvation Temp 550°C, Desolvation Gas Flow 1000 L/h, Cone Gas Flow 150 L/h, Nebuliser 7.0

Bar, Source Temp 150°C. Authentic amino acid standards were purchased from Sigma-Aldrich

and used to establish MRM conditions and calibration curves. The monitored parent/daughter

ions, fragmentation collision energy (CE), and retention time window for each amino acid are

listed in Supplementary Table 4. The acquired MRM data was processed using the software

application Skyline 21.2 (MacCoss Lab Software).

1251

1252   ***In-silico* model building of KCa3.1(SK4)-Calmodulin complex and simulation.**

1253

To study the impact of SK4s arginine dimethylation on the binding of calmodulin, we simulated

the SK4 channel:Calmodulin complex in a membrane layer via all-atom molecular dynamics (MD),

which is considered a computational microscope(4). The Cryo-EM structure of the human SK4

channel:Calmodulin complex state (PDB:6CNN(5)) was used as the starting structure. Thisa

tetramer compleconsistsng of four SK4 monomers, four calmodulins (CAMs), three calcium ions

bound to each calmodulin (CAM) (total 1,2) and four K^+ ions bound to the channel selectivity

filter.

Initially, protein was prepared was using the molecular operating environment (MOE) software.

The missing residues in the cryo-EM structure were modeled based on the human AlphaFold

structure (AF-015554-F1). Protonate3D module in MOE was used to assign appropriate

protonation states at pH 7.4 and then energy minimized. Further, simulation system was

prepared using CHARMM-GUI web server(6-8). Protein was placed in the POPE membrane bilayer

and protein orientation was obtained using the PPM web server(9). The protein-membrane

bilayer complex was immersed in the TIP3P(10) water box of size ≈150 X 150 X 230 $Å^3$ with an

edge distance of 40 Å on all sides. KCl salt was added to balance the charges and maintain a

physiological salt concentration of 0.15M. Overall, each wildtype simulation system contains four

SK4 monomers, four CAMs, 12 Ca^{2+} ions, ≈355 K^+ ions, ≈367 Cl^- ions, ≈618 POPE lipids, and

≈128,144 TIP3P waters. The total numbers of atoms in this system are ≈498,944. Two more

systems were generated via a protocol similar to that explained above, except that in one system,

R352 of the SK4 monomer was symmetrically dimethylated and in the second system, R352 was

asymmetrically dimethylated. Dimethylation was introduced using the CHARMM-GUI web

server. Note that arginine's at position 352 in all four SK4 monomers were dimethylated in both

systems.

Further, all three systems were simulated using AMBER20/22 MD software(11). The MD

simulation was performed in three steps. In step-1, energy minimization was carried out to

remove potential steric clashes in two stages. In stage 1, each system was energy-minimized for

5000 steps, while positional restraints were applied to the protein and membrane: the first 2500

steps with the steepest-descent algorithm and the remaining 2500 steps using the conjugate

gradient algorithm(12). In stage 2, the second round of minimization was carried out with all

restraints removed in 50,000 steps. The steepest-descent algorithm(13) was used for the first

25,000 steps, while the conjugate gradient algorithm was used for the remaining 25,000 steps.

In step-2, equilibration was carried in six stages: (1) equilibrated for 125 ps with a time step of

1fs while applying positional restraints on protein (K=10 kcal/mol.Å^2) and membrane (K=2.5

kcal/mol.Å^2), (2) equilibrated for 125 ps with a time step of 1fs w while applying positional

restraints on the protein (K=5.0 kcal/mol.Å^2) and membrane (K=2.5 kcal/mol.Å^2), (3) equilibrated

for 125 ps with a time step of 1fs while applying positional restraints on the protein (K=2.5

kcal/mol.Å^2) and membrane (K=1.0 kcal/mol.Å^2), (4) equilibrated for 500 ps with a time step of 2

fs while applying positional restraints on the protein (K=1.0 kcal/mol.Å^2) and membrane (K=0.5

kcal/mol.Å^2), (5) equilibrated for 500 ps with a time step of 2 fs while applying positional

restraints on the protein (K=0.5 kcal/mol.Å^2) and membrane (K=0.1 kcal/mol.Å^2) and (6)

equilibrated for 500 ps with a time step of 2 fs while applying weak positional restraints on the

protein (K=0.5 kcal/mol.Å^2) and restraints on the rest of the system were completely removed.

The first two equilibration simulations were performed under NVT conditions, whereas the last

four equilibration simulations were performed under NPT conditions. In step-3, production

simulations were carried out under periodic boundary conditions using the NPT ensemble, during

which the temperature (310 K) and pressure (1 atm) were kept constant. Three replicates of each

system were simulated, and each replicate was simulated for ≈1 μs. Langevin integrator and

Monte Carlo barostat were used($14, 15$) to conduct NPT simulations, whereas the SHAKE

algorithm was used to restrain bonds with hydrogens(16). Particle-mesh Ewald (PME) method

was used to treat long-ranged electrostatic interactions(17) and the nonbonded cutoff was 12Å

(force switching = 10Å). CHARM36 all-atom additive force field($18, 19$) was used to treat the

entire system. Overall, nine MD simulations were conducted across three systems, with each

system comprising three replicates that each ran for approximately 1μs. Consequently, the total

MD simulation data accumulated for analysis amounts to approx. 9 μs, derived from the sum of

simulation times for all systems and replicates (3 systems X 3 replicates X ≈1μs/per replicate =

≈9μs). Two frames per nanosecond were used for the data analysis. Visual molecular dynamics

(VMD)(20) was used for visualization and graphic generation, whereas Tcl was used for data

analysis. Gnuplot was used to generate the time series plots.

1312

ATAC-seq

1313

OT-I CD8$^+$ T cells were activated with 2.5ng/ml SIINFEKL in either 0.1 mM Met or 0.03 mM Met

with restoration of Met to 0.1 mM in 0.03 mM condition post 30 min of activation and incubated

for 24 hrs. KCa3.1^{WT} and KCa3.1^{R350A} cells were activated by 2.5ng/ml SIINFEKL for 24 hrs. Cells

were processed for ATAC-Seq after 24 hrs of activation. Briefly, 50,000 cells were washed with 1

ml cold PBS by centrifugation at 500 g for 5 min at 4°C. Cells were then resuspended in 50 ml cold

cell lysis buffer (10 mM Tris pH 7.4, 10 mM NaCl, 3 mM MgCl$_2$, and 0.1% NP-40). Isolated nuclei

were pelleted at 1,000 × g for 10 min at 4°C, resuspended in 50 µl transposition reaction mixture

(25 µl TD buffer, 2.5 µl TDE1, 22.5 µl ddH$_2$O) (20034197, Illumina), and incubated at 42°C for 45

min. DNA was purified using a MiniElute PCR purification kit (cat. 28004, Qiagen). Samples were

then processed for library barcoding and amplification with Q5 High-Fidelity 2× Master Mix (cat.

M0492S, NEB). Prepared libraries were sent for sequencing after quantification using Qubit and

size distribution as determined using an Agilent 4200 TapeStation.

Analysis of ATAC-seq data

1326

Paired-end sequencing reads were trimmed using Trim Galore (version 0.5.0)

(https://github.com/FelixKrueger/TrimGalore) with default parameters. Reads were aligned to the

reference mouse mm10 assembly using Bowtie 2 (version 2.3.5.1)(*21*) with settings --end-to-end -

-very-sensitive -X 2000. The resulting alignments, recorded in a BAM file, were sorted, indexed,

and marked for duplicates with Picard MarkDuplicates function (version 2.19.0). Afterward, the

BAM file was filtered with SAMtools (version 1.9)(*22*), BamTools (version 2.5.1)(*23*) and scripts

of nf-core/chipseq(*24*) to discard reads, mates that were unmapped, PCR/optical duplicates, not

primary alignments, mapped to multiple locations, or mapped to ENCODE blacklisted

regions(*25*); only reads mapped in proper pair were kept (-F 1804 -f 2 -q 30). The alignments were

shifted by deepTools alignmentSieve(*26*) with --ATACshift and default parameters. Nucleosome-

free reads (fragment length < 109 bp) were separated from the BAM files with SAMtools (version

1.9). MACS (version 2.1.2)(*27*) was used to call peaks from the BAM files with narrowPeak

setting, --extsize 200, and recommended mappable genome size (default value for other

parameters). Chromatin accessibility signal was normalized by scaling to 1 million mapped reads

using BEDTools (version 2.27.1)(*28*) and bedGraphToBigWig (version 377)(*29*) and visualized

as heatmaps using deepTools plotHeatmap (version 3.2.1)(*26*).

Differentially accessibility analysis was performed using DiffBind(*30*) (v.3.4.7) (summit = 50,

normalized by trimmed mean of M values(*31*) (TMM) approach, library size estimated by reads

mapped to "background" regions, the genomic regions (binned into 15000 bp) overlapped with

peaks). Peaks were annotated to the nearest genes with annotatePeak function in R package

ChIPseeker(*32*) (v.1.30.3) at the gene level using the default parameters. Genes were ranked by

fold change (log2 transformed) of associated peaks located within +/- 1.5 kb of transcription start

sites (if a gene was associated with multiple peaks, the mean was used). Gene Set Enrichment

Analysis(*33*) (GSEA) was then performed against MSigDB collections with R package

clusterProfiler(*34*) (v.4.2.2). The common differentially accessible regions (DAR) between ATAC-

Seq of KCa3.1WT and KCa3.1R350A and T cells activated in 0.1 mM or 0.03 mM Met were

obtained by the overlapping of the DARs (at least 1 bp). Enrichment analysis was performed using

the over-representation approach against MSigDB using the R package clusterProfiler.

CUT&RUN

CUT&RUN experiments were performed as previously described(*35*) with slight modifications.

Purified OT-I CD8[+] T cells were activated with 2.5ng/ml SIINFEKL in 0.1 mM and 0.03 mM Met for

30 min. Met was restored to 0.1 mM post 30 min and cultured for 24 hrs. Cells were washed with

cold PBS and dead cells were removed using Easysep dead cell removal kit (cat. 17899, StemCell).

Cells (2×10^5) were washed twice with wash buffer (20 mM HEPES (H3375, Sigma-Aldrich],

150 mM NaCl (cat. AM9760G, Invitrogen), 0.5 mM spermidine (cat. S0266, Sigma-Aldrich), and

protease inhibitor cocktail (cat. 5056489001, Sigma-Aldrich)), resuspended and bound to

concanavalin-A coated magnetic beads (cat. BP531, Bang Laboratories). The resuspension buffer

was removed using a magnetic stand and beads were resuspended in 100 μl antibody buffer

(20 mM HEPES, 150 mM NaCl, 0.5 mM spermidine, 0.01% digitonin (cat. 300410, Millipore),

2 mM EDTA (cat. AM9260G, Invitrogen), and protease inhibitor cocktail). Primary antibodies

(1:100) were added to the samples and incubated overnight at 4 °C. Next, samples were washed

twice with cold Dig-wash buffer (20 mM HEPES, 150 mM NaCl, 0.5 mM spermidine, 0.01%
digitonin, and protease inhibitor) and pAG-MNase (cat. 123461, Addgene) was added and rotated
at 4 °C for 1 h. Samples were washed twice and resuspended in 50 µl Dig-wash buffer and
CaCl$_2$ (2 µl of 100 mM, cat. 2115, Sigma-Aldrich) was added followed by brief vortexing and
incubation on ice for 30 min. Next, 50 µl of 2× STOP buffer (340 mM NaCl, 20 mM EDTA, 4 mM
EGTA (cat. E3889, Sigma-Aldrich) and 100 µg/ml RNase A (cat. EN0531, Thermo Fisher), and
50 µg/ml GlycoBlue (cat. AM9515, Invitrogen)) was added and mixed by gentle vortexing and
incubated for 30 min at 37 °C to release CUT&RUN fragments. Fragmented DNA was purified
using the NEB Monarch PCR&DNA Cleanup Kit (cat. T1030S, New England Biolabs). DNA libraries
were prepared using the NEBNext Ultra II DNA Library Prep Kit (cat. E7645S, New England Biolabs)
and purified with AMPure SPRI beads (cat. B23318, Beckman-Coulter). Prepared libraries were
quantified using Qubit and size distribution was determined with a Agilent 4200 TapeStation
analysis before paired-end sequencing.

CUT&RUN data processing

Paired-end sequencing reads were trimmed using Trim Galore (v.0.5.0)
(https://github.com/FelixKrueger/TrimGalore) with default parameters. The reads were then
aligned to the reference mouse mm10 assembly using Bowtie 2 (v.2.3.5.1)(*21*) with settings --end-
to-end --very-sensitive --no-mixed --no-discordant -q --phred33 -I 10 -X 700. The resulting
alignments, recorded in the BAM file, were sorted, indexed, and marked for duplicates using the
Picard MarkDuplicates function (v.2.19.0) (Picard toolkit. *Broad Institute GitHub Repository*,
2019). Next, the BAM file was filtered with SAMtools (v.1.9)(*22*), BamTools (v.2.5.1)(*23*) and
scripts of nf-core/chipseq(*24*) to discard reads, including mates that were unmapped, PCR/optical
duplicates, non-primary alignments, reads that mapped to multiple locations or to ENCODE-
blacklisted regions(*25*), or with more than 4 mismatches (-F 0x004 -F 0x008 -F 0x0100 -F 0x0400
-f 0x001 -q 1). MACS (v.2.1.2)(*27*) was used to call peaks from the BAM file with IgG control
and recommended mappable genome size (the default values were used for the other parameters).
NarrowPeak mode was used for NFAT1, H3K27me3, and H3K4me3. Binding signal was
normalized by scaling to 1 million mapped reads using BEDTools (version 2.27.1)(*28*) and

bedGraphToBigWig (version 377)(*29*) and visualized as heatmaps using deepTools plotHeatmap

(version 3.2.1)(*26*)

In the CUT&RUN experiment with spike-in E. coli, two modifications were made: i) a hybrid

reference of mouse mm10 and E. coli ASM584v2, and ii) signals were normalized by scaling to

per million reads mapped to E. coli.

Differential binding analysis was performed using DiffBind (v.3.4.7) (summit = 75 (for NFAT1) and

100 (for H3K27me3 and K3K4me3 modifications), normalized by the trimmed mean of M values

(TMM) approach library size estimated by reads mapped to "background" regions. Peaks were

annotated to the nearest genes with the annotatePeak function in R package ChIPseeker

(v.1.30.3) at the gene level using default parameters. Genes were ranked by fold-change (log2

transformed; if a gene was associated with multiple peaks, the mean was used). GSEA was then

performed against MSigDB collections with R package clusterProfiler (v.4.2.2).

RNA-seq

The isolated CD8$^+$ TIL were washed once with cold PBS and pelleted at 1500 rpm at 4°C. Total

RNA was isolated using the Direct-zol RNA Microprep Kit (cat. R2061, Zymo Research) according

to the manufacturer's instructions and quantified using an Agilent 4200 TapeStation. Libraries

were prepared using the KAPA RNA HyperPrep Kit with RiboErase (HMR) (cat. 08098131702,

Roche) and purified by AMPure SPRI beads (cat. B23318, Beckman-Coulter). Libraries were

quantified and size distribution was determined using a Agilent 4200 TapeStation before paired-

end sequencing was performed.

RNA-seq data processing

Paired-end sequencing reads were mapped by the pipeline of St. Jude Center for Applied

Bioinformatics.   The   reads   were   trimmed   using   Trim   Galore   (version   0.5.0)

(https://github.com/FelixKrueger/TrimGalore) with default parameters. Reads were aligned to

the reference mouse mm10 assembly plus ERCC spike-in sequences using STAR (version

2.7.5a)(*36*). The resulting alignments, recorded in BAM file, were sorted, indexed, and marked

for duplicates with Picard MarkDuplicates function (version 2.19.0). Transcript quantification was

calculated using RSEM(*37*). Differential gene expression analysis was carried out with DESeq2

with Wald test (default parameters)(*38*). To examine if global changes in gene expression were

present, the RUVg function in the RUVSeq package was used for normalization with ERCC spike-

in followed by DESeq2 according to the RUVSeq manual(*39*). Gene set enrichment analysis (GSEA)

was performed against the MSigDB database(*40*) with R package clusterProfiler (version

4.2.2)(*34*) with the genes ranked by the Wald statistic from DESeq2 analysis.

1429

Quantitative Proteomics by tandem mass tag mass spectrometry (TMT-MS)

CD8$^+$ T cells were isolated from WT mice and activated with anti-CD3 antibody by crosslinking

with IgG for 30 min. Activated and untreated cells were washed once with ice-cold PBS and flash-

frozen in liquid nitrogen. Proteomic profiling of the whole proteome and methylome was carried

out using a previously optimized protocol(*41*) with modifications. Briefly, the cells were lysed in

8M Urea lysis buffer using pulse sonication, and approximately 100 μg of protein per sample was

digested with Lys-C (Wako, 1:100 w/w) at 21 °C for 2 h, followed by dilution to reduce urea to

2M, and further digestion with trypsin (Promega, 1:50 w/w) at 21 °C overnight. The protein

digests were acidified (trifluoroacetic acid to 1%), centrifuged at 21,000 x g for 10 min at 4°C to

remove any insoluble material, desalted with Sep-Pak C18 cartridge (Waters), and dried by

Speedvac. Each sample was resuspended in 50 mM HEPES (pH 8.5), TMT labeled, mixed equally,

desalted, and fractionated by offline HPLC (Agilent 1220) using basic-pH reverse phase LC (Waters

XBridge C18 column, 3.5 μm particle size, 4.6 mm x 25 cm, 180 min gradient, 80 fractions).

Fractions were aliquoted (5% v/v) for whole proteome and methylome (95% v/v) profiling. The

methylome fractions were further concatenated into 10 fractions for antibody-based sequential

mono-methyl-arginine and di-methyl-arginine peptide enrichment.

1446

Antibody-based Sequential Methylome Enrichment

Each concatenated fraction (approximately 1 mg) was resuspended in 400 μL ice-cold IAP buffer

(50mM MOPS, pH 7.2, 10 mM sodium phosphate, and 50 mM NaCl) and centrifuged at 21,000 ×

g for 10 min at 4°C to remove any insoluble material. The peptides were then incubated with

mono-methyl-Arginine antibody beads (mono-me-R, 12235S Cell Signaling Technology) at an

antibody-to-peptide ratio of 1:20 (w/w, optimized through a pilot experiment) for 2 hr at 4°C with

gentle end-over-end rotation. The antibody beads were then collected by a brief centrifugation,

washed 3 times with 1 mL ice-cold IAP buffer and twice with 1 mL ice-cold PBS, while the

supernatants were carefully removed and used for the sequential di-methyl-Arginine peptide

enrichment (di-me-R symmetric, cat.13563S Cell Signaling Technology). Peptides were eluted

from the beads twice at room temperature with 50 µL of 0.15% TFA, dried, and analyzed by LC-

MS/MS.

1459

LC-MS/MS Analysis

Each sample was dried, reconstituted, and analyzed by LC-MS/MS (a CoAnn 75 µm x 30 cm

column, packed with 1.9 µm C18 resin from Dr. Maisch GmbH) interfaced with the Orbitrap

Fusion MS (Thermo Fisher). LC settings included an 80 min gradient of 15%-40% buffer B (70%

ACN, 2.5% DMSO, and 0.1% FA) with buffer A (2.5% DMSO, and 0.1% FA) at a flow rate of ~0.25

µL /min flow rate. MS settings included data-dependent (3 s cycle) mode with a survey scan in

the Orbitrap (60,000 resolution, scan range 410–1600 m/z, 1×10^6 AGC target, and 50 ms

maximal ion time), followed by sequential isolation of abundant ions in a 3 s duty cycle, with

fragmentation by higher-energy collisional dissociation (HCD, 38 normalized collision energy),

and high-resolution detection of MS/MS ions in the Orbitrap (60,000 resolution, 1×10^5 AGC

target, 105 ms maximal ion time,1.0 m/z isolation window, and 20 s dynamic exclusion).

1471

Database Search and TMT Quantification

The MS/MS spectra were searched against the UniProt Mouse protein database (version

2020.04.22) using the COMET algorithm (v2018.013(*42*) **)** with the JUMP software suite(*43*)**.**

Search parameters included MS1 mass tolerance of 20 ppm and MS/MS of 0.02 Da, fully tryptic,

static mass shift for the TMT16 tags (+304.2071453) and carbamidomethyl modification of

57.02146 on cysteine, dynamic mass shift for Met oxidation (+15.99491), for mono-methyl-

Arginine (+14.01565), and for di-methyl-Arginine (+28.0313), maximal missed cleavage (n = 3),

and maximal dynamic modifications per peptide (n=5). All matched MS/MS spectra were filtered

by mass accuracy and matching scores to reduce false discovery rate (FDR) below 1% for either

proteins (whole proteome analysis) or peptides (methylome analysis), based on the target-decoy

strategy(*44, 45*).

1483

TMT quantification analysis was performed as previously reported(*46*) with the following

modifications: (i) extracting reporter ion intensities from each PSM; (ii) correcting the intensities

according to the isotopic distribution of each TMT reagent; (iii) removing PSMs of very low

intensities (e.g., minimum value of 1,000 and median value of 5,000); (iv) normalizing sample

loading bias with the trimmed median intensity of all PSMs; (v) calculating the mean-centered

intensities across samples (e.g., relative intensities between each sample and the mean), (vi)

summarizing protein relative intensities by averaging related PSMs; (vii) finally deriving protein

absolute intensities by multiplying the relative intensities by the grand-mean of the three most

highly abundant PSMs. (vii) Protein fold change and p-values of different comparisons were

calculated based on the protein intensities, using a log transformation and moderated t-test with

the limma R package(*47, 48*).

1495

Quantification and statistical analysis

Data were plotted and analysed using GraphPad Prism (GraphPad Software, v.9.2.0). Statistical

significance was calculated using unpaired one- or two-tailed Student's *t*-tests. Two-way ANOVA

was performed to compare tumor growth curves. The log-rank (Mantel–Cox) test was performed

to compare the mouse survival curves. For Fig. 4A, a linear mixed effects model with log10(y) as

the response variable, group as a fixed-effect predictor variable, and mouse as a random-effect

predictor variable was fit to the data. This model represents data with cells as random samples

from each mouse. The results showed that the mean log10(y) values of the two mouse groups

differed by -0.131 units (95% CI: -0.163, -0.0999; p = 6.98 10^{-15}). Statistical significance was set at

P < 0.05. Data are presented as mean ± SEM. No statistical methods were used to determine

sample size.

1507