Phytoplasma Targeting of MADS-Box Factor SVP Suppresses Leaf Responses to Insect Vector Males, Promoting Female Attraction and Colonization.

Authors: Zigmunds Orlovskis[1*¤], Archana Singh[1¤], Adi Kliot[1¤], Weijie Huang[1¤] and Saskia A. Hogenhout[1*]

[1] John Innes Centre, Norwich Research Park, NR4 7UH, United Kingdom

*correspondence: zigmunds.orlovskis@biomed.lu.lv; saskia.hogenhout@jic.ac.uk

[¤] current addresses:

ZO, Latvian Biomedical Research and Study Centre, Rātsupītes 1k-1, Rīga, LV-1067, Latvia.

AS, Department of Plant Sciences, University of Cambridge, Downing Street, Cambridge, CB2 3EA, UK.

AK, Department of Entomology, ARO, the Volcani Center, HaMaccabim Road 68, PO Box 15159, Rishon LeZion 7505101, Israel.

WH, Key Laboratory of Plant Molecular Genetics, CAS Center for Excellence in Molecular Plant Sciences, Shanghai Institute of Plant Physiology & Ecology, Chinese Academy of Sciences, Shanghai 200032, China.

OrchidIDs:

ZO, 0000-0001-8583-9814

AS, 0000-0002-5027-4582

AK, 0000-0002-0370-0817

WH, 0000-0001-5208-6713

SH, 0000-0003-1371-5606

Keywords:

Phytoplasma, bacteria, mollicutes, plant pathogen, obligate intracellular parasites, effector protein, SAP54/PHYL1, extended phenotype, host modulation, plant development, plant immunity, biotic stress response, MADS-box transcription factor, insect vector, leafhopper, *SHORT VEGETATIVE PHASE, SVP.*

Abstract

Obligate parasites often trigger significant changes in their hosts to facilitate transmission to new hosts. The molecular mechanisms behind these hyper-extended phenotypes - where genetic information of the parasite modulates the biology of one host to facilitate transmission through another host - remain largely unclear. This study explores the role of the virulence protein SAP54, produced by parasitic phytoplasmas, in attracting leafhopper vectors. SAP54 is responsible for the induction of leaf-like flowers in phytoplasma-infected plants. However, we previously demonstrated that the insects were attracted to leaves and the leaf-like flowers were not required. Here we made the surprising discovery that leaf exposure to leafhopper males is required for the attraction phenotype, suggesting a leaf response that distinguishes leafhopper sex in the presence of SAP54. In contrast, this phytoplasma effector alongside leafhopper females discourages further female colonization. We demonstrate that SAP54 effectively suppresses biotic stress response pathways in leaves exposed to the males. Critically, the host plant MADS-box transcription factor SHORT VEGETATIVE PHASE (SVP) emerges as a key element in the female leafhopper preference for plants exposed to males, with SAP54 promoting the degradation of SVP. This preference extends to female colonization of male-exposed *svp null* mutant plants over those not exposed to males. Our research underscores the dual role of the phytoplasma effector SAP54 in host development alteration and vector attraction - integral to the phytoplasma life cycle. Importantly, we clarify how SAP54, by targeting SVP, heightens leaf vulnerability to leafhopper males, thus facilitating female attraction and subsequent plant colonization by the insects. This study not only provides insights into the long reach of single parasite genes in extended phenotypes, but also opens avenues for understanding how transcription factors that regulate plant developmental processes intersect with and influence plant-insect interactions.

Introduction

Parasites often excel in altering the development and behaviour of their hosts, a trait essential for their survival and propagation. The trait is particularly pronounced in obligate parasites, which have substantial control over their hosts, earning them the nickname 'puppet masters'. The phenomenon exemplifies the concept of extended phenotypes (originally coined by Dawkins, 1982), where the impact of an organism's genes extends beyond its own physical form and affects other organisms. This concept is especially evident in obligate parasites that depend on alternate hosts for transmission, as they often do not only alter the conditions of their immediate hosts but also have far-reaching effects on other organisms in the ecosystem.

For instance, the protozoan parasite *Taxoplasma gondii* changes the behaviour of their intermediate rodent hosts by reducing their innate fear for cats, which are the definite hosts where the parasite undergoes sexual reproduction to produce oocysts (Tong *et al.*, 2021). Similarly, the rust fungus *Puccinia monoica* induces its host plant, *Boechera stricta*, to produce 'pseudoflowers'. These structures mimic real flowers and attract pollinators with their scent and sugary rewards, an essential strategy for the fungus to spread its spores between plants (Roy, 1993; Cano *et al.*, 2013). Although advancements have been made in identifying parasite virulence factors, our understanding of the specific host processes that are commandeered to produce these extended phenotypes is still developing.

Parallel to these discoveries are the studies on the extended phenotypes of phytoplasmas, which have revealed that their manipulative abilities stem from particular genes within these pathogens (Sugio *et al.*, 2011b; Tomkins *et al.*, 2018; Huang *et al.*, 2020; Wang *et al.*, 2024). Phytoplasmas cause disease in crops, ornamentals and native plants worldwide (Kumari *et al.*, 2019) and are often dependent on sap-feeding insects, including leafhoppers, plant hoppers and psyllids, for transmission (Weintraub & Beanland, 2006). As obligate bacterial parasites, phytoplasmas frequently trigger the emergence of unusual plant structures such as leaf-like floral parts (phyllody) and the excessive growth and clustering of leaves and branches (witch's broom) (Lee *et al.*, 2000; Al-Subhi *et al.*, 2018; Kumari *et al.*, 2019). These alterations not only compromise plant health but also promote attraction and colonization of insect vectors that are primarily responsible for phytoplasma spread and transmission (Sugio *et al.*, 2011a; Frost *et al.*, 2013;

MacLean *et al.*, 2014; Orlovskis & Hogenhout, 2016; Clements *et al.*, 2021; Al-Subhi

et al., 2021; Huang & Hogenhout, 2022). The responsible phytoplasma genes

encode for effector molecules that, once inside the plant cell cytoplasm, target and

typically disrupt or degrade essential plant transcription factors involved in growth,

development, and defence (review by Wang *et al.*, 2024 and references therein as

well as Liu *et al.*, 2023; Suzuki *et al.*, 2024; Correa Marrero *et al.*, 2024; Yan *et al.*,

2024; Zhang *et al.*, 2024). This molecular interference exemplifies the extended

phenotype reach, affecting not just the host appearance but also its physiological

and biological processes.

The interaction between the Aster Yellows strain Witches Broom (AY-WB)

phytoplasma and its vector, the aster leafhopper *Macrosteles quadrilineatus*, offers

insights into the complex interplay between parasites and hosts. The effector protein

secreted AY-WB protein (SAP) 11 binds to and destabilizes class II TCP

transcription factors - this action leads to changes in leaf shapes and stem

proliferation, reminiscent of witches' brooms, and altered root architecture

resembling the hairy roots found in infected plants (Bai *et al.*, 2009; Sugio *et al.*,

2011a, 2014; Lu *et al.*, 2014; Chang *et al.*, 2018; Pecher *et al.*, 2019). Additionally,

this effector reduces plant jasmonic acid and salicylic acid-mediated defence

responses, alters volatile organic compounds, and promotes the reproduction rates

of the insect vectors (Sugio *et al.*, 2011a; Lu *et al.*, 2014; Tan *et al.*, 2016). Another

effector, SAP05, recruits the 26S proteasome component RPN10, instigating the

breakdown of SPL and GATA transcription factors (Huang *et al.*, 2021; Liu *et al.*,

2023; Yan *et al.*, 2024; Zhang *et al.*, 2024). This process is linked to the typical leaf

and stem proliferations of witches' brooms in phytoplasma-infected plants. The

SAP05-mediated degradation of SPLs, but not GATAs, increases the attractiveness

of these plants to insect vectors (Huang *et al.*, 2021).

However, the relationship between phytoplasma SAP54/PHYL1 effectors

inducing leaf-like flowers and attracting insects seems more intricate. These

effectors target and disrupt MADS-box transcription factors, akin to animal HOX

genes, leading to a transformation of flowers into leaf-like structures, alterations that

mirror the phyllody and virescence symptoms of phytoplasma infection (MacLean et

al., 2011; MacLean et al., 2014; Wang *et al.*, 2024; Suzuki et al., 2024; Correa

Marrero et al., 2024). SAP54 recruits plant 26S proteasome shuttle factors called

RADIATION SENSITIVE 23 (RAD23) to break down these transcription factors

(MacLean et al., 2014; Kitazawa et al., 2022; Suzuki et al., 2024). Leafhoppers
prefer plants that stably produce SAP54 for reproduction and this preference
phenotype requires the presence of RAD23, which is essential for both breaking
down MADS-box transcription factors and inducing leaf-like flowers (MacLean et al.,
2014). However, the increased reproduction of insect vectors does not depend on
the presence of leaf-like flowers. Our past research showed that insects prefer
SAP54 plants during the vegetative developmental phase, before flowers emerge,
and even flowering SAP54 plants upon removal of the leaf-like flowers (Orlovskis &
Hogenhout, 2016). Therefore, it remains unclear whether plant MADS-box
transcription factors play a role in the leafhopper attraction phenotype.

MADS-box transcription factors have been predominantly studied in relation to
regulation of floral transition and flower organ development. Nonetheless, factors like
SUPPRESSOR OF OVEREXPRESSION OF CO 1 (SOC1) and short vegetative
phase (SVP) are also expressed in leaves and other organs during the *A. thaliana*
vegetative developmental phase, and regulate processes other than orchestrating
the flowering process, such as plant defence responses, as has been shown for
SOC1 (Li *et al.*, 2020). Based on these insights, we hypothesize that leafhopper
attraction is influenced by MADS-box transcription factors involved in processes
other than flowering.

Here, we analysed the factors influencing leafhopper colonization preference
in SAP54 plants. Surprisingly, we found that colonization preference on SAP54
plants only occurs in the presence of males on leaves during choice tests. Both
SAP54 and males are essential for female preference, as females did not exhibit a
preference for male-exposed versus non-exposed control plants, nor for SAP54
plants versus control plants in the absence of males. In contrast, female-only
presence on SAP54 plants deters colonization by other females. We noted a clear
downregulation of biotic stress response pathways in male-exposed SAP54 leaves,
contrasting with the upregulation observed in female-exposed SAP54 leaves and
control plants exposed to both males and females. Furthermore, we found that the
MADS-box transcription factor SVP is essential for female preference in male-
exposed SAP54 plants. Females also prefer colonizing male-exposed *svp null*
mutant plants over non-exposed ones. Our research sheds light on how the

conserved MADS-box transcription factor SVP influences leaf susceptibility to male
herbivorous insects, promoting female attraction and colonization.

163
164

Results

Attraction of SAP54 plants to fecund female leafhoppers relies on the
simultaneous presence of leafhopper males on these plants.

Given our previous finding that SAP54 promotes fecundity of the AY-WB
phytoplasma leafhopper vector *M. quadrilineatus*, we wished to further investigate
what aspect of the insect-plant interaction is affected by SAP54. Females feed and
lay eggs, while males only feed, and it is known that plants can induce different plant
defence responses to insect feeding and egg laying (Little *et al.*, 2007). Hence, the
first experiment was to assess the SAP54 effect on males and females separately
using choice tests. In these tests, the insects were given a choice to feed and lay
eggs on 35S:SAP54 or 35S:GFP transgenic plants (henceforth referred to as SAP54
and GFP plants) for 5 days, then adult leafhoppers were removed. Immediately after,
plants from each choice test were individually caged to avoid the hatching nymphs
from moving between plants. The number of progeny (nymphs) were counted 14
days later. When these choice tests were done with both males and females, the
leafhoppers produced more progeny on SAP54 than on GFP plants (**Figure 1A, B;**
treatment 1; Figure supplement 1), confirming previous findings (MacLean *et al.*,
2014; Orlovskis & Hogenhout, 2016). However, to our surprise, choice tests with
fecund females alone, without males, did not result in more leafhopper progeny on
the SAP54 versus GFP plants (**Figure 1A, B; treatment 2; Figure supplement 1**).
The females also produced more progeny on the SAP54 compared to GFP plants
when the males were caged and were not in direct contact with females (**Figure 1A,**
B; treatment 4; Figure supplement 1). In contrast, the females did not produce
more progeny on the SAP54 plants when males were left on the plant for only 48
hours and removed before the choice test with females alone (**Figure 1A, B;**
treatment 5; Figure supplement 1) or when the plants were exposed to the
conspecific females in the clip cages during fecund female-only choice tests (**Figure**
1A, B; treatment 6; Figure supplement 1). In the latter choice test, there was a

reproduction preference for the GFP plants instead. These results indicate that the presence of males on the leaves are required for the leafhopper preference to reproduce more on SAP54 versus GFP plants. Importantly, we found no obvious difference in progeny when fecund females were given a choice between (non-transgenic) wild type plants with and without males **(Figure supplement 2)**, indicating that SAP54 is required for the modulation of plant processes that results in the male-dependent reproduction preference of females.

Quantification of honeydew excretions provide a measure of how much phloem fluids the insects acquire from their host plant and therefore is a proxy for measuring feeding preferences (Hong & Rumei, 1993; Ammar *et al.*, 2013; Cameron *et al.*, 2014). Honeydew secretion tests showed that fecund females secreted more honeydew on SAP54 plants in the presence of males in clip-cages (**Figure 1A,C; treatment 4; Figure supplement 1**), whereas no increase in honeydew secretions of these females was observed on the SAP54 plants in absence of males (**Figure 1A,C; treatment 2; Figure supplement 1**) or when males were removed from the clip-cages prior to the choice test with females alone (**Figure 1A,C; treatment 5; Figure supplement 1**), indicating that females ingest a greater volume of plant sap from SAP54 than from GFP plants when males are present. There were slight increases in honeydew secretions on SAP54 versus GFP plants when the choice tests were done with both sexes, but these differences were not significant (**Figure 1A,C; treatment 1; Figure supplement 1**), in agreement with behaviour studies showing that female feeding is interrupted when in direct contact with males possibly due to mating (Beanland *et al.*, 1999). Females did not produce more honeydew on the SAP54 plants when other conspecific females were caged on the same plant (**Figure 1A,C; treatment 6; Figure supplement 1),** indicating their enhanced feeding activity is specifically dependent on the presence of males. Males alone did not secrete more honeydew on SAP54 versus GFP plants (**Figure 1A,C; treatment 3; Figure supplement 1**), indicating that the females are primarily responsible for the increased honeydew production on SAP54 plants.

Together, the fecundity and honeydew secretion data indicate that attraction of SAP54 plants to fecund female leafhoppers relies on the simultaneous presence of leafhopper males on the plants. The females produce more progeny and feed more on these plants only when males are present, even when the males are physically separated from the females.

Figure 1. *M. quadrilineatus* **leafhopper preference to reproduce and feed on SAP54**
versus GFP plants is dependent on leaf exposure to leafhopper males. A. Experimental
design of 6 choice tests (treatments) with 10 male and/or 10 female insects, as indicated, on
6 weeks old *A. thaliana* rosettes. Dashed circles indicate clip-cages and the arrow the
removal of males before the start of the choice test. Each choice test (treatment) is placed in
a separate cage. **B**. Percentages (%) of nymphs found on SAP54 versus GFP plants.
Horizontal bars in B, C indicate the mean ± 1 SEM. *p < 0.05, ***p<0,001. The entire series
of choice tests 1 to 6 were performed in parallel and repeated independently 3 times for
progeny count and 2 times for honeydew quantification - data presented in B and C include
the pooled results of the independent choice test series.

The online version of this article includes the following source data and figure supplement(s)
for figure 1:

Figure supplement 1: Data distributions of independent repeats that were used to generate
graphs displayed in B and C.

Figure supplement 2: *Macrosteles quadrilineatus* female leafhoppers show no preference
for *A. thaliana* Col-0 wild-type plants exposed to conspecific male leafhoppers.

Figure supplement 3: Female *M. quadrilineatus* preference for male-exposed SAP54 plants
is unlikely to involve long-distance cues.

Female leafhopper preference for male-exposed SAP54 plants unlikely
involves long-distance cues.

To investigate if the fecund females are attracted to the SAP54 plants via
volatile cues released by male-exposed SAP54 plants or mating calls from males
that may be perceived by females from a distance, we established a sticky-trap
assay that capture females before they access the plants themselves. Interestingly,
female leafhoppers were equally likely captured on GFP and SAP54 plants when
both of these plants were placed in an odour and sound permeable black container,

equipped with either transparent or green sticky traps (**Figure supplement 3A, choice tests 1 and 2**), suggesting that volatile or acoustic cues emitted from male-exposed GFP or SAP54 plants do not bias female landing choice at a distance when identical visual cues are present. To test whether the assay works, we compared different colour sticky traps as a positive control for insect visual choice. By doing so, female leafhoppers were more likely to be captured on green versus transparent sticky traps placed over the black containers, regardless of whether the green sticky traps were placed over male exposed GFP or SAP54 plants (**Figure supplement 3A, choice tests 3 and 4**), indicating that visual cues from the green traps were the primary determinant for female landing choice. Female insects also did not show an obvious landing preference for wild-type plants colonized by males compared to insect-free plants (**Figure supplement 3B)**. These results indicate that female choice for SAP54 versus GFP plants does not appear to involve plant volatiles that are released by the SAP54 plants in the absence or presence of males on the leaves and that the females do not appear to respond to potential mating calls of the males that may be perceived by the females from a distance. Therefore, these results indicate that female preference for the male-exposed SAP54 versus GFP plants is dependent on immediate access of the females to the leaves of SAP54 plants.

SAP54 plants display a dramatic altered transcriptional response specifically to male leafhoppers.

To further investigate how the combination of SAP54 and male-exposure may affect the preference of fecund females to reproduce on SAP54 versus GFP plants, we conducted RNA-seq experiments and compared the transcriptional responses of leaves of SAP54 and GFP plants that were exposed to five individual *M. quadrilineatus* females or males in clip cages or to clip cages without insects (controls). Each treatment consisted of 4 biological replicates (**Figure supplement 4A**). After normalization, transcripts with coverage of <1 FPKM in all replicates in at least one of the treatments were removed, leaving 16'307 genes (**Figure supplement 4B**). In a separate analysis, the data were analysed for differentially expressed genes (DEGs) that display significant changes in transcription levels between any two treatments, amounting to 6'957 DEGs (**Figure supplement 4B**). Considering both categories, a total of 17'153 genes were identified

(**Supplementary file 1**) of which 6'111 genes were shared between the categories (**Figure supplement 4B**). Based on these 17'153 genes, we performed multiple discriminant analysis (MDA) to visualize the separation between the treatment groups. While most treatments largely clustered together, SAP54_MALE_2 and GFP_MALE_2 appear to be switched based on the groupings of the other 3 replicates in the SAP54_male and GFP_male treatments (**Figure supplement 4C**). Analyses of the normalized reads that map to GFP versus GFP-SAP54 ruled out that the plant genotypes were accidentally swapped between the SAP54_MALE_2 and GFP_MALE_2 treatments or that the GFP or GFP-SAP54 were not expressed in the leaves (**Figure supplement 4D**). Moreover, plots of the median FRKM of all transcripts versus the GFP or GFP-SAP54 read counts did not flag the SAP54_MALE_2 and GFP_MALE_2 treatments as different from the others (**Figure supplement 4E**). Therefore, the apparent SAP54_MALE_2 and GFP_MALE_2 outliers may represent inherent variation of the biological system. Nonetheless, the MDA plot shows that in at least three out of four replicates per treatment, the leaf transcriptional responses are clearly distinct in male-treated SAP54 plants versus GFP plants, and more so than for the non-exposed and female-exposed SAP54 plants versus GFP plants (**Figure supplement 4C**).

When all samples were considered, a total of 136 DEGs were identified (**Figure supplement 5A**). Of these, 79 DEGs were derived from male-exposed leaves of SAP54 plants versus those of GFP plants and 69 DEGs from the non-exposed leaves of SAP54 versus those of GFP plants, with 12 DEGs overlapping between these two categories, whereas no DEGs were identified from female-exposed leaves of SAP54 versus those of GFP plants (**Figure supplement 5A**). This suggests that SAP54 has a bigger impact on leaf responses to males than to females. This was corroborated with the PCA analysis showing that insect exposure, male or female, of leaves of either SAP54 or GFP plants explained 54% of the variance (PC1), whereas the male-exposed SAP54 leaves are predominantly responsible for the 25% variance of PC2 among the insect-exposed treatments (**Figure supplement 5B**). The impact of SAP54 on male-exposed leaves was more obvious when three outliers were removed (**Figure supplement 5C,D**). Because the goal of this experiment was to assess how leaf responses of male-exposed plants differ from other treatments, we excluded these outliers, as including these obscures the transcriptional signal derived from the other three samples.

To further assess how SAP54 modulates leaf responses specifically to males,
we compared DEGs between male or female exposed leaves of GFP versus those
of SAP54 plants (**Figure 2A**; **Supplementary file 2**). The leaf response of GFP
plants to female exposure amounted to 2'375 DEGs as opposed to 909 male-
responsive DEGs of which the vast majority (894 DEGs) overlap with those of males
in the GFP plants (**Figure 2A**; **Supplementary file 2, table tabs A-B**). A higher
transcriptional response to females may be expected, because whereas both
females and males feed, females also lay eggs on leaves, and plants induce defence
responses associated with egg laying of insects (Stahl *et al.,* 2020; Little *et al.*,
2007). In contrast, leaves of SAP54 plants display dramatically different
transcriptional response to the insects, and with 2'456 male-specific and only 571
female-specific DEGs, the leaves particularly exhibit an altered transcriptional
response to leafhopper males (**Figure 2B; Figure supplement 6A; Supplementary**
file 2, table tabs C-D). This is striking particularly given that only minimal
transcriptional differences (266 DEGs) are observed between leaves of SAP54 and
GFP plants that were not exposed to insects (**Figure supplement 6A;**
Supplementary file 2, table tab E). These data suggest that SAP54 modulates leaf
responses of plants that are challenged by leafhopper herbivory, particularly
leafhopper males.

SAP54 plants exposed to males display significant downregulation of biotic
stress responses.

To better understand how SAP54 may alter leaf responses to the insects, *A.*
thaliana genes were binned into functional groups and these groups were analysed
for enrichment of DEGs among treatments using the MapMan built-in Wilcoxon rank
sum test with Benjamini-Hochberg correction (Usadel *et al.*, 2005). Leaves of GFP
plants exposed to females show enrichment for DEGs in 30 functional groups
compared to those of non-exposed GFP plants, as opposed to only 2 groups in GFP
plants exposed to males (**Supplementary file 3**). However, strikingly, leaves of the
SAP54 plants exposed to males show enrichment of DEGs in 42 functional groups
compared to those of non-exposed SAP54 plants, as opposed to only 11 groups in
leaves of SAP54 plants exposed to females (**Supplementary file 3**). The majority of
the functional groups, and particularly in leaves of the male-exposed SAP54 plants,
included genes with known functions in biotic stress responses, hormone

metabolism, signalling, secondary metabolism and other plant defence-related processes (**Supplementary file 3**). Moreover, bins enriched for genes in respiration (glycolysis) and cell wall modification functions were uniquely enriched in the leaves of male-exposed SAP54 plants (**Supplementary file 3**). In the no-insect/cage-only SAP54 versus GFP treatments, only 1 functional group (biotic stress responses involving plant defensins) was significantly enriched in the leaves (**Supplementary file 3**) and the majority of genes were upregulated in leaves of the SAP54 plants (**Figure supplement 6B**). Therefore, SAP54 alone has a relatively minor impact on leaf responses. Visualization of the fold change of these genes in MapMan graphs focused on genes with roles in plant responses to pathogen and pest attack (Usadel *et al.*, 2005) show upregulation of the majority of biotic stress response DEGs in leaves of female- and male-exposed GFP and those of female-exposed SAP54 plants, as opposed to a prominent downregulation of the majority of DEGs in leaves of male-exposed SAP54 plants (**Figure 2C and 2D; Supplementary file 4**). Specifically, multiple genes with roles in abiotic stress, cell wall modification, proteolysis, respiratory burst, secondary metabolism and defence signalling (MAPKs) were downregulated in the male-exposed SAP54 leaves. Together these results indicate that SAP54 dramatically alters leaf biotic stress response to leafhoppers, and downregulates the majority of this response in the presence of males.

Figure 2. SAP54 plants display a dramatically altered leaf response to male leafhoppers by transcriptionally downregulating the majority of biotic stress and plant defence related processes. **A, B**. Euler-Venn diagrams illustrating DEGs in leaves of GFP plants (A) and SAP54 plants (B) exposed to female leafhoppers compared to male leafhoppers, versus leaves of plants in the control group (cage-only, non-exposed plants). **C, D**. MapMan diagrams of *A. thaliana* DEGs involved in biotic stress. Pathways are indicated and each square is a gene with red versus green shades illustrating the level of up- or downregulation, respectively, of the gene in insect-exposed versus cage-only, non-exposed leaves of GFP (**C**) and SAP54 (**D**) plants.

The online version of this article includes the following source data and figure supplement(s) for figure 2:

Figure supplement 4. Experimental design and selection of transcripts for downstream analysis.

Figure supplement 5. Biological variation and role of outliers in separation of treatments and identification of differentially expressed genes.

Figure supplement 6. The cage-only SAP54 vs cage-only GFP treatments show few biotic stress DEGs.

Supplementary file 1. FPKM and differential expression values of 17'153 differentially expressed genes (DEGs) included in the response analyses of plants to SAP54 vs GFP and male vs female leafhopper exposure.

Supplementary file 2. IDs and log2-fold changes of DEGs of male and female *M. quadrilineatus* leafhopper-exposed GFP and SAP54 plants compared to insect-free GFP plants.

Supplementary file 3. MapMan build-in functional bins enriched for DEGs in male and
female *M. quadrilineatus* leafhopper-exposed GFP and SAP54 plants compared to insect
free GFP plants.

Supplementary file 4. Source data for generating Fig. 2CD - Enrichment statistics of biotic
stress bins and fold-change of DEGs in each bin.

We wished to further characterise which components of defence pathways
were predominantly affected in male-exposed leaves of SAP54 plants. To achieve
this, we performed a *de-novo* manual curation of *A. thaliana* defence signalling
module, integrating published literature and public TAIR database annotations for
known families of membrane receptors & receptor-like kinases (Shiu & Bleecker,
2001), cytoplasmic receptors such as NLR proteins (Hofberger *et al.*, 2014; Kroj *et*
al., 2016; Sarris *et al.*, 2016), CDPK-SnRK superfamily (Hrabak *et al.*, 2003), MAP
kinases cascade (Asai *et al.*, 2002; Jonak, 2002) as well as known salicylic acid,
jasmonic acid and ethylene biosynthesis and signalling genes (van Verk, 2010).
These manually curated defence signalling modules are provided here as **Figure**
supplement 7 and **Supplementary file 5,** and were imported into the MapMan tool
for visualisation and DEG analyses (**Figure 3**). Genes for LRR cell surface receptors,
CC-NBS-LRRs, RLCKs, MAPKs, JA biosynthesis and ET biosynthesis and signalling
were largely down regulated in leaves of the male-exposed SAP54 plants compared
those of the other treatments (**Figure 3).** Among the DEG-enriched biotic stress or
defence signalling pathway bins, the one for LRR receptors is most enriched and
most of these transcripts were downregulated in male-exposed leaves of SAP54
plants compared to the other treatments (**Supplementary file 6)**.

Figure 3. Genes involved in *A. thaliana* defence responses are predominantly down regulated in male-exposed leaves of SAP54 plants. MapMan diagrams with the manually curated bins for plant cell-surface receptors, NLRs, RLCKs, MAPKs and hormone biosynthesis and signalling proteins involved in plant responses to biotic stress. The number of DEGs belonging to each entity are indicated as squares above or adjacent to each protein category with red versus green shades illustrating the level of up- or downregulation, respectively, of the gene in insect-exposed versus cage-only, non-exposed leaves as shown in the upper panel.

The online version of this article includes the following source data and figure supplement(s) for figure 3:

Figure supplement 7: Manually drawn MapMan image for defence signalling pathway visualisation.

Supplementary file 5: Manually curated and assigned defence signalling bins for MapMan import.

Supplementary file 6: Functional bins for manually annotated defence genes enriched for DEGs in male and female *M. quadrilineatus* leafhopper-exposed GFP and SAP54 plants compared to insect free GFP plants.

Given our finding that females prefer to reproduce on male-exposed SAP54 leaves over those of male-exposed GFP plants in the choice experiments, and don't prefer the female-exposed leaves of SAP54 plants (**Figure 1**), we also compared leaf transcriptional profiles of the male-exposed and female-exposed SAP54 versus GFP treatments. Leaves of SAP54 plants respond more to males (2'857 male-specific DEGs) than to females (179 female-specific DEGs) compared to those of GFP plants with 957 DEGs shared between the two exposures (**Figure 4A**). Strikingly, none of the functional categories (bins) are enriched for DEGs in the female exposure treatment, as opposed to 51 categories in the male exposure treatment (**Supplementary file 7**). The 957 shared DEGs do not show enrichment for functional categories, though removing the shared DEGs from the male dataset by considering only the 2'857 male-specific DEGs reduced the enriched categories for the male treatment to 41. Among the latter DEG-enriched categories were protein synthesis, signalling involving receptor kinases, cell wall modification, hormone metabolism, including those of jasmonate and ethylene, and secondary metabolism of defence compounds. MapMan graphs of biotic stress responses show downregulation of all 957 shared DEGs and the majority of 2'857 male-specific DEGs, and only few downregulated genes in the 179 female-specific DEGs (**Figure 4B; Supplementary file 8**). Among the biotic stress response genes, those that are significantly enriched and display the greatest fold-change among male-specific

DEGs in leaves of SAP54 plants encode cell surface receptors of LRR and WAKL family and MAP kinases, as well as those involved the ethylene synthesis and signalling, protein degradation and isoprenoid (relating to lignin, glucosinolate and flavonoid biosynthesis) pathways (**Figure 4B; Supplementary file 8**). In summary, protein translational processes and defence pathways predominantly downregulate in male-exposed leaves of SAP54 plants compared to those of GFP plants, and this response was not obvious in the female-exposed leaves. Therefore, leaf defence responses of SAP54 plants are downregulated specifically in response to male exposure and this could explain female preference for the male-exposed SAP54 plants.

Figure 4. Biotic stress response genes are predominantly downregulated in male-exposed SAP54 versus GFP plants. Plant biotic stress is among the most enriched bins with male-specific responses in SAP54 leaves.

The online version of this article includes the following source data and figure supplement(s) for figure 4:

Supplementary file 7: MapMan build-in functional bins enriched for DEGs in SAP54 versus GFP plants with or without exposure to male and female *M. quadrilineatus* leafhoppers.

Supplementary file 8: Biotic stress bins from MapMan build-in and manually curated defence signalling pathway bins enriched for DEGs in GFP and SAP54 plants with or without exposure to male and female *M. quadrilineatus* leafhoppers.

The MADS-box transcription factor SVP is required for female preference of male-exposed SAP54 plants.

MADS-box transcription factors (MTFs) are predominantly investigated for their involvement in regulation of flowering. However, how MTFs may affect leaf responses to biotic stress, such as leafhopper colonization, is less well investigated. SAP54 mediates the degradation of many MTFs via the 26S proteasome by recruiting RAD23 proteins, which are 26S proteasome shuttle factors, and RAD23 proteins are required for both MTF degradation and leafhopper preference for SAP54 plants (MacLean *et al.*, 2014a). Therefore, we wished to investigate if the SAP54-mediated degradation of specific MTFs is required for the female preference for male-exposed SAP54 plants. The 17'153 leaf transcripts identified earlier (**Figure supplement 4B**) were filtered for the presence of 107 known MTFs (de Folter *et al.*, 2005) identifying 20 MTFs that are expressed in *A. thaliana* leaves independent from the presence of SAP54 (**Supplementary file 9**). The majority of the MTFs show differential expression among the male or female-exposed or cage-only leaves of SAP54 and GFP plants, including SVP, SOC1 and several of the MAFs (**Supplementary file 9**). The most striking included MAF5, which showed substantial downregulation in the male-exposed versus cage-only leaf responses of GFP plants and substantial upregulation in leaves that had combined exposure to SAP54 and leafhoppers (**Supplementary file 9**). Others, such SVP and AGL24, were mostly downregulated in the presence of insects or SAP54 and upregulated in the combined presence of insects and SAP54 (**Supplementary file 9**). The MADS-box transcription factor genes had similar expression levels between female-exposed and male-exposed leaves of SAP54 vs GFP plants, except MAF4 that was substantially downregulated in leaves of male-exposed SAP54 vs GFP plants and substantially upregulated in female-exposed SAP54 vs GFP plants (**Supplementary file 9**). Therefore, MADS-box transcription factor genes are differentially expressed in leaves upon exposure to insects and/or presence of SAP54.

We and others previously found that SAP54 directly interacts with many
MADS-box transcription factors and mediates their degradation via recruiting RAD23
(MacLean *et al.*, 2014b; Kitazawa *et al.*, 2022; Suzuki *et al.*, 2024). Here we found
that SAP54 mediated the degradation of several MADS-box transcription factors for
which we observed corresponding genes to be expressed in leaves (**Figure 5A**).
SAP54 interacts directly with the majority of these (**Figure 5B**). However, it does not
appear to interact directly with SVP (**Figure 5B**). Nonetheless, SVP interacts with
other MTFs, including SOC1 and AGL6, which interact with SAP54 (**Figure 5B**) and
SVP forms heteromeric complexes with SOC1, FUL, AGL6, AGL24 and several
other MTFs (de Folter *et al.*, 2005; Gregis *et al.*, 2006, 2009; Balanzà *et al.*, 2014;
Mateos *et al.*, 2015). Moreover, SAP54 mediates degradation of SOC1 and other
MTFs via the 26S proteasome (MacLean *et al.*, 2014). Hence, SAP54-mediated
degradation of SVP could occur indirectly via SAP54 interactions with SOC1, AGL6
and other MADS-box transcription factors that form multimeric complexes with SVP.

Figure 5. SAP54 interacts with multiple MADS-box transcription factors and mediates
their degradation. A. Western blots showing degradation of MTFs in the presence of
SAP54 in *A. thaliana* protoplasts. **B**. Yeast two-hybrids assays with GAL4-activation (AD)
and GAL4-DNA binding (BD) domains fused to the test proteins. -L-W-H-A denote
auxotrophic SD media lacking leucine, tryptophan, histidine or adenine, conditionally
supplemented with 3-Amino-1,2,4-triazole (3AT). EV, empty vector control.

Next, we conducted leafhopper choice tests with *null* mutants for several
MADS-box transcription factor genes. These revealed that the leafhoppers preferred
to reproduce on the *svp* and *maf5* mutants, whereas no clear preferences of the
leafhoppers were observed for *A. thaliana agl24, sep4, maf1, maf4, ful* and *soc1*
mutants compared to wild-type plants (**Figure 6A; Figure supplement 8**). Moreover,
leafhopper egg-laying preference was abolished when SAP54 was introduced into
the *svp* mutant background, but not when introduced into the *maf5* mutant
background (**Figure 6B; Figure supplement 8**), suggesting that that plant SVP is
the dominant MADS-box transcription factor required for the SAP54-dependent
leafhopper preference. In parallel, choice tests were done with AY-WB-infected wild-
type and mutant plants, and the leafhoppers did not display the colonization
preference for both *svp* and *maf5* mutants (**Figure 6C; Figure supplement 8**),
indicating that AY-WB-infected wild type and mutant plants are equally attractive to
the leafhoppers. While females showed a preference for male-exposed *svp* and
maf5 mutants versus male-exposed wild-type plants (**Figure 6A**) this preference for
svp and *maf5* mutants was abolished in the absence of leafhopper males (**Figure
5D; Figure supplement 8**). Transcripts for *svp* and *maf5* were detected, and
upregulated, in leaves in the combined presence of SAP54 and leafhoppers (**Figure
5E; Supplementary file 9**). Together, these data provide evidence that female
preference for male-exposed SAP54 plants is dependent upon the presence of SVP,
while MAF5 plays a role in female choice independently from SAP54.

561
Figure 6. The MADS-box transcription factor SVP is required for female preference to
reproduce on male-exposed SAP54 plants. A-C. Choice test with equal numbers of 10
males and 10 females on wild type and MTF *null* mutant *A. thaliana* rosettes (**panel A**),
35S:GFP-SAP54 *A. thaliana* rosettes (**panel B**) and AY-WB phytoplasma-infected plants
(**panel C**). **D.** Choice tests with 10 females on *A. thaliana svp* and *maf5 null* mutants without
males. The entire series of choice tests depicted in panels **A-D** were conducted in parallel
and repeated independently 2 times. ** p<0.01, *** p<0.001. Bars are ±1 SEM. **E.**

Expression levels of *SVP and MAF5* in leaves among treatments showing that SVP and MAF5 are upregulated in male-exposed SAP54 plants.

The online version of this article includes the following source data and figure supplement(s) for figure 6:

Figure supplement 8: Data distributions of independent repeats that were used to generate graphs displayed A-D.

Supplemental file 9. Fold-expression changes of MADS-box transcription factor genes insect-exposed SAP54 vs GFP leaves.

SVP differentially regulates plant responses to herbivores.

Considering the role of SVP in leafhopper reproduction preference, we conducted transcriptional analyses to assess how leaves of *svp* mutant plants respond to male and female leafhopper exposure and compared this to male or female exposed SAP54 plants (**Figure supplement 9A)**. A comparable number of DEGs was found in leaves exposed to both males and females, with 464 DEGs uniquely responding to females, 347 to males, and 460 to both sexes (**Figure supplement 9B**). Many of biotic stress related DEGs in male or female exposed *svp* mutant were upregulated (**Figure supplement 9C**). When searching for functions that were enriched with the 924 DEGs in the female exposed *svp* mutant versus female exposed wild-type plants, we found significant enrichment for abiotic stress and cell wall genes (**Supplementary file 10)** as well as DUF26 family cell surface receptors (**Supplementary file 11**). Furthermore, the 807 DEGs in *svp* leaves exposed to males compared to male colonized wild-type plants were predominantly involved in sulphur-containing glucosinolates and cell wall associated proteins (**Supplementary files 10 and 11**). Hence, the DEGs linked to cell wall structure and modulation were prominently enriched in leaves exposed to both male and female leafhoppers. There was no clear enrichment of functional categories among the 347 male-specific DEGs in *svp* mutant (**Supplementary files 10 and 11**). These findings suggest that SVP influences leaf responses to leaf-colonizing insects in a manner that is partially specific to the gender of the leafhoppers.

We further wished to explore the differentially expressed genes (DEGs) that are common between the male-exposed leaves of SAP54 and *svp* mutant plants. Firstly, we analysed the overlap between male-specific (not up- or down-regulated by female) SAP54 expressing plants and the *svp* mutant, where we found only 49 DEGs (**Figure supplement 9D**). Many of these genes display similar up- or down-

regulation in SAP54 and *svp* plants, and some of the most downregulated DEGs include cytochrome P450s such as CYP81F, CYP78A as well as glutamate receptor GLR3 and calcium-binding transcription factor MYC5 (NIG1) (**Supplementary file 12, Table Tab A**).

However, due to the relatively small number (49) of shared DEGs for full pathway characterisation, we analysed functions encoded by all 155 shared DEGs in male exposed SAP54 and *svp* plants, including those that overlap with the female regulated DEGs in those plants (**Figure supplement 9E**). The 155 DEGs showed significant GO term enrichment associated with various biological responses, including responses to stimuli, oxygen-containing compounds, lipids, hormones, and calcium signalling (**Supplementary file 12, Table Tab B**), suggesting common functions in biotic stress regulation between SAP54 and SVP in male exposed plants. Moreover, we also examined the full spectrum of MapMan biological functions enriched by DEGs in leaves of male-exposed SAP54 (3814 DEGs) and *svp* mutant (807 DEGs) plants separately before looking for the overlap in functions (rather than gene identities first). This comparison revealed that the similarities between the enriched functions in male exposed SAP54 and *svp* mutant plants predominantly pertain to cell wall modification and secondary (isoprenoid) metabolism (**Supplementary file 12, Table, Tab C**). Considering that female insects are more attracted to the male-exposed leaves of SAP54 and *svp* mutant plants for feeding and oviposition, alterations in cell wall structure and changes in secondary (isoprenoid) metabolism may potentially play significant roles in influencing female preferences.

Discussion

Here, we investigated colonization preferences of the leafhopper vector *M. quadrilineatus*, the predominant vector of AY-WB phytoplasma, for plants that produce the AY-WB effector SAP54. We made the surprising discovery that female preference for SAP54 plants only occurs in the presence of males. Both SAP54 and males are necessary for this preference, as females did not show a preference in their absence. As well, SAP54 plants exposed to males display significant downregulation of biotic stress responses. We identified the MADS-box transcription

factor SVP to be essential for this female preference to male-exposed SAP54 plants.
Leafhopper females exhibited a preference for reproducing on *svp null* mutant
plants, a behaviour that is also contingent upon the presence of leafhopper males,
and aligning with our observation that SAP54 mediates the degradation of SVP in
leaves. Therefore, by characterizing a parasite effector, we revealed the crucial role
of SVP in leaf susceptibility to male herbivorous insects, enhancing female attraction
and colonization.

This work built on our previous findings that the leaf-like flowers are not
required for the insect colonization preference and the insect vectors are
predominantly attracted to the leaves and not flowers of plants (Orlovskis &
Hogenhout, 2016). Here we independently confirm these data via experiments that
show a preference of the insects for *A. thaliana* rosettes before the onset of
flowering. Therefore, a morphological change in the host may not always be
responsble for the extended phenotype effect on the alternative host, which is in this
case the insect vector. It is possible that the attraction of insect vectors to leaves is
the primary function of SAP54, and the induction of the leaf-like flowers an
evolutionary side-effect of SAP54 adaptation to target MTFs that leads to
degradation of SVP. However, the prevention of seed production through the
formation of leaf-like flowers likely is likely to present an advantage to phytoplasma.
AY-WB phytoplasma commonly infects annual plants, where leaves naturally
senesce and die off after flowering and seed production. It is probable that
prolonging the longevity of their plant hosts is advantageous for phytoplasmas,
particularly considering that they are not known to be seed-transmitted. Instead,
phytoplasmas rely on leafhoppers, which feed on vegetative tissues, for
transmission.

We found that SAP54 mediates the destabilization of SVP. However, we did
not find evidence of a direct interaction between SAP54 and SVP in our Y2H
analyses. An explanation of this apparent discrepancy is that MTF proteins exhibit a
propensity to form diverse complexes, including homo- and hetero-dimers, as well as
higher order ternary and quaternary complexes (Davies *et al.*, 1996; Egea-Cortines,
1999; Hartmann *et al.*, 2000; Honma & Goto, 2001; Immink *et al.*, 2002; de Folter *et
al.*, 2005). Notably, SVP is known to interact with other MTFs such as SOC1, FUL,
and AGL24 (de Folter *et al.*, 2005; Gregis *et al.*, 2009; Balanzà *et al.*, 2014), and we

previously established that SAP54 interacts with and facilitates the degradation of multiple MTFs, including SOC1, FUL, and AGL24 (MacLean *et al.*, 2014), as is also shown for FUL and AGL24 herein (**Figure 5**). Moreover, our research showed that these three MTFs are expressed in leaves, with SOC1 and AGL24 showing elevated expression in SAP54 plants exposed to males (**Supplementary file 9**). Hence, the destabilization of SVP by SAP54 may occur through its involvement with various SVP-containing protein complexes. This intricate interaction network also provides a possible explanation for the absence of a leafhopper preference phenotype in *soc1*, *agl24*, and *ful* mutant plants. It suggests that the interaction of SAP54 with multiple SVP-containing ternary and quaternary MTFs, leading to their degradation, might be essential to trigger the observed phenotype.

SVP is recognized as a conserved regulator of flowering time across a variety of eudicot species, as evidenced by several studies (Sun *et al.*, 2016; Li *et al.*, 2016). It is expressed in the leaves during the vegetative growth stages of plants (Zheng *et al.*, 2009; Li *et al.*, 2016; Wu *et al.*, 2017), which correlates with the wide host range of AY-WB phytoplasmas. While the primary role of SVP and its homologs has been associated with flowering, their involvement in additional biological processes beyond this remains underexplored (Liu et al., 2018). However, emerging research has begun to shed light on the roles of SVP outside of flowering: for instance, it acts as a positive regulator of age-related resistance against bacterial pathogens in *Arabidopsis thaliana* (Wilson *et al.*, 2017) and serves as a negative regulator of various JASMONATE ZIM-domain (JAZ) genes during the vegetative growth phase (Gregis *et al.*, 2013). Our findings contribute to this expanding body of knowledge, suggesting that plant immunity may be intricately controlled by developmental regulators like SVP, which possesses multiple, diverse functions throughout plant development (Berry & Argueso, 2022, and references therein). The observation that female leafhoppers prefer to reproduce on both SAP54 and *svp* mutant plants exclusively in the presence of leafhopper males underscores a sophisticated regulatory effect of SVP and other MADS-box transcription factors targeted by SAP54 on plant responses to the different sexes of the herbivores. This intricate interaction is detailed in the mechanism proposed in **Figure 7**.

Figure 7. Proposed model for SVP-dependent enhancement of insect vector colonisation of phytoplasma-infected host plants. SAP54 recruits proteasome shuttle protein RAD23 to degrade MTFs in 26S proteasome dependent manner (MacLean *et al.*, 2014). This destabilises SVP which is required for recruitment female leafhoppers to lay eggs on male-exposed leaves during vegetative growth phase. In addition, SAP54-induced perturbations in the functioning of MTF network leads to the generation of leaf-like flowers after the phase transition.

Considering that certain phytoplasmas can modify plant volatile production (Mayer *et al.*, 2008a,b; Rid *et al.*, 2016; Tan *et al.*, 2016), we examined whether male-induced volatile cues play a role in attracting females. Our results suggest that female leafhopper preference for laying eggs on male-exposed SAP54 plants is unlikely to depend on volatiles originating from the males or induced by them (**Figure supplement 3**). Thus, it is plausible that females are guided to male-occupied SAP54 and *svp* mutant plants through a different mechanism, possibly involving changes in leaf responses that are specifically induced to male insects. Male-exposed leaves of SAP54 and *svp* mutant plants shared transcriptional changes of genes involved cell wall structure processes and changes in secondary (isoprenoid) metabolism. As such, these could influence female leafhoppers' host plant selection and their egg-laying preferences on plants exposed to males.

Males and females of *M. quadrilineatus*, the primary vector of AY-WB phytoplasma, exhibit distinct movement and feeding behaviours, as documented by

Beanland et al. (1999). Moreover, the leafhoppers, especially females, live longer on aster yellows infected plants compared to non-infected plants (Beanland *et al.*, 2000). Here we show that the combination of the aster yellows phytoplasma effector SAP54 and males attract females in a manner that depends on SVP. Moreover, plant biotic stress and defence responses are largely suppressed in male (compared to female) exposed SAP54 plants, and SVP also displays distinct plant transcriptional responses to male and female leafhoppers. Nevertheless, the mechanisms how male and female leafhoppers, in combination with SAP54, differentially affect plant defence responses will need further investigation. While the differences in male and female behaviour may play a role (Beanland *et al.*, 1999), there is also the possibility that effectors in the saliva of sap-feeding insects modulate plant processes, including defence (Snoeck *et al.*, 2022), and sex-specific differences exist in the proteome of sap-feeding insect saliva (Miao *et al.*, 2018), potentially influencing plant response. Moreover, plant immune responses initiated by the molecular patterns associated with insect eggs (EAMPs) that bear similarities to pattern-triggered immunity (PTI) activated by bacterial pathogens (Gouhier-Darimont *et al.*, 2013) may play a role in the observed gender-specific differences.

Pathogens uniquely exploit sexual dimorphism in their hosts to enhance their dispersal and survival in nature. For example, the bacterial parasite infecting the crustacean *Daphnia magna* selectively uses male hosts for horizontal transmission, while exploiting females for internal replication and spore production, thus facilitating widespread disease transmission (Nørgaard *et al.*, 2019). Toll signalling pathway contributes to the varied resistance observed between male and female *Drosophila* to both Gram-positive and Gram-negative bacteria (Duneau *et al.*, 2017). Similarly, the endosymbiotic bacterium *Wolbachia* often exhibits distinct effects on the embryonic development of male and female gametes, showcasing another instance of pathogens leveraging host sexual differences (Serbus *et al.*, 2008). Recent discovery identified a novel *Wolbachia* factor Oscar which interacts with host protein Masculinizer to inhibit masculinization and lead to male-killing in two insect species (Katsuma *et al.*, 2022). Our study further expands the mechanistic understanding about insect gender specific factors targeted by parasites.

In our research, we have uncovered a mechanism whereby the phytoplasma effector SAP54 targets a conserved host regulator, SVP, to attract female leafhopper

vectors to produce more progeny on plants exposed to leafhopper males. Given that phytoplasmas rely on leafhoppers for acquisition during feeding and subsequent transmission to other plants, SAP54 likely promotes the abundance of phytoplasma-carrying insect vectors and facilitates phytoplasma transmission. This illustrates how a bacterial gene extends its influence beyond its primary plant host to modulate the host choice and reproductive behaviour of the insect vector. Such findings underscore the complexity of tripartite plant-insect-microbe interactions, wherein parasitic genes exhibit what could be termed 'hyper-extended phenotypes,' manipulating the biology of one host to facilitate transmission through another host. This research advances our understanding of plant-microbe interactions and resonates with the concept of the Extended Evolutionary Synthesis (EES), as articulated by Hunter (2018). Within the EES framework, parasitic genes like SAP54 evolve adaptively through both natural selection and direct biochemical interactions with plant regulators, as well as through the extended functions these regulators serve in interactions with insect vectors. In this study, we found that the extended role includes enhancing the likelihood of male vectors to attract females for reproduction.

Materials and methods

Generation of plants for insect choice experiments

Generation of 35S:GFP-SAP54 and 35S:GFP transgenic *Arabidopsis* lines was done according to methods described in (MacLean *et al.*, 2011, 2014). *ful-1* line was provided by Lars Ostergaard Lab and described in (Gu *et al.*, 1998). *soc-1* and *sep4-1* by Richard Immink Lab and described in (Immink *et al.*, 2012). *maf4-2 and maf5-3* seeds were provided by Hao You Lab and described in (Shen *et al.*, 2014). *maf1* (also known as *flm-3)* was provided by Claus Schwechheimer Lab and described in (Lutz *et al.*, 2015). *svp1-41* was provided by Martin Karter Lab and described in (Hartmann *et al.*, 2000). *agl24* (N595007, SALK_095007) was obtained from The Nottingham Arabidopsis Stock Centre (NASC). Phytoplasma-infected plants were generated identical to methods described in (MacLean *et al.*, 2014). All plants were grown under short days (10h/14h light/dark) identical to methods described in (Orlovskis & Hogenhout, 2016)

Insect oviposition choice assays

All insect choice experiments were performed identical to methods described in (Orlovskis & Hogenhout, 2016) with the following modifications. Female-only oviposition choice experiments described in **Fig. 1** (*experiment 2*) were set up by releasing 10 *M. quadrilineatus* females, while the mixed-sex choice assay (experiment 1) was done with 10 male and 10 female insects. Five male leafhoppers were confined on test and another five on control plants using transparent clip-cage (2 cm diameter) before the release of 10 females in *experiments 3* and *4* displayed in **Fig. 1**. For the *experiment 5* in **Fig. 1**, five females were restricted on test and five on control plants prior the release of another 10 females in the choice cage. Notably, single-sex insects were selected from mixed-sex population of *M. quadrilineatus* stock. Adults were removed 5 days after release, nymphs counted 14 d after adult removal. Nymph counts analyzed using paired t-test.

Insect olfactory choice assay

Test and control plants were placed in a black (non-transparent) plastic container 12cm x 10cm x 10cm (H x W x D) with a perforated topside, permitting diffusion of plant odors but rendering plants invisible from the outside of the container. A colourless or a coloured sticky landing platform (OECO, Kimpton, UK) was fitted on the top of each container but not covering the perforations. 20 *M. quadrilineatus* females were released in the center of the arena (transparent polycarbonate 62cm x 30cm x 41cm (H x W x D)) equal distance from the test and the control containers. When coming in contact with the sticky landing platform, insects stuck permanently to its surface. Within the following 1h under the ambient room light the majority of insects made their first landing choice, and were counted. Each experiment utilized clean choice arena, new containers and landing platforms to avoid any possible residual semiochemicals left by insects from the previous experiment which could bias the leafhopper choice in the next experiment. In selected experiments male leafhoppers were contained in clip-cages on the test plants placed inside the containers.

Western blotting

Proteins were separated on 12.5% (w/v) SDS-PAGE gels and transferred to 0.45 mm Protran BA85 nitrocellulose membranes (Whatman, UK) using the BioRad

minigel and blotting systems following standard protocols. Blotted membranes were
incubated in blocking buffer (5% w/v) milk powder in 1X phosphate buffered saline
and 0.1% (v/v) Tween-20) with primary antibody at 4°C overnight. Peroxidase-
conjugated anti-rabbit or anti-mouse secondary antibody (Sigma Aldrich) was added
to washed blots and incubated at room temperature for 4h. Bound antibodies were
detected using Immobilon Western Chemiluminescent HRP Substrate (Millipore, UK)
when exposed to Super RX film (Fujifilm, Germany). Protein loading was visualized
using Ponceau S solution (0.1% (w/v) in 5% acetic acid).

Yeast-two-hybrid assays and Protoplast assays

The Matchmaker Gold Yeast Two-Hybrid System (Clonetech) and the
DUALhybrid system (Dualsystems Biotech) were used for detecting protein-protein
interactions in yeasts. Yeast transformation were carried out following
manufacturers' instructions. Successful transformants grew on medium lacking
leucine and tryptophan (-L -W). Medium lacking leucine, tryptophan, histidine and
Adenine (-L-W-H-A) or Medium lacking leucine, tryptophan, histidine and
supplemented with 10 mM 3-AT (-L-W-H +10 mM 3-AT) were used to examine
protein-protein interactions. The growth of yeast colonies was scored after 5 days of
incubation at 28°C. Protoplast assays were carried out according to (Yoo *et al.*,
2007). Briefly, mesophyll protoplasts isolated from 4 weeks-old Arabidopsis leaves
were resuspended at 4 × 105 ml−1 in MMG solution. 300 μl protoplast solution were
then transfected with 24 μg plasmids (12 μg each for double transformation) using
PEG–calcium-mediated transfection. Protoplasts were kept at 22 °C for 16 hours in
dark and then harvested for western blot analysis.

Generation of plants for RNA-sequencing

The experiment used 8-weeks old *Arabidopsis thaliana* (Col-0) plants
ectopically expressing Aster Yellows phytoplasma strain Withes' Broom effector
SAP54 (35S:GFP-SAP54) or a control construct (35S:GFP). Plants were grown at
short day photoperiod (10h/14h light/dark). Single fully expanded leaf of each plant
was exposed to either 5 male or 5 female insects by placing them in a transparent
2cm diameter clip-cage which confines the leafhoppers to the leaf and prevents from
escaping. An empty clip-cage was placed on a leaf for no-insect control. In order to
prevent any insects from the room to land on the experimental plants and confound
the results, we placed all plants with clip-caged female leafhoppers in a sealed larger

cage separate from plants with clip-caged males or plants with empty clip-cages. Thus, all replicate plants with male, female or no-insect exposed leaves were maintained in similar micro-environment. The experiment with clip-caged males, females and empty clip-cages was done at the same time on the same shelf within the growth room to ensure similar conditions across all treatments. Plant tissue samples were collected 48h after exposure to insects and placed straight into liquid N_2. Insect number and exposure time of 48h was previously experimentally optimised by measuring number of feeding sites and eggs laid per clip-cage. The experiment with clip-caged male or female plants on *svp* mutant and *wild-type* Col-0 plants was performed at a different time than the experiment with 35S:GFP-SAP54 and 35S:GFP plants but with identical setup.

Total RNA extraction for RNA-seq and quality control

The collected leaf area enclosed by the clip-cage (~2 cm diam.) was stored at -80°C for subsequent RNA extraction using QIAGEN Plant RNeasy kit following the manufacturer's instructions. Initial RNA integrity was assessed by gel electrophoresis (1% Agarose) visualisation of ribosomal bands in the extracted dsRNA as well as ssRNA following 65°C denaturation and immediate transfer on ice to prevent hybridisation. Total RNA concentration and quality was assessed using Nanodrop (Thermofisher). Total ≥2µg of each RNA sample at ≥50ng/µL concentration, 260/280 ratio between 1.9 and 2.1, and 260/230 ratio between 1.5 and 2.0 was submitted for library preparation and RNA-sequencing.

RNA sequencing, read alignment and differential expression analysis

Experimental design consisted of 4 biological replicates for male- or female-exposed or insect-free transgenic 35S:GFP-SAP54 plants and transgenic 35S:GFP control plants as well as 3 biological replicates for *svp* null mutant and wild-type col-0 control plants. cDNA library construction for SAP54 and GFP was performed according to the IlluminaTruSeq protocol, followed by sequencing on Illumina HiSeq2000 platform and pooling 4 randomized libraries per lane, with 50bp single-end reads and 25M read coverage per sample, whereas *svp* null mutant and wild-type libraries were prepared with paired-end reads. Reads were trimmed for low quality and adapter contamination using Trim Galore! v0.4.0 with default settings (Krueger, 2012). Trimmed reads were aligned to *A. thaliana* reference genome

(TAIR10) by HISat2 (Pertea *et al.*, 2016). We identified differentially expressed
transcripts with DESeq2 package in R (version 1.2.10) (Love *et al.*, 2014) using
expression count generated by Kallisto v0.42.3 (Bray *et al.*, 2016). Independent
filtering was employed in in DESeq2 to remove very low expressing transcripts on
the basis of normalized counts. Genes were considered differentially expressed if
they had a *p* value less than 0.05 after accounting for a 5% FDR according to the
Benjamini-Hochberg procedure and if a fold-change in expression of at least 2 was
observed.

Functional analysis of transcriptome data

Enrichment of GO biological functions and GO molecular functions for DEGs
were calculated using Fisher exact test (PANTHER)
(http://geneontology.org/docs/go-enrichment-analysis). Graphical visualisation for
functional catagorisation of differentially regulated transcripts was performed using
MapMan 3.5.1R2 functional annotation tool (Thimm *et al.,* 2004). The tool,
supporting resources and annotation database available from
http://mapman.gabipd.org/web/guest/home. Statistical analysis on enrichment of
differentially changed pathways was performed using Wilcoxon rank test and
Benjamini-Hochberg (BH) p-value correction (Usadel, 2005). Methods for manual
creation of new pathways and categorical filtering of expression data were adopted
from (Usadel *et al.*, 2009) and explained in sections below. The graphical overview
of defence signalling cascades in **Figure 3** and *Figure supplement 7* were manually
drawn and loaded as separate pathway image files into MapMan. The mapping file
(**Supplementary file 5**) containing a list of gene identifiers and descriptions
manually assigned to new functional bins previously absent from the MapMan
annotation was generated from published literature and the public TAIR database
(www.arabidopsis.org/browse/genefamily) for membrane-located receptor-like
kinases and their classes (Shiu and Bleecker, 2001), cytoplasmic receptors such as
NLR proteins (Hofberger et al., 2014; Kroj *et al.,* 2016; Sarris *et al.*, 2016), CDPK-
SnRK superfamily (Hrabak *et al.,* 2003), MAP kinases cascade (Asai *et al.*, 2002;
Jonak *et al.,* 2002) as well as SA, JA and ET biosynthesis and signalling genes (van
Verk, 2010).

Data availability

RNA-seq raw data (fastq) are uploaded to NCBI under BioProject: PRJNA1090849.

Author contributions (The CRedit Roles and Number):

ZO: conceptualization (1), data curation (2) and formal analysis (3), investigation (4), methodology (5), data validation (11), statistical analysis and visualisation (12), writing, reviewing and editing the original manuscript (13,14)

WH: investigation (4), methodology (5), data visualisation (12), review of the original manuscript (14)

AK: data validation (11), review of the original manuscript (14)

AS: data curation (2), review of the original manuscript (14)

SH: conceptualization (1), data curation (2) and formal analyses (3), funding acquisition (4), supervision (10), data validation (11), writing, reviewing and editing the original manuscript (13,14).

Funding

This work was supported by Human Frontier Science Program grant RGP0024/2015 (to SH), the European Research Council and UK Research and Innovation (UKRI) Engineering and Physical Sciences Research Council grant EP/X024415/1 (to SH), BBSRC (BBS/E/J/000PR9797 and BBS/E/JI/230001B), and a BBSRC student fellowship (to ZO).

Acknowledgements

We thank Lars Ostergaard (JIC) for providing *ful-1* seeds, Richard Immink (University of Wageningen) for giving *soc-1* and *sep4-1* seeds, Hao Yu (Zhejiang University) for *maf4-2* and *maf5-3* seeds, Claus Schwechheimer for *maf1* seeds (Technical University of Munich), and Martin Kater (University of Milan) for *svp1-41* seeds. We thank the JIC Entomology and Insectary Platform staff for maintaining the leafhopper and phytoplasma stocks and the JIC Horticultural services staff for growth and maintenance of the plants used in this study. We also thank Sam Mugford (JIC) for technical assistance throughout the project. Allyson MacLean (now University of

Ottawa), Vera Thole, Enrico Coen (JIC) deserve personal gratitude for their advice and discussions as members of Z. Orlovskis PhD thesis supervisory team. We are grateful to Robert Sablowski (JIC) and Sebastian Schornack (University of Cambridge) for constructive feedback on thesis work that helped to conceptualize the study herein.

Competing interest: The authors declare that no competing interests exist.

Literature Cited

Al-Subhi AM, Al-Sadi AM, Al-Yahyai RA, Chen Y, Mathers T, Orlovskis Z, Moro G, Mugford S, Al-Hashmi KS, Hogenhout SA. **2021**. Witches' Broom Disease of Lime Contributes to Phytoplasma Epidemics and Attracts Insect Vectors. *Plant Disease* **105**: 2637–2648.

Al-Subhi AM, Hogenhout SA, Al-Yahyai RA, Al-Sadi AM. **2018**. Detection, Identification, and Molecular Characterization of the 16SrII-D Phytoplasmas Infecting Vegetable and Field Crops in Oman. *Plant Disease* **102**: 576–588.

Ammar E-D, Alessandro R, Shatters Jr RG, Hall DG. **2013**. Behavioral, Ultrastructural and Chemical Studies on the Honeydew and Waxy Secretions by Nymphs and Adults of the Asian Citrus Psyllid Diaphorina citri (Hemiptera: Psyllidae). *PLoS ONE* **8**.

Asai T, Tena G, Plotnikova J, Willmann MR, Chiu W-L, Gomez-Gomez L, Boller T, Ausubel FM, Sheen J. **2002**. MAP kinase signalling cascade in Arabidopsis innate immunity. *Nature* **415**: 977–983.

Bai X, Correa VR, Toruño TY, Ammar E-D, Kamoun S, Hogenhout SA. **2009**. AY-WB Phytoplasma Secretes a Protein That Targets Plant Cell Nuclei. *Molecular Plant-Microbe Interactions* **22**: 18–30.

Balanzà V, Martínez-Fernández I, Ferrándiz C. **2014**. Sequential action of FRUITFULL as a modulator of the activity of the floral regulators SVP and SOC1. *Journal of Experimental Botany* **65**: 1193–1203.

Beanland L, Hoy CW, Miller SA, Nault LR. **1999**. Leafhopper (Homoptera: Cicadellidae) Transmission of Aster Yellows Phytoplasma: Does Gender Matter? *Environmental Entomology* **28**: 6.

Beanland L, Hoy CW, Miller SA, Nault LR. **2000**. Influence of Aster Yellows Phytoplasma on the Fitness of Aster Leafhopper (Homoptera: Cicadellidae). *Annals of the Entomological Society of America* **93**: 271–276.

Berry HM, Argueso CT. **2022**. More than growth: Phytohormone-regulated
transcription factors controlling plant immunity, plant development and plant
architecture. *Current Opinion in Plant Biology* **70**: 102309.

Bray NL, Pimentel H, Melsted P, Pachter L. **2016**. Near-optimal probabilistic RNA-
seq quantification. *Nature biotechnology* **34**: 525–527.

Cameron R, Lang EB, Alvarez JM. **2014**. Use of Honeydew Production to
Determine Reduction in Feeding by Bemisia tabaci (Hemiptera: Aleyrodidae) Adults
When Exposed to Cyantraniliprole and Imidacloprid Treatments. *Journal of*
Economic Entomology **107**: 546–550.

Cano LM, Raffaele S, Haugen RH, Saunders DGO, Leonelli L, MacLean D,
Hogenhout SA, Kamoun S. **2013**. Major Transcriptome Reprogramming Underlies
Floral Mimicry Induced by the Rust Fungus Puccinia monoica in Boechera stricta (J-
H Yu, Ed.). *PLoS ONE* **8**: e75293.

Chang SH, Tan CM, Wu C-T, Lin T-H, Jiang S-Y, Liu R-C, Tsai M-C, Su L-W,
Yang J-Y. **2018**. Alterations of plant architecture and phase transition by the
phytoplasma virulence factor SAP11. *Journal of Experimental Botany* **69**: 5389-5401.

Clements J, Bradford BZ, Garcia M, Piper S, Huang W, Zwolinska A, Lamour K,
Hogenhout S, Groves RL. **2021**. 'Candidatus Phytoplasma asteris' subgroups
display distinct disease progression dynamics during the carrot growing season.
PloS One **16**: e0239956.

Correa Marrero M, Capdevielle S, Huang W, Al-Subhi AM, Busscher M,
Busscher-Lange J, van der Wal F, de Ridder D, van Dijk ADJ, Hogenhout SA, *et*
al. **2024**. Protein interaction mapping reveals widespread targeting of development-
related host transcription factors by phytoplasma effectors. *The Plant Journal: For*
Cell and Molecular Biology **117**: 1281–1297.

Davies B, Egea-Cortines M, de Andrade Silva E, Saedler H, Sommer H. **1996**.
Multiple interactions amongst floral homeotic MADS box proteins. *The EMBO*
Journal **15**: 4330–4343.

Dawkins R. **1982**. *The extended phenotype*. Oxford university press Oxford.

Duneau DF, Kondolf HC, Im JH, Ortiz GA, Chow C, Fox MA, Eugénio AT, Revah
J, Buchon N, Lazzaro BP. **2017**. The Toll pathway underlies host sexual
dimorphism in resistance to both Gram-negative and Gram-positive bacteria in
mated Drosophila. *BMC Biology* **15**: 124.

Egea-Cortines M. **1999**. Ternary complex formation between the MADS-box
proteins SQUAMOSA, DEFICIENS and GLOBOSA is involved in the control of floral
architecture in Antirrhinum majus. *The EMBO Journal* **18**: 5370–5379.

Krueger, F. (2012). Trim Galore: a wrapper tool around Cutadapt and FastQC to
consistently apply quality and adapter trimming to FastQ files, with some extra
functionality for MspI-digested RRBS-type (Reduced Representation Bisufite-Seq)
libraries. *http://www. bioinformatics. babraham. ac. uk/projects/trim_galore/.(Date of*
access: 28/04/2016).

De Folter S, Immink RGH, Kieffer M, Parenicová L, Henz SR, Weigel D, Busscher M, Kooiker M, Colombo L, Kater MM, *et al.* 2005. Comprehensive interaction map of the Arabidopsis MADS Box transcription factors. *The Plant Cell.* **17**: 1424-1433.

Frost KE, Esker PD, Van Haren R, Kotolski L, Groves RL. 2013. Factors influencing aster leafhopper (Hemiptera: Cicadellidae) abundance and aster yellows phytoplasma infectivity in Wisconsin carrot fields. *Environmental Entomology* **42**: 477–490.

Gouhier-Darimont C, Schmiesing A, Bonnet C, Lassueur S, Reymond P. 2013. Signalling of Arabidopsis thaliana response to Pieris brassicae eggs shares similarities with PAMP-triggered immunity. *Journal of Experimental Botany* **64**: 665– 674.

Gregis V, Andrés F, Sessa A, Guerra RF, Simonini S, Mateos JL, Torti S, Zambelli F, Prazzoli GM, Bjerkan KN, *et al.* 2013. Identification of pathways directly regulated by SHORT VEGETATIVE PHASE during vegetative and reproductive development in Arabidopsis. *Genome Biology* **14**: R56.

Gregis V, Sessa A, Colombo L, Kater MM. 2006. AGL24, SHORT VEGETATIVE PHASE, and APETALA1 redundantly control AGAMOUS during early stages of flower development in Arabidopsis. *The Plant cell* **18**: 1373–1382.

Gregis V, Sessa A, Dorca-Fornell C, Kater MM. 2009. The Arabidopsis floral meristem identity genes AP1, AGL24 and SVP directly repress class B and C floral homeotic genes. *The Plant journal : for cell and molecular biology* **60**: 626–37.

Gu Q, Ferrandiz C, Yanofsky MF, Martienssen R. 1998. The FRUITFULL MADS-box gene mediates cell differentiation during Arabidopsis fruit development. *Development (Cambridge, England)* **125**: 1509–1517.

Hartmann U, Höhmann S, Nettesheim K, Wisman E, Saedler H, Huijser P. 2000. Molecular cloning of SVP: a negative regulator of the floral transition in Arabidopsis. *The Plant Journal* **21**: 351–360.

Hofberger JA, Zhou B, Tang H, Jones JD, Schranz ME. 2014. A novel approach for multi-domain and multi-gene family identification provides insights into evolutionary dynamics of disease resistance genes in core eudicot plants. *BMC Genomics* **15**:966.

Hong L, Rumei X. 1993. Studies on honeydew excretion by greenhouse whitefly, Trialeurodes vaporariorum (Westw.) on its host plant, Cucumis sativus. *Journal of Applied Entomology* **115**: 43–51.

Honma T, Goto K. 2001. Complexes of MADS-box proteins are sufficient to convert leaves into floral organs. *Nature* **409**: 525–529.

Hrabak EM, Chan CWM, Gribskov M, Harper JF, Choi JH, Halford N, Kudla J, Luan S, Nimmo HG, Sussman MR, *et al.* 2003. The Arabidopsis CDPK-SnRK Superfamily of Protein Kinases. *Plant Physiology* **132**: 666–680.

Huang W, Hogenhout SA. **2022**. Interfering with plant developmental timing promotes susceptibility to insect vectors of a bacterial parasite. *bioRxiv* 2022.03.30.486463; doi: https://doi.org/10.1101/2022.03.30.486463.

Huang W, MacLean AM, Sugio A, Maqbool A, Busscher M, Cho S-T, Kamoun S, Kuo C-H, Immink RGH, Hogenhout SA. **2021**. Parasitic modulation of host development by ubiquitin-independent protein degradation. *Cell* **184**: 5201-5214.e12.

Huang W, Reyes-Caldas P, Mann M, Seifbarghi S, Kahn A, Almeida RPP, Béven L, Heck M, Hogenhout SA, Coaker G. **2020**. Bacterial Vector-Borne Plant Diseases: Unanswered Questions and Future Directions. *Molecular Plant* **13**: 1379–1393.

Hunter P. **2018**. The revival of the extended phenotype. *EMBO reports* **19**: e46477.

Immink RGH, Gadella TWJ, Ferrario S, Busscher M, Angenent GC. **2002**. Analysis of MADS box protein-protein interactions in living plant cells. *Proceedings of the National Academy of Sciences* **99**: 2416–2421.

Immink RGH, Posé D, Ferrario S, Ott F, Kaufmann K, Valentim FL, Folter S de, Wal F van der, Dijk ADJ van, Schmid M, *et al.* **2012**. Characterization of SOC1's Central Role in Flowering by the Identification of Its Upstream and Downstream Regulators. *Plant Physiology* **160**: 433–449.

Jonak C. **2002**. Complexity, Cross Talk and Integration of Plant MAP Kinase Signalling. *Current Opinion in Plant Biology* **5**: 415–424.

Katsuma S, Hirota K, Matsuda-Imai N, Fukui T, Muro T, Nishino K, Kosako H, Shoji K, Takanashi H, Fujii T, *et al.* **2022**. A Wolbachia factor for male killing in lepidopteran insects. *Nature Communications* **13**: 6764.

Kitazawa Y, Iwabuchi N, Maejima K, Sasano M, Matsumoto O, Koinuma H, Tokuda R, Suzuki M, Oshima K, Namba S, *et al.* **2022**. A phytoplasma effector acts as a ubiquitin-like mediator between floral MADS-box proteins and proteasome shuttle proteins. *The Plant Cell* **34**: 1709–1723.

Kroj T, Chanclud E, Michel-Romiti C, Grand X, Morel J-B. **2016**. Integration of decoy domains derived from protein targets of pathogen effectors into plant immune receptors is widespread. *New Phytologist* **210**: 618–626.

Kumari S, Nagendran K, Rai AB, Singh B, Rao GP, Bertaccini A. **2019**. Global Status of Phytoplasma Diseases in Vegetable Crops. *Frontiers in Microbiology* **10**: 1349.

Lee I-M, Davis RE, Gundersen-Rindal DE. **2000**. Phytoplasma: Phytopathogenic Mollicutes. *Annual Review of Microbiology* **54**: 221–255.

Li P, Lu Y-J, Chen H, Day B. **2020**. The Lifecycle of the Plant Immune System. *Critical reviews in plant sciences* **39**: 72–100.

Li Z, Zeng S, Li Y, Li M, Souer E. **2016**. Leaf-Like Sepals Induced by Ectopic Expression of a SHORT VEGETATIVE PHASE (SVP)-Like MADS-Box Gene from the Basal Eudicot *Epimedium sagittatum*. *Frontiers in Plant Science* **7**: 1461.

Little D, Gouhier-Darimont C, Bruessow F, Reymond P. **2007**. Oviposition by Pierid Butterflies Triggers Defense Responses in Arabidopsis. *Plant Physiology* **143**: 784–800.

Liu Q, Maqbool A, Mirkin FG, Singh Y, Stevenson CEM, Lawson DM, Kamoun S, Huang W, Hogenhout SA. **2023**. Bimodular architecture of bacterial effector SAP05 that drives ubiquitin-independent targeted protein degradation. *Proceedings of the National Academy of Sciences of the United States of America* **120**: e2310664120.

Love MI, Huber W, Anders S. **2014**. Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. *Genome biology* **15**: 550.

Lu Y-T, Li M-Y, Cheng K-T, Tan CM, Su L-W, Lin W-Y, Shih H-T, Chiou T-J, Yang J-Y. **2014**. Transgenic Plants That Express the Phytoplasma Effector SAP11 Show Altered Phosphate Starvation and Defense Responses. *Plant Physiology* **164**: 1456–1469.

Lutz U, Posé D, Pfeifer M, Gundlach H, Hagmann J, Wang C, Weigel D, Mayer KFX, Schmid M, Schwechheimer C. **2015**. Modulation of Ambient Temperature-Dependent Flowering in Arabidopsis thaliana by Natural Variation of FLOWERING LOCUS M. *PLOS Genetics* **11**: e1005588.

MacLean AM, Orlovskis Z, Kowitwanich K, Zdziarska AM, Angenent GC, Immink RGH, Hogenhout SA. **2014**. Phytoplasma Effector SAP54 Hijacks Plant Reproduction by Degrading MADS-box Proteins and Promotes Insect Colonization in a RAD23-Dependent Manner (D Wagner, Ed.). *PLoS Biology* **12**: e1001835.

MacLean AM, Sugio A, Makarova OV, Findlay KC, Grieve VM, Toth R, Nicolaisen M, Hogenhout S a. 2011. Phytoplasma Effector SAP54 Induces Indeterminate Leaf-Like Flower Development in Arabidopsis Plants. *Plant Physiology* **157**: 831–841.

Mateos JL, Madrigal P, Tsuda K, Rawat V, Richter R, Romera-branchat M, Fornara F, Schneeberger K, Krajewski P, Coupland G. **2015**. Combinatorial activities of SHORT VEGETATIVE PHASE and FLOWERING LOCUS C define distinct modes of flowering regulation in Arabidopsis. *Genome Biology* 16:31. doi: 10.1186/s13059-015-0597-1.

Mayer CJ, Vilcinskas A, Gross J. **2008a**. Phytopathogen Lures Its Insect Vector by Altering Host Plant Odor. *Journal of Chemical Ecology* **34**: 1045–1049.

Mayer CJ, Vilcinskas A, Gross J. **2008b**. Pathogen-induced Release of Plant Allomone Manipulates Vector Insect Behavior. *Journal of Chemical Ecology* **34**: 1518–1522.

Miao Y-T, Deng Y, Jia H-K, Liu Y-D, Hou M-L. **2018**. Proteomic analysis of watery saliva secreted by white-backed planthopper, Sogatella furcifera (X-W Wang, Ed.). *PLOS ONE* **13**: e0193831.

Nørgaard LS, Phillips BL, Hall MD. **2019**. Can pathogens optimize both transmission and dispersal by exploiting sexual dimorphism in their hosts? *Biology Letters* **15**: 20190180.

Orlovskis Z, Hogenhout SA. **2016**. A Bacterial Parasite Effector Mediates Insect Vector Attraction in Host Plants Independently of Developmental Changes. *Frontiers in Plant Science* **7:** 885.

Pecher P, Moro G, Canale MC, Capdevielle S, Singh A, MacLean A, Sugio A, Kuo C-H, Lopes JRS, Hogenhout SA. **2019**. Phytoplasma SAP11 effector destabilization of TCP transcription factors differentially impact development and defence of Arabidopsis versus maize. *PLOS Pathogens* **15**: e1008035.

Pertea M, Kim D, Pertea GM, Leek JT, Salzberg SL. **2016**. Transcript-level expression analysis of RNA-seq experiments with HISAT, StringTie and Ballgown. *Nature protocols* **11**: 1650–1667.

Rid M, Mesca C, Ayasse M, Gross J. **2016**. Apple Proliferation Phytoplasma Influences the Pattern of Plant Volatiles Emitted Depending on Pathogen Virulence. *Frontiers in Ecology and Evolution* **3**.

Roy BA. **1993**. Floral mimicry by a plant pathogen. *Nature* **362**: 56–58.

Sarris PF, Cevik V, Dagdas G, Jones JDG, Krasileva KV. **2016**. Comparative analysis of plant immune receptor architectures uncovers host proteins likely targeted by pathogens. *BMC Biology* **14**: 8.

Serbus LR, Casper-Lindley C, Landmann F, Sullivan W. **2008**. The Genetics and Cell Biology of Wolbachia-Host Interactions. *Annual Review of Genetics* **42**: 683–707.

Shen L, Thong Z, Gong X, Shen Q, Gan Y, Yu H. **2014**. The putative PRC1 RING-finger protein AtRING1A regulates flowering through repressing MADS AFFECTING FLOWERING genes in Arabidopsis. *Development (Cambridge, England)* **141**: 1303–12.

Shiu S-H, Bleecker AB. **2001**. Receptor-like kinases from Arabidopsis form a monophyletic gene family related to animal receptor kinases. *Proceedings of the National Academy of Sciences* **98**: 10763–10768.

Snoeck S, Guayazán-Palacios N, Steinbrenner AD. **2022**. Molecular tug-of-war: Plant immune recognition of herbivory. *The Plant Cell* **34**: 1497–1513.

Sugio A, Kingdom HN, MacLean AM, Grieve VM, Hogenhout SA. **2011a**. Phytoplasma protein effector SAP11 enhances insect vector reproduction by manipulating plant development and defense hormone biosynthesis. *Proceedings of the National Academy of Sciences* **108**: E1254–E1263.

Sugio A, MacLean AM, Hogenhout SA. **2014**. The small phytoplasma virulence effector SAP11 contains distinct domains required for nuclear targeting and CIN-TCP binding and destabilization. *New Phytologist* **202**: 838–848.

Sugio A, MacLean AM, Kingdom HN, Grieve VM, Manimekalai R, Hogenhout SA. **2011b**. Diverse Targets of Phytoplasma Effectors: From Plant Development to Defense Against Insects. *Annual Review of Phytopathology* **49**: 175–195.

Sun L-M, Zhang J-Z, Hu C-G. **2016**. Characterization and Expression Analysis of PtAGL24, a SHORT VEGETATIVE PHASE/AGAMOUS-LIKE 24 (SVP/AGL24)-Type MADS-Box Gene from Trifoliate Orange (Poncirus trifoliata L. Raf.). *Frontiers in Plant Science* **7**: 823.

Suzuki M, Kitazawa Y, Iwabuchi N, Maejima K, Matsuyama J, Matsumoto O, Oshima K, Namba S, Yamaji Y. **2024**. Target degradation specificity of phytoplasma effector phyllogen is regulated by the recruitment of host proteasome shuttle protein. *Molecular Plant Pathology* **25**: e13410.

Tan CM, Li C-H, Tsao N-W, Su L-W, Lu Y-T, Chang SH, Lin YY, Liou J-C, Hsieh L-C, Yu J-Z, *et al.* **2016**. Phytoplasma SAP11 alters 3-isobutyl-2-methoxypyrazine biosynthesis in *Nicotiana benthamiana* by suppressing *NbOMT1*. *Journal of Experimental Botany* **67**: 4415–4425.

Tomkins M, Kliot A, Marée AF, Hogenhout SA. **2018**. A multi-layered mechanistic modelling approach to understand how effector genes extend beyond phytoplasma to modulate plant hosts, insect vectors and the environment. *Current Opinion in Plant Biology* **44**: 39–48.

Tong WH, Pavey C, O'Handley R, Vyas A. **2021**. Behavioral biology of Toxoplasma gondii infection. *Parasites & Vectors* **14**: 77.

Usadel B, Nagel A, Thimm O, Redestig H, Blaesing OE, Palacios-Rojas N, Selbig J, Hannemann J, Piques MC, Steinhauser D, *et al.* **2005**. Extension of the Visualization Tool MapMan to Allow Statistical Analysis of Arrays, Display of Coresponding Genes, and Comparison with Known Responses. *Plant Physiology* **138**: 1195–1204.

Verk, MC van. (2010, June 15). WRKY transcription factors involved in salicylic acid-induced defense gene expression. Retrieved from https://hdl.handle.net/1887/15688

Wang R, Bai B, Li D, Wang J, Huang W, Wu Y, Zhao L. **2024a**. Phytoplasma: A plant pathogen that cannot be ignored in agricultural production—Research progress and outlook. *Molecular Plant Pathology* **25**: e13437.

Wang R, Bai B, Li D, Wang J, Huang W, Wu Y, Zhao L. **2024b**. Phytoplasma: A plant pathogen that cannot be ignored in agricultural production-Research progress and outlook. *Molecular Plant Pathology* **25**: e13437.

Weintraub PG, Beanland L. **2006**. INSECT VECTORS OF PHYTOPLASMAS. *Annual Review of Entomology* **51**: 91–111.

Wilson DC, Kempthorne CJ, Carella P, Liscombe DK, Cameron RK. **2017**. Age-
Related Resistance in *Arabidopsis thaliana* Involves the MADS-Domain
Transcription Factor SHORT VEGETATIVE PHASE and Direct Action of Salicylic
Acid on *Pseudomonas syringae*. *Molecular Plant-Microbe Interactions* **30**: 919–929.

Wu R, Tomes S, Karunairetnam S, Tustin SD, Hellens RP, Allan AC, Macknight
RC, Varkonyi-Gasic E. **2017**. SVP-like MADS Box Genes Control Dormancy and
Budbreak in Apple. *Frontiers in Plant Science* **8:** 477.

Yan X, Yuan X, Lv J, Zhang B, Huang Y, Li Q, Ma J, Li Y, Wang X, Li Y, *et al.*
2024. Molecular basis of SAP05-mediated ubiquitin-independent proteasomal
degradation of transcription factors. *Nature Communications* **15**: 1170.

Yoo S-D, Cho Y-H, Sheen J. **2007**. Arabidopsis mesophyll protoplasts: a versatile
cell system for transient gene expression analysis. *Nature protocols* **2**: 1565–1572.

Zhang L, Du Y, Qu Q, Zheng Q. **2024**. Structure basis for recognition of plant
Rpn10 by phytoplasma SAP05 in ubiquitin-independent protein degradation.
iScience **27**: 108892.

Zheng Y, Ren N, Wang H, Stromberg AJ, Perry SE. **2009**. Global Identification of
Targets of the *Arabidopsis* MADS Domain Protein AGAMOUS-Like15. *The Plant Cell*
21: 2563–2577.

1239

1240

Figure supplements:

Figure supplement 1. Data distributions of independent repeats that were used to
generate graphs displayed in Fig. 1 B and C. Experimental design for choice tests
(treatments) 1 to 6 represented in **Figure 1** of the main text is shown in **panel A**. Each
choice test was performed in a separate choice cage (arena) but simultaneously with the
other choice tests in the same room. Boxplots show variation among repeated experiments
for progeny (**panel B)** and feeding (honeydew excretion) data (**panel C)**. Each data point in
panel B represents oviposition choice of 10 female insects. Each data point in panel C
corresponds to feeding choice of 20 adult insects within a single choice arena. No
preference for either test or control plant in a choice cage is represented by the 50%
reference line. Survived progeny or feeding preference for SAP54 plants is characterised by
the skewed distribution above the 50% reference line. Each reproduction choice experiment
consisted of 6 choice arenas and was performed independently 3 times (Repeat1, Repeat2,
Repeat3 with corresponding bar colours), involving a total of 540 insects. Each feeding
choice experiment consisted of 6 choice arenas and was performed independently 2 times
(Repeat1 and Repeat2 with corresponding bar colours), involving a total of 480 insects).
Paired t-test statistics for combined repeated experiments are summarised in the table below
the boxplots.

1260

1261

Figure supplement 2. *Macrosteles quadrilineatus* **female leafhoppers show no preference for** *A. thaliana* **Col-0 wild-type plants exposed to conspecific male leafhoppers**. **A**. In each choice arena 10 female leafhoppers were allowed to choose to lay eggs between insect-free *A. thaliana* Col-0 plants (with an empty clip-cage) and Col-0 plants with 10 male insects confined in clip-cages. Eggs laid by females were counted over the entire plant. Bars are 1 standard error of the mean. **B.** This experiment was independently repeated 3 times with combined total of 180 female insects. Paired t-test was performed on combined dataset considering all repeated experiments (t=1.09; p=0.325).

1270

1271

1272

1273

Figure supplement 3. Female *M. quadrilineatus* preference for male-exposed SAP54 plants are unlikely to involve long-distance cues. Choice tests were performed in separate cages, inside which two test plants were placed in non-transparent black plastic boxes that permit plant volatiles to escape but conceal the plants inside. Transparent or green sticky landing platforms were placed over each box (horizontal dashed bars). Twenty (20) female insects were released in a choice arena and females sticking to the transparent or sticky landing platforms were recorded after 1h. Bars indicate the percentage of recaptured females on each trap type. Boxplots show data distribution for recaptured females on SAP54 plants. **A**. Each choice test contained 4 choice arenas (cages) corresponding to a single datapoint in the boxplot. Each test was repeated independently 2 times. Females preferred the green over transparent platforms regardless of plant identity in within the trap. Figure shows the results of combined repeated choice tests. Choice test 1, $t_7=1.521$, p=0.172; Choice test 2, $t_7=0.226$, p=0.828; Choice test 3, $t_7=21.826$, p<0.001; Choice test 4, $t_7=21.104$, p<0.001. **B**. Females do not show preference for platforms of cages that contain male-exposed plants over insect free plants. Figure shows the results of single choice test with 4 choice arenas. Choice test 1, $t_3=0.245$, p=0.822; Choice test 2, $t_3=0.322$, p=0.769; Choice test 3, $t_3=0.302$, p=0.783; Choice test 4; $t_3=0.555$, p=0.617.

1291

1292

Figure supplement 4. Experimental design and selection of transcripts for
downstream analysis. **A**. Five (5) male or 5 female *M. quadrilineatus* individuals were
placed within a clip-cage onto a single rosette leaf of 35S:GFP-SAP54 or 35S:GFP plants.
Empty clip-cages without insects served as controls. **B**. Mapped SAP54 reads plotted
against GFP reads and colour coded for treatments (m=male; f=female; n=no insect) on
GFP and SAP54 plants. **C**. Multi-dimensional analysis (MDA) plot demonstrates grouping of
cDNA libraries according to treatment. **D.** Transcripts with normalized read count (FRKM) ≥1
in any of the sequenced libraries (10'196) and significantly differentially expressed

transcripts (DEGs) from any of the treatment pairwise comparisons (6947) were considered for downstream analysis (total=17'153 transcripts). **E**. Median of all transcript FRKM plotted against GFP reads and colour coded for treatments (m=male; f=female; n=no insect) on GFP and SAP54 plants.

1306

1307

Figure supplement 5. Biological variation and role of outliers in separation of
treatments and identification of differentially expressed genes. Euler-Venn diagrams
illustrating DEGs that differentiate SAP54 and GFP plants exposed to female or male or no
leafhoppers when outliers are retained (**A**) or removed (**B**). Principal component analysis
(PCA) plots illustrate variation within and among treatments when outliers are retained (**C**) or
removed (**D**). Arrows in panel **C** indicate the outliers (SAP54_male, GFP_male,
SAP54_female) that were removed in panel **D**.

1315

1316

1317

1318

Figure supplement 6. The cage-only SAP54 vs cage-only GFP treatments show a limited number of biotic stress DEGs. A. Euler-Venn diagrams illustrating DEGs in SAP54 plants exposed to female or male or no leafhoppers compared to no insect exposed (empty cage-only) GFP plants. **B.** Mapman diagram of DEGs involved in biotic stress in the cage-only SAP54 vs cage-only GFP plants. Pathways are indicated and each square is a gene with red versus green shades illustrating the level of up- or downregulation.

1325

1326

Figure supplement 7. Manually drawn MapMan image for defence signalling pathway visualisation.

1327

1328

1329

Figure supplement 8. SVP enhances female egg-laying preference for male exposed plants in phytoplasma and SAP54-dependent manner. MTF mutants *maf5* and *svp* display preferential leafhopper reproduction over wild-type plants in mixed-sex choice tests **(A).** Preferential leafhopper reproduction on SAP54 plants is abolished in *svp* but not *maf5* mutant (**B**). Preferential leafhopper reproduction on *svp* and *maf5* mutants is abolished in AY-WB infected plants **(C)**. *svp* displays preferential leafhopper reproduction over wild-type plants in presence of males but not when males are removed (**D**). Each datapoint represents reproductive choice of 10 female and 10 male insects within single choice arena. Experiment consists of 6 choice arenas. Each experiment was repeated independently 2 times. No preference for either test or control plant in a choice cage is represented by the 50% reference line. Oviposition preference for test or control plant is characterised by significant deviation from the 50% reference line. All pairwise comparisons done with paired t-tests by combining all datapoints from the two repeated experiments for each choice test. Test statistics summarised in table (**E**).

1346

Figure supplement 9. Comparison between male and female colonized *svp* plants reveal SVP dependent insect regulated biotic stress genes. Pairwise comparisons are schematically depicted and abbreviated in **panel A** and correspond to the Venn diagrams and MapMan biotic stress graphs in **panels B-E**. The two Venn diagrams in **panel B** display the differences in the magnitude of response to male and female leafhopper exposure of SAP54 plants and *svp* mutants compared to insect-exposed GFP and wild-type controls, respectively. **Panel C** represents log2 fold change of biotic stress DEGs in male-and female-exposed *svp* mutants according to the MapMan annotation. 8 vs 6 includes male 347+460 DEGs, 7 vs 5 includes female 464+460 DEGs, and male specific 8 vs 6 includes male 347DEGs. **Panel D** displays the overlap between 49 male-specific (not regulated in female) DEGs of SAP54 plants and *svp* mutants while the **panel E** shows 155 DEGs that are shared by male-exposed SAP54 and male-exposed *svp* plants regardless of their regulation in female-exposed SAP54 or *svp* plants.

Includes the following source data and figure supplement(s) for Figure supplement 9:

Supplementary file 10: DEG encoded function enrichment for all MapMan built-in bins in female- or male-exposed *svp* plants vs female- or male-exposed wild type plants.

Supplementary file 11: DEG encoded function enrichment for MapMan built-in built-in biotic stress and manually designed defence signalling pathways in female- or male-exposed *svp* plants vs female- or male-exposed wild type plants.

Supplementary file 12: GO-term and MapMan functional enrichment of shared 155 DEGs between ''male-exposed SAP54 vs male-exposed GFP'' comparison and ''male-exposed *svp* vs male exposed wild type'' comparison.

1369

1370

Supplementary files:

Supplementary file 1. Supplementary Table 1. FPKM and differential expression values of 17'153 genes included in the response analyses of plants to SAP54 vs GFP and male vs female leafhopper exposure.

Supplementary file 2. Supplementary Table 2. IDs and log2-fold changes of differentially expressed genes (DEGs) of male and female *M. quadrilineatus* leafhopper-exposed GFP and SAP54 plants compared to insect free GFP plants.

Tab A: GFP plants exposed to caged females versus cages alone.

Tab B: GFP plants exposed to caged males versus cages alone.

Tab C: SAP54 plants exposed to caged females versus cages alone.

Tab D: SAP54 plants exposed to caged males versus cages alone.

Tab E: SAP54 versus GFP plants exposed to cages alone.

Supplementary file 3. Supplementary Table 3. MapMan build-in functional bins enriched for DEGs in male and female *M. quadrilineatus* leafhopper-exposed GFP and SAP54 plants compared to insect-free GFP plants.

Supplementary file 4. Supplementary Table 4. MapMan build-in biotic stress functional bins enriched for DEGs in male and female *M. quadrilineatus* leafhopper-exposed GFP and SAP54 plants compared to insect-free/cage-only GFP plants.

Tab A: MapMan build-in biotic stress functional bins enriched for DEGs. *p*-values based on Wilcoxon rank test with BH correction. Bins highlighted in bold are enriched for DEGs. Individual transcript identities within significantly enriched bins are provided in the tabs B, C, D, E.

Tab B: list of DEGs within significantly enriched biotic stress bins for GFP plants exposed to caged females versus cages alone.

Tab C: list of DEGs within significantly enriched biotic stress bins for GFP plants exposed to caged males versus cages alone.

Tab D: list of DEGs within significantly enriched biotic stress bins for SAP54 plants exposed to caged females versus cages alone.

Tab E: list of DEGs within significantly enriched biotic stress bins for SAP54 plants exposed to caged males versus cages alone.

Supplementary file 5. Supplementary Table 5. Manually curated and assigned defence signalling bins for MapMan import.

Tab A, Functional bin categories.

Tab B, IDs of genes assigned to each of the functional bins listed in Tab A. "Type T" means "transcript", nomenclature used according to in-built MapMan pathway files.

Supplementary file 6. Supplementary Table 6. Functional bins for manually annotated defence genes enriched for DEGs in male and female *M. quadrilineatus* leafhopper-exposed GFP and SAP54 plants compared to insect free GFP plants.

Tab A: Functional bins for manually annotated defense genes enriched for DEGs. P-values based on Wilcoxon rank test with BH correction. Bins highlighted in bold are enriched for DEGs. Individual transcript identities within significantly enriched bins are provided in the tabs B, C, D, E.

Tab B: list of DEGs within significantly enriched biotic stress bins for GFP plants exposed to caged females versus cages alone.

Tab C: list of DEGs within significantly enriched biotic stress bins for GFP plants exposed to caged males versus cages alone.

Tab D: list of DEGs within significantly enriched biotic stress bins for SAP54 plants exposed to caged females versus cages alone.

Tab E: list of DEGs within significantly enriched biotic stress bins for SAP54 plants exposed to caged males versus cages alone.

Supplementary file 7. Supplementary Table 7. MapMan build-in functional bins enriched for DEGs in SAP54 versus GFP plants with or without exposure to male and female *M. quadrilineatus* leafhoppers.

Supplementary file 8. Supplementary Table 8. Biotic stress bins from MapMan build-in and manually curated defence signalling pathway bins enriched for DEGs in GFP and SAP54 plants with or without exposure to male and female *M. quadrilineatus* leafhoppers.

Supplementary file 9. Supplementary Table 9. Fold-expression changes of MADS-box transcription factor genes insect-exposed SAP54 vs GFP leaves.

Tab A: List of 20 genes encoding MADS-box transcription factors that are expressed in leaves in the (insect-exposed) SAP54 and GFP plants.

Tab B: List of all 107 genes annotated as MADS-box transcription factors in *A. thaliana* (de Folter et al., 2005).

1443

1444

1445 **Supplementary file 10**. Supplementary Table 10. DEG encoded function
1446 enrichment for all MapMan built-in bins in female- or male-exposed *svp* plants vs
1447 female- or male-exposed wild type plants.

1448 *p*-values based on Wilcoxon rank test and have BH correction.

1449

Supplementary file 11. Supplementary Table 11. DEG encoded function
enrichment for MapMan built-in built-in biotic stress and manually designed defence
signalling pathways in female- or male-exposed *svp* plants vs female- or male-
exposed wild type plants.

Tab A: biotic stress and defence signalling pathway bins enriched with female- or
male-specific DEGs. *p*-value based on Wilcoxon rank test and BH correction.

Tab B: list if DEGs within enriched bins for clip-caged females on *svp* plants vs clip-
caged females on wild type plants.

Tab C: list if DEGs within enriched bins for clip-caged males on *svp* plants vs clip-
caged males on wild type plants.

Tab D: since no enriched bins found for male-specific DEGs in tab A, no transcripts
highlighted here.

1462

Supplementary file 12. Supplementary Table 12. GO-term and MapMan functional
enrichment on shared 155 DEGs between ''male-exposed SAP54 vs male-exposed
GFP'' comparison and ''male-exposed *svp* vs male-exposed wild type'' comparison.

Tab A: GO-term enrichment with 155 DEGs shared by male-exposed SAP54 plants
(versus male-exposed GFP plants) and male-exposed *svp* plants (vs male-exposed
wild type plants) and depicted in Figure supplement 9D.

Tab B: gene names, encoded function description and log2(fold) change for the 155
DEGs analysed in tab A and depicted in Figure supplement 9D.

Tab C: Common functions identified between (1) MapMan function enrichment for
3816 DEGs on male-exposed SAP54 plants vs male-exposed GFP plants and (2)
MapMan function enrichment for 807 DEGs on male-exposed *svp* plants vs male-
exposed wild type plants; Venn diagram of the DEGs in Figure supplement 9D.

1475

A Treatment

B

35S:GFP-SAP54   *35S:GFP*

Macrosteles quadrilineatus nymphs (%)

C

35S:GFP-SAP54   *35S:GFP*

Macrosteles quadrilineatus honeydew secretion (%)

Plant selection by female leafhoppers ♀

Insect vector egg-laying

Male leafhoppers colonise plants ♂

Insect-induced plant responses

MTF Network remodelling

Leaf-like flowers

Plant Cell

Vegetative functions

Flowering and reproduction

?

SVP

26S proteasome

MTFs

Phytoplasma

SAP54

RAD23

T M