Modeling Riboswitches: The impact of SAM concentration on the folding of the SAM-II riboswitch

Osama Alaidi [1,2]

Affiliations

[1] Biocomplexity for Research and Consulting, Cairo, Egypt.

[2] Department of Pharmaceutical Sciences, University of Tennessee Health Science Center, Memphis, TN, 38163, United States.

Author email

osama.alaidi@gmail.com

ORCID: Osama Alaidi: 0000-0002-1292-6789

ABSTRACT

Several mechanistic (thermodynamic) models have been developed for the folding of SAM-II riboswitch as a function of SAM and magnesium concentrations. For each model, the model parameters (equilibrium constants) have been determined from experimental (apparent) binding data, based on the underlying assumptions of the model. The predicted titration curves computed from the different models were calculated and compared with actual experimental observation of the fraction of the RNA forming a pseudoknot at specific concentration of the ligands. Strikingly, only one of the six models correctly predicts the experimental findings, confirming the dominant mechanism of the riboswitch function. More interestingly, the latter mechanism is found to be the most efficient compared to the other possible mechanisms. The study sheds light on the cognate ligand conformational capture mechanism of the SAM-II riboswitch in the presence of specific concentrations of magnesium ions. The presented mathematical and thermodynamic framework, as well as the inferred equilibrium constants, provide foundations for making accurate quantitative prediction of the SAM-II riboswitch ensemble populations as a function of SAM and magnesium concentrations. The mechanistic linked equilibria model can be generalized to describe other thermodynamically driven riboswitches and hence facilitate identifying RNA intermediates that can be leveraged for small molecule drug design.

KEYWORDS

SAM-II riboswitch, modeling riboswitches, ligand-induced conformational changes, conformational capture mechanism, thermodynamics, SAM.

1   INTRODUCTION

Riboswitches are RNA sensor elements, primarily present in bacterial mRNA, that aid in gene regulation of small molecule metabolism on both the transcriptional and translational levels [1]. To date, more than fifty classes of riboswitches have been confirmed, and thousands of novel riboswitches are predicted in bacterial genomes [2]. The SAM-II riboswitch is a class of riboswitches that binds to the metabolite; SAM (S-Adenosyl-Methionine), leading to termination of the ribosomal translation of the downstream genes that are involved in the metabolism of SAM [3,4]. The mechanism by which the RNA switches its conformation from ON to OFF and vice versa, in the absence and the presence of SAM, respectively, is yet to be fully understood [5–10]. Understanding such mechanism is crucial to be able to control the function of the riboswitch by means of SAM or other designed small molecules [11,12]. It is likely that riboswitches share common functional mechanisms. Consequently, structural and mechanistic insights from one riboswitch may be observed in other classes of riboswitches, and hence the importance of mechanistic studies on simple model riboswitches such as the SAM-II riboswitch from the Sargasso Sea metagenome that controls the mRNA translation of the *metX* gene [13,3–7,14,8,9].

In the first part of this series of studies, the impact of magnesium ions on the folding of the SAM-II riboswitch was investigated. In this manuscript, mechanistic models, by which the riboswitch RNA conformational changes may take place in response to its cognate ligand (SAM), are presented and applied to predict titration curves for the SAM-II riboswitch. The models build on the general framework used in our previous work [10], and generally combines RNA structure prediction along with a systems biology approach to enable the prediction of the fraction of riboswitch bound and/or folded at equilibrium, at various ligand concentrations. By deriving and applying linked equilibria models that carry various hypotheses, different possible mechanisms by which the ligand may induce its effect were examined. Comparing the results from these models indicated that one of the mechanisms is more plausible than the others and hence likely to be the dominant mechanism.

2   METHODS

2.1   Predicting ligand concentration-dependent SAM-II riboswitch folding (titration curves)

Titration curves provide a means to compare model predictions with experimental findings. Doubtlessly, the accuracy of the predictions made, based on a certain model, reflects the accuracy of the underlying assumptions of the model. In this study, mechanistic linked equilibria models were developed and used to examine the effect of SAM on the SAM-II riboswitch population in the presence of Mg^{2+} ions. The derived models that were used for the calculations in this study are tabulated in **Table I**, along with the major assumptions made in each model. Previously studied models describing the impact of magnesium ions on the riboswitch (designated as *Mg^{2+} binding models*) are also listed in the table for clarity.

The presented six SAM-binding models in this study assume distinct mechanisms, and differ from each other based on whether if: (1) the presence of Mg^{2+} is essential for SAM binding, (2) the Mg^{2+} concentration influence the fraction of the finally folded state, (3) tertiary interactions are essential for the conformational capture mechanism to take place, (4) more than one mechanistic pathway may play a role in the apparent binding. Each of the developed models therefore examines a certain set of assumptions or a proposed plausible mechanism. The model that correctly predicts an experimental observable, is that whose underlying assumptions are likely to be correct. Hence, the comparison of the modeling results with experimental values can provide an indication of which model is likely to be valid.

The usual workflow in using such mathematical models was followed here. First, the analytical/mathematical model was constructed based on a plausible mechanism. Obviously, any mechanism will intrinsically have certain assumptions. Second, the model parameters were determined based on a few measured apparent equilibrium constants obtained from the literature. Third, a titration curve was computed from each model and were compared to an experimentally observed/measured values. The model which gave values that are most consistent with experimental observations, was considered the one that most likely to occur, and hence effectively represents the mechanism of the riboswitch function.

The presented models are listed and compared in **Table I**. Models were named according to the presence and absence of Mg^{2+} and SAM in the model. For example, *Model 0* (**Fig. S1**) is a model that simply describes the riboswitch folding in the absence of both Mg^{2+} and SAM,

whereas Mg^{2+} *Model* is a model that describes the riboswitch folding in the presence of Mg^{2+} and in the absence of SAM. The latter models are mentioned here for clarity and/or comparison and have been described in our previous study [10]. The SAM-binding models *(Models 1 to 6)* depict six distinct mechanistic models that can describe the riboswitch folding in the presence of SAM, and in which the effect of Mg^{2+} may (either implicitly or explicitly) or may not be taken into account, depending on the specific scheme and the underlaying assumptions of the model.

The SAM-binding models are schematically illustrated in the top panels of **Fig. 1-6** and described in detail in the following sections of the manuscript. Moreover, as elaborated in the following sections, the equilibrium constants for each model are inferred based on the apparent experimentally determined binding affinities, and when necessary, from the cognate thermodynamic cycles shown in bottom panels in **Fig. 1, 3, 5** and **6**.

Model name	SAM binding	Presence of Mg^{2+}	Effect of Mg^{2+} concentration	Tertiary interactions play a role in SAM conformational selection	Alternative SAM binding pathways	Reference
Model 0	-	-	-	Not applicable	Not applicable	Reference [10]. Also briefly described in this manuscript (Supplementary methods).
Mg^{2+} *Model*	-	+	+	Not applicable	Not applicable	Reference [10]
Model 1	+	+	+	+	+	This manuscript
Model 2	+	+	+	+	-	This manuscript
Model 3	+	+	+	-	+	Derived in reference [10]
Model 4	+	-	-	+	-	This manuscript
Model 5	+	+	-	-	+	This manuscript
Model 6	+	-	-	-	+	This manuscript

TABLE I. Summary of the models constructed to describe the mechanism of the SAM-II riboswitch. The models vary in their underlying assumptions regarding the role of tertiary interactions in the conformational capture mechanism, the role of alternative SAM-binding pathways and whether the model considers the presence and concentration effects of Mg^{2+}. The *Model 0* and the Mg^{2+} *Model,* are listed here for clarity (and have been described in our previous

manuscript) and they represent, respectively, the description of the RNA in the absence of both ligands (Mg^{2+} and SAM) and a model that describe the impact of Mg^{2+} on the RNA, in the absence of SAM. In the table above, the - and + signs are used to indicate whether the feature or assumption is considered by in the model. *SAM Models 1* to *6* are pictorially illustrated in the top panels of **Fig. 1** - **6**, respectively.

2.2 Models for quantitative prediction of RNA-ligand interactions for the SAM-II riboswitch

2.2.1 Definitions of the RNA species (states) and parameters (equilibrium constants) used in the models

The following definitions or nomenclature of the RNA species and model parameters (equilibrium constants) are, for consistency, used in all models, though not all the models necessarily have all these species and parameters. The following nomenclature is therefore used throughout this manuscript.

1- **Total and free concentrations:** Like in the Mg^{2+} binding model described in our previous manuscript [10], in all models presented in this study, the total RNA, Mg^{2+} ions and SAM concentrations are designated as R_{tot} , Mg_{tot} and S_{tot}, respectively. Whereas the free (unbound) Mg^{2+} ions and SAM are designated as Mg and S, respectively.

2- **RNA species (states) concentrations:** In general, RNA species with a pseudoknot formed are considered to be in the OFF state (inaccessible ribosome binding site) whereas RNA species without a pseudoknot formed are in the ON state (accessible ribosome binding site). The RNA conformers in the models are defined as,

R_0 is the concentration of the unbound *binding incompetent* RNA (i.e. RNA conformers in which the SAM binding contacts are not accessible). Such RNA states/species do not show the specific RNA secondary structure, nor they are bound with any ligands.

R is the concentration of the unbound intermediate. This class of conformers has the minimal secondary structure required for SAM binding (i.e. which allows the SAM binding contacts to be accessible) but does not have a pseudoknot.

$R^{`}$ is the RNA concentration without a pseudoknot. Hence, this class of RNA is composed of more than one state, namely, $R_0$ and $R$ (i.e. $R^{`} = R_0 + R$).

R_{ter} is the concentration of the riboswitch RNA with a nucleated pseudoknot but not bound to Mg^{2+} nor SAM.

R_{terMg} is a state structurally similar R_{ter} except that it is bound with Mg^{2+}.

R_{terE} , is the concentration of RNA with a nucleated pseudoknot that effectively enters the subsequent reaction, regardless of its binding with Mg^{2+}. This class of RNA is effectively a mixture of states, i.e. $R_{terE} = R_{ter} + R_{terMg}$.

I is the concentration of the SAM bound intermediate (SAM bound riboswitch with or without the pseudoknot, depending on the model).

C is the concentration of the SAM bound *final and most stable state of the riboswitch* (SAM bound riboswitch with the pseudoknot fully formed). Depending on the model, this class of conformers may or may not be bound to Mg^{2+} or may represent a mixture of both. It is important to note that only species I and C are bound to SAM.

3- *Models' equilibrium constants:*

The equilibrium constants for the RNA conversions, are defined as,

K_{conv1} is the equilibrium constant for the conversion from R_0 to R.
K_{conv2} is the equilibrium constant for the conversion from R to R_{ter}.

To reduce the number of parameters in the tested models as much as possible, and to ease the incorporation of experimental data, the following apparent equilibrium constants are defined:

K_{ter} is the apparent equilibrium constant for the conversion from $R^`$ to R_{ter}.
K_{terE} is the apparent equilibrium constant for the conversion from R to R_{terE}.
Hence, both K_{ter} and K_{terE} are apparent equilibrium constants.

As described above, each of $R^`$ and R_{terE} are composed of more than one state. The equilibrium constants for RNA binding, are defined as:

K_{Mg} is the equilibrium binding association constant of Mg^{2+} to species R_{ter}.
K_{bind} is the equilibrium binding association constant of SAM to species R.
K_{bindE} is the equilibrium binding association constant of SAM to species R_{terE}.
$K_{bindter}$ is the equilibrium binding association constant of SAM to species R_{ter}.
K_{bindMg} is the equilibrium binding association of SAM to species R_{terMg}.

K_{fold} is the equilibrium association constant for the folding of the intermediate I to form C.

In the absence of any ligands (**Fig. S1**, *Model 0*), the equilibrium constants for RNA conversion can be simply obtained from RNA folding probabilities, as follows.

$K_{conv1} = \frac{P_R}{P_{R_0}}$, represents the ratio of probabilities of conformers R and R_0, in the *absence* of SAM and absence of tertiary interactions. Hence, the calculation can be performed exclusively based on secondary structure energetics.

$K_{conv2} = \frac{P_{R_{ter}}}{P_R}$, represents the ratio of probabilities of R_{ter} and R, in the *absence* of SAM. The probabilities of these populations do not rely on any ligands, and the value for the equilibrium constant was obtained from the previously reported two-step Mg^{2+} binding model [10].

$K_{ter} = \frac{P_{R_{ter}}}{P_{R^`}}$ , represents the ratio of probabilities of $R_{ter}$ and $R^`$, in the *absence* of SAM. The probabilities of these populations also do not rely on any ligands, and the value for the equilibrium constant has also been obtained from the previously reported two-step Mg^{2+} binding model [10].

$K_{terE} = \frac{P_{R_{terE}}}{P_R}$ or $K_{terE} = \frac{R_{terE}}{R \,.\, Mg}$ (depending on the model, reflecting if the model considers or ignores the effect of magnesium), represents the ratio of probabilities of R_{terE} and R, in the *absence* of SAM. The ratio between these two populations relies on the concentration of Mg^{2+} ions. The probabilities and equilibrium constant can also be obtained from a two-step Mg^{2+} binding model [10].

For each model, the binding and folding parameters are determined from the apparent experimental binding constants reported in the literature, with the help of thermodynamic cycles if needed. The latter equilibrium constants may include, K_{bind}, $K_{bindter}$, K_{bindMg}, K_{bindE} and K_{fold}. Thereafter, a quantitative prediction of the concentration of the bound and folded (i.e. with the pseudoknot formed) riboswitch (i.e. species I and/or C, depending on the model) is obtained as a function of the equilibrium constants and the total concentrations of RNA and ligands (Mg^{2+} and SAM). To enable this, it was essential to establish links between the latter equilibrium constants and the literature apparent binding constants.

4- *Experimental equilibrium constants:*

Ka_{ITC} and Ka_{FS} denotes two apparent experimental binding association constants previously reported in the literature that are frequently used in this work, derived based on Isothermal

calorimetry (ITC) [4] and Fluorescence spectroscopy (FS) [6], respectively. Whereas the former defines binding in the usual ratio between the bound and free RNA in the presence of the ligand, the second monitors the changes in the RNA folding. More specifically, the latter method monitors the changes (increase or decrease) in fluorescence intensity from a 2-Aminopurine inserted at a specific site (as an indicator for the increase or decrease of a state), that arise from local interactions (many of which are tertiary interactions) associated with RNA folding, as a function of the ligand concentration [15–17]. Effectively, such technique monitors the fraction change in the emitted fluorescent signal, at a specific wavelength range, as the cognate ligand is added. In their study on the SAM-II riboswitch, Haller *et al.* monitored the increase in the fluorescent signal associated with the increase of SAM concentration at several bases that undergo changes associated with the pseudoknot formation [6]. Since the equilibrium constant from such method is based on monitoring the change in the fraction of RNA that gives off a signal at the specific wavelength range, and that the ensemble contains a fraction of RNA that already has a pseudoknot prior to adding SAM as well as a fraction that remains unfolded, then measured constant represent, the ratio between the concentration of the folded & bound RNA relative to the concentration of folded & unbound RNA, taking the ligand concentration, that is, the reaction scheme and equilibrium constant, respectively, can be written as:

$$Folded\ \&\ unbound\ RNA\ (gives\ initial\ FS\ signal) + SAM \xrightleftharpoons[Ka_{FS}]{} Folded\ \&\ bound\ RNA\ (increased\ FS\ signal)$$

and $Ka_{FS} = \dfrac{Folded\ \&\ bound\ RNA}{Folded\ \&\ unbound\ RNA \times free\ SAM} = \dfrac{Bound\ RNA\ species\ with\ a\ pseudoknot}{Unbound\ RNA\ species\ with\ a\ pseudoknot\ \times free\ SAM}$.

In the latter expression, the concentrations of the folded and bound RNA species will depend on the specific scheme for each model. As shown in the remaining parts of the methods, by specifying those RNA species for each model, links between the experimental measurements and the model were established, and the model parameters were inferred.

2.2.2   *Model 1: A ligand-binding model for RNA that considers the effect of both Mg^{2+} ions and tertiary interactions*

The model, designated as *Model 1*, is schematically illustrated in **Fig. 1** (top panel). In this model, the cognate ligand (SAM) binds to all molecules with a nucleated pseudoknot, whether bound with Mg^{2+} ions or not. Hence, this model considers the effect of both Mg^{2+} concentration and tertiary interactions, and the complete folding can be achieved through either magnesium-dependent or a magnesium-independent pathway.

FIG. 1. *Model 1* **and its cognate thermodynamic cycle.** The figure shows the functional mechanism of the SAM-II riboswitch. The top panel illustrates a graphical reaction scheme for the proposed model designated as *Model 1*. The related Thermodynamic cycle is shown in the bottom panel. The model assumes a conformational capture mechanism that is conditioned by the presence of a pseudoknot that gets stabilized by the binding of SAM. The latter binding can occur with or without the prior binding of Mg^{2+}. That is, in this mechanism, SAM binds to two different forms of the partially formed pseudoknot structures (i.e. two intermediates), hence induces the complete RNA folding and SAM trapping.

Determination of Model 1 parameters

The overall (apparent) dissociation constant which was determined experimentally using Isothermal calorimetry (ITC) and reported by Gilbert *et al* [4]. describes the following reaction,

$$Unbound\ RNA + SAM \xrightleftharpoons[Ka_{ITC}]{} bound\ RNA$$

$$[R_0 + R + R_{ter} + R_{terMg}] + [S] \xrightleftharpoons[Ka_{ITC}]{} [I + C]$$

$$[R^` + R_{ter} + R_{terMg}] + [S] \xrightleftharpoons[Ka_{ITC}]{} [I + C]$$

Kd_{ITC} was experimentally determined to be $0.67 \pm 0.06\ \mu M$. Hence, this experimental dissociation constant $Kd_{ITC} = \frac{1}{Ka_{ITC}}$, describes the following equilibrium,

$$Ka_{ITC} = \frac{Bound\ RNA}{Unbound\ RNA\ \times free\ SAM}$$

$$= \frac{[I + C]}{[R_0 + R + R_{ter} + R_{terMg}] \times [S]}$$

$$= \frac{[I + C]}{[R^` + R_{ter} + R_{terMg}] \times [S]}$$

$$= \frac{\left[\left(\frac{1}{K_{fold}.Mg}\right).C + C\right]}{\left[\frac{R_{ter}}{K_{ter}} + \frac{R_{terMg}}{K_{Mg}.Mg} + R_{terMg}\right] \times [S]}$$

$$= \frac{C\left[\left(\frac{1}{K_{fold}.Mg}\right) + 1\right]}{R_{terMg} \times S \times \left[\frac{1}{K_{ter}K_{Mg}.Mg} + \frac{1}{K_{Mg}.Mg} + 1\right]}$$

$$= \frac{C \times \beta_1}{R_{terMg} \times S \times \alpha_1}$$

Thus,

$$Ka_{ITC} = K_{bindMg}\frac{\beta_1}{\alpha_1} \qquad\qquad \text{Eq. 1}$$

where, K_{bindMg} is the SAM binding association constant of R_{terMg}, $\beta_1 = \left(\frac{1}{K_{fold}.Mg}\right) + 1$ and, $\alpha_1 = \left(\frac{1}{K_{ter}K_{Mg}.Mg} + \frac{1}{K_{Mg}.Mg} + 1\right)$. The value of α_1 is given and depends on the Mg^{2+} ions concentration, whereas β_1 is unknown and cannot be determined from the thermodynamic cycle alone.

Similarly (and alternatively),

$$Ka_{ITC} = \frac{Bound\ RNA}{Unbound\ RNA\ \times free\ SAM}$$

$$= \frac{[I + C]}{[R_0 + R + R_{ter} + R_{terMg}] \times [S]}$$

$$= \frac{[I + C]}{[R` + R_{ter} + R_{terMg}] \times [S]}$$

$$= \frac{[I + K_{fold}.Mg.I]}{\left[\frac{R_{ter}}{K_{ter}} + R_{ter} + K_{Mg}.Mg.R_{ter}\right] \times [S]}$$

$$= \frac{I\left[K_{fold}.Mg + 1\right]}{R_{ter} \times S \times \left[\frac{1}{K_{ter}} + K_{Mg}.Mg + 1\right]}$$

$$= \frac{I \times \beta_2}{R_{ter} \times S \times \alpha_2}$$

Thus,

$$Ka_{ITC} = K_{bindter}\ \frac{\beta_2}{\alpha_2} \qquad\qquad \text{Eq. 2}$$

where, $K_{bindter}$ is the SAM binding association constant of R_{ter}, $\alpha_2 = \left(\frac{1}{K_{ter}} + K_{Mg}.Mg + 1\right)$ and $\beta_2 = (K_{fold}.Mg + 1)$.

The previously reported FS studies monitor RNA folding in the presence of the ligand relative to its absence [6]. In other words, the apparent binding constant determines the fraction of RNA species with a pseudoknot and bound to the ligand compared to the fraction with a pseudoknot but without the ligand. Therefore, binding constants from such FS studies, represent the following reaction,

$$Folded\ \&\ unbound\ RNA + SAM \underset{Ka_{FS}}{\Longleftrightarrow} Folded\ \&\ bound\ RNA$$

$$[R_{ter} + R_{terMg}] + [S] \underset{Ka_{FS}}{\Longleftrightarrow} [I + C]$$

$$Ka_{FS} = \frac{Bound\ RNA\ species\ with\ a\ pseudoknot}{Unbound\ RNA\ species\ with\ a\ pseudoknot\ \times free\ SAM}$$

$$= \frac{[I + C]}{\left[R_{ter} + R_{terMg}\right] \times [S]}$$

$$= \frac{\left[I + K_{fold}.Mg.I\right]}{\left[R_{ter} + K_{Mg}.Mg.R_{ter}\right] \times [S]}$$

$$= \frac{I\left[K_{fold}.Mg + 1\right]}{R_{ter} \times S \times \left[K_{Mg}.Mg + 1\right]}$$

$$= \frac{I \times \beta_2}{R_{ter} \times S \times \epsilon}$$

Thus,

$$Ka_{FS} \quad = K_{bindter} \; \frac{\beta_2}{\epsilon} \qquad\qquad \text{Eq. 3}$$

where, $K_{bindter}$ is the SAM binding association constant of R_{ter}, $\epsilon = \left(K_{Mg}.Mg + 1\right)$ and $\beta_2 = \left(K_{fold}.Mg + 1\right)$.

For convenience and validation, by substitution for β_2 (from Eq. 2) into Eq. 3, a relation between Ka_{FS} and Ka_{ITC} can be established,

$$Ka_{FS} = K_{bindter} \; \frac{\beta_2}{\epsilon} = K_{bindter} \frac{Ka_{ITC} \; \alpha_2}{\epsilon \; K_{bindter}} = \; Ka_{ITC} \; \frac{\alpha_2}{\epsilon} \qquad\qquad \text{Eq.4}$$

Since, K_{Mg} and K_{ter} is known from magnesium data [10], α_2 and ϵ can be computed, and hence knowing one of the two constants, either Ka_{ITC} or Ka_{FS} can determine the other. Each of the latter two constants are determined by two of the unknown constants.

To determine each of K_{fold}, $K_{bindter}$ and K_{bindMg} a third relation from the thermodynamic cycle is needed. From the cycle in **Fig. 1** (bottom panel), we have the following,

$$\Delta G_{fold} + \Delta G_{bindter} - \Delta G_{bindMg} - \Delta G_{Mg} = 0$$

$$\Delta G_{fold} = \Delta G_{bindMg} + \Delta G_{Mg} - \Delta G_{bindter}$$

By dividing both sides by $-RT$ and taking the exponents of both sides, we obtain,

$$e^{\frac{-\Delta G_{fold}}{RT}} = e^{\frac{-\Delta G_{bindMg}}{RT} + \frac{-\Delta G_{Mg}}{RT} - \frac{-\Delta G_{bindter}}{RT}}$$

$$e^{\frac{-\Delta G_{fold}}{RT}} = \frac{e^{\frac{-\Delta G_{bindMg}}{RT}} \cdot e^{\frac{-\Delta G_{Mg}}{RT}}}{e^{\frac{-\Delta G_{bindter}}{RT}}}$$

$$K_{fold} = \frac{K_{bindMg}}{K_{bindter}} K_{Mg}$$

By substitution of K_{bindMg} (from Eq. 1) and $K_{bindter}$ (from Eq. 3) in terms of experimentally measured constants (Ka_{ITC} and Ka_{FS}), we obtain,

$$K_{fold} = \frac{Ka_{ITC} \ \alpha_1 \ \beta_2}{Ka_{FS} \ \epsilon \ \beta_1} K_{Mg}$$

$$K_{fold} \frac{\beta_1}{\beta_2} = \frac{Ka_{ITC} \ \alpha_1 \ K_{Mg}}{Ka_{FS} \ \epsilon} = \frac{\sigma}{\rho}$$

Then by the substitution for β_2 and β_1 (as defined in Eq. 1 and Eq. 2), and solving for K_{fold}, the latter constant can be determined,

$$K_{fold} = \frac{\frac{\sigma}{Mg} - \rho}{\rho Mg - \sigma} \qquad \text{Eq. 6}$$

Where, $\sigma = Ka_{FS} \ \epsilon$ and $\rho = Ka_{ITC} \ \alpha_1 \ K_{Mg}$. Once K_{fold} is determined from Eq. 6, K_{bindMg} and $K_{bindter}$ can be determined through Eq. 1-3.

Given the equilibrium constants, the expression for C (given by Eq 1.15, in the **Supplementary methods** and tabulated in **Table S2**) as well as all stable RNA species in the model can be calculated as a function of SAM concentration (see model derivation in the **Supplementary methods**). The *fraction bound* (corresponding to the total probability of bound RNA at a given SAM concentration (S_{tot})), as well as the *fraction folded*, *fraction both bound & folded* and *fraction binding competent*, can be obtained from,

$$fraction \ bound = \frac{I + C}{R_{tot}}$$

$$fraction \ folded \ (i.e. with \ a \ pseudoknot) = \frac{R_{ter} + R_{terMg} + I + C}{R_{tot}}$$

$$fraction \ bound \ and \ folded \ = \frac{I + C}{R_{tot}}$$

$$fraction \ of \ binding \ competent \ RNA \ = \frac{R_{ter} + R_{terMg} + I + C}{R_{tot}}$$

2.2.3 Model 2: A ligand-binding model for RNA that considers the effect of both Mg²⁺ ions and tertiary interactions in which the primary ligand can only bind if Mg²⁺ is bound

This model, designated as *Model 2*, is schematically illustrated in **Fig. 2**. In this model, the cognate ligand (SAM) binds exclusively to molecules with a nucleated pseudoknot and that are bound with Mg^{2+} ions. Hence this model considers the effect of both Mg^{2+} and tertiary interactions, however, assumes that the complete folding of the RNA is only achievable though a single magnesium-dependent pathway.

FIG. 2. A schematic representation of *Model 2*. The figure illustrates a graphical reaction scheme for the proposed model designated as *Model 2*. The model assumes that SAM binds via a conformational capture mechanism. The occurrence of the latter mechanism is conditioned by both the presence of pseudoknot and the binding of Mg^{2+}.

Determination of Model 2 parameters

Model parameters can be obtained in a similar way to the previous model except that the intermediate I is absent i.e. $I = 0$ and hence the factor $\beta_1 = 1$. Briefly, the overall (apparent) dissociation constant which was determined experimentally using Isothermal calorimetry (ITC), and reported by Gilbert *et al.* [4] describes the following reaction,

$$Unbound\ RNA + SAM \underset{Ka_{ITC}}{\overset{}{\rightleftharpoons}} bound\ RNA$$

$$\left[R_0 + R + R_{ter} + R_{terMg}\right] + [S] \underset{Ka_{ITC}}{\overset{}{\rightleftharpoons}} [C]$$

$$\left[R` + R_{ter} + R_{terMg}\right] + [S] \underset{Ka_{ITC}}{\overset{}{\rightleftharpoons}} [C]$$

Kd_{ITC} was experimentally determined to be $0.67 \pm 0.06\ \mu M$. Since this experimental dissociation constant $Kd_{ITC} = \frac{1}{Ka_{ITC}}$, describes the following equilibrium,

$$Ka_{ITC} = \frac{Bound\ RNA}{Unbound\ RNA\ \times\ free\ SAM}$$

$$= \frac{[C]}{[R_0 + R + R_{ter} + R_{terMg}] \times [S]}$$

$$= \frac{[C]}{[R` + R_{ter} + R_{terMg}] \times [S]}$$

$$= \frac{[C]}{\left[\frac{R_{ter}}{K_{ter}} + \frac{R_{terMg}}{K_{Mg}.Mg} + R_{terMg}\right] \times [S]}$$

$$= \frac{C}{R_{terMg} \times S \times \left[\frac{1}{K_{ter}K_{Mg}.Mg} + \frac{1}{K_{Mg}.Mg} + 1\right]}$$

$$= \frac{C}{R_{terMg} \times S \times \alpha_1}$$

Thus,

$$Ka_{ITC} \quad = \quad \frac{K_{bindMg}}{\alpha_1} \qquad\qquad \text{Eq. 7}$$

where, K_{bindMg} is the SAM binding association constant of R_{terMg}, and $\alpha_1 = \left(\frac{1}{K_{ter}K_{Mg}.Mg} + \frac{1}{K_{Mg}.Mg} + 1\right)$. The value of α_1 is given and it depends on the Mg^{2+} ions concentration along with other equilibrium constants.

As in the previous model, again the reported FS studies monitor RNA folding in the presence of the ligand relative to its absence. That is, the apparent binding constant determines the fraction of RNA species with a pseudoknot and bound to the ligand compared to the fraction with a pseudoknot but without the ligand considering the ligand concentration. Therefore, if this model is true, the binding constants from FS studies, would represent the following reaction,

$$Folded\ \&\ unbound\ RNA + SAM \xrightleftharpoons[Ka_{FS}]{} Folded\ \&\ bound\ RNA$$

$$[R_{ter} + R_{terMg}] + [S] \xrightleftharpoons[Ka_{FS}]{} [I + C]$$

$$Ka_{FS} = \frac{Bound\ RNA\ species\ with\ a\ pseudoknot}{Unbound\ RNA\ species\ with\ a\ pseudoknot\ \times\ free\ SAM}$$

$$= \frac{[C]}{[R_{ter} + R_{terMg}] \times [S]}$$

$$= \frac{[C]}{\left[\frac{R_{terMg}}{K_{Mg}.Mg} + R_{terMg}\right] \times [S]}$$

$$= \frac{[C]}{R_{terMg} \times S \times \left[\frac{1}{K_{Mg}.Mg} + 1\right]}$$

$$Ka_{FS} = K_{bindMg} \frac{1}{\left[\frac{1}{K_{Mg}.Mg} + 1\right]} = K_{bindMg} \cdot \gamma \qquad \text{Eq. 8}$$

Thus, for the purpose of validation, a relation between Ka_{FS} and Ka_{ITC} can be established,

$$Ka_{FS} = Ka_{ITC} \cdot \alpha_1 \cdot \gamma \qquad \text{Eq. 9}$$

where, $\gamma = \frac{1}{\left[\frac{1}{K_{Mg}.Mg} + 1\right]}$.

Given the equilibrium constants (K_{Mg} and K_{ter}) and after computing K_{bindMg} (from Eq. 8), the expression for C (Eq. 2.6, in the **Supplementary methods** and tabulated in **Table S2**) as well as all stable RNA species in the model can be calculated as a function SAM concentration (see *Model 2* derivation in **Supplementary methods**). The *fraction bound, fraction folded, fraction both bound & folded* and *fraction binding competent*, for this model, can be obtained from,

$$fraction\ bound = \frac{C}{R_{tot}}$$

$$fraction\ folded\ (i.e.\ with\ a\ pseudoknot) = \frac{R_{ter} + R_{terMg} + C}{R_{tot}}$$

$$fraction\ bound\ and\ folded = \frac{C}{R_{tot}}$$

$$fraction\ of\ binding\ competent\ RNA = \frac{R_{terMg} + C}{R_{tot}}$$

2.2.4 Model 3: A ligand-binding model for RNA that ignores the effect of tertiary interactions in the conformational capture mechanism but considers the effect of Mg^{2+} ions

This model, designated as *Model 3*, is schematically illustrated in **Fig. 3** (top panel). In this model, unlike the previous two models, the conformational selection relies on the formation of secondary structure only, and not on tertiary interactions. However, the model considers the effect of Mg^{2+} concentration explicitly. In other words, this model is similar to *Model 5* except that here the explicit Mg^{2+} concentration is taken into account.

FIG. 3. *Model 3* and its cognate thermodynamic cycle. The top panel illustrates a graphical reaction scheme for the proposed model designated as *Model 3*. The related Thermodynamic cycle is shown in the bottom panel. The model considers the differential binding between the

conformers with and without the pseudoknot as well as the effect of Mg^{2+} concentration. However, any RNA that contains the known secondary structure is assumed to bind with SAM.

Determination of Model 3 parameters

Assuming this model, the overall (apparent) dissociation constant using Isothermal calorimetry (ITC) which was reported by Gilbert *et al.* [4] ($Kd_{ITC} = 0.67 \pm 0.06\ \mu M$) would describe the following reaction,

$$Unbound\ RNA + SAM \xrightleftharpoons[Ka_{ITC}]{} bound\ RNA$$

$$[R_0 + R + R_{ter} + R_{terMg}] + [S] \xrightleftharpoons[Ka_{ITC}]{} [I + C]$$

$$[R_0 + R + R_{terE}] + [S] \xrightleftharpoons[Ka_{ITC}]{} [I + C]$$

Hence, this experimental dissociation constant describes the following equilibrium by,

$$Ka_{ITC} = \frac{Bound\ RNA}{Unbound\ RNA\ \times free\ SAM}$$

$$= \frac{[I + C]}{[R_0 + R + R_{ter} + R_{terMg}] \times [S]}$$

$$= \frac{[I + C]}{[R_0 + R + R_{terE}] \times [S]}$$

$$= \frac{\left[\left(\frac{1}{K_{fold}.Mg}\right).C + C\right]}{\left[\left(\frac{1}{K_{conv1}K_{terE}.Mg}\right)R_{terE} + \left(\frac{1}{K_{terE}.Mg}\right)R_{terE} + R_{terE}\right] \times [S]}$$

$$= \frac{C\left(\frac{1}{K_{fold}.Mg} + 1\right)}{R_{terE} \times S \times \left(\frac{1}{K_{conv1}K_{terE}.Mg} + \frac{1}{K_{terE}.Mg} + 1\right)}$$

$$= \frac{C \times \beta_1}{R_{terE} \times S \times \alpha_1}$$

Thus,

$$Ka_{ITC} = K_{bindE} \frac{\beta_1}{\alpha_1} \qquad\qquad \text{Eq. 10}$$

where, K_{bindE} is the SAM binding association constant of R_{terE}, $\beta_1 = \left(\frac{1}{K_{fold} \cdot Mg} + 1\right)$ and ,

$\alpha_1 = \left(\frac{1}{K_{conv1} \, K_{terE} \cdot Mg} + \frac{1}{K_{terE} \cdot Mg} + 1\right)$. The value of α_1 is given, and depends on the Mg^{2+}

ions concentration, whereas β_1 is unknown and cannot be determined from the thermodynamic

cycle alone.

Similarly,

$$Ka_{ITC} = \frac{Bound\ RNA}{Unbound\ RNA\ \times\ free\ SAM}$$

$$= \frac{[I + C]}{[R_0 + R + R_{ter} + R_{terMg}] \times [S]}$$

$$= \frac{[I + C]}{[R_0 + R + R_{terE}] \times [S]}$$

$$= \frac{[I + K_{fold} \cdot I \cdot Mg]}{\left[\frac{1}{K_{conv1}} R + R + K_{terE} \cdot Mg \cdot R\right] \times [S]}$$

$$= \frac{I(K_{fold} \cdot Mg + 1)}{R \times S \times \left(\frac{1}{K_{conv1}} + K_{terE} \cdot Mg + 1\right)}$$

$$= \frac{I \times \beta_2}{R \times S \times \alpha_2}$$

Thus,

$$Ka_{ITC} = K_{bind} \frac{\beta_2}{\alpha_2} \qquad\qquad \text{Eq. 11}$$

where, K_{bind} is the SAM binding association constant of R, $\alpha_2 = \left(\frac{1}{K_{conv1}} + K_{terE} \cdot Mg + 1\right)$ and $\beta_2 = \left(K_{fold} \cdot Mg + 1\right)$.

As in previous models, we consider that the previously reported FS studies monitored the RNA folding in the presence of the ligand relative to its absence [6]. In other words, the apparent binding constant determines the fraction of RNA species with a fully formed pseudoknot in the presence of the ligand relative to the ones with partially formed pseudoknot in the absence of the ligand. Therefore, the binding constants deduced from these FS studies, represent the following reaction,

$$\text{Folded \& unbound RNA} + \text{SAM} \underset{Ka_{FS}}{\Longleftrightarrow} \text{Folded \& bound RNA}$$

$$[R_{ter} + R_{terMg}] + [S] \underset{Ka_{FS}}{\Longleftrightarrow} [C]$$

$$[R_{terE}] + [S] \underset{Ka_{FS}}{\Longleftrightarrow} [C]$$

$$Ka_{FS} = \frac{Bound\ RNA\ species\ with\ a\ pseudoknot}{Unbound\ RNA\ species\ with\ a\ pseudoknot\ \times\ free\ SAM}$$

$$= \frac{[C]}{[R_{ter} + R_{terMg}] \times [S]} = \frac{[C]}{[R_{terE}] \times [S]} = K_{bindE}$$

Eq. 12

From Eq. 10 and Eq. 12, it can be observed that,

$$Ka_{ITC} = Ka_{FS}\ \frac{\beta_1}{\alpha_1}$$

$$\beta_1 = \frac{Ka_{ITC}\ \alpha_1}{Ka_{FS}} = 1 + \frac{1}{K_{fold}\ .\ Mg}$$

Thus,

$$Kd_{fold} = \left[\frac{Ka_{ITC}\ \alpha_1}{Ka_{FS}} - 1\right] Mg$$

$$= \left[\frac{Ka_{ITC}\ \alpha_1}{Ka_{FS}} - 1\right] Mg$$

Where, $\alpha_1 = \left(\frac{1}{K_{conv1}\ K_{terE}\ .\ Mg} + \frac{1}{K_{terE}\ .\ Mg} + 1\right).$ Eq. 13

Based on the Mg^{2+} binding model [10] we can therefore determine K_{terE} as follows,

$$K_{terE} = \frac{R_{terE}}{R \times Mg}$$

$$= \frac{R_{ter} + R_{terMg}}{R \times Mg}$$

$$= \frac{K_{conv2}.R + K_{conv2}.R.Mg.K_{Mg}}{R \times Mg}$$

$$= K_{conv2} \frac{1}{Mg} + K_{conv2}.K_{Mg}$$

$$= K_{conv2} \left(\frac{1}{Mg} + K_{Mg}\right)$$

Eq. 14

Since K_{conv2} and K_{Mg} are known, K_{terE} can be calculated. Consequently, Kd_{fold} for this model can be computed based on Ka_{ITC} and Ka_{FS}. Kd_{fold} can also be obtained with the aid of the thermodynamic cycle shown in **Fig. 3**). As in all models, the calculated values for the equilibrium constants are based on the model assumptions.

As in the previous models, after computing the equilibrium constants, the expression for C (given by Eq. 3.15, in the **Supplementary methods** and tabulated in **Table S2**) as well as all stable RNA species in the model can be calculated as a function of SAM concentration (see model derivation in the **Supplementary methods**). The *fraction bound, fraction folded*, *fraction both bound & folded* and *fraction binding competent* RNA, for this model, can be obtained from,

$$fraction\ bound = \frac{I + C}{R_{tot}}$$

$$fraction\ folded\ (i.e.with\ a\ pseudoknot) = \frac{R_{terE} + C}{R_{tot}}$$

$$fraction\ bound\ and\ folded\ = \frac{C}{R_{tot}}$$

$$fraction\ of\ binding\ competent\ RNA\ = \frac{R + R_{terE} + I + C}{R_{tot}}$$

2.2.5 Model 4: A ligand-binding model for RNA that considers the effect of tertiary interactions but ignores the role of Mg^{2+}

This model, designated as *Model 4*, is schematically illustrated in **Fig. 4**. This model assumes a conformational selection mechanism in which SAM binds only with RNA that possess a nucleate pseudoknot, hence, the effect of tertiary interactions is considered in this model. However, the binding of a single ligand, SAM, is taken into account and the effect of Mg^{2+} binding is assumed to be negligible. In effect, the model represents a simple classical two step conformational capture mechanism in the absence of Mg^{2+}. The equations for this model are derived in detailed in reference [10], except that the ligand used here is SAM. Therefore, the model assumes that only a single magnesium-independent pathway, for the riboswitch binding and folding, is possible.

FIG. 4. A Mg^{2+} independent, single pathway SAM binding Model (*Model 4*). The figure illustrates a graphical reaction scheme for the proposed model designated as *Model 4*. The model assumes a conformational capture mechanism based on the presence of a pseudoknot and assumes that the effect of Mg^{2+} concentration is negligible.

From the scheme and the definitions of $K_{bindter}$ and the definition of K_{FS} used in the above models, it can be seen that $K_{bindter} \approx K_{FS}$. Knowing the equilibrium constants the expression for the species *I* can be computed from Eq. 4.4 in the **Supplementary methods** (also tabulated in **Table S2**). All other stable RNA species in the model can also be calculated as a function of SAM concentration (see model derivation in the **Supplementary methods**). Thus, the *fraction bound, fraction folded*, *fraction both bound & folded* and *fraction binding competent* RNA, for this model are given by,

$$fraction\ bound\ = \frac{I}{R_{tot}}$$

$$fraction\ folded\ (i.e\ with\ a\ pseudoknot) = \frac{R_{ter} + I}{R_{tot}}$$

$$fraction\ bound\ and\ folded\ = \frac{I}{R_{tot}}$$

$$fraction\ of\ binding\ competent\ RNA\ = \frac{R_{ter} + I}{R_{tot}}$$

2.2.6   Model 5: A ligand-binding model for RNA that assumes ligand recognition is not influenced by tertiary interactions and ignores the concentration effect of Mg^{2+}

This model, designated as *Model 5,* is schematically illustrated in **Fig. 5** (top panel). In this model, the concentration effect of Mg^{2+} is assumed to be negligible and the observed effect of Mg^{2+} can be implicitly taken into account using an apparent equilibrium constant (K_{bindE}). Moreover, like *Model 3*, it is assumed that all RNA molecules with plausible secondary structures can bind with SAM, regardless of the presence of tertiary interactions. Nonetheless, the model takes into account that the binding affinities of the conformers that have tertiary interactions may differ from those that lack such interactions.

Fig. 5 schematically illustrates *Model 5* for SAM binding and folding of the SAM-II riboswitch. This model assumes that SAM binding is not influenced by the concentration of Mg^{2+} ions and assumes that the formation of the correct secondary structure is enough to form binding competent conformers. As described in section 2.2.1, the equilibrium constants for the model shown in (**Fig. 5**) are defined as follows. K_{conv1}, K_{terE}, K_{bind}, K_{bindE} and K_{fold} are the equilibrium constants for the conversion between (R_0 and R), between (R and R_{terE}), binding of the intermediates, binding of pseudoknot containing structures and folding of the bound intermediates (formation of pseudoknot in the presence of the ligand), respectively.

FIG. 5. *Model 5* **and its cognate thermodynamic cycle.** The top panel illustrates a graphical reaction scheme for the proposed model designated as *Model 5*. The related Thermodynamic cycle is shown in the bottom panel. The model considers the differential binding between the conformers with and without the pseudoknot but assumes that the effect of Mg^{2+} concentration is negligible.

Determination of Model 5 parameters

To be able to apply this model, the equilibrium constants K_{conv1} , K_{terE}, K_{bind}, K_{bindE} and K_{fold} needs to be estimated from existing experimental values. The equilibrium constant K_{conv1} can be obtained from the calculated binding energies and from the probabilities of the binding *competent* species (R) and binding *incompetent* (R_0) species. The model considers the effect of differential binding of riboswitch with pseudoknot (R_{terE}) and intermediate which is

the riboswitch without a pseudoknot (R) into account. Since Mg^{2+} concentration is assumed negligible (or with constant effect), K_{terE} can be calculated as the ratio between $\frac{R_{terE}}{R}$.

To determine K_{terE} at a given fixed magnesium concentration it is useful to not that,

$$K_{terE} = \frac{RNA\ with\ the\ accessible\ binding\ contacts\ and\ with\ pseudoknot\ in\ the\ presence\ of\ Mg}{RNA\ with\ the\ accessible\ binding\ contacts\ and\ without\ pseudoknot\ in\ the\ presence\ of\ Mg}$$

$$= \frac{fraction\ binding\ competent\ at\ a\ given\ Mg\ concentration}{R}$$

$$= \frac{R_{terE}}{R} = \frac{R_{ter} + R_{terMg}}{R} = \frac{R_{ter}}{R} + \frac{R_{terMg}}{R} = K_{conv2} + \left(K_{conv2} \times \frac{R_{terMg}}{R_{ter}} \right)$$

where, K_{conv2} and $\frac{R_{terMg}}{R_{ter}}$ can be computed based on Mg^{2+} data and model, as shown in our initial study [10], at specific Mg^{2+} concentrations (the total Mg^{2+} concentration is assumed to be 2 mM). The accessible contacts in the above equation refer to the population with the correct secondary structure. This assumes that the riboswitch will not be affected by the magnesium concentration (i.e. Mg^{2+} concentration has no effect on the R_{terE}). The consequence of the latter assumption is that the fraction having a pseudoknot in the absence of SAM is fixed and generally independent of the Mg^{2+} concentration. The latter approximation/assumption may become valid if, for example, the consumption of R_{terE} is very limited i.e. if the binding of R_{terE} is very weak.

To ease the link between the model parameters and the measured (apparent) experimental data, we define K_2 be the equilibrium determined from experimentally measured probabilities (presence and absence of pseudoknot), in the presence of a fixed Mg^{2+} concentration. Thus, from the definition of equilibrium of K_{terE} (Eq. 5.2 in the **Supplementary methods**, and illustrated in **Fig. 5**) we have,

$$K_2 = \frac{RNA\ with\ pseudoknot}{RNA\ without\ pseudoknot}$$

$$= \frac{R_{terE}}{R`} = \frac{R_{terE}}{R_0 + R} = \frac{R_{terE}}{R\left(\frac{1}{K_{conv1}}+1\right)} = K_{terE} \frac{1}{\left(\frac{1}{K_{conv1}}+1\right)} \qquad \text{Eq. 15}$$

Like in previous models, we make use of the fact that the overall (apparent) dissociation constant was determined experimentally using Isothermal calorimetry (ITC), reported by Gilbert *et al.* [4], Kd_{ITC} was experimentally determined to be $0.67 \pm 0.06\ \mu M$.

Assuming this model, the latter apparent reaction equilibrium constant would describe the following reaction,

$$Unbound\ RNA + SAM \xrightleftharpoons[Ka_{ITC}]{} bound\ RNA$$

$$[R_0 + R + R_{terE}] + [S] \xrightleftharpoons[Ka_{ITC}]{} [I + C]$$

$$[R` + R_{terE}] + [S] \xrightleftharpoons[Ka_{ITC}]{} [I + C]$$

Hence, this experimental dissociation constant describes the following equilibrium by,

$$Ka_{ITC} = \frac{Bound\ RNA}{Unbound\ RNA\ \times free\ SAM}$$

$$= \frac{[I + C]}{[R_0 + R + R_{terE}] \times [S]}$$

$$= \frac{[I + C]}{[R` + R_{terE}] \times [S]}$$

$$= \frac{\left[\left(\frac{1}{K_{fold}}\right) . C + C\right]}{\left[\left(\frac{1}{K_2}\right) . R_{terE} + R_{terE}\right] \times [S]}$$

$$= \frac{C\left(\frac{1}{K_{fold}} + 1\right)}{R_{terE} \times S\left(\frac{1}{K_2} + 1\right)}$$

$$= \frac{C \times \beta_1}{R_{terE} \times S \times \alpha_1}$$

Thus,

$$Ka_{ITC} = K_{bindE}\ \frac{\beta_1}{\alpha_1} \qquad\qquad \text{Eq. 16}$$

where, K_{bindE} is the SAM binding association constant of R_{terE} , $\alpha_1 = \left(\frac{1}{K_2} + 1\right)$ and $\beta_1 = \left(\frac{1}{K_{fold}} + 1\right)$. It is worth noting that α_1 can be re-written in terms of K_{terE} and K_{conv1} by substituting for K_2 (Eq. 15) to yield in $\alpha_1 = \frac{1}{K_{terE}\ K_{conv1}} + K_{terE} + 1$. The value of K_{conv1} is readily available from base pairing probabilities and K_{terE} can be computed at a specific Mg^{2+}

concentration. Thus, the value of α_1 can be obtained at an assumed Mg^{2+} concentration, whereas β_1 is unknown and cannot be determined from the thermodynamic cycle alone.

Similarly (and alternatively),

$$Ka_{ITC} = \frac{Bound\ RNA}{Unbound\ RNA\ \times free\ SAM}$$

$$= \frac{[I + C]}{[R_0 + R + R_{terE}] \times [S]}$$

$$= \frac{[I + C]}{[R^{`} + R_{terE}] \times [S]}$$

$$= \frac{[I + K_{fold}.I]}{[R + K_{terE}.R] \times [S]}$$

$$= \frac{I(1 + K_{fold})}{R \times S(1 + K_{terE})}$$

$$= \frac{I \times \beta_2}{R \times S \times \alpha_2}$$

Thus,

$$K_{ITC} = K_{bind}\ \frac{\beta_2}{\alpha_2} \qquad \text{Eq. 17}$$

where, K_{bind} is the SAM binding association constant of R, $\alpha_2 = (1 + K_{terE})$ and $\beta_2 = (1 + K_{fold})$.

It should be noted that, the value of the parameter K_{bindE} in this model is considered to be equivalent to the constant that has been reported experimentally by Haller *et al* [6], that is we have $K_{bindE} \approx K_{FS}$. The latter was used in subsequent calculations. Thus, from the above expression for Ka_{ITC}, K_{fold} is given by,

$$K_{fold} = \frac{1}{\frac{Kd_{bindE}.\alpha_1}{Kd_{ITC}} - 1} = \frac{1}{\frac{Kd_{FS}.\alpha_1}{Kd_{ITC}} - 1} \qquad \text{Eq. 18}$$

Hence, $Kd_{fold} = \frac{1}{K_{fold}}$ and $dG_{fold} = -RT \ln(K_{fold})$ can be obtained from Eq. 18. Since three of the four parameters are known, then the fourth can be determined through a thermodynamic cycle.

Hence, we construct a thermodynamic cycle (**Fig. 5**, bottom panel), *where*,

$$\Delta G_{bind} + \Delta G_{fold} + (-\Delta G_{bindE}) + (-\Delta G_{Conv2}) = 0$$

By rearrangement, dG_{bind} can be computed from,

$$\Delta G_{bind} = -\Delta G_{fold} - (-\Delta G_{bindE}) - (-\Delta G_{Conv2})$$

As all other models, given the equilibrium constants, the expression for C (given by Eq. 5.15, in the **Supplementary methods** and tabulated in **Table S2**) as well as all stable RNA species can be calculated as a function of SAM concentration. The *fraction bound, fraction folded, fraction both bound & folded* and *fraction binding competent* RNA, for this model, can be obtained from,

$$fraction\ bound = \frac{I + C}{R_{tot}}$$

$$fraction\ folded\ (i.e.\ with\ a\ pseudoknot) = \frac{R_{terE} + C}{R_{tot}}$$

$$fraction\ bound\ and\ folded = \frac{C}{R_{tot}}$$

$$fraction\ of\ binding\ competent\ RNA = \frac{R + R_{terE} + I + C}{R_{tot}}$$

2.2.7 *Model 6: A ligand-binding model for RNA that assuming that SAM recognition is not influenced by tertiary interactions and occurs in the absence of Mg²⁺ ions*

The model, designated as *Model 6*, is schematically illustrated in **Fig. 6** (top panel). In this model, the impact of Mg^{2+} is assumed negligible, and the magnesium bound species are not represented in the model. Despite that the latter assumption contradicts previously reported experimental data, it is still instructive to derive and study this model for the purpose of comparisons with other models, as well as, for confirming the most probable mechanism by excluding other possibilities. As in the previous model, here, the tertiary interactions are

assumed to have no major role in SAM recognition. In other words, it is assumed that all RNA molecules with plausible secondary structures (and not necessarily any specific tertiary interactions) can bind with SAM in the absence of magnesium.

FIG. 6. *Model 6* **and its cognate thermodynamic cycle.** The top panel illustrates a graphical reaction scheme for the proposed model designated as *Model 6.* The related Thermodynamic cycle is shown in the bottom panel. The model assumes that SAM can bind to all RNA that has the minimal secondary structure. In other words, SAM binding occurs in the presence or absence of tertiary interactions, with no effect of Mg^{2+}.

Determination of Model 6 parameters

Again, as in other models, that fact that the overall (apparent) dissociation constant had been determined experimentally using Isothermal calorimetry (ITC), and was reported by Gilbert *et*

al [4] is used to determine the model parameters. In the latter study, the authors reported a dissociation constant of Kd_{ITC} of $0.67 \pm 0.06\ \mu M$.

If this model, *Model 6,* was true, then the latter apparent reaction association equilibrium constant $(Ka_{ITC} = \frac{1}{Kd_{ITC}})$ would describe the following reaction,

$$Unbound\ RNA + SAM \underset{Ka_{ITC}}{\Longleftrightarrow} bound\ RNA$$

$$[R_0 + R + R_{ter}] + [S] \underset{Ka_{ITC}}{\Longleftrightarrow} [I + C]$$

$$[R^{\grave{}} + R_{ter}] + [S] \underset{Ka_{ITC}}{\Longleftrightarrow} [I + C]$$

Hence, the experimental association constant Ka_{ITC} would be then described by the following equilibrium,

$$Ka_{ITC} = \frac{Bound\ RNA}{Unbound\ RNA\ \times free\ SAM}$$

$$= \frac{[I + C]}{[R_0 + R + R_{ter}] \times [S]}$$

$$= \frac{[I + C]}{[R^{\grave{}} + R_{ter}] \times [S]}$$

By substituting for the equilibrium constants, as defined in the **Supplementary methods**, Equations S6.1 to S6.5, we therefore obtain,

$$= \frac{\left[\left(\frac{1}{K_{fold}}\right).C + C\right]}{\left[(\frac{1}{K_{ter}}).R_{ter} + R_{ter}\right] \times [S]}$$

$$= \frac{C\left(^{1}/_{K_{fold}} + 1\right)}{R_{ter} \times S\left(^{1}/_{K_{ter}} + 1\right)}$$

$$= \frac{C \times \beta_1}{R_{ter} \times S \times \alpha_1}$$

Thus,

$$Ka_{ITC} = K_{bindter} \frac{\beta_1}{\alpha_1} \qquad \text{Eq. 19}$$

where, $K_{bindter}$ is the SAM binding association constant of R_{ter} , $\alpha_1 = \left(\frac{1}{K_{ter}} + 1\right)$ and $\beta_1 = \left(\frac{1}{K_{fold}} + 1\right)$. The value of α_1 is given, and according to the Mg^{2+} ions concentration, whereas β_1 is unknown and cannot be determined from the thermodynamic cycle alone.

Similarly (and alternatively),

$$Ka_{ITC} = \frac{Bound\ RNA}{Unbound\ RNA\ \times free\ SAM}$$

$$= \frac{[I + C]}{[R_0 + R + R_{ter}] \times [S]}$$

$$= \frac{\left[I + K_{fold}.I\right]}{[(1/K_{conv1})R + R + K_{conv2}.R] \times [S]}$$

$$= \frac{I\left(K_{fold} + 1\right)}{R \times S\left((1/K_{conv1}) + K_{conv2} + 1\right)}$$

$$= \frac{I \times \beta_2}{R \times S \times \alpha_2}$$

Thus,

$$Ka_{ITC} = K_{bind} \frac{\beta_2}{\alpha_2} \qquad \text{Eq. 20}$$

where, K_{bind} is the SAM binding association constant of R, $\alpha_2 = \left((1/K_{conv1}) + K_{conv2} + 1\right)$ and $\beta_2 = \left(K_{fold} + 1\right)$.

As in the previous model, the value of $K_{bindter} \approx K_{FS}$ is a parameter that has been reported experimentally by Haller *et al* [6], which was used in subsequent calculations. Hence, based on the above expression for Ka_{ITC} , K_{fold} is given by,

$$K_{fold} = \frac{1}{\frac{Kd_{bindter} \cdot \alpha_1}{Kd_{ITC}} - 1} = \frac{1}{\frac{Kd_{FS} \cdot \alpha_1}{Kd_{ITC}} - 1} \qquad \text{Eq. 21}$$

Hence, $Kd_{fold} = \frac{1}{K_{fold}}$ and $dG_{fold} = -RT\ln(K_{fold})$ can be computed from Eq. 21. Since three $(\Delta G_{fold}, \Delta G_{bindter}\ and\ \Delta G_{Conv2})$ of the four parameters are known, then the fourth can be determined through a thermodynamic cycle.

Hence, we construct a thermodynamic cycle (**Fig. 6**, bottom panel), *where*,

$$\Delta G_{bind} + \Delta G_{fold} + (-\Delta G_{bindter}) + (-\Delta G_{Conv2}) = 0$$

By rearrangement, dG_{bind} is given by,

$$\Delta G_{bind} = -\Delta G_{fold} - (-\Delta G_{bindter}) - (-\Delta G_{Conv2})$$

Given the equilibrium constants, the expression for C (given by Eq. 6.15, in the **Supplementary methods** and tabulated in **Table S2**) as well as all stable RNA species can be calculated as a function of SAM concentration. The *fraction bound, fraction folded, fraction both bound & folded* and *fraction binding competent* RNA, for this model, can be obtained from,

$$fraction\ bound = \frac{I + C}{R_{tot}}$$

$$fraction\ folded\ (i.e.with\ a\ pseudoknot) = \frac{R_{ter} + C}{R_{tot}}$$

$$fraction\ bound\ and\ folded = \frac{C}{R_{tot}}$$

$$fraction\ of\ binding\ competent\ RNA = \frac{R + R_{ter} + I + C}{R_{tot}}$$

3   RESULTS AND DISCUSSION

3.1   Determining model parameters (equilibrium constants) enables computing titration curves for the individual RNA species of the different models

To be able to predict the corresponding titration curves for each model, the model parameters were determined as described in the **Methods** and **Supplementary methods** sections). *Model 1* depends on five parameters (equilibrium constants): $K_{ter}, K_{Mg}, K_{bindter}, K_{bindMg}$ and K_{fold} (**Fig. 1,** top panel). As illustrated in **Fig. 1** (top panel) and in **Fig. S1**, K_{ter} is the apparent association equilibrium constant for the formation of the pseudoknot in the absence of Mg^{2+} ions. The latter constant depends on both K_{conv2} and K_{conv1} (**Fig. S1**). In the absence of tertiary

interactions, SAM and Mg^{2+} ions, the only two accessible states of the system are R_0 and R. The equilibrium ratio between the two spices is controlled by constant K_{conv1}. Hence, the latter constant can be determined from secondary structure calculations, by calculating the ratio between the probability of conformers that form a closed loop (i.e. the P1 helix is formed, denoted as species R) and those that do not have the P1 helix (denoted as species R_0). The calculations were performed using the method previously reported by Alaidi *et al.* [18,10]. Further, using the Mg^{2+} model described in our previous study [10], K_{ter} and K_{Mg} were determined. Thereafter, using one of the experimentally reported constants Kd_{ITC} [4] and Kd_{FS} [6], along with the thermodynamic cycle shown in **Fig. 1** (bottom panel), the remaining parameters ($K_{bindter}$, K_{bindMg} and K_{fold}) were determined as described in detail in the methods section. The determined equilibrium constants and/or corresponding free energies for this model are listed in **Table II**. The determination of the model parameters for *Models 2* and *Model 4* (**Fig. 2** and **4**), however, is much simpler and more straightforward compared to the other models. The two models generally resemble the 2 or 3 step equilibrium models [10] and their parameters a can be directly determined without the need to construct a thermodynamic cycle (as described in the **Methods** section).

Model 1	
Constant	**Value**
Given K_{Mg} from *Mg Model* [10]	878
Given K_{ter} based on *Mg Model* [10]	0.0819
Given Kd_{ITC} from experiment [4]	6.70E-07
Parameters in the presence of Mg^{2+} (kcal/mol):	
dG_{Mg}	-4.016
dG_{ter}	1.483
dG_{fold}	-3.696
$dG_{bindter}$	-9.608
dG_{bindMg}	-9.289
dG_{FS} (calculated from model)	-9.425

TABLE II. *Model 1 parameters.* The table lists the parameters computed for *Model 1*. Despite that SAM binding energies, in the presence or absence of Mg^{2+} (via dG_{bindMg} and $dG_{bindter}$, respectively) clearly play the major role in stabilizing the final folded conformer, a significant energetic contribution to the stability comes from both the Mg^{2+} binding and the Mg^{2+} mediated tertiary interactions (via dG_{ter} and dG_{fold}, respectively) that leads to pseudoknot stabilization

or zipping of the binding pocket. It should be noted that K_{ter} can be expressed in terms of K_{conv1} and K_{conv2} as described in detail in the *Mg Model* [10]. The free energies were calculated from the corresponding equilibrium constants $\Delta G_i = RT \ln(Kd_i) = -RT \ln(Ka_i)$, at room temperature (298.15 Kelvin).

On the contrary, determining the parameters *Models 3, 5* and *6* (**Fig. 3**, **5** and **6**, top panels) are not as straightforward, due to the differential binding affinities between the RNA species. The latter models, assume that tertiary interactions influence the degree of binding with SAM. These models therefore resemble a conformational capture mechanism that is based on differential binding. This may occur when SAM is loosely selective for conformers with tertiary interactions (nucleated pseudoknot), though still all conformers with correct secondary structure bind with significant affinity to SAM. More specifically, the models account for the difference in binding affinity between the riboswitch with and without the nucleated pseudoknot as well as the difference in the folding in the presence of the ligand compared to its absence. In other words, the model considers that both riboswitches lacking the pseudoknot (with the binding contacts accessible and are in close proximity, enforced by a stable P1 stem and L1 loop) and riboswitch having a nucleated pseudoknot bind to SAM, with different affinities. This discrepancy in the binding, in turn, results in a difference in the energy of the formation of the pseudoknot (i.e. a difference in the folding energy) in the presence and absence of the ligand.

Given the limited available experimental data, as in *Model 1*, constructing thermodynamic cycles was needed to determine all the parameters for these models (**Fig. 3**, **5** and **6**, bottom panels). In *Models 3* and *5*, as in previous attempts [10], it was assumed that Mg^{2+} level is almost constant in the cell. On the one hand, in *Model 5* the system is modeled at a fixed Mg^{2+} concentration. On the other hand, *Model 3* makes it possible to study the system at different concentrations of Mg^{2+}. In the latter models, it is assumed that the formation of the P1 helix allows SAM contacts as well as the residues that form the pseudoknot in the loop (L1) to be in close proximity to the ligand. Hence, all structures with the expected RNA secondary structure are assumed to be binding competent. Based on the previous experimental data and models [7,8,10], the equilibrium value K_{conv2} at 2 mM Mg^{2+} can be obtained. Further, K_{bindE} or $K_{bindter}$ has been experimentally measured by FS experiments. From this value and from the experimentally measured overall binding Kd_{ITC} the equilibrium constant K_{fold} can be determined (see methods for detailed derivation). Finally, K_{bind} can be calculated with the aid

of the corresponding thermodynamic cycles (**Fig. 3, 5** and **6**, bottom panels). **Table S1** lists all parameters calculated and used for *Models 2-6*.

One key concept we have used in this study is expressing the apparent binding in terms of the binding constants that are specific to each model. Hence, the observed *fraction bound* can incorporate different species or conformers that vary according to the model. This allowed us to establish links between the apparent experimental measured quantities with the different conformers proposed in the model, enabling the computation of the model equilibrium constants. If the model correctly reflects the underlaying mechanism, then it should be able to predict other independent experimental observations. For instance, here we used binding data, to compute the parameters for all models. As a result, the fraction bound is the same, and certain individual species may be the same, but the fraction folded is the key difference. We then tested the predicted fraction folded against the experimentally observed fraction that formed a pseudoknot at a specific concentration of the SAM and Mg^{2+}. Further, we have taken into account the methods by which binding was measured. Specifically, whereas ITC measurements evaluate the Kd based on the bound and free molecules, several previously reported FS-based apparent Kd for riboswitches [6,19] were based on the local or global RNA structural changes, observed as An increase in the fluorescence signal change, upon ligand binding. Hence, the interpretation of the apparent Kd by each technique is different.

3.2 The comparison of predicted titration curves from the SAM binding models against experimental observations reveals that *Model 1* is the most probable model by which the SAM-II riboswitch function

For each model, the fractions of *individual RNA species*, as well as the *fraction bound*, the *fraction folded*, the *fraction bound & folded,* and the *fraction of binding competent RNA* have been determined as a function of SAM. To be able to determine which model truly represents how the SAM-II riboswitch works, we compared an experimentally observed fraction at certain conditions with the one predicted from various models. Specifically, it has been shown experimentally by single molecule FRET, that in the presence of 10 µM SAM (depicted by the vertical dashed line in **Fig. 7A**) and 2 mM Mg^{2+}, increased the folded by ten-fold [6]. Since in the absence of both ligands, the fraction of conformers that forms a pseudoknot is ~ 9.3-9.5% (as reported in [8], and ~ 9.34% after fitting to Mg^{2+} data as discussed [10]), then the expected fraction having a partially or fully formed pseudoknot (*fraction folded*) would be ~ 93-95 % (specifically, 93.4%). **Fig. 7** shows the *fraction folded*, as well as the *fraction bound*, the *Bound*

and folded fraction and the *fraction of binding competent RNA* as a function of increasing SAM concentration (panels **A-D,** respectively), at the experimental level of Mg^{2+}. The only models that give a *fraction folded* that is very similar to the latter FRET experiment are *Model 1* and *Model 2,* predicting that 94.64% (and 94.62% when accounting for non-specific Mg^{2+}) of the RNA population is folded (with a pseudoknot) at these experimental conditions, as opposed to 83.18%, 85.05%, 81.84% and 34.35% (depicted by the horizontal dashed line in **Fig. 7A**) by *Models 3-6*, respectively.

At this experimental Mg^{2+} concentrations both *Model 1* and *Model 2* overlap. However, at lower Mg^{2+} concentrations, as shown in **Fig. 7**, panels **E** and **F**, the two models give significantly different results. The presence of an alternative binding pathway in *Model 1* makes the folding in the presence of SAM less sensitive to the level of Mg^{2+}, as opposed to *Model 2* in which the pseudoknot can only be formed in the presence of Mg^{2+}. Several studies, however, have reported that at high concentrations of SAM most of the SAM-II RNA still form a pseudoknot and becomes folded, even in the absence of Mg^{2+} [6,7]. The latter findings are therefore consistent with *Model 1* and not with *Model 2*. Since *Model 1* also incorporates the previously discussed concentration dependent *Mg^{2+} Model* and its deduced parameters [10] (unlike *Models 3,4,6*), it is the only model that is qualitatively and quantitatively consistent with structural (population) changes observed in the absence of any ligands, after adding Mg^{2+} only, after adding SAM only and after adding both Mg^{2+} and SAM. Such data was collected using multiple techniques [4,6–8].

Thus, *Model 1* provides the best explanation to the experimental qualitative and quantitative measurements both at high (or in the presence) and at low (or in the absence) of Mg^{2+} concentrations, compared to the other Models. Strikingly, the comparison between the models and experimental data illustrated that natural riboswitches exhibits the most efficient mechanism, at various levels of Mg^{2+}, allowing the RNA to act as a highly efficient sensor and with a broad sensitivity-range. Thus, in the following sections of this manuscript, only *Model 1* will be discussed further.

FIG 7. Titration curves for the SAM-II riboswitch assuming different models. The curves were simulated using a total SAM concentration of 10 μM. Panels **A** to **D**, show the simulations results at a fixed total magnesium concentration of 2.0 mM using the different models. The panels show the computed *fraction folded*, *fraction bound*, *the bound and folded fraction* and the *fraction of binding competent RNA*, respectively, for all SAM *Models 1-6*. The dashed red and black lines (in panel **A**) depict the level of SAM and corresponding folded RNA in the previously reported single molecule FRET experiment [6] that is used to contrast with the predictions from the models. Panels **E** and **F**, show the computed *fraction folded* and *fraction bound*, respectively, at a much lower total magnesium concentration (20 μM) using the same models. All simulations were performed using an RNA concentration of 0.5 μM.

3.3 The binding and folding mechanism of the SAM-II riboswitch

Model 1 shows that the aptamer pseudoknot of the SAM-II riboswitch, is initially stabilized by the magnesium ion to form a partially stable aptamer in an open conformation, paving the way for SAM to bind. The binding of SAM then closes the binding pocket (forms a closed aptamer conformation, as observed in the crystal structure [4]) and further stabilizes the aptamer, essentially sequestering the ribosome binding site, and making it inaccessible for the Ribosome by a large free energy barrier. An alternative pathway for the riboswitch binding and folding may occur, in which SAM binds before magnesium. In either case, it is evident that the binding of both molecules is cooperative in nature (**Fig. 8C**). Since magnesium ions are usually present in the cell during transcription, the dominant product is dependent on the Mg concentration. In the presence of 2 mM magnesium, which was used in several *in vitro* experiments and also a relevant cell concentration (see [10] and references therein for a detailed discussion), approximately half of the SAM bound, and folded population is also bound to Mg^{2+} (depicted as species C) and the other half is Mg^{2+} free (depicted species I). Thus, the fact that the crystal structure of the SAM-II riboswitch did not show a bound Mg^{2+} ion [4], whereas a very similar SAM-V riboswitch [20] showed a bound Mg^{2+} ion, can be explained by the notion that both states co-exist, even in the abundance of SAM.

FIG. 8. **Titration curves for individual RNA species as a function of SAM concentration from *Model 1*.** The plots in panels **A** and **B** illustrate the fractions and concentrations of the

different RNA species computed from *Model 1*, as a function of total SAM concentration at a fixed total Mg^{2+} concentration of 2.0 mM and 0.5 μM RNA. Similar titration curves for individual RNA species computed based on this and other models discussed in this study (*Model 2* to *6*) are illustrated in **Fig. S2**. The plots in panels **C** and **D** illustrate the *fraction bound* and *fraction folded* of the riboswitch as a function of SAM concentration, for different concentrations of Mg^{2+} ions, using the same model (*Model 1*). The latter titration curves illustrate that the magnesium level determines the sensitivity range of the riboswitch to its cognate ligand (SAM), by setting a threshold for the fraction of the already folded riboswitch.

As shown previously based on Mg^{2+} data, the formation of the R_{ter} is not favorable, with a free energy of 1.483 kcal/mol. R_{ter}, however, can be stabilized by the binding of Mg^{2+} to form R_{terMg} with a -4.016 kcal/mol. SAM can bind to both R_{ter} and R_{terMg} with a stabilizing binding energies of -9.608 and -9.289 kcal/mol, respectively, to form I and C, which represent the open and closed folded aptamer conformations. Finally, a Mg^{2+} ion may further stabilize I (the open conformation) with a free energy of -3.696 kcal/mol by zipping the binding pocket, converting it to the closed conformation C. The expected SAM binding using single molecule FS (dG_{FS}) based on *Model 1* is -9.425 kcal/mol, which is very close to the experimentally reported value -9.350 kcal/mol at room temperature. Notably, this model seems to the most efficient compared to the other models (**Fig. 7A**). That is, the fraction folded (i.e. forming a pseudoknot and hence essentially sequestering the ribosome binding site) can reach the highest at expected levels of SAM and Mg^{2+}.

Interestingly, the concentration of Mg^{2+} achieves a certain level of folding prior to SAM binding, allowing the riboswitch to strongly respond to SAM only if the latter exceeds a certain threshold concentration. Such property is probably to be desirable in the cell and had been often discussed in transcriptional or kinetically driven riboswitches [21,22]. In other words, Mg^{2+} controls the sensitivity range of the riboswitch to the primary ligand. **Fig. 8B** and **C** show the *fraction folded* and *fraction bound* of the RNA (simulated using *Model 1*) as a function of SAM for different Mg^{2+} levels. It is evident that the sensitivity range of the riboswitch decreases as Mg^{2+} ions increase.

FIG 9. The free energy landscape of the SAM-II riboswitch. The figure illustrates the free energy landscape of the SAM-II riboswitch with corresponding standard free energy differences between a give state and the neighbouring one are shown on the arrows. The initial unfolded or incorrectly folded state (R_0) here acts as the reference. The values of the free energy differences shown on the figure are tabulated in **Table II** and are further discussed in the text.

3.4 Prediction of the fraction of riboswitch folded as a function of both Mg^{2+} and SAM: *The development of a ligand-induced conformational shift simulation method*

The determined binding and folding free energies (which are the model parameters) allow us to construct a standard binding and folding free energy landscape of the system (**Fig. 9**), which provides insights about the system, at standard conditions. Such energies can be thought of as the molar free energy differences, at standard room temperature and pressure, and at saturating concentrations of the ligands. It is evident from the figure that one can determine the standard free energy difference between any two states of the system, for example R and I, even if the conversion between these states must go through one or more intermediate.

More importantly, the model can be used to quantitatively predict the probabilities of all species in the ensemble at varying concentrations of two or three reactants. For example, the three-dimensional (3D) titration curves in **Fig. S3** and **Fig. S4,** respectively, illustrate the changes of the various fractions (or concentrations) of RNA species as a function of 1) SAM and Mg^{2+} concentrations, at a fixed total RNA concentration, and 2) total RNA and Mg^{2+} concentrations, at a fixed total SAM concentration. This study therefore shows a case where concentration dependent mechanistic/mathematical models were combined with conformer populations (ensembles calculations based on secondary structure energetics) and linked to multiple experimental ligand binding studies to provide a rich and detailed understanding of the system. It is worth noting that the methods described can also be used to compute ligand titration curves

at different transcription lengths. More generally, when basic individual conformer probabilities along with models for single or multiple Mg^{2+} ions [10,18] are considered, it is possible to directly apply or adapt the mathematical framework of the models in this study to any RNA system (including other riboswitches) that undergoes structural changes upon ligand binding.

3.5 Implications on targeting Riboswitches for therapeutics

Noncoding RNA has been proposed as potential drug target in prokaryotic and eukaryotic cells [23,24]. However, to date, no rationally designed small molecules that target non-ribosomal RNA exists in the market. Achieving such a goal is hindered by many factors including the fact that our understanding of RNA functional mechanisms remains limited. Riboswitches are one of the RNA classes that have the potential to be perceived as targets for small molecule drugs [11,12,25–27]. Arguably, understanding riboswitch functional mechanisms has direct implications on the strategy to undertake if one wishes to design small molecules that can specifically target riboswitches in bacteria and fungi. One obvious strategy is to find molecules that can mimic, and subsequently block, the cognate ligands from binding. Novel strategies for targeting RNA may involve, for instance, targeting stable intermediates, cryptic pockets, or allosteric binding sites. Thus, more broadly, thermodynamic mechanisms highlight various aspects of RNA functions, and which can facilitate targeting RNA with small molecule inhibitors.

The mechanism deduced in this study for the SAM-II riboswitch, described in *Model 1*, illustrates the importance of tertiary interactions in the cognate ligand recognition and the critical role of these interactions in producing and stabilizing the finally folded and functional aptamer. Only the RNA with previously nucleated tertiary interactions will produce the fully folded riboswitch. The fact that SAM binds specifically to only molecules with a nucleated or fully formed pseudoknot indicates the importance of such intermediates as drug targets and that inhibiting the tertiary interactions can be a strategy for drug design. These conclusions for the SAM-II riboswitch are consistent with the several studies that have shown that mutations that inhibit the formation of major tertiary interactions (such as those that lead to the formation of a pseudoknot) perturb the normal function of riboswitches [28–31], suggesting the possibility that such mechanism is general and plausible for other riboswitch classes.

4   CONCLUSION

This study deconvolutes the mechanism by which the SAM-II riboswitch functions. In the latter mechanism, RNA with a pre-existing pseudoknot binds tightly with SAM, in the presence or the absence of magnesium ions, and hence become more stable, eventually dominating the RNA population. The level of magnesium ions sets the concentration threshold and range in which the riboswitch can be responsive to SAM. It is evident that, such mechanism may be similar for other riboswitches, paving the way for practical and novel avenues in drug design. The presented analytical/mathematical framework, based on linked equilibria model, which combines secondary structure energetics with ligand binding, though specifically used here to quantitatively model the SAM-II Riboswitch, is yet applicable to other riboswitches. Moreover, the presented models and methods that failed to describe the behavior of this riboswitch, are still generally applicable to other RNA that undergoes conformational changes in the presence of one or more ligands. We anticipate that the mechanisms and tools, contributed via this work, will advance our understanding of RNA function, and facilitate the ultimate goal of targeting RNA by small molecules.

5   SUPPLEMENTARY MATERIAL

Supplementary material including Supplementary methods, figures and tables are available online.

6   AUTHOR DECLARATIONS

Conflict of Interest

The author declares that he has shares in Biocomplexity for Research and Consulting, which is a for-profit company.

7   DATA AVAILABILITY

The data that supports the findings of this article are available within the article and its supplementary material.

8   REFERENCES

[1] A. Roth, and R.R. Breaker, "The structural and functional diversity of metabolite-binding riboswitches," Annu Rev Biochem **78**, 305–334 (2009).
[2] K. Kavita, and R.R. Breaker, "Discovering riboswitches: the past and the future," Trends in Biochemical Sciences **48**(2), 119–141 (2023).

[3] K.A. Corbino, J.E. Barrick, J. Lim, R. Welz, B.J. Tucker, I. Puskarz, M. Mandal, N.D. Rudnick, and R.R. Breaker, "Evidence for a second class of S-adenosylmethionine riboswitches and other regulatory RNA motifs in alpha-proteobacteria," Genome Biol **6**(8), R70 (2005).

[4] S.D. Gilbert, R.P. Rambo, D. Van Tyne, and R.T. Batey, "Structure of the SAM-II riboswitch bound to S-adenosylmethionine," Nat Struct Mol Biol **15**(2), 177–182 (2008).

[5] J.M. Kelley, and D. Hamelberg, "Atomistic basis for the on–off signaling mechanism in SAM-II riboswitch," Nucleic Acids Research **38**(4), 1392–1400 (2010).

[6] A. Haller, U. Rieder, M. Aigner, S.C. Blanchard, and R. Micura, "Conformational capture of the SAM-II riboswitch," Nature Chemical Biology **7**(6), 393–400 (2011).

[7] B. Chen, X. Zuo, Y.-X. Wang, and T.K. Dayie, "Multiple conformations of SAM-II riboswitch detected with SAXS and NMR spectroscopy," Nucleic Acids Res **40**(7), 3117–3130 (2012).

[8] B. Chen, R. LeBlanc, and T.K. Dayie, "SAM-II Riboswitch Samples at least Two Conformations in Solution in the Absence of Ligand: Implications for Recognition," Angew Chem Int Ed Engl **55**(8), 2724–2727 (2016).

[9] S. Roy, H. Lammert, R.L. Hayes, B. Chen, R. LeBlanc, T.K. Dayie, J.N. Onuchic, and K.Y. Sanbonmatsu, "A magnesium-induced triplex pre-organizes the SAM-II riboswitch," PLoS Comput Biol **13**(3), e1005406 (2017).

[10] O. Alaidi, "Modeling Riboswitches: The impact of magnesium ions concentration on the folding of the SAM-II riboswitch," Co-submitted, (2024).

[11] K.F. Blount, and R.R. Breaker, "Riboswitches as antibacterial drug targets," Nat Biotechnol **24**(12), 1558–1564 (2006).

[12] K.F. Blount, J.X. Wang, J. Lim, N. Sudarsan, and R.R. Breaker, "Antibacterial lysine analogs that target lysine riboswitches," Nat Chem Biol **3**(1), 44–49 (2007).

[13] J.C. Venter, K. Remington, J.F. Heidelberg, A.L. Halpern, D. Rusch, J.A. Eisen, D. Wu, I. Paulsen, K.E. Nelson, W. Nelson, D.E. Fouts, S. Levy, A.H. Knap, M.W. Lomas, K. Nealson, O. White, J. Peterson, J. Hoffman, R. Parsons, H. Baden-Tillson, C. Pfannkoch, Y.-H. Rogers, and H.O. Smith, "Environmental genome shotgun sequencing of the Sargasso Sea," Science **304**(5667), 66–74 (2004).

[14] X. Xue, W. Yongjun, and L. Zhihong, "Folding of SAM-II riboswitch explored by replica-exchange molecular dynamics simulation," J Theor Biol **365**, 265–269 (2015).

[15] J.M. Jean, and K.B. Hall, "2-Aminopurine fluorescence quenching and lifetimes: Role of base stacking," Proceedings of the National Academy of Sciences **98**(1), 37–41 (2001).

[16] R. Rieder, K. Lang, D. Graber, and R. Micura, "Ligand-Induced Folding of the Adenosine Deaminase A-Riboswitch and Implications on Riboswitch Translational Control," ChemBioChem **8**(8), 896–902 (2007).

[17] K. Lang, R. Rieder, and R. Micura, "Ligand-induced folding of the thiM TPP riboswitch investigated by a structure-based fluorescence spectroscopic approach," Nucleic Acids Res **35**(16), 5370–5378 (2007).

[18] O. Alaidi, and F. Aboul-Ela, "Statistical mechanical prediction of ligand perturbation to RNA secondary structure and application to riboswitches," J Comput Chem **41**(16), 1521–1537 (2020).

[19] B. Heppell, S. Blouin, A.-M. Dussault, J. Mulhbacher, E. Ennifar, J.C. Penedo, and D.A. Lafontaine, "Molecular insights into the ligand-controlled organization of the SAM-I riboswitch," Nat Chem Biol **7**(6), 384–392 (2011).

[20] L. Huang, and D.M.J. Lilley, "Structure and ligand binding of the SAM-V riboswitch," Nucleic Acids Res **46**(13), 6869–6879 (2018).

[21] J.K. Wickiser, M.T. Cheah, R.R. Breaker, and D.M. Crothers, "The Kinetics of Ligand Binding by an Adenine-Sensing Riboswitch," 13404–13414 (2005).

[22] J.K. Wickiser, W.C. Winkler, R.R. Breaker, and D.M. Crothers, "The speed of RNA transcription and metabolite binding kinetics operate an FMN riboswitch.," Molecular Cell **18**(1), 49–60 (2005).

[23] C. Sheridan, "First small-molecule drug targeting RNA gains momentum.," Nat Biotechnol **39**(1), 6–8 (2021).

[24] J.L. Childs-Disney, X. Yang, Q.M.R. Gibaut, Y. Tong, R.T. Batey, and M.D. Disney, "Targeting RNA structures with small molecules," Nat Rev Drug Discov **21**(10), 736–762 (2022).

[25] K.E. Deigan, and A.R. Ferré-D'Amaré, "Riboswitches: discovery of drugs that target bacterial gene-regulatory RNAs," Acc Chem Res **44**(12), 1329–1338 (2011).

[26] C.E. Lünse, A. Schüller, and G. Mayer, "The promise of riboswitches as potential antibacterial drug targets," Int J Med Microbiol **304**(1), 79–92 (2014).

[27] Y.W. Ham, D.J. Humphreys, S. Choi, and D.L. Dayton, "Rational design of SAM analogues targeting SAM-II riboswitch aptamer," Bioorg Med Chem Lett **21**(17), 5071–5074 (2011).

[28] C. Lu, F. Ding, A. Chowdhury, V. Pradhan, J. Tomsic, W.M. Holmes, T.M. Henkin, and A. Ke, "SAM recognition and conformational switching mechanism in the Bacillus subtilis yitJ S box/SAM-I riboswitch.," Journal of Molecular Biology **404**(5), 803–18 (2010).

[29] A. Lussier, L. Bastet, A. Chauvier, and D.A. Lafontaine, "A kissing loop is important for btuB riboswitch ligand sensing and regulatory control," J Biol Chem **290**(44), 26739–26751 (2015).

[30] U. Rieder, K. Lang, C. Kreutz, N. Polacek, and R. Micura, "Evidence for pseudoknot formation of class I preQ1 riboswitch aptamers," Chembiochem **10**(7), 1141–1144 (2009).

[31] D.J. Klein, T.E. Edwards, and A.R. Ferré-D'Amaré, "Cocrystal structure of a class I preQ1 riboswitch reveals a pseudoknot recognizing an essential hypermodified nucleobase," Nat Struct Mol Biol **16**(3), 343–344 (2009).