ErbB2/HER2 governs CDK4 inhibitor sensitivity and timing and

irreversibility of G1/S transition by altering c-Myc and cyclin D function

Ayaka Nagasato-Ichikawa[1], Ken Murakami[1], Kazuhiro Aoki[2,3], and Mariko Okada[1]*

[1]Laboratory for Cell Systems, Institute for Protein Research, Osaka University, Suita, Osaka 565-0871, Japan

[2]Exploratory Research Center on Life and Living Systems (ExCELLS), National Institutes of Natural Sciences, Okazaki, Japan; National Institute for Basic Biology, National Institutes of Natural Sciences, Okazaki, Japan; Department of Basic Biology, School of Life Science, SOKENDAI (The Graduate University for Advanced Studies), Okazaki 444-8787, Japan

[3]Graduate School of Biostudies, Kyoto University, Kyoto 606-8501, Japan

*Corresponding author: mokada@protein.osaka-u.ac.jp

Competing interests: The authors declare that they have no competing interests.

Abstract

Cell cycle entry and the irreversible transition from the G1 to S phase are crucial for

mammalian cell proliferation. Among the ErbB family, the ErbB2/HER2 receptor is a key

driver of cancer growth. However, the quantitative mechanisms underlying the ErbB2-

mediated G1/S transition remain unclarified. Here, we performed an extensive time-course

analysis of high and low ErbB2-expressing breast cancer cells to describe the regulatory

mechanisms of the ErbB2-mediated G1/S transition. Live-cell imaging using cell cycle

reporters revealed that the G1/S transition occurs 20 h after ErbB2 activation, driven

primarily by the cyclin D1/CDK4–RB axis. Hsp90 is regulated by CDK4 activity and

controls the stability of ErbB2 protein in a time-dependent manner. CDK4 inhibitor treatment

arrested the cell cycle in most cells; a subpopulation showed a 25-h delay in G1/S entry

associated with enhanced c-Myc activation. In high ErbB2-expressing cells, CDK4 inhibition

led to c-Myc overactivation, a rapid decrease in cyclin D1 expression, and cell cycle arrest.

Overall, we demonstrate how ErbB2 receptor levels modulate the roles of cyclin D1 and c-

Myc in the G1/S transition and suggest that variations in ErbB2 levels within breast cancer

tissues confer heterogeneous sensitivity to CDK4 inhibitors, potentially complicating

treatment.

Main text:

Introduction

The initiation of the cell cycle and subsequent transition from the G1 to S phase are regulated

by a complex molecular network involving the formation of a cyclin D1/CDK4 protein

complex (*1–3*). Understanding the regulation of the G1/S transition by mitogenic signals is

fundamental for comprehending cell biology and developing cancer treatments that overcome

drug resistance. For example, based on the important role of cyclin D1 overexpression for

enhancing breast cancer cell proliferation (*4*), various cell cycle inhibitors targeting CDK4

have been developed to treat breast cancer (*5*, *6*).

Breast cancer can be classified into four subtypes based on the expression of estrogen

receptors (ERs), progesterone receptors, and ErbB2 (HER2) receptors: luminal A (ER$^+$ and

HER2$^-$), luminal B (ER$^+$ and HER2$^+$), ErbB2/HER2-positive (ER$^-$ and HER2$^+$), and triple-

negative (ER$^-$ and HER2$^-$) (*7–9*). CDK4 inhibitors (CDK4is) have been used to treat the

luminal A subtype (*5*, *10*), but long-term administration has led to drug resistance (*11–17*).

ErbB2 tyrosine kinase activity, commonly elevated in malignant breast cancer, increases

cyclin D1 levels (*18*, *19*). Stimulation of ErbB2 receptors activates the MAPK/ERK and

PI3K/AKT signaling pathways, leading to the rapid induction of early-response genes such as

c-fos and *c-myc* (*20–22*)—key regulators of cyclin D1 and CDK4 expression (*22–24*). Both

c-Myc and cyclin D1 are involved in the G1/S transition, though they appear to have distinct

roles (*25*). Therefore, we speculated that the small population of high ErbB2-expressing cells

in the luminal A subtype may contribute to CDK4i resistance by rewiring the G1/S network

during the early stages of drug treatment. However, the molecular mechanisms by which

ErbB2 affects the G1/S transition following CDK4i treatment remain unclear.

The G1/S transition typically involves signal-dependent and -independent processes, which

are distinguished by the restriction (R) point (*2*, *26*). The mitogenic signal primarily regulates

the expression and activation of cyclin D1/CDK4, which phosphorylates retinoblastoma

protein (RB) prior to the R point. Thus, this signal-dependent process is primarily driven by

the cyclin D1/CDK4-RB axis (*2*). The phosphorylation of RB (p-RB) activates E2F1,

promoting the expression of its target genes, including cyclin E. Cyclin E then forms a

complex with CDK2, which further phosphorylates p-RB, resulting in a fully phosphorylated

state (pp-RB). Fully activated pp-RB enhances E2F1 activity, creating a mitogen-independent

cyclin E/CDK2–RB–E2F1 positive feedback loop (*27*, *28*). However, the expression and

stability of c-Myc are modulated by external signals (*29*), and c-Myc directly influences

E2F1 promoter activity (*30*). Consequently, the core machinery of cyclin E/CDK2–RB–E2F1

is considered to be indirectly regulated by mitogenic signals.

ERK controls the expression of c-Myc and cyclin D1 (*31*, *32*), and AKT regulates their

protein stability (*31*, *33*). However, p27 expression, which plays a crucial role in the

accumulation of cyclin D1 and CDK4 during the G1 phase (*34*), is repressed by c-Myc (*35*).

Since the ErbB2 receptor activates the ERK and AKT pathways at different magnitudes (*36*),

different levels of ErbB2 may modulate the core machinery of the G1/S network and,

consequently, the timing of cell cycle entry through modulation of cyclin D1 and c-Myc.

Therefore, investigating the ratio of c-Myc to cyclin D1 at the single-cell level may elucidate

their specific functions in the G1/S transition and the regulatory role of the ErbB2 receptor.

In the present study, we used representative luminal A subtype cell lines (MCF-7 and T47D),

which express low levels of the ErbB1 receptor (EGFR) and high levels of the ErbB3

receptor (*22*, *36*, *37*). ErbB2 is an orphan receptor that is activated by heterodimerization with

ErbB3, ErbB4, or EGFR (*21*). Stimulation with the ErbB3 ligand, heregulin (HRG), triggers

the heterodimerization of ErbB2 and ErbB3, thereby activating the ERK and AKT pathways

(*36*). We conducted time-course protein phosphorylation profiling after HRG stimulation,

fluorescent ubiquitination-based cell cycle indicator (FUCCI) live-cell imaging for cell cycle

dynamics, and transcriptomic/epigenetic studies to identify key transcription factors during

CDK4 inhibition. We evaluated the role of ErbB2 in regulating the timing and irreversibility

of the G1/S transition, revealing the molecular mechanism underlying heterogeneous CDK4

inhibitor sensitivity.

Results

G1/S transition occurs approximately 20 h after HRG stimulation

First, we used FUCCI (*38*) live-cell imaging to analyze the cell cycle dynamics in MCF-7 or

T47D cells after HRG stimulation (Fig. 1A) and analyzed the distribution of each cell cycle

phase in the presence or absence of HRG for up to 72 h in MCF-7 cells (Fig. 1B–D). HRG

stimulation began to significantly increase the proportion of S/G2/M-phase cells after 24 h

compared to the control. The number of nuclei, serving as a proxy of cell number, increased

1.4-fold after 48 h, suggesting that HRG stimulation enhances cell proliferation. Conversely,

control cells showed a gradual increase in the proportion of S/G2/M cells, but there was no

increase in the number of nuclei, suggesting cell cycle arrest during the S/G2/M phase. To

further examine the times of transition for each cell cycle phase, we performed single-cell

tracking and quantified the distribution of the cell cycle phases (Fig. 1E; Fig. S1A, B). The

G1/S transition occurred at 19.0 ± 7.2 h after HRG stimulation. In comparison, the S/G2 and

M/G1 transitions occurred at 32.5 ± 9.3 h and 39.4 ± 8.7 h, respectively (Fig. 1F). These

findings suggested that most cells entered the S phase approximately 20 h after HRG

stimulation, with some intercellular variability.

CDK4 and cyclin D1 proteins are expressed in a time-dependent manner at the G1/S

transition

We further examined the signaling dynamics of the ErbB2, AKT, and ERK pathways (*21, 39*)

following HRG stimulation (Fig. 2A, B). p-ErbB2, p-AKT, and p-ERK levels decreased after

peaking within 30 min, and remained present until the S/G2 transition point (32.5 ± 9.3 h).

Although c-Fos protein levels peaked within 2 h and quickly reduced to basal levels, p-c-Fos

showed similar dynamics to those of p-ERK, a critical regulator of p-c-Fos (*22*). Following

ErbB2 activation, p-AKT, p-ERK, and p-c-Fos levels remained consistent whereas c-Myc and

p-c-Myc (Ser^{62}) levels decreased around the time of the G1/S transition (19.0 ± 7.2 h).

We estimated the timing of the R point based on the dynamics of cell cycle regulatory

proteins (Fig. 2C, D; Fig. S2A, B). The protein levels of cyclin D1 and CDK4 started to

increase shortly after HRG stimulation, peaked at approximately 20 h (G1/S transition point),

and then remained consistent until the S/G2 transition (Fig. 2D), consistent with the trend in

p-RB (Thr^{373}) expression (*40*). p-RB (Ser$^{807/811}$) showed a monotonic increase similar to

cyclin E (Fig. S2A, B), suggesting the involvement of CDK2-mediated S phase progression

(Fig. S2A, B). Since the R point is found just before the G1/S transition, we estimated that it

occurs at 12–16 h, when the E2F1 protein level is highly elevated. Thus, our results

substantiate that cyclin D1/CDK4 maintains RB phosphorylation near the R point, which is a

critical initial step toward the G1/S transition associated with HRG-induced ErbB2

activation.

A population subset enters the cell cycle even after CDK4 inhibition

We confirmed the involvement of cyclin D1/CDK4 in the HRG-stimulated G1/S transition.

Cells were treated with CDK4i at 12 h after HRG stimulation (before the R point; Fig. S3A,

B). As ERK activates cyclin D1 expression and AKT inhibits its degradation (*41*), we used

AKT inhibitor (AKTi) and MEK inhibitor (MEKi), which have comparable inhibitory effects

on cyclin D1 expression (Fig. S3A, B), as CDK4is. We examined the proportions of cells in

each cell cycle phase at 32 h and 72 h after HRG stimulation (Fig. 3A, B). At 32 h, fewer

cells treated with CDK4i (13.8%), AKTi (7.05%), or MEKi (13.8%) were in the S/G2/M

phase compared to non-treated cells (29.2%). At 72 h, AKTi (9.85%) and MEKi (5.50%)

reduced the proportion of cells in the S/G2/M phase, but CDK4i (17.7%) did not. At this time

point, the number of nuclei was significantly reduced by AKTi and MEKi but not CDK4i

(Fig. 3C). This finding indicated that AKTi and MEKi induced the growth of most cells,

whereas a significant subpopulation of cells were not affected by CDK4i, suggesting that

additional mechanisms maintain cell cycle functionality in the absence of CDK4.

To investigate the distinct mechanisms associated with CDK4i, we analyzed the correlations

between the times to the G1/S and M/G1 transitions up to 72 h for the different treatments

(Fig. 3D). Cells non-responsive to CDK4i showed significantly delayed G1/S transition times

(49.5 ± 13.2) of ~25 h compared to the control (24.7 ± 5.7), which tended to be longer than

under AKTi (39.6 ± 14.8) and MEKi (26.6 ± 11.5; Fig. 3D). We analyzed the slope (s),

representing the effect on the subsequent cell cycle duration after G1/S transition, and

intercept (i), representing the time to the M/G1 transition, and observed a lower slope (s =

0.56) and higher intercept (i = 37) under CDK4i than under AKTi (s = 1.0, i = 19), MEKi (s =

0.95, i = 17), and control conditions (s = 1.0, i = 17). Thus, Pearson's correlation coefficients

(R) were significantly lower for CDK4i (0.76) than for AKTi (0.91), MEKi (0.95), and

control conditions (0.84). This observation implied that the duration of the G1/S to G2/M

phase might not be uniformly controlled only by CDK4i and phase-specific regulation might

occur.

The CDK4i condition showed a clear negative correlation between the time to G1/S transition

and duration of the S/G2/M phase (Fig. 3E). We divided the cell population into the early and

late groups based on the median time of each G1/S transition (CDK4i, 49.5 ± 13.2; AKTi,

39.6 ± 14.8; MEKi, 26.6 ± 11.5; Control, 24.7 ± 5.7) and compared the duration of each S,

G2, and M phase (Fig. S3C). The mean duration of S/G2/M differed significantly between

the early (22.3 ± 7.2) and late (15.8 ± 2.4) groups, with a 6-h difference observed in the

CDK4i groups. Thus, in this CDK4i-specific response, the majority of CDK4i non-

responsive cells showed a long G1/S and short S/G2/M phase (Fig. 4A). We observed that

CDK4i non-responsive T47D cells (another luminal A cell line), which normally express

ErbB2 at a level three times higher than MCF-7 cells (Fig. S3D, E), also required a long time

(32.6 ± 12.2 h) to reach the G1/S transition compared to the control (21.62 ± 6.2 h; Fig. S3F).

CDK4i enhances c-Myc transcriptional activity via H3K27Ac modification

Given the distinct cell cycle kinetics observed under CDK4 inhibition (Fig. 4A), we

performed transcriptome and epigenetic analyses to investigate the molecular mechanisms

responsible for cell cycle regulation post CDK4 inhibition. To elucidate the changes in gene

expression patterns, we added CDK4i 12 h after HRG stimulation and performed a time-

course analysis of mRNA expression [0 (control), 2, 4, 8, 12, 16, 20, 24, and 28 h after HRG]

as well as H3K27Ac and H3K4me1 modification [12 (control), 16, 20, 24, and 28 h after

HRG]. First, active transcription factors were predicted at each time point with DoRothEA

using RNA-seq data (*42*) (Fig. S4A). mRNAs corresponding to key transcription factors,

such as *Myc*, AP-1 family (*Jun* and *Fos*), *EGR1*, *NFKB1*, and *STAT3*, which are known to

promote cell cycle progression (*21*, *23*, *25*, *43*, *44*), were highly upregulated 16 h after HRG

stimulation.

Motif-enrichment analysis was performed to confirm the binding of these transcription

factors to the promoter regions. Here, we defined the regions without overlap between the

H3K27Ac and H3K4me1 peak signals as the active promoter regions (*45*, *46*). E-box

sequences (CANNTG) (*47*) were enriched in the promoter regions at all time points

following CDK4i treatment (Fig. 4B; Fig. S4B, C). Given that c-Myc binds with E-box

sequences (*48*), we focused on its role in G1/S transition. Control cells showed lower c-Myc

protein and p-c-Myc (Ser^{62}) levels near the R point (12–16 h after HRG) than during the early

time point (0–2 h after HRG; Fig. 2A, B). Additionally, gene set enrichment analysis (GSEA)

using normalized enrichment scores (*49*) did not show any significant enrichment of c-Myc-

target genes during 12–16 h (Fig. 4C). However, CDK4i induced a significant enrichment of

c-Myc-target genes at 16 h after HRG stimulation, followed by a downregulation at 24–28 h

compared to that in the control (Fig. 4D). We analyzed the temporal expression of cell cycle-

related c-Myc-target genes.

Focusing on genes exhibiting a higher abundance of H3K27Ac modifications in their

promoter than their enhancer region (Fig. S4D). *CCNE1* expression was significantly

enhanced 16 h after HRG stimulation and CDK4i treatment, whereas the expression of

CCND1, *CDK1*, *CDC25A*, *E2F2*, *E2F3*, and *CDKN1B* showed delayed upregulation during

20–28 h. These findings suggested that CDK4 inhibition alters the H3K27Ac levels in the

promoter regions of c-Myc-target genes near the R point, promoting the expression of cell

cycle-related genes. Therefore, when the G1/S transition, originally regulated by the cyclin

D1 and CDK4 complex, is inhibited by CDK4i, it is maintained by an alternative pathway

facilitated by c-Myc activity.

We confirmed whether the delayed G1/S entry following CDK4i treatment was due to c-Myc

activation by testing the effects of an inhibitor of c-Myc transcriptional activity (EN4). In

both MCF-7 and T47D cells after 30 h, the combination of CDK4i and EN4 increased the

proportion of G1-phase cells and decreased the number of S/G2/M-phase cells in a dose-

dependent manner (Fig. 4E–H), while no change was observed with EN4 alone (Fig. S4E–H).

These results suggested that the transcriptional activity of c-Myc promotes the delayed G1/S

entry and escape from cell cycle arrest in response to CDK4 inhibition.

To further validate these findings using clinical data, we retrospectively analyzed the

microarray data from patients with luminal A subtype breast cancer who received long-term

(16 weeks, $n = 23$) or short-term (4 weeks, $n = 26$) CDK4/6i therapy (*50*). *PCNA*, a target

gene for both c-Myc and E2F involved in DNA replication during G1/S transition, showed a

0.5 $\log_2$ fold change (FC) between the long-term and short-term treatment groups (Fig. 4I).

PCNA mRNA expression increased after CDK4i treatment from 4 to 12 h (16–24 h after

HRG) relative to the control (Fig. 4J). Collectively, these findings suggested that CDK4

inhibition-induced epigenetic modifications activated c-Myc-target genes, ultimately

promoting cell cycle maintenance in cancer cells (Fig. 4K).

CDK4i induces ErbB2 degradation and delays G1/S transition

c-Myc is known to promote G1/S transition and cell proliferation (*25*, *35*). Therefore, if the

transcriptional activity of c-Myc is enhanced by CDK4i at 16 h, the time to G1/S transition is

unlikely to be extended; however, the G1/S transition was prolonged in our study (Fig. 3D, E;

Fig. 4A). We found that the levels of phosphorylated ErbB2, ERK, and AKT, which induce *c-*

Myc mRNA expression and stabilize the c-Myc protein, decreased after CDK4 inhibition,

leading to a slight reduction in c-Myc protein level (Fig. 5A, B). Although the expression of

ErbB2 was lower in MCF-7 and T47D cells than in HER2-positive cells, growth in these cell

lines is mainly dependent on ErbB2-ErbB3 heterodimer activation after HRG stimulation

given their low expression of EGFR and ErbB4 (*37*). This finding suggested that the CDK4i-

induced downregulation of the ErbB2 pathway may prolong the G1/S transition in MCF-7

and T47D cells (Fig. 3D; Fig. S3F). To investigate this mechanism, we focused on the

Hsp90-chaperone system, particularly with respect to the stability and maturation of nascent

ErbB2. As WT MCF-7 cells naturally express low levels of ErbB2, we constructed MCF-7

cells with moderate overexpression of ErbB2 (Fig. S5A, B). After confirming the unaltered

protein expression of ErbB2, PY100 (pan-tyrosine phosphorylation for estimating p-ErbB2

level), and Hsp90 under CDK4i (Fig. S5C–G), we examined the co-localization patterns

using Mander's and Pearson's coefficients. Compared to control conditions, CDK4i reduced

the co-localization of Hsp90 with ErbB2 or PY100 (Fig. 5C–F). No significant difference

was observed in the total number of nuclei (Fig. S5H, I). These results suggested that CDK4i

inhibits the interaction between Hsp90 and ErbB2, leading to ErbB2 misfolding. In addition,

ErbB2 protein expression was reduced 20 h after HRG stimulation (Fig. 5G, H; Fig. S5J);

however, *ErbB2* mRNA levels did not change regardless of CDK4i treatment (Fig. S5K).

Therefore, we investigated whether the downregulation of ErbB2 was caused by its protein

degradation using the proteasome inhibitor MG132. Upon MG132 treatment, ErbB2 protein

expression remained unchanged (Fig. 5G, H). This observation indicated that CDK4i disrupts

the interaction between Hsp90 and ErbB2, triggering ErbB2 degradation (Fig. S5L). To

confirm this, we treated cells with the Hsp90 inhibitor geldanamycin at a concentration that

suppressed ErbB2 expression by 40%, similar to the effect of CDK4i (Fig. S5M, N). As a

result, cell cycle progression grounded to a halt, and there was no substantial delay in the

G1/S transition (Movie S1). Therefore, the loss of ErbB2 protein stability following CDK4i

treatment may impair the regulation of c-Myc-mediated G1/S transition, even if c-Myc is

activated by epigenetic modification, thereby delaying the G1/S transition (Fig. 5I).

High ErbB2 expression induces cell cycle arrest by repressing cyclin D1 activity via c-

Myc under CDK4 inhibition

We speculated that ErbB2 protein instability contributes to the delay in G1/S entry under

CDK4i and thus could be counteracted by ErbB2 overexpression. To test this hypothesis, we

generated MCF-7 cells expressing varying levels of ErbB2, from moderate to high (Fig. S5A,

B), and analyzed their cell cycle phase distribution using live-cell imaging (Fig. 5J).

After CDK4i treatment, the proportion of S/G2/M-phase cells with high ErbB2 expression

was significantly reduced (OE-Moderate: 14.9%, OE-High: 12.5%; Fig. 5K), indicating cell

cycle arrest. We examined the protein expression of cyclin D1 and p-RB, which are critical

for initiating cell cycle entry (Fig. 5A, B). We observed a significant reduction in both cyclin

D1 and p-RB levels. Therefore, we investigated the cause of the decreased cyclin D1 levels.

The p27 protein reportedly promotes the assembly of the cyclin D1/CDK4 complex in the G1

phase (*34*), while its gene expression is inhibited by p-AKT and c-Myc (*35*). Our data

revealed that p27 protein expression decreased after CDK4i treatment, whereas p-AKT and c-

Myc protein levels tended to increase post treatment. By contrast, p-ERK levels remained

unchanged under different conditions. These findings suggested that, in ErbB2-

overexpressing cells, the enhanced p-AKT–c-Myc axis antagonizes the protein expression of

p27 and cyclin D1, leading to cell cycle arrest in the absence of CDK4 activity.

Reversible cell cycle arrest is maintained by c-Myc transcriptional activity

Drug resistance is closely linked to a reversible cell cycle arrest state regulated by residual

survival signals (*11*). In ErbB2-overexpressing cells, a cell cycle arrest state is induced after

CDK4 inhibition, but this may be reversed by the survival signals from c-Myc. In CDK4i-

treated cells with high ErbB2 expression, CDK4i removal increased cyclin D1, p27, and p-

RB levels, while c-Myc expression remained unchanged (Fig. 6A, B). We analyzed the c-

Myc phosphorylation (Ser^{62}) levels to confirm its elevated transcriptional activity in cell

cycle arrested cells. Single-cell analysis revealed that CDK4i depletion decreased the ratio of

phosphorylated c-Myc (Ser^{62}) to cyclin D1 (p-c-Myc/cyclin D1) and increased the ratio of S-

phase cells from 18.7% to 47.5% (Fig. 6C). ErbB2-overexpressing cells maintained a higher

p-c-Myc/cyclin D1 ratio than wild-type cells (Fig. S6A), suggesting that sustained cell cycle

arrest may result from persistent c-Myc transcriptional activation.

To investigate whether blocking c-Myc transcriptional activity could prevent reversible cell

cycle arrest, we treated cells with EN4 and CDK4i for 8 h, washed the inhibitors, and

analyzed cell cycle distribution at 52 h (72 h after HRG stimulation; Fig. 6D, E). The

combined treatment increased G1-phase and reduced S/G2/M-phase cell numbers, indicating

impaired cell cycle re-entry. CDK4i reduced p27 (*CDKN1B*) mRNA levels, which were

restored by co-treatment with EN4 (Fig. 6F, G), suggesting direct c-Myc regulation of p27

gene expression. Overall, these data supported the hypothesis that, in cells overexpressing

ErbB2, CDK4 inhibition promotes reversible cell cycle arrest due to an excessive increase in

c-Myc transcriptional activity and rapid decrease in cyclin D1 protein expression, which is

attributed to the reduction in p27 gene and protein expression.

Given the association between stable cell cycle arrest and cellular (*51*, *52*) or ErbB2-induced

senescence (*53*), we examined the senescence phenotype. Senescence associated β-

galactosidase (SA-β-Gal) staining intensity after 6 d of CDK4i treatment showed an inverse

correlation with ErbB2 expression. EN4 (4 μM) enhanced SA-β-Gal staining in WT MCF-7

cells beyond the effect of CDK4i alone but showed no additional effect in T47D or high

ErbB2-expressing MCF-7 cells (Fig. 6H, I). This finding was consistent with cell line-

specific sensitivity, since 2 nM EN4 induced cell cycle arrest in WT MCF-7 cells, whereas 2

μM was required for WT T47D cells (Fig. 4E, F). Our results suggested that ErbB2 levels

modulate the irreversibility of the G1/S transition under CDK4i treatment, and the addition of

a c-Myc inhibitor supports the irreversible state (Fig. 6J).

Discussion

Our findings reveal that ErbB2 receptor levels modulate the relative roles of cyclin D1 and c-

Myc in the G1/S transition, ultimately providing two distinct pathways for cell cycle

progression or arrest in the presence of CDK4i.

The luminal A subtype accounts for 30% of breast cancer cases. Although these patients

typically receive standard hormone therapy, CDK4/6is are also commonly used in cases of

more advanced or metastatic tumors (5, 54). However, their success is limited by

heterogeneous drug responses. Notably, an ErbB2-overexpressing subpopulation has been

identified within luminal A breast cancer cells, which were initially characterized by low

ErbB2 expression (55). Given that these subpopulations occur relatively infrequently, at

approximately 7% of the total population (56), identifying differences in cell cycle regulation,

especially in terms of ErbB2 expression, is critical for improving cancer treatment. Long-

term CDK4i treatment, commonly used for luminal A breast cancer, activates alternative

pathways, such as PI3K/AKT/mTOR, androgen receptor, and Hippo signaling (13, 14, 16,

17), along with transcriptional reprogramming via non-coding RNAs (15, 57), promoting the

development of drug resistance. To mitigate this resistance, pathways that may act at the very

early step of cell cycle entry must be inhibited. Accordingly, we investigated how ErbB2

levels within cells of the same luminal A subtype affect the core cell cycle regulatory network

and CDK4 inhibitor sensitivity (Fig. 7A, B). In MCF-7 cells under HRG stimulation, CDK4

RNA levels increased and remained elevated through R-point and S-phase entry, whereas

CDK6 RNA levels were quite low compared to those of CDK4 and tended to decrease after

HRG stimulation (Fig. S4D). Therefore, we focused on the effect of CDK4 inhibitors rather

than CDK4/6 inhibitors.

The temporal dynamics of the cell cycle regulators and live-cell imaging analysis showed that

the roles of cyclin D1 and c-Myc, key regulators of the G1/S transition, vary depending on

the expression status of ErbB2 under CDK4i treatment. This variation ultimately affects the

timing and irreversibility of the G1/S transition. In MCF-7 cells, a representative luminal A

breast cancer cell line with low ErbB2 expression, the G1/S transition is normally regulated

by the cyclin D1/CDK4 complex via Hsp90 (Fig. 5C–E; Fig. S5C–O). Although CDK

inhibitors are known to sequester Hsp90 and CDK4/6 (58), our results suggested that the

function of Hsp90 is complemented by CDK4 activity and contributes to stabilizing the

ErbB2 protein. This positive feedback mechanism provides valuable insights into the signal-

dependent complexity of cell cycle regulation. In the non-responsive subpopulations to

CDK4 inhibition, the G1/S transition persists, but with delays due to epigenetic modification

and increased c-Myc transcriptional activity and ErbB2 degradation (Fig. 4; Fig. 7A).

However, the excessive transcriptional activity of c-Myc promotes cell cycle arrest (Fig. 6A–

G) by rapidly reducing cyclin D1 and p27 levels. The expression of p27, which recruits

cyclin D1 to the nucleus during the G1 phase, is also repressed by AKT and c-Myc (35).

Therefore, in high ErbB2-expressing MCF-7 cells, with high levels of AKT and c-Myc

activity, CDK4 inhibition increases free cyclin D1 levels, rendering it more susceptible to

degradation (59–61). This process generates a hypophosphorylated RB state in ErbB2-

overexpressing MCF-7 cells compared to WT cells, leading to cell cycle arrest due to a

decrease in free E2F now bound to c-Myc (Fig. 7B).

c-Myc regulates the cell survival pathway (35, 62). In ErbB2-overexpressing MCF-7 cells,

cell cycle arrest was reversible after CDK4 inhibition (Fig. 6D, E), suggesting that the

transcriptional activity of c-Myc promotes sustained cell survival in this context. Co-

treatment with inhibitors targeting the transcriptional activity of c-Myc and CDK4 may be

necessary to induce irreversible cell cycle arrest (Fig. 6H, I). However, at the same

concentration at which the Myc inhibitor promoted cellular senescence in WT MCF-7 cells,

the percentage of senescent WT T47D and ErbB2-overexpressing cells did not change

compared to that in the respective control conditions. For drug response assessments, the

threshold at which c-Myc switches from supporting cell survival to inducing irreversible cell

cycle arrest in relation to ErbB2 levels could inform the development of more effective

molecular-targeted treatment strategies for breast cancer.

Overall, our results suggest that the c-Myc-to-cyclin D1 ratio provides valuable quantitative

insights into the varying sensitivity of cancer cells to CDK4is, without relying solely on

observable cell proliferation rates or single biomarker gene analysis.

Materials and methods

Reagents

The MEKi PD0325901 was purchased from FUJIFILM Wako Pure Chemical Corp. (Osaka,

Japan); CDK4i V was purchased from Calbiochem (San Diego, CA, USA); AKTi VIII was

purchased from Cayman Chemical (Ann Arbor, MI, USA); EN4 (Myci) was purchased from

Selleck Chemicals (Houston, TX, USA); geldanamycin was purchased from FUJIFILM

Wako Pure Chemical Corp.; blasticidin was purchased from Kaken Pharmaceutical (Tokyo,

Japan); Lipofectamine LTX was purchased from Thermo Fisher Scientific (Waltham, MA,

USA); and HRG was procured from R&D Systems (Minneapolis, MN, USA).

ErbB2 overexpression

MCF-7 (ATCC, HTB-22) cells overexpressing full-length human ErbB2 (HER2) were

established as follows: Human ErbB2 cDNA in the pCMV6-XL5 vector was purchased from

OriGene (Catalog No: TC128161, Rockville, MD, USA). The cDNA fragment was cloned

into the vector pCMV-6-Neo vector (OriGene). The plasmid was then transfected into MCF-7

cells using Lipofectamine LTX (Invitrogen, Carlsbad, CA, USA) according to the

manufacturer's protocol, and stable transfectants were selected with G418 (neomycin).

Colonies derived from single cells compartmentalized in 96-well plates were isolated and

examined. ErbB2 expression was examined via western blotting to group cells with moderate

or high ErbB2 expression (as described below for protein extraction and immunoblotting).

Cell culture

MCF-7, ErbB2-overexpressing MCF-7, and T47D (ATCC, HTB-133) cells were cultured in

Dulbecco's modified Eagle's medium (DMEM, Nacalai Tesque, Kyoto, Japan) supplemented

with 10% (*v/v*) fetal bovine serum (FBS, Sigma-Aldrich, Saint Louis, MO, USA) and 1%

penicillin–streptomycin (100 IU/ml penicillin, 100 μg/ml streptomycin; Nacalai Tesque) at

37 °C in a humidified 5%-CO_2 atmosphere. Cells were routinely tested using a CycleavePCR

Mycoplasma Detection Kit (Takara bio, CY-232); no mycoplasma contamination was

detected.

FUCCI transfection

MCF-7 and ErbB2-overexpressing MCF-7 cells stably expressing the FUCCI biosensor were

generated using a PiggyBac transposase system (*63*). Lipofectamine LTX (Thermo Fisher

Scientific) was used to transfect the FUCCI plasmid (pPBbsr2-H2B-iRFP-P2A-mScarlet-I-

hGem-P2A-PIP-tag-NLS-mNeonGreen; https://benchling.com/s/seq-

LPZ1tLdpgnpJIYOG2ujR) along with hyPBase (https://benchling.com/s/seq-

oGkw53b41IZqvzF5yQ9K) into MCF-7 and ErbB2-overexpressing MCF-7 cells, following

the manufacturer's protocol (*64*). Cells were suspended in a drug-free medium for 48 h and

treated with blasticidin (10 μg/ml) to separate those stably expressing the plasmid. Colonies

derived from single cells compartmentalized in 96-well plates were selected by microscopic

live-cell observation of Geminin and PIP-tag fluorescence.

Protein extraction and immunoblotting

MCF-7, T47D, and ErbB2-overexpressing (moderate or high) MCF-7 cells were seeded in

six-well plates (450 000 cells/well) and cultured in 10% FBS containing DMEM for 48 h. The

medium was changed to FBS-free phenol red-free DMEM (FUJIFILM Wako Pure Chemical

Corp.) containing 1% penicillin–streptomycin (100 IU/ml penicillin, 100 μg/ml streptomycin;

Nacalai Tesque) and 1% 100 mmol/L sodium pyruvate solution (FUJIFILM Wako Pure

Chemical Corp.), and the cells were cultured for 17 h. The medium was changed again, and

the cells were cultured for 3 h. After administration of 10 nM HRG (R&D Systems) for 12 h,

the cells were independently treated with CDK4i and dimethyl sulfoxide (DMSO). Serum

was not included under conditions where HRG was added. Total cell lysates were prepared in

RIPA buffer (Nacalai Tesque, 1648834) containing protease inhibitors (11836153001, Roche

Rotkreuz, Switzerland) and phosphatase inhibitors (Roche, 4906837001) at 4 °C. Cells were

vortexed for 5 min and centrifuged at $16 000 \times g$ and 4 °C for 15 min. The supernatant was

collected and denatured by heating at 95 °C with sample buffer (5×) and 2-mercaptoethanol

(Nacalai Tesque, 21418-55) for 5 min. The proteins were subjected to SDS polyacrylamide

gel electrophoresis (Nacalai Tesque, 1306644), transferred to polyvinylidene fluoride

membranes (Merck, Darmstadt, Germany), and incubated with 10% blocking buffer (Nacalai

Tesque, 03953-95) prepared in Tris-buffered saline/Tween 20 (TBS-T; T9142; Takara Bio,

Shiga, Japan) for 1 h at 25 °C. Membranes were probed overnight with antibodies against

AKT (1:1 000; #4685; Cell Signaling Technology, Danvers, MA, USA), p-AKT (Thr^{308}; 1:1

000; Cell Signaling Technology; #2965), ERK (1:1 000; Cell Signaling Technology; #9102),

p-ERK (Thr^{202} Tyr^{204}; 1:1 000; Cell Signaling Technology; #4370), HER2/ErbB2 (1:1 000;

Cell Signaling Technology; #2165), p-HER2/ErbB2 (Thr$^{1221/1222}$; 1:1 000; Cell Signaling

Technology; #2249), p-c-Fos (Ser[374]; 1:1000; #ab55836; Abcam, Cambridge, UK), c-Fos

(1:200; #sc-166940; Santa Cruz Biotechnology, Dallas, TX, USA), p-c-Myc (Ser[62]; 1:1000;

Abcam; #ab51156), c-Myc (1:200; Santa Cruz; #sc-40), E2F-1 (1:1000; Cell Signaling

Technology; #3742), p-RB (Ser[807/811]; 1:1000; Cell Signaling Technology; #8516), p-RB

(Thr[373]; 1:1000; Abcam; #ab52975), RB1 (1:1000; #SPM353; NOVUS biologicals,

Centennial, CO, USA), cyclin D1 (1:1000; Cell Signaling Technology; #2978), cyclin E

(1:200; Santa Cruz; #sc-247), cyclin A (1:200; Santa Cruz; #sc-271682), CDK4 (1:1000;

Santa Cruz; #DSC-35), CDK2 (1:1000; Santa Cruz; #D-12), P27 Kip1 (1:1000; Cell

Signaling Technology; #3686), and GAPDH (1:4000; 10494-1-AP; Proteintech, Boston, MA,

USA) in TBS-T including 10% blocking buffer at 4 °C. They were then incubated with

horseradish peroxidase (HRP)-conjugated secondary antibodies, HRP-conjugated anti-mouse

IgG (1:5000; Cell Signaling Technology; #7076) and HRP-conjugated anti-rabbit IgG

(1:5000; Cell Signaling Technology; #7074), for 1 h at room temperature. The membranes

were developed using a chemiluminescent HRP substrate (Merck). Protein bands were

detected using an Amersham Imager 600 UV (GE Healthcare, Eindhoven, Netherlands) and

visualized using the ImageJ-Fiji software (v.2.14.0/1.54f, NIH, Bethesda, MD, USA).

qRT-PCR

qRT-PCR was conducted as previously described (*65*). RNA was isolated from cells using

NucleoSpin RNA Plus (Cat. No. 740984.250; Macherey-Nagel, Duren, Germany) according

to the manufacturer's protocol. cDNA was synthesized using ReverTra Ace qPCR RT Master

Mix (TOYOBO, Osaka, Japan), and quantitative PCR was performed using a KOD SYBR

qPCR kit (TOYOBO) on a CFX96 Real-Time PCR System (Bio-Rad Laboratories, Hercules,

CA, USA). The ΔΔCq method was used to quantify gene expression, using *RPL27*

expression as an internal reference (*66*). The primers used for real-time PCR are listed in

Table S1.

Cell cycle analysis using the FUCCI biosensor

Cell cycle analysis was conducted as previously described (*65*). Time-lapse microscopy was

performed using an IN Cell Analyzer 2500HS (GE Healthcare, Chicago, IL, USA) equipped

with a CFI S Plan Fluor ELWD dry objective lens 20× (NA: 0.45) at an excitation

wavelength of 475 nm and emission wavelength of 511 nm for PIP-tag-mNeonGreen, and an

excitation wavelength of 542 nm and emission wavelength of 597 nm for mScarlet-hGem.

Cells were maintained at 37 °C in a humidified 5%-CO_2 atmosphere and imaged at 20-min

intervals for 72 h. Image processing and quantification of cell cycle phase distribution were

performed using ImageJ-Fiji software (v.2.14.0/1.54f, NIH). Masks of the mScarlet-hGem

and PIP-tag-mNeonGreen signals were obtained by applying a fixed threshold using

"Analyze Particles" and automatically counted as ROIs. Cell cycle phases were determined in

the following manner: G1 phase, PIP-tag-positive; S phase, hGem-positive; G2/M phase,

PIP-tag- and hGem-positive. The average brightness of mScarlet and mNeonGreen was

obtained using cell-tracking software (Olympus, Tokyo, Japan). The threshold parameters for

mScarlet and mNeonGreen were set manually for each cell line. We employed an algorithm

for estimating the cell cycle phase at the single-cell level using R packages [brunnermunzel

(v.1.4.1), multimode (v.1.4), rlist (v.0.4.6.1), and tidyverse (v.1.3.0)]. The duration of each

cell cycle phase was obtained from the single-cell data.

Nucleus count

The number of nuclei was quantified based on the fluorescence of the FUCCI probes, and

signals for a nuclear marker (histone H2B), mScarlet-hGem, and PIP-tag-mNeonGreen were

used to determine cell proliferation. Image processing and nuclear ratio quantification were

performed using the ImageJ-Fiji software (v.2.14.0/1.54f, NIH). Masks of the merged nuclear

marker, mScarlet-hGem, and PIP-tag-mNeonGreen signals were obtained by applying a fixed

threshold using "Analyze Particles" and automatically counted as ROIs. The nuclear ratio

was calculated by dividing the number of nuclei at the end of observation by that at the

beginning. Thus, a nuclear ratio near 1.0 indicated potential cell cycle arrest, above 1.0

indicated cell proliferation, and below 1.0 indicated apoptosis.

Immunostaining and quantitative analysis of images

MCF-7 and ErbB2-overexpressing (moderate or high) cells were seeded in 12-well plates

(200 000 cells/well) and cultured in 10% FBS containing DMEM for 48 h. The medium was

changed to FBS-free phenol red-free DMEM (FUJIFILM Wako Pure Chemical Corp.)

containing 1% penicillin–streptomycin (100 IU/ml penicillin, 100 μg/ml streptomycin;

Nacalai Tesque) and 1% 100 mmol/L sodium pyruvate solution (FUJIFILM Wako Pure

Chemical Corp.), and the cells were cultured for 17 h. The medium was changed again, and

the cells were cultured for 3 h. After administration of 10 nM HRG (R&D Systems) for 12 h,

the cells were independently treated with CDK4i and DMSO. Serum was not included under

conditions where HRG was added. Cells were washed twice with PBS at the indicated times,

fixed for 2 min with pre-prepared methanol at –20 °C, washed twice with Blocking One

(Nacalai Tesque, 03953-95), and incubated overnight at 4 °C with Blocking One. The next

day, cells were incubated overnight at 4 °C with primary antibodies diluted in 10% goat

serum (Thermo Fisher Scientific, 16210064)/Blocking One, washed five times with PBS, and

incubated with fluorescent dye-conjugated secondary antibody, 2 μg/ml DAPI (Nacalai

Tesque, 11034-56), and 0.5 μg/ml CellMask (Thermo Fisher Scientific, H32721) for 1 h at

room temperature. Cells were washed five times with PBS, and images were acquired using

the IN Cell Analyzer 2500HS (GE Healthcare Life Science) at 20× magnification. At least

three images were collected for each treatment, containing at least 100 cells each, for

quantification using the ImageJ-Fiji software (v.2.14.0/1.54f, NIH) and CellProfiler (v.4.2.1).

Co-localization analysis for Hsp90 with either ErbB2 or PY100 was performed using the

"Coloc 2" plugin in the ImageJ-Fiji software. Cell fractionation was performed on CellMask-

stained images using CellProfiler, and the mean intensity was calculated for p-c-Myc, cyclin

D1, and DAPI in each cell. The ratio of phosphorylated c-Myc (Ser[62]) to cyclin D1 (p-c-

Myc/cyclin D1) for each cell was plotted over the DAPI intensity. The DAPI intensity

indicating the G1/S transition was determined based on the percentage of cells that were in

the S/G2 /M phase 24 h after HRG stimulation. In WT MCF-7 cells, 22.4% of cells were in

the S/G2/M phase, at which time the DAPI intensity was 0.105. In MCF-7 cells

overexpressing ErbB2 (high), 25.4% of the cells were in the S/G2/M phase, at which time the

DAPI intensity was 0.0775.

Analysis of senescence associated β-galactosidase-positive cells

SA-b-gal staining was conducted as previously described (*65*). Briefly, senescent cells were

stained using a Senescence β-Galactosidase Staining Kit (#9860; Cell Signaling Technology)

according to the manufacturer's instructions. Images were acquired with a BZ-9000

microscope (Keyence, Osaka, Japan) at 20× magnification. The percentage of SA-β-gal-

positive cells was calculated from images using CellProfiler (v.4.2.1). The total number of

cells was determined by visualizing nuclear segmentation with DAPI staining. The number of

SA-β-gal-stained cells was identified using the UnmixColors module. The percentage of SA-

β-gal-positive cells was calculated by dividing the number of SA-β-gal-stained cells by the

total number of nuclei in each field of view.

RNA-seq

FUCCI-expressing MCF-7 cells were seeded in six-well plates (450 000 cells/well) and

cultured in 10% FBS containing DMEM for 48 h. The medium was changed to FBS-free

phenol red-free DMEM, the cells were cultured for 17 h, the medium was changed again, and

the cells were cultured for 3 h. After administration of 10 nM HRG for 12 h, the cells were

independently treated with CDK4i and DMSO. Serum was not included under conditions

where HRG was added. Cells were washed twice with PBS at the indicated times. Total RNA

was isolated using NucleoSpin RNA Plus (Macherey-Nagel; Cat. No. 740984.250) at the

specified time. Library construction and RNA-seq were performed on an Illumina sequencing

platform (Rhelixa Tokyo, Japan). Poly(A) RNA was prepared with a Poly(A) mRNA

Magnetic Isolation Module (E7490; New England Biolabs, Ipswich, MA, USA), and the

sequencing library was generated using a NEBNext Ultra II Directional RNA Library Prep

Kit (New England Biolabs, E7760). RNA-seq was performed on a NovaSeq 6000 (Illumina,

San Diego, CA, USA) with 150-bp paired-end reads. Raw reads were quality-checked using

FastQC (v.0.11.9) and trimmed using TrimGalore (v.0.6.0) (*67*). Trimmed reads were mapped

using STAR (v.2.7.9a) (*68*) with GRCh38 as the reference genome. featureCounts (v.1.6.4)

(*69*) was used to construct the count matrix with the "-p -B -t exon -g gene_name"

parameters. Raw counts were converted to transcripts per kilobase million and corrected

using EdgeR (v.3.36.0) (*70*). Principal component analysis was performed using the prcomp

function in R (v.4.1.1) (*71*). GSEA was performed using the fgsea package (v.1.20.0) (*72*) and

the hallmark gene set in the Human MSigDB Collections (*49*).

ChIP-seq

FUCCI-expressing MCF-7 cells were seeded in 100-mm dishes (2 300 000 cells/dish) and

cultured in 10% FBS containing DMEM for 48 h. The medium was changed to FBS-free

phenol red-free DMEM, the cells were cultured for 17 h, the medium was changed, and the

cells were cultured for 3 h. After administration of 10 nM HRG for 12 h, the cells were

independently treated with CDK4i and DMSO. Cells were washed twice with PBS at the

indicated times. Cells were fixed with 1% formaldehyde (Thermo Fisher Scientific) for 5

min. Fragmented DNA was obtained using a SimpleChIP Plus Enzymatic Chromatin IP Kit

(Cell Signaling Technology, 9005) and M220 Focused-ultrasonicator (Covaris, Woburn, MA,

USA). Immunoprecipitated DNA was obtained using an Auto iDeal ChIP-seq kit for Histones

×24 (Diagenode, Liege, Belgium) on an SX-8G platform (Diagenode). Next, 2 µg anti-

H3K27Ac antibody (Diagenode) and 2 µg anti-H3K4me1 antibody (Diagenode) were mixed

with 10 µl DiaMag Protein A-coated magnetic beads, and IP reactions with DNA were

performed for 12 h. De-crosslinking was conducted using 4 µl of 5 M NaCl and 1 µl of 20

mg/ml proteinase K (Thermo Fisher Scientific) for 4 h at 60 °C. Library construction and

RNA-seq were performed using the Illumina sequencing platform (Rhelixa). A sequencing

library was generated using a NEBNext Ultra II DNA Library Prep Kit for Illumina (New

England Biolabs, 7645). ChIP-seq was performed on the NovaSeq 6000 (Illumina) with 150-

bp paired-end reads. H3K27Ac ChIP-seq and H3K4me1 data were processed using the same

protocol as for RNA-seq. FastQC (v.0.11.9) was used for quality control. Trimming was

performed using Trimmomatic (v.0.39) (*73*) with the following parameters:

ILLUMINACLIP, TruSeq2-SE; fa, 2:30:10; LEADING, 30; TRAILING, 30;

SLIDINGWINDOW, 4:15; MINLEN, 30; HEADCROP, 2. Reads were mapped using

Bowtie2 (v.2.4.4) (*74*) with GRCh38 as the reference genome, and duplicate reads were

excluded using Picard (v.2.26.6) (*75*). Bigwig files were created from the generated BAM

files using deepTools (v.3.5.1; bamCoverage function) (*76*). Peak regions were identified

using HOMER (v.4.11) (*77*) with the "-style histone -L 0 -fdr 0.00005" parameters. All

detected peak regions were annotated to their nearest neighbor genes using HOMER

annotatePeaks.pl (v.4.11).

Promoter and enhancer detection

Promoter regions were defined as those in the H3K27Ac peak that did not overlap with the

H3K4me1 peak and were within 250-bp of the transcription start site (TSS, at the center of

the peak). Enhancer regions were defined as those in the H3K27Ac peak overlapping with the

H3K4me1 peak and were more than 250-bp from the TSS. The TSS was obtained from the

Ensembl database using BiomaRt (v.2.50.3) (*78*). The bed files containing the promoter and

enhancer regions for all time points were merged, and the time-course of the coverage values

for H3K27Ac ChIP-seq was calculated using HOMER annotatePeaks.pl (v.4.11). The

estimated coverage values were used to detect promoters and enhancers affected by CDK4i.

Promoters that gained or lost signal were defined as those with coverage values of $|\log_2 FC| >$

0.3 for inhibitor and control conditions. Enhancers that gained or lost signals were defined as

those with coverage values changing by $|\log_2 FC| > 0.3$ and target gene expression changing

by $|\log_2 FC| > 0.2$ in the same direction as the change in enhancer activity.

Motif analysis of ChIP-seq data

Motif analysis was performed using MEME Suite SEA (v.5.4.1) (*79*). The motif files were

obtained from the JASPAR 2022 database

(JASPAR2022_CORE_vertebrates_nonredundant_v2) (*80*). Peak regions containing motifs

were detected using MEME Suite FIMO (v.5.4.1). The analysis was performed using default

parameters.

Transcription factor enrichment analysis

Transcription factor analysis was conducted using DoRothEA (v.1.6.0) (*42*) with the

"run_viper" function. We used regulon data with "confidence" levels of A and B. Target genes

were selected based on their changes in expression (FDR < 0.01) between the control and

inhibitor treatments at each time point.

Clinical data analysis

We used microarray gene expression data (GEO GSE93204) for biopsy samples from the

NeoPalAna clinical trial (NCT01723774) (*50*) to identify differentially expressed genes based

on *P*-values adjusted using the Benjamini–Hochberg method to control the false discovery

rate in GEO2R.

Statistical analysis

Statistical analysis, data visualization, and Kolmogorov–Smirnov normality testing were

performed in the R software (v.4.2.1; The R Foundation, Vienna, Austria). For normally

593 distributed data, the Student's *t*-test and Welch's two-sample *t*-test were used after confirming

594 equal variances using the F-test for comparisons between two groups. Comparisons between

595 more than two groups were performed using one-way analysis of variance followed by the

596 Tukey's post-hoc test. For non-normally distributed data, a Kruskal–Wallis test or Mann–

597 Whitney U test was used. $P < 0.05$ was considered significant.

599 **Acknowledgments:**

600 We thank all members of the Laboratory for Cell Systems at the Institute for Protein

601 Research for the discussions regarding our research and especially Mr. Kyoichi Ebata for

602 helping to design the cell cycle analysis. We also thank Mr. Hideya Aragaki from Olympus

603 Corporation for providing the cell-tracking software.

605 **Funding:**

606  JSPS KAKENHI (18H04031, MO; 21K15503, ANI)

607  JST CREST Program (JPMJCR21N3, MO)

608  Uehara Memorial Foundation (MO)

609  Kaneko Narita Research Grant from the Protein Research Foundation (ANI)

610  New Field Development Support Program from the Institute for Protein Research (ANI)

611  Joint Research Grant from ExCELLS (MO, KA)

613 **Contributions:**

614  Conceptualization: ANI, MO

615  Methodology: ANI, KM, KA, MO

616  Validation: ANI

617  Formal analysis: ANI, KM

Investigation: ANI, KM

Resources: ANI, KA, MO

Visualization: ANI, KM

Writing – Original Draft: ANI, KM, MO

Writing – Review & Editing: ANI, KM, KA, MO

Supervision: ANI, MO

Project administration: ANI, MO

Funding acquisition: ANI, KA, MO

Data availability: The sequence data for RNA-seq and ChIP-seq reported in this paper have

been deposited in the DNA Data Bank of Japan with accession number ESUB001718. The

code is available at https://github.com/okadalabipr/HER2_Cellcycle. All other data will be

made available upon reasonable request.

References

1.   Zetterberg A, Larsson O. Kinetic analysis of regulatory events in G1 leading to

proliferation or quiescence of Swiss 3T3 cells. Proc Natl Acad Sci U S A. 1985; 82:

5365–5369.

2.   Rubin SM, Sage J, Skotheim JM. Integrating old and new paradigms of G1/S control.

Mol Cell. 2020; 80: 183–192.

3.   Matthews HK, Bertoli C, de Bruin RAM. Cell cycle control in cancer. Nat Rev Mol Cell

Biol. 2022; 23: 74–88.

4.   Yu Q, Geng Y, Sicinski P. Specific protection against breast cancers by cyclin D1

ablation. Nature. 2001; 411: 1017–1021.

5.   Adon T, Shanmugarajan D, Kumar HY. CDK4/6 inhibitors: A brief overview and

prospective research directions. RSC Adv. 2021; 11: 29227–29246.

6.   Asghar U, Witkiewicz AK, Turner NC, Knudsen ES. The history and future of targeting

cyclin-dependent kinases in cancer therapy. Nat Rev Drug Discov. 2015; 14: 130–146.

7.   Weigelt B, Peterse JL, van, van 't Veer LJ.. Breast cancer metastasis: Markers and

models. Nat Rev Cancer. 2005; 5: 591–602.

8.   Perou CM, Sørlie T, Eisen MB, van de Rijn M, Jeffrey SS, Rees CA, et al. Molecular

portraits of human breast tumours. Nature. 2000; 406: 747–752.

9.   Sørlie T, Perou CM, Tibshirani R, Aas T, Geisler S, Johnsen H, et al. Gene expression

patterns of breast carcinomas distinguish tumor subclasses with clinical implications.

Proc Natl Acad Sci U S A. 2001; 98: 10869–10874.

10.   Finn RS, Dering J, Conklin D, Kalous O, Cohen DJ, Desai AJ, et al. PD 0332991, a

selective cyclin D kinase 4/6 inhibitor, preferentially inhibits proliferation of luminal

estrogen receptor-positive human breast cancer cell lines in vitro. Breast Cancer Res.

2009; 11: R77.

11.   Lüönd F, Tiede S, Christofori G. Breast cancer as an example of tumour heterogeneity

and tumour cell plasticity during malignant progression. Br J Cancer. 2021; 125: 164–

175.

12.   McGranahan N, Swanton C. Clonal heterogeneity and tumor evolution: Past, present,

and the future. Cell. 2017; 168: 613–628.

13.   Herrera-Abreu MT, Palafox M, Asghar U, Rivas MA, Cutts RJ, Garcia-Murillas I, et al.

Early adaptation and acquired resistance to CDK4/6 inhibition in estrogen receptor–

positive breast cancer. Cancer Res. 2016; 76: 2301–2313.

14.   Knudsen ES, Kumarasamy V, Ruiz A, Sivinski J, Chung S, Grant A, et al. Cell cycle

plasticity driven by mTOR signaling: Integral resistance to CDK4/6 inhibition in

patient-derived models of pancreatic cancer. Oncogene. 2019; 38: 3355–3370.

668 15. Cai Z, Shi Q, Li Y, Jin L, Li S, Wong LL, et al. LncRNA EILA promotes CDK4/6

669   inhibitor resistance in breast cancer by stabilizing cyclin E1 protein. Sci Adv. 2023; 9:

670   eadi3821.

671 16. Ji W, Shi Y, Wang X, He W, Tang L, Tian S, et al. Combined androgen receptor blockade

672   overcomes the resistance of breast cancer cells to palbociclib. Int J Biol Sci. 2019; 15:

673   522–532.

674 17. Li Z, Razavi P, Li Q, Toy W, Liu B, Ping C, et al. Loss of the FAT1 tumor suppressor

675   promotes resistance to CDK4/6 inhibitors via the Hippo pathway. Cancer Cell. 2018; 34:

676   893–905.e8.

677 18. Ross JS, Fletcher JA. The HER-2/neu oncogene in breast cancer: Prognostic factor,

678   predictive factor, and target for therapy. Oncologist. 1998; 3: 237–252.

679 19. Lenferink AE, Busse D, Flanagan WM, Yakes FM, Arteaga CL. ErbB2/neu kinase

680   modulates cellular p27(Kip1) and cyclin D1 through multiple signaling pathways.

681   Cancer Res. 2001; 61: 6583–6591.

682 20. Moasser MM. The oncogene HER2: Its signaling and transforming functions and its role

683   in human cancer pathogenesis. Oncogene. 2007; 26: 6469–6487.

684 21. Yarden Y, Sliwkowski MX. Untangling the ErbB signalling network. Nat Rev Mol Cell

685   Biol. 2001; 2: 127–137.

686 22. Nakakuki T, Birtwistle MR, Saeki Y, Yumoto N, Ide K, Nagashima T, et al. Ligand-

687   specific c-Fos expression emerges from the spatiotemporal control of ErbB network

688   dynamics. Cell. 2010; 141: 884–896.

689 23. Burch PM, Yuan Z, Loonen A, Heintz NH. An extracellular signal-regulated kinase 1-

690   and 2-dependent program of chromatin trafficking of c-Fos and Fra-1 is required for

691   cyclin D1 expression during cell cycle reentry. Mol Cell Biol. 2004; 24: 4696–4709.

692 24. Keller UB, Old JB, Dorsey FC, Nilsson JA, Nilsson L, MacLean KH, et al. Myc targets

Cks1 to provoke the suppression of p27^{Kip1}, proliferation and lymphomagenesis. EMBO

J. 2007; 26: 2562–2574.

25. Dong P, Maddali MV, Srimani JK, Thélot F, Nevins JR, Mathey-Prevot B, et al. Division

of labour between Myc and G1 cyclins in cell cycle commitment and pace control. Nat

Commun. 2014; 5: 4750.

26. Pardee AB. A restriction point for control of normal animal cell proliferation. Proc Natl

Acad Sci U S A. 1974; 71: 1286–1290.

27. Yao G, Tan C, West M, Nevins JR, You L. Origin of bistability underlying mammalian

cell cycle entry. Mol Syst Biol. 2011; 7: 485.

28. Spencer SL, Cappell SD, Tsai F-C, Overton KW, Wang CL, Meyer T. The proliferation-

quiescence decision is controlled by a bifurcation in CDK2 activity at mitotic exit. Cell.

2013; 155: 369–383.

29. Murphy LO, MacKeigan JP, Blenis J. A network of immediate early gene products

propagates subtle differences in mitogen-activated protein kinase signal amplitude and

duration. Mol Cell Biol. 2004; 24: 144–153.

30. Leone G, Sears R, Huang E, Rempel R, Nuckolls F, Park CH, et al. Myc requires

distinct E2F activities to induce S phase and apoptosis. Mol Cell. 2001; 8: 105–113.

31. Lee T, Yao G, Nevins J, You L. Sensing and integration of Erk and PI3K signals by Myc.

PLOS Comput Biol. 2008; 4: e1000013.

32. Chambard J-C, Lefloch R, Pouysségur J, Lenormand P. ERK implication in cell cycle

regulation. Biochim Biophys Acta. 2007; 1773: 1299–1310.

33. Muise-Helmericks RC, Grimes HL, Bellacosa A, Malstrom SE, Tsichlis PN, Rosen N.

Cyclin D expression is controlled post-transcriptionally via a phosphatidylinositol 3-

kinase/Akt-dependent pathway. J Biol Chem. 1998; 273: 29864–29872.

34. Larrea MD, Liang J, Da Silva T, Hong F, Shao SH, Han K, et al. Phosphorylation of

p27Kip1 regulates assembly and activation of cyclin D1-Cdk4. Mol Cell Biol. 2008; 28: 6462–6472.

35. García-Gutiérrez L, Delgado MD, León J. MYC oncogene contributions to release of cell cycle brakes. Genes. 2019; 10: 244.

36. Nagashima T, Shimodaira H, Ide K, Nakakuki T, Tani Y, Takahashi K, et al. Quantitative transcriptional control of ErbB receptor signaling undergoes graded to biphasic response for cell differentiation. J Biol Chem. 2007; 282: 4045–4056.

37. Jeong H, Kim J, Lee Y, Seo JH, Hong SR, Kim A. Neuregulin-1 induces cancer stem cell characteristics in breast cancer cell lines. Oncol Rep. 2014; 32: 1218–1224.

38. Grant GD, Kedziora KM, Limas JC, Cook JG, Purvis JE. Accurate delineation of cell cycle phase transitions in living cells with PIP-FUCCI. Cell Cycle. 2018; 17: 2496–2516.

39. Imoto H, Okada M. Signal-dependent regulation of early-response genes and cell cycle: A quantitative view. Curr Opin Syst Biol. 2019; 15: 100–108.

40. Chung M, Liu C, Yang HW, Köberlin MS, Cappell SD, Meyer T. Transient hysteresis in CDK4/6 activity underlies passage of the restriction point in G1. Mol Cell. 2019; 76: 562–573.e4.

41. Mueller S, Huard J, Waldow K, Huang X, D'Alessandro LA, Bohl S, et al. T160-phosphorylated CDK2 defines threshold for HGF dependent proliferation in primary hepatocytes. Mol Syst Biol. 2015; 11: 795.

42. Garcia-Alonso L, Holland CH, Ibrahim MM, Turei D, Saez-Rodriguez J. Corrigendum: Benchmark and integration of resources for the estimation of human transcription factor activities. Genome Res. 2021; 31: 745.

43. Joyce D, Albanese C, Steer J, Fu M, Bouzahzah B, Pestell RG. NF-κB and cell-cycle regulation: The cyclin connection. Cytokine Growth Factor Rev. 2001; 12: 73–90.

44. Hirano T, Ishihara K, Hibi M. Roles of STAT3 in mediating the cell growth,

differentiation and survival signals relayed through the IL-6 family of cytokine

receptors. Oncogene. 2000; 19: 2548–2556.

45. Ernst J, Kheradpour P, Mikkelsen TS, Shoresh N, Ward LD, Epstein CB, et al. Mapping

and analysis of chromatin state dynamics in nine human cell types. Nature. 2011; 473:

43–49.

46. Road Map Epigenomics Consortium, Kundaje A, Meuleman W, Ernst J, Bilenky M, Yen

A, et al. Integrative analysis of 111 reference human epigenomes. Nature. 2015; 518:

317–330.

47. Liu Z, Venkatesh SS, Maley CC. Sequence space coverage, entropy of genomes and the

potential to detect non-human DNA in human samples. BMC Genomics. 2008; 9: 509.

48. Guo J, Li T, Schipper J, Nilson KA, Fordjour FK, Cooper JJ, et al. Sequence specificity

incompletely defines the genome-wide occupancy of Myc. Genome Biol. 2014; 15: 482.

49. Subramanian A, Tamayo P, Mootha VK, Mukherjee S, Ebert BL, Gillette MA, et al.

Gene set enrichment analysis: A knowledge-based approach for interpreting genome-

wide expression profiles. Proc Natl Acad Sci U S A. 2005; 102: 15545–15550.

50. Ma CX, Gao F, Luo J, Northfelt DW, Goetz M, Forero A, et al. NeoPalAna:

Neoadjuvant palbociclib, a cyclin-dependent kinase 4/6 inhibitor, and anastrozole for

clinical stage 2 or 3 estrogen receptor-positive breast cancer. Clin Cancer Res. 2017; 23:

4055–4065.

51. Gorgoulis V, Adams PD, Alimonti A, Bennett DC, Bischof O, Bishop C, et al. Cellular

senescence: Defining a path forward. Cell. 2019; 179: 813–827.

52. Huang W, Hickson LJ, Eirin A, Kirkland JL, Lerman LO. Cellular senescence: The

good, the bad and the unknown. Nat Rev Nephrol. 2022; 18: 611–627.

53. Trost TM, Lausch EU, Fees SA, Schmitt S, Enklaar T, Reutzel D, et al. Premature

senescence is a primary fail-safe mechanism of ERBB2-driven tumorigenesis in breast

carcinoma cells. Cancer Res. 2005; 65: 840–849.

54. Polk A, Kolmos IL, Kümler I, Nielsen DL. Specific CDK4/6 inhibition in breast cancer:

A systematic review of current clinical evidence. ESMO Open. 2016; 1: e000093.

55. Turner KM, Yeo SK, Holm TM, Shaughnessy E, Guan J-L. Heterogeneity within

molecular subtypes of breast cancer. Am J Physiol Cell Physiol. 2021; 321: C343–C354.

56. Prat A, Perou CM. Deconstructing the molecular portraits of breast cancer. Mol Oncol.

2011; 5: 5–23.

57. Ji W, Zhang W, Wang X, Shi Y, Yang F, Xie H, et al. c-myc regulates the sensitivity of

breast cancer cells to palbociclib via c-myc/miR-29b-3p/CDK6 axis. Cell Death Dis.

2020; 11: 760.

58. Hallett ST, Pastok MW, Morgan RML, Wittner A, Blundell KLIM, Felletar I, et al.

Differential regulation of G1 CDK complexes by the Hsp90-Cdc37 chaperone system.

Cell Rep. 2017; 21: 1386–1398.

59. Alao JP. The regulation of cyclin D1 degradation: Roles in cancer development and the

potential for therapeutic invention. Mol Cancer. 2007; 6: 24.

60. Guiley KZ, Stevenson JW, Lou K, Barkovich KJ, Kumarasamy V, Wijeratne TU, et al.

p27 allosterically activates cyclin-dependent kinase 4 and antagonizes palbociclib

inhibition. Science. 2019; 366: eaaw2106.

61. Sears R, Nuckolls F, Haura E, Taya Y, Tamai K, Nevins JR. Multiple Ras-dependent

phosphorylation pathways regulate Myc protein stability. Genes Dev. 2000; 14: 2501–

2514.

62. McMahon SB. MYC and the control of apoptosis. Cold Spring Harb Perspect Med.

2014; 4: a014407.

63. Yusa K, Zhou L, Li MA, Bradley A, Craig NL. A hyperactive piggyBac transposase for

mammalian applications. Proc Natl Acad Sci U S A. 2011; 108: 1531–1536.

64. Tabata S, Matsuda K, Nagai K, Izumi Y, Takahashi M, Motomura Y, et al. Nuclear

NFκB activity balances purine metabolism in cellular senescence. bioRxiv. 2023.

https://doi.org/10.1101/2023.04.18.536673.

65. Tabata S, Matsuda K, Soeda S, Nagai K, Izumi Y, Takahashi M, et al. NFκB dynamics-

dependent epigenetic changes modulate inflammatory gene expression and induce

cellular senescence. FEBS Journal. 2024; 291: 4951–4968.

66. Livak KJ, Schmittgen TD. Analysis of relative gene expression data using real-time

quantitative PCR and the $2^{-\Delta\Delta CT}$ method. Methods. 2001; 25: 402–408.

67. Martin M. Cutadapt removes adapter sequences from high-throughput sequencing reads.

EMBnet J. 2011; 17: 10–12.

68. Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, et al. STAR: Ultrafast

universal RNA-seq aligner. Bioinformatics. 2013; 29: 15–21.

69. Liao Y, Smyth GK, Shi W. featureCounts: An efficient general purpose program for

assigning sequence reads to genomic features. Bioinformatics. 2014; 30: 923–930.

70. Robinson MD, McCarthy DJ, Smyth GK. edgeR: A Bioconductor package for

differential expression analysis of digital gene expression data. Bioinformatics. 2010;

26: 139–140.

71. R Core Team. A Language and Environment for Statistical Computing. (R Foundation

for Statistical Computing, 2021). https://www.R-project.org/.

72. Korotkevich G, Sukhov V, Budin N, Shpak B, Artyomov MN, Sergushichev A. Fast

gene set enrichment analysis. bioRxiv. 2021. https://doi.org/10.1101/060012.

73. Bolger AM, Lohse M, Usadel B. Trimmomatic: A flexible trimmer for Illumina

sequence data. Bioinformatics. 2014; 30: 2114–2120.

74. Langmead B, Salzberg SL. Fast gapped-read alignment with Bowtie 2. Nat Methods.

2012; 9: 357–359.

75.   Broad Institute. Picard, 2024. http://broadinstitute.github.io/picard.

76.   Ramírez F, Ryan DP, Grüning B, Bhardwaj V, Kilpert F, Richter AS, et al. deepTools2: A

next generation web server for deep-sequencing data analysis. Nucleic Acids Res. 2016;

44: W160–W165.

77.   Heinz S, Benner C, Spann N, Bertolino E, Lin YC, Laslo P, et al. Simple combinations

of lineage-determining transcription factors prime cis-regulatory elements required for

macrophage and B cell identities. Mol Cell. 2010; 38: 576–589.

78.   Durinck S, Spellman PT, Birney E, Huber W. Mapping identifiers for the integration of

genomic datasets with the R/Bioconductor package biomaRt. Nat Protoc. 2009; 4: 1184–

1191.

79.   Bailey TL, Johnson J, Grant CE, Noble WS. The MEME suite. Nucleic Acids Res. 2015;

43: W39–W49.

80.   Castro-Mondragon JA, Riudavets-Puig R, Rauluseviciute I, Lemma RB, Turchi L,

Blanc-Mathieu R, et al. JASPAR 2022: The 9th release of the open-access database of

transcription factor binding profiles. Nucleic Acids Res. 2022; 50: D165–D173.

Fig. 1. G1/S transition occurs approximately 20 h after HRG stimulation. (A–F) MCF-7
cells stably expressing FUCCI. (**A**) Experimental scheme for live-cell imaging using FUCCI
biosensor. (**B**) Images were captured every 20 min up to 72 h in conditions with or without
10 nM HRG. Scale bar: 100 μm. (**C**) Percentage distribution of each cell cycle phase (%) in
each image; $n = 3$ fields per image. (**D**) The number of nuclei increased with each passing
hour relative to the cell number at 0 h. (**E**) Cell cycle phase dynamics estimated based on
Geminin and PIP fluorescence for individual cells followed up to 72 h; $n = 79$. (**F**) Histogram
of cell cycle transition times for cells that completed the cell cycle within 72 h ($n = 77$), based
on the results in (E). Red dotted lines indicate the mean ± SE of each transition time. In (C)
and (D), *$P < 0.05$, **$P < 0.01$, ***$P < 0.001$ (with respect to control and HRG; t-test).

Fig. 2. Time-dependent expression of CDK4 and cyclin D1 in G1/S transition. (A–D) Dynamic changes in the expression or phosphorylation of individual proteins in MCF-7 cells stably expressing FUCCI, detected by western blotting up to 36 h after treatment with 10 nM HRG. (**A, B**) ErbB2 pathway proteins. (**C, D**) Proteins regulating G1/S transition. Protein and phosphorylation levels were normalized relative to those of GAPDH; *n* = 3.

Fig. 3. A population subset enters the cell cycle even after CDK4 inhibition. (**A**) Each
inhibitor (250 nM CDK4i; 600 nM AKTi; 500 nM MEKi) was used to treat the FUCCI stably
expressing MCF-7 cells 12 h after HRG stimulation. Images were acquired every 20 min
from 16 to 72 h after HRG stimulation. Scale bar: 100 μm. (**B**) Percentage distribution of
each cell cycle phase (%) in the overall image at 32 h and 72 h after HRG stimulation; $n = 3$.
(**C**) Increasing number of nuclei from 16 h to 72 h after HRG stimulation; $n = 3$. (**D**) Data
from cells that completed the M/G1 transition within 72 h. Control, $n = 96$; CDK4i, $n = 52$;
AKTi, $n = 42$; MEKi, $n = 75$. Times to G1/S (x-axis) and M/G1 (y-axis) transitions, with
slope (*s*) and intercept (*i*) obtained from the regression equation. (**E**) Comparison of
Pearson's coefficients using data from (D). In (B) to (C), *$P < 0.05$, **$P < 0.01$, ***$P < 0.001$ (with respect to control and each inhibitor; *t*-test).

Fig. 4. CDK4i enhances c-Myc transcriptional activity via H3K27Ac modification. (**A**) Changes in cell-cycle distribution following CDK4 inhibition. (**B–H, J**) MCF-7 cells stably expressing FUCCI. (**B**) Motif-enrichment analysis of H3K27Ac modifications in the promoter region. E-box sequences (CANNTG) were enriched in the promoter region at all time points in the CDK4i group compared to the control group. (**C**) GSEA was performed using the Myc-target gene set (*n* = 58) obtained from the Human MSigDB Collections. Comparison between 0–2 h after HRG stimulation (NES = 1.6693, *P* = 0.004143) and near the R point (12–16 h; NES = –0.8963, *P* = 0.6763). (**D**) GSEA was performed using the Myc-target gene set (C). Comparison between CDK4i and control conditions every 4 h (16 h: NES = 1.39358, *P* = 0.03597; 20 h: NES = 0.8508581, *P* = 0.7883; 24 h: NES = –0.9007113, *P* = 0.6327; 28 h: NES = –1.180552, *P* = 0.2277). (**E–H**) After stimulation with 10 nM HRG for 11.5 h, cells were treated with 0 (DMSO)/0.5/1.0/2.0 nM EN4 (E, F) or 0 (DMSO)/1/2 mM

EN4 (G, H) for 30 min. Subsequently, MCF-7 (**E, F**) and T47D cells (**G, H**) were treated

with 250 nM CDK4i. Scale bar: 100 μm. The percentage distribution of each cell cycle phase

over time was quantified (**F, H**). (**I**) PCNA expression in microarray data from patients with

luminal A subtype breast cancer undergoing long-term (16 weeks, $n = 23$) or short-term (4

weeks, $n = 26$) treatment with CDK4/6i (adj. $P = 0.000132$). (**J**) *PCNA* mRNA expression in

RNA-seq data. (**K**) Increased c-Myc activity induces cell cycle re-entry to promote E2F

target gene expression. In (F) and (H), *$P < 0.05$ (with respect to control and each inhibitor;

t-test). In (I), *$P < 0.05$ (calculated using the Benjamini–Hochberg method). In (J), *$P < 0.05$

(calculated using EdgeR).

Fig. 5. High ErbB2 expression induces cell cycle arrest by repressing cyclin D1

expression via c-Myc under CDK4 inhibition. (**A, B**) WT MCF-7 and ErbB2-

overexpressing MCF-7 cells (high levels) were treated with CDK4i or DMSO 12 h after

HRG stimulation and harvested 24 h after HRG stimulation. Western blotting of (A) p-

ErbB2, p-RB (Ser$^{807/811}$), p-RB (Thr^{373}), RB, p-AKT (Thr^{308}), AKT, p-ERK (Thr^{202}/Thr^{204}),

ERK, c-Myc, cyclin D1, and p27 expression (in control cells, i.e., DMSO treatment) after

normalization with GAPDH expression (B). (**C–F**) ErbB2-overexpressing cells (moderate)

were individually treated with 250 nM CDK4i or DMSO (no treatment) 12 h after stimulation

with 10 nM HRG and fixed 16 h after stimulation with 10 nM HRG (CDK4i or DMSO

addition 4 h later). Hsp90 and ErbB2 were co-immunostained. Scale bar: 25 μm (C). Co-

localization rate of Hsp90 and ErbB2 based on Mander's tM1 and Pearson's R values

determined using Coloc 2 plugin of Image J; $n = 10$ (D). Hsp90 and PY100 were co-

immunostained. Scale bar: 25 μm (E). Co-localization rate of Hsp90 and PY100 based on

Mander's tM2 and Pearson's R values; $n = 10$ (F). (**G, H**) FUCCI stably expressing MCF-7

cells were treated with CDK4i or DMSO (control) as well as 10 mM MG132 12 h after

stimulation with 10 nM HRG and collected 8 h later. (**G**) Expression of ErbB2 and GAPDH

determined by western blotting. (**H**) ErbB2 expression normalized to GAPDH expression; n

$= 3$. (**I**) In CDK4i-unresponsive WT MCF-7 cells, the loss of ErbB2 protein stability impairs

the regulation of c-Myc-mediated G1/S transition, leading to the delayed G1/S transition. (**J**)

WT MCF-7 and ErbB2-overexpressing cells (moderate and high levels) were individually

treated with 250 nM CDK4i or DMSO (no treatment) 12 h after stimulation with 10 nM

HRG, and images were acquired every 20 min until 72 h after stimulation. Scale bar: 100 µm. (**K**) Images in (J) were analyzed to determine the proportion (%) of each cell cycle phase in the entire set of images; $n = 3$. In (B) and (H), $*P < 0.05$ (Tukey's test). In (D) and (F), $*P < 0.05$ (with respect to control and CDK4i; t-test).

Fig. 6. Reversible cell cycle arrest is maintained under c-Myc transcriptional activity.
(**A–G**) MCF-7 cells overexpressing ErbB2 (high levels). (**A, B**) Cells were treated with 250
nM CDK4i 12 h after HRG stimulation, subjected to a wash/no-wash treatment 4 h later, and
harvested 40 h after HRG stimulation. Western blotting results showing the expression of p-
RB, c-Myc, cyclin D1, p27, and GAPDH. After individually normalizing the proteins relative
to GAPDH, the ratio to the no-wash condition was quantified; *n* = 3. (**C**) Washing operations
were performed under the same conditions as in (A, B); cells were fixed 40 h after HRG
stimulation and immunostained with p-c-Myc and cyclin D1 antibodies together with DAPI;
n = 3 859. A higher value on the y-axis indicates that the amount of p-c-Myc in one cell was
greater than that of cyclin D1. The circle shows the vertex of each histogram, and the red
dotted line indicates the G1/S transition time point. (**D, E**) Cells were pre-treated for 30 min
with/without 64 nM c-Myc inhibitor (EN4) 11.5 h after HRG stimulation. The cells were then
incubated with 250 nM CDK4i for 8 h, washed, and imaged 52 h later (72 h after HRG

stimulation, D). (**E**) Percentage distributions of G1 and S/G2/M phases based on the images in (D); $n = 3$. Scale bar: 100 μm. (**F**) Cells were treated with 250 nM CDK4i or DMSO (control) 12 h after HRG stimulation and p27 mRNA levels up to 20 h after HRG stimulation were examined by qPCR; $n = 3$. (**G**) After the cells were treated with or without EN4 under the same conditions as in (D–E), RNA was collected 8 h after treatment with inhibitors, and *CDKN1B* mRNA levels were examined by qPCR; $n = 3$. (**H**) WT MCF-7, WT T47D, and ErbB2 (high) OE MCF-7 cells were pre-treated for 30 min with/without 4 mM c-Myc inhibitor (EN4) 11.5 h after HRG stimulation, incubated with 250 nM CDK4i for 6 d, and SA-β-Gal staining was assessed. (**I**) Percentage SA-β-Gal-stained cells based on the images in (H); $n = 12$. (**J**) ErbB2 levels modulate the irreversibility of the G1/S transition under CDK4i treatment. In (B), (E), (F), (G), (I) $*P < 0.05$ (with respect to control and each inhibitor; t-test).

1022

1023

Fig. 7. Proposed model the ErbB2 receptor regulation during G1/S transition via cyclin D1 and c-Myc in CDK4i non-responsive cells. (**A**) In low ErbB2-expressing cells, CDK4 inhibition leads to reduced pathway activity due to ErbB2 degradation via Hsp90 dis-colocalization. This attenuates the transcriptional activity of c-Myc activated by epigenetic changes. Consequently, a subpopulation of cells unresponsive to CDK4i exhibits a delayed G1/S transition. (**B**) In high ErbB2-expressing cells, increased AKT activity leads to increased c-Myc expression. Subsequently, the expression of p27, which is critical for the stability of cyclin D1/CDK4 during the G1 phase, is suppressed by c-Myc, resulting in cell cycle arrest after CDK4 inhibition. This arrest is reversibly maintained by the strong transcriptional activity of c-Myc, ultimately sustaining survival signals.

1034

1035