Varroa mites escape the evolutionary trap of haplodiploidy

Nurit Eliash[1, 2, 3, 4], Tetsuya Endo[4], Xia Hua[5], J. Spencer Johnston[6], Maeva A. Techer[6, 7], Valerie

R. Holmes[6], Juliana Rangel[6], Huoqing Zheng[7], Evan P. Economo[1,8] and Alexander S.

Mikheyev[4, 9]

[1] Biodiversity and Biocomplexity Unit, Okinawa Institute of Science and Technology

Graduate University, Onna-son, Okinawa, Japan

[2] Shamir Research Institute, Israel

[3] University of Haifa, Israel

[4] Ecology and Evolution Unit, Okinawa Institute of Science and Technology

Graduate University, Onna-son, Okinawa, Japan

[5] Australian National University, Mathematical Sciences Institute, Canberra, Australia 2600

[7] Department of Entomology, Texas A&M University, College Station, Texas, USA

[6] Behavioral Plasticity Research Institute, NSF-BII, USA

[7] College of Animal Sciences, Zhejiang University, Hangzhou 310058, China

[8] Department of Entomology, University of Maryland, College Park, MD, USA

[9] Australian National University, Research School of Biology, Canberra, Australia 2600

*Corresponding author: Nurit Eliash, nurite@gri.org.il

Keywords: haplodiploid, parasite evolution, automixis, varroa

Abstract

Genetic diversity is essential for populations adapting to environmental changes. Due to genetic bottlenecks invasive species often have reduced genetic diversity. However, they must quickly adapt to changes in new environments, potentially including anthropogenic countermeasures. This paradox raises a fundamental question: how do species adapt to changes while having low genetic diversity? This tension is more pronounced for some species, for example parasites go through additional bottlenecks between hosts and haplodiploid species have a lower effective population size, as males can inherit and transmit only half of their mother's genome. To investigate how species with such traits reconcile diversity and adaptability, we examined inheritance in the Varroa mite (*Varroa destructor*), a well-studied invasive parasite of honey bees fitting all of the above criteria. By following the flow of alleles across three-generation pedigrees we found that Varroa, so far believed to be haplodiploid, is actually not. Rather, it has a unique reproductive system in which females clonally produce functionally diploid sons. While males lose somatic DNA during development, they can transmit either maternal allele to their daughters with equal probability. Modeling shows Varroa loses heterozygosity slower under typical sib-mating conditions, relative to haplodiploidy. This permits a greater effective population size relative to a purely haplodiploid system and increased evolutionary potential. This reversion to diploidy is a singular example of escaping the 'evolutionary trap' of haplodiploidy, which is believed to be an evolutionary stable end state. Plasticity in reproductive systems could be more common than assumed, and may potentially explain the remarkable resilience and high adaptivity of Varroa and other invasive parasites.

Significance

Varroa mites have driven the collapse of honey bee populations since their worldwide spread in the middle of the 20th century. Despite repeated genetic bottlenecks, Varroa has adapted to diverse environments and has overcome many pesticides. Using pedigree analysis, we found that Varroa re-evolved diplodiploid reproduction from an evolutionary history of haplodiploidy. Diplodiploidy permits a higher effective population size and evolutionary potential, likely facilitating Varroa's ongoing success. Females produce males clonally, passing on their entire genomes. Varroa is a singular exception to the theoretically

and empirically observed rule that, once evolved, haplodiploidy is an evolutionarily stable

end state (an 'evolutionary trap'). Novel mechanistic studies of even well-known organisms

can lead to surprising insights into the evolutionary plasticity of reproductive systems.

Introduction

Genetic diversity is the basis for biodiversity, being required for species to evolve in

response to environmental changes (1). For a given selection coefficient, the effectiveness

of natural selection *vs.* genetic drift is determined by the effective population size (2).

Selection dominates at higher population sizes, yet in small populations, beneficial alleles

can be lost due to drift. Biological invasions are often accompanied by genetic bottlenecks

that reduce effective population size. Parasites are particularly at risk, as they often

experience bottlenecks as part of their life cycle: an introduction to a new host often

involves one or a few individuals, and the host-parasite population dynamics also involve

frequent and drastic changes in the population's size (3). In addition, because of difficulties

involved in finding unrelated mates within a host, many parasites frequently inbreed,

which further decreases genetic variability (4–6). Although all the above scenarios are

expected to decrease the effective population size, by definition, invasive parasites are

successful species that are highly adaptable to different environments, hosts, and in many

cases, the chemical pesticides used to control them (7, 8). This 'genetic paradox of invasion'

(9, 10) raises the question of how invasive species with expected low genetic diversity are

so successful.

A few solutions have been suggested for this paradox. These include physiological

explanations (11–13), and multiple invasions that compensate for the initial low genetic

diversity (13–15). Occasionally, organisms evolve unusual reproductive strategies that help

them avoid inbreeding. For example, in some ants' males and females constitute genetically

distinct clonal lineages that combine to produce sterile workers but otherwise do not

intermix (16, 17). Detecting such mechanisms requires multi-generation-controlled

crosses, which are often difficult if not impossible, especially in non-domesticated

organisms, suggesting that many unusual reproductive modes remain to be discovered.

Here we examined the reproductive biology of the Varroa mite (*Varroa destructor*), a global
invasive parasite of honey bees. The mites experienced a range of genetic bottlenecks. They
went through a genetic bottleneck when switching from their original host, the eastern
honey bee (*Apis cerana*) to the western honey bee (*Apis mellifera*) (18). They also lose
genetic diversity when colonizing new hives. Lastly, they routinely sib-mate in small
families (typically one male and three to four females) (19, 20). This mating behavior
further increases the strength of genetic drift and makes natural selection less effective.
Varroa is believed to be haplodiploid, which further reduces the effective population size
relative to diplodiploid systems (21, 22). Yet, Varroa shows high adaptability: it efficiently
switches between host species (23–25), and can rapidly evolve resistance to pesticides via
various mechanisms (26). While a high level of adaptability explains the mite's successful
global spread, the mechanisms underlying this adaptability remain puzzling in the light of
regular genetic bottlenecks and routine inbreeding.
98
Varroa is believed to be haplodiploid with fertilized eggs developing into diploid females,
while unfertilized eggs develop into haploid males (20, 27–29). Haplodiploidy was inferred
from two karyotyping studies (30, 31). However, as those studies examined Varroa
embryos, sex could not be directly determined. Häußermann et al. (20) found that unmated
females can produce males, consistent with arrhenotokous parthenogenesis *vs.* paternal
genome elimination (PGE), which is found in many other mites. Yet, while consistent with
haplodiploidy, these lines of evidence are indirect.
106
Haplodiploidy is believed to be evolutionarily stable, sometimes being referred to as an
'evolutionary trap', suggesting limited potential for an evolutionary reversion to diploidy
(32, 33). Cruickshank and Thomas (34) used phylogenetic reconstruction to infer that
Varroa evolved from diploid ancestors, via intermediate pseudo-arrhenotokous species
Note that while Varroa was included in this study, its haplodiploidy was assumed based on
the fact that all members of its family are haplodiploid. However, the taxon sampling in this
study was sparse and until recently, the phylogenetic placement of Varroa was uncertain,
opening conclusions to criticism (35). However, a recent phylogenetic analysis of Varroa
placed it as a subfamily within Laelapidae allowing a more targeted comparison (36). Of

the 19 Laelapid sexually reproducing mites in the Tree of Sex database (37) only a single

one is not haplodiploid – *Dicrocheles phalaenodectes*, a parasite inhabiting the ear canal of

some Noctuid moths. However, the placement of *Dicrocheles* in Laelapidae is based solely

on morphological data, and has been debated (38); it is certainly only distantly related to

Varroa. Thus, based on the sum total of the evidence, haplodiploidy was ancestral to

Varroa.

Theory suggests that reversal to diploidy from arrhenotoky is unlikely (33), and, indeed, no

reversals of arrhenotokous sex determination are known to date (39). Yet, despite strong

circumstantial evidence suggesting the existence of arrhenotokous parthenogenesis in

Varroa, no study has examined Varroa's mode of genetic inheritance, making its success in

spite of genetic handicaps somewhat of a mystery. Here, we explicitly examine this species'

mode of genetic inheritance using multi-generational pedigrees.

Results

Three lines of evidence suggest that Varroa males are functionally diploid: [1] they inherit

two maternal alleles evident in their genotypes (Fig 1), [2] they randomly transmit these

genotypes to their daughters (Fig 1B), and [3] both alleles are expressed in somatic cells

(Fig 2). Interestingly, both flow cytometric and karyotyping investigations suggest that

adult females have twice as much genetic material and chromosomes as males (Fig S1 and

S2). This suggests that during development half the chromosomes are eliminated in males.

Males are produced by diploid arrhenotoky preserving their mothers' genotype.

Adult males can carry up to two alleles (Figure 1, Fig S3). Specifically, males inherit their

mothers' genotype, in contrast to their sisters, which are sexually produced (Fig 1).

Males transmit their two alleles to their daughters with equal probability

Consistently, males' maternal inheritance was also observed for the heterozygous variants.

Looking at the cross between a heterozygous F1 female (AB) and a homozygous F1 male

(AA), we found that 97% of the variants in F2 males maintained their mother's

heterozygosity (Fig 1A and 1C). Furthermore, males not only inherit two alleles, but they

also randomly transmit each allele to their daughters. Looking at the cross between a

homozygous female (AA) and a heterozygous F1 male (AB), we observed that about ~50%

of the variants in F2 females were homozygous (like mother) and the rest were

heterozygous (like father) (Fig 1B and 1D). F2 male offspring of the same cross were

homozygous (AA), just like their mother.

Males carry and express two alleles

Varroa males not only inherit and transmit two alleles, but also express both alleles, as

evident from RNA-seq data of adult female mites and their sons (Fig 2, Table S1). The two

sexes show similar heterozygosity proportions in their RNA, across five families (mean

difference = 0.022 ± 0.10, paired t-test: t = 0.46, df = 4, P = 0.672).

Ploidy assessment suggest females have twice genetic material of males

Flow cytometric and karyotyping investigations suggest that females have twice as much

genetic material as males (Fig S1 and S2). While karyotyping data were in general error-

prone, analysis of eight male and seven female immature mites shows that males had

significantly fewer chromosomes (mean = 8.79 ± 0.92 chromosomes) than the expected

diploid number of 14 (t-test: t = -6.59, df = 7, P = 0.0003; Fig S1). Flow cytometric studies of

75 samples showed that females had diploid peaks (2C) and males have the major peak at

1C, indicating that the bulk of the cells are haploid (Fig S2).

Discussion

In this study, we found that *V. destructor* has evolved from ancestral haplodiploidy (34) to a

novel reproductive strategy with diploid arrhenotokous males that are clones of their

mothers (for a scheme of the hypothesized mode of genetic inheritance see Fig 3). This strategy

has both short-term and long-term potential benefits for Varroa, each overcoming a

different evolutionary 'paradox'. In the short-term, the production of clonal males increases

the females' fitness by assuring transmission of both of her genetic variants. This strategy

can overcome the "paradox of sex" by blending the benefits of sexual and asexual

reproduction. Namely, we found that females can produce sons containing both copies of

their genes, but female eggs are produced meiotically. In the long term, functional

diplodiploidy can potentially increase the effective population size (vs. haplodiploidy) (Fig

S4). A higher effective population size imparts greater evolutionary potential, and indeed

Varroa has been remarkably adaptable despite repeated bottlenecks exemplifying the

"genetic paradox of invasion".

Another evolutionary surprise from this study is the effective loss of haploid arrhenotoky.

Once evolved, asymmetric inheritance appears to be in an evolutionarily stable state,

sometimes referred to as an 'evolutionary trap' (33). Obstacles to re-evolving Mendelian

inheritance are diverse. One obstacle is mechanistic, in that males would need to re-evolve

meiosis, which was always thought to be difficult (40). However, recent work from asexual

stick insects indicates that male meiotic regulatory mechanisms are preserved in the

females (41), so re-evolving male meiosis may be easier than expected. The other obstacle

is evolutionary, as examined in detail by Bull (42). Females reach a fitness optimum in

haplodiploid systems, being able to control male production and thus the sex ratio. Thus,

re-emergence of diploid males would have to be a male trait, and biparental diploid males

must carry mutations that allow them to produce diploid sons. However, given the level of

arrhenotokous female control over fertilization and sex ratios, these traits are theoretically

unlikely to spread (42). Indeed, the only evolutionary feasible transition from

haplodiploidy appears to be towards parthenogenesis, though these are short-lived (43).

Yet, Varroa overcame the constraint on the production of diploid sons by having females

solely control the sex ratio, as well as diploid male production. The production of meiotic

sperm and the evolution of diploid arrhenotokous male production illustrate how the

unforeseen complexity of biological mechanisms may challenge long-standing elegant

mathematical models.

The mechanism by which sex is determined in Varroa remains unclear, but other

facultatively parthenogenetic systems provide insights into how it may occur. For instance,

Cape honey bees (*Apis mellifera capensis*) store new eggs in their oviducts arrested in

metaphase I, while meiosis II takes place only after oviposition (44). If the egg is not

fertilized, it develops into a female by central fusion. If it is, a diploid female is sexually

produced. A similar mechanism may be at play in Varroa, except that unfertilized eggs

undergo central fusion to develop into males while preserving their mother's genotype (Fig

1). Unlike in honey bees, where sex is controlled genetically, an epigenetic factor would be

required for sex determination in Varroa.

While Varroa males are a mosaic of their mother's genotype, they eliminate one copy from

each cell during development (Fig S1 and S2). The mechanisms for genome elimination are

unclear, though genome elimination appears to take place rapidly, as early male nymphal

stages are already haploid. Indeed, elimination may happen during development in the

mother's ovaries, which would explain findings by two previous karyotyping studies that

found the presence of haploid embryos (30, 31). Haploidy in somatic tissues may maintain

the dosage of male-specific genes, which have evolved in a haploid context. Given that the

maternal genotype is preserved (Fig 1), the elimination process appears to be random, and

the male somatic genotype is likely chimeric. Unusually, spermatogenesis in Varroa is an

ongoing process during the adult stage (45, 46). This suggests that genome elimination

happens only in the male somatic tissues while germline remains diploid. Varroa genome

elimination differs from either symbiont-induced or paternal genome elimination

(reviewed in (47, 48)) as it does not arise from a reproductive conflict between the sexes

and thus does not favor the maternal genome.

Conclusions

Our work underscores the importance of comprehensive experimental inquiries into

diverse reproductive mechanisms, even in organisms that appeared to be well studied.

Recent investigations have unveiled a broader spectrum of reproductive strategies than

were previously recognized (40), exemplifying the dynamic nature of biological systems.

Yet, the discovery of a novel reproductive system, such as that in Varroa, can have wide-

ranging fundamental and applied consequences. For instance, Varroa's haplodiploidy is

assumed when considering how it evolves in response to countermeasures, such as

acaricides, breeding honey bees for parasite tolerance, and meiotic drive (49–51).

Furthermore, Varroa has emerged as a compelling model for the study of the evolutionary

dynamics of reproductive systems, offering insights into mechanisms for overcoming

evolutionary constraints and the intricacies of sex determination. These findings open

promising avenues for future research, contributing to a deeper understanding of

evolutionary innovation and reproductive plasticity in Varroa and beyond.

Materials and methods

Honey bees and mites

For the pedigree study: Mites and honey bees (*Apis mellifera* L.) were obtained from

colonies at the Onna village apiary and the experimental apiary of Okinawa Institute of

Science and Technology (OIST) in Okinawa, Japan. All colonies used in the experiment were

maintained without any treatment against Varroa and received sugar solution and 70%

pollen cakes as supplemental feed when necessary. Experiments were conducted between

July and November 2020.

For comparative transcriptomic analysis of Varroa gene expression heterozygosity

in mothers and sons: Mites were collected from *A. mellifera* colonies at the Zhejiang

University, China in November 2024. A total of ten mites were collected from five families:

a mother and her son from each family.

For the flow cytometry study: Mites were freshly collected from drones and worker

brood frames of *A. mellifera* colonies at the Texas A&M University Honey Bee Lab apiary

(Janice and John G. Thomas Honey Bee Facility). Mites were collected in June, July, and

December 2022. Capped cells containing red-eye pupae (day 15) to emerging adults (day

19-20) were carefully opened and inspected for single infestation, following standard

methods and procedures (52). The developmental stages and identification of vigorous

specimens were conducted using a soft and fine paintbrush within a petri dish under a

Leica macro-binocular.

For the karyotyping study: Varroa mites were collected from a drone brood frame

obtained from Onna village apiary, Okinawa, Japan, between January-February 2022. Mite

eggs and nymphs were collected from single-infested cells directly into a PBS buffer, to

preserve their vitality, and fixed within 3 hours after collection. First laid eggs, presumably

developing into males ((52) and personal observations), were collected from prepupa (~3

days post capping). Second and third laid eggs, presumably developing into female mites,

were collected from a pupa in later developmental stages. Protonymphs and deutonymphs

were collected from a pink eye pupa bee brood stage (~8 days post capping). The sex of

the nymphs was determined based on the shape of the end part of the ideosoma: males

have a more rectangular shape, while female nymphs have a perfect round shape (53).

Artificial infestation for Varroa pedigree construction

To obtain a three-generational mite pedigree (F0, F1, and F2), we used a semi-natural

infestation system. This started with the collection of a naturally infesting mated Varroa

female (F0). Her adult daughter (F1) was then artificially introduced into a honey bee pre-

pupal cell and allowed to commence the reproductive phase of its life cycle. The F1 mite

and her offspring (F2) were then collected, at the end of the reproductive cycle. For a

detailed description of the experimental design, see Fig 4 and the sections below. Using this

approach, we could genotype all members of the pedigree, except for the male that mated

with the F0 female.

Collection of naturally infesting families (F0 and F1 generations)

To obtain mites from the first two generations (F0 and F1), naturally infested families were

collected from within the cells of 18- to 20-day-old worker bee pupae. At this

developmental stage of the honey bee host, there is typically one dark brown female mite

(the foundress, F0), one adult male (the foundress's son, F1), at least one light-colored

female (the first foundress's daughter, F1), a few nymphs females, and an egg (27). We

determined each mite's sex and developmental stage according to external morphology and

cuticle coloring based on (53) for the color index, please see Fig S5. To ensure that all

offspring belonged to a single family, we thoroughly examined the cell content to confirm

that it was infested by a single foundress mite and that no other adults were present.

Except for the adult female daughters (F1), the rest of the family members were preserved

in a separate microcentrifuge tube kept at -20 °C for later DNA extraction. The adult F1

daughters were easily distinguished from the F0 foundress mite by their lighter cuticle

color (color index 0-2, see Fig S5), and were assumed to have already copulated with their

brother, the F1 male, by the time of the collection (20). Next, the F1 young adult females

were prepared to be artificially introduced into a freshly capped 5th instar larva cell to start

their first reproductive cycle.

Based on our and others' observations (54), the survival and fecundity of young adult mites

increase if they stay for a few days on an adult nurse bee before starting their first

reproductive cycle. This incubation period is presumably imitating the "dispersal stage", in

which the adult female mite infests an adult honey bee (54, 55). Therefore, the collected

young adult females (F1 generation) spent three days on a nurse bee before infestation.

However, during the experiment, we noticed that the very young adult daughter's cuticle

was light and fragile, and most of the mites did not survive when they were transferred

directly from the pupae onto a nurse bee. Therefore, we let the young mites first feed on a

white-eyed pupa in a gelatin capsule for one to four days until their cuticle was sclerotized

and darker (color index 3-4, see Fig S5). After this intermediate period, the mites were

marked on their dorsal idiosoma (PC-1MR POSCA, Japan), and transferred onto an adult

nurse bee in a cage for another three days, for the above-mentioned "dispersal stage". In all

incubation periods, the mites were kept in a controlled dark environment, imitating hive

conditions (34-34.5 °C, 60-75% RH).

Artificial infestation of F1 female mites

After the artificial "dispersal phase", the mites were brushed off the bees using a fine paint-

brush following anesthesia by a 20 seconds exposure to CO_2, 0.2mPa at 10 L/hour, and

placed on a moist filter paper to avoid desiccating till infestation. The mites were then

introduced to a worker bee cell, one mite per cell, as described by (52). In detail, six hours

before infestation, cells containing a 5th instar larva (about to be sealed) were marked on

an acetate sheet attached to a frame from a mite-free hive. The frame was then placed back

into the hive so that the bees could seal the cells. Five hours later the frame was taken into

a warm room (25-28 °C) for the F1 females to be introduced into the freshly sealed cells.

For the artificial infestation, a precise incision (3-5 mm) was made at the margin of the cell-

capping using a surgical blade (N 11). After the incision was made, the cell-capping was

slightly lifted, just enough for the mite to crawl under the capping and into the cell. The

mite was then placed at the entrance of the cell using a moist fine paintbrush, with its head

facing the cell opening. After the mite had entered the cell, it moved under the larva to the

cell bottom. The cap was then resealed by gently pushing it down using the back end of the

paintbrush. The infested frame was then incubated in a dark controlled environment (34-

34.5 °C, 60-75% RH), and emerging bees were removed daily.

Collection of an artificially infested family (F2 generation)

Ten days after the F1 mites' infestation, the cells were opened, and their contents were

examined. The family members' sex, developmental stage (adult, nymph, or egg), and

viability were recorded. To assure the parenthood of the offspring (F2), double-infested

cells and unmarked mites were not included in the final analysis. In total, out of 78

successful infestations, 14 were double-infested by a non-marked mite. Each individual was

snap-frozen in a separate tube at -20 °C for later DNA extraction. In total, we collected 223

mites belonging to 30 families. Each family was composed of a foundress adult female (F0),

her male and female adult offspring (F1), and their F2 offspring (at least one male/female

mite in various developmental stages).

Genomic analysis of the Varroa pedigree

DNA extraction, library preparation, and sequencing

Genomic DNA was extracted from each single mite following the protocol described

previously (25). In brief, after surface sterilization using absolute ethanol, each mite was

crushed in a 1.5 mL microcentrifuge tube dipped in liquid nitrogen. The crushed mite was

then processed using QIAamp DNA Micro Kit (© Qiagen) following the manufacturer's

instructions and eluted in 20 µL of water. Total dsDNA quantity was measured using a

Qubit™ 4 Fluorometer with an Invitrogen dsDNA HS Assay Kit, and DNA quality was

verified using a Bioanalyzer 2100 with high-sensitivity DNA kits (Agilent, Japan). Short-

inserts of 150-bp paired-end libraries were prepared for each individual using a Nextera

XT DNA Library Preparation Kit (Illumina, Japan) and 16 PCR cycles. cDNA was then

cleaned and size-selected using CA-magnetic beads (Dynabeads® MyOne Carboxylic Acid,

Invitrogen), and 11-11.5% PEG 6000 (Sigma-Aldrich, Japan). The library quality was

measured using a Qubit™ 4 Fluorometer with an Invitrogen dsDNA HS Assay Kit, and its

size was assessed using a 4200 Tapestation with a high-sensitivity D5000 kit (Agilent,

Japan). Libraries were run on a NovaSeq6000 in 150 bp x 2 paired-end mode (Illumina,

Japan) at the OIST Sequencing Center. All biosamples are available in the Sequence Read

Archive (SRA) under the accession number PRJNA794941. Biosample details including sex,

developmental stage, pedigree information, and sequencing coverage, are provided in

Table S2.

Read mapping, variant calling, and filtering, of genomic data

We mapped all reads to the *V. destructor* reference genome [GCF_002443255.1] (18) using

the very sensitive mode of NextGenMap v0.5.0 (56). Reads were sorted and duplicates were

removed using SAMtools (57). We called variants against the reference genome using

FreeBayes v1.1.0 (58) with the following parameters: minimum alternative allele counting

= 5, minimum alternative allele fraction = 0.2, minimum mapping quality = 8, minimum

base quality = 5, and using the four best SNP alleles. The variant calling resulted in a total of

73,903,468 variated sites. As our population consisted of closely related individuals (mite

families from the same apiary, in a high-inbreeding system), we did not expect to detect

many variants. Therefore, we set the minor allele frequency threshold to 0.2, which

includes sites with a minor allele frequency of 20% or more, to exclude any rare variants

due to mutations or possible mapping errors. We included variants only for the seven

chromosomes (excluding the mitochondrial DNA and unscaffolded regions). In addition, we

filtered for the following site quality parameters: we included only bi-allelic sites, a depth

range between 16-40, and maximum missing data per site = 0.5. The variants' quality

distribution showed a clear bimodal distribution with a cutoff at ~10,000 (Fig S6). To

ensure the quality of the variants for the genetic inheritance analysis, we chose a strict

filter of minimum quality of 15,000. Filters were applied using VCFtools v0.1.12b (59)

resulting in 33,925 final variants in 223 individuals. The pipeline from fastq files down to

the VCF file was constructed using Snakemake (60), and codes are available and

reproducible on GitHub [https://github.com/nurit-eliash/Varroa-pedigree-study]. For

further security of site quality, we applied two additional site-quality parameters to

account for biases in alternate allele representation and to ensure high-confidence

genotype calls based on allele frequency and quality. Variant allele frequency (VAF) was

calculated as $VAF = AO/DP$, where AO (Alternate Observations) represents the number of

reads supporting the alternate allele, and DP (Depth) is the total read depth at the site.

Similarly, Quality Allele Frequency (QAF) was computed as $QAF = QA/QA+QO$, where QA

(Quality of Alternate Allele) is the sum of Phred-scaled base quality scores for reads

supporting the alternate allele, and QO (Quality of Reference Allele) is the sum of Phred-

scaled base quality scores for reads supporting the reference allele. To minimize errors in

genotyping due to sequencing artifacts or depth-related biases, we applied genotype-

specific thresholds for VAF and QAF based on observation of values from F2 females of

which parents and grandmother genotypes are fixed, and so their own genotype is

assumed with high confidence (Fig S7, S8 and S9 for QAF and VAF distribution, in

genotypes AA, BB and AB, in accordance). The thresholds (Table 1), ensure that

homozygous reference (AA) sites exhibit low alternate allele support (VAF and QAF < 0.01),

heterozygous (AB) sites maintain a balanced allele distribution (0.2< VAF < 0.8, 0.2 < QAF <

0.8), and homozygous alternate (BB) sites demonstrate high alternate allele support (VAF

and QAF > 0.9). This filtering approach enhances the accuracy of genotype classification

while reducing the inclusion of low-confidence variant calls.

Table 1. Quality Allele Frequency (QAF) and Variant allele frequency (VAF) threshold and

their ideal values used for sites' filtration in the genotype flow of mites' pedigree.

Genotype	VAF threshold	Ideal value, interpretation
AA (Homozygous Reference)	< 0.01	~0, Strong reference support
AB (Heterozygous)	0.2 - 0.8	~0.5, Balanced allele representation
BB (Homozygous Alternate)	> 0.9	~1, Strong alternate support

Genotype validation using Sanger sequencing

Several genotype calls obtained from whole genome sequencing were validated by Sanger

sequencing using the same genomic DNA extracted. For the validation, we randomly

selected three genes, located on different chromosomes, each containing at least one

variant site. In total, we re-sequenced 14 individual mites from four families on six variant

sites. The sequenced region was located on the exons of three genes: alpha-mannosidase

2x-like (LOC111243621 on chromosome NW_019211455.1), 5-oxoprolinase-like

(LOC111252938, on chromosome NW_019211459.1), and probable E3 ubiquitin-protein

ligase RNF144A-A (LOC111254119, on chromosome NW_019211460.1). Primers were

designed with the NCBI primer design tool (utilizing Primer3 and BLAST), with default

parameters and product size set to 300-1,000 bp. Primer sequence, product length, and

gene accession numbers are provided in Table S3. The size of the PCR product was checked

by running in 1% agarose gel (135V, 20 min), purified, and size selected using the solid

phase reverse immobilization (SPRI) method. The purified PCR products were Sanger

sequenced on a ThermoFisher SeqStudio Genetic Analyzer using the original reverse

primer and BigDye® Direct Cycle Sequencing kit (Thermo Fisher Scientific), following the

manufacturer's instructions. We then aligned the reverse complementary sequences to the

Varroa reference genome (GCF_002443255.1), to detect and validate the variant sites using

Geneious Prime 2022.1.1. (https://www.geneious.com). For sequence alignments and the

validated sites please see Fig S10.

Ploidy assessment

Flow cytometry

We used flow cytometry of the Varroa mite nuclei to assess the male and female mites'

ploidy during development from nymphs to adults. Live mites were freshly collected and

snap-frozen in liquid nitrogen before being processed. Samples were ground using 15

strokes of the "A" pestle in a 2 mL Kontes Dounce tissue grinder and filtered through a

40μm nylon mesh. Each sample consisted of an individual mite: for immature mites (larvae

and nymphs), a whole larval body, and for adult mites, dissected somatic or germinal

tissues. The DNA in the nuclei released by grinding was stained for 2 hr under dark

refrigeration with 25 μg/mL propidium iodide. The mean red PI fluorescence of the stained

nuclei was quantified using a Beckman-Coulter (Brea, CA) CytoFlex flow cytometer with a

solid-state laser emitting at 488 nm. As a standard, we used nuclei from the brain of

Drosophila virilis, 1C =328 Mbp (61). The total quantity of DNA in the sample was

calculated as the ratio of red fluorescence of the sample/standard (mean red fluorescence

of the 2C peak of the sample divided by the mean fluorescence of the 2C peak of the

standard), times the 1C amount of DNA in the standard. In total, 75 samples were evaluated

of different sex and developmental stages. We scored these samples to produce biological

replicates, to calculate the standard error of the mean genome size estimate. To identify the

ploidy level of a Cytometric DNA peak (62), genome size is reported as 1C, the mean

amount of DNA in Mb in one copy of a single complete genome.

Karyotyping

Sample fixation and staining: Procedure was based on the air-dry method by (63). In

brief, in a Colchicine-hypotonic solution (0.005 %), embryos were dissected from the eggs

while nymph mites were punctured to allow the solution to penetrate the inner tissues. The

samples were incubated in the Colchicine-hypotonic solution for 20-60 mins to inflate cells,

and increase mitosis events. The samples were then incubated in 60% 1:1 Acetic-ethanol

for 30-60 mins, quickly macerated then treated with 1:1 Acetic-ethanol for 1-2 mins

followed by 100% Glacial acetic acid. The samples were left to dry in room temperature for

one day then stained for 10 mins in 3% Giemsa solution in Sorensen's Phosphate buffer,

and covered with cover slip before visualization.

Karyotyping and data analysis: Chromosomes karyotyping was done by imaging the

fixed samples under a Nikon Ti2 fully motorized inverted microscope equipped with a

linear encoded motorized xy stage, controlled by NIS Elements software. A white LED was

used as a diascopic light source, and pictures were taken using a Nikon DS-Qi2 camera.

Cells undergoing mitosis (in both anaphase and metaphase) were used to determine the

chromosome count in each sample. All imaging was done at the OIST Imaging Section. For

examples of imaging of males and females please see Fig S11 and S12 in accordance.

K-mer distribution

We used the Smudgeplot method (64) to visualize and estimate the ploidy of male and

female genomes by analyzing heterozygous k-mer pair, from the raw reads of a male

Varroa mite (BioProject PRJDB6279, runs DRR121854 and DRR124352 for male and

females respectively).

Transcriptomic analysis of Varroa mother and son

Total RNA was extracted from each single mite and sequenced by Metware Biotechnology

Co., Ltd. (Wuhan, China) following an in-house protocol. RNA integrity was assessed using a

Qubit™ 4 Fluorometer (Invitrogen) and a Qsep400 Bioanalyzer. For mRNA enrichment,

polyadenylated RNA was isolated using Oligo(dT) beads, followed by fragmentation with

fragmentation buffer. First-strand cDNA was synthesized using random hexamers and

reverse transcriptase, while second-strand synthesis was carried out with dNTPs and DNA

polymerase I. The resulting double-stranded cDNA was purified, subjected to end-repair, A-

tailing, and adapter ligation, followed by PCR amplification. Library quality was assessed

using a Qubit™ 4 Fluorometer with an Invitrogen dsDNA HS Assay Kit, and insert size

distribution was confirmed using a Bioanalyzer (Agilent, Japan). The final libraries were

pooled and sequenced on an Illumina platform in paired-end mode. All biosamples are

available in the Sequence Read Archive (SRA) under the accession number PRJNA1216898.

Biosample details including sex, family, and mapped reads, are provided in Table S1.

RNA-seq reads were mapped to the *V. destructor* reference genome

[GCF_002443255.2_Vdes_3.0] using HISAT2 v2.2.1 in default settings (65). The mapping

rate varied between samples, with uniquely mapped reads ranging from ~5.88% to

15.71% of the total reads. Clean reads were obtained after quality filtering using fastq

v0.23.2 (66), with adapters removed, reads containing more than 10% N bases discarded,

and bases with quality scores below 20 trimmed. Variant calling was performed using

GATK (67), and SNPs and InDels were annotated with ANNOVAR (68). To ensure high-

confidence variant sites, we filtered out low-quality SNPs and retained only sites with

adequate coverage and quality metrics, although the exact thresholds were not specified in

the company report. We observed thousands of exonic variants per sample, including both

synonymous and nonsynonymous SNVs, as well as stop-gain mutations. The resulting

variant dataset included annotations for exonic, 5' and 3' UTR regions.

Mathematical modeling

We followed the classical approach used by (69) to mathematically model declines in heterozygosity due to inbreeding within haplodiploid systems. This approach involves constructing path diagrams representing genetic transmission paths from ancestors to descendants and quantifying genetic relatedness through path coefficients. By enumerating paths and calculating their coefficients, we derived two mathematical relationships describing the expected declines in heterozygosity per generation under the reproductive life cycle of Varroa. One for the reproductive system with arrhenotokous diploid males and the other for haploid males. These two equations allowed a direct comparison between the two reproductive systems. Figure 4 plots the two equations in terms of how the rate of decline in heterozygosity changes with the degree of inbreeding, which is measured by the probability of co-founding by females of the same mother. Mathematical details are in Supporting Information 1.

Statistical analysis

All analyses were carried out in the R statistical environment (70). All bioinformatic analyses and figures are available and reproducible directly from the online supplementary data [https://rpubs.com/Nurit_Eliash/1288713].

Acknowledgments

ASM was supported by a Future Fellowship from the Australian Research Council (FT160100178). HZ was supported by the earmarked fund for CARS (CARS-44). We wish to thank local beekeepers of Okinawa, Japan: Tsuto Arakaki, Takashi Ikemiya, and Ryo Kirino, from the Agricultural Environment Coordinator in Onna-village, for their help in providing Varroa-infested colonies, and Ms. Saori Chappell, OIST, for coordinating the sample collection and administrative helping throughout. At OIST, we thank the Imaging Section for providing access to microscopes and Dr. Paolo Barzaghi for support, Jan Moran and the HPC team for their support. We wish to thank Olliver Stewart, ANU, for helping in chromosome karyotyping and Alison McAfee for providing the beautiful illustrations of Varroa mites and honey bees incorporated in the manuscript figures. At Texas A&M, we

would like to extend our warm thanks to Jordan Twombly Ellis for readily sharing some infested frames to collect the last needed Varroa. We also thank Ms. Cristina de Morais Cianci from the Editorial Office of Genetics and Molecular Biology journal, for kindly providing a copy of the fundamental research on Varroa cytogenetics (30). We also wish to thank Yoav Ram for commenting on the earlier draft of the manuscript. Finally, we wish to thank Mikheyev and Economo lab members for their continuous support and discussion throughout.

References

1. D. H. Reed, R. Frankham, Correlation between fitness and genetic diversity. *Conserv. Biol.* **17**, 230–237 (2003).

2. R. Kliman, B. Sheehy, J. Schultz, Genetic drift and effective population size. *Nature Education* **1**, 3 (2008).

3. A. Papkou, C. S. Gokhale, A. Traulsen, H. Schulenburg, Host-parasite coevolution: why changing population size matters. *Zoology (Jena)* **119**, 330–338 (2016).

4. R. Frankham, K. Ralls, Inbreeding leads to extinction: Conservation biology. *Nature* **392**, 441–442 (1998).

5. D. Spielman, B. W. Brook, D. A. Briscoe, R. Frankham, Does inbreeding and loss of genetic diversity decrease disease resistance? *Conserv. Genet.* **5**, 439–448 (2004).

6. D. Charlesworth, J. H. Willis, The genetics of inbreeding depression. *Nat. Rev. Genet.* **10**, 783–796 (2009).

7. R. C. Venette, W. D. Hutchison, Invasive insect species: Global challenges, strategies & opportunities. *Front. Insect Sci.* **1**, 650520 (2021).

8. J. A. Siddiqui, *et al.*, Insights into insecticide-resistance mechanisms in invasive species: Challenges and control strategies. *Front. Physiol.* **13**, 1112278 (2022).

9. F. W. Allendorf, L. L. Lundquist, Introduction: Population biology, evolution, and control of invasive species. *Conserv. Biol.* **17**, 24–30 (2003).

10. R. Frankham, Resolving the genetic paradox in invasive species. *Heredity* **94**, 385 (2005).

11. A. K. Sakai, *et al.*, The population biology of invasive species. *Annu. Rev. Ecol. Syst.* **32**, 305–332 (2001).

12. C. M. Suehs, L. Affre, F. Médail, Invasion dynamics of two alien *Carpobrotus* (Aizoaceae) taxa on a Mediterranean island: II. Reproductive strategies. *Heredity (Edinb.)* **92**, 550–556 (2004).

13. F. Manfredini, M. Arbetman, A. L. Toth, A potential role for phenotypic plasticity in invasions and declines of social insects. *Front. Ecol. Evol.* **7** (2019).

14. L. M. Chevin, A. A. Hoffmann, Evolution of phenotypic plasticity in extreme environments.
Philos. Trans. R. Soc. Lond. B Biol. Sci. **372** (2017).

15. M. A. Tonione, N. Reeder, C. C. Moritz, High genetic diversity despite the potential for stepping-
stone colonizations in an invasive species of gecko on Moorea, French Polynesia. *PLoS One* **6**,
e26874 (2011).

16. M. O. Miyakawa, A. S. Mikheyev, QTL mapping of sex determination loci supports an ancient
pathway in ants and honey bees. *PLoS Genet.* **11**, e1005656 (2015).

17. O. Rey, B. Facon, J. Foucaud, A. Loiseau, A. Estoup, Androgenesis is a maternal trait in the
invasive ant *Wasmannia auropunctata*. *Proc. Biol. Sci.* **280**, 20131181 (2013).

18. M. A. Techer, *et al.*, Divergent evolutionary trajectories following speciation in two
ectoparasitic honey bee mites. *Communications Biology* **2**, 357 (2019).

19. E. Frey, H. Schnell, P. Rosenkranz, Invasion of *Varroa destructor* mites into mite-free honey bee
colonies under the controlled conditions of a military training area. *J. Apic. Res.* **50**, 138–144
(2011).

20. C. K. Häußermann, *et al.*, Reproductive parameters of female *Varroa destructor* and the impact
of mating in worker brood of *Apis mellifera*. *Apidologie* **51**, 342–355 (2020).

21. R. A. Metcalf, J. C. Marlin, G. S. Whitt, Low levels of genetic heterozygosity in hymenoptera.
Nature **257**, 792–794 (1975).

22. A. Moro, *et al.*, Adaptive population structure shifts in invasive parasitic mites, *Varroa*
destructor. *Ecol. Evol.* **11**, 5937–5949 (2021).

23. B. P. Oldroyd, Coevolution while you wait: *Varroa jacobsoni*, a new parasite of western
honeybees. *Trends Ecol. Evol.* **14**, 312–315 (1999).

24. V. Dietemann, *et al.*, Population genetics of ectoparasitic mites *Varroa* spp. in Eastern and
Western honey bees. *Parasitology* 1–11 (2019).

25. M. A. Techer, J. M. K. Roberts, R. A. Cartwright, A. S. Mikheyev, The first steps toward a global
pandemic: Reconstructing the demographic history of parasite host switches in its native
range. *Mol. Ecol.* **31**, 1358–1374 (2022).

26. G. A. Mitton, *et al.*, More than sixty years living with *Varroa destructor*: a review of acaricide
resistance. *Int. J. Pest Manage.* 1–18 (2022).

27. P. Rosenkranz, P. Aumeier, B. Ziegelmann, Biology and control of *Varroa destructor*. *J. Invertebr.*
Pathol. **103**, 96–119 (2010).

28. K. S. Traynor, *et al.*, *Varroa destructor*: A complex parasite, crippling honey bees worldwide.
Trends Parasitol. **36**, 592–606 (2020).

29. F. Nazzi, Y. Le Conte, Ecology of *Varroa destructor*, the major ectoparasite of the western honey
bee, *Apis mellifera*. *Annu. Rev. Entomol.* **61**, 417–432 (2016).

30. J. Steiner, S. D. Pompolo, C. S. Takahashi, L. S. Goncalves, Cytogenetics of the acarid *Varroa*

jacobsoni. Rev. Bras. Genet. **4**, 841–844 (1982).

31. A. Ruijter, N. Pappas, Karyotype and sex determination of *Varroa jacobsoni* Oud in *Varroa*
Jacobsoni Oud. Affecting Honey Bees : Present Status and Needs, R. Cavalloro, Ed. (Rotterdam:
Published for the Commission of the European Communities, 1983), pp. 41–44.

32. D. Bachtrog, *et al.*, Sex determination: why so many ways of doing it? *PLoS Biol.* **12**, e1001899
(2014).

33. J. J. Bull, Sex determining mechanisms: an evolutionary perspective. *Experientia* **41**, 1285–1296
(1985).

34. R. H. Cruickshank, R. H. Thomas, Evolution of haplodiploidy in *dermanyssine* mites (Acari:
Mesostigmata). *Evolution* **53**, 1796–1803 (1999).

35. H. Blackmon, N. B. Hardy, L. Ross, The evolutionary dynamics of haplodiploidy: Genome
architecture and haploid viability. *Evolution* **69**, 2971–2978 (2015).

36. J. Oh, *et al.*, Molecular phylogeny reveals Varroa mites are not a separate family but a subfamily
of Laelapidae. *Sci. Rep.* **14**, 13994 (2024).

37. Tree of Sex Consortium, Tree of Sex: a database of sexual systems. *Sci. Data* **1**, 140015 (2014).

38. G. J. DE Moraes, *et al.*, Catalogue of the free-living and arthropod-associated *Laelapidae*
canestrini (Acari: Mesostigmata), with revised generic concepts and a key to genera. *Zootaxa*
5184, 1–509 (2022).

39. B. B. Normark, The evolution of alternative genetic systems in insects. *Annu. Rev. Entomol.* **48**,
397–423 (2003).

40. L. Ross, A. J. Mongue, C. N. Hodson, T. Schwander, Asymmetric inheritance: The diversity and
Evolution of non-Mendelian reproductive strategies. *Annu. Rev. Ecol. Evol. Syst.* **53**, 1–23
(2022).

41. G. Forni, B. Mantovani, A. S. Mikheyev, A. Luchetti, Parthenogenetic stick insects exhibit
signatures of preservation in the molecular architecture of male reproduction. *Genome Biol.*
Evol. **16** (2024).

42. J. J. Bull, An advantage for the evolution of male haploidy and systems with similar genetic
transmission. *Heredity* **43**, 361–381 (1979).

43. M. Neiman, S. Meirmans, P. G. Meirmans, What can asexual lineage age tell us about the
maintenance of sex? *Ann. N. Y. Acad. Sci.* **1168**, 185–200 (2009).

44. K. Sasaki, Y. Obara, Egg activation and timing of sperm acceptance by an egg in honeybees (*Apis*
mellifera L.). *Insectes Soc.* **49**, 234–240 (2002).

45. G. Alberti, H. Hanel, Fine structure of the genital system in the bee parasite, *Varroa jacobsoni*
(Gamasida: Dermanyssina) with remarks on spermiogenesis, spermatozoa and capacitation.
Exp. Appl. Acarol. **2**, 63–104 (1986).

46. C. K. Häußermann, B. Ziegelmann, P. Rosenkranz, Spermatozoa production in male *Varroa*

destructor and its impact on reproduction in worker brood of *Apis mellifera*. *Exp. Appl. Acarol.* **74**, 43–54 (2018).

47. D. Dedukh, A. Krasikova, Delete and survive: strategies of programmed genetic material elimination in eukaryotes. *Biol. Rev. Camb. Philos. Soc.* **97**, 195–216 (2022).

48. M. Herbette, L. Ross, Paternal genome elimination: patterns and mechanisms of drive and silencing. *Curr. Opin. Genet. Dev.* **81**, 102065 (2023).

49. A. Moro, T. Blacquière, D. Panziera, V. Dietemann, P. Neumann, Host-parasite co-evolution in real-time: Changes in honey bee resistance mechanisms and mite reproductive strategies. *Insects* **12**, 120 (2021).

50. N. R. Faber, A. B. Meiborg, G. R. Mcfarlane, G. Gorjanc, B. A. Harpur, A gene drive does not spread easily in populations of the honey bee parasite *Varroa destructor*. *Apidologie (Celle)* **52**, 1112–1127 (2021).

51. A. L. Beaurepaire, *et al.*, Population genetics of ectoparasitic mites suggest arms race with honeybee hosts. *Sci. Rep.* **9**, 11355 (2019).

52. V. Dietemann, *et al.*, Standard methods for varroa research. *J. Apic. Res.* **52**, 1–54 (2013).

53. C. R. Büchler, C. Costa, F. Mondet, N. Kezic, M. Kovacic, Screening for low Varroa mite reproduction (SMR) and recapping in European honey bees. (2017). Available at: https://www.beebreeding.net/wp-content/uploads/2017/11/RNSBB_SMR-recapping_protocol_2017_09_11.pdf [Accessed 15 January 2023].

54. X. Xie, Z. Y. Huang, Z. Zeng, Why do Varroa mites prefer nurse bees? *Sci. Rep.* **6**, 28228 (2016).

55. V. Piou, J. Tabart, V. Urrutia, J. L. Hemptinne, A. Vétillard, Impact of the phoretic phase on reproduction and damage caused by *Varroa destructor* (Anderson and Trueman) to its host, the European honey bee (*Apis mellifera* L.). *PLoS One* **11**, e0153482 (2016).

56. F. J. Sedlazeck, P. Rescheneder, A. von Haeseler, NextGenMap: fast and accurate read mapping in highly polymorphic genomes. *Bioinformatics* **29**, 2790–2791 (2013).

57. H. Li, *et al.*, The Sequence Alignment/Map format and SAMtools. *Bioinformatics* **25**, 2078–2079 (2009).

58. E. Garrison, G. Marth, Haplotype-based variant detection from short-read sequencing. *arXiv [q-bio.GN]* (2012).

59. P. Danecek, *et al.*, Twelve years of SAMtools and BCFtools. *Gigascience* **10**, giab008 (2021).

60. J. Köster, S. Rahmann, Snakemake--a scalable bioinformatics workflow engine. *Bioinformatics* **28**, 2520–2522 (2012).

61. J. S. Johnston, A. Bernardini, C. E. Hjelmen, Genome size estimation and quantitative cytogenetics in insects. *Methods Mol. Biol.* **1858**, 15–26 (2019).

62. J. Rangel, K. Strauss, K. Seedorf, C. E. Hjelmen, J. S. Johnston, Endopolyploidy changes with age-related polyethism in the honey bee, *Apis mellifera*. *PLoS One* **10**, e0122208 (2015).

63. H. T. Imai, A manual for ant chromosome preparations (an improved air-drying method) and
Giemsa staining. *Chromosome Sci.* **19**, 57–66 (2016).

64. T. R. Ranallo-Benavidez, K. S. Jaron, M. C. Schatz, GenomeScope 2.0 and Smudgeplot for
reference-free profiling of polyploid genomes. *Nat. Commun.* **11**, 1432 (2020).

65. D. Kim, B. Langmead, S. L. Salzberg, HISAT: a fast spliced aligner with low memory
requirements. *Nat. Methods* **12**, 357–360 (2015).

66. S. Chen, Y. Zhou, Y. Chen, J. Gu, fastp: an ultra-fast all-in-one FASTQ preprocessor.
Bioinformatics **34**, i884–i890 (2018).

67. A. McKenna, *et al.*, The Genome Analysis Toolkit: a MapReduce framework for analyzing next-
generation DNA sequencing data. *Genome Res.* **20**, 1297–1303 (2010).

68. K. Wang, M. Li, H. Hakonarson, ANNOVAR: functional annotation of genetic variants from high-
throughput sequencing data. *Nucleic Acids Res.* **38**, e164 (2010).

69. S. Wright, Inbreeding and homozygosis. *Proc. Natl. Acad. Sci. U. S. A.* **19**, 411–420 (1933).

70. R. C. Team, R: A language and environment for statistical computing. (2013).

Main figures:

Figure 1. Inheritance patterns in *Varroa destructor* mites shows that males inherit two

alleles from their mothers and transmit both of them to their daughters with equal

probability. Females are produced sexually.

Panels A and B show the genotype flow of sites on the largest chromosome, NW_019211454,

for a representative family spanning three generations: foundress grandmother (F0 female),

her offspring (F1 female and male), and their F2 offspring (F2 female and male). A: Family

478, crossing a heterozygous F1 female (AB) with a homozygous F1 male (AA), showing the

genotypes of their F2 female and male offspring across 33 sites on chromosome NW_019211454. B:

Family 534, crossing a homozygous F1 female (AA) with a heterozygous F1 male (AB), showing the

genotypes of their F2 female and male offspring across 14 sites on chromosome NW_019211454.

Panels C and D show the pooled genotype proportions in F2 females and males from all

families, across all sites on all seven chromosomes, for the same cross types as in A and B,

respectively.

C: Pooled data for the cross of heterozygous F1 female (AB) × homozygous F1 male (AA): 7 F2

females (11,207 sites) and 12 F2 males (16,515 sites). D: Pooled data for the cross of homozygous

F1 female (AA) × heterozygous F1 male (AB): 11 F2 females (4,999 sites) and 10 F2 males (2,536

sites).

F2 females show a ~1:1 ratio of AA:AB genotypes, close to sexual reproduction, while F2 males

predominantly inherit their mother's genotype, suggesting maternal inheritance. The data is

filtered using strict quality thresholds (VAF < 0.01 for AA, 0.2-0.8 for AB, > 0.9 for BB;

corresponding QAF thresholds) to ensure high-confidence genotype calls.

As expected from AA × AB crosses, no BB genotypes were observed. F0 males were not collected for

the pedigree, as they remain in the cell when the honey bee emerges. However, based on the

genotypes of the F0 and F1 offspring, we can infer the F0 male genotype: In the cross of F1 AB

female × AA male, the F0 male must have at least one B allele (indicated as "?/B" in panel A). In the

cross of F1 AA female × AB male, the F0 male must have at least one A allele (indicated as "?/A" in

panel B). Colors: Orange = AA, Green = AB, Blue = BB. Note: "?" indicates an inferred but unobserved

allele.

Figure 2. RNA-seq analysis of heterozygous site proportions in male and female mites. Each point represents the mean proportion of heterozygous sites per individual mite, with error bars showing 95% confidence intervals and lines connecting mother-son pairs. The two sexes show similar heterozygosity proportions in their RNA, across five families (mean difference = 0.022 ± 0.10, paired t-test: t = 0.46, df = 4, P = 0.672). The y-axis shows the proportion of heterozygous sites as a percentage of total sites analyzed. Males are much smaller than females and result in lower-quality sequencing libraries, possibly explaining their greater variability. Note, that while males and females are clones, we do not expect identical heterozygosities in transcriptomic data because different genes are expressed in the two sexes.

Figure 3. Hypothesized mode of genetic inheritance in varroa male and female mites.

Females are produced via diploid sexual reproduction, while males are produced via diploid

arrhenotoky. To illustrate the maternal and paternal alleles inheritance in the two sexes, we show

an example of a cross between a homozygous female and a heterozygous male, and their possible

female and male offspring (as in Figure 1). While the eggs start out as diploid in both sexes, males

lose half their genome during development (Figs S1 and S2).

Semi *in vivo* varroa rearing

Figure 4. Pedigree experimental design. A three-generation Varroa pedigree (F0-in red, F1-in

orange, and F2-in yellow) of 30 families, was constructed by the following steps: **A.** Collecting a

naturally infested family: foundress female mite (F0), and her two adult offspring: male and female

F1 mites (F0 male's genotype was not possible to study because the F0 female was previously

mated, typically by a single male); **B.** "Incubation phase": (B1) Transferring the young F1 female

onto a bee pupa for 1-4 days, till cuticle sclerotized and darkened; (B2) Transferring F1 female onto

a nurse adult bee for another 3 days; **C.** Introducing the matured F1 female to a freshly capped 5th

instar larva cell; **D.** F1 female reproducing inside the bee cell; **E.** 10 days post introduction,

collecting the artificially infested family: F1 female and her F2 offspring. The F2 generation included

a son and one or more daughter nymphs. This design provided known parents and genotypes for

the F1 and F2 generations, allowing a detailed examination of inheritance.