Title: Transcription elongation is dictated by single residues in the histone core domain

Authors: Zachary H. Harvey,[1,*] Kathryn M. Stevens,[2,3] Tobias Warnecke,[2,3,4,**] Frédéric

Berger[1,***]

Affiliations:

[1]Gregor Mendel Institute, Austrian Academy of Sciences, Vienna, Austria

[2]MRC Laboratory of Medical Sciences, London, United Kingdom

[3]Institute of Clinical Sciences, Imperial College London, London, United Kingdom

[4]Trinity College, Oxford, United Kingdom

*Correspondence: zachary.harvey@gmi.oeaw.ac.at

**Correspondence: tobias.warnecke@lms.mrc.ac.uk

***Correspondence: frederic.berger@gmi.oeaw.ac.at

Abstract: The chromatin fiber is thought to suppress transcription in eukaryotes by acting as

a structural barrier. However, once begun, transcription can readily proceed on chromatin,

suggesting this model is insufficient. Here, we establish that the ultra-conserved core domain

of the ancestral histone H2A.Z dictates transcription elongation via direct interaction of its loop

2 region with the RNAPII subunit Spt6, rather than biophysical changes to chromatin.

Interrogating H2A.Z sequences representing more than a billion years of eukaryotic evolution

in a single synthetic host, we show that Spt6 can distinguish even single-residue substitutions

within their loop 2, driving either super-repressed or -activated transcriptional states. Our

results place the histone core domain at the origin of eukaryotic gene expression, establishing

its transformative power to shape transcription.

Main Text

Eukaryotic RNA polymerase II (RNAPII) transcribes mRNA from a chromatin fiber organized into repeating units of ~147 bp of DNA wrapped around a multi-subunit protein complex, the nucleosome (*1*). An enormous body of work has established the importance of nucleosomes in regulating transcription (*2-7*) in development (*8, 9*), disease (*10-12*), and in response to stress (*13*). The prevailing model is that nucleosomes suppress transcription as structural obstacles to RNAPII, and that this conflict must be alleviated to achieve context-specific gene expression (*14-18*). However, recent reports have established that RNAPII has sophisticated mechanisms for navigating nucleosomes without disassembly during transcription elongation (*19, 20*), and once transcription is initiated, eukaryotic RNAPII can readily traverse a chromatinized template (*21-23*), suggesting that the chromatin fiber is not inherently and uniformly suppressive of transcription.

One explanation for the role of the chromatin fiber in shaping transcription is through changes in its composition via incorporation of alternative histone forms, referred to as variants (*4, 5, 24-28*). Consistent with this view, the suite of histone variants found within the transcriptional unit varies according to transcriptional state (*29-31*), with distinct profiles for either repressed (*4, 5*) or activated (*2-4, 11, 26, 32-54*) chromatin domains. Further, histone variants often have divergent biophysical properties *in vitro* (*41, 42*). However, most are highly structurally similar (*55, 56*), and how these physical properties impact transcription *in vivo* remains unresolved.

To address whether and how changes in the chromatin fiber impact transcription, we developed a synthetic complementation paradigm using the histone H2A variant H2A.Z as a model. H2A.Z is an essential feature of eukaryotic chromatin, and has been tightly linked to both transcription activation (*2, 3, 26, 32, 38, 47, 49*) and repression (*11, 33, 48, 50-53*) in different species. We reasoned that we could leverage divergent observations of H2A.Z

function across species as a natural toolkit to generate functional variation in the chromatin fiber. We interrogated H2A.Z sequence variation representing more than a billion years of eukaryotic evolution in a single experimentally tractable synthetic system, the fission yeast *Schizosaccharomyces pombe*, replacing the endogenous *S. pombe* H2A.Z (Pht1) with H2A.Z sequences from diverse eukaryotes. Remarkably, we identify that the RNAPII subunit Spt6 directly reads the primary sequence of the loop 2 (L2) region of the H2A.Z core domain, with even single amino acid changes to its core motif being sufficient to dictate transcription elongation. Together, our findings indicate that an ancient interaction between H2A.Z and Spt6 informs RNAPII directly of the nature of the chromatin template, providing insight into its fundamental role in shaping eukaryotic gene expression, and establishing the nucleosome core as a powerful source of evolutionary innovation.

H2A.Z sequence encodes its transcriptional function

To build our toolkit of H2A.Z sequences, we first constructed a phylogenetic tree of all H2A sequences from diverse eukaryotes in the EukProt database (Fig. S1, S2A, S3; Table S1; Data File S1; Methods). As previously suggested from analyses of smaller datasets (*40, 57*), we found that H2A.Z forms a monophyletic clade that spans nearly the entirety of known eukaryotic diversity (Fig. S2B, S3A). Almost all sequences in this clade, but very few outside of it, harbor the signature DEELD motif located in a C-terminal region of H2A.Zs referred to as the docking domain (DD) (*40*). A small cluster of sequences, mostly from the unicellular flagellate *Carpediemonas membranifera* (phylum Metamonada), branch just outside the H2A.Z clade, contain a near-perfect, but nevertheless variable, DEELD motif and might represent rare, divergent forms of H2A.Z (Fig. S2B). Other H2A variants either have a more restricted phylogenetic distribution (e.g. macroH2A and H2A.W) or cannot be traced to a single ancestral sequence, appearing paraphyletic (e.g. H2A.X) (Fig. S3). Importantly, we found

H2A.Z orthologs in nearly every eukaryotic genome, even in deep-branching clades like

Metamonada that were previously thought to lack H2A.Z (Fig. S2B) (*58*). Thus, the

phylogenetic distribution points to H2A.Z being present in the last eukaryotic common

ancestor (LECA), and broadly retained throughout the evolution of eukaryotes. These findings

support the view that H2A.Z is the only H2A variant whose ancestry can be traced back to the

LECA.

We selected nine H2A.Z sequences from across the phylogenetic tree (Fig. S2B and S4A,

Data File S2). The sequences were selected to maximize the number of eukaryotic supergroups

represented, and their total sequence diversity. Although they span more than a billion years of

evolution, these representative H2A.Z proteins were ~65–90% identical in their core domain

(excluding highly divergent N- and C-terminal tail regions; Fig. S4B), highlighting their high

degree of conservation.

To test whether transcription is impacted by sequence diversity in these nine

representative H2A.Z proteins, we synthesized *S. pombe* codon-optimized versions of the nine

H2A.Z ortholog coding sequences, and generated yeast lines in which the endogenous gene

encoding H2A.Z (*pht1*) was replaced by one of these orthologs (Fig. 1A). We then assessed

whether these replacement lines phenocopied the wild-type (WT) gene, resembled a H2A.Z

deletion mutant (*pht1⁻*), or differed from both.

We assessed global transcriptional changes in the replacement lines by using strand-

specific mRNA-seq. Consistent with prior reports (*2*), *pht1⁻* cells exhibited modest

transcriptional changes, affecting ~10% of detected transcripts (Fig. S4C). By contrast, many

H2A.Z replacement lines showed changes in ≥20% of detected transcripts when compared with

WT (Fig. S4C). To investigate this further, we examined how similar the transcriptomes of the

nine H2A.Z replacement lines were to either WT or *pht1⁻* using principal component analysis.

Replacement lines differed markedly both from WT and *pht1⁻* (Fig. 1B), suggesting that

H2A.Zs from different species are not simply gradations of total loss of function alleles, but

give rise to sequence-specific gain-of-function states.

We next asked whether transcriptional variation in H2A.Z from different species had

consequence at the level of phenotype. To do this, we assessed the growth of H2A.Z

replacement lines using a plate-based assay in the presence of stress conditions targeting

metabolism (rapamycin (*59*)), protein folding (low-dose guanidinium hydrochloride (*60*)), cell

wall biogenesis (the anti-fungal drug clotrimazole (*61*)), fermentation (ethanol (*62*)), or either

directly (doxorubicin (*63*), formamide(*64*)) or indirectly (caffeine (*65*)) transcription and

chromatin (Fig. 1C). Consistent with functional variation at the level of transcriptomes, the

growth of all nine replacement lines differed substantially from WT and *pht1*⁻ cells across all

tested conditions (Fig. 1C), with most lines growing more slowly than *pht1*⁻ cells. Notably,

greater sequence divergence of replacement orthologs from *S. pombe* Pht1 correlated with

weaker growth in several stress conditions, with the strongest correlation being for conditions

known to be linked directly to transcription (doxorubicin, formamide) or stresses known to

require a transcriptional response (ethanol (*62*)) (Fig. 1D, S4D). By contrast, we saw no strong

correlation between the level of H2A.Z expression and growth (Fig. 1D, S4D), indicating that

these effects were not attributable to differential expression of H2A.Z but likely were related

to differences in the primary sequence. Together, these data link primary sequence variation in

H2A.Z from different species to variation in its function and confirm a tight linkage between

H2A.Z and transcription.

The L2 region of H2A.Z determines its function

Histone proteins are composed of a core histone-fold domain, characterized by a nearly

sequence-invariant bundle of three alpha helices (α1, α2, and α3), bounded by mostly

unstructured N- and C-terminal tails and separated by two unstructured loop regions, L1 and

L2 (Fig. 2A) (*1*). To pinpoint which sequence domains of H2A.Z are responsible for changes

in transcriptional output across the set of orthologs tested above, we first examined sequence

conservation across the full H2A.Z dataset (Fig. S3), calculating sequence identity along the

length of H2A.Z using a 3-residue sliding window average (Fig. 2A). As expected, we observed

high conservation of the core alpha-helical bundle and relatively lower conservation of the tails

and the L1 (Fig. 2A). Surprisingly, however, we found that parts of the L2 region, specifically

around the core motif KDLKVK, were as highly conserved as the core α-helices (Fig. 2A).

This was especially striking when compared with the H2A variant H2A.X, which was notably

less conserved in this region.

The unexpectedly high sequence conservation of the L2 region suggested it might play a

role in determining the global changes in transcription we observed in the replacement *S.*

pombe lines. To test this, we engineered a series of chimeric histone genes whereby the N, C,

L1, L2, or docking domain (DD) (*40*) regions of *S. pombe pht1* were substituted into histone

H2A (*hta1*) (Fig. 2B, S5A). We expressed each of these chimeric genes by knocking them into

the *pht1* locus and assayed their ability to complement the *pht1*⁻ null allele by measuring the

growth of the chimeric lines under selected stress conditions based on prior experiments with

the representative full-length H2A.Z replacement lines (Fig. 1C). The chimeric histone genes

fell into two groups across all phenotypes tested: those that partially complemented the *pht1*⁻

mutant (those encoding H2A/Z^{DD}, H2A/Z^{N}, and H2A/Z^{L1}) and those that fully complemented

the *pht1*⁻ mutant in all tested conditions (those encoding H2A/Z^{L2} and H2A/Z^{C}), making them

indistinguishable from WT (Fig. 2C). These data indicate that both the L2 region and the C-

terminal tail play key roles in determining H2A.Z function.

To investigate whether sequence variation in the L2 region of H2A.Z alone is sufficient

to explain the transcriptional differences in our full-length H2A.Z replacement lines, we looked

for a correlation between transcriptional divergence and sequence divergence from WT *S.*

pombe in any of the structurally defined primary sequence regions of H2A.Z. Although the

short L2 affords limited resolution, we observed a strong correlation between L2 sequence

divergence and transcriptional divergence ($\rho = 0.63$, $p < 10^{-3}$, Fig. 2D). By contrast, the

sequence divergence of the C-terminal tail, which fully complemented the *pht1⁻* mutant in

growth assays, did not correlate with transcriptional divergence ($\rho = 0.056$, $p = 0.8$, Fig. 2D).

There was also a correlation between the sequence divergence of the N-terminal tail and

transcriptional divergence ($\rho = 0.5$, $p < 10^{-2}$; Fig. S5B). This was likely due to co-variation

with the L2 sequence, as there was a tight correlation between sequence variation in these two

regions ($\rho = 0.77$, $p < 10^{-5}$, Fig. S5B). Together, these data show that the L2 region of H2A.Z

is an important contributor to its function.

Transcription elongation factor Spt6 binds H2A.Z L2

The L2 region of H2A.Z is largely solvent accessible (*55*), particularly during

transcription when the DNA is unwrapped from the nucleosome (*19, 20*). We therefore

hypothesized that L2 might impact transcription by physically interacting with proteins

involved in transcription, rather than by influencing histone–DNA contacts, as is the case for

the L2 region of histone H3 (*1, 66*). To identify proteins in the *S. pombe* nuclear proteome that

might interact with the L2 region of Pht1, we used AlphaFold2 Multimer (*67*) to predict

physical interactions. We first performed a virtual interaction screen of WT Pht1 L2 against

the *S. pombe* annotated proteome (*68, 69*) (Fig. 3A), identifying 247 proteins with an interface

predicted template modelling (IPTM) score threshold of >0.5 (Table S2). To further refine

potential interactors, we reasoned that L2 interactors responsible for driving functional

variation in transcriptional output must vary in their interaction in accordance with their L2

primary sequence. We thus selected 17 unique single-residue variants of the L2 core motif from

our database of H2A.Z sequences (Fig. S3, Data File S1), performing an AlphaFold2 virtual

interaction screen with correspondingly mutated Pht1 L2 sequences against a subset of 500

nuclear proteins based on their IPTM score with WT Pht1 L2 (Fig. 3B, S6A, Table S3). To

assess differences in predicted binding across Pht1 mutants, we then calculated the variance in

IPTM score for each candidate interactor protein (Fig. 3C). In most cases, the variance was

very low, suggesting that the candidate interacting proteins do not respond to variation in L2

(Fig. 3C, S6A, Table S3). Among the candidate interactors with the highest variance, the

transcription elongation factor Spt6 (Fig. 3D–E, Table S3) was the only component of RNAPII

(*70, 71*). It interacts with nucleosomes (*72-74*), had highly robust folding models across

multiple simulations (Fig. S6B), and closely matched experimentally derived structures (Fig.

S6C).

We first experimentally validated the AlphaFold2-predicted interaction between Pht1

and Spt6, using glutathione-S-transferase (GST) as negative control. H2A was previously

reported to bind to Spt6 and was used as positive control (*75*). We expressed and purified

recombinant *S. pombe* GST-tagged Pht1 and Hta1, and His-tagged Spt6 in *Escherichia coli*

(Fig. 3F) and assessed binding *in vitro* to nickel resin (binds the His tag). Pht1 bound Spt6 (Fig.

3G), confirming the Pht1–Spt6 interaction. Further, we noted a qualitative difference in the

amount of Hta1 and Pht1 bound to Spt6, suggesting that the two histone proteins might interact

differently with Spt6. To verify this finding, we repeated Spt6-Pht1 pulldown experiments

under more stringent binding conditions, where the amount of bound Pht1 was considerably

less than that of Hta1 (Fig. S6D), reinforcing the suggestion Hta1 and Pht1 may interact with

Spt6 differently. Together, these data establish a physical interaction between Spt6 and Pht1.

To test whether the L2 region was sufficient to mediate the Pht1–Spt6 interaction, we

synthesized a biotinylated peptide corresponding to this region of *S. pombe* Pht1 and assayed

its binding to His-tagged Spt6 (Fig. 3H). As a negative control, we used a peptide

corresponding to the docking domain (DD) of *S. pombe* Pht1, which is adjacent to the L2 region

and important for the interaction of Pht1 with H3 and H4 (*55*) and Swr1-C (*76*). Whereas the DD peptide did not bind Spt6, the peptides corresponding to the L2 region of Hta1 or Pht1 bound Spt6 (Fig. 3I), consistent with the results of our experiments with the full-length proteins. These data establish that the L2 region alone is sufficient to explain the interaction of Pht1 with Spt6.

To test whether sequence variation in the L2 region of H2A.Z alters its binding to Spt6, we synthesized L2 peptides from five of the original set of representative H2A.Z sequences and assayed their binding to *S. pombe* Spt6, as above. Consistent with the sequence of L2 determining its interaction with Spt6, we observed variable amount of each peptide bound to Spt6 across species (Fig. 3I), with *A. godoyi*, *T. globosa*, and *P. marinus* H2A.Z L2 peptides having the greatest qualitative difference relative to *S. pombe* Pht1 L2. Although there were clear differences between peptides derived from different species, we did not observe a strong association between Spt6-binding and sequence similarity (Fig. 3I), suggesting that specific sequence features of the peptides, rather than overall sequence divergence, must govern their association with Spt6. Indeed, when we compared the sequence of the three species' L2 peptides that had discernable differences relative to Pht1 to those that did not, we noted that all of them had substitutions within the L2 core motif (Fig. S6F), especially at position L83.

To investigate the role of sequence variation in the core motif around L83 in determining the binding of Pht1 L2 to Spt6, we further examined AlphaFold2 models. Pht1 was predicted to bind to the Spt6 death-like domain (DLD) (Fig. 3D). This domain, whose function has remained uncharacterized, is the most highly conserved feature of the Spt6 surface (*77*), suggesting an important functional role. To assess this further, we gathered Spt6 sequences from the same EukProt database we used previously to identify H2A.Zs, calculating sequence conservation of the DLD across known eukaryotic diversity (Fig. 3J). In the DLD large numbers of hydrophobic residues were not only highly conserved in absolute terms (Fig.

3J), but also relative to other hydrophobic residues in Spt6 ($p < 0.005$, Wilcoxon test; Fig. S6F).

These residues were organized into a hydrophobic groove, many of which contacted the Pht1

L2 core motif in AlphaFold2 models (Fig. 3E, S6E). Further, these contacts were predicted to

vary according to the H2A.Z L2 sequence, both when examining models for L2s from different

species (Fig. S6E), as well as Pht1 mutants to the L83 position we identified earlier (Fig. 3E),

suggesting a rationale for why sequence variation at this Pht1 position might affect binding.

To test the contribution of Pht1 L2 sequence variation, particularly at position L83, to

the Pht1–Spt6 interaction in the context of the DLD hydrophobic groove, we synthesized three

mutated Pht1 L2 peptides with either more hydrophobic (L83F) or more hydrophilic (L83M,

L83N) substitutions and tested their binding to Spt6. The more hydrophilic L83N substitution,

which corresponds to Hta1 residue N75, does not recapitulate the binding of full-length Hta1

(Fig. 3D), reinforcing our earlier conclusion that numerous other primary sequences

differences in Hta1 (Fig. S5A) must contribute to its binding. Whereas L83F mutant L2 peptide

bound Spt6 similarly to WT Pht1 peptide, Spt6 bound both more hydrophilic substitutions

(L83M/N) to a much lesser extent than WT Pht1 L2 (Fig. 3I). These relative binding abilities

corresponded to their hydrophobicities, and are thus consistent with AlphaFold2 predictions

that the L2 region of Pht1 interacts with the hydrophobic groove of the DLD of Spt6 (Fig. 3D).

Together, these findings suggest that the hydrophobic L2 region of H2A.Z determines its

binding to the DLD of Spt6.

H2A.Z L2 encodes transcription elongation states

Since Spt6 is a transcription elongation factor (*70, 71*), we hypothesized that sequence

variation in the L2 region of H2A.Z might be sufficient to affect transcription elongation,

thereby explaining species-specific differences in reported H2A.Z function. To test this, we

first developed a growth-based reporter assay using the uracil biosynthesis gene *ura4*. In this

assay, the amount of transcription of *ura4* is directly related to the amount of growth in media

lacking uracil (Fig. 4A). Assaying a growth-based reporter has the advantage that it can be used

to test multiple conditions and it is not easily confounded by global transcriptional changes.

We first confirmed that our reporter gene accumulates Pht1 by ChIP-qPCR (Fig. S7A). We

then introduced point mutations corresponding to those found in the L2 core motif of our nine

representative H2A.Z orthologs (Fig. S4A, S6F) into the endogenous *pht1* (H2A.Z) gene in *S.*

pombe (Fig. 4B). Whereas *pht1⁻* cells gave rise to a modest increase in reporter expression (Fig.

4C), supporting a repressive function of H2A.Z in *S. pombe* as in some other model systems

(*11, 48, 50, 53*), half of L2 mutants had a significant increase (by ~1.3–1.5 fold for L83F, L83N

and L83Q) or decrease (by ~2 fold for L83M) in reporter expression. These results were not

confounded by general fitness defects in the mutant strains (Fig. S7B), and all the mutants were

epistatic to Swc2, an essential subunit of the Swr1-C chromatin remodeling complex, which

incorporates H2A.Z into nucleosomes (*78, 79*) (Fig. S7C). Together these results establish that

reporter expression differences are driven by Pht1 presence at the reporter gene and suggest

that L2 sequence variation is sufficient to impact transcription.

265   We next tested whether differences in reporter expression across Pht1 mutant lines were

specifically attributable to transcription elongation through two orthogonal approaches. First,

focusing on the four Pht1 mutant lines with the strongest differences in reporter expression

relative to WT (L83M, L83N, K86Q, and L83F), we made use of the drug mycophenolic acid

(MPA) to specifically interfere with RNAPII elongation by lowering cellular GTP levels (*80*).

In this experiment, if a mutant has increased or decreased RNAPII elongation relative to WT,

then it should be resistant or sensitive to MPA, respectively. Treatment of WT cells with MPA

caused a ~40% reduction in growth relative to the control (Fig. 4D). By contrast, *pht1⁻* and the

Pht1 mutants L83N, K86Q, and L83F were more resistant to MPA than the WT (~25–35%

reduction in growth; Fig. 4D), consistent with increased transcription elongation in these lines.

By contrast, the single mutant that had decreased reporter expression (L83M) was also the most

sensitive to MPA treatment (~60% reduction, $p_{adj} < 10^6$, Fig. 4D), confirming its reduction in

transcription elongation activity. Together, these data establish that Pht1 L2 mutants are

sufficient to impact elongating RNAPII.

Finally, as an orthogonal measure to our reporter-based assays, we probed the impact of

Pht1 L2 mutants on transcription elongation by measuring nascent transcription using precision

run-on sequencing (qPRO-seq) (*81, 82*). Sequencing nascent transcripts allowed us to position

RNAPII in the transcriptional unit and to distinguish between promoter-proximal paused

RNAPII and gene body-located RNAPII participating in active elongation across all expressed

genes, albeit without accurately quantifying specific parameters of RNAPII activity. Consistent

with results of our reporter assays, differential expression analysis of gene body-located qPRO-

seq counts in the WT and *pht1*[-] lines showed a modest increase in active elongation by RNAPII

at ~100 genes in *pht1*[-] cells (Fig. 4E). Similar analyses of the Pht1 point mutant lines identified

three classes of mutant. In one class, comprising half of the tested mutants (L83M, L83G,

A80S, and V85A), we observed essentially no change in RNAPII elongation, when compared

with WT at the level of differentially transcribed genes (Fig. 4E). Notably, this class included

the single super-repressive L2 mutant (L83M), suggesting it is likely linked to Spt6's function

as part of transcription termination (*70*). The second class comprised a single mutant (K84H),

which had a modest increase in RNAPII engaging in elongation, very similar to that of *pht1*[-]

(Fig. 4E). The third class comprised three mutants (L83N, K86Q, and L83F) in which we

observed a robust increase in RNAPII engaging in elongation at ~50% more genes than in *pht1*[-]

(Fig. 4E), consistent with their impact on reporter gene transcription. Together, these data

establish that sequence variation in the core L2 region of H2A.Z is sufficient to transform its

transcriptional impact on the genome.

For much of the past 50 years, investigations into variation in the core histone fold have

focused on its interaction with DNA and the stability of the nucleosome (*26, 41, 42, 83-87*).

Our findings, by contrast, highlight the importance of the histone core domain for genomic

regulation through protein–protein interactions (*66, 88, 89*). Given the extreme depth of

conservation of not only H2A.Z and Spt6, but also the entire chromatin and transcription

apparatus across eukaryotic evolution (*90*), it seems likely that the nucleosome core of

transcriptional regulation originated with eukaryogenesis and diversified with histone variants.

That we uncover sequence variation in just seven residues of the L2 core motif of H2A.Z can

give rise to enormous phenotypic complexity at the level of transcriptional output and beyond,

it is likely that many protein–protein interactions with the core domain of other histone variants

remain to be discovered. These will not only broaden our understanding of the importance of

the evolution of the nucleosome, but also, given the prevalence of transcription elongation

defects in cancers (*91, 92*), provide a new targets to address human disorders.

References and Notes

313   1.   K. Luger, A. W. Mader, R. K. Richmond, D. F. Sargent, T. J. Richmond, Crystal
314   structure of the nucleosome core particle at 2.8 A resolution. *Nature* **389**, 251-260
315   (1997).

316   2.   M. D. Meneghini, M. Wu, H. D. Madhani, Conserved histone variant H2A.Z protects
317   euchromatin from the ectopic spread of silent heterochromatin. *Cell* **112**, 725-736
318   (2003).

319   3.   C. D. Allis *et al.*, hv1 is an evolutionarily conserved H2A variant that is preferentially
320   associated with active genes. *Journal of Biological Chemistry* **261**, 1941-1948 (1986).

321   4.   R. Yelagandula *et al.*, The histone variant H2A.W defines heterochromatin and
322   promotes chromatin condensation in Arabidopsis. *Cell* **158**, 98-109 (2014).

323   5.   C. Costanzi, J. R. Pehrson, Histone macroH2A1 is concentrated in the inactive X
324   chromosome of female mammals. *Nature* **393**, 599-601 (1998).

325   6.   B. D. Giaimo, F. Ferrante, A. Herchenrother, S. B. Hake, T. Borggrefe, The histone
326   variant H2A.Z in gene regulation. *Epigenetics Chromatin* **12**, 37 (2019).

327   7.   V. Turinetto, C. Giachino, Multiple facets of histone variant H2AX: a DNA double-
328   strand-break marker with several biological functions. *Nucleic Acids Res* **43**, 2489-
329   2498 (2015).

330   8.   M. Perino, G. J. Veenstra, Chromatin Control of Developmental Dynamics and
331   Plasticity. *Dev Cell* **38**, 610-620 (2016).

332   9.   R. P. Ghosh, B. J. Meyer, Spatial Organization of Chromatin: Emergence of
333   Chromatin Structure During Development. *Annu Rev Cell Dev Biol* **37**, 199-232
334   (2021).

335   10.   A. M. Valencia, C. Kadoch, Chromatin regulatory mechanisms and therapeutic
336   opportunities in cancer. *Nat Cell Biol* **21**, 152-161 (2019).

337   11.   D. G. Berta *et al.*, Deficient H2A.Z deposition is associated with genesis of uterine
338   leiomyoma. *Nature*, (2021).

339   12.   R. S. Greenberg, H. K. Long, T. Swigut, J. Wysocka, Single Amino Acid Change
340   Underlies Distinct Roles of H2A.Z Subtypes in Human Syndrome. *Cell* **178**, 1421-
341   1436 e1424 (2019).

342   13.   A. Weiner *et al.*, High-resolution chromatin dynamics during a yeast stress response.
343   *Mol Cell* **58**, 371-386 (2015).

14. D. Tremethick, K. Zucker, A. Worcel, The transcription complex of the 5 S RNA
gene, but not transcription factor IIIA alone, prevents nucleosomal repression of
transcription. *J Biol Chem* **265**, 5014-5023 (1990).

15. G. Almouzni, M. Mechali, A. P. Wolffe, Transcription complex disruption caused by
a transition in chromatin structure. *Mol Cell Biol* **11**, 655-665 (1991).

16. J. N. Hirschhorn, S. A. Brown, C. D. Clark, F. Winston, Evidence that SNF2/SWI2
and SNF5 activate transcription in yeast by altering chromatin structure. *Genes Dev* **6**,
2288-2298 (1992).

17. O. I. Kulaeva, F. K. Hsieh, H. W. Chang, D. S. Luse, V. M. Studitsky, Mechanism of
transcription through a nucleosome by RNA polymerase II. *Biochim Biophys Acta*
1829, 76-83 (2013).

18. B. R. Cairns, The logic of chromatin architecture and remodelling at promoters.
Nature **461**, 193-198 (2009).

19. H. Ehara, T. Kujirai, M. Shirouzu, H. Kurumizaka, S. I. Sekine, Structural basis of
nucleosome disassembly and reassembly by RNAPII elongation complex with FACT.
Science **377**, eabp9466 (2022).

20. T. Kujirai *et al.*, Structural basis of the nucleosome transition during RNA polymerase
II passage. *Science* **362**, 595-598 (2018).

21. B. Wasylyk, G. Thevenin, P. Oudet, P. Chambon, Transcription of in vitro assembled
chromatin by Escherichia coli RNA polymerase. *J Mol Biol* **128**, 411-440 (1979).

22. J. A. Knezetic, D. S. Luse, The presence of nucleosomes on a DNA template prevents
initiation by RNA polymerase II in vitro. *Cell* **45**, 95-104 (1986).

23. Y. Lorch, J. W. LaPointe, R. D. Kornberg, Nucleosomes inhibit the initiation of
transcription but allow chain elongation with the displacement of histones. *Cell* **49**,
203-210 (1987).

24. P. B. Talbert, M. P. Meers, S. Henikoff, Old cogs, new tricks: the evolution of gene
expression in a chromatin context. *Nat Rev Genet* **20**, 283-297 (2019).

25. S. Henikoff, M. M. Smith, Histone variants and epigenetics. *Cold Spring Harb*
Perspect Biol **7**, a019364 (2015).

26. C. M. Weber, S. Ramachandran, S. Henikoff, Nucleosomes are context-specific,
H2A.Z-modulated barriers to RNA polymerase. *Mol Cell* **53**, 819-830 (2014).

27. P. B. Talbert, S. Henikoff, Histone variants--ancient wrap artists of the epigenome.
Nat Rev Mol Cell Biol **11**, 264-275 (2010).

28. S. B. Hake, C. D. Allis, Histone H3 variants and their potential role in indexing
mammalian genomes: the "H3 barcode hypothesis". *Proc Natl Acad Sci U S A* **103**,
6428-6435 (2006).

29. D. Jiang, F. Berger, Variation is important: Warranting chromatin function and
dynamics by histone variants. *Curr Opin Plant Biol* **75**, 102408 (2023).

30. D. Jiang *et al.*, The evolution and functional divergence of the histone H2B family in
plants. *PLoS Genet* **16**, e1008964 (2020).

31. A. Herchenrother, T. M. Wunderlich, J. Lan, S. B. Hake, Spotlight on histone H2A
variants: From B to X to Z. *Semin Cell Dev Biol* **135**, 3-12 (2023).

32. H. Zhang, D. N. Roberts, B. R. Cairns, Genome-wide dynamics of Htz1, a histone
H2A variant that poises repressed/basal promoters for activation through histone loss.
Cell **123**, 219-231 (2005).

33. B. Guillemette *et al.*, Variant histone H2A.Z is globally localized to the promoters of
inactive yeast genes and regulates nucleosome positioning. *PLoS Biol* **3**, e384 (2005).

34. R. M. Raisner *et al.*, Histone variant H2A.Z marks the 5' ends of both active and
inactive genes in euchromatin. *Cell* **123**, 233-248 (2005).

35. A. Barski *et al.*, High-resolution profiling of histone methylations in the human
genome. *Cell* **129**, 823-837 (2007).

36. D. Zilberman, D. Coleman-Derr, T. Ballinger, S. Henikoff, Histone H2A.Z and DNA
methylation are mutually antagonistic chromatin marks. *Nature* **456**, 125-129 (2008).

37. T. N. Mavrich *et al.*, Nucleosome organization in the Drosophila genome. *Nature*
453, 358-362 (2008).

38. C. D. Allis, C. V. Glover, J. K. Bowen, M. A. Gorovsky, Histone variants specific to
the transcriptionally active, amitotically dividing macronucleus of the unicellular
eucaryote, Tetrahymena thermophila. *Cell* **20**, 609-617 (1980).

39. M. H. West, W. M. Bonner, Histone 2A, a heteromorphous family of eight protein
species. *Biochemistry* **19**, 3238-3245 (1980).

40. H. S. Malik, S. Henikoff, Phylogenomics of the nucleosome. *Nat Struct Biol* **10**, 882-
891 (2003).

41. A. Osakabe *et al.*, Histone H2A variants confer specific properties to nucleosomes
and impact on chromatin accessibility. *Nucleic Acids Res* **46**, 7675-7685 (2018).

42. N. Horikoshi, T. Kujirai, K. Sato, H. Kimura, H. Kurumizaka, Structure-based design
of an H2A.Z.1 mutant stabilizing a nucleosome in vitro and in vivo. *Biochem Biophys*
Res Commun **515**, 719-724 (2019).

43. Z. Wen, L. Zhang, H. Ruan, G. Li, Histone variant H2A.Z regulates nucleosome
unwrapping and CTCF binding in mouse ES cells. *Nucleic Acids Res* **48**, 5939-5952
(2020).

44. M. Adam, F. Robert, M. Larochelle, L. Gaudreau, H2A.Z is required for global
chromatin integrity and for recruitment of RNA polymerase II under specific
conditions. *Mol Cell Biol* **21**, 6270-6279 (2001).

45. M. S. Santisteban, T. Kalashnikova, M. M. Smith, Histone H2A.Z Regulates
Transcription and Is Partially Redundant with Nucleosome Remodeling Complexes.
Cell **103**, 411-422 (2000).

46. M. S. Santisteban, M. Hang, M. M. Smith, Histone variant H2A.Z and RNA
polymerase II transcription elongation. *Mol Cell Biol* **31**, 1848-1860 (2011).

47. L. A. Stargell *et al.*, Temporal and spatial association of histone H2A variant hv1 with
transcriptionally competent chromatin during nuclear development in Tetrahymena
thermophila. *Genes Dev* **7**, 2641-2651 (1993).

48. C. Mylonas, C. Lee, A. L. Auld, Cisse, II, L. A. Boyer, A dual role for H2A.Z.1 in
modulating the dynamics of RNA polymerase II initiation and elongation. *Nat Struct*
Mol Biol **28**, 435-442 (2021).

49. C. D. Allis, Y. S. Ziegler, M. A. Gorovsky, J. B. Olmsted, A conserved histone
variant enriched in nucleoli of mammalian cells. *Cell* **31**, 131-136 (1982).

50. D. Coleman-Derr, D. Zilberman, Deposition of histone variant H2A.Z within gene
bodies regulates responsive genes. *PLoS Genet* **8**, e1002988 (2012).

51. B. Carter *et al.*, The Chromatin Remodelers PKL and PIE1 Act in an Epigenetic
Pathway That Determines H3K27me3 Homeostasis in Arabidopsis. *Plant Cell* **30**,
1337-1352 (2018).

52. B. Jamge *et al.*, Histone variants shape chromatin states in Arabidopsis. *Elife* **12**,
(2023).

53. A. Thakar, P. Gupta, W. T. McAllister, J. Zlatanova, Histone variant H2A.Z inhibits
transcription in reconstituted nucleosomes. *Biochemistry* **49**, 4018-4026 (2010).

54. K. Ahmad, S. Henikoff, The histone variant H3.3 marks active chromatin by
replication-independent nucleosome assembly. *Mol Cell* **9**, 1191-1200 (2002).

55. R. K. Suto, M. J. Clarkson, D. J. Tremethick, K. Luger, Crystal structure of a
nucleosome core particle containing the variant histone H2A.Z. *Nat Struct Biol* **7**,
1121-1124 (2000).

56. H. Tachiwana *et al.*, Structures of human nucleosomes containing major histone H3
variants. *Acta Crystallogr D Biol Crystallogr* **67**, 578-583 (2011).

57. P. B. Talbert *et al.*, A unified phylogeny-based nomenclature for histone variants.
Epigenetics Chromatin **5**, 7 (2012).

58. M. C. Dalmasso, W. J. Sullivan, Jr., S. O. Angel, Canonical and variant histones of
protozoan parasites. *Front Biosci (Landmark Ed)* **16**, 2086-2105 (2011).

59. J. Heitman, N. R. Movva, M. N. Hall, Targets for cell cycle arrest by the
immunosuppressant rapamycin in yeast. *Science* **253**, 905-909 (1991).

60. Y. O. Chernoff, S. L. Lindquist, B. Ono, S. G. Inge-Vechtomov, S. W. Liebman, Role
of the chaperone protein Hsp104 in propagation of the yeast prion-like factor [psi+].
Science **268**, 880-884 (1995).

61. P. D. Crowley, H. C. Gallagher, Clotrimazole as a pharmaceutical: past, present and
future. *J Appl Microbiol* **117**, 611-617 (2014).

62. W. Samakkarn, K. Ratanakhanokchai, N. Soontorngun, Reprogramming of the
Ethanol Stress Response in Saccharomyces cerevisiae by the Transcription Factor
Znf1 and Its Effect on the Biosynthesis of Glycerol and Ethanol. *Appl Environ
Microbiol* **87**, e0058821 (2021).

63. F. Yang, C. J. Kemp, S. Henikoff, Doxorubicin enhances nucleosome turnover around
promoters. *Curr Biol* **23**, 782-787 (2013).

64. R. Hoyos-Manchado *et al.*, RNA metabolism is the primary target of formamide in
vivo. *Scientific reports* **7**, 15895 (2017).

65. S. Torres-Garcia *et al.*, Epigenetic gene silencing by heterochromatin primes fungal
resistance. *Nature* **585**, 453-458 (2020).

66. K. M. Stevens *et al.*, Histone variants in archaea and the evolution of combinatorial
chromatin complexity. *Proceedings of the National Academy of Sciences* **117**, 33384-
33395 (2020).

67. J. Jumper *et al.*, Highly accurate protein structure prediction with AlphaFold. *Nature*
596, 583-589 (2021).

68. M. A. Harris *et al.*, Fission stories: using PomBase to understand
Schizosaccharomyces pombe biology. *Genetics* **220**, (2022).

69. C. The Gene Ontology, The Gene Ontology Resource: 20 years and still GOing
strong. *Nucleic Acids Res* **47**, D330-D338 (2019).

70. A. Narain *et al.*, Targeted protein degradation reveals a direct role of SPT6 in RNAPII
elongation and termination. *Mol Cell* **81**, 3110-3127 e3114 (2021).

71. K. Zumer *et al.*, Two distinct mechanisms of RNA polymerase II elongation
stimulation in vivo. *Mol Cell* **81**, 3096-3109 e3098 (2021).

72. A. Bortvin, F. Winston, Evidence that Spt6p controls chromatin structure by a direct
interaction with histones. *Science* **272**, 1473-1476 (1996).

73. S. M. McDonald, D. Close, H. Xin, T. Formosa, C. P. Hill, Structure and biological
importance of the Spn1-Spt6 interaction, and its regulatory role in nucleosome
binding. *Mol Cell* **40**, 725-735 (2010).

74. M. W. Adkins, J. K. Tyler, Transcriptional activators are dispensable for transcription
in the absence of Spt6-mediated chromatin reassembly of promoter regions.
Molecular cell **21**, 405-416 (2006).

75. C. Jeronimo, S. Watanabe, C. D. Kaplan, C. L. Peterson, F. Robert, The Histone
Chaperones FACT and Spt6 Restrict H2A.Z from Intragenic Locations. *Mol Cell* **58**,
1113-1123 (2015).

76. J. Hong *et al.*, The catalytic subunit of the SWR1 remodeler is a histone chaperone for
the H2A.Z-H2B dimer. *Mol Cell* **53**, 498-505 (2014).

77. D. Close *et al.*, Crystal structures of the S. cerevisiae Spt6 core and C-terminal
tandem SH2 domain. *J Mol Biol* **408**, 697-713 (2011).

78. W. H. Wu *et al.*, Swc2 is a widely conserved H2AZ-binding module essential for
ATP-dependent histone exchange. *Nat Struct Mol Biol* **12**, 1064-1071 (2005).

79. E. Luk *et al.*, Stepwise histone replacement by SWR1 requires dual activation with
histone H2A.Z and canonical nucleosome. *Cell* **143**, 725-736 (2010).

80. R. J. Shaw, J. L. Wilson, K. T. Smith, D. Reines, Regulation of an IMP
dehydrogenase gene and its overexpression in drug-sensitive transcription elongation
mutants of yeast. *J Biol Chem* **276**, 32905-32916 (2001).

81. G. T. Booth, I. X. Wang, V. G. Cheung, J. T. Lis, Divergence of a conserved
elongation factor and transcription regulation in budding and fission yeast. *Genome*
Res **26**, 799-811 (2016).

82. J. Judd *et al.*, A rapid, sensitive, scalable method for Precision Run-On sequencing
(PRO-seq). *bioRxiv*, (2020).

83. C. Bonisch, S. B. Hake, Histone H2A variants in nucleosomes and chromatin: more
or less stable? *Nucleic Acids Res* **40**, 10719-10741 (2012).

84. S. Chakravarthy, Y. Bao, V. A. Roberts, D. Tremethick, K. Luger, Structural
characterization of histone H2A variants. *Cold Spring Harb Symp Quant Biol* **69**,
227-234 (2004).

85. Y. Arimura *et al.*, Structural basis of a nucleosome containing histone
H2A.B/H2A.Bbd that transiently associates with reorganized chromatin. *Scientific*
reports **3**, 3510 (2013).

86. N. Horikoshi, Y. Arimura, H. Taguchi, H. Kurumizaka, Crystal structures of
heterotypic nucleosomes containing histones H2A.Z and H2A. *Open Biol* **6**, (2016).

87. H. Kohestani, J. Wereszczynski, Effects of H2A.B incorporation on nucleosome
structures and dynamics. *Biophys J* **120**, 1498-1509 (2021).

88. A. K. Michael *et al.*, Cooperation between bHLH transcription factors and histones
for DNA access. *Nature* **619**, 385-393 (2023).

89. S. Sanulli *et al.*, HP1 reshapes nucleosome core to promote phase separation of
heterochromatin. *Nature* **575**, 390-394 (2019).

90. X. Grau-Bove *et al.*, A phylogenetic and proteomic reconstruction of eukaryotic
chromatin evolution. *Nat Ecol Evol* **6**, 1007-1023 (2022).

91. V. Modur *et al.*, Defective transcription elongation in a subset of cancers confers
immunotherapy resistance. *Nat Commun* **9**, 4410 (2018).

92. T. E. Miller *et al.*, Transcription elongation factors represent in vivo cancer
dependencies in glioblastoma. *Nature* **547**, 355-359 (2017).
529

Acknowledgements: We thank the entire Berger group and Antoine Hocher, for their

insightful, considerate, and helpful discussions, as well as Carol Featherstone from Life

Science Editors and Matt Watson for editorial assistance during the preparation of the

manuscript. We especially thank Zdravko Lorković, Elin Axelsson-Ekker, and Chung Hyun

Cho for their invaluable assistance preparing antibodies, biochemical assays, and

bioinformatic analyses. We also thank John T. Lis for advice with qPRO-seq. We thank the

Molecular Biology Service and Media Kitchen for a constant supply of plates, basic reagents,

and cloning and sequencing services, as well as the Peptide Synthesis Service. Additionally,

we thank the Vienna BioCenter Core Facilities, in particular the Next Generation Sequencing

facility for their advice and swift handling of all our requests and the Protein Technologies

facility for their expertise, helpful discussions, and recombinant protein production.

Funding:

Gregor Mendel Institute, Austrian Academy of Sciences Core Funding (FB)

Österreichischer Wissenschaftsfonds FWF TAI304 (FB)

Österreichischer Wissenschaftsfonds FWF ESP213 (ZHH)

EMBO Postdoctoral Fellowship ALTF169-2020 (ZHH)

United Kingdom Research and Innovation's Medical Research Council MC_A658_5TY40

(TW)

Author contributions: Conceptualization: ZHH, FB, and TW; Methodology: ZHH, KS, and

TW; Validation: ZHH and KS; Formal Analysis: ZHH, KS, and TW; Investigation: ZHH,

KS, and TW; Data Curation: ZHH, KS, and TW; Writing – original draft: ZHH, TW, and FB;

Visualization: ZHH, KS, and TW; Supervision: FB, TW; Project Administration: ZHH;

Funding Acquisition: ZHH, FB, and TW.

Competing interests: Z.H.H. and F.B. are inventors on a patent submitted by the Gregor

Mendel Institute at the Austrian Patent Office (A50392/2024) pertaining to the modification

of transcription and fitness by engineering H2A.Z L2.

Data and materials availability: PRO-seq pipeline and scripts used for data analysis are

available as a container on Github, and all other scripts used in analysis are available upon

request to the lead contact. NGS data (PRO-seq, mRNA-seq) are deposited in GEO under

accession number GSE265749 and GSE265775, respectively. For material request (e.g.

strains), please contact the lead (zachary.harvey@gmi.oeaw.ac.at).

Supplementary Materials

Materials and Methods

Figs. S1 to S7

Supplementary Data Tables S1 to S4

Supplementary Data File S1 to S3

References ()

FIGURES AND FIGURE LEGENDS

Figure 1. Phylogenetically diverse H2A.Z sequences drive distinct gain-of-function

transcriptional states. A, Experimental scheme describing the replacement of endogenous *S.*

pombe pht1 by representative H2A.Zs taken from the H2A phylogeny (Fig. S2B, S3A). **B**,

Principal component analysis (PCA) for the top 500 variable genes from mRNA-seq data

generated for H2A.Z replacement lines. **C**, Growth phenotypes for H2A.Z replacement lines

in various stress conditions, annotated with the sequence distance of each H2A.Z from that of

S. pombe Pht1 (which has a sequence distance of zero). Plotted values are the mean of four

biological replicates. Growth is quantified as the area under the curve (AUC) from respective

growth curves. H2A.Zs are annotated with their sequence distance relative to *S. pombe* Pht1,

with zero being Pht1 and larger numbers indicating greater divergence. **D**, Example

Spearman's correlations between the growth from (C) with either sequence distance (as in C)

of each H2A.Z relative to WT *S. pombe*, or the expression of the respective H2A.Z (from

mRNA-seq).

Figure 2. The L2 region of H2A.Z is a key determinant of its function. A, Mean pairwise

sequence identity of H2A.Z and H2A.X residues across eukaryotes, averaged using a 3-amino

acid sliding window. The L2 region is highlighted in grey on the histone secondary structure,

with the core domain we identify indicated. Dashed lines indicate the N- and C-terminal tails

that were not included in this analysis due to variable length and resulting gappy alignment. **B**,

Experimental scheme for generation of Hta1/Pht1 chimeras. **C**, Mean growth (AUC) for four

biological replicates in each indicated condition, which are as in Fig. 1B. **D**, Spearman's

correlation analysis between the sequence divergence from *S. pombe* Pht1 of either the L2 or

C-terminal tail region and principal component 1 from mRNA-seq analysis from Fig. 1B. In

this analysis, the *S. pombe* Pht1 sequence corresponds to a value of zero, with a larger number

corresponding to a greater divergence. See also Fig. S5B.

Figure 3. The L2 region of H2A.Z binds to transcription elongation factor Spt6. A, Scheme explaining the AlphaFold2 virtual interaction screen against *S. pombe* Pht1. **B,** Scheme explaining the AlphaFold2 approach used to calculate the IPTM variance for 17 Pht1 L2 variants, which is plotted in **C**. The reduced nuclear proteome was defined by the top 500 WT *S. pombe* H2A.Z interactors from (A). **D,** AlphaFold2 predicted structure of the Pht1 L2 interaction with Spt6, with the the death-like-domain (DLD) highlighted with shading. **E,** Example AlphaFold2 structures of *S. pombe* Pht1, Hta1, and Pht1 mutants. Spt6 DLD is colored with a lipophilicity surface, and each H2A position 83 (e.g. L83 for Pht1) is labelled to highlight its interaction with the Spt6 hydrophobic groove. **F,** Experimental scheme for *in vitro* interaction assays of full-length GST-Pht1, GST-Hta1, with 14xHis-Spt6(282-1159). **G,** Coomassie-stained SDS-PAGE gel of Ni-NTA-pulldowns of recombinant *S. pombe* GST-Pht1 or GST-Hta1 with *S. pombe* His-Spt6(282-1159). Detection done with colloidal Coomassie staining. **H,** Experimental scheme showing the pulldown experiment modified for biotinylated peptides. **I,** Binding assay of Pht1-derived peptides to *S. pombe* 14xHis-Spt6(282-1159) immobilized on Ni-NTA resin. Detection was done either by colloidal Coomassie staining (for 14xHis-Spt6 loading control) or by a streptavidin-HRP conjugate (for biotinylated peptides). Representative of five experiments, each with similar results. **J,** Conservation analysis of Spt6 DLD. Conservation is calculated as the mean pairwise amino acid identity across Spt6 orthologs (see Methods). The location of hydrophobic residues are highlighted in red, with those predicted to interact with Pht1 L2 further indicated by an asterisk (*). The following residues are considered hydrophobic for the purpose of this analysis: A, V, I, L, F, M, W, Y.

Figure 4. H2A.Z L2 sequence variation explains species-specific effects on transcription.

A, Experimental scheme explaining the *ura4$^+$* reporter gene. **B,** Experimental scheme describing the generation of H2A.Z point mutants. **C,** Growth of H2A.Z point mutant strains bearing the transcriptional reporter from (D) in EMM-Uracil. The mean of WT is represented with a dashed line. Each boxplot represents 12 biological replicates, adjusted *p*-values (p_{adj}) are calculated using a two-way ANOVA followed by a post-hoc Tukey test. **D,** Growth of indicated *S. pombe* Pht1 mutants in the presence of 12.5 µM mycophenolic acid (MPA) expressed as the ratio of their paired growth in the absence of the drug. Data are 9 biological replicates and the experiment was repeated three times with similar results. Boxplots represent the median bounded by quartiles. Adjusted *p*-values (p_{adj}) are calculated using a two-way ANOVA followed by a post-hoc Tukey test. **E,** Differential count analysis for gene body qPRO-seq signal calculated using DESeq2. Red line, diagonal (Y=X) line drawn to aid in interpretation. Statistically significant ($p_{adj}<0.1$) changes are indicated in blue.