A novel expectation-maximization approach to infer general diploid selection from time-series genetic data

Adam G. Fine[1,2] and Matthias Steinrücken[1,3]

[1]Department of Ecology and Evolution, University of Chicago
[2]Graduate Program in Biophysical Sciences, University of Chicago
[3]Department of Human Genetics, University of Chicago

May 10, 2024

Abstract

Detecting and quantifying the strength of selection is a main objective in population genetics. Since selection acts over multiple generations, many approaches have been developed to detect and quantify selection using genetic data sampled at multiple points in time. Such time series genetic data is commonly analyzed using Hidden Markov Models, but in most cases, under the assumption of additive selection. However, many examples of genetic variation exhibiting non-additive mechanisms exist, making it critical to develop methods that can characterize selection in more general scenarios. Thus, we extend a previously introduced expectation-maximization algorithm for the inference of additive selection coefficients to the case of general diploid selection, in which heterozygote and homozygote fitnesses are parameterized independently. We furthermore introduce a framework to identify bespoke modes of diploid selection from given data, as well as a procedure for aggregating data across linked loci to increase power and robustness. Using extensive simulation studies, we find that our method accurately and efficiently estimates selection coefficients for different modes of diploid selection across a wide range of scenarios; however, power to classify the mode of selection is low unless selection is very strong. We apply our method to ancient DNA samples from Great Britain in the last 4,450 years, and detect evidence for selection in six genomic regions, including the well-characterized LCT locus. Our work is the first genome-wide scan characterizing signals of general diploid selection.

1 Introduction

Genetic variation that confers a fitness advantage to an organism over its peers increases in frequency in the population over time until eventual fixation, and this process of selection ultimately forms the basis of adaptation. Detecting evidence of this adaptive process and quantifying its strength is a fundamental problem in evolutionary biology, with applications ranging from finding mutations critical to early hominid evolution (Bustamante et al. 2005) to predicting tumor growth (Bignell et al. 2010). In population genetics, many methods to detect signatures of past adaptive events in contemporary population genomic data have thus been developed (Nielsen 2005; Vitti et al. 2013; Lachance and Tishkoff 2013).

However, since selection acts over multiple generations, genetic data observed at several time-points throughout allows for more accurate quantification of selective processes than present-day

samples alone. Recent technological advances have enabled researchers to collect such time series genetic data genome-wide. One main source of time series genetic data is ancient DNA (aDNA), that is, genetic material extracted from deceased individuals in humans or other species (Orlando et al. 2021). Next-generation sequencing has enabled collecting genetic data from large numbers of ancient samples, particularly through the development of techniques such as hybridization enrichment (Hofreiter et al. 2015). Another major source of temporal genetic data are experimental evolution studies (Barghi et al. 2019). Contemporary experimental evolution studies use next-generation sequencing technologies on several biological replicates in evolve and resequence (E&R) experiments to obtain high-quality estimates of temporal allele frequency changes at many loci throughout the genome (Schlötterer et al. 2015). These datasets present unprecedented opportunities to detect and characterize the adaptive processes that shape genomic variation (Malaspinas 2016; Dehasque et al. 2020).

Observing the true underlying population allele frequency trajectory as it changes over time would allow for highly accurate characterization of the underlying selective processes. However, in data obtained in practice, genetic variation is often only assessed for a set of individuals sampled at a finite number of time points. Quantifying the strength of selection therefore involves modeling the action of selection, genetic drift, and other population genetic processes on the unobserved trajectory of the population allele frequency, and considering sampled data as imprecise observations of this underlying trajectory.

A commonly used framework for analysis of time series data that readily accommodates this uncertainty are Hidden Markov Models (HMMs, Bollback et al. 2008). In these HMMs, the underlying population allele frequency evolves according to a Markov process, the Wright–Fisher model, and samples are modeled as binomial observations given the underlying population allele frequency. This HMM framework has been used to estimate a variety of parameters: The additive selection coefficient s (Bollback et al. 2008), the time at which a beneficial mutation arose (Malaspinas et al. 2012), or the effective population size N_e (Wang 2001). Within this HMM framework, Mathieson and McVean (2013) introduced an expectation-maximization (EM) approach, which can be used to estimate additive selection coefficients, as well as migration rates between sub-populations. For reviews of HMM-based approaches to estimate selection coefficients and comparisons between methods in aDNA and E&R analyses, see Tataru et al. (2017) and Vlachos et al. (2019), respectively.

Most implementations of the aforementioned HMM approach presented in the literature to date are designed to only detect additive selection, but many examples of genetic variation exhibiting non-additive mechanisms exist. In humans, non-additive targets of selection range from the classical case of the heterozygote advantage conferred by one copy of the sickle-cell allele (Gemmell and Slate 2006; Hedrick 2012) to recent evidence of pervasive dominance in an analysis of data from the UK Biobank (Palmer et al. 2023). Additionally, stabilizing selection on complex traits, which is believed to be wide-spread in humans (Sanjak et al. 2018), manifests as underdominant selective dynamics at the loci effecting the trait (Barton 1986; de Vladar and Barton 2014; Simons et al. 2018).

Here, we thus extend the EM approach of Mathieson and McVean (2013) to estimate selection coefficients under a general diploid model, that is, when the fitness values of homozygous and heterozygous genotypes are independently parameterized. We furthermore develop a novel framework for identifying the best-fitting mode of selection for a given temporal dataset. While some methods to estimate general diploid selection have been presented in the literature (Steinrücken et al. 2014; Foll et al. 2015; Schraiber et al. 2016; Iranmehr et al. 2017; Taus et al. 2017; Cheng and Steinrücken 2023), they do not explicitly address the statistical problem of using the data to distinguish between

different modes of selection. Furthermore, none of these methods have been applied to genome-wide data from human populations. To our knowledge, our analysis is the first that characterizes recent general diploid selection in humans from ancient DNA data on a full-genome scale.

The remainder of this article is organized as follows. In Section 2, we outline our iterative EM algorithm for efficiently estimating general diploid selection coefficients, as well as the statistical procedure for inferring the most likely mode of diploid selection. In Section 3, we apply our algorithm and inference framework in a wide range of simulated scenarios to assess its accuracy. We find that our method is generally well-powered to detect selection and estimate its strength, however, power to classify the mode of selection is limited. In Section 4, we then perform a genome-wide scan for recent diploid selection in the human genome, using publicly available ancient DNA data from individuals that lived in Great Britain in the last 4,450 years (Mallick et al. 2024, version 54.1) and discuss six genomic regions that show signals of recent selection. Lastly, we discuss future directions in Section 5. Our method `EMSel` (EM algorithm for detecting Selection) and the scripts to generate the figures in this manuscript are available online at `https://github.com/steinrue/EMSel`.

2 Methods

2.1 Parameterizing general diploid selection

Throughout this article, we consider selection acting at a given biallelic locus in a diploid population of constant size N_e. The dynamics of the population allele frequency at this locus can be described by the discrete Wright–Fisher model, where we denote by A and a the two alleles at the locus, and by $Y_t \in [0,1]$ and $1 - Y_t$ the random population-level frequency of the A and a allele in generation $t \in \{1, \dots T\}$, respectively. Suppose that the relative fitness of individuals with genotypes AA, Aa, and aa is $1 + s_2$, $1 + s_1$, and 1, respectively. Given these fitness values, if the frequency of A alleles in the current generation is $Y_t = p$, then the allele frequency in the next generation Y_{t+1} is given by $\frac{1}{2N_e}$ times a binomially distributed random number with $2N_e$ draws and success probability $p' := p + p(1 - p)\big(s_1(1 - 2p) + s_2 p\big)$ for small s_1 and s_2. This can further be approximated by a normal distribution, where $Y_{t+1} \sim \mathcal{N}\big(p', \frac{1}{2N_e} p(1 - p)\big)$, which is commonly referred to as the Wright–Fisher diffusion (Ewens 2004, Chapter 5.3).

We use the term *general diploid selection* for the case of arbitrary $s_1 > -1$ and $s_2 > -1$, that is, the fitness values for the homozygotes and heterozygtes are uncoupled. Many bespoke modes of selection correspond to constrained diploid selection, where the possible values of s_1 and s_2 are restricted. We consider the following modes of selection:

- **Additive**: $s_1 = \frac{1}{2}s_2$. The fitness is proportional to the number of copies of the A allele. Also referred to as **haploid** or **genic** selection.

- **Dominant**: $s_1 = s_2$. Any number of copies of the A allele confers the full fitness effect.

- **Recessive**: $s_1 = 0$. Only individuals homozygous for the A allele have a fitness effect.

- **Overdominance**: $s_1 > \max\{0, s_2\}$. Heterozygous individuals have the highest fitness.

- **Underdominance**: $s_1 < \max\{0, s_2\}$. Heterozygous individuals have the lowest fitness.

Additive, dominant, recessive selection are one-parameter modes, with $s := s_2 = 2s_1$, $s := s_1 = s_2$, and $s := s_2$ $(s_1 = 0)$, respectively. *Over-* and *underdominance* are two-dimensional sub-spaces, but

Figure 1: Two-dimensional space of general diploid selection and the five bespoke selection modes we consider: *Additive*, *dominant*, *recessive*, *over-* and *underdominance*. Colors indicate the sub-space of the respective mode. 50 replicates simulated under each mode, as well as their mean, are plotted to illustrate the characteristic shapes of the trajectories in each mode.

we frequently use the version $s := s_1$ $(s_2 = 0)$ as a one-parameter selection mode that describes complete *over-* or *underdominance*. We also refer to the combination of the latter two as *heterozygote difference*. While the one-parameter modes for complete *over-* and *underdominance* are more convenient for simulations, combining them into the one-dimensional mode *heterozygote difference* when inferring parameters avoids statistical artifacts. In Figure 1, we show a diagram of the different modes of selection as sub-spaces of the full two-dimensional parameter space of general diploid selection.

2.2 Hidden Markov Model for inferring diploid selection coefficients

2.2.1 Derivation of the EM-HMM algorithm

To derive our novel method for inferring general diploid selection coefficients from sampled time series genetic data we extend the method developed by Mathieson and McVean (2013) for *additive* selection. If the exact frequency trajectory of the focal allele $p_1, \ldots, p_T \in [0, 1]$ over T generations is known, then the normal approximation to the Wright-Fisher process can be used to define the likelihood of this trajectory as

$$L_{s_1,s_2}^{(C)}(p_1, \ldots, p_T) := \mathbb{P}_{s_1,s_2}\{Y_t \in dp_t \, \forall \, 1 \le t' \le T\}. \tag{1}$$

Here, we indicate the parameters of interest, the selection coefficients s_1 and s_2, in the subscript, the superscript (C) indicates the model where the population allele frequency in each generation takes values in the continuous range $[0, 1]$. Without loss of generality, the focal allele is the A allele. Maximizing this likelihood yields a maximum likelihood estimator (MLE) of the selection coefficients, and in the case of *additive* selection, this estimator has been presented by Watterson (1982). In Sections S.1.1 and S.1.2 in the Supplementary Material, we revisit Watterson's formula for the *additive* MLE and extend the result to general diploid selection, which yields the estimators

$$\widehat{s_1} = \frac{(p_T - p_1) \sum_t p_t^3 q_t - \sum_t p_t(p_{t+1} - p_t) \sum_t p_t^2 q_t}{\sum_t p_t q_t \sum_t p_t^3 q_t - \left(\sum_t p_t^2 q_t\right)^2}, \text{ and}$$

$$\widehat{s_2} = \frac{\sum_t p_t q_t(1 - 2p_t) \sum_t p_t(p_{t+1} - p_t) - (p_T - p_1) \sum_t p_t^2 q_t(1 - 2p_t)}{\sum_t p_t q_t \sum_t p_t^3 q_t - \left(\sum_t p_t^2 q_t\right)^2},$$

with $q_t := 1 - p_t$.

However, as discussed in Section 1, the data often consists of a certain number of individuals sampled from the population at certain points in time, and thus the full trajectory of the population allele frequency is in general not known. To efficiently integrate over this uncertainty, Bollback et al. (2008) introduced an HMM framework, where the population allele frequency at a given time is the hidden state, evolving according to the Wright-Fisher model, and the sampled genotypes are the observations. Specifically, assume that we have sampled the population at times $1 \leq t_1, \ldots, t_K \leq T$. At each timepoint, the data $o_{t_i} := (n_{t_i}, a_{t_i})$ consists of the number of haplotypes sampled at this time n_{t_i}, and the number of observed focal alleles a_{t_i}. For convenience, we set $n_t := 0$ and $a_t := 0$ at times where no data is observed, and we denote the random variable associated with sampling the data at t by O_t. In addition, we discretize the population allele frequency into M hidden states to allow efficient computation: $\mathcal{G} := \{g_0 = 0, g_1, \ldots, g_{M-1} = 1\}$, with $g_i \in [0, 1]$, and $g_{i-1} < g_i$. The hidden state in generation t is then the discretized population allele frequency at t, which we denote by $F_t \in \mathcal{G}$.

To apply the standard HMM framework (Rabiner 1989; Bishop 2006, Ch. 13.2), we must define initial probabilities $\mathbb{P}_{s_1, s_2}\{F_1 = g_i\}$, transition probabilities, and emission probabilities. For now, we assume that the initial probabilities are given. These can either be fixed or estimated, and we provide details on an estimation procedure in Section 2.2.3. For the transition probabilities from hidden state g_i to hidden state g_j, we use the normal approximation to the Wright-Fisher process and define

$$\mathbb{P}_{s_1, s_2}\{F_{t+1} = g_j | F_t = g_i\} := \int_{(g_j + g_{j-1})/2}^{(g_j + g_{j+1})/2} \phi\big(x; \mu_i(s_1, s_2), \tfrac{1}{2N_e} g_i(1 - g_i)\big) dx,$$

where $\mu_i(s_1, s_2) := g_i + g_i(1 - g_i)(s_1(1 - 2g_i) + s_2 g_i)$ for general diploid selection and $\phi(x; \mu, \sigma^2)$ denotes the density of a normal distribution with mean μ and variance σ^2. Here, we also set $g_{-1} := -\infty$ and $g_M := \infty$, which assigns all the probability mass to transition outside of the interval $[0, 1]$ to the respective boundary points. Lastly, the emission probabilities to observe a_t focal alleles in generation t are given by the binomial distribution

$$a_t \sim \text{Bin}(n_t, F_t),$$

with n_t draws and success probability F_t, the respective population allele frequency. Figure 2A depicts a schematic of this HMM.

The *forward-backward algorithm* (Rabiner 1989; Bishop 2006, Ch. 13.2) can then be used to obtain the posterior probability over the hidden states in generation t

$$\gamma_{s_1,s_2}(t,i) := \mathbb{P}_{s_1,s_2}\{F_t = g_i | O_{t'} = o_{t'} \,\forall\, 1 \le t' \le T\}$$

conditional on the observed data and given selection coefficients s_1 and s_2, and the joint posterior of the hidden states

$$\xi_{s_1,s_2}(t,i,j) := \mathbb{P}_{s_1,s_2}\{F_t = g_i, F_{t+1} = g_j | O_{t'} = o_{t'} \,\forall\, 1 \le t' \le T\}.$$

Based on these posterior distributions, posterior expectations

$$\mathbb{E}^{(D)}_{s_1,s_2}\left[h(F_t) \big| O_{t'} = o_{t'} \,\forall\, 1 \le t' \le T\right] = \sum_{i=0}^{M} \gamma_{s_1,s_2}(t,i) h(g_i)$$

and

$$\mathbb{E}^{(D)}_{s_1,s_2}\left[h(F_t, F_{t+1}) \big| O_{t'} = o_{t'} \,\forall\, 1 \le t' \le T\right] = \sum_{i=0}^{M}\sum_{j=0}^{M} \xi_{s_1,s_2}(t,i,j) h(g_i, g_j),$$

can be computed for arbitrary functions $h(\cdot)$ and $h(\cdot,\cdot)$ of the marginal frequency and joint marginal frequencies, respectively. Here, the superscript (D) indicates the use of the model with the discretized allele frequencies.

To find the maximum likelihood estimate (MLE) of the diploid selection coefficients $\widehat{s_1}$ and $\widehat{s_2}$ under this HMM, we use the EM algorithm (Rabiner 1989; Bishop 2006, Ch. 13.2), similar to Mathieson and McVean (2013) in the *additive* case. We refer to this algorithm as an EM-HMM algorithm. The algorithm starts with an initial parameter estimate $(s_1^{(0)}, s_2^{(0)})$. At iteration k, the algorithm then computes $\gamma_{s_1^{(k)},s_2^{(k)}}(t,i)$ and $\xi_{s_1^{(k)},s_2^{(k)}}(t,i,j)$ (E-step), and updates the parameter estimates by maximizing the conditional log-likelihood using

$$\left(s_1^{(k+1)}, s_2^{(k+1)}\right) = \operatorname*{argmax}_{s_1,s_2 \in \mathbb{R}} \mathbb{E}^{(D)}_{s_1^{(k)},s_2^{(k)}}\left[\ln L^{(C)}_{s_1,s_2}(F_1,\ldots,F_T) \big| O_t = o_t \,\forall\, 1 \le t \le T\right], \qquad (2)$$

(M-step) until the estimates converge. Note that, under the binomial model, the emission probabilities are independent of s_1 and s_2 and do not need to be considered explicitly in this update step. Similar to the derivation of the MLE for a given trajectory from equation (1), equation (2) yields

$$s_1^{(k+1)} = \frac{\left(\mathbb{E}[F_T] - \mathbb{E}[F_1]\right) \sum_t \mathbb{E}\left[F_t^2 H_t\right] - \sum_t \mathbb{E}\left[F_t(F_{t+1} - F_t)\right] \sum_t \mathbb{E}\left[F_t H_t\right]}{\sum_t \mathbb{E}\left[H_t\right] \sum_t \mathbb{E}\left[F_t^2 H_t\right] - \left(\sum_t \mathbb{E}\left[F_t H_t\right]\right)^2},$$

$$s_2^{(k+1)} = \frac{\sum_t \mathbb{E}\left[H_t(1 - 2F_t)\right] \sum_t \mathbb{E}\left[F_t(F_{t+1} - F_t)\right] - \left(\mathbb{E}[F_T] - \mathbb{E}[F_1]\right) \sum_t \mathbb{E}\left[F_t H_t(1 - 2F_t)\right]}{\sum_t \mathbb{E}\left[H_t\right] \sum_t \mathbb{E}\left[F_t^2 H_t\right] - \left(\sum_t \mathbb{E}\left[F_t H_t\right]\right)^2},$$

where $H_t := F_t(1 - F_t)$, and we denote the discretized conditional expectation by

$$\mathbb{E}[\,\cdot\,] = \mathbb{E}^{(D)}_{s_1^{(k)},s_2^{(k)}}\left[\,\cdot\,\big| O_t = o_t \,\forall\, 1 \le t \le T\right]$$

for brevity. See Section S.2.2 in the Supplementary Material for details of the derivation.

Note that this approach is not an exact EM algorithm, since it combines the conditional expectation $\mathbb{E}^{(D)}[\cdot]$ computed under the discretized model with maximization of the likelihood $L^{(C)}(\cdot)$ in the continuous model, similar to Mathieson and McVean (2013). The reason for this hybrid approach is that while the conditional log-likelihood can be maximized in the continuous setting using the diploid generalization of Watterson's MLE, the posterior expectations cannot readily be computed. On the other hand, in the discretized model, the posterior expectations can be computed, but the conditional log-likelihood cannot be maximized analytically. The hybrid approach, however, is computationally tractable and yields highly accurate estimates.

2.2.2 Constrained optimization for bespoke selection modes

In addition to estimating unconstrained diploid selection coefficients s_1 and s_2, we want to estimate selection coefficients in the one-parameter selection modes *additive*, *recessive*, *dominance*, and *heterozygote difference*. To this end, we introduce constraints on s_1 and s_2 using the framework of Lagrange multipliers (e.g. Hoffmann et al. 2010, Ch. 7.5). Denoting the likelihood by L_{s_1,s_2}, the optimization problem can be formulated as maximizing the conditional log-likelihood $\mathbb{E}[\ln L_{s_1,s_2}]$ in equation (2) subject to $g(s_1, s_2) = 0$ for an arbitrary function $g(\cdot, \cdot)$, which can be solved by introducing the Lagrangian $\mathcal{L}_{s_1,s_2,\lambda} = \mathbb{E}[\ln L_{s_1,s_2}] - \lambda g(s_1, s_2)$ and solving $\nabla_{s_1,s_2,\lambda}\mathcal{L} = 0$.

All aforementioned one-parameter selection modes of interest can be expressed as a linear constraint, that is, $as_1 - bs_2 = 0$ for suitable $a, b \in \mathbb{R}$. In the Lagrange multiplier formalism, we can thus set $g(s_1, s_2) = as_1 - bs_2$ and solve the constrained optimization problem to obtain the MLE under the respective mode of selection. In Section S.2.3 of the Supplementary Material, we derive explicit expressions for the update rules for s_1 and s_2 for arbitrary a, b. As an example, Figure 2B shows the iterations of the *additive*, *recessive*, *dominance*, and unconstrained general diploid selection EM-HMM algorithms on a dataset simulated under *additive* selection.

2.2.3 Estimation of the initial allele frequency distribution

The discretized distribution of the initial allele frequency can be fixed for the analysis or estimated as well. When estimating it, we fit a beta distribution to the initial frequency, as it is a flexible distribution with only two parameters, denoted by (α, β), which avoids potential over-parametrization. If we assume that the initial distribution does not depend on the selection coefficients, then the discretized version of the EM update rule given in equation (2) becomes

$$
\begin{aligned}
\left(\alpha^{(k+1)}, \beta^{(k+1)}\right) &:= \operatorname*{argmax}_{\alpha,\beta \in \mathbb{R}_{>0}} \mathbb{E}\left[\ln L^{(D)}_{\alpha,\beta}(F_1)\right] \\
&= \operatorname*{argmax}_{\alpha,\beta \in \mathbb{R}_{>0}} \sum_{m=0}^{M-1} \gamma_{\theta^{(k)}}(0, m) \ln p(m; \alpha, \beta)
\end{aligned}
\tag{3}
$$

for the parameters (α, β). Here, $L^{(D)}_{\alpha,\beta}(\cdot)$ denotes the likelihood of a discretized beta distribution, the conditional expectation $\mathbb{E}[\cdot]$ is now parameterized by $\theta^{(k)} := \left(\alpha^{(k)}, \beta^{(k)}, s_1^{(k)}, s_2^{(k)}\right)$, and $p(m; \alpha, \beta)$ is the integral of a beta distribution with parameters (α, β) over the m-th discretization interval. At each iteration k of the EM-HMM algorithm, we solve equation (3) numerically to update α and β alongside the selection coefficients. Since the EM update for the selection coefficients already requires computing $\gamma^{(D)}_{\theta^{(k)}}(t, i)$, the extra step of updating the initial condition comes at minimal

Figure 2: A) Schematic of an HMM. Each "stoplight" represents a haploid sample at the given time, with a certain number of focal alleles. Also plotted are three possible allele frequency trajectories through the hidden state space. Trajectories with population frequencies closer to the frequencies in the samples are given more weight when computing the expected values in the E-step of the algorithm. B) Log-likelihood surface and path of the EM-HMM optimization under each mode for a given replicate simulated under *additive* selection.

computational cost. We observed that estimating the initial distribution improves the accuracy of the selection coefficient estimation, see Figure S.11 in the Supplementary Material.

2.2.4 Additional implementation details

Analyzing data using our EM-HMM algorithm requires choosing the number of hidden states or discretization intervals M, and where to place the discretization points $\mathcal{G} := \{g_0 = 0, g_1, ..., g_{M-1} = 1\}$. A common choice for the discretization points is to space them equidistantly, that is $g_i = \frac{i}{M-1}$. However, we observed that this discretization did not perform well when selection was strong, see Section S.5 in the Supplementary Material for details. We thus decided to use the Chebychev nodes $g_i := \frac{1}{2} + \frac{1}{2} \cos\left(\frac{2\pi i}{2*(M-1)}\right)$ for $0 \leq i \leq M - 1$. These nodes are often used for numerical integration, since they mitigate Runge's phenomenon (Mathews and Fink 2003, Ch. 4.5), and indeed improve performance of our algorithm. Intuitively, when using equidistant spacing, the variance of the normal distribution for the transition used in the M-step becomes smaller than the size of the discretization intervals near the boundary, and consequently, the density at the discretization points does not approximate the probability mass in the interval well. Using the Chebychev nodes effectively increases the density of discretization points near the boundary and decreases it in the interior, mitigating this problem.

We analyzed simulated data using different choices of the number of hidden states M in Section 3.3.1. We observed that the inference does not perform well when using less than 250 hidden states, but is stable for higher values. To balance accuracy of the EM-HMM algorithm and computational efficiency, we thus choose $M = 500$ hidden states in all analyses. Unless stated otherwise, we initialize the selection coefficients for the EM-HMM algorithm with $s_1^{(0)} = s_2^{(0)} = 0$ and set the starting parameters for the initial distribution to $\alpha^{(0)} = \beta^{(0)} = 1$, which corresponds to the uniform

distribution. For all analyses, we use the convergence criterion that the difference in log-likelihood between iteration k and iteration $k+1$ must be less than 10^{-3}.

We noticed that, especially for simulations under neutrality, the EM-HMM for many replicates would return $s_1^{(0)} = s_2^{(0)} = 0$ as the MLE, but the actual MLE would deviate slightly from 0, with a log-likelihood increase $< 10^{-3}$. Nonetheless, this would result in many replicates reporting a log-likelihood ratio of exactly 1, and consequently, p-values around 1 were not well calibrated. To avoid potentially unwanted distortions to the distribution of p-values, we required the EM-HMM to perform at least 5 iterations before stopping, taking the maximum log-likelihood obtained in the first 5 iterations if further iterations do not increase the log-likelihood. In practice, this has minimal effect on replicates simulated under selection, since the EM-HMM requires more than 5 steps to converge in these cases.

2.3 Distinguishing between modes of selection

2.3.1 P-values for a single alternative

Before describing our approach to the full problem of inferring the mode of selection with multiple alternatives, we first outline the solution to the task of rejecting neutrality in the case of a one-parameter alternative mode of selection. For a given replicate and one-parameter selection mode, we use the EM-HMM algorithm to compute the MLEs $(\hat{s}, \hat{\alpha}, \hat{\beta})$, as well the log-likelihood ll_s for these parameters. Moreover, we compute the MLEs $(\hat{\alpha_0}, \hat{\beta_0})$ and log-likelihood ll_0 under neutrality ($s = 0$). Treating the parameters of the initial distribution as nuisance parameters, we then perform standard likelihood-ratio testing using the test likelihood-ratio statistic $D = 2(ll_s - ll_0)$. As the sample size goes to infinity, Wilks' theorem (Wilks 1938) implies that D should be χ^2 distributed with one degree of freedom under the null hypothesis $s = 0$, which we denote by $\chi^2(1)$. However, each replicate consists only of one set of temporal samples, so we are not operating in the asymptotic regime of Wilks' theorem, and have no theoretical guarantee regarding the distribution of D. Nonetheless, in Section 3.2.2, we show that if we compute p-values using $p = 1 - \mathbb{P}\{\chi^2(1) < D\}$, assuming a $\chi^2(1)$ distribution, then the resulting p-values in the neutral case do approximately follow a uniform distribution, and thus the asymptotic formula provides a good approximation. P-values computed this way can thus be used accept or reject the null hypothesis of neutrality for a given dataset, under the given one-parameter alternative.

2.3.2 Multiple alternatives

To infer the mode of selection with multiple alternatives, we need to compare models with different numbers of parameters: One parameter in the constrained case, two in the unconstrained case. A natural criterion to use would thus be the Akaike information criterion (Bozdogan 1987): AIC $= 2k - 2ll$, where k is the number of parameters of the model, and ll is the log-likelihood at the MLE. However, we found that the penalty under the AIC of one log-likelihood unit per model parameter yielded a very high false positive rate for neutral replicates. The AIC has two additional disadvantages for our problem: first, AIC values cannot readily be integrated into a hypothesis testing framework since we cannot rely on asymptotic results for its distribution, as discussed in Section 2.3.1. Second, simply choosing the model with the highest AIC cannot provide guarantees for statistical properties such as the false discovery rate (FDR).

To address these issues with the AIC, we developed the following simulation-based procedure to infer the mode of selection and assess significance for a target dataset with a given sampling

scheme. Broadly, the method finds an optimal value k_{opt} to penalize the two-parameter modes of selection, and fits the distribution of likelihood-ratios between the highest-likelihood amongst non-neutral modes and neutrality to a generalized gamma distribution. If the p-value from this fit is significant, we classify the mode of selection according to the non-neutral mode with the highest-likelihood. In detail, our procedure is as follows:

1. Simulate replicates that match the sampling times t_k, sample sizes n_{t_k}, the total number of generations T, and the distribution of the initial frequency for a target dataset (known or empirically estimated). Moreover, simulate replicates under neutrality, each one-parameter mode of selection *additive*, *recessive*, *dominant*, complete *over-* and complete *underdominance*. For the selection coefficient s and the effective population size N_e, choose a range of combinations that cover the range expected in the analysis. See Section 4.2 for an example of such a set of simulations for a target dataset.

2. For each simulated replicate, use the EM-HMM algorithm under all constrained modes of selection, *additive*, *recessive*, *dominant*, and the unconstrained mode, to obtain the MLEs of the selection coefficients in each case and corresponding log-likelihood values ll_0 ($s_1 = s_2 = 0$), ll_{add}, ll_{rec}, ll_{dom}, and ll_{uncon}, respectively. Note that the parameters of the initial distribution are again treated as nuisance parameters for this inference.

3. Using the simulated *non-neutral* replicates only (the magenta region in Figure 3), determine the optimal penalty k_{opt} for the unconstrained (two-parameter) mode. To this end, classify each replicate according to $\max\{ll_{\mathrm{add}}, ll_{\mathrm{rec}}, ll_{\mathrm{rec}}, ll_{\mathrm{uncon}} - k\}$, with the unconstrained mode penalized by k. The simulations for *over-* and *underdominance* are considered correctly classified if the highest likelihood is $ll_{\mathrm{uncon}} - k$ and the estimated s is located in the correct region of diploid selection space. Varying the penalty k, maximize the number of correctly classified replicates over all non-neutral replicates to determine k_{opt}. In practice, we plot the number of correctly classified replicates against k and choose k_{opt} to maximize this graph after suitable interpolation.

4. For all simulated *neutral* replicates (the green region in Figure 3), compute the test statistic $\delta_{ll} := 2(\max\{ll_{\mathrm{add}}, ll_{\mathrm{dom}}, ll_{\mathrm{rec}}, ll_{\mathrm{uncon}} - k_{\mathrm{opt}}\} - ll_0)$. Fit a generalized gamma distribution to the distribution of the δ_{ll} values under neutrality that are greater than the median value.

5. Estimate the MLE parameters for all constrained modes and the unconstrained mode on the target dataset using the EM-HMM. Compute δ_{ll} for the target dataset using the k_{opt} determined from the simulated data in step 3.

6. Use the generalized gamma distribution estimated in step 4 to obtain p-values for the target dataset based on the δ_{ll} statistic computed in step 5. Reject neutrality based on these p-values at a given significance threshold.

7. If neutrality is rejected, classify the mode of selection according to $\max\{ll_{\mathrm{add}}, ll_{\mathrm{dom}}, ll_{\mathrm{rec}}, ll_{\mathrm{uncon}} - k_{\mathrm{opt}}\}$. In the case that the unconstrained model is the most likely model, classify as *over-dominant* if $\widehat{s_1} > \max(0, \widehat{s_2})$, *underdominant* if $\widehat{s_1} < \min(0, \widehat{s_2})$, or *indeterminate* in the other cases.

The unconstrained EM-HMM has an extra degree of freedom compared to the one-parameter EM-HMMs, and thus penalizing its output is necessary. We thus developed the procedure outlined

Inferred True	H_0	H_1	$H_2 - k$
H_0 (Neutral)		Step 4	
H_1 (Add., Dom., Rec.) H_2 (Over/Underdom.)		Step 3	

Figure 3: Abstract representation of the confusion matrix components used in key steps of the procedure to infer the mode of selection. In step 3, we adjust the penalty k on the likelihood of the unconstrained EM-HMM model to maximize the number of correctly inferred replicates - that is, the sum of the values on the diagonal of the magenta region. In step 4, we numerically fit the distribution of the test statistic δ_{ll} for all simulated neutral replicates (green region) to a generalized gamma distribution.

above for determining the optimal penalty k_{opt} in a specific scenario, and use it to obtain well-calibrated p-values. If the simulations to determine k_{opt} are prohibitive, we recommend using a value of $k_{\text{opt}} = 1.5$, which produced acceptable p-values across a range of simulations. Figure 3 shows an abstract confusion matrix, highlighting the values used in different steps of the procedure.

Unlike for the likelihood-ratio statistic D, we found that the δ_{ll} statistic was not well-described by a χ^2 distribution with different degrees of freedom. To address this, we fit the δ_{ll} values to a generalized gamma distribution (Stacy 1962), which we denote by G. Since we are only interested in low p-values, we further improve the fit of this generalized gamma distribution by fitting only the δ_{ll} values above the median, denoted by $\overline{\delta_{ll}}$. Specifically, the p-values for the median-generalized gamma fit have the form:

$$p = \begin{cases} \frac{1}{2}\big(1 - \mathbb{P}\{G \leq \delta_{ll} - \overline{\delta_{ll}}\}\big), & \text{if } \delta_{ll} > \overline{\delta_{ll}}, \\ 1 - \frac{\delta_{ll}}{2*\overline{\delta_{ll}}}, & \text{if } \delta_{ll} \leq \overline{\delta_{ll}}, \end{cases}$$

where the p-values for $\delta_{ll} \leq \overline{\delta_{ll}}$ are linearly interpolated from 0.5 to 1.

3   Simulation Study

To assess the performance of our EM-HMM algorithm to estimate selection coefficients and infer the correct mode of selection across a variety of selection regimes, we simulated datasets under several combinations of parameters and exhibit the accuracy of the inference.

3.1   Simulation parameters

We simulated population allele frequency trajectories under the discrete Wright-Fisher model for a given number of generations and sampled a certain number of haplotypes at given times from a binomial distribution with success probability given by the respective population allele frequency. Specifically, we simulated datasets under neutrality ($s_1 = s_2 = 0$), each one-parameter selection mode (*additive*, *recessive*, *dominant*), where we set $s = s_2$, and complete *over-* or *underdominance* with $s = s_1$ and $s_2 = 0$. We simulated using selection coefficients $s \in \{0.005, 0.01, 0.025, 0.05\}$ and the length of the simulated trajectory in generations $T \in \{101, 251, 1001\}$. We considered two different types of initial conditions: 1) The population allele frequency is initialized at a fixed value $p \in \{0.005, 0.25\}$ for each replicate, and 2) the initial frequency of each replicate is drawn from the set $i \in \{\frac{1}{2N_e}, \dots, \frac{2N_e-1}{2N_e}\}$ with probability proportional to $\frac{1}{i}$. The latter is the stationary distribution

for neutral segregating sites under the Poisson Random Field model (Sawyer and Hartl 1992), and thus corresponds to selection from standing variation. We conditioned the simulated data on the focal allele A not being lost or fixed, depending on the mode of selection. Specifically, for *additive*, *dominant*, and *recessive* selection, we conditioned on A not being fixed at the first generation and not being lost at the last generation; for *overdominance*, the simulation is conditioned on A segregating in the last generation; for *underdominance*, we conditioned on A segregating in the first generation.

For all simulations, the population size was set to $N_e = 10,000$. the sampling scheme consisted of $K = 11$ equidistantly-spaced points in time, where the first and last point align with the start and the end of the simulated trajectory, respectively. At each time, we sampled $n_{t_k} = 50$ haploids. For each combination of parameters, we simulated 500 replicates. For each replicate, we performed inference using the unconstrained diploid EM algorithm as well as the four constrained modes.

In Section 3.2.1, Section 3.2.2, and Section 3.2.3, we present results for the simulations with 251 generations and initial allele frequency fixed to $p = 0.25$. The results for other combinations of number of generations and initial allele frequency distribution can be found in Section S.3 of the Supplemental Material. In general, the results are similar, with the exception of extreme cases: For *recessive* or *underdominance* with low initial frequencies, the selected alleles is often lost, resulting in inaccurate estimates.

3.2 Simulation results

3.2.1 Estimating selection coefficients

In Figure 4A, we present the distribution of the estimated selection coefficient $\hat{s}$ across replicates simulated under the *additive*, *recessive*, *dominant*, *overdominant*, and *underdominant* mode. Estimates of the selection coefficients are in general accurate and unbiased. For intermediate-to-strong selection, overdominance has the highest variance in $\hat{s}$ estimates. This occurs because the overdominance replicates reach stationarity ($p = 0.5$) quickly, resulting in fewer informative samples, and the variance of the estimates is increased.

3.2.2 Validation of single-alternative p-values

Next, we report the single-alternative p-values obtained via the method described in Section 2.3.1 for all four modes: *additive*, *recessive*, *dominant*, and *heterozygote difference*. Figure 4B shows a Q-Q plot, where we plot the empirical p-values against their expected value for replicates simulated under neutrality. For all selection modes, the points follow the line $y = x$ closely, indicating that the p-values are well-calibrated, and that the distribution of the likelihood-ratio statistic D is well-approximated by a $\chi^2(1)$ distribution.

Additionally, we report the performance of the single-alternative tests in terms of the area under the curve (AUC) of a receiver-operator characteristic curve. An AUC value of 0.5 indicates no power to distinguish neutral replicates from non-neutral replicates, whereas an AUC value of 1 indicates perfect discrimination. Figure 4C reports the AUC values each simulated selection mode over the range of selection parameters s. We observe that in these scenarios, the method is well-powered to reject the null hypothesis for $s \geq 0.01$. AUC values are lower in the case of recessive selection, since 251 generations can be insufficient for replicates at low frequency to escape the drift-dominated regime.

Figure 4: A) Boxplot of $\hat{s}$ for 500 replicates simulated under each selection mode. Whiskers extend to 2.5% and 97.5%-tiles. Number of estimates outside plotting range indicated above the plot. Estimates are generally unbiased and have low variance, with the exception of overdominance. B) Q-Q plot of $-\log_{10}(\mathrm{p})$ against $\mathbb{E}[-\log_{10}(\mathrm{p})]$ of single-alternative tests for neutral replicates. Inset shows same plot for raw values. The p-values are well-calibrated under all modes of selection. C) Table of AUC values based on likelihood-ratios for each selection mode and selection strength simulated. For $s > 0.01$, AUC values are near 1, indicating perfect discrimination between neutral and non-neutral replicates. D) Q-Q plot of $-\log_{10}(\mathrm{p})$ against $\mathbb{E}[-\log_{10}(\mathrm{p})]$ for the δ_{ll} statistic, assuming different distributions. Generalized gamma yields the best calibration. For all simulations, number of generations is $T = 251$ and initial condition is fixed at frequency $p = 0.25$.

Figure 5: Confusion matrix of the results when inferring the mode of selection, with a p-value threshold of 0.05 to reject neutrality. Cells with an orange border represent correct classification. The correct model is inferred for the majority of replicates for $s = 0.05$, with the exception of *underdominant*. For all simulations, number of generations is $T = 251$ and initial condition is fixed at frequency $p = 0.25$.

3.2.3 Validation of selection mode inference

We furthermore computed p-values for the simulated datasets in the case of multiple alternatives, following the procedure outlined in Section 2.3.2. A Q-Q plot of the p-values against their expected values is shown in Figure 4D. P-values computed using the generalized gamma distribution fall closer to the line $y = x$ compared to the $\chi^2(1)$ or $\chi^2(2)$ distribution, indicating a better calibration.

We then tested the ability of our approach to correctly identify the mode of selection. Figure 5 shows the confusion matrix resulting from inferring the mode of selection with a p-value significance threshold of 0.05. Each row of the confusion matrix represents all replicates simulated under a particular mode of selection. The numbers in the corresponding column indicate the fraction of replicates where a particular mode is inferred. Values on the diagonal reflect identification of the correct mode, whereas off-diagonal values reflect replicates where the mode is not correctly inferred. For $s = 0.05$, neutrality is rejected in most cases and the correct model is inferred for a majority of replicates, except for the simulations under *underdominance*. For weaker selection, $s = 0.01$, neutrality is rejected for over 50% of the replicates, indicating that power to reject neutrality exists, but accuracy to correctly infer the mode of selection is limited.

Lastly, we investigate accuracy of the MLE of the selection parameters, given that the mode was

Figure 6: Strip plots of inferred against true s, conditioned on the correct model being chosen among multiple alternatives. In the neutral case, the plot shows the neutral replicates classified into certain modes. The number over each strip indicates the number of replicates correctly classified, out of 500 total simulated. For $s = 0.005$ and neutrality, the inferred values are strongly biased, due to a "winner's curse" effect. For all simulations, number of generations is $T = 251$ and initial condition is fixed at frequency $p = 0.25$.

correctly classified. Figure 6 shows the distribution of $\hat{s}$ estimates for replicates that were correctly classified as *additive*, *dominant*, as *recessive* selection, as well as $\hat{s}$ for neutral replicates that were classified incorrectly. For larger s, correctly classified replicates are more tightly clustered around the true value, with far fewer outliers compared to the distribution using all replicates. For lower values of s and neutrality, the $\hat{s}$ estimates for replicates classified as non-neutral are biased. This is primarily due to a "winner's curse" effect, where neutrality is only rejected for extreme replicates with extreme MLEs.

3.3 Robustness of estimates

3.3.1 Parameter variation

Next, we investigated how estimation accuracy is affected if we vary parameters that were fixed in the previous simulation study. We varied the number of times a sample is taken $K \in \{2, 5, 11, 35, 101\}$, the number of haploid samples at each timepoint $n_{t_k} = \{6, 20, 50, 100, 200\}$, the effective population size $N_e \in \{10^2, 10^3, 10^4, 10^5, 10^6\}$, and the number of hidden states in the HMM $M \in \{100, 250, 500, 1000, 2000\}$. We varied one parameter at a time, and fixed the others to $K = 11$, $n_{t_k} = 50$, $N_e = 10,000$, and $M = 500$. We fixed the selection coefficient s to 0.025 for all simulations, set the total number of generations to $T = 251$, and set the initial distribution to a fixed frequency at $p = 0.25$. Boxplots of the estimated selection coefficients for *additive*, *recessive*, and *dominant* selection are shown for each scenario in Figure 7. Similar to Mathieson and McVean (2013) in the *additive* case, we find that the accuracy does not depend strongly on the sampling scheme except for very extreme values, such as sampling 6 haplotypes per timepoint or only sam-

pling at the beginning and end of the trajectory. However, the boxplots are tighter for larger sample sizes and more sampling times. This implies that both variance due to finite sampling and variance in the true allele frequency contribute to uncertainty of the selection coefficients estimates. Similarly, the accuracy of the estimates are comparable for 250, 500, 1000, and 2,000 hidden states. However, for 100 hidden states, the estimates are strongly biased downwards. We thus recommend using $M = 500$ to balance accuracy with computational efficiency.

3.3.2 Misspecification of selection mode

To investigate the accuracy of the estimates when the mode of selection is incorrect, we re-analyzed a subset of the data under scenarios in which we misspecify the mode. We consider two cases: (1) Simulating under *additive*, *dominant*, *recessive*, *over-*, or *underdominant* selection and analyzing under *additive* selection, and (2) simulating under *additive* selection and analyzing under *additive*, *dominant*, *recessive*, or *heterozygote difference*. Figures 8A and 8B depict boxplots of the MLEs $\hat{s}$ under both of these misspecification scenarios.

When simulating under each mode of selection and analyzing under the *additive* EM-HMM, the estimates of $\hat{s}$ are biased for all modes, most strongly for overdominant selection at high s. Alleles undergoing strong overdominant selection quickly reach a stationary frequency of $p = 0.5$. Since the alleles do not continue to increase in frequency, the *additive* EM-HMM thus underestimates the true selection strength. In the case when we simulate under *additive* selection and analyze under each selection mode, the $\hat{s}$ estimates are not strongly biased for *recessive* and *dominant*, but reverse sign for *heterozygote difference* at high selection strengths. In this case, the allele frequency increases beyond $p = 0.5$, so the heterozygote difference EM-HMM changes from *overdominance* to *underdominant* selection. Overall, the biases in the estimates $\hat{s}$ indicate that using the incorrect one-parameter EM-HMM provides inaccurate estimates of s.

We furthermore investigated the power to reject or accept neutrality in the single-alternative testing framework described in Section 2.3.1 under each of these misspecification scenarios. The resulting AUCs are presented in Figures 8C and 8D. For the case of analyzing other modes under *additive* selection, AUC values are slightly lower than the values in Figure 4C, but still substantially greater than 0.5. For the *additive* datasets analyzed using the incorrect selection modes, the AUC values do not differ substantially between different modes. In both misspecification scenarios, the one-parameter *additive* EM-HMM has sufficient power to accurately reject or fail to reject neutrality for moderate-to-high selection coefficients. Thus, if the goal is only to identify non-neutral evolution, it can be sufficient to analyze given data using the *additive* EM-HMM only.

We also performed simulations with $N_e = 10,000$, and analyzed these simulations using $N_e \in \{5,000, 10,000, 20,000\}$. We present the results in Section S.4 in the Supplementary Material, and find that while the estimates of the selection coefficients are unbiased and the variability is not affected strongly, the p-values for smaller $N_e = 5,000$ are more conservative and the p-values for larger $N_e = 20,000$ are less conservative.

4 Ancient DNA dataset from Great Britain

4.1 GB aDNA dataset

Having demonstrated the ability of our approach to accurately characterize selection in simulated data, we applied our method to time series genetic data obtained from human ancient DNA. To

Figure 7: Boxplots for data simulated and inferred under *additive*, *dominant*, and *recessive* selection for different parameter combinations. Whiskers extend to 2.5% and 97.5%-tiles. We vary: A) Number of samples taken at each timepoint, B) Number of timepoints sampled, C) Effective population size, D) Number of hidden states in the HMM. Except for the lowest parameters, estimates of the selection coefficient are unbiased. Width of boxplots decrease as number of samples and sampling timepoints increases. For all simulations, number of generations is $T = 251$ and initial condition is fixed at frequency $p = 0.25$. When not varied, parameters used are $K = 11$ sampling times, $n_{t_k} = 50$ samples at each timepoint, $N_e = 10,000$, and $M = 500$ hidden states.

Figure 8: A) Boxplots of $\hat{s}$ for 500 replicates simulated under each selection mode and analyzed using *additive* selection. The estimates are mostly inaccurate. B) Boxplots of $\hat{s}$ for 500 replicates simulated under *additive* selection and analyzed under each selection mode. Again, the estimates are not accurate if the mode is incorrect. For both sets of boxplots, whiskers extend to 2.5% and 97.5%-tiles. C) Table of AUC values for data simulated under each selection mode using likelihood-ratios obtained assuming *additive* selection. Values are lower compared to analyzing under the correct mode of selection, but still show substantial power to reject neutrality. D) Table of AUC values for data simulated under *additive* selection using likelihood-ratios obtained assuming each possible selection mode. The values are very similar across selection modes. For all simulations, number of generations is $T = 251$ and initial condition is fixed at frequency $p = 0.25$.

this end, we extracted the genotype information from a subset of the individuals in the Allen Ancient DNA Resource (AADR), version v54.1, which is a frequently updated repository that aims at comprising most currently published ancient DNA datasets (Mallick et al. 2024).

Our method assumes that the data originates from a single panmictic population. We thus follow a rationale similar to Mathieson and Terhorst (2022) and restrict our analyses to samples from Great Britain in the last 4,450 years. We manually removed samples that were not from the mainland. Restricting to this geographic area and time window, it is unlikely that the data we analyze substantially violates the assumption of a single panmictic population, since the last major admixture event into Great Britain is estimated to have occurred before 4,450 years ago (Olalde et al. 2018; Chintalapati et al. 2022), although evidence for some more recent gene flow has been presented (Patterson et al. 2022; Gretzinger et al. 2022). Figure S.12 in the Supplementary Material shows a map with sample locations and times. Moreover, Figure S.13 in the Supplementary Material depicts two PCA plots, demonstrating that the samples cluster with modern European individuals on a global scale, but do not exhibit evidence of strong structure on a local scale.

When analyzing samples that were genotyped using the 1240K capture assay and samples genotyped using whole genome sequencing together, we noticed spurious signals of selection, see Figure S.16 in the Supplementary Material. As a conservative approach, we thus only analyzed samples that were genotyped using 1240K capture, and excluded the 174 whole genome samples in this geographic area and timeframe from our dataset. We note that this conservative approach also removed the present-day samples. Our resulting dataset, henceforth referred to as the GB aDNA dataset, thus comprises 504 ancient pseudo-haploid samples genotyped using the 1240K assay and spanning 125 generations, when using a generation time of 28.1 (Moorjani et al. 2016). These individuals are a subset of the individuals published by Olalde et al. (2018), Patterson et al. (2022), and Gretzinger et al. (2022).

We furthermore applied three filters to each SNP in the dataset. First, each SNP must have genotyped samples at two or more timepoints. Second, each SNP must have more than 50 (10% of 504) samples genotyped in total. Third, the minor allele frequency when pooling all samples at a given SNP across time must be greater than 0.05. We expect only SNPs that pass these filters to yield reliable signals of selection. In total, out of the 1,150,638 SNPs available, 743,417 (64.6%) pass these three filters and were used in the final analysis.

4.2 Data-matched simulations

To assess the accuracy of our method in the specific context of the GB aDNA dataset, and to calibrate our procedures to compute p-values and infer the mode of selection, we simulated data that matches the sampling scheme and timeframe of the GB aDNA dataset. Specifically, we simulated an allele frequency trajectory under a particular mode of selection for $T = 125$ generations assuming $N_e = 10,000$, and sampled from it according to the sampling times and sizes of the GB aDNA dataset. In addition, we randomly omitted samples with probability equal to the fraction of missingness at a randomly selected SNP in the GB aDNA dataset, to emulate the same degree of missing data. We provide a histogram of the fraction of samples missing for each SNP in the GB aDNA dataset in Figure S.14 in the Supplementary Material.

As before, we simulated using selection coefficients $s \in \{0, 0.005, 0.01, 0.025, 0.05\}$. To generate the initial frequency for each simulated replicate, we first estimated the parameters α and β for the beta distribution of the initial frequency under neutrality ($s = 0$) at each SNP in the GB aDNA dataset that passes our filters, as described in Section 2.2.3. Figure S.15 in the Supplementary

Material shows a histogram of the means of the initial distributions estimated for each SNP. For each simulated replicate, we then chose one SNP uniformly at random, and set the initial frequency of the replicate equal to the mean of the chosen SNP. This procedure ensures that the initial frequency distribution of the simulated data matches the GB aDNA dataset closely, and captures any potential biases, for example, due to ascertainment of the 1240K SNP set. To further ensure that the simulated replicates match the GB aDNA dataset, we apply the same three SNP-based filtering criteria, and only keep replicates that pass all filters.

We simulated 10,000 replicates under neutrality and for each s under each of the five one-parameter selection modes. We then computed the MLEs for the selection coefficients using our EM-HMM for each replicate, with the mode of selection in the analysis matched to the simulated mode, and show the distribution of the MLEs $\hat{s}$ in Figure 9A. As for the simulated datasets in Section 3.1, the estimates of the selection coefficients are largely unbiased. However, unlike in the simulated datasets, for strong selection, the data-matched replicates have a slight downward bias. This is likely due to the fact that the data-matched simulation comprise half the number of generations as the simulation study presented in Figure 4A. The simulated datasets with $T = 100$ generations show a similar, though less pronounced, downward bias at high s, see Figure S.3A in the Supplementary Material.

Next, we applied our procedure to test for a single alternative and our procedure to infer the mode of selection to the data-matched simulations. For the test of a single alternative, Figure 9B shows a Q-Q plot of the p-values computed from the likelihood-ratio statistic D assuming a $\chi^2(1)$ distribution against their expected value. We again observe that these p-values are well-calibrated. As in Section 3.2.2, we computed AUC values to assess the single-alternative on the data-matched simulations, which are shown in Figure 9C. For $s \geq 0.025$, AUC values are between 0.8 and 1, with the exception of *overdominant* selection. In general, the AUC values are lower than in Figure 4C; for example, for $s = 0.05$ in the case of *overdominance*, the AUC is 0.81 for the data-matched simulations, compared to 0.99 for the simulations in Section 3.1. This is likely due to the initial distribution that we used for the data-matched simulations, which has more weight close to $p = 0$, combined with the reduced number of generations. Both of these properties result in smaller cumulative changes in allele frequency, and thus power to distinguish from neutral replicates is reduced.

For multiple alternatives, Figure 9D shows the p-values computed from the statistic δ_{ll} under $\chi^2(1)$, $\chi^2(2)$ and the generalized gamma distributions, plotted against the expected value. The δ_{ll} statistic is again poorly-described by a $\chi^2(1)$ distribution, but the generalized gamma distribution, and to a lesser extent, the $\chi^2(2)$ distribution, provides a good calibration of the p-values in the tail of the distribution. We thus again proceed with the generalized gamma distribution.

In Figure 10, we summarize the inference of the selection mode for all replicates of the data-matched simulations using the procedure described in Section 2.3.2; again using a p-value threshold of 0.05 to reject neutrality. For $s = 0.05$, we successfully reject neutrality for a large fraction of replicates, up to 95% of cases when simulating under *additive* selection, but the correct model is only inferred in a approximately 30% of the cases. For the lower selection strength $s = 0.01$, neutrality is only rejected in a small percentage of the simulations. Moreover, *overdominant* selection is inferred for many replicates regardless of the selection mode that the data is simulated under. *Overdominance* is primarily distinguished from other modes of selection if allele frequency trajectories remain at intermediate frequencies. However, for simulations with a low number of generations, allele frequencies often fail to increase beyond frequencies of 0.5, regardless of the selection mode.

Figure 9: A) Boxplots of MLEs $\hat{s}$ for all one-parameter selection modes. Each boxplot shows 500 replicates out of the 10,000 simulated. Whiskers extend to 2.5% and 97.5%-tiles. Estimates are largely unbiased for small s, and slightly biased downward for large s. B) Q-Q plot of $-\log_{10}$(p-value) against $\mathbb{E}[-\log_{10}$(p-value)] for single-alternative tests obtained using the $\chi^2(1)$ distribution. Inset shows raw p-value against expected p-value. As for the simulated datasets, p-values are well calibrated. C) Table of AUC values for data-matched simulations using likelihood-ratios for each one-parameter selection mode. D) Q-Q plot of $-\log_{10}$(p-value) against $\mathbb{E}[-\log_{10}$(p-value)] for multiple-alternative test statistic δ_{ll} using $\chi^2(1)$, $\chi^2(2)$, and generalized gamma distributions. The generalized gamma distribution provides the best fit. For all simulated replicates, the number of generations is $T = 125$ and the initial frequency is randomly drawn from the initial frequencies estimated in the GB aDNA dataset.

Figure 10: Confusion matrix for procedure to infer selection mode applied to data-matched simulations, using a p-value threshold of 0.05. Each cell represents the fraction of replicates that were classified as a particular mode. Performance is worse than for the simulations in Section 3.1 - the correct model is only inferred for the plurality of replicates for $s = 0.05$. For all simulations, $10,000$ replicates are simulated under a given selection mode and strength, the number of generations is $T = 125$ and the initial frequency is randomly drawn from the set of initial frequencies in the GB aDNA dataset.

Thus, *overdominance* can often not be distinguished from the other modes. As in the simulation study presented in Section 3.2.3, we find that we can detect non-neutral evolution, but power to infer the correct mode of selection is low.

Lastly, Figure 11 shows the distribution of the inferred selection coefficient, conditional on inferring the correct mode of selection among multiple alternatives, as well as the distribution of the inferred selection coefficient for neutral replicates classified as non-neutral. For $s > 0.01$, estimated selection coefficients are similar to the unconditional estimates, indicating that our model inference procedure does not strongly bias the estimates in this parameter regime. However, for $s = 0.005$ and $s = 0.01$, most inferred coefficients are higher than the true value. As with the simulated data in Section 3.2.3, the data simulated with lower selection coefficients exhibit a "winner's curse" phenomenon, where only replicates with extreme allele frequency changes are classified as non-neutral, and consequently $\hat{s}$ is also large.

Figure 11: Strip plots of $\hat{s}$ against true s, conditioned on inferring non-neutrality for the neutral replicates and on inferring the correct selection mode among multiple alternatives for non-neutral replicates. Each strip plot is for 500 replicates randomly chosen from the 10,000 simulated. Number above each strip indicates number of replicates correctly classified. For $s = 0.005$ and $s = 0.01$, the inferred values are biased upwards, due to the "winner's curse". For all simulations, the number of generations is $T = 125$ and the initial frequency is randomly drawn from the distribution estimated in the GB aDNA dataset.

4.3 Analysis of the GB aDNA dataset

4.3.1 Significant signals of selection in the GB aDNA dataset

We then applied our EM-HMM algorithm and procedure to infer the mode of selection to all 743,417 SNPs in the GB aDNA dataset that pass our filters. From the time series genetic data at each SNP, we estimated the MLE using our EM-HMM for all one-parameter modes of selection and the unconstrained two-parameter MLE, assuming $N_e = 10,000$. Here, we primarily describe the results under the *additive* mode as well as the results of the procedure to infer the mode of selection. Results for the other one-parameter modes, as well as Q-Q plots of the p-values for all modes, can be found in Section S.7 of the Supplementary Material.

Figure 12 shows a Manhattan plot of the p-values for the single-alternative likelihood-ratio test computed using the *additive* EM-HMM. We also indicate the significance threshold obtained from applying the Benjamini–Hochberg (BH) procedure (Benjamini and Hochberg 1995) to correct for multiple testing at a false discovery rate (FDR) of $\alpha = 0.05$. In Section S.8 of the Supplementary Material, we compute the same p-values when permuting the sampling times in the dataset, demonstrating that the enrichment of low p-values we observe is a reliable signal in the data. Furthermore, we observe clusters of low p-values on Chromosomes 2, 5, and 6. This clustering of p-values is expected, since due to genetic hitchhiking (Smith and Haigh 1974), SNPs that are in proximity to an actual target of selection, and thus in linkage disequilibrium (LD) with the target, will also exhibit non-neutral dynamics.

In addition to these clusters, we observe several isolated SNPs with p-values exceeding the

Figure 12: Manhattan plot of p-values obtained from the likelihood-ratio test under the *additive* mode of selection at all SNPs in the GB aDNA dataset. The significance threshold is obtained via the Benjamini–Hochberg procedure with an FDR of $\alpha = 0.05$. We observe clusters of significant p-values on chromosomes 2, 5, and 6, as wells as several isolated signals.

BH threshold, but no surrounding SNPs show evidence of selection, a pattern that would not be expected under genetic hitchhiking. We thus do not believe that these isolated SNPs correspond to true signals of selection and they are likely artifacts in the dataset. However, several other regions have a SNP whose p-value exceeds the BH threshold, and multiple nearby SNPs exhibiting low p-values, thus giving some support that these potentially correspond to real signals of selection. Figures 13A and 13B show the p-values in a genomic region with low p-values surrounding a significant SNP on chromosome 5 and a region surrounding a SNP with a spuriously low p-value on chromosome 7, respectively.

To remove spurious SNPs while keeping significant SNPs in regions that show additional support at surrounding SNPs, we post-process the p-values using a modified version of Brown's method (Brown 1975) for combining non-independent p-values. Specifically, we consider overlapping windows of 50 SNPs around each analyzed SNP. For each of these overlapping windows, we compute the negative sum of the logarithm of the p-values, including the focal SNP. We then fit the parameters of a scaled χ^2 distribution to these sums, resulting in a post-processed p-value for each SNP. We then apply the BH method to this new set of p-values to obtain a second BH threshold for each SNP. Isolated SNPs, such as the SNP in Figure 13B no longer exceed the corresponding BH threshold, while regions containing a significant SNP and additional support, such as the region in Figure 13A, have a broad peak exceeding the corresponding BH threshold.

After applying this post-processing procedure, we group significant p-values into distinct regions of significance. For a region to be considered significant, we require that at least one SNP within the region must have both a raw p-value and a post-processed p-value each exceeding its respective BH threshold. The post-processing procedure broadens p-value peaks; we therefore take each contiguous block of post-processed p-values exceeding the corresponding BH threshold as a separate candidate region. Under this criterion, for the one-dimensional *additive* EM-HMM algorithm, there are 8 distinct candidate regions. Table 1 lists these regions (in hg19 coordinates), any genes overlapping

Figure 13: A) Manhattan plot of p-values in genomic region on chromosome 5, centered around SNPs with p-value exceeding BH threshold. Surrounding SNPs exhibit low p-values, as expected under genetic hitchhiking. Post-processed p-values exceed the respective BH threshold. B) Manhattan plot of p-values in genomic region on chromosome 7, centered around SNP with p-value exceeding BH threshold. Surrounding SNPs do not show evidence of selection. Post-processed p-values do not show any significant signal. C) Binned allele frequency trajectories for 20 SNPs centered around SNP with lowest p-value in Figure 13A. Nearby SNPs show correlated allele frequency change, indicative of genetic hitchhiking and true signal. Size of points indicates number of samples; color hue indicates genomic position: red smaller, blue larger. D) Binned allele frequency trajectories for 20 SNPs centered around SNP with lowest p-value in Figure 13B. Nearby SNPs do not exhibit correlated allele frequency change, suggesting spurious signal at lead SNP.

Chr.	Genomic region (hg19)	Gene(s)	Lead SNP	Ref	Alt	# signif. SNPs		$-\log_{10} p_{min}$	$\hat{s}(p_{min})$
						Raw	Post		
2	135,161,631-137,087,324	LCT	rs4988235	G	A	63	395	16.11	0.080 (0.056, 0.101)
2	137,164,476-137,419,024	LCT	rs580879	C	T	1	76	5.24	-0.044 (-0.060, -0.024)
4	38,680,186-38,806,462	TLR10/1/6	rs10008492	C	T	1	54	7.09	-0.047 (-0.062, -0.028)
5	33,854,740-34,004,707	SLC45A2	rs16891982	C	G	4	51	7.08	0.049 (0.030, 0.066)
5	131,465,688-131,675,046	SLC22A4	rs7727544	C	T	2	67	6.58	0.042 (0.024, 0.057)
6	31,008,368-31,091,197	HLA	rs2535317	C	T	3	151	6.40	-0.046 (-0.062, -0.025)
7	100,212,254-100,361,675	TFR2	rs4434553	A	G	1	27	6.35	0.044 (0.024, 0.060)
**16	70,771,279-71,330,074	HYDIN	rs79233902	T	G	3	51	17.38	0.255**

Table 1: Genomic regions identified as significant under *additive* selection. We indicate genes that overlap the region, number of SNPs exceeding the BH threshold pre- and post-processing, $-\log_{10} p_{min}$, and $\hat{s}$ for the derived allele at the SNP with $-\log_{10} p_{min}$. The asterisks at the HYDIN locus indicate that this signal is likely an artifact.

the region, the rsID of the SNP corresponding to the minimal p-value in the region, the reference and alternative alleles at this SNP, the number of significant SNPs (pre- and post-processing) in each region, the negative logarithm of the minimal p-value in the region, and the additive $\hat{s}$ for the derived allele at the SNP with the lowest p-value with confidence intervals. Confidence intervals were obtained by simulating 1000 replicates matching the sampling scheme, estimated initial frequency, and estimated selection coefficient of each lead SNP, then bias-correcting all $\hat{s}$ values and taking the 0.025 and 0.975 quantiles of the simulated replicates. Additionally, we present Manhattan plots of the p-values and the binned allele frequency trajectories in each significant genomic region in Section S.9.

4.3.2 Functional genetic variation in significant genomic regions

With the exception of TFR2, all genomic regions that we detect as significant by applying our *additive* EM-HMM have previously been identified as targets of selection, although not all have been identified as such in populations from Great Britain. In the following paragraphs, we will discuss each of these regions in the context of the relevant literature.

The strongest signal in our analysis is the well-characterized LCT locus, which has been identified as a target of selection in populations from Great Britain (Mathieson and Terhorst 2022; Nait Saada et al. 2020; Field et al. 2016) and broader Western European populations (Bersaglieri et al. 2004; Itan et al. 2009; Mathieson, Lazaridis, et al. 2015). The lead SNP in our analysis, rs4988235, has previously been identified as the strongest signal of selection at the LCT locus (Itan et al. 2009; Peter et al. 2012; Mathieson and Terhorst 2022), and the derived allele at this SNP has been linked to the ability to digest lactase into adulthood (Enattah et al. 2002). Our estimated selection coefficient of 0.080 (CI: [0.056, 0.101]) is in line with other estimates provided in the literature (Peter et al. 2012; Mathieson and Terhorst 2022; Bersaglieri et al. 2004). Although the derived allele at this SNP has been linked to lactase persistence, recent studies argue that the introduction of milk consumption predates the increase in frequency of this allele, and that the recent strong selective pressure perhaps results from later famines where the allele proved beneficial (Burger et al. 2020; Evershed et al. 2022). Our approach also identified a secondary region of significance near the LCT locus which is likely a result of genetic hitchhiking.

Genomic variation in two of our candidate regions, TLR10/1/6 and HLA, is involved in immune

regulation. Polymorphisms in the TLR10/1/6 gene cluster have been linked to incidence of several cancers, as well as tuberculosis and leprosy (Purdue et al. 2009; Sun et al. 2006; Ma et al. 2007; Wong et al. 2010). The TLR10/1/6 locus has been identified as a target of selection in a previous study using ancient DNA (Mathieson, Lazaridis, et al. 2015), although this work pooled samples from West Eurasia. Similarly, using a dataset of present-day individuals, Barreiro et al. (2009) found that the TLR genes, with the exception of TLR1, experience strong negative selection, and that the TLR10/1/6 cluster has undergone recent selection in non-African populations.

The HLA locus spans a large region on chromosome 6 and encodes a set of highly polymorphic genes critical for the function of the innate immune system. Previous studies have identified multiple signals of selection in this region (Mathieson and Terhorst 2022; Field et al. 2016), and the SNPs we identify overlap with one of these signals. Individual loci within the HLA region are believed to be under balancing or frequency-dependent selection to increase allelic diversity (Hedrick and Thomson 1983; Bronson et al. 2013). For that reason, it is somewhat surprising that genomic scans for positive selection, here and in the literature (Mathieson and Terhorst 2022; Field et al. 2016), find signals at these loci. Additionally, our scan for *heterozygote difference*, which includes *overdominance*, a model of balancing selection, reveals no significant signals in the HLA region. The likely explanation is that several alleles changing in frequency are detected as positive selection, but the short time horizon considered here is not sufficient to detect alleles under long-term balancing selection.

The SLC45A2 locus has also been well-characterized as a target of selection in European populations (Lao et al. 2007; Beleza et al. 2013). Polymorphism at this locus is associated with differences in hair and skin pigmentation (Soejima and Koda 2007; Hysi et al. 2018). Studies using ancient DNA data from both Western Europe broadly and Great Britain specifically have identified the SLC45A2 locus as a target of selection (Mathieson, Lazaridis, et al. 2015; Mathieson and Terhorst 2022). In addition, our estimated selection coefficient at the lead SNP matches values obtained from both analyses of present-day and ancient DNA: Our MLE is $s = 0.049$ (CI: [0.030, 0.066]), Beleza et al. (2013) estimate s in the range of 0.04 to 0.05, and Mathieson and Terhorst (2022) estimate $s = 0.043$.

The gene SLC22A4 is contained within the larger IBD5 locus, which consists of a group of genes with polymorphisms linked to gastrointestinal disorders such as Crohn's disease (Fisher et al. 2006). Huff et al. (2012) found that genetic variants in SLC22A4 increases absorption of the antioxidant ergothioneine and show signals of positive selection, likely due to the low amounts of ergothioneine in early Neolithic farmer diets. Furthermore, they argue that variants linked to Crohn's disease likely increased in frequency via genetic hitchhiking. The SLC22A4/IBD5 locus was also identified as a target of selection using ancient DNA from Western Europeans (Mathieson, Lazaridis, et al. 2015).

Genetic variation at the TFR2 locus has not been identified as a target of selection using either contemporary or ancient genomic samples. Figure S.29 in the Supplementary Material shows that the allele frequency at the lead SNP and surrounding SNPs shift in concert indicating that this is potentially a real target of selection rather than a false signal. Mutations at the TFR2 locus cause type 3 hereditary hemochromatosis, which is characterized by abnormally high systemic iron levels (Girelli et al. 2002). In addition, a haplotype that includes the variant at the lead SNP we identify as under selection has been correlated with Parkinson's disease (Rhodes et al. 2014).

The HYDIN locus contains the SNP with the most significant p-value in our dataset. However, upon inspection of the p-values in this genomic region, provided in Figure S.30A in the Supplemen-

tary Material, we observe that the locus indeed contains three SNPs with very low p-values, but also two 100 kbp regions without any SNPs. These empty regions are a result of our filtering procedure, which removed a large number of SNPs with minor allele frequency below 0.05. In addition, the significant SNPs at the HYDIN locus have an extremely low binned minor allele frequency at all but the last timepoint, see Figure S.30B in the Supplementary Material. Due to the low number of SNPs with substantial frequencies, unusual allele frequency trajectories, and lack of a clear signal of hitchhiking, we believe that the signal of selection at the HYDIN locus is spurious.

Lastly, we more explicitly compare the results of our analysis to those of Mathieson and Terhorst (2022), a recent study that identified targets of selection in a temporal dataset similar to ours. The authors analyze present-day and ancient DNA samples from the AADR localized to England, dated to under 4,450 years BP, and find signals of selection in five genomic regions. Three regions (LCT, SLC45A2, HLA) are identified in our study as well, three regions (TLR10/1/6, SLC22A4, TFR2) are only identified in our analysis, and two regions (DHCR7, HERC2) are only identified by Mathieson and Terhorst. Of the loci identified in both studies, the estimates of the selection coefficients at the most significant SNPs largely agree: the coefficients in the LCT, SLC45A2, and HLA regions are 0.080 (CI: [0.056, 0.101]), 0.049 (CI: [0.030, 0.066]), and 0.046 (CI: [0.025, 0.062]) in our study and 0.064, 0.043, and 0.046 in Mathieson and Terhorst's study, respectively. Figures S.31 and S.32 in the Supplementary Material show the p-values computed using our *additive* EM-HMM and binned allele frequency trajectories in both regions identified as significant by Mathieson and Terhorst (2022) that are not significant in our analysis. At the DHCR7 locus on chromosome 11, the post-processed p-value are close to exceeding the BH threshold, but no raw p-value reaches significance. The HERC2 locus on chromosome 15 also exhibits low p-values, although to a lesser degree than DHCR7. In contrast to our conservative approach of only analyzing samples genotyped using the 1240K assay, Mathieson and Terhorst also analyze samples genotyped using whole genome data (including present-day samples), use a method that has the potential to detect selection where the *additive* coefficient changes over time, and use a different post-processing when combining signals at neighboring SNPs. Thus, a perfect alignment of signals should not be expected.

4.3.3   Inferring the mode of selection in the GB aDNA dataset

In addition to the *additive* EM-HMM, we analyzed the GB aDNA dataset under each of the other three one-parameter modes of selection – *dominant*, *recessive*, and *heterozygote difference*, see Section S.7 in the Supplementary Material, where we provide full Manhattan plots for each one-parameter mode of selection as well as tables analogous to Table 1. Additionally, we provide a table of genome-wide Spearman rank correlation coefficients between the log-likelihood ratio statistics of all one-parameter modes for the top 1% of SNPs by additive p-value in Figure S.21 of the Supplementary Material. Both the *dominant* and the *recessive* EM-HMM have high correlation with the *additive* EM-HMM, and identify the same regions as the *additive* analysis, although the LCT locus is not split into two distinct regions in the *dominant* analysis.

In contrast, the *heterozygote difference* EM-HMM has a lower correlation with the other modes and only identifies LCT, the two SLC loci, and HYDIN as significant. The inability of this mode to detect selection at the HLA locus is at first somewhat surprising, given that balancing selection or frequency-dependent selection should result in a dynamic similar to *overdominant* selection. However, the signature that would provide strong support for these types of selection would be alleles that remain at intermediate frequencies longer than expected under neutrality, and thus the short time-horizon considered here is likely not sufficient.

We also used the procedure detailed in Section 2.3.2 to infer the most likely mode of selection at each SNP. Specifically, we used the $10{,}000$ neutral and $200{,}000$ non-neutral replicates from the data-matched simulations detailed in Section 4.2 to identify k_{opt} and fit the generalized gamma distribution to the test statistic δ_{ll}. We computed δ_{ll} for all SNPs in the GB aDNA dataset to obtain a p-value for each locus. We then used the BH procedure to determine a p-value threshold at an FDR of $\alpha = 0.05$. SNPs that did not exceed the threshold were classified as neutral. SNPs exceeding the threshold were classified by the mode of selection with the highest log-likelihood, appropriately correcting for the difference in the parametrizations using k_{opt}.

Despite several significant regions identified using the one-parameter modes, the mode inference procedure only identifies LCT and HYDIN as significant. We do note, however, that when using a less stringent FDR of $\alpha = 0.1$, two more regions, TLR10/1/6 and SLC45A2, also exceed the BH thresholds. As detailed in Section 4.3.2, we believe that the signal at the HYDIN locus is spurious. Figure 14 shows the resulting raw p-values at the LCT locus, with significant SNPs colored by their inferred selection mode. Out of the 31 SNPs exceeding the BH threshold in this region, 18 are classified as *additive*, but the other SNPs show different selection modes. The fact that a majority of loci in this region are classified as *additive* could indicate weak support for the hypothesis that LCT is evolving under *additive* selection. However, the inferred mode at the lead SNP rs4988235 is "indeterminate", and the selection coefficients $(\hat{s_1}, \hat{s_2})$ inferred using the unconstrained diploid EM-HMM are $(0.055, 0.073)$, indicating that the selection dynamics would be intermediate between *additive* and *dominance*. We do caution against over-interpretation, as the simulation study in Section 4.2 shows that identifying the mode of selection is in general challenging.

We believe that, given the difficulty of inferring the mode of selection, the fact that our procedure is conservative in identifying significant regions is desirable. Additionally, we note that the one-parameter mode with the highest log-likelihood ratio at the lead SNP for each identified region is as follows: LCT – *dominance*, TLR10/1/6 – *additive*, SLC45A2 – *heterozygote difference*, SLC22A4 – *dominance*, HLA – *recessive*, and TFR2 – *additive*, although we again caution against over-interpretation of these results.

5 Discussion

In this work, we presented a novel method to compute maximum likelihood estimates (MLEs) for general diploid selection coefficients from time series genetic data. To this end, we extended the framework of Mathieson and McVean (2013) for the *additive* case and derived an EM-HMM algorithm to estimate the parameters of diploid selection. We show that the diploid EM-HMM framework can also be constrained to bespoke one-parameter models of selection via the method of Lagrange multipliers. We furthermore introduced a novel likelihood-based procedure, motivated by the AIC, for inferring the best fitting diploid mode of selection from temporal data between *additive*, *recessive*, *dominant*, and *over-* or *underdominant* selection. To our knowledge, our study is the first to address the statistical problem of inferring the mode of selection from given time series genetic data. To further improve power to detect selection and remove spurious signals, we also introduced a procedure based on Brown's method to combine p-values across linked loci.

Using simulation studies, we show that the estimated selection coefficients are accurate across a range of selection parameters, population parameters, and sampling schemes. However, we find that determining the mode of selection from time series data is challenging, and only yields reliable results when selection is strong. We also demonstrate that assuming *additive* selection when

Figure 14: Manhattan plot of raw p-values at the LCT locus. P-values are computed using the procedure to identify the mode of selection described in Section 2.3.2, and significant SNPs are colored by inferred mode. The majority of SNPs in this region are classified as *additive*, although the lead SNP is classified as indeterminate.

analyzing data simulated under different modes of selection yields comparable power to reject neutrality as when the data is simulated under *additive* selection, implying that analyzing given data assuming *additive* selection may be sufficient for selection scans. However, the estimated selection coefficients are inaccurate when the mode is misspecified. We apply our method to time series genetic data obtained from 504 ancient individuals in the AADR from Great Britain dated to under 4,450 BP, and identify six genomic regions with signals of selection. These regions, except TFR2, have been identified as targets of selection in previous studies, and we discuss them in the context of the relevant literature. These regions are identified as significant under multiple directional selection modes (*additive, recessive, dominant*). When classifying the mode of selection from the data, however, the results are inconclusive: We find that a majority of the SNPs at the LCT locus provide evidence for *additive* selection, but the lead SNPs is classified as "indeterminate," and the point estimate is intermediate between *additive* and *dominant* selection.

Note that our algorithm identifies the MLEs from a given temporal datasets with high accuracy and computes the correct likelihoods. Thus, the statistical properties of the MLEs and the likelihood-ratio tests exhibited in our simulation studies in Section 3 and Section 4.2 are not exclusive to our method but characterize the MLEs and the power of the respective tests in general under the given population genetic model, regardless whether our method or a different method is used for the computation. We characterize the statistical properties in a range of scenarios, but if these scenarios do not cover the exact scheme encountered in a specific dataset, our simulation framework can be readily modified to characterize the statistical properties in the respective scenario. Naturally, the power to identify and characterize selection does depend on the exact sampling scheme: Strong selection can be readily identified, even when samples are limited to a short time period. However, weaker selection requires sampling more data over a longer time period.

Our method computes the MLEs of general diploid selection parameters, and we believe that this is useful to researchers in at least two regards: (1) Our approach can be used to infer the mode of selection from a given temporal dataset. While we demonstrate that selection needs to be fairly strong for reliable classification, our framework can be used to characterize statistical power in a given scenario, and determine whether additional samples at potentially additional timepoints are necessary. (2) If the selection mode operating on the genetic variants in a given temporal dataset is known a priori, our method enables researchers to estimate the selection coefficients accurately under the correct model. We demonstrate that assuming the wrong mode of selection yields inaccurate estimates.

In practice, we recommend the following approach to analyze given time series genetic data, potentially at a large number of loci: If computational resources are limited, researchers should apply the *additive* EM-HMM to obtain MLEs of the additive selection coefficient at each locus, and use standard likelihood-ratio testing to identify outliers. As detailed in Section 3.3.2, the likelihood-ratio test under *additive* selection can identify non-neutral replicates, even if the mode of selection is misspecified. The analysis of the 743,417 SNPs for 504 samples over 125 generations in our GB aDNA dataset took roughly 15 hours parallelized across chromosomes. If additional computational resources are available, we recommend analyzing the data under each bespoke one-parameter selection mode, as well as the unconstrained mode, to characterize signals in the data that are not correctly described by *additive* selection. The results can then also be used to identify the mode of selection from the data, but as we demonstrate, accuracy is limited. In addition, we strongly recommend performing a data-matched simulation study, as presented in Section 4.2. Such data-matched simulations enable exact characterization of the statistical power and accuracy of the approach in the specific scenario, and is also required to fine-tune the parameters for inferring the mode of selection.

As for similar approaches, the HMM underlying our method assumes that the population is panmictic, and violations of this assumption can dilute signals or introduce spurious signals. Future work to address this shortcoming could proceed along at least two possible avenues: (1) Controlling for population structure by using a Principal Component Analysis as a covariate in the estimation procedure directly (Luu et al. 2017; Ju and Mathieson 2021), or (2) explicitly including the population structure and exchange of migrants in the underlying population genetic model (Joseph and Pe'er 2019).

Moreover, our approach estimates selection coefficients from temporal data at the focal locus only, and does not incorporate the allele frequency dynamics at linked loci. In our analysis of the GB aDNA dataset, we do leverage signals across loci using a novel post-processing approach to combine p-values in a genomic window. This post-processing can reduce signal-to-noise ratios in genomic scans for selection in general. However, incorporating the genetic variation at multiple SNPs using a proper likelihood model for the multi-locus dynamics under the Wright-Fisher process has the potential to account for chromosomal linkage more accurately and result in a more robust inference (Terhorst et al. 2015; He et al. 2020).

While we focus on analyzing time series genetic data at a single locus in this study, the capacity of our method to characterize selection modes more general than *additive* only also has potential benefits when studying polygenic selection on complex traits: In models of stabilizing polygenic selection around around an optimal trait, the genetic variants affecting the trait experience *underdominant* selection dynamics, which can be readily addressed using our framework.

Acknowledgements

We want to thank Xinyi Li, Xiaoheng Cheng, Constanza de la Fuente, and Maanasa Raghavan for helpful comments on the method and the data analysis. Moreover, we thank Jeremy Berg, Maryn Carlson and Rowan Hart for comments on the manuscript. In addition, we thank the members of the Raghavan, Berg, and Novembre labs for valuable feedback throughout the project. A.F. was supported by the Department of Education, Graduate Assistance in Areas of National Need (GAANN), Grant #P200A210054, and this work was supported by the National Institute of General Medical Sciences (NIGMS) of the National Institutes of Health under award R01GM146051.

References

Barghi, Neda, Raymond Tobler, Viola Nolte, Ana Marija Jakšić, François Mallard, Kathrin Anna Otte, Marlies Dolezal, Thomas Taus, Robert Kofler, and Christian Schlötterer (2019). "Genetic Redundancy Fuels Polygenic Adaptation in Drosophila". *PLOS Biology* 17(2), e3000128.

Barreiro, Luis B., Meriem Ben-Ali, Hélène Quach, Guillaume Laval, Etienne Patin, Joseph K. Pickrell, Christiane Bouchier, Magali Tichit, Olivier Neyrolles, Brigitte Gicquel, Judith R. Kidd, Kenneth K. Kidd, Alexandre Alcaïs, Josiane Ragimbeau, Sandra Pellegrini, Laurent Abel, Jean-Laurent Casanova, and Lluís Quintana-Murci (2009). "Evolutionary Dynamics of Human Toll-Like Receptors and Their Different Contributions to Host Defense". *PLOS Genetics* 5(7), e1000562.

Barton, Nick (1986). "The Maintenance of Polygenic Variation through a Balance between Mutation and Stabilizing Selection". *Genetical Research* 47(3), pp. 209–216.

Beleza, Sandra, António M. Santos, Brian McEvoy, Isabel Alves, Cláudia Martinho, Emily Cameron, Mark D. Shriver, Esteban J. Parra, and Jorge Rocha (2013). "The Timing of Pigmentation Lightening in Europeans". *Molecular Biology and Evolution* 30(1), pp. 24–35.

Benjamini, Yoav and Yosef Hochberg (1995). "Controlling the False Discovery Rate: A Practical and Powerful Approach to Multiple Testing". *Journal of the Royal Statistical Society: Series B (Methodological)* 57(1), pp. 289–300.

Bersaglieri, Todd, Pardis C. Sabeti, Nick Patterson, Trisha Vanderploeg, Steve F. Schaffner, Jared A. Drake, Matthew Rhodes, David E. Reich, and Joel N. Hirschhorn (2004). "Genetic Signatures of Strong Recent Positive Selection at the Lactase Gene". *American Journal of Human Genetics* 74(6), pp. 1111–1120.

Bignell, Graham R. et al. (2010). "Signatures of Mutation and Selection in the Cancer Genome". *Nature* 463, pp. 893–898.

Bishop, Christopher M. (2006). *Pattern Recognition and Machine Learning.* Information Science and Statistics. New York: Springer.

Bollback, Jonathan P., Thomas L. York, and Rasmus Nielsen (2008). "Estimation of 2Nes from Temporal Allele Frequency Data". *Genetics* 179(1), pp. 497–502.

Bozdogan, Hamparsum (1987). "Model Selection and Akaike's Information Criterion (AIC): The General Theory and Its Analytical Extensions". *Psychometrika* 52(3), pp. 345–370.

Bronson, Paola G., Steven J. Mack, Henry A. Erlich, and Montgomery Slatkin (2013). "A Sequence-Based Approach Demonstrates That Balancing Selection in Classical Human Leukocyte Antigen (HLA) Loci Is Asymmetric". *Human Molecular Genetics* 22(2), pp. 252–261.

Brown, Morton B. (1975). "A Method for Combining Non-Independent, One-Sided Tests of Significance". *Biometrics* 31(4), pp. 987–992.

Burger, Joachim et al. (2020). "Low Prevalence of Lactase Persistence in Bronze Age Europe Indicates Ongoing Strong Selection over the Last 3,000 Years". *Current Biology* 30(21), 4307–4315.e13.

Bustamante, Carlos D., Adi Fledel-Alon, Scott Williamson, Rasmus Nielsen, Melissa Todd Hubisz, Stephen Glanowski, David M. Tanenbaum, Thomas J. White, John J. Sninsky, Ryan D. Hernandez, Daniel Civello, Mark D. Adams, Michele Cargill, and Andrew G. Clark (2005). "Natural Selection on Protein-Coding Genes in the Human Genome". *Nature* 437, pp. 1153–1157.

Cheng, Xiaoheng and Matthias Steinrücken (2023). "diplo-locus: A lightweight toolkit for inference and simulation of time-series genetic data under general diploid selection". *bioRxiv*. https://doi.org/10.1101/2023.10.12.562101.

Chintalapati, Manjusha, Nick Patterson, and Priya Moorjani (2022). "The Spatiotemporal Patterns of Major Human Admixture Events during the European Holocene". *eLife* 11, e77625.

Dehasque, Marianne, María C. Ávila-Arcos, David Díez-del-Molino, Matteo Fumagalli, Katerina Guschanski, Eline D. Lorenzen, Anna-Sapfo Malaspinas, Tomas Marques-Bonet, Michael D. Martin, Gemma G. R. Murray, Alexander S. T. Papadopulos, Nina Overgaard Therkildsen, Daniel Wegmann, Love Dalén, and Andrew D. Foote (2020). "Inference of Natural Selection from Ancient DNA". *Evolution Letters* 4(2), pp. 94–108.

De Vladar, Harold P. and Nick Barton (2014). "Stability and Response of Polygenic Traits to Stabilizing Selection and Mutation". *Genetics* 197(2), pp. 749–767.

Enattah, Nabil Sabri, Timo Sahi, Erkki Savilahti, Joseph D. Terwilliger, Leena Peltonen, and Irma Järvelä (2002). "Identification of a Variant Associated with Adult-Type Hypolactasia". *Nature Genetics* 30(2), pp. 233–237.

Evershed, Richard P. et al. (2022). "Dairying, Diseases and the Evolution of Lactase Persistence in Europe". *Nature* 608, pp. 336–345.

Ewens, Warren J. (2004). *Mathematical Population Genetics*. 2nd ed. Vol. I. Theoretical Introduction. Springer.

Field, Yair, Evan A Boyle, Natalie Telis, Ziyue Gao, Kyle J. Gaulton, David Golan, Loic Yengo, Ghislain Rocheleau, Philippe Froguel, Mark I. McCarthy, and Jonathan K. Pritchard (2016). "Detection of Human Adaptation during the Past 2000 Years". *Science* 354, pp. 760–764.

Fisher, Sheila A., Jochen Hampe, Clive M. Onnie, Mark J. Daly, Christine Curley, Shaun Purcell, Jeremy Sanderson, John Mansfield, Vito Annese, Alastair Forbes, Cathryn M. Lewis, Stefan Schreiber, John D. Rioux, and Christopher G. Mathew (2006). "Direct or Indirect Association in a Complex Disease: The Role of SLC22A4 and SLC22A5 Functional Variants in Crohn Disease". *Human Mutation* 27(8), pp. 778–785.

Foll, Matthieu, Hyunjin Shim, and Jeffrey D. Jensen (2015). "WFABC: a Wright–Fisher ABC-based approach for inferring effective population sizes and selection coefficients from time-sampled data". *Molecular Ecology Resources* 15(1), pp. 87–98.

Gemmell, Neil J. and Jon Slate (2006). "Heterozygote Advantage for Fecundity". *PLOS ONE* 1(1), e125.

Girelli, Domenico, Claudia Bozzini, Antonella Roetto, Federica Alberti, Filomena Daraio, Romano Colombari, Oliviero Olivieri, Roberto Corrocher, and Clara Camaschella (2002). "Clinical and Pathologic Findings in Hemochromatosis Type 3 Due to a Novel Mutation in Transferrin Receptor 2 Gene". *Gastroenterology* 122(5), pp. 1295–1302.

Gretzinger, Joscha et al. (2022). "The Anglo-Saxon Migration and the Formation of the Early English Gene Pool". *Nature* 610, pp. 112–119.

He, Zhangyi, Xiaoyang Dai, Mark Beaumont, and Feng Yu (2020). "Detecting and Quantifying Natural Selection at Two Linked Loci from Time Series Data of Allele Frequencies with Forward-in-Time Simulations". *Genetics* 216, pp. 521–541.

Hedrick, Philip W (2012). "What Is the Evidence for Heterozygote Advantage Selection?" *Trends in Ecology & Evolution* 27(12), pp. 698–704.

Hedrick, Philip W and Glenys Thomson (1983). "Evidence for balancing selection at HLA". *Genetics* 104(3), pp. 449–456.

Hoffmann, Laurence D., Gerald L. Bradley, and Kenneth H. Rosen (2010). *Applied Calculus for Business, Economics, and the Social and Life Sciences.* Expanded 10th ed. / Laurence D. Hoffmann, Gerald L. Bradley. New York, NY: McGraw-Hill.

Hofreiter, Michael, Johanna L. A. Paijmans, Helen Goodchild, Camilla F. Speller, Axel Barlow, Gloria G. Fortes, Jessica A. Thomas, Arne Ludwig, and Matthew J. Collins (2015). "The Future of Ancient DNA: Technical Advances and Conceptual Shifts". *BioEssays* 37(3), pp. 284–293.

Huff, Chad D., David J. Witherspoon, Yuhua Zhang, Chandler Gatenbee, Lee A. Denson, Subra Kugathasan, Hakon Hakonarson, April Whiting, Chadwick T. Davis, Wilfred Wu, Jinchuan Xing, W. Scott Watkins, Michael J. Bamshad, Jonathan P. Bradfield, Kazima Bulayeva, Tatum S. Simonson, Lynn B. Jorde, and Stephen L. Guthery (2012). "Crohn's Disease and Genetic Hitchhiking at IBD5". *Molecular Biology and Evolution* 29(1), pp. 101–111.

Hysi, Pirro G. et al. (2018). "Genome-Wide Association Meta-Analysis of Individuals of European Ancestry Identifies New Loci Explaining a Substantial Fraction of Hair Color Variation and Heritability". *Nature Genetics* 50(5), pp. 652–656.

Iranmehr, Arya, Ali Akbari, Christian Schlötterer, and Vineet Bafna (2017). "Clear: Composition of Likelihoods for Evolve and Resequence Experiments". *Genetics* 206(2), pp. 1011–1023.

Itan, Yuval, Adam Powell, Mark A. Beaumont, Joachim Burger, and Mark G. Thomas (2009). "The Origins of Lactase Persistence in Europe". *PLOS Computational Biology* 5(8), e1000491.

Joseph, Tyler A. and Itsik Pe'er (2019). "Inference of Population Structure from Time-Series Genotype Data". *The American Journal of Human Genetics* 105(2), pp. 317–333.

Ju, Dan and Iain Mathieson (2021). "The evolution of skin pigmentation-associated variation in West Eurasia". *Proceedings of the National Academy of Sciences* 118(1), e2009227118.

Lachance, Joseph and Sarah A. Tishkoff (2013). "Population Genomics of Human Adaptation". *Annual Review of Ecology, Evolution, and Systematics* 44(1), pp. 123–143.

Lao, O., J. M. de Gruijter, K. van Duijn, A. Navarro, and M. Kayser (2007). "Signatures of Positive Selection in Genes Associated with Human Skin Pigmentation as Revealed from Analyses of Single Nucleotide Polymorphisms". *Annals of Human Genetics* 71, pp. 354–369.

Luu, Keurcien, Eric Bazin, and Michael G. B. Blum (2017). "pcadapt: an R package to perform genome scans for selection based on principal component analysis". *Molecular Ecology Resources* 17(1), pp. 67–77.

Ma, Xin, Yuhua Liu, Brian B. Gowen, Edward A. Graviss, Andrew G. Clark, and James M. Musser (2007). "Full-Exon Resequencing Reveals Toll-Like Receptor Variants Contribute to Human Susceptibility to Tuberculosis Disease". *PLOS ONE* 2(12), e1318.

Malaspinas, Anna-Sapfo (2016). "Methods to Characterize Selective Sweeps Using Time Serial Samples: An Ancient DNA Perspective". *Molecular Ecology* 25(1), pp. 24–41.

Malaspinas, Anna-Sapfo, Orestis Malaspinas, Steven N. Evans, and Montgomery Slatkin (2012). "Estimating Allele Age and Selection Coefficient from Time-Serial Data". *Genetics* 192(2), pp. 599–607.

Mallick, Swapan, Adam Micco, Matthew Mah, Harald Ringbauer, Iosif Lazaridis, Iñigo Olalde, Nick Patterson, and David Reich (2024). "The Allen Ancient DNA Resource (AADR) a curated compendium of ancient human genomes". *Scientific Data* 11(1), p. 182.

Mathews, John and Kurtis Fink (2003). *Numerical Methods Using Matlab*. 4th edition. Upper Saddle River, N.J: Pearson.

Mathieson, Iain, Iosif Lazaridis, et al. (2015). "Genome-Wide Patterns of Selection in 230 Ancient Eurasians". *Nature* 528, pp. 499–503.

Mathieson, Iain and Gil McVean (2013). "Estimating Selection Coefficients in Spatially Structured Populations from Time Series Data of Allele Frequencies". *Genetics* 193(3), pp. 973–984.

Mathieson, Iain and Jonathan Terhorst (2022). "Direct Detection of Natural Selection in Bronze Age Britain". *Genome Research*, gr.276862.122.

Moorjani, Priya, Sriram Sankararaman, Qiaomei Fu, Molly Przeworski, Nick Patterson, and David Reich (2016). "A Genetic Method for Dating Ancient Genomes Provides a Direct Estimate of Human Generation Interval in the Last 45,000 Years". *Proceedings of the National Academy of Sciences* 113(20), pp. 5652–5657.

Nait Saada, Juba, Georgios Kalantzis, Derek Shyr, Fergus Cooper, Martin Robinson, Alexander Gusev, and Pier Francesco Palamara (2020). "Identity-by-Descent Detection across 487,409 British Samples Reveals Fine Scale Population Structure and Ultra-Rare Variant Associations". *Nature Communications* 11(1), p. 6130.

Nielsen, Rasmus (2005). "Molecular Signatures of Natural Selection". *Annual Review of Genetics* 39(1), pp. 197–218.

Olalde, Iñigo et al. (2018). "The Beaker Phenomenon and the Genomic Transformation of Northwest Europe". *Nature* 555, pp. 190–196.

Orlando, Ludovic, Robin Allaby, Pontus Skoglund, Clio Der Sarkissian, Philipp W. Stockhammer, María C. Ávila-Arcos, Qiaomei Fu, Johannes Krause, Eske Willerslev, Anne C. Stone, and Christina Warinner (2021). "Ancient DNA Analysis". *Nature Reviews Methods Primers* 1(1), pp. 1–26.

Palmer, Duncan S., Wei Zhou, Liam Abbott, Emilie M. Wigdor, Nikolas Baya, Claire Churchhouse, Cotton Seed, Tim Poterba, Daniel King, Masahiro Kanai, Alex Bloemendal, and Benjamin M. Neale (2023). "Analysis of Genetic Dominance in the UK Biobank". *Science* 379, pp. 1341–1348.

Patterson, Nick et al. (2022). "Large-Scale Migration into Britain during the Middle to Late Bronze Age". *Nature* 601, pp. 588–594.

Peter, Benjamin M., Emilia Huerta-Sanchez, and Rasmus Nielsen (2012). "Distinguishing between Selective Sweeps from Standing Variation and from a De Novo Mutation". *PLOS Genetics* 8(10), e1003011.

Purdue, Mark P. et al. (2009). "A Pooled Investigation of Toll-like Receptor Gene Variants and Risk of Non-Hodgkin Lymphoma". *Carcinogenesis* 30(2), pp. 275–281.

Rabiner, L.R. (1989). "A Tutorial on Hidden Markov Models and Selected Applications in Speech Recognition". *Proceedings of the IEEE* 77(2), pp. 257–286.

Rhodes, Shannon L., Daniel D. Buchanan, Ismaïl Ahmed, Kent D. Taylor, Marie-Anne Loriot, Janet S. Sinsheimer, Jeff M. Bronstein, Alexis Elbaz, George D. Mellick, Jerome I. Rotter, and

Beate Ritz (2014). "Pooled Analysis of Iron-Related Genes in Parkinson's Disease: Association with Transferrin". *Neurobiology of Disease* 62, pp. 172–178.

Sanjak, Jaleal S., Julia Sidorenko, Matthew R. Robinson, Kevin R. Thornton, and Peter M. Visscher (2018). "Evidence of directional and stabilizing selection in contemporary humans". *Proceedings of the National Academy of Sciences* 115(1), pp. 151–156.

Sawyer, S. A. and D. L. Hartl (1992). "Population Genetics of Polymorphism and Divergence". *Genetics* 132(4), pp. 1161–1176.

Schlötterer, C., R. Kofler, E. Versace, R. Tobler, and S. U. Franssen (2015). "Combining Experimental Evolution with Next-Generation Sequencing: A Powerful Tool to Study Adaptation from Standing Genetic Variation". *Heredity* 114(5), pp. 431–440.

Schraiber, Joshua G., Steven N. Evans, and Montgomery Slatkin (2016). "Bayesian Inference of Natural Selection from Allele Frequency Time Series". *Genetics* 203(1), pp. 493–511.

Simons, Yuval B., Kevin Bullaughey, Richard R. Hudson, and Guy Sella (2018). "A population genetic interpretation of GWAS findings for human quantitative traits". *PLOS Biology* 16(3), e2002985.

Smith, John Maynard and John Haigh (1974). "The Hitch-Hiking Effect of a Favourable Gene". *Genetics Research* 23(1), pp. 23–35.

Soejima, Mikiko and Yoshiro Koda (2007). "Population Differences of Two Coding SNPs in Pigmentation-Related Genes SLC24A5 and SLC45A2". *International Journal of Legal Medicine* 121(1), pp. 36–39.

Stacy, E. W. (1962). "A Generalization of the Gamma Distribution". *The Annals of Mathematical Statistics* 33(3), pp. 1187–1192.

Steinrücken, Matthias, Anand Bhaskar, and Yun S. Song (2014). "A Novel Spectral Method for Inferring General Diploid Selection from Time Series Genetic Data". *The Annals of Applied Statistics* 8(4), pp. 2203–2222.

Sun, Jielin, Fredrik Wiklund, Fang-Chi Hsu, Katarina Bälter, S. Lilly Zheng, Jan-Erik Johansson, Baoli Chang, Wennuan Liu, Tao Li, Aubrey R. Turner, Liwu Li, Ge Li, Hans-Olov Adami, William B. Isaacs, Jianfeng Xu, and Henrik Grönberg (2006). "Interactions of Sequence Variants in Interleukin-1 Receptor–Associated Kinase4 and the Toll-Like Receptor 6-1-10 Gene Cluster Increase Prostate Cancer Risk". *Cancer Epidemiology, Biomarkers & Prevention* 15(3), pp. 480–485.

Tataru, Paula, Maria Simonsen, Thomas Bataillon, and Asger Hobolth (2017). "Statistical Inference in the Wright–Fisher Model Using Allele Frequency Data". *Systematic Biology* 66(1), e30–e46.

Taus, Thomas, Andreas Futschik, and Christian Schlötterer (2017). "Quantifying Selection with Pool-Seq Time Series Data". *Molecular Biology and Evolution* 34(11), pp. 3023–3034.

Terhorst, Jonathan, Christian Schlötterer, and Yun S. Song (2015). "Multi-Locus Analysis of Genomic Time Series Data from Experimental Evolution". *PLOS Genetics* 11(4), e1005069.

Vitti, Joseph J., Sharon R. Grossman, and Pardis C. Sabeti (2013). "Detecting Natural Selection in Genomic Data". *Annual Review of Genetics* 47(1), pp. 97–120.

Vlachos, Christos, Claire Burny, Marta Pelizzola, Rui Borges, Andreas Futschik, Robert Kofler, and Christian Schlötterer (2019). "Benchmarking Software Tools for Detecting and Quantifying Selection in Evolve and Resequencing Studies". *Genome Biology* 20(1), p. 169.

Wang, J. (2001). "A Pseudo-Likelihood Method for Estimating Effective Population Size from Temporally Spaced Samples". *Genetical Research* 78(3), pp. 243–257.

Watterson, G. A. (1982). "Testing Selection at a Single Locus". *Biometrics* 38(2), p. 323.

Wilks, S. S. (1938). "The Large-Sample Distribution of the Likelihood Ratio for Testing Composite Hypotheses". *The Annals of Mathematical Statistics* 9(1), pp. 60–62.

Wong, Sunny H. et al. (2010). "Leprosy and the Adaptation of Human Toll-Like Receptor 1". *PLOS Pathogens* 6(7), e1000979.