Epigenetic upregulation of carotid body angiotensin signaling increases blood pressure.

Fengli Zhu M.Sc.[1], Zhuqing Wang Ph.D.[1], Kayla Davis B.Sc.[1], Hayden McSwiggin B.Sc.[1],

Jekaterina Zyuzin Ph.D.[1], Jie Liu Ph.D.[1], Wei Yan M.D. Ph.D.[1, 2, 5], Virender K. Rehan M.D.[1, 3, 5],

Nicholas Jendzjowsky Ph.D.[1, 4, 5*]

[1]The Lundquist Institute for Biomedical Innovation at Harbor-UCLA Medical Center, Torrance,
CA 90502, USA
[2]Division of Metabolic Diseases and Translational Genomics, Harbor-UCLA Medical Center,
Torrance, CA 90502
[3]Division of Neonatology, Harbor-UCLA Medical Center, Torrance, CA 90502, USA
[4] Division of Respiratory and Critical Care Medicine and Physiology, Harbor-UCLA Medical
Center, Torrance, CA 90502, USA
[5]David Geffen School of Medicine, UCLA, Los Angeles, CA 90073, USA

Short title: Epigenetic programming of carotid bodies

*Corresponding Author: Nicholas Jendzjowsky, PhD
Rm 209 Martin Research Building,
The Lundquist Institute for Biomedical Innovation at Harbor-UCLA Medical Center
Division of Respiratory and Critical Care Medicine and Physiology
David Geffen School of Medicine, UCLA
1124 W Carson Street
Torrance, CA
90502-2006
Office:323-457-1663
Email: nicholas.jendzjowsky@lundquist.org
24

25   **Total word count: 8853**

Abstract

Epigenetic changes can be shaped by a wide array of environmental cues as well as maternal health and behaviors. One of the most detrimental behaviors to the developing fetus is nicotine exposure. Perinatal nicotine exposure remains a significant risk factor for cardiovascular health and in particular, hypertension. Increased basal carotid body activity and excitation are significant contributors to hypertension. This study investigated the epigenetic changes to carotid body activity induced by perinatal nicotine exposure resulting in carotid body-mediated hypertension. Using a rodent model of perinatal nicotine exposure, we show that angiotensin II type 1 receptor is upregulated in the carotid bodies of nicotine-exposed offspring. These changes were attributed to an upregulation of genetic promotion as DNA methylation of angiotensin II type 1 receptor occurred within intron regions, exemplifying an upregulation of genetic transcription for these genes. Nicotine increased angiotensin signaling *in vitro*. Carotid body reactivity to angiotensin was increased in perinatal nicotine-exposed offspring compared to control offspring. Further, carotid body denervation reduced arterial pressure as a result of suppressed efferent sympathetic activity in perinatal nicotine-exposed offspring. Our data demonstrate that perinatal nicotine exposure adversely affects carotid body afferent sensing, which augments efferent sympathetic activity to increase vasoconstrictor signaling and induce hypertension. Targeting angiotensin signaling in the carotid bodies may provide a way to alleviate hypertension acquired by adverse maternal uterine environments in general and perinatal nicotine exposure in particular.

Abstract word count: 229

Key words: Carotid body, sympathetic nervous activity, angiotensin, nicotine, epigenetic, developmental programming

Introduction

Adverse developmental programming is directly related to cardiovascular disease in later life. The epigenetic modifications that occur in response to intra-uterine alterations have lasting effects and are known to alter cardiovascular structure and function in response to several environmental factors and maternal behaviors[1–5]. One of the most detrimental maternal behaviors during the perinatal period, which can deleteriously predispose offspring to cardiovascular diseases, is nicotine exposure, typically in the form of tobacco product use[6].

Perinatal tobacco smoke/nicotine exposure has been shown to augment blood pressure and increase the risk of hypertension in later life[2,6,7]. Arteriolar wall thickening and increased vasoconstrictor signaling are thought to be the primary causes of perinatal smoke/nicotine exposure-induced hypertension[8–10]. Nicotine stimulates the renin-angiotensin-aldosterone system[11] and increases both angiotensin-converting enzyme activity[12,13] as well as angiotensin II type 1 receptor (AgtR1) expression and activity[12–14] within the aorta[15–17] and mesenteric arterioles[16,18]. Such arteriolar reactivity would likely implement a hypertensive state.

Aberrant sympathetic nerve activity is significantly tied to cardiovascular disease and a prime mediator of hypertension in a multitude of cardiometabolic diseases[19,20]. Carotid bodies are polymodal sensors that regulate sympathetic nervous activity to control vasoconstriction to regulate tissue oxygen/nutrient delivery[21,22]. Indeed, hypertension appears to be critically dependent on tonic input from carotid bodies to augment sympathetic tone[19,20]. Carotid body activity of nicotine-exposed offspring is not suppressed by hyperoxia, which typically quiesces carotid body activity, suggesting an augmented basal sensitization[23,24]. Given their primary role in stimulating arteriolar vasoconstriction[19,20] and their susceptibility to nicotine exposure[23,24], the carotid bodies are likely susceptible to epigenetic modulation because of developmental

programming[25–27], which would lead to an elevation of tonic activity. Therefore, we investigated

the hypothesis that carotid bodies are susceptible to epigenetic hypersensitization which results

in increased blood pressure in perinatal nicotine-exposed offspring. To test our hypothesis, we

engaged modern unbiased sequencing of RNA and DNA methylation of carotid bodies, tested

signaling pathways *in vitro* and *in vivo,* and assessed the necessity of carotid bodies for

heightened arterial pressure in perinatal nicotine-exposed pups.

Methods

Data availability

RNA sequencing and DNA bisulfite sequencing data have been deposited and can be found in

the NIH SRA (Bioproject ID: PRJNA1106547). All other data are found in the supporting data

and supplementary files and can be further made available upon request to the corresponding

author.

Animal models, care and ethical standards

First-time pregnant Sprague-Dawley (SD) rat dams were purchased from Charles River

(Hollister, CA) at gestational day 3. SD mothers were subject to subcutaneous nicotine injections

(2mg/kg, Millipore Sigma, PHR2532) from gestational day 7 to post-natal day (PND) 21[28–30].

Control mothers received PBS (vehicle). Animals were housed with controlled temperature (21 $\pm$

2 °C), humidity (55 $\pm$ 10%) and 12h light-dark cycle. Pups were weaned on post-natal day

(PND) 21 and housed in pairs. Animals had unlimited access to food and water. Experiments on

pups took place from PND56-80. All experimental procedures were approved by The Lundquist

Institute for Biomedical Innovation at Harbor UCLA Medical Center IACUC board, protocol #

31659.

Assignment of control or nicotine to mothers was block randomized. Unbiased transcriptomic

and genomic sequencing was based on previous data acquired by neuronal sequencing results. *In*

vitro and *in vivo* power calculations were based on an estimated effect size of 50% difference

with assumed unified standard deviation across groups.

PC12 culture

PC12 cells were cultured in RPMI 1640 (Gibco # 11875093) supplemented with 10% Horse

Serum (Gibco Cat. #26050-088) and 5% fetal bovine serum (Gibco Cat. #F2442), 0.4µg/ml

dexamethasone (Millipore Sigma Cat. # D4902) and 1% penicillin/streptomycin (Millipore

Sigma Cat. # P0781). Cells were seeded at $3x10^5$ cells in 1ml of media per well (24 well plate,

qPCR) or 2500 cells in 150µl of media per well (96 well plate calcium imaging).

Next-generation RNA sequencing

On PND 56, animals were sedated (5% isoflurane, balance O2), exsanguinated and carotid

bodies gross dissected and placed in a sylgard-coated dissecting dish filled with Ham's F12

media. Carotid bodies were freed from the carotid bifurcation under a dissecting microscope.

Five carotid body pairs (n=10 carotid bodies from 5 rats) from each group and sex (n=5 rats per

group, per sex) were lysed and RNA was isolated using the easy-spin Total RNA extraction kit

(Boca scientific, Cat. # 17221). RNA purity was tested with Nanodrop spectrophotometer

(Thermofisher Cat. # ND2000USCAN) and deemed to be of appropriate quality when A260/280

was between 1.99 and 2.10. RNA sequencing was then conducted by an unbiased commercial

vendor (Novogene corporation, Sacramento CA), where further RNA purity was analyzed (RIN

>7) and downstream library construction took place.

Whole genome DNA-bisulfite sequencing

Carotid bodies were dissected as above. Five carotid body pairs (n=10 carotid bodies from 5 rats)

from each group and sex (n=5 rats per group, per sex) were lysed and DNA was isolated using

the QIAamp UCP DNA Micro Kit (Qiagen, Cat. # 56204). DNA purity was deemed sufficient

with A260/280 between 1.75-1.85. Bisulfite conversion and downstream whole-genome DNA-

bisulfite sequencing were conducted by (Novogene Corporation, Sacramento, CA).

RNA qPCR

For carotid bodies, an additional group of n=3 samples (n=5 rats per sample) from each group

was harvested as above, snap frozen and stored until analysis. Brain and kidneys were harvested

from each group of pups (one tissue per pup was deemed as n=1). For PC12 cells were washed in

PBS and then immediately lysed in lysis buffer and converted into cDNA. N=3 wells were

analyzed for each condition for each probe and five separate experiments were run. RNA was

converted to cDNA using the Tetro cDNA synthesis kit (Meridian life sciences, Cat. #

Bio65043). qPCR was conducted with Taqman specific probes. AgtR1a (Thermofisher

Rn02758772_s1), AgtR2 (Thermofisher Rn00560677_s1), angiotensinogen (Thermofisher

RN00593114_m1), angiotensin-converting enzyme (Thermofisher Rn00561094_m1), renin

(Thermofisher Rn02586313_m1), TH (Thermofisher Rn00562500_m1), endothelial PAS

domain-containing protein 1/ hypoxia inducible factor 2 alpha (EPAS1/Hif2α, Rn00576515_m1)

and TRPV1 (Thermofisher Rn00583117_m1). All experimental genes were evaluated in

reference to hypoxanthine-guanine phosphoribosyltransferase (HPRT Thermofisher

Rn01527840_m1). The qPCR reaction was completed using Taqman fast advanced master mix

enzyme (Thermofisher Cat. # 4444557) and the QuantStudio 3 thermocycler (Thermofisher Cat.

#A28567). Delta CT was calculated in reference to HPRT, and comparisons between groups

were made using an unpaired two-sided t-test.

Methylated DNA immunoprecipitation-qPCR (MeDIP-qPCR)

DNA was isolated from carotid bodies (n=4 per group, each n is comprised of carotid bodies

from 5 rats), brains and kidneys (n=4 rats per group, one tissue is n=1) and PC12 cells (1 well is

n=1). Carotid body and PC12 DNA was isolated using the QIAamp UCP DNA Micro Kit

(Qiagen, Cat. # 56204). Brain and kidney DNA was isolated using the Puregene Tissue kit

(Qiagen, Cat. # 158063). MeDIP was performed using the Methylated DNA

immunoprecipitation kit (Cat.# ab117133, Abcam) following the procedures described

previously[31]. Briefly, 100μL of the antibody buffer and 1μL anti-5-methylcytosine or reference

IgG antibody were added into wells and incubated at room temperature for 1 hour. During the

incubation, 1μg of genomic DNA was sheared between 200 and 1000 bp in the reaction buffer

using the Covaris M220 ultrasonicator, followed by denaturation at 95 °C for 2 minutes, then

placed on ice until use. An aliquot of 5 μL of the denatured DNA was saved as the input DNA.

The strip wells bound with antibodies were washed with 150μL of the antibody buffer and 150μL

of the wash buffer once, followed by incubation with the sheared DNA at room temperature for

2 hours. The antibody-enriched DNA was eluted in the DNA release buffer containing proteinase

K followed by DNA purification. qPCR was performed using Fast SYBR™ Green Master Mix

(Cat.# 4385612, Thermo Fisher Scientific) to identify DNA methylation levels. Primer sequence-

AgtR1: Forward TACCTAAACATAGTAAAAGCCAAACACA; AgtR1: Reverse

TATCCTGTTGATCTCTTTTTGTTGTCTG. ΔCt was calculated as Methylated AgtR1 DNA –

Total AgtR1 DNA and was used to plot the data[31,32]. Comparisons between groups were made

using an unpaired two-sided t-test.

Immunohistochemistry

Carotid bodies were gross dissected as above. Then they were cleaned from connective tissue but

left attached to the surrounding carotid artery bifurcation as previously[33]. Tissues were fixed in

4% paraformaldehyde for 1-2 hours at 4°C, then cryopreserved in 30% sucrose overnight at 4°C,

dried embedded in optimal cutting temperature compound (OCT) and cut at 14μm sections on a

cryostat. Sections were then washed (PBS), permeabilized (Triton x 20%) blocked with 10%

goat serum (Abcam, Cat. # ab7481), and incubated with antibodies for tyrosine hydroxylase (TH,

Millipore Sigma ZMS1033, clone 20/40/15), AgtR1 (Millipore Sigma AB15552, polyclonal) and

TRPV1 (Alomone Labs ACC-030, polyconal) for 24hours at 4°C and subsequently stained with

appropriate secondaries (Jackson Laboratories Goat-anti rabbit Cy3 111-165-003, Cy5 111-005-

003 and Goat-anti mouse 488 115-545-003) for 90 minutes and then counterstained with DAPI

(Millipore Sigma MBD0015) and coverslipped with prolong diamond antifade mounting media

(Thermofisher #P36970). Slides were imaged using the Leica Thunder imaging acquisition

system.

Calcium imaging

PC12 cells were separated into media (control), stimulated with nicotine (Sigma Cat. # 6019-06-

3, 50μg/ml), angiotensin II (Tocris Cat. # 1158, 5μM), nicotine + angiotensin II, angiotensin II +

losartan (Tocris Cat. # 3798, 3μM), nicotine + angiotensin II + losartan, angiotensin II +

AMG9810 (TRPV1 antagonist Tocris Cat. # 2316, 10μM), nicotine + angiotensin II +

AMG9810, or nicotine + angiotensin II + losartan + AMG9810, for 1 hour and then incubated

with Calcium Orange AM methyl ester (Invitrogen Cat. # C3015) + 0.01% Pluronic acid

(Invitrogen Cat. #F-127) for 20 minutes, washed in media and then imaged in the Incucyte

imaging system (Sartorius) using the Neural imaging parameters (Ex: 549λm/ Em: 576λm, 60s)

for 1 hour. The frequency of methyl ester excitation bursts was averaged for each well to express

a single value. Data were analyzed with One-way ANOVA using the Holm-Sidak post-hoc test.

In vivo carotid body reactivity

Rats from control or nicotine groups (as above) were tested on PND 56-58. Rats were

anesthetized with 5% isoflurane (balance room air) using the Kent Scientific Somnosuite

anesthesia system. The femoral artery and vein were cannulated, and the arterial line was

instrumented to a pressure transducer to measure continuous arterial pressure (AD instruments

MLT0670) using the PowerLab data acquisition console (AD instruments PL3516/P). The vein

was then connected to a syringe pump (Fisherscientific Cat. # 14831200) to induce a

15mg/kg/min infusion of alfaxan (Victor Medical Cat. # 1629020)[34,35]. Then the jugular vein was

cannulated to deliver bolus injections of Angiotensin II (Tocris Cat. # 1158 1mg/kg) and sodium

cyanide (NaCN, Fisherscientific Cat. # AA1213722, 200µg/kg). These injections were repeated

following bilateral carotid body denervation. Briefly, the neck area was shaved, and a midline

incision exposed the carotid artery. Then glands were pushed aside, and the carotid bodies were

identified within the carotid bifurcation, underneath the occipital artery. The carotid sinus nerve

was confirmed as it was traced to the glossopharyngeal nerve, then denervated, as previous[34].

Mean arterial pressure was averaged at the 60s preceding each bolus injection and a 60s average

during the peak reactive response to each bolus injection. Data were analyzed by Two-way

ANOVA (Group x Carotid body intact vs. denervated) for Angiotensin II or NaCN injection.

Holm-Sidak post-hoc test determined group differences.

Serum angiotensin concentration

Blood was collected after induction of IV anesthetic but before any drug infusion from the above

experiment from the femoral vein. Blood was left to coagulate for 20 minutes on ice then spun at

4,000rpm for 10 minutes at 4°C. Serum was drawn, aliquoted, snap-frozen, and stored at -80°C

until analysis. Angiotensin ELISA (Raybiotech #EIAR-ANGII-1) was used per manufacturer's

instructions. Groups (Control pups vs. perinatal nicotine-exposed pups) were analyzed with two-

sided unpaired t-test.

Arterial pressure assessment between groups

Rats from control or perinatal nicotine-exposed mothers were handled daily following weaning

at PND 21 and habituated to the tail plethysmography system (IITC Lifesciences). Three-minute

averages of tail cuff-measured blood pressure was averaged to attain a single value for each

animal. Data were compared using an unpaired two-sided t-test.

Carotid body-mediated changes in arterial pressure

Perinatal nicotine-exposed rats were prepared for telemetric implant between PND 56-60. Rats

were anesthetized with 5% isoflurane (balance O_2) as induction and maintained on a surgical

plane with 2-3% isoflurane. The abdominal area was isolated, shaved, and treated with 3x

betadyne and 3x 70% alcohol. A midline incision was made, and the descending aorta below the

renal arteries and renal nerves were isolated. The Kaha telemetry dual sympathetic nerve activity

/pressure (TRM56SP, AD Instruments) were instrumented around the renal nerves using

KwikGard dental impression material to isolate electrodes for renal sympathetic nerve activity

(RSNA) recording. The pressure transducer was inserted into the aorta as per manufacturer's

instructions to measure continuous pressure measurement. The surgical area was sewn in layers,

and the skin was secured with surgical staples. Rats were given buprenorphine (0.03mg/kg SQ)

2x daily for three days and allowed to recover for a total of 5 days before commencing recording.

Pressure and RSNA were recorded for 1 week; then rats were anesthetized as above; the neck

area was isolated, shaved, and treated with 3x betadine and 3x 70% alcohol. A midline incision

was made on the neck, and the carotid bodies were identified behind the occipital arteries. The

carotid sinus nerve was severed bilaterally, and the skin was sewn with subcuticular sutures.

Buprenorphine was delivered as above, and recording commenced for another week following

recovery. Arterial pressure waveforms delineated heart rate, systolic, diastolic, mean arterial

pressure, pulse pressure, and respiratory rate. RSNA was filtered, rectified, and integrated using

LabChart 8. Data were binned into daily averages and analyzed pre-post with a two-sided paired

t-test.

Statistics

Specific statistical tests are described for each experiment above. Data were averaged and

analyzed for each experiment as above. Data were analyzed using Graphpad v10.

Results

Epigenetic changes in the carotid body in response to perinatal nicotine exposure

To investigate possible genetic changes expounded by adverse maternal behaviors, we

implemented a standardized model of perinatal-nicotine exposure[28–30]. Given that the carotid

bodies are prime drivers of hypertension in self-actualized cardiometabolic disease states[22,36], we

conducted an unbiased transcriptomic screen of carotid bodies harvested from perinatal nicotine-

exposed pups on PND 56, 5 weeks after weaning and the final dose of nicotine delivered to the

mother. We pooled together carotid bodies from 5 pups in each group of nicotine-exposed and

saline control pups (Figure 1a). Samples were analyzed by sex, but no sex difference was found.

A total of 922 genes were differentially expressed between conditions, with 685 up-regulated,

237 down-regulated, and 19196 unchanged transcripts (Figure 1b, c). AgtR1a was significantly

upregulated in nicotine-exposed offspring compared to control (0.69 Log2 change, p_{adj}=0.04).

Angiotensin II type 2 receptor (AgtR2) was downregulated by perinatal nicotine exposure

(Log2FC=-3.246, p_{adj} =6.65x10^{-12}). No differences in angiotensinogen (Log2FC=0.233, p_{adj}

=0.693), angiotensin-converting enzyme (Log2FC=-0.071, p_{adj}=0.871), or renin (Log2FC=-

2.248, p_{adj}=0.645) were demonstrated (Supplementary data 2). Gene set enrichment analysis

(GSEA) was performed on the differentially expressed gene, revealing a vast number of highly

significant GO pathways, including GO:0004930 (normalized enrichment score =1.539;

Figure1d, Supplementary data 1), which included AgtR1a.

To validate these findings, we conducted immunohistochemistry and qPCR with taqman probes

to identify possible upregulation of potential signaling pathways in the carotid bodies between

perinatal nicotine exposed and control pups. We show that in relation to HPRT, AgtR1a was

significantly upregulated in perinatal nicotine-exposed offspring compared to control. AgtR2 and

TRPV1 transcription was not significantly increased (Supplementary figure 1).

To test whether changes in angiotensin signaling occurred outside the carotid bodies, we tested

whether RNA transcripts of AgtR1a, AgtR2, angiotensinogen, angiotensin-converting enzyme, or

renin were altered in the kidney and brains of perinatal nicotine exposed pups compared to

controls. We observed that angiotensinogen and angiotensin-converting enzymes were

significantly upregulated in the brains and kidneys of perinatal nicotine-exposed rats compared

to the control, and renin was downregulated in the kidneys of perinatal nicotine-exposed rats

compared to the control with no change in the brain (Supplementary figure 2). In contrast, both

AgtR1a and AgtR2 were unchanged in the brains and kidneys of perinatal nicotine-exposed rats

compared to control rats (Supplementary figure 2).

Given that our goal was to elucidate possible epigenetic differences responsible for mediating a

neurogenic form of hypertension, we probed further and conducted whole genome DNA bisulfite

sequencing[37] of the carotid bodies from perinatal nicotine-exposed and control pups on PND 56

(Figure 2a). Samples were analyzed by sex, but no sex difference was found. Mapping the whole

genome of the carotid bodies between groups revealed that the predominant changes to DNA

methylation occurred in the promoter (suppression) and intron/exon regions (enhancement,

Figure 2b). Methylation was distributed throughout the genome where a differential set of Genes

were altered predominantly on the cytosine-guanine (CG) and cytosine-

arginine/threonine/guanine-guanine (CHG) regions (Figure 2b). GO analyses of whole genes

(Figure 2d) and promoter regions (Figure 2e) revealed similar GO terms as with transcriptomic

changes. Importantly, a significant difference in methylation AgtR1a between groups was

demonstrated with a difference of 0.448319 between perinatal nicotine exposed and control rats

(Supplementary data 3). AgtR1 was orthogonally confirmed with immunoprecipitation of

methylated DNA compared to total DNA amplifying AgtR1 to assess global methylation (Figure

2c). AgtR1 DNA was not significantly methylated in brains but was increased in the kidneys of

perinatal nicotine-exposed rats compared to controls (Supplementary figure 3). Given that these

methylation changes occurred in regions associated with the upregulation of angiotensin GPCR

signaling, we wanted to investigate the functional outcomes of this pathway further. The

combined transcriptomic changes and DNA methylation of the whole genome point to an

upregulation of angiotensin signaling within the carotid bodies of perinatal nicotine-exposed

offspring.

Proteins for AgtR1, TRPV1, and tyrosine hydroxylase (a standard marker of carotid body glomus

cells) were all present in sections of carotid bodies harvested from perinatal nicotine-exposed

offspring where AgtR1 is on glomus cells, and TRPV1 appears to oppose glomus cells on the

post-synaptic terminals[38] (Supplementary figure 1), However, our resolution is insufficient to

determine the localization of glomus cells definitively. It is possible that these antibodies may

suffer from this lack of specificity in rats. As such, our staining may not reflect the accurate

location of AgtR1 in rat carotid body glomus cells. Nevertheless, given our RNAseq, qPCR, and

DNA methylation data, along with others who have orthogonally demonstrated the presence of

AgtR1 in glomus cells[38–41], we are confident of the expression and upregulation of AgtR1 in the

carotid bodies of perinatal nicotine exposed rats.

Nicotine upregulates AgtR1 activity

To show that nicotine itself is responsible for AgtR1 upregulation, we used a cell culture model.

PC12 cells have been routinely used as a surrogate carotid body glomus cell system[42–44]. Six

hours of nicotine exposure significantly upregulated AgtR1a gene expression (Figure 3a).

Differences in AgtR2 and TRPV1 were not demonstrated (Figure 3a). Differences in

angiotensinogen, angiotensin converting enzyme, renin, and regulatory genes of EPAS1 (Hif2α)

and tyrosine hydroxylase (TH- Figure 3a) were not demonstrated between conditions in relation

to the housekeeping gene HPRT. DNA methylation of AgtR1 was confirmed with the DNA

methylation pulldown assay (Figure 3b).

Using calcium imaging, we investigated how nicotine would augment glomus cell excitation. We

show that with 1 hour of exposure to nicotine, PC12 cells increased their level of calcium spike

frequency. The addition of angiotensin increased PC12 cell firing frequency to a level similar to

nicotine. Importantly, nicotine combined with angiotensin did not further increase PC12 cell

excitation. Further, losartan reduced calcium activity in PC12 cells when delivered to angiotensin

or nicotine + angiotensin-stimulated cells concurrently. The reduction in calcium flux was also

reduced with TRPV1 antagonist (AMG9810), and in combination with AMG9810 and Losartan

(Figure 3c). In conclusion, nicotine has a stimulatory effect on PC12 and, quite likely, carotid

body glomus cells to increase the expression and excitation of the angiotensin signaling pathway via AgtR1 and TRPV1.

Angiotensin acutely increases arterial pressure

To assess acute carotid body reactivity to angiotensin, pups from perinatal nicotine-exposed or control mothers were anesthetized and given a bolus angiotensin injection into the jugular vein before and after acute carotid body denervation (Figure 4 a, b). In response to jugular vein injected angiotensin (circulates through the pulmonary circulation, then encounters the carotid bodies), arterial pressure was significantly elevated in nicotine-exposed offspring compared to controls (Figure 4c, Supplementary figure 4). In response to sodium cyanide, which blocks mitochondrial complex III and has been routinely used as a carotid body stimulant, there was no significant difference between groups (Figure 4d, Supplementary figure 4). Following carotid body denervation, arterial pressure was suppressed and did not respond to angiotensin or sodium cyanide as vigorously (Figure 4c, d, Supplementary figure 4). The reactivity to angiotensin and NaCN were attributed to the carotid bodies, however, given that boluses were delivered through the jugular vein, off-target effects cannot be ruled out.

Importantly, serum harvested from these rats prior to any drug delivery demonstrated that circulating levels of angiotensin were not altered between groups (Figure 4e), similar to previous results[18]. It is possible that our analysis may be affected by background proteins in our sample, despite the ability of angiotensin II ELISAs to discern various forms of angiotensin and cleaved fragments[45].

These data demonstrate a significant increase in carotid body excitability in response to angiotensin *in vivo* and suggest a hyperactive angiotensin signaling pathway within the carotid bodies of perinatal nicotine-exposed offspring.

Carotid bodies are responsible for elevated blood pressure in perinatal nicotine-exposed

offspring

We evaluated arterial pressure between quiescent conscious control and nicotine-exposed

offspring using tail-cuff plethysmography on PND 56. Indeed, arterial pressure was higher in

nicotine-exposed offspring (Figure 5a). To definitively demonstrate that the carotid bodies were

primarily involved in perinatal nicotine-exposed hypertension, we implanted indwelling

telemetry to measure real-time arterial pressure from the aorta and efferent sympathetic activity

directed to the renal nerve between PND 56-60 (Figure 5b, Supplementary figure 5). We show

that within the same animal, carotid body denervation drastically reduces arterial pressure and

RSNA (Figure 5b, Supplementary figure 5). The reduction in arterial pressure with carotid body

denervation was predominantly due to the suppression of systolic pressure, which significantly

reduced pulse pressure (Supplementary figure 5). The reductions of RSNA as a result of carotid

body denervation also coincide with a reduced respiratory rate, similar to other models of

hypertension[19]. We attribute our changes in response to carotid body denervation to a reduction

of RSNA as measured and demonstrated by others[19,20]. However, we did not assess the precise

contribution of the carotid bodies to RSNA levels as previous[19]. Nevertheless, we note the

association of RSNA to respiratory rate, which indicates that the reduction of respiratory rate

following carotid body denervation would be tied to the fall in RSNA. These data demonstrate

that carotid body basal activity and excitation contribute to the hypertensive state in perinatal

nicotine-exposed offspring.

Discussion

Epigenetic changes in response to adverse maternal programming lead to cardiovascular

sequelae in later life. Hypertension has been a consistent outcome associated with many adverse

intra-uterine insults ranging from undernutrition[46,47], overnutrition[47,48], hypoxia[49,50], and stress[51].

The risk of hypertension is thought to be significantly increased in perinatal nicotine-exposed

offspring. The current data demonstrate that epigenetic changes in angiotensin signaling in the

carotid bodies mediate hypertension in perinatal nicotine-exposed offspring.

We show, for the first time, that epigenetic modifications in response to adverse maternal

programming in the form of nicotine exposure increase angiotensin, angiotensin II type 1a

receptor transcripts, and angiotensin II type 1 receptor DNA intron methylation, demonstrating

an increase in angiotensin signaling in the carotid bodies. These epigenetic changes *in vivo* are

confirmed *in vitro* as we show that nicotine directly increases AgtR1 gene expression, AgtR1

DNA methylation, and AgtR1 stimulation in PC12, glomus-like cells, in culture.

Importantly, the difference in epigenetic changes to AgtR1 was predominantly confined to

carotid bodies as the brain did not show an increase in gene expression or DNA methylation,

whereas kidneys only showed increased DNA methylation between perinatal nicotine exposed

and control pups. Angiotensinogen and angiotensin-converting enzyme gene expression was

upregulated in the brains and kidneys, and renin was increased in the kidneys of perinatal

nicotine exposed compared to control pups. In line with previous investigations, blood pressure

reactivity to angiotensin is increased through a neurogenic mechanism involving carotid body-

mediated excitation of efferent sympathetic activity. Finally, we showed, for the first time, that

carotid body denervation reduced blood pressure in perinatal nicotine-exposed rats alongside

efferent sympathetic activity.

Clinical data demonstrate that hypertension is likely in offspring that experienced intra-

uterine/perinatal smoke exposure[2,7,52–56]; however, the age, sex, and potential for additional

confounding lifestyle factors make it difficult to tease out this association, and measurements

beyond adolescence have not been completed to date. Pre-clinical data using rodent models show

that vascular reactivity to angiotensin via angiotensin II type 1 receptor gene expression, protein

expression, and signaling increases in aortae and mesenteric arterioles following prenatal and

perinatal nicotine exposure[14,15,18]. Accordingly, increased angiotensin II type 1 receptor activity,

RNA, and DNA methylation[57,58] have been consistent findings in response to perinatal nicotine-

exposed offspring in vascular tissues. Our data show that additional sensory neural systems,

which are significantly tied to hypertension[20], are also augmented and induce hypertension in

perinatal nicotine-exposed rats.

To date, scant data are available regarding the effects of nicotine exposure or perinatal nicotine

exposure on carotid body activity and resulting autonomic reflex disturbances[23,59]. Carotid bodies

are tasked with a multitude of homeostatic control mechanisms that regulate nutrient and oxygen

delivery to tissues[21]. Their role in blood pressure regulation is highlighted by genetic[22] and

epigenetic[27] modifications demonstrating the ability to increase basal activity and result in

augmented blood pressure. Specifically, the spontaneously hypertensive rat (SHR) has

augmented carotid body basal activity, which translates to alterations in breathing, blood pressure

regulation, and glucose homeostasis[22]. Importantly, these carotid body changes appear to hinge

on a significant difference in transcriptomic profile compared to their normotensive genetic

counterpart, the Wistar-Kyoto (WKY) rat[22]. Further, and particularly salient to the current

argument, carotid body denervation in SHR reduced arterial pressure and respiratory modulation,

whereas carotid body denervation in WKY rats had no effect[19]. Importantly, evidence

demonstrates that carotid body basal excitation can change via epigenetic modifications that

suppress redox disinhibition[27] to alter basal regulatory mechanisms such as breathing regulation.

A consistent finding is that intrauterine conditions with low oxygen[23] or hyperleptinemia[60] lead

to augmented carotid body basal activity. Therefore, it is possible that deficiencies in the

intrauterine environment augment arterial pressure in offspring due to the carotid body-regulated

mechanism. Our data are the first to demonstrate that adverse maternal programming leads to

carotid body epigenetic changes in AgtR1a and, importantly, the first to demonstrate that these

epigenetic changes occur as a result of genetic upregulation via DNA methylation, which results

in carotid body excitation. This carotid body excitation appears to be due to increased

angiotensin signaling within the carotid bodies due to increases in carotid body micro signaling

domains. Our *in vitro* experiments with PC12 cells demonstrate that epigenetic changes in

carotid body angiotensin signaling directly result from nicotine exposure. Further, the reactivity

of PC12 cells was regulated by nicotine to increase AgtR1 signaling. AgtR1 signaling in the

carotid bodies has been tied to a TRPV1-mediated mechanism in response to repeated hypoxic

bouts[38]. Given that losartan and TRPV1 blockade suppressed calcium signaling in PC12 cells,

with no further reduction in calcium excitation with dual inhibition, we suggest that a similar

mechanism is responsible for the augmentation of basal carotid body excitation. These data then

fit with the increased carotid body reactivity to angiotensin in an anesthetized preparation and the

consistent reduction in arterial pressure with carotid body denervation in conscious perinatal

nicotine-exposed offspring.

Conclusion

Hypertension resulting from perinatal nicotine exposure is due to epigenetic modifications within

the carotid bodies. The increased AgtR1-TRPV1 signaling increases efferent sympathetic activity

to augment arterial pressure. Future investigations should target neurogenic signaling

mechanisms to alleviate hypertension due to adverse epigenetic programming.

Perspectives

Carotid body denervation has shown clinical utility, as human clinical trials have shown that unilateral carotid body denervation results in a meaningful reduction in human blood pressure[61,62]. In fact, the reduction of systolic and diastolic arterial pressure with carotid body unilateral denervation[62,63] appears to be similar to renal denervation[64,65]. Targeting the carotid bodies offers the advantage of suppressing the first step of reflex blood pressure regulation (i.e., sensory afferents[64,65]. Therefore, we suggest that a targeted suppression or reversal of angiotensin signaling in the carotid bodies and renin-angiotensinogen and angiotensin-converting enzyme in the kidneys would help alleviate increased blood pressure and cardiac afterload and limit unwanted side effects by standard drug therapies or invasive procedures.

Acknowledgements and sources of funding

This project was supported by T32IP4707 of the Regents of the University of California Tobacco-Related Diseases Research Program (TRDRP). NJ is supported by NIH grant R21AI159221, R56AI175328, UCLA CTSI UL1TR001881-01 and, T32KT4708 of the Regents of the University of California TRDRP. WY is supported by NIH grant P50HD098593 and R01HD099924. VR is supported by NIH grant R01HL151769 and IR00737, T32IP5044, T32IR5048, and T32IR5365 of the Regents of the University of California TRDRP. The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health or the UCOP TRDRP. Figures, 1a, 2a, 4a and 6a,b were completed with Biorender under a CC-BY-NC-ND license.

Author contributions

F.Z. Z.W.& J.K. contributed to experimental design, conducted experiments and analyzed data.

H.M, K.D. and J.L. contributed to experimental execution and data analysis. W.Y. and V.R.

assisted with experimental design and analysis. N.J. designed the study, conducted experiments,

analyzed data, and prepared the manuscript and figures. All authors agree on the manuscript.

Disclosures

All authors do not declare competing interests.

References

1. Higa R, Leonardi ML, Jawerbaum A. Intrauterine Programming of Cardiovascular Diseases
in Maternal Diabetes. Front Physiol [Internet]. 2021 [cited 2024 Jan 11];12. Available from:
https://www.frontiersin.org/articles/10.3389/fphys.2021.760251

2. Parada-Ricart E, Luque V, Zaragoza M, Ferre N, Closa-Monasterolo R, Koletzko B, Grote
V, Gruszfeld D, Verduci E, Xhonneux A, Escribano J. Effect of maternal smoking during
pregnancy on child blood pressure in a European cohort. Sci Rep. Nature Publishing
Group; 2022 Oct 15;12(1):17308.

3. Palinski W. Effect of Maternal Cardiovascular Conditions and Risk Factors on Offspring
Cardiovascular Disease. Circulation. American Heart Association; 2014 May
20;129(20):2066–2077.

4. Dasinger JH, Davis GK, Newsome AD, Alexander BT. Developmental Programming of
Hypertension. Hypertension. American Heart Association; 2016 Oct;68(4):826–831.

5. Alexander BT, Dasinger JH, Intapad S. Fetal Programming and Cardiovascular Pathology.
Compr Physiol. 2015 Apr;5(2):997–1025. PMCID: PMC4772789

6. Taylor L, Kelly B, Leeson P. Maternal Smoking and Infant Cardiovascular Physiology.
Hypertension. American Heart Association; 2010 Mar;55(3):614–616.

7. Kooijman MN, Bakker H, Franco OH, Hofman A, Taal HR, Jaddoe VWV. Fetal Smoke
Exposure and Kidney Outcomes in School-Aged Children. Am J Kidney Dis Off J Natl
Kidney Found. 2015 Sep;66(3):412–420. PMID: 25641064

8. Nordenstam F. Prenatal nicotine exposure was associated with long-term impact on the
cardiovascular system and regulation—Review. Acta Paediatr. John Wiley & Sons, Ltd;
2021 Sep 1;110(9):2536–2544.

9.   Nordenstam F, Norman M, Caidahl K, Wickström R. Arterial Stiffness and Carotid Intima-
Media Thickness in Children Exposed to Smokeless Tobacco in Fetal Life. J Am Heart
Assoc. Wiley; 2024 Jan 16;13(2):e9128.

10.   Spindel ER, McEvoy CT. The Role of Nicotine in the Effects of Maternal Smoking during
Pregnancy on Lung Development and Childhood Respiratory Disease. Implications for
Dangers of E-Cigarettes. Am J Respir Crit Care Med. 2016 Mar 1;193(5):486–494. PMCID:
PMC4824926

11.   Oakes JM, Fuchs RM, Gardner JD, Lazartigues E, Yue X. Nicotine and the renin-
angiotensin system. Am J Physiol - Regul Integr Comp Physiol. 2018 Nov 1;315(5):R895–
R906. PMCID: PMC6295500

12.   Ramalingam A, Budin SB, Mohd. Fauzi N, Ritchie RH, Zainalabidin S. Angiotensin II Type I
Receptor Antagonism Attenuates Nicotine-Induced Cardiac Remodeling, Dysfunction, and
Aggravation of Myocardial Ischemia-Reperfusion Injury in Rats. Front Pharmacol [Internet].
2019 [cited 2024 Jan 25];10. Available from:
https://www.frontiersin.org/articles/10.3389/fphar.2019.01493

13.   Fried ND, Morris TM, Whitehead A, Lazartigues E, Yue X, Gardner JD. Angiotensin II type
1 receptor mediates pulmonary hypertension and right ventricular remodeling induced by
inhaled nicotine. Am J Physiol Heart Circ Physiol. 2021 Apr 1;320(4):H1526–H1534.
PMCID: PMC8260386

14.   Mao C, Zhang H, Xiao D, Zhu L, Ding Y, Zhang Y, Wu L, Xu Z, Zhang L. Perinatal nicotine
exposure alters AT1 and AT2 receptor expression pattern in the brain of fetal and offspring
rats. Brain Res. 2008 Dec 3;1243:47–52. PMCID: PMC2664080

15.   Xiao D, Huang X, Lawrence J, Yang S, Zhang L. Fetal and Neonatal Nicotine Exposure
Differentially Regulates Vascular Contractility in Adult Male and Female Offspring. J
Pharmacol Exp Ther. American Society for Pharmacology and Experimental Therapeutics;
2007 Feb 1;320(2):654–661. PMID: 17068201

16.   Gao YJ, Holloway AC, Su LY, Takemori K, Lu C, Lee RMKW. Effects of fetal and neonatal
exposure to nicotine on blood pressure and perivascular adipose tissue function in adult
life. Eur J Pharmacol. 2008 Aug 20;590(1):264–268.

17.   Paxton WG, Marrero MB, Klein JD, Delafontaine P, Berk BC, Bernstein KE. The
angiotensin II AT1 receptor is tyrosine and serine phosphorylated and can serve as a
substrate for the src family of tyrosine kinases. Biochem Biophys Res Commun. 1994 Apr
15;200(1):260–267. PMID: 7513159

18.   Tao H, Rui C, Zheng J, Tang J, Wu L, Shi A, Chen N, He R, Wu C, Li J, Yin X, Zhang P,
Zhu Z, Tao J, Xiao J, Mao C, Xu Z. Angiotensin II-mediated vascular changes in aged
offspring rats exposed to perinatal nicotine. Peptides. 2013 Jun 1;44:111–119.

19.   Abdala AP, McBryde FD, Marina N, Hendy EB, Engelman ZJ, Fudim M, Sobotka PA,
Gourine AV, Paton JF. Hypertension is critically dependent on the carotid body input in the
spontaneously hypertensive rat. J Physiol. 2012 Sep 1;590(Pt 17):4269–77. PMID:
22687617

20. McBryde FD, Abdala AP, Hendy EB, Pijacka W, Marvar P, Moraes DJ, Sobotka PA, Paton
JF. The carotid body as a putative therapeutic target for the treatment of neurogenic
hypertension. Nat Commun. 2013;4:2395.

21. Zera T, Moraes DJA, da Silva MP, Fisher JP, Paton JFR. The Logic of Carotid Body
Connectivity to the Brain. Physiology. 2019 Jun 5;34(4):264–282. PMID: 31165684

22. Pauza AG, Thakkar P, Tasic T, Felippe I, Bishop P, Greenwood MP, Rysevaite-Kyguoliene
K, Ast J, Broichhagen J, Hodson DJ, Salgado HC, Pauza DH, Japundzic-Zigon N, Paton
JFR, Murphy D. GLP1R Attenuates Sympathetic Response to High Glucose via Carotid
Body Inhibition. Circ Res. 2022 Mar 4;130(5):694–707. PMCID: PMC8893134

23. Stéphan-Blanchard E, Chardon K, Léké A, Delanaud S, Djeddi D, Libert JP, Bach V, Telliez
F. In Utero Exposure to Smoking and Peripheral Chemoreceptor Function in Preterm
Neonates. Pediatrics. American Academy of Pediatrics; 2010 Mar 1;125(3):e592–e599.
PMID: 20176675

24. Argacha JF, Xhaët O, Gujic M, Adamopoulos D, Beloka S, Dreyfuss C, Degaute JP, van de
Borne P. Nicotine increases chemoreflex sensitivity to hypoxia in non-smokers. J
Hypertens. 2008 Feb;26(2):284–294. PMID: 18192843

25. Holgert H, Hökfelt T, Hertzberg T, Lagercrantz H. Functional and developmental studies of
the peripheral arterial chemoreceptors in rat: effects of nicotine and possible relation to
sudden infant death syndrome. Proc Natl Acad Sci U S A. 1995 Aug 1;92(16):7575–7579.
PMCID: PMC41382

26. Prieto-Lloret J, Caceres AI, Obeso A, Rocher A, Rigual R, Agapito MT, Bustamante R,
Castañeda J, Perez-Garcia MT, Lopez-Lopez JR, Gonzalez C. Ventilatory responses and
carotid body function in adult rats perinatally exposed to hyperoxia. J Physiol. 2004 Jan
1;554(Pt 1):126–144. PMCID: PMC1664733

27. Nanduri J, Prabhakar NR. Epigenetic regulation of Carotid Body Oxygen Sensing: Clinical
Implications. Adv Exp Med Biol. 2015;860:1–8. PMCID: PMC4870818

28. Liu J, Yu C, Doherty TM, Akbari O, Allard P, Rehan VK. Perinatal nicotine exposure-
induced transgenerational asthma: Effects of reexposure in F1 gestation. FASEB J Off
Publ Fed Am Soc Exp Biol. 2020 Sep;34(9):11444–11459. PMCID: PMC7839813

29. Xiao D, Huang X, Li Y, Dasgupta C, Wang L, Zhang L. Antenatal Antioxidant Prevents
Nicotine-Mediated Hypertensive Response in Rat Adult Offspring. Biol Reprod. 2015
Sep;93(3):66. PMCID: PMC4710188

30. Wongtrakool C, Wang N, Hyde DM, Roman J, Spindel ER. Prenatal Nicotine Exposure
Alters Lung Function and Airway Geometry through α7 Nicotinic Receptors. Am J Respir
Cell Mol Biol. 2012 May;46(5):695–702. PMCID: PMC3359906

31. Wang Z, McSwiggin H, Newkirk SJ, Wang Y, Oliver D, Tang C, Lee S, Wang S, Yuan S,
Zheng H, Ye P, An W, Yan W. Insertion of a chimeric retrotransposon sequence in mouse
Axin1 locus causes metastable kinky tail phenotype. Mob DNA. 2019 May 3;10(1):17.
PMID: 31073336

32.   Lisanti S, von Zglinicki T, Mathers JC. Standardization and quality controls for the
568   methylated DNA immunoprecipitation technique. Epigenetics. Taylor & Francis; 2012 Jun
569   1;7(6):615–625. PMID: 22507898

570   33.   Jendzjowsky NG, Roy A, Wilson RJA. Asthmatic allergen inhalation sensitises carotid
571   bodies to lysophosphatidic acid. J Neuroinflammation. 2021 Aug 31;18:191. PMCID:
572   PMC8408927

573   34.   Jendzjowsky NG, Roy A, Barioni NO, Kelly MM, Green FHY, Wyatt CN, Pye RL, Tenorio-
574   Lopes L, Wilson RJA. Preventing acute asthmatic symptoms by targeting a neuronal
575   mechanism involving carotid body lysophosphatidic acid receptors. Nat Commun. 2018
576   Oct 2;9(1):4030. PMID: 30279412 PMCID: PMC6168495

577   35.   Marshall JM. Interactions between local dilator and sympathetic vasoconstrictor influences
578   in skeletal muscle in acute and chronic hypoxia. Exp Physiol. 2015 Dec 1;100(12):1400–
579   1411.

580   36.   Conde SV, Sacramento JF, Guarino MP. Carotid body: a metabolic sensor implicated in
581   insulin resistance. Physiol Genomics. 2018 01;50(3):208–214. PMID: 29373079

582   37.   Kurdyukov S, Bullock M. DNA Methylation Analysis: Choosing the Right Method. Biology.
583   2016 Jan 6;5(1):3. PMCID: PMC4810160

584   38.   Roy A, Farnham MMJ, Derakhshan F, Pilowsky PM, Wilson RJA. Acute intermittent
585   hypoxia with concurrent hypercapnia evokes P2X and TRPV1 receptor-dependent sensory
586   long-term facilitation in naïve carotid bodies. J Physiol. :n/a-n/a.

587   39.   Schultz HD. Angiotensin and Carotid Body Chemoreception in Heart Failure. Curr Opin
588   Pharmacol. 2011 Apr;11(2):144–149. PMCID: PMC3075378

589   40.   Lam SY, Liu Y, Ng KM, Liong EC, Tipoe GL, Leung PS, Fung ML. Upregulation of a local
590   renin-angiotensin system in the rat carotid body during chronic intermittent hypoxia. Exp
591   Physiol. 2014 Jan;99(1):220–231. PMID: 24036592

592   41.   M-L Fung null, Lam SY, Dong X, Chen Y, Leung PS. Postnatal hypoxemia increases
593   angiotensin II sensitivity and up-regulates AT1a angiotensin receptors in rat carotid body
594   chemoreceptors. J Endocrinol. 2002 May;173(2):305–313. PMID: 12010638

595   42.   Hockman D, Adameyko I, Kaucka M, Barraud P, Otani T, Hunt A, Hartwig AC, Sock E,
596   Waithe D, Franck MCM, Ernfors P, Ehinger S, Howard MJ, Brown N, Reese J, Baker CVH.
597   Striking parallels between carotid body glomus cell and adrenal chromaffin cell
598   development. Dev Biol. 2018 May 25; PMID: 29807017

599   43.   Dixon DN, Loxley RA, Barron A, Cleary S, Phillips JK. Comparative studies of PC12 and
600   mouse pheochromocytoma-derived rodent cell lines as models for the study of
601   neuroendocrine systems. In Vitro Cell Dev Biol Anim. 2005 Aug;41(7):197–206. PMID:
602   16223334

603   44.   Zhu WH, Conforti L, Czyzyk-Krzeska MF, Millhorn DE. Membrane depolarization in PC-12
604   cells during hypoxia is regulated by an O2-sensitive K+ current. Am J Physiol. 1996
605   Aug;271(2 Pt 1):C658-665. PMID: 8770007

45. Chappell MC. Biochemical evaluation of the renin-angiotensin system: the good, bad, and absolute? Am J Physiol - Heart Circ Physiol. American Physiological Society; 2016 Jan 1;310(2):H137. PMID: 26475588

46. Mariano VS, Boer PA, Gontijo JAR. Fetal Undernutrition Programming, Sympathetic Nerve Activity, and Arterial Hypertension Development. Front Physiol [Internet]. 2021 [cited 2024 Feb 6];12. Available from: https://www.frontiersin.org/journals/physiology/articles/10.3389/fphys.2021.704819

47. Eriksson J, Forsén T, Tuomilehto J, Osmond C, Barker D. Fetal and Childhood Growth and Hypertension in Adult Life. Hypertension. American Heart Association; 2000 Nov;36(5):790–794.

48. Cho NH, Silverman BL, Rizzo TA, Metzger BE. Correlations between the intrauterine metabolic environment and blood pressure in adolescent offspring of diabetic mothers. J Pediatr. 2000 May 1;136(5):587–592.

49. Walton SL, Bielefeldt-Ohmann H, Singh RR, Li J, Paravicini TM, Little MH, Moritz KM. Prenatal hypoxia leads to hypertension, renal renin-angiotensin system activation and exacerbates salt-induced pathology in a sex-specific manner. Sci Rep. Nature Publishing Group; 2017 Aug 15;7(1):8241.

50. Zhou J, Xiao D, Hu Y, Wang Z, Paradis A, Mata-Greenwood E, Zhang L. Gestational Hypoxia Induces Preeclampsia-Like Symptoms via Heightened Endothelin-1 Signaling in Pregnant Rats. Hypertension. American Heart Association; 2013 Sep;62(3):599–607.

51. Costa TJ, De Oliveira JC, Giachini FR, Lima VV, Tostes RC, Bomfim GF. Programming of Vascular Dysfunction by Maternal Stress: Immune System Implications. Front Physiol [Internet]. 2022 [cited 2024 Feb 6];13. Available from: https://www.frontiersin.org/journals/physiology/articles/10.3389/fphys.2022.787617

52. Mourino N, Varela-Lema L, Ahluwalia JS, Rey-Brandariz J, Candal-Pedreira C, Ruano-Ravina A, Vila-Farinas A, Torres A, Pérez-Rios M. Maternal smoking in pregnancy and blood pressure during childhood and adolescence: a meta-analysis. Eur J Pediatr. 2023;182(5):2119–2132. PMCID: PMC10175379

53. He B, Zhang Q, Guo Y, Ao Y, Tie K, Xiao H, Chen L, Xu D, Wang H. Prenatal smoke (Nicotine) exposure and offspring's metabolic disease susceptibility in adulthood. Food Chem Toxicol. 2022 Oct 1;168:113384.

54. Hacke C, Weisser B. Effects of Parental Smoking on Exercise Systolic Blood Pressure in Adolescents. J Am Heart Assoc. Wiley; 2015 May 11;4(5):e001936.

55. Aryanpur M, Yousefifard M, Oraii A, Heydari G, Kazempour-Dizaji M, Sharifi H, Hosseini M, Jamaati H. Effect of passive exposure to cigarette smoke on blood pressure in children and adolescents: a meta-analysis of epidemiologic studies. BMC Pediatr. 2019 May 21;19:161. PMCID: PMC6528314

56. Liu SH, Liu B, Sanders AP, Saland J, Wilson KM. Secondhand smoke exposure and higher blood pressure in children and adolescents participating in NHANES. Prev Med. 2020 May;134:106052. PMCID: PMC8025403

57. Li Y, Xiao D, Dasgupta C, Xiong F, Tong W, Yang S, Zhang L. Perinatal Nicotine Exposure
Increases Vulnerability of Hypoxic-Ischemic Brain Injury in Neonatal Rats: Role of
Angiotensin II Receptors. Stroke J Cereb Circ. 2012 Sep;43(9):2483–2490. PMCID:
PMC3429721

58. Demura M, Demura Y, Takeda Y, Saijoh K. Dynamic regulation of the angiotensinogen
gene by DNA methylation, which is influenced by various stimuli experienced in daily life.
Hypertens Res. Nature Publishing Group; 2015 Aug;38(8):519–527.

59. Stéphan-Blanchard E, Bach V, Telliez F, Chardon K. Perinatal nicotine/smoking exposure
and carotid chemoreceptors during development. Respir Physiol Neurobiol. 2013 Jan
1;185(1):110–119. PMID: 22743051

60. Dos Santos E, Sacramento JF, Melo BF, Conde SV. Carotid Body Dysfunction in Diet-
Induced Insulin Resistance Is Associated with Alterations in Its Morphology. Adv Exp Med
Biol. 2018;1071:103–108. PMID: 30357740

61. Narkiewicz K, Ratcliffe LEK, Hart EC, Briant LJB, Chrostowska M, Wolf J, Szyndler A,
Hering D, Abdala AP, Manghat N, Burchell AE, Durant C, Lobo MD, Sobotka PA, Patel NK,
Leiter JC, Engelman ZJ, Nightingale AK, Paton JFR. Unilateral Carotid Body Resection
in Resistant Hypertension: A Safety and Feasibility Trial. JACC Basic Transl Sci. 2016 Aug
1;1(5):313–324.

62. Niewinski P, Janczak D, Rucinski A, Tubek S, Engelman ZJ, Piesiak P, Jazwiec P,
Banasiak W, Fudim M, Sobotka PA, Javaheri S, Hart ECJ, Paton JFR, Ponikowski P.
Carotid body resection for sympathetic modulation in systolic heart failure: results from
first-in-man study. Eur J Heart Fail. 2017;19(3):391–400.

63. Schlaich M, Schultz C, Shetty S, Hering D, Worthley S, Delacroix S, Reddy V, Sievert H,
Zeller T, Noory E, Boehm M, Mahfoud F, Malek F, Kmonicek P, Neuzil P. 1416Transvenous
carotid body ablation for resistant hypertension: main results of a multicentre safety and
proof-of-principle cohort study. Eur Heart J. 2018 Aug 1;39(suppl_1):ehy565.1416.

64. Sesa-Ashton G, Nolde JM, Muente I, Carnagarin R, Lee R, Macefield VG, Dawood T, Sata
Y, Lambert EA, Lambert GW, Walton A, Kiuchi MG, Esler MD, Schlaich MP. Catheter-
Based Renal Denervation: 9-Year Follow-Up Data on Safety and Blood Pressure
Reduction in Patients With Resistant Hypertension. Hypertension. American Heart
Association; 2023 Apr;80(4):811–819.

65. Jami O, Tijani Y, Et-tahir A. Device-Based Therapy for Resistant Hypertension: An Up-to-
Date Review. High Blood Press Cardiovasc Prev. Nature Publishing Group;
2022;29(6):537. PMID: 36178479

Figure legends

Figure 1. Perinatal nicotine exposure alters RNA transcripts associated with angiotensin

signaling in the carotid body. a) Transcriptomic study design. Nicotine or saline was delivered

to the mother on estrous day 7 (E7) to post-natal day (PND) 21. Bilateral carotid body (CB)

samples were micro-dissected from perinatal nicotine or saline (control) exposed rats on PND

56. (n=5 rats per group- tissues were pooled to yield sufficient RNA for analysis) Two male and

Two female groups per condition. **b)** Principal component analysis (PCA) plot showing distinct

separation between nicotine and control groups. PCA1 was group and PCA2 was sex. **c)**

Heatmap showing overall differences in RNA distribution. **d)** Gene set enrichment analysis

(GSEA) plot for G-protein-coupled receptor activity (0004930) where AgtR1a was significantly

increased in nicotine-exposed offspring compared to controls.

Figure 2. Perinatal nicotine exposure alters DNA methylation in the carotid body. a) DNA-

methylation study design. Nicotine or saline was delivered to the mother on estrous day 7 (E7) to

post-natal day (PND) 21. Bilateral carotid body (CB) samples were micro-dissected from

perinatal nicotine or saline (control) exposed rats on PND 56. (n=5 rats per group- tissues were

pooled to yield sufficient DNA for analysis). One male and one female group per condition. **b)**

Heatmap displaying cytosine-guanine (C-G) and cytosine-arginine/threonine/guanine-guanine

(CHG) methylation differences in carotid bodies between nicotine and saline-exposed rats.

Mapping of methylation changes is confined to the CG and CHG regions as per the heat maps. **c)**

Immunoprecipitation of methylated DNA compared to whole DNA demonstrates an increase of

methylated AgtR1 DNA in the carotid bodies of nicotine exposed compared to control offspring.

ΔCT was used to compare differences between groups (lower value equals greater amount of

methylation). **d)** Enriched gene ontology (GO) terms for C-G methylated DNA between nicotine-

exposed and control pups. **e)** Dot plot showing the gene number and enrichment factor of GO

pathways between nicotine-exposed and control pups.

Figure 3. Nicotine increases AgtR1 pathway in cultured PC12 cells. a) PC12 cells (rat adrenal

pheochromocytoma cells, resemble carotid body glomus cells) were cultured for 6 hours with

and without 50μg/ml nicotine (similar concentration to a single dose of nicotine used *in vivo*), **a)**

angiotensin II type 1a receptor (AgtR1a) was increased by nicotine (T_{28}=2.299) The calculated

increase of AgtR1a was 100.3% in comparison to control. Angiotensin II type 2 receptor

(AgtR2) (T_{28}=0.5307), transient receptor potential vanilloid 1 (TRPV1) (T_{28}=0.2589),

angiotensin converting enzyme (ACE, T_{16}=0.4231), angiotensinogen (T_{16}=1.056), renin

(T_{16}=0.1950), and regulatory genes of endothelial PAS domain-containing protein 1/ hypoxia

inducible factor 2 alpha (EPAS1/Hif2α; T_{28}=0.3774) and Tyrosine hydroxylase (TH- common

PC12 and glomus cell marker, T_{28}=0.6274) were unchanged. Data presented as ΔCT compared to

Hypoxanthine-guanine phosphoribosyltransferase (HPRT) housekeeping gene. **b)** AgtR1 DNA

was hypermethylated in PC12 cells following 6 hours of nicotine exposure. (T_{10}=3.881). **c)** PC12

cells were cultured for 1 hour with media (control) nicotine (Nic, 50μg/ml), Angiotensin (Ang-

50ng/ml), Nic + Ang, Ang+Losartan (AII1R blockade 3μM), Nic+Ang+Losartan, Ang+TRPV1

antagonist (TRPV1x, AMG9810 10μM), Nic+Ang+TRPV1x, Ang+Losartan+TRPV1x, and

Nic+Ang+Losartan+TRPV1x and imaged for calcium flux using Incucyte calcium imaging with

Calcium Orange (forms Calcium esther complex- non-ratiometric imaging). One-way ANOVA

($F_{9,54}$=5.225, p<0.0001) Holm-Sidak post-hoc test, p-values inset in figure.

Figure 4. Carotid body reactivity to angiotensin is increased in perinatal nicotine-exposed

pups compared to control with unaltered serum angiotensin. a) Diagram of *in vivo*

experimental preparation- serum was isolated before injection of angiotensin or sodium cyanide

(NaCN). **b)** A representative trace of arterial pressure of vehicle and perinatal nicotine-exposed

pups on post-natal day (PND) 56 in response to a bolus jugular vein IV injection of angiotensin.

c) The absolute increase in arterial pressure in response to angiotensin in vehicle (grey) and

perinatal-nicotine exposed pups (red) before and following carotid body denervation F_1,

$_{11}$=11.34). **d)** The absolute increase in arterial pressure in response to NaCN in vehicle (grey) and

perinatal-nicotine exposed pups (red) before and following carotid body denervation F_1,

$_{11}$=2.364). **e)** Serum was harvested before administration of bolus injections (T_{14}=0.8077).

Figure 5. Perinatal nicotine-exposed rats have increased mean arterial pressure mediated

by carotid body augmentation of sympathetic activity. a) Mean arterial pressure (MAP), as

assessed by tail-cuff plethysmography, was augmented in nicotine-exposed offspring compared

to controls (T_{20}=7.964). N=11 per group, 2 independent experiments. **b)** Perinatal nicotine-

exposed pups were implanted with radio telemetry to measure arterial pressure and renal

sympathetic nerve activity (RSNA) on PND 56 as per Figure1a. Arterial pressure and renal

sympathetic nerve activity was recorded for 7 days, and then carotid bodies were denervated and

recordings recommenced for an additional 7 days. Raw traces of arterial pressure (AP) and

efferent renal sympathetic nerve activity (vasoconstrictor outflow to kidneys) before carotid body

denervation (Carotid body intact- BEFORE) and following carotid body denervation (Carotid

body denervated- AFTER) in a representative perinatal nicotine exposed rat using telemetry.

Summary statistics in n=5 perinatal nicotine rats comparing the absolute change of MAP and

RSNA. MAP (T_4=3.568), RSNA (T_4=4.762).

Figure 1.

Figure 2.

Figure 3.

Figure 4.

Figure 5.

a

Mean Arterial Pressure (mmHg)

p < 0.0001

Con Nic

b

Before

Arterial Pressure (mmHg)

SNA filtered (µV)

2s

After

Arterial Pressure(mmHg)

SNA integrated(µV)

MAP (mmHg)

p = 0.0234

CB intact CB denervated

RSNA (µV)

p = 0.0089

CB intact CB denervated