LAGOS-US LANDSAT: Remotely sensed water quality estimates for U.S. lakes over 4 ha from 1984 to 2020

Authors

Patrick J. Hanly[1,*], Katherine E. Webster[1], and Patricia A. Soranno[1,2]

Affiliations

[1] Department of Fisheries and Wildlife, Michigan State University, MI, USA

[2] Department of Integrative Biology, Michigan State University, MI, USA

* Corresponding author: hanlypat@msu.edu

June 3, 2025

Abstract

Broad-scale, long-term water quality (WQ) studies are critical for understanding increasing pressures on inland waters but remain rare due to cost. The LAGOS-US LANDSAT dataset provides 37-year remote sensing-derived WQ estimates for thousands of U.S. lakes $\geq$ 4 ha (1984–2020). WQ estimates use machine-learning models with Landsat imagery and ground-truthed Water Quality Portal data (LAGOS-US LIMNO). The dataset includes: (a) 45.9 million whole-lake water reflectance (six bands and 15 band ratios); (b) 740,627 matchups from 13,756 lakes with in situ data for six WQ variables: chlorophyll, Secchi depth, true color, dissolved organic carbon, total suspended solids, or turbidity; and (c) predictions for each WQ variable across lake-time combinations with quality imagery. Two random forest models were fit for each variable: Holdout-data (75/25 spatially representative train-test split) and Full-data (trained on all data). Variance explained for the Full-data predictions ranged from 20.7% for TSS to 63.7% for Secchi depth. Imagery underwent cloud and pixel quality control, and workflow components were validated guiding future research.

Background & Summary

Contemporary studies of lake water quality (WQ) at regional, continental, and global scales are constrained by limited in situ monitoring data. This scarcity of comprehensive and consistent data hinders our ability to assess how WQ has responded to recent global changes, including growing pressures from land-use change and increasing climate variability. Additional approaches are needed to address these data gaps. We present LAGOS-US LANDSAT, a dataset that includes the WQ observations, and also the underlying data, full documentation, and model code to enable not only research using these data, but to enable the development of similar long term datasets in other locations using this method. Our dataset leverages the expanding Landsat image archive of remotely sensed images that can provide estimates of WQ in regions of the globe or time periods that have been poorly sampled. Our objective was to build a database of long-term observations of six water quality variables in tens of thousands of lakes $\geq$ 4 ha in the conterminous US from 1984 - 2020 using remotely sensed imagery from the Landsat SR data product. We used Landsat because it is the longest-running earth-observing satellite platform and has been used extensively in methods development for measuring inland WQ[1-8].

Previous studies, particularly on US and Chinese lakes, have advanced methods for data integration and WQ modeling to leverage Landsat for broad-scale studies[1,7,11–13] (**Table 1**). For example, AquaSat[7] is an open-access dataset for the conterminous US containing match-ups between surface reflectances from Landsat SR and in situ WQ observations collected within one day. Predictive models for specific WQ components, such as Secchi depth[11] and trophic categorization based on the Nutrient-Color Paradigm[13], have also been created at the US national scale. These databases are supported by well-documented and fully reproducible methods that enable other researchers to both use their data and build new applications off them. Other broad-scale data products have more limited reusability, such as ground-truth Secchi depth predictions provided as 5-year averages for over 10,000 Chinese lakes across the Landsat platform period[14]. Additional global datasets that focus more on addressing scientific questions rather than producing reusable data products have been published for Secchi depth[15] and for algal bloom metrics[15–18]. However, none of these studies provides long-term estimates of multiple WQ variables or for lakes as small as 4 ha.

Our study and data release[19] builds on this prior work to advance the use of Landsat SR for broad-scale studies of inland waters. Although using Landsat and ground-truthed WQ observations to predict WQ in unsampled lakes has a long history, many workflow components remain untested at the scale of

thousands of lakes. We present results from these tests to guide future applications. Our advancements
include: (1) testing computationally expensive components of the workflow to determine when and where
they are needed to improve predictions; (2) creating a single data product with six WQ variables in
addition to the reflectances, match-ups with in situ data, ground-truthed predictions, and data quality
flags; and, (3) creating and testing the validity of using Landsat SR for a broader range of WQ variables.
Our final data product meets the FAIR data principles of findability, accessibility, interoperability, and
reusability (20) and we include validation of predictions, methods, and recommendations for specific
processing steps.
Our dataset provides ground-truthed estimates for six WQ variables from 1984 to 2020, derived
from over 45 million images from Landsat 5, 7, and 8 for 136,977 lakes, with a median of over 121,000
lakes sampled annually. In addition to the WQ estimates, we include: five data flags for users; reflectance
values on all individual bands and 2-band ratios for all lake-date combinations; Landsat platform
identifiers and quality metrics; and our matchup dataset that includes matchups within 0 to 7 days in
either direction. The dataset is fully interoperable with the LAGOS-US research platform, which provides
millions of data points on hundreds of lake features relevant to WQ including delineated watersheds,
regional characteristics, land use, lake depth, and other variables[21–23].
The six WQ variables are chlorophyll a (CHL), Secchi depth, true color, dissolved organic carbon
(DOC), total suspended solids (TSS), and turbidity. Chlorophyll a (CHL; μg/L) estimates suspended
phytoplankton biomass in the pelagic zone, corrected for pheophytin, serving as a key measure of lake
productivity. Secchi depth (meters) measures water clarity using a disk lowered until no longer visible.
True water color (true color) measures the colored fractions of dissolved organic material using filtered
water samples, reported in platinum cobalt units (PCU). Dissolved organic carbon (DOC; mg/L)
measured from filtered water, includes both colored and non-colored fractions of organic carbon. Total
suspended solids (TSS; mg/L) include silt, clay, sand, algae, and plant material suspended in water.
Turbidity, another measure of water clarity, measures light scattering due to suspended particles and is
measured in Nephelometric Turbidity Units (NTU). Together, these variables provide a comprehensive
snapshot of WQ status and biological functioning in inland waters, reflecting eutrophication pressures,
land-water interactions, climate change, hydrologic forcing, and food web dynamics. While CHL, true
color, DOC, TSS, and turbidity represent specific components of dissolved or suspended particulate
matter, Secchi depth provides an integrated measure of all components. Full definitions, source
information, and method descriptions for these six variables are described in the documentation of
methods for the LAGOS-US LIMNO data product we used for our in situ WQ data[10].
104
105

Methods
We developed a reproducible workflow for generating long-term, remotely sensed water quality
(WQ) predictions for 136,977 lakes ≥ 4 hectares across the conterminous United States (1984–2020).
Following established approaches, we used the Landsat Surface Reflectance (SR) data product to develop
ground-truthed WQ predictions over time. We used Google Earth Engine (GEE) to extract reflectances
for over 100,000 lakes across the 37-year period. Because many methodological components for inland
water remote sensing remain untested, we evaluated several key workflow components to guide future
researchers using Landsat SR data in GEE for large-scale WQ predictions. Our overall approach,
illustrated in **Figure 1** for an individual WQ variable, was repeated for all six variables. Detailed methods
for each step follow.

(1) Process Landsat SR scenes

117
We used the Tier 1, Level-2 Landsat Surface Reflectance product (Collection 1), which employs
the Landsat Ecosystem Disturbance Adaptive Processing System (LEDAPS)[24] to atmospherically correct
scenes from Landsat 5 (TM) and 7 (ETM+) and the Land Surface Reflectance Code (LaSRC)[25] to
atmospherically correct scenes from Landsat 8 (OLI). We extracted surface reflectance values for all
137,465 LAGOS-US lakes ≥ 4 hectares in surface area[21], from 1984-2020 (1985-2020 complete years)
using GEE (**Figure 1,** step 1). We selected 4 ha as our minimum lake size because any smaller area
results in large numbers of lakes with pixels that have mixed water and land components, making it
challenging to capture true water-only pixels. Because lakes were run as separate instances and the default
average concurrent tasks allowed on GEE at the time of execution was two, the total runtime for all lakes
took over a year to complete and increases substantially with lake size in our method where all pixels in a
polygon are extracted.
For each lake and for each sensor overpass and band, we extracted the median, minimum, and
standard deviation of individual pixel reflectance values, and band ratios calculated at the per-pixel level
before a lake-wide median was extracted. The potential area for extraction was fixed to the static
polygons from LAGOS-US LOCUS for the full time series. We removed pixels coded as clouds, cloud
shadows, snow/ice, or fill data, retaining only clear-sky pixels over water using the CFMASK-derived
pixel quality attributes[26] in the Landsat Pixel Quality Assessment Band. To enable historical time series
analyses across the full study period we harmonized bands across the TM, ETM+, and OLI sensors, for
which we used complete handoff following the launch of a new sensor, using published surface
reflectance transformations[6] prior to reflectance extraction.
As part of our data product, we created the data table, *LAGOS_US_LANDSAT_compiledRS*,
which contains output from the steps described in **Figure 1a** to aid in later steps and to make data
available to other users for their modeling purposes in the future. Each row in the data table represents a
unique lake-scene combination and includes: median lake-wide values for the 6 bands available for all
sensors, pixel-specific ratios for all possible band combinations (n=15), the percent cloud cover of each
scene, the total number of pixels extracted, and a 3BDA-like "KIVU" algorithm applied to the individual
pixels[27]. We also created data flags for: (1) any pixel or any band with negative reflectances; (2) any lake
median reflectance that is negative; (3) the percent pixels retrieved for a lake; and (4) duplicate scenes on
the same day for a lake (rare, but present due to overlapping overpasses). Our final data product
represents 136,977 lakes (99.6% of LAGOS-US lakes ≥ 4 hectares) that had at least one set of Landsat
reflectances and provides 45,867,023 reflectance sets of lake and overpass combinations through time.
We retrieved an average of 48.0% of all pixels within a lake polygon, with 37,529,125 lake-scene
combinations meeting or exceeding our 10 percent pixel threshold (81.8%). Finally, 4,877,886 lake
reflectance sets had shared calendar days for a lake due to scene overlap (1.1%).
152
(2) Build a matchup dataset for six water quality variables
Our matchup dataset **(Figure 1, step 2)** is based on the LAGOS-US LIMNO data product[10] that
contains in situ data extracted primarily from the Water Quality Portal and three sampling years from the
USEPA National Lakes Assessment. We conducted quality control analyses on each of the six WQ
variable matchup sets and made the following data decision. First, for CHL, we removed negative values
and extremely large values > 800 ug/L that are rarely measured in pelagic zones of lakes. Second, if a
lake had multiple measurements for a given WQ variable on a calendar day, the median value was

calculated prior to matching with remote sensing data. Third, because the few observations for true color > 100 PCU appeared to skew the distribution and were from a small number of primary data sources, we removed them from the training set, recognizing that our resulting models would likely not be able to capture the most extreme high range of true color. Nevertheless, because lakes are considered tannic if they have values > 20 PCU[28] and highly colored lakes were rare, we made the decision to apply a data limit in order to produce more accurate training models. The ranges of data for all other variables were less skewed and were not trimmed.

After merging the remote sensing dataset with the LIMNO dataset on a calendar date basis (with a maximum matchup window of +/- 7 days), we had 723,206 matchups of lake-date combinations with at least one observation for any of the six WQ variables across 13,756 LAGOS-US lakes. The number of matchups (including multiple values by date per lake) by WQ variable were: CHL = 238,248; Secchi depth = 666,060, DOC = 8,269; true color = 37,839; TSS = 52,677; turbidity = 68,573. Matchup lakes were widely distributed across the conterminous US with high densities of observations related to regional lake density and monitoring program specifications (**Figure 2**). Although substantial matchup data exist across the study period, changing monitoring intensity and resulted in an uneven temporal distribution, with matchup numbers peaking around 2008 (**Figure 3**). The complete set of matchup data that are within +/- 7 days is provided in the data table *LAGOS_US_LANDSAT_matchups*.

(3) Create the data test sets for each water quality variable

We tested major methodological components related to image and pixel processing, modeling, and matchup windows (**Figure 1, step 3**). Using the matchup dataset, we conducted exploratory assessments of key workflow components for training models for all six water quality variables. Because methodological decisions were interrelated, analyses were not performed in linear order. While most tests included all water quality variables, computational constraints in GEE limited some tests to one to three variables. For example, using a randomly selected lake matchup dataset of 500 lakes with in situ DOC data, we ran a set of 40 training models that varied the image and pixel processing steps, transformations, and matchup windows. In addition to identifying the best-performing model type (**Figure 4**), we tested the influence on prediction quality for the following model inputs (**See Figure 1, step 3**): 5 types of pixel retrievals; 2 types of band ratio calculations; 2 types of scene quality indicators; and inclusion of all bands or all bands plus all ratios. Because many components that we tested have not been analyzed in any published studies to date, the outcomes of these tests are informative not only for our final model selection but also for future studies using the Landsat SR platform for broadscale water quality studies.

Table 2 summarizes seven main outcomes from our workflow tests for building large-scale remote-sensing models using the Landsat platform. All tests were conducted for all water quality variables and lakes, except the first two steps, which were either computationally prohibitive or unnecessary for the full dataset.

Image and pixel processing

1. **Pixel-retrieval calculated at the whole-lake scale performed best**. The lake center with a buffer performed close to the whole-lake estimates while the lake center calculated using the Chebyshev point was consistently better than the lake centroid (**Figure 5a**). This test was extremely resource-intensive and so we only tested it for a subset of lakes and for a subset of water quality variables (CHL, Secchi depth, DOC).

2. **Calculating band ratios at the pixel level was always better,** i.e., with lower prediction error, than first calculating lake-wide or buffer-wide band averages and then calculating ratios from those (Figure 5a). Because this step must also be conducted within GEE, it was critical for consideration early in the workflow.

3. **Removal of negative reflectances and scenes > 50% clouds was best (except for DOC).** The results for DOC were likely due to the lower sample size of the matchup data. DOC (**Table 2**; **Figure 5a**). However, the differences in this test were not large, so when sample sizes are low, retaining some scenes with poorer quality may be better because sample size is increased. In general, these poor-quality images were relatively rare, so the gain in predictive performance was not large across many lakes.

Matchups

4. **The optimal matchup window ranged from 1 to 7 days across the WQ variables and should always be tested.** We found the optimal matchup window differed by WQ variable and was likely due to the inherent daily variability in the WQ variable (**Table 2; Figure 5b)**. For example, true color is a more temporally stable variable for many lakes than CHL, TSS, and turbidity which can all respond readily to large precipitation or wind events day to day. Consequently, the optimal matchup window for the latter 3 variables was 1 day but was 7 days for Secchi depth, true color, and DOC. Nevertheless, we encourage users to test this important part of the workflow in future applications, and the results may be specific to the type of lakes being modeled.

Modeling components

5. **The machine learning model performed best**. We tested three types of models that have been used by other studies using the CHL dataset: linear models with all bands plus band ratios; the three-band KIVU retrieval algorithm[29] that was previously found to be relatively effective for inland water chlorophyll retrieval[27]; and random forest regression models using all bands plus band ratios. We found that the random forest regression model, applied using the 'randomForest' package in R[30] far outperformed the other two models with over two times reduction in error compared to the other two models (**Figure 4**). We found similar gains in performance for Secchi depth data (data not shown). Based on this test that was conducted only for CHL and that is backed by prior studies demonstrating the superior performance of machine learning models[11], all of the remaining tests were performed only using the machine learning model.

6. **Using all bands and all band ratios outperformed RGB-only models**. Although we present initial results only for DOC (**Figure 5a**), we retained the testing of these model components throughout the process. The variable importance plots for the final models for all WQ variables demonstrate that although the non-RGB bands were never the top predictor, ratios containing one of these bands were often the second or third most important variable (see **Figure 6**).

7. **Log-transformed in situ data resulted in better-performing models** than their non-transformed versions (**Table 2**, final column). Because the distributions of most of our WQ variables were left-skewed and more data-rich in the lower ranges, the models performed better when the transformations resulted in greater granularity at the lower ends of the data distributions.

(4) Conduct two-stage modeling: Holdout-data validation and Full-data prediction

We implemented a two-stage modeling approach to balance the needs of different users while
maximizing predictive accuracy. This dual approach provides both traditional error metrics through a
spatially-stratified holdout validation procedure and optimal predictions using all available data. While
we recommend the Full-data model predictions for most applications due to their superior coverage of
spatial and temporal variability, we recognize that some users require traditional train-test validation
metrics for comparison with other studies or regulatory compliance.

For our first modeling approach, we created a Holdout-data model test set based on a 75%
training and 25% test split to provide external validation metrics. We used the Generalized Random
Tessellation Stratified (GRTS) sampling method, implemented in the `spsurvey` R package[31] to create
spatially representative splits. For our second modeling approach, we constructed a Full-data model
leveraging the entire in situ dataset to maximize predictive accuracy across the full range of spatial and
temporal variability in lake water quality. Because random forest models internally use out-of-bag (OOB)
samples to assess prediction error, we used OOB variance explained as our primary performance metric.
Prior studies have shown that OOB estimates provide a reliable alternative to test sets when sufficient
trees are grown[32-33]. This method avoids sensitivity to a single train-test split, a known issue in ecological
modeling[34]. We recommend the Full-data model and its OOB variance-explained for most ecological
applications. The Holdout-data model is provided for users requiring traditional train-test error metrics.

Both modeling approaches followed identical workflow steps for each WQ variable. Our
approach used a modified grid search to explore combinations of data filtering and workflow components,
aiming to identify optimal model–data pairings for WQ prediction. We fit random forest regression
models to each WQ Full-data dataset. The components varied included: matchup window (0, 1, 3, 5, or 7
days); modeling inputs (RGB, all bands, all bands plus whole-lake ratios, all bands plus pixel ratios); and
data transformation (original units or $\log_{10}$-transformed). After selecting the best-performing model from
these 40 random forest models based on out-of-bag (OOB) variance explained we sequentially applied
two additional filters: first, removing high cloud-cover scenes from the training data; and second,
excluding matchups with any negative reflectance values in any band. In total, 42 models were fit
separately for each water quality variable. The best-performing model for each variable was then selected
to generate the final predictions. We then applied the same data filtering and modeling components within
the Holdout-data model to better characterize the performance of the Full-data model for each WQ
variable.

Table 2 details the final model specifications and workflow steps with data decisions specified in
each row. Final models were based on in situ training data that were $\log_{10}$-transformed (except for Secchi
depth). Variance explained of predicted versus in situ data, a valuable model diagnostic, ranged from a
low of 20.7% for TSS to a high of 63.7% for Secchi depth, ranging between 42-52% for the remaining
WQ variables. Thus, except for TSS, the variance explained is quite high and certainly suitable for broad-
scale water quality studies. All random forest models included all bands and ratios. We found that the top
predictors were always a 2-band ratio based on variable importance factors calculated for each WQ model
(**Figure 6)** like other studies demonstrating the importance of the blue/red ratio and blue/green ratios for
Secchi depth and CHL. However, fewer studies have also included the SWIR bands, which were also
quite important. Interestingly, the top predictor for true color was the Green/Red ratio, which was also the
second highest ratio for closely related DOC, although, for DOC, the top predictor was Blue/Red, which
was also top for Secchi depth and CHL. These results make ecological sense since DOC is related to both
true color (which is the colored fraction of DOC) as well as CHL which is known to track with the non-

colored fraction of DOC. Turbidity and TSS have similar dominant predictors, although the contributions of the other wavelengths and ratios differ suggesting they reflect distinct WQ features.

Our data product includes five quality-control flags, enabling users to filter data according to their study requirements:

1. Negative reflectance (any band) - flags lake-date combination where any band has negative value for at least one pixel
2. Negative median lake reflectance (any band) - flags lake-date combination where any whole-lake retrieval has a negative median value
3. Percent of lake pixels with quality reflectances - flags scenes with <10% of maximum lake pixels containing quality reflectances (e.g., due to cloud cover)
4. Duplicate predictions - identifies calendar dates with two satellite overpasses resulting in multiple predictions
5. QAQC_recommend - integrates the previous flags (2-4), removing data with negative median reflectance values and selecting the duplicate date prediction with the greatest number of pixels, to identify highest quality predictions that users can easily query to easily meet these criteria

Data Records

The LAGOS-US LANDSAT data product[19] consists of the following four data tables that include all components of our workflow described in **Figure 1**.

LAGOS_US_LANDSAT_data_description.csv

This table is the combined data dictionary for the three tables below that describes and defines variables included in each of the four data tables. For each variable, we include the variable name, description, data type, and column index.

LAGOS_US_LANDSAT_compiledRS.rds

This data table, in the form of an R Data Serialization (RDS) file, provides reflectance extraction output generated from Google Earth Engine that was extracted from a single lake at a specific time. Each row is a single LAGOS lake and Landsat scene combination. This data table contains 57 variables related to whole-lake summaries of reflectance values calculated for lake water pixels on all individual bands (min, max, median, standard deviation) and all 2-band ratios in addition to the KIVU calculation. Ratios are first calculated at the pixel level (referred to as pixel-wise in the data dictionary), and then the median is calculated across all lake-water pixels. We also include scene identifier information from the Landsat platform (satellite name, sensing time, scene ID, etc), quality estimates obtained from the Landsat platform (e.g., % cloud cover of the scene, image quality indicators), earth-sun distance, solar angles, etc.

LAGOS_US_LANDSAT_matchups.csv

This data table includes all possible matchups within 0 to +/- 7 days between Landsat overpasses and in situ data obtained from the LAGOS-US LIMNO[10] data module for the six WQ variables (CHL, Secchi depth, True color, DOC, TSS, Turbidity). For reference, we include the WQ variables from individual dates and the associated reflectance values and scene identifier information from the above dataset. If any date had more than one WQ estimate, we took the median value, however, more than one estimate was extremely rare, and the vast majority of data were from a single sampling event per day. We also include

333   the matchup window value for each WQ in situ value so that future users can select the matchup window.
334   Note that not all WQ variables are sampled on the same dates, so the sample sizes vary by WQ variable.
335
LAGOS_US_LANDSAT_Predictions_v1_QAQC.csv
The predicted WQ data from the **Full-data model** that uses all available data for predictions for the
selected matchup window. As described above in step 4 of our workflow, we include five data flags
including the field QAQC_recommend in which we apply our criteria for quality predictions that users
can select for a highest-quality version of the dataset. This dataset includes predictions using all data
within the model-selected matchup window to minimize prediction errors.
342
LAGOS_US_LANDSAT_TT.zip
The predicted WQ data from a **Holdout-data model** per WQ variable that is based on 75% train and 25%
test datasets that is spatially balanced using the Generalized Random Tessellation Stratified (GRTS)
algorithm. This dataset has predictions alongside measurements of prediction error from the training
datasets using OOB variance explained and MAPE on the test datasets for each WQ variable. These
datasets are useful for creating comparisons with other machine learning algorithms and for the
generation of test data error estimates that are unseen by any tree in the modeling process.
350

Technical Validation

The validity of using Landsat 5, 7, and/or 8 sensors for inferring water quality in lakes has been
supported by decades of studies conducted on individual and combination of sensors[3,11,35–39]. In addition,
the Landsat Collection 1 SR product has been noted for its suitability for machine learning estimation of
CHL across the US[40]. However, challenges remain that careful methods development can address. For
example, the spectral signals of WQ variables are not always independent; for example, colored dissolved
organic matter and chlorophyll have overlapping absorption spectra[41]. Further, Landsat imagery may have
lower accuracy for estimating WQ values in oligotrophic relative to eutrophic waters[42]. Therefore, we
used a series of analyses to characterize the level of uncertainty in the LAGOS-US Landsat data and
identify potential biases. In addition to quantifying numerous fit statistics and conducting validation that
supports the use of the six models that we developed (**Table 2**), we also investigated the ecological
interpretability of the predicted water quality estimates such as by comparing in situ and predicted
individual lake time series.
364

Model prediction error
In addition to examining the overall model fits that quantify model error across the whole dataset,
we examine quantiles of predictions to examine whether errors differ across the range of the matchup
dataset for both the Holdout-data (**Figure 7**) and Full-data (**Figure 8**) models. We found that in most
comparisons, 90% of the data were close to the 1:1 line and that differences between predicted and in situ
data are not that far off ecologically, especially when one considers predictions of water quality were
made in over 100,000 lakes across 34 years. The model with the poorest performance was TSS (**Table 2**),
which had the largest bias in predictions, with the median TSS being underpredicted relative to in situ
data. Plots of error by binned values of predicted values also show that TSS has higher errors, particularly
for high TSS lakes. We also found error by binned values for DOC to also be particularly high for the
highest DOC values (> 16 mg/L). A study of global lake DOC estimates predicted the average DOC was

3.9 mg/L, with a maximum value of 27.0 mg/L[43]. Therefore, while our modeled DOC values are more
accurate for values < 16 mg/L, predictions at rarely observed higher values have more error.
Both sets of prediction errors provide similar conclusions where the OOB variance explained by
the Holdout-data and Full-data models were substantially identical for each WQ variable. MAPE values
were generally within the 30% range. Despite these differences, model predictions for the same 75%
GRTS subset used in both frameworks were highly consistent, with R^2 values ≥ 0.92 across all WQ
variables (**Figure 9b**). This consistency supports the reliability of both modeling approaches and affirms
the utility of the dataset for large-scale ecological inference, provided users account for known sources of
error using the supplied QA flags.
385
Model validation using individual lake data
To evaluate model performance at the individual lake scale, we analyzed lakes with dense, high-
quality time series of in situ data and satellite imagery for three different WQ variables. First, we looked
at Secchi depth data for Lake Okeechobee, a well-studied lake in Florida that contained 10,158 matchups
within the ±7 window with satellite overpasses with 2,989 Secchi depth measurements. In this
comparison, higher prediction errors occurred in concert with known issues with imagery data that we
have flagged for in our database. Secchi depth was overestimated when the proportion of lake pixels
available was low (< 10%; **Figure 10a**), an effect that is removed by eliminating such predictions from
analysis. Second, in the same set of data from Lake Okeechobee, we found that negative reflectance
values also led to prediction errors (**Figure 10 b-c).** Removing date-image combinations flagged with
these errors results in the majority of the predicted versus in situ data falling close to the 1:1 line (**Figure
10d).**
Further validation of the LAGOS-US LANDSAT dataset comes from a lake which has
documented changes in water quality following a deliberate management intervention. The eutrophic
Lake McCarrons in Minnesota was treated in 2004 with aluminum salts to reduce internal phosphorus
loading, and, consequently, decreased algal growth[44]. The comparison between predicted and *in-situ* data
for CHL suggested first, a close relationship near the 1:1 line (**Figure 11a**). Second, the alignment
between the in situ data and predicted values improved when a narrower matchup window was applied
(**Figure 11b-c**). However, the most striking was that the timing of the dramatic decrease in the in situ
CHL was matched exactly in the predicted values from the remote-sensing models (**Figure 11d**). This
lake thus presents an excellent case study to examine the application of our data product to individual lake
studies beyond our primary goal of facilitating regional and continental-scale research.
Finally, we demonstrate how our remote-sensing-based prediction dataset can be used to extend
time series in lakes for which in situ data are missing. For example, DOC is not sampled as commonly as
other water quality variables and yet is an important regulator of the ecology and biogeochemistry of
inland waters, particularly related to carbon budgets[45]. Beaver Lake, Arkansas has a robust time series
beginning in 2002 for in situ DOC data that aligns well with predicted DOC (**Figure 12a**). With our
remote-sensing predictions, we can extend that time series back in time an additional 18 years (**Figure
12b)**. These individual lake studies lend strong support for the potential for greatly expanding the
estimates of water quality for individual inland lakes across the US in lakes with incomplete records or
that have never been studied to date. Being able to estimate water quality for lakes across the entire US is
an urgent priority to ensure that people have equitable access to good water quality regardless of
socioeconomic status, race, ethnicity, and other factors. A recent study has shown that in situ estimates of
water quality from government agencies and community science programs are not equitably distributed

across all communities based on race and ethnicity[46]. Remotely sensed water quality predictions such as
ours can help fill this gap in sampling.
422
423

Usage Notes

Data quality flags and filtering

In addition to the data values, our prediction dataset includes multiple measures of quality that provide
users with the information needed to apply more lenient or stricter data quality filters: (1) if any pixel in
any band for a lake retrieval had a negative reflectance value, (2) if any band for the whole lake retrieval
had a negative median reflectance value, (3) the percent of pixels retrieved out of the maximum number
of pixels ever retrieved for the lake across the prediction time series, and (4) if the retrieval for a lake
shares the same calendar date as other retrievals due to overlapping scenes. The first two flags represent
overcorrection over lake water pixels or other scene artifacts while the third provides an estimate of the
extent of the lake surface represented in the predictions. Of all predictions, 3,145,912 (6.9%) were flagged
for any negative reflectance pixel while 410,065 (0.9%) were flagged for median negative reflectance
across the whole lake. Finally, because a lake can be within multiple Landsat row and path combinations
due to scene overlap, we additionally flag predictions that have multiple retrievals for a single calendar
date since users may want to treat these data differently than predictions for a single day. To summarize
across these data flags, we include a variable called *QAQC_recommend* that identifies (with a value =
TRUE) what we define to be research-ready and good quality data based on the following criteria:
removal of retrievals with any median negative reflectance for any band, a threshold of least 10% of
pixels being retrieved per lake, and removal of lower quality duplicate day predictions. The dataset
filtered in this way includes a total sample size of 35,794,821 lake and date combinations and includes
78.0% of the total prediction sets. Another layer of quality control that users may want to consider is to
drop predictions for DOC that exceed 16 mg/L and TSS that exceed 80 mg/L (**Figure 7**).
445

Spatial and temporal distribution of predictions

The location and density of predictions for the six WQ variables are widely-distributed across the
conterminous US (**Figure 13**). Due to cloudiness that is higher in the eastern US compared to the western
US, the highest densities of prediction data occur in spatial banding patterns that result where Landsat
overpasses overlap, and are more likely to contain cloud-free scenes that are known to occur with the
Landsat platform[47]. A decrease in the spatial banding of the data can be seen from east to west and is
likely a function of regional cloud patterns across the US. Because our dataset began with all lakes ≥ 4 ha,
it is close to a census population of lakes, with the majority of lakes having at least 1 scene. Observations
are also distributed across time (**Figure 14**), with peak numbers of scenes occurring in April to May and
then October through November - again, likely a result of general patterns of clouds. The number of
images extracted through time increases through time as well, with the largest retrievals per year
occurring after around 2015 that contain overlapping satellites that increase the potential for cloud-free
scenes. Nevertheless, over 70,000 lakes across the US contain at least one summer water quality estimate
across the full time period, resulting in an unprecedented time series dataset for lake WQ.
The ranges of the matchup dataset include broad ranges of all six WQ variables (**Figure 15;
Table 3).** The prediction dataset generally had narrower distributions of the data as shown by the quantile
plots that showed less spread in the data. However, given that we know that most lakes in the US have
never been sampled (Shuvo et al. 2023), we would not have expected the ranges of the matchup dataset to

match the prediction dataset closely given the biases in water quality sampling programs[48,49]. Examining

sampling sizes through time is also valuable. The median number of observations per lake is 257 (**Table**

3); the median lake-year combination with WQ observations is 37; and the median number of lakes per

year with observations is 121,812.

Water quality variable distributions and correlations

Although there is overlap in reflectances that these six WQ variables are related to (e.g,

overlapping absorption spectra for CDOM and chlorophyll documented by Olmanson et al. 2015), the

dominant factors that these WQ variables represent are distinct enough that they are predicted by different

bands and band ratios (**Figure 6**), and, despite very high sample sizes, they are not highly correlated with

the exception of Secchi depth which is highly correlated to everything except true color (**Figure 16**). In

fact, these correlations match expected ecological relationships. For example, Secchi depth and turbidity,

both measures of water clarity, are the most strongly correlated in the dataset (r=0.90). In contrast, the

correlation between true color, which measures dissolved fractions, and TSS, which measures suspended

particles, is the lowest among all comparisons (r=0.3).

To examine relationships among the variables further, we also applied a common threshold for

eutrophication that separates oligo-mesotrophic lakes from eu-hypereutrophic lakes (CHL ≤ 7 ug/l) and

examined relationships among CHL, Secchi depth, and DOC. The patterns (**Figure 17)** make ecological

sense based on our current understanding of key limnological relationships [28,50]. First, DOC is positively

related to true color in both sets of lakes, but DOC at a given color value tends to be lower in oligo-

mesotrophic lakes than in eu-hypereutrophic lakes. These latter lakes have the highest DOC values in the

dataset, particularly where with true color > 20 suggesting high contributions from both non-colored DOC

from high algal growth and colored DOC from tannins that occur in high color waters. Second, Secchi

depth versus true color also diverges in the lakes in the two trophic classes, with a far clearer relationship

between the two variables in oligo-mesotrophic lakes compared to eu-hypereutrophic lakes suggesting a

shift in control of water clarity from color to algal biomass. Finally, the importance of Secchi depth and

its relationships with all other variables is demonstrated in **Figure 18**. Secchi depth is the most commonly

sampled WQ variable in US lakes[10], and its inclusion in nutrient modeling of lakes has been found to

decrease prediction error[51,52]. Being able to predict Secchi depth in an even broader range of lakes has

great potential for improving water quality modeling at broad scales.

Filtering and combining predictions for temporal analysis

Users can filter and combine LAGOS-US LANDSAT predictions to support diverse modeling

frameworks and research questions. For example, in Soranno et al.[53], retained only lake-date

combinations with the highest quality data based on the QAQC_recommend flag, and lakes that had at

least eight valid chlorophyll-a observations per summer over a 34-year period, excluding lakes with more

than one consecutive year missing. These criteria resulted in annual summaries of lake CHL in 24,452

lakes that were used in advanced time series models that require dense, temporally regular observations.

Therefore, users conducting time series analyses to address other types of research questions might

consider: (1) setting minimum observation thresholds per lake-year, (2) excluding pre-2000 data due to

lower scene availability, (3) restricting analyses to ice-free months (e.g., April–October), and (4) applying

the QAQC_recommend flag for highest quality predictions. The data density of predictions per year for

lakes is shown in **Figure 19**.

Summary
LAGOS-US LANDSAT is the first published dataset of remotely-sensed, ground-truthed estimates of
multiple WQ variables providing near-census coverage of lakes ≥4 ha (>100,000 lakes) at the continental
scale that also includes:
● Coverage of small lakes (≥4 ha), with 50% of observations from lakes <9 ha
● Whole-lake water pixel extractions of all bands
● Complete workflow products in a single, integrated dataset including data quality flags
● Full interoperability with the LAGOS-US research platform
The LAGOS-US platform provides comprehensive geospatial and temporal datasets for macroscale lake
studies (watershed delineations, land use/cover, geology, soils, climate, stream connections). This
LANDSAT module adds 37 years of water quality data on > 100,000 lakes, enabling unprecedented long-
term analyses of lake ecosystems at macroscales.

520
521
522

Code Availability

524 The entire code, including the code used for extracting reflectances in Google Earth Engine, is available
525 in Zenodo at https://doi.org/10.5281/zenodo.11110283[54]

526
527

Acknowledgments

We thank the many contributors who created the LAGOS-US research platform, especially those
researchers who created the LAGOS-US LIMNO data module, and the original data providers for this
rich source of in situ data without which this data product would not be possible. This research was
funded by the National Science Foundation (DEB-1638679). PAS also acknowledges support from the
National Institute of Food and Agriculture, Hatch Project 176820. This material is based upon work
supported by (while serving at) the National Science Foundation from 2019-2023 for PAS on her
independent research and development time. Thanks also to Yvonne Allen, Spatial Ecologist, US FWS-
Southeast Region for the idea of using the Google Earth Engine platform for inland aquatic studies of
water quality parameters and for a very helpful review of an earlier draft.

Author contributions

540 PJH had the initial idea and developed and implemented the analytical approach for this dataset. All
541 authors contributed to making decisions about the modeling approach as well as outlining and writing the
542 article. PJH created all figures and tables except for Figure 1 and Table 1-2 (PAS) and Figure 14, 16-18,
543 and Table 3 (KEW).

Competing interests

546 The authors declare no competing interests.

Figures

Figure 1. Overview of the four major steps used to create LAGOS-US LANDSAT. See text for full description of all steps. The steps are described for an individual water quality variable (e.g., CHL). Because the same dataset in step 1 is used for all models, steps 2-4 are repeated for each water quality variable.

Figure 2. Maps of the location of lakes in the LAGOS-US LANDSAT matchup dataset by water quality variable. For display purposes, we show the matchup data that are within ±7 calendar days of the LAGOS-US LIMNO in situ observations for each of the six water quality variables which represents the maximum possible matchup dataset for each variable and the maximum sample sizes and what was tested in our modeling workflow. See results in Table 2 for the final matchup windows used for each variable as well as the final sample sizes used. The map outlines are NEON region borders.

570
571 **Figure 3.** Histogram of the number of matchups by year for the matchup dataset. The matchup window
572 includes all dates within ±7 calendar days of the LAGOS-US LIMNO in situ observations for all water
573 quality variables as a sum across all variables within each year.
574
575
576
577

578
Figure 4. Results from the three model approaches tested for the CHL model. The left panel shows the
reverse cumulative distribution of the absolute value of residuals of the three models using log10-
transformed in situ CHL data and using matchups within ±1 day. See text for model details. 'Random
forest' is a random forest regression model that includes all bands and band ratios; 'Linear model' is a
linear multiple regression model with all bands and band ratios; and 'KIVU' is a 3-band algorithm
calculated at the pixel level for each lake. The right panel shows boxplots of these same absolute values
of residuals with the red dot indicating the RMSE: random forest = 0.23, linear model = 0.51, 'KIVU' =
0.54.
587

588
589
Figure 5. Results from testing different components of the workflow related to image and pixel
processing for a subset of the DOC dataset. (a) A plot of variance explained for each training model run
conducted on $\log_{10}$-transformed in situ data that included the indicated components. Results are plotted
for the 5 different ways that reflectance is summarized by lake (see text). 'Whole-lake Ratios' are ratios
calculated on median band data, whereas 'Pixel Ratios' are those ratios first calculated at the pixel-level
and then the median value is calculated; 'All Bands' includes all bands; 'Overcorrection Filter' is the flag
for when there is negative median reflectance. (b) A plot of % OOB variance explained by matchup
window and the 5 different ways that reflectance is summarized by lake. Models are based on a 500-lake
subset that contained DOC matchups, with some lakes being sampled more than once (N = 1,348).
599

601
602
603 **Figure 6.** The full-data model variable importance (mean decrease Gini) for all predictors in the random
604 forest regression models for each water quality variable. Orange bars are individual bands, and purple
605 bars are ratios between two bands.

612
613
Figure 7. Fit statistic plots for the **Holdout-data model** of each of the six water quality variables in
original units. The top six plots show predicted versus in situ quantile lines for the median quantile
(black) and the 5th, 25th, 75th, and 95th quantiles (shaded colored lines). The maximum range of the X
and Y axes is trimmed to the 90th percentile of in situ data (CHL = 53 ug/L; Secchi depth = 5.3 m; true
color = 63 PCU; DOC = 13 mg/L; Turbidity = 19 NTU; TSS = 133 mg/L). The dashed red line is the 1:1
line. The internal random forest OOB variance explained from the model training data as well as the r-
squared value of the holdout data are shown. The bottom 6 plots show the absolute percent error between
predicted and in situ values in the model training dataset by intervals of predicted values for the training
dataset. The red dashed line is the threshold identified as above which predictions are thought to be less
reliable. Outliers are not plotted.
624
625

Figure 8. Fit statistic plots for the **Full-data predictive model** of each of the six water quality variables in original units. The top six plots show predicted versus in situ quantile lines for the median quantile (black) and the 5th, 25th, 75th, and 95th quantiles (shaded colored lines). The maximum range of the X and Y axes is trimmed to the 90th percentile of in situ data (CHL = 53 ug/L; Secchi depth = 5.3 m; true color = 63 PCU; DOC = 13 mg/L; Turbidity = 19 NTU; TSS = 133 mg/L). The dashed red line is the 1:1 line. The bottom 6 plots show the absolute percent error between predicted and in situ values in the model training dataset by intervals of predicted values for the training dataset. The red dashed line is the threshold identified as above which predictions may be less reliable.

Figure 9. Comparison of the predictions from the Holdout-data and Full-data models for the same 75% of the data split into spatially representative test and train samples using Generalized Random Tessellation Stratified (GRTS) with the associated r-squared value for each water quality variable. The 1:1 line is represented by a solid black line.

658
659
660
Figure 10. Analysis demonstrating pixel QAQC decisions and relationship to predictions. Predicted
versus in situ Secchi depth (m) for a single lake in the dataset (Lake Okeechobee, FL) matchups in our
training dataset (N = 2,989). (a) Predicted versus in situ Secchi depth (m) for all matchups color-coded by
the percent of the lake's pixels retrieved for that overpass out of the maximum ever retrieved across the
1984-2021 time series. (b) Predicted versus in situ Secchi depth coded by whether any satellite band for
the overpass had a negative value for any lake pixel. (c) Predicted versus in situ Secchi depth coded by
whether any satellite band for the overpass had a median reflectance value that was negative (note, this
flag is part of the final prediction QAQC and is one of three components of our combined QAQC
recommendation for good quality predictions). (d) Plot of predicted versus in situ Secchi depth after
removing predictions with < 25% of a lake's pixels retrieved or with negative median band reflectance.
The 1:1 line is in red. R-squared values are presented prior to application of flags (a) and after removing
predictions with < 25% of a lake's pixels retrieved or with negative median band reflectance (d).
673

679
Figure 11. Predicted versus in situ CHL for Lake McCarrons, MN, which received alum treatment in
October 2004. (a) Predicted versus in situ CHL across all matchup windows, color coded by the number
of calendar days between the satellite overpass and the in situ measurement. The 1:1 line is in red. (b) The
time series for the full matchup dataset for both in situ (green) and predicted (orange) CHL data. (c) The
time series for the ±1 matchup dataset for both in situ (green) and predicted (orange) CHL. (d) Locally
estimated scatterplot smoothing (LOESS) for the in situ and predicted CHL for the ±1 matchup dataset.
The date of the alum treatment is shown as a vertical blue line for (c) and (d).

688
689 **Figure 12.** Plots showing DOC data for Beaver Lake, AR. (a) Predicted versus in situ DOC. (b) Predicted
690 and in situ DOC through time.
691

693
694
695 **Figure 13.** Map showing the total number of predictions for all 6 water quality variables by lake from
696 1984-2020. We include the prediction sets that have passed our QC filter (i.e., those without negative
697 median reflectance for any band and that contain a minimum of 10% of the lake's maximum pixels). The
698 black borders show NEON region delineations.

704
705
706 **Figure 14.** Plots showing the number of predictions in the final dataset, which applies to both predictive
707 models individually. The left panel shows thousands of predictions per day by day of the year within each
708 month. The panel on the right shows thousands of predictions by year. All data have had the minimum
709 quality flags applied.

710
711

712
713 **Figure 15.** Plots of quantiles for each of the six water quality variables from the matchup dataset (left
714 plots) and from the Full-data model prediction dataset (right plots). Black lines are median values.
715

719 **Figure 16.** Correlation matrix of predicted values from the Full-data model for the six water quality
720 variables. Data points were randomly selected single summer (June - September) predicted values for
721 each year from 1984-2020 for the 20,191 lakes with 37 years of data. Note, all WQ variables were from
722 the same scene for a lake year. Spearman rank correlations were based on $\log_{10}$ transformed data. All p
723 values << 0.00001. Color is true color.
724

726
Figure 17. Density plots of predicted values from the Full-data model for true color plotted against three
of the water quality variables related to nutrient and color status. Both axes are plotted on a $\log_{10}$ scale.
Data points were randomly selected single summer (June - September) predictions for each year from
1984-2020 for the 20,191 lakes with 37 years of data. Note, all WQ variables were from the same scene
for a lake year. Columns represent all lakes and lakes were placed into two trophic classifications based
on CHL with values $\leq$ 7 ug/L classified as oligo-mesotrophic and values > 7 ug/L as eutrophic-
hypereutrophic.
734
735
736

747 **Figure 18.** Density plots of Secchi depth against all other water quality variables from the Full-data
748 model predictions. Both axes are plotted on a $\log_{10}$ scale. Data points were randomly selected single
749 summer (June - September) predictions for each year from 1984-2020 for the 20,191 lakes with 37 years
750 of data. Note, all WQ variables were from the same scene for a lake year.
751
752
753
754

755
Figure 19. Density plot of the number of lakes for a given number of mean predictions per year and the
total number of years of data. These values represent the maximal data amount for lakes in the prediction
datasets prior to the application of QAQC flags. The Full-data and Holdout-data models have the identical
number of predictions presented here.

Figure Legends

Figure 1. Overview of the four major steps used to create LAGOS-US LANDSAT. See text for full
description of all steps. The steps are described for an individual water quality variable (e.g., CHL).
Because the same dataset in step 1 is used for all models, steps 2-4 are repeated for each water quality
variable.

Figure 2. Maps of the location of lakes in the LAGOS-US LANDSAT matchup dataset by water quality
variable. For display purposes, we show the matchup data that are within ±7 calendar days of the
LAGOS-US LIMNO in situ observations for each of the six water quality variables which represents the
maximum possible matchup dataset for each variable and the maximum sample sizes and what was tested
in our modeling workflow. See results in Table 2 for the final matchup windows used for each variable as
well as the final sample sizes used. The map outlines are NEON region borders.

Figure 3. Histogram of the number of matchups by year for the matchup dataset. The matchup window
includes all dates within ±7 calendar days of the LAGOS-US LIMNO in situ observations for all water
quality variables as a sum across all variables within each year.

Figure 4. Results from the three model approaches tested for the CHL model. The left panel shows the
reverse cumulative distribution of the absolute value of residuals of the three models using log10-
transformed in situ CHL data and using matchups within ±1 day. See text for model details. 'Random
forest' is a random forest regression model that includes all bands and band ratios; 'Linear model' is a

linear multiple regression model with all bands and band ratios; and 'KIVU' is a 3-band algorithm
calculated at the pixel level for each lake. The right panel shows boxplots of these same absolute values
of residuals with the red dot indicating the RMSE: random forest = 0.23, linear model = 0.51, 'KIVU' =
0.54.
788
Figure 5. Results from testing different components of the workflow related to image and pixel
processing for a subset of the DOC dataset. (a) A plot of variance explained for each training model run
conducted on $\log_{10}$-transformed in situ data that included the indicated components. Results are plotted
for the 5 different ways that reflectance is summarized by lake (see text). 'Whole-lake Ratios' are ratios
calculated on median band data, whereas 'Pixel Ratios' are those ratios first calculated at the pixel-level
and then the median value is calculated; 'All Bands' includes all bands; 'Overcorrection Filter' is the flag
for when there is negative median reflectance. (b) A plot of % OOB variance explained by matchup
window and the 5 different ways that reflectance is summarized by lake. Models are based on a 500-lake
subset that contained DOC matchups, with some lakes being sampled more than once (N = 1,348).
798
799   **Figure 6.** The full-data model variable importance (mean decrease Gini) for all predictors in the random
800   forest regression models for each water quality variable. Orange bars are individual bands, and purple
801   bars are ratios between two bands.
802
Figure 7. Fit statistic plots for the **Holdout-data model** of each of the six water quality variables in
original units. The top six plots show predicted versus in situ quantile lines for the median quantile
(black) and the 5th, 25th, 75th, and 95th quantiles (shaded colored lines). The maximum range of the X
and Y axes is trimmed to the 90th percentile of in situ data (CHL = 53 ug/L; Secchi depth = 5.3 m; true
color = 63 PCU; DOC = 13 mg/L; Turbidity = 19 NTU; TSS = 133 mg/L). The dashed red line is the 1:1
line. The internal random forest OOB variance explained from the model training data as well as the r-
squared value of the holdout data are shown. The bottom 6 plots show the absolute percent error between
predicted and in situ values in the model training dataset by intervals of predicted values for the training
dataset. The red dashed line is the threshold identified as above which predictions are thought to be less
reliable. Outliers are not plotted.
813
Figure 8. Fit statistic plots for the **Full-data predictive model** of each of the six water quality variables
in original units. The top six plots show predicted versus in situ quantile lines for the median quantile
(black) and the 5th, 25th, 75th, and 95th quantiles (shaded colored lines). The maximum range of the X
and Y axes is trimmed to the 90th percentile of in situ data (CHL = 53 ug/L; Secchi depth = 5.3 m; true
color = 63 PCU; DOC = 13 mg/L; Turbidity = 19 NTU; TSS = 133 mg/L). The dashed red line is the 1:1
line. The bottom 6 plots show the absolute percent error between predicted and in situ values in the model
training dataset by intervals of predicted values for the training dataset. The red dashed line is the
threshold identified as above which predictions may be less reliable.
822
823   **Figure 9.** Comparison of the predictions from the Holdout-data and Full-data models for the same 75%
824   of the data split into spatially representative test and train samples using Generalized Random
825   Tessellation Stratified (GRTS) with the associated r-squared value for each water quality
826   variable. The 1:1 line is represented by a solid black line.
827
Figure 10. Analysis demonstrating pixel QAQC decisions and relationship to predictions. Predicted
versus in situ Secchi depth (m) for a single lake in the dataset (Lake Okeechobee, FL) matchups in our
training dataset (N = 2,989). (a) Predicted versus in situ Secchi depth (m) for all matchups color-coded by
the percent of the lake's pixels retrieved for that overpass out of the maximum ever retrieved across the
1984-2021 time series. (b) Predicted versus in situ Secchi depth coded by whether any satellite band for
the overpass had a negative value for any lake pixel. (c) Predicted versus in situ Secchi depth coded by

whether any satellite band for the overpass had a median reflectance value that was negative (note, this
flag is part of the final prediction QAQC and is one of three components of our combined QAQC
recommendation for good quality predictions). (d) Plot of predicted versus in situ Secchi depth after
removing predictions with < 25% of a lake's pixels retrieved or with negative median band reflectance.
The 1:1 line is in red. R-squared values are presented prior to application of flags (a) and after removing
predictions with < 25% of a lake's pixels retrieved or with negative median band reflectance (d).
840

Figure 11. Predicted versus in situ CHL for Lake McCarrons, MN, which received alum treatment in
October 2004. (a) Predicted versus in situ CHL across all matchup windows, color coded by the number
of calendar days between the satellite overpass and the in situ measurement. The 1:1 line is in red. (b) The
time series for the full matchup dataset for both in situ (green) and predicted (orange) CHL data. (c) The
time series for the ±1 matchup dataset for both in situ (green) and predicted (orange) CHL. (d) Locally
estimated scatterplot smoothing (LOESS) for the in situ and predicted CHL for the ±1 matchup dataset.
The date of the alum treatment is shown as a vertical blue line for (c) and (d).
848

849   **Figure 12.** Plots showing DOC data for Beaver Lake, AR. (a) Predicted versus in situ DOC. (b) Predicted
850   and in situ DOC through time.
851

852   **Figure 13.** Map showing the total number of predictions for all 6 water quality variables by lake from
853   1984-2020. We include the prediction sets that have passed our QC filter (i.e., those without negative
854   median reflectance for any band and that contain a minimum of 10% of the lake's maximum pixels). The
855   black borders show NEON region delineations.
856

857   **Figure 14.** Plots showing the number of predictions in the final dataset, which applies to both predictive
858   models individually. The left panel shows thousands of predictions per day by day of the year within each
859   month. The panel on the right shows thousands of predictions by year. All data have had the minimum
860   quality flags applied.
861

862   **Figure 15.** Plots of quantiles for each of the six water quality variables from the matchup dataset (left
863   plots) and from the Full-data model prediction dataset (right plots). Black lines are median values.
864

865   **Figure 16.** Correlation matrix of predicted values from the Full-data model for the six water quality
866   variables. Data points were randomly selected single summer (June - September) predicted values for
867   each year from 1984-2020 for the 20,191 lakes with 37 years of data. Note, all WQ variables were from
868   the same scene for a lake year. Spearman rank correlations were based on log_{10} transformed data. All p
869   values << 0.00001. Color is true color.
870

Figure 17. Density plots of predicted values from the Full-data model for true color plotted against three
of the water quality variables related to nutrient and color status. Both axes are plotted on a log_{10} scale.
Data points were randomly selected single summer (June - September) predictions for each year from
1984-2020 for the 20,191 lakes with 37 years of data. Note, all WQ variables were from the same scene
for a lake year. Columns represent all lakes and lakes were placed into two trophic classifications based
on CHL with values $\leq$ 7 ug/L classified as oligo-mesotrophic and values > 7 ug/L as eutrophic-
hypereutrophic.
878

879   **Figure 18.** Density plots of Secchi depth against all other water quality variables from the Full-data
880   model predictions. Both axes are plotted on a log_{10} scale. Data points were randomly selected single
881   summer (June - September) predictions for each year from 1984-2020 for the 20,191 lakes with 37 years
882   of data. Note, all WQ variables were from the same scene for a lake year.
883

884   **Figure 19.** Density plot of the number of lakes for a given number of mean predictions per year and the
885   total number of years of data. These values represent the maximal data amount for lakes in the prediction
886   datasets prior to the application of QAQC flags. The Full-data and Holdout-data models have the identical
887   number of predictions presented here.

Tables

Table 1. Synthesis of published studies that use Landsat for macroscale studies of lake water quality.

Characteristics of macroscale lake studies using Landsat							OA data product includes					
Country or spatial extent	Name of resource	Temporal extent (maximum possible)	Min lake size	Number of lakes	Aquatic Variables of interest	Study type	Reflectances	Match-ups	Ground-truthed predictions	Calculated values from reflectances	Methods documentation	Reference
Global	NA	1982-2018	> 100 ha	176,032	Algal blooms (FAI)	*Analysis*: Calculations from reflectances				✓	No (Only methods section; no code)	Fang et al. 2022
Global	**Global Bloom Database**	1982-2019	> 10 ha	248,243	Algal blooms (FAI)	*Analysis*: Calculations from reflectances				✓	No (Only methods section; no code)	Hou et al. 2022
Global	NA	2019	> 1 ha	NA	Secchi	*Model development*					No (Only methods section; no code)	Song et al. 2022
Global	NA	1984-2022	NA	> 300	Secchi	*Model development*	✓		✓		Partial (Docum. & code available, but not reproducible)	Maciel et al. 2023
China	**Annual inland water clarity dataset of China**	1984-2018 (5-yr means)	> 1 ha	10,814	Secchi	*Dataset*: Ground-truthed predictions			✓		Partial	Tao et al. 2022
China	**CNLTSI**	1984-2023	> 100 ha	2,693	Trophic State Index (TSI)	*Dataset:*			✓		Partial (Docum. & code available, but not reproducible)	Hu et al. 2024
Continental US	**AquaSat**	1984-2019	NA	Sites, not lakes	TSS, DOC, CDOM, CHL, Secchi	*Dataset*: Reflectances (6 bands); Match-ups	✓	✓			✓	Ross et al. 2019
Continental US	**LimnoSat-US**	1984-2020	> 10 ha	56,792	NA	*Dataset*: Reflectances (6 bands)	✓				✓	Topp et al. 2020
Continental US	NA	1984-2018; 5-yr means	> 10 ha	14,000	Secchi	*Analysis*: Ground-truthed predictions	✓		✓		✓	Topp et al. 2021
Continental US	NA	1984-2020	> 10 ha	26,000	Spectral color	*Analysis*: Calculations	✓			✓	✓	Topp et al. 2021
Continental US	**LTS-US**	1984-2020	> 10 ha	55,662	Trophic state	*Dataset*: Modeled predictions			✓		✓	Meyer et al. 2023
Continental US	**LAGOS-US LANDSAT**	1984-2020	> 4 ha	106,000	Secchi, CHL, DOC, True color, TSS, Turbidity	*Dataset*: Reflectances (6 bands); matchups; ground-truthed predictions	✓	✓	✓		✓	*This study*

893 **Table 2.** LAGOS-US LANDSAT workflow decisions and fit statistics for the Full-data model for each water quality predictive model.

| | Workflow components tested and selected | | | | | | | | | | | | Final model fit statistics |
| | Image and pixel processing by lake | | | | | Modeling steps | | Matchup training data | | | | | |
Variable	Pixel retrievals	Band ratio calculation	Scene quality: *Negative reflectances*	Scene quality: *Scene clouds > 50%*	In situ data transformed	# of bands and ratios	Training model type	Matchup window *(± days)*	Final matchups *(N)*	Variance explained *(%)*	RMSE	MAPE (%)	MAPE (%) *Improvement of log$_{10}$ model*
CHL	Whole-lake	Pixel-specific	Removed	Removed	Log$_{10}$	All bands & ratios	Random forest	1	43,755	44.1	0.43	23.3	(+9.8)
Secchi	Whole-lake	Pixel-specific	Removed	Removed	none	All bands & ratios	Random forest	7	586,368	63.7	1.23	22.8	--
True color	Whole-lake	Pixel-specific	Removed	Removed	Log$_{10}$	All bands & ratios	Random forest	7	33,194	48.6	0.26	7.5	(+12.7)
DOC	Whole-lake	Pixel-specific	Retained	Retained	Log$_{10}$	All bands & ratios	Random forest	7	7,466	42.0	0.24	10.2	(+6.1)
TSS	Whole-lake	Pixel-specific	Removed	Removed	Log$_{10}$	All bands & ratios	Random forest	1	10,845	20.7	0.62	27.0	(+16.3)
Turbidity	Whole-lake	Pixel-specific	Removed	Removed	Log$_{10}$	All bands & ratios	Random forest	1	58,999	52.1	0.37	8.8	(+20.9)

Table 3. Statistical summary table of predictions for the six water quality variables, along with summaries of predictions per lake, and per year. Lake area was from LAGOS-US LOCUS[23].

Variable	Units	min	q5	q25	q50	q75	q95	max
Water quality variables (37,251,456 predictions)								
CHL	μg/L	0.04	3.1	5.7	8.7	14.0	29.1	299.9
True color	PCU	0.1	8.7	16.4	23.9	32.4	44.7	88.1
DOC	mg/L	0.5	2.8	4.3	5.7	7.1	9.7	67.7
Turbidity	NTU	0.05	1.2	2.4	4.4	9.3	21.2	197.8
Secchi depth	meters	0.02	0.7	1.2	1.9	2.7	4.3	59.3
TSS	mg/L	0.5	4.8	9.1	14.2	24.3	51.6	675.9
Lake-related (136,877 lakes)								
Predictions per lake	N	1	32	160	255	361	568	1377
Lake–years with predictions*	N	1	13	32	37	37	37	37
Lake water area	ha	4	4	6	9	19	116	338,950
Year-related (37 years)								
Number of lakes per year	N	108,158	110,454	117,689	121,821	123,719	125,928	126,177
Number of predictions per year	N	534,405	604,508	938,970	1,021,304	1,066,120	1,317,022	1,455,641

*The percentages of lakes with 37 and 30+ years with at least 1 summer observation (June-September) are, respectively, 14.8% and 51.6%

References

907   1.   Topp, S. N., Pavelsky, T. M., Jensen, D., Simard, M. & Ross, M. R. V. Research Trends in the Use of Remote Sensing for Inland Water Quality Science: Moving Towards Multidisciplinary Applications. *Water* **12**, 169 (2020).

2.   Adjovu, G. E., Stephen, H., James, D. & Ahmad, S. Overview of the Application of Remote Sensing in Effective Monitoring of Water Quality Parameters. *Remote Sens.* **15**, 1938 (2023).

3.   Kloiber, S. M., Brezonik, P. L. & Bauer, M. E. Application of Landsat imagery to regional-scale assessments of lake clarity. *Water Res.* **36**, 4330–4340 (2002).

4.   Olmanson, L. G. *et al.* Regional measurements and spatial/temporal analysis of CDOM in 10,000+ optically variable Minnesota lakes using Landsat 8 imagery. *Sci. Total Environ.* **724**, 138141 (2020).

5.   Hicks, B. J., Stichbury, G. A., Brabyn, L. K., Allan, M. G. & Ashraf, S. Hindcasting water clarity from Landsat satellite images of unmonitored shallow lakes in the Waikato region, New Zealand. *Environ. Monit. Assess.* **185**, 7245–7261 (2013).

6.   Roy, D. P. *et al.* Characterization of Landsat-7 to Landsat-8 reflective wavelength and normalized difference vegetation index continuity. *Remote Sens. Environ.* **185**, 57–70 (2016).

7.   Ross, M. R. V. *et al.* AquaSat: A Data Set to Enable Remote Sensing of Water Quality for Inland Waters. *Water Resour. Res.* **55**, 10012–10025 (2019).

8.   Yang, H. *et al.* A Review of Remote Sensing for Water Quality Retrieval: Progress and Challenges. *Remote Sens.* **14**, 1770 (2022).

9.   Pollard, A. I., Hampton, S. E. & Leech, D. M. The Promise and Potential of Continental-Scale Limnology Using the U.S. Environmental Protection Agency's National Lakes Assessment. *Limnol. Oceanogr. Bull.* **27**, 36–41 (2018).

10.   Shuvo, A. K. *et al.* LAGOS-US LIMNO: Data module of surface water chemistry from 1975-2021 for lakes in the conterminous U.S. https://doi.org/10.6073/PASTA/2C58F5A50AB813919F99CC1F265F271C (2023).

11.   Topp, S. N. *et al.* Multi-decadal improvement in US Lake water clarity. *Environ. Res. Lett.* **16**, 055025 (2021).

12.   Topp, S. N. *et al.* Shifting Patterns of Summer Lake Color Phenology in Over 26,000 US Lakes. *Water Resour. Res.* **57**, e2020WR029123 (2021).

13.   Meyer, M. F. *et al.* National-scale remotely sensed lake trophic state from 1984 through 2020. *Sci. Data* **11**, 77 (2024).

14.   Tao, H. *et al.* A Landsat-derived annual inland water clarity dataset of China between 1984 and 2018. *Earth Syst. Sci. Data* **14**, 79–94 (2022).

15.   Song, K. *et al.* Climatic versus Anthropogenic Controls of Decadal Trends (1983–2017) in Algal Blooms in Lakes and Reservoirs across China. *Environ. Sci. Technol.* **55**, 2929–2938 (2021).

16.   Fang, C. *et al.* Global divergent trends of algal blooms detected by satellite during 1982–2018. *Glob. Change Biol.* **28**, 2327–2340 (2022).

17.   Hou, X. *et al.* Global mapping reveals increase in lacustrine algal blooms over the past decade. *Nat. Geosci.* **15**, 130–134 (2022).

18.   Maciel, D. A. *et al.* Towards global long-term water transparency products from the Landsat archive. *Remote Sens. Environ.* **299**, 113889 (2023).

19.   Hanly, P. J., Webster, K. E. & Soranno, P. A. LAGOS-US LANDSAT: Data module of remotely-sensed water quality estimates for U.S. lakes over 4 ha from 1984 to 2020. *Environ. Data Initiat.* https://doi.org/10.6073/pasta/128700feb3bbc3ffe5800e7b232bd81f (2024).

20. Wilkinson, M. D. *et al.* The FAIR Guiding Principles for scientific data management and
stewardship. *Sci. Data* **3**, 160018 (2016).
21. Cheruvelil, K. S. *et al.* LAGOS-US LOCUS v1.0: Data module of location, identifiers, and physical
characteristics of lakes and their watersheds in the conterminous U.S. *Limnol. Oceanogr. Lett.* **6**,
270–292 (2021).
22. Smith, N. J., Webster, K. E., Rodriguez, L. K., Cheruvelil, K. S. & Soranno, P. A. LAGOS-US GEO
v1.0: Data module of lake geospatial ecological context at multiple spatial and temporal scales in the
conterminous U.S. https://doi.org/10.6073/PASTA/0E443BD43D7E24C2B6ABC7AF54CA424A
(2022).
23. Smith, N. J., Webster, K. E., Rodriguez, L. K., Cheruvelil, K. S. & Soranno, P. A. LAGOS-US
LOCUS v1.0: Data module of location, identifiers, and physical characteristics of lakes and their
watersheds in the conterminous U.S.
https://doi.org/10.6073/PASTA/E5C2FB8D77467D3F03DE4667AC2173CA (2021).
24. Masek, J. G. *et al.* A Landsat Surface Reflectance Dataset for North America, 1990–2000. *IEEE*
Geosci. Remote Sens. Lett. **3**, 68–72 (2006).
25. Vermote, E., Justice, C., Claverie, M. & Franch, B. Preliminary analysis of the performance of the
Landsat 8/OLI land surface reflectance product. *Remote Sens. Environ.* **185**, 46–56 (2016).
26. Foga, S. *et al.* Cloud detection algorithm comparison and validation for operational Landsat data
products. *Remote Sens. Environ.* **194**, 379–390 (2017).
27. Boucher, J., Weathers, K. C., Norouzi, H. & Steele, B. Assessing the effectiveness of Landsat 8
chlorophyll *a* retrieval algorithms for regional freshwater monitoring. *Ecol. Appl.* **28**, 1044–1054
(2018).
28. Webster, K. E. *et al.* An empirical evaluation of the nutrient-color paradigm for lakes. *Limnol.*
Oceanogr. **53**, 1137–1148 (2008).
29. Brivio, P. A., Giardino, C. & Zilioli, E. Determination of chlorophyll concentration changes in Lake
Garda using an image-based radiative transfer code for Landsat TM images. *Int. J. Remote Sens.* **22**,
487–502 (2001).
30. Liaw, A. & Wiener, M. Classification and Regression by randomForest. *R News* **2**, 18–22.
31. Dumelle, M., Kincaid, T., Olsen, A. R., & Weber, M. spsurvey: spatial sampling design and analysis
in R. *Journal of Statistical Software*. **105**, 1-29 (2023).
32. Breiman, L. Random forests. Machine learning. **45**, 5-32 (2001).
33. Liaw, A., & Wiener, M. Classification and regression by randomForest. R news. **2**, 18-22 (2002).
34. Rokach, L. Decision forest: Twenty years of research. Information fusion, **27**, 111-125 (2016).
35. Carpenter, D. J. & Carpenter, S. M. Modeling inland water quality using Landsat data. *Remote Sens.*
Environ. **13**, 345–352 (1983).
36. Lillesand, T. M. Use of Landsat Data to Predict the Trophic State of Minnesota Lakes. *Photogramm.*
Eng. (1983).
37. Brezonik, P., Menken, K. D. & Bauer, M. Landsat-based Remote Sensing of Lake Water Quality
Characteristics, Including Chlorophyll and Colored Dissolved Organic Matter (CDOM). *Lake*
Reserv. Manag. **21**, 373–382 (2005).
38. Olmanson, L. G., Bauer, M. E. & Brezonik, P. L. A 20-year Landsat water clarity census of
Minnesota's 10,000 lakes. *Remote Sens. Environ.* **112**, 4086–4097 (2008).
39. Paltsev, A. & Creed, I. F. Multi-decadal changes in phytoplankton biomass in northern temperate
lakes as seen through the prism of landscape properties. *Glob. Change Biol.* **28**, 2272–2285 (2022).

40. Mamun, M., & Yang, X. Advancing lakes algal chlorophyll estimation in the contiguous USA: A
comparative study of machine learning models and satellite data. *Ecological Informatics*, **87**, 103087
(2025).

41. Olmanson, L. G., Brezonik, P. L. & Bauer, M. E. Remote sensing for regional lake water quality
assessment: capabilities and limitations of current and upcoming satellite systems. *Adv. Watershed*
Sci. Assess. 111–140 (2015).

42. Markogianni, V., Kalivas, D., Petropoulos, G. & Dimitriou, E. An Appraisal of the Potential of
Landsat 8 in Estimating Chlorophyll-a, Ammonium Concentrations and Other Water Quality
Indicators. *Remote Sens.* **10**, 1018 (2018).

43. Toming, K. *et al.* Predicting lake dissolved organic carbon at a global scale. *Sci. Rep.* **10**, 8471
(2020).

44. James, W. F. & Bischoff, J. M. Sediment aluminum:phosphorus binding ratios and internal
phosphorus loading characteristics 12 years after aluminum sulfate application to Lake McCarrons,
Minnesota. *Lake Reserv. Manag.* **36**, 1–13 (2020).

45. Toming, K. *et al.* Predicting lake dissolved organic carbon at a global scale. *Sci. Rep.* **10**, 8471
(2020).

46. Diaz Vazquez, J., McCullough, I. M., Haite, M., Soranno, P. A. & Cheruvelil, K. S. US lakes are
monitored disproportionately less in communities of color. *Front. Ecol. Environ.* (In Press).

47. Wulder, M. A. *et al.* The global Landsat archive: Status, consolidation, and direction. *Remote Sens.*
Environ. **185**, 271–283 (2016).

48. Wagner, T. *et al.* Quantifying sample biases of inland lake sampling programs in relation to lake
surface area and land use/cover. *Environ. Monit. Assess.* **141**, 131–147 (2008).

49. Stanley, E. H. *et al.* Biases in lake water quality sampling and implications for macroscale research.
Limnol. Oceanogr. **64**, 1572–1585 (2019).

50. Leech, D. M., Pollard, A. I., Labou, S. G. & Hampton, S. E. Fewer blue lakes and more murky lakes
across the continental U.S.: Implications for planktonic food webs. *Limnol. Oceanogr.* **63**, 2661–
2680 (2018).

51. Wagner, T. & Schliep, E. M. Combining nutrient, productivity, and landscape-based regressions
improves predictions of lake nutrients and provides insight into nutrient coupling at macroscales.
Limnol. Oceanogr. **63**, 2372–2383 (2018).

52. Wagner, T. *et al.* Increasing accuracy of lake nutrient predictions in thousands of lakes by leveraging
water clarity data. *Limnol. Oceanogr. Lett.* **5**, 228–235 (2020).

53. Soranno, P. A., Hanly, P. J., Webster, K. E., Wagner, T., McDonald, A., Shuvo, A., ... & Spence
Cheruvelil, K. Abrupt changes in algal biomass of thousands of US lakes are related to climate and
are more likely in low-disturbance watersheds. *Proceedings of the National Academy of Sciences*,
122(9), e2416172122 (2025).

54. Hanly, P. J. Code for generating Landsat reflectance time series, water chemistry matchups, and
machine learning models to predict water chemistry for US lakes.
Zenodo. https://doi.org/10.5281/zenodo.14164583 (2024)

1033
1034
1035
1036