Regulatory T cell-derived enkephalin gates nociception

Élora Midavaine[1], Beatriz C. Moraes[1], Jorge Benitez[1], Sian R. Rodriguez[1], Joao M. Braz[1], Nathan P. Kochhar[1], Walter L. Eckalbar[1], Ana I. Domingos[2], John E. Pintar[3], Allan I. Basbaum[1*], Sakeen W. Kashem[4,5*]

Affiliations:
[1]Department of Anatomy, University of California San Francisco, San Francisco, California, USA.
[2]Department of Physiology, Anatomy and Genetics, University of Oxford, Oxford, UK.
[3]Department of Neuroscience and Cell Biology, Rutgers Robert Wood Johnson Medical School, Piscataway, NJ, USA.
[4]Department of Dermatology, University of California San Francisco, San Francisco, CA, USA.
[5]Dermatology, Veterans Affairs Medical Center, San Francisco, California, USA.
*These authors contributed equally

Correspondence:
Allan I. Basbaum,
Chair and Professor, Department of Anatomy
University of California San Francisco
1550 Rock Hall, Room 345A, San Francisco, California, USA 94158
Email: allan.basbaum@ucsf.edu

Lead correspondence:
Sakeen W. Kashem
Assistant Professor, Dermatology
University of California San Francisco
1701 Divisadero, 3rd Floor, San Francisco, CA 94115
Email: sakeen.kashem@ucsf.edu

Keywords:
Pain, regulatory T cells, neuroimmunology, sex dimorphism, opioid, sensory neurons, enkephalin, meninges

Summary

T cells have emerged as sex-dependent orchestrators of pain chronification but the sexually dimorphic mechanisms by which T cells control pain sensitivity is not resolved. Here, we demonstrate an influence of regulatory T cells (Tregs) on pain processing that is distinct from their canonical functions of immune regulation and tissue repair. Specifically, meningeal Tregs (mTregs) express the endogenous opioid, enkephalin, and mTreg-derived enkephalin exerts an antinociceptive action through a presynaptic opioid receptor signaling mechanism that is dispensable for immunosuppression. We demonstrate that mTregs are both necessary and sufficient to suppress mechanical pain sensitivity in female, but not male, mice, with this modulation reliant on sex hormones. These results uncover a fundamental sex-specific, and immunologically-derived endogenous opioid circuit for nociceptive regulation with critical implications for pain biology.

Highlights

1. Gating of allodynia by meningeal Tregs is sex hormone-dependent

3. Treg-derived enkephalin modulates mechanical pain sensitivity, not inflammation

4. Delta opioid receptor on MrgprD$^+$ sensory neuron mediates pain processing by mTregs

Introduction

Pain prevalence is significantly higher in women across multiple pain conditions including nerve injury-induced neuropathic pain, musculoskeletal pain, fibromyalgia and migraine[1]. Gender disparities in pain are further evidenced by notable changes in chronic pain severity during hormonal gender affirming care[2]. Here, we identified a previously unknown immunological mechanism that underlies sex differences in both acute and chronic pain regulation.

Regulatory T cells (Tregs) are a subset of CD4$^+$ T cells characterized by their immunosuppressive function and the expression of the X-linked master regulatory transcriptional factor *Foxp3*. In addition to their critical function in limiting inflammation, Tregs are also major contributors to wound healing, they regulate stem cell turnover, maintain metabolic homeostasis, facilitate placental implantation and promote maternal-fetal tolerance[8–16]. In the context of nervous system injury, Tregs mitigate pro-inflammatory cytokine interferon-□ (IFN-□)-driven mechanical pain hypersensitivity, suppress microglia-driven nociceptive processing and can reduce astrogliosis. Additionally, they contribute to improving remyelination, thereby promoting tissue repair[17–20]. However, whether and how Tregs can directly alter neuronal activity is still unknown.

Here, we demonstrate that meningeal regulatory T cells (mTregs) are essential contributors to baseline mechanical sensitivity, and to the inhibition of mechanical pain hypersensitivity (allodynia) after nerve injury, but only in female mice. Using a well-established spared-nerve injury (SNI) model of neuropathic pain, we further show that expanding mTregs can reduce nociceptive processing independently of Treg tissue repair programs. We find that mTregs produce the endogenous opioid met-enkephalin, with female mice exhibiting increased numbers of enkephalinergic mTregs. mTreg-derived enkephalin is required for the suppression of mechanical pain hypersensitivity, but not inflammation. This distinction reveals a novel Treg function that differs from their well-established roles in immunological restraint and tissue repair.

The involvement of distinct opioid receptors in the peripheral (PNS) and central nervous systems (CNS) in the regulation of different pain modalities is well documented. Delta (δOR) and mu (μOR) opioid receptors are differentially expressed among neuronal subsets and regulate mechanical or thermal pain processing, respectively[21–23]. Enkephalin can bind to both δOR and μOR, but preferentially engages the δOR. Here we demonstrate that the anti-allodynic effects derived from mTregs are mediated by the δOR that is expressed on the *Mas*-related G protein-coupled receptor member D (MrgprD) subset of unmyelinated primary sensory neurons.

Given the pronounced sex differences observed in Treg-mediated suppression of nociceptive processing, we further explored the determinants of this sex selectivity. Our findings indicate that gonadal hormones, rather than sex chromosomes, play a key role in modulating Treg suppression of nociceptive thresholds. Thus, we propose a novel mechanism by which Tregs mediate the suppression of nociception, regulated by sex-specific hormonal influences.

Results

Sex-specific suppression of nociceptive thresholds by meningeal Tregs

As the immune system has emerged as a central determinant driving sex differences in pain sensitivity, we sought to investigate its contribution to nociceptive processing[24]. Tregs are a fundamental cell type essential for maintaining and restoring tissue homeostasis. Here we initially focused on identifying Treg localization and function within nervous system tissues. Using confocal microscopy and flow cytometry on whole meningeal sheets, dorsal root ganglia (DRG), sciatic nerves, spinal cords (SC) and brain, we localized Tregs to the meninges of the CNS, and to the leptomeninges of the DRG **(Figure 1A-C)**. To enhance the sensitivity of Treg detection within these organs, we utilized the bright reporter signals from the double reporter $Foxp3^{eGFP-Cre-ERT2}$;$Rosa26^{LSL-tdTomato}$ mice. Notably, and consistent with previous reports we observed a more pronounced localization of Tregs in the lumbar and caudal segments of the spinal cord meninges **(Figure 1A)** [25]. In the DRG, as for most leukocytes other than resident macrophages, Tregs predominated in the leptomeninges, proximal to the dorsal roots entry zone, with sparse presence within the DRG parenchyma **(Figure 1B)**[26]. By intravenous administration of a fluorescently-labeled CD45 antibody and using flow cytometry, we distinguished intravascular Tregs from tissue Tregs **(Figure S1A)**[27]. We quantified the numbers of non-vascular, tissue Tregs in various organs within the nervous and lymphoid systems. We will further refer to the SC meningeal and DRG leptomeningeal Tregs as mTregs **(Figure 1C)**. We observed minimal localization of Tregs in peripheral nerves and did not detect any Tregs within the parenchyma of the CNS in young, uninjured mice. We observed nearly equivalent numbers of tissue Tregs between male and female mice across tissue **(Figure 1D)**[18].

To assess the feasibility of site-specific depletion of mTregs, we performed intrathecal (IT) injections of pegylated diphtheria toxin (pegDT) in $Foxp3$-DTR mice[28]. Although an IT injection of Evan's blue rapidly spreads through the SC meninges and DRG, into the brain and to the draining cervical and lumbar lymph nodes, pegylated fluorescently labeled molecules (pegDyLight 650) remain restricted to the SC meninges and to the DRG **(Figure S2A-B)**. Consistently, a single 20 ng dose of pegDT IT selectively depleted >90% of SC and DRG mTregs in both male and female mice, but spared Tregs located in the brain meninges, draining lymph nodes, spleen, and peripheral nerves **(Figure 1E-F)**. Importantly, $Foxp3$-DTR mice subjected to repeated IT administrations of pegDT do not exhibit the significant weight loss, splenomegaly and mortality that typically develops following systemic autoimmunity in $Foxp3$-DTR mice induced by repeated intraperitoneal (IP) injections of diphtheria toxin (DT) **(Figure S2C-F)**. Clearly, the pegDT IT system offers a novel method for selective depletion of mTregs while avoiding systemic inflammation.

We next evaluated behavioral outcomes in mice following mTreg depletion. A single dose of pegDT IT induced a profound and prolonged decrease in mechanical thresholds in naïve female but not male $Foxp3$-DTR mice **(Figure 1G-H)**. Importantly, mechanical thresholds in wildtype (WT) C57BL/6 mice treated with pegDT IT or $Foxp3$-DTR mice treated with vehicle IT did not differ, ruling out pegDT or IT injections as the cause of the sex-specific allodynia **(Figure S3A)**. In addition to evaluating mechanical

hypersensitivity, which is conveyed by mechanosensitive unmyelinated and myelinated
primary afferent nerve fibers, we also assessed mice for noxious heat sensitivity
mediated by Trpv1[+] nociceptors, cold sensitivity that is mediated by Trpm8[+] nociceptors,
pin prick sensitivity mediated by Aδ afferents and brush responses mediated by Aβ
fibers. Although depletion of mTregs selectively induced mechanical allodynia in
females, it did not impact any other sensory modality. Motor function tests using the
rotarod also did not differ in either sex **(Figure 1I and Figure S3A-H)**. We conclude that
mTregs selectively suppress mechanical thresholds in a sex-dependent manner,
effectively preventing mechanical allodynia in a previously uninjured state.
102
Expansion of mTregs alleviates injury-induced mechanical allodynia
independently of tissue repair
In addition to exploring mTreg role in mechanical sensitivity in uninjured mice, we
investigated whether mTreg can suppress allodynia following nerve injury. Using a well-
established spared nerve injury (SNI) model of neuropathic pain, we transected and
ligated the common peroneal and tibial nerve branches of the sciatic nerve, sparing the
sural nerve. This model induces chronic, unremitting, and permanent mechanical
hypersensitivity with a non-healing neuroma formation four weeks after the injury
(Figure 2A-B)[29,30]. As mice with SNI exhibit mechanical thresholds at the limit of
detection with commercially available von Frey filaments, we conducted single fiber
testing using the lowest available 0.008 g von Frey filament. Again, mTreg depletion
increased allodynia following SNI in females, but not in males **(Figure 2C-D)**.
115
We next asked whether expanding mTregs could alleviate the mechanical allodynia
independently of tissue repair. Tregs express the high affinity interleukin-2 receptor, IL-
2Rα, and low-doses of IL-2 can effectively expand Tregs in mice, a therapeutic
approach that has been used to treat autoimmune diseases in humans[31]. IT injections of
low-dose IL-2 successfully expanded mTregs in both male and female mice **(Figure**
2F). However, although mTreg expansion promoted significant anti-allodynia in SNI
female mice, it did not exhibit a similar effect in males **(Figure 2G-H)**. It is noteworthy
that acute IT injection of IL-2, in uninjured mice, did not increase nociceptive thresholds,
suggesting that an IL-2 or Treg-based therapy could selectively improve neuropathic
pain without affecting basal nociceptive processing **(Figure S4A-B)**. IL-2 injections and
mTreg expansion in mice with SNI likewise did not alter noxious cold or heat sensitivity,
again highlighting the specificity of the sensory modality modulation by mTreg in the
context of neuropathic pain **(S4C-E)**.
129
Gonadal hormones, not sex chromosomes determine sex-selective, anti-
nociceptive function of mTregs
Foxp3 is an X-linked gene, some of which escape X-inactivation. Moreover, random X-
inactivation has been suggested to be potentially altered during inflammatory state in
females[32–34]. To test whether sex chromosomes dosage contributes to our observed
phenotype, we used the Four Core Genotypes (FCG) mouse model in which gonadal
sex in mice is independent of sex chromosomes[35,36]. FCG mice harbor a deficiency in
the sex determining region Y protein (*Sry*) on the Y chromosome and instead feature an
autosomal transgenic insertion of *Sry.* This genetic configuration enables the

discrimination of sex chromosome dose influence from the contribution of gonadal hormones **(Figure 2I)**. Both XX and XY chromosome gonadal female mice displayed mTreg-mediated alleviation of mechanical allodynia after SNI; XX- and XY- gonadal male mice did not **(Figure 2J-L)**. Similarly, after mTreg depletion in *Foxp3*-DTR mice crossed to the FCG system, we found that female specific gonadal hormones, but not sex chromosome, mediate the mTreg suppression of nociceptive thresholds in the absence of injury **(Figure 2M-N)**. Based on our findings in both uninjured and chronic injury states, we conclude that there is a profound and consistent sex hormone-dependent contribution of mTregs to the modulation mechanical pain sensitivity.

Regulatory T cells express the endogenous opioid peptide enkephalin

To investigate the molecular mechanisms though which Tregs suppress nociceptive thresholds, we first interrogated public genomic resources. We hypothesized that meningeal tissue Tregs could exhibit an activated lymphoid Treg phenotype rather than a resting Treg phenotype. Tregs have increased expression of the *Penk* gene, which encodes for *Proenkephalin*, a peptide precursor of both Met- and Leu-enkephalin, in various tissues including the nervous system, in both mice and humans[19,37–39]. Here we re-analyzed raw public RNA-seq data of activated Tregs, resting Tregs, as well as activated and resting CD4[+] Foxp3[-] conventional T cells (Tconv)[40]. Strikingly, in activated versus resting Tregs, we observed a significant upregulation of *Penk* expression **(Figure 3A)**. We also investigated other opioid ligand and receptor genes but only recorded a very sparse expression of other opioid-related genes among the CD4[+] T cell subsets **(Figure 3B)**. Based on our prior experience defining mechanical sensitivity through enkephalin-δOR signaling[21], we therefore focused on Treg expression of *Penk.* By ATAC-seq analysis, we observed open chromatin regions of the *Penk* locus in activated Tregs, but not in other CD4[+] T cell subsets. This open chromatin was similar to the open chromatin, promoter and enhancer regions of the developing forebrain, an established enkephalin-producing area of the murine CNS **(Figure 3C)**[41]. By analyzing the raw dataset from the Immunological Genome Project[42], we also explored *Penk* expression within cell populations of the immune system. We observed a strikingly greater *Penk* expression in Tregs, compared to other immune cells (**Figure 3D)**. Furthermore, *Penk* expression in Tregs increase significantly following stimulation with IL-2, compared to other common gamma chain cytokines (**Figure 3E).**

To establish whether Tregs indeed produce the endogenous opioid peptide enkephalin, we screened commercially available anti-Met-enkephalin antibodies. Met-enkephalin was chosen over leu-enkephalin as the latter can be cleaved from both proenkephalin and prodynorphin peptides[43]. These antibodies were validated using *Penk*[-/-] mice as negative controls. Figure 3F shows that mTregs produce met-enkephalin, but meningeal CD4[+] T cells and lymphoid Tregs produce very low levels even after cytokine stimulation (**Figure 3F)**. We validated this finding by generating *Penk*[Cre]*;Rosa26*[tdTomato] mice, which fate-labeled enkephalinergic cells. Consistent with our antibody finding, we observed very similar number of enkephalinergic lineage (tdTomato positive) mTregs in naïve mice. Very few lymphoid or intravascular Tregs were tdTomato labeled **(Figure 3G)**. Most interestingly, female mice exhibited significantly greater numbers of enkephalin-positive Tregs in the meninges, but not in the lymphoid organs. This

185 distinction suggests that Treg fate and function variation across the sexes may be organ
186 system specific **(Figure 3H)**.
187
mTreg-derived enkephalin is required for suppressing nociceptive processing
Using $Penk^{Cre};Rosa26^{tdTomato}$ mice, we next investigated enkephalin lineage positive
cells in the meninges and the DRG. Interestingly, we observed a significant increase in
the representation of mTregs in the tdTomato-positive enkephalin subpopulation
compared to tdTomato-negative cells **(Figure 4A-B)**. In order to manipulate the
enkephalin-producing immune cells, we generated bone marrow chimeric mice by
transplanting $Penk^{Cre};Rosa26^{DTR}$ bone marrow into irradiated CD45.2 congenically
marked WT mice **(Figure 4C)**. This strategy enables a selective DT-induced depletion
of hematopoietic enkephalinergic cells, that spares depletion of non-hematopoietic
enkephalinergic cells of the nervous system and the stroma. Importantly, $CD4^+$ T cells
of the meninges are predominantly bone marrow-derived and exhibit a tissue circulatory
characteristic rather than acquiring tissue residency[44]. Consistently, we found that
mTregs are indeed bone marrow-derived, similar to lymphoid Tregs, and differ from
spinal microglia and skin Langerhans cells, which are host-derived **(Figure S5A-C)**.
pegDT IT administrations in $PenkDTR^{\Delta heme}$ chimeric mice decreased the number of
mTregs and led to profound mechanical hypersensitivity in female, but not male mice, in
both uninjured and nerve injured states **(Figure 4D-G)**. We conclude that blood-derived
meningeal enkephalinergic cells gate mechanical hypersensitivity in females.
206
Having established a female-specific contribution of the bone marrow-derived
enkephalin system, we next used female mice to dissect the mechanism of pain
regulation by mTregs. To establish whether bone marrow-derived enkephalin is required
for the regulation of nociceptive thresholds, we generated bone marrow chimeras in
which $Penk$ deficient bone marrow is transplanted into irradiated hosts, thus generating
$Penk^{\Delta heme}$ mice. As predicted, these $Penk^{\Delta heme}$ mice display decreased nociceptive
thresholds during uninjured state, compared to vehicle-injected $Penk^{\Delta heme}$ mice, which
supports our conclusion that hematopoietic cell-derived enkephalin controls basal
mechanical sensitivity but only in females **(Figure 4H)**.
216
We recognize that recombination-based selective ablation of enkephalin on Tregs,
using $Foxp3^{Cre}$ or $Foxp3^{Cre-ERT2}$ has multiple limitations and caveats. These include
systemic targeting of Tregs, including enkephalinergic Tregs in the skin, potential
compensatory $Penk$ regulation upon constitutive ablation, potential side effects of
tamoxifen, potential stochastic deletion of $Penk$ outside of Tregs in homozygote
Cre/Cre-ERT2 mice, and the impact of random X-inactivation on heterozygous mice[37].
In light of these concerns, we also generated mixed bone marrow chimeras using 1:1
ratio of $Foxp3$-DTR and $Penk^{-/-}$ bone marrow and implanted these chimeras in irradiated
WT mice. Intrathecal injection of pegDT into these mice results in ablation of $FoxP3$-
DTR Tregs; the remaining Tregs are left deficient for $Penk$ ($Penk^{\Delta mTreg}$ mice) while
preserving other immune cell types. Importantly, this approach circumnavigated
potential depletion of any previously unrecognized non-hematopoietic cells that express
$Foxp3$. At baseline, uninjected mixed chimeric mice had similar mechanical thresholds
as $WT^{\Delta heme}$ control chimeras. As predicted, pegDT IT injection led to mechanical

hypersensitivity in uninjured $Penk^{\Delta mTreg}$ but not $WT^{\Delta heme}$ control mice and exacerbated
nerve injury-induced hypersensitivity **(Figure 4I-J)**. Having shown that IL-2-induced
mTreg expansion and expression of enkephalin alleviates neuropathic pain, we next
investigated whether this could be mediated by the δOR, the preferred receptor for
enkephalin. In these studies, we co-administered IL-2 and naltrindole, a selective
antagonist of the δOR, and observed that IL-2-induced anti-allodynia was abolished
(Figure 4K-L). We conclude that mTreg-derived enkephalin is required for suppressing
mechanical pain hypersensitivity and that this suppression is mediated by the δOR.
239
Treg-derived enkephalin is dispensable for immune suppression
Previously, Tregs have been shown to suppress hyperalgesia following nerve injury by
suppressing IFN-□-induced primary afferent sensitization[17]. Thus, we hypothesized that
a potential mechanism by which Treg-derived enkephalin mediates the suppression of
nociceptive thresholds involves modulation of immunological responses. To address
this possibility, we tested the nociceptive thresholds of immunodeficient $Rag2^{-/-}$ mice
which are missing both T and B cells and compared them to immunocompetent
littermates. In these studies, we mated $Rag2^{+/-}$ mice and were surprised to observe
decreased nociceptive thresholds in the $Rag2^{-/-}$ offspring compared to their $Rag2^{+/+}$ and
$Rag2^{+/-}$ littermates. This finding suggests that there may be a mechanism of Treg-
mediated control of nociceptive thresholds, which is independent of exaggerated
lymphocyte-driven inflammation **(Figure 5A)**. Furthermore, consistent with this
conclusion, depleting macrophages through liposomal clodronate administration did not
reverse the mechanical allodynia observed in female mice deficient in mTreg **(Figure
5B)**[45].
255
We also used the $Penk$DTR$^{\Delta heme}$ bone marrow chimeric mice to deplete all bone
marrow-derived enkephalin lineage cells and assessed their contribution to the
regulation of immune responses. Unlike $Foxp3$-DTR mice, we observed no changes in
mouse weight or spleen size in $Penk$DTR$^{\Delta heme}$ bone marrow chimeric mice chronically
injected with systemic DT **(Figure S5D-E)**. Furthermore, we also did not observe any
specific alterations in CD4$^+$ T cell cytokine production after nerve injury in the meninges
or lymphoid organs. We conclude, therefore that peripheral enkephalin does not
contribute to T cell-driven inflammatory responses **(Figure S5F-G)**.
264
We next investigated the contribution of Treg-derived enkephalin in the regulation of
conventional T cell proliferation. We assessed T cell suppression capacity by co-
culturing naïve conventional CD4$^+$ T cells with either WT $Penk^{+/+}$ or $Penk^{-/-}$ Tregs. We
observed no difference in the suppressive capacity of $Penk^{-/-}$ Treg compared to control
Tregs **(Figure 5C-D)**. Next, we transplanted equal amounts of CD45.1 $Penk^{+/+}$ or
CD45.2 $Penk^{-/-}$ CD4$^+$ T cells into $Rag2^{-/-}$ mice and performed SNI to measure chimerism
of congenic markers among CD4$^+$ T cells. We did not observe a competitive advantage
or disadvantage amongst $Penk^{-/-}$ CD4$^+$ T cells across various tissues **(Figure 5E-G)**.
Restimulating harvested T cells from distinct tissues with PMA/Ionomycin revealed no
differences in T cell differentiation across Tregs, T helper 1 (Th1) and Th17 subsets
between the $Penk$-sufficient or deficient T cells **(Figure 5H and Figure S5H-I)**. In
addition, we noticed no difference in weight, health, or spleen size between $Penk^{\Delta heme}$

and WT^{Δheme} bone marrow chimeric mice further revealing that peripheral enkephalin
has a very limited, if any, role in the suppressing of systemic inflammatory responses
(Figure S5J-K).
280
Finally, using an adoptive transfer-based graft versus host disease (GVHD) model, we
assessed whether Treg-derived enkephalin is required for suppressing immune
responses. As expected, we observed a profound Th1 response in mice transferred with
activated Tconv alone. However, mice that received additional transfers of either
Penk$^{+/+}$ or *Penk*$^{-/-}$ Tregs equally suppressed Th1 responses, reduced GVHD severity
and mitigated weight dysregulation **(Figure 5 I-J and S5L-N)**. In summary, we conclude
that Treg-derived enkephalin does not contribute to any inflammatory response restraint
mechanism. Rather, we conclude that Tregs can suppress pain sensitivity through a
mechanism that is independent of their function in immunosuppression.
290
Delta opioid receptor signaling on MrgprD$^+$ primary afferent DRG neurons is
required for the anti-allodynic function of mTregs
Enkephalin is a potent agonist at the δOR, and, to a lesser extent the μOR. In our
previous studies, we demonstrated the divergence of expression and function of δOR
and μOR in mediating distinct pain modalities. Specifically, δOR is expressed on
nonpeptidergic IB4$^+$ unmyelinated as well as myelinated primary afferents and
selectively regulates mechanical thresholds and nerve injury-induced mechanical
hypersensitivity[21]. Conversely, the μOR is expressed on Trpv1$^+$ nociceptors and
selectively regulates thermal hyperalgesia. In addition, a spinal δOR can dampen
mechanical hypersensitivity by inhibiting the excitability of somatostatin-positive dorsal
horn interneurons[23].
302
To assess the requirement of PNS or CNS δOR circuits in coordinating the anti-
allodynic effect of mTregs, we intravenously injected *Oprd1*$^{+/+}$ control or *Oprd1*$^{fl/fl}$ mice
with AAV.PHP.S-CAG-Cre or AAV.PHP.eB-CAG-Cre. This approach selectively
introduces Cre recombinase and targets deletion of δOR into to the PNS (DRG) or CNS
(spinal cord and brain), respectively **(Figure 6A).** Three weeks after the AAV injection,
we performed SNI and four weeks later administered IL-2. Mice selectively lacking δOR
in the PNS lost the capacity to respond to the anti-allodynic effect of IL-2, but the effects
of IL-2 were preserved in mice lacking δOR in the CNS **(Figure 6B-C)**. We conclude
that a sensory neuron-expressed, presynaptic δOR coordinates mTreg suppression of
mechanical pain hypersensitivity.
313
Next, we identified the specific sensory neuron subset that coordinates the anti-
nociception mediated by mTreg-derived enkephalin. Previous studies of δOR
expression on DRG sensory neurons using *Oprd1*^{eGFP} reporter mice revealed that
approximately half of the reporter-positive cells in the DRG are myelinated neurons,
while approximately 36% are IB4$^+$ non-peptidergic neurons expressing the MrgprD
receptor. Using established single cell RNA sequencing resources, we found that the
MrgprD$^+$ subset of DRG sensory neurons not only expresses *Oprd1*, but also other
receptors for Treg ligands, including *Il10ra and Icosl*, which have been implicated in
suppression of pain thresholds[46–48] **(Figure 6D)**. The total proportion of sensory neurons

expressing the *Oprd1* transcript matches previous data using *Oprd1*[eGFP] reporter mice
(Figure 6E). Our subsequent flow cytometry-based profiling of *Oprd1*[eGFP] reporter
expression on DRG cells confirmed GFP expression specifically on IB4[+] CD45[-] Thy1[+]
sensory neurons, which corresponds to the MrgprD[+] nonpeptidergic nociceptive neuron
population. Importantly, the flow analysis confirmed the absence of GFP expression on
broadly defined CD45[+] CD90.2[+] IB4[-] lymphoid cells and CD45[+] CD11b[+] CD90.2[-] IB4[-]
myeloid cells **(Figure 6F)**. We also confirmed the absence of GFP expression on
microglia as well as on immune cells profiled from the draining lymph nodes (data not
shown)[23]. Based on this selective Oprd1 expression profile, we generated mice in which
MrgprD[+] neurons lack the δOR (*MrgprD*[Cre-ERT2];*Oprd1*[fl/fl]). Female *MrgprD*[Cre-
ERT2];*Oprd1*[fl/fl] mice, but not their male counterparts, exhibited exaggerated mechanical
hypersensitivity after SNI compared to tamoxifen-injected sex-matched littermate
controls **(Figure 6G-H)**. Female mice lacking δOR on MrgprD[+] sensory neurons and
treated with IL-2 IT four weeks following SNI displayed a complete deficiency in IL-2
anti-allodynic efficacy **(Figure 6I-J)**. We conclude that, the enkephalin receptor δOR,
expressed specifically by MrgprD[+] sensory neurons, mediates the anti-nociceptive
function of mTregs.
340
341
<u>Discussion</u>
In this report, we describe a novel, sexually dimorphic mechanism for pain regulation by
the immune system. Using a range of site-selective targeting strategies to deplete or
expand Tregs within the recently recognized borders of the nervous system, specifically
the meninges, we find that meningeal Tregs (mTregs) can profoundly modulate
mechanical hypersensitivity. Strikingly, this pain regulatory function of mTregs is sex-
specific and controlled by gonadal hormones. Although proenkephalin expression by
Tregs has been observed in sequencing studies, its functional relevance to nociception
had not been explored. Here, we demonstrate that enkephalin secreted by mTregs acts
on δ-opioid receptors (δOR) on primary sensory neurons to selectively modulate
mechanical sensitivity. Our findings provide the first mechanism of Treg-mediated
suppression of nociception and establish regulatory T cells as key sentinels of pain
homeostasis.
355
To assess for sex differences in transcriptional identify after nerve injury, primary
afferent neurons, including those expressing MrgprD, have recently been sorted for
deep RNA sequencing[52]. Surprisingly, very few differences were found between the
sexes, suggesting a lack of a strong intersection of sex and injury in the transcriptional
identity of peripheral neurons. The lack of transcriptional differences may also suggest
that non-neuronal cells may be a primary determinant of sex-selective pain modulation.
Indeed, preclinical research has demonstrated the involvement of T cells in driving pain
phenotypes in female mice[53]. Additionally, human leukocyte antigen (HLA) risk alleles
have been identified for human chronic pain conditions, further suggesting a potential
role for T cells in pain modulation[24,54].
366
Regulatory T cells display broad tissue supportive roles that extend beyond their
originally described function in suppressing inflammation[8,11,12,20]. A major advantage of

our analysis is that we utilized a site-selective mTreg ablation strategy that preserved
peripheral Tregs and avoided systemic inflammation. Using this approach, we identify a
novel, sex-specific mechanism by which Tregs modulate nociceptor activity to regulate
pain sensitivity in the context of health and nerve injury. Although, proenkephalin-
expressing Tregs have been identified in various tissues, their functional assessment
has been limited[19,37,38]. Somewhat paradoxically, a recent pre-print study demonstrated
a small but statistically significant decrease in basal heat sensitivity in both female and
male mice conditionally depleted for *Penk* expression in systemic Tregs. Mechanical
thresholds and other sensory modalities were however not examined[55]. In distinct
contrast, we uncover a sensory modality selective function of mTreg that is consistent
with prior findings of δOR agonism, namely providing relief of mechanical but not heat
pain[21,56]. Whether mTregs tonically restrain nociception is difficult to conclude. Our
finding of increased basal sensitivity in $Rag2^{-/-}$ or $Penk^{\Delta heme}$ mice, is supportive of this
hypothesis, however, alternative possibilities exist. It is conceivable that mTreg
deficiency could lead to inflammation, which may alter nociceptor sensitivity.
Additionally, the endogenous opioid signaling pathway may have a role in opioid
induced analgesia. It is significant that naloxone, a non-selective opioid receptor
antagonist IV injection, does not induce pain in healthy individuals. Whether
experiments in uninjured female mice have been performed is unclear[57].
388
Our observations suggest that sex hormones, rather than sex chromosomes, are the
main drivers of mTreg-induced anti-nociception. While the specific sex hormones
involved in regulating mTreg function in pain remain unclear, previous studies have
implicated estrogen and progesterone in modulating neuropathic pain associated with
SNI[49]. Estrogen administration increases *Penk* expression in the whole spinal cord[58].
Additionally, in CD4$^+$ T cells, the estrogen receptor engages conserved non-coding
sequences (CNS) in *Foxp3* enhancer regions[59]. Importantly, we reveal an anti-
nociceptive role for regulatory T cells that is distinct from their well-established functions
in immune suppression and tissue repair. Furthermore, we demonstrate that this
mechanism operates within nervous system tissues, at a site distant from the peripheral
nerve injury, highlighting the immune system's remarkable ability to modulate
nociception.

415
416
417

Methods

Mice

All mouse experiments were approved by UCSF Institutional Animal Care and Use Committee and conducted in accordance with the guidelines established by the Institutional Animal Care and Use Committee and Laboratory Animal Resource Center. All mice experiments were performed on age-matched adult male and female mice at a starting age between 8 and 14 weeks old. Littermate controls were used for all experiments when feasible. Mice were bred in-house and backcrossed over 10 generations to C56BL/6 breeders obtained from Jackson labs. Experimental mice were co-housed to maintain the same microbiome. They were maintained in a temperature (21°C) - and light (12h light/dark cycle)-controlled environment and were provided with food and water *ad libidum*. The following mouse strains are used: C57BL/6J (JAX #000664), *Foxp3*-DTR (B6.129(Cg)-Foxp3tm3 (Hbegf/GFP)Ayr/J, JAX# 016958), *Foxp3*$^{eGFP-Cre-ERT2}$ (Foxp3tm9 (EGFP/cre/ERT2)Ayr/J; JAX#016961), Ai9 (B6.Cg-*Gt(ROSA)26Sor*$^{tm14(CAG-tdTomato)Hze}$/J; JAX#007914), Four Core Genotypes (B6.Cg-Tg(Sry)2Ei Srydl1Rlb T(XTmsb4x-Hccs;Y)1Dto/ArnoJ; JAX#010905), Penk-IRES2-Cre(B6;129S-Penktm2(cre)Hze/J; JAX#025112), Rosa26-ISL-iDTR (CBy.B6-Gt(ROSA)26Sortm1(HBEGF)Awai/J;JAX#008040),CD45.1 (B6.SJL-Ptprca Pepcb/BoyJ; JAX# 002014), B6C3F1/J (JAX#100010), *Rag2*$^{-/-}$ (B6.Cg-Rag2tm1.1Cgn/J; JAX#008449), Mrgprd-CreERT2 (Mrgprdtm1.1(cre/ERT2)Wql/J; JAX# 031286), Oprd1fl/fl (B6;129-Oprd1tm1.1Cgrf/KffJ; JAX# 030075), DOR-eGFP (B6;129S2-Oprd1tm2Kff/J; JAX#029012). *Penk*$^{-/-}$ were kindly provided by Dr. John Pintar on a C57BL/6 background (MGI 3628668)[60]. *Foxp3*-DTR (X-linked) was mated with male Four Core Genotypes XY^{Sry-}Sry^{Tg} and the X$^{Foxp3-DTR}$YSry$^-$Sry^{Tg} male mice were mated to homozygous *Foxp3*-DTR female mice.

Bone marrow transplantation

CD45 mismatched host recipient mice were irradiated at 550 cGy twice, 5 hours apart and injected retro-orbitally with 5x10^6 cells from the bone marrow of CD45.1 WT, *Penk*^{Cre};*Rosa26*^{DTR} *Penk*$^{-/-}$ or a 1:1 mix of *Foxp3*-DTR and *Penk*$^{-/-}$ sex-matched 7-10 week old mice. Mice were kept on doxycycline chow for the first week (Bioserv #S3888) and chimerism was assessed at 8 weeks post-transplant.

Pharmacological interventions

Animals were randomly assigned to vehicle control or treatment group. Pegylated diphtheria toxin (pegDT) was a generous gift from Ana I. Domingos and generated as previously described[61]. pegDT (20 ng) or corresponding phosphate-buffered saline (PBS) control were injected intrathecally (IT) in a volume of 5μL in naive mice below lumbar level L4. All intrathecal injections were performed in non-anesthetized, lightly restrained mice and injections were validated by a sudden flick of the tail. Of note, the 5-10 μL injection distributes predominantly to the lumbo-sacral cord given lidocaine IT injection at that segment paralyzes the hindpaws but not forepaws. Non-pegylated

diphtheria toxin (30 ng/g, Sigma Cat#322326) or corresponding saline control were
administered in a volume of 200 µl every three days intraperitoneally (IP). IL-2 (0.1 µg,
Peprotech, Cat# 212-12) or PBS vehicle control were administered daily for three
consecutive days IT. Selective δOR agonist, [D-Ala2]-Deltorphin 2 (15 µg, Abcam Cat
#ab120708, CAS 122752-16-3) and selective δOR antagonist, naltrindole (5 µg, Sigma
Cat# N115) administrations were performed 30 min before behavior experiments.
467
468 **Tamoxifen injections**
469 *Mrgprd*[CreERT2];*Oprd1*[fl/fl] and *Foxp3*[eGFP-Cre-ERT2];*Rosa26*[tdTomato] mice were injected IP with
470 tamoxifen (Sigma Cat #5648) 100 mg/kg in corn oil (Sigma Cat #8267) for five
471 consecutive days to induce Cre-mediated recombination.
472
Intrathecal dye tracing
Naive mice were intrathecally injected with 5µL of Evans Blue 1% solution (Sigma, Cat
#E2129) or pegylated DyLight 650-4xPEG NHS Ester (Thermo Fischer, Cat #62274).
24 hours post-injection, mice were anesthetized with avertin and euthanized by
decapitation. Spinal cord and brain meninges, spinal cord, brain, dorsal root ganglia and
trigeminal ganglia, sciatic nerve and lymph nodes were assessed for dye uptake.
479
480 **Animal behavior**
481 For all behavioral tests, the experimenter was blind to genotype and treatment and
482 performed during the light cycle. The project utilized both male and female
483 experimenters but a predominant number of experiments were performed by a female
484 investigator[62].
485
von Frey measurement of mechanical hypersensitivity
Mice were acclimatized once to the von Frey apparatus for two hours. The lateral
plantar surface of the ipsilateral and contralateral hind paws (sural innervation) was
stimulated with von Frey hairs of logarithmically increasing stiffness (Stoelting Cat #
58011). Animals were habituated on a wire mesh for 1 hour, after which they were
tested with von Frey filaments (0.008, 0.02, 0.04, 0.07, 0.16, 0.4, 0.6, 1, 1.4, 2, 4 and 8
g) using the Dixon up–down method[63,64]. The von Frey hairs were held for 3 sec with
intervals of several minutes between each stimulation. For the Dixon up-down method
we recorded 2 days of baseline mechanical sensitivity which were averaged. After SNI,
nearly all mice reached a 50% paw withdrawal threshold of the lowest filament (0.008
g), thus we utilized a single fiber method of testing to achieve resolution of allodynia
severity. Mice were stimulated 10 times with the 0.008 g filament. The filament was
applied for 3 seconds and the number of positive responses across the 10 stimulations
were registered as percent nociceptive responses.
500
501 **Hargreaves measurement of heat hypersensitivity**
502 Mice were acclimatized for 30 min in plexiglass cylinders. The mice were then placed on
503 the glass of a Hargreaves apparatus and the latency to withdraw the paw from the heat
504 source was recorded. Each paw was tested three times and latencies were averaged
505 over the trials.
506

Acetone induced cold sensitivity

Mice were habituated for 60 min on a mesh in plexiglass cylinders. A syringe was used to spray 50 µl of acetone (Thermo Scientific Cat # 423240010) onto the plantar surface of the paw and the behaviors were video recorded for 30 seconds after each trial using a Sony HDR-CX440 camera. The left hind paw was tested five times and positive responses included withdrawals, shakes, licks and jumps. Results are displayed as the total number of behaviors across the five trials.

Tail flick measurement of heat hypersensitivity

Mice were placed in a restrainer and 2 cm of the tip of the tail was submerged in a 52°C water bath. The latency (seconds) to withdraw the tail from the water was recorded. A cut-off of 15 s was set to prevent tissue damage and testing was performed with intervals of several minutes between each stimulation. Mice were tested three times and withdrawal latencies were averaged.

Hot plate measurement of heat hypersensitivity

Mice were acclimated to the testing environment as described above. The hot plate temperature was set to 52°C. The mouse was placed on the plate and the latency to shake, lick or bite a hindpaw was scored. A cut-off of 20 s was set to prevent tissue damage.

Pin prick withdrawal test

Mice were habituated for 60 min on a mesh in plexiglass cylinders. A 27G needle was gentle applied onto the hindpaws, with minutes between each stimulation for a total of 5 stimulations per paw. Mice were scored as follow: 1: brief withdrawal, 2: mice lifting their paw for 1 sec, 3: mice lifting their paw for 2 sec or shake, 4: lick. Results are displayed as the total number of behaviors across the five trials.

Brush withdrawal test

Mice were habituated for 60 min on a mesh in plexiglass cylinders. A 5-0 brush was gently applied onto the hindpaws, with minutes between each stimulation for a total of 5 stimulations per paw. Mice were scored as follow: 1: brief withdrawal, 2: mice lifting their paw for 1 sec, 3: mice lifting their paw for 2 sec or shake, 4: lick. Results are displayed as the total number of behaviors across the five trials.

Rotarod

Mice were acclimatized to the testing room and trained by placing them on an accelerating rotarod for a maximum of 60 seconds at low speed, three times with training taking place on two consecutive days. Latency to fall was measured with a cutoff of 300 seconds. The procedure was repeated three times and latencies averaged across trials.

Spared Nerve injury

We employed an established, robust and reliable spared nerve injury (SNI) model to induce a chronic neuropathic injury[65]. This model utilizes non-healing surgical intervention on two branches of the sciatic nerve (the common peroneal and tibial

branches), while sparing the third branch (the sural branch) for sensory testing on the
lateral portion of the hindpaw. Briefly, mice were anesthetized with isoflurane (3% for
induction and 1.5% for maintenance, mixed with oxygen). The fur on the left hind leg
was shaved and disinfected with 3 passages of alcohol and iodine solution,
alternatively. A 1 cm incision was performed on the upper thigh skin, near the division
point of the sciatic nerve. A 2% lidocaine solution was applied and the biceps femoris
muscle was gently separated through a blunt opening to reveal the sciatic nerve's
common peroneal, tibial, and sural branches. The common peroneal and tibial nerves
were ligated with non-dissolvable 8-0 silk sutures (Fine Science Tools Cat # 12052-08).
Subsequently, a 2 mm segment from both the common peroneal and tibial nerves was
transected, ensuring the sural nerve remained undisturbed. The muscle and the skin
were stitched using 6–0 sutures (Henry Schein Surgical suture Cat #101-2636), and the
skin was further sealed with a tissue adhesive (3M Vetbond Cat # 1469SB), after an
ethanol solution application. Mice were kept on heating pad until they regained
consciousness and demonstrated stable, balanced locomotion. Mice were transferred
into their home cage and observed meticulously for the next two days.
569
Immunohistochemistry
Avertin-anesthetized mice were transcardially perfused with 10 ml of 1× PBS followed
by 30 ml of 4% paraformaldehyde (PFA, Thermo Scientific Cat # 119690010) diluted in
PBS. After perfusion, spinal cord, sciatic nerves, lymph nodes, spleens, brains and
DRG were collected, postfixed in 4% PFA solution at 4°C for 5 h and then
cryoprotected in 30% sucrose in PBS at 4°C.
576
Spinal meninges were harvested from fixed spinal cords. Spinal cords were transferred
in PBS and meninges were gently peeled into a single sheet onto a microscope slide
after a longitudinal hemisection of the spinal cord. Brain meninges were similarly
harvested from the skull. Frozen tissues were embedded at –35°C in O.C.T. compound
and 30 µm transverse spinal cord sections were generated using a Leica SM220R
sliding microtome and 20 µm DRG sections were generated using a cryostat (Thermo
Fisher Scientific) on SuperFrost Plus slides. Spinal cord sections were processed as
free-floating. Sections were blocked (10% NGS, 1% BSA, 0.05% Tween-20, 0.1% Triton
X-100 in PBS) and incubated in 0.3 M glycine containing 0.2% Tween 20. Sections
were labeled in blocking buffer for 24 hours at 4°C. Slides were coverslipped with
Fluoromount-G (Thermo Fisher Scientific). Fluorescence images were acquired using
an Olympus FV3000 confocal microscope and quantified using ImageJ (Fiji).
589
Tissue clearing
Whole DRG or spinal cords from *Foxp3*[eGFP-Cre-ERT2];Ai9 mice were cleared after PFA
fixation using SHIELD tissue clearing (LifeCanvas PCK-500)[66]. Tissues were washed in
PBS then processed as previously described[66]. Briefly, the tissues were incubated in
epoxy solution (SHIELD OFF) for 10 hours at 4°C with gentle shaking then incubated
overnight at 37°C in SHIELD ON-Epoxy solution for epoxy crosslinking. DRG were then
further incubated in SHIELD ON solution for 10h and delipidated for two days (DRG) to
five days (spinal cord) at 45°C with shaking then washed with PBS. Whole mount DRG

and spinal cords were acquired in FocusClear reflexive index matching solution
(CelExplorer, FC-102).
600
Tissue digestion
Mice were injected intravenously with 50 µl of anti-ARTC2 nanobody (Biolegend Cat #
149802) in 200 µL of PBS 30 minutes before euthanasia to protect Treg during harvest
from purinergic-mediated cell death[67]. 5 minutes before harvest mice were injected
intravenously with 6 µg of FITC-conjugated anti-CD45 antibody to label blood immune
cells in 200 µL of PBS [27]. Avertin-anesthetized mice were decapitated, and spinal cord
meninges, brain meninges, L4-6 DRG, sciatic nerves, lymph nodes, brains and spleens
were harvested. Spinal cord meninges, brain meninges, DRG, sciatic nerves and brains
were crushed with the back of a 3ml syringe in a serrated 24 well plate and triturated in
digestion media (Liberase TM (0.208 WU/ml) (Roche Cat # 054010200001) and DNase
I (40 ug/ml) (Sigma Cat # DN25) in 1.0 ml cRPMI (RPMI supplemented with 10%
(vol/vol) fetal bovine serum (FBS), 1% (vol/vol) HEPES, 1% (vol/vol) Sodium Pyruvate,
1% (vol/vol) penicillin-streptomycin). They were digested for 30 min at 37°C, 220 RPM
and triturated every 15 minutes. Digested samples were again triturated and passed
over a 40 µm cell strainer and any remaining tissue pieces mashed through the cell
strainer. Cell strainers were flushed with staining media (PBS w/o Mg^{2+} and
Ca^{2+} supplemented with 3% FBS, 2 mM EDTA and 0.05% NaN_3). Single-cell
suspensions were centrifuged at 500 g at 4°C, washed and resuspended in staining
media. Spleens and lymph node immune cells were obtained by mashing the tissues
over a 40 µm cell strainer and washed with staining media.
621
Cell stimulation
Isolated single cell suspensions were incubated for 4 hr at 37°C in complete IMDM
(supplemented with 10% FBS, 2 mM L-glutamine, 10 mM HEPES, 1% sodium pyruvate,
1% penicillin-streptomycin, 50 µM 2-mercaptoethanol) with (phorbol 12-myristate 13-
acetate) PMA, Ionomycin in the presence of Brefeldin A and Monensin (Tonbo, TNB-
4975).
628
Flow cytometry
Single-cell suspensions were stained in a 96 well plate. Briefly, they were washed with
250 µL of staining media and stained with viability dye (1:500) and cell surface
antibodies (1:100) in 100 µL of staining media with Fc shield (1:100). Samples were
washed twice in staining media and stained for intracellular cytokines or for Foxp3
according to manufacturer recommendations with BD Cytofix/Cytoperm
Fixation/Permeabilization Kit (BD 554714, AB_2869008). Samples were further stained
with conjugated intracellular antibodies (1:100) overnight at 4°C in BD
permeabilization/wash buffer. Samples were washed with permeabilization/wash buffer
twice and resuspended in 200 µL of staining media. For visualization of both tdTomato
reporter signal and intranuclear Foxp3 signal, cells were fixed in 200 µL of freshly
prepared 2% formaldehyde (EM grade) in PBS for 60 minutes exactly and washed with
eBioscience Permeabilization buffer (eBioscience Foxp3 perm-kit) and stained
overnight in 1x eBioscience Permeabilization buffer at 4°C[68]. Cells were washed twice
in the buffer and resuspended in staining media. For Met-enkephalin staining, we

screened multiple commercially available antibodies and selected an antibody that
showed positive staining in wildtype mTreg but not *Penk*[-/-] mTreg. The antibody was
conjugated to fluorescent phycoerythrin with Lightning-link conjugation kit (ab102918)
and utilized after cell stimulation and intracellular cytokine staining. Cells were counted
with 50 µL of counting beads (Thermo Fisher Scientific Cat # C36950) and samples
were analyzed using a BD FACSCanto2 or BD FACS Aria Fusion flow cytometer (BD
Biosciences). Positive and negative selection gates were set using fluorescence minus
unstained cells. For negative control of enkephalin staining, *Penk*[-/-] samples were used
for gating. Fluorescence intensity distribution was analyzed using FlowJo 10 software
(BD Biosciences). Antibodies for flow cytometry are listed in the resource table. Lineage
exclusion markers include viability dye, CD11b (to exclude myeloid cells), B220 (to
exclude B cells), Ter119 (to exclude red blood cells).
656
Cell sorting
Spleens were mashed on a cell strainer and cells were pelleted at 500 g for 10 min at
4°C. Cells were stained for viability and lineage exclusion markers, in addition to CD45,
CD4, CD45RB to stain naïve Tconv and CD25 to stain Tregs. Cells were pelleted,
washed and incubated with CD4 negative selection beads and purified on a LS
magnetic column (Miltenyi Biotec). Cells were double sorted in BD FACS Aria Fusion for
Singlet, Live, CD45[+] CD4[+] CD45RB[+] CD25[-] for Tconv or Singlet, Live, CD45[+] CD4[+]
CD25[+] CD45RB[-] for Treg into complete IMDM.
665
T cell suppression assay
Sorted Tconv from lymphoid organs of CD45.1 female mice were labeled with cell trace
violet (Thermo-Fisher, Scientific #C34571) according to manufacturer instructions.
0.25×10^5 Tconv per well were cultured with distinct dilution of Treg from CD45.2 WT or
Penk deficient lymphoid organs. Cells were then washed and resuspended with mouse
anti-CD3/CD28 Dynabeads at a 1:1 ratio of beads to Tconv. Cells were incubated for 96
hours in a humidified incubator at 37°C. Cells were washed and resuspended in staining
media and suppression ratio was calculated by dividing the percent proliferated cells
from incubated Tconv[+] Treg samples by percent proliferated cells from Tconv only
samples[69].
676
677 **T cell adoptive transfer**
678 *Competition assay*
679 1×10^6 negatively selected bulk CD4[+] T cells from WT female and *Penk* deficient
680 lymphoid organs were transplanted into female *Rag2*[-/-] mice. SNI was performed and
681 organs were collected 28 days later for cell stimulation and flow cytometry.
682
Graft versus host disease (GVHD)
GVHD was established as previously described. Briefly, WT or *Penk* deficient female
mice bone marrow was transplanted into MHC mismatched B6C3F1/J female mice to
activate T cells. 0.25×10^6 sorted WT Tconv were transplanted into male *Rag2*[-/-] mice to
induce chronic GVHD either without Tregs or in the presence of 0.125 $\times 10^6$ WT or *Penk*
deficient Tregs. Mice were measured for weight changes and GVHD score. Mice were
euthanized then harvested for cytokine secretion assay by flow cytometry[70,71]. GVHD

scoring is as follows: 0 = no signs of GVHD, 1 = visible signs of GVHD (hunching,
lethargy, ruffled fur), 2 = no weight gain, 3 = 0-5% weight loss, 4 = >5% weight loss.
One Tconv mouse did not survive for harvesting for cytokine stimulation.
693
Analysis of sequencing data
Bulk RNA-seq
Raw files GSM4677053-064 from GEO dataset GSE154680, and all files from GSE
GSE180020 were gathered and aligned using STAR for uniquely mapped reads
(outFilterMultimapNmax 1–outFilterMatchNmin 30–alignIntronMin 20–alignIntronMax 1-
0000). Data was annotated with GENCODE GRCm38/mm10 genome assembly. Raw
count tables were normalized by median of ratios method with DESeq2 package from
Bioconductor to analyze for differential expression.
702
ATAC-seq
Fastq files were gathered from SRR12264679-94 from GSE154680. Raw reads were
mapped to the mouse mm10 genome assembly using STAR alignment (--
outFilterMismatchNoverLmax 0.04 --outFilterMismatchNmax 999 --
alignSJDBoverhangMin 1 --outFilterMultimapNmax 1 --alignIntronMin 20 --
alignIntronMax 1000000 --alignMatesGapMax 1000000). Bam files were generated by
STAR. PCR duplicates were removed by Picard, and peak calling performed using
MACS2 (--keep-dup 1 --bw 500 -n output --nomodel --extsize 400 --slocal 5000 --llocal
100000 -q 0.01) PMID: 22936215). To generate bigwig files for ATAC-seq datasets, all
aligned bam files were merged by condition using samtools merge. Bedtools
genomecov was run to covert the merged bam files into a bedgraph files. Finally,
bedGraphToBigWig (ucsc-tools/363) was used to generate the bigwig files displayed on
browser tracks using the IGV browser and compared to existing encode ATAC and
Chip-Seq peaks.
717
scRNA-seq
Fastq files were gathered from GEO from datasets GSE139088 GSE201653 and initial
counts were obtained using the Cell Ranger pipeline[48,72]. Using Seurat v4, individual
cells were removed from the data set if they had fewer than 1000 discovered
genes/features, fewer than 1000 UMI or greater than 10% reads mapping to
mitochondrial genes. 2000 variable genes were found for each normalized library, and
anchors were selected for integration with dimensionality of each dataset set at 30. Glial
cells noted for the markers of *Sparc* and *Mpz* and non-neuronal cells lacking the
expression of Avil were excluded. Variable genes were identified from the merged
dataset, and PCA and UMAP were ran to generate new UMAP coordinates with a
dimensionality of 30 and clustering was performed with a resolution of 0.5.
Findallmarkers function utilizing a Wilcoxon rank-sum test was used to find cluster
specific markers and annotation was performed as recently established[48]. Number of
Oprd1 expressing cells were defined by a threshold of non-zero expression.
732
Statistical analysis
Statistical analysis was performed using GraphPad Prism 9 software. Data are
presented as mean ± SEM. Differences pre- and post-injection within a single group

were assessed using a Wilcoxon matched-pairs signed rank test. Differences between
two groups were assessed using a Mann-Whitney test. Statistical analysis for multiple
comparisons were performed using Kruskal-Wallis test followed by Dunn's multiple
comparison test or a Two-Way ANOVA followed by Sidak's multiple comparison test.
$p < 0.05$ (*), $p < 0.01$ (**), $p < 0.001$ (***).
741

Acknowledgement: We thank Dr. Dena Dubal and her laboratory for the FCG mouse,
Dr. Kevin Yackle for $Penk^{Cre}$ mice, Dr. Ari Molofsky for $Foxp3^{Cre-ERT2}Rosa26^{TdTomato}$
mice, Dr. Amynah Pradhan for $Oprd1^{eGFP}$ mice and Dr. Mike Ansonoff for assistance
with $Penk^{-/-}$ tissue isolation. We thank additional members of the Basbaum laboratory
and UCSF ImmunoX for critical feedback. Funding for this work was supported by
grants Canadian Institute of Health Research (CIHR) (to É.M.), the Fonds de Recherche
en Santé-Québec (to É.M.), the Dermatology Foundation (Career Development Award
to S.W.K.), the Sandler Foundation PBBR (to S.W.K), Grunfeld Scholar Award from
SFVAMC (to S.W.K), T32AR007175-44 (to S.W.K), NIH NSR35NS097306 (to A.I.B.)
and Open Philanthropy (to A.I.B.). Figures were generated with BioRender.com.

752
753　**Disclosures:** Authors have no conflicts of interests to declare.
754
755　**Author contributions:** É.M. and S.W.K. designed experiments. É.M., B.C.M, J.B.,
756　S.R. N.P.K. W.L.E. and S.W.K. performed experiments, data analysis or visualization.
757　J.E.P. and A.I.D. provided critical reagents or tools. S.W.K, É.M. and A.I.B., acquired
758　funding and provided supervision. É.M, A.I.B and S.W.K. wrote the manuscript.
759
760　**Material availability and requests:** No new reagents, original code or original genomic
761　datasets were generated. Requests for reagents or mice can be sent to
762　sakeen.kashem@ucsf.edu or allan.basbaum@ucsf.edu.
763
764　**Diversity, equity, and inclusion statement:** Authors support diversity and inclusion
765　values. At least one author, including the lead author, self-identifies as a woman. At
766　least one author identifies as an under-represented minority and/or as an immigrant.

Figure Legends

Figure 1. mTreg suppress mechanical pain hypersensitivity in female mice.
Representative whole mount maximum projection confocal microscopy image of the (A) lumbar spinal cord meninges and (B) DRG showing Tregs (green-red: yellow) and nerves (autofluorescence, red) in $Foxp3^{eGFP-CreERT2};Rosa26^{tdTomato}$ reporter mice. Inlet showing DRG magnification. Scale bar represents 100 μm in A) and 150 μm in B). Arrows indicate Tregs. (C) Total number of weight-adjusted tissue Tregs across organs, in both sexes combined. (D) Relative number of tissue Tregs from male (white) and female (black) mice per organ. 100% represents mean number of female Tregs per organ. Comparison is made between each individual organ. (E) Representative concatenated flow cytometry plots of tissue Treg after a single intrathecal (IT) injection of 20 ng of pegylated diphtheria toxin (pegDT). (F) Relative quantifications of tissue Treg depletion 2 days after a single IT pegDT injection across organs. 100% represents mean number of tissue Tregs in IT vehicle-injected mice per organ. (G and H) 50% paw withdrawal thresholds measured using von Frey filaments before (day 0) and after a single dose of 20 ng of IT pegDT or vehicle in female (G) or male (H) $Foxp3$-DTR mice. (I) Summary of significant behavioral differences comparing IT pegDT- and control-injected female and male mice. Total number of mice for G-I is presented in Figure S3. ScMg= Spinal cord meninges, BrMg= Brain meninges, LN= Lymph nodes, ns = not significant, $*p<0.05$, $**p<0.01$, $***p<0.001$. **Related to Figure S1-S3.**

Figure 2. Expanding mTreg alleviates nociception dependent on sex hormones and independent of tissue repair.
(A) Schematic representation of the spared nerve injury (SNI) surgery. (B) Long-term assessment of mechanical thresholds in mice following SNI surgery (both sexes combined, no difference between the sexes). n = 7-8 mice per group. (C and D) Percent response to 0.008 g von Frey filament in mice with SNI (day 0) and treated with IT pegDT or vehicle every 4 days. n = 8 per group for females and 9-10 per group for males. (E) Schematic representation of mTreg expansion in mice 4 weeks after SNI by 3 IT injections of low-dose IL-2 (0.1 μg). (F) Total mTreg number in meninges after low-dose IL-2 or vehicle IT injections (both sexes combined, no differences between the sexes). (G and H) Nociceptive thresholds of females (G) and male (H) mice given low-dose IL-2 or vehicle IT 4 weeks after SNI. (I) Schematic representation of the mating strategy of Four Core Genotypes (FCG) $FoxP3$-DTR mice demonstrating resulting XX and XY females and XX and XY male mice. (J and K) Nociceptive thresholds of FCG female (J) and male (K) mice following low-dose IL-2 or vehicle IT injections 4 weeks after SNI. (L) Anti-nociceptive efficacy determined as post IL-2/vehicle injection threshold divided by baseline mechanical threshold in male and female mice with XX (white) or XY (pink) chromosomes. (M) Nociceptive thresholds of FCG $FoxP3$-DTR female and male mice following a single IT pegDT or vehicle injection. (N) Percent baseline nociceptive thresholds determined as post pegDT/vehicle injection threshold divided by baseline mechanical threshold in male and female mice with XX (white) or XY (pink) chromosomes. ns = not significant, $*p<0.05$, $**p<0.01$,$***p<0.001$. **Related to Figure S4.**

Figure 3. mTregs express and produce enkephalin.
(A) Volcano plot of transcription fold change of activated Treg (aTreg) vs resting Treg
(rTreg) and (B) heatmap of relative log2 expression value of aTreg, rTreg and activated
and resting $CD4^+$ $CD25^-$ conventional T cells (aTconv and rTconv) from public dataset
GSE154680 (n=3). (C) Averaged ATAC sequencing (ATACseq) of open chromatin
accessibility peaks on the *Penk* locus in different T cell subsets (n=4 per group,
GSE154680), compared to ATACseq, and histone modification Chip-Seq from public
ENCODE dataset of the p0 developing forebrain, a known enkephalinergic region. (D
and E) Log2 values of *Penk* expression by different unstimulated immune cell types
from Immgen dataset GSE180020. (E) Treg *Penk* expression fold change after cytokine
stimulation compared to vehicle control. (F) Representative PMA:Ionomycin stimulated
mTregs, meningeal $CD4^+$ T cells (mCD4) from WT mice or *Penk*$^{-/-}$ mTreg (control). (G)
Representative flow cytometry plots of Tregs from meninges or secondary lymphoid
organs (SLO) from *Penk*^{Cre}*Rosa26*^{tdtomato} mice. Pink represents non-vascular, tissue
Tregs from transgenic *Penk* lineage reporter mice. Gray represents vascular Treg in
reporter mice while Blue corresponds to tissue Treg from non-transgenic control mice.
(H-I) Number of enkephalin lineage fate reporter positive tissue Tregs in (H) meninges
and (I) secondary lymphoid organs (SLO) in male and female mice. NK: Natural Killer
cells, Tgd: γδ T cells, Mo: Monocytes, MF.rp: Red pulp macrophages, CD4T: $CD4^+$ T
cells, CD8T: $CD8^+$ T cells, B.fo: splenic follicular B cells, DC8: $CD8^+$ dendritic cells,
pDC: splenic plasmacytoid, MF.pc: peritoneal macrophages, MC: myeloid cells, B.mz:
splenic marginal zone B cells, ns = not significant, *$p<0.05$, **$p<0.01$,***$p<0.001$.

Figure 4. mTreg-derived enkephalin controls nociceptive thresholds.
(A) Representative flow cytometry plots of Cre positive (pink) or Cre negative (blue)
$CD45^+$ non-vascular cells from the meninges and the DRG, combined, from
Penk^{Cre}*Rosa26*^{tdTomato} mice. (B) Flow plot shows representative tdTomato negative and
right plot shows tdTomato positive leukocytes, demonstrating the more pronounced
mTreg representation in the enkephalinergic fate cell population. (C) Schematic
representation of bone marrow transplants to generate a global depletion of
enkephalinergic cells, a depletion of hematopoietic-derived enkephalin or a depletion of
Treg-derived enkephalin, respectively. (D-G) Bone marrow chimera of
Penk^{Cre}*Rosa26*^{DTR} → irradiated WT recipients. Nociceptive thresholds after a single
pegDT (pink) or vehicle (white) IT injection in female (D) and male (E) mice. n= 5 per
group. Nociceptive thresholds after SNI and pegDT (pink) or vehicle (white) IT injection
in female (F) and male (G) mice. (H) Nociceptive thresholds at baseline of female WT →
WT (black) or *Penk*$^{-/-}$ → WT (white) bone marrow chimeras. (I) Female *Foxp3*-DTR +
Penk$^{-/-}$ (1:1) → WT mice and tested for nociceptive thresholds after pegDT (pink) or
vehicle (white) IT, n=10 per group. (J) Nociceptive thresholds in SNI mice after pegDT
(pink) or vehicle (white) IT injections. (K) WT SNI female mice given low-dose IL-2 and
naltrindole. (L) Nociceptive efficacy calculated as percent compared to baseline
threshold. ns = not significant, *$p<0.05$, **$p<0.01$,***$p<0.001$.

Figure 5. Treg-derived enkephalin is dispensable for suppressing inflammation.
(A) Baseline nociceptive thresholds of uninjured $Rag2^{+/+, +/- \text{ or } -/-}$ female mice. (B)
Nociceptive thresholds of female $Foxp3$-DTR mice injected with pegDT IT + IV
clodronate (pink) or control (white) liposomes showing peripheral macrophages do not
mediate the nociception induced by mTreg depletion, n=5 per group. (C) Representative
flow cytometry histograms of proliferated conventional T cells (Tconv) alone (pink) or
4:1 with WT Tregs (gray), $Penk^{-/-}$ Tregs (yellow), or unstimulated, un-proliferated cell
trace violet-stained control (blue). Histogram shows cells that have not proliferated. (D)
Suppression of Tconv cell proliferation by different concentrations of WT Tregs (white)
or $Penk^{-/-}$ Tregs (yellow). (E) Schematic representation of competition experiment
showing 1:1 transfer of WT or $Penk^{-/-}$ T cells into $Rag2^{-/-}$ mice. SNI surgery was
performed and organs were harvested 4 weeks later for F-H. (F) Equal competition of
$Penk$ sufficient CD45.1 and $Penk$ deficient CD45.2 CD4$^+$ T cells in the meninges
represented as a concatenated flow cytometry plot, n=4 per group. Representative flow
cytometric plots. (G) Pooled proportion of $Penk$ sufficient CD45.1 and $Penk$ deficient
CD45.2 CD4$^+$ T cells in different organs, n=4 per genotype. (H) Percent of FoxP3$^+$, IFN-
□$^+$ and IL-17A$^+$ CD4$^+$ T cells from G. (I) Representative flow cytometric plots of cytokine
secretion by CD4$^+$ T cells after GVHD induced by transfer of pre-activated Tconv alone
or combined with $Penk^{+/+}$ or $Penk^{-/-}$ Tregs. (J) Weight curves of GVHD mice, n=3-4 per
group. ns = not significant, *p<0.05, **p<0.01,***p<0.001. **Related to Figure S5.**
894
Figure 6. δOR on MrgprD$^+$ sensory neurons is required for mTreg mediated anti-
nociception.
(A) Schematic representation of AAV-induced ablation of $Oprd1$ in the PNS (B-C) or
CNS. (B) Nociceptive thresholds of female mice lacking $Oprd1$ in the PNS after mTreg
expansion compared to controls. (C) Anti-nociceptive efficacy of mTreg expansion. (D)
No difference in nociceptive thresholds in female mice lacking $Oprd1$ in the CNS after
mTreg expansion compared to controls. (E) Heatmap of row normalized expression
from DRG sensory neurons clusters from combined GSE139088 and GSE201653. (F)
Proportions of sensory neuron clusters expressing $Oprd1$ from E. (G) Representative
flow cytometry plot of δOR-GFP (green) expression on IB4$^+$ MrgprD$^+$ DRG sensory
neurons compared to cells from non-reporter mice (purple). Overlaid are lymphoid
CD45$^+$ CD90.2$^+$ cells and myeloid CD45$^+$ CD11b$^+$ cells from the DRG. (H-I) Percent
response to 0.008 g von Frey fiber stimulation after SNI in female (H) or male (I) mice
conditionally lacking δOR on MrgprD$^+$ neurons (pink) or controls (white). (J) Nociceptive
thresholds after mTreg expansion in female mice lacking $Oprd1$ in MrgprD$^+$ neurons
compared to controls. (K) Anti-nociceptive efficacy of mTreg expansion. SA-LTMR=
Slowly adapting low-threshold mechanoreceptor, RA-LTMR= rapidly-adapting low-
threshold mechanoreceptor, MrgprD= Mas-related G protein-coupled receptor D, Prop.=
proprioceptor, SST: somatostatin, Trpm8= Transient receptor potential cation channel
subfamily M member 8, TAM= Tamoxifen, cKO= Conditional KO, ns = not significant,
*p<0.05, **p<0.01,***p<0.001.
916
917
918
919

920
 Supplemental Figure Legends

922
923 **Figure S1. Flow cytometric gating strategy of meningeal Tregs.**
924 (A) Gating strategy to quantify tissue Treg numbers. Numbers indicate gate frequency.
925 Mice were injected IT with ARTC2 nanobody to minimize Treg apoptosis and injected IV
926 with anti-CD45 FITC (pink) antibody or vehicle injected (blue) to label vascular immune
927 cells.

928
Figure S2. pegDT IT injection avoids systemic inflammation and weight loss in
***Foxp3*-DTR mice.**
(A) Evan's blue staining after IT injection showing diffusion into the cerebellum, the
olfactory bulb, the cervical and lumbar lymph nodes, the spinal cord meninges and the
lumbar DRG. (B) pegDyLight650 IT injection exhibits a more limited diffusion than
Evan's blue. (C) Weight curves of *Foxp3*-DTR mice injected with IT pegDT, IP DT or IT
vehicle every 3 days demonstrating a lack of weight loss after site-selective Treg
ablation. Arrows represent DT injections. (D and E) Representative images of spleen
sizes and spleen weights from mice in (C). (F) Survival curves of mice in (C). ns = not
significant, *p<0.05, **p<0.01,***p<0.001.

939
940 **Figure S3. mTreg depletion selectively induces mechanical hypersensitivity in**
941 **female mice.**
942 (A) von Frey, (B) Hargreaves, (C) hotplate, (D) tail flick, (E) acetone, (F) pinprick, (G)
943 brush and (G) rotarod behavioral testing in *FoxP3*-DTR mice injected with a single dose
944 of IT pegDT. n = 4-20 mice per group. ns = not significant,****p<0.0001.

945
Figure S4. mTreg expansion selectively improves mechanical hypersensitivity in
injured female mice.
(A) mTreg expansion using IL-2 IT injections induces no changes in nociceptive
thresholds in uninjured naive male and female mice. mTreg expansion in nerve-injured
mice induces no changes in (B) acetone, (C) Hargreaves and (D) hotplate behavioral
testing. ns = not significant.

952
Figure S5. Functional characterization of enkephalin from CD4$^+$ T cells.
(A) Chimerism of meningeal Tregs and spinal cord microglia in PenkDTR$^{\Delta heme}$ mice and
(B) pooled chimerism comparing Tregs in the meninges and in the lymphoid organs
compared to tissue macrophages of the spinal cord (microglia) and the epidermis
(Langerhans cells, LC). (C) mTreg number after IP DT injection in PenkDTR$^{\Delta heme}$ mice.
(D) Weight curves of PenkDTR$^{\Delta heme}$ mice injected with IP DT every three days showing
peripheral penk ablation doesn't induce weight loss. (E) Unaltered spleen weight and
(F) meningeal and (G) spleen CD4$^+$ T cell populations after IP DT. Unaltered CD4 T cell
populations in the (H) spleen and (I) in the nerve and unaltered (J) mouse weight and
(K) spleen weight. (L) Graft Versus Host Disease (GVHD) score in *Rag2$^{-/-}$* mice injected
with pre-activated Tconv alone or with *Penk$^{+/+}$* or *Penk$^{-/-}$* Tregs, n=3-4 mice per group.
(M) IFN-□$^+$ and (N) IL-17$^+$ CD4$^+$ T cells after GVHD induction. ns = not significant,
*p<0.05, ***p<0.001.

References

1. Mogil, J.S. (2012). Sex differences in pain and pain inhibition: multiple explanations of a controversial phenomenon. Nat. Rev. Neurosci. *13*, 859–866. 10.1038/nrn3360.

2. Aloisi, A.M., Bachiocco, V., Costantino, A., Stefani, R., Ceccarelli, I., Bertaccini, A., and Meriggiola, M.C. (2007). Cross-sex hormone administration changes pain in transsexual women and men. Pain *132 Suppl 1*, S60–S67. 10.1016/j.pain.2007.02.006.

3. Petrovski, B.É., Vetvik, K.G., Lundqvist, C., and Eberhard-Gran, M. (2018). Characteristics of menstrual versus non-menstrual migraine during pregnancy: a longitudinal population-based study. J. Headache Pain *19*, 27. 10.1186/s10194-018-0853-3.

4. Negro, A., Delaruelle, Z., Ivanova, T.A., Khan, S., Ornello, R., Raffaelli, B., Terrin, A., Reuter, U., Mitsikostas, D.D., and European Headache Federation School of Advanced Studies (EHF-SAS) (2017). Headache and pregnancy: a systematic review. J. Headache Pain *18*, 106. 10.1186/s10194-017-0816-0.

5. Melhado, E.M., Maciel, J.A., and Guerreiro, C.A.M. (2007). Headache during gestation: evaluation of 1101 women. Can. J. Neurol. Sci. J. Can. Sci. Neurol. *34*, 187–192. 10.1017/s0317167100006028.

6. Gintzler, A.R. (1980). Endorphin-Mediated Increases in Pain Threshold During Pregnancy. Science *210*, 193–195. 10.1126/science.7414330.

7. Ray-Griffith, S.L., Wendel, M.P., Stowe, Z.N., and Magann, E.F. (2018). Chronic pain during pregnancy: a review of the literature. Int. J. Womens Health *10*, 153–164. 10.2147/IJWH.S151845.

8. Burzyn, D., Kuswanto, W., Kolodin, D., Shadrach, J.L., Cerletti, M., Jang, Y., Sefik, E., Tan, T.G., Wagers, A.J., Benoist, C., et al. (2013). A Special Population of Regulatory T Cells Potentiates Muscle Repair. Cell *155*, 1282–1295. 10.1016/j.cell.2013.10.054.

9. Astarita, J.L., Dominguez, C.X., Tan, C., Guillen, J., Pauli, M.L., Labastida, R., Valle, J., Kleinschek, M., Lyons, J., and Zarrin, A.A. (2023). Treg specialization and functions beyond immune suppression. Clin. Exp. Immunol. *211*, 176–183. 10.1093/cei/uxac123.

10. Samstein, R.M., Josefowicz, S.Z., Arvey, A., Treuting, P.M., and Rudensky, A.Y. (2012). Extrathymic generation of regulatory T cells in placental mammals mitigates maternal-fetal conflict. Cell *150*, 29–38. 10.1016/j.cell.2012.05.031.

11. Arpaia, N., Green, J.A., Moltedo, B., Arvey, A., Hemmers, S., Yuan, S., Treuting, P.M., and Rudensky, A.Y. (2015). A Distinct Function of Regulatory T Cells in Tissue Protection. Cell *162*, 1078–1089. 10.1016/j.cell.2015.08.021.

12. Ali, N., Zirak, B., Rodriguez, R.S., Pauli, M.L., Truong, H.-A., Lai, K., Ahn, R., Corbin, K., Lowe, M.M., Scharschmidt, T.C., et al. (2017). Regulatory T Cells in Skin Facilitate Epithelial Stem Cell Differentiation. Cell *169*, 1119-1129.e11. 10.1016/j.cell.2017.05.002.

13. Bapat, S.P., Myoung Suh, J., Fang, S., Liu, S., Zhang, Y., Cheng, A., Zhou, C., Liang, Y., LeBlanc, M., Liddle, C., et al. (2015). Depletion of fat-resident Treg cells prevents age-associated insulin resistance. Nature *528*, 137–141. 10.1038/nature16151.

14. Salvany-Celades, M., Van Der Zwan, A., Benner, M., Setrajcic-Dragos, V., Bougleux Gomes, H.A., Iyer, V., Norwitz, E.R., Strominger, J.L., and Tilburgs, T. (2019). Three Types of Functional Regulatory T Cells Control T Cell Responses at the Human Maternal-Fetal Interface. Cell Rep. *27*, 2537-2547.e5. 10.1016/j.celrep.2019.04.109.

15. Robertson, S.A., Care, A.S., and Moldenhauer, L.M. (2018). Regulatory T cells in embryo implantation and the immune response to pregnancy. J. Clin. Invest. *128*, 4224–4235. 10.1172/JCI122182.

16. Rowe, J.H., Ertelt, J.M., Xin, L., and Way, S.S. (2012). Pregnancy imprints regulatory memory that sustains anergy to fetal antigen. Nature *490*, 102–106. 10.1038/nature11462.

17. Davoli-Ferreira, M., de Lima, K.A., Fonseca, M.M., Guimarães, R.M., Gomes, F.I., Cavallini, M.C., Quadros, A.U., Kusuda, R., Cunha, F.Q., Alves-Filho, J.C., et al. (2020). Regulatory T cells counteract neuropathic pain through inhibition of the Th1 response at the site of peripheral nerve injury. Pain *161*, 1730–1743. 10.1097/j.pain.0000000000001879.

18. Kuhn, J.A., Vainchtein, I.D., Braz, J., Hamel, K., Bernstein, M., Craik, V., Dahlgren, M.W., Ortiz-Carpena, J., Molofsky, A.B., Molofsky, A.V., et al. (2021). Regulatory T-cells inhibit microglia-induced pain hypersensitivity in female mice. eLife *10*, e69056. 10.7554/eLife.69056.

19. Ito, M., Komai, K., Mise-Omata, S., Iizuka-Koga, M., Noguchi, Y., Kondo, T., Sakai, R., Matsuo, K., Nakayama, T., Yoshie, O., et al. (2019). Brain regulatory T cells suppress astrogliosis and potentiate neurological recovery. Nature *565*, 246–250. 10.1038/s41586-018-0824-5.

20. Dombrowski, Y., O'Hagan, T., Dittmer, M., Penalva, R., Mayoral, S.R., Bankhead, P., Fleville, S., Eleftheriadis, G., Zhao, C., Naughton, M., et al. (2017). Regulatory T cells promote myelin regeneration in the central nervous system. Nat. Neurosci. *20*, 674–680. 10.1038/nn.4528.

21. Scherrer, G., Imamachi, N., Cao, Y.-Q., Contet, C., Mennicken, F., O'Donnell, D., Kieffer, B.L., and Basbaum, A.I. (2009). Dissociation of the opioid receptor mechanisms that control mechanical and heat pain. Cell *137*, 1148–1159. 10.1016/j.cell.2009.04.019.

22. François, A., Low, S.A., Sypek, E.I., Christensen, A.J., Sotoudeh, C., Beier, K.T., Ramakrishnan, C., Ritola, K.D., Sharif-Naeini, R., Deisseroth, K., et al. (2017). A Brainstem-Spinal Cord Inhibitory Circuit for Mechanical Pain Modulation by GABA and Enkephalins. Neuron *93*, 822-839.e6. 10.1016/j.neuron.2017.01.008.

23. Wang, D., Tawfik, V.L., Corder, G., Low, S.A., François, A., Basbaum, A.I., and Scherrer, G. (2018). Functional Divergence of Delta and Mu Opioid Receptor Organization in CNS Pain Circuits. Neuron *98*, 90-108.e5. 10.1016/j.neuron.2018.03.002.

24. Sorge, R.E., Mapplebeck, J.C.S., Rosen, S., Beggs, S., Taves, S., Alexander, J.K., Martin, L.J., Austin, J.-S., Sotocinal, S.G., Chen, D., et al. (2015). Different immune cells mediate mechanical pain hypersensitivity in male and female mice. Nat. Neurosci. *18*, 1081–1083. 10.1038/nn.4053.

25. Othy, S., Jairaman, A., Dynes, J.L., Dong, T.X., Tune, C., Yeromin, A.V., Zavala, A., Akunwafo, C., Chen, F., Parker, I., et al. (2020). Regulatory T cells suppress Th17 cell Ca2+ signaling in the spinal cord during murine autoimmune neuroinflammation. Proc. Natl. Acad. Sci. *117*, 20088–20099. 10.1073/pnas.2006895117.

26. Maganin, A.G., Souza, G.R., Fonseca, M.D., Lopes, A.H., Guimarães, R.M., Dagostin, A., Cecilio, N.T., Mendes, A.S., Gonçalves, W.A., Silva, C.E.A., et al. (2022). Meningeal dendritic cells drive neuropathic pain through elevation of the kynurenine metabolic pathway in mice. J. Clin. Invest. *132*. 10.1172/JCI153805.

27. Anderson, K.G., Mayer-Barber, K., Sung, H., Beura, L., James, B.R., Taylor, J.J., Qunaj, L., Griffith, T.S., Vezys, V., Barber, D.L., et al. (2014). Intravascular staining for discrimination of vascular and tissue leukocytes. Nat. Protoc. *9*, 209–222. 10.1038/nprot.2014.005.

28. Pereira, M.M.A., Mahú, I., Seixas, E., Martinéz-Sánchez, N., Kubasova, N., Pirzgalska, R.M., Cohen, P., Dietrich, M.O., López, M., Bernardes, G.J.L., et al. (2017). A brain-sparing diphtheria toxin for chemical genetic ablation of peripheral cell lineages. Nat. Commun. *8*, 14967. 10.1038/ncomms14967.

29. Minarelli, J., Davis, E.L., Dickerson, A., Moore, W.C., Mejia, J.A., Gugala, Z., Olmsted-Davis, E.A., and Davis, A.R. (2019). Characterization of neuromas in peripheral nerves and their effects on heterotopic bone formation. Mol. Pain *15*, 1744806919838191. 10.1177/1744806919838191.

30. Xie, W., Strong, J.A., and Zhang, J.-M. (2017). Active Nerve Regeneration with Failed Target Reinnervation Drives Persistent Neuropathic Pain. eNeuro *4*. 10.1523/ENEURO.0008-17.2017.

31. Graßhoff, H., Comdühr, S., Monne, L.R., Müller, A., Lamprecht, P., Riemekasten, G., and Humrich, J.Y. (2021). Low-Dose IL-2 Therapy in Autoimmune and Rheumatic Diseases. Front. Immunol. *12*, 648408. 10.3389/fimmu.2021.648408.

32. Syrett, C.M., Paneru, B., Sandoval-Heglund, D., Wang, J., Banerjee, S., Sindhava, V., Behrens, E.M., Atchison, M., and Anguera, M.C. (2019). Altered X-chromosome inactivation in T cells may promote sex-biased autoimmune diseases. JCI Insight *4*. 10.1172/jci.insight.126751.

33. Pyfrom, S., Paneru, B., Knox, J.J., Cancro, M.P., Posso, S., Buckner, J.H., and Anguera, M.C. (2021). The dynamic epigenetic regulation of the inactive X chromosome in healthy human B cells is dysregulated in lupus patients. Proc. Natl. Acad. Sci. U. S. A. *118*, e2024624118. 10.1073/pnas.2024624118.

34. Jiwrajka, N., Toothacre, N.E., Beethem, Z.T., Sting, S., Forsyth, K.S., Dubin, A.H., Driscoll, A., Stohl, W., and Anguera, M.C. (2023). Impaired dynamic X-chromosome inactivation maintenance in T cells is a feature of spontaneous murine SLE that is exacerbated in female-biased models. J. Autoimmun. *139*, 103084. 10.1016/j.jaut.2023.103084.

35. Arnold, A.P. (2020). Four Core Genotypes and XY* mouse models: Update on impact on SABV research. Neurosci. Biobehav. Rev. *119*, 1–8. 10.1016/j.neubiorev.2020.09.021.

36. Arnold, A.P., and Chen, X. (2009). What does the "four core genotypes" mouse model tell us about sex differences in the brain and other tissues? Front. Neuroendocrinol. *30*, 1–9. 10.1016/j.yfrne.2008.11.001.

37. Shime, H., Odanaka, M., Tsuiji, M., Matoba, T., Imai, M., Yasumizu, Y., Uraki, R., Minohara, K., Watanabe, M., Bonito, A.J., et al. (2020). Proenkephalin+ regulatory T cells expanded by ultraviolet B exposure maintain skin homeostasis with a healing function. Proc. Natl. Acad. Sci. *117*, 20696–20705. 10.1073/pnas.2000372117.

38. Galván-Peña, S., Leon, J., Chowdhary, K., Michelson, D.A., Vijaykumar, B., Yang, L., Magnuson, A.M., Chen, F., Manickas-Hill, Z., Piechocka-Trocha, A., et al. (2021). Profound Treg perturbations correlate with COVID-19 severity. Proc. Natl. Acad. Sci. *118*, e2111315118. 10.1073/pnas.2111315118.

39. Sjaastad, L.E., Owen, D.L., Joo, S., Knutson, T.P., O'Connor, C.H., McCluskey, B., LaRue, R.S., Langlois, R.A., and Farrar, M.A. (2022). Influenza infection recruits distinct waves of regulatory T cells to the lung that limit lung resident IgA+ B cells. Preprint at bioRxiv, 10.1101/2022.09.19.508325 10.1101/2022.09.19.508325.

40. van der Veeken, J., Glasner, A., Zhong, Y., Hu, W., Wang, Z.-M., Bou-Puerto, R., Charbonnier, L.-M., Chatila, T.A., Leslie, C.S., and Rudensky, A.Y. (2020). The transcription factor Foxp3 shapes regulatory T cell identity by tuning the activity of trans-acting intermediaries. Immunity *53*, 971-984.e5. 10.1016/j.immuni.2020.10.010.

41. Gong, S., Zheng, C., Doughty, M.L., Losos, K., Didkovsky, N., Schambra, U.B., Nowak, N.J., Joyner, A., Leblanc, G., Hatten, M.E., et al. (2003). A gene expression atlas of the central nervous system based on bacterial artificial chromosomes. Nature *425*, 917–925. 10.1038/nature02033.

42. Baysoy, A., Seddu, K., Salloum, T., Dawson, C.A., Lee, J.J., Yang, L., Gal-oz, S., Ner-Gaon, H., Tellier, J., Millan, A., et al. (2023). The interweaved signatures of common-gamma-chain

cytokines across immunologic lineages. J. Exp. Med. *220*, e20222052.
10.1084/jem.20222052.

43. Zamir, N., Palkovits, M., Weber, E., Mezey, E., and Brownstein, M.J. (1984). A dynorphinergic pathway of Leu-enkephalin production in rat substantia nigra. Nature *307*, 643–645. 10.1038/307643a0.

44. Cugurra, A., Mamuladze, T., Rustenhoven, J., Dykstra, T., Beroshvili, G., Greenberg, Z.J., Baker, W., Papadopoulos, Z., Drieu, A., Blackburn, S., et al. (2021). Skull and vertebral bone marrow are myeloid cell reservoirs for the meninges and CNS parenchyma. Science *373*, eabf7844. 10.1126/science.abf7844.

45. Weisser, S.B., van Rooijen, N., and Sly, L.M. (2012). Depletion and Reconstitution of Macrophages in Mice. J. Vis. Exp. JoVE, 4105. 10.3791/4105.

46. Sankaranarayanan, I., Tavares-Ferreira, D., Mwirigi, J.M., Mejia, G.L., Burton, M.D., and Price, T.J. (2023). Inducible co-stimulatory molecule (ICOS) alleviates paclitaxel-induced neuropathic pain via an IL-10-mediated mechanism in female mice. J. Neuroinflammation *20*. 10.1186/s12974-023-02719-8.

47. Laumet, G., Bavencoffe, A., Edralin, J.D., Huo, X.-J., Walters, E.T., Dantzer, R., Heijnen, C.J., and Kavelaars, A. (2020). Interleukin-10 resolves pain hypersensitivity induced by cisplatin by reversing sensory neuron hyperexcitability. PAIN *161*, 2344. 10.1097/j.pain.0000000000001921.

48. Qi, L., Iskols, M., Shi, D., Reddy, P., Walker, C., Lezgiyeva, K., Voisin, T., Pawlak, M., Kuchroo, V.K., Chiu, I.M., et al. (2024). A mouse DRG genetic toolkit reveals morphological and physiological diversity of somatosensory neuron subtypes. Cell *187*, 1508-1526.e16. 10.1016/j.cell.2024.02.006.

49. Rosen, S.F., Ham, B., Drouin, S., Boachie, N., Chabot-Dore, A.-J., Austin, J.-S., Diatchenko, L., and Mogil, J.S. (2017). T-Cell Mediation of Pregnancy Analgesia Affecting Chronic Pain in Mice. J. Neurosci. Off. J. Soc. Neurosci. *37*, 9819–9827. 10.1523/JNEUROSCI.2053-17.2017.

50. Baron, S.A., and Gintzler, A.R. (1984). Pregnancy-induced analgesia: effects of adrenalectomy and glucocorticoid replacement. Brain Res. *321*, 341–346. 10.1016/0006-8993(84)90190-2.

51. Munoz-Suano, A., Kallikourdis, M., Sarris, M., and Betz, A.G. (2012). Regulatory T cells protect from autoimmune arthritis during pregnancy. J. Autoimmun. *38*, J103–J108. 10.1016/j.jaut.2011.09.007.

52. Barry, A.M., Zhao, N., Yang, X., Bennett, D.L., and Baskozos, G. (2023). Deep RNA-seq of male and female murine sensory neuron subtypes after nerve injury. PAIN *164*, 2196. 10.1097/j.pain.0000000000002934.

53. Midavaine, É., Côté, J., Marchand, S., and Sarret, P. (2021). Glial and neuroimmune cell choreography in sexually dimorphic pain signaling. Neurosci. Biobehav. Rev. *125*, 168–192. 10.1016/j.neubiorev.2021.01.023.

54. van Rooijen, D.E., Roelen, D.L., Verduijn, W., Haasnoot, G.W., Huygen, F.J.P.M., Perez, R.S.G.M., Claas, F.H.J., Marinus, J., van Hilten, J.J., and van den Maagdenberg, A.M.J.M. (2012). Genetic HLA Associations in Complex Regional Pain Syndrome With and Without Dystonia. J. Pain *13*, 784–789. 10.1016/j.jpain.2012.05.003.

55. Aubert, N., Purcarea, M., Fornier, M., Cagnet, L., Naturel, M., Casrouge, A., Dietrich, G., Dieu-Nosjean, M.-C., and Marodon, G. (2024). Enkephalin-mediated modulation of basal somatic sensitivity by regulatory T cells in mice. eLife *13*. 10.7554/eLife.91359.1.

56. R, B., Vl, T., D, W., A, F., C, S., Sa, S., P, C., C, B., C, C., K, S., et al. (2014). Delta opioid receptors presynaptically regulate cutaneous mechanosensory neuron input to the spinal cord dorsal horn. Neuron *81*. 10.1016/j.neuron.2014.01.044.

57. G, S., Ta, A., F, S., C, S., G, S.-J., G, G., G, M., and H, S. (1988). Naloxone does not alter the perception of pain induced by electrical and thermal stimulation of the skin in healthy humans. Pain *34*. 10.1016/0304-3959(88)90122-4.

58. A, A., M, H., Al, H., O, H., and A, B. (1999). Estrogen-induced alterations of spinal cord enkephalin gene expression. Pain *83*. 10.1016/s0304-3959(99)00109-8.

59. Adurthi, S., Kumar, M.M., Vinodkumar, H.S., Mukherjee, G., Krishnamurthy, H., Acharya, K.K., Bafna, U.D., Uma, D.K., Abhishekh, B., Krishna, S., et al. (2017). Oestrogen Receptor-α binds the FOXP3 promoter and modulates regulatory T-cell function in human cervical cancer. Sci. Rep. *7*, 17289. 10.1038/s41598-017-17102-w.

60. Nitsche, J.F., Schuller, A.G.P., King, M.A., Zengh, M., Pasternak, G.W., and Pintar, J.E. (2002). Genetic Dissociation of Opiate Tolerance and Physical Dependence in δ-Opioid Receptor-1 and Preproenkephalin Knock-Out Mice. J. Neurosci. *22*, 10906–10913. 10.1523/JNEUROSCI.22-24-10906.2002.

61. Pereira, M.M.A., Mahú, I., Seixas, E., Martinéz-Sánchez, N., Kubasova, N., Pirzgalska, R.M., Cohen, P., Dietrich, M.O., López, M., Bernardes, G.J.L., et al. (2017). A brain-sparing diphtheria toxin for chemical genetic ablation of peripheral cell lineages. Nat. Commun. *8*, 14967. 10.1038/ncomms14967.

62. Sorge, R.E., Martin, L.J., Isbester, K.A., Sotocinal, S.G., Rosen, S., Tuttle, A.H., Wieskopf, J.S., Acland, E.L., Dokova, A., Kadoura, B., et al. (2014). Olfactory exposure to males, including men, causes stress and related analgesia in rodents. Nat. Methods *11*, 629–632. 10.1038/nmeth.2935.

63. Dixon, W.J. (1965). The Up-and-Down Method for Small Samples. J. Am. Stat. Assoc. *60*, 967–978. 10.2307/2283398.

64. Chaplan, S.R., Bach, F.W., Pogrel, J.W., Chung, J.M., and Yaksh, T.L. (1994). Quantitative assessment of tactile allodynia in the rat paw. J. Neurosci. Methods *53*, 55–63. 10.1016/0165-0270(94)90144-9.

65. Shields, S.D., Eckert, W.A., and Basbaum, A.I. (2003). Spared nerve injury model of neuropathic pain in the mouse: a behavioral and anatomic analysis. J. Pain *4*, 465–470. 10.1067/S1526-5900(03)00781-8.

66. Park, Y.-G., Sohn, C.H., Chen, R., McCue, M., Yun, D.H., Drummond, G.T., Ku, T., Evans, N.B., Oak, H.C., Trieu, W., et al. (2018). Protection of tissue physicochemical properties using polyfunctional crosslinkers. Nat. Biotechnol. 10.1038/nbt.4281.

67. Borges da Silva, H., Wang, H., Qian, L.J., Hogquist, K.A., and Jameson, S.C. (2019). ARTC2.2/P2RX7 Signaling during Cell Isolation Distorts Function and Quantification of Tissue-Resident CD8+ T Cell and Invariant NKT Subsets. J. Immunol. *202*, 2153–2163. 10.4049/jimmunol.1801613.

68. Heinen, A.P., Wanke, F., Moos, S., Attig, S., Luche, H., Pal, P.P., Budisa, N., Fehling, H.J., Waisman, A., and Kurschus, F.C. (2014). Improved method to retain cytosolic reporter protein fluorescence while staining for nuclear proteins. Cytom. Part J. Int. Soc. Anal. Cytol. *85*, 621–627. 10.1002/cyto.a.22451.

69. Collison, L.W., and Vignali, D.A.A. (2011). In Vitro Treg Suppression Assays. Methods Mol. Biol. Clifton NJ *707*, 21–37. 10.1007/978-1-61737-979-6_2.

70. Schroeder, M.A., and DiPersio, J.F. (2011). Mouse models of graft-versus-host disease: advances and limitations. Dis. Model. Mech. *4*, 318–333. 10.1242/dmm.006668.

71. Hess, N.J., Hudson, A.W., Hematti, P., and Gumperz, J.E. (2020). Early T Cell Activation Metrics Predict Graft-versus-Host Disease in a Humanized Mouse Model of Hematopoietic Stem Cell Transplantation. J. Immunol. *205*, 272–281. 10.4049/jimmunol.2000054.

72. N, S., K, F., K, L., De, W., Am, K., and Dd, G. (2020). The emergence of transcriptional identity in somatosensory neurons. Nature *577*. 10.1038/s41586-019-1900-1.

A. CD45+iv^neg, Cre+, Cre−; FSC vs tdTomato; 7.44%

B. tdTomato− (Cre+) 3.57%; tdTomato+ (Cre+) 32.8%; CD4 vs Foxp3

C. Irradiated host WT

Donors:
D-G: *Penk*^Cre;*Rosa26*^DTR
F: *Penk*^+/+ or −/−
G-H: *Foxp3*-DTR + *Penk*^−/− (1:1)

D. ♀ Nociceptive threshold (g) vs Time (days): *, **, ns

E. ♂ Nociceptive threshold (g) vs Time (days): ns, ns, ns

F. SNI Nociceptive response (%) ♀ ***

G. SNI Nociceptive response (%) ♂ ns

○ *Penk*DTR^Δheme + Veh IT   ● *Penk*DTR^Δheme + pegDT IT
Depletion of enkephalinergic hematopoietic cells

H. Nociceptive threshold (g) ♀ ***; ● WT^Δheme ○ *Penk*^Δheme
hematopoietic enkephalin deficiency

I. Nociceptive threshold (g) vs Time (days): *, **; ○ Control^ΔmTreg ● *Penk*^ΔmTreg ♀
mTreg specific enkephalin deficiency

J. SNI Nociceptive response (%) **

K. Nociceptive threshold (g); IL-2 + Veh ** ; IL-2 + δOR ant ns; ♀ WT SNI POD30 + IL-2 + Veh / δOR antagonist

L. IL-2 anti-nociceptive efficacy (%) *; ○ Veh ● δOR ant