1 **Plant phenotypes as trajectories: 38-yr monitoring reveals that shape of lifetime**

2 **fecundity schedule is under selection in a long-lived shrub**

4 Carlos M. Herrera [1,2]

5 [1] *Estación Biológica de Doñana, Consejo Superior de Investigaciones Científicas*

6 *(CSIC), Américo Vespucio 26, E-41092 Sevilla, Spain*

7 [2] e-mail: herrera@ebd.csic.es

10 Short title: Plant phenotypes as trajectories

13 Key words: demography; individual variation; *Lavandula latifolia*; life history; lifetime

14 fecundity schedule; plant phenotype; plant senescence; phenotypic selection.

Abstract

Defining a phenotype is sometimes problematic in the case of modularly-built,

nonunitary organisms with indeterminate growth such as plants. This paper presents a

proof of concept of the evolutionary significance of considering lifetime trajectories of

individual plants as a component of their phenotypes. Size and inflorescence production

were monitored for the whole reproductive lifespans of $N = 128$ individuals of the

Mediterranean woody shrub *Lavandula latifolia* (Lamiaceae) over a 38-year period to

address the following questions: Did individuals vary in lifetime trajectories of size and

fecundity?, and Were parameters describing individual trajectories significant predictors

of cumulative lifetime reproduction? Individuals differed widely in lifetime fecundity

and in every parameter describing lifetime fecundity schedule (age at first and last

reproduction, longevity, and mean, variance, skewness and kurtosis of lifetime temporal

distribution of inflorescence production). Phenotypic selection analysis revealed

significant relationships between parameters of individual lifetime schedules and a

surrogate of relative fitness of individuals (cumulative lifetime production of

inflorescences divided by the average for all individuals). Significant selection gradients

involved positive and negative directional selection, as well as instances of nonlinear

selection, which showed that plants with certain shapes of lifetime fecundity schedules

had fitness advantage over others. The notion of plant phenotypes as trajectories was

strongly supported by the combined findings that individuals differed greatly in their

"appearances" with regard to the way in which size and fecundity unfolded over

lifetime, and that selection efficaciously "saw" such variation.

Introduction

Defining the phenotype of an organism is central to understanding the links between selection and potential evolutionary change, and how selection "sees" and "filters out" the variants of individual appearance that commonly coexist in natural populations of the same species. As stressed by Lewontin (1974, p. 19), "it is the evolution of the phenotype that interests us … [because] … what are ultimately to be explained are the myriad and subtle changes in size, shape, behavior, and interactions with other species that constitute the real stuff of evolution". Selecting discrete or continuous measurable traits of individuals for obtaining a representation of their appearance is a trivial matter when one deals with unitary, non-modularly built organism with determinate growth such as most animals, but it is somewhat problematic in the case of modularly-built, nonunitary organisms with indeterminate growth such as most plants (Chitwood and Topp, 2015; Herrera, 2024). For example, defining the leaf or seed phenotype of an individual plant which can bear up to thousands of nonidentical copies of each of these kinds of homologous, reiterated structures poses a conceptual challenge, and it has been proposed that intraplant distribution in quantitative traits of reiterated structures should be considered a relevant dimension of its phenotype or appearance (Herrera, 2024).

One additional, insufficiently recognized difficulty to define plant phenotypes is that single-value traits of individuals (e.g., plant size), or features of the subindividually variable organs (e.g., leaf traits), can vary in the short term or over a plant's lifetime due to ontogenetic processes (Herrera, 2009; Kulbaba et al., 2017; Steppe et al., 2011; Harder et al., 2019). In organisms with long life cycles such as perennial plants phenotypic traits will change with age, and development can be highly variable among individuals (Coleman et al., 1995; Chitwood and Topp, 2015). This points to the need of somehow incorporating a temporal dimension into the description of individual plant

phenotypes for those traits that experience discernible ontogenetic changes over the

plant's lifetime (e.g., size, chemical, structural and functional traits of single leaves;

Goodger et al., 2006; Greenwood et al., 2008; Steppe et al., 2011; Ji et al., 2021).

Whenever individuals in a plant population differ in the properties of their lifetime

trajectories for a given phenotypic trait, the *shape* of the trajectory will in itself be an

element of the phenotype (my usage here of "phenotypes as trajectories" bears no

relationship to the concept of "phenotypic trajectories" used to depict evolutionary

changes of species or populations in phenotypic space; Adams and Collyer, 2009;

Billman et al., 2014).

This paper presents a proof of concept of the evolutionary significance of

considering lifetime trajectories as part of individual plants' phenotypes. The study is

based on data obtained by monitoring the growth and reproduction of a large sample of

individuals of the Mediterranean woody shrub *Lavandula latifolia* (Lamiaceae) over

their whole reproductive lifespans. The shape of lifetime reproductive schedule is taken

here as a particular instance of phenotypic trajectory. The following specific questions

will be addressed: (1) Did *L. latifolia* individuals differ substantially with regard to

descriptive parameters of their lifetime size and fecundity trajectories, so that it makes

sense to incorporate such trajectories into the description of their phenotypes ?; and (2)

If individual differences in lifetime trajectory did exist, were descriptive parameters of

individual trajectories significant predictors of cumulative lifetime reproduction, so that

the existence of phenotypic selection on lifetime fecundity schedules can be inferred ?

Results will provide affirmative answers to these two questions by showing that, when

monitored over their entire reproductive lifetimes, individuals varied widely in size and

fecundity trajectories, and that such variation was related to differential lifetime

fecundity. Although some results presented here are also relevant in the light of plant

demography and life history evolution (see Discussion), most of these aspects will not

be considered in detail, as the main purpose of this article is to highlight the conceptual

significance of considering the shape of lifetime trajectories as a phenotypic component

in long-lived plants.

Materials and methods

Study plant

Lavandula latifolia is an evergreen shrub inhabiting forest clearings and edges in mid-

elevation woodlands of the eastern Iberian Peninsula (see Fig. A1 in Appendix A of

Herrera and Jovani, 2010, and Fig. 1 in Alonso et al., 2018, for photographs). The

species reproduces exclusively by seeds, which are small (~1 mg) and lack special

mechanisms for dispersal. Plants have a main taproot and a short (usually <15 cm) main

woody stem, and can be accurately aged by counting growth rings. Branching is

dichasial, generally conforming to Leeuwenberg's development model (a sympodial

succession of equivalent sympodial units, each of which is orthotropic and determinate

in its growth; Hallé et al., 1978; Hallé 1986). This branching pattern leads to crowns of

adult plants being made up of distinct leaf clusters borne by short stems, many of which

produce one terminal inflorescence in early summer. Each of these leaf clusters

represents a "module" according to Hallé's (1986, p. 78) definition ("the leafy axis in

which the entire sequence of aerial differentiation is carried out, from the initiation of

the meristem that builds up the axis to the sexual differentiation of its apex"). This

produces an architecturally-mediated relationship between the number of inflorescences

produced by an individual shrub at a given season and the total number of actively

growing modules, as well as a close correlation between current leaf biomass and

annual inflorescence production by individual plants. In the population studied, yearly

seed production by adult plants was predicted by the number of inflorescences ($R^2 =$

0.44, Herrera, 1991), and individual variation in average per-seed viability (emergence

rate and seedling survival) was comparatively narrow in relation to the broad

differences in inflorescence production (Herrera, 2000). Annual inflorescence

production can therefore be used as a reasonable proxy for fecundity comparisons both

across individuals and across years within individuals. Further details on natural history,

reproductive biology and demography of *L. latifolia* in the study area can be found in

Herrera (1991), Herrera and Jovani (2010), Herrera and Bazaga (2016), Alonso et al.

(2018) and Herrera et al. (2021).

120   *Field methods*

This study was conducted during 1986-2023 at the "Aguaderillos-2" site of Herrera

(1988, 1991), located at 1220 m elevation in a mixed woodland of *Pinus nigra* and

Quercus rotundifolia in the Sierra de Cazorla, Jaén province, southeastern Spain (site

coordinates 37.9612ºN, 2.8847ºW). A permanent 200-m^2 plot was established there in

July 1986 which remained undisturbed for the whole duration of this study. In August

1986 all *L. latifolia* plants within the plot bearing ≥10 leaves (roughly ≥ 2 yr in age)

were mapped, tagged and measured (see below), and the number of current year's

inflorescences counted. Only the subset of plants which at the time of tagging in 1986

were nonreproductive juveniles and reproduced in two or more years over 1987-2023

will be considered here ($N = 128$). All these plants had already died by the summer of

2023, hence the data used here encompass their complete reproductive lifespans. All

plants were checked every summer (August-September) between 1987-2023. On each

occasion, individuals found dead were digged up, collected, and their age at the time of

death determined by ring counting. For each individual that remained alive, the number

of inflorescences produced in the current year was counted and, on the even years, their

size was assessed by measuring the two major axes of the canopy and then computing

the area of the horizontal projection by assimilating the shape to an ellipse. In senescent

plants with discontinuous canopies comprising two or more living sectors (see

photographs in Fig. A4, Appendix A of Herrera and Jovani, 2010, for examples),

measurements were taken and areas computed separately for each living sector, and

then summed up to obtain a single size estimate for the individual.

142   *Data analysis*

The birth year of every monitored individual was determined by substracting the age at

death from the year of death, and this information was then used to retrospectively

compute the age of every plant in each study year. These data were then merged with

those on yearly fecundity (number of inflorescences) and size measurements (estimated

canopy area) to obtain individual lifetime profiles of size and fecundity in relation to

current age. As will be shown in the results, size and fecundity lifetime profiles were

closely correlated, and only those for fecundity will be analyzed here in detail.

The lifetime fecundity schedule of each plant was characterized quantitatively by

combining two sets of parameters, hereafter named "landmark" and "shape" lifetime

parameters. The landmark parameters refer to individual life history attributes from the

classical "generalized triangular reproductive function" (Lewontin, 1965; Harper, 1977):

age at first reproduction, age at last reproduction, and age at death ("total lifespan"

hereafter). The shape parameters summarize numerically the individual trajectories by

taking advantage of the fact that, when complete lifetime data are used, individual

curves of yearly fecundity versus age actually represent frequency distributions of the

age at which reproductive units (inflorescences in the present instance) were produced.

The first four statistical moments of such frequency distributions (mean, variance,

skewness, kurtosis) were computed for each plant and provided the shape parameters of

lifetime fecundity schedules. The intensity of linear (directional, β_i) and quadratic

(stabilizing or disruptive, γ_i) selection gradients on lifetime fecundity schedules was

investigated by adopting a phenotypic selection approach, based on regressing total

inflorescences produced over individual lifetime against landmark and shape parameters

and their quadratic transforms (Lande and Arnold, 1983; Svensson, 2023).

All statistical analyses reported in this paper were carried out using the R

environment (R Core Team, 2023). Average population-level trajectories of plant size

and annual inflorescence production in relation to plant age were estimated by fitting

random-slopes, random-intercepts linear mixed-effect models to plant area and

inflorescence number data (both log-transformed), with age and age-squared as fixed-

effect predictors and individual plant as random factor. The population average for the

within-plant relationship over lifetime between plant size and inflorescence production

was evaluated by fitting a random-slopes, random-intercepts linear mixed-effect model

to the data with plant size as fixed-effect predictor and plant as random factor. The

function lmer in the package LME4 (Bates et al., 2015) was used for fitting mixed-effect

models, and predicted shapes of the relationships and associated confidence intervals

were obtained by applying to each fitted model the function ggpredict from the

GGEFFECTS package (Lüdecke, 2018). Marginal R^2 (variance explained by fixed factors)

and conditional R^2 (variance explained by both fixed and random factors) of mixed-

effect models (Nakagawa and Schielzeth, 2013) were obtained with function

r2_nakagawa in the PERFORMANCE package (Lüdecke et al., 2021).

Phenotypic selection on descriptive parameters of individual lifetime fecundity

schedule was assessed by fitting a linear model to the data using function lm in the

STATS package. For the estimation of selection coefficients (β_i, γ_i) and their standard

errors, the response variable was the cumulative production of inflorescences over each

individual's lifetime divided by the average of this value for all individuals in the

sample ("relative fitness", Lande and Arnold, 1983). The predictors were the three

landmark and four shape schedule parameters defined above plus their quadratic

transforms, all of which were included simultaneously in a single multiple regression

model. All variables in the model except the response were scaled to mean zero and

standard deviation unity, so that the estimated model parameters represented

standardized estimates (Lande and Arnold, 1983). Individual lifetime fecundity departed

substantially from normality, which invalidated asymptotical statistical significance

tests of parameters based on their standard errors from this model. To evaluate the

statistical significance of predictors, the model was re-run after transforming fecundity

values logarithmically, which satisfactorily normalized residuals, and significance of

parameter estimates (i.e., β_i or $\gamma_i \neq 0$) was then evaluated with ordinary F tests using the

Anova function in the car package (Fox and Weisberg, 2019).

199   **Results**

200   *Population-level relationships between size, fecundity and age*

After statistically accounting for individual heterogeneity by including plant as a

random factor in the model, there existed a close linear relationship at the population

level between yearly inflorescence production and current plant size (area alive) over an

individual plant's lifetime (Figure 1A; Chi-square = 316.14, df = 1, $P < $ 1e-06). The

model fit well the data, and the rough similarity between marginal (0.598) and

conditional (0.788) R^2 values suggests that individual variation in the size-fecundity

relationship was only a secondary source of variance in yearly inflorescence production.

208   As would be anticipated from the close relationship between size and fecundity over

individual plant lifetime, the model-estimated, population-level relationship linking

plant size (Figure 1B; marginal $R^2 = 0.353$, conditional $R^2 = 0.852$) and yearly

inflorescence production (Figure 1C; marginal $R^2 = 0.237$, conditional $R^2 = 0.820$) with

plant age were virtually identical. On average, both size and yearly inflorescence

production first increased steadily until plants reached ~15 years and then declined

steadily until death. The substantial difference between marginal and conditional R^2 in

each of these two models provides an indirect indication that the among-plant

differences subsumed into the random terms were substantial and contributed most

variance to the response variables in the models, as detailed in the following section.

Given the close similarity between lifetime patterns for size and inflorescence

production, only the latter will be considered hereafter.

220   *Individual lifetime fecundity schedules and parameters*

The lifetime reproductive schedules of all *L. latifolia* plants studied are plotted side by

side in Figure 2 to graphically highlight the extraordinary multivariate diversity of

individual lifetime fecundity trajectories. Individual schedules included, for example,

broad-flat ones with slowly increasing production of inflorescences over all the

reproductive lifespan; broad-weakly convex ones denoting long reproductive lifespans

with a shallow peak at some intermediate age; or narrow-convex shapes with

reproduction concentrated over short reproductive spans. Schedule diversity involved

variation in age at first reproduction (range = 2-17 years, interquartile range IQR = 5-7),

age at last reproduction (range = 6-42 years, IQR = 18-24), and total lifespan (range 10-

44 years, IQR = 20-26). Across plants, age at first reproduction was not significantly

correlated with either age at last reproduction or total lifespan (r_s = 0.106 and 0.144, *P* =

0.23 and 0.11, respectively, *N* = 128, Spearman rank correlation). The extensive

individual variation in shape of lifetime fecundity schedules (Figure 2) was also

reflected in the broad frequency distributions of the four shape parameters considered

(Figure 3).

The diversity of lifetime fecundity schedules represented in the set of monitored

plants is well synthesized by the broad two-dimensional scatter of the empirical

cumulative distributions of fecundity against age when data points for all individuals are

plotted together (Figure 4). The cumulative proportion of inflorescences produced up to

a given age, or the age required to achieve a given proportion of cumulative lifetime

fecundity, were both extremely variable among plants. For example, some individuals

had already produced 50% of their eventual lifetime inflorescences before being 10

years old, while others only achieved that cumulative fecundity threshold at ages ≥20.

At age 20 some plants had produced <25% of lifetime inflorescences while many had

already produced >75%.

246   *Phenotypic selection on lifetime fecundity schedule*

Individual plants differed widely in lifetime fecundity (range = 3-2810 inflorescences,

IQR = 86-406). After transformation to individual relative fitness, lifetime fecundity

was significantly predicted by the landmark and shape parameters of lifetime fecundity

schedules ($F_{14, 113} = 30.6$, $P < 2.2e-06$, adjusted $R^2 = 0.791$; multiple regression of log-

transformed relative fitness against standardized landmark and shape parameters). Each

descriptive parameter of the fecundity schedule was a significant predictor of lifetime

inflorescence production after statistically controlling for the combined effects of the

rest of parameters (Table 1). In the case of landmark parameters only the linear terms

had significant effects on lifetime fecundity: age at last reproduction was directly related

to lifetime fecundity, while age at first reproduction and total lifespan were both

inversely related. These relationships denote that plants starting to reproduce earlier in

life, discontinuing reproduction at a later age, and/or having shorter longevities tended

to be at relative fecundity advantage. In the case of shape parameters there were both

linear and quadratic significant effects. Lifetime schedule curves with lower variance

and kurtosis (i.e., narrower curves) were at relative fecundity advantage. Significant

quadratic effects had either positive (variance, skewness) or negative (mean) signs

(Table 1), revealing instances of stabilizing and disruptive selection, respectively.

264   **Discussion**

265   *Population average versus individual lifetime schedules*

Monitoring the whole reproductive lifespans of a large sample of *L. latifolia* plants of

known birth date has revealed a well-defined average lifetime fecundity schedule at the

population level. It involved a steep increase in size and reproduction up to reaching a

peak about midway in lifetime (~15 years), followed by a steep decline ending up in the

plant's death (~30-40 years). This two-stage lifetime pattern was parsimoniously

accounted for by a shift in the dynamic balance between module formation and module

death, in accordance with the notion of the plant individual as a population of semi-

autonomous modules (White, 1979; Watkinson and White, 1985). During the first phase

the number of living modules and inflorescences produced annually increased steadily

with age, while a senescence period (meant here as age-dependent decline in size and

reproduction) ensued during the second phase. In this period, inflorescence production

declined and living area shrank because yearly production of new modules did not

compensate for the increasing annual mortality of modules. The present results for *L.*

latifolia (see also Herrera and Jovani, 2010) contribute to the accumulating evidence

suggesting that gradual individual senescence is probably more frequent in natural

populations of perennial plants than previously thought (Silvertown et al., 2001;

Munné-Bosch, 2008, 2015; Roach and Smith, 2020).

The population average for the lifetime fecundity schedule was only weakly

representative of the broad range of individual schedules actually occurring in the

sample of *L. latifolia* plants studied, whose general shapes, temporal landmarks and

shape descriptors varied widely among individuals. The lifetime fecundity schedule

represents a keystone component in the demography of populations and the evolution of

life history traits (Lewontin, 1965; Harper, 1977; Stearns, 1992), yet I was unable to

locate any published research reporting individual lifetime fecundity schedules for

woody plants. Instances of broad individual variability in supra-annual patterns of fruit

production have been frequently reported for tropical and temperate woody plants (e.g.,

Janzen, 1978; Crawley and Long, 1995; Herrera, 1998; Munilla and Guitián, 2014,

Pérez-Ramos et al., 2014), but these data are poor indicators of individual lifetime

schedules because they ordinarily refer to plants of unknown age and comprise periods

much shorter than the presumed reproductive lifespan. It is not possible therefore to

know the extent to which the large individual variability found here for *L. latifolia* is a

general characteristic of individual lifetime schedules of woody plants.

The practical difficulties inherent to collecting the data needed for elucidating

individual fecundity schedules of long-lived plants have been emphasized many times

(Harper and White, 1974; Janzen, 1978; Clarke, 1992; Salguero-Gómez et al., 2015).

For example, the modal duration of studies included in the initial version of the

comprehensive COMPADRE database on plant demography was "4 years,

corresponding to the length of an average PhD project, as well as that of most funding

agencies" (Salguero-Gómez et al. 2015, p. 213). The modal duration of studies

subsequently declined to 3 years in the most recent version (COMPADRE 6.23.5.0,

accessed 8 March 2024), where only 0.85% of the studies included ($N = 8823$) were on

trees/shrubs and had a duration ≥25 years. Since 62% of the angiosperms are woody

species (Zanne et al., 2014), their underrepresentation in demographic studies speaks of

the practical difficulties noted above and denotes a considerable bias in the demographic

information available for plants. Statistical methods devised to circumvent limitations

imposed by unknown plant age and fragmentary fecundity data can successfully

estimate average lifetime schedules and other demographic parameters at the population

or species levels (Clarke, 1992; Silvertown et al., 2001; Jones et al., 2014; Wenk and

Falster, 2015; Caswell, 2019). These estimates are valid for comparative purposes

(Silvertown et al., 2001; Jones et al., 2014; Wenk and Falster, 2015; Qiu et al., 2021),

but the lack of empirical information on the magnitude of individual variation in

lifetime parameters prevents formal statistical testing of differences between species or

populations, since uncertainty around average estimates cannot be evaluated (van

Daalen and Caswell, 2017). Most importantly, average schedules for groups of

individuals do not allow for tests of the relationships between individual fitness and

features of lifetime fecundity schedules which have been historically hypothesized to

underly life history evolution (see references in Introduction).

Selection on lifetime fecundity schedule

The phenotypic selection analysis revealed strong, significant relationships between all

landmark and shape parameters of individual lifetime schedules and the surrogate used

here to estimate the relative fitness of *L. latifolia* individuals (cumulative lifetime

production of inflorescences divided by the average for all individuals). Selection

gradients on landmark parameters involved positive directional selection on age at last

reproduction, and negative directional selection on age at first reproduction and total

lifespan, which revealed that individuals combining early start of reproduction, long

reproductive lifespan, and short total lifespan were at relative fecundity advantage.

Selection on shape parameters involved both directional (negative on variance and

kurtosis) and quadratic (stabilizing on the mean, disruptive on variance and kurtosis)

components, again reflecting that everything else being equal certain shapes of lifetime

fecundity schedules had relative fitness advantage over others. Selection on the set of

descriptive parameters of lifetime schedules was consistently strong, as revealed by the

high absolute values of β_i (median $|\beta_i| = 1.44$, range = 0.22-1.66) and γ_i (median $|\gamma_i| =$

0.25, range = 0.03-0.85). These figures fall in the uppermost quantiles of the

distributions of (absolute) selection gradients previously reported for all types of traits

in natural or seminatural plant populations, including life history features (see

Kingsolver et al., 2001; Siepielski et al., 2009; Kingsolver and Diamond, 2011; Caruso

et al. 2018, 2020, for comprehensive reviews of phenotypic selection studies). It is not

possible to elucidate whether the higher selection gradients found here in relation to

most previous phenotypic selection studies stem from the type of traits considered, the

lifetime nature of the fitness surrogate data (lack of estimates of lifetime fitness is "a

major fault of most studies of natural selection"; Endler, 1986, p. 162), or some

combination of both.

The usual caveats raised on phenotypic selection approaches, particularly those

questioning inferences of causality drawn from trait-fitness relationships (Svensson,

2023), apply also to the present study, since it is not possible to identify the proximate

selective mechanism(s) accounting for the strong selection observed on landmark and

shape parameters of individual lifetime fecundity schedules. Furthermore, the

phenotypic selection analysis conducted here was based on a proxy of fecundity rather

than on actual fecundity data (i.e., number of offspring produced). Keeping these

caveats in mind, it is worth noting that results of this study are among the few instances

to date of empirical support to some central tenets of life history evolution theory in a

within-population context. Regarding landmark parameters, the strong negative

directional selection on age at first reproduction found here agrees with the selective

value consistently attached to reproductive precocity in models of life history evolution,

particularly in the case of species with colonizing, expanding populations (Cole, 1954;

Lewontin, 1965; Stearns, 1976, 1992; Harper, 1977) such as *L. latifolia* and, more

generally, Mediterranean woody plants whose populations are exposed to recurrent

wildfires (Guiote and Pausas, 2023). Positive directional selection on reproductive

lifespan also agrees with some assumptions of life history theory (Stearns, 1992). Shape

parameters of lifetime fecundity schedules have been rarely considered in theoretical

formulations of life history evolution, but selection on mean, variance and skewness

found here are implicit in some of the hypothesized scenarios envisaged by Lewontin

(1965) under his "generalized triangular reproductive function".

Phenotypes as trajectories

The significance of ontogenetic changes in long-lived plants has been previously

considered in a variety of ecological and evolutionary contexts, but there have been few

attempts at placing the phenomenon in the conceptual framework of an expanded

definition of plant phenotypes (but see, e.g., Harder et al., 2004; Herrera, 2009;

Chitwood and Topp, 2015; Kulbaba et al., 2017; Harder et al., 2019; Herrera et al.,

2021; Barton, 2024). The set of landmark and shape features used here to characterize

the lifetime schedules of growth and reproduction of individual *L. latifolia* plants

represent a description of the way in which individual genotypes unfolded over its

lifetime with regard to two important facets of its external (i.e., phenotypic) appearance,

namely plant size and annual number of inflorescences. The heterogeneous assortment

of shapes depicted in Figure 2 portrays an assembly of "visible and measurable

appearances" (Wanscher, 1975, p. 144) of monitored plants, and hence are genuine

elements of their phenotypes. Further support to the notion of phenotypes as trajectories

was provided by the finding that selection was somehow able to "see" the variation

existing among individuals, so that these individual differences were eventually related

to lifetime fecundity. The possible evolutionary significance of selection on phenotypic

trajectories will obviously depend on whether the causal mechanisms responsible for

individual differences in lifetime schedules have some heritable basis (van Daalen and

Caswell, 2017). Given the considerable evidence supporting a role of genetic and

(heritable) epigenetic control on individual development, growth and reproduction in

plants (Bogan et al, 2024; Mitchell⬚Olds, 1996a,b; Turck and Coupland, 2013; Kottler

et al., 2018; Herrera et al., 2019), and the considerable genetic and epigenetic diversity

exhibited by the marked population of *L. latifolia* studied (Herrera and Bazaga, 2016), it

seems plausible to speculate that observed individual differences in lifetime trajectories

were at least in part due to heritable factors. Unfortunately, the long time required for

estimating the heritability of lifetime schedule parameters using quantitative genetics

methods based on controlled crosses and parent-offspring correlations will in fact render

such heritability estimates unknowable for most woody plants.

While the concepts of gene and genotype have been a recurrent topic of discussion

throughout the history of genetics (Roll-Hansen, 2014), the description and use of the

closely associated concept of phenotype have remained remarkably stable and

undebated since its inception more than a century ago (Wanscher, 1975; Herrera, 2024).

There are reasons, however, for revising the concept of phenotype in the case of

modularly-built, nonunitary organisms with indeterminate growth such as plants and

some colonial animals (Chitwood and Topp, 2015; Herrera, 2024). On one side, there is

the need of accommodating instantaneous intraindividual variability in features of

reiterated organs into the concept of phenotype by treating such variation as trait

distributions rather than a simple average (Herrera, 2024). On the other hand, plants that

grow, develop and reproduce over long periods demand the incorporation of individual

lifetime trajectories as a critical temporal dimension of their respective phenotypes, as

motivated in the Introduction to this article and supported by the results presented.

Understanding the genotype-phenotype relationship was a major goal of the research

agenda in genetics and evolutionary biology during the last century (Sturtevant, 1965;

Lewontin, 1974; Falconer, 1981; Noble, 2015). This research tradition paid most

attention to unraveling the genetic underpinnings of phenotypic traits and the

mechanisms whereby genetic background accounted for the organisms' appearance. In

contrast, little effort has been historically devoted to scrutinize whether the original

Johannsenian definition of phenotype (Johannsen 1909, 1911) suited equally well to all

kinds of organisms or, alternatively, should in certain groups be superseded by a more

elaborate concept (Herrera, 2024, and present study). An updated definition of

phenotype should incorporate the ontogenetic, developmental and architectural

peculiarities of organisms, particularly the multiplicity of nonidentical structures

produced by single genotypes and the protracted period of steady phenotypic unfolding

arising from ontogenetic shifts, growth schedules and internal modular dynamics.

Acknowledgments

People helping with plot maintenance and plant monitoring over nearly four decades, or

providing ideas and discussion on this project, are too numerous to remember or

mention individually. Among them, special thanks are due to Manolo Carrión (†), Pedro

Jordano, Roger Jovani, Luis López-Soria and Dori Ramírez. I am indebted to Conchita

Alonso, Mónica Medrano and Juli Pausas for valuable suggestions which significantly

improved an earlier version of the manuscript. Consejería de Medio Ambiente, Junta de

Andalucía, provided permission to work in Sierra de Cazorla along with invaluable

facilities. While writing this article I benefitted from grants from Ministerio de Ciencia,

Innovación y Universidades (PID2022-141530NB-C22, DISTEPIC) and Generalitat de

Valencia (PROMETEO-2021/040, FocScales).

Data and Code Accessibility Statement

All the raw data and scripts required to replicate the analyses reported in this study will

be deposited in figshare upon manuscript acceptance.

Conflict of interest

439 The author declares he has no conflict of interest.

440 **ORCID**

441 Carlos M. Herrera https://orcid.org/0000-0003-2452-3888

Literature cited

Adams, D. C., and M. L. Collyer. 2009. A general framework for the analysis of

phenotypic trajectories in evolutionary studies. *Evolution* 63:1143–1154.

Alonso, C., R. Pérez, P. Bazaga, M. Medrano, and C. M. Herrera. 2018. Within-plant

variation in seed size and inflorescence fecundity is associated with epigenetic

mosaicism in the shrub *Lavandula latifolia* (Lamiaceae). *Annals of Botany*

121:153–160.

Barton, K. E. 2024. The ontogenetic dimension of plant functional ecology. *Functional*

Ecology 38:98–113.

Bates, D., M. Maechler, B. Bolker, and S. Walker. 2015. Fitting linear mixed-effects

models using lme4. *Journal of Statistical Software* 67:1–48.

Billman, E. J., J. E. Rasmussen, J. C. Creighton, J. B. Johnson, and M. C. Belk. 2014. A

multivariate approach to the analysis of within lifetime variation in life history.

Methods in Ecology and Evolution 5:797–805.

Bogan, S. N., and S. V. Yi. 2024. Potential role of DNA methylation as a driver of

plastic responses to the environment across cells, organisms, and populations.

Genome Biology and Evolution 16. Doi: 10.1093/gbe/evae022

Caruso, C. M., K. E. Eisen, R. A. Martin, and N. Sletvold. 2018. A meta-analysis of the

agents of selection on floral traits. *Evolution* 73:4–14.

Caruso, C. M., H. Maherali, and R. A. Martin. 2020. A meta-analysis of natural

selection on plant functional traits. *International Journal of Plant Sciences*

181:44–55.

Caswell, H. 2019. Sensitivity analysis: matrix methods in demography and ecology.

Springer, Cham, Switzerland.

Chitwood, D. H., and C. N. Topp. 2015. Revealing plant cryptotypes: defining

meaningful phenotypes among infinite traits. *Current Opinion in Plant Biology*

24:54–60.

Clarke, P. J. 1992. Predispersal mortality and fecundity in the grey mangrove

(*Avicennia marina*) in southeastern Australia. *Australian Journal of Ecology*

17:161–168.

Cole, L. C. 1954. The population consequences of life history phenomena. *Quarterly*

Review of Biology 29:103–137.

Coleman, J. S., K. D. M. McConnaughay, and D. D. Ackerly. 1994. Interpreting

phenotypic variation in plants. *Trends in Ecology and Evolution* 9:187–191.

Crawley, M. J., and C. R. Long. 1995. Alternate bearing, predator satiation and seedling

recruitment in *Quercus robur* L. *Journal of Ecology* 83:683–696.

Endler, J. A. 1986. Natural selection in the wild. Princeton University Press, Princeton,

New Jersey, USA.

Falconer, D. S. 1981. Introduction to quantitative genetics. Second edition. Longmans,

London, UK.

Fox, J., and S. Weisberg 2019. An R companion to applied regression. Third edition.

Sage, Thousand Oaks, California,USA.

Goodger, J. Q. D., R. M. Gleadow, and I. E. Woodrow. 2006. Growth cost and

ontogenetic expression patterns of defence in cyanogenic *Eucalyptus* spp. *Trees*

20:757–765.

Greenwood, M. S., M. H. Ward, M. E. Day, S. L. Adams, and B. J. Bond. 2008. Age-

related trends in red spruce foliar plasticity in relation to declining productivity.

Tree Physiology 28:225–232.

Guiote, C., and J. G. Pausas. 2023. Fire favors sexual precocity in a Mediterranean pine.

Oikos 2023:e09373.

Hallé, F. 1986. Modular growth in seed plants. *Philosophical Transactions of the Royal*

Society of London B, Biological Sciences 313:77–87.

Hallé, F., R. A. A. Oldeman, and P. B. Tomlinson. 1978. Tropical trees and forests. An

architectural analysis. Springer-Verlag, Berlin.

Harder, L. D., C. Y. Jordan, W. E. Gross, and M. B. Routley. 2004. Beyond

floricentrism: the pollination function of inflorescences. *Plant Species Biology*

19:137–148.

Harder, L. D., M. M. Strelin, I. C. Clocher, M. W. Kulbaba, and M. A. Aizen. 2019.

The dynamic mosaic phenotypes of flowering plants. *New Phytologist* 224:1021–

1034.

Harper, J. L. 1977. Population biology of plants. Academic Press, London, UK.

Harper, J. L., and J. White. 1974. The demography of plants. *Annual Review of Ecology*

and Systematics 5:419–463.

Herrera, C. M. 1988. Variation in mutualisms: the spatio-temporal mosaic of a

pollinator assemblage. *Biological Journal of the Linnean Society* 35:95–125.

Herrera, C. M. 1991. Dissecting factors responsible for individual variation in plant

fecundity. *Ecology* 72:1436–1448.

Herrera C. M. 1998. Population-level estimates of interannual variability in seed

production: What do they actually tell us? *Oikos* 82:612–616.

Herrera, C. M. 2000. Individual differences in progeny viability in *Lavandula latifolia*:

a long-term field study. *Ecology* 81:3036–3047.

Herrera, C. M. 2009. Multiplicity in unity. Plant subindividual variation and interactions

with animals. University of Chicago Press, Chicago, Illinois, USA.

Herrera, C. M. 2024. Plant phenotypes as distributions: Johannsen's beans revisited.

American Naturalist 203:219–229.

Herrera, C. M., and R. Jovani. 2010. Log-normal distribution of individual lifetime

fecundity: insights from a 23-yr study. *Ecology* 91:422–430.

Herrera, C. M., and P. Bazaga. 2016. Genetic and epigenetic divergence between

disturbed and undisturbed subpopulations of a Mediterranean shrub: a 20-year

field experiment. *Ecology and Evolution* 6:3832–3847.

Herrera, C. M., M. Medrano, R. Pérez, P. Bazaga, and C. Alonso. 2019. Within-plant

heterogeneity in fecundity and herbivory induced by localized DNA

hypomethylation in the perennial herb *Helleborus foetidus*. *American Journal of*

Botany 106:798–806.

Herrera, C. M., P. Bazaga, R. Pérez, and C. Alonso. 2021. Lifetime genealogical

divergence within plants leads to epigenetic mosaicism in the shrub *Lavandula*

latifolia (Lamiaceae). *New Phytologist* 231:2065–2076.

Janzen, D. H. 1978. Seeding patterns of tropical trees. Pages 83-128 in P. B. Tomlinson

and M. H. Zimmermann, editors. Tropical trees as living systems. Cambridge

University Press, Cambridge, UK.

Ji, M., G. Jin, and Z. Liu. 2021. Effects of ontogenetic stage and leaf age on leaf

functional traits and the relationships between traits in *Pinus koraiensis*. *Journal*

of Forestry Research 32:2459–2471.

Johannsen, W. 1909. Elemente der exakten Erblichkeitslehre. Gustav Fischer, Jena,

Germany.

Johannsen, W. 1911. The genotype conception of heredity. *American Naturalist*

45:129–159.

Jones, O. R., A. Scheuerlein, R. Salguero-Gomez, C. G. Camarda, R. Schaible, B. B.

Casper, J. P. Dahlgren, J. Ehrlén, M. B. García, E. S. Menges, P. F. Quintana-

Ascencio, H. Caswell, A. Baudisch, and J. W. Vaupel. 2014. Diversity of ageing

across the tree of life. *Nature* 505:169–173.

Kingsolver, J. G., H. E. Hoekstra, J. M. Hoekstra, D. Berrigan, S. N. Vignieri, C. E.

Hill, A. Hoang, P. Gibert, and P. Beerli. 2001. The strength of phenotypic

selection in natural populations. *American Naturalist* 157:245–261.

Kingsolver, J. G., and S. E. Diamond. 2011. Phenotypic selection in natural

populations: what limits directional selection? American Naturalist 177:346–357.

Kottler, E. J., A. VanWallendael, and S. J. Franks. 2018. Experimental treatment with a

hypomethylating agent alters life history traits and fitness in *Brassica rapa*.

Journal of Botany 2018:7836845.

Kulbaba, M. W., I. C. Clocher, and L. D. Harder. 2017. Inflorescence characteristics as

function valued traits: analysis of heritability and selection on architectural

effects. *Journal of Systematics and Evolution* 55:559–565.

Lande, R., and S. J. Arnold. 1983. The measurement of selection on correlated

characters. *Evolution* 37:1210–1226.

Lewontin, R. C. 1965. Selection for colonizing ability. Pages 77–94 in The genetics of

colonizing species, edited by H. G. Baker and G. L. Stebbins. Academic Press,

New York, New York, USA.

Lewontin, R. C. 1974. The genetic basis of evolutionary change. Columbia University

Press, New York, New York, USA.

Lüdecke, D. 2018. ggeffects: Tidy data frames of marginal effects from regression

models. *Journal of Open Source Software* 3:772.

Lüdecke, D., M. S. Ben-Shachar, I. Patil, P. Waggoner, and D. Makowski. 2021.

performance: An R package for assessment, comparison and testing of statistical

models. *Journal of Open Source Software* 6:3139.

Mitchell□Olds, T. 1996a. Genetic constraints on life□history evolution:

quantitative□trait loci influencing growth and flowering in *Arabidopsis thaliana*.

Evolution 50:140–145.

Mitchell□Olds, T. 1996b. Pleiotropy causes long□term genetic constraints on

life□history evolution in *Brassica rapa*. *Evolution* 50:1849–1858.

Munné-Bosch, S. 2008. Do perennials really senesce? *Trends in Plant Science* 13:216–

220.

Munné-Bosch, S. 2015. Senescence: is it universal or not? *Trends in Plant Science*

20:713–720.

Munilla, I., and J. Guitián. 2014. Long-term individual-level variation of reproductive

features in *Sorbus aucuparia*, a fleshy-fruited tree. *Trees* 28:1489–1496.

Nakagawa, S., and H. Schielzeth. 2013. A general and simple method for obtaining R^2

from generalized linear mixed-effects models. *Methods in Ecology and Evolution*

4:133–142.

Noble, D. 2015. Evolution beyond neo-Darwinism: a new conceptual framework.

Journal of Experimental Biology 218:7–13.

Pérez-Ramos, I. M., C. Aponte, L. V. García, C. M. Padilla-Díaz, and T. Marañón.

2014. Why is seed production so variable among individuals? A ten-year study

with oaks reveals the importance of soil environment. *PLoS ONE* 9:e115371.

Qiu, T., M.-C. Aravena, R. Andrus, D. Ascoli, Y. Bergeron, R. Berretti, M.

Bogdziewicz, T. Boivin, R. Bonal, T. Caignard, et al. 2021. Is there tree

senescence? The fecundity evidence. *Proceedings of the National Academy of*

Sciences USA 118:e2106130118.

R Core Team (2023). R: A language and environment for statistical computing. R

Foundation for Statistical Computing, Vienna, Austria.

Roach, D. A., and E. F. Smith. 2020. Life-history trade-offs and senescence in plants.

Functional Ecology 34:17–25.

Roll-Hansen, N. 2014. The holist tradition in twentieth century genetics. Wilhelm

Johannsen's genotype concept. *Journal of Physiology* 592:2431–2438.

Salguero-Gómez, R., O. R. Jones, C. R. Archer, Y. M. Buckley, J. Che-Castaldo, H.

Caswell, D. Hodgson, A. Scheuerlein, D. A. Conde, E. Brinks, et al. 2015. The

COMPADRE Plant Matrix Database: an open online repository for plant

demography. *Journal of Ecology* 103:202–218.

Siepielski, A. M., J. D. DiBattista, and S. M. Carlson. 2009. It's about time: the

temporal dynamics of phenotypic selection in the wild. *Ecology Letters* 12:1261–

1276.

Silvertown, J., M. Franco, and R. Perez-Ishiwara. 2001. Evolution of senescence in

iteroparous perennial plants. *Evolutionary Ecology Research* 3:393–412.

Stearns, S. C. 1976. Life-history tactics: a review of the ideas. *Quarterly Review of*

Biology 51:3–47.

Stearns, S. C. 1992. The evolution of life histories. Oxford University Press, Oxford,

UK.

Steppe, K., Ü. Niinemets, and R. O. Teskey. 2011. Tree size- and age-related changes in

leaf physiology and their influence on carbon gain. Pages 235-253 in F. C.

Meinzer, B. Lachenbruch, and T. E. Dawson, editors. Size- and age-related

changes in tree structure and function. Springer, Dordrecht, The Netherlands.

Sturtevant, A. H. 1965. A history of genetics. Harper and Row, New York, New York,

USA.

Svensson, E. I. 2023. Phenotypic selection in natural populations: what have we learned

in 40 years? *Evolution* 77:1493–1504.

Turck, F., and G. Coupland. 2013. Natural variation in epigenetic gene regulation and

its effects on plant developmental traits. *Evolution* 68:620–631.

van Daalen, S. F., and H. Caswell. 2017. Lifetime reproductive output: individual

stochasticity, variance, and sensitivity analysis. *Theoretical Ecology* 10:355–374.

Wanscher, J. H. 1975. The history of Wilhelm Johannsen's genetical terms and concepts

from the period 1903 to 1926. *Centaurus* 19:125–147.

Watkinson, A. R., and J. White. 1985. Some life-history consequences of modular

construction in plants. *Philosophical Transactions of the Royal Society of London,*

Series B 313:31–51.

Wenk, E. H., and D. S. Falster. 2015. Quantifying and understanding reproductive

allocation schedules in plants. *Ecology and Evolution* 5:5521–5538.

White, J. 1979. The plant as a metapopulation. *Annual Review of Ecology and*

Systematics 10:109–145.

Zanne, A. E., D. C. Tank, W. K. Cornwell, J. M. Eastman, S. A. Smith, R. G. FitzJohn,

D. J. McGlinn, B. C. O'Meara, A. T. Moles, P. B. Reich et al. 2014. Three keys to

the radiation of angiosperms into freezing environments. *Nature* 506:89–92.

Table 1. Standardized linear and quadratic selection gradients on the seven parameters used to describe the lifetime fecundity schedules of

Lavandula latifolia plants studied ($N = 128$). Statistical significant terms ($P < 0.05$) are shown in bold type.

	Terms in linear model					
	Linear			Quadratic		
Fecundity schedule parameter	β_i	SE	*P*-value	γ_i	SE	*P*-value
Landmark parameters:						
Age at first reproduction	**-1.439**	0.237	**2.8E-08**	+0.192	0.049	0.32
Age at last reproduction	**+1.442**	0.994	**4.2E-05**	+0.848	0.364	0.50
Total lifespan	**-0.512**	0.778	**0.012**	-0.253	0.370	0.53
Shape parameters:						
Mean	+1.525	0.485	0.11	**-0.665**	0.126	**0.024**
Variance	**-1.661**	0.469	**5.5E-05**	**+0.251**	0.098	**0.0051**
Skewness	+0.216	0.149	0.78	**+0.287**	0.115	**0.022**
Kurtosis	**-0.582**	0.226	**0.0059**	-0.029	0.048	0.18

Legends to figures

Figure 1. Model-predicted, average population-level estimates of the relationships

linking current plant size with yearly inflorescence production (A), and plant age with

size (B) and yearly inflorescence production (C), over the lifetime of individual plants.

Predictions were obtained by fitting random-slopes, random-intercepts linear mixed-

effect models to the data in which the variable on the vertical axis was the response

variable, the variable on the horizontal axis was the fixed-effect predictor (the quadratic

transform was also included in the case of B and C), and plant was included as random

factor. In B and C the response variable was log-transformed for the analyses, and

model predictions were back-transformed to the original scale of measurement. Grey

areas represent standard errors around estimates.

Figure 2. Lifetime reproductive schedules of individual *Lavandula latifolia* plants that

were monitored for their whole reproductive lifespans (only the $N = 119$ plants in the

sample that flowered on at least 5 years are shown). Each panel corresponds to a

different individual (identified by its field tag), and shows lifetime variation in yearly

inflorescence production (orange-filled area under thin black lines), smoothed lifetime

fecundity trajectory (thick orange line, fitted to yearly data using a gam smoother), and

total lifespan (horizontal blue bar). Plants are ordered by increasing lifetime fecundity

(orange area, from left to right, and top to bottom). Note logarithmic scale on vertical

axis.

Figure 3. Frequency distributions of the four shape parameters describing individual

lifetime fecundity curves in the set of *Lavandula latifolia* plants studied ($N = 128$).

Shape parameters are the four moments of the frequency distributions of the ages at

which lifetime inflorescences were produced by each plant (mean, variance, skewnes

and kurtosis).

Figure 4. Cumulative density distributions of the number of inflorescences produced

over lifetime by individual *Lavandula latifolia* plants. To avoid cluttering and highlight

the broad individual variability represented in the sample of plants studied, lines

connecting points for individual plants are omitted. For any given age, each point

corresponds to a different plant, and the vertical scatter illustrates individual variation in

the proportion of total lifetime inflorescences that were produced up to and including

that age. A small random noise has been added to the horizontal and vertical coordinates

of data points.

Figure 1

Figure 2

Figure 3

Figure 4