1

Intradental mechano-nociceptors serve as sentinels that prevent tooth damage

Elizabeth A. Ronan[1], Akash R. Gandhi[1], Brian S. C. Constantinescu[1], Karin H. Uchima Koecklin[1,2], Mak E. Guenther[1], Maximilian Nagel[3], Kaitlyn J. Blumberg[1], Tomer Stern[1], Peng Li[1,2], Joshua J. Emrick[1,*]

[1]Department of Biologic and Materials Sciences & Prosthodontics, School of Dentistry, University of Michigan, Ann Arbor, MI, USA.

[2]Life Sciences Institute, University of Michigan, Ann Arbor, MI, USA

[3]Sensory Cells and Circuits Section, National Center for Complementary and Integrative Health, Bethesda, MD, USA

*To whom correspondence should be addressed: jjemrick@umich.edu

ABSTRACT

Pain is the anticipated output of the trigeminal sensory neurons that innervate the tooth's vital interior[1,2]; however, the contribution of intradental neurons to healthy tooth sensation has yet to be defined. Here, we employ in vivo Ca^{2+} imaging to identify and define a population of myelinated high-threshold mechanoreceptors (intradental HTMRs) that detect superficial structural damage of the tooth and initiate jaw opening to protect teeth from damage. Intradental HTMRs remain inactive when direct forces are applied to the intact tooth but become responsive to forces when the structural integrity of the tooth is compromised, and the dentin or pulp is exposed. Their terminals collectively innervate the inner dentin through overlapping receptive fields, allowing them to monitor the superficial structures of the tooth. Indeed, intradental HTMRs detect superficial enamel damage and encode its degree, and their responses persist in the absence of either PIEZO2 or $Na_v1.8$[3,4]. Optogenetic activation of intradental HTMRs triggers a rapid, jaw opening reflex via contraction of the digastric muscle. Taken together, our data indicate that intradental HTMRs serve as sentinels that guard against mechanical threats to the tooth, and their activation results in physical tooth separation to minimize irreversible structural damage. Our work provides a new perspective on the role of intradental neurons as protective rather than exclusively pain-inducing and illustrates additional diversity in the functions of interoreceptors.

INTRODUCTION

Mammals instinctively use their teeth for mechanical behaviors including chewing to initiate digestion and biting for prey capture and self-defense. The protection and maintenance of enamel and teeth are essential when teeth do not regenerate as in humans. Loss of teeth is associated with poorer diet and nutritional status[5] as well as early mortality[6]. The somatosensory neurons within the trigeminal ganglion extend processes into the oral and

craniofacial tissues, including the inner tooth. Broadly, activation of primary sensory cells and

their downstream neuronal targets produces outputs including sensorimotor reflexes[7,8] and

perceptions of touch, temperature, body position, and pain. While we assume that the

activation of tooth innervation is the source of substantial pain[9], whether tooth-innervating

somatosensory neurons (hereafter intradental neurons) might contribute to the protection of

tooth structure in health has not been explored.

Historically, intradental innervation is thought to arise from both myelinated (Aβ/Aδ) and

unmyelinated (C-type) trigeminal sensory neurons. However, myelinated intradental neurons

predominate based on immunohistochemical labeling[10], and transcriptomic analyses indicating

the majority of intradental neurons express *S100b*, a marker of myelinated somatosensory

neurons[11,12]. While electrophysiological experiments have demonstrated that fibers with

conduction velocities in the Aδ range spike in extreme conditions when the inner tooth has

been exposed[13], the molecular identity of intradental neurons and their capacity to respond in

an intact tooth has not been explored. The combination of transcriptomic analyses with in vivo

functional imaging provides a new ability to relate molecular identity with cell function[7,11,14–18].

Indeed, these tools have revealed the basis for touch sensation in the skin[12,19], and provide

great potential for defining the role of intradental neurons.

Here, we develop a strategy to identify intradental neurons in the trigeminal ganglion

without damaging the tooth, then monitor their response profiles to mechanical stimulation. We

determined most responding neurons are large diameter and express transcriptomic markers

of myelinated mechano-nociceptors (i.e., *S100b*, *Scn10a*, and *Calca*). These intradental

neurons do not encode any direct forces applied to the intact outer tooth but are activated once

the dentin has been exposed. Intradental terminal endings extend into the inner dentin

providing an anatomical basis for their observed dentin-response profile. Importantly, we show

that intradental neurons represent a population of high-threshold mechanonociceptors

(HTMRs) that collectively respond to superficial damage to the tooth. Damage responses

persist in the absence of *Piezo2* further indicating intradental neurons represent HTMRs.

Furthermore, direct optogenetic activation of intradental HTMRs induces digastric muscle

contractions and evokes a protective jaw opening response that physically separates the

opposing teeth. This work provides a new perspective on intradental neuron function by

providing the sensory foundation for tooth protection, which involves the detection of superficial

damage and the reflexive protection of the tooth.

RESULTS

Identification of large diameter intradental neurons in vivo through electrical stimulation

of the intact tooth

To enable functional evaluation of intradental neurons, we adapted an in vivo trigeminal

Ca^{2+} imaging technique in mice to visualize real-time activity of sensory neurons[7,20] while

stimulating individual molars (Figure S1A-C). Past work in large mammals indicates that

electrical stimulation applied to exposed dentin can elicit single unit responses within the

alveolar nerve[21]. Furthermore, the electric pulp test is routinely used in clinical dentistry to

assess tooth sensation in human patients. Thus, we reasoned that low voltage pulses

delivered to the occlusal surface of individual intact molar teeth may identify corresponding

intradental neurons. Electrical stimulation of molar 1 or molar 2 (2-4 V, 0.6 Hz, 200 ms, Figure

1A, see Methods) resulted in stereotyped phasic, transient Ca^{2+} responses from separate

groups of trigeminal sensory neurons (Figure 1B-D, S1C) indicating discrete sensory inputs

from each tooth. In each experimental iteration, we identified pulsing neurons (9-21 neurons,

n=10, Figure 1E) on the surface of the ganglion (Figure S1C) with large diameters (46.4 ± 0.1

µm, n=99 neurons, Figure 1F) consistent with the upper range of somas of Aδ myelinated

sensory neurons. These neurons also responded to direct application of Endo-Ice to the tooth

surface, another clinical assay used to test human tooth pulp sensitivity[22] (Figure S1D). To

confirm that electrical stimulation preferentially activates intradental neurons, we performed

retrograde labeling (CTB-AF555) to ipsilateral mandibular molars 1 and 2[23] prior to Ca^{2+}

imaging. This allowed us to confirm that the majority of CTB+ neurons demonstrated pulsed

responses (81/111, 4 mice, Figure 1G-I). Taken together we have established an in vivo

method to identify intradental neurons within an intact tooth enabling molecular and functional

evaluation.

Intradental neurons express markers of putative HTMRs

We next sought to determine the molecular identity of electrically-activated intradental

neurons. Our prior retrograde labeling opens the possibility that intradental neurons represent

mechanoreceptors based on their transcriptomic classification[23]. To this end, we performed

Ca^{2+} imaging followed by post-hoc in situ hybridization of whole mount dissected trigeminal

ganglion[12]. We used a defined panel of probes (*S100b*, *Scn10a*, *Calca*, and *Mrgprd*) to

delineate mechanoreceptor subtypes[12]. We utilized an AAV-based strategy to induce

simultaneous GCaMP6f and mCherry expression in trigeminal sensory neurons. mCherry

expression provided reliable guideposts for image alignment (Figure S1H) to match in situ

hybridization staining (Figure S1I) with fluorescence responses from Ca^{2+} imaging. In line with

our prior work, we determined that most responding intradental neurons expressed *Scn10a,*

Calca, and *S100b* (31/47) suggesting these cells are nociceptive large diameter, myelinated

HTMRs[12] (Figure 1J). Conversely, intradental neurons lacked expression of *Mrgprd* (1/47), a

marker of unmyelinated C-type HTMRs[12]. Furthermore, 11/47 intradental neurons were

marked by only *S100b*, suggesting few are likely to represent low-threshold mechanoreceptors

(LTMRs). Together these results demonstrate that intradental neurons captured by our

paradigm primarily represent myelinated nociceptive HTMRs.

Intradental neurons respond to direct force after enamel removal

As intradental neurons express transcriptional markers of HTMRs, we next set out to

determine their responses to mechanical forces directed to the intact tooth. We first tested

whether intradental neurons respond to a range of noxious to innocuous forces applied to the

enamel surface (0.4 - 20 g. Interestingly, we found that none of the intradental neurons as

identified by electrical stimulation were activated by the mechanical forces tested (Figure 2A-

B). As pressure applied to exposed inner tooth elicits responses in humans in the clinical

setting[24], we reasoned that intradental neurons might detect these mechanical forces in

exposed teeth. To test this, we evaluated responses to application of force before and after

exposing the dental pulp. As expected, we found that intradental neurons were not activated

by the mechanical force on the intact tooth (Figure 2D). Interestingly, in the same tooth after

the dental pulp was exposed, a portion of intradental neurons now responded to mechanical

stimulation (9/54, 17% of electrical responders, Figure 2D, F). To our surprise, in teeth where

only the dentin was exposed, direct mechanical force now elicited responses from the majority

of intradental neurons (32/41, 78% of electrical responders, Figure 2E, F). From these

experiments, we conclude that intradental neurons are HTMRs (intradental HTMRs) that do

not contribute to any direct forces applied to the intact tooth but respond to mechanical force

after the removal of enamel.

Intradental HTMRs terminate in the tooth pulp and inner dentin

Our experiments testing direct force on tooth structures suggest the presence of

terminal endings within the dentin, as mechanical stimulation of exposed dentin recruits a large

number of intradental neurons. We applied genetic strategies relying on *S100b* and *Scn10a* to

drive fluorescent labeling of intradental HTMRs to visualize their terminal endings. While

S100b delineates myelinated somatosensory neurons, it is also broadly expressed in the

nervous system as well as non-neuronal peripheral cells. To restrict fluorescent labeling to

S100b+ somatosensory neurons we crossed *S100b*-Cre into a Cre-dependent neuronal

specific reporter strain (*Snap*25-LSL-2A-eGFP). We found dense GFP+ innervation of the

coronal pulp as well as parallel GFP+ terminals penetrating to the inner 3rd of the dentin

(Figure S2A-I). As *Scn10a* expression is highly restricted to sensory neurons, we repeated

immunohistochemical experiments in teeth obtained from *Scn10a*-Cre; Ai32 mice (*Scn10a*-

Cre; LSL-YFP). Here, we observed remarkably similar YFP+ innervation patterns within the

pulp and inner dentin (Figure S2J-O). While these studies indicate that dentin is uniformly

innervated by endings stemming from intradental neurons, the endings of individual neurons

were not yet discerned. To generate sparse labeling of terminal endings we turned to AAV-

based strategies that offer lower transduction efficiency[25] (Figure S3A-D). Here, AAV injections

into either *S100b*-Cre and *Scn10a*-Cre mice achieved labeling of few intradental neurons to

reveal terminal patterns within the dentin. Fluorescently-labeled fibers (*S100b*-tdT and *Scn10a*-

GFP) were localized to only a portion of the overall dentin typically within a single pulp horn

(Figure 2G, H) and exhibited tuft-like patterning penetrating into non-contiguous dentinal

tubules. These findings indicate that single neurons target several tubules and multiple

neurons are likely to innervate a local region of the inner dentin. With our sparse labeling, we

also noted that terminal morphology appeared as a series of puncta running in parallel along

the length of dentin tubules (Figure 2G, H). Taken together, we establish that intradental

HTMRs, as defined by expression of *S100b* and *Scn10a*, contribute intermingled terminals to

create dense innervation of the inner dentin.

Intradental HTMRs function to detect tooth enamel damage

Intradental HTMRs respond to mechanical stimulation following dentin exposure, providing input *after* the inner layers of the tooth have been exposed. We next sought to explore if these neurons also activate *while* molar teeth are damaged. To this end, we monitored responses of trigeminal sensory neurons while simultaneously damaging the enamel surface using a dental bur. We found that most intradental HTMRs responded to cutting of the surface enamel (64/70, 91%, n = 4). We next tested a range of vibration frequencies (50 - 200 Hz) applied to the tooth. Minimal activity was observed from intradental HTMRs (2/52, 3%, Figure S4B-E) indicating these cells detect enamel cutting. Taken together, these experiments indicate that intradental HTMRs encode damaging mechanical stimulation of individual teeth.

Intradental HTMRs detect superficial frictional damage of the tooth surface and encode damage severity

The cutting of the exterior tooth produced overt damage of the enamel layer (Figure S4A) and triggered a response from intradental HTMRs. Does the response profile of these neurons provide information about the degree of tooth damage? To this end we sought to introduce a stimulus that creates lesser damage by removing the carbide flutes from the dental bur. This "fluteless" bur produces friction, but not perceptible damage (Figure S4A). Remarkably, friction on the surface of the tooth triggered activation of almost half of the intradental HTMRs identified by electrical stimulation (98/205, 47%, n = 7 mice, Figure 3A, B). Subsequently, the cutting stimulus evoked responses in these 98 neurons in addition to recruiting responses from 58 additional intradental HTMRs (156/205, 76%, n = 7 mice, Figure 3C). Interestingly, friction triggered smaller magnitude responses compared to cutting,

suggesting that intensity of activation increases with degree of tooth damage (Figure 3H). Post

hoc in situ confirmed that intradental HTMRs that respond to tooth cutting were largely

Scn10a+ (22/27 neurons, 82%, n = 4 mice, Figure 3E). We found that cutting and friction

produced equivalent, minimal warming of the inner tooth (<3ºC, Figure S4F-H) that is unlikely

to trigger any response and cannot account for our observed differences between response

magnitude. These results demonstrate that intradental HTMRs detect superficial frictional

damage and encode the degree of damage.

Damage responses of intradental HTMRs are independent of *Piezo2* or *Scn10a* (Na$_v$1.8)

Our previous work indicated that intradental neurons express the mechanosensitive ion

channel *Piezo2*[23] that underlies mechanosensation in the skin[12,26]. Is PIEZO2 required for

intradental HTMR activation by enamel cutting? To investigate this and circumvent lethality of

global *Piezo2* deletion, we examined neuron responses to cutting in mice with AAV-mediated

conditional knock out of *Piezo2* (Piezo2-cKO). To confirm recombination in sensory neurons

and functional loss of PIEZO2[12,17], we ensured Ca^{2+} responses to innocuous cheek brush were

lost while responses to cheek pinch were maintained (Figure S4I). However, even in the

context of Piezo2-cKO, we observed no significant change in the proportion of intradental

HTMRs responding to cutting (Figure 3F, H). These findings indicate that *Piezo2* is not

required for the neuronal detection of enamel damage, further supporting their identity as

intradental HTMRs.

Intradental HTMRs are marked by *Scn10a* expression, encoding the pore-forming alpha

subunit of Na$_v$1.8 which contributes to noxious mechanical pain[4]. We next sought to determine

whether *Scn10a* is required for intradental HTMR sensory transduction in response to enamel

damage. To this end, we examined neuronal responses in the well-established knock in/knock

out Scn10a-Cre driver strain[27]. As expected, *Scn10a*-Cre enabled capture of intradental

HTMRs with nearly equivalent counts and proportion of cutting responders compared to initial

unbiased AAV-mediated recombination (Figure S5A-C). However, in homozygotes where

Scn10a is replaced by Cre expression, we found no reduction in the proportion of intradental

HTMRs responding to enamel cutting compared to other strains (Figure 3G, H). These

experiments indicate that the *Scn10a* gene product ($Na_v1.8$) is not required for the transduction

of damage responses. Taken together, these experiments suggest a role for an unknown high-

threshold mechanical receptor in intradental HTMRs that detects superficial damage of the

teeth.

Activation of intradental HTMRs induces digastric muscle contraction and initiate

protective jaw opening

Considering teeth are at risk for damage during regular function, we next sought to

determine how activation of the intradental HTMRs might drive protective responses.

Somatosensory HTMRs are known to initiate reflexes to protect structures from damage[7].

Interestingly, low intensity electrical stimulation of teeth in larger mammals has been shown to

elicit electromyogram (EMG) activity in some muscles of mastication[28]. Indeed, we first

confirmed that electrical stimulation of the inferior alveolar nerve (IAN) elicited EMG activity in

the anterior digastric muscle (Figure S6A) which contributes to jaw opening (i.e., mandibular

deflection). We next sought to determine if selective activation of intradental neurons was

capable of inducing EMG activity in the anterior digastric muscle. To this end, we relied on

AAV-Cre transduction of intradental neurons via cavitations in the molar teeth to initiate the

expression of channelrhodopsin-2 (ChR2) in a Cre-dependent effector strain (Ai32, Figure 4A,

S6B). Activation of intradental neurons expressing ChR2+ using blue light stimulation (15 ms

pulse) of the trigeminal ganglion produced an immediate, transient waveform in the ipsilateral

anterior digastric EMG (Figure 4B-C, S6C, E-G; 0.97 +/- 0.3 mV amplitude). Alternatively, we

found no EMG activity in response to blue light stimulation when intradental neurons were

ChR2-. The ipsilateral masseter muscle that contributes to closing the jaw also failed to

demonstrate EMG activity after light stimulation in either condition (Figure 4B, S6E). Prolonged

blue light stimulation (100 ms pulse) elicited similar waveforms in ipsilateral digastric EMG

(Figure S6D-G; p>0.05). Of note, changes in EMG activity were never observed at light offset

(Figure 4B, S6C,D). Taken together, these data indicate that somatic activation of intradental

neurons leads to electrical activity of the anterior digastric muscle.

The anterior digastric muscles function to open the jaw by depressing the mandible[29].

To directly determine if collective activation of intradental HTMRs initiates overt jaw movement,

we next employed a genetic strategy to drive ChR2 expression in these cells using *Scn10a*-

Cre. For these experiments, we selectively activated intradental HTMRs by directing our

optical fiber onto a single intact molar tooth. Here, we found that pulses of blue light (470 nm,

1s) elicited a rapid, brief mandibular deflection (10 ms onset, reversing after 30 ms, Figure 4E).

Paralleling EMG activity, light offset had no effect on jaw position. We determined that

repeated blue light pulses (10 pulses, 0.5 Hz, 1s) induced a series of mandibular deflections

entrained to light onset (Figure 4F) that did not diminish between pulse trains (Figure 4F-G,

S6H), average deflection 305 +/- 37 vs 289 +/- 49 µm, p>0.05). Conversely, green light (545

nm) failed to produce any jaw movements despite identical stimulating frequency and duration

(10 pulses, 0.5 Hz, 1s). Importantly, blue light stimulation of molar-adjacent tissues (i.e. hard

palate or oral vestibule) also failed to produce any jaw movements (Figure 4F). Taken

together, these experiments reveal that activation of intradental HTMRs drives a rapid jaw

opening that physically disengages the teeth from an insult.

DISCUSSION

Mammalian teeth are specialized internal organs that endure a range of

mechanosensory stimuli during biting and mastication. While intradental neurons are known to

evoke pain through their activation, a biological role has not been defined for this interoceptive

innervation using modern approaches. Here, we establish an in vivo functional imaging

approach to identify intradental neurons in an intact tooth, define their responses to

mechanical damage of the tooth, and determine that their targeted activation leads to jaw

opening. Our experiments reveal a uniform population of intradental HTMRs that 1) detect

forces applied to the exposed dentin as well as damage of the superficial enamel, 2) do not

contribute to the detection of innocuous forces encountered by the tooth, 3) target the inner

dentin with intermingled terminals providing anatomical evidence supporting their coordinated

response to stimuli, 4) rely on a Piezo2-independent mechanism to transduce damage

responses, and 5) drive digastric muscle contraction and a protective jaw opening reflex.

Protective reflexes have been described for the knee (patellar reflex), cornea (eye-blink

reflex) and the jaw (jaw-jerk reflex) that occur within 100 ms latencies. Here, we describe an

extremely rapid reflex (5-15 ms) that is initiated following the activation of terminals of

intradental HTMRs within the dentin. We predict that the intradental HTMRs respond when the

tooth makes unexpected contact with hard foods (i.e., cartilage or bone) or opposing teeth.

Based on our data, activation of intradental HTMRs within a single tooth during mastication

would be sufficient to drive a reflex response that would halt jaw closure. Intradental HTMRs

efficiently respond to damage of the tooth indicating they may be tuned for its detection;

however, the precise component of the damage stimulus that triggers intradental HTMR

responses will be difficult to determine considering the miniature size of the mouse tooth. We

expect that comparing structural changes in the tooth caused by vibration and cutting could

reveal the activating component.

Recent work indicates that *Calca*+ circumferential-HTMRs (circ-HTMRs) innervating hair

follicles in skin mediate a nocifensive response[7]. Intradental HTMRs also represent a

population of *Calca*/CGRP+ population of somatosensory neurons that mediate a reflexive,

protective response. While CGRP+ neurons have been further subdivided within the skin

based on their expression patterns, none feature terminal morphology exhibited by the

intradental HTMRs[19]. Functionally, intradental HTMRs share the response profile of some

subsets of CGRP+ HTMRs including *Smr2*+ and *Bmpr1b*+ somatosensory neurons. We

anticipate that intradental HTMRs may diverge in their thermal response profiles considering

their position within the insulated tooth structure. It is yet to be determined if intradental

HTMRs represent a new subclass of CGRP+ HTMRs that innervate and protect other

vulnerable tissue structures by engages withdrawal reflexes.

Intradental neurons have been previously suggested to function as LTMRs[30] which is

consistent with their expression of Piezo2. On the contrary, our data directly shows that

intradental neurons fail to respond to direct forces or vibration applied to the intact tooth. It is

highly probable that light forces applied to the teeth are transduced by LTMRs innervating

surrounding periodontal (i.e., tooth surrounding) tissues. Future studies investigating the

molecular identity and innervation targets within surrounding periodontium tissues will shed

insight into how direct mechanical forces applied to the tooth are sensed.

We determined that damage responses by intradental neurons are independent of the

canonical mechanoreceptor PIEZO2 as well as a nociceptive transducer $Na_v1.8$. Our data

suggest that blocking either ion channel is unlikely to prevent activation of the intradental

neurons as they are structurally damaged. This is in contrast to a recent clinical trial that

supports the effectiveness of inhibition of $Na_v1.8$ to diminish post-surgical inflammatory pain[31].

Instead, we propose the existence of unidentified molecular sensors for high threshold

mechanical stimulation of the tooth. Single-cell sequencing of intradental neurons[32] toward

screening and identification of novel high-threshold mechanoreceptors and nociceptive

transducers will provide additional targets for next-generation dental anesthetics that can be

validated with our approach.

While some evidence suggests that the tooth pulp receives innervation from C-fiber

nociceptors that contribute to forms of tooth-related pain, our work to-date suggests these are

unlikely to contribute to the protective reflex response. Our findings do not exclude the

possibility that C-fibers may be present deep in the tooth pulp where they could contribute to

differential aspects of tooth physiology such as in the context of inflammation. Clinically, we

predict intradental HTMRs may underlie acute pain during biting as part of cracked tooth

syndrome[33]. Further work will be needed to determine if responses of intradental HTMRs are

altered by the inflammatory milieu. For instance, intradental HTMRs may exhibit lower

threshold responses to mechanical stimulation in the context of tooth decay, and begin to

contribute to the exquisite pain associated with the colloquial toothache.

While our study indicates that intradental HTMRs respond to damage and initiate an

important protective reflex, others behavioral outputs resulting from their activation has yet to

be explored. We anticipate that excessive or persistent activation of intradental HTMRs may

be sufficient to elicit pain behaviors in mice, and could be the basis for defining additional

orofacial movements that indicate the experience of pain. Further experiments will be

necessary to understand if the jaw-opening reflex and pain are necessarily concurrent or may

be uncoupled in freely-behaving animals.

REFERENCES

1. Henry, M. A. & Hargreaves, K. M. Peripheral Mechanisms of Odontogenic Pain. *Dental Clinics of North America* **51**, 19–44 (2007).

2. Renton, T. Dental (Odontogenic) Pain. *Rev Pain* **5**, 2–7 (2011).

3. Murthy, S. E. *et al.* The mechanosensitive ion channel Piezo2 mediates sensitivity to mechanical pain in mice. *Sci Transl Med* **10**, eaat9897 (2018).

4. Akopian, A. N. *et al.* The tetrodotoxin-resistant sodium channel SNS has a specialized function in pain pathways. *Nat Neurosci* **2**, 541–548 (1999).

5. Xu, K.-H. *et al.* Association of Tooth Loss and Diet Quality with Acceleration of Aging: Evidence from NHANES. *Am J Med* **136**, 773-779.e4 (2023).

6. Koka, S. & Gupta, A. Association between missing tooth count and mortality: A systematic review. *J Prosthodont Res* **62**, 134–151 (2018).

7. Ghitani, N. *et al.* Specialized Mechanosensory Nociceptors Mediating Rapid Responses to Hair Pull. *Neuron* **95**, 944-954.e4 (2017).

8. Barik, A., Thompson, J. H., Seltzer, M., Ghitani, N. & Chesler, A. T. A Brainstem-Spinal Circuit Controlling Nocifensive Behavior. *Neuron* **100**, 1491-1503.e3 (2018).

9. Slade, G. D. Epidemiology of dental pain and dental caries among children and adolescents. *Community Dent Health* **18**, 219–227 (2001).

10. Byers, M. R. & Cornel, L. M. Multiple complex somatosensory systems in mature rat molars defined by immunohistochemistry. *Archives of Oral Biology* **85**, 84–97 (2018).

11. Nguyen, M. Q., Wu, Y., Bonilla, L. S., von Buchholtz, L. J. & Ryba, N. J. P. Diversity amongst trigeminal neurons revealed by high throughput single cell sequencing. *PLoS ONE* **12**, e0185543 (2017).

12. von Buchholtz, L. J. *et al.* Decoding Cellular Mechanisms for Mechanosensory

Discrimination. *Neuron* **109**, 285-298.e5 (2021).

13. Närhi, M. V., Hirvonen, T. J. & Hakumäki, M. O. Activation of intradental nerves in the dog

to some stimuli applied to the dentine. *Arch Oral Biol* **27**, 1053–1058 (1982).

14. von Buchholtz, L. J., Lam, R. M., Emrick, J. J., Chesler, A. T. & Ryba, N. J. P. Assigning

transcriptomic class in the trigeminal ganglion using multiplex in situ hybridization and

machine learning. *Pain* **161**, 2212–2224 (2020).

15. Sharma, N. *et al.* The emergence of transcriptional identity in somatosensory neurons.

Nature **577**, 392–398 (2020).

16. Usoskin, D. *et al.* Unbiased classification of sensory neuron types by large-scale single-cell

RNA sequencing. *Nat Neurosci* **18**, 145–153 (2015).

17. Servin-Vences, M. R. *et al.* PIEZO2 in somatosensory neurons controls gastrointestinal

transit. *Cell* **186**, 3386-3399.e15 (2023).

18. Lam, R. M. *et al.* PIEZO2 and perineal mechanosensation are essential for sexual function.

Science **381**, 906–910 (2023).

19. Qi, L. *et al.* A DRG genetic toolkit reveals molecular, morphological, and functional diversity

of somatosensory neuron subtypes. 2023.04.22.537932 Preprint at

https://doi.org/10.1101/2023.04.22.537932 (2023).

20. Yarmolinsky, D. A. *et al.* Coding and Plasticity in the Mammalian Thermosensory System.

Neuron **92**, 1079–1092 (2016).

21. Virtanen, A., Närhi, M., Huopaniemi, T. & Hirvonen, T. Thresholds of intradental A-and C-

nerve fibres in the cat to electrical current pulses of different duration. *Acta Physiologica*

Scandinavica **119**, 393–398 (1983).

22. Chen, E. & Abbott, P. V. Dental Pulp Testing: A Review. *Int J Dent* **2009**, 365785 (2009).

23. Emrick, J. J., von Buchholtz, L. J. & Ryba, N. J. P. Transcriptomic Classification of Neurons

Innervating Teeth. *J Dent Res* **99**, 1478–1485 (2020).

24. Wanachantararak, S., Ajcharanukul, O., Vongsavan, N. & Matthews, B. Effect of cavity

depth on dentine sensitivity in man. *Arch Oral Biol* **66**, 120–128 (2016).

25. Byers, M. R. & Calkins, D. F. Trigeminal sensory nerve patterns in dentine and their

responses to attrition in rat molars. *Arch Oral Biol* **129**, 105197 (2021).

26. Szczot, M. *et al.* PIEZO2 mediates injury-induced tactile pain in mice and humans. *Sci*

Transl Med **10**, eaat9892 (2018).

27. Stirling, C. L. *et al.* Nociceptor-specific gene deletion using heterozygous NaV1.8-Cre

recombinase mice. *PAIN* **113**, 27 (2005).

28. Närhi, M., Virtanen, A., Hirvonen, T. & Huopaniemi, T. Comparison of electrical thresholds

of intradental nerves and jaw-opening reflex in the cat. *Acta Physiol Scand* **119**, 399–403

(1983).

29. Perry, S. K. & Emrick, J. Trigeminal somatosensation in the temporomandibular joint and

associated disorders. *Front. Pain Res.* **5**, (2024).

30. Dong, W. K., Chudler, E. H. & Martin, R. F. Physiological properties of intradental

mechanoreceptors. *Brain Research* **334**, 389–395 (1985).

31. Jones, J. *et al.* Selective Inhibition of NaV1.8 with VX-548 for Acute Pain. *N Engl J Med*

389, 393–405 (2023).

32. Lee, P. R. *et al.* Transcriptional profiling of dental sensory and proprioceptive trigeminal

neurons using single-cell RNA sequencing. *Int J Oral Sci* **15**, 1–14 (2023).

33. Lynch, C. D. & McConnell, R. J. The cracked tooth syndrome. *J Can Dent Assoc* **68**, 470–

475 (2002).

ACKNOWLEDGEMENTS

This work is supported by K22DE029779 (J.J.E.) and T32DE007057 (E.A.R.). We thank Dr.

Nicholas Ryba for his feedback throughout this project and valuable comments on this

manuscript. We thank Dr. Nima Ghitani for his technical assistance with in vivo imaging. We

thank Dr. Alexander Chesler, Dr. Lauren Surface, and members of the Emrick lab for helpful

comments and suggestions. We are grateful to donors of knockin mice for *S100b*-Cre knockin

mice (Ryba); *Piezo2*-cKO knockin mice (Chesler); and *Snap25*-LSL-2A-EGFP-D knockin mice

(Dr. Claire LePichon).

AUTHOR CONTRIBUTIONS

All authors gave final approval and agreed to be accountable for all aspects of the work. J.J.E.

and E.A.R. conceptualized and directed the study. E.A.R., A.R.G., and J.J.E. performed Ca^{2+}

imaging experiments with acquisition help from M.N. and B.S.C.C., and assistance in analysis

from B.S.C.C. and T.S.. E.A.R., M.E.G., A.R.G., B.S.C.C, K.J.B., and J.J.E. performed survival

surgeries and histological studies. K.H.U.K. performed optogenetic stimulation and EMG

measurements, and analyzed the data. E.A.R., B.S.C.C., and J.J.E. performed optogenetic

stimulation and mandibular deflection measurements. E.A.R., A.R.G., and J.J.E. interpreted

the data, wrote the manuscript and generated figures with input from all other authors.

DECLARATION OF INTERESTS

414 Authors declare no competing interests.

 FIGURES AND FIGURE LEGENDS

Figure 1. Functional imaging of intradental neurons reveals a large-diameter population

of putative myelinated HTMRs.

(A) Schematic of methodology used to record in vivo sensory neurons in the trigeminal

ganglion (TG) in response to stimulation of mouse molar tooth in the oral cavity. Intradental

neurons were identified via delivery of low voltage pulses (4V, 0.6 Hz, 200 ms) to individual

mandibular molars. Data in (A-E) were obtained from adult Ai95(RCL-GCaMP6f)-D (Ai95D)

mice injected postnatally (P0-P3) with AAV9-Cre.

(B-D) Voltage stimulation enabled identification of intradental neurons innervating a single molar. (B) Imaging field showing merged single frames of Ca^{2+} response ($\Delta F/F0$). Neurons with responses entrained to voltage pulses to either molar 1 (m1, red) or molar 2 (m2, cyan). Scale bar, 100µm. (C) Example raw traces of GCaMP6f signal from identified m1 (red) and m2 (cyan) intradental neurons. Numbers correspond to neurons circled in (B). Bars to the right of traces indicate 20 $\Delta F/F0$. (D) Example heatmap from single TG showing in vivo GCaMP6f responses of 18 intradental neurons to voltage pulses. Neurons were grouped according to molar target of stimulation (m1 or m2). Stimuli used are indicated above the heatmap. Traces in (C) are designated neurons shown in (D).

(E) Bar graph showing the mean number of neurons per trial that responded to ≥2 voltage pulses, designating them as intradental. Plotted individual data points represent the number of responding neurons in separate imaging trials. Error bars indicate the SEM. n = 10 mice.

(F) Bar graph showing the mean diameter of neurons that responded to ≥2 voltage pulses, designating them as intradental. Plotted individual data points represent the number of responding neurons in separate imaging trials. Bar shows the mean value, and error bars indicate the SEM. n = 99 neurons from 7 mice.

(G-I) Retrograde labeling marks intradental neurons that are activated by electrical stimulation of the molars. Retrograde tracer CTB-AF555 was applied to small cavitations in mandibular molars (m1/m2) 16 hr prior to functional imaging. (G) Sample images depicting overlap of CTB labeling with Ca^{2+} response ($\Delta F/F0$) of neurons responding to applied voltage pulses. Scale bar, 50µm. Left panel: CTB labeling; Middle panel: single frame ($\Delta F/F0$) from recording during response to voltage pulse; Right panel: Merge. (H) Example heatmap showing responses of combined m1/m2 CTB+ neurons to voltage pulses. (I) Bar graph showing the proportion of CTB+ TG neurons that respond to ≥2 voltage pulses. Plotted individual data points represent

the calculated proportion of CTB+ cells that respond to ≥2 voltage pulses for each ganglion.

Bar shows the mean and error bars indicate the SEM. n = 4 mice. CTB, cholera toxin B-

subunit. Data in (G-I) were obtained from *S100b*-Cre; Ai95D mice. See Figure S1E-G for

validation of *S100b*-Cre; Ai95D intradental neuron responses.

(J) Alignment of in vivo functional imaging responses with multiplexed whole-mount ISH

enables molecular classification of intradental neurons. Data for all experiments were obtained

from Ai95(RCL-GCaMP6f)-D (Ai95D) mice injected postnatally (P0-P3) with AAV9-hSyn1-

mCherry-2A-iCre, enabling mCherry expression to serve as guideposts to align functional

imaging with ISH images. (Left) Multiplexed whole-mount ISH staining following alignment to in

vivo GCaMP6f fluorescence. Probes: s100 calcium-binding protein B (*S100b*, blue), alpha-

Nav1.8 (*Scn10a*, green), calcitonin gene-related peptide (CGRP; *Calca*, red), mas-related G

protein-coupled receptor member D (*Mrgprd*, white). Plus sign in bottom right of each subpanel

indicates positive staining. (Right) Vertical bar (parts of a whole) indicates number and

percentage of neurons expressing selected genes in intradental neurons. *S100b, Scn10a,* and

Calca mark the majority of functionally-responding intradental neurons designating these as

large diameter, myelinated nociceptors indicative of HTMRs (66.0%, 31/47 responding

neurons). Most other neurons were *S100b*-positive labeled cells indicating large-diameter

myelinated cells (23.4%, 11/47 responding neurons). n = 5 mice, comprising 47 intradental

neurons.

Figure 2. Intradental neurons are HTMRs that encode direct force after enamel removal

and innervate the dental pulp and inner dentin

(A-C) Intradental neurons do not respond to direct forces applied to the intact tooth. (A)

Example trace (top) and associated heatmap (bottom) showing absence of intradental neuron

response to direct forces applied to the intact tooth, although they maintained responsiveness

to Endo-Ice. Stimuli used are indicated above the traces. Bar to the right of trace indicates 10

$\Delta F/F0$. (B) Bar graph showing proportion of intradental neurons that respond to direct force.

Plotted individual data points represent the proportion of force responsive/intradental neurons

per trial. Bar shows the mean and error bars indicate the SEM. n = 3 mice, comprising 46 cells.

(C) Bar graph showing proportion of intradental neurons that respond to Endo-Ice. Plotted

individual data points represent the proportion of Endo-Ice responsive/intradental neurons per

trial. Bar shows the mean and error bars indicate the SEM. n = 3 mice, comprising 46 cells.

(D-F) Intradental neurons respond to direct forces following structural damage of the tooth. (D)

Heatmap showing response profiles of 54 intradental neurons from 3 mice when direct force is

applied to the tooth while intact (left) and following pulp exposure via removal of overlying

enamel and dentin (right). (E) Example heatmap showing the response of intradental neurons

to direct force applied to exposed dentin. (F) Bar graph showing proportion of direct force

response/intradental neurons before and after two forms of tooth structural damage. Plotted

individual data points represent the proportion of force responsive/intradental neurons per trial.

Bar shows the mean and error bars indicate the SEM. n = 6 mice for intact enamel, n = 6 mice

for pulp exposed, n = 3 mice for dentin exposed. p-values from one-way ANOVA with Tukey

correction indicated with the graph. Data for D-F were obtained from Ai95(RCL-GCaMP6f)-D

(Ai95D) mice injected postnatally (P0-P3) with AAV9-Cre.

(G) Representative image of immunostained neuronal terminals merged with brightfield image

of dentin tubules in demineralized molars from *S100b*-Cre mice receiving TG injection of

AAV9-flex-tdTomato to induce sparse labeling of trigeminal sensory neurons. Enamel, a fully

mineralized structure, has been removed completely by demineralization. This experiment was

repeated in n = 5 mice. Scale bar: 20 µm.

(H) Representative image of immunostained neuronal terminals merged with brightfield image

of dentin tubules in demineralized molars from *Scn10a*-Cre mice injected using a Cre-

dependent AAV to induce sparce labeling of terminals (see methods). Enamel, a fully

mineralized structure, has been removed completely by demineralization. This experiment was

repeated in n = 4 mice with additional littermate and no primary controls. Scale bar: 20 µm.

Figure 3. Intradental HTMRs detect enamel damage, independent of *Piezo2* or *Scn10a*

(Na$_v$1.8).

(A-E) Intradental HTMRs sense cutting and superficial frictional damage of tooth enamel. (A)

Example traces showing intradental neuron response to enamel cutting and friction applied to

individual molars. Stimuli used are indicated above the traces. Bars to the right of traces

indicate 20 ΔF/F0. (B) Example heatmap containing traces shown in (A). (C) Bar graph

showing proportion of friction and cutting responsive intradental neurons identified by electrical

stimulation. Plotted individual data points represent the proportion for each trial. Bar shows the

mean and error bars indicate the SEM. n = 7 mice, comprising 205 total cells, p<0.01,

Unpaired t-test. (D) Line graph depicting magnitude of Ca^{2+} responses of intradental neurons

to friction versus cutting. Individual points represent calculated area under the curve (AUC)

during designated stimulation for each cell. n = 23 cells, p<0.0001, Paired t-test. (E) Vertical

bar (parts of a whole) indicates number and percentage of cutting responsive neurons that

express *Scn10a* (81.4%, 22/27 cutting responsive neurons, n = 4 mice). Results in E are

cutting responses from primary data also used in Figure 1J.

(F) Example heatmap showing stereotypical intradental HTMR responses to cutting persist in

Piezo2-cKO mice.

(G) Example heatmap showing stereotypical intradental HTMR responses to cutting persist in

Scn10a-KO mice.

(H) Bar graph depicting the proportion of intradental HTMRs that respond to enamel cutting in

wildtype, *Piezo2*-cKO, and *Scn10a*-KO mice. Plotted individual data points represent the

calculated proportion of enamel cutting responsive/intradental neurons. Bar shows the mean,

and error bars indicate the SEM. Intradental neurons were imaged from n = 6 or 7 adult mice

per genotype. No significant differences were observed with $p > 0.05$ using One-way ANOVA

with Tukey's correction.

Figure 4. Optogenetic activation of intradental HTMRs induces digastric muscle activity

and initiates a jaw opening reflex

(A-C) Optogenetic activation of intradental neurons induces activity in the anterior digastric

muscle EMG. (A) Schematic demonstrating the experimental approach to selectively activate

intradental neurons while recording EMG from the anterior digastric and masseter muscles.

Channelrhodopsin-2 was selectively expressed in right mandibular intradental neurons

innervating M1 and M2 molars via tooth injection of AAV6/2-hEF1a-iCre into Ai32(RCL-

ChR2(H134R)/EYFP) mice. An optical fiber was implanted over the right trigeminal ganglia to

selectively activate intradental neurons while EMG signals of the ipsilateral anterior digastric or

masseter muscles were recorded. (B) Example traces demonstrating EMG activity elicited by

optogenetic activation of intradental neurons (15 ms, 473 nm). EMG activity is observed in the

ipsilateral anterior digastric muscle, but not masseter muscle. Mice lacking channelrhodopsin-2

expression in intradental neurons show no changes in EMG activity for either the anterior

digastric or the masseter muscles in response to blue light stimulation; n = 3 mice for all

conditions. (C) Graph depicting the amplitude of activity of digastric muscle EMG in response

to blue light stimulation (15 ms, 473 nm). Plotted individual data points represent the peak-to-

peak amplitude of the elicited EMG trace. Error bar indicates SD. n = 3 mice per condition.

Significant differences were observed between experimental (ChR2+) and control animals

(ChR2-) with $p<0.0001$ using Unpaired t-Test.

(D-G) Optogenetic activation of intradental neuron terminals in the intact tooth results in

mandibular deflection (i.e. jaw opening). (D) Schematic demonstrating the experimental

approach. Channelrhodopsin-2 expression (Ai32) was driven by *Scn10a*-Cre. An optical fiber

(500 µm) was positioned perpendicular to the surface of the first maxillary molar tooth or

adjacent oral structures (hard palate or oral vestibule). Deflection was determined by

monitoring the position of maxillary and mandibular incisors (see Methods). (E) Traces of

average measured mandibular deflection in response to a single blue-light pulses (470 nm, 1s)

directed to the molar tooth. Black trace represents the average (n = 4 mice). Multicolored

traces represent average measurements from each mouse (n = 10 traces/mouse). Inset shows

brief timescale of trace from (E) demonstrating kinetics of mandibular deflection in relation to

light onset. (F) Average trace of measured mandibular deflection in response to trains of blue

or green light pulses (470nm or 545nm, 10 pulses, 0.5 Hz, 1s) directed at oral structures.

Sequential blue light pulses directed at the molar tooth (black trace), but not adjacent

structures (light and dark grey traces) induced mandibular deflection after each blue light

onset. No jaw movements were observed in response to green light stimulation. (G) Line graph

depicting average mandibular deflection in response to sequential blue light pulse train

stimulation of molar. Average deflection from each mouse is plotted as multicolored individual

points and represents mean peak amplitude and error bars show SEM of mandibular deflection

during stimulation. n = 10 measurements per mouse per train related to traces shown in (F).

Black points represent the average of n = 4 mice. $p > 0.05$ using Unpaired t-Test.