Caspase-11 drives macrophage hyperinflammation in models of Polg-related mitochondrial disease

Jordyn J. VanPortfliet[1,2], Yuanjiu Lei[2,3], Muthumeena Ramanathan[1], Camila Guerra Martinez[2], Jessica Wong[1], Tim J. Stodola[1], Brian R. Hoffmann[1], Kathryn Pflug[4], Raquel Sitcheran[4], Stephen C. Kneeland[1], Stephen A. Murray[1], Peter. J. McGuire[5], Carolyn L. Cannon[2], A. Phillip West[1,2,*]

[1]The Jackson Laboratory, Bar Harbor, Maine 04609, USA

[2]Department of Microbial Pathogenesis and Immunology, College of Medicine, Texas A&M University, Bryan, Texas 77807, USA

[3]Department of Pathology, Yale School of Medicine, New Haven, CT 06520, USA

[4]Department of Cell Biology and Genetics, College of Medicine, Texas A&M University, Bryan, Texas 77807, USA

[5]Metabolism, Infection and Immunity Section, National Human Genome Research Institute, National Institutes of Health, Bethesda, Maryland, 20892, USA

* Correspondence to: phillip.west@jax.org

<u>Abstract</u>

Mitochondrial diseases (MtD) represent a significant public health challenge due to their heterogenous clinical presentation, often severe and progressive symptoms, and lack of effective therapies. Environmental exposures, such bacterial and viral infection, can further compromise mitochondrial function and exacerbate the progression of MtD. Infections in MtD patients more frequently progress to sepsis, pneumonia, and other detrimental inflammatory endpoints. However, the underlying immune alterations that enhance immunopathology in MtD remain unclear, constituting a key gap in knowledge that complicates treatment and increases mortality in this vulnerable population. Here we employ in vitro and in vivo approaches to clarify the molecular and cellular basis for innate immune hyperactivity in models of polymerase gamma (Polg)-related MtD. We reveal that type I interferon (IFN-I)-mediated upregulation of caspase-11 and guanylate-binding proteins (GBPs) increase macrophage sensing of the opportunistic microbe *Pseudomonas aeruginosa* (PA) in *Polg* mutant mice. Furthermore, we show that excessive cytokine secretion and activation of pyroptotic cell death pathways contribute to lung inflammation and morbidity after infection with PA. Our work sheds new light on innate immune dysregulation in MtD and reveals potential targets for limiting infection- and inflammation-related complications in Polg-related MtD.

Introduction

Mitochondrial diseases (MtD) are the most common inborn error of metabolism and are caused by deleterious variants in genes that function in oxidative phosphorylation (OXPHOS) and mitochondrial DNA (mtDNA) maintenance[1]. MtD can present at any age, vary in severity, and display clinical heterogeneity with differing degrees of organ system involvement[2,3]. Although the genetic basis of MtD is becoming clearer, the underlying mechanisms that cause pathology, as well as the contribution of environmental factors to disease progression, are less clear. Notably, viral and bacterial infections may unmask MtD or hasten the stepwise progression of disease by triggering metabolic decompensation, a dangerous deterioration in mitochondrial function[4]. Clinical evidence suggests that patients with MtD are highly susceptible to infection by opportunistic respiratory pathogens. Moreover, pneumonia and microbial sepsis are the primary causes of early death in pediatric patients, and some adults, with MtD[5,6]. Thus, there is ample evidence highlighting the damaging toll of infection in MtD[7]. However, the underlying immune alterations that drive immunopathology and prevent proper control of infection represent a significant knowledge gap that complicates treatment and enhances morbidity and mortality in this vulnerable patient population.

In addition to their bioenergetic functions, mitochondria are key metabolic and signaling hubs in the mammalian immune system[8]. Mitochondria are also important sources of damage-associated molecular patterns (DAMPs), such as mtDNA, which engage innate immune signaling when released from the organelle and are linked to inflammation in an increasing number of human diseases[9,10]. There is growing appreciation for a bidirectional relationship between MtD and inflammation, where mitochondrial dysfunction in MtD activates the innate immune system leading to inflammatory responses that further compromise mitochondrial

activity[11]. In agreement with this hypothesis, elevated inflammatory cytokines have been

observed in patients with Polymerase gamma (Polg)-related Alpers-Huttenlocher syndrome[12],

Barth syndrome[13], and Friedreich ataxia[14,15]. Animal models have also revealed innate immune

system dysregulation as a frequent feature of MtD. For example, the $Ndufs4^{-/-}$ mouse model of

neurodegeneration and Leigh syndrome exhibits increased circulating pro-inflammatory

cytokines along with macrophage and microglial activation[16–19].

We recently uncovered that the $Polg^{D257A}$ mutator model of mtDNA instability exhibits a

hyperinflammatory innate immune status due to chronic engagement of the cyclic GMP-AMP

synthase (cGAS)—Stimulator of Interferon Genes (STING)—type one interferon (IFN-I) axis[20].

Notably, we reported that persistent IFN-I signaling in these mice causes widespread repression

of the antioxidant and anti-inflammatory transcription factor nuclear factor erythroid 2-related

factor 2 (Nrf2)[21,22] in myeloid immune cells and organs of the $Polg^{D257A}$ mouse. Consequently,

$Polg^{D257A}$ mice are more sensitive to septic shock caused by intraperitoneal injection of a high

dose of lipopolysaccharide (LPS)[20]. Although $Polg^{D257A}$ mutator mice do not phenocopy any

particular human MtD, these animals present pathology overlapping the MtD spectrum,

including hearing loss, cardiomyopathy, progressive anemia, and lymphopenia[23–25]. Moreover,

increased susceptibility to septic shock and elevated inflammatory and IFN-I signatures are

common features of both $Polg$ mutator mice and patients with diverse MtD[11].

Overall, our prior work provided foundational evidence linking mitochondrial

dysfunction to hyperinflammatory innate immune signaling in the $Polg^{D257A}$ model of MtD.

However, there remains a critical need to expand these findings into relevant infection models,

given the high rate of infections that frequently lead to innate immune hyperactivity and sepsis in

MtD. Pseudomonas aeruginosa (PA) is a common opportunistic pathogen of concern in MtD, as

infection can lead to sepsis and pneumonia[6]. PA is a robust activator of innate immunity in the

respiratory tract, and lung colonization by PA drives persistent recruitment of monocytes and

neutrophils leading to pneumonia, lung injury, and/or systemic bacterial spread[26,27]. PA is

especially problematic in MtD due to the prevalence of antibiotic resistant strains, abundance in

hospitals, and high rate of recurrence[26,28,29]. Thus, it is important to define how PA and other

opportunistic bacteria engages the innate immune system in MtD to define immunomodulatory

strategies that boost anti-bacterial immunity, while limiting hyperinflammation that may hasten

metabolic decompensation and disease progression.

In this study, we utilized in vitro and in vivo PA infection models to characterize the

underlying innate immune mechanisms governing hyperinflammation in $Polg^{D257A}$ mice. We also

report a new mouse model that incorporates a pathogenic $Polg$ variant observed in humans. We

reveal that both $Polg$ mutant models exhibit excessive macrophage inflammasome activation,

cytokine secretion, and pyroptotic cell death, which contribute to inflammation, acute lung

injury, and morbidity after infection with PA. These findings further advance knowledge of how

mitochondrial dysfunction impacts innate immune responses during bacterial infection and Polg-

related diseases and reveals innate immune targets for limiting infection- and inflammation-

related complications in MtD.

Results

$Polg^{D257A}$ mutant macrophages exhibit increased IFN-I and pro-inflammatory responses

after bacterial challenge

Our recent work uncovered that elevated IFN-I signaling augments innate immune

activity and inflammation in $Polg^{D257A}$ mice challenged with Toll-like receptor (TLR) agonist,

LPS[20]. We hypothesized that the hyperinflammatory phenotype of $Polg^{D257A}$ mice might be

further exacerbated during bacterial infection. Therefore, we challenged bone marrow-derived

macrophages (BMDMs) with *Pseudomonas aeruginosa* strain O1 (PAO1) (Fig. 1a) and

performed RNA sequencing (RNAseq) and pathway analysis (Fig. 1b). Similar to BMDMs

treated with LPS[20], PAO1-challenged $Polg^{D257A}$ BMDMs displayed heightened expression of

pro-inflammatory cytokines, IFN-I, interferon-stimulated genes (ISGs), and innate immune

signaling factors (Fig. 1c-d; Supplementary Fig. 1a). Gene Set Enrichment Analysis confirmed

PAO1-infected $Polg^{D257A}$ BMDMs exhibit more robust pro-inflammatory cytokine, chemokine,

and IFN-I gene expression relative to wild-type (WT) BMDMs (Supplementary Fig. 1b).

Additional experiments substantiated RNAseq datasets, revealing heightened expression of ISGs

after challenge of $Polg^{D257A}$ BMDMs with PAO1 between six to twenty four hours (Fig. 1e-f).

Proteomic analysis of PAO1-challenged BMDMs revealed strong concurrence with RNAseq

data, showing elevated IFN-I and inflammatory responses in $Polg^{D257A}$ BMDMs (Fig. 1g-h,

Supplementary Fig. 1c). Moreover, multi-analyte cytokine analysis by LegendPlex showed a

global elevation of proinflammatory cytokines produced by $Polg^{D257A}$ peritoneal macrophages

(PerMacs) six hours after PAO1 challenge (Fig. 1i) and ELISA showed a more rapid and

sustained release of these cytokines over a one-day time course (Fig. 1j). In sum, our data

highlight that $Polg^{D257A}$ macrophages are hyperactivated by both LPS and PAO1 challenge.

$Polg^{D257A}$ mutant macrophages exhibit potentiated pyroptotic cell death upon bacterial

challenge

As we observed increased IL-1β secretion from $Polg^{D257A}$ PerMacs (Fig. 1i, k), we next

investigated inflammatory cell death pathways during PAO1 infection. RNAseq and pathway

analysis revealed significant enrichment of genes involved in inflammasome signaling,

immunogenic cell death, and death receptor signaling in $Polg^{D257A}$ BMDMs after PAO1

challenge (Supplementary Fig. 1d-e). Pyroptotic cell death proteins were also highly upregulated

in proteomic analysis (Fig. 1l). To further confirm a role for pyroptosis in hyperinflammation

and increased cell death in $Polg^{D257A}$ macrophages, we incubated BMDMs with Ac-YVAD-cmk

(AcYVAD), a selective caspase-1 inhibitor. AcYVAD nearly abrogated IL-1β secretion from

$Polg^{D257A}$ macrophages, while also reducing the levels of inflammasome-independent

inflammatory cytokines TNFα and IL-6 upon PAO1 challenge (Fig. 2a). In contrast, QVD-OPH,

a more potent inhibitor of apoptotic caspases, had minor effects on PAO1-induced cytokine

secretion in $Polg^{D257A}$ BMDMs (Fig. 2a). Moreover, we did not observe striking differences in

necroptotic signaling proteins during PAO1 challenge, and necrostatin-1 failed to blunt elevated

cytokine secretion from $Polg^{D257A}$ BMDMs (Supplementary Fig. 2a-b).

Consistent with cell death gene expression patterns, $Polg^{D257A}$ BMDMs more rapidly

became propidium iodide (PI) positive during PAO1 challenge (Fig. 2b). $Polg^{D257A}$ PerMacs

stained weakly for the nuclear and plasma membrane dye CellMask HCS Deep Red and

exhibited rapid condensation of nuclei after PAO1 challenge (Fig. 2c-d). Reduced CellMask

staining was also observed in ASC positive WT BMDMs after LPS priming and nigericin

treatment (Supplementary Fig. 2c-d), indicating that low CellMask staining marks cells

undergoing pyroptosis. $Polg^{D257A}$ BMDMs also released more lactate dehydrogenase (LDH) after

PAO1 challenge (Supplementary Fig. 2e). We also noted elevated release of PAO1 into the

culture supernatant from $Polg^{D257A}$ BMDMs, concomitant with lower numbers of intracellular

bacteria (Supplementary Fig. 3a). Reduced intracellular bacteria was not the result of impaired

phagocytosis, as $Polg^{D257A}$ BMDMs displayed no impairment in uptake of LPS coated

fluorescent microspheres (Supplementary Fig. 3b-c). AcYVAD treatment was sufficient to blunt

pyroptotic phenotypes of $Polg^{D257A}$ macrophages (Fig. 2b-d) after incubation with PAO1, further

supporting a caspase-1-dependent cell death pathway.

In addition to activating NLRP3, PA can also activate the NLRC4 canonical

inflammasome and the non-canonical caspase-11 inflammasome[27,30–32]. Culture of $Polg^{D257A}$

BMDMs with non-pathogenic *E. coli* (DH5α), which lack flagellin and therefore can only

activate NLRP3 or caspase-11, also promoted pyroptotic cell death (Supplementary Fig. 3d-e)

and increased IL-1β secretion (Supplementary Fig. 3f), while boosting TNFα and IL-6 secretion

(Supplementary Fig. 3g). Collectively, these data indicate that elevated cell death and cytokine

secretion in $Polg^{D257A}$ macrophages during bacterial challenge are not due to PAO1 virulence

factors but are likely caused by potentiated activation of NLRP3 and/or caspase-11

inflammasomes.

Elevated IFN-I enhances expression and activation of caspase-11 and caspase-1 in $Polg^{D257A}$

macrophages

In WT macrophages, PAO1 mainly engages the NLRC4 inflammasome, while activation

of NLRP3 and caspase-11 inflammasomes occur at later stages of infection[30,33–35]. Given

elevated pyroptosis in PAO1-infected $Polg^{D257A}$ BMDMs, we next performed isolated challenge

experiments to evaluate hyperactivation of particular inflammasome pathways in $Polg^{D257A}$

macrophages. $Polg^{D257A}$ BMDMs treated with LPS alone did not exhibit increased PI uptake,

suggesting that priming is not sufficient to potentiate cell death (Supplementary Fig. 4a).

Surprisingly, LPS priming followed by flagellin transfection to activate the NLRC4

inflammasome revealed that $Polg^{D257A}$ macrophages exhibit reduced PI uptake and IL-1β

secretion (Supplementary Fig. 4b-c). LPS priming followed by the addition of ATP also revealed

modestly impaired NLRP3 inflammasome activation in $Polg^{D257A}$ BMDMs, as measured by PI

uptake, IL-1β secretion, and caspase-1 cleavage (Supplementary Fig. 4d-f). LPS priming

followed by nigericin treatment revealed similar impairments in NLRP3 inflammasome activity

(Supplementary Fig. 3g-i). OXPHOS has recently been shown to be vital for NLRP3 activation

in response to nigericin and other stimuli[36]. $Polg^{D257A}$ BMDMs displayed dysfunctional

mitochondrial morphology by TEM as well as reduced basal and maximal respiration at rest and

after LPS challenge (Supplementary Fig. 4j-m), suggesting that reduced canonical NLRP3

activation is likely due to OXPHOS impairment in $Polg^{D257A}$ BMDMs.

Although $Polg^{D257A}$ BMDMs exhibited reduced NLRP3 and NLRC4 responses when

triggered with pure inflammasome activators, mutant BMDMs displayed elevated caspase-1 and

Gasdermin D (GSDMD) cleavage, as well as increased caspase-11 and IL-1β expression, after

incubation with PAO1 as compared with WT controls (Fig. 2e). Moreover, AcYVAD treatment

inhibited the hypersecretion of IL-1β and other pro-inflammatory cytokines in $Polg^{D257A}$

BMDMs challenged with PAO1 for 24 hours (Fig. 2f-g). Notably, IL-1α secretion was markedly

higher in $Polg^{D257A}$ macrophages after PAO1 challenge (Fig. 1i and 2f). IL-1α is an alarmin that

feeds forward to augment production of other cytokines, and the robust secretion of IL-1α from

myeloid cells involves caspase-11[37–39]. Thus, given elevated caspase-11 expression in $Polg^{D257A}$

BMDMs, we next explored its involvement in cell death and cytokine secretion. After LPS

priming followed by LPS transfection, $Polg^{D257A}$ BMDMs rapidly became PI positive and

secreted increased levels of IL-1β (Fig. 2h-i). Both phenotypes could be markedly inhibited by

treatment with AcYVAD, indicating that heightened activation of caspase-11 feeds forward to

activate caspase-1 more robustly in $Polg^{D257A}$ BMDMs (Fig. 2h-i). Inhibition of NLRP3 with

MCC950 reduced cell death and hyperinflammation in $Polg^{D257A}$ BMDMs after PAO1 challenge

(Supplementary Fig. 5a-d). NLRP3-dependent caspase-1 activation can drive GSDMD to

permeabilize mitochondrial membranes[40]. Therefore, we isolated mitochondria from

unchallenged and PAO1-challenged BMDMs and examined GSDMD accumulation in

mitochondria-enriched fractions. Although there was no difference in the mitochondrial

accumulation of GSDMD between genotypes, we observed markedly increased GSDMD

cleavage in whole cell lysates of $Polg^{D257A}$ BMDMs rapidly upon PAO1 challenge

(Supplementary Fig. 5e).

As caspase-11 activation requires IFN-I priming[41] and $Polg^{D257A}$ macrophages and mice

exhibit elevated IFN-I signaling[20], we reasoned that IFN-I is responsible for caspase-11

expression and activation in $Polg^{D257A}$ BMDMs. In agreement, we observed potentiated caspase-

11 mRNA and protein after PAO1 infection (Fig. 1l, 2e, 2j). Notably, IFN alpha receptor

(IFNAR) blockade using the MAR1-5A3 monoclonal antibody markedly reduced PI uptake after

caspase-11 activation by transfected LPS (Fig. 2k). Collectively, these data indicate that the IFN-

I-dependent potentiation of caspase-11 drives non-canonical caspase-1 activation and

hyperinflammation in $Polg^{D257A}$ macrophages exposed to PAO1.

Elevated IFN-I signaling increases GBP expression and inflammation in $Polg^{D257A}$

macrophages.

We next hypothesized that elevated IFN-I may contribute to the hyperinflammatory

response of $Polg^{D257A}$ macrophages during PAO1 infection. Priming macrophages with an anti-

IFNAR antibody prior to PAO1 challenge reduced cell death and IL-1β release in $Polg^{D257A}$

BMDMs (Fig. 3a-c). This blockade also reduced production of NF-κB-dependent cytokines and

ISG expression at the RNA level over a day time course (Fig 3d-e).

Prior work has shown that IFN-I signaling during bacterial infection drives the expression

of guanylate binding proteins (GBPs)[42], which facilitate innate immune responses and non-

canonical inflammasome activation by disrupting membranes of pathogen-containing vacuoles,

or pathogens themselves, to liberate LPS or other PAMPs[31,43–45]. Consistent with elevated IFN-I

signaling, RNAseq analysis revealed more robust GBP expression in $Polg^{D257A}$ BMDMs after

challenge with PAO1 (Supplementary Fig. 5f). Proteomics analysis also showed elevated GBPs,

with marked increases in GBP2 and GBP5 in $Polg^{D257A}$ BMDMs six hours post challenge with

PAO1 (Supplementary Fig. 5g). Immunofluorescence microscopy revealed increased GBP5

expression in resting $Polg^{D257A}$ BMDMs (Supplementary Fig. 5h-i), as well as during PAO1

challenge (Fig. 3f-g). Notably, we observed a more robust colocalization of GBP5 with PAO1 in

$Polg^{D257A}$ macrophages after staining with an antibody specific for PA LPS. Moreover, GBP5

staining was more aggregated on PAO1-LPS positive bacteria than in WT BMDMs (Fig. 3h).

These findings suggest that elevated IFN-I in $Polg^{D257A}$ macrophages augments GBP expression,

likely enhancing sensing of LPS via caspase-11 that contributes to potentiated cell death and pro-

inflammatory cytokine release.

Caspase-11 activation, accentuated by GBP expression, controls pyroptosis and subsequent

inflammation in $Polg^{D257A}$ macrophages

Our data suggest that potentiation of caspase-11-mediated pyroptosis drives excessive

cell death and hyperinflammation in $Polg^{D257A}$ macrophages challenged with PAO1. Therefore,

we hypothesized that elevated levels GBPs may contribute to the enhanced release of

phagolysosomal PAO1 LPS for sensing by caspase-11 in the cytoplasm. We next used siRNA to

silence caspase-11 or GBP2 and GBP5 in WT and $Polg^{D257A}$ BMDMs to repress this pathway.

Knockdown efficiency of siRNAs was confirmed by qPCR of RNA at the time of infection

(Supplementary Fig. 6a). Notably, caspase-11 silencing reduced elevated cell death, heightened

cleavage of both GSDMD caspase-1, and increased secretion of IL-1 cytokines from $Polg^{D257A}$

BMDMs 2 hours post challenge with PAO1, while having little effect on WT BMDMs (Fig 4a-

d). Similar results were obtained by silencing GBP2 and GBP5 (Fig. 4a-d). Furthermore,

heightened caspase-11 association with LPS in $Polg^{D257A}$ BMDMs was reduced upon either

caspase-11 or GBP silencing, although to a lesser extent after GBP knockdown (Fig. 4e).

Caspase-11 or GBP knockdown lowered hypersecretion of proinflammatory cytokines in

$Polg^{D257A}$ BMDMs at the twenty-four hour timepoint (Fig. 4f). Finally, we utilized an LPS-

deficient PAO1 mutant to further investigate caspase-11 activation. The $algC$ gene product is

important for synthesis of the LPS core, and therefore this mutant displays altered lipid A levels

and lower LPS overall[46], which we confirmed by LPS staining intensity as measured with an

anti-PAO1 LPS antibody (Supplementary Fig. 6b). Indeed, $Polg^{D257A}$ BMDMs challenged with

PAO1-$algC$ displayed reduced cell death, inflammasome activation, IL-1β release, IFN-I

secretion, and pro-inflammatory cytokine production (Supplementary Fig. 6c-g), further

supporting that increased detection of PAO1 LPS by caspase-11 is a predominant driver of

hyperinflammatory phenotypes in $Polg^{D257A}$ macrophages.

Increased intracellular availability of LPS enhances caspase-11 activation and initiation of

IL-1 signaling in $Polg^{D257A}$ macrophages

Transmission electron microscopy (TEM) revealed that while $Polg^{D257A}$ BMDMs have

similar nuclear and cellular area at baseline (Supplementary Fig. 7a), $Polg^{D257A}$ BMDMs present

condensed nuclei and enlarged phagolysosomes 2 hours post challenge with PAO1 (Fig. 5a-b

and Supplementary Fig. 7c). Mitochondrial dysfunction can drive lysosomal defects and

leakiness[47–49], and we observed abnormal phagolysosomes in $Polg^{D257A}$ BMDMs by TEM two

hours after PAO1 (Fig. 5a-b). We also observed increased presence of vesicles adjacent to PAO1

in $Polg^{D257A}$ phagolysosomes, which resembled bacterial outer membrane vesicles (OMVs)[50,51].

At eight hours post PAO1 challenge, $Polg^{D257A}$ BMDMs contained more intact bacteria with

OMVs that were contained in phagolysosomes with poorly defined membranes (Supplementary

Fig. 7-e). This was accompanied by increased PAO1 protein expression as assessed by

proteomics analysis (Supplementary Fig. 7f). Oxidative stress is a critical driver of PA OMV

generation, and we noted increased reactive oxygen species[20] and nitrite production from

$Polg^{D257A}$ BMDMs after PAO1 challenge (Supplementary Fig. 7g). Therefore, we hypothesized

that caspase-11 hyperactivity in $Polg^{D257A}$ macrophages may be due to increased sensing of LPS

on OMVs or lingering bacteria, leading to more robust recruitment and activation of caspase-11.

Quantitation of IF imaging showed a greater intensity of caspase-11 and GBP5 colocalizing with

PAO1 in $Polg^{D257A}$ BMDMs (Fig. 5c-d), which was not observed after challenge with LPS-

deficient PAO1-*algC* (Supplementary Fig. 7h-i). To determine if these phenotypes were in part

due to LPS uptake and/or degradation kinetics, we utilized a fluorescently labeled LPS. Notably,

WT and $Polg^{D257A}$ BMDMs took up comparable amounts of LPS after LPS priming followed by

LPS transfection at acute timepoints (Supplementary Fig. 8a-b). However, we observed more

fluorescent LPS in the cytosol seven hours after LPS transfection of LPS primed BMDMs

(Supplementary Fig. 8c-d), as well as significantly more caspase-11 associated with LPS by IF

analysis (Supplementary Fig. 8e-f). WT and $Polg^{D257A}$ BMDMs contained similar levels of

fluorescent LPS signal immediately after passive uptake from the culture media (Supplementary

Fig. 8g-h), and $Polg$ mutant BMDMs did not release IL-1β at early timepoints (Supplementary

Fig. 8k). However, 48 hours after the addition of LPS to the cell culture media, $Polg^{D257A}$

BMDMs had a marked increase of cell-associated LPS (Supplementary Fig. 8i-j) and released

small amounts of IL-1β (Supplementary Fig. 8k). Together, these data suggest that elevated

levels of GBPs and caspase-11 in $Polg^{D257A}$ macrophages, coupled with increased bacterial

OMVs and LPS in the cytosol, promote enhanced caspase-11 activation, leading pyroptosis and

hyperinflammation.

In WT macrophages, Caspase-11 and caspase-1 are not directly involved in the

generation of nuclear-factor kappa B (NF-κB)-dependent cytokines such as TNFα and IL-6.

Therefore, we hypothesized that elevated caspase-11 activity may potentiate autocrine or

paracrine signaling via the early release of alarmins that hyperactive PA-induced IFN-I or NF-

κB. A kinetic stimulation study revealed increased phosphorylation of factors involved in IFN-I

signaling (i.e. TBK1, STAT1) in $Polg^{D257A}$ BMDMs at early timepoints after PAO1 challenge, as

well as elevated phospho-IκBα levels at early timepoints (Supplementary Fig. 9a). As mentioned

above, IL-1α release is associated with caspase-11 activation and GSDMD cleavage[52]. IL-1α

signaling through the IL-1 receptor (IL-1R) activates NF-κB and induces transcription of pro-IL-

1β, inflammasome components including NLRP3, and other proinflammatory cytokines

including TNFα and IL-6 [53–55]. To determine if early caspase-11-dependent IL-1α signaling feeds

forward to potentiate cytokine secretion in $Polg^{D257A}$ BMDMs at later timepoints, we used a

neutralizing antibody to block IL-1R signaling. Notably, the IL-1R antibody did not reduce

elevated IL-1 cytokine release, cell death, or GSDMD and caspase-11 cleavage in $Polg^{D257A}$

BMDMs two hours post challenge (Fig. 5e-f, Supplementary Fig. 9b-c). However, IL-1R

blockade was sufficient to markedly decrease IL-1 cytokines, IL-6, and TNFα from $Polg^{D257A}$

BMDMs 24 hours post PAO1 challenge (Fig. 5g-h). Overall, these data indicate that the rapid,

caspase-11-dependent release of IL-1α after PAO1 challenge drives feed forward NF-κB and

IFN signaling to potentiate inflammatory cytokine production in $Polg^{D257A}$ macrophages.

A novel $Polg^{R292C}$ mouse model incorporating a deleterious patient variant exhibit

macrophage mtDNA depletion, elevated IFN-I responses, increased pyroptosis, and hyper-

inflammation after challenge with PAO1

To determine if our findings in $Polg^{D257A}$ macrophages are more broadly applicable to

other models of OXPHOS deficiency due to mtDNA perturbations, we leveraged a novel model

of Polg-related mitochondrial disease harboring a single amino acid missense mutation

(Arg292Cys) in exon 4 of the exonuclease domain of Polg (Supplementary Fig. 10a), which is

homologous to the human Arg309Cys (c.925C>T) pathogenic variant[56,57]. We generated these

mice using CRISPR/Cas9 editing of ES cells and verified germline transmission of the mutation

by sequencing (Supplementary Fig. 10b). The Arg309Cys variant alone or in combination with

other *Polg* variants causes reduced mtDNA copy number, mtDNA deletions, and severe

phenotypes including mitochondrial myopathy, encephalopathy, lactic acidosis and stroke-like

episodes (MELAS), peripheral neuropathy, ataxia, and epilepsy in patients. Although being born

below expected mendelian ratios, homozygous $Polg^{R292C}$ mice are viable and display a reduced

mitochondrial DNA copy number and elevated ISGs in multiple tissues (Supplementary Fig.

10c-e).

BMDMs derived from $Polg^{R292C}$ mice exhibited mtDNA depletion at rest (Supplementary

Fig. 11a), as well as lower numbers of nucleoids and fragmented mitochondrial networks after

stimulation with LPS (Supplementary Fig. 11b-c). In line with our data from the $Polg^{D257A}$

mutator mice, $Polg^{R292C}$ BMDMs displayed a constitutive elevation of ISGs at rest and

potentiated ISG expression after LPS stimulation (Supplementary Fig. 10d-e). Using

extracellular flux analysis, we also noted markedly reduced basal and maximal respiration after

LPS stimulation (Supplementary Fig. 11f-h), indicating that $Polg^{R292C}$ macrophages exhibit

OXPHOS deficiencies resembling those of $Polg^{D257A}$ BMDMs (Supplementary Fig. 4l-m).

After challenge with PAO1, $Polg^{R292C}$ BMDMs exhibited increased ISG and GBP

expression (Fig. 6a), while also secreting more cytokines over a twenty-four-hour time course

(Fig. 6b). Similar to $Polg^{D257A}$ BMDMs, $Polg^{R292C}$ exhibited elevated rates of PAO1-induced cell

death, IL-1 cytokine secretion, and proinflammatory cytokine production, which were blunted by

treatment with caspase-1 inhibitor AcYVAD (Fig. 6c-e, Supplementary Fig. 12) or IFNAR

blockade (Fig. 6f-h). Finally, silencing of caspase-11 in $Polg^{R292C}$ BMDMs (Fig. 6i) alleviated

acute cell death (Fig. 6j-k), IL-1 release (Fig. 6l), and secretion of IL-6 and TNFα at late

timepoints (Fig. 6m). These data suggest that elevated IFN-I, hyperinflammation, and pyroptosis

in response to PAO1 are common features of macrophages with deficient Polg exonuclease

activity, including the MtD-relevant Arg292Cys variant.

Intratracheal instillation of a non-lethal dose of PAO1 elicits greater lung cytokine

production and alterations in lung immune cell populations in $Polg^{D257A}$ mice

Patients with primary MtD often experience recurrent viral and bacterial infections of the

respiratory tract, including infections by PA[5]. These events can trigger metabolic

decompensation and can progress to systemic inflammatory endpoints and/or sepsis. However,

few studies have examined infection kinetics and lung immune responses in Polg-related

mitochondrial disorders. We therefore employed intratracheal instillation protocols to administer

LPS or PAO1 into the lungs of $Polg^{D257A}$ mutator mice. Histological analysis of resting $Polg^{D257A}$

lungs did not reveal obvious differences in morphology or cellular composition (Supplementary

Fig. 13a-c), although consistent with our earlier findings[20], basal expression of ISGs were

elevated compared to WT littermates (Supplementary Fig. 13d). After LPS was instilled into

lungs through cannulation of the trachea, $Polg^{D257A}$ mice secreted more TNFα and IL-6 at all

timepoints and increased IL-1β at the 12-hour timepoint in the bronchial alveolar lavage fluid

(BAL) compared to WT mice (Supplementary Fig. 13e).

To next characterize lung innate immune responses to PAO1 infection in the $Polg^{D257A}$

mice, we challenged mice with a non-lethal dose of PAO1[58]. Both WT and $Polg^{D257A}$ mice

infected with $1x10^4$ colony forming units (CFU) of PAO1 via intratracheal instillation lost

weight initially but returned to normal weight (Supplementary Fig. 14a) and exhibited no

culturable bacteria (data not shown) 3 days post infection. Flow cytometric analysis of lung cells

in BAL fluid revealed an overall decrease in the percentage of CD45$^+$ immune cells at rest and

after infection, consistent with general leukopenia noted in $Polg^{D257A}$ mice[20,25] (Supplementary

Fig. 14b-c). Analysis of specific immune cell markers on lung CD45$^+$ cells revealed few changes

in BAL populations (Supplementary Fig. 14d-l) between uninfected WT and $Polg^{D257A}$ mice,

although there were more SiglecF$^+$ and F4/80$^+$ cells at rest perhaps indicating elevated numbers

of lung macrophages (Supplementary Fig. 14e,g). Six hours post infection, there was a marked

reduction in SiglecF$^+$ cells in $Polg^{D257A}$ mice relative to WTs and an increase in CD3$^+$ T cells

(Supplementary Fig. 14e,i). Cells labeling with the neutrophil marker, Ly6G, were dramatically

increased at 6 hours, although there were no differences between WT and $Polg^{D257A}$ mice

(Supplementary Fig. 14k). At 48 hours post infection, neutrophils remained the most

predominant CD45$^+$ population in both genotypes, but we observed a reversion toward increased

numbers of cells staining positive for macrophage markers SiglecF$^+$ and F4/80$^+$ in $Polg^{D257A}$

mice (Supplementary Fig. 14e,g,k). Despite modest changes in cellular composition of the BAL

fluid, $Polg^{D257A}$ mice had elevated numbers of CD45$^+$ cells expressing TNFα both at rest and six

hours post infection (Supplementary Fig. 14m), consistent with increased TNFα production after

LPS challenge.

Multianalyte analysis of the BAL fluid using Legendplex assays revealed 2-4 fold

increases in lung pro-inflammatory cytokines in $Polg^{D257A}$ mice 6 hours post PAO1 infection

(Fig. 7a, Supplementary Fig. 15a). Consistent with resolution of infection in both genotypes,

cytokine differences in the BAL of WT and $Polg^{D257A}$ mice were less pronounced at 48 hours

post infection (Supplementary Fig. 15b). However, IFNβ and inflammatory cytokines associated

with the skewing and differentiation of T cell Th1 (IL-27) and Th17 subsets (IL-23) remained

elevated in $Polg^{D257A}$ mice in an IFNAR-dependent fashion (Supplementary Fig. 15b). Analysis

of lung protein extracts revealed elevated caspase-1 cleavage, as well as increased caspase-11,

NLRC4, and NLRP3 expression in $Polg^{D257A}$ mice 48 hours post intratracheal instillation of

1x10^4 CFU of PAO1. GSDMD, GBP5, and other ISGs were also elevated in $Polg^{D257A}$ lungs,

consistent with sustained IFNβ expression observed by Legendplex (Supplementary Fig. 15c).

Given that recurrent respiratory tract infections are often observed in patients with MtD,

we next infected mice with 1x10^4 CFU PAO1, followed by a 7-day rest period, then re-infected

with 1x10^4 CFU PAO1. Histological analysis of lungs 24 hours following re-infection revealed

monocyte and granulocyte infiltration into lungs, areas of vascular hemorrhage, and more eosin

positive cells in $Polg^{D257A}$ lungs relative to WT (Supplementary Fig. 16a-b). Lung IL-12 family

cytokines and IFNβ were significantly higher in $Polg^{D257A}$ mice at the 24-hour harvest point

following re-infection, while other cytokines were trending higher in $Polg^{D257A}$ mice, although

they did not reach statistical significance (Supplementary Fig. 16c). We also observed higher

bacterial burdens in the lungs and spleens of $Polg^{D257A}$ mice compared to WT controls

(Supplementary Fig. 16d). Together, these data indicate that $Polg^{D257A}$ mice exhibit

hyperinflammation and impaired antibacterial immunity upon re-infection with PAO1.

PAO1 elicits greater caspase-11 expression, increases inflammation, and reduces survival

of *Polg* mutant mice

We next utilized a higher dose of PAO1 to further assess innate immune responses *Polg*

mutant mice. RNAseq of lung tissue collected six hours after intratracheal instillation of $1x10^5$

CFU PAO1 revealed a striking upregulation of many inflammatory-, IFN-, and programmed cell

death-related pathways in $Polg^{D257A}$ mice relative to WT littermates (Fig. 7b-c, Supplementary

Fig. 17a). In agreement with macrophage data, transcriptional signatures of pyroptosis were

apparent, with *Casp11*, *Nlrp3*, *Il1a*, and *Il1b* some of the most abundantly enriched transcripts in

$Polg^{D257A}$ versus WT lung lysates (Fig. 7d). $Polg^{R292C}$ lung homogenates also displayed

upregulation in transcriptional pathways indicative of hyperinflammation, elevated IFN

signaling, and potentiated pyroptotic cell death 6 hours post PAO1 infection (Supplementary Fig.

17b-d). Western blotting of $Polg^{D257A}$ lung homogenates reinforced RNAseq data, showing

increased protein expression of pyroptosis factors including caspase-11 and cleaved GSDMD

(Fig. 7e), as well as a depletion in OXPHOS Complex I and IV subunits 6 hours post infection

(Supplementary Fig. 17e). Moreover, $Polg^{D257A}$ mice exhibited elevated bacterial burdens in the

lung, liver, and spleen (Fig. 7f). Increased bacterial dissemination was accompanied by elevated

plasma proinflammatory cytokines, as well as elevated ISG expression with reduced OXPHOS

in $Polg^{D257A}$ liver homogenates (Supplementary Fig. 17f-g). We next utilized the PAO1-$algC$

mutant to further define whether reduced LPS sensing via caspase-11 would alter response to

infection in $Polg^{D257A}$ mice. Compared to WT PAO1, PAO1-$algC$ did not disseminate (Fig.

7g), or trigger increased expression of pyroptotic proteins and cytokines in $Polg^{D257A}$ lungs

relative to WTs (Fig. 7h, Supplementary Fig. 17h). Overall, these data suggest $Polg$ mutant mice

display increased caspase-11-medated sensing of PAO1 LPS, elevated bacterial burdens, and

hyperinflammatory responses during acute PAO1 infection

At later timepoints, $Polg^{D257A}$ lungs displayed elevated neutrophilic inflammation and

overall inflammatory scores via histological analysis (Fig. 7i-j, Supplementary Fig. 17i), despite

harboring similar bacterial burdens to WT mice (Supplementary Fig. 17j). However, caspase-11

and IL-1β remained elevated in lung lysates of $Polg^{D257A}$ mice compared to WT littermates

twenty hours post infection (Fig. 7k). Using a comprehensive health scoring system to track

breathing, hydration, body appearance, and activity, we noted that $Polg^{D257A}$ mice appeared

sicker and continued to decline in health, whereas WT animals recovered from infection (Fig.

8l). Even though we observed a steady decline in the health of $Polg^{D257A}$ mice at 48 hours after

infection, PAO1 were largely cleared in both $Polg^{D257A}$ and WT animals (Supplementary Fig.

17k). However, $Polg^{D257A}$ mice sustained elevated ISG expression in peripheral tissues,

suggesting sustained IFN-I signaling (Supplementary Fig. 17l-m). Finally, we observed that a

greater number of $Polg^{D257A}$ mice reached a moribund state after infection with $1x10^5$ CFU of

PAO1 relative to WT littermates (Fig. 8m). Overall, these in vivo data suggest that $Polg^{D257A}$

mice are more prone to excessive inflammation and acute lung injury following infection with

PAO1.

<u>Discussion</u>

Patients with MtD experience a greater frequency of bacterial and viral infections that

can rapidly progress to sepsis or systemic inflammatory response syndrome (SIRS), leading to

premature death[5]. Recurrent infections can also accelerate MtD progression by promoting

metabolic decompensation, a life-threatening condition that hastens deterioration of

mitochondrial function[59–61]. Polg is the sole mitochondrial DNA polymerase, and variants in the

POLG gene are a prevalent cause of MtD[62]. There are several clinical reports of bacterial or viral

infections unmasking Polg-related MtD symptoms and hastening disease progression[63,64].

Moreover, one study revealed the presence of inflammatory cytokines in the cerebrospinal fluid

of a patient with Polg-related MtD[12,19]. As cytokine-mediated toxicity is an important driver of

metabolic decompensation[7], there is a critical need to understand immune responses in patients

with Polg-related disorders and other MtD to manage infections and inflammation-related

complications. This study builds upon our prior work to characterize innate immune responses in

mouse models of MtD[20,65] and leverages bacterial infection, cytokine profiling, and in vitro and

ex vivo cellular assays to define molecular and cellular nodes governing hyperinflammatory

immune phenotypes in two models of Polg-related MtD (Supplementary Fig. 18).

Our prior work revealed that elevated IFN-I responses in $Polg^{D257A}$ mutator mice are

mediated by chronic mtDNA release and cGAS-STING signaling[20]. Using our new, disease

relevant $Polg^{R292C}$ mouse model, we observed increased IFN-I signatures basally and after

challenge with PAO1, similar to $Polg^{D257A}$. Analogous variants in human POLG at R309 affect

proofreading activity[66], and therefore we hypothesize that the $Polg^{R292C}$ exonuclease mutation

also impairs replication fidelity, leading to mtDNA release and chronic IFN-I signaling. A recent

study reported that fibroblasts from patients carrying the *POLG1* c.2243G>C, p.W748S

mitochondrial recessive ataxia syndrome (MIRAS) allele display reduced IFN-I signaling in

response to viral infection and nucleic acid stimuli[67]. The MIRAS variant affects POLG

holoenzyme processivity but not the exonuclease domain, and thus impaired exonuclease activity

leading to mtDNA mutations and deletions may selectively trigger IFN-I in the $Polg^{D257A}$ and

$Polg^{R292C}$ backgrounds. Despite reduced IFN-I, mice homozygous for the MIRAS variant do

exhibit increased necroptosis, elevated liver immune cell infiltration, and potentiated

inflammatory cytokine secretion during viral infection[67]. Thus, inflammatory cell death and

hyperinflammation are common features in both patient cells and mouse models of Polg-related

MtD.

Our data reveal that IFN-I signaling is a key driver of caspase-11-dependent pyroptosis

and hyperinflammation in *Polg* mutant macrophages. Pre-treatment with an anti-IFNAR

neutralizing antibody was sufficient to reduce both augmented PI uptake after LPS transfection

and cytokine hypersecretion after challenge with PAO1. IFN-I is critical to license caspase-11

expression[41], but also to enhance expression of interferon-inducible GTPases like GBPs, which

help to expose cytosolic LPS for detection[31,68–71]. We have found that macrophages from both

the $Polg^{D257A}$ and our new MtD-relevant $Polg^{R292C}$ model exhibit enhanced expression of

multiple GBPs at rest and after challenge with PAO1. Moreover, we observed that GBP5

strongly co-localizes with intracellular PAO1 and caspase-11 in $Polg^{D257A}$ BMDMs. GBPs are

important in orchestrating caspase-11 activation to bacterial outer membrane vesicles

(OMVs)[50,51] and can localize to damaged cell membranes[45]. Although the mechanisms by which

GBPs sense intracellular pathogens and facilitate caspase-11 activation are not fully understood,

silencing of caspase-11 or GBP2 and GBP5 markedly reduced pyroptosis and the secretion of IL-

1 and other cytokines from both *Polg* mutant BMDMs. TEM imaging also revealed altered

PAO1-containing phagolysosomal membranes and increased bacteria-associated vesicles in

Polg^D257A BMDMs. Thus, it is likely that GBPs and caspase-11 more readily sense PAO1 LPS

due to phagolysosmal leakiness, or elevated OMV generation, yet more work is required to fully

define the underlying mechanisms.

506   Our data reveal that *Polg* mutant macrophages undergo rapid pyroptosis upon PAO1

challenge in a NLRP3- and caspase-1-dependent fashion. Although caspase-1 cleavage is

elevated in *Polg*^D257A compared to WT macrophages during PAO1 challenge and its inhibition is

sufficient to block augmented PA-induced cell death and IL-1β secretion, we found that

Polg^D257A BMDMs are less responsive to direct activation of the NLRP3 and NLRC4

inflammasomes by their respective ligands. In contrast, activation of the caspase-11

inflammasome via LPS priming followed by transfection of LPS into the cytosol drove more PI

uptake and IL-1β expression in *Polg*^D257A BMDMs. Therefore, sensing of cytosolic PA LPS by

the non-canonical inflammasome leads to more robust activation of the NLRP3 for caspase-1

dependent cell death and cytokine secretion. Notably, a PAO1-*algc* mutant lacking robust LPS

synthesis did not hyper-stimulate pyroptosis and cytokine secretion, providing further support for

caspase-11 as the predominant driver of inflammatory cell death in *Polg* mutant mice.

518   It is interesting to note that treatment of *Polg* mutant macrophages with caspase-1

inhibitor dramatically reduced nearly all cytokine levels in culture supernatants after challenge

with PAO1. These include inflammasome-independent cytokines such as TNFα, IL-6, IFNγ, and

IL-12 family members as well as the DAMP IL-1α, which does not require caspase-1 cleavage

for secretion but is downstream of caspase-11 and GSDMD[37,72]. This is in contrast with cytokine

secretion in WT BMDMs, where AcYVAD treatment only reduced IL-1β levels during

challenge with PAO1. Similarly, anti-IFNAR blockade or caspase-11 and GBP knockdown

reduced hyper-secretion of IL-1β levels and reduced IL-6 and TNFα in *Polg* macrophages, while

having marginal effects on WT BMDM cytokine profiles. While blocking the IL-1R did not

reduce early cell death in *Polg* mutant macrophages, it was sufficient to blunt elevated cytokines

a day after PAO1 challenge. This suggests that early IL-1α release downstream of capsase-11

and GSDMD synergizes with IFN-I signaling to sustain hyperinflammation in PAO1-challenged

Polg mutant BMDMs. Our data do not definitively rule out contributions for canonical

inflammasome-mediated IL-1β in potentiating NF-κB-mediated cytokine release. However, the

link between caspase-11 activation and IL-1α suggest that the early release of this alarmin may

prime neighboring cells for activation of canonical inflammasomes, further promoting increased

cell death and inflammation in *Polg* mutant macrophages.

Although it is appreciated that bacterial and viral infections are problematic in MtD, few

studies have examined innate immune responses in relevant mouse models of disease. Using an

intratracheal instillation model, we found that $Polg^{D257A}$ mice exhibit hyperactive innate immune

responses resulting in elevated cytokine production, changes in lung immune cell dynamics

recruitment, and increased morbidity following PAO1 infection. Notably, we observed elevated

expression of inflammasome proteins including caspase-11, ISGs including GBPs, and increased

caspase-1 cleavage in $Polg^{D257A}$ lung homogenates after infection. Remarkably, RNAseq analysis

of PA infected lungs from 3-month-old $Polg^{R292C}$ mice also revealed increased pyroptosis,

interferon, and hyperinflammatory signaling. These data agree with our in vitro findings

implicating IFN-I as an overarching regulator of caspase-1/11 activation and hyperinflammation

in our models. Additional studies are warranted to define the cellular populations that govern

lung hyperinflammation and to determine the degree to which caspase-11 shapes inflammatory

signaling and morbidity in *Polg* mutant mice in vivo.

While *Polg*^D257A^ mice had higher bacterial burdens in the lung and peripheral tissues at

acute timepoints, they ultimately managed to control the infection by 20 hours post infection and

cleared bacteria by 48 hours. Despite this, lungs from *Polg*^D257A^ mice were more inflamed at 20

hours post infection and the mice had more severe sickness scores, resulting in higher rates of

morbidity. This suggests that the adverse outcomes observed in *Polg* mutant animals may be due

to chronic and damaging hyperinflammation as opposed to impaired bacterial clearance.

Therefore, therapeutic targeting of IFN-I and caspase-11 may represent promising avenues to

restore proper immune balance patients with Polg-related MtD. Although our work does not

include clinical data, two recent publications revealed elevated IFN-I signatures in the whole

blood and peripheral blood mononuclear cells (PBMCs) from patients with diverse MtD disease

variants[11,73]. Moreover, gene ontology analysis revealed heightened IL-1β signatures in MtD

PBMCs[11]. Our study provides important new insight into the innate immune mechanisms

governing infection-related hyperinflammation in mouse models of MtD. The consistency of our

discoveries across two different *Polg* mutant models strengthens the relevance of our findings to

human disease and suggest that IFN-I and inflammasome hyperactivity may be critical

determinants of increased rates of sepsis and SIRS in patients with MtD.

Figures and Legends

Fig. 1: *Polg^{D257A}* **mutator macrophages exhibit increased IFN-I and pro-inflammatory responses after challenge with LPS or** *Pseudomonas aeruginosa***.**

(a) Workflow for preparing RNA from *Polg* mutant BMDMs after *Pseudomonas aeruginosa* (PA) strain O1 (PAO1) challenge.

(b) Ingenuity Pathway Analysis (IPA) of RNA-seq from bone marrow-derived macrophages (BMDMs) showing Z-scores of pathways in *Polg^{D257A}* homozygous mutant BMDMs compared to wild-type (WT) after LPS or PAO1 challenge. hpc=hours post challenge. N=3 biological replicates/genotype/condition.

(c) Log^2 fold changes of representative genes in IPA pathways from **(b)** in *Polg^{D257A}* BMDMs compared to WT after LPS or PAO1.

(d) Protein abundance of interferon-stimulated genes (ISGs) in BMDMs exposed to PAO1 at a multiplicity of infection (MOI) of 5 at the indicated hours post challenge (hpc).

(e) qRT-PCR analysis of ISGs in WT and *Polg^{D257A}* BMDMs after PAO1. N=3 biological replicates, 2 technical replicates/genotype/condition.

(f) Western blot analysis of interferon stimulated genes (ISGs) over time course infection.

(g) Ingenuity pathway analysis showing z-scores of upregulated pathways by proteomics from *Polg^{D257A}* (PAO1/Un)/WT (PAO1/Un) BMDM. N=4 biological replicates

(h) IPA Upstream Regulator analysis showing Z-scores in *Polg^{D257A}* (PAO1/Un)/WT (PAO1/Un).

(i) Legendplex pro-inflammatory cytokines analysis of PerMac supernatants 24 hours post challenge (hpc). N=3 biological replicates

(j) Proinflammatory cytokines in supernatant of peritoneal macrophages (PerMacs) measured by ELISA. MOI= multiplicity of infection, N=2 biological replicates per time point.

(k) IL-1β in supernatant of peritoneal macrophages (PerMacs) measured by ELISA. MOI=

multiplicity of infection, N=2 biological replicates per time point.

(l) Ingenuity pathway analysis of proteomics from $Polg^{D257A}$ (PAO1, 8hpc/Un)/WT (PAO1, 8

hpc/Un) BMDM proteins in the Pyroptotic Signaling Pathway. N=4 biological replicates

Statistics: (e, i & j) Two-Way ANOVA. *P<0.05, **P<0.01, ***P<0.001, ****P<0.0001 Error

bars represent SEM.

Fig. 2

a Polg^D257A BMDM, PAO1 MOI=5
■ +DMSO ■ +AcYVAD ■ +QVD-OPh

IL-1β   IL-6   TNFα

b BMDM, PAO1 MOI=5
● WT
■ Polg^D257A
○ WT+AcYVAD
□ Polg^D257A +AcYVAD

c PerMac, PAO1 MOI = 5, 2 hpc
■ WT
■ Polg^D257A
□ WT +AcYVAD
■ Polg^D257A +AcYVAD

d PerMac, PAO1 MOI = 5, 2 hpc

WT+DMSO — Merge / DAPI / PAO1-GFP / CellMask — 10 μm
WT +AcYVAD — Merge / DAPI / PAO1-GFP / CellMask — 10 μm
Polg^D257A +DMSO — Merge / DAPI / PAO1-GFP / CellMask — 10 μm
Polg^D257A +AcYVAD — Merge / DAPI / PAO1-GFP / CellMask — 10 μm

e BMDM, PAO1 MOI= 5
1   1.5 hpc
WT / Polg^D257A / WT / Polg^D257A

CASPASE-1
p20
CASPASE-11
GSDMD
GSDMD-NT
IL-1β
HSP60

■ WT
■ Polg^D257A
□ WT +AcYVAD
■ Polg^D257A +AcYVAD

f BMDM, PAO1 MOI=5, LegendPlex, 24 hpc
IL-1β   IL-1α

g TNFα   IL-6   IFNγ   IL-23   IL-10   IL-27   IL-12p70

BMDM, LPS Priming + LPS transfection

h PI Positive Cells (10^3)
● WT
■ Polg^D257A
○ WT+AcYVAD
□ Polg^D257A +AcYVAD

i IL-1β — 9 hpc

j Casp11 — Un / PA, 6 hpc

k PI Positive Cells (10^3)
● WT IgG
○ WT αIFNAR
■ Polg^D257A αIgG
□ Polg^D257A αIFNAR

Fig. 2: Caspase-11 and caspase-1 drive pyroptosis and inflammation in *Polg^{D257A}* macrophages challenged with PAO1.

(**a**) ELISA analysis of inflammatory cytokines in BMDM supernatant after PAO1 or PAO1 plus caspase inhibitors. N= 2 biological replicates/genotype/condition/timepoint.

(**b**) Cytation5 image analysis of PI positive BMDMs after PAO1 or PAO1 plus caspase-1 inhibitor. N= 6 biological replicates/genotype/condition.

(**c**) Percent PerMacs dying after PAO1 and PAO1 plus caspase-1 inhibitor from IF (**d**). N=4 IF images/genotype/condition.

(**d**) Representative IF of PerMacs after PAO1 or PAO1 plus caspase inhibitor. Cells were stained with DAPI to mark the nucleus, CellMask to mark the total cell volume, and anti-GFP to mark GFP tagged PAO1. White arrows highlight dying cells characterized by condensed nuclei and loss of CellMask staining.

(**e**) Western blot analysis of lysates from BMDMs challenged with PAO1.

(**f**) LegendPlex IL-1 cytokine analysis of BMDM supernatant 24 hpc. All undefined comparisons between WT and WT+AcYVAD are not significant. N=3 biological replicates/genotype/condition.

(**g**) LegendPlex pro-inflammatory cytokine analysis of BMDM supernatant 24 hpc. All undefined comparisons between WT and WT+AcYVAD are not significant. N=3 biological replicates/genotype/condition.

(**h**) Cytation5 image quantification of PI positive BMDMs after LPS priming plus LPS transfection and LPS priming plus LPS transfection plus caspase-1 inhibitor. N= 3-5 biological replicates/genotype/condition.

(i) ELISA analysis of IL-1β in BMDM supernatant after LPS priming plus LPS transfection in

the presence or absence of caspase-1 inhibitor. N=4 biological replicates

(j) qRT-PCR analysis of Caspase-11 RNA expression in BMDMs after PAO1. N=3 biological

replicates, 3 technical replicates/genotype/condition.

(k) Cytation5 image quantification of PI positive cells in BMDMs after LPS priming plus LPS

transfection and incubated with αIFNAR blocking antibody or αIgG control. N= 4-6 biological

replicates/genotype/condition.

Statistics: (c, f, g & i) One-Way ANOVA (a, b, h, j & k) Two-Way ANOVA. *P<0.05,

**P<0.01, ***P<0.001, ****P<0.0001 Error bars represent SEM.

Fig. 3

a BMDM, PAO1 MOI=5, 2 hpc

- WT+IgG
- WT+αIFNAR
- Polg^{D257A}+IgG
- Polg^{D257A}+αIFNAR

Cell Death

b
WT IgG — Merge / DAPI / PAO1-LPS / CellMask — 10 μm
WT αIFNAR — Merge / DAPI / PAO1-LPS / CellMask — 10 μm
Polg^{D257A} IgG — Merge. / DAPI / PAO1-LPS / CellMask — 10 μm
Polg^{D257A} αIFNAR — Merge / DAPI / PAO1-LPS / CellMask — 10 μm

c BMDM, PAO1 MOI=5, 24 hpc
IL-1β

d IL-6

TNFα

e BMDM, PAO1 MOI=5, qPCR, 6 hpc
Zbp1   Stat1   Ifit3   Gbp9   Ifit1   Gbp3
Relative Expression (fold)

f BMDM, PAO1 MOI=5, 2 hpc
WT — Merge / GBP5/PAO1-LPS / GBP5 / PAO1-LPS — 10 μm
Polg^{D257A} — Merge / GBP5/PAO1-LPS / GBP5 / PAO1-LPS — 10 μm

g GBP5 intensity
Sum Intensity/Area (10^5)
2 hpc
- WT
- Polg^{D257A}

h PAO1-LPS co-localized with GBP5
GFP localized with GBP5 (sum intensity, 10^4)
2 hpc

Fig. 3: Elevated interferon signaling contributes to elevated GBP expression and hyperinflammation in *Polg^{D257A}* macrophages.

(a) Percent BMDMs dying after PAO1 and PAO1 plus anti-IFNAR blockade from **(b)** representative IF. N=5-6 IF images/genotype/condition.

(c) ELISA analysis of pyroptotic cytokines in BMDM supernatant after PAO1 or PAO1 plus anti-IFNAR blockade. N= 2 biological replicates/genotype/condition.

(d) ELISA analysis of inflammatory cytokines in BMDM supernatant after PAO1 or PAO1 plus anti-IFNAR blockade. N= 2 biological replicates/genotype/condition.

(e) qRT-PCR analysis of ISGs in WT and *Polg^{D257A}* BMDMs after PAO1 and PAO1 plus anti-IFNAR blockade. N=3 technical replicates/genotype/condition.

(f) Representative IF of BMDMs 2 hpc with PAO1 at a multiplicity of infection (MOI) of 5. Cells were fixed, permeabilized, and stained with DAPI to mark the nucleus, anti-LPS antibody to stain PAO1 LPS, and anti-GBP5 antibody.

(g) Sum intensity of GBP5 per cell area and **(h)** PAO1-LPS localized with GBP5 in BMDMs after PAO1 in **(f)** IF. N=7-8 images/genotype.

Statistics: (g & h) Student's t-test (a, c, d, & e) One-Way ANOVA. *P<0.05, **P<0.01, ***P<0.001, ****P<0.0001 Error bars represent SEM.

Fig. 4

Fig. 4: Caspase-11 activation, accentuated by GBPs, induces cell death and subsequent inflammation in *Polg^{D257A}* macrophages.

(a) Percent BMDMs dying after being given an siRNA control or siRNA to knock down caspase-11 or GBPs and then challenged with PAO1 from **(b)** representative IF. N=8-9 IF images/genotype/condition.

(c) Western blot analysis of lysates from BMDMs challenged with PAO1 after siRNA knock down of caspase-11.

(d) ELISA analysis of IL-1 cytokines in BMDM supernatant after being given an siRNA control or siRNA to knock down caspase-11 or GBPs and then challenged with PAO1. N= 3-4 biological replicates/genotype/condition.

(e) Quantification of caspase-11 associated with LPS in BMDMs after knock down of caspase-11 or GBPs by siRNA and then challenge with PAO1. N=5 IF images/genotype/condition.

(f) ELISA analysis of inflammatory cytokines in BMDM supernatant after being given an siRNA control or siRNA to knock down caspase-11or GBPs and then challenged with PAO1. N= 4 biological replicates/genotype/condition.

Statistics: (a, d, e & f) One-Way ANOVA. $*P<0.05$, $**P<0.01$, $***P<0.001$, $****P<0.0001$ Error bars represent SEM. Comparisons between WT siCtrl and WT siCaspase-11 or WT siGBP2,5 that are not shown are n.s.

Fig. 5

a BMDM, PAO1 MOI=5, 2 hpc

Phagolysosome | PAO1/phagolysosome | % Phagolyososomes with OMVs

Fig. 5: Elevated levels of caspase-11 binding to LPS contributes to potentiated pyroptosis in *Polg^{D257A}* macrophages.

(a) Quantitation of PAO1 and lysosomes in BMDMs from (b) TEM images. Black arrows highlight outer membrane derived vesicles. N= 9 images/condition.

(c) Quantification of capase-11 and GBP5 intensity in the area containing caspase-11, GBP5, and PAO1 from representative IF images (d). N=10-12 IF images/genotype/condition.

(i) ELISA analysis of IL-1 cytokines in BMDM supernatant after PAO1 or PAO1 plus anti-IL-1R blockade 2 hpi. N= 4 biological replicates/genotype/condition.

(j) Percent BMDMs dying after PAO1 and PAO1 plus anti-IL-1R blockade from (**Supplementary Fig. 9b**) representative IF. N=11-13 IF images/genotype/condition.

(k) ELISA analysis of IL-1 cytokines in BMDM supernatant after PAO1 or PAO1 plus anti-IL-1R blockade 24 hpi. N= 4 biological replicates/genotype/condition.

(l) ELISA analysis of inflammatory cytokines in BMDM supernatant after PAO1 or PAO1 plus anti-IL-1R blockade 24 hpi. N= 4 biological replicates/genotype/condition.

Statistics: (a, c, e, & g) Student's t-test (i, j, k, & l) Two-Way ANOVA. *P<0.05, **P<0.01, ***P<0.001,

Fig. 6

Fig. 6: Macrophages incorporating a deleterious patient mutation ($Polg^{R292C}$) exhibit mtDNA depletion, elevated IFN-I responses, increased pyroptosis, and hyper-inflammation after challenge with PAO1

(a) qRT-PCR analysis of ISGs in BMDMs after PAO1, N=3 technical replicates/genotype.

(b) ELISA analysis of inflammatory cytokines in BMDM supernatant after PAO1. N= 4 biological replicates/genotype/timepoint.

(c) Percent BMDMs dying after PAO1 and PAO1 plus caspase-1/11 inhibitor. N=7 IF images/genotype/condition.

(d) ELISA analysis of IL-1 cytokines in BMDM supernatant after PAO1 or PAO1 plus caspase-1/11 inhibitor. N= 2 biological replicates/genotype/condition.

(e) ELISA analysis of proinflammatory cytokines in BMDM supernatant after PAO1 or PAO1 plus caspase-1/11 inhibitor. N= 2 biological replicates/genotype/condition.

(f) Percent BMDMs dying after PAO1 and PAO1 plus anti-IFNAR blockade from. N=7 IF images/genotype/condition.

(g) ELISA analysis of IL-1 cytokines in BMDM supernatant after PAO1 or PAO1 plus anti-IFNAR blockade. N= 2 biological replicates/genotype/condition.

(h) ELISA analysis of pro-inflammatory cytokines in BMDM supernatant after PAO1 or PAO1 plus anti-IFNAR blockade. N= 2 biological replicates/genotype/condition.

(i) Quantification of caspase-11 in BMDMs after knock down by siRNA and PAO1 challenge. N=5 IF images/genotype/condition.

(k) Percent BMDMs dying after PAO1 and PAO1 plus caspase-11 knock down by siRNA from

(j) representative IF. N=8 IF images/genotype/condition.

(l) ELISA analysis of IL-1 cytokines in BMDM supernatant after PAO1 or PAO1 caspase-11

knock down by siRNA. N= 4 biological replicates/genotype/condition.

(m) ELISA analysis of pro-inflammatory cytokines in BMDM supernatant after PAO1 or PAO1

caspase-11 knock down by siRNA. N= 2 biological replicates/genotype/condition.

Statistics: (a) Students t-test. (b) Two-Way ANOVA. (c, d, e, f, g, h, i, k, l, & m) One-Way

ANOVA. *P<0.05, **P<0.01, ***P<0.001, ****P<0.0001 Error bars represent SEM.

Fig. 7

a

BAL cytokines, LegendPlex,
1×10^4 CFU PAO1, 6 hpi

IL-1β   TNFα   IL-23   IFNβ

● WT   ■ Polg^{D257A}

b

Ingenuity Pathway Analysis of RNA-Seq,
Polg^{D257A}/WT lung, 1×10^5 CFU PAO1, 6 hpi

Pathogen Induced Cytokine Storm Signaling Pathway
Interleukin-1 family signaling
IL-6 Signaling
Interferon gamma signaling
iNOS Signaling
Interferon alpha/beta signaling
Macrophage Classical Activation Signaling Pathway
Role of PRRs in Recognition of Bacteria and Viruses
Pyroptosis
Death Receptor Signaling
Macrophage Alternative Activation Signaling Pathway
Caspase activation via Death Receptors in the presence of ligand

z score

c

Polg^{D257A}/WT lung,
1×10^5 CFU PAO1, 6 hpi

IPA Upstream Regulators

d

Pyroptosis   Caspase activation

e

Lung, 1×10^5 CFU PAO1, 6 hpi

WT   Polg^{D257A}

CASPASE-11
GSDMD
GSDMD-NT
NLRP3
ASC
IFIT3
HSP60

Caspase-11

GSDMD-NT

f

CFU, 1×10^5 CFU PAO1, 6 hpi

Lung   Liver   Spleen

● WT   ■ Polg^{D257A}

g

CFU, 1×10^5 CFU PAO1-*algC*, 6 hpi

Lung   Liver   Spleen

● WT   ■ Polg^{D257A}

h

BAL Cytokines, ELISA, 1×10^5 CFU PAO1-*algC*, 6 hpi

IL-1β   TNFα   IFNβ   IL-6   IL-1α

● WT   ■ Polg^{D257A}

i

Lung, 1×10^5 CFU PAO1, 20 hpi

WT   Polg^{D257A}

100 μm   100 μm

j

1×10^5 CFU PAO1, 20 hpi

Total Inflammation   Neutrophilic Inflammation

● WT   ■ Polg^{D257A}

k

1×10^5 CFU PAO1, 20 hpi

WT   Polg^{D257A}

CASPASE-11
IL-1β
HSP60

l

1×10^5 CFU PAO1

Health Score

m

1×10^5 CFU PAO1

Survival

Fig. 7: Intratracheal instillation of PAO1 into lungs elicits greater inflammation, caspase-11 expression, and death of *Polg^{D257A}* mutant mice.

(a) LegendPlex pro-inflammatory cytokine analysis of bronchial alveolar lavage (BAL) 6 hours post infection (hpi) with 1×10^4 colony forming units (CFU) of PAO1. N=5 mice/genotype

(b) Ingenuity Pathway Analysis (IPA) of RNA-seq from lung showing Z-scores of pathways in *Polg^{D257A}* homozygous mutant BMDMs compared to wild-type (WT) after PAO challenge. N=3 biological replicates/genotype/condition.

(c) IPA Upstream Regulator analysis showing Z-scores in *Polg^{D257A}* homozygous mutant lungs compared to WT after 6 hour PAO1 challenge.

(d) Log^2 fold changes of representative genes in IPA pathways from (**a**) in *Polg^{D257A}* lungs compared to WT after PAO1.

(e) Protein expression or WT and *Polg^{D257A}* lung homogenates 6 hpi with 1×10^5 CFU of PAO1 and densitometry.

(f) Colony counts from homogenized tissue infected with 1×10^5 PAO1 for 6 hours. N=4 mice/genotype.

(g) Colony counts from homogenized tissue infected with 1×10^5 PAO1-*algC* for 6 hours. N=4 mice/genotype.

(h) ELISA analysis of pro-inflammatory cytokines in BAL of mice infected with 1×10^5 PAO1-*algC* for 6 hours. N=4 mice/genotype.

(i) Representative hematoxylin and eosin (H&E) staining of lung sections 20 hpi with 1×10^5 CFU of PAO1.

(j) Graphs of pathology scoring from H&E stained lungs in (**i**) histology. N=5 mice/genotype

(k) Protein expression or WT and *Polg^{D257A}* lung homogenates 20 hpi with 1×10^5 CFU of PAO1.

(l) Illness scoring of WT and *Polg^{D257A}* mice over PAO1 infection time course. N=4-5

mice/genotype

(m) Kaplan-Meier survival curve after infection with $1x10^5$ CFU of PAO1. N= 4-5

mice/genotype

Statistics: (a, e, f, g, h, & j) Students t-test (l) Two-Way ANOVA (m) Kaplan-Meijer Survival

Curve. *P<0.05, **P<0.01, ***P<0.001, ****P<0.0001 Error bars represent SEM.

Supplementary Fig. 1

Supplementary Fig. 1: *Polg^{D257A}* **macrophages exhibit elevated IFN-I and pro-inflammatory**

gene expression when challenged with LPS or PAO1.

(a) Log^2 fold changes of representative genes in IPA pathways from (**Fig. 1b**) in *Polg^{D257A}*

BMDMs compared to WT after challenge with LPS or PAO1. N=3 biological

replicates/genotype/condition.

(b) Gene Set Enrichment (GSEA) pathways of RNA-seq data in WT ang *Polg ^{D257A}* BMDMs

after challenge with PAO1. Heatmaps represent fold changes of each group relative to WT

PAO1 challenged BMDMs. N=3 biological replicates/genotype/condition.

(c) Ingenuity pathway analysis of proteomics from *Polg^{D257A}* (PAO1, 8hpc/Un)/WT (PAO1, 8

hpc/Un) BMDM proteins in upregulated pathways from **Fig 1g**. N=4 biological replicates

(d) Ingenuity Pathway Analysis (IPA) of RNA-seq from bone marrow-derived macrophages

(BMDMs) showing Z-scores of pathways related to cell death signaling in *Polg^{D257A}* mutant

BMDMs compared to wild-type (WT) after LPS or PAO1 challenge. N=3 biological

replicates/genotype/condition.

(e) Log^2 fold changes of representative genes in IPA pathways from **(d)** in *Polg^{D257A}* BMDMs

compared to WT after LPS or PAO1.

Supplementary Fig. 2

a BMDM, PAO1 MOI=5

b Polg^{D257A} BMDM, PAO1 MOI=5

WT+DMSO
Polg^{D257A}+DMSO
WT+nec-1
Polg^{D257A}+nec-1

c BMDM   Baseline   LPS priming + Nigericin

d

e BMDM, PAO1 MOI=5
Released LDH

Supplementary Fig. 2: *Polg*[D257A] **macrophages exhibit increased inflammatory cell death when exposed to bacteria.**

(**a**) Western blot of BMDM protein lysates after nec-1 priming plus PAO1 for protein involved in necroptosis.

(**b**) ELISA analysis of inflammatory cytokines in BMDM supernatant after PAO1 or PAO1 plus necroptosis inhibitors. N= 2 biological replicates/genotype/condition/timepoint.

(**c**) Representative IF of BMDMs unchallenged or primed with LPS for 4 hours and stimulated with nigericin for 1 hour to activate the NLRP3 inflammasome. Cells are stained with DAPI to mark the nucleus, CellMask to mark total cell volume, and anti-ASC antibody to mark puncta.

(**d**) Number of ASC puncta per cell in cells with high and low CellMask staining in (**a**) IF. N= 7 images/condition.

(**e**) Percent LDH released from BMDMs during PAO1 challenge. N=3 biological replicates/genotype/timepoint.

Statistics: (d) Students t-test (b & e) Two-way ANOVA. *P<0.05, **P<0.01, ***P<0.001, ****P<0.0001 Error bars represent SEM.

Supplementary Fig. 3

Supplementary Fig. 3: *Polg^{D257A}* **macrophages have a greater release of PAO1 that is not due to a deficiency in phagocytosis or PAO1 virulence factors.**

(a) PAO1 colony forming units obtained from BMDM supernatant or detergent lysates 2 and 4 hours post challenge (hpc). N=4 biological replicates/genotype/timepoint.

(b) Representative IF of BMDMs phagocytosing LPS coated, fluorescent microspheres. Cells were stained with DAPI to mark the nucleus, CellMask to label the cell volume, and anti-LAMP1 to mark phagolysosomes.

(c) Beads per cell quantified from **(e)** IF. N= 100 cells/genotype.

(d) Percent BMDMs dying after challenge with DH5α **(e)** IF. N=7 IF images/genotype.

(e) Representative IF of BMDMs stained with DAPI to mark the nucleus, CellMask to mark the total cell volume, and anti-LPS to mark DH5α bacteria. White arrows highlight dying cells characterized by condensed nuclei and loss of CellMask staining.

(f) ELISA analysis of IL-1β in BMDM supernatants after challenge with *Escherichia coli* strain DH5α. N= 4 biological replicates/genotype/timepoint.

(g) ELISA analysis of inflammatory cytokines in BMDM supernatants after challenge with *Escherichia coli* strain DH5α. N= 4 biological replicates/genotype/timepoint.

Statistics: (c & d) Students t-test (a, f & g) Two-way ANOVA. *P<0.05, **P<0.01, ***P<0.001, ****P<0.0001 Error bars represent SEM.

Supplementary Fig. 4

Supplementary Fig. 4: Assessment of roles for canonical inflammasome activity and

metabolism in driving increased cell death and pro-inflammatory cytokine secretion in

***Polg^{D257A}* macrophages.**

(a) Cytation5 imaging quantification of PI positive cells in BMDMs after LPS priming for 4

hours. N= 3 biological replicates, 5 technical replicates/genotype.

(b) Cytation5 imaging quantification of PI positive cells in BMDMs after LPS priming plus PA-

flagellin transfection. N= 5 biological replicates/genotype.

(c) ELISA analysis of IL-1β in BMDM supernatant after LPS priming plus PA-flagellin

transfection. N= 2 biological replicates.

(d) Cytation5 imaging quantification of PI positive cells in BMDMs after LPS priming plus

ATP. N= 6 biological replicates/genotype.

(e) ELISA analysis of IL-1β in BMDM supernatant after LPS priming plus ATP. N= 2 biological

replicates.

(f) Western blot of BMDM protein lysates after LPS priming plus ATP.

(g) Cytation5 imaging quantification of PI positive cells in BMDMs after LPS priming plus

nigericin. N= 6 biological replicates/genotype.

(h) ELISA analysis of IL-1β in BMDM supernatant after LPS priming plus nigericin. N= 2

biological replicates.

(i) Western blot of BMDM protein lysates after LPS priming plus nigericin.

(j) Representative TEM images of mitochondria from BMDMs.

(k) & **(l)** Seahorse analysis of BMDMs unchallenged or treated with overnight LPS.

Extracellular acidification rate (ECAR) **(k)** and oxygen consumption rate (OCR) **(l)** are plotted.

N=3 biological replicates and 3-4 technical replicates.

(m) Quantification of Seahorse OCR measurements. N=3 biological replicates and 3-4 technical

replicates.

Statistics: (b, e, & h) Two-Way ANOVA (m) One-Way ANOVA (a, c, f, & i) Students t-test.

*$P<0.05$, **$P<0.01$, ***$P<0.001$, ****$P<0.0001$ Error bars represent SEM.

Supplementary Fig. 5

Supplementary Fig. 5: The NLRP3 inflammasome is downstream of caspase-11 activation in *Polg*^{D257A} macrophages.

(**a**) Cytation5 imaging quantification of PI positive cells in BMDMs with PAO1 or PAO1 plus MCC950. N= 6 biological replicates/genotype/timepoint.

(**c**) Percent BMDMs dying after PAO1 and PAO1 plus NLRP3 inhibitor, MCC950, from (**b**) representative IF. N=8-9 IF images/genotype/condition.

(**d**) ELISA analysis of pyroptotic cytokines in BMDM supernatant after PAO1 or PAO1 plus NLRP3 inhibitor, MCC950. N= 2 biological replicates/genotype/condition.

(**e**) Western blot of GSDMD localization during PAO1 challenge.

(**f**) RNA-seq from BMDMs showing induction of Gbps and Caspase-11 RNA 6 hpc with PAO1. N=3 biological replicates/genotype/condition.

(**g**) GBP expression in *Polg*^{D257A} (8hr PAO1/Un)/WT (8hr PAO1/Un) BMDMs by proteomics. N=4 biological replicates/genotype/condition.

(**h**) Representative IF of unchallenged BMDMs stained with DAPI to mark the nucleus and anti-GBP5 antibody.

(**i**) Sum intensity of GBP5 per cell area in resting BMDMs in (**h**) IF. N=5-6 images/genotype

Statistics: (f, & i) Student's t-test (c, & d) One-way ANOVA (a) Two-way ANOVA. *P<0.05, **P<0.01, ***P<0.001, ****P<0.0001 Error bars represent SEM.

Supplementary Fig. 6

Supplemental Fig. 6: The PAO1 *algC* mutant displays decreased levels of LPS and does not robustly engage the caspase-11 inflammasome or induce potentiated cell death and hyperinflammation in *Polg*[D257A] macrophages.

(a) qRT-PCR quantification of siRNA knock down efficiency in BMDMs. N=3 technical replicates/genotype.

(b) Quantification of LPS associated with PAO1 and total LPS staining in BMDMs after PAO1 or PAO1- *algC* challenge by IF. N=6-7 IF images/genotype/condition.

(c) Percent BMDMs dying after PAO1 and PAO1- *algC* from **(d)** representative IF. N=10 IF images/genotype/condition.

(e) Western blotting of proteins involved in pyroptosis in BMDMs after PAO1 or PAO1- *algC* challenge.

(f) ELISA analysis of IL-1 cytokines in BMDM supernatant after PAO1 or PAO1-*algC*. N= 2 biological replicates/genotype/condition.

(g) ELISA analysis of proinflammatory cytokines in BMDM supernatant after PAO1 or PAO1- *algC*. N= 2 biological replicates/genotype/condition.

Statistics: (a & b) Student's t-test (c, f, & g) One-way ANOVA. $*P<0.05$, $**P<0.01$, $***P<0.001$, $****P<0.0001$ Error bars represent SEM.

Supplementary Fig. 7

a BMDM, Baseline

Nucleus | Whole Cell

WT ■ Polg^D257A

b WT | Polg^D257A

2 μm | 2 μm

c BMDM, PAO1 MOI=5, 2 hpc
Nucleus

d BMDM, PAO1 MOI=5, 8 hpc
Bacteria/Cell | Bacteria/Phagolysosome

WT ■ Polg^D257A

WT
1μm

e Polg^D257A

1μm | 400 nm

Polg^D257A

1μm | 400 nm

Polg^D257A

1μm | 400 nm

f Top 25 Upregulated
PAO1 Proteins 8 hpc
Polg^D257A/WT BMDM

pvdF, dcd, gltP, pvdA, ptsN, atpA, exaC, acpP1, atpF, hupB, rpmC, gor, rpmE, atpH, lldD, hmp, lptF, katA, rpsA, braC, atpD, sdhA, eagT6, sucC, lptD

log2 ratio

2 | 8 hpc

g BMDM, PAO1 MOI=5, 24 hpc

Nitric Oxide

WT ■ Polg^D257A

24 hpc

h BMDM, PAO1, 2 hpc
Caspase-11with PAO1-algC

WT ■ Polg^D257A

i WT + PAO1-algC | Polg + PAO1-algC

Merge | CASPASE-11 | Merge | CASPASE-11

5 μm PAO1-AlgC | GBP5 | 5 μm PAO1-AlgC | GBP5

Supplemental Fig. 7: *Polg^{D257A}* macrophages have more PAO1 and PAO1 proteins present

at late timepoints.

(a) Quantitation of nucleus and cellular area of BMDMs from **(b)** TEM images of BMDMs at

rest. N= 14-15 images/condition.

(c) Quantitation of nuclear area of BMDMs from **(Fig. 6b)** TEM images. N= 9 images/condition.

(d) Quantification of bacteria in BMDMs and their lysosomes 8 hours after PAO1 by **(e)** TEM.

N=15-20 cells/genotype.

(f) Proteomic analysis of PAO1 proteins in *Polg^{D257A}* homozygous mutant BMDMs compared to

WT 8 hours after PAO1 challenge. N=4 biological replicates/genotype/condition.

(g) Quantification of nitric oxide in BMDMs 24 hpc with PAO1. N=3 biological

replicates/genotype.

(h) Quantification of capase-11 associated with PAO1-GFP or PAO1-*algC* mutant from

representative IF images **(i)**. N=6 IF images/genotype/condition.

Statistics: (a, c, d, g, & h) Students t-test. *P<0.05, **P<0.01, ***P<0.001, ****P<0.0001 Error

bars represent SEM.

Supplementary Fig. 8

a BMDM, LPS + tLPS, 1.5 hpc

LPS Area — ns

LPS Intensity — ns

b WT · CellMask/DAPI/LPS · Polg^{D257A} · CellMask/DAPI/LPS · 10 µm

c BMDM, LPS + tLPS, 7 hpc

LPS Area — ****

LPS Intensity — ****

d WT · CellMask/DAPI/LPS · Polg^{D257A} · CellMask/DAPI/LPS · 10 µm

e BMDM, LPS + tLPS, 7 hpc

Caspase-11 — ****

Caspase-11 with LPS — **

WT · Polg^{D257A}

f WT · Merge · CASPASE-11 · Polg^{D257A} · Merge · CASPASE-11 · DAPI · LPS · 5 µm

g BMDM, LPS, 2 hpc · WT · Polg^{D257A}

LPS Area — ns

LPS Intensity — ns

h WT · CellMask/DAPI/LPS · Polg^{D257A} · CellMask/DAPI/LPS · 5 µm

i BMDM, LPS, 48 hpc

LPS Area — ***

LPS Intensity — ***

j WT · CellMask/DAPI/LPS · Polg^{D257A} · CellMask/DAPI/LPS · 5 µm

k BMDM, LPS, 48 hpc

IL-1β · WT · Polg^{D257A}

Limit of detection

Supplemental Fig. 8: *Polg*^{D257A} **macrophages uptake similar levels of LPS but maintain higher levels of LPS and more caspase-11 association with LPS at later challenge timepoints.**

(a) Quantification of LPS per cell in BMDMs after 4 hours LPS priming then 1.5 hours LPS transfection from **(b)** IF. N=10-11 IF images/genotype/condition.

(c) Quantification of LPS per cell in BMDMs after 4 hours LPS priming then 7 hours LPS transfection from **(d)** IF. N=10-12 IF images/genotype/condition.

(e) Quantification of caspase-11 intensity and caspase-11 associated with LPS in BMDMs after 4 hours LPS priming then 7 hours LPS transfection from **(f)** IF. N=7 IF images/genotype/condition.

(g) Quantification of LPS per cell in BMDMs 2 hours post LPS challenge from **(h)** IF. N=8-9 IF images/genotype/condition.

(i) Quantification of LPS per cell in BMDMs 48 hours post LPS challenge from **(j)** IF. N=10 IF images/genotype/condition.

(k) ELISA analysis of IL-1β in BMDM supernatant after LPS. N= 2 biological replicates/genotype/timepoint.

Statistics: (a, c, e, & g) Students t-test (k) Two-way ANOVA. *P<0.05, **P<0.01, ***P<0.001, ****P<0.0001 Error bars represent SEM.

Supplementary Fig. 9

a Immune signaling in BMDM, PAO1 MOI=5

b BMDM, PAO1 MOI=5, 2 hpc

c BMDM, PAO1 MOI=5, 1.5 hpc

Supplementary Fig. 9: *Polg*^{D257A} **BMDMs exhibit enhanced NF-κB signaling that is**

downstream of the IL-1R when challenged with PAO1.

(a) Western blot of signaling in BMDMs during a PAO1 time course.

(b) Representative images of cell death quantitation in **Fig. 5f**.

(c) Western blot analysis of lysates from BMDMs challenged with PAO1 and PAO1 plus anti-

IL-1R blockade 1.5 hpc.

Supplementary Fig. 10

Supplemental Fig. 10: Mice incorporating a deleterious patient mutation ($Polg^{R292C}$) exhibit

mtDNA depletion and elevated IFN-I responses.

(a) Generation of the $Polg^{R292C}$ mouse.

(b) Chromatograph from Sanger sequencing of DNA from heterozygous $Polg^{R292C}$ mouse

verifying germline incorporation of mutation.

(c) qPCR analysis of mtDNA abundance in tissues. N= 4-6 mice/genotype.

(d) Western blot analysis of ISG, IFIT3, in lung lysates from $Polg^{R292C}$ mice.

(e) Western blot analysis of ISGs in spleen lysates from $Polg^{R292C}$ mice.

Statistics: (c) Student's t-test. *P<0.05, **P<0.01, ***P<0.001, ****P<0.0001 Error bars

represent SEM.

Supplementary Fig. 11

a BMDM, unchallenged

Non-numpt

D-loop CytB

b BMDM, LPS, 24 hpc

c BMDM, LPS, 24 hpc

Number of Nucleoids   Nucleoid DNA Intensity   Nucleoid Area

d BMDM, unchallenged

Stat1 *Ifit1* *Zbp1* *Gbp2* *Gbp4* *Gbp9*

e BMDM, LPS, 8 hpc

Stat1 *Ifit1* *Zbp1* *Gbp2* *Gbp4* *Gbp9* *Casp11*

f ECAR

g OCR

h

Maximal Respiration   Basal Respiration   Glycolytic Capacity

Supplemental Fig. 11: Macrophages incorporating a deleterious patient mutation

(***Polg*^{R292C}**) exhibit mtDNA depletion, elevated IFN-I responses, and altered metabolism.**

(a) qPCR analysis of mtDNA abundance in BMDMs. N= 3 technical triplicates/genotype.

(b) Representative IF of nucleoids in BMDMs stained with anti-PDH antibody to label

mitochondria, anti-DNA antibody to label nucleoids, and anti-TFAM antibody.

(c) Quantification of nucleoids per cell, mean intensity of DNA signal per nucleoid, and nucleoid

area per cell from BMDMs in **(b)** IF. N=40-50 cells/genotype

(d) qRT-PCR analysis of ISGs in unchallenged *Polg*^{R292C} BMDMs. N=3 biological replicates, 3

technical replicates/genotype.

(e) qRT-PCR analysis of ISGs in *Polg*^{R292C} BMDMs 8 hours post challenge (hpc) with LPS. N=

3 technical replicates/genotype.

(f) & **(g)** Seahorse analysis of BMDMs unchallenged or treated with overnight LPS.

Extracellular acidification rate (ECAR) **(f)** and oxygen consumption rate (OCR) **(g)** are plotted.

N=2 biological replicates and 5-6 technical replicates.

(h) Quantification of Seahorse OCR measurements. N=2 biological replicates and 4-5 technical

replicates.

Statistics: (a, c, d, & e) Students t-test (h) One-Way ANOVA. *$P<0.05$, **$P<0.01$, ***$P<0.001$,

****$P<0.0001$ Error bars represent SEM.

Supplementary Fig. 12

a BMDM, PAO1 MOI=5, 2 hpc

Supplemental Fig. 12: AcYVAD and anti-IFNAR blockade reduces cell death in *Polg^{R292C}*

macrophages

(a) Representative IF of dying cells characterized by condensed nuclei and loss of cell mask

staining (white arrows) after PAO1 and PAO1 plus caspase-1/11 inhibitor, AcYVAD, quantified

in **Fig. 6c**.

(b) Representative IF of dying cells characterized by condensed nuclei and loss of cell mask

staining (white arrows) after PAO1 and PAO1 plus anti-IFNAR blockade quantified in **Fig. 6f**.

Supplementary Fig. 13

a Lung, unchallenged

WT — Polg^D257A

○ Eosin⁺
○ Eosin⁻

b Lungs, unchallenged

Genotype	Mouse ID	Total Inflammation	Neutrophilic Inflammation	Mononuclear Inflammation
WT	1	minimal	0	1
	2	minimal	0	1
	3	none	0	0
Polg^D257A	1	none	0	0
	2	none	0	0
	3	none	0	0

c ● WT ■ Polg^D257A

d Lungs, unchallenged

STAT1
IFIT3
HSP60

e BAL Cytokines, 0.5 mg/kg LPS

IL-1β   TNFα   IL-6

━ WT ━ Polg^D257A

Supplementary Fig. 13: *Polg* **mutant mice exhibit elevated IFN-I and inflammatory**

responses to LPS, but do not exhibit basal differences in lung morphology or immune cell

abundance.

(a) Representative histology images of hematoxylin and eosin (H&E) staining of WT and

Polg^{D257A} lungs, with an overlay of eosin positive cells using QuPath software.

(b) Pathology assessment of unchallenged lungs.

(c) QuPath analysis of eosin$^+$ cells in the lungs from **(c)**. N=3 mice/genotype

(d) Protein expression of ISGs in unchallenged WT and *Polg*^{D257A} lungs homogenates.

(e) ELISA analysis of inflammatory cytokines in bronchioalveolar lavage of LPS challenged WT

and *Polg*^{D257A} lungs. N=2 mice/genotype/timepoint.

Statistics: (c) Students t-test (e) Two-Way ANOVA. *P<0.05, **P<0.01, ***P<0.001,

****P<0.0001 Error bars represent SEM.

Supplementary Fig. 14

Supplementary Fig. 14: Intratracheal instillation of PAO1 alters immune cell composition

in *Polg* mutant mouse lungs.

(a) Weight change in WT and $Polg^{D257A}$ mice after infection with $1x10^4$ PAO1. N=4

mice/genotype.

(b) CD45 expression on cells from bronchioalveolar lavage (BAL) from uninfected and PAO1

infected WT and $Polg^{D257A}$ mice. N=5-9 mice/genotype/timepoint.

(c) Gates used to quantify CD45 expression in **(b)**.

(d) Percent of $CD45^+$ cells expressing CD11c from the BAL of uninfected and PAO1 infected

lungs. N=5-9 mice/genotype/timepoint.

(e) Percent of $CD45^+$ cells expressing SiglecF from the BAL of uninfected and PAO1 infected

lungs. N=5-9 mice/genotype/timepoint.

(f) Percent of $CD45^+$ cells expressing MHCII from the BAL of uninfected and PAO1 infected

lungs. N=5-9 mice/genotype/timepoint.

(g) Percent of $CD45^+$ cells expressing F4/80 from the BAL of uninfected and PAO1 infected

lungs. N=5-9 mice/genotype/timepoint.

(h) Percent of $CD45^+$ cells expressing CD11b from the BAL of uninfected and PAO1 infected

lungs. N=5-9 mice/genotype/timepoint.

(i) Percent of $CD45^+$, $CD11b^-$ cells expressing CD3 from the BAL of uninfected and PAO1

infected lungs. N=5-9 mice/genotype/timepoint.

(j) Percent of $CD45^+$, $CD11b^-$ cells expressing CD19 from the BAL of uninfected and PAO1

infected lungs. N=5-9 mice/genotype/timepoint

(k) Percent of $CD45^+$ expressing Ly6G from the BAL of uninfected and PAO1 infected lungs.

N=5-9 mice/genotype/timepoint.

(l) Percent of CD45$^+$ cells expressing Ly6C from the BAL of uninfected and PAO1 infected

lungs. N=5-9 mice/genotype/timepoint.

(m) CD45$^+$ cells with cell associated TNFα expression from BAL of uninfected and PAO1

infected lungs. N=5-9 mice/genotype/timepoint.

Statistics: (a) Two-way ANOVA (b, & d-m) Students t-test. *P<0.05, **P<0.01, ***P<0.001,

****P<0.0001 Error bars represent SEM.

1013

Supplementary Fig. 15

a BAL cytokines, LegendPlex, 1x10⁴ CFU PAO1, 6 hpi

b BAL Cytokines, LegendPlex 1x10⁴ CFU PAO1, 48 hpi

c Lung, 1x10⁴ CFU PAO1, 48 hpi

1014

Supplementary Fig. 15: A non-lethal dose of PAO1 induces hyperinflammation and activation of pyroptosis in the lungs of *Polg^{D257A}* mice.

(**a**) LegendPlex pro-inflammatory cytokine analysis of bronchial alveolar lavage (BAL) 6 hours post infection (hpi) with 1×10^4 colony forming units (CFU) of PAO1. N=5 mice/genotype

(**b**) LegendPlex pro-inflammatory cytokine analysis of bronchioalveolar lavage (BAL) from WT, *Polg^{D257A}*, WT IFNAR$^{-/-}$, and *Polg^{D257A}* IFNAR$^{-/-}$ lungs infected with 1×10^4 PAO1 for 48 hours. All undefined comparisons are not significant. N=5-6 mice/genotype.

(**c**) Western blot of ISGs and proteins involved in pyroptosis from homogenized lung tissue from mice infected with 1×10^4 PAO1 for 48 hours.

Statistics: (a) Students t-test (b) One-way ANOVA. *P<0.05, **P<0.01, ***P<0.001, ****P<0.0001 Error bars represent SEM.

Supplementary Fig. 16

a Lung, 1x10⁴ CFU PAO1, 7 dpi + 24 hpi

b

c BAL Cytokines, LegendPlex, 1x10⁴ CFU PAO1, 7 dpi + 24 hpi

d CFU, 1x10⁴ CFU PAO1, 7 dpi + 24 hpi

Supplementary Fig. 16: *Polg^D257A* mice exhibit hyperinflammation in response to infection

and reinfection with PAO1.

(a) Representative histology images of hematoxylin and eosin (H&E) staining of WT and

Polg^D257A lungs after initial infection with 1×10^4 PAO1, followed by 7 days rest, then reinfection

with 1×10^4 PAO1. Overlay of eosin positive cells was done using QuPath software.

(b) QuPath analysis of eosin$^+$ cells in the lung from **(a)** histology. N=6 mice/genotype.

(c) LegendPlex pro-inflammatory cytokine analysis of BAL from WT and *Polg^D257A* lungs after

initial infection with 1×10^4 PAO1, followed by 7 days rest, then reinfection with 1×10^4 PAO1.

N=5 mice/genotype.

(d) Colony counts from homogenized tissue after initial infection with 1×10^4 PAO1, followed by

7 days rest, then reinfection with 1×10^4 PAO1. N=5 mice/genotype.

Statistics: (b, c, & d) Students t-test. *P<0.05, **P<0.01, ***P<0.001, ****P<0.0001. Error bars

represent SEM.

1040

1041

Supplementary Fig. 17

a Polg^D257A/WT lung, RNA sequencing, 1x10^5 CFU PAO1, 6 hpi

Pathogen Induced Cytokine Storm

IL-1 Signaling

IFN gamma Signaling

b Ingenuity Pathway Analysis of RNA-Seq, Polg^R292C/WT lung, 1x10^6 CFU PAO1, 6 hpi

Pathogen Induced Cytokine Storm Signaling Pathway
Interferon alpha/beta signaling
ISGylation Signaling Pathway
Cytosolic sensors of pathogen-associated DNA
Macrophage Classical Activation Signaling Pathway
iNOS Signaling
Pyroptosis Signaling Pathway
IL-6 Signaling
Interferon Signaling
Immunogenic Cell Death Signaling Pathway
Death Receptor Signaling
TNFR1 Signaling
IL-1 Signaling
STAT3 Pathway

c Polg^R292C/WT lung, RNAseq 1x10^6 CFU PAO1, 6 hpi
IPA Upstream Regulators

d Polg^R292C/WT lung, RNAseq 1x10^6 CFU PAO1, 6 hpi
Interferon Signaling

Pyroptosis

e Lung, 1x10^5 CFU PAO1, 6 hpi

WT   Polg^D257A

CV-ATPA
CIII-UQCRC2
CIV-MTCO1
CII-SDHB
CI-NDUFB8
HSP60

f Plasma Cytokine, 1x10^5 CFU PAO1, 6 hpi

IL-6   IL-12p70

g Liver, 1x10^5 CFU PAO1, 6 hpi

WT   Polg^D257A

IFIT3
HSP60

WT   Polg^D257A

CV-ATPA
CIII-UQCRC2
CIV-MTCO1
CI-NDUFB8

h Lung, 1x10^5 CFU PAO1-algC, 6 hpi

WT   Polg^D257A

GSDMD
GSDMD-NT
CASPASE-11
NLRP3
IFIT3
HSP60

i Histology analysis, 1x10^5 CFU PAO1, 20 hpi

Genotype	ID	Total Inflammation	Neutrophilic Inflammation	Mononuclear Inflammation
WT	1	minimal	1	1
	2	minimal	0	1
	3	mild	2	1
	4	moderate	3	0
	5	mild	2	0
PolG	1	mild/moderate	2	0
	2	moderate	3	0
	3	moderate	3	0
	4	moderate	3	2
	5	severe	4	1

j CFU, 1x10^5 CFU PAO1, 20 hpi

Lung   Liver   Spleen

k CFU, 1x10^5 CFU PAO1, 48 hpi

Lung   Liver   Spleen

l Spleen, 1x10^5 CFU PAO1, 48 hpi

WT   Polg^D257A

IFIT3
GAPDH

m Liver, 1x10^5 CFU PAO1, 48 hpi

WT   Polg^D257A

IFIT3
GAPDH

1042

Supplementary Fig. 17: *Polg* **mutant mice exhibit increased IFN-I, pyroptotic cell death, and pro-inflammatory responses after challenge with PAO1.**

(a) Fold changes of representative genes in IPA pathways from **Fig. 7b** in $Polg^{D257A}$ lungs compared to WT after LPS or PAO1.

(b) Ingenuity Pathway Analysis (IPA) of RNAseq from lungs showing Z-scores of pathways in $Polg^{R292C}$ lungs compared to WT after PAO1 challenge. N=3 biological replicates/genotype/condition.

(c) IPA Upstream Regulator analysis showing Z-scores in $Polg^{R292C}$ lung compared to WT after PAO1 challenge.

(d) Log^2 fold changes of representative genes in IPA pathways from (b) in $Polg^{R292C}$ lungs compared to WT after PAO1.

(e) Western blot of proteins involved in oxidative phosphorylation in the lung 6 hours post infection (hpi) with $1x10^5$ colony forming units (CFU) of PAO1.

(f) LegendPlex of circulating pro-inflammatory cytokine in the plasma 6 hpi with $1x10^5$ CFU of PAO1. N=4 mice/genotype

(g) Western blot of ISG and proteins involved in oxidative phosphorylation in the liver 6 hpi with $1x10^5$ CFU of PAO1.

(h) Western blot of ISG and proteins involved in pyroptosis in the lung 6 hours post infection with $1x10^5$ CFU of PAO1-*algC*.

(i) Summary of pathologist scoring of H&E stained lungs infected with $1x10^5$ CFU PAO1 at 20 hpi. N=5 mice/genotype.

(j) Colony counts from homogenized tissue infected with $1x10^5$ CFU PAO1 at 20 hpi. N=5 mice/genotype.

(k) Colony counts from homogenized tissue infected with $1x10^5$ CFU PAO1 at 48 hpi. N=4

mice/genotype.

(l) Western blot of ISGs from homogenized liver and **(m)** spleen from mice infected with $1x10^5$

CFU PAO1 at 48 hpi.

Statistics: (f, j, & k) Students t-test. *P<0.05, **P<0.01, ***P<0.001, ****P<0.0001. Error bars

represent SEM.

1072

Supplementary Fig. 18: Elevated IFN-I primes macrophages to undergo caspase-11 non-canonical inflammasome activation and drives hyperinflammatory signaling, resulting in lung injury in mice with *Polg*-related mitochondrial dysfunction.

Mitochondrial DNA instability in exonuclease-deficient *Polg* mutant mice triggers chronic IFN-I signaling that enhances expression of caspase-11, GBPs, and other ISGs **(a)**. Inhibitors and genetic approaches (maroon arrows) revealed that GBPs accumulate on bacteria and augment activation of the caspase-11 non-canonical inflammasome, resulting in pyroptosis and the GSDMD-dependent release of IL-1α **(b)**. IL-1α signals through the IL-1R to further upregulate the expression of inflammasome mediators including NLRP3 and pro-IL-1β **(c)**, leading to the release of mature IL-1β **(d)** and DAMPs **(e)** that feed forward to sustain and enhance IFN-I and IL-1R signaling, elevating release of other pro-inflammatory cytokines such as TNFα and IL-6 **(f)**. Elevated rates of pyroptotic cell death and hyperinflammation in the lung of *Polg* mutant mice contribute to acute lung injury and systemic inflammation resulting in decreased survival after infection.

Acknowledgements

We thank Robbie Moore at the Texas A&M College of Medicine Analytical Cytometry Core and Will Schott and Danielle Littlefield in the JAX Flow Cytometry Service for technical assistance. We thank Pete Finger in the JAX Histopathology and Microscopy Sciences Service for assistance preparing macrophage samples for TEM imaging. We acknowledge the JAX Center for Precision Genetics (JCPG) for generating the *Polg*$^{R292C/R292C}$ mouse line. We would like to acknowledge the Mass Spectrometry and Protein Chemistry Service within Protein Sciences at The Jackson Laboratory for their guidance and support with the mass spectrometry analyses

1099   performed in this study. Finally, we thank Vijay Rathinam for protocols and helpful discussions

1100   on non-canonical inflammasome biology.

1101

Funding

This research was supported by awards W81XWH-17-1-0052 to A.P.W. and W81XWH-20-1-

0150 to C.C. and A.P.W. from the Office of the Assistant Secretary of Defense and for Health

Affairs through the Peer Reviewed Medical Research Programs. Additional support was

provided by NIH grants R01HL148153 to A.P.W., U54OD030187 to S.A.M, and a gift from the

POLG Foundation. J.J.V. was supported by NIH NRSA F31AI179168 and Y. Lei was supported

by American Heart Association Predoctoral Fellowship grant 825908.

1109

Author contributions

A.P.W and J.J.V. designed experiments, analyzed data, and wrote the manuscript. Y.L. assisted

with animal experiments and sample processing. M.R. assisted with sample processing. C.G.M.

performed inflammasome challenge experiments. J.W. assisted with pathology scoring of

histology samples. T.J.S and B.R.H. performed proteomics and analysis. C.P. and R.S.

maintained mouse lines and generated BMDMs. S.C.K. and S.A.M. generated the $Polg^{R292C}$

mouse model and provided samples. P.J.M. and C.L.C. assisted with data analysis and

interpretation. A.P.W. conceived the project and provided overall direction.

1118

1119   **Competing interests**

1120   The authors declare that they have no competing interests.

1121

 Data and materials availability

 All data needed to evaluate the conclusions drawn herein are present in the paper and

 Supplementary Materials. Raw data will be provided upon request. The RNA sequencing and

 proteomic datasets generated through this work will be deposited in the Gene Expression

 Omnibus or PRIDE database, respectively, after acceptance of the paper for publication.

 Methods

 Mouse Husbandry and Strains

 C57BL/6J (JAX Stock # 000664), $Polg^{D257A/D257A}$ (JAX Stock # 017341), and $Ifnar1^{-/-}$

 (JAX Stock # 028256) mice were obtained from The Jackson Laboratory and bred and

 maintained in multiple vivaria at Texas A&M University and The Jackson Laboratory.

 Experimental $Polg^{+/+}$ (WT) and $Polg^{D257A/D257A}$ littermates were obtained from $Polg^{D257A/+}$

 maintenance colony crosses of male $Polg^{D257A/+}$ to female $Polg^{+/+}$. This breeding strategy was

 adopted to limit the accumulation and maternal transmission of mutant mtDNA molecules to

 heterozygous breeders. $Polg^{D257A/D257A}$ mice and littermate controls used for macrophage

 experiments were aged for 12-16 months. $Polg^{D257A/D257A}$ mice and littermate controls used for in

 vivo infections were aged from 7-12 months.

 The $Polg^{R292C/R292C}$ mouse line (B6N(Cg)-$Polg^{em5(F290F,R292C)Lutzy/Mmjax}$) was generated

 using CRISPR/Cas9 endonuclease-mediated genome editing in mouse embryonic stem cells

 (mESCs). A single guide RNA (CTCCCATCCACAGGCTGCGC) targeting exon 4 of $Polg$ was

 cloned into pX459, which expresses Cas9 nuclease. This plasmid was co-delivered to JM8A3

 ESCs via electroporation with a single-stranded oligonucleotide donor (ssODN) that included the

 missense R292C (CGC>TGC) and an additional silent mutation (F290F; TTC>TTT) to reduce

recutting of the targeted allele[74]. Following puromycin selection, resistant clones were screened

for proper targeting and to exclude integration of the pX459 plasmid. Multiple clones were

selected for expansion and karyotyping, and euploid clones were injected into C57BL/6J-albino

blastocysts (B6(Cg)-Tyrc-2J/J; JAX STOCK# 000058) to produce chimeras. Chimeras were bred

to C57BL/6NJ (JAX Stock #005304) mice and a single germline clone was selected for

expansion and characterization. The line is available through the MMRRC repository (071656-

JAX). All $Polg^{R292C/R292C}$ mice and littermate controls used for macrophage experiments were

aged for 5-8 months. $Polg^{R292C/R292C}$ mice and littermate controls used for in vivo experiments

were aged for 3-4 months. Breeding colonies of $Polg^{R292C}$ mice were maintained similarly to

$Polg^{D257A}$ models. All animal experiments were approved by the Institutional Animal Care and

Use Committees (IACUC) at Texas A&M University and The Jackson Laboratory.

1156

In vivo PAO1 infection

7-12 month old mice were moved into BSL2 procedure space and allowed to acclimate

for 1-2 days. Mice were anesthetized with an intraperitoneal injection of 100-10 mg/kg

Ketamine-Xylazine cocktail. Mice were then intubated with a 24G catheter. Sterile PBS (sham),

0.5 mg/kg LPS (E. Coli serotype 0111:B4; Invivogen), or the specified dose of PAO1-GFP was

pipetted into the catheter in a 25-30 µL volume. A bolus of air was delivered using a syringe to

ensure the dose evenly coated the lungs. Mice were kept warm until fully recovered from

anesthesia and monitored daily until experimental timepoints were reached.

1165

Bronchial alveolar lavage fluid collection

Mice were euthanized using intraperitoneal injection of 200 mg/kg Pentobarbital Sodium

to preserve the lung environment and structure of the tracheal. Upon confirmation of death the

chest cavity and throat were opened. The trachea was canulated using a 23G needle threaded

with tubing and the lung was washed with 3mL cold Hanks' Balanced Salt Solution (HBSS,

Thermo Fisher Scientific, 14175-129) containing 100 µM EDTA (VMR Life Science, E522) and

used in subsequent flow cytometry and LegendPlex assays.

1173

Flow Cytometry

Bronchial alveolar lavage fluid (BAL) was spun at 350 g for 5 minutes to pellet the cells.

Cells were transferred to a V bottom plate for staining. Cells were stained with viability dye then

Fc receptors were blocked with anti-mouse CD16/CD32 (2.4G2) antibody. Surface proteins were

stained with antibodies specified in Table 6. Cells were permeabilized with Foxp3/Transcription

Factor Staining Buffer Kit (TNB-0607-KIT, Tonbo) and stained with antibodies against

intracellular proteins. Flow cytometry analysis was done using a 5L Cytek Aurora.

1181

1182   **Colony forming unit counts from macrophages and tissues**

1183   BMDMs or tissues were homogenized in 0.1% NP40 in sterile PBS using an immersion

1184   homogenizer for 15 seconds per tissue. Serial dilutions were plated on soy agar containing 100

1185   µg/mL carbenicillin. Plates were incubated at 37°C overnight and colonies were counted.

1186

Histology

Tissues were collected in cassettes, then were washed in PBS, incubated for 24 hours in

10% formalin, and transferred to 70% ethanol. Tissue embedding, sectioning, and staining were

performed at AML Laboratories in Jacksonville, FL or at the histology core at Texas A&M

University. Images of hematoxylin and eosin (H&E) staining were collected using an Olympus

slide scanner with a 20X objective. Slides were analyzed and scored by a pathologist, Dr. Jessica

Wong, at the Jackson Laboratory in Sacramento, CA. Briefly, semi-quantitative histologic

scoring from 0 to 4 severity was performed on neutrophilic and mononuclear cell inflammation.

0 = no inflammation, 1 = minimal inflammation including individual cells or small clusters, 2 =

mild inflammation including small clusters and increased perivascular and peribronchiolar

cuffing, 3 = moderate inflammation including multifocal to coalescent areas and perivascular

cuffing, and 4 = severe inflammation including coalescent to patchy-diffuse inflammation.

1199

Body Condition Scoring

Mice were monitored three times daily and were assigned a body condition score based

on an additive score of appearance, behavior, hydration, and respiration. A score of 0

corresponds to a healthy mouse and a score of 10 corresponds to a moribund animal requiring

euthanasia. Animals were given an appearance score of 0 if normal, 1 if they lacked grooming or

ruffled fur, 2 if they displayed piloerection, and 3 if they displayed hunched posture. Animals

were given a behavior score of 0 if they were socially active, 1 if they had decreased mobility, 2

if they showed increased isolation and only moved when prodded, and 10 if they did not

ambulate with prodding. Animals were given a hydration score of 0 if the skin tent immediately

returned to normal, 1 if the tent took 1-2 seconds to return to normal, 2 if the tent took greater

than 2 seconds to return to normal, and 3 if this occurred accompanied by sunken and dry eyes.

Animals were given a respiration score of 0 if they displayed a normal breathing pattern, 1 if

1212 they had an increased breathing rate with an abdominal component, 3 if their breathing appeared

1213 labored, and 10 if their breathing was agonal.

1214

1215 **Cell Culture**

1216    L929 cells were obtained from ATCC and maintained in DMEM (Sigma-Aldrich,

1217 D5796) and 10% fetal bovine serum (FBS; VWR, 97068-085). Primary bone marrow-derived

1218 macrophages (BMDMs) were isolated from femurs of 6-12 month old WT, $Polg^{D257A}$ and 2-6

1219 month old $Polg^{R292C}$ mice as described[20]. Briefly, whole femurs were crushed in media and

1220 passed through a 70 µm filter. Red blood cells were lysed through osmosis by resuspending in

1221 ACK lysis buffer (Quality Biological, 118-156-101) for 2 minutes. Remaining cells were passed

1222 through a 40 µm filter, counted, and plated on non-tissue culture treated dishes. Macrophages

1223 were differentiated over for 7 days using 30% L929 conditioned medium. Peritoneal

1224 macrophages (PerMacs) were collected from the peritoneal cavity via gavage 4 days after

1225 intraperitoneal injection of 3% brewer thioglycolate medium (Sigma-Aldrich, B2551) and plated

1226 in in medium with 10% FBS. Unless stated, 6 x 10^5 BMDMs and 1.2x10^6 PerMacs per

1227 millimeter were used in in vitro experiments. Cells were plated and left to adhere overnight prior

1228 to challenge.

1229

1230 ***Pseudomonas aeruginosa* (PAO1) in vitro infection**

1231    GFP-expressing PAO1 was provided by Dr. Carolyn Cannon and was grown on Soy agar

1232 plates (BD, 211043) with 100 µg/mL carbenicillin (Dot Scientific, 46000-5) overnight from a

1233 glycerol stock. A single colony was incubated in 5 mL Mueller Hinton broth (BD, 275730) with

1234 carbenicillin at 37 □C spinning overnight. Overnight culture was back diluted 1:5 and grown for

2 hours until bacteria reached log-phase. Bacteria were washed with sterile phosphate-buffered

saline (PBS) and opsonized for 20 minutes in mouse serum (Fisher, PI31880). After two more

washed with PBS macrophages were infected at the corresponding MOI and spun at 250 g for 5

minutes to synchronize infection. For inhibitor experiments, cells were primed with 10 µM Ac-

YVAD-cmk (Invivogen, inh-yvad), 10 µM Q-VD-OPh (Millipore Sigma, 551476), or 10 µM

MCC950 (InvivoGen inh-mcc) for 3 hours prior to challenge. For antibody blockade

experiments, cells were primed with 20 µg/mL anti-IFNAR antibody (Bio X Cell, BE0241), anti-

IL-1R (Bio X Cell, BE0256), or anti-IgG control (Bio X Cell, BE0083 or BE0091) antibodies for

4 hours prior to challenge. One hour after infection, wells were washed and treated with 300

µg/mL gentamicin (VWR, 0304) for 30 minutes to inactivate extracellular bacteria. The

remainder of the experiment was completed with media supplemented with 50 µg/mL

gentamicin and inhibitor or blocking antibody when described.

1247

Escherichia coli **(E. coli) in vitro infection**

E. Coli containing empty pcDNA3.1 vector for antibiotic resistance was grown on LB

agar plates (Fisher, DF0401-17) with 100 µg/mL carbenicillin (Dot Scientific, 46000-5)

overnight from a glycerol stock. A single colony was incubated in 5 mL LB broth (Fisher,

BP1427-500) with carbenicillin at 37 □C spinning overnight. Overnight culture was back diluted

1:5 and grown for 2 hours until bacteria reached log-phase. Bacteria were washed with sterile

phosphate-buffered saline (PBS) and opsonized for 20 minutes in mouse serum. After two more

washed with PBS macrophages were infected at the corresponding MOI and spun at 250 g for 5

minutes to synchronize infection. One hour after infection, wells were washed and treated with

300 µg/mL gentamicin (VWR, 0304) for 30 minutes to kill extracellular bacteria. The rest of the

experiment was completed in DMEM + 10% FBS supplemented with 50 µg/mL gentamicin.

1259

Inflammasome challenge

BMDMs were primed for 4 hours with 20 ng/mL LPS-B5 Ultrapure (InvivoGen, tlrl-

pb5lps). When used, caspase inhibitor Ac-YVAD-cmk (InvivoGen, inh-yvad) was added an hour

after LPS, to allow for 3 hour priming, at 10 μM concentration. To trigger the non-canonical and

NLRC4 inflammasomes respectively, polyethyleneimine (Alfa Aesar, 43896) was used to

transfect 2 μg/mL LPS[75] and 10 ng/mL PA flagella[76] into the cytosol. To trigger the NLRP3

inflammasome, 5 μM ATP (Sigma, A6419) or 40 μM Nigericin (Cayman Chemical Company,

11437) was added to the media. When used, 20 µg/mL anti-IFNAR antibody (Bio X Cell,

BE0241) or anti-IgG control (Bio X Cell, BE0083) were added to cells 5 hours prior to LPS

priming and added again upon LPS transfection.

1270

Immunoblotting

Cells and tissues were lysed in 1% NP-40 lysis buffer supplemented with protease

inhibitor and then centrifuged at 15,000 rpm for 10 min at 4 □C and the supernatant was

collected as the cellular lysate. A bicinchoninic acid assay (BCA) protein assay (Thermo Fisher

Scientific, PI23235), was used to quantify protein and 10 to 40 μg total protein were loaded into

10% SDS-polyacrylamide gels and transferred onto 0.22 μM polyvinylidene difluoride

membranes. Membranes were dried to return them to a hydrophobic state and then incubated in

primary antibodies diluted in 1x PBS containing 1% casein overnight at 4 □C rolling.

Membranes were washed and incubated with horseradish peroxidase (HRP)- conjugated

secondary antibodies for 1 hour at room temperature, shaking. Membranes were developed with

Luminata Crescendo Western HRP Substrate (Millipore, WBWR0500). Primary antibody

information can be found in Table 1.

1283

LegendPlex multianalyte cytokine analysis

Supernatants from macrophages were subjected to centrifugation prior to the assay to

remove detached cells and debris from the sample. The assay was completed using the

LegendPlex Mouse Inflammation Panel (BioLegend, 740446) according to the included protocol.

Samples were analyzed by flow cytometry using a 5L Cytek Aurora and the BioLegend analysis

software. Samples were normalized to total protein content of each sample determined by a

bicinchoninic acid assay (BCA) protein assay and MFI from the blank control beads was

subtracted to exclude background.

1292

Enzyme-linked immunosorbent assay (ELISA)

Supernatants were collected from macrophages plated on glass coverslips. Macrophages

were stained for IF analysis while supernatants were subjected to centrifugation to remove

detached cells and debris from the sample. Samples were stored at -80C until use in this assay

and were thawed on ice upon use. The assay was completed in 96-well half area clear, flat

bottom plates (USA Scientific, 5667-5061) and wells were incubated overnight at 4□C in

capture antibodies. Blocking was completed for one hour with PBS containing 10% FBS at room

temperature. Standards and samples were added to wells for two hours. All sample dilutions

were made in blocking buffer. Next, wells were incubated in detection antibody for one hour.

Avidin-HRP and trimethyl boron (TMB) were used for quick color development. Plates were

washed with PBS containing 0.05% Tween 20 between each step. Capture and detection

antibodies were used in the following pairs from BioLegend (Table 3). Samples were normalized

to corresponding nuclei counts from 20X IF images taken on an EDLIPSE Ti2 (Nikon). Nuclei

counts were determined using the automated bright spot counting function in NIS-Elements on

DAPI and an average of 3 images was used to normalize ELISA. Normalization of BAL from

PAO1-*algC* infected mice was normalized to protein concentration. ELISA concentrations were

determined using an epoch plate reader (BioTek) and BioTek Gen5 software. Normalized results

were graphed in GraphPad.

1311

Quantitative polymerase chain reaction

Relative gene expression with measured by qRT-PCR in cells. Total cellular RNA was

isolated using Quick-RNA microprep kit (Zymo Research, R1051). Approximately 0.5 µg of

RNA was used to make cDNA with XLT qScript cDNA Synthesis Kit (Quantabio, 95047).

cDNA was then used in a qPCR assay using PerfeCTa Green FastMix (Quantabio, 95071). Two

to three biological replicates and two to three technical replicates were normalized to *Rpl37*

cDNA using the $2^{-\Delta\Delta CT}$ method. All primer sequences can be found in Table 4.

1319

Immunofluorescence (IF) imaging

Cells were seeded at 2.5×10^5 cell/well in a 24 well plate and allowed to grow overnight

on glass coverslips sterilized with ethanol. After challenge or infection, cells were washed in

PBS and fixed with 4% paraformaldehyde for 20 minutes. Next cells were permeabilized with

0.1% Triton X-100 in PBS for 5 minutes, blocked with PBS containing 5% FBS for 30 minutes,

stained with primary antibodies (table 2) for 1 hour, stained with secondary antibodies for 1

hour, and stained with 200 ng/mL 4',6-diamidino-2-phenylindole (DAPI) (ThermoFisher

Scientific, 62247) for 2 minutes. When described, coverslips were stained with 2 µg/mL

CellMask (ThermoFisher Scientific, C10046) for 10 minutes after secondary antibody staining

and washed multiple times in PBS before DAPI staining. Coverslips were mounted with

ProLong Diamond Antifade Mountant (Invitrogen, MP33025) and sealed with nail polish. Cells

were imaged with a 40x or 60x oil-immersed objective with an ECLIPSE Ti2 widefield

microscope (Nikon) and analyzed with the NIS-Elements AR 5.21.02 software.

Quantification was done from multiple intensity projection (MIP) images generated from

Z-stacks. MIPs were made and analysis was done using NIS-Elements AR 5.21.02. Specifically,

cell death was determined using the automated bright spot counting function in NIS-Elements on

DAPI to count the number of nuclei. Dying cells were manually determined by identifying cells

with condensed nuclei and low CellMask staining. Percentage of dying cells were determined

and graphed in GraphPad.

Protein intensities were quantified by thresholding a mask over each channel and

exporting sum intensities and area of masks. The sum intensities were normalized to area or cell

numbers where described. When normalized to cell number, cells were counted using the

automated bright spot counting function in NIS-Elements on DAPI. To quantify intensities of

colocalized proteins, a mask was generated over each protein. Next, a new mask was made for

the overlap of these channels and the intensity and area of each channel in this mask was

exported and graphed.

To quantify nucleoids, whole cells were outlines and made into a region of interest (ROI)

and these ROIs were numbered. A thresholding mask was made on the channels with TFAM and

DNA staining and channel intensities were determined for these masks and graphed. A new

mask was made based on the overlap of TFAM and DNA. The area of this mask and number of

spots per ROI were exported and graphed in GraphPad. Graphs and statistics were generated in

GraphPad. Huygens Essential software (v21.4.0) was used to deconvolve representative images

from z-stacks taken with a 60X objective in oil immersion.

1353

1354 **Lactate dehydrogenase release**

1355 Lactate dehydrogenase (LDH) was measured in the supernatant of PAO1 challenged

1356 macrophages using a LDH-Cytox Assay Kit (Biolegend, 426401) according to manufacturer's

1357 recommendations. A set of cells were treated with 1% Triton X-100 and the supernatant was

1358 used to determine total cellular LDH.

1359

PI uptake

BMDMs in 96 well half area glass bottom imaging plates were seeded at 6×10^4 cells/well

in 50 µL of Phenol-Red free DMEM + 10% FBS. The next day, cells were subjected to

inflammasome challenge or infection with PAO1. Propidium iodide (PI, Alfa Aesar, J66584)

was added to media and plates were immediately imaged every ten minutes using a BioTek

Cytation V imager which maintains 37 □C and 5% CO_2. PI positive nuclei were identified and

counted using BioTek Gen5 software. Control wells treated with PI and Triton X-100 (to

permeabilize cells) to for normalization. The difference between PI positive cells compared to

the initial reading were graphed in line graphs using GraphPad Prism.

1369

Cell fractionation and mitochondrial enrichment

Macrophages were lifted from cell culture dishes in PBS using a cell scraper. Cells were washed

with PBS and then resuspended in Resuspension buffer (10 mM NaCl, 1.5 mM CaCl2, 10 mM

Tris-Hcl, pH 7.5) with phosphatase and protease inhibitors. A dounce homogenizer was used to

break cell membranes and liberate organelles. Mitochondria were bound to magnetic beads

coated with TOMM20 antibodies isolated using a mitochondria isolation kit (Miltenyi Biotec,

120-009-492). The remaining cell components and unbound mitochondria were pelleted and run

as the cellular fraction. All samples were lysed in 1% SDS lysis buffer and run on a western blot

for analysis.

1379

Seahorse assay

Mitochondrial respiration and glycolysis in BMDMs were measured using a Seahorse XFe96

Analyzer. Macrophages were plates at a denisity of $5x10^5$ cells/well in DMEM supplemented

with 10% FBS and 5% L929 in an Agilent Seahorse XF96 Cell Culture Microplate. When

described, 20 ng/mL LPS was added 16 hours prior to analysis. Cells were placed in XF assay

medium immediately before analysis. ECAR and OCR measurements were recorded through

sequential addition of 10 mM glucose, 1.5 µM oligomycin, 1.5 µM carbonyl cyanide p-

trifluoromethoxy phenylhydrazone with 1 mM sodium pyruvate, and finally 833 nM antimycin

A plus 833 nM rotenone. Data was analyzed and calculations were made according to[77].

1389

siRNA knock down in bone marrow derived macrophages

Bone marrow was isolated from mice as described above. On day four, macrophages were

moved to 10 cm dishes. On day five, macrophages were transfected with 25 nM of Negative

Control (51-01-14-03), mm.Ri.Casp4.13.1, mm.Ri.Gbp2.13.1, or mm.Ri.Gbp5.13.1 DsiRNAs

(IDT) using Lipofectamine RNAiMAX (ThermoFisher Scientific, 13778100) in OPTIMEM

(Gibco, 51985-034) according to company recommendations. Six hours after transfection, media

was removed and replaced with fresh DMEM containing 10% FBS, 5% L929, and 20 µg/mL

anti-IFNAR antibody (Bio X Cell, BE0241) to reduce IFN-I-mediated toxicity due to siRNA

transfection. A day after transfection, cells were replated in cell culture plates in macrophage

media without anti-IFNAR antibody, then used in infection experiments the next day as

described above.

1401

mtDNA copy number analysis

DNA was isolated from cell pellets or crushed tissues by boiling for 30 minutes in 50

mM NaOH lysis buffer and then neutralizing using 1 M Tris, pH8 solution. DNA was diluted to

2 ng/uL and utilized in qRT-PCR reaction (described above) with mtDNA specific primers. Two

to three biological replicates and two to three technical replicates were normalized to nuclear

encoded *Tert* using the $2^{-\Delta\Delta CT}$ method. All primer sequences can be found in Table 5.

1408

Transmission electron microscopy

Macrophages were lifted from cell culture dishes and pellets were fixed for 1 hour in 4%

PFA. After fixation, cells were rinsed in 0.1M cacodylate buffer and then post-fixed in 1%

osmium tetroxide (OsO_4) in 0.1 M cacodylate buffer for 1 hour at room temperature. Cells were

rinsed again in the same buffer then spun down into pellets with 2% liquid agarose in distilled

water and placed in a 4 C refrigerator for 30 minutes, allowing the agarose to harden. After

trimming the agarose cell pellets, samples were transferred through graded ethanol (40% ETOH

to 100% ETOH), propylene oxide, and then infiltrated with 1:1 propylene oxide:Embed 812

resin (Electron Microscopy Sciences, Hatfield, PA) overnight, 1:3 propylene oxide:Embed 812

resin for 4 hours, and finally 100% Embed 812 resin overnight. Tissues were embedded in 100%

resin in beam capsules and cured at 65°C for 48 hours prior to sectioning. Thin sections (90 nm)

were cut on a Leica EM UC7 ultramicrotome (Leica Microsystems, Buffalo Grove, IL.) with a

diamond knife, placed onto 300 mesh copper grids and stained using 2% uranyl acetate and

Reynolds lead citrate. Samples were evaluated at 80 kV using a JEOL JM-1230 transmission

electron microscope (JEOL, Tokyo, Japan) and images collected with an AMT Nanosprint15

MK-11 sCMOS digital camera (Advanced Microscopy Techniques, Woburn, MA.). Image

quantification of cellular and PAO1 areas were completed in using NIS-Elements AR 5.21.02.

1426

RNA sequencing and bioinformatic analysis

Total cellular RNA was prepared from uninfected BMDMs and infected BMDMs using

Quick-RNA microprep kit (Zymo Research, R1051) and used for the next-generation RNA-

sequencing procedure at Texas A&M University Bioinformatics Core. BaseSpace Sequence Hub

(Illumina) was used for RNA seq analysis. Briefly, Spliced Transcripts Alignment to a Reference

(STAR) algorithm of RNA-seq Alignment V2.0.0 software was used to align the results to

reference genome Mus musculus/mm10 (RefSeq), and then RNA-seq Differential Expression

V1.0.0 software was used to obtain raw gene expression files and determine statistically

significant changes in gene expression in *Polg* mutant macrophages relative to WT. Ingenuity

Pathway Analysis software (QIAGEN) was used to identify gene families and putative upstream

regulators in the datasets. Gene Set Enrichment Analysis (GSEA) Software (Broad Institute) was

used to generate enrichment plots and examine pathway enrichment. Heatmaps were generated

using GraphPad Prism. Datasets of LPS-treated *Polg*[D257A] BMDMs used in Fig. 1-2 and

Supplementary Fig. 1 were previously reported[20] and are available under Gene Expression

Omnibus number GSE171960. Normalized gene counts from PAO1-infected *Polg*[D257A] BMDMs

are supplied for review in a supplemental data spreadsheet (see **Dataset 1** Excel file). These data

will be deposited into the Gene Expression Omnibus upon acceptance of the paper for

publication.

RNA sequencing of lungs was performed by isolating total cellular RNA from infected

lungs using a Direct-zol RNA miniprep Plus kit (Zymo Research, R2072). Next-generation

RNA-sequencing was performed by The Jackson Laboratory Genomic Technologies Core.

Partek Flow (v12.3.0) was used for RNA seq analysis. Briefly, Spliced Transcripts Alignment to

a Reference (STAR) algorithm of RNA-seq Alignment was used to align the results to reference

genome Mus musculus/mm10 (RefSeq), and then RNA-seq Differential Expression software

analysis was used to obtain raw gene expression files and determine statistically significant

changes in gene expression in *Polg* mutant macrophages relative to WT. Ingenuity Pathway

Analysis software (QIAGEN) was used to identify gene families and putative upstream

regulators in the datasets. Heatmaps were generated using GraphPad Prism. Normalized gene

counts from PAO1-infected *Polg^{D257A}* lungs and PAO1-infected *Polg^{R292C}* lungs are supplied for

review in a supplemental data spreadsheet (see **Dataset 2** Excel file). These data will be

deposited into the Gene Expression Omnibus upon acceptance of the paper for publication.

1458

In vitro proteomic and bioinformatic analysis

Cell sample protein extraction and quantification.

The adherent cells were washed on the six-well culture plate with PBS (+calcium and

magnesium) three times. After washing the cells, 250 µL of 6M Urea in 50 mM HEPES, pH 8.0

was added to each well. Cells were scraped from the plate and the solution was transferred to a 2

mL microcentrifuge tube. Samples were then snap frozen in liquid nitrogen and transferred to

the Mass Spectrometry and Protein Chemistry Service at The Jackson Laboratory for further

processing. Each tube had a pre-chilled 5 mm stainless steel bead (QIAGEN) added and were

lysed with the Tissue Lyser II (QIAGEN) for 2 minutes at 30 1/s. A magnet was used to remove

the stainless steel beads and each sample was further lysed using a 1 mL syringe with a 23-gauge

needle for 10 passes. Samples were then ice-waterbath sonicated at 37 Hz (100% power;

Firsherbrand FB11207) for 5 minutes (30 seconds on, 30 seconds off for five cycles). The cell

lysates were then centrifuged at 21,000 x g at 4°C for 15 minutes and the supernatant was

transferred to a new 1.5 mL microcentrifuge tube. A small fraction of the sample was then used

to create a 1:50 dilution (sample:50 mM HEPES) and a microBCA assay (Thermo) was run for

protein quantification according to the manufacturer protocol.

For each of the quantified cellular protein lysates, 20 µg of protein was diluted in 50 mM

HEPES, pH 8.0 to bring the urea concentration less than 2M for the trypsin digest. Each sample

was then reduced with 10 mM dithiothreitol (final concentration) for 30 minutes at 37oC for 30

minutes in a ThermoMixer (500 rpm agitation). Samples were then cooled to room temperature

and alkylated with 15 mM iodoacetamide for 20 minutes at 21oC in a ThermoMixer (500 rpm

agitation).   Sequence grade modified trypsin (Promega) was then add at a 1:40 ratio

(trypsin:protein) to each sample for digestion at 37oC for 20 hours.   Millipore C18 zip-tips

(Millipore, ZTC18S096) were then used to desalt and purify the peptide digests according to the

manufacturer protocol[78,79]. All purified peptide samples were then dried using a vacuum

centrifuge. Dried peptide digests were reconstituted in 20 µL of 98% H2O/2% ACN with 0.1%

formic acid via pipetting up and down 10 times, followed by vortexing at max speed for 30

seconds. Reconstituted samples were centrifuged at low speed for 15 seconds on a tabletop

minicentrifuge to collect the liquid from the walls in the bottom of the microcentrifuge tube. All

liquid was then transferred to a mass spec vial (Thermo) and placed in the Vanquish Neo

(Thermo) liquid chromatography autosampler set to 4oC.

1490

Liquid Chromatography Mass Spectrometry Analysis

Data independent acquisition mass spectrometry analysis (DIA-MS) was performed on a

Thermo Orbitrap Astral mass spectrometer coupled to the Vanquish Neo liquid chromatography

system with an EASY-Spray column (PepMap Neo C18, 75 µm x 150 mm, #ES75150PN) in the

Mass Spectrometry and Protein Chemistry Service at The Jackson Laboratory. Each sample had

two technical replicate runs consisting of an 8 µL injection of reconstituted sample. Before each

sample injection the separation column underwent fast equilibration in the method using pressure

control at 1500 bar and an equilibration factor of three column volumes. The trapping column

underwent fast wash and equilibration using a was factor of 100, an automatic equilibration

factor, and flow control at 120 uL/minute. The method gradient was 25 minutes after the column

equilibration using Buffer A (100% H2O with 0.1% formic acid) and Buffer B (100%

acetonitrile with 0.1% formic acid) with a flow rate of 600 nL/min. The full gradient consisted of

2% B to 4% B from 0-2.5 minutes, 4% B to 6% B from 2.5-4.5 minutes, 6% B to 30% B from

4.5-22.5 minutes, and 30% B to 90% B from 22.5-24 minutes while ramping to 1 µL/minute

flow rate in the last step. The column was then washed with 90% B at 1 µL/minute from 24-25

minutes. Orbitrap Astral settings included a static spray voltage using the NSI source at 2000 V

in positive mode, an ion transfer tube temperature of 325oC, an expected peak width of 10

seconds, advanced peak determination, a default charge state of 2, EASY-IC lock mass

correction, and RunStart mode. Full scan settings in positive mode included an orbitrap

resolution of 240,000, scan range of 380-980 m/z, RF lens at 60%, normalized AGC target of

500% with an absolute AGC value of 5e6, and a maximum injection time of 5 ms. DIA method

settings included a precursor mass range of 380-980 m/z, an isolation window of 2 m/z with no

overlap, normalized HCD collision energy of 25%, detector type set to Astral with a scan range

of 150-2000 m/z, RF lens at 60%, normalized AGC target of 500% with an absolute AGC value

of 5e4, maximum injection time of 5 ms, and loop control set at 0.6 seconds.

1516

DIA-MS data analysis

All Orbitrap Astral RAW data files underwent automatic processing through the Thermo

Ardia Server and were directly sent to the Proteome Discoverer 3.1 workstation connected to the

system. The RAW data files for each sample were searched against the UniProtKB Mus

musculus (sp_tr_incl_isoforms TaxID=10090; v2024-05-01) protein database in Proteome

Discoverer (version 3.1) using the CHIMERYS Ardia Server searching algorithm with the

inderys 3.0 fragmentation prediction model.   Search settings included trypsin digestion, a

maximum missed cleavage of one, peptide length of 7-30 amino acids, peptide charge from 1-4,

a fragment mass tolerance of 10 ppm, a dynamic oxidation of methionine modification (+15.995

Da), a static carbamidomethlyl modification of cysteine (+57.021 Da), and a false discovery rate

of less than 0.05 for all matches; all other settings were set to manufacturer default

recommendations. Total peptide amount was then utilized to normalize the ion signal in the

samples. Quantification analyses settings included the use of unique peptides for quantification

and a protein abundance-based ratio calculation was used for the summed abundances, followed

by a background based t-test. Strict parsimony principle was then applied for protein grouping

and master proteins were reported. The data was also run with no imputation and low abundance

resampling imputation for comparison during the analysis process.   The full dataset can be

found in the Proteome Exchange PRIDE upload XXXXXXXXXX (number will go here when

published, see **Dataset 3** Excel file).

1536

Statistical Analyses

To ensure rigor and reproducibility, experiments were repeated at least three times with

multiple biological replicates. Error bars displayed throughout the manuscript represent the mean

+/- s.e.m. Statistical significance between 2 groups was determined via unpaired, two-tailed

Student's t-test or Mann-Whitney test, and differences between 3 or more groups was determined

by two-way ANOVA with appropriate post-test. Survival plots were compared by the Log-rank

test. Sample sizes were chosen by started methods to ensure adequate power, and no

randomization or blinding was used for animal studies. Significance was established as * $p<0.05$;

** $p<0.01$; *** $p<0.001$; **** $p<0.0001$; or not significant (n.s.).

1546

1547 **<u>Tables</u>**

1548 **Table 1: Western blot antibodies**

Protein	Company	Catalog
ASC	Cell Signaling Technology	67824
CASPASE-1	Adipogen	AG-20B0042-C100
CASPASE-11	Cell Signaling Technology	14340
GAPDH	ProteinTech	60004-1-Ig
GBP5	Cell Signaling Technology	67798
GSDMD	Abcam	ab209845
GSDME	Cell Signaling Technology	88874S
HSP60	Santa Cruz Biotechnology	Sc-1052
IFIT3	Ganes Sen, Cleveland Clinic	N/A
IL-1β	Cell Signaling Technology	12242
IRF1	Cell Signaling Technology	8478
IRG1	Cell Signaling Technology	17805
NLRC4	Abcam	Ab201792
NLRP3	Cell Signaling Technology	15101
OXPHOS	Abcam	ab110413

pSTAT1	Cell Signaling Technology	7649
pTBK1	Cell Signaling Technology	5483
pIκBα	Cell Signaling Technology	2859
TBK1	Cell Signaling Technology	3504
STAT1	Cell Signaling Technology	9172
ZBP1	Adipogen	AG-20B-0010

1549

Table 2: Immunofluorescence microscopy antibodies

Protein	Company	Catalog
CASPASE-11	Cell Signaling Technology	14340
GFP	Thermo Fisher Scientific	NC9777966
LPS	Thermo Fisher Scientific	PA173178
PAO1	Thermo Fisher Scientific	PA173116
GBP5	Cell Signaling Technology	67798
TFAM	Proteintech	23996-1-AP
PDH	Abcam	Ab110333
DNA	Sigma Millipore	CBL186
ASC	Cell Signaling Technology	67824

1551

Table 3: ELISA antibodies (Biolegend)

Protein	Catalog (Binding)	Catalog (Detection)
TNFα	506302	506312
IL-1β	503514	515801
IL-6	504502	504602
IL-1α	503208	512503
IFNβ	519202	508105

1553

Table 4: Primer sequences for qRT-PCR analysis

Gene	F Primer 5'→ 3'	R Primer 5'→ 3'
Cxcl10	CCAAGTGCTGCCGTCATTTTC	GGCTCGCAGGGATGATTTCAA
Ifit1	CAAGGCAGGTTTCTGAGGAG	GACCTGGTCACCATCAGCAT
Casp11	TGTCATCTCTTTGATATATTCCTGAAG	CAAGGTTGCCCGATCAAT
Gbp2	CAGCATAGGAACCATCAACCA	TCTACCCCACTCTGGTCA
Gbp4	CAGGCTCTACATGGACATGAGG	TGCCTGCAAGATGGAACTCTCG
Gbp9	TCGTGGCTATTGTTGGATTG	GGTCTGAGACTGCACAGTGGG
Rpl37	CATCCTTTGGTAAGCGTCGCA	TGGCACTCCAGTTATACTTCCT

Stat1	CGCGCATGCAACTGGCATATAACT	ATGCTTCCGTTCCCACGTAGACTT
Zbp1	TCAAAGGGTGAAGACATGACAGAC	GTGGGTGGCTTCAGAGCTT
Ifit3	TTCCCAGCAGCACAGAAAC	AAATTCCAGGTGAAATGGCA
Gbp3	GGAAATCCTACCTCATGAATCG	TCCAGATACCCTTGGTTTCG

1555

1556 **Table 5: Primers for mtDNA copy number analysis by qPCR**

Gene	F Primer 5'→ 3'	R Primer 5'→ 3'
mt-Non-numpt	CTAGAAACCCCGAAACCAA	CCAGCTATCACCAAGCTCGT
Mt-D-loop	AATCTACCATCCTCCGTGAAACC	TCAGTTTAGCTACCCCCAAGTTTAA
Mt-Cytb	GCTTTCCACTTCATCTTACCATTTA	TGTTGGGTTGTTTGATCCTG
nuc Tert	CTAGCTCATGTGTCAAGACCCTCTT	GCCAGCACGTTTCTCTCGTT

1557

1558 **Table 6: Flow cytometry antibody panel**

Antigen	Fluorophore	Laser	Company	Catalog/ Ref #
CD45	BUV395	UV2	BD	564279
Ly6C	BV650	V11	BioLegend	128049
MHCII	BV711	V13	BioLegend	107643
CD19	BV605	V10	BioLegend	115539
CD3	APC-fire810	R8	BioLegend	100267
F4/80	VF450	V3	TONBO Biosciences	75-4801-U025
PA01	FITC	B2		
CD11c	PE-Cy5	YG5	TONBO Biosciences	55-0114-U100
TNF	PE-Cy7	YG9	Biolegend	506324
Ly6G	PE	YG1	TONBO Biosciences	50-1276-U100
Siglec-F	APC	R1	Biolegend	155508
CD11b (M1/70)	APC-Cy7	R7	TONBO Biosciences	25-0112-U100
CD326	PE/Dazzel 594	YG3	Biolegend	118235
Viability	Ghost Dye 710	R4	TONBO Biosciences	13-0871-T100

1559

1560

<u>**References**</u>

1. Gorman, G. S. *et al.* Mitochondrial diseases. *Nat Rev Dis Primers* **2**, 16080 (2016).

2. Suomalainen, A. & Battersby, B. J. Mitochondrial diseases: the contribution of organelle

stress responses to pathology. *Nat Rev Mol Cell Biol* **19**, 77–92 (2018).

3. Russell, O. M., Gorman, G. S., Lightowlers, R. N. & Turnbull, D. M. Mitochondrial

Diseases: Hope for the Future. *Cell* **181**, 168–188 (2020).

4. Kruk, S. K. *et al.* Vulnerability of pediatric patients with mitochondrial disease to vaccine-

preventable diseases. *J Allergy Clin Immunol Pract* **7**, 2415-2418.e3 (2019).

5. Walker, M. A. *et al.* Predisposition to infection and SIRS in mitochondrial disorders: 8

years' experience in an academic center. *J Allergy Clin Immunol Pract* **2**, 465–468, 468.e1

(2014).

6. Eom, S. *et al.* Cause of Death in Children With Mitochondrial Diseases. *Pediatr Neurol* **66**,

82–88 (2017).

7. Jestin, M. *et al.* Mitochondrial disease disrupts hepatic allostasis and lowers the threshold for

immune-mediated liver toxicity. *Molecular Metabolism* **37**, 100981 (2020).

8. West, A. P., Shadel, G. S. & Ghosh, S. Mitochondria in innate immune responses. *Nat Rev*

Immunol **11**, 389–402 (2011).

9. West, A. P. Mitochondrial dysfunction as a trigger of innate immune responses and

inflammation. *Toxicology* **391**, 54–63 (2017).

10. West, A. P. & Shadel, G. S. Mitochondrial DNA in innate immune responses and

inflammatory pathology. *Nat Rev Immunol* **17**, 363–375 (2017).

11. Warren, E. B. *et al.* Inflammatory and interferon gene expression signatures in patients with

mitochondrial disease. *J Transl Med* **21**, 331 (2023).

12. Hasselmann, O. *et al.* Cerebral folate deficiency and CNS inflammatory markers in Alpers

disease. *Mol Genet Metab* **99**, 58–61 (2010).

13. Wilson, L. D., Al-Majid, S., Rakovski, C. S. & Md, C. D. Higher IL-6 and IL6:IGF Ratio in

Patients with Barth Syndrome. *J Inflamm* **9**, 25 (2012).

14. Khan, W. *et al.* Neuroinflammation in the Cerebellum and Brainstem in Friedreich Ataxia:

An [^{18} F]☐ FEMPA PET Study. *Movement Disorders* **37**, 218–224 (2022).

15. Nachun, D. *et al.* Peripheral blood gene expression reveals an inflammatory transcriptomic

signature in Friedreich's ataxia patients. *Human Molecular Genetics* **27**, 2965–2977 (2018).

16. Jin, Z., Wei, W., Yang, M., Du, Y. & Wan, Y. Mitochondrial Complex I Activity Suppresses

Inflammation and Enhances Bone Resorption by Shifting Macrophage-Osteoclast

Polarization. *Cell Metabolism* **20**, 483–498 (2014).

17. Yu, A. K. *et al.* Mitochondrial complex I deficiency leads to inflammation and retinal

ganglion cell death in the Ndufs4 mouse. *Human Molecular Genetics* **24**, 2848–2860 (2015).

18. Stokes, J. C. *et al.* Leukocytes mediate disease pathogenesis in the Ndufs4(KO) mouse

model of Leigh syndrome. *JCI Insight* **7**, e156522 (2022).

19. Hanaford, A. & Johnson, S. C. The immune system as a driver of mitochondrial disease

pathogenesis: a review of evidence. *Orphanet J Rare Dis* **17**, 335 (2022).

20. Lei, Y. *et al.* Elevated type I interferon responses potentiate metabolic dysfunction,

inflammation, and accelerated aging in mtDNA mutator mice. *Sci Adv* **7**, eabe7548 (2021).

21. Saha, S., Buttari, B., Panieri, E., Profumo, E. & Saso, L. An Overview of Nrf2 Signaling

Pathway and Its Role in Inflammation. *Molecules* **25**, 5474 (2020).

22. Nezu, M., Suzuki, N. & Yamamoto, M. Targeting the KEAP1-NRF2 System to Prevent

Kidney Disease Progression. *Am J Nephrol* **45**, 473–483 (2017).

23. Trifunovic, A. *et al.* Premature ageing in mice expressing defective mitochondrial DNA

polymerase. *Nature* **429**, 417–423 (2004).

24. Kujoth, G. C. *et al.* Mitochondrial DNA mutations, oxidative stress, and apoptosis in

mammalian aging. *Science* **309**, 481–484 (2005).

25. Chen, M. L. *et al.* Erythroid dysplasia, megaloblastic anemia, and impaired lymphopoiesis

arising from mitochondrial dysfunction. *Blood* **114**, 4045–4053 (2009).

26. Thomsen, K. *et al.* Animal models of chronic and recurrent Pseudomonas aeruginosa lung

infection: significance of macrolide treatment. *APMIS* **130**, 458–476 (2022).

27. Lavoie, E. G., Wangdi, T. & Kazmierczak, B. I. Innate immune responses to Pseudomonas

aeruginosa infection. *Microbes Infect* **13**, 1133–1145 (2011).

28. Lister, P. D., Wolter, D. J. & Hanson, N. D. Antibacterial-resistant Pseudomonas aeruginosa:

clinical impact and complex regulation of chromosomally encoded resistance mechanisms.

Clin Microbiol Rev **22**, 582–610 (2009).

29. Lambert, M.-L. *et al.* Clinical outcomes of health-care-associated infections and

antimicrobial resistance in patients admitted to European intensive-care units: a cohort study.

Lancet Infect Dis **11**, 30–38 (2011).

30. Sutterwala, F. S. *et al.* Immune recognition of Pseudomonas aeruginosa mediated by the

IPAF/NLRC4 inflammasome. *J Exp Med* **204**, 3235–3245 (2007).

31. Balakrishnan, A., Karki, R., Berwin, B., Yamamoto, M. & Kanneganti, T.-D. Guanylate

binding proteins facilitate caspase-11-dependent pyroptosis in response to type 3 secretion

system-negative Pseudomonas aeruginosa. *Cell Death Discov* **4**, 3 (2018).

32. Arlehamn, C. S. L. & Evans, T. J. Pseudomonas aeruginosa pilin activates the

inflammasome. *Cell Microbiol* **13**, 388–401 (2011).

33. Minns, M. S. *et al.* NLRP3 selectively drives IL-1β secretion by Pseudomonas aeruginosa

infected neutrophils and regulates corneal disease severity. *Nat Commun* **14**, 5832 (2023).

34. Kroken, A. R. *et al.* Intracellular replication of Pseudomonas aeruginosa in epithelial cells

requires suppression of the caspase-4 inflammasome. *mSphere* **8**, e0035123 (2023).

35. Ramadan, A. *et al.* The NLRP3 Inflammasome Is Required for Protection Against

Pseudomonas Keratitis. *Invest Ophthalmol Vis Sci* **64**, 11 (2023).

36. Billingham, L. K. *et al.* Mitochondrial electron transport chain is necessary for NLRP3

inflammasome activation. *Nat Immunol* **23**, 692–704 (2022).

37. Kumari, P., Russo, A. J., Wright, S. S., Muthupalani, S. & Rathinam, V. A. Hierarchical cell-

type-specific functions of caspase-11 in LPS shock and antibacterial host defense. *Cell*

Reports **35**, 109012 (2021).

38. Benjamin, J. T. *et al.* IL-1α and Not IL-1β Drives IL-1R1–Dependent Neonatal Murine

Sepsis Lethality. *The Journal of Immunology* **201**, 2873–2878 (2018).

39. Pyrillou, K., Burzynski, L. C. & Clarke, M. C. H. Alternative Pathways of IL-1 Activation,

and Its Role in Health and Disease. *Front. Immunol.* **11**, 613170 (2020).

40. Miao, R. *et al.* Gasdermin D permeabilization of mitochondrial inner and outer membranes

accelerates and enhances pyroptosis. *Immunity* **56**, 2523-2541.e8 (2023).

41. Rathinam, V. A. K. *et al.* TRIF Licenses Caspase-11-Dependent NLRP3 Inflammasome

Activation by Gram-Negative Bacteria. *Cell* **150**, 606–619 (2012).

42. Kim, B.-H. *et al.* A family of IFN-γ-inducible 65-kD GTPases protects against bacterial

infection. *Science* **332**, 717–721 (2011).

43. Kim, B.-H. *et al.* Interferon-induced guanylate-binding proteins in inflammasome activation

and host defense. *Nat Immunol* **17**, 481–489 (2016).

44. Shenoy, A. R. *et al.* GBP5 promotes NLRP3 inflammasome assembly and immunity in

mammals. *Science* **336**, 481–485 (2012).

45. Coers, J. Sweet host revenge: Galectins and GBPs join forces at broken membranes. *Cellular*

Microbiology **19**, e12793 (2017).

46. Coyne, M. J., Russell, K. S., Coyle, C. L. & Goldberg, J. B. The Pseudomonas aeruginosa

algC gene encodes phosphoglucomutase, required for the synthesis of a complete

lipopolysaccharide core. *J Bacteriol* **176**, 3500–3507 (1994).

47. Deus, C. M., Yambire, K. F., Oliveira, P. J. & Raimundo, N. Mitochondria-Lysosome

Crosstalk: From Physiology to Neurodegeneration. *Trends Mol Med* **26**, 71–88 (2020).

48. Newman, L. E. *et al.* Mitochondrial DNA replication stress triggers a pro-inflammatory

endosomal pathway of nucleoid disposal. *Nat Cell Biol* **26**, 194–206 (2024).

49. Nguyen, M. *et al.* Parkinson's genes orchestrate pyroptosis through selective trafficking of

mtDNA to leaky lysosomes. Preprint at https://doi.org/10.1101/2023.09.11.557213 (2023).

50. Santos, J. C. *et al.* LPS targets host guanylate☐binding proteins to the bacterial outer

membrane for non☐canonical inflammasome activation. *The EMBO Journal* **37**, e98089

(2018).

51. Finethy, R. *et al.* Inflammasome Activation by Bacterial Outer Membrane Vesicles Requires

Guanylate Binding Proteins. *mBio* **8**, e01188-17 (2017).

52. Tsuchiya, K. *et al.* Gasdermin D mediates the maturation and release of IL-1α downstream

of inflammasomes. *Cell Rep* **34**, 108887 (2021).

53. Weber, A., Wasiliew, P. & Kracht, M. Interleukin-1 (IL-1) pathway. *Sci Signal* **3**, cm1

(2010).

54. Malik, A. & Kanneganti, T.-D. Function and regulation of IL-1α in inflammatory diseases

and cancer. *Immunol Rev* **281**, 124–137 (2018).

55. Dinarello, C. A., Simon, A. & van der Meer, J. W. M. Treating inflammation by blocking

interleukin-1 in a broad spectrum of diseases. *Nat Rev Drug Discov* **11**, 633–652 (2012).

56. Lam, C.-W. *et al.* Novel POLG mutation in a patient with sensory ataxia, neuropathy,

ophthalmoparesis and stroke. *Clin Chim Acta* **448**, 211–214 (2015).

57. Amiot, A. *et al.* Frequency of mitochondrial defects in patients with chronic intestinal

pseudo-obstruction. *Gastroenterology* **137**, 101–109 (2009).

58. Kinoshita, M. & Sawa, T. Serological immune response against major secretory proteins

expressed by Pseudomonas aeruginosa upon intratracheal infection in mice. *Respir Investig*

60, 318–321 (2022).

59. Edmonds, J. L. *et al.* The Otolaryngological Manifestations of Mitochondrial Disease and the

Risk of Neurodegeneration With Infection. *Arch Otolaryngol Head Neck Surg* **128**, 355

(2002).

60. Lee, W. S. & Sokol, R. J. Mitochondrial Hepatopathies: Advances in Genetics, Therapeutic

Approaches, and Outcomes. *The Journal of Pediatrics* **163**, 942–948 (2013).

61. The Mitochondrial Medicine Society *et al.* A modern approach to the treatment of

mitochondrial disease. *Curr Treat Options Neurol* **11**, 414–430 (2009).

62. Rahman, S. & Copeland, W. C. POLG-related disorders and their neurological

manifestations. *Nat Rev Neurol* **15**, 40–52 (2019).

63. Gaudó, P. *et al.* Infectious stress triggers a POLG-related mitochondrial disease.

Neurogenetics **21**, 19–27 (2020).

64. Al-Zubeidi, D. *et al.* Fatal Human Herpesvirus 6–Associated Encephalitis in Two Boys With

Underlying POLG Mitochondrial Disorders. *Pediatric Neurology* **51**, 448–452 (2014).

65. Torres-Odio, S. *et al.* Loss of Mitochondrial Protease CLPP Activates Type I IFN Responses

through the Mitochondrial DNA-cGAS-STING Signaling Axis. *J Immunol* **206**, 1890–1900

(2021).

66. Buchel, G. *et al.* Structural basis for DNA proofreading. *Nat Commun* **14**, 8501 (2023).

67. Kang, Y. *et al.* Ancestral allele of DNA polymerase gamma modifies antiviral tolerance.

Nature **628**, 844–853 (2024).

68. Liu, B. C. *et al.* Constitutive Interferon Maintains GBP Expression Required for Release of

Bacterial Components Upstream of Pyroptosis and Anti-DNA Responses. *Cell Reports* **24**,

155-168.e5 (2018).

69. Dickinson, M. S. *et al.* LPS-aggregating proteins GBP1 and GBP2 are each sufficient to

enhance caspase-4 activation both in cellulo and in vitro. *Proc. Natl. Acad. Sci. U.S.A.* **120**,

e2216028120 (2023).

70. Pilla, D. M. *et al.* Guanylate binding proteins promote caspase-11–dependent pyroptosis in

response to cytoplasmic LPS. *Proc. Natl. Acad. Sci. U.S.A.* **111**, 6046–6051 (2014).

71. Meunier, E. *et al.* Caspase-11 activation requires lysis of pathogen-containing vacuoles by

IFN-induced GTPases. *Nature* **509**, 366–370 (2014).

72. Yazdi, A. S. & Drexler, S. K. Regulation of interleukin 1α secretion by inflammasomes:

Table 1. *Ann Rheum Dis* **72**, ii96–ii99 (2013).

73. Keshavan, N., Mhaldien, L., Gilmour, K. & Rahman, S. Interferon Stimulated Gene

Expression Is a Biomarker for Primary Mitochondrial Disease. *Ann Neurol* **96**, 1185–1200

(2024).

74. Pettitt, S. J. *et al.* Agouti C57BL/6N embryonic stem cells for mouse genetic resources. *Nat*

Methods **6**, 493–495 (2009).

75. Bezbradica, J. S., Coll, R. C. & Boucher, D. Activation of the Non-canonical Inflammasome

in Mouse and Human Cells. in *The Inflammasome* (ed. Abdul-Sater, A. A.) vol. 2459 51–63

(Springer US, New York, NY, 2022).

76. Kamajaya, L. J. & Boucher, D. Gasdermin D Cleavage Assay Following Inflammasome

Activation. in *The Inflammasome* (ed. Abdul-Sater, A. A.) vol. 2459 39–49 (Springer US,

New York, NY, 2022).

77. Van den Bossche, J., Baardman, J. & de Winther, M. P. J. Metabolic Characterization of

Polarized M1 and M2 Bone Marrow-derived Macrophages Using Real-time Extracellular

Flux Analysis. *J Vis Exp* 53424 (2015) doi:10.3791/53424.

78. Zhang, B. *et al.* Metabolic fitness of IgA+ plasma cells in the gut requires DOCK8. *Mucosal

Immunol* **17**, 431–449 (2024).

79. Thatcher, K. *et al.* Temporal Progression of Aortic Valve Pathogenesis in a Mouse Model of

Osteogenesis Imperfecta. *J Cardiovasc Dev Dis* **10**, 355 (2023).

1735

1736