Caspase-11 drives macrophage hyperinflammation in models of Polg-related mitochondrial disease

Jordyn J. VanPortfliet[1,2], Yuanjiu Lei[2,3], Camila Guerra Martinez[2], Jessica Wong[1], Kathryn Pflug[4], Raquel Sitcheran[4], Stephen C. Kneeland[1], Stephen A. Murray[1], Peter. J. McGuire[5], Carolyn L. Cannon[2], A. Phillip West[1,2,*]

[1]The Jackson Laboratory, Bar Harbor, Maine 04609, USA

[2]Department of Microbial Pathogenesis and Immunology, School of Medicine, Texas A&M University, Bryan, Texas 77807, USA

[3]Department of Pathology, Yale School of Medicine, New Haven, CT 06520, USA

[4]Department of Cell Biology and Genetics, School of Medicine, Texas A&M University, Bryan, Texas 77807, USA

[5]Metabolism, Infection and Immunity Section, National Human Genome Research Institute, National Institutes of Health, Bethesda, Maryland, 20892, USA

* Correspondence to: phillip.west@jax.org

Abstract

Mitochondrial diseases (MtD) represent a significant public health challenge due to their heterogenous clinical presentation, often severe and progressive symptoms, and the lack of effective therapies. Environmental exposures, such bacterial and viral infection, can further compromise mitochondrial function and exacerbate the progression of MtD. Infections in MtD patients more frequently progress to sepsis, pneumonia, and other detrimental inflammatory endpoints. However, the underlying immune alterations that enhance immunopathology in MtD remain unclear, constituting a key gap in knowledge that complicates treatment and increases mortality in this population. Here we employ in vitro and in vivo approaches to clarify the molecular and cellular basis for innate immune hyperactivity in models of polymerase gamma (Polg)-related MtD. We reveal that type I interferon (IFN-I)-mediated upregulation of caspase-11 and guanylate-binding proteins (GBPs) increase macrophage sensing of the opportunistic microbe *Pseudomonas aeruginosa* (PA) in Polg mutant mice. Furthermore, we show that excessive macrophage cytokine secretion and pyroptotic cell death contribute to lung inflammation and morbidity after infection with PA. Our work sheds new light on innate immune dysregulation in MtD and reveals potential targets for limiting infection- and inflammation-related complications in Polg-related MtD.

Introduction

Mitochondrial diseases (MtD) are the most common inborn error of metabolism and are caused by deleterious variants in genes that function in oxidative phosphorylation (OXPHOS) and mitochondrial DNA (mtDNA) maintenance[1]. MtD can present at any age, vary in severity, and display clinical heterogeneity with differing degrees of organ system involvement[2,3]. Although the genetic basis of MtD is becoming more clear, the underlying mechanisms that cause pathology, as well as the contribution of environmental factors to disease progression, are less clear. Notably, viral and bacterial infections may unmask MtD or hasten the stepwise progression of disease by triggering metabolic decompensation, a dangerous deterioration in mitochondrial function[4]. Clinical evidence suggests that patients with MtD are highly susceptible to infection by opportunistic respiratory pathogens. Moreover, pneumonia and microbial sepsis are the primary causes of early death in pediatric patients, and some adults, with MtD[5,6]. Thus, there is ample evidence highlighting the damaging toll of infection in MtD[7]. However, the underlying immune alterations that drive immunopathology and prevent proper control of infection represent a significant knowledge gap that complicates treatment and enhances morbidity and mortality in this vulnerable patient population.

In addition to their bioenergetic functions, mitochondria are key metabolic and signaling hubs in the mammalian immune system[8]. Mitochondria are also important sources of damage-associated molecular patterns (DAMPs), such as mtDNA, which engage innate immune signaling when released from the organelle and are linked to inflammation in a growing number of human diseases[9,10]. There is growing appreciation for a bidirectional relationship between MtD and inflammation, where mitochondrial dysfunction in MtD activates the innate immune system leading to inflammatory responses that further compromise mitochondrial activity[11]. In

agreement with this hypothesis, elevated inflammatory cytokines have been observed in patients with Polymerase gamma (Polg)-related Alpers-Huttenlocher syndrome[12], Barth syndrome[13], and Friedreich ataxia[14,15]. Animal models have also revealed innate immune system dysregulation as a frequent feature of MtD. For example, the $Ndufs4^{-/-}$ mouse model of neurodegeneration and Leigh syndrome exhibits increased circulating pro-inflammatory cytokines along with macrophage and microglial activation[16–19].

We recently uncovered that the Polg^{D257A} mutator model of mtDNA instability exhibits a hyperinflammatory innate immune status due to chronic engagement of the cyclic GMP-AMP synthase (cGAS)—Stimulator of Interferon Genes (STING)—type one interferon (IFN-I) axis[20]. Notably, we reported that persistent IFN-I signaling in these mice causes widespread repression of the antioxidant and anti-inflammatory transcription factor nuclear factor erythroid 2-related factor 2 (Nrf2)[21,22] in myeloid immune cells and organs of the Polg^{D257A} mouse. Consequently, Polg^{D257A} mice are more sensitive to septic shock caused by intraperitoneal injection of a high dose of lipopolysaccharide (LPS)[20]. Although Polg^{D257A} mutator mice do not phenocopy any particular human MtD, these animals present pathology overlapping the MtD spectrum, including hearing loss, cardiomyopathy, progressive anemia, and lymphopenia[23–25]. Moreover, increased susceptibility to septic shock and elevated inflammatory and IFN-I signatures are common features of both Polg mutator mice and patients with diverse MtD[11].

Overall, our prior work provided foundational evidence linking mitochondrial dysfunction to hyperinflammatory innate immune signaling in the Polg^{D257A} model of MtD. However, there remains a critical need to expand these findings into relevant infection models, given the high rate of infections that frequently lead to innate immune hyperactivity and sepsis in MtD. *Pseudomonas aeruginosa* (PA) is a common opportunistic pathogen of concern in MtD as

infection can lead to sepsis and pneumonia[6]. PA is a robust activator of innate immunity in the

respiratory tract, and lung colonization by PA drives persistent recruitment of monocytes and

neutrophils leading to pneumonia, lung injury, and/or systemic bacterial spread[26,27]. PA is

especially problematic in MtD due to the prevalence of antibiotic resistant strains, abundance in

hospitals, and high rate of recurrence[26,28,29]. Thus, it is important to define how PA and other

opportunistic bacteria engages the innate immune system in MtD to define immunomodulatory

strategies that boost anti-bacterial immunity, while limiting hyperinflammation that may hasten

metabolic decompensation and disease progression.

In this study, we utilized in vitro and in vivo PA infection models to characterize the

underlying innate immune mechanisms governing hyperinflammation in Polg[D257A] mice. We also

report a new mouse model that incorporates a pathogenic Polg variant observed in humans. We

reveal that both Polg mutant models exhibit excessive macrophage inflammasome activation,

cytokine secretion, and pyroptotic cell death, which contribute to inflammation, acute lung

injury, and morbidity after infection with PA. These findings further advance knowledge of how

mitochondrial dysfunction impacts innate immune responses during bacterial infection and Polg-

related diseases and reveals innate immune targets for limiting infection- and inflammation-

related complications in MtD.

Results

Polg mutant macrophages exhibit increased IFN-I and pro-inflammatory responses after

bacterial challenge

Our recent work uncovered that elevated IFN-I and repression of Nrf2 signaling augment

innate immune activity and inflammation in Polg[D257A] mice challenged with Toll-like receptor

(TLR) agonist, LPS[20]. We hypothesized that the hyperinflammatory phenotype of Polg[D257A] mice

might be further exacerbated during bacterial infection. Therefore, we challenged bone marrow-

derived macrophages (BMDMs) with *Pseudomonas aeruginosa* strain O1 (PAO1) (Fig. 1a) and

performed RNA sequencing (RNAseq) and pathway analysis (Fig. 1b). Similar to BMDMs

treated with LPS[20], PAO1-challenged Polg[D257A] BMDMs displayed heightened expression of

pro-inflammatory cytokines, IFN-I, interferon-stimulated genes (ISGs), and innate immune

signaling factors (Fig. 1c; Supplementary Fig. 1a). Gene Set Enrichment Analysis confirmed

PAO1-infected Polg[D257A] BMDMs exhibit more robust pro-inflammatory cytokine, chemokine,

and IFN-I gene expression relative to wild-type (WT) BMDMs (Supplementary Fig. 1b).

Additional experiments substantiated RNAseq datasets, revealing heightened expression of ISGs

and pro-inflammatory cytokine secretion in PAO1-infected Polg[D257A] peritoneal macrophages

(PerMacs) and BMDMs over a one-day time course (Fig. 1d-e). Prior work from our lab showed

that elevated IFN-I responses in Polg[D257A] mice suppress Nrf2-mediated antioxidant and anti-

inflammatory signaling[20]. We noted consistent activation of IFN-I signaling factors and a

coordinated downregulation of Nrf2 target genes in PAO1-infected Polg[D257A] macrophages (Fig.

1f-g). Addition of dimethyl fumarate (DMF), an anti-inflammatory and Nrf2 stabilizing

agent[30,31], dramatically reduced IFN-I and inflammatory gene signatures in Polg[D257A]

macrophages during PAO1 challenge, while also restoring expression of Nrf2 target genes (Fig.

1b-c, f-h and Supplementary Fig. 1). Notably, DMF treatment reduced the hyperproduction of

multiple pro-inflammatory cytokines in Polg[D257A] PerMacs at both 8 and 24 hours post challenge

(hpc) (Fig. 1i). In sum, our data highlight that Polg[D257A] macrophages are hyperactivated by LPS

and PAO1 and reveal that DMF inhibits inflammation and restores Nrf2 levels in Polg mutant

cells.

Polg mutant macrophages exhibit potentiated cell death upon bacterial challenge

As we observed increased IL-1β secretion from Polg^{D257A} PerMacs, we next investigated

inflammatory cell death pathways during PAO1 infection. RNAseq and pathway analysis

revealed significant enrichment of genes involved in inflammasome signaling, immunogenic cell

death, and death receptor signaling in Polg^{D257A} BMDMs after PAO1 challenge (Fig. 2a-b).

Consistent with cell death gene expression patterns in BMDMs, Polg^{D257A} PerMacs stained

weakly with the nuclear and plasma membrane dye CellMask HCS Deep Red and exhibited

rapid condensation of nuclei after PAO1 challenge (Fig. 2c-d). Reduced CellMask staining was

also observed in ASC positive WT BMDMs after LPS priming and nigericin treatment

(Supplementary Fig. 2a-b), indicating that low CellMask staining marks cells undergoing

pyroptosis. In line with increased pyroptotic cell death, Polg^{D257A} BMDMs more rapidly became

propidium iodide (PI) positive (Fig. 2e) and secreted more IL-1β after incubation with PAO1

(Fig. 2f). We also noted elevated release of PAO1 into the culture supernatant of Polg^{D257A}

BMDMs, concomitant with lower numbers of intracellular bacteria (Supplementary Fig. 2c).

Reduced intracellular bacteria was not the result of impaired phagocytosis, as Polg^{D257A} BMDMs

displayed no impairment in uptake of LPS coated fluorescent microspheres (Supplementary Fig.

2d-e).

DMF is a robust inhibitor of the NLRP3 inflammasome[32], and addition of DMF during

PAO1 infection was sufficient to reduce genes governing cell death and heightened pyroptotic

phenotypes in Polg^{D257A} macrophages (Fig. 2). In addition to activating NLRP3, PA can also

activate the NLRC4 canonical inflammasome and the non-canonical caspase-11

inflammasome[27,33–35]. Culture of Polg^{D257A} BMDMs with non-pathogenic *E. coli* (DH5α), which

lack flagellin and therefore can only activate NLRP3 or caspase-11, also promoted increased IL-

1β, TNFα, and IL-6 secretion (Supplementary Fig. 2f) and pyroptotic cell death (Supplementary

Fig. 2g-h). Collectively, these data indicate that elevated cell death and cytokine secretion in

Polg^{D257A} macrophages during bacterial challenge are not due to PAO1 virulence factors, but are

likely caused by potentiated activation of NLRP3 and/or caspase-11 inflammasomes.

Elevated IFN-I potentiates expression and activation of the non-canonical caspase-11
inflammasome in Polg mutant macrophages

To further confirm a role for pyroptosis in hyperinflammation and increased cell death in

Polg mutant macrophages, we incubated BMDMs with Ac-YVAD-cmk (AcYVAD), a selective

caspase-1 inhibitor. AcYVAD nearly abrogated IL-1β secretion from Polg^{D257A} macrophages

while also decreasing TNFα and IL-6 during PAO1 challenge (Fig. 3a). In contrast, QVD-OPH,

a more potent inhibitor of apoptotic caspases, had minor effects on PAO1-induced cytokine

secretion in Polg^{D257A} BMDMs (Fig. 3a). AcYVAD also reduced PI uptake in Polg^{D257A} BMDMs

(Fig. 3b) and blunted pyroptotic phenotypes of Polg^{D257A} PerMacs (Fig. 3c-d) after incubation

with PAO1. Polg^{D257A} BMDMs treated with LPS alone did not exhibit increased PI uptake,

suggesting that priming is not sufficient to potentiate cell death (Supplementary Fig. 3a).

Given elevated pyroptosis in PAO1-infected Polg^{D257A} BMDMs, we next performed

isolated challenge experiments to determine which inflammasome pathways were hyperactivated

in mutant macrophages. Surprisingly, LPS priming followed by flagellin transfection to activate

the NLRC4 inflammasome revealed that Polg^{D257A} macrophages exhibit reduced PI uptake and

IL-1β secretion (Supplementary Fig. 3b-c). LPS priming followed by the addition of ATP also

revealed modestly impaired NLRP3 inflammasome activation in Polg^{D257A} BMDMs, as

measured by PI uptake, IL-1β secretion, and caspase-1 cleavage (Supplementary Fig. 3d-f). LPS

priming followed by nigericin treatment revealed similar impairments in NLRP3 inflammasome

activity (Supplementary Fig. 3g-i). OXPHOS has recently been shown to be vital for NLRP3

activation in response to nigericin and other stimuli[36]. Polg[D257A] BMDMs displayed reduced

basal and maximal respiration at rest and after LPS challenge (Supplementary Fig. 3j-l),

suggesting that reduced canonical inflammasome activation is likely due to OXPHOS

impairment in Polg[D257A] BMDMs.

Although Polg[D257A] BMDMs exhibited reduced NLRP3 and NLRC4 responses when

triggered with pure inflammasome activators, mutant BMDMs displayed elevated caspase-1 and

Gasdermin D (GSDMD) cleavage, as well as increased caspase-11 and IL-1β expression, after

incubation with PAO1 as compared with WT controls (Fig. 3e). Moreover, AcYVAD treatment

inhibited the hypersecretion of IL-1β and other pro-inflammatory cytokines in Polg[D257A]

BMDMs challenged with PAO1 for 24 hours (Fig. 3f). Notably, IL-1α secretion was markedly

higher in Polg[D257A] macrophages after PAO1 challenge (Fig. 1i and 3f). IL-1α is an alarmin that

feeds forward to augment production of other cytokines, and the robust secretion of IL-1α from

myeloid cells involves caspase-11[37–39]. Thus, given elevated caspase-11 expression in Polg[D257A]

BMDMs, we next explored its involvement in cell death and cytokine secretion. After LPS

priming followed by LPS transfection, Polg[D257A] BMDMs rapidly became PI positive and

secreted increased levels of IL-1β (Fig. 3g-h). Both phenotypes could be markedly inhibited by

treatment with AcYVAD, indicating that heightened activation of caspase-11 triggers NLRP3

inflammasome and caspase-1 activation more robustly in Polg[D257A] BMDMs (Fig. 3g-h). As

caspase-11 activation requires IFN-I priming[40] and Polg[D257A] macrophages and mice exhibit

elevated IFN-I signaling[20], we reasoned that IFN-I is responsible for caspase-11 expression and

activation in Polg^{D257A} BMDMs. In agreement, we observed potentiated caspase-11 mRNA and

protein after PAO1 infection (Fig. 3e, i). Notably, IFN alpha receptor (IFNAR) blockade using

the MAR1-5A3 monoclonal antibody markedly reduced PI uptake after caspase-11 activation

(Fig. 3j), while also lowering pro-inflammatory cytokine hyperproduction in PAO1-challenged

Polg^{D257A} BMDMs (Supplementary Fig. 3m). Collectively, these data indicate that the IFN-I-

dependent potentiation of caspase-11 mediates caspase-1-dependent pyroptosis and

hyperinflammation in Polg^{D257A} macrophages exposed to PAO1.

Increased GBP expression and targeting to intracellular PAO1 in Polg^{D257A} macrophages

Prior work has shown that IFN-I signaling during bacterial infection drives the expression

of guanylate binding proteins (GBPs)[41], which facilitate innate immune responses and non-

canonical inflammasome activation by disrupting membranes of pathogen-containing vacuoles,

or pathogens themselves, to liberate LPS or other PAMPs[34,42–44]. Consistent with elevated IFN-I

signaling, RNA-seq analysis revealed more robust GBP expression in Polg^{D257A} BMDMs after

challenge with PAO1 (Fig. 4a). Immunofluorescence microscopy revealed increased GBP5

expression in resting Polg^{D257A} BMDMs (Fig. 4b-c), as well as during PAO1 challenge (Fig. 4d-

e). Notably, we observed a more robust colocalization of GBP5 with PAO1 challenged Polg^{D257A}

macrophages after staining with an antibody specific for PA LPS (Fig. 4d, f). Intracellular PAO1

in Polg^{D257A} BMDMs colocalized strongly with GBP5 and appeared more aggregated with

bacteria than in WT BMDMs. These findings suggest that elevated IFN-I in Polg^{D257A}

macrophages augments GBP expression, likely enhancing sensing of LPS via caspase-11 that

contributes to potentiated cell death and pro-inflammatory cytokine release.

A novel Polg^{R292C} mouse model incorporating a deleterious patient mutation exhibits macrophage mtDNA depletion, elevated IFN-I responses, increased pyroptosis, and hyper-inflammation after challenge with PAO1

To determine if our findings in Polg^{D257A} macrophages are more broadly applicable to other models of OXPHOS deficiency due to mtDNA perturbations, we leveraged a novel model of Polg-related mitochondrial disease harboring a single amino acid missense mutation (Arg292Cys) in exon 4 of the exonuclease domain of Polg (Fig. 5a), which is homologous to the human Arg309Cys (c.925C>T) pathogenic variant[45,46]. We generated these mice using CRISPR/Cas9 editing of ES cells and verified germline transmission of the mutation by sequencing (Fig. 5b). The Arg309Cys variant alone or in combination with other Polg variants causes reduced mtDNA copy number, mtDNA deletions, and severe phenotypes including mitochondrial myopathy, encephalopathy, lactic acidosis and stroke-like episodes (MELAS), peripheral neuropathy, ataxia, and epilepsy in patients. Although being born below expected mendelian ratios, homozygous Polg^{R292C} mice are viable (data not shown, data to be reported in subsequent manuscript). BMDMs derived from Polg^{R292C} mice did exhibit mtDNA depletion at rest (Fig. 5c), as well as lower numbers of nucleoids and fragmented mitochondrial networks after stimulation with LPS (Fig. 5d-e). In line with our data from the Polg^{D257A} mutator mice, Polg^{R292C} BMDMs displayed a constitutive elevation of ISGs at rest and potentiated ISG expression after LPS stimulation (Supplementary Fig. 4a-b). Moreover, Polg^{R292C} BMDMs exhibited increased ISG and GBP expression after challenge with PAO1 (Fig. 5f), while also exhibiting more pyroptosis (Fig. 5g-h) and pro-inflammatory cytokine secretion (Fig. 5i). These data suggest that elevated IFN-I, hyperinflammation, and cell death in response to PAO1 are

common features of macrophages harboring Polg exonuclease domain mutations, including the

mitochondrial disease-relevant Arg292Cys variant.

Intratracheal instillation of a non-lethal dose of PAO1 elicits greater lung cytokine

production and modest changes in lung immune cell populations in Polg mutant mice

Patients with primary MtD often experience recurrent viral and bacterial infections of the

respiratory tract, including infections by PA[5]. These events can trigger metabolic

decompensation and can progress to systemic inflammatory endpoints and/or sepsis. However,

few studies have examined infection kinetics and lung immune responses in Polg-related

mitochondrial disorders. We therefore employed intratracheal instillation protocols to administer

LPS or PAO1 into the lungs of Polg^{D257A} mutator mice. Histological analysis of resting Polg^{D257A}

lungs did not reveal obvious differences in morphology or cellular composition (Supplementary

Fig. 4c-e), although consistent with our earlier findings[20], basal expression of ISGs were elevated

compared to WT littermates (Supplementary Fig. 4f). After LPS was instilled into lungs through

cannulation of the trachea, Polg^{D257A} mice secreted more TNFα and IL-6 at all timepoints and

increased IL-1β at the 12 hour timepoint in the bronchial alveolar lavage fluid (BAL) compared

to WT mice (Supplementary Fig. 4g).

To next characterize lung innate immune responses to PAO1 infection in the Polg^{D257A}

mice, we challenged mice with a non-lethal dose of PAO1[47]. Both WT and Polg^{D257A} mice

infected with 1×10^4 colony forming units (CFU) of PAO1 via intratracheal instillation lost

weight initially, but returned to normal weight (Supplementary Fig. 5a) and exhibited no

culturable bacteria (data not shown) 3 days post infection. Flow cytometric analysis of lung cells

in BAL fluid revealed an overall decrease in the percentage of CD45$^+$ immune cells at rest and

after infection, consistent with general leukopenia noted in Polg[D257A] mice[20,25] (Supplementary

Fig. 5b-c). Analysis of specific immune cell markers on lung CD45[+] cells revealed few changes

in BAL populations (Supplementary Fig. 5d-l) between uninfected WT and Polg[D257A] mice,

although there were more SiglecF[+] and F4/80[+] cells at rest perhaps indicating elevated numbers

of lung macrophages (Supplementary Fig. 5e,g). Six hours post infection, there was a marked

reduction in SiglecF[+] cells in Polg[D257A] mice relative to WTs and an increase in CD3[+] T cells

(Supplementary Fig. 5e,i). Cells labeling with the neutrophil marker, Ly6G, were dramatically

increased at 6 hours, although there were no differences between WT and Polg[D257A] mice

(Supplementary Fig. 5k). At 48 hours post infection, neutrophils remained the most predominant

CD45[+] population in both genotypes, but we observed a reversion toward increased numbers of

cells staining positive for macrophage markers SiglecF[+] and F4/80[+] in Polg[D257A] mice

(Supplementary Fig. 5e,g,k).

Despite modest changes in cellular composition of the BAL fluid, Polg[D257A] mice had

elevated numbers of CD45[+] cells expressing TNFα both at rest and six hours post infection

(Supplementary Fig. 5m), consistent with increased TNFα production after LPS challenge.

Multianalyte analysis of the BAL fluid using Legendplex assays revealed 2-4 fold increases in

lung pro-inflammatory cytokines in Polg[D257A] mice 6 hours post PAO1 infection (Fig. 6a).

Consistent with resolution of infection in both genotypes, cytokine differences in the BAL of

WT and Polg[D257A] mice were less pronounced at 48 hours post infection (Supplementary Fig. 6a).

However, IFNβ and inflammatory cytokines associated with the skewing and differentiation of T

cell Th1 (IL-27) and Th17 subsets (IL-23) remained elevated in Polg[D257A] mice in an IFNAR-

dependent fashion (Supplementary Fig. 6a). Analysis of lung protein extracts revealed elevated

caspase-1 cleavage, as well as increased caspase-11, NLRC4, and NLRP3 expression in

Polg^{D257A} mice 48 hours post intratracheal instillation of 1×10^4 CFU of PAO1. GSDMD, GBP5,

and other ISGs were also elevated in Polg^{D257A} lungs, consistent with sustained IFNβ expression

observed by Legendplex (Fig. 6b).

Given that recurrent respiratory tract infections are often observed in patients with MtD,

we next infected mice with 1×10^4 CFU PAO1, followed by a 7-day rest period, then re-infected

with 1×10^4 CFU PAO1. Histological analysis of lungs 24 hours following re-infection revealed

monocyte and granulocyte infiltration into lungs, areas of vascular hemorrhage, and more Eosin

positive cells in Polg^{D257A} lungs relative to WT (Supplementary Fig. 6b-c). Lung IL-12 family

cytokines and IFNβ were significantly higher in Polg^{D257A} mice at the 24 hour harvest point

following re-infection, while other cytokines were trending higher in Polg^{D257A} mice, although

they did not reach statistical significance (Supplementary Fig. 6d). We also observed higher

bacterial burdens in the lungs and spleens of Polg^{D257A} mice compared to WT controls

(Supplementary Fig. 6e). Altogether, these data indicate that Polg^{D257A} mice exhibit

hyperinflammation and impaired antibacterial immunity upon re-infection with PAO1.

Intratracheal infection with a higher dose of PAO1 elicits greater lung immune cell

infiltration, caspase-11 expression, and death of Polg mutant mice

We next utilized a higher dose of PAO1 to expand on low-dose data and assess mortality

in Polg^{D257A} mice. Upon intratracheal instillation of 1×10^5 CFU PAO1, Polg^{D257A} mutant mice

displayed elevated neutrophilic inflammation and overall lung inflammatory scores via

histological analysis (Fig. 6c-e) at 20 hours post infection, despite no differences in bacterial

burden (Supplementary Fig. 6f). We also observed heightened expression of caspase-11 in lung

lysates of Polg^{D257A} mutant mice (Fig. 6f), consistent with our findings in macrophages and lungs

exposed to lower doses of PAO1 (Fig. 6a-b). Moreover, pro-IL-1β and the mature cytokine were

elevated in Polg^{D257A} lungs 20 hours after infection with 1×10^5 CFU of PAO1 (Fig. 6f). Finally,

we observed that a greater number of Polg^{D257A} mice reached a moribund state after infection

with this higher dose of PAO1 relative to WT littermates (Fig. 6g). Overall, these in vivo data

suggest that Polg^{D257A} mice are more prone to excessive inflammation and acute lung injury

following infection with PAO1. We therefore conclude that macrophage and myeloid cell

hyperinflammation driven by caspase-11 and other innate immune pathways promotes lung

damage that alters antibacterial responses and increases mortality in the Polg^{D257A} model of

mitochondrial dysfunction (Fig. 6h).

<u>Discussion</u>

Patients with MtD experience a greater frequency of bacterial and viral infections that

can rapidly progress to sepsis or systemic inflammatory response syndrome (SIRS), leading to

premature death[5]. Recurrent infections can also accelerate MtD progression by promoting

metabolic decompensation, a life-threatening condition that hastens deterioration of

mitochondrial function[48–50]. Polg is the sole mitochondrial DNA polymerase, and mutations in

POLG are the most prevalent single-gene cause of MtD[51]. There are several clinical reports of

bacterial or viral infections unmasking Polg-related MtD symptoms and hastening disease

progression[52,53]. Moreover, one study revealed the presence of inflammatory cytokines in the

cerebrospinal fluid of a patient with Polg-related MtD[12,19]. As cytokine-mediated toxicity is an

important driver of metabolic decompensation[7], there is a critical need to understand immune

responses in patients with Polg-related disorders and other MtD to manage infections and

inflammation-related complications.

The present study builds upon our prior work to characterize innate immunity in mouse

models of MtD[20,54] and leverages bacterial infection, cytokine profiling, histological analyses,

and in vitro and ex vivo cellular assays to define macrophage inflammatory dynamics in

Polg[D257A] mutator mice. We also utilize macrophages from Polg[R292C] mice that carry a mutation

observed in patients to broaden our findings into a more faithful model of Polg-related MtD.

Macrophages from Polg[D257A] mutator and our novel Polg[R292C] mice exhibit elevated ISG

expression and secrete more pro-inflammatory cytokines after challenge with PA. Notably,

inclusion of DMF, an immunomodulatory and Nrf2-stabilizing agent, during PAO1 challenge

significantly reduced IFN-I and pro-inflammatory signatures. These findings are consistent with

our previous work implicating Nrf2 pathway inhibition as a major driver of hyperinflammatory

signatures in Polg mutant macrophages and mice[20]. We also observed that DMF reduced PA-

induced cell death and hypersecretion of IL-1α and IL-1β in Polg mutant macrophages. By

activating Nrf2 and/or inhibiting NF-κB, DMF reduces transcription of multiple pro-

inflammatory cytokine genes including *Il1b*[55,56]. However, DMF can also directly impair NLRP3

activation and GSDMD-mediated pyroptosis[57,58], leading us to directly examine inflammasome

activation in our Polg mutant models.

Our data reveal that Polg mutant macrophages have an increased propensity to undergo

rapid pyroptosis upon PAO1 challenge in a caspase-1-dependent fashion. Although caspase-1

cleavage is elevated in Polg[D257A] over WT macrophages during PAO1 challenge and its

inhibition is sufficient to block augmented PA-induced cell death and IL-1β secretion, we found

that Polg[D257A] BMDMs are less responsive to direct activation of the NLRP3 and NLRC4

inflammasomes by their respective ligands. In contrast, activation of the non-canonical caspase-

11 inflammasome via LPS priming followed by transfection of LPS into the cytosol drove more

PI uptake and IL-1β expression in Polg[D257A] BMDMs. Caspase-11 expression is significantly

elevated in Polg mutant BMDMs, collectively indicating that non-canonical inflammasome

activation by cytosolic PA LPS leads to more robust activation of the NLRP3 inflammasome for

caspase-1 dependent cell death and cytokine secretion. We hypothesize that increased caspase-

11-dependent cleavage of GSDMD may lead to more rapid permeabilization of mitochondrial

membranes in Polg mutant BMDMs, leading to augmented release of oxidized mtDNA or other

mitochondrial DAMPs that potentiate activation of NLRP3[59–61]. It is also possible that

potentiated release of mitochondrial RNA or mtDNA, or increased vita-PAMP accumulation in

Polg BMDMs, may enhance caspase-11 and NLRP3 interaction, leading to elevated pyroptosis

and pro-inflammatory cytokine secretion[62]. Finally, it is interesting to note that treatment of Polg

mutant macrophages with caspase-1 inhibitor dramatically reduced most cytokine levels in

culture supernatants after challenge with PAO1 (**Fig. 3f**).   These included inflammasome-

independent cytokines such as TNFα, IL-6, IFNγ, and IL-12 family members as well as the

DAMP IL-1α, which does not require caspase-1 cleavage but is downstream of caspase-11 and

GSDMD[37,63]. This is in contrast with cytokine secretion in WT BMDMs, where AcYVAD

treatment only reduced IL-1β levels during challenge with PAO1. Overall, these data suggest key

roles for caspase-11 and caspase-1 in the broad hyperinflammatory phenotype of Polg mutant

mice.

We have also found that elevated IFN-I signaling is a key driver of caspase-11-dependent

pyroptosis and hyperinflammation in Polg mutants. Inclusion of anti-IFNAR neutralizing

antibodies was sufficient to reduce both augmented PI uptake after LPS transfection and

cytokine hypersecretion after challenge with PAO1. Our prior work revealed that increased IFN-

I signaling in Polg[D257A] macrophages depends on chronic cGAS-STING signaling, and here we

show that Polg[R292C] macrophages also exhibit increased expression of ISGs at rest and after

challenge with PAO1[20]. Thus, alterations to IFN-I signaling appear to be a common feature in

Polg mutant mice, human fibroblasts harboring the *POLG1* c.2243G>C, p.W748S variant[64], and

peripheral blood mononuclear cells from patients with diverse MtD mutations[11]. IFN-I is critical

to license caspase-11 expression[40], but also to enhance expression of interferon-inducible

GTPases like GBPs, which help to expose cytosolic LPS for detection[34,65–68]. We have found that

macrophages from both the Polg[D257A] and our new MtD-relevant Polg[R292C] macrophages exhibit

enhanced expression of many GBPs at rest and after challenge with PAO1. Moreover, we

observed that GBP5 strongly co-localizes to intracellular PAO1 in Polg mutant BMDMs,

suggesting that IFN-I mediated triggering of GBPs potentiates LPS sensing by caspase-11 to

augment cytokine secretion.

Although it is appreciated that bacterial and viral infections are problematic for the MtD

population due to their propensity to trigger systemic hyperinflation and disseminated disease,

few studies have examined immune responses in relevant mouse models of MtD. Using an

intratracheal instillation model, we find that Polg[D257A] mice exhibit hyperactive innate immune

responses resulting in elevated cytokine production, changes in lung immune cell dynamics

recruitment, and increased susceptibility to death following PAO1 infection. Notably, we

observed elevated expression of inflammasome proteins, including caspase-11, ISGs including

GBPs, and increased caspase-1 cleavage in Polg[D257A] lung homogenates after infection. These

data agree with our in vitro findings implicating IFN-I as an overarching regulator of caspase-

1/11 activation and hyperinflammation in our models. Additional studies are warranted to define

the cellular populations that govern lung hyperinflammation and to determine the degree to

which caspase-11 shapes inflammatory signaling and morbidity in Polg mutant mice in vivo.

Overall, our study provides important new insight into the innate immune mechanisms governing

infection-related hyperinflammation in mouse models Polg MtD and furthers our understanding

of why patients with MtD are more prone to developing sepsis and SIRS following infection.

DMF and/or other therapeutics targeting IFN-I and caspase-11 may therefore represent

promising avenues to restore proper immune balance in the MtD population.

Figures and Legends

Fig. 1: Polg^{D257A} mutator macrophages exhibit increased IFN-I and pro-inflammatory

responses after challenge with LPS or *Pseudomonas aeruginosa*.

(a) Workflow for preparing RNA from Polg BMDMs after *Pseudomonas aeruginosa* (PA)

strain O1 (PAO1) challenge.

(b) Ingenuity Pathway Analysis (IPA) of RNA-seq from bone marrow-derived macrophages

(BMDMs) showing Z-scores of pathways in Polg^{D257A} homozygous mutant BMDMs compared

to wild-type (WT) after LPS, PAO1, or PAO1 plus dimethyl fumarate (DMF) challenge.

hpc=hours post challenge. N=3 biological replicates/genotype/condition.

(c) Log^2 fold changes of representative genes in IPA pathways from (**b**) in Polg^{D257A} BMDMs compared

to WT after LPS, PAO1, or PAO1 plus DMF.

(d) Protein abundance of interferon-stimulated genes (ISGs) in BMDMs exposed to PAO1 at a

multiplicity of infection (MOI) of 5 at the indicated hours post challenge (hpc).

(e) Cytokines in supernatant of peritoneal macrophages (PerMacs) measured by ELISA. MOI=

multiplicity of infection, N=3 biological replicates per time point.

(f) IPA Upstream Regulator analysis showing Z-scores in Polg^{D257A} homozygous mutant

BMDMs compared to WT after LPS, PAO1, or PAO1 plus dimethyl fumarate (DMF) challenge.

(g) Log^2 fold changes Nrf2 target genes from IPA pathway **(b)** in Polg^{D257A} BMDMs compared to WT

after LPS, PAO1, or PAO1 plus DMF.

(h) qRT-PCR analysis of ISGs in WT and Polg^{D257A} BMDMs after PAO1 or PAO1 plus DMF. N=3

biological replicates, 3 technical replicates/genotype/condition.

(i) Legendplex pro-inflammatory cytokines analysis of PerMac supernatants 24 hours post

challenge (hpc). N=3 biological replicates

Statistics: (e, h, & i) Two-Way ANOVA. *P<0.05, **P<0.01, ***P<0.001, ****P<0.0001 Error

bars represent SEM.

Fig. 2: Polg^{D257A} mutant macrophages exhibit potentiated cell death upon PAO1

challenge.

(a) Ingenuity Pathway Analysis (IPA) of RNA-seq from bone marrow-derived macrophages

(BMDMs) showing Z-scores of pathways related to cell death signaling in Polg^{D257A}

homozygous mutant BMDMs compared to wild-type (WT) after LPS, PAO1, or PAO1 plus

dimethyl fumarate (DMF) challenge. hpc=hours post challenge. N=3 biological

replicates/genotype/condition.

(b) Log^2 fold changes of representative genes in IPA pathways from **(a)** in Polg^{D257A} BMDMs compared

to WT after LPS, PAO1, or PAO1 plus DMF.

(c) Representative immunofluorescent (IF) images of peritoneal macrophages (PerMacs) after

PAO1 or PAO1 plus DMF. Cells were stained with DAPI to mark the nucleus, CellMask to mark

plasma membranes, and anti-GFP antibody to mark GFP-tagged PAO1. White arrows highlight

dying cells characterized by condensed nuclei and loss of CellMask staining.

(d) Percent of PerMacs dying after PAO1 and PAO1 plus DMF in **(c)** IF. N= 5 IF

images/genotype/condition.

(e) Cytation5 Image analysis of PI positive BMDMs after PAO1 or PAO1 plus DMF. N= 3

biological replicates, 5 technical replicates/genotype/condition.

(f) LegendPlex pro-inflammatory cytokines analysis of PerMac supernatants at 8 and 24 hpc in

the presence or absence of DMF. N=3 biological replicates/genotype/condition.

Statistics: (e & f) Two Way ANOVA (d) One Way ANOVA. *P<0.05, **P<0.01, ***P<0.001,

****P<0.0001 Error bars represent SEM.

Fig. 3: Caspase-11 and caspase-1 drive pyroptosis and inflammation in Polg mutant

macrophages challenged with PAO1.

(a) ELISA analysis of inflammatory cytokines in BMDM supernatant after PAO1 or PAO1 plus

caspase inhibitors. N= 3 biological replicates/genotype/condition.

(b) Cytation5 image analysis of PI positive BMDMs after PAO1 or PAO1 plus caspase-1

inhibitor. N= 3 biological replicates, 5 technical replicates/genotype/condition.

(c) Percent PerMacs dying after PAO1 and PAO1 plus caspase-1 inhibitor from IF **(d)**. N=4 IF

images/genotype/condition.

(d) Representative IF of PerMacs after PAO1 or PAO1 plus caspase inhibitor. Cells were stained

with DAPI to mark the nucleus, CellMask to mark the total cell volume, and anti-GFP to mark

GFP tagged PAO1. White arrows highlight dying cells characterized by condensed nuclei and

loss of CellMask staining.

(e) Western blot analysis of lysates from BMDMs challenged with PAO1.

(f) LegendPlex pro-inflammatory cytokines analysis of BMDM supernatant 24 hpc. All

undefined comparisons between WT and WT+AcYVAD are not significant. N=3 biological

replicates/genotype/condition.

(g) Cytation5 image quantification of PI positive BMDMs after LPS priming plus LPS

transfection and LPS priming plus LPS transfection plus caspase-1 inhibitor. N= 3 biological

replicates, 5 technical replicates/genotype/condition.

(h) ELISA analysis of IL-1β in BMDM supernatant after LPS priming plus LPS transfection in

the presence or absence of caspase-1 inhibitor. N=3 biological replicates

(i) qRT-PCR analysis of Caspase-11 RNA expression in BMDMs after PAO1. N=3 biological

replicates, 3 technical replicates/genotype/condition.

(j) Cytation5 image quantification of PI positive cells in BMDMs after LPS priming plus LPS

transfection and incubated with αIFNAR blocking antibody or αIgG control. N= 3 biological

replicates, 5 technical replicates/genotype/condition.

Statistics: (c, f, & h) One-Way ANOVA (a, b, g, i & j) Two-Way ANOVA. *P<0.05, **P<0.01,

***P<0.001, ****P<0.0001 Error bars represent SEM.

Fig. 4: Increased GBP expression and targeting to intracellular PAO1 in Polg^{D257A}

macrophages.

(a) RNA-seq from BMDMs showing induction of Gbps and Caspase-11 RNA 6 hours post

challenge (hpc) with PAO1. N=3 biological replicates/genotype/condition.

(b) Representative IF of unchallenged BMDMs stained with DAPI to mark the nucleus and anti-

GBP5 antibody.

(c) Sum intensity of GBP5 per cell area in resting BMDMs in (**b**) IF. N=7 images/genotype

(d) Representative IF of BMDMs 2 hpc with PAO1 at a multiplicity of infection (MOI) of 5.

Cells were fixed, permeabilized, and stained with DAPI to mark the nucleus, anti-LPS antibody

to stain PAO1 LPS, and anti-GBP5 antibody.

(f) Sum intensity of GBP5 per cell area in BMDMs after PAO1 in **(d)** IF. N=7 images/genotype.

(e) Sum intensity of PAO1-LPS localized with GBP5 in BMDMs after PAO1 in **(d)** IF. N=7

images/genotype.

Statistics: (a, c, e, & f) Student's t-test. *P<0.05, **P<0.01, ***P<0.001, ****P<0.0001 Error

bars represent SEM.

Fig. 5: Macrophages incorporating a deleterious patient mutation (Polg^{R292C}) exhibit

mtDNA depletion, elevated IFN-I responses, increased pyroptosis, and hyper-inflammation

after challenge with PAO1

(a) Generation of the Polg^{R292C} mouse.

(b) Chromatograph from Sanger sequencing of DNA from heterozygous Polg^{R292C} mouse

verifying germline incorporation of mutation.

(c) qPCR analysis of mtDNA abundance in BMDMs. N= 3 biological replicates, 3 technical

triplicates/genotype.

(d) Representative IF of nucleoids in BMDMs stained with anti-PDH antibody to label

mitochondria, anti-DNA antibody to label nucleoids, and anti-TFAM antibody.

(e) Quantification of nucleoids per cell, mean intensity of DNA signal per nucleoid, and nucleoid

area per cell from BMDMs in **(d)** IF. N=40-50 cells/genotype

(f) qRT-PCR analysis of ISGs in BMDMs after PAO1, N=3 biological replicates, 3 technical

replicates/genotype.

(g) Representative IF of BMDMs after PAO1 stained with DAPI to mark the nucleus, CellMask

to mark the total cell volume, and anti-GFP to mark GFP tagged PAO1. White arrows highlight

dying cells characterized by condensed nuclei and loss of CellMask staining.

(h) Percent BMDMs dying after PAO1 in **(g)** IF. N=7 IF images/genotype.

(i) ELISA analysis of inflammatory cytokines in BMDM supernatant after PAO1. N= 3

biological replicates.

Statistics: (c, e, f, & h) Students t-test. (i) Two-Way ANOVA. *P<0.05, **P<0.01, ***P<0.001,

****P<0.0001 Error bars represent SEM.

Fig. 6: Intratracheal instillation of PAO1 into lungs elicits greater inflammation, caspase-

11 expression, and death of Polg mutant mice.

(a) LegendPlex pro-inflammatory cytokine analysis of bronchial alveolar lavage (BAL) 6 hours

post infection (hpi) with $1x10^4$ colony forming units (CFU) of PAO1. N=5 mice/genotype

(b) Protein expression or WT and Polg^{D257A} lung homogenates 48 hpi with $1x10^4$ CFU of PAO1.

(c) Representative hematoxylin and eosin (H&E) staining of lung sections 20 hpi with $1x10^5$

CFU of PAO1.

(d) Pathology scoring of lung sections from **(c)**.

(e) Graphs of pathology scoring from H&E stained lungs in **(d)** histology. N=5 mice/genotype

(f) Protein expression or WT and Polg^{D257A} lung homogenates 20 hpi with $1x10^5$ CFU of PAO1.

(g) Kaplan-Meier survival curve after infection with $1x10^5$ CFU of PAO1. N= 7 mice/genotype

(h) Graphical abstract depicting findings of the study.

Statistics: (a & e) Students t-test (g) Kaplan-Meijer Survival Curve. *P<0.05, **P<0.01,

***P<0.001, ****P<0.0001 Error bars represent SEM.

Supplementary Fig. 1: Polg^{D257A} macrophages exhibit elevated IFN-I and pro-inflammatory

gene expression when challenged with LPS or PAO1.

(a) Log^2 fold changes of representative genes in IPA pathways from (**Fig. 1b**) in Polg^{D257A}

BMDMs compared to WT after challenge with LPS, PAO1, or PAO1 plus DMF. N=3 biological

replicates/genotype/condition.

(b) Gene Set Enrichment (GSEA) pathways of RNA-seq data in WT ang Polg^{D257A} BMDMs

after challenge with PAO1 and PAO1 plus DMF. Heatmaps represent fold changes of each group

relative to WT PAO1 infected. N=3 biological replicates/genotype/condition.

Supplementary Fig. 2: Polg^{D257A} macrophages exhibit increased inflammatory cell death when

exposed to bacteria.

(a) Representative IF of BMDMs unchallenged or primed with LPS for 4 hours and stimulated with

nigericin for 1 hour to activate the NLRP3 inflammasome. Cells are stained with DAPI to mark the

nucleus, CellMask to mark total cell volume, and anti-ASC antibody to mark puncta.

(b) Number of ASC puncta per cell in cells with high and low CellMask staining in **(a)** IF. N= 7

images/condition.

(c) PAO1 colony forming units obtained from BMDM supernatant or detergent lysates 2 and 4 hours post

challenge (hpc). N=4 biological replicates.

(d) Beads per cell quantified from **(e)** IF. N= 100 cells/genotype.

(e) Representative IF of BMDMs phagocytosing LPS coated, fluorescent microspheres. Cells were

stained with DAPI to mark the nucleus, CellMask to label the cell volume, and anti-LAMP1 to mark

phagolysosomes.

(f) ELISA analysis of inflammatory cytokines in BMDM supernatants after challenge with

Escherichia coli strain DH5α. N= 3 biological replicates.

(g) Percent BMDMs dying after challenge with DH5α **(h)** IF. N=7 IF images/genotype.

(h) Representative IF of BMDMs stained with DAPI to mark the nucleus, CellMask to mark the total cell

volume, and anti-LPS to mark DH5α bacteria. White arrows highlight dying cells characterized by

condensed nuclei and loss of CellMask staining.

Statistics: (c & f) Two-way ANOVA (b, d, & g) Students t-test. *P<0.05, **P<0.01,

***P<0.001, ****P<0.0001 Error bars represent SEM.

Supplementary Fig. 3: Assessment of roles for canonical inflammasome activity, metabolism, and

IFN-I dependence in driving increased cell death and pro-inflammatory cytokine secretion in Polg

mutant macrophages.

(a) Cytation5 imaging quantification of PI positive cells in BMDMs after LPS priming for 4 hours. N= 3

biological replicates, 5 technical replicates/genotype.

(b) Cytation5 imaging quantification of PI positive cells in BMDMs after LPS priming plus PA-flagellin

transfection. N= 3 biological replicates, 5 technical replicates/genotype.

(c) ELISA analysis of IL-1β in BMDM supernatant after LPS priming plus PA-flagellin transfection. N=

3 biological replicates.

(d) Cytation5 imaging quantification of PI positive cells in BMDMs after LPS priming plus ATP. N= 3

biological replicates, 5 technical replicates/genotype.

(e) ELISA analysis of IL-1β in BMDM supernatant after LPS priming plus ATP. N= 3 biological

replicates.

(f) Western blot of BMDM protein lysates after LPS priming plus ATP.

(g) Cytation5 imaging quantification of PI positive cells in BMDMs after LPS priming plus nigericin. N=

3 biological replicates, 5 technical replicates/genotype.

(h) ELISA analysis of IL-1β in BMDM supernatant after LPS priming plus nigericin. N= 3 biological

replicates.

(i) Western blot of BMDM protein lysates after LPS priming plus nigericin.

(j) & **(k)** Seahorse analysis of BMDMs unchallenged or treated with overnight LPS.

Extracellular acidification rate (ECAR) **(j)** and oxygen consumption rate (OCR) **(k)** are plotted.

N=3 biological replicates and 5 technical replicates.

(l) Quantification of Seahorse OCR measurements. N=3 biological replicates and 5 technical

replicates.

(m) ELISA analysis of inflammatory cytokines in BMDM supernatant after PAO1 challenge in the

presence of αIFNAR neutralizing antibody or αIgG. N= 3 biological replicates.

Statistics: (b, e, & h) Two-Way ANOVA (l & m) One-Way ANOVA (a, c, f, & i) Students t-test.

*P<0.05, **P<0.01, ***P<0.001, ****P<0.0001 Error bars represent SEM.

Supplementary Fig. 4: Polg mutant mice exhibit elevated IFN-I and inflammatory

responses to LPS, but do not exhibit basal differences in lung morphology or immune cell

abundance.

(a) qRT-PCR analysis of ISGs in unchallenged Polg^{R292C} BMDMs. N=3 biological replicates, 3 technical

replicates/genotype.

(b) qRT-PCR analysis of ISGs in Polg^{R292C} BMDMs 8 hours post challenge (hpc) with LPS. N=3

biological replicates, 3 technical replicates/genotype.

(c) Representative histology images of hematoxylin and eosin (H&E) staining of WT and Polg^{D257A} lungs,

with an overlay of eosin positive cells using QuPath software.

(d) Pathology assessment of unchallenged lungs.

(e) QuPath analysis of eosin$^+$ cells in the lungs from **(c)**. N=3 mice/genotype

(f) Protein expression of ISGs in unchallenged WT and Polg^{D257A} lungs homogenates.

(g) ELISA analysis of inflammatory cytokines in bronchioalveolar lavage of LPS challenged WT

and Polg^{D257A} lungs. N=2 mice/genotype/timepoint.

Statistics: (g) Two-Way ANOVA (a, b, & e) Students t-test. *P<0.05, **P<0.01, ***P<0.001,

****P<0.0001 Error bars represent SEM.

Supplementary Fig. 5: Intratracheal instillation of PAO1 alters immune cell composition in

Polg mutant mouse lungs.

(a) Weight change in WT and Polg^{D257A} mice after infection with 1×10^4 PAO1. N=5

mice/genotype.

(b) CD45 expression on cells from bronchioalveolar lavage (BAL) from uninfected and PAO1

infected WT and Polg^{D257A} mice. N=5-9 mice/genotype/timepoint.

(c) Gates used to quantify CD45 expression in **(b)**.

(d) Percent of CD45$^+$ cells expressing CD11c from the BAL of uninfected and PAO1 infected

lungs. N=5-9 mice/genotype/timepoint.

(e) Percent of CD45$^+$ cells expressing SiglecF from the BAL of uninfected and PAO1 infected

lungs. N=5-9 mice/genotype/timepoint.

(f) Percent of CD45$^+$ cells expressing MHCII from the BAL of uninfected and PAO1 infected

lungs. N=5-9 mice/genotype/timepoint.

(g) Percent of CD45$^+$ cells expressing F4/80 from the BAL of uninfected and PAO1 infected

lungs. N=5-9 mice/genotype/timepoint.

(h) Percent of CD45$^+$ cells expressing CD11b from the BAL of uninfected and PAO1 infected

lungs. N=5-9 mice/genotype/timepoint.

(i) Percent of CD45$^+$, CD11b$^-$ cells expressing CD3 from the BAL of uninfected and PAO1

infected lungs. N=5-9 mice/genotype/timepoint.

(j) Percent of CD45$^+$, CD11b$^-$ cells expressing CD19 from the BAL of uninfected and PAO1

infected lungs. N=5-9 mice/genotype/timepoint

(k) Percent of CD45$^+$ expressing Ly6G from the BAL of uninfected and PAO1 infected lungs.

N=5-9 mice/genotype/timepoint.

(l) Percent of CD45$^+$ cells expressing Ly6C from the BAL of uninfected and PAO1 infected

lungs. N=5-9 mice/genotype/timepoint.

(m) CD45$^+$ cells with cell associated TNFα expression from BAL of uninfected and PAO1

infected lungs. N=5-9 mice/genotype/timepoint.

Statistics: (a) Two-way ANOVA (b, & d-m) Students t-test. *P<0.05, **P<0.01, ***P<0.001,

****P<0.0001 Error bars represent SEM.

Supplementary Fig. 6: Polg mutant mice exhibit hyperinflammation in response to

infection and reinfection with PAO1.

(a) LegendPlex pro-inflammatory cytokine analysis of bronchioalveolar lavage (BAL) from WT,

Polg^{D257A}, WT IFNAR$^{-/-}$, and Polg^{D257A} IFNAR$^{-/-}$ lungs infected with 1x10^4 PAO1 for 48 hours.

All undefined comparisons are not significant. N=5 mice/genotype.

(b) Representative histology images of hematoxylin and eosin (H&E) staining of WT and Polg^{D257A} lungs

after initial infection with 1x10^4 PAO1, followed by 7 days rest, then reinfection with 1x10^4 PAO1.

Overlay of eosin positive cells was done using QuPath software.

(c) QuPath analysis of eosin$^+$ cells in the lung from **(b)** histology. N=5 mice/genotype.

(d) LegendPlex pro-inflammatory cytokine analysis of BAL from WT and Polg^{D257A} lungs after

initial infection with 1x10^4 PAO1, followed by 7 days rest, then reinfection with 1x10^4 PAO1. N=5-6

mice/genotype.

(e) Colony counts from homogenized tissue after initial infection with $1x10^4$ PAO1, followed by 7

days rest, then reinfection with $1x10^4$ PAO1. N=5 mice/genotype.

(f) Colony counts from homogenized tissue infected with $1x10^5$ PAO1. N=5 mice/genotype.

Statistics: (a) Two-way ANOVA (c, d, e, & f) Students t-test. *P<0.05, **P<0.01, ***P<0.001,

****P<0.0001. Error bars represent SEM.

Acknowledgements

We thank R. Moore at the Texas A&M School of Medicine Analytical Cytometry Core for flow

cytometry assistance.

Funding

This research was supported by awards W81XWH-17-1-0052 to A.P.W. and W81XWH-20-1-

0150 to C.C. and A.P.W. from the Office of the Assistant Secretary of Defense and for Health

Affairs through the Peer Reviewed Medical Research Programs. Additional support was

provided by NIH grants R01HL148153 to A.P.W., U54OD030187 to S.A.M, and a gift from the

POLG Foundation. J.J.V. was supported by NIH NRSA F31AI179168 and Y. Lei was supported

by American Heart Association Predoctoral Fellowship grant 825908.

Author contributions

A.P.W and J.J.V. designed experiments, analyzed data, and wrote the manuscript. Y.L. assisted

with animal experiments and sample processing. C.G.M. performed inflammasome challenge

experiments. J.W. assisted with pathology scoring of histology samples. C.P. and R.S.

maintained animals and generated macrophages. S.C.K. and S.A.M. generated the $Polg^{R292C}$

mouse model and provided samples. P.J.M. and C.L.C. assisted with data analysis and

interpretation. A.P.W. conceived the project and provided overall direction.

Competing interests

The authors declare that they have no competing interests.

Data and materials availability

All data needed to evaluate the conclusions drawn herein are present in the paper and

Supplementary Materials. Raw data will be provided upon request. The RNA sequencing

datasets generated through this work will be deposited in the Gene Expression Omnibus after

acceptance of the paper for publication.

Methods

Mouse Husbandry and Strains

C57BL/6J (JAX Stock # 000664), $Polg^{D257A/D257A}$ (JAX Stock # 017341), and $Ifnar1^{-/-}$

(JAX Stock # 028256) mice were obtained from The Jackson Laboratory and bred and

maintained in multiple vivaria at Texas A&M University and The Jackson Laboratory.

Experimental $Polg^{+/+}$ and $Polg^{D257A/D257A}$ littermates were obtained from $Polg^{D257A/+}$ maintenance

colony crosses of male $Polg^{D257A/+}$ to female $Polg^{+/+}$. This breeding strategy was adopted to limit

the accumulation and maternal transmission of mutant mtDNA molecules to heterozygous

breeders.

The $Polg^{R292C/R292C}$ mouse line (B6N(Cg)-Polgem5(F290F,R292C)Lutzy/Mmjax) was

generated using CRISPR/Cas9 endonuclease-mediated genome editing in mouse embryonic stem

cells (mESCs). A single guide RNA (CTCCCATCCACAGGCTGCGC) targeting exon 4 of Polg

was cloned into pX459, which expresses Cas9 nuclease. This plasmid was co-delivered to

JM8A3 ESCs via electroporation with a single-stranded oligonucleotide donor (ssODN) that

included the missense R292C (CGC>TGC) and an additional silent mutation (F290F;

TTC>TTT) to reduce recutting of the targeted allele. Following puromycin selection, resistant

clones were screened for proper targeting and to exclude integration of the pX459 plasmid.

Multiple clones were selected for expansion and karyotyping, and euploid clones were injected

into C57BL/6J-albino blastocysts (B6(Cg)-Tyrc-2J/J; JAX STOCK# 000058) to produce

chimeras. Chimeras were bred to C57BL/6NJ (JAX Stock #005304) mice and a single germline

clone was selected for expansion and characterization. The line is available through the MMRRC

repository (071656-JAX).

All animal experiments were approved by the Institutional Animal care and Use

Committees (IACUC) at Texas A&M University and The Jackson Laboratory.

Cell Culture

L929 cells were obtained from ATCC and maintained in DMEM (Sigma-Aldrich,

D5796) and 10% fetal bovine serum (FBS; VWR, 97068-085). Primary bone marrow-derived

macrophages (BMDMs) were isolated from femurs of 6-12 month old WT, Polg^{D257A} and 2-6

month old Polg^{R292C} mice as described[20]. Briefly, whole femurs were crushed in media and

passed through a 70 µm filter. Red blood cells were lysed through osmosis by resuspending in

ACK lysis buffer (Quality Biological, 118-156-101) for 2 minutes. Remaining cells were passed

through a 40 µm filter, counted, and plated on non-tissue culture treated dishes. Macrophages

were differentiated over for 7 days using 30% L929 conditioned medium. Peritoneal

macrophages (PerMacs) were collected from the peritoneal cavity via gavage 4 days after

intraperitoneal injection of 3% brewer thioglycolate medium (Sigma-Aldrich, B2551) and plated

in in medium with 10% FBS. Unless stated, 6 x 10^5 BMDMs and 1.2x10^6 PerMacs per

millimeter were used in in vitro experiments. Cells were plated and left to adhere overnight prior

to challenge.

Pseudomonas aeruginosa **(PAO1) in vitro infection**

GFP expressing PAO1 was provided by the Cannon laboratory and was grown on Soy

agar plates (BD, 211043) with 100 µg/mL carbenicillin (Dot Scientific, 46000-5) overnight from

a glycerol stock. A single colony was incubated in 5 mL Mueller Hinton broth (BD, 275730)

with carbenicillin at 37 □C spinning overnight. Overnight culture was back diluted 1:5 and

grown for 2 hours until bacteria reached log-phase. Bacteria were washed with sterile phosphate-

buffered saline (PBS) and opsonized for 20 minutes in mouse serum (Fisher, PI31880). After two

more washed with PBS macrophages were infected at the corresponding MOI and spun at 250 g

for 5 minutes to synchronize infection. For inhibitor experiments, cells were primed with 10 µM

Ac-YVAD-cmk (Invivogen, inh-yvad) or 10 uM Q-VD-OPh (Millipore Sigma, 551476) for 3

hours prior to challenge. One hour after infection, wells were washed and treated with 300

µg/mL gentamicin (VWR, 0304) for 30 minutes to inactivate extracellular bacteria. The rest of

the experiment was completed in media supplemented with 50 µg/mL gentamicin and 50 µM

dimethyl fumarate (DMF; Sigma-Aldrich, 242926) or inhibitor when described.

RNA sequencing and bioinformatic analysis

Total cellular RNA was prepared from uninfected BMDMs, infected BMDMs, and

infected BMDMs treated with 50 μM DMF using Quick-RNA microprep kit (Zymo Research,

R1051) and used for the next-generation RNA-sequencing procedure at Texas A&M University

Bioinformatics Core. BaseSpace Sequence Hub (Illumina) was used for RNA seq analysis.

Briefly, Spliced Transcripts Alignment to a Reference (STAR) algorithm of RNA-seq Alignment

V2.0.0 software was used to align the results to reference genome Mus musculus/mm10

(RefSeq), and then RNA-seq Differential Expression V1.0.0 software was used to obtain raw

gene expression files and determine statistically significant changes in gene expression in Polg

macrophages relative to WT. Ingenuity Pathway Analysis software (QIAGEN) was used to

identify gene families and putative upstream regulators in the datasets. Gene Set Enrichment

Analysis (GSEA) Software (Broad Institute) was used to generate enrichment plots and examine

pathway enrichment. Heatmaps were generated using GraphPad Prism. Datasets of LPS-treated

Polg[D247A] BMDMs used in Fig. 1-2 and Supplementary Fig. 1 were previously reported[20] and are

available under Gene Expression Omnibus number GSE171960. Raw gene counts from PAO1-

infected Polg[D247A] BMDMs are supplied for review in a supplemental data spreadsheet. These

data will be deposited into the Gene Expression Omnibus upon acceptance of the paper for

publication.

Escherichia coli (E. coli) in vitro infection

E. Coli containing empty pcDNA3.1 vector for antibiotic resistance was grown on LB

agar plates (Fisher, DF0401-17) with 100 μg/mL carbenicillin (Dot Scientific, 46000-5)

overnight from a glycerol stock. A single colony was incubated in 5 mL LB broth (Fisher,

BP1427-500) with carbenicillin at 37 □C spinning overnight. Overnight culture was back diluted

1:5 and grown for 2 hours until bacteria reached log-phase. Bacteria were washed with sterile

phosphate-buffered saline (PBS) and opsonized for 20 minutes in mouse serum. After two more

washed with PBS macrophages were infected at the corresponding MOI and spun at 250 g for 5

minutes to synchronize infection. One hour after infection, wells were washed and treated with

300 μg/mL gentamicin (VWR, 0304) for 30 minutes to kill extracellular bacteria. The rest of the

experiment was completed in DMEM + 10% FBS supplemented with 50 μg/mL gentamicin.

Inflammasome challenge

BMDMs were primed for 4 hours with 20 ng/mL LPS-B5 Ultrapure (InvivoGen, tlrl-

pb5lps). When used, caspase inhibitor Ac-YVAD-cmk (InvivoGen, inh-yvad) was added an hour

after LPS, to allow for 3 hour priming, at 10 μM concentration. To trigger the non-canonical and

NLRC4 inflammasomes respectively, polyethyleneimine (Alfa Aesar, 43896) was used to

transfect 2 μg/mL LPS[69] and 100 μg/mL PA flagella[70] into the cytosol. To trigger the NLRP3

inflammasome, 5 μM ATP (Sigma, A6419) or 40 μM Nigericin (Cayman Chemical Company,

11437) was added to the media. When used, 20 µg/mL anti-IFNAR antibody (Bio X Cell,

BE0241) or anti-IgG control (Bio X Cell, BE0083) were added to cells 5 hours prior to LPS

priming and added again upon LPS transfection.

Immunoblotting

Cells and tissues were lysed in 1% NP-40 lysis buffer supplemented with protease

inhibitor and then centrifuged at 15,000 rpm for 10 min at 4 □C and the supernatant was

collected as the cellular lysate. A bicinchoninic acid assay (BCA) protein assay (Thermo Fisher

Scientific, PI23235), was used to quantify protein and 10 to 40 μg total protein were loaded into

10% SDS-polyacrylamide gels and transferred onto 0.22 μM polyvinylidene difluoride

membranes. Membranes were dried to return them to a hydrophobic state and then incubated in

primary antibodies diluted in 1x PBS containing 1% casein overnight at 4 □C rolling.

Membranes were washed and incubated with horseradish peroxidase (HRP)- conjugated

secondary antibodies for 1 hour at room temperature, shaking. Membranes were developed with

Luminata Crescendo Western HRP Substrate (Millipore, WBWR0500). Primary antibody

information can be found in Table 1.

LegendPlex multianalyte cytokine analysis

Supernatants from macrophages were subjected to centrifugation prior to the assay to

remove detached cells and debris from the sample. The assay was completed using the

LegendPlex Mouse Inflammation Panel (BioLegend, 740446) according to the included protocol.

Samples were analyzed by flow cytometry using a 5L Cytek Aurora and the BioLegend analysis

software. Samples were normalized to total protein content of each sample determined by a

bicinchoninic acid assay (BCA) protein assay and MFI from the blank control beads was

subtracted to exclude background.

Enzyme-linked immunosorbent assay (ELISA)

Supernatants were collected from macrophages plated on glass coverslips. Macrophages

were stained for IF analysis while supernatants were subjected to centrifugation to remove

detached cells and debris from the sample. Samples were stored at -80C until use in this assay

and were thawed on ice upon use. The assay was completed in 96-well half area clear, flat

bottom plates (USA Scientific, 5667-5061) and wells were incubated overnight at 4□C in

capture antibodies. Blocking was completed for one hour with PBS containing 10% FBS at room

temperature. Standards and samples were added to wells for two hours. All sample dilutions

were made in blocking buffer. Next, wells were incubated in detection antibody for one hour.

Avidin-HRP and trimethyl boron (TMB) were used for quick color development. Plates were

washed with PBS containing 0.05% Tween 20 between each step. Capture and detection

antibodies were used in the following pairs from BioLegend: Il-1β (Cat # 503514, 515801);

TNFα (Cat # 506310, 506312) and IL-6 (Cat # 504501, 504601). Samples were normalized to

corresponding nuclei counts from 20X IF images taken on an EDLIPSE Ti2 (Nikon). Nuclei

counts were determined using the automated bright spot counting function in NIS-Elements on

DAPI and an average of 3 images was used to normalize ELISA. ELISA concentrations were

determined using an epoch plate reader (BioTek) and BioTek Gen5 software. Normalized results

were graphed in GraphPad.

Quantitative polymerase chain reaction

Relative gene expression with measured by qRT-PCR in cells. Total cellular RNA was

isolated using Quick-RNA microprep kit (Zymo Research, R1051). Approximately 0.5 µg of

RNA was used to make cDNA with XLT qScript cDNA Synthesis Kit (Quantabio, 95047).

cDNA was then used in a qPCR assay using PerfeCTa Green FastMix (Quantabio, 95071). Two

to three biological replicates and two to three technical replicates were normalized to *Rpl37*

cDNA using the $2^{-\Delta\Delta CT}$ method. All primer sequences can be found in Table 2.

Immunofluorescence (IF) imaging

Cells were seeded at $2.5x10^5$ cell/well in a 24 well plate and allowed to grow overnight

on glass coverslips sterilized with ethanol. After challenge or infection, cells were washed in

PBS and fixed with 4% paraformaldehyde for 20 minutes. Next cells were permeabilized with

0.1% Triton X-100 in PBS for 5 minutes, blocked with PBS containing 5% FBS for 30 minutes,

stained with primary antibodies for 1 hour, stained with secondary antibodies for 1 hour, and

stained with 200 ng/mL 4',6-diamidino-2-phenylindole (DAPI) (ThermoFisher Scientific,

62247) for 2 minutes. When described, coverslips were stained with 2 µg/mL CellMask (Thermo

Fisher Scientific, C10046) for 10 minutes after secondary antibody staining and washed multiple

times in PBS before DAPI staining. Coverslips were mounted with ProLong Diamond Antifade

Mountant (Invitrogen, MP33025) and sealed with nail polish. Cells were imaged with a 40x or

60x oil-immersed objective with an ECLIPSE Ti2 widefield microscope (Nikon) and analyzed

with the NIS-Elements AR 5.21.02 software.

Quantification was done from multiple intensity projection (MIP) images generated from

Z-stacks. MIPs were made and analysis was done using NIS-Elements AR 5.21.02. Specifically,

cell death was determined using the automated bright spot counting function in NIS-Elements on

DAPI to count the number of nuclei. Dying cells were manually determined by identifying cells

with condensed nuclei and low CellMask staining. Percentage of dying cells were determined

and graphed in GraphPad.   To quantify nucleoids, whole cells were outlines and made into a

region of interest (ROI) and these ROIs were numbered. A thresholding mask was made on the

channels with TFAM and DNA staining and channel intensities were determined for these masks

and graphed. A new mask was made based on the overlap of TFAM and DNA. The area of this

mask and number of spots per ROI were exported and graphed in GraphPad. For GBP

quantification, cells were counted using the automated bright spot counting function in NIS-

Elements on DAPI. A thresholding mask was made on GBP5 and LAMP1 and total intensities of

each of these channels were exported. Next, a new mask was made for the overlap of these

channels and the intensity of each channel in this mask was exported and graphed. Graphs and

statistics were generated in GraphPad. Huygens Essential software (v21.4.0) was used to

deconvolve representative images from z-stacks taken with a 60X objective in oil immersion.

PI uptake

BMDMs in 96 well half area glass bottom imaging plates were seeded at 6×10^4 cells/well

in 50 µL of Phenol-Red free DMEM + 10% FBS. The next day, cells were subjected to

inflammasome challenge or infection with PAO1. Propidium iodide (PI, Alfa Aesar, J66584)

was added to media and plates were immediately imaged every ten minutes using a BioTek

Cytation V imager which maintains 37 □C and 5% CO_2. PI positive nuclei were identified and

counted using BioTek Gen5 software. Control wells treated with PI and Triton X-100 (to

permeabilize cells) to for normalization. The difference between PI positive cells compared to

the initial reading were graphed in line graphs using GraphPad Prism.

mtDNA copy number analysis

DNA was isolated from cells by boiling for 30 minutes in 50 mM NaOH lysis buffer and

then neutralizing using 1 M Tris, pH8 solution. DNA was diluted to 2 ng/uL and utilized in qRT-

PCR reaction (described above) with mtDNA specific primers. Two to three biological replicates

and two to three technical replicates were normalized to nuclear encoded *Tert* using the $2^{-\Delta\Delta CT}$

method. All primer sequences can be found in Table 3.

In vivo PAO1 infection

  6-7 month old mice were moved into BSL2 procedure space and allowed to acclimate for 1-2 days. Mice were anesthetized with an intraperitoneal injection of 100-10 mg/kg Ketamine-Xylazine cocktail. Mice were then intubated with a 24G catheter. Sterile PBS (sham), 0.5 mg/kg LPS (E. Coli serotype 0111:B4; Invivogen), or the specified dose of PAO1-GFP was pipetted into the catheter in a 25-30 µL volume. A bolus of air was delivered using a syringe to ensure the dose evenly coated the lungs. Mice were kept warm until fully recovered from anesthesia and monitored daily until experimental timepoints were reached.

Bronchial alveolar lavage fluid collection

  Mice were euthanized using intraperitoneal injection of 200 mg/kg Pentobarbital Sodium to preserve the lung environment and structure of the tracheal. Upon confirmation of death the chest cavity and throat were opened. The trachea was canulated using a 23G needle threaded with tubing and the lung was washed with 3mL cold Hanks' Balanced Salt Solution (HBSS, Thermo Fisher Scientific, 14175-129) containing 100 µM EDTA (VMR Life Science, E522) and used in subsequent flow cytometry and LegendPlex assays.

Flow Cytometry

  Bronchial alveolar lavage fluid (BAL) was spun at 350 g for 5 minutes to pellet the cells. Cells were transferred to a v bottom plate for staining. Cells were stained with viability dye then Fc receptors were blocked with anti-mouse CD16/CD32 (2.4G2) antibody. Surface proteins were stained with antibodies specified in Table 4. Cells were permeabilized with Foxp3/Transcription

Factor Staining Buffer Kit (TNB-0607-KIT, Tonbo) and stained with antibodies against

intracellular proteins. Flow cytometry analysis was done using a 5L Cytek Aurora.

Colony forming unit counts from macrophages and tissues

BMDMs or tissues were homogenized in 0.1% NP40 in sterile PBS using an immersion

homogenizer for 15 seconds per tissue. Serial dilutions were plated on soy agar containing 100

μg/mL carbenicillin. Plates were incubated at 37°C overnight and colonies were counted.

Histology

Tissues were collected in cassettes, then were washed in PBS, incubated for 24 hours in

10% formalin, and transferred to 70% ethanol. Tissue embedding, sectioning, and staining were

performed at AML Laboratories in Jacksonville, FL or at the histology core at Texas A&M

University. Images of hematoxylin and eosin (H&E) staining were collected using an Olympus

slide scanner with a 20X objective. Slides were analyzed and scored by a pathologist, Dr. Jessica

Wong, at the Jackson Laboratory in Sacramento, CA.

Statistical Analyses

To ensure rigor and reproducibility, experiments were repeated at least three times with

multiple biological replicates. Error bars displayed throughout the manuscript represent the mean

+/- s.e.m. Statistical significance between 2 groups was determined via unpaired, two-tailed

Student's t-test or Mann-Whitney test, and differences between 3 or more groups was determined

by two-way ANOVA with appropriate post-test. Survival plots were compared by the Log-rank

test. Sample sizes were chosen by started methods to ensure adequate power, and no

randomization or blinding was used for animal studies. Significance was established as * p<0.05;

** p<0.01; *** p<0.001; **** p<0.0001; or not significant (ns).

Tables:

Table 1: Western blot antibodies

Protein	Company	Catalog
IFIT3	Ganes Sen, Cleveland Clinic	N/A
STAT1	Cell Signaling Technology	9172
GAPDH	ProteinTech	60004-1-Ig
HSP60	Santa Cruz Biotechnology	Sc-1052
CASPASE-11	Cell Signaling Technology	14340
GSDMD	Abcam	ab209845
NLRP3	Cell Signaling Technology	15101
CASPASE-1	Adipogen	AG-20B0042-C100
ZBP1	Adipogen	AG-20B-0010
IL-1β	Cell Signaling Technology	12242
NLRC4	Abcam	Ab201792
GBP5	Cell Signaling Technology	67798

944 **Table 2: Primer sequences for qRT-PCR analysis**

Gene	F Primer 5'→ 3'	R Primer 5'→ 3'
Cxcl10	CCAAGTGCTGCCGTCATTTTC	GGCTCGCAGGGATGATTTCAA
Ifit1	CAAGGCAGGTTTCTGAGGAG	GACCTGGTCACCATCAGCAT
Casp11	TGTCATCTCTTTGATATATTCCTGAAG	CAAGGTTGCCCGATCAAT
Gbp2	CAGCATAGGAACCATCAACCA	TCTACCCCACTCTGGTCA
Gbp4	CAGGCTCTACATGGACATGAGG	TGCCTGCAAGATGGAACTCTCG
Gbp9	TCGTGGCTATTGTTGGATTG	GGTCTGAGACTGCACAGTGGG
Rpl37	CATCCTTTGGTAAGCGTCGCA	TGGCACTCCAGTTATACTTCCT
Stat1	CGCGCATGCAACTGGCATATAACT	ATGCTTCCGTTCCCACGTAGACTT
Zbp1	TCAAAGGGTGAAGACATGACAGAC	GTGGGTGGCTTCAGAGCTT

946 **Table 3: Primers for mtDNA copy number analysis by qPCR**

Gene	F Primer 5'→ 3'	R Primer 5'→ 3'
mt Non-	CTAGAAACCCCGAAACCAA	CCAGCTATCACCAAGCTCGT

numpt		
mt D-loop	AATCTACCATCCTCCGTGAAACC	TCAGTTTAGCTACCCCCAAGTTTAA
mt Cytb	GCTTTCCACTTCATCTTACCATTTA	TGTTGGGTTGTTTGATCCTG
nuc tert	CTAGCTCATGTGTCAAGACCCTCTT	GCCAGCACGTTTCTCTCGTT

948 **Table 4: Flow cytometry antibody panel**

Antigen	Fluorophore	Laser	Company	Catalog/ Ref #
CD45	BUV395	UV2	BD	564279
Ly6C	BV650	V11	BioLegend	128049
MHCII	BV711	V13	BioLegend	107643
CD19	BV605	V10	BioLegend	115539
CD3	APC-fire810	R8	BioLegend	100267
F4/80	VF450	V3	TONBO Biosciences	75-4801-U025
PA01	FITC	B2		
CD11c	PE-Cy5	YG5	TONBO Biosciences	55-0114-U100
TNF	PE-Cy7	YG9	Biolegend	506324
Ly6G	PE	YG1	TONBO Biosciences	50-1276-U100
Siglec-F	APC	R1	Biolegend	155508
CD11b (M1/70)	APC-Cy7	R7	TONBO Biosciences	25-0112-U100
CD326	PE/Dazzel 594	YG3	Biolegend	118235
Viability	Ghost Dye 710	R4	TONBO Biosciences	13-0871-T100

<u>References</u>

1. Gorman, G. S. *et al.* Mitochondrial diseases. *Nat. Rev. Dis. Primer* **2**, 16080 (2016).

2. Suomalainen, A. & Battersby, B. J. Mitochondrial diseases: the contribution of organelle stress

responses to pathology. *Nat. Rev. Mol. Cell Biol.* **19**, 77–92 (2018).

3. Russell, O. M., Gorman, G. S., Lightowlers, R. N. & Turnbull, D. M. Mitochondrial Diseases: Hope for

the Future. *Cell* **181**, 168–188 (2020).

4. Kruk, S. K. *et al.* Vulnerability of pediatric patients with mitochondrial disease to vaccine-preventable

diseases. *J. Allergy Clin. Immunol. Pract.* **7**, 2415-2418.e3 (2019).

5. Walker, M. A. *et al.* Predisposition to infection and SIRS in mitochondrial disorders: 8 years'

experience in an academic center. *J. Allergy Clin. Immunol. Pract.* **2**, 465–468, 468.e1 (2014).

6. Eom, S. *et al.* Cause of Death in Children With Mitochondrial Diseases. *Pediatr. Neurol.* **66**, 82–88

(2017).

7. Jestin, M. *et al.* Mitochondrial disease disrupts hepatic allostasis and lowers the threshold for

immune-mediated liver toxicity. *Mol. Metab.* **37**, 100981 (2020).

8. West, A. P., Shadel, G. S. & Ghosh, S. Mitochondria in innate immune responses. *Nat. Rev. Immunol.*

11, 389–402 (2011).

9. West, A. P. Mitochondrial dysfunction as a trigger of innate immune responses and inflammation.

Toxicology **391**, 54–63 (2017).

10.   West, A. P. & Shadel, G. S. Mitochondrial DNA in innate immune responses and inflammatory

pathology. *Nat. Rev. Immunol.* **17**, 363–375 (2017).

11.   Warren, E. B. *et al.* Inflammatory and interferon gene expression signatures in patients with

mitochondrial disease. *J. Transl. Med.* **21**, 331 (2023).

12.   Hasselmann, O. *et al.* Cerebral folate deficiency and CNS inflammatory markers in Alpers

disease. *Mol. Genet. Metab.* **99**, 58–61 (2010).

13.   Wilson, L. D., Al-Majid, S., Rakovski, C. S. & Md, C. D. Higher IL-6 and IL6:IGF Ratio in Patients

with Barth Syndrome. *J. Inflamm.* **9**, 25 (2012).

14.   Khan, W. *et al.* Neuroinflammation in the Cerebellum and Brainstem in Friedreich Ataxia: An [[18]

F]- FEMPA PET Study. *Mov. Disord.* **37**, 218–224 (2022).

15.   Nachun, D. *et al.* Peripheral blood gene expression reveals an inflammatory transcriptomic

signature in Friedreich's ataxia patients. *Hum. Mol. Genet.* **27**, 2965–2977 (2018).

16.   Jin, Z., Wei, W., Yang, M., Du, Y. & Wan, Y. Mitochondrial Complex I Activity Suppresses

Inflammation and Enhances Bone Resorption by Shifting Macrophage-Osteoclast Polarization. *Cell*

Metab. **20**, 483–498 (2014).

17.   Yu, A. K. *et al.* Mitochondrial complex I deficiency leads to inflammation and retinal ganglion cell

death in the Ndufs4 mouse. *Hum. Mol. Genet.* **24**, 2848–2860 (2015).

18.   Stokes, J. C. *et al.* Leukocytes mediate disease pathogenesis in the Ndufs4(KO) mouse model of

Leigh syndrome. *JCI Insight* **7**, e156522 (2022).

19.   Hanaford, A. & Johnson, S. C. The immune system as a driver of mitochondrial disease

pathogenesis: a review of evidence. *Orphanet J. Rare Dis.* **17**, 335 (2022).

20.   Lei, Y. *et al.* Elevated type I interferon responses potentiate metabolic dysfunction,

inflammation, and accelerated aging in mtDNA mutator mice. *Sci. Adv.* **7**, eabe7548 (2021).

21.   Saha, S., Buttari, B., Panieri, E., Profumo, E. & Saso, L. An Overview of Nrf2 Signaling Pathway

and Its Role in Inflammation. *Mol. Basel Switz.* **25**, 5474 (2020).

22.   Nezu, M., Suzuki, N. & Yamamoto, M. Targeting the KEAP1-NRF2 System to Prevent Kidney

Disease Progression. *Am. J. Nephrol.* **45**, 473–483 (2017).

23.   Trifunovic, A. *et al.* Premature ageing in mice expressing defective mitochondrial DNA

polymerase. *Nature* **429**, 417–423 (2004).

24.   Kujoth, G. C. *et al.* Mitochondrial DNA mutations, oxidative stress, and apoptosis in mammalian

aging. *Science* **309**, 481–484 (2005).

25.   Chen, M. L. *et al.* Erythroid dysplasia, megaloblastic anemia, and impaired lymphopoiesis arising

from mitochondrial dysfunction. *Blood* **114**, 4045–4053 (2009).

26.   Thomsen, K. *et al.* Animal models of chronic and recurrent Pseudomonas aeruginosa lung

infection: significance of macrolide treatment. *APMIS Acta Pathol. Microbiol. Immunol. Scand.* **130**,

458–476 (2022).

27.   Lavoie, E. G., Wangdi, T. & Kazmierczak, B. I. Innate immune responses to Pseudomonas

1006   aeruginosa infection. *Microbes Infect.* **13**, 1133–1145 (2011).

1007   28.   Lister, P. D., Wolter, D. J. & Hanson, N. D. Antibacterial-resistant Pseudomonas aeruginosa:

1008   clinical impact and complex regulation of chromosomally encoded resistance mechanisms. *Clin.*

1009   *Microbiol. Rev.* **22**, 582–610 (2009).

1010   29.   Lambert, M.-L. *et al.* Clinical outcomes of health-care-associated infections and antimicrobial

1011   resistance in patients admitted to European intensive-care units: a cohort study. *Lancet Infect. Dis.*

1012   **11**, 30–38 (2011).

1013   30.   Hayashi, G. *et al.* Dimethyl fumarate mediates Nrf2-dependent mitochondrial biogenesis in mice

1014   and humans. *Hum. Mol. Genet.* **26**, 2864–2873 (2017).

1015   31.   Robledinos-Antón, N., Fernández-Ginés, R., Manda, G. & Cuadrado, A. Activators and Inhibitors

1016   of NRF2: A Review of Their Potential for Clinical Development. *Oxid. Med. Cell. Longev.* **2019**,

1017   9372182 (2019).

1018   32.   Hoyle, C., Green, J. P., Allan, S. M., Brough, D. & Lemarchand, E. Itaconate and fumarate

1019   derivatives inhibit priming and activation of the canonical NLRP3 inflammasome in macrophages.

1020   *Immunology* **165**, 460–480 (2022).

1021   33.   Sutterwala, F. S. *et al.* Immune recognition of Pseudomonas aeruginosa mediated by the

1022   IPAF/NLRC4 inflammasome. *J. Exp. Med.* **204**, 3235–3245 (2007).

1023   34.   Balakrishnan, A., Karki, R., Berwin, B., Yamamoto, M. & Kanneganti, T.-D. Guanylate binding

1024   proteins facilitate caspase-11-dependent pyroptosis in response to type 3 secretion system-negative

1025   Pseudomonas aeruginosa. *Cell Death Discov.* **4**, 66 (2018).

1026   35.   Arlehamn, C. S. L. & Evans, T. J. Pseudomonas aeruginosa pilin activates the inflammasome. *Cell.*

1027   *Microbiol.* **13**, 388–401 (2011).

1028   36.   Billingham, L. K. *et al.* Mitochondrial electron transport chain is necessary for NLRP3

1029   inflammasome activation. *Nat. Immunol.* **23**, 692–704 (2022).

1030   37.   Kumari, P., Russo, A. J., Wright, S. S., Muthupalani, S. & Rathinam, V. A. Hierarchical cell-type-

1031   specific functions of caspase-11 in LPS shock and antibacterial host defense. *Cell Rep.* **35**, 109012

1032   (2021).

1033   38.   Benjamin, J. T. *et al.* IL-1α and Not IL-1β Drives IL-1R1–Dependent Neonatal Murine Sepsis

1034   Lethality. *J. Immunol.* **201**, 2873–2878 (2018).

1035   39.   Pyrillou, K., Burzynski, L. C. & Clarke, M. C. H. Alternative Pathways of IL-1 Activation, and Its

1036   Role in Health and Disease. *Front. Immunol.* **11**, 613170 (2020).

1037   40.   Rathinam, V. A. K. *et al.* TRIF Licenses Caspase-11-Dependent NLRP3 Inflammasome Activation

1038   by Gram-Negative Bacteria. *Cell* **150**, 606–619 (2012).

1039   41.   Kim, B.-H. *et al.* A family of IFN-γ-inducible 65-kD GTPases protects against bacterial infection.

1040   *Science* **332**, 717–721 (2011).

1041   42.   Kim, B.-H. *et al.* Interferon-induced guanylate-binding proteins in inflammasome activation and

1042   host defense. *Nat. Immunol.* **17**, 481–489 (2016).

1043   43.   Shenoy, A. R. *et al.* GBP5 promotes NLRP3 inflammasome assembly and immunity in mammals.

1044   *Science* **336**, 481–485 (2012).

1045   44.   Coers, J. Sweet host revenge: Galectins and GBPs join forces at broken membranes. *Cell.*

1046   *Microbiol.* **19**, e12793 (2017).

1047   45.   Lam, C.-W. *et al.* Novel POLG mutation in a patient with sensory ataxia, neuropathy,

1048   ophthalmoparesis and stroke. *Clin. Chim. Acta Int. J. Clin. Chem.* **448**, 211–214 (2015).

1049   46.   Amiot, A. *et al.* Frequency of mitochondrial defects in patients with chronic intestinal pseudo-

1050   obstruction. *Gastroenterology* **137**, 101–109 (2009).

47.   Kinoshita, M. & Sawa, T. Serological immune response against major secretory proteins

expressed by Pseudomonas aeruginosa upon intratracheal infection in mice. *Respir. Investig.* **60**, 318–

321 (2022).

48.   Edmonds, J. L. *et al.* The Otolaryngological Manifestations of Mitochondrial Disease and the Risk

of Neurodegeneration With Infection. *Arch. Otolaryngol. Neck Surg.* **128**, 355 (2002).

49.   Lee, W. S. & Sokol, R. J. Mitochondrial Hepatopathies: Advances in Genetics, Therapeutic

Approaches, and Outcomes. *J. Pediatr.* **163**, 942–948 (2013).

50.   The Mitochondrial Medicine Society *et al.* A modern approach to the treatment of mitochondrial

disease. *Curr. Treat. Options Neurol.* **11**, 414–430 (2009).

51.   Rahman, S. & Copeland, W. C. POLG-related disorders and their neurological manifestations.

Nat. Rev. Neurol. **15**, 40–52 (2019).

52.   Gaudó, P. *et al.* Infectious stress triggers a POLG-related mitochondrial disease. *neurogenetics*

21, 19–27 (2020).

53.   Al-Zubeidi, D. *et al.* Fatal Human Herpesvirus 6–Associated Encephalitis in Two Boys With

Underlying POLG Mitochondrial Disorders. *Pediatr. Neurol.* **51**, 448–452 (2014).

54.   Torres-Odio, S. *et al.* Loss of Mitochondrial Protease CLPP Activates Type I IFN Responses

through the Mitochondrial DNA-cGAS-STING Signaling Axis. *J. Immunol. Baltim. Md 1950* **206**, 1890–

1900 (2021).

55.   Kobayashi, E. H. *et al.* Nrf2 suppresses macrophage inflammatory response by blocking

proinflammatory cytokine transcription. *Nat. Commun.* **7**, 11624 (2016).

56.   Seidel, P. *et al.* Dimethylfumarate inhibits NF-κB function at multiple levels to limit airway

smooth muscle cell cytokine secretion. *Am. J. Physiol.-Lung Cell. Mol. Physiol.* **297**, L326–L339 (2009).

57.   Hu, H. *et al.* Dimethyl fumarate covalently modifies Cys673 of NLRP3 to exert anti-inflammatory

effects. *iScience* **27**, 109544 (2024).

58.   Humphries, F. *et al.* Succination inactivates gasdermin D and blocks pyroptosis. *Science* **369**,

1633–1637 (2020).

59.   Miao, R. *et al.* Gasdermin D permeabilization of mitochondrial inner and outer membranes

accelerates and enhances pyroptosis. *Immunity* **56**, 2523-2541.e8 (2023).

60.   Miao, N. *et al.* Oxidized mitochondrial DNA induces gasdermin D oligomerization in systemic

lupus erythematosus. *Nat. Commun.* **14**, 872 (2023).

61.   Weindel, C. G. *et al.* Mitochondrial ROS promotes susceptibility to infection via gasdermin D-

mediated necroptosis. *Cell* **185**, 3214-3231.e23 (2022).

62.   Moretti, J. *et al.* Caspase-11 interaction with NLRP3 potentiates the noncanonical activation of

the NLRP3 inflammasome. *Nat. Immunol.* **23**, 705–717 (2022).

63.   Yazdi, A. S. & Drexler, S. K. Regulation of interleukin 1α secretion by inflammasomes: Table 1.

Ann. Rheum. Dis. **72**, ii96–ii99 (2013).

64.   Kang, Y. *et al.* Ancestral allele of DNA polymerase gamma modifies antiviral tolerance. *Nature*

628, 844–853 (2024).

65.   Liu, B. C. *et al.* Constitutive Interferon Maintains GBP Expression Required for Release of

Bacterial Components Upstream of Pyroptosis and Anti-DNA Responses. *Cell Rep.* **24**, 155-168.e5

(2018).

66.   Dickinson, M. S. *et al.* LPS-aggregating proteins GBP1 and GBP2 are each sufficient to enhance

caspase-4 activation both in cellulo and in vitro. *Proc. Natl. Acad. Sci.* **120**, e2216028120 (2023).

67.   Pilla, D. M. *et al.* Guanylate binding proteins promote caspase-11–dependent pyroptosis in

response to cytoplasmic LPS. *Proc. Natl. Acad. Sci.* **111**, 6046–6051 (2014).

68.   Meunier, E. *et al.* Caspase-11 activation requires lysis of pathogen-containing vacuoles by IFN-

induced GTPases. *Nature* **509**, 366–370 (2014).

69.   Bezbradica, J. S., Coll, R. C. & Boucher, D. Activation of the Non-canonical Inflammasome in

1099   Mouse and Human Cells. in *The Inflammasome* (ed. Abdul-Sater, A. A.) vol. 2459 51–63 (Springer US,

1100   New York, NY, 2022).

1101   70.   Kamajaya, L. J. & Boucher, D. Gasdermin D Cleavage Assay Following Inflammasome Activation.

1102   in *The Inflammasome* (ed. Abdul-Sater, A. A.) vol. 2459 39–49 (Springer US, New York, NY, 2022).

1103

1104

Fig. 1

Fig. 2

a Ingenuity Pathway Analysis of RNA-Seq
Polg^{D257A}/WT BMDM, 6 hpc

Significantly changed pathways
determined by IPA

Z Score

Death Receptor Signaling
Inflammasome Pathway
Immunogenic Cell Death Signaling
IL-1 Signaling

LPS PA PA+DMF

b

Inflammasome Signaling
Pycard, Nod2, Nlrc4, Nfkb2, Myd88, Il1b, Il18, Casp8, Casp4, Casp1, Aim2
LPS PA PA+DMF

Immunogenic Cell Death
Tlr3, Pycard, P2ry2, Nlrp3, Il1b, Ifnar1, Dapk3, Cxcl10, Cflar, Casp8, Casp4, Casp3, Atg16l1, Anxa1
LPS PA PA+DMF

Death Receptor Signaling
Traf2, Tnfsf15, Tnfsf10, Tiparp, Tbk1, Tank, Rel, Parp2, Nfkbib, Nfkbia, Nfkb1, Map2k4, Ikbkb, Fas, Daxx, Casp8, Casp4, Casp3, Casp2, Apaf1, Actb
LPS PA PA+DMF

Log2 F.C.

c PerMac, PAO1 MOI = 5, 2 hpc

WT
Merge | DAPI
PAO1-GFP | CellMask
20 μm

Polg^{D257A}
Merge | DAPI
PAO1-GFP | CellMask
20 μm

WT +DMF
Merge | DAPI
PAO1-GFP | CellMask
20 μm

Polg^{D257A} +DMF
Merge | DAPI
PAO1-GFP | CellMask
20 μm

d BMDM, PAO1 MOI=5, 2 hpc

% of CellMask$^+$ Cells with Condensed Nuclei/image

****   ****

WT
Polg^{D257A}
WT+DMF
Polg^{D257A}+DMF

e BMDM, PAO1 MOI=5

Number of PI+ Cells (10^4)

****

WT
Polg^{D257A}
WT+DMF
Polg^{D257A}+DMF

0 30 60 90 120 mpc

f IL-1β

MFI (10^4)

**   ***   ****
***   ****   ****

WT
WT+DMF
Polg^{D257A}
Polg^{D257A}+DMF

8   24 hpc

Fig. 3

Fig. 4

a **RNA-Seq BMDM, 6 hpc**

WT PA/WT Un ■ Polg^{D257A} PA/Polg^{D257A} Un ■

Gbp2 *Gbp2b* *Gbp3* *Gbp4* *Gbp5* *Gbp6* *Gbp7* *Gbp8* *Gbp9* *Gbp10* *Gbp11* *Caspase11*

Relative Induction (fold)

b **BMDM, Unchallenged**

WT — Merge — 10 μm — GBP5

Polg^{D257A} — Merge — 10 μm — GBP5

d **BMDM, PAO1 MOI=5, 2 hpc**

WT — Merge — 10 μm — GBP5 — PAO1-LPS — GBP5/PAO1-LPS

Polg^{D257A} — Merge — 10 μm — GBP5 — PAO1-LPS — GBP5/PAO1-LPS

c **GBP5 intensity**

WT ■ Polg^{D257A} ■

Sum Intensity/Area (10^4)

Uninfected

e **GBP5 intensity**

Sum Intensity/Area (10^5)

2 hpc

f **PAO1-LPS co-localized with GBP5**

GFP localized with GBP5 (sum intensity, 10^4)

2 hpc

Fig. 5

a Generation of Polg^R292C mouse

R292C

| NTD | Exonuclease | Spacer | Palm/Fingers |

CRISPR guide to introduce **R292C** mutation in exon 4 of the exonuclease domain of Polg, mimicking human **R309C** pathogenic mutation.

CRISPR guide was cloned into plasmid pX459 expressing Cas9. Plasmid and an oligonucleotide donor with the R292C mutation and a silent F290F mutation (to limit guide re-cutting) were electroporated into ES cells.

Embryonic Stem (ES) Cells

Puro resistant ES cells were screened for proper targeting and exclusion of pX459.

Euploid clones injected into C57BL/6J-albino blastocysts to produce chimeras

Chimeras were bred into C57BL/6NJ and a single germline clone was selected for expansion.

b Polg^R292C/+ sequence validation

Polg^+   F Q R S
         TTCAG TGCAGC

Polg^R292C F Q C S
           TTCAG CGCAGC
→ R292C mutation
→ F290F Silent mutation

c BMDM, unchallenged

Non-numpt

WT / Polg^R292C

mtDNA Abundance (%)

D-loop / CytB

d BMDM, LPS, 24 hpc

WT / Polg^R292C

Merge / TFAM / DNA
5 um
PDH/TFAM/DNA

e BMDM, LPS, 24 hpc

WT / Polg^R292C

Number of Nucleoids — Nucleoids/Cell (#)
Nucleoid DNA Intensity — DNA Mean Intensity/Cell (10^3)
Nucleoid Area — Nucleoid Area/Cell

f Polg^R292C BMDM, PAO1 MOI=5, 8 hpc

Stat1 / Ifit1 / Gbp2
Zbp1 / Gbp4 / Gbp9

Relative Expression (fold)

g Polg^R292C BMDM, PAO1 MOI=5, 2 hpc

WT
Merge / DAPI
PAO1-LPS / CellMask
10 μm

Polg^R292C
Merge / DAPI
PAO1-LPS / CellMask
10 μm

h % of CellMask- Cells with Condensed Nuclei/Image

WT / Polg^R292C

i

IL-1β — WT / Polg^R292C
ng/mL

TNFα
ng/mL

IL-6
ng/mL

2 8 24 hpc

Fig. 6

a BAL cytokines, LegendPlex, 1x10⁴ CFU PAO1, 6 hpi

IL-1β ** | TNFα * | IFNβ ** | IL-23 ** | IL-10 * | IL-27 *

● WT
■ Polg^D257A

IL-17α ** | IL-12p70 ** | GM-CSF * | IFNγ 0.077 | IL-1α 0.177 | MCP-1 0.220 | IL-6 0.286

b Lung, 1x10⁴ CFU PAO1, 48 hpi

WT   Polg^D257A

CASPASE1
p20
CASPASE 11
NLRC4
NLRP3
GSDMD
GBP5
IFIT3
STAT1
GAPDH

c Lung, 1x10⁵ CFU PAO1, 20 hpi

WT   Polg^D257A

d Histology analysis, 1x10⁵ CFU PAO1, 20 hpi

Genotype	ID	Total Inflammation	Neutrophilic Inflammation	Mononuclear Inflammation
WT	1	minimal	1	1
	2	minimal	0	1
	3	mild	2	1
	4	moderate	3	0
	5	mild	2	0
PolG	1	mild/moderate	2	0
	2	moderate	3	0
	3	moderate	3	0
	4	moderate	3	2
	5	severe	4	1

e 1x10⁵ CFU PAO1, 20 hpi

Total Inflammation * | Neutrophilic Inflammation *

● WT
■ Polg^D257A

f 1x10⁵ CFU PAO1, 20 hpi

WT   Polg^D257A

CASPASE-11
IL-1β
HSP60

g 1x10⁵ CFU PAO1

Percent Survival vs dpi

h

Mitochondrial Disease (Polg mutation)
↑ IFN-I
↓ Nrf2
DMF

Supplementary Fig. 1

a BMDM — Unchallenged / LPS priming + Nigericin

WT: Merge, ASC, CellMask, DAPI — 10 µm

b ASC puncta/Cell (CellMask high/low); CellMask Intensity/Area (ASC −/+) ****

c PAO1 in Supernatant; PAO1 in Macrophages — CFU (10⁴), 2 / 4 hpc *** **

d WT / Polg^D257A — Beads/Cell n.s.; % Cells with beads/Image n.s.

e BMDM + LPS coated microspheres, 30 min — WT / Polg^D257A: Merge, DAPI, Sphere, LAMP1 — 20 µm

f BMDM, DH5α MOI=5 — IL-1β, TNFα, IL-6 (ng/mL) vs WT / Polg^D257A, 2 4 8 24 hpc ****

g % of CellMask⁻ Cells with Condensed Nuclei/Image — WT / Polg^D257A ****

h BMDM, DH5α MOI=5, 2 hpc — WT / Polg^D257A: Merge, DH5α-LPS, CellMask, DAPI — 10 µm

Supplementary Fig. 4

a BMDM, unchallenged

Stat1 *Ifit1* *Zbp1* *Gbp2* *Gbp4* *Gbp9*

Relative Expression (fold)

WT / Polg^R292C

b BMDM, LPS, 8 hpc

Stat1 *Ifit1* *Zbp1* *Gbp2* *Gbp4* *Gbp9* *Casp11*

Relative Expression (fold)

c Lung, unchallenged

WT — H&E stain / Eosin intensity overlay

Polg^D257A — H&E stain / Eosin intensity overlay

Eosin+ / Eosin-

d Lungs, unchallenged

Genotype	Mouse ID	Total Inflammation	Neutrophilic Inflammation	Mononuclear Inflammation
WT	1	minimal	0	1
	2	minimal	0	1
	3	none	0	0
Polg	1	none	0	0
	2	none	0	0
	3	none	0	0

e WT / Polg^D257A

% Eosin+ Cells — ns

Eosin+ Cells/Area — ns

f Lungs, unchallenged

WT / Polg^D257A / WT IFNAR / Polg^D257A IFNAR

STAT1
IFIT3
HSP60

g BAL Cytokines, 0.5 mg/kg LPS

IL-1β / TNFα / IL-6

pg/mL

WT / Polg^D257A

4 6 12 hpc

Supplementary Fig. 5

a BAL Cytokines,1x10⁴ CFU PAO1

Weight

→ WT
→ Polg^D257A

% of D0 Weight

-10, -8, -6, -4, -2, 0

0 1 2 3 4 dpi

b

Uninfected   6 hpi   48 hpi

CD45+   % of Live Cells   *, **, **

CD45-   % of Live Cells   *, **, **

c

FSC-A

WT

CD45- 67.5 | CD45+ 31.7
CD45- 37.1 | CD45+ 62.7
CD45- 27.8 | CD45+ 71.6

Polg^D257A

CD45- 70.9 | CD45+ 28.2
CD45- 54.4 | CD45+ 45.4
CD45- 35.0 | CD45+ 64.0

CD45

d CD11c+   % of CD45+ Cells

Uninfected   6 hpi   48 hpi

ns, ns, ns

e SiglecF+   % of CD45+ Cells   *, *, **

f MHCII+   % of CD45+ Cells   *, ns, ns

g F4/80+   % of CD45+ Cells   *, ns, **

h CD11b+   % of CD45+ Cells   ns, ns, ns

i CD3+   % of CD45+ Cells

Uninfected   6 hpi   48 hpi

ns, *, ns

j CD19+   % of CD45+ Cells   ns, ns, ns

k Ly6G+   % of CD45+ Cells   ns, ns, ns

l Ly6C+   % of CD45+ Cells   ns, ns, ns

m CD45+, TNF+   % of CD45+ Cells   *, *, ns

■ WT ■ Polg^D257A

a BAL Cytokines, LegendPlex 1x10⁴ CFU PAO1, 48 hpi

b Lung, 1x10⁴ CFU PAO1, 7 dpi + 24 hpi

c

d BAL Cytokines, LegendPlex 1x10⁴ CFU PAO1, 7 dpi + 24 hpi

e CFU, 1x10⁴ CFU PAO1, 7 dpi + 24 hpi

f CFU, 1x10⁵ CFU PAO1, 20 hpi