1

Active vision in freely moving marmosets using head-mounted eye tracking

Vikram Pal Singh[1*], Jingwen Li[1], Jude Mitchell[2], Cory Miller[1,3]

[1] Cortical Systems & Behavior Lab, University of California San Diego
[2] Department of Brain and Cognitive Science, University of Rochester
[3] Neurosciences Graduate Program, University of California San Diego

*Corresponding Author: vpsingh@ucsd.edu

30
31 **Abstract:**
32
Our understanding of how vision functions as primates actively navigate the real-world is
remarkably sparse. As most data have been limited to chaired and typically head-restrained
animals, the synergistic interactions of different motor actions/plans inherent to active sensing –
e.g. eyes, head, posture, movement, etc. - on visual perception are largely unknown. To address
this considerable gap in knowledge, we developed an innovative wireless head-mounted eye
tracking system called **CEREBRO** for small mammals, such as marmoset monkeys. Our system
performs **C**hair-free **E**ye-**Re**cording using **B**ackpack mounted mic**RO**controllers. Because eye
illumination and environment lighting change continuously in natural contexts, we developed a
segmentation artificial neural network to perform robust pupil tracking in these conditions.
Leveraging this innovative system to investigate active vision, we demonstrate that although
freely-moving marmosets exhibit frequent compensatory eye movements equivalent to other
primates, including humans, the predictability of the visual system is enhanced when animals are
freely-moving relative to when they are head-fixed. Moreover, despite increases in eye/head-
motion during locomotion, gaze stabilization actually improved over periods when the monkeys
were stationary. Rather than impair vision, the dynamics of gaze stabilization in freely-moving
primates has been optimized over evolution to enable active sensing during natural exploration.

Main:

Primate vision has been the subject of intense study for many decades and is arguably the most well understood neural system in the simian brain (Felleman and Van Essen 1991; A. J. Parker and Newsome 1998). And yet, our understanding of primate vision is incomplete. Like all sensory systems, the primate vision evolved to overcome the challenges of actively moving, exploring and engaging with the objects, individuals and the environment from different perspectives (P. R. L. Parker et al. 2022; Miller et al. 2022; Ngo et al. 2022). While the processes of visual encoding have been extensively studied with head-restrained subjects observing stimuli presented on a screen, details of how vision functions as primates actively move through and explore the real-world are remarkably limited. The tacit expectation being that the visual processes discovered in head-fixed preparations reflect the core of sensory encoding that will generalize to freely-moving animals once mechanisms to stabilize gaze are considered. Similar to all vertebrates, the head and eyes coordinate their respective movements in primates to enable a stable percept of visual inputs (Guitton 1992; Martinez-Trujillo, Wang, and Crawford 2003; Angelaki and Hess 2005; Angelaki and Cullen 2008). Because data on this complement of mechanisms has been limited to chaired animals unable to locomote, the synergistic effects of different motor actions – e.g. eyes, head, posture, movement, etc. - on primate visual perception and cognition during active exploration of the world are almost entirely unknown. The principal bottleneck being technical. Previous studies have achieved limited precision for projecting the gaze of free-moving primates into visual scenes (Shepherd and Platt 2006; Milton, Shahidi, and Dragoi 2020; Mao et al. 2021) and as a result have not detailed the stability nor the eye and head dynamics that contribute to gaze during free-motion. Here we introduce a method that precisely quantifies eye-movements and can accurately project the gaze of a primate into scenes as individuals freely explore an environment.

Recent work with mice demonstrate that eye-tracking systems can be miniaturized and mounted to the head in order to provide insight into vision during natural visual behavior (Michaiel, Abe, and Niell 2020; Meyer et al. 2018; Wallace et al. 2013; Meyer, O'Keefe, and Poort 2020) , but these systems are not well suited for comparable studies in primates for at least two critical reasons. First, systems in mice rely on a tether which restricts the 3D mobility of primates. Second, they lack the precision and temporal resolution needed to accurately characterize high-resolution vision of primates. To address this challenge, we developed an innovative head-mounted eye tracking system to enable the study of active, natural visual behaviors, and related neural processes, in freely-moving marmosets. Our system - **CEREBRO** - allows for **C**hair-free **E**ye-**Re**cording using **B**ackpack mounted mic**RO**controllers at a speed and resolution needed to accurately quantify the visual behavior and underlying neural mechanisms of natural, active vision in primates. Using CEREBRO we confirmed that freely-moving marmosets exhibit frequent compensatory eye movements that enable them to stabilize gaze when viewing real world scenes consistent with previous studies in body restrained animals. By using this innovative system, however, we discovered that gaze stabilization and predictability was enhanced when the monkeys were moving naturally despite increases in eye/head-motion during locomotion. This suggests that previously unreported synergistic mechanisms for gaze stabilization are not only

109 integral to primate active vision in the real-world but can be enhanced for greater compensation
110 as animals move naturally through their environment.
111
112
113
114 **Results:**
115
116 *CEREBRO* **is a head mounted eye tracking system for freely moving marmosets:**
117
The small body size of marmosets (~300-400g) necessitated certain design considerations when
developing CEREBRO. The first of these decisions is related to the weight and wearability of the
system itself. Based on previous experience, the estimated 60g weight of the complete system
would not be feasible to be entirely situated on animal's head without affecting its visual behavior.
To resolve this issue, we separated the system into two separate, but integrated hardware
submodules: the **Head-piece** and the **Backpack**. The *Head-piece* includes the camera assembly
and scaffold fitted to animals' head *(fig 1a)*; and the *Backpack* includes the backend electronics
for camera synchronizing, image acquisition and local data storage on the custom designed
printed circuit boards (PCBs) *(fig1b)*. Both the PCB and the head-piece IR LED are powered using
a 600 mAh, Lithium-Polymer (Li-Po) battery that is housed inside the backpack.

The second consideration for this system was the cameras used here, as we sought to accurately
quantify the high-resolution vision of primates. Commercially available parallel communication
camera modules such as OV7725, OV2640, etc. have low frame rates (< 30hz) which are not
ideal for eye tracking in primates. Better camera modules with faster frame rate (60 or 90hz) and
HD resolution (1080p) are available but they do not follow parallel communication protocol. Rather
these cameras use a camera serial interface called MIPI CSI-2 (Mobile Industry Processor
Interface Camera Serial Interface). We also wanted to use a microprocessor as the core of the
system to get better temporal resolution of the incoming data. Since STM32H750 can only support
a parallel camera interface, we used a STMIPID02 MIPI CSI-2 deserializer to deserialize CSI
communication from a MIPI CSI-2 camera sensor to DCMI in order to use cameras with sufficient
recording frame rate and resolution needed here (Supplementary fig 3b). For our current system,
we used OV4689 camera modules with two MIPI lanes and customized Flex cable length of 15
cm (SincereFirst, Guangzhou, China).

141
Head-Piece Module. The head-piece weighs ~20g and comprises five different components (*fig*
1a): (1) the scaffold: a curved metal tube that serves as an anchor for all the pieces, (2) eye-cam:
an HD MIPI camera (90fps) with Visible light filter and a Macro lens for looking at the eye, (3)
world cam: an HD MIPI camera (60fps) for capturing the world scene in front of the camera, (4)
an IR LED to illuminate the eye, and (5) a strategically placed Hot mirror to image the eye. The
mechanical parts for the head-piece are made up of a Titanium alloy (Ti-6Al-4V) due to strength
and weight considerations, and are held to the scaffold using M2 and M4 screw sets (see
Supplementary Information 1 for detailed assembly description). Each piece is custom designed
using FreeCAD and 3D printed using Direct-LASER-Sintering (DLS). The eye camera, IR LED
and the hot mirror are positioned to achieve a direct image of the eye with optimal illumination.

To achieve reliable eye-tracking in a fully unrestrained animal, the eye camera's position requires
to be fixed with respect to the animal's eye. This is accomplished by attaching 2 vertical headposts
(5 mm diameter, 1 cm tall cylinders with a flat cut on one side) on the animals' head (*fig 1c*). The
two headposts restrict any rotation or translational movement of the headpiece and allows the
headpiece to be placed at the same position for every recording session.

Backpack Module. The backpack module weighs 40g and comprises two Customized PCBs, as
well as the casing and harness worn by the animals (*fig 1b*). Each camera has its own dedicated
PCB (*fig 1d*). Both cameras (eye cam and world cam) have long flex cables (15 cm) and are
operated by the PCB.

Customized Printed Circuit Boards. The PCB used here is an embedded system that runs on
an STM32H750 microprocessor (*fig 1d*). STM32H750 is a 32-bit Arm Cortex-M7 core processor
running up to 480 MHz and is better suited for interfacing with camera sensors since it embeds a
digital camera interface (DCMI). Each PCB can be programmed and debugged using the Serial
Wire Debug (SWD) mode. The PCBs store the raw camera data on a locally mounted, readily
available UHS-1 (Ultra High Speed) class, SD card. The binary files are decoded offline to get an
AVI format video of the stream and a CSV file with timestamp and IMU data. Two PCBs are
stacked on top of each other and are enclosed in a custom designed 3D printed casing with a
custom designed back harness (described below). To measure camera synchronized movements
(acceleration and orientation) of head, the boards interface with an inertial measurement unit
(IMU). The PCB boards also feature a 0.96" SPI TFT display which allows the users to preview
the captured camera image and ensure the placement of the head-piece at the start of the
session.

Eye-Tracking System Wearability. When fully configured on a marmoset (fig 1e) the CEREBRO
allows a full range of motions characteristic of the marmoset's natural behavior, while monitoring
its position in the environment, eye position, and view of the scene (*fig 1f*). Overall, CEREBRO
weighs ~60 g (headpiece module: 20g + backpack module: 40g), which is comparable to the
weight of two infant marmosets that an adult would normally carry on its back. To test the influence
of this added weight on mobility, we compared marmosets' behavior both with and without
CEREBRO in multiple sessions (30 min) of freely-moving exploration in a large open arena, (200
cm x 100 cm x 240 cm). The animals carried an IR reflective 4mm bead on their backpacks in the
arena and the body movement was tracked using motion tracking system *"OptiTrack"* with a
resolution of less than 1mm at 120 Hz. We defined locomotion as any movement of the animal
greater than 5 cm/sec (i.e. moving more than 20-30% the typical marmoset body length in under
one second). Using this threshold criterion, we quantified the percentage of time animals spend
moving through the arena. Over 12 sessions spread across months, we observed no significant
difference (*fig 1g*) in the time animals spent locomoting vs stationary and scanning the
environment (p=0.83). Even though the animals moved for the same amount of time, there is still
a possibility that the weight can impede the speed of the animal. Upon investigation, we did not
observe any significant difference in either the distance covered by the animals (p=0.569) or the
average speed of the animal with and without CEREBRO (p=0.5692). This leads us to conclude

that our system does not significantly impede the animals' movement and can be comfortably
used with a freely behaving animal.

Fast, efficient and accurate pupil detection:

In the traditional experimental paradigms that have dominated primate neuroscience, animals are
head-fixed and the light source can be controlled. The consistency of the light allows for highly
accurate eye tracking to be achieved by thresholding the dark iris observed under infra-red light
from a brightly illuminated image and the pupil detected either using its centroid or center of the
ellipse fit. A critical challenge for eye-tracking in freely moving animals is the lack of control over
the illumination. As an individual moves through any real-world environment external lighting
conditions and shadows are constantly changing. This point is illustrated by our application of
thresholding for pupil detection of a freely-moving marmoset (*Fig 2a*), which can lead to labeling
of shadows at the edge of the eye rather than the pupil. Previous studies that examined gaze in
free-moving primates relied on conventional pupil thresholding (Shepherd and Platt 2006; Milton,
Shahidi, and Dragoi 2020; Mao et al. 2021), which limited the precision that could be obtained.
To achieve accurate primate eye-tracking in natural, freely-moving conditions, an alternative
method for pupil detection was needed.

Artificial Neural Networks (ANNs) offer an alternative approach to pupil detection under real-world
conditions because they rely on features of an image and not necessarily the grayscale values,
and therefore are resistant to brightness changes and/or shadows/occlusions. Such approaches
have been successfully applied using commercial software, such as DeepLabCut (DLC), to track
the pupil of freely-moving mice (Michaiel, Abe, and Niell 2020). To further optimize pupil tracking
in freely-moving marmosets, here we developed a custom semantic segmentation ANN called
UNet (*fig 2b*) (Ronneberger, Fischer, and Brox 2015) that yielded superior performance in these
conditions. Whereas classical convolutional networks are classifiers which output a single class
label with a confidence value, the UNet architecture achieves this by assigning a class label to
each pixel of the image. The core logic for the UNet implemented here was to supplement a
classical convolutional network (contracting path) with successive upscaled layers such that the
final output image is of the same dimension with object of interest segmented out as white (>0)
and background set to black (0). This workflow allows to detect pupil features robust from lighting
and noise (*fig 2b*). With a fully trained network, we reliably detected pupils across various sessions
and different animals. The network is easily trainable with fewer epochs and a training dataset of
only 250-500 images (*fig 2c*). The data preparation and training of the ANN is explained in detail
in *Supplementary information 4*, which also includes a user-friendly GUI (*Supplementary fig 4b*).

Accurately calculating the gaze point from the world camera necessitates that the pupil position
from the eye camera be calibrated to the real-world position. To this end, we developed the
following procedure. The animal is chaired and head-fixed in front of a computer screen placed
~35 cm directly in front of the subject. Because marmosets naturally look at faces (Mitchell,
Reynolds, and Miller 2014), 1-12 small marmoset faces are presented as calibration targets at
multiple locations on the display monitor (*fig 2d*). A custom designed graphical user interface was
used to adjust for scaling and offset in horizontal and vertical axes by a human operator offline.

This allowed us to reliably map pupil eye position to screen coordinates (*fig 2e*), which when head-
free, generalized to world coordinates in front of the marmoset. The GUI provides functionality to
project the estimated eye position, based on the manually defined gain and center parameters,
onto the video of the scene camera in order to evaluate the quality of the eye calibration and
iteratively refine the scaling and offset parameters to better fit the eye position data. The
transformed raw eye positional data can then be exported in a format convenient for further
analysis. To compare the system against other pupil-based eye trackers we computed the root-
mean-square (RMS) stability of eye position during stable fixation epochs when animals were
head-fixed. We find that across 7 recording sessions the RMS stability was 0.05 (+/- 0.0012 std).
These estimates provide a lower bound on the system precision that rivals head-stabilized pupil-
based eye trackers.

247
248 **Electrophysiology in combination with CEREBRO:**
249
CEREBRO was designed explicitly to create a tool to investigate the neurobiology of natural,
active vision in freely-moving monkeys. As such, integral to its design was the capacity to
simultaneously record neural activity, eye behavior, and the visual scene of the animal, as each
are integral to this broader ambition. We tested the validity and sufficiency of the system to this
end by performing experiments with CEREBRO while activity of single neurons was recorded with
chronically implanted multi-electrode arrays (N-form, Modular Bionics) in the visual cortex (V1/V2)
of the marmosets. These experiments sought to (1) estimate visual tuning properties of neurons
in primate visual cortex, i.e. receptive field mapping, as well as orientation and spatial frequency
tuning (Emerson et al. 1987; DeAngelis, Ohzawa, and Freeman 1993; Ringach, Sapiro, and
Shapley 1997; Livingstone, Pack, and Born 2001), using CEREBRO in more traditional head-
fixed paradigms , so as to demonstrate the accuracy of our eye-tracking system by replicating
these classic effects, and (2) obtain eye, head and body behavior, visual scene, and activity of
single neurons simultaneously in freely-moving paradigm to demonstrate the capacity of
CEREBRO in investigating primate active vision.

264
To recapitulate the tuning properties of V1 neurons, subjects were head-fixed while wearing
CEREBRO and presented with the following stimuli: flashing dots for receptive field mapping, and
drifting gratings with different orientations and spatial frequencies for tuning properties (*fig. 3a*;
see online methods). Critically, marmosets were allowed to free-view the video screen during
stimulus presentation and offline corrections for eye position enable accurate reconstruction of
visual properties following a recently developed free-viewing approach (Yates et al. 2023). A key
difference here from the previous study with head-fixed marmosets is that the visual input is
obtained by the world camera's view along with the estimated eye position from CEREBRO
instead of what was known to be displayed on the screen, thus validating that these methods
could generalize to real-world stimuli. Results from three example neurons demonstrate visual
receptive fields estimated at the peak visual latency (*fig. 3b top row*, Online Methods). The
orientation tuning at the peak visual latency (*fig. 3b bottom row*, Online Methods) estimated from
the CEREBRO world camera image (in red) is shown in comparison to the tuning curves
estimated from the known stimuli presented on the screen (in blue). These results replicate classic
findings about the properties of neurons in early visual cortex thereby demonstrating that our eye-

tracking system and calibration approach can accurately record neural responses in response to
visual stimuli on the primate retina.
Following this, the animal was allowed to actively explore a 200cm x 100 cm arena decorated
with various visual stimuli. Activity of single neurons was continuously recorded in the period of
head-restraint and freely-moving condition; body and head movements were simultaneously
recorded using *OptiTrack* system (*fig. 3c*, Online Methods). The comparison of spike waveforms
in head-restraint versus freely-moving condition among 5 example neurons demonstrates the
stability of neural recording throughout the session (*fig. 3d*). Within the same session, the eye
behavior and the neural activity (spike rate) change between the head restrained vs freely moving
animal. We explore some of these behaviors in the following sections.
290
291
292 **Visual behavior of freely moving marmosets:**
293
The primary motivation to develop CEREBRO was to precisely quantify the characteristics of eye-
movements and gaze in freely-moving, naturally behaving primates. To this end, we recorded
visual behavior in marmosets wearing CEREBRO as they explored an open rectangular arena
(*fig 4a*). As marmosets do not continuously move when in open-field environments, we
distinguished between the following two behavioral states in these test sessions to determine
whether differences in visual behavior emerged: a) "*stationary*" - the monkey was seated or
standing and visually scanning the environment without physically changing locations (b)
"*locomotion*" - the monkey was physically moving and changing its position in the environment
(Online Methods). In a typical session, a marmoset remains stationary for extended periods at
fixed locations (*occupancy map in fig 4a*), and then moved between those locations (*gray traces,
fig 4a*). We conjectured that the gaze dynamics likely differs between these two behavioral states,
as each differs in motoric demands and exploratory function.
306
In a freely moving primate, visual exploration is accomplished by changing gaze which can be
defined as a sum of head and eye movements (*fig 4b*). Stable eye positions in this context are
rare because even when an animal fixates at a fixed position in the scene – referred to here as a
gaze-fixation - the eyes must still move smoothly to compensate for any head-movements for
retinal stabilization. These compensatory eye movements reflect the vestibular ocular reflex
(VOR) that well conserved across species and normally engaged to reduce retinal motion that is
due to head and body motion (Wallman et al. 1982; Angelaki and Hess 2005; Angelaki and Cullen
2008; Cullen 2011, 2012). These compensatory movements correlate negatively with the head-
movement velocity to subtract its effect and achieve stable gaze. By contrast, during rapid gaze-
shifts, equivalent to traditional saccades in the head-fixed case, the VOR is suppressed and there
is a combination of conjugate head and eye movements along the same direction, with eye
velocity reversing at the end of the rapid shift as VOR is restored and gaze is again stabilized by
compensatory eye movements compensating for continuing head velocity. The inset in Figure 4b
illustrates this point as the sum of eye and head position (gaze) exhibits steps in position with
stable periods in between. While head position is continuously changing, the eye position exhibits
saw-toothed type patterns in which a jump in position is followed by decay backwards that
compensates for the change in head position. To distinguish epochs of rapid gaze shifts and

compensatory movements, we set a gaze velocity threshold of ±200 °/sec. Examination of eye
movements during gaze-shifts and gaze-fixations revealed a clear trend of negative correlation
between head and eye movements for fixation periods reflecting compensatory eye movements
to stabilize gaze (*purple, fig 4c*) in a freely-moving marmoset. For the gaze velocity threshold set,
we find compensatory movements are well separated from other rapid gaze shifts (*green, fig 4c*).

A core feature of mammalian ocular-motor behavior is known as the main sequence; a
characteristic linear relation between the amplitude and peak velocity of eye movements. To test
whether the main sequence is evident in a freely-moving primate, and characterize it with respect
to head and gaze, we quantified the main sequence and amplitude-duration relationship. For
conjugate gaze shifts we observed a characteristic pattern wherein eye velocity reached a peak
velocity more quickly than head-velocity. As head-velocity decayed with a long tail, eye velocity
reversed direction in order to counter-act the head-velocity and stabilize gaze (Fig 4d). This
pattern was evident both for small and large gaze shifts, with the duration of the gaze shift being
longer for larger gaze shifts. To quantify these patterns more accurately, we next plotted the peak
velocity, latency to peak, and duration of the eye, head, and gaze components as a function of
gaze amplitude (Fig 4e-h). Although the main sequence was evident in freely-moving marmosets
here, evidence suggested differences in how the eye and head components contribute to it as a
function of gaze amplitude (Fig 4e). Whereas the peak velocity of eye movements saturated for
gaze shifts of roughly 10 degrees in amplitude around 400 degrees/sec, the peak velocity of head
shifts continued to increase even for the largest measured shifts out to 80 degrees. The latency
of the peak eye velocity always leads peak head velocity as a function of gaze amplitude with
each growing as a function of shift amplitude (Fig. 4f). The peak of the gaze velocity follows more
closely with eye velocity for small shifts (<36 degrees) and with head-velocity for larger shifts. The
duration of gaze shifts follows a roughly linear relation with gaze amplitude (Fig. 4g). In summary,
the components of eye and head shifts differ in their contribution to gaze shifts depending on the
amplitude of the shift, with eye velocity rapidly saturating in its maximum velocity for shifts of about
10 degrees in size after which the slower initiating head-shift contributes more to the total gaze
movement. As the gaze amplitude becomes larger, the contributions of the head increase linearly
to the overall shifts while the eye saturates in its contribution after about 20 visual degrees (*fig
4h*).

To determine how visual behavior differed between behavioral contexts we compared the eye-
movements of marmosets when head-fixed and freely-moving, distinguishing between instances
when individuals were visually scanning the environment while 'stationary' and instances when
animals were actively moving through the world by 'locomotion' in the latter context (*fig 5a*). As
demonstrated in *fig 5b*, differences in marmosets' visual behavior when head-fixed and both
freely-moving contexts were stark, with the eye movements being notably dynamic in freely-
moving contexts reflecting VOR adjustments for self-motion. We quantified these differences
further by calculating the approximate entropy of marmoset eye movements in each of these three
contexts. This analysis estimates the randomness of a time series where higher value of
approximate entropy means a system that is more random and vice versa – thereby allowing a
comparison of the collective consistency and predictability of the visual behavior. Overall, we
observed that marmoset eye-movements in the head-restrained condition exhibited a significantly

lower entropy than both the freely-moving conditions (stationary and locomotion; *fig 5c*),
suggesting that eye movements in a freely moving animal is more chaotic/random. However, the
approximate entropy for "gaze" of freely-moving animals – the combination of both head and eye-
movements – a different pattern emerged. Here the approximate entropy in both freely-moving
contexts was in fact lower than in the head-fixed context. This suggests that the synergistic
combination of head and eye movements in a freely-moving animal yields a more predictable
visual behavior than when head-fixed, a likely computational optimization of the visual system to
stabilize visual perception in animals as they naturally move and explore the world. The statistics
of maximal speed and amplitude of the eye movement shows a significant faster and larger eye
movement in the freely-moving context (*fig 5d*).

In addition to significant contrasts in visual behavior between head-fixed and freely-moving
contexts, we also observed differences in the latter when marmosets were stationary or
locomoting. Specifically, marmosets make significantly more gaze shifts per second (*fig 5e*) with
greater speeds (*fig 5f*) and larger amplitudes and maximal speed (*fig 5g*) during locomotion than
during stationary phases. This pattern raises the question if the gaze of the animal is less stable
during locomotion given the larger gaze shift amplitudes. To investigate that we calculated at the
Root-mean-square stabilization of the gaze during gaze fixations during locomotion and stationary
contexts. Analyses revealed that gaze was on average about 3 times more stable than head
movements during gaze fixations (*fig 5h*), consistent with compensatory eye movements help
correct for head-motion in these periods. When comparing the stability of gaze to that of head,
however, we observed that although head movements were less stable during locomotion gaze
stability surprisingly improved during locomotion (*fig 5h*), despite the increase in head-motion.
Thus, while the head is clearly less stable during locomotion, compensatory eye movements
appear to provide better stabilization achieving more stable epochs of gaze fixation than even the
sedentary phase. This highlights the importance and potential context-dependence of gaze
stabilization, wherein it appears to be enhanced to achieve greater stability during locomotion.

Discussion:

Here we quantified the active visual behavior of a freely-moving primate by developing an
innovative head-mounted eye-tracking system. Our system achieves several technical
innovations that enabled accurate quantification of head and eye movements in a small bodied
(~300-400g) monkey – common marmosets - with the resolution and speed needed to accurately
quantify primate visual behavior in real-world contexts. Specifically, this system achieves high-
speed recording of the eye (90 FPS) and world (63 FPS), as well as applies an innovative solution
to overcoming the challenge that frequent changes in lighting cause for pupil detection by
leveraging a segmentation neural network. As CEREBRO was designed to seamlessly integrate
with wireless neural recording methods, this system is poised to not only elucidate the dynamics
of active visual behaviors but the supporting neural processes. While findings here recapitulate
the core features of conjugate gaze movements from prior studies using head-free - but chair-
restrained - macaques and extend those to freely-moving primates, analyses also revealed
several related characteristics of visual behavior that have not been observed previously. Perhaps

most notably, we observed that the entropy of the visual system was significantly better when
animals were freely-moving than when head fixed, and that the stability of the visual gaze
improved during locomotion. Our technical innovations described here enable us to examine
visual processing in the evolutionary context to which it has been optimized, wherein coordinated
head-eye movements act to stabilize the visual system during natural exploration, critical features
that cannot be recapitulated in the absence of ethological movements.
418
The contribution of eye and head movements towards conjugate gaze shifts in freely-moving
marmosets were qualitatively similar that of chair restrained but head-free macaques and other
mammals (Martinez-Trujillo, Wang, and Crawford 2003; Freedman and Sparks 1997; Morasso,
Bizzi, and Dichgans 1973; Tomlinson and Bahra 1986; Guitton, Douglas, and Volle 1984;
Collewijn 1977; Goossens and Van Opstal 1997). Although broadly similar, marmosets did exhibit
some quantitative differences relative to macaques, potentially due to their smaller head-size. For
example, marmoset the eye movement velocity and amplitude saturate for much smaller gaze
shifts around 10-20 degrees, after which the head contributes to the bulk of the shift in gaze (Fig.
4h), whereas in the macaque this transition does not occur until one reaches much larger gaze
shifts between 20-40 degrees (Freedman and Sparks 1997). Moreover, most gaze shifts in
macaques under 20 degrees in size are predominantly driven by shifts in eye position and not the
head, while in the marmoset the same gaze shifts would have significant head-movement
components. Only the smallest of gaze shifts under 5-10 degrees in the marmoset, a range
comparable to the movements made in head-fixed marmosets (Fig 5d), are dominated by
changes in eye position after which the head would normally make significant contributions. These
differences likely reflect an efficiency tradeoff related to smaller head-size in marmosets
(Malinzak, Kay, and Hullar 2012).
436
Because we recorded marmoset eye-movements in head-fixed and freely-moving conditions, this
novel dataset can both be compared to prior studies and extend our understanding of visual
behavior in this New World primate. As in previous experiments (Mitchell, Reynolds, and Miller
2014), when head-fixed the oculomotor range is relatively fixed within about 10 visual degrees.
But when marmoset is freely-moving that range increases to roughly 20 degrees (Fig. 5d),
suggesting the limited oculomotor range when head-fixed represents more of a motor preference
than a physical limitation. In a previous study of chaired but head-free marmosets, a paradigm
was used to evoke large gaze shifts involving up to 180-degree head rotation (Pandey, Simhadri,
and Zhou 2020). Although similar large shifts were rare in the study here, a similar linear relation
between peak head velocity and gaze shifts that peak near a velocity of 750 $^\circ$/sec for an 80-
degree gaze shifts were evident in freely-moving marmosets suggesting that this aspect of head-
gaze control are relatively invariant to the form of the task. A direct comparison of the two freely-
moving contexts here revealed a number of notable differences including larger amplitude gaze
shifts with higher maximum velocities when locomoting (Fig. 5, d-g). By contrast, when stationary
and visually scanning the environment, marmoset biased to smaller gaze shifts indicating that
motor demands of actively moving through space likely drives differences in how the head and
eyes coordinate to stabilize the visual field.
454

Our understanding of primate vision is almost entirely based on studies employing conventional
head-fixed paradigms.   While it has long been assumed that primate visual behavior in
conventional, head-fixed paradigms reflects the core features of the system and - once movement
is compensated - be representative of the system in more ethological contexts, there have been
few explicit tests.   Quantifying primate visual behavior with the innovative eye-tracking system
here afforded the powerful opportunity to examine this assumption more directly is indeed
accurate. Indeed, we observed broad qualitative similarities in many features of conjugate eye
movements between head-fixed and freely-moving contexts. However, it was also apparent that
certain assumptions about results in the more conventional paradigm may not be strictly true. The
first being that gaze stability was not worsened despite greater eye and head movements during
locomotion, and in fact, was slightly enhanced. We observed that the approximate entropy of the
marmoset visual system was significantly lower (i.e. improved) when individuals were freely-
moving than head fixed suggesting that the collective visual system is more consistent and
predictable when marmosets are moving and exploring the environment. Likewise, a comparison
of freely-moving marmosets when stationary – i.e. visually scanning - and locomoting revealed
that the stability of visual gaze - as measured by RMS – was significantly better when animals
were locomoting. In other words, despite an increase in the number, speed and length of the gaze
shifts when locomoting, vision actually became more stable.   These findings contradict the
intuition that greater movement freedom would introduce chaos and instability to vision. Rather,
as evidenced here, the coordinated movements of head and eyes have been optimized to
accommodate this self-motion.

The context-dependence of visual stability between the stationary scanning and locomotion states
could reflect changes in VOR at the neural level, as well as other positional strategies that
optimize the VOR system (Vidal, Graf, and Berthoz 1986; Graf et al. 1995). Such context-
dependence of gaze control has long been appreciated from head-mounted eye tracking studies
in humans (Land and Lee 1994; Hayhoe and Ballard 2014), and recently includes free motion in
natural contexts (Matthis, Yates, and Hayhoe 2018). These strategies likely reflect species
specific adaptations to movement. For example, the natural movement statistics of head-position
in free-moving primates is known to differ substantially from that of rodents (Carriot et al. 2014).
However, technical limitations for using free-moving primates in neural investigations, particularly
the precision of conventional pupil thresholding in those contexts, has kept the number of studies
relatively limited (Shepherd and Platt 2006; Milton, Shahidi, and Dragoi 2020; Mao et al. 2021).
The methods introduced here provide a basis to study gaze during free-motion with non-human
primates, even as small as the marmoset (~300-400g). In future work the system could be
expanded to address other features of oculomotor control that could be especially relevant to free-
motion these contexts, including vergence and torsional eye movements (Tweed, Haslwanter,
and Fetter 1998; Ong and Haslwanter 2010). The current eye tracking system offers an
opportunity to study how computational constraints and movement strategies during active vision
act to achieve stability and provide high acuity vision in different contexts, including locomotion
and navigation through the environment. The necessity over evolution to stabilize vision in the
freely-moving context is the basal state of all organisms, including human and nonhuman
primates.
498

The sensory and motor systems of animals co-evolved. Amongst the most significant selective
forces acting on animals is to move and interact with the world, and to do so requires sensory
feedback. This ground truth inherently couples sensory and motor systems. Despite this, the
study of the primate visual system has largely ignored considerations of movement, assuming
instead that a head-restrained animal only able to move their eyes was representative of vision.
Leveraging our innovative head-mounted eye-tracking system for marmosets, we provide
compelling evidence that the coordinated actions of eye and head movements are optimized to
stabilize primate vision and increase its predictability. These patterns emphasize the significance
in considering the ethological relevance of how primates are tested in vision studies. Evidence
suggests that neural responses in head-fixed paradigms are not necessarily predictive of how the
same single neurons – or population ensembles – respond in naturalistic contexts (Jovanovic et
al. 2022; McMahon et al. 2015). Indeed, investigations of vision in freely-moving mice suggest
that at least some elements of neural activity in this context are distinct (P. R. L. Parker et al.
2022; Chaplin and Margrie 2020). Our innovative eye-tracking system can be leveraged in
marmosets to precisely examine the primate visual system during natural, freely-moving
behaviors (Ngo et al. 2022; Shaw, Wang, and Mitchell 2023) and address a suite of foundational
questions that we have as of yet been unable to address.
516

Online Methods:

Fitting the animal with CEREBRO:

The head assembly for CEREBRO is custom fitted to every animal. For the initial fitting, the animal is anesthetized using a combination of Ketamine (dose:20 mg/kg) and Acepromazine (dose: 0.75 mg/kg). All the parts (*Supplementary fig 1b*) are adjusted for best view of the eye and the front camera and fastened using set screws.

Multiple system synchronization:

Our Neural recording systems, motion trackers and CEREBRO were all synchronized using an ESP32 microcontroller based custom board. CEREBRO and the motion tracking system sent TTL signals for different mode shifts that were recorded by the synchronization microcontroller. For synchronizing the neural signal, we sent a TTL pulse (100 mV |100 m) to the electrophysiology system (in recording mode) before putting it on the animal. The timestamp of this signal was logged by the sync microcontroller and since this signal was also logged into the real data stream of the electrophysiology system, we can easily align the time of this pulse in the data to other systems (CEREBRO and motion tracking). The board was connected to a computer via serial communication and the timing data was logged on to a console terminal (CoolTerm).

Animal subjects:

The development and testing of CEREBRO were performed on two common marmosets (monkey M, male; monkey S, female). Both subjects were group housed and were 2 years old at the time of implant. Monkey M had a chronic implant in left V1 and monkey S had bilateral chronic implants in both left and right V1. All surgeries and experiments were performed in the Cortical Systems and Behavior Laboratory at University of California, San Diego (UCSD), and approved by the UCSD Institutional Animal Care and Use Committee in accordance with National Institute of Health standards for care and use of laboratory animals.

Electrophysiology:

Neural activity was recorded with multi-electrode arrays (N-form, Modular Bionics, Berkeley CA) chronically implanted in V1. The N-form array has 64 channels in total embedded in 16 shanks located in a 4x4 grids evenly spaced 0.25 mm apart. Each shank has 4 iridium oxide electrodes located at 0.5, 0.375, 0.25, and 0 mm from the tip. Implantations followed the standard surgical procedures for chronically implanted arrays in primates. In each recording session, a wireless Neurologger (SpikeLog-64, Deuteron Technologies) was hosted in a 3.5 cm (width) x 2.5 cm (height) x 1.2 cm (depth) protecting case and connected to the array to record the extracellular voltage at 32 kHz. Spike sorting was performed offline using Kilo sort (Pachitariu et al. 2016)and manually curated using the graphic user interface Phy.

Receptive field and tuning property of V1 neurons:

For the receptive field mapping, marmosets were head-fixed and freely viewed the stimulus of
120 randomly positioned black and white flashing dots generated at a frequency of 10 Hz for 4
minutes. Eye calibration was performed offline with the GUI provided in this paper. We then used
the calibrated eye position on the world camera images to estimate the visual input on the retina,
and mapped the receptive fields using forward correction (Zhao et al. 2020). For the orientation
and spatial frequency tuning of the V1 neurons, we presented the full-field drifting gratings with
12 orientations (evenly-spaced 30-degree intervals) and 3 spatial frequencies (0.5, 1, 2
cycles/degree). This information was used in calculating the tuning curves with the information
from the screen; when calculating the tuning curves with the information from CEREBRO, we
applied Hough transform on the world camera images to detect the lines and further calculated
the orientation and spatial frequency of the gratings.
574
575 Head and body movement tracking using OptiTrack:
576
The head and body movement of the animals were recorded using OptiTrack- an image-based
motion tracking system. For the head movements, the added 3 IR reflective beads (12mm in
diameter) on the head-assembly of the animal. For the body, we added a single IR bead on to the
backpack. The arena was fitted with 10 OptiTrack cameras strategically placed to cover every
view of the arena. The cameras were calibrated before each session and sessions were
performed only for calibrations where the error for the worst camera was less than 0.1 mm. The
data collected from the session was manually curated to remove stray/false markers and any
gaps were filled with linear interpolation.
585
586 Locomotion behavior analysis:
587
Locomotion is detected with body and head tracking data from OptiTrack. We first calculated the
velocity of the body IR bead and set a threshold of 3 cm/s after applying a 0.2 Hz low-pass filter
to select the period when the monkey had a fast body movement. However, not all the fast body
movement came from locomotion; it could happen when the monkey made a body turn, or a
sudden change of posture. We add two other conditions: 1) the duration of the period is larger
than 3s; 2) the position of the head is lower than 18 cm. These two conditions are determined by
manually matching the data with the video. After the locomotion period is detected, the rest of the
time is defined as sedentary.

608 Data Availability:
609
610 The GUI for UNET segmentation-based pupil detection can be found at
611 *https://github.com/Vickey17/UNET_implementation_V2*. The STL designs for the headpiece and
612 its material description can be downloaded at
613 *https://github.com/Vickey17/CEREBRO_HeadPiece*. Data for the paper can be obtained upon
614 request from the authors.
615
616
617
618
619 **Acknowledgements:**
620

621 We thank Alex Huk and Cris Niell for their invaluable insights throughout the development of this
622 system. We thank Brian Corneil for his valuable comments on the manuscript. This work was
623 supported by grants from The BRAIN Initiative (R01 NS118457 & U01 NS116377) and AFOSR
624 (FA9550-19-1-0357)

639   References:
640

Angelaki, Dora E., and Kathleen E. Cullen. 2008. "Vestibular System: The Many Facets of a
Multimodal Sense." *Annual Review of Neuroscience* 31: 125–50.
Angelaki, Dora E., and Bernhard J. M. Hess. 2005. "Self-Motion-Induced Eye Movements:
Effects on Visual Acuity and Navigation." *Nature Reviews. Neuroscience* 6 (12): 966–76.
Carriot, Jérome, Mohsen Jamali, Maurice J. Chacron, and Kathleen E. Cullen. 2014. "Statistics
of the Vestibular Input Experienced during Natural Self-Motion: Implications for Neural
Processing." *The Journal of Neuroscience: The Official Journal of the Society for
Neuroscience* 34 (24): 8347–57.
Chaplin, Tristan A., and Troy W. Margie. 2020. "Cortical Circuits for Integration of Self-Motion
and Visual-Motion Signals." *Current Opinion in Neurobiology* 60 (February): 122–28.
Collewijn, H. 1977. "Eye- and Head Movements in Freely Moving Rabbits." *The Journal of
Physiology* 266 (2): 471–98.
Cullen, Kathleen E. 2011. "The Neural Encoding of Self-Motion." *Current Opinion in
Neurobiology* 21 (4): 587–95.
———. 2012. "The Vestibular System: Multimodal Integration and Encoding of Self-Motion for
Motor Control." *Trends in Neurosciences* 35 (3): 185–96.
DeAngelis, G. C., I. Ohzawa, and R. D. Freeman. 1993. "Spatiotemporal Organization of
Simple-Cell Receptive Fields in the Cat's Striate Cortex. I. General Characteristics and
Postnatal Development." *Journal of Neurophysiology* 69 (4): 1091–1117.
Emerson, R. C., M. C. Citron, W. J. Vaughn, and S. A. Klein. 1987. "Nonlinear Directionally
Selective Subunits in Complex Cells of Cat Striate Cortex." *Journal of Neurophysiology*
58 (1): 33–65.
Felleman, D. J., and D. C. Van Essen. 1991. "Distributed Hierarchical Processing in the Primate
Cerebral Cortex." *Cerebral Cortex* 1 (1): 1–47.
Freedman, E. G., and D. L. Sparks. 1997. "Eye-Head Coordination during Head-Unrestrained
Gaze Shifts in Rhesus Monkeys." *Journal of Neurophysiology* 77 (5): 2328–48.
Goossens, H. H., and A. J. Van Opstal. 1997. "Human Eye-Head Coordination in Two
Dimensions under Different Sensorimotor Conditions." *Experimental Brain Research.
Experimentelle Hirnforschung. Experimentation Cerebrale* 114 (3): 542–60.
Graf, W., C. de Waele, P. P. Vidal, D. H. Wang, and C. Evinger. 1995. "The Orientation of the
Cervical Vertebral Column in Unrestrained Awake Animals. II. Movement Strategies."
Brain, Behavior and Evolution 45 (4): 209–31.
Guitton, D. 1992. "Control of Eye-Head Coordination during Orienting Gaze Shifts." *Trends in
Neurosciences* 15 (5): 174–79.
Guitton, D., R. M. Douglas, and M. Volle. 1984. "Eye-Head Coordination in Cats." *Journal of
Neurophysiology* 52 (6): 1030–50.
Hayhoe, Mary, and Dana Ballard. 2014. "Modeling Task Control of Eye Movements." *Current
Biology: CB* 24 (13): R622-8.
Jovanovic, Vladimir, Adam Ryan Fishbein, Lisa de la Mothe, Kuo-Fen Lee, and Cory Thomas
Miller. 2022. "Behavioral Context Affects Social Signal Representations within Single
Primate Prefrontal Cortex Neurons." *Neuron* 110 (8): 1318-1326.e4.
Land, M. F., and D. N. Lee. 1994. "Where We Look When We Steer." *Nature* 369 (6483): 742–
44.
Livingstone, M. S., C. C. Pack, and R. T. Born. 2001. "Two-Dimensional Substructure of MT
Receptive Fields." *Neuron* 30 (3): 781–93.
Malinzak, Michael D., Richard F. Kay, and Timothy E. Hullar. 2012. "Locomotor Head
Movements and Semicircular Canal Morphology in Primates." *Proceedings of the
National Academy of Sciences of the United States of America* 109 (44): 17914–19.

Mao, Dun, Eric Avila, Baptiste Caziot, Jean Laurens, J. David Dickman, and Dora E. Angelaki. 2021. "Spatial Modulation of Hippocampal Activity in Freely Moving Macaques." *Neuron* 109 (21): 3521-3534.e6.

Martinez-Trujillo, Julio C., Hongying Wang, and J. Douglas Crawford. 2003. "Electrical Stimulation of the Supplementary Eye Fields in the Head-Free Macaque Evokes Kinematically Normal Gaze Shifts." *Journal of Neurophysiology* 89 (6): 2961–74.

Matthis, Jonathan Samir, Jacob L. Yates, and Mary M. Hayhoe. 2018. "Gaze and the Control of Foot Placement When Walking in Natural Terrain." *Current Biology: CB* 28 (8): 1224-1233.e5.

McMahon, David B. T., Brian E. Russ, Heba D. Elnaiem, Anastasia I. Kurnikova, and David A. Leopold. 2015. "Single-Unit Activity during Natural Vision: Diversity, Consistency, and Spatial Sensitivity among AF Face Patch Neurons." *The Journal of Neuroscience: The Official Journal of the Society for Neuroscience* 35 (14): 5537–48.

Meyer, Arne F., John O'Keefe, and Jasper Poort. 2020. "Two Distinct Types of Eye-Head Coupling in Freely Moving Mice." *Current Biology: CB* 30 (11): 2116-2130.e6.

Meyer, Arne F., Jasper Poort, John O'Keefe, Maneesh Sahani, and Jennifer F. Linden. 2018. "A Head-Mounted Camera System Integrates Detailed Behavioral Monitoring with Multichannel Electrophysiology in Freely Moving Mice." *Neuron* 100 (1): 46-60.e7.

Michaiel, Angie M., Elliott Tt Abe, and Cristopher M. Niell. 2020. "Dynamics of Gaze Control during Prey Capture in Freely Moving Mice." *ELife* 9 (July). https://doi.org/10.7554/eLife.57458.

Miller, Cory T., David Gire, Kim Hoke, Alexander C. Huk, Darcy Kelley, David A. Leopold, Matthew C. Smear, Frederic Theunissen, Michael Yartsev, and Cristopher M. Niell. 2022. "Natural Behavior Is the Language of the Brain." *Current Biology: CB* 32 (10): R482–93.

Milton, Russell, Neda Shahidi, and Valentin Dragoi. 2020. "Dynamic States of Population Activity in Prefrontal Cortical Networks of Freely-Moving Macaque." *Nature Communications* 11 (1): 1–10.

Mitchell, Jude F., John H. Reynolds, and Cory T. Miller. 2014. "Active Vision in Marmosets: A Model System for Visual Neuroscience." *The Journal of Neuroscience: The Official Journal of the Society for Neuroscience* 34 (4): 1183–94.

Morasso, P., E. Bizzi, and J. Dichgans. 1973. "Adjustment of Saccade Characteristics during Head Movements." *Experimental Brain Research. Experimentelle Hirnforschung. Experimentation Cerebrale* 16 (5): 492–500.

Ngo, Victoria, Julia C. Gorman, María Fernanda De la Fuente, Antonio Souto, Nicola Schiel, and Cory T. Miller. 2022. "Active Vision during Prey Capture in Wild Marmoset Monkeys." *Current Biology: CB* 32 (15): 3423-3428.e3.

Ong, James K. Y., and Thomas Haslwanter. 2010. "Measuring Torsional Eye Movements by Tracking Stable Iris Features." *Journal of Neuroscience Methods* 192 (2): 261–67.

Pachitariu, Marius, Nicholas Steinmetz, Shabnam Kadir, Matteo Carandini, and Harris Kenneth D. 2016. "Kilosort: Realtime Spike-Sorting for Extracellular Electrophysiology with Hundreds of Channels." *BioRxiv*. bioRxiv. https://doi.org/10.1101/061481.

Pandey, Swarnima, Sravanthi Simhadri, and Yi Zhou. 2020. "Rapid Head Movements in Common Marmoset Monkeys." *IScience* 23 (2): 100837.

Parker, A. J., and W. T. Newsome. 1998. "Sense and the Single Neuron: Probing the Physiology of Perception." *Annual Review of Neuroscience* 21: 227–77.

Parker, Philip R. L., Elliott T. T. Abe, Emmalyn S. P. Leonard, Dylan M. Martins, and Cristopher M. Niell. 2022. "Joint Coding of Visual Input and Eye/Head Position in V1 of Freely Moving Mice." *Neuron* 110 (23): 3897-3906.e5.

Ringach, D. L., G. Sapiro, and R. Shapley. 1997. "A Subspace Reverse-Correlation Technique for the Study of Visual Neurons." *Vision Research* 37 (17): 2455–64.

Ronneberger, Olaf, Philipp Fischer, and Thomas Brox. 2015. "U-Net: Convolutional Networks for Biomedical Image Segmentation." *ArXiv [Cs.CV]*. arXiv. https://doi.org/10.48550/ARXIV.1505.04597.

Shaw, Luke, Kuan Hong Wang, and Jude Mitchell. 2023. "Fast Prediction in Marmoset Reach-to-Grasp Movements for Dynamic Prey." *Current Biology: CB* 33 (12): 2557-2565.e4.

Shepherd, Stephen V., and Michael L. Platt. 2006. "Noninvasive Telemetric Gaze Tracking in Freely Moving Socially Housed Prosimian Primates." *Methods* 38 (3): 185–94.

Tomlinson, R. D., and P. S. Bahra. 1986. "Combined Eye-Head Gaze Shifts in the Primate. I. Metrics." *Journal of Neurophysiology* 56 (6): 1542–57.

Tweed, D., T. Haslwanter, and M. Fetter. 1998. "Optimizing Gaze Control in Three Dimensions." *Science* 281 (5381): 1363–66.

Vidal, P. P., W. Graf, and A. Berthoz. 1986. "The Orientation of the Cervical Vertebral Column in Unrestrained Awake Animals. I. Resting Position." *Experimental Brain Research. Experimentelle Hirnforschung. Experimentation Cerebrale* 61 (3): 549–59.

Wallace, Damian J., David S. Greenberg, Juergen Sawinski, Stefanie Rulla, Giuseppe Notaro, and Jason N. D. Kerr. 2013. "Rats Maintain an Overhead Binocular Field at the Expense of Constant Fusion." *Nature* 498 (7452): 65–69.

Wallman, J., J. Velez, B. Weinstein, and A. E. Green. 1982. "Avian Vestibuloocular Reflex: Adaptive Plasticity and Developmental Changes." *Journal of Neurophysiology* 48 (4): 952–67.

Yates, Jacob L., Shanna H. Coop, Gabriel H. Sarch, Ruei-Jr Wu, Daniel A. Butts, Michele Rucci, and Jude F. Mitchell. 2023. "Detailed Characterization of Neural Selectivity in Free Viewing Primates." *Nature Communications* 14 (1): 3656.

Zhao, Yuan, Jacob L. Yates, Aaron J. Levi, Alexander C. Huk, and Il Memming Park. 2020. "Stimulus-Choice (Mis)Alignment in Primate Area MT." *PLoS Computational Biology* 16 (5): e1007614.

774 Figure 1:

775
776

Fig1: Head mounted eye-tracker assembly

(a) 3D render depicting different parts of the head-piece. The Ti- alloy scaffolding is custom
designed and fabricated using DLS **(b)** 3D printed backpack that holds the 2 PCBs and the
battery. **(c)** The animals are fitted with two 1cm headposts that act as anchors for holding the
head-piece. **(d)** Custom PCB board which drives the camera sensors, IMU and saves the data to
a locally mounted SD card. **(e)** A marmoset wearing the fully assembled system (CEREBRO). **(f)**
View from the 3 different cameras (eye camera(pink), world camera(yellow) and the external
arena camera (cyan)) during a freely-moving session. **(g)** Bar graphs plot the % movement (top)
average distance travelled (bottom-left) and average speed in freely-moving marmosets with
(purple) and without (blue) CEREBRO.

787
788   Figure 2:

789
790

Fig2: Iris/Pupil detection using segmentation Artificial Neural Network.

(a) Conventional thresholding approach for pupil detection fails in scenes with variable light intensities. **(b)** Architecture of the segmentation neural network to crop out the iris. **(c)** The performance of the ANN matches closely to that of human annotation with relatively low training epochs (10 in this example) and relatively small amount of training data (~250 samples) **(d)** For calibrating the eye and world camera, the animals are presented with marmoset faces on the screen in a predetermined layout. **(e)**The calibrated eye movements can be overlaid on the world scene to determine the visual scene falling on the retina of the animal.

808   Figure 3:

809
810

Fig3: Eye tracking coupled electrophysiology in freely moving marmosets.
(a) To validate use of CEREBRO with electrophysiology, subjects were presented with orientation
grating and receptive field stimuli in a classical head fixed preparation. **(b)** The receptive field (top)
and orientation tuning curve (bottom) is shown for three representative neurons recorded in
marmoset V1 while using CEREBRO. **(c)** Example raster of 30+ single neurons in marmoset V1
while subjects are head-restrained (left) and freely-moving (right) while wearing CEREBRO. The
head (blue) and body (purple) speed of the marmoset using CEREBRO is plotted below the freely-
moving raster **(d)** Example spike waveforms from five exemplar neurons demonstrate that we
were able to stably record from same neurons in head-restraint vs freely moving conditions.

Figure 4:

821
Fig4.: Gaze characteristics in freely moving marmosets.

(a) Representative session of activity pattern for marmosets in an open arena (200 cm x 100 cm
x 240 cm). Grey dotted lines indicate periods of locomotion, while heat maps plot stationary
occupancy. **(b)** Plots the estimate of the animal's gaze (blue) with CEREBRO. Horizontal eye
movement (red) and head yaw (green) are also plotted. Inset magnifies a 2s period of time. **(c)**
Horizontal eye movement and head yaw in freely-moving marmosets are shown. Green dots
indicate gaze fixations, while purple dots indicate gaze shifts. **(d-h):** eye-movements (red), head-
movements (green) and gaze-shifts (red) in freely-moving marmosets. **(d)** Plots the position (top)
and velocity (bottom) of eye-movements, head-movements and gaze-shifts <10degrees (left) and
>10degrees (right) **(e)** Shows the maximum speed and gaze amplitude. Plots the latency to max
speed (f), duration (g) and amplitude (h) as a function of gaze amplitude

Figure 5:

835
Fig5.: Characteristics of eye movements in freely moving marmosets.
(a) Illustration depicts the two typical modes of visual behavior in freely-moving marmosets:
locomotion. (left) and stationary (right). **(b)** Representative marmoset eye-movements in three
behavioral contexts: head-fixed (top - grey line), freely-moving locomotion (middle – red line) and
freely-moving stationary (bottom – black). **(c)** Plots the approximate entropy of visual behavior
across contexts. Eye-movements during the head-fixed (grey), locomotion (red) and stationary
(black) are shown, while gaze during locomotion (brown) and stationary (dark green) is shown to
the right. Gaze reflects the conjugate movements of the head and eyes. *** indicates p<.0001 and
n.s. – not significant. **(d)** Plots the maximum speed (left) and probability (right) as a function of
eye amplitude in the head-fixed (grey), stationary (black) and locomotion (red) contexts. **(e)** A
comparison of gaze shifts per second when freely-moving marmosets were either stationary
(black) or locomoting (red). **(f)** Plots probability of gaze speed in the stationary (black) and
locomotion(red) contexts. **(g)** Compares the maximum speed (left) and probability (right) of gaze
shift across different gaze amplitudes in the stationary (black) and locomotion (red) contexts. **(h)**
Quantifies RMS stability in five behavioral contexts- head-fixed (blue), gaze sedentary (black) and
locomotion (red), head-movements in sedentary (black dotted line) and locomotion (red dotted
line). Left - Plots the probability of RMS stability in various contexts. Right – Shows the median
RMS stability across contexts. *p<.01.