1

Active vision in freely moving marmosets using head-mounted eye tracking.

Vikram Pal Singh[1, *], Jingwen Li[1], Kana Dawson[1], Jude F. Mitchell[2], Cory T. Miller[1,3, **]

[1] Cortical Systems & Behavior Lab, University of California San Diego, San Diego, California, USA
[2] Department of Brain and Cognitive Science, University of Rochester, Rochester, New York, USA
[3] Neurosciences Graduate Program, University of California San Diego, San Diego, California, USA

*Correspondence: vpsingh@ucsd.edu
**Correspondence: corymiller@ucsd.edu

Abstract:

Our understanding of how vision functions as primates actively navigate the real-world is remarkably sparse. As most data have been limited to chaired and typically head-restrained animals, the synergistic interactions of different motor actions/plans inherent to active sensing – e.g. eyes, head, posture, movement, etc. - on visual perception are largely unknown. To address this considerable gap in knowledge, we developed an innovative wireless head-mounted eye tracking system called **CEREBRO** for small mammals, such as marmoset monkeys. Our system performs **C**hair-free **E**ye-**Re**cording using **B**ackpack mounted mic**RO**controllers. Because eye illumination and environment lighting change continuously in natural contexts, we developed a segmentation artificial neural network to perform robust pupil tracking in these conditions. Leveraging this innovative system to investigate active vision, we demonstrate that although freely-moving marmosets exhibit frequent compensatory eye movements equivalent to other primates, including humans, the predictability of the visual behavior (gaze) is higher when animals are freely-moving relative to when they are head-fixed. Moreover, despite increases in eye/head-motion during locomotion, gaze stabilization remains steady because of an increase in VOR gain during locomotion. These results demonstrate the efficient, dynamic visuo-motor mechanisms and related behaviors that enable stable, high-resolution foveal vision in primates as they explore
the natural world.

Significance Statement:

Vision is arguably the most thoroughly understood of all neural systems in the primate brain. Yet there is little known about how vision functions in real-world contexts in which individuals freely move and explore an environment. This dearth in knowledge is largely due to the lack of technology that can accurately track eye-movements in freely-moving individuals with the speed and resolution needed to quantify primate vision. Here we developed an innovative wireless head-mounted eye-tracking system for marmosets that meets these technical needs and enabled us to quantify facts of primate vision in a manner not previously possible, including a set of discoveries that are likely to transform our understanding of this keystone system.

Keywords:
Marmosets, Active-vision, eye-tracking, gaze stabilization

Introduction:

Primate vision has been the subject of intense study for many decades and is arguably the most well understood neural system in the simian brain (1, 2). And yet, our understanding of primate vision is incomplete. Like all sensory systems, primate vision evolved in response to the challenges inherent to actively moving, exploring and engaging with objects, individuals and the environment from different perspectives (3–5). While the processes of visual encoding have been extensively studied with head-restrained subjects observing stimuli presented on a screen, details of how vision functions as primates actively move through and explore the real-world are remarkably limited. Similar to all vertebrates, the head and eyes coordinate their respective movements in primates to enable a stable percept of visual inputs (6–10). Previous studies in chaired but head-free macaques demonstrate how the eye and head coordinate to stabilize gaze but have yet to address similar coordination in freely-moving primates. And in marmosets, though we have some understanding that their oculomotor range is more restricted than macaques when head-fixed (11) and that their head-position can shift more rapidly than humans or macaques (12), we still lack descriptions of the head-eye coordination for gaze shifts in freely-moving scenarios. Because data on this complement of mechanisms has been limited to chaired animals unable to locomote, the synergistic effects of different motor actions – e.g. eyes, head, posture, movement, etc. - on primate visual perception and cognition during active exploration of the world are almost entirely unknown. The principal bottleneck being technical. Several previous studies have sought to bridge this gap and achieved limited precision for projecting the gaze of free-moving non-human-primates (NHPs) into visual scenes (13–15), but due to technical constraints of these methods, no studies have been able to quantify the stability of visual gaze nor the eye and head dynamics during freely moving active exploration. Here we overcome these obstacles and introduce a method that precisely quantifies eye-movements and can accurately project the gaze of a NHP into scenes as individuals freely explore an environment.

Recent work with mice demonstrates that eye-tracking systems can be miniaturized and mounted to the head in order to provide insight into vision during natural visual behavior (16–19), but these systems are not well suited for comparable studies in primates for at least two critical reasons. First, systems in mice rely on a tether which restricts the 3D mobility of primates. Second, they lack the precision and temporal resolution needed to accurately characterize high-resolution vision of primates. To address this challenge, we developed an innovative head-mounted eye tracking system to enable the study of active, natural visual behaviors, and related neural processes, in freely-moving marmosets. Our system - **CEREBRO** - allows for **C**hair-free **E**ye-**Re**cording using **B**ackpack mounted mic**RO**controllers at a speed and resolution needed to accurately quantify the visual behavior and underlying neural mechanisms of natural, active vision in primates. Using CEREBRO we confirmed that freely-moving marmosets exhibit frequent compensatory eye movements that enable them to stabilize gaze when viewing real world scenes consistent with previous studies in body restrained animals. By using this innovative system, however, we discovered that visual gaze stabilization and predictability was enhanced when the monkeys were moving naturally despite increases in eye/head-motion during locomotion. This suggests that previously unreported synergistic

121 mechanisms for gaze stabilization are not only integral to primate active vision in the real-world
122 but can be enhanced for greater compensation as animals move naturally through their
123 environment.
124
125
126
127 **Results:**
128
129 *CEREBRO **is a head mounted eye tracking system for freely moving marmosets:***
130
The small body size of marmosets (~300-400g) necessitated certain design considerations
when developing CEREBRO. The first of these decisions is related to the weight and wearability
of the system itself. Based on previous experience, the estimated 60g weight of the complete
system would not be feasible to be entirely situated on animal's head without affecting its visual
behavior. To resolve this issue, we separated the system into two separate, but integrated
hardware submodules: the **Head-piece** and the **Backpack**. The *Head-piece* includes the
camera assembly and scaffold fitted to the animals' head *(fig 1a)*; and the *Backpack* includes
the backend electronics for camera synchronizing, image acquisition and local data storage on
the custom designed printed circuit boards (PCBs) *(fig1b)*. Both the PCB and the head-piece IR
LED are powered using a 600 mAh, Lithium-Polymer (Li-Po) battery that is housed inside the
backpack. OV4689 camera modules (SincereFirst, Guangzhou, China) were used to record the
eye and the world scene. For more details about camera interface see Materials & Methods
(Camera and communication interface).
144
Head-Piece Module. The head-piece weighs ~20g and comprises five different components (*fig*
1a): (1) the scaffold: a curved metal tube that serves as an anchor for all the pieces, (2) eye-
cam: an HD MIPI camera (90fps) with Visible light filter and a Macro lens for looking at the eye,
(3) world cam: an HD MIPI camera (60fps) for capturing the world scene in front of the camera,
(4) an IR LED to illuminate the eye, and (5) a strategically placed Hot mirror to image the eye.
The mechanical parts for the head-piece are made up of a Titanium alloy (Ti-6Al-4V) due to
strength and weight considerations, and are held to the scaffold using M2 and M4 screw sets
(see Supplementary Information 1 for detailed assembly description). To achieve reliable eye-
tracking in a fully unrestrained animal, the eye camera's position requires to be fixed with
respect to the animal's eye. This is accomplished by attaching 2 vertical headposts (5 mm
diameter, 1 cm tall cylinders with a flat cut on one side) on the animals' head (*fig 1c*). The two
headposts restrict any rotation or translational movement of the headpiece and allow the
headpiece to be placed at the same position for every recording session.
158
159 **Backpack Module.** The backpack module weighs 40g and comprises two Customized PCBs,
160 as well as the casing and harness worn by the animals (*fig 1b*). Each camera has its own
161 dedicated PCB (*fig 1d*). Both cameras (eye cam and world cam) have long flex cables (15 cm)
162 and are operated by the PCB.
163

164   **Customized Printed Circuit Boards.** The PCB used here is an embedded system that runs on
165   an STM32H750 microprocessor (*fig 1d*). The custom PCB consists of a 0.96" SPI TFT display to
166   preview camera frames and store data image stream and IMU data on a SD card. For more
167   details of the PCB and data storage logic, see Materials & Methods (Printed Circuit Board and
168   data storage).
169

Eye-Tracking System Wearability. When fully configured on a marmoset (fig 1e) the
CEREBRO allows a full range of motions characteristic of the marmoset's natural behavior,
while monitoring its position in the environment, eye position, and view of the scene (*fig 1f*).
Overall, CEREBRO weighs ~60 g (headpiece module: 20g + backpack module: 40g), which is
comparable to the weight of two infant marmosets that an adult would normally carry on its
back. The habituation protocol for adjusting the animals to carrying the backpack and the head-
piece is discussed in Materials & Methods (Habituation of animals to backpack). To test the
influence of this added weight on mobility, we compared marmosets' behavior both with and
without CEREBRO in multiple sessions (30 min) of freely-moving exploration in a large open
arena, (200 cm x 100 cm x 240 cm). The animals were tracked using Optitrack imaging systems
(see Methods). Using the velocity threshold of 5 cm/sec, we quantified the percentage of time
animals spend moving through the arena. Over 11 sessions spread across months, we
observed no significant difference (*fig 1g*) in the time animals spent locomoting vs stationary
and scanning the environment ($p=0.83$). Even though the animals moved for the same amount
of time, there is still a possibility that the weight can impede the speed of the animal. Upon
investigation, we did not observe any significant difference in either the distance covered by the
animals ($p=0.569$) or the average speed of the animal with and without CEREBRO ($p=0.5692$).
This leads us to conclude that our system does not significantly impede the animals' movement
and can be comfortably used with a freely behaving animal.

189
Fast, efficient and accurate pupil detection:

In the traditional experimental paradigms that have dominated primate neuroscience, animals
are head-fixed and the light source to illuminate the eye can be controlled. The consistency of
the light allows for highly accurate eye tracking to be achieved by thresholding the dark iris
observed under infra-red light from a brightly illuminated image and the pupil detected either
using its centroid or center of the ellipse fit. A critical challenge for eye-tracking in freely moving
animals is the lack of control over the illumination. As an individual moves through any real-
world environment external lighting conditions and shadows are constantly changing. This point
is illustrated by our application of thresholding for pupil detection of a freely-moving marmoset
(*Fig 2a*), which can lead to labeling of shadows at the edge of the eye rather than the pupil.
Previous studies that examined gaze in free-moving primates relied on conventional pupil
thresholding (13–15), which limited the precision that could be obtained. To achieve accurate
primate eye-tracking in natural, freely-moving conditions, an alternative method for pupil
detection was needed.

Artificial Neural Networks (ANNs) offer an alternative approach to pupil detection under real-
world conditions because they rely on features of an image and not necessarily the grayscale
values, and therefore are resistant to brightness changes and/or shadows/occlusions. Such
approaches have been successfully applied using commercial software, such as DeepLabCut
(DLC), to track the pupil of freely-moving mice (16). To further optimize pupil tracking in freely-
moving marmosets, here we developed a custom semantic segmentation ANN called UNet (*fig*
2b) (20) that yielded superior performance in these conditions. This workflow allows to detect
pupil features robust from lighting and noise (*fig 2b*). With a fully trained network, we reliably
detected pupils across various sessions and different animals. The network is easily trainable
with fewer epochs for marmoset eyes and a training dataset of only 250-500 images (*fig 2c*).
The data preparation, training of the ANN, post-processing, and the use of a user-friendly GUI
are explained in detail in *Supplementary information 4* (*Supplementary fig 4b*). This UNet
approach closely follows an architecture used previously to segment the pupil from images of
human eyes (e.g. RITnet (21). We compared the performance of UNet to the RITnet model
(Supplementary Fig. 5). The UNet performed better at identifying the pupil from marmoset eye
images, including novel views that varied in position and lighting, but was less robust on human
eye images, which is consistent with the respective emphasis in the two models' training sets.
Future modifications of UNet could include additional architecture features of the RITnet model
to potentially improve performance.

Accurately calculating the gaze point from the world camera necessitates that the pupil position
from the eye camera be calibrated to the real-world position. To this end, we developed the
following procedure. The animal is chaired and head-fixed in front of a computer screen placed
~35 cm directly in front of the subject. Because marmosets naturally look at faces (11), 1-12
small marmoset faces are presented as calibration targets at multiple locations on the display
monitor (*fig 2d*). A custom designed graphical user interface was used to adjust for scaling and
offset in horizontal and vertical axes by a human operator offline. This allowed us to reliably
map pupil eye position to screen coordinates (*fig 2e*), which when head-free, generalized to
world coordinates in front of the marmoset. The GUI for pupil detection can be found at
https://github.com/Vickey17/UNET_implementation_V2. To compare the system against other
pupil-based eye trackers we computed the root-mean-square (RMS) stability of eye position
during stable fixation epochs when animals were head-fixed. We find that across 7 recording
sessions the RMS stability was 0.05 deg (+/- 0.0012 std). These estimates provide a lower
bound on the system precision that rivals head-stabilized pupil-based eye trackers such as
Arrington eye trackers. To estimate our accuracy with respect to the visual stimuli on the world
camera, we created an error estimation algorithm (see Materials & Methods: Error estimation for
eye in world calibration) wherein we measure the distance between the calibrated eye position
and the visual stimulus (a marmoset face; *fig 2f*). For comparing CEREBRO with the Arrington
eye tracker, subjects were head-restrained and presented with marmoset faces on a monitor
placed 35 cm from the animal. We observed that CEREBRO had significantly better accuracy
as compared to the Arrington eye tracker system (*fig 2g*).

To estimate the accuracy of eye/gaze position in a freely moving animal, we designed a novel
paradigm (Fig 3a). Details of the setup are described in the Materials & Methods (Freely moving
eye calibration setup). Briefly, the testing arena consists of a plexiglass box with a perch for the

animals to sit on. All sides of the box are painted with black acrylic to make it completely opaque
except the front panel which is covered with an IR dichroic filter that allows IR light to pass for
video recording of the animal inside while blocking the visible light for what the animal can see.
A small window (10 x 10 cm) is cut in the filter to allow the animal visual access to see the
screen outside. While the CEREBRO equipped animals sit in the dark, a top-mounted
galvanometric LASER system draws geometric lines and points on a screen placed ~75 cm
from the box. Each shape is drawn for a duration of 3 sec followed by a dark period of 3 sec.
The LASER draws subject's gaze to these shapes, thereby allowing an estimate of the eye
target calibration and its error. In a second set of trials, we also used a single point that moved
slowly (<5 visual degrees per second) along random trajectories. In either set of saccade or
smooth pursuit trials, the gaze is projected onto the coordinates of the screen through the world
view camera and error computed as the nearest distance to a point or line within 3.5 visual
degree maximum bound to register looking at the feature (*fig 3b*). Projected eye position was
found to track geometric shapes defined by points and lines (*top panels, fig 3c*) and during
pursuit of the point (*bottom panels, fig 3c*). To compare accuracy against the head-restrained
preparation we presented the same stimuli to head-fixed marmosets. We observe that errors
cluster below 2 visual degrees in all conditions, well away the maximum bound for registering
looking at a feature, and the median accuracy remained less than 1 degree in both saccade
trials for geometric shapes (*fig 3d*) and smooth pursuit trials for a moving point (*fig 3e*). The
median error was higher for both trial types in the freely-moving compared to head-restrained
condition with an increase in error by 1.98 times for saccade trials and 1.4 times for pursuit trials
(Wilcoxon rank sum test, p<0.0001 saccade; p=0.006 smooth pursuit), but still remained under
1 visual degree, validating an upper bound of the system's accuracy.

270
271 **Electrophysiology in combination with CEREBRO:**
272
CEREBRO was designed explicitly to create a tool to investigate the neurobiology of natural,
active vision in freely-moving monkeys. As such, integral to its design was the capacity to
simultaneously record neural activity, eye behavior, and the visual scene of the animal, as each
are integral to this broader ambition. We tested the validity and sufficiency of the system to this
end by performing experiments with CEREBRO while activity of single neurons was recorded
with chronically implanted multi-electrode arrays (N-form, Modular Bionics) in the primary visual
cortex (V1) of the marmosets. These experiments sought to (a) estimate visual tuning
properties of neurons in primate visual cortex, i.e. receptive field mapping, as well as orientation
and spatial frequency tuning (22–25), using CEREBRO in more traditional head-fixed
paradigms, so as to demonstrate the accuracy of our eye-tracking system by replicating these
classic effects, and (b) obtain eye, head and body behavior, visual scene, and activity of single
neurons simultaneously in freely-moving paradigm to demonstrate the capacity of CEREBRO in
investigating primate active vision.
286
287 To recapitulate the tuning properties of V1 neurons, subjects were head-fixed while wearing
288 CEREBRO and presented with the following stimuli: flashing dots for receptive field mapping,
289 and drifting gratings with different orientations and spatial frequencies for tuning properties (*fig.
290 4a*; see online methods). Critically, marmosets were allowed to free-view the video screen

during stimulus presentation and offline corrections for eye position enable accurate reconstruction of visual properties following a recently developed free-viewing approach (26). A key difference here from the previous study with head-fixed marmosets is that the visual input is obtained by the world camera's view along with the estimated eye position from CEREBRO instead of what was known to be displayed on the screen, thus validating that these methods could generalize to real-world stimuli. Results from three example neurons demonstrate visual receptive fields estimated at the peak visual latency (*fig. 4b top row*, Online Methods). The orientation tuning at the peak visual latency (*fig. 4b bottom row*, Online Methods) estimated from the CEREBRO world camera image. These results replicate classic findings about the properties of neurons in early visual cortex thereby demonstrating that our eye-tracking system and calibration approach can accurately record neural responses in response to visual stimuli on the primate retina.

Following this, the animal was allowed to actively explore a 200cm x 100 cm arena decorated with various visual stimuli. Activity of single neurons was continuously recorded in the period of head-restraint and freely-moving condition; body and head movements were simultaneously recorded using *OptiTrack* system (*fig. 4c*, Online Methods). The comparison of spike waveforms in head-restraint versus freely-moving condition among 5 example neurons demonstrates the stability of neural recording throughout the session (*fig. 4d*). Within the same session, the eye behavior and the neural activity (spike rate) change between the head restrained vs freely moving animal. We explore some of these behaviors in the following sections.

Visual behavior of freely moving marmosets:

We compared the eye behavior of a head-restrained marmoset with that of a freely moving marmoset. In the head-restrained context, the animal was presented with a series of naturalistic images (see Materials & Methods). During the freely-moving context, the animal was placed in an open arena and allowed to explore the environment. Consistent with previous studies (11), analysis in the head-restrained context revealed that the eye in head position i.e. the horizontal and vertical eye position spanned +/- 5 visual degrees for 2SD of all the positions (fig 5a) and the eye position distribution for saccades vs fixation were identical in this case (fig 5b & c). However, the results for freely-moving marmoset are starkly different. Firstly, the eye in head position is much more limited in a freely moving animal spanning only +/- 2.5 visual degrees (fig 5d). Moreover, most of the eye positions at the edges of the range were involved in gaze shifts and not gaze fixations (fig 5e & f). These findings suggest that during stable epochs of gaze the head-position provides a reliable estimate of gaze position. The primary motivation to develop CEREBRO was to precisely quantify the characteristics of eye-movements and gaze in freely-moving, naturally behaving primates. To this end, we recorded visual behavior in marmosets wearing CEREBRO as they explored an open rectangular arena (*fig 6a*). As marmosets do not continuously move when in open-field environments, we distinguished between the following two behavioral states in these test sessions to determine whether differences in visual behavior emerged: a) "*stationary"* - the monkey was seated or standing and visually scanning the environment without physically changing locations (b) "*locomotion*" - the monkey was physically

335 moving and changing its position in the environment (Online Methods). In a typical session, a
336 marmoset remains stationary for extended periods at fixed locations (*occupancy map in fig 6a*),
337 and then moved between those locations (*gray traces, fig 6a*). We conjectured that the gaze
338 dynamics likely differs between these two behavioral states, as each differs in motoric demands
339 and exploratory function.
340
In a freely moving primate, visual exploration is accomplished by changing gaze which can be
defined as a sum of head and eye movements (*fig 6b*). Stable eye positions in this context are
rare because even when an animal fixates at a fixed position in the scene – referred to here as
a gaze-fixation - the eyes must still move smoothly to compensate for any head-movements for
retinal stabilization. These compensatory eye movements reflect the vestibular ocular reflex
(VOR) that well conserved across species and normally engaged to reduce retinal motion that is
due to head and body motion (8, 9, 27–29). These compensatory movements correlate
negatively with the head-movement velocity to subtract its effect and achieve stable gaze. By
contrast, during rapid gaze-shifts, equivalent to traditional saccades in the head-fixed case, the
VOR is suppressed and there is a combination of conjugate head and eye movements along the
same direction, with eye velocity reversing at the end of the rapid shift as VOR is restored and
gaze is again stabilized by compensatory eye movements compensating for continuing head
velocity. In our current study, we have primarily focused on the horizontal eye movements and
head yaw since it has been demonstrated that horizontal pursuit eye movements are more
accurate and symmetric than vertical ones in primates (30, 31). The inset in Figure 6b illustrates
this point as the sum of eye and head position (gaze) exhibits steps in position with stable
periods in between. While head position is continuously changing, the eye position exhibits saw-
toothed type patterns in which a jump in position is followed by decay backwards that
compensates for the change in head position. To distinguish epochs of rapid gaze shifts and
compensatory movements, we set a gaze velocity threshold of ±200 °/sec. Examination of eye
movements during gaze-shifts and gaze-fixations revealed a clear trend of negative correlation
between head and eye movements for fixation periods reflecting compensatory eye movements
to stabilize gaze (*purple, fig 6c*) in a freely-moving marmoset. For the gaze velocity threshold
set, we find compensatory movements are well separated from other rapid gaze shifts (*green,*
fig 6c).
366
A core feature of mammalian ocular-motor behavior is known as the main sequence; a
characteristic linear relation between the amplitude and peak velocity of eye movements. To
test whether the main sequence is evident in a freely-moving primate, and characterize it with
respect to head and gaze, we quantified the main sequence and amplitude-duration
relationship. For conjugate gaze shifts we observed a characteristic pattern wherein eye velocity
reached a peak velocity more quickly than head-velocity. As head-velocity decayed with a long
tail, eye velocity reversed direction in order to counter-act the head-velocity and stabilize gaze
(*fig 6d*). This pattern was evident both for small and large gaze shifts, with the duration of the
gaze shift being longer for larger gaze shifts. To quantify these patterns more accurately, we
next plotted the peak velocity, latency to peak, and duration of the eye, head, and gaze
components as a function of gaze amplitude (*fig 6e-h*). Although the main sequence was
evident in freely-moving marmosets here, evidence suggested differences in how the eye and

head components contribute to it as a function of gaze amplitude (*fig 6e*). Whereas the peak
velocity of eye movements saturated for gaze shifts of roughly 10 degrees in amplitude around
400 degrees/sec, the peak velocity of head shifts continued to increase even for the largest
measured shifts out to 80 degrees. The latency of the peak eye velocity always leads peak head
velocity as a function of gaze amplitude with each growing as a function of shift amplitude (*fig.*
6f). The peak of the gaze velocity follows more closely with eye velocity for small shifts (<36
degrees) and with head-velocity for larger shifts. The duration of gaze shifts follows a roughly
linear relation with gaze amplitude (*fig. 6g*). In summary, the components of eye and head shifts
differ in their contribution to gaze shifts depending on the amplitude of the shift, with eye velocity
rapidly saturating in its maximum velocity for shifts of about 10 degrees in size after which the
slower initiating head-shift contributes more to the total gaze movement. As the gaze amplitude
becomes larger, the contributions of the head increase linearly to the overall shifts while the eye
saturates in its contribution after about 20 visual degrees (*fig 6h*).
392
To determine how visual behavior differed between behavioral contexts we compared the eye-
movements of marmosets when head-fixed and freely-moving, distinguishing between instances
when individuals were visually scanning the environment while 'stationary' and instances when
animals were actively moving through the world by 'locomotion' in the latter context (*fig 7a*). As
demonstrated in *fig 7b*, differences in marmosets' visual behavior when head-fixed and both
freely-moving contexts were stark, with the eye movements being notably dynamic in freely-
moving contexts reflecting VOR adjustments for self-motion. We quantified these differences
further by calculating the approximate entropy of marmoset eye movements in each of these
three contexts (32). This analysis estimates the randomness of a time series where higher value
of approximate entropy means a system that is more random and vice versa – thereby allowing
a metric for quantification of the visual behavior as a time series. Overall, we observed that
marmoset eye-movements in the head-restrained condition exhibited a significantly lower
entropy than both the freely-moving conditions (stationary and locomotion; *fig 7c*), suggesting
that eye movements in a freely moving animal is more chaotic/random. However, the
approximate entropy for "gaze" of freely-moving animals – the combination of both head and
eye-movements – a different pattern emerged. Here the approximate entropy in both freely-
moving contexts was in fact lower than in the head-fixed context. This suggests that the
synergistic combination of head and eye movements in a freely-moving animal yields a less
random/chaotic scanning of the visual environment than when head-fixed, a likely computational
optimization of the visual system to sample the scene in animals as they naturally move and
explore the world populated with elements with varying salience. The statistics of maximal
speed and amplitude of the eye movement shows a significant faster and larger eye movement
in the freely-moving context (*fig 7d*).
416
In addition to significant contrasts in visual behavior between head-fixed and freely-moving
contexts, we also observed differences in the latter when marmosets were stationary or
locomoting. Specifically, marmosets make significantly more gaze shifts per second (*fig 7e*) with
greater speeds (*fig 7f*) and larger amplitudes and maximal speed (*fig 7g*) during locomotion than
during stationary phases. This pattern raises the question if the gaze of the animal is less stable
during locomotion given the larger gaze shift amplitudes. To investigate that, we calculated the

Root-mean-square (RMS) stabilization of the gaze during gaze fixations during locomotion and
stationary contexts (16). We use RMS as a measure to quantify the deviation of gaze and head
yaw during stabilization periods (gaze fixations). A smaller RMS during fixation epochs indicates
a more stable fixation while a higher value for RMS will indicate poor stabilization. Analyses
revealed that gaze was on average about 3 times more stable than head movements during
gaze fixations (*fig 7h*). When comparing the stability of gaze to that of head, however, we
observed that although head movements were less stable during locomotion gaze stability
surprisingly improved during locomotion (*fig 7h*), despite the increase in head-motion. Thus,
while the head is clearly less stable during locomotion, compensatory eye movements appear to
provide better stabilization achieving more stable epochs of gaze fixation than even the
sedentary phase. This highlights the importance and potential context-dependence of gaze
stabilization, wherein it appears to be enhanced to achieve greater stability during locomotion.
435
436 **Discussion:**
437
Here we quantified the active visual behavior of a freely-moving primate by developing an
innovative head-mounted eye-tracking system. Our system achieves several technical
innovations that enabled accurate quantification of head and eye movements in a small bodied
(~300-400g) monkey – common marmosets - with the resolution and speed needed to
accurately quantify primate visual behavior in real-world contexts. Specifically, this system
achieves high-speed recording of the eye (90 FPS) and world (63 FPS), as well as applies an
innovative solution to overcoming the challenge that frequent changes in lighting cause for pupil
detection by leveraging a segmentation neural network. As CEREBRO was designed to
seamlessly integrate with wireless neural recording methods, this system is poised to not only
elucidate the dynamics of active visual behaviors but the supporting neural processes. While
findings reported here in freely-moving primates recapitulate the core features of conjugate
gaze movements from prior studies using head-free - but chair-restrained – macaques (33, 34)
and extend those to freely moving primates, analyses also revealed several related
characteristics of visual behavior that have not been observed previously. Perhaps most
notably, we observed that the entropy of the gaze behavior was significantly lower when
animals were freely moving than when head fixed, and that the stability of the visual gaze
improved during locomotion. Overall CEREBRO opens the door to examining active primate
visual processing in the same naturalistic, freely-moving contexts in which the system has been
optimized over evolution, wherein coordinated head-eye movements act to stabilize vision
during natural exploration, critical features that cannot be recapitulated in the absence of
ethological movements.
459
CEREBRO is designed to allow for studying visual behavior coupled with neural responses in a
freely moving animal. This will enable investigation of the neural correlates of active vision. To
that end, we tested our system during neural recordings to characterize orientation tuning and
receptive fields using classical reverse correlation methods during head-fixation. As
demonstrated in Fig. 4, we were able to recover clear receptive fields and orientation tuning of
V1/V2 neurons using eye tracking from CEREBRO in combination with the view from the world
camera. In the future, this system will more broadly enable us to investigate the role of saccade

related modulation in head-fixed vs head-free contexts, and more so, to obtain receptive fields
in freely moving animals. Further, the video streams from our world camera will provide
statistics for how the visual scene and optic flow changes during free motion and help to expand
our understanding of the natural scene statistics and how they are processed in vision.
471
More generally, it will be possible to estimate visual receptive field properties up to the accuracy
of the gaze tracking in freely moving scenarios. Our estimates of gaze accuracy, though less
accurate in the free-moving than head-fixed conditions (fig 3d and 3e), are still roughly
comparable and on median under 1 visual degree. Thus, reconstruction of visual receptive
fields, as was shown for the head-fixed condition using the scene camera (fig 4b), should also
be feasible for free-moving cases given similar flashed visual stimuli. However, a key challenge
during free-motion is that the statistics of the visual input are radically different and instead
controlled by the animal, which on one hand provides a novel and exciting aspect of active
vision but on the other hand pose challenges for fitting receptive field models due to the
spatiotemporal correlations found during natural motion and with real visual scenes. These
issues can be addressed by exploiting computational approaches that are highly data-efficient.
Indeed, the receptive field mapping of V1 neurons in freely moving mice, where visual receptive
fields are much larger than primates, has been achieved with a Generalized Linear Model
(GLM) approach (3). Similar methods could be applied in free-moving marmosets, especially if
the visual areas under study have receptive field sizes larger than gaze tracking error, as in the
peripheral visual field.
488
The contribution of eye and head movements towards conjugate gaze shifts in freely-moving
marmosets were qualitatively similar that of chair restrained but head-free macaques and other
mammals (7, 33, 35–39). Although broadly similar, marmosets did exhibit some quantitative
differences relative to macaques, potentially due to their smaller head-size. For example,
marmoset eye movement velocity and amplitude saturate for much smaller gaze shifts around
10-20 degrees, after which the head contributes to the bulk of the shift in gaze (Fig. 4h),
whereas in the macaque this transition does not occur until one reaches much larger gaze shifts
between 20-40 degrees (33). Moreover, most gaze shifts in macaques under 20 degrees in size
are predominantly driven by shifts in eye position and not the head, while in the marmoset the
same gaze shifts would have significant head-movement components. Only the smallest of
gaze shifts under 5-10 degrees in the marmoset, a range comparable to the movements made
in head-fixed marmosets (*fig 6d*), are dominated by changes in eye position after which the
head would normally make significant contributions. These differences likely reflect an efficiency
tradeoff related to smaller head-size in marmosets (40).
503
Because we recorded marmoset eye-movements in head-fixed and freely-moving conditions,
this novel dataset can both be compared to prior studies and extend our understanding of visual
behavior in this New World primate. As in previous experiments (11), when head-fixed the
oculomotor range is relatively fixed within about 10 visual degrees. But when marmoset is
freely-moving that range increases to roughly 20 degrees (*fig. 7d*), suggesting the limited
oculomotor range when head-fixed represents more of a motor preference than a physical
limitation. In a previous study of chaired but head-free marmosets, a paradigm was used to

evoke large gaze shifts involving up to 180-degree head rotation (12). Although similar large
shifts were rare in the study here, a similar linear relation between peak head velocity and gaze
shifts that peak near a velocity of 750 $^{\circ}$/sec for an 80-degree gaze shifts were evident in freely-
moving marmosets suggesting that this aspect of head-gaze control are relatively invariant to
the form of the task. A direct comparison of the two freely-moving contexts here revealed a
number of notable differences including larger amplitude gaze shifts with higher maximum
velocities when locomoting (*fig. 7, d-g*). By contrast, when stationary and visually scanning the
environment, marmoset biased to smaller gaze shifts indicating that motor demands of actively
moving through space likely drives differences in how the head and eyes coordinate to stabilize
the visual field.
521
Our understanding of NHP vision is almost entirely based on studies in which subjects are
head-restrained. Quantifying primate visual behavior with the innovative eye-tracking system
here afforded the powerful opportunity to examine whether our assumptions about natural vision
are accurate. Indeed, we observed broad qualitative similarities in many features of conjugate
eye movements between head-fixed and freely-moving contexts. However, it was also apparent
that certain assumptions about results in the more conventional paradigm may not be strictly
true. The first being that gaze stability was not worsened despite greater eye and head
movements during locomotion, and in fact, was slightly enhanced. We observed that the
approximate entropy of the marmoset visual system was significantly lower (i.e. improved) when
individuals were freely-moving than head fixed suggesting that the collective visual system is
more consistent and predictable when marmosets are moving and exploring the environment.
Likewise, a comparison of freely-moving marmosets when stationary – i.e. visually scanning -
and locomoting revealed that the stability of visual gaze - as measured by RMS – was
significantly better when animals were locomoting. In other words, despite an increase in the
number, speed and length of the gaze shifts when locomoting, vision actually became more
stable. These findings show that the coordinated movements of head and eyes have been
optimized to accommodate self-motion in a diversity of ethological contexts.
539
The context-dependence of visual stability between the stationary scanning and locomotion
states could reflect changes in VOR at the neural level, as well as other positional strategies
that optimize the VOR system (41, 42). Such context-dependence of gaze control has long been
appreciated from head-mounted eye tracking studies in humans (43, 44), and recently includes
free motion in natural contexts (45). The technical parallels between these eye-tracking systems
in human experiments and CEREBRO allows for comparisons between these primate species,
as well as between marmosets and other taxa. In humans, for example, the gain for angular
VOR (aVOR) is a function of locomotion speed(46, 47). Those findings are consistent with the
RMS stabilization of gaze in marmosets being maintained, or even slightly improved, during
locomotion as compared to when they are stationary and scanning the scene (Fig. 7h). An
increase in VOR gain during locomotion would explain how stability is maintained even though
head and gaze movements increase in frequency and amplitude during locomotion (Fig. 7e,f,g).
Further, we measured the negative correlation between horizontal head and eye velocity during
periods of stable fixation to estimate the VOR gain directly (Supplementary Fig. 8). We find that

VOR gain increases during locomotion as compared to stationary scanning, consistent with prior
human studies and the maintenance of RMS stability during locomotion.
556
Another parallel across humans and marmosets is the change in the relative contribution of
head and eye movements to gaze shifts with increasing gaze amplitude(48). We show that
larger head amplitudes lead to less eye contribution (Fig 6h) wherein the eye amplitude
saturates at ~10 deg and then the putative target is achieved with help of head-movement. In
humans the reliance on head-movements does not typically occur until much larger gaze shifts
are required, but this also can depend critically on the task context, including the speed of
locomotion(46). The natural movement statistics of head-position in free-moving primates,
however, is known to differ substantially from that of rodents (49), suggesting that at least some
mechanisms for visual stabilizations may not be evident across all vertebrates or mammals. In
future work the system could be expanded to address other features of oculomotor control that
could be especially relevant to free-motion in these contexts in primates and other taxa,
including vergence and torsional eye movements (50, 51). The current eye tracking system
offers an opportunity to study how computational constraints and movement strategies during
active vision act to achieve stability and provide high acuity vision in different contexts, including
locomotion and navigation through the environment.
572
From the outset, CEREBRO was explicitly designed to allow for studying visual behavior
coupled with neural responses in freely moving animals. This enables us to investigate the
neural correlates of active vision. To that end, we tested our system on classical visual
correlates like orientation tuning and receptive fields to assess the quality of responses
collected by our system with the conventional methods. As demonstrated in our findings (fig 4),
we were able to obtain clear receptive fields and orientation tuning of V1/V2 neurons using just
the eye and the world camera of our system. However, quantifying the visual information at the
level of the retina and its representation in early visual cortex presents particular analytic
challenges beyond the resolution of the eye-tracking system. In contrast to head-fixed
paradigms in which controlled visual stimuli can be presented to map receptive fields in visual
cortex, the natural environment has highly correlated spatiotemporal features in the visual input,
as well as low contrast in a large number of images (e.g. when the animal looks at the ceiling or
the floor). These issues cause insufficient statistical power in the collected samples and require
more advanced computational approaches to resolve that were beyond the scope of this study.
Fortunately, a recent elegant experiment that addressed this issue in mice (3) provides a
roadmap that we will pursue in future studies Additional fundamental questions will also be
pursued in future experiments such as dissecting the role of visual flow on the neural population
to understand how optical flow aids in separating foreground and background elements as
previously studied in some of the human experiments (52). Indeed, the innovative system
described here eliminates a critical bottleneck for the field that opens the door to studies of
natural active vision in primates that may expand our understanding of the underlying neural
mechanisms that support our preeminent sensory domain.
595
Despite its potential, the innovative eye-tracking system described here is not without its
limitations. For example, like most primates, marmosets are arboreal. This 3D environment

presents significant but distinct challenges to the visual system compared to the 2D
environment tested here. However, it is not clear how this wearable technology may affect their
movement in more precarious environments, and by extension the data and our interpretations
of those data. Moreover, one of the key plans for future research is to leverage this system to
investigate social perception and its underlying neural mechanisms (e.g. face patches) but it is
not yet clear if the hardware on the head would affect how other marmosets perceive and
interact with a test subject, which could have similar biases on the resultant data. Finally, we
have made significant efforts to quantify the precision of the marmoset gaze targets in a freely
moving context, but it is important to recognize that these are estimations. Although the same
can be said of all eye-tracking systems, the inherent complexity and variability of real- world
environments have the potential to introduce more error in our quantification than in more
controlled experimental settings.
610
The sensory and motor systems of animals co-evolved. Amongst the most significant selective
forces acting on animals is to move and interact with the world, and to do so requires sensory
feedback. This ground truth inherently couples sensory and motor systems. The animals'
behavior can be thought of as an outcome filtered by the environment and the animals' potential
actions (53). For a freely moving animal this "affordance landscape" can be very dynamic given
the exteroceptive (sensory feedback), proprioceptive and interoceptive feedback. These parallel
and complementary feedback processes work in synergy to guide relevant behavior of the
animal such as decision-making. It has been argued that perception is not just about creating an
internal representation of the world based on sensory inputs, but instead filters responses
relevant to the environment and animals' internal state (54). Despite this, the study of the
primate visual system has largely ignored considerations of movement and relationships
between the sensory inputs and affordances. Head-restrained experiments limit the
sensorimotor and "affordance landscape" of an animal that potentially bias results in a way that
differs from naturalistic behaviors in which an animal acts as an agent. Freely moving animals
express continuous behaviors based on the elements in the environment, history of previous
actions and internal state which is starkly different from a trial-based structure and lead to many
interesting discoveries about computation strategies in the brain. Leveraging our innovative
head-mounted eye-tracking system for marmosets, we provide compelling evidence that the
coordinated actions of eye and head movements are optimized to stabilize primate vision and
increase its predictability. These patterns emphasize the significance in considering the
ethological relevance of how primates are tested in vision studies. Evidence suggests that
neural responses in head-fixed paradigms are not necessarily predictive of how the same single
neurons – or population ensembles – respond in naturalistic contexts (55–57). Indeed,
investigations of vision in freely-moving mice suggest that at least some elements of neural
activity in this context are distinct (3, 58). Our innovative eye-tracking system can be leveraged
in marmosets to precisely examine the primate visual system during natural, freely-moving
behaviors (5, 59) and address a suite of foundational questions that we have as of yet been
unable to investigate with sufficient quantitative rigor.
639
640
641

Acknowledgements:

650
651
652 We thank Alex Huk and Cris Niell for their invaluable insights throughout the development of this
653 system. We thank Brian Corneil for his valuable comments on an earlier version of the
654 manuscript. This work was supported by grants from the NIH/BRAIN Initiative (R01 NS118457 &
655 U01 NS116377) and AFOSR (FA9550-19-1-0357) to CTM and by a Kavli Institute for Brain and
656 Mind Postdoctoral Award to JL.
657

Author contributions:

659 Conceptualization: VPS, JL, JM, and CM; Methodology: VPS; Software: VPS; Formal analysis:
660 VPS and JL; Investigation: VPS and JL; Resources: VPS, JL, CM; Writing: VPS, JL, JM, CM;
661 Visualization: VPS, JL, JM, CM; Supervision: JM, CM; Funding acquisition: CM
662
663

Declaration of Interest:

665 VPS is inventor on provisional patent application no: US 20230404467 A1 filed by the Regents
666 of the University of California entitled "Head Mounted Camera and Eye Track System for
667 Animals."

 RESOURCE AVAILABILITY:

Lead Contact:
Further information and requests for resources and reagents should be directed to and will be
fulfilled by the lead contact, Cory Miller (corymiller@ucsd.edu).

Materials availability:
The STL designs for the headpiece and its material description can be downloaded at
https://github.com/Vickey17/CEREBRO_HeadPiece.

Data and code availability:
- Head, eye and body locomotion data has been deposited at https://doi.org/10.5061/dryad.8gtht76xb
- The GUI for UNet segmentation-based pupil detection can be found at https://github.com/Vickey17/UNET_implementation_V2.

Materials & Methods:

1. Experimental model and study participant details:

The development and testing of CEREBRO were performed on two common marmosets
(monkey M, male; monkey S, female). Both subjects were group housed in home cages (76 cm
x 76 cm x 81 cm) and were 2 years old at the time of implant. Monkey M had a chronic implant
in left V1 and monkey S had bilateral chronic implants in both left and right V1. All surgeries and
experiments were performed in the Cortical Systems and Behavior Laboratory at University of
California, San Diego (UCSD), and approved by the UCSD Institutional Animal Care and Use
Committee in accordance with National Institute of Health standards for care and use of
laboratory animals.

700
701

2. Fitting the animal with CEREBRO:

The head assembly for CEREBRO is custom fitted to every animal. For the initial fitting, the
animal is anesthetized using a combination of Ketamine (dose:20 mg/kg) and Acepromazine
(dose: 0.75 mg/kg). All the parts (*Supplementary fig 1b*) are adjusted for best view of the eye
and the front camera and fastened using set screws.

708
709

3. Habituation of animals to backpack:

Subjects were habituated to carry backpacks for 1-3 months. This was done in habituation

sessions by placing the backpack on an individual and incrementally increasing weight every week by 5 gm increments up to 50 gm. The length of these sessions was increased in parallel until the animals did not exhibit any duress (i.e. pulling or biting the straps) while wearing the backpack. Significant Individual differences were evident in the duration of this habituation period, but all ultimately were habituated to wearing the backpack. While habituation to the backpack was necessary, the marmosets were minimally affected by the head-piece. In other words, marmosets rarely grabbed the hot mirrors used to track pupils. The only instance when this happened was a single occasion when the IR LED wire accidently slipped and was in a monkey's peripheral view, and a separate occasion when an animal was chaired and a few drops of the liquid reward accidentally spilled on the hot mirror.

4. Multiple system synchronization:

Our Neural recording systems, motion trackers and CEREBRO were all synchronized using an ESP32 microcontroller based custom board. CEREBRO and the motion tracking system sent TTL signals for different mode shifts that were recorded by the synchronization microcontroller. For synchronizing the neural signal, we sent a TTL pulse (100 mV |100 m) to the electrophysiology system (in recording mode) before putting it on the animal. The timestamp of this signal was logged by the sync microcontroller and since this signal was also logged into the real data stream of the electrophysiology system, we can easily align the time of this pulse in the data to other systems (CEREBRO and motion tracking). The board was connected to a computer via serial communication and the timing data was logged on to a console terminal (CoolTerm).

5. Electrophysiology:

Neural activity was recorded with multi-electrode arrays (N-form, Modular Bionics, Berkeley CA) chronically implanted in V1. The N-form array has 64 channels in total embedded in 16 shanks located in a 4x4 grids evenly spaced 0.25 mm apart. Each shank has 4 iridium oxide electrodes located at 0.5, 0.375, 0.25, and 0 mm from the tip. Implantations followed the standard surgical procedures for chronically implanted arrays in primates. In each recording session, a wireless Neurologger (SpikeLog-64, Deuteron Technologies) was hosted in a 3.5 cm (width) x 2.5 cm (height) x 1.2 cm (depth) protecting case and connected to the array to record the extracellular voltage at 32 kHz. Spike sorting was performed offline using Kilo sort (60)and manually curated using the graphic user interface Phy.

6. Receptive field and tuning property of V1 neurons:

For the receptive field mapping, marmosets were head-fixed and freely viewed the stimulus of 120 randomly positioned black and white flashing dots generated at a frequency of 10 Hz for 4 minutes. Eye calibration was performed offline with the GUI provided in this paper. We then used the calibrated eye position on the world camera images to estimate the visual input on the retina, and mapped the receptive fields using forward correction (61). For the orientation and

spatial frequency tuning of the V1 neurons, we presented the full-field drifting gratings with 12
orientations (evenly spaced 30-degree intervals) and 3 spatial frequencies (0.5, 1, 2
cycles/degree). This information was used in calculating the tuning curves with the information
from the screen; when calculating the tuning curves with the information from CEREBRO, we
applied Hough transform on the world camera images to detect the lines and further calculated
the orientation and spatial frequency of the gratings.
764
765 7. Head and body movement tracking using OptiTrack:
766
The head and body movement of the animals were recorded using OptiTrack- an image-based
motion tracking system (www.optitrack.com). For the head movements, the added 3 IR
reflective beads (12mm in diameter) on the head-assembly of the animal. For the body, we
added a single IR bead on to the backpack. The arena was fitted with 10 OptiTrack cameras
strategically placed to cover every view of the arena. The cameras were calibrated before each
session and sessions were performed only for calibrations where the error for the worst camera
was less than 0.1 mm. The data collected from the session was manually curated to remove
stray/false markers and any gaps were filled with linear interpolation.
775
776 8. Locomotion behavior analysis:
777
Locomotion is detected with body and head tracking data from OptiTrack. We first calculated the
velocity of the body IR bead and set a threshold of 5 cm/s (i.e. moving more than 20-30% the
typical marmoset body length in under one second) after applying a 0.2 Hz low pass filter to
select the period when the monkey had a fast body movement. However, not all the fast body
movement came from locomotion; it could happen when the monkey made a body turn, or a
sudden change of posture. We add two other conditions: 1) the duration of the period is larger
than 3s; 2) the position of the head is lower than 18 cm. These two conditions are determined
by manually matching the data with the video. After the locomotion period is detected, the rest
of the time is defined as sedentary.
787
788 9. Freely moving eye calibration setup:
789
Freely moving eye calibration paradigm consisted of a clear-plexiglass box (dimension: 30 cm x
30 cm x 40 cm). All sides of the box except the front panel were painted with black acrylic paint
rendering it opaque. The front of the box was covered with IR long pass filter (OPTIR 1.0 NG
305 X 100, Newark) with a 10 cm x 10 cm window cut out in the middle at the height of 24 cm
from the floor. The inside of the box had a wooden perch for subjects to sit on while looking out
the window. The top of the box was lined with IR led strips (Near Infrared 850 nm NIR Single
Chip DC12V 24W, 360DigitalSignage). The IR LEDs enabled video recording of subject's
behavior in the box since the IR dichroic allows the IR to pass through. Two synchronized FLIR
cameras were mounted on the box. One camera was positioned to continuously monitor the
subject's behavior and the second camera was positioned to record the projection screen. A 2-
axis galvanometric mirror driven LASER system (20Kpps speed Laser Galvo laser set,
Wonsung Store) was installed on top of the box. This was used to draw various geometric

shapes on the screen. All the shapes were generated in Python as plots with no axis. These
plots were saved as SVG files and converted into ILD files using LASEROS software. The ILD
files were loaded onto the galvanometric system using an SD card.
805
806 10. Error estimation for eye in world calibration:
807
Eye/gaze data were calibrated (transformed to match the world scene) for a select number of
trials. Following the initial process, the same transformations were applied to remaining trials.
The eye/gaze position for these trials was overlaid on the world scene and exported as a video
file. This video file was analyzed using a custom python script. Instances when the eye position
was within 3.5 degrees of a visual feature and the residence time for the eye position for that
location exceeded 60 ms were considered valid instances of looking at the feature. For such
valid instances, the closest distance (radius of the smallest circle that touches the visual feature
with eye position as the center) between the eye position and the closest visual feature (in
degrees) was used as the estimate of the error associated with our eye calibration. This places
an upper bound on the error as the animal's actual eye position would not exactly match the
center of the object even when viewing it due to the object's extent, as well as fixational eye
movements and drift.

11. Camera and communication interface:

OV4689 camera modules with two MIPI lanes and customized Flex cable length of 15 cm
(SincereFirst, Guangzhou, China) was used both as the eye camera and the world camera. The
cameras operated on a serial interface called MIPI CSI-2 (Mobile Industry Processor Interface
Camera Serial Interface). Since microprocessors are the core of the system a STMIPID02 MIPI
CSI-2 deserializer was used to deserialize CSI communication from a MIPI CSI-2 camera
sensor to DCMI and improve the temporal resolution of the incoming data.

12. Printed Circuit Board and data storage:
828
Our custom PCBs use STM32H750 which is a 32-bit Arm Cortex-M7 core processor running up
to 480 MHz. It is better suited for interfacing with camera sensors since it embeds a digital
camera interface (DCMI). Each PCB can be programmed and debugged using the Serial Wire
Debug (SWD) mode. The PCBs store the raw camera data on a locally mounted, UHS-1 (Ultra
High Speed) class, SD card. The binary files are decoded offline to get an AVI format video of
the stream and a CSV file with timestamp and IMU data. Two PCBs are stacked on top of each
other and are enclosed in a custom designed 3D printed casing with a custom designed back
harness . To measure camera synchronized movements (acceleration and orientation) of head,
the boards interface with an inertial measurement unit (IMU). The PCB boards also feature a
0.96" SPI TFT display which allows the users to preview the captured camera image and ensure
the placement of the head-piece at the start of the session.
840
841

References:

1. D. J. Felleman, D. C. Van Essen, Distributed hierarchical processing in the primate cerebral
cortex. *Cereb. Cortex* **1**, 1–47 (1991).

2. A. J. Parker, W. T. Newsome, Sense and the single neuron: probing the physiology of
perception. *Annu. Rev. Neurosci.* **21**, 227–277 (1998).

3. P. R. L. Parker, E. T. T. Abe, E. S. P. Leonard, D. M. Martins, C. M. Niell, Joint coding of
visual input and eye/head position in V1 of freely moving mice. *Neuron* **110**, 3897-3906.e5
(2022).

4. C. T. Miller, *et al.*, Natural behavior is the language of the brain. *Curr. Biol.* **32**, R482–R493
(2022).

5. V. Ngo, *et al.*, Active vision during prey capture in wild marmoset monkeys. *Curr. Biol.* **32**,
3423-3428.e3 (2022).

6. D. Guitton, Control of eye-head coordination during orienting gaze shifts. *Trends Neurosci.*
15, 174–179 (1992).

7. J. C. Martinez-Trujillo, H. Wang, J. D. Crawford, Electrical stimulation of the supplementary
eye fields in the head-free macaque evokes kinematically normal gaze shifts. *J.*
Neurophysiol. **89**, 2961–2974 (2003).

8. D. E. Angelaki, B. J. M. Hess, Self-motion-induced eye movements: effects on visual acuity
and navigation. *Nat. Rev. Neurosci.* **6**, 966–976 (2005).

9. D. E. Angelaki, K. E. Cullen, Vestibular system: the many facets of a multimodal sense.
Annu. Rev. Neurosci. **31**, 125–150 (2008).

10. R. J. Skyberg, C. M. Niell, Natural visual behavior and active sensing in the mouse. *Curr.*
Opin. Neurobiol. **86**, 102882 (2024).

11. J. F. Mitchell, J. H. Reynolds, C. T. Miller, Active vision in marmosets: a model system for
visual neuroscience. *J. Neurosci.* **34**, 1183–1194 (2014).

12. S. Pandey, S. Simhadri, Y. Zhou, Rapid Head Movements in Common Marmoset Monkeys.
iScience **23**, 100837 (2020).

13. S. V. Shepherd, M. L. Platt, Noninvasive telemetric gaze tracking in freely moving socially
housed prosimian primates. *Methods* **38**, 185–194 (2006).

14. R. Milton, N. Shahidi, V. Dragoi, Dynamic states of population activity in prefrontal cortical
networks of freely-moving macaque. *Nat. Commun.* **11**, 1–10 (2020).

15. D. Mao, *et al.*, Spatial modulation of hippocampal activity in freely moving macaques.
Neuron **109**, 3521-3534.e6 (2021).

16. A. M. Michaiel, E. T. Abe, C. M. Niell, Dynamics of gaze control during prey capture in
freely moving mice. *Elife* **9** (2020).

17. A. F. Meyer, J. Poort, J. O'Keefe, M. Sahani, J. F. Linden, A Head-Mounted Camera
System Integrates Detailed Behavioral Monitoring with Multichannel Electrophysiology in
Freely Moving Mice. *Neuron* **100**, 46-60.e7 (2018).

18. D. J. Wallace, *et al.*, Rats maintain an overhead binocular field at the expense of constant
fusion. *Nature* **498**, 65–69 (2013).

19. A. F. Meyer, J. O'Keefe, J. Poort, Two Distinct Types of Eye-Head Coupling in Freely
Moving Mice. *Curr. Biol.* **30**, 2116-2130.e6 (2020).

20. O. Ronneberger, P. Fischer, T. Brox, U-Net: Convolutional Networks for Biomedical Image
Segmentation. *arXiv [cs.CV]* (2015).

21. A. K. Chaudhary, *et al.*, RITnet: Real-time semantic segmentation of the eye for gaze
tracking. *arXiv [cs.CV]* (2019).

22. R. C. Emerson, M. C. Citron, W. J. Vaughn, S. A. Klein, Nonlinear directionally selective
subunits in complex cells of cat striate cortex. *J. Neurophysiol.* **58**, 33–65 (1987).

23. G. C. DeAngelis, I. Ohzawa, R. D. Freeman, Spatiotemporal organization of simple-cell
receptive fields in the cat's striate cortex. I. General characteristics and postnatal
development. *J. Neurophysiol.* **69**, 1091–1117 (1993).

24. D. L. Ringach, G. Sapiro, R. Shapley, A subspace reverse-correlation technique for the
study of visual neurons. *Vision Res.* **37**, 2455–2464 (1997).

25. M. S. Livingstone, C. C. Pack, R. T. Born, Two-dimensional substructure of MT receptive
fields. *Neuron* **30**, 781–793 (2001).

26. J. L. Yates, *et al.*, Detailed characterization of neural selectivity in free viewing primates.
Nat. Commun. **14**, 3656 (2023).

27. J. Wallman, J. Velez, B. Weinstein, A. E. Green, Avian vestibuloocular reflex: adaptive
plasticity and developmental changes. *J. Neurophysiol.* **48**, 952–967 (1982).

28. K. E. Cullen, The neural encoding of self-motion. *Curr. Opin. Neurobiol.* **21**, 587–595
(2011).

29. K. E. Cullen, The vestibular system: multimodal integration and encoding of self-motion for
motor control. *Trends Neurosci.* **35**, 185–196 (2012).

30. N. Takeichi, *et al.*, Directional asymmetry in smooth ocular tracking in the presence of
visual background in young and adult primates. *Exp. Brain Res.* **149**, 380–390 (2003).

31. R. E. Kettner, H. C. Leung, B. W. Peterson, Predictive smooth pursuit of complex two-
dimensional trajectories in monkey: component interactions. *Exp. Brain Res.* **108**, 221–235
(1996).

32. A. Delgado-Bonal, A. Marshak, Approximate Entropy and Sample Entropy: A
Comprehensive Tutorial. *Entropy* **21** (2019).

33. E. G. Freedman, D. L. Sparks, Eye-head coordination during head-unrestrained gaze shifts
in rhesus monkeys. *J. Neurophysiol.* **77**, 2328–2348 (1997).

34. H. K. Arora, *et al.*, Eye-head-hand coordination during visually guided reaches in head-
unrestrained macaques. *J. Neurophysiol.* **122**, 1946–1961 (2019).

35. P. Morasso, E. Bizzi, J. Dichgans, Adjustment of saccade characteristics during head
movements. *Exp. Brain Res.* **16**, 492–500 (1973).

36. R. D. Tomlinson, P. S. Bahra, Combined eye-head gaze shifts in the primate. I. Metrics. *J.*
Neurophysiol. **56**, 1542–1557 (1986).

37. D. Guitton, R. M. Douglas, M. Volle, Eye-head coordination in cats. *J. Neurophysiol.* **52**,
1030–1050 (1984).

38. H. Collewijn, Eye- and head movements in freely moving rabbits. *J. Physiol.* **266**, 471–498
(1977).

39. H. H. Goossens, A. J. Van Opstal, Human eye-head coordination in two dimensions under
different sensorimotor conditions. *Exp. Brain Res.* **114**, 542–560 (1997).

40. M. D. Malinzak, R. F. Kay, T. E. Hullar, Locomotor head movements and semicircular canal
morphology in primates. *Proc. Natl. Acad. Sci. U. S. A.* **109**, 17914–17919 (2012).

41. P. P. Vidal, W. Graf, A. Berthoz, The orientation of the cervical vertebral column in
unrestrained awake animals. I. Resting position. *Exp. Brain Res.* **61**, 549–559 (1986).

42. W. Graf, C. de Waele, P. P. Vidal, D. H. Wang, C. Evinger, The orientation of the cervical
vertebral column in unrestrained awake animals. II. Movement strategies. *Brain Behav.*
Evol. **45**, 209–231 (1995).

43. M. F. Land, D. N. Lee, Where we look when we steer. *Nature* **369**, 742–744 (1994).

44. M. Hayhoe, D. Ballard, Modeling task control of eye movements. *Curr. Biol.* **24**, R622-8
(2014).

45. J. S. Matthis, J. L. Yates, M. M. Hayhoe, Gaze and the Control of Foot Placement When
Walking in Natural Terrain. *Curr. Biol.* **28**, 1224-1233.e5 (2018).

46. H. Dietrich, M. Wuehr, Strategies for gaze stabilization critically depend on locomotor
speed. *Neuroscience* **408**, 418–429 (2019).

47. S. E. Haggerty, W. M. King, The interaction of pre-programmed eye movements with the
Vestibulo-Ocular Reflex. *Front. Syst. Neurosci.* **12**, 4 (2018).

48. D. Guitton, M. Volle, Gaze control in humans: eye-head coordination during orienting
movements to targets within and beyond the oculomotor range. *J. Neurophysiol.* **58**, 427–
459 (1987).

49. J. Carriot, M. Jamali, M. J. Chacron, K. E. Cullen, Statistics of the vestibular input
experienced during natural self-motion: implications for neural processing. *J. Neurosci.* **34**,
8347–8357 (2014).

50. D. Tweed, T. Haslwanter, M. Fetter, Optimizing gaze control in three dimensions. *Science*
281, 1363–1366 (1998).

51. J. K. Y. Ong, T. Haslwanter, Measuring torsional eye movements by tracking stable iris
features. *J. Neurosci. Methods* **192**, 261–267 (2010).

52. P. R. MacNeilage, Z. Zhang, G. C. DeAngelis, D. E. Angelaki, Vestibular facilitation of optic
flow parsing. *PLoS One* **7**, e40264 (2012).

53. G. Pezzulo, P. Cisek, Navigating the affordance landscape: Feedback control as a process
model of behavior and cognition. *Trends Cogn. Sci.* **20**, 414–424 (2016).

54. P. Cisek, J. F. Kalaska, Neural mechanisms for interacting with a world full of action
choices. *Annu. Rev. Neurosci.* **33**, 269–298 (2010).

55. V. Jovanovic, A. R. Fishbein, L. de la Mothe, K.-F. Lee, C. T. Miller, Behavioral context
affects social signal representations within single primate prefrontal cortex neurons. *Neuron*
110, 1318-1326.e4 (2022).

56. D. B. T. McMahon, B. E. Russ, H. D. Elnaiem, A. I. Kurnikova, D. A. Leopold, Single-unit
activity during natural vision: diversity, consistency, and spatial sensitivity among AF face
patch neurons. *J. Neurosci.* **35**, 5537–5548 (2015).

57. M. C. Rose, B. Styr, T. A. Schmid, J. E. Elie, M. M. Yartsev, Cortical representation of
group social communication in bats. *Science* **374**, eaba9584 (2021).

58. T. A. Chaplin, T. W. Margrie, Cortical circuits for integration of self-motion and visual-motion
signals. *Curr. Opin. Neurobiol.* **60**, 122–128 (2020).

59. L. Shaw, K. H. Wang, J. Mitchell, Fast prediction in marmoset reach-to-grasp movements
for dynamic prey. *Curr. Biol.* **33**, 2557-2565.e4 (2023).

60. M. Pachitariu, N. Steinmetz, S. Kadir, M. Carandini, H. Kenneth D., Kilosort: realtime spike-
sorting for extracellular electrophysiology with hundreds of channels. *bioRxiv* (2016).

61. Y. Zhao, J. L. Yates, A. J. Levi, A. C. Huk, I. M. Park, Stimulus-choice (mis)alignment in
primate area MT. *PLoS Comput. Biol.* **16**, e1007614 (2020).

Figures and legends:

Fig1: Head mounted eye-tracker assembly

(a) (a) 3D render depicting different parts of the head-piece. The Ti- alloy scaffolding is custom
designed and fabricated using DLS **(b)** 3D printed backpack that holds the 2 PCBs and the
battery. **(c)** The animals are fitted with two 1cm headposts that act as anchors for holding the
head-piece. **(d)** Custom PCB board which drives the camera sensors, IMU and saves the data
to a locally mounted SD card**. (e)** A marmoset wearing the fully assembled system (CEREBRO).
(f) View from the 3 different cameras (eye camera(pink), world camera(yellow) and the external
arena camera (cyan)) during a freely-moving session. **(f)** Animals wearing CEREBRO exhibited
no impairment in their free- moving characteristics such as percent movement, distance covered
and speed.

1002
Fig2: Iris/Pupil detection using segmentation Artificial Neural Network.
(a) Conventional thresholding approach for pupil detection fails in scenes with variable light
intensities. **(b)** Architecture of the segmentation neural network to crop out the iris. **(c)** The
performance of the ANN matches closely to that of human annotation with relatively low training
epochs (10 in this example) and relatively small amount of training data (~250 samples) **(d)** For
calibrating the eye and world camera, the animals are presented with marmoset faces on a 120
Hz LCD monitor in a predetermined layout. **(e)** The calibrated eye movements can be overlaid
on the world scene to determine the visual scene falling on the retina of the animal. (f) An
illustration of the error estimation algorithm. **(g)** Accuracy comparison between Arrington eye
tracker system and CEREBRO.
1013

1014

Fig3.: Eye/gaze accuracy in various experimental setups using CEREBRO.

(a) Schematic for the experimental setup used to test accuracy of eye calibration in unrestrained
animal. The setup consists of a dark box with a perch. The front of the box is covered with an IR
longapss filter that doesn't allow visible light to pass except for a small window shown with red
dashed line in the inset image. A galvanometric LASER system draws different geometric
shapes on a screen for the animal to observe. **(b)** Schematic illustrating error estimation
algorithm for freely moving eye calibration. **(c)** Maximum intensity projection of stimulus and eye
in world position (red markers) for saccades and smooth pursuit trials in freely moving and
head-restrained setup. **(d)** Accuracy comparison for eye calibration error for head restrained
and freely moving animals during saccade trials and **(e)** smooth pursuit trails.

1025

1026

1027

1028

1029

1030

1031

1032

Fig4: Eye tracking coupled electrophysiology in freely moving marmosets.

(a) To validate use of CEREBRO with electrophysiology, subjects were presented with orientation grating and receptive field stimuli in a classical head fixed preparation. **(b)** The receptive field (top) and orientation tuning curve (bottom) is shown for three representative neurons recorded in marmoset V1 while using CEREBRO. **(c)** Example raster of 30+ single neurons in marmoset V1 while subjects are head-restrained (left) and freely-moving (right) while wearing CEREBRO. The head (blue) and body (purple) speed of the marmoset using CEREBRO is plotted below the freely-moving raster **(d)** Example spike waveforms from five exemplar neurons demonstrate that we were able to stably record from same neurons in head-restraint vs freely moving conditions.

Fig5.: Characteristics of eye/gaze behavior in various experimental setups using CEREBRO. **(a)** Heatmap of eye position in a restrained marmoset with dashed lines marking 2 SD mark. **(b & c)** distribution of horizontal and vertical eye position split by fixation (purple) vs saccade (green) with corresponding dashed lines representing 2-SD of data variation. **(d)** Heatmap of eye position in an unrestrained marmoset. **(e & f)** distribution of horizontal and vertical eye position split by gaze fixation(purple) and gaze shifts (green). Color-matched dashed lines represent 2-SD of variation in data.

1064

Fig6.: Gaze characteristics in freely moving marmosets.

(a) Representative session of activity pattern for marmosets in an open arena (200 cm x 100 cm x 240 cm). Grey dotted lines indicate periods of locomotion, while heat maps plot stationary occupancy. **(b)** Plots the estimate of the animal's gaze (blue) with CEREBRO. Horizontal eye movement (red) and head yaw (green) are also plotted. Inset magnifies a 2s period of time. **(c)** Horizontal eye movement and head yaw in freely-moving marmosets are shown. Green dots indicate gaze fixations, while purple dots indicate gaze shifts. **(d-h):** eye-movements (red), head-movements (green) and gaze-shifts (red) in freely-moving marmosets. **(d)** Plots the position (top) and velocity (bottom) of eye-movements, head-movements and gaze-shifts <10degrees (left) and >10degrees (right) **(e)** Shows the maximum speed and gaze amplitude. Plots the latency to max speed (f), duration (g) and amplitude (h) as a function of gaze amplitude

Fig7.: Characteristics of horizontal eye movements in freely moving marmosets.

(a) An illustration of the gaze shifts in a freely behaving marmoset in two different states of
locomotion(left) and stationary(right). (b) Animals exhibited noticeably different eye behavior in
various contexts i.e. chaired, locomotion and stationary. **(c)** Horz. eye movements have
significantly less "approximate entropy" in head-fixed context as compared to freely moving.
Although horz. gaze has the least approximate entropy of all the contexts. This shows that a
freely moving gaze is computationally more regular than a head-fixed context. (d) Freely moving
animal exhibits larger range of horz. eye amplitude and the speed of horz. eye movements is
higher during locomotion than stationary state. Marmosets make more gaze shifts (e) during
locomotion with higher speed (f) and larger amplitudes (g). (h) RMS stabilization of head(yaw)
and horz. gaze shows that despite poorer stability because of head movements, gaze remains
remarkably stable during locomotion and stationary epochs.
1093