Title: MLT-11 is a transient apical extracellular matrix component required for cuticle patterning and function

James Matthew Ragle[1], Ariela Turzo[1], Max T. Levenson[1], Keya Jonnalagadda[1], Anton Jackson[1], An A. Vo[1], Vivian T. Pham[1] and Jordan D. Ward[1‡].

[1]Department of Molecular, Cell, and Developmental Biology, University of California-Santa Cruz, Santa Cruz, CA 95064, USA.

‡Author for correspondence.

Corresponding author email: jward2@ucsc.edu

Keywords: *C. elegans*, molting, Kunitz domain, apical extracellular matrix, protease inhibitor

Running Title: MLT-11 and *C. elegans* development

ABSTRACT

Apical extracellular matrices (aECMs) are associated with all epithelia and form a protective layer against biotic and abiotic threats in the environment. Despite their importance, we lack a deep understanding of their structure and dynamics in development and disease. *C. elegans* molting offers a powerful entry point to understanding developmentally programmed aECM remodeling. A transient matrix is formed in embryos and at the end of each larval stage, presumably to pattern the new cuticle. Focusing on targets of NHR-23, a key transcription factor which drives molting, we identified the Kunitz family protease inhibitor gene *mlt-11* as an NHR-23 target. We identified NHR-23-binding sites that are necessary and sufficient for epithelial expression. *mlt-11* is necessary to pattern every layer of the adult cuticle, suggesting a broad patterning role prior to the formation of the mature cuticle. MLT-11::mNeonGreen::3xFLAG transiently localized to the aECM in the cuticle and embryo. It was also detected in lining openings to the exterior (vulva, rectum, mouth). Reduction of *mlt-11* function disrupted the barrier function of the cuticle. Tissue-specific RNAi suggested *mlt-11* activity is primarily necessary in seam cells and we observed alae and seam cell fusion defects upon *mlt-11* inactivation. Predicted *mlt-11* null mutations caused fully penetrant embryonic lethality and elongation defects suggesting *mlt-11* also plays an important role in patterning the embryonic sheath. Finally, we found that *mlt-11* inactivation suppressed the blistered cuticle phenotype of mutants of *bli-4* mutants, a subtilisin protease gene but did not affect BLI-4::sfGFP expression. These data could suggest that MLT-11 may be necessary to assure proper levels of BLI-4 activity.

INTRODUCTION

Specialized extracellular matrices cover the apical surface of all epithelial cells and form the skin in almost all animals [1–3]. These apical extracellular matrices (aECMs) also line the lumen of internal tubular epithelia to form a protective layer against biotic and abiotic threats [1–3]. Despite their importance, understanding the structure and dynamics of aECM components in development and disease remains challenging.

C. elegans is emerging as a powerful model to study aECM structure and remodeling.

They have a collagen-based aECM (cuticle) so understanding its assembly may provide

insight into mammalian skin biology [4]. The components of the cuticle are secreted by

hypodermal and seam cells and are assembled in distinct layers [4–6]. The cuticle

secreted by the hypodermal syncytium form circumferential ridges called annuli separated

from one another by furrows [4]. In adults an inner basal layer contains two fibrous sub-

layers angled in opposite directions and an outer cortical layer [5]. A fluid-filled medial

layer contains hollow, nano-scale struts built from the BLI-1 and BLI-2 collagens [5,7]. All

layers are composed of extensively cross-linked collagens. The cortical layer also

contains cuticlins, proteins which remain in the insoluble fraction after cuticle solubilization

[8]. The cortical layer is covered by the epicuticle, a poorly understood structure that is

thought to be a lipid bilayer covered by a glycoprotein rich surface coat [9–11]. There can

be stage-specific variations in cuticle structure. Only adults have a medial layer, and both

L1 and adult cuticles contain alae, lateral longitudinal ridges secreted by seam cells in the

epidermis [5,12].

Fig. 1. The *C. elegans* adult cuticle has multiple layers with unique structures and collagen content. Cartoon model of the *C. elegans* adult cuticle sublayers (epicuticle, cortical, medial and basal), topological features (annuli and furrows). The localization of proteins used to mark specific layers of the cuticle in this study is indicated (CUT-2, BLI-1, ROL-6 and COL-19)[7,8,49].

During each larval stage animals must build a new aECM underneath the old one and separate the old aECM (apolysis) that is subsequently shed (ecdysis) [13]. A specialized, transient structure known as the pre-cuticle is thought to pattern the new cuticle and is then endocytosed [14]. Pre-cuticle components include zona

pellucida proteins, lipocalins, fibrillin, extracellular leucine-rich repeat proteins, and

hedgehog-related proteins [14–19]. The sheath is a similar structure in embryos which

ensures embryonic membrane integrity and directs force during elongation [20–23]. The

vulval aECM has recently been shown to be highly dynamic, and specialized aECMs also

line the rectum and excretory system [15,16,24].

A major question is how is the new aECM constructed and the old one shed during
molting? A poorly understood genetic oscillator is thought to coordinate the expression of
cuticle components and processing enzymes [25,26]. The molting cycle must also be
tightly linked to developmental progression with homologs of circadian rhythm proteins
(LIN-42/PER, NHR-23/ROR) playing important roles in both the molting and
developmental timers [27–36]. Study of NHR-23-regulated genes revealed an enrichment
in predicted protease and protease inhibitors [32,33]. Proteases are required for molting
or ecdysis in both *C. elegans* and parasitic nematodes, presumably by promoting
apolysis, though some are thought to function in collagen processing [37–44]. Protease
inhibitors have been implicated in molting through RNAi screening, and have been
suggested to suppress ecdysis [13,44,45]. MLT-11 is a predicted protease inhibitor in the
Kunitz family and *mlt-11 (RNAi)* causes molting defects [44]. *mlt-11* mRNA oscillates,
peaking mid-molt and its expression is regulated by NHR-23, yet *mlt-11* remains poorly
characterized [25,26,31–35,44].

Here we show that MLT-11 is secreted and transiently localizes to the aECM over the
hypodermis and seam cells, before being endocytosed. It also localizes to the rectal
epithelium, excretory duct lumen, and vulval lumen. Reduction of *mlt-11* function in larvae
causes developmental delay, failure to apolyse and a loss of the cuticle barrier function.
Additionally, cuticle collagens in each layer of the mature adult cuticle are aberrantly
localized suggesting *mlt-11* has a broad role in patterning the developing cuticle. *mlt-11*
activity in seam cells is necessary for molting and a normal developmental rate. Like
embryonic sheath components, it is secreted into the extracellular space and surrounds
the exterior of the embryo before being endocytosed prior to hatching. *mlt-11* null alleles
cause embryonic lethality, characterized by disorganization of adherens junctions. *mlt-*
11 inactivation suppresses the blistered cuticle phenotypes in mutants of the
subtilisin/kexin (PCSK) family protease, *bli-4*. Together, this work provides the first insight
into how MLT-11 functions to promote embryogenesis and aECM integrity during molting.
Our identification of BLI-4 as a potential MLT-11 substrate is an entry point into
understanding mechanistically how MLT-11 promotes molting.

RESULTS

Multiple NHR-23 occupied genomic regions regulate *mlt-11* expression

Our previous study found an enrichment in protease and protease inhibitors in genes

regulated by the NHR-23 nuclear hormone receptor transcription factor [32]. *mlt-11* is a

poorly characterized predicted protease inhibitor that is *nhr-23*-regulated with four NHR-

23 ChIP-seq peaks upstream of its transcriptional start site [32,33,44,46]. Consistent with

potential direct NHR-23 binding to these regions, there are GGGTCA or AGGTCA half

sites that NHR-23 binds *in vitro* (Fig. 2A)[35]. To determine the contribution of these NHR-

23 occupied regions to *mlt-11* expression we turned to promoter reporters. We first

created a single-copy *mlt-11p::NLS::mNeonGreen* promoter reporter including the

sequences of peaks 1-3 (-2.8 kb *mlt-11p*) (Fig. 2A). Similar to a previous promoter

reporter study [44], we observed mNeonGreen expression in hypodermal and seam cells,

as well as novel expression in the developing rectal and vulval epithelial cells, and

developing embryo (Fig. 2B,C). Expression in embryos was first detected at the bean

stage in posterior epithelial cells and persisted through the 3-fold stage and expression

spread anteriorly (Fig. 2C). The previously characterized *mlt-11* promoter reporter

contained the the sequence of peak 3 and part of peak 4 [44], but lacked the sequences

from peaks 1 and 2 . We therefore generated an equivalent promoter reporter to Frand et

al. (2005) as well as a larger promoter fragment (-5.2 kb *mlt-11p*) containing the

sequences from peaks 1-4. Similar expression timing and intensity was seen in all three

reporters (Fig. 2D).

Given that the Frand promoter and our -2.8 kb promoter overlapped in the region

containing the NHR-23 peak 3 sequences, we hypothesized that this region may play a

key role in regulating *mlt-11* expression. We therefore tested whether the sequence

contained in each NHR-23 ChIP-seq peak, as well as two candidate enhancers identified

by ATAC-seq were sufficient to drive the expression of a reporter containing a *pes-10*

minimal promoter. This reporter displays undetectable mNG expression in the absence

of an additional *cis*-regulatory element (Fig. 2E). We detected reporter expression

containing peaks 3 and 4 in hypodermal cells with peak 3 being the most robust (Fig. 2E).

Additionally, we detected expression in seam cells with the regulatory element from peaks

Fig. 2. *mlt-11* **is expressed in larval and embryonic epidermal cells.** (A) Genome browser track of *mlt-11* promoter depicting NHR-23 ChIP-seq peaks; relative genomic locations of NHR-23 binding motifs; relative genomic locations of *mlt-11* promoter reporters used in this study; relative genomic locations of candidate *cis*-regulatory elements from the *mlt-11* promoter inserted into a *pes-10* minimal promoter in this study; conservation calculated across 26 nematode species. (B) Expression pattern of a *-2.8kb mlt-11p::mNG::tbb-2 3'UTR* promoter reporter in hypodermal cells of L4 worms. White arrows indicate seam cells (top), rectal epithelial cells (middle) and vulval cells (bottom) and yellow arrows indicate hypodermal cells near rectum. Images are representative of 40 animals examined over two biological replicates. (C) Time course of *-2.8kb mlt-11p::mNG::tbb-2 3'UTR* expression in embryos with stages determined by embryo morphology. (D) Hypodermal expression pattern in L4s of three single-copy integrated promoter reporters. One reporter uses the same sequence used by Frand *et al.* (2005) and the other two use -2.8 kb and 5.2 kb of sequence upstream of the *mlt-11* transcriptional start site (TSS). (E) Hypodermal expression pattern in L4s of single-copy integrated promoter reporters with the *pes-10* minimal promoter elements combined with *mlt-11* promoter sequence fragments corresponding to either NHR-23 ChIP-seq or peaks or ATAC-seq open chromatin regions. (F) Hypodermal expression pattern in L4 larvae of the indicated single-copy integrated promoter reporters. Reporter strains were fed either control or *nhr-23(RNAi)* knockdown conditions. Scale bars are 20 μm in B, D, E and F and 10 μm in C.

2, 3 and 4 (Fig. 2E). We could detect no expression of reporters containing peak 1 nor

ATAC-seq 1 or 2 (Fig. 2E). To test whether NHR-23 regulated expression of the *mlt-11*

peak 3 pes-10::mNeonGreen promoter reporter, we performed *nhr-23* RNAi. RNAi of *nhr-*

23 reduced expression of both full length (Fig. 2F) and *mlt-11 peak 3 pes-*

10::mNeonGreen promoter reporters (Fig. 2F). These data indicate that NHR-23

regulates *mlt-11* and that the DNA sequences in NHR-23 ChIP-seq peaks 3 and 4 play

an important role in this regulation.

MLT-11 transiently localizes to epithelial aECMs

To gain insight into how MLT-11 regulates molting, we analyzed spatiotemporal MLT-11

expression using vulval morphology in L4 larvae for precise staging [47]. We knocked an

mNeonGreen::3xFLAG cassette into the 3' end of the 7th exon producing an internal

translational fusion that labels all described *mlt-11* isoforms (Fig. 3A). MLT-

11::mNeonGreen::3xFLAG (MLT-11::mNG) was detected internally in punctate and

tubular structures in the hypodermis that are likely lysosomal based on previous

colocalization studies of a *mlt-11::mScarlet* knock-in and the precuticle component *noah-*

1 (Fig. 3B)[32,48]. MLT-11::mNG is also transiently detected in the aECM of hyp7 and

seam cells, first appearing in furrows of the aECM (larval sub-stage L4.3) and then

localizing to annuli in L4.5 (Fig. 3B). By L4.9, localization in the aECM diminished beyond

observable levels and we only observed internal punctate expression (Fig. 3B). MLT-

11::mNG was seen in the aECM of seam cells from early to mid-L4, localizing to alae and

then disappeared in stage L4.6, while loss of MLT-11::mNG in the annuli began in stage

L4.6 and was absent in the cuticle by stage L4.9. (Fig. 3B). The transient localization of

MLT-11 in the cuticle followed by endocytosis is reminiscent of the pre-cuticle component

NOAH-1, which we showed to be internalized in late L4 [32]. In the aECM of the

developing vulva, MLT-11::mNG was observed faintly beginning at L4.3, intensifying until

L4.5 and remaining localized through the end of the L4 larval stage (Fig. 3C). It appeared

most intensely in the aECM of the inner vulva, adjacent to vulC, D, E and F (Fig. 3C).

MLT-11::mNG was also observed in the epithelial aECM of other external-facing orifices:

the buccal cavity, excretory duct and rectum (Fig. 3D-F). Together, these data indicate

that MLT-11 lines openings to the exterior and transiently associates with the cuticle

aECM, similar to precuticle components [24,32].

Fig. 3. MLT-11 localizes to the aECM in seam and hypodermal cells and interfacial orifices. (A) Schematic of the interna; *mlt-11::mNG::3xFLAG* knock-in. (B) Developmental time course of MLT-11::mNG in the aECM and internal vesicles of seam and hypodermal cells in the indicated L4 substages. (C) Developmental time course of MLT-11::mNG in the vulva of the indicated L4 substages. (D) Representative images of MLT-11::mNG lining the lumen of the buccal cavity, excretory duct and rectum. Images are representative of 40 animals examined over two biological replicates. Scale bars are 20 µm.

***mlt-11* knockdown causes defective aECM structure and function**

Our *mlt-11::mNG* translational fusion strain provided an entry point into understanding

how the candidate NHR-23 regulatory elements (Fig. 2) in the *mlt-11* promoter regulate

MLT-11 expression. Using CRISPR/Cas9, we individually deleted the sequence from

each NHR-23 ChIP-seq peak (Fig. 4A) in the MLT-11::mNG background and observed

expression of the translational fusion. MLT-11::mNG expression was comparable
between the control and strains with deletions of peaks 1 or 2 (Fig. 4B). However MLT-
11::mNG expression was reduced following deletion of peaks 3 or 4 (Fig. 4B). Deletion
of both peaks 3 and 4 in the same strain resulted in severely reduced expression of MLT-
11::mNG (Fig. 4B), suggesting that together these sequences are necessary for activation
of expression of *mlt-11* by NHR-23. We also scored developmental rate, size, and motility

Fig. 4. Disruption of *mlt-11* expression by RNAi or promoter deletion causes abnormalities in cuticle structure. (A) Genome browser track of *mlt-11* promoter depicting NHR-23 ChIP-seq peaks; relative genomic locations of *mlt-11* promoter sequence deletions; an H3K4me3 ChIP-seq track to indicate potential enhancers; conservation calculated across 26 nematode species; relative genomic locations of NHR-23 binding motifs. (B) Representative images of MLT-11::mNG::3xFLAG in mid-stage L4 larvae with endogenous deletion of the indicated promoter regions. (C-F) Representative images of mNG::3xFLAG fused to ROL-6 (C), BLI-1 (D), CUT-2 (E) or COL-19 (F) in the cuticles of mid-stage L4 larvae in control (full promoter) or *mlt-11* expression abrogated (peak 3Δ, peaks 3+4Δ, *mlt-11(RNAi)*) backgrounds. (G) (DIC images of adult alae in wild-type animals, *mlt-11(peaks 3+4Δ)* deletion mutants, and *mlt-11(RNAi)* animals. Scale bars are 10 μm in B-F and 5 μm H.

187   (Fig. S1). Interestingly, peak 2 Δ, peak 3+4 Δ and *mlt-11(RNAi)* animals exhibited

188   comparable developmental delay (Fig. S1A), but peak 2Δ animals had wild type motility

189   and size (Fig. S1B, C). In contrast, peak 3+4 Δ and *mlt-11(RNAi)* animals exhibited a

190   significant motility defect with the RNAi causing a stronger phenotype and a comparable

191   smaller body size (Fig. S1B,C).

Supplemental Fig. 1. *mlt-11* expression abrogation causes developmental delay, motility defects and smaller bodies. Worms of the indicated genotypes and treatment (control or *mlt-11(RNAi))* were allowed to lay eggs for two hours, then removed from the plates. Embryos were allowed to develop for 51 hours, then scored for (A) developmental stage (L3 or L4 substage), (B) thrashes per minute and (C) body length and width.

Given that *mlt-11* expression is regulated by NHR-23 and that NHR-23 depletion causes

aECM aberrations, we next wished to see how *mlt-11* inactivation impacted cuticle

structure and function. The *C. elegans* cuticle is a complex structure that is composed of

multiple layers (Figure 1) [4].   In the adult cuticle, the basal layer is closest to the

hypodermis, the cortical layer is furthest from the hypodermis and a fluid-filled medial
layer is between them. Each layer contains a unique complement of collagen proteins
that provide structure to the cuticle [13]. Given that knockdown of *mlt-11* by RNAi (Frand
et al., 2005) causes gross molting defects, we surmised that *mlt-11* could have a role in
establishing or maintaining the structure of the cuticle, in apolysis or ecdysis. To test a
role in cuticle structure, we depleted *mlt-11* by RNAi or deleted promoter elements in
strains carrying translational mNG fusions to proteins that mark the basal layer (ROL-6,
COL-19), medial layer (BLI-1), and cortical layer (CUT-2) [7,8,49].

ROL-6 is a collagen that localizes to tight longitudinal bands (furrows) in the basal layer
of the adult cuticle (Fig. 4C)[32,50]. It also resides in patches along the junction of
opposing annuli above seam cells (Fig. 4C). Following deletion of peak 3 or peaks 3+4,
ROL-6::mNG localized to both furrows and annuli. Gaps were observed between annuli
and patches were still seen, but smaller and fewer in number over seam cells. Following
RNAi knockdown of *mlt-11*, annuli were separated by larger and more numerous gaps.
Localization of ROL-6::mNG was absent over seam cells and opposing annuli at the
junction were completely separated from each other and often broken at their termini (Fig.
4C). COL-19::mNG, another basal layer marker and adult-specific collagen normally
localizes to alae and annuli, but following *mlt-11* RNAi we observed a reduction in
expression and a loss of localization over the seam cells (Fig. 4F)[51].

BLI-1 is a structural collagen found in the medial layer of the adult cuticle, providing
connection between the basal and cortical layers in the form of vertical struts [7,52]. BLI-
1::mNG localized in punctae organized in circumferential rows across the hyp7 cuticle,
but is absent over seam cells (Fig. 4D). Deletion of *mlt-11* peak 3 led to gaps between
the rows and inconsistency in the size of punctae (Fig. 4D). Deletion of peaks 3 and 4
together or RNAi knockdown caused further disorganization of BLI-1::mNG rows and
mislocalization of punctae in seam cuticle (Fig. 4D).

Similarly, CUT-2, a cuticlin in the cortical layer [8,53], had localization patterns that were
altered following knockdown of *mlt-11*. CUT-2::mNG displayed aberrant localization over

the seam cells and a fibrous pattern over seam cells following *mlt-11* RNAi (Fig. 4E).

Together, this suggests *mlt-11* is necessary for proper formation or patterning of each

layer of the adult cuticle.

Given the aberrant localization of ROL-6, BLI-1, CUT-2, and COL-19 over the seam cells,

we next examined alae morphology. The lateral alae are cuticle ridges formed by the

interaction between the actin cytoskeleton in epithelial cells and an extracellular

provisional matrix [12,54]. Three continuous longitudinal ridges span the midline of lateral

surfaces of adult worms (Fig. 4G). Deletion of *mlt-11* peaks 3 and 4 resulted in

discontinuous alae. Knockdown of *mlt-11* with RNAi caused the formation of spotty

stretches of discontinuous and disorganized alae (Fig. 4G). It is evident that *mlt-11* is

necessary for accurate development of cuticle structures derived from both hypodermal

and seam cells.

Reduced *mlt-11* expression disrupts the cuticle permeability barrier

The cuticle functions as a barrier to protect *C. elegans* from environmental threats such

as pathogens and xenobiotics. Gaps and disorganization of multiple layers of the *C.*

elegans cuticle following *mlt-11* knockdown suggest that this cuticle barrier function may

be disrupted, similar to what we observed in NHR-23-depleted animals [32]. To test the

cuticle barrier function, we incubated control, promoter deletion mutants and *mlt-11(RNAi)*

animals with the cuticle impermeable, cell membrane permeable Hoechst 33258 dye and

scored animals with stained nuclei. In wild-type control and single peak deletion animals

we observed no Hoechst staining, while we observed staining in 91% of peak 3+4

deletion mutants and 93% of *mlt-11(RNAi)* worms (Fig. 5A). This barrier defect was

comparable to the *bus-8* positive control strain [55]. Additionally, double deletion of peaks

3+4 as well as *mlt-11* RNAi-induced expression of an *nlp-29::GFP* promoter reporter

generally activated by infection, acute stress, and physical damage to the cuticle (Fig. 5B)

[56,57]. Together, these indicate *mlt-11* is necessary for the barrier function of the cuticle.

Fig. 5. *mlt-11* knockdown causes defective aECM structure and function. (A,B) Representative images of mid-stage L4 larvae in animals of the indicated genotype/treatment. (A) Worms were washed and incubated with the cuticle impermeable/membrane permeable Hoechst 33258 dye before being imaged. (B) Worms carried an *nlp-29p::GFP* reporter activated by infection, acute stress, and physical damage to the cuticle [56,57]. Three biological replicates were performed and over 40 animals scored. Scale bars are 20 µm.

mlt-11 activity is necessary in seam cells for larval development

As we observed aberrant cuticle protein localization in the aECM over both seam and hypodermal cells, we were curious whether *mlt-11* expressed in these cells may be playing distinct functions and turned to a tissue-specific RNAi approach (Fig. 6A). *mlt-11* knockdown in JDW371, a tissue-specific RNAi strain that restricts knockdown to seam, hypodermal, and intestinal cells [32], phenocopied *mlt-11* RNAi in wildtype animals with respect to developmental delay and molting defects (Fig. 6B-D). *mlt-11* RNAi in QK52, a hypodermal and seam cell-specific RNAi strain, produced less penetrant developmental delay and molting defects (Fig. 6B-D). Notably, *mlt-11* RNAi in a hypodermal-specific

Fig. 6. *mlt-11* is necessary in seam cells for molting and larval development. (A) Tissue-specific RNAi strains used. A timed egg lay of animals of the indicated genotype were performed on control or *mlt-11 (RNAi)* plates and phenotypes were scored three days later. (B) Representative images of animals of the indicated genotype grown on control or *mlt-11* RNAi. The red box highlights conditions that produced smaller larvae with molting defects. (C) Developmental delay was scored in the indicated strains grown on control or *mlt-11* RNAi plates and classified as a failure to reach adulthood after 72 hours of growth. (D) Molting defects were scored in the indicated strains on control or *mlt-11* RNAi. Scored defects included animals dragging cuticles, ecdysis failure and cuticle corsets. Tissue-specific RNAi data is from two independent replicates.

271   RNAi strain, JDW510, produced no developmental delay or molting defects, suggesting

272   *mlt-11* activity is necessary in seam cells and perhaps, in a limited capacity, in other

273   hypodermal cells (Fig. 6B-D).

Fig. 7. MLT-11 localizes to the embryonic sheath and is necessary for proper hypodermal development during embryogenesis. (A) Developmental time course of MLT-11::mNG in the aECM and internal vesicles of seam and hypodermal cells in the indicated embryonic substages. Yellow arrows indicate MLT-11::mNG has been endocytosed to internal vesicles at the end of elongation. (B) Endogenous deletions of the signal sequence and Kunitz domains 3-10 (*mlt-11Δ*) are homozygous inviable and so were balanced and maintained as heterozygotes. These heterozygotes were plated as adults and allowed to lay eggs for 2 hours, then removed. The resulting embryos were allowed to develop for 24 hours and then scored for hatching. (C) Representative orthogonal projection of a balanced *mlt-11Δ* heterozygous embryo and a homozygous *mlt-11Δ* embryo expressing DLG-1::mNG labeled adherens junctions in the pharynx, intestine, and hypodermis. White arrows indicate invaginations in the hypodermal cells. Scale bars are 20 μm.

***mlt-11* is a component of the embryonic sheath and is required for elongation**

Most of our analyses of *mlt-11* was carried out in L4 larvae as they developed the adult

and final cuticle. Abrogation of *mlt-11* expression led to wide-spread cuticle defects

suggesting MLT-11 may play a larger role in patterning the cuticle as opposed to being a
component of the cuticle itself. Indeed, we never saw MLT-11 shed in the cuticle, rather
it was endocytosed, likely to lysosomes (Fig. 3B). This idea is supported by the timing of
MLT-11::mNG expression and localization at the intermolt when the pre-cuticle or
provisional matrix is present. A similar scenario plays out during embryonic
morphogenesis when a ball of cells transitions to an elongated tube. The embryonic
sheath serves as a framework upon which the L1 cuticle is built [21,22,42]. The precuticle
components are endocytosed and recycled for subsequent rounds of cuticle synthesis.
Given the embryonic *mlt-11* promoter reporter expression we observed (Fig. 2C), we
imaged MLT-11::mNG in embryos from prior to elongation through to just prior to hatching
(Fig. 7A). Expression was detected in the posterior of bean stage embryos, quickly
encompassing the entire embryo by comma stage. MLT-11::mNG localizes to the aECM
of the animals and is endocytosed prior to hatching, similar to other precuticle
components [20].

Loss of other precuticle components causes embryonic lethality or morphogenesis
defects [20,23]. However, we had not observed embryonic defects with our promoter
element deletion strains, the *rol-9* allele, or our RNAi experiments. However, as RNAi
causes reduction-of-function, we turned to deletion alleles. MLT-11 is predicted to be a
large protein (234-341 kDA) with a signal sequence, a thyroglobulin domain, and 10
Kunitz protease inhibitor domains (Fig. 3A). Eight of the 10 Kunitz domains contain all 6
conserved cysteine residues that are necessary to form three disulfide bonds critical for
stabilizing the domain (Fig. S2)[58]. To prevent MLT-11 secretion, we deleted the signal
sequence. We also deleted the coding region downstream from the *rol-9(sc148)* indel,
removing Kunitz domains 3-10 (*mlt-11Δ*; Fig. 3A). We were unable to homozygose these
alleles, indicating embryonic lethality or sterility. As *mlt-11* is on the far right arm of
chromosome V in an area lacking classic genetic balancer chromosomes, we used a
myo-2p::GFP::unc-54 3'UTR insertion approximately 40 kb away to balance the deletions.
We never observed viable progeny from balanced mutant worms that lacked GFP in the
pharynx, but these balanced worms produced roughly 25% dead embryos, a rate
expected for a homozygous lethal mutation (Fig. 7B).

Supplemental Fig. 2. Alignment of MLT-11 homologs. MLT-11 homologs from the indicated nematode species were aligned using Clustal Omega. The length in amino acids of each homolog follows the species and homolog name. To the left and right of the alignment are amino acid positions of the end residues for each protein. Blue shading indicates conserved sequences and the histogram at the bottom depicts the degree of conservation with a consensus sequence listed below. The positions of the *C. elegans* MLT-11 signal sequence, thyroglobulin domain, and ten Kunitz domains are indicated. We note that we cut off the extended *P. pacificus* C-terminus (amino acids 2626-3742) since no sequence aligned to it as all proteins terminated at the *C. elegans* MLT-11 stop codon. No predicted motifs are found in the *P. pacificus* C-terminus.

We next examined the nature of the embryonic lethality in *mlt-11* deletion mutants using

a DLG-1::mNG allele to mark adherens junctions [59]. In control embryos, DLG-1::mNG

labeled adherens junctions in the pharynx, intestine, and hypodermis (Fig. 7C).

Hypodermal cells stretched around the circumference of and down the length of the coiled

embryo in relatively straight lines. In contrast, *mlt-11Δ* embryos matched to the same

stage displayed severe disorganization. The pharynx and foregut adherens junctions

appeared wild type, but the remainder of the junctions were disorganized and there was

evidence of invaginations in the hypodermis (Fig. 7C). These data implicate the signal

sequence and sequence within the Kunitz 3-10 region as being essential for hypodermal

organization during embryogenesis (Fig. 7C).

mlt-11(RNAi) suppresses the blister phenotype of _bli-4_ hypomorphic adults

The expression timing of _mlt-11_ and deletion phenotypes were reminiscent of the

precuticle components _noah-1_ and _noah-2_, and the subtilisin/kexin family protease _bli-4_

[20,42]. _noah-1_ and _noah-2_ mutants are essential for embryonic integrity and elongation,

while _bli-4_ null animals initially elongate but then retract and collapse [20,42]. We were

particularly intrigued by the similarity of MLT-11 to BLI-4 because both proteins are

expressed in larvae and embryos and essential for viability at both stages. MLT-11 is

endocytosed (Fig. 7A) while BLI-4 is ingested [42]. We hypothesized that MLT-11 could

conceivably inhibit BLI-4. In this model, a reduction in _bli-4_ activity could be rescued by

reduction in _mlt-11_ levels. _bli-4(e937)_ is a hypomorphic allele that causes a blistered

cuticle in adult larvae (Fig. 8A)[43]. Strikingly, _mlt-11_ completely suppressed the blistered

phenotype. To test whether _mlt-11_ inactivation affected BLI-4 levels or localization, we

depleted _mlt-11_ RNAi in a _bli-4::sfGFP_ strain. There was no significant change in BLI-

4::sfGFP expression or localization in _mlt-11(RNAi)_ compared to the control animals (Fig.

8B). These data indicate that reduction of MLT-11 levels compensates for reduction of

bli-4 function.

Fig. 8. *mlt-11(RNAi)* suppresses a reduction-of-function allele of *bli-4*. (A) Representative DIC images of *bli-4(e937)* adults grown on control or *mlt-11* RNAi plates. A large blister (separation of the cortical layer from the body of the worm) is indicated with an arrow. Over 200 animals were scored over two independent experiments. *bli-4(e937)* mutants displayed blistering in ~90% of adults similar to what Thacker *et al.* (1995) reported, which *mlt-11* RNAi completely suppressed the blister phenotype (B) Representative images of BLI-4::sfGFP seen in internal hypodermal punctae in control and *mlt-11(RNAi)* mid-late stage L4 larvae. Images are representative of >40 animals scored over two independent experiments. Scale bars are 20 μm.

DISCUSSION

Despite their importance, how aECMs are built and dynamically remodeled during development and disease remains poorly understood. Using the *C. elegans* cuticle as a model aECM we show that *mlt-11* is directly regulated by the NHR-23 transcription factor. MLT-11::mNG transiently localizes to the embryonic and larval aECM before endocytosis and trafficking to a compartment that is most likely lysosomal. MLT-11::mNG also lines openings to the exterior (vulva, buccal cavity, excretory duct, rectal epithelium). *mlt-11* inactivation affects the localization of translational fusions marking the basal (ROL-6::mNG, COL-19), medial (BLI-1::mNG), and cortical layers (CUT-2) of the cuticle indicating a broad role for MLT-11 in patterning this aECM. In agreement with these broad defects in aECM component localization, *mlt-11* inactivation causes loss of the epithelial barrier and activation of an epidermal damage reporter. Tissue-specific RNAi data indicates that *mlt-11* may act primarily in seam cells, though a role in the hypodermis is also supported. Alleles predicted to cause a loss of *mlt-11* function cause embryonic lethality with severe disorganization of hypodermal adherens junctions. *mlt-11*

inactivation suppresses the blistering phenotype caused by the *e937* mutation in the *bli-*
4 subtilisin/kexin family protease. Together, this work suggests that MLT-11 may be part
of the pre-cuticle and functions to coordinate proper assembly of the aECM.

mlt-11 is an NHR-23-regulated gene required for patterning of all cuticle layers
mlt-11 was identified as an *nhr-23*-regulated gene through studies using promoter
reporter and a microarray analysis of genes misregulated in L2 animals following *nhr-23*
RNAi [33,44]. NHR-23 ChIP-seq data identified four NHR-23 occupied regions 5' to the
transcription start site, and we showed that peaks 3 and 4 drive robust reporter expression
and NHR-23 controls the expression of a *mlt-11 peak 3* reporter. While it is not clear
whether NHR-23 regulates expression of these reporters through direct DNA binding or
whether the regulation is indirect, there are consensus sites that NHR-23 is known to bind
within or adjacent to these peak sequences (Fig. 2) [35]. Consistent with the reporter data,
deletion of the endogenous peak sequences revealed that peaks 3 and 4 produced the
strongest effect on MLT-11::mNG expression and a double deletion of peaks 3 and 4
results in even stronger reduction in expression. Despite robust hypodermal expression
and localization, our tissue-specific RNAi suggests that the seam cells are an important
site of *mlt-11* expression (Fig. 3,4,6). We consistently saw the most severe RNAi
phenotypes in tissue-specific RNAi strains using a seam cell enriched promoter and no
phenotypes from a hypodermal/intestinal-specific RNAi strain (Fig. 6)[32]. Consistent with
these data, we observed aberrant expression of ROL-6::mNG, CUT-2::mNG, and COL-
19::mNG over the seam cells (Fig. 4). We also see gaps in the alae following *mlt-11*
inactivation. Moreover, BLI-1 is normally excluded from the aECM over the seam cells
but invades this area following *mlt-11* knockdown (Fig. 4). These data were reminiscent
of our work on NHR-23 as depletion of this factor causes aberrant aECM formation over
the seam cells and *nhr-23* activity also was necessary in the seam cells for developmental
progression [32]. *nhr-23* and *lin-42* both regulate the molting timer and the heterochronic
pathway and these two timers need to be tightly coordinated [27–31,36]. Pharmacologic
uncoupling through nicotinic agonists results in animal death through rupturing when
animals molt before seam cell division is completed [60]. Exploring the tissue-specificity

and coordination of the molting and heterochronic timers and how the aECM might impact
these timers is an important future direction.
386

A *mlt-11* allelic series

Following *cis*-regulatory element deletion in the *mlt-11* promoter, deletion of *mlt-11* coding
sequence, and RNAi we observed a range of phenotypic severity: i) embryonic lethality;
ii) molting defects and developmental delay; iii) cuticle organization defects; and iv) cuticle
barrier defects. These mutants and *mlt-11(RNAi)* comprise an allelic series to link *mlt-11*
levels to specific phenotypes. Embryonic development seems to require the least MLT-
11 activity as we only encountered embryonic lethality when we prevented MLT-11
secretion through signal sequence deletion, or deleted the majority of the protein (Fig. 7).
Motility was also less sensitive to MLT-11 levels as we needed RNAi depletion to observe
phenotypes (Fig. S1). BLI-1 localization, developmental rate and the cuticle barrier
displayed intermediate sensitivity to MLT-11 levels as peak 3+4 deletion caused defects,
as did RNAi (Fig. 4, 5, 6). Finally ROL-6 organization seemed highly sensitive to MLT-11
levels, as peak 3 deletion caused a reduction of MLT-11 levels and aberrant ROL-6
localization especially over the seam cells (Fig. 4). MLT-11 has 10 predicted protease
inhibitor domains most of which contain residues necessary for Kunitz inhibitor activity
[58,61], which could buffer reduction of levels. It will be important to perform careful
structure-function analyses to determine the role of specific Kunitz domains and other
sequences in mediating this allelic series.
405

Is MLT-11 a pre-cuticle component?

The hallmarks of *bona fide* precuticle components are that they are secreted, transiently
localize in the aECM before being endocytosed and degraded in lysosomes [14–19].
Precuticle components are also found in the embryonic sheath and are required for
embryonic viability [20,23]. We found that MLT-11 is secreted and endocytosed in both
embryos and larvae (Fig. 3, 7). Predicted nulls or strong reduction-of-function alleles
cause embryonic death during elongation which is strikingly similar to *noah-1* and *noah-*
2 mutants [20]. *mlt-11(RNAi)* animals display gaps in alae similar to precuticle mutants
[12], and we observed defective localization of mNG translational fusions marking each

layer of the adult cuticle (Fig. 4). If MLT-11 is indeed a precuticle component it would be

the first putative protease inhibitor to localize to this structure. What is MLT-11 doing in

this transient matrix? It could function as a protease inhibitor restraining protease

substrates until it is endocytosed, releasing the protease to digest aECM substrates.

Alternatively, MLT-11 could function as a scaffold to coordinate aECM and precuticle

assembly. The MLT-11 dynamics raise several questions. How is it endocytosed? It is

internalized in L4.7-L4.8 stage larvae while NOAH-1 is endocytosed in L4.9 larvae, so

there appears to be intriguing temporal regulation of endocytosis (Fig. 3) [32]. Additionally

MLT-11 localization in the aECM over the alae is lost in L4.6 stage larvae, earlier than

endocytosis from the hypodermal aECM. Understanding the mechanisms controlling

precuticle internalization will provide an important entry point into aECM dynamics during

development.

***mlt-11* and *bli-4* genetically interact**

Our working model is that MLT-11 is a protease inhibitor that restrains the activity of

cognate protease(s). In looking for candidate substrates we looked for protease genes

that are essential and expressed in both embryos and larvae. While Kunitz domains are

typically serine protease inhibitors, in the flatworm *Fasciola hepatica* an atypical Kunitz

domain protein inhibits cysteine proteases, indicating a broader spectrum of potential

substrates [62]. BLI-4 was an intriguing candidate. Reduction-of-function alleles cause

either larval lethality or blistering, while nulls cause embryonic lethality but later in

development than *mlt-11* nulls (Fig. 7) [43,43,63]. *bli-4(0)* mutants complete elongation

but then the embryo retracts and loses organization [42]. BLI-4::sfGFP is secreted in

embryos and ingested rather than endocytosed. We reasoned that reducing *mlt-11*

activity might suppress *bli-4* hypomorphs by rebalancing protease activity, similar to how

inactivation of two antagonistic developmental regulators restores correct cell fates in the

C. elegans somatic gonad precursor cells [64]. When we inactivated *mlt-11* by RNAi we

completely suppressed the blister phenotype of *bli-4(e937)* hypomorphs. This effect was

not through regulating BLI-4 levels or localization as we saw no obvious changes in BLI-

4::sfGFP. These data could reflect a direct interaction between BLI-4 and MLT-11. In this

model, some *mlt-11* phenotypes could come from aberrant BLI-4 activity. Alternatively,

the interaction could be indirect with MLT-11 broadly affecting aECM organization. For
example, some suppressors of the *bli-1* blistered phenotype are collagen mutants that
are thought to make a more larval-like cuticle and eliminate the fluid-filled medial layer
[7]. Testing MLT-11 and BLI-4 for a direct interaction, exploring the nature of the
suppression, and testing for altered processing of predicted BLI-4 substrates are all
important future directions.

Future perspective

This work provides an entry point into understanding how a predicted protease inhibitor
regulates the structure and function of the *C. elegans* aECM. Going forward, it will be
critical to determine the biochemical activity of MLT-11 to determine what function(s) it
plays in the aECM before internalization and what its substrates are. As proteases are
important targets to combat parasitic nematode infections, understanding how they are
regulated during development by endogenous protease inhibitors will be critical to
develop novel approaches to combat this group of devastating pathogens.

MATERIALS AND METHODS

Strains and culture

C. elegans were cultured as originally described [65], except worms were grown on
MYOB media instead of NGM. MYOB agar was made as previously described [66].

Strains created by injection in the Ward Lab and used in this study:

Name	Genotype
JDW359	*ezIs2 III [fkh-6::GFP + unc-119(+)] ; him-8(e1489) IV ; F46B3.7(gk5359[loxP + myo-2p::GFP::unc-54 3' UTR + rps-27p::neoR::unc-54 3' UTR + loxP]) V.*
JDW371	*jsTi1493 {mosL loxP [wrdSi72(SCMp::pes-10delta::rde-1 CDS+3'UTR)] FRT3::mosR} IV ; rde-1(ne300) V [32]*

JDW380	*jsTi1493 {mosL loxP [wrdSi72(mlt-11(-2.8kb)p::mNeonGreen(dpi)::tbb-2 3'UTR)] FRT3::mosR} IV*
JDW387	*mlt-11(wrd82[mlt-11 Kunitz 3-10 deletion]) , oxTi633 [eft-3p::tdTomato::H2B::unc-54 3'UTR] / F46B3.7(gk5359[loxP + myo-2p::GFP::unc-54 3' UTR + rps-27p::neoR::unc-54 3' UTR + loxP]) V*
JDW389	*bli-1(wrd84[linker::mNeonGreen::3xFLAG::linker]) [32]*
JDW401	*mlt-11(wrd82[mlt-11 Kunitz 3-10 deletion]) , oxTi633 [eft-3p::tdTomato::H2B::unc-54 3'UTR] / F46B3.7(gk5359[loxP + myo-2p::GFP::unc-54 3' UTR + rps-27p::neoR::unc-54 3' UTR + loxP]) V. dlg-1(cp301[dlg-1::mNG-C1^3xFlag]) X*
JDW458	*rol-6(wrd117[rol-6::C-term mNeonGreen (dpi)::3xFLAG::linker]) [32]*
JDW510	*jsTi1493 {mosL loxP [wrdSi97(suro-1p::rde-1 CDS+3'UTR)] FRT3::mosR} IV ; rde-1 (ne300) V* (Johnson *et al.,* 2022) [32]
JDW511	*ezIs2 [fkh-6::GFP + unc-119(+)] / + III ; him-8(e1489) / + IV ; mlt-11(wrd126[mlt-11 signal sequence deletion...mNG::3xFLAG]) / F46B3.7(gk5359[loxP + myo-2p::GFP::unc-54 3' UTR + rps-27p::neoR::unc-54 3' UTR + loxP]) V*
JDW523	*mlt-11(wrd141[mlt-11 signal sequence deletion]) , oxTi633 [eft-3p::tdTomato::H2B::unc-54 3'UTR + Cbr-unc-119(+)] V*
JDW535	*jsSi1579 jsSi1706 jsSi1726 wrdSi103[loxP myo-2p::NLS::mNeonGreen, rps-0p HygR, loxP mlt-11p peak 3::pes-10 minimal promoter::mNeonGreen(dpi)::PEST::3xFLAG::tbb-2 3'UTR FRT3] II*
JDW541	*mlt-11(wrd153[mNG::3xFLAG internal]) V*
JDW551	*mlt-11(wrd158[mlt-11 promoter chip-seq peak 3 deletion]), oxTi633 [eft-3p::tdTomato::H2B::unc-54 3'UTR + Cbr-unc-119(+)] V*
JDW558	*mlt-11(wrd153[mNG::3xFLAG internal], wrd161[promoter chip-seq peak 2 deletion]) V*
JDW559	*mlt-11(wrd153[mNG::3xFLAG internal], wrd162[promoter chip-seq peak 3 deletion]) V*
JDW655	*cut-2(wrd233{cut-2::C-terminal modular linker::mNG::3xFLAG::linker]) V*

JDW663	*mlt-11(wrd153[mNG::3xFLAG internal], mlt-11(wrd237[mlt-11 promoter chip-seq peak 4 deletion]) V*
JDW666	*jsSi1579 jsSi1706 jsSi1726 wrdSi108[loxP myo-2p::NLS::mNeonGreen, rps-0p HygR, loxP mlt-11p peaks 1, 2 and 3::mNeonGreen(dpi)::tbb-2 3'UTR FRT3] II*
JDW667	*jsSi1579 jsSi1706 jsSi1726 wrdSi109[loxP myo-2p::NLS::mNeonGreen, rps-0p HygR, loxP mlt-11p peak 1::pes-10 minimal promoter::mNeonGreen(dpi)::tbb-2 3'UTR FRT3] II*
JDW668	*jsSi1579 jsSi1706 jsSi1726 wrdSi110[loxP myo-2p::NLS::mNeonGreen, rps-0p HygR, loxP mlt-11p peak 2::pes-10 minimal promoter::mNeonGreen(dpi)::tbb-2 3'UTR FRT3] II*
JDW669	*jsSi1579 jsSi1706 jsSi1726 wrdSi111[loxP myo-2p::NLS::mNeonGreen, rps-0p HygR, loxP mlt-11p peak 4::pes-10 minimal promoter::mNeonGreen(dpi)::tbb-2 3'UTR FRT3] II*
JDW670	*jsSi1579 jsSi1706 jsSi1726 wrdSi112[loxP myo-2p::NLS::mNeonGreen, rps-0p HygR, loxP mlt-11p enhancer 1::pes-10 minimal promoter::mNeonGreen(dpi)::tbb-2 3'UTR FRT3] II*
JDW671	*jsSi1579 jsSi1706 jsSi1726 wrdSi113[loxP myo-2p::NLS::mNeonGreen, rps-0p HygR, loxP mlt-11p enhancer 2::pes-10 minimal promoter::mNeonGreen(dpi)::tbb-2 3'UTR FRT3] II*
JDW687	*jsSi1579 jsSi1706 jsSi1726 wrdSi114[loxP myo-2p::NLS::mNeonGreen, rps-0p HygR, loxP mlt-11p(-5.2 kb)::mNeonGreen(dpi)::3xFLAG::PEST::tbb-2 3'UTR FRT3] II*
JDW696	*mlt-11(wrd153[mNG::3xFLAG internal], wrd268[mlt-11 promoter chip-seq peak 1 deletion]) V*
JDW703	*mlt-11(wrd153[mNG::3xFLAG internal], wrd162[promoter chip-seq peak 3 deletion]), wrd249[mlt-11 promoter chip-seq peak 4 deletion]) V*
JDW709	*mlt-11(wrd162[promoter chip-seq peak 3 deletion], wrd252[mlt-11 promoter chip-seq peak 4 deletion]) V*
JDW710	*rol-6(wrd253[rol-6::C-term mNeonGreen (dpi)::3xFLAG::linker]) II; mlt-11(wrd158[mlt-11 promoter chip-seq peak 3 deletion]) , oxTi633 [eft-3p::tdTomato::H2B::unc-54 3'UTR + Cbr-unc-119(+)] V*
JDW711	*jsSi1579 jsSi1706 jsSi1726 wrdSi115[loxP myo-2p::NLS::mNeonGreen, rps-0p HygR, loxP empty control::pes-10 minimal promoter::mNeonGreen(dpi)::tbb-2 3'UTR FRT3] II*

JDW713	*bli-1(wrd254[bli-1::internal mNeonGreen (dpi)::3xFLAG::linker]) II; mlt-11(wrd158[mlt-11 promoter chip-seq peak 3 deletion]) , oxTi633 [eft-3p::tdTomato::H2B::unc-54 3'UTR + Cbr-unc-119(+)] V*
JDW715	*rol-6(wrd255[rol-6::C-term mNeonGreen (dpi)::3xFLAG::linker]) II; mlt-11(wrd162[promoter chip-seq peak 3 deletion], wrd252[mlt-11 promoter chip-seq peak 4 deletion]) V*
JDW719	*jsSi1579 jsSi1706 jsSi1726 wrdSi116[loxP myo-2p::NLS::mNeonGreen, rps-0p HygR, loxP mlt-11p (Frand promoter -1977-5300 bp)::mNeonGreen (dpi)::3xFLAG::PEST::tbb-2 3'UTR FRT3] II*
JDW743	*bli-1(wrd254[bli-1::internal mNeonGreen (dpi)::3xFLAG::linker]) II; mlt-11(wrd158[mlt-11 promoter chip-seq peak 3 deletion]) , mlt-11(wrd288[mlt-11 promoter chip-seq peak 4 deletion]) , oxTi633 [eft-3p::tdTomato::H2B::unc-54 3'UTR + Cbr-unc-119(+)] V*
JDW750	*frIs7 [nlp-29p::GFP + col-12p::DsRed] IV ; mlt-11(wrd294[mlt-11 promoter chip-seq peak 3 deletion]) V*
JDW757	*frIs7 [nlp-29p::GFP + col-12p::DsRed] IV ; mlt-11(wrd294[mlt-11 promoter chip-seq peak 3 deletion] + wrd295[mlt-11 promoter chip-seq peak 4 deletion]) V*

467 **Strains provided by the *Caenorhabditis* Genetics Center:**

CB6193[67]	*bus-8B(e2885) X [55]*
CB937	*bli-4(e937) I [67]*
DZ325	*ezIs2 III [fkh-6::GFP + unc-119(+)] ; him-8(e1489) IV [68]*
EG7968	*unc-119(ed3) III; oxTi633 [eft-3p::tdTomato::H2B::unc-54 3'UTR + Cbr-unc-119(+)] V [69]*
JU2039	*mfIs70 [lin-31p::rde-1 + myo2p::GFP] IV ; rde-1(ne219) V [70]*
LP598	*dlg-1(cp301[dlg-1::mNG-C1^3xFlag]) X [59]*
MGH171	*alxIs9 [vha-6p::sid-1::SL2::GFP] sid-1(qt9) V ; alxIs9 [71]*
N2	*Wild-type[56]*
IG274	*frIs7 [nlp-29p::GFP + col-12p::DsRed] IV*

QK52	*rde-1(ne219) V ; xkIs99(wrt-2p::rde-1::unc-54 3'UTR) [71]*
VC4276	*F46B3.7(gk5359[loxP + myo-2p::GFP::unc-54 3' UTR + rps-27p::neoR::unc-54 3' UTR + loxP]) V [72]*
VP303	*rde-1(ne219) V ; kbIs7[nhx-2p::rde-1 + rol-6(su1006)]) [73]*
WM118	*rde-1(ne300) V ; neIs9(neIs9 [myo-3::HA::RDE-1 + rol-6(su1006)]) X [74]*
NM5548	*jsSi1579 jsSi1706 jsSi1726[loxP myo-2p FRT nlsCyOFP myo-2 3' mex-5p FLP D5 glh-2 3' FRT3] II [75]*

Other strains

PHX4625 *col-19::mNG(cyb4625) X* was a gift from Dr. Andrew Chisholm and will be described elsewhere. ARF403 *aaaIs24[egl-18p::utrnch::gfp::unc-54 3'UTR, ttx-3::gfp]* was previously published and a gift from Ali Frand and Meera Sundaram [12].

Genome Editing

All plasmids used are listed in Table S1. Annotated plasmid sequence files are provided in File S1. Specific cloning details and primers used are available upon request. JDW380 *jsTi1493 {mosL loxP [wrdSi72(mlt-11(-2.8kb)p::mNeonGreen(dpi)::tbb-2 3'UTR)] FRT3::mosR} IV* was created by recombination-mediated cassette exchange (RMCE)[76]. A 2.8 kb *mlt-11* promoter fragment was initially Gibson cloned into the *NLS::mScarlet (dpi)::tbb-2 3'UTR* vector pJW1841 [77] to generate pJW1934. The mScarlet cassette was then replaced with mNeonGreen (dpi) to generate pJW2229. The *mlt-11p (-2.8kb) mNeonGreen (dpi)-tbb-2 3'UTR* fragment was PCR amplified from pJW2229 and Gibson cloned into *Sph*I-HF+*Spe*I-HF double digested RMCE integration vector pLF3FShC to produce pJW2337. This vector was integrated into NM5179 and the SEC was excised as previously described [76].

A pJW2361 *mNeonGreen(dpi)::3xFLAG::PEST-tbb-2 3'UTR* vector for SapTrap with ATG and GTA connectors was constructed by linearizing pJW2322 [48] by PCR and Gibson cloning in PCR amplified *mNeonGreen (dpi)::3xFLAG* from pJW2172 and a

linker::PEST::tbb-2 3'UTR from pJW1836 [77]. The pJW2286 *mlt-11p (-2.8 kb)* promoter

for SapTrap cloning was previously described [48]. The *mlt-11* promoter fragment from

Frand *et al.* (2005) and the *mlt-11p (-5.3 kb)* promoter fragment were PCR amplified from

a fosmid containing the *mlt-11* gene and Gibson cloned into linearized pJW2286 to make

pJW2451 and pJW2457, respectively. These *mlt-11* promoter plasmids were SapTrap

cloned with pJW2361 and pNM4216 to generate insertion vectors for rRMCE [75,78]. The

remaining promoter reporters were constructed by SapTrap cloning and rapid RMCE

[75,78]. A pJW2365 *pes-10 minimal promoter::mNeonGreen(dpi):: 3xFLAG::PEST-tbb-2*

3'UTR vector for SapTrap with ATG and GTA connectors was constructed by linearing

pJW2361 and Gibson cloning in a *pes-10* minimal promoter amplified from pJW1947 [77].

The ATAC-seq and NHR-23 ChIP-seq peak DNA sequences were PCR amplified from

pJW2337 and Gibson cloned into pDONR221 with TGG and ATG connectors for

SapTrap. These plasmids were then SapTrap cloned with pJW2365 and pNM4216 to

generate insertion vectors for rRMCE [75,78].

mlt-11 promoter and coding region deletion strains were created by injection of Cas9

ribonucleoprotein complexes (RNPs)[79,80] [700 ng/µl IDT Cas9, 115 ng/µl each crRNA

and 250 ng/µl IDT tracrRNA], oligonucleotide repair template (110 ng/µl) and pSEM229

co-injection marker (25 ng/µl)[81] for screening into strain EG7968. Where possible, we

selected "GGNGG" crRNA targets as these have been the most robust in our hand and

support efficient editing [82]. F1s expressing the co-injection marker were isolated to lay

eggs and screened by PCR for the deletion. For lethal mutations, F2 progeny of a verified

F1 deletion mutant were crossed to JDW359 males expressing *myo-2::GFP* to genetically

balance the mutation. Genotyping primers are provided in Table S2. JDW389 was created

by injection of RNPs [700 ng/µl IDT Cas9, 115 ng/µl crRNA and 250 ng/µl IDT tracrRNA]

and a dsDNA repair template (25-50 ng/ul) created by PCR amplification of a plasmid

template into N2 animals [79,80](Table S1). PCR products were melted to boost editing

efficiency, as previously described [83]. For the *mlt-11* internal knock-in, the

mNeonGreen::3xFLAG cassette was inserted in an unstructured region of exon 7. The

JDW655 *cut-2*::mNG::3xFLAG knock-in used a pJW2332 repair template, which will be

described elsewhere (J.M.R. and J.D.W., unpublished). Sequences of CRISPR/Cas9-

mediated genome edits are provided in File S2. crRNAs used are provided in Table S3

and repair template oligos for deletions are provided in Table S4. F1 progeny were

screened by mNeonGreen expression.

Imaging

Synchronized animals were collected from MYOB, control, or auxin plates by either

picking or washing off plates. For washing, 1000 µl of M9 + 2% gelatin was added to the

plate or well, agitated to suspend animals in M9+gelatin, and then transferred to a 1.5 ml

tube. Animals were spun at 700xg for 1 min. The media was then aspirated off and

animals were resuspended in 500µl M9 + 2% gelatin with 5 mM levamisole. 12 µl of

animals in M9 +gel with levamisole solution were placed on slides with a 2% agarose pad

and secured with a coverslip. For picking, animals were transferred to a 10 µl drop of

M9+5 mM levamisole on a 2% agarose pad on a slide and secured with a coverslip.

Synchronized embryos were picked to a 2% agarose pad on a slide, secured with a

sealed coverslip and imaged once an hour. Heterozygous Kunitz 3-10Δ embryos were

identified by initial expression of a pharyngeal myo-2p::GFP::unc-54 3'UTR reporter on

the balancing wildtype chromosome. Images were acquired using a Plan-Apochromat

40x/1.3 Oil DIC lens or a Plan-Apochromat 63x/1.4 Oil DIC lens on an AxioImager M2

microscope (Carl Zeiss Microscopy, LLC) equipped with a Colibri 7 LED light source and

an Axiocam 506 mono camera. Acquired images were processed through Fiji software

(version: 2.0.0- rc-69/1.52p). For direct comparisons within a figure, we set the exposure

conditions to avoid pixel saturation of the brightest sample and kept equivalent exposure

for imaging of the other samples.

For the embryo imaging timecourse (Fg. 7A), 15-20 mlt-11(wrd153[mNG::3xFLAG

internal]) gravid adults were plated on a 6 cm plate seeded with OP50 and allowed to lay

eggs for ~3 hours. Adults were removed and embryos were collected with an eyelash pick

for imaging. A microscope slide was coated with a 1 mm thick agarose pad made with

2% agarose in milli-Q water. Embryos were transferred to the agarose pad and coated

with 15 uL of an M9+gelatin solution. A coverslip was added and the edges were sealed

with nail polish to prevent liquid evaporation. Slides were imaged at 100x magnification

on the above described Zeiss Axioimager. Embryo positions were marked via the 'Tiles'
feature and snaps were taken of each embryo every 15 minutes for 4 hours or until the
slide began to appear dry. Exposure times were kept constant throughout the timecourse.
Embryo stages imaged ranged from ~100 cell stage to 3-fold, with at least 10 embryos
observed for each stage across three replicate experiments.
557
558 **RNAi Knockdown**
559 RNA interference experiments were performed as in Johnson *et al.* (2023). Control RNAi
560 used an empty L4440. The *mlt-11 (RNAi)* vector was streaked from the Ahringer library
561 [84]. Vector sequences are provided in File S1.
562
563 **Hoechst staining**
564 Hoechst 33258 staining was performed as described previously [85], except that we used
565 10 μg/ml of Hoechst 33258 as previously described [86]. Two biological replicates were
566 performed examining 50 animals per experiment. Representative images were taken with
567 equivalent exposures using a 63× Oil DIC lens, as described in the imaging section.
568

Acknowledgements
We would like to thank Prof. David Fay and John C. Clancy for helpful conversations. We
thank Krista Myles, Patricia Bliatout, Zoe Johnson, Javier Hernandez Lopez, Zoie Reyna,
Emma Cadena, Valarie Hallin, and Olivia Vedar for research support. We thank Prof. Ali
Frand and Prof. Michael Nonet for strains and Prof. Meera Sundaram and Prof. Andrew
Chisholm for strains and helpful discussion. Some strains were provided by the
Caenorhabditis Genetics Center, which is funded by the NIH Office of Research
Infrastructure Programs [P40 OD010440].

577
578 **Competing interests**
579 The authors declare no competing or financial interests.
580
Author Contributions
Conceptualization: J.M.R., J.D.W., M.T.L.

Methodology: J.M.R., A.T., M.T.L., K.J., A.J., A.A.V., V.T.P., J.D.W.

Validation: J.M.R., A.T., M.T.L., K.J., A.J., A.A.V., V.T.P., J.D.W.

Formal analysis:J.M.R., A.T., M.T.L., K.J., A.J., A.A.V., V.T.P., J.D.W.

Resources: J.M.R., J.D.W.

Data curation: J.M.R., J.D.W.

Writing - original draft: J.M.R., J.D.W.

Writing - review & editing: J.M.R., A.T., M.T.L., K.J., A.J., A.A.V., V.P., J.D.W.

Supervision: J.M.R., J.D.W.

Project administration: J.D.W.

Funding acquisition: J.D.W.

Funding

This work was funded by the National Institutes of Health (NIH) National Institute

of General Medical Sciences (NIGMS) [R00GM107345 and R01 GM138701] to J.D.W.

REFERENCES

1. Li Zheng S, Adams JG, Chisholm AD. Form and function of the apical extracellular matrix:

new insights from *Caenorhabditis elegans, Drosophila melanogaster*, and the vertebrate

inner ear. Fac Rev. 2020;9: 27. doi:10.12703/r/9-27

2. Diller RB, Tabor AJ. The Role of the Extracellular Matrix (ECM) in Wound Healing: A

Review. Biomimetics. 2022;7: 87. doi:10.3390/biomimetics7030087

3. Bonnans C, Chou J, Werb Z. Remodelling the extracellular matrix in development and

disease. Nat Rev Mol Cell Biol. 2014;15: 786–801. doi:10.1038/nrm3904

4. Page AP, Johnstone IL. The cuticle. WormBook Online Rev C Elegans Biol. 2007; 1–15.

doi:10.1895/wormbook.1.138.1

5. Edgar RS, Cox GN, Kusch M, Politz JC. The Cuticle of *Caenorhabditis elegans*. J Nematol.

1982;14: 248–258.

6. Cox GN, Kusch M, Edgar RS. Cuticle of *Caenorhabditis elegans*: its isolation and partial

characterization. J Cell Biol. 1981;90: 7–17. doi:10.1083/jcb.90.1.7

7. Adams JRG, Pooranachithra M, Jyo EM, Zheng SL, Goncharov A, Crew JR, et al.

Nanoscale patterning of collagens in *C. elegans* apical extracellular matrix. Nat Commun.

2023;14: 7506. doi:10.1038/s41467-023-43058-9

8. Ristoratore F, Cermola M, Nola M, Bazzicalupo P, Favre R. Ultrastructural immuno-

localization of CUT-1 and CUT-2 antigenic sites in the cuticles of the nematode

Caenorhabditis elegans. J Submicrosc Cytol Pathol. 1994;26: 437–443.

9.   Peixoto CA, De Souza W. Freeze-fracture and deep-etched view of the cuticle of

Caenorhabditis elegans. Tissue Cell. 1995;27: 561–568. doi:10.1016/s0040-

8166(05)80065-5

10.   Blaxter ML. Cuticle surface proteins of wild type and mutant *Caenorhabditis elegans*. J Biol

Chem. 1993;268: 6600–6609.

11.   Bada Juarez J, O'Rourke D, Judge P, Liu L, Hodgkin J, Watts A. Lipodisqs for eukaryote

lipidomics with retention of viability: Sensitivity and resistance to Leucobacter infection

linked to C.elegans cuticle composition. Chem Phys Lipids. 2019;222. Available:

https://ora.ox.ac.uk/objects/uuid:4d3c3aa1-3f12-443c-8ef0-c3647759832c

12.   Katz SS, Barker TJ, Maul-Newby HM, Sparacio AP, Nguyen KCQ, Maybrun CL, et al. A

transient apical extracellular matrix relays cytoskeletal patterns to shape permanent

acellular ridges on the surface of adult *C. elegans*. PLOS Genet. 2022;18: e1010348.

doi:10.1371/journal.pgen.1010348

13.   Lažetić V, Fay DS. Molting in *C. elegans*. Worm. 2017;6: 1–20.

doi:10.1080/21624054.2017.1330246

14.   Cohen JD, Sundaram MV. *C. elegans* Apical Extracellular Matrices Shape Epithelia. J Dev

Biol. 2020;8: 23. doi:10.3390/jdb8040023

15.   Cohen JD, Flatt KM, Schroeder NE, Sundaram MV. Epithelial Shaping by Diverse Apical

Extracellular Matrices Requires the Nidogen Domain Protein DEX-1 in *Caenorhabditis*

elegans. Genetics. 2019;211: 185–200. doi:10.1534/genetics.118.301752

16.   Gill HK, Cohen JD, Ayala-Figueroa J, Forman-Rubinsky R, Poggioli C, Bickard K, et al.

Integrity of Narrow Epithelial Tubes in the C. elegans Excretory System Requires a

Transient Luminal Matrix. PLoS Genet. 2016;12: e1006205.

doi:10.1371/journal.pgen.1006205

17.   Flatt KM, Beshers C, Unal C, Cohen JD, Sundaram MV, Schroeder NE. Epidermal

Remodeling in *Caenorhabditis elegans* Dauers Requires the Nidogen Domain Protein

DEX-1. Genetics. 2019;211: 169–183. doi:10.1534/genetics.118.301557

18.   Forman-Rubinsky R, Cohen JD, Sundaram MV. Lipocalins Are Required for Apical

Extracellular Matrix Organization and Remodeling in Caenorhabditis elegans. Genetics.

2017;207: 625–642. doi:10.1534/genetics.117.300207

19.   Serra ND, Sundaram MV. Transcytosis in the development and morphogenesis of

epithelial tissues. EMBO J. 2021;n/a: e106163. doi:10.15252/embj.2020106163

20.   Vuong-Brender TTK, Suman SK, Labouesse M. The apical ECM preserves embryonic
integrity and distributes mechanical stress during morphogenesis. Development. 2017;144:
4336–4349. doi:10.1242/dev.150383

21.   Priess JR, Hirsh DI. *Caenorhabditis elegans* morphogenesis: The role of the cytoskeleton
in elongation of the embryo. Developmental Biology. 1986. pp. 156–173.

22.   Costa M, Draper BW, Priess JR. The role of actin filaments in patterning the
Caenorhabditis elegans cuticle. Dev Biol. 1997;184: 373–384. doi:10.1006/dbio.1997.8530

23.   Kelley M, Yochem J, Krieg M, Calixto A, Heiman MG, Kuzmanov A, et al. FBN-1, a fibrillin-
related protein, is required for resistance of the epidermis to mechanical deformation
during *C. elegans* embryogenesis. eLife. 2015;4: 1629. doi:10.7554/eLife.06565

24.   Cohen JD, Sparacio AP, Belfi AC, Forman-Rubinsky R, Hall DH, Maul-Newby H, et al. A
multi-layered and dynamic apical extracellular matrix shapes the vulva lumen in
Caenorhabditis elegans. Shen K, Sengupta P, Sherwood DR, editors. eLife. 2020;9:
e57874. doi:10.7554/eLife.57874

25.   Hendriks G-J, Gaidatzis D, Aeschimann F, Großhans H. Extensive oscillatory gene
expression during *C. elegans* larval development. Mol Cell. 2014;53: 380–392.
doi:10.1016/j.molcel.2013.12.013

26.   Meeuse MW, Hauser YP, Morales Moya LJ, Hendriks G, Eglinger J, Bogaarts G, et al.
Developmental function and state transitions of a gene expression oscillator in
Caenorhabditis elegans. Mol Syst Biol. 2020;16: e9975. doi:10.15252/msb.20209498

27.   Jeon M, Gardner HF, Miller EA, Deshler J, Rougvie AE. Similarity of the *C. elegans*
Developmental Timing Protein LIN-42 to Circadian Rhythm Proteins. Science. 1999;286:
1141–1146. doi:10.1126/science.286.5442.1141

28.   Tennessen JM, Gardner HF, Volk ML, Rougvie AE. Novel heterochronic functions of the
Caenorhabditis elegans period-related protein LIN-42. Dev Biol. 2006;289: 30–43.
doi:10.1016/j.ydbio.2005.09.044

29.   Monsalve GC, Van Buskirk C, Frand AR. LIN-42/PERIOD controls cyclical and
developmental progression of *C. elegans* molts. Curr Biol CB. 2011;21: 2033–2045.
doi:10.1016/j.cub.2011.10.054

30.   Spangler RK, Ashley GE, Braun K, Wruck D, Ramos-Coronado A, Ragle JM, et al. A
conserved chronobiological complex times *C. elegans* development. bioRxiv; 2024. p.
2024.05.09.593322. doi:10.1101/2024.05.09.593322

31.   Hayes GD, Frand AR, Ruvkun G. The *mir-84* and *let-7* paralogous microRNA genes of
Caenorhabditis elegans direct the cessation of molting via the conserved nuclear hormone

receptors NHR-23 and NHR-25. Dev Camb Engl. 2006;133: 4631–4641.

doi:10.1242/dev.02655

32.   Johnson LC, Vo AA, Clancy JC, Myles KM, Pooranachithra M, Aguilera J, et al. NHR-23

activity is necessary for *C. elegans* developmental progression and apical extracellular

matrix structure and function. Dev Camb Engl. 2023;150: dev201085.

doi:10.1242/dev.201085

33.   Kouns NA, Nakielna J, Behensky F, Krause MW, Kostrouch Z, Kostrouchova M. NHR-23

dependent collagen and hedgehog-related genes required for molting. Biochem Biophys

Res Commun. 2011;413: 515–520. doi:10.1016/j.bbrc.2011.08.124

34.   Kostrouchova M, Krause M, Kostrouch Z, Rall JE. Nuclear hormone receptor CHR3 is a

critical regulator of all four larval molts of the nematode *Caenorhabditis elegans*. Proc Natl

Acad Sci U S A. 2001;98: 7360–7365. doi:10.1073/pnas.131171898

35.   Kostrouchova M, Krause M, Kostrouch Z, Rall JE. CHR3: a *Caenorhabditis elegans*

orphan nuclear hormone receptor required for proper epidermal development and molting.

Development. 1998;125: 1617–1626.

36.   Patel R, Galagali H, Kim JK, Frand AR. Feedback between a retinoid-related nuclear

receptor and the *let-7* microRNAs controls the pace and number of molting cycles in *C.*

elegans. eLife. 2022;11: e80010. doi:10.7554/eLife.80010

37.   Davis MW, Birnie AJ, Chan AC, Page AP, Jorgensen EM. A conserved metalloprotease

mediates ecdysis in *Caenorhabditis elegans*. Development. 2004;131: 6001–6008.

doi:10.1242/dev.01454

38.   Hashmi S, Zhang J, Oksov Y, Lustigman S. The *Caenorhabditis elegans* cathepsin Z-like

cysteine protease, Ce-CPZ-1, has a multifunctional role during the worms' development. J

Biol Chem. 2004;279: 6035–6045. doi:10.1074/jbc.M312346200

39.   Kim TH, Kim YJ, Cho JW, Shim J. A novel zinc-carboxypeptidase SURO-1 regulates

cuticle formation and body morphogenesis in *Caenorhabditis elegans*. FEBS Lett.

2011;585: 121–127. doi:10.1016/j.febslet.2010.11.020

40.   Stepek G, McCormack G, Birnie AJ, Page AP. The astacin metalloprotease moulting

enzyme NAS-36 is required for normal cuticle ecdysis in free-living and parasitic

nematodes. Parasitology. 2011;138: 237–248. doi:10.1017/S0031182010001113

41.   Stepek G, McCormack G, Page AP. Collagen processing and cuticle formation is catalysed

by the astacin metalloprotease DPY-31 in free-living and parasitic nematodes. Int J

Parasitol. 2010;40: 533–542. doi:10.1016/j.ijpara.2009.10.007

42.   Birnbaum SK, Cohen JD, Belfi A, Murray JI, Adams JRG, Chisholm AD, et al. The

718   proprotein convertase BLI-4 promotes collagen secretion prior to assembly of the

719   *Caenorhabditis elegans* cuticle. PLOS Genet. 2023;19: e1010944.

720   doi:10.1371/journal.pgen.1010944

721 43. Thacker C, Peters K, Srayko M, Rose AM. The *bli-4* locus of *Caenorhabditis elegans*

722   encodes structurally distinct kex2/subtilisin-like endoproteases essential for early

723   development and adult morphology. Genes Dev. 1995;9: 956–971.

724   doi:10.1101/gad.9.8.956

725 44. Frand AR, Russel S, Ruvkun G. Functional Genomic Analysis of *C. elegans* Molting. PLoS

726   Biol. 2005;3: e312. doi:10.1371/journal.pbio.0030312

727 45. Stepek G, McCormack G, Page AP. The kunitz domain protein BLI-5 plays a functionally

728   conserved role in cuticle formation in a diverse range of nematodes. Mol Biochem

729   Parasitol. 2010;169: 1–11. doi:10.1016/j.molbiopara.2009.08.005

730 46. Gerstein MB, Lu ZJ, Van Nostrand EL, Cheng C, Arshinoff BI, Liu T, et al. Integrative

731   Analysis of the *Caenorhabditis elegans* Genome by the modENCODE Project. Sci N Y NY.

732   2010;330: 1775–1787. doi:10.1126/science.1196914

733 47. Mok DZL, Sternberg PW, Inoue T. Morphologically defined sub-stages of *C. elegans* vulval

734   development in the fourth larval stage. BMC Dev Biol. 2015; 1–8. doi:10.1186/s12861-015-

735   0076-7

736 48. Clancy JC, Vo AA, Myles KM, Levenson MT, Ragle JM, Ward JD. Experimental

737   considerations for study of *C. elegans* lysosomal proteins. G3 GenesGenomesGenetics.

738   2023; jkad032. doi:10.1093/g3journal/jkad032

739 49. Peixoto CA, de Melo JV, Kramer JM, de Souza W. Ultrastructural Analyses of the

740   *Caenorhabditis elegans rol-6 (su1006)* Mutant, Which Produces Abnormal Cuticle

741   Collagen. J Parasitol. 1998;84: 45–49. doi:10.2307/3284528

742 50. Kim TH, Kim DH, Nam HW, Park SY, Shim J, Cho JW. Tyrosylprotein sulfotransferase

743   regulates collagen secretion in *Caenorhabditis elegans*. Mol Cells. 2010;29: 413–418.

744   doi:10.1007/s10059-010-0049-4

745 51. Thein MC, McCormack G, Winter AD, Johnstone IL, Shoemaker CB, Page AP.

746   *Caenorhabditis elegans* exoskeleton collagen COL-19: an adult-specific marker for

747   collagen modification and assembly, and the analysis of organismal morphology. Dev Dyn

748   Off Publ Am Assoc Anat. 2003;226: 523–539. doi:10.1002/dvdy.10259

749 52. Tong A, Lynn G, Ngo V, Wong D, Moseley SL, Ewbank JJ, et al. Negative regulation of

750   *Caenorhabditis elegans* epidermal damage responses by death-associated protein kinase.

751   Proc Natl Acad Sci U S A. 2009;106: 1457–1461. doi:10.1073/pnas.0809339106

53.   Lassandro F, Sebastiano M, Zei F, Bazzicalupo P. The role of dityrosine formation in the
crosslinking of CUT-2, the product of a second cuticlin gene of *Caenorhabditis elegans*.
Mol Biochem Parasitol. 1994;65: 147–159. doi:10.1016/0166-6851(94)90123-6

54.   Cox GN, Staprans S, Edgar RS. The cuticle of *Caenorhabditis elegans*: II. Stage-specific
changes in ultrastructure and protein composition during postembryonic development. Dev
Biol. 1981;86: 456–470. doi:10.1016/0012-1606(81)90204-9

55.   Partridge FA, Tearle AW, Gravato-Nobre MJ, Schafer WR, Hodgkin J. The *C. elegans*
glycosyltransferase BUS-8 has two distinct and essential roles in epidermal
morphogenesis. Dev Biol. 2008;317: 549–559. doi:10.1016/j.ydbio.2008.02.060

56.   Pujol N, Cypowyj S, Ziegler K, Millet A, Astrain A, Goncharov A, et al. Distinct innate
immune responses to infection and wounding in the *C. elegans* epidermis. Curr Biol.
2008;18: 481–489. doi:10.1016/j.cub.2008.02.079

57.   Zugasti O, Ewbank JJ. Neuroimmune regulation of antimicrobial peptide expression by a
noncanonical TGF-β signaling pathway in *Caenorhabditis elegans* epidermis. Nat
Immunol. 2009;10: 249–256. doi:10.1038/ni.1700

58.   Ranasinghe S, McManus DP. Structure and function of invertebrate Kunitz serine protease
inhibitors. Dev Comp Immunol. 2013;39: 219–227. doi:10.1016/j.dci.2012.10.005

59.   Heppert JK, Pani AM, Roberts AM, Dickinson DJ, Goldstein B. A CRISPR Tagging-Based
Screen Reveals Localized Players in Wnt-Directed Asymmetric Cell Division. Genetics.
2018;208: 1147–1164. doi:10.1534/genetics.117.300487

60.   Ruaud A-F, Bessereau J-L. Activation of nicotinic receptors uncouples a developmental
timer from the molting timer in *C. elegans*. Development. 2006;133: 2211–2222.
doi:10.1242/dev.02392

61.   Ascenzi P, Bocedi A, Bolognesi M, Spallarossa A, Coletta M, Cristofaro R, et al. The
Bovine Basic Pancreatic Trypsin Inhibitor (Kunitz Inhibitor): A Milestone Protein. Curr
Protein Pept Sci. 2003;4: 231–251. doi:10.2174/1389203033487180

62.   Smith D, Cwiklinski K, Jewhurst H, Tikhonova IG, Dalton JP. An atypical and functionally
diverse family of Kunitz-type cysteine/serine proteinase inhibitors secreted by the helminth
parasite *Fasciola hepatica*. Sci Rep. 2020;10: 20657. doi:10.1038/s41598-020-77687-7

63.   Thacker C, Srayko M, Rose AM. Mutational analysis of bli-4/kpc-4 reveals critical residues
required for proprotein convertase function in C. elegans. Gene. 2000;252: 15–25.
doi:10.1016/s0378-1119(00)00211-0

64.   Asahina M, Valenta T, Silhánková M, Korinek V, Jindra M. Crosstalk between a nuclear
receptor and beta-catenin signaling decides cell fates in the *C. elegans* somatic gonad.

Dev Cell. 2006;11: 203–211. doi:10.1016/j.devcel.2006.06.003

65. Brenner S. The genetics of *Caenorhabditis elegans*. Genetics. 1974;77: 71–94.

66. Church DL, Guan KL, Lambie EJ. Three genes of the MAP kinase cascade, *mek-2, mpk-*

1/sur-1 and *let-60 ras*, are required for meiotic cell cycle progression in *Caenorhabditis*

elegans. Development. 1995;121: 2525–2535.

67. Peters K, McDowall J, Rose AM. Mutations in the *bli-4* (I) locus of *Caenorhabditis elegans*

disrupt both adult cuticle and early larval development. Genetics. 1991;129: 95–102.

doi:10.1093/genetics/129.1.95

68. Chang W, Tilmann C, Thoemke K, Markussen F-H, Mathies LD, Kimble J, et al. A forkhead

protein controls sexual identity of the *C. elegans* male somatic gonad. Development.

2004;131: 1425–1436. doi:10.1242/dev.01012

69. Frøkjær-Jensen C, Davis MW, Sarov M, Taylor J, Flibotte S, LaBella M, et al. Random and

targeted transgene insertion in *Caenorhabditis elegans* using a modified *Mos1* transposon.

Nat Methods. 2014;11: 529–534. doi:10.1038/nmeth.2889

70. Barkoulas M, van Zon JS, Milloz J, Van Oudenaarden A, Félix M-A. Robustness and

Epistasis in the *C. elegans* Vulval Signaling Network Revealed by Pathway Dosage

Modulation. Dev Cell. 2013;24: 64–75. doi:10.1016/j.devcel.2012.12.001

71. Melo JA, Ruvkun G. Inactivation of conserved *C. elegans* genes engages pathogen- and

xenobiotic-associated defenses. Cell. 2012;149: 452–466. doi:10.1016/j.cell.2012.02.050

72. Au V, Li-Leger E, Raymant G, Flibotte S, Chen G, Martin K, et al. CRISPR/Cas9

Methodology for the Generation of Knockout Deletions in *Caenorhabditis elegans*. G3.

2019;9: 135–144. doi:10.1534/g3.118.200778

73. Espelt MV, Estevez AY, Yin X, Strange K. Oscillatory Ca2+ signaling in the isolated

Caenorhabditis elegans intestine: role of the inositol-1,4,5-trisphosphate receptor and

phospholipases C beta and gamma. J Gen Physiol. 2005;126: 379–392.

doi:10.1085/jgp.200509355

74. Watts JS, Harrison HF, Omi S, Guenthers Q, Dalelio J, Pujol N, et al. New Strains for

Tissue-Specific RNAi Studies in *Caenorhabditis elegans*. G3. 2020;10: 4167–4176.

doi:10.1534/g3.120.401749

75. Nonet ML. Rapid generation of *Caenorhabditis elegans* single-copy transgenes combining

recombination-mediated cassette exchange and drug selection. Genetics. 2023;224:

iyad072. doi:10.1093/genetics/iyad072

76. Nonet ML. Efficient Transgenesis in *Caenorhabditis elegans* Using Flp Recombinase-

Mediated Cassette Exchange. Genetics. 2020;215: 903–921.

doi:10.1534/genetics.120.303388

77.   Ashley GE, Duong T, Levenson MT, Martinez MAQ, Johnson LC, Hibshman JD, et al. An

expanded auxin-inducible degron toolkit for *Caenorhabditis elegans*. Genetics. 2021;217:

iyab006. doi:10.1093/genetics/iyab006

78.   Schwartz ML, Jorgensen EM. SapTrap, a Toolkit for High-Throughput CRISPR/Cas9 Gene

Modification in *Caenorhabditis elegans*. Genetics. 2016;202: 1277–1288.

doi:10.1534/genetics.115.184275

79.   Paix A, Wang Y, Smith HE, Lee C-YS, Calidas D, Lu T, et al. Scalable and versatile

genome editing using linear DNAs with microhomology to Cas9 Sites in *Caenorhabditis*

elegans. Genetics. 2014;198: 1347–1356. doi:10.1534/genetics.114.170423

80.   Paix A, Folkmann A, Rasoloson D, Seydoux G. High Efficiency, Homology-Directed

Genome Editing in *Caenorhabditis elegans* Using CRISPR-Cas9 Ribonucleoprotein

Complexes. Genetics. 2015;201: 47–54. doi:10.1534/genetics.115.179382

81.   El Mouridi S, Peng Y, Frøkjær-Jensen C. Characterizing a strong pan-muscular promoter

(P*mlc-1*) as a fluorescent co-injection marker to select for single-copy insertions.

MicroPublication Biol. 2020;2020. doi:10.17912/micropub.biology.000302

82.   Farboud B, Meyer BJ. Dramatic Enhancement of Genome Editing by CRISPR/Cas9

Through Improved Guide RNA Design. Genetics. 2015;199: 959–971.

doi:10.1534/genetics.115.175166

83.   Ghanta KS, Mello CC. Melting dsDNA Donor Molecules Greatly Improves Precision

Genome Editing in *Caenorhabditis elegans*. Genetics. 2020;216: 643–650.

doi:10.1534/genetics.120.303564

84.   Kamath RS, Fraser AG, Dong Y, Poulin G, Durbin R, Gotta M, et al. Systematic functional

analysis of the *Caenorhabditis elegans* genome using RNAi. Nature. 2003;421: 231–237.

doi:10.1038/nature01278

85.   Moribe H, Yochem J, Yamada H, Tabuse Y, Fujimoto T, Mekada E. Tetraspanin protein

(TSP-15) is required for epidermal integrity in *Caenorhabditis elegans*. J Cell Sci.

2004;117: 5209–5220. doi:10.1242/jcs.01403

86.   Ward JD, Mullaney B, Schiller BJ, He LD, Petnic SE, Couillault C, et al. Defects in the *C.*

elegans acyl-CoA Synthase, *acs-3*, and Nuclear Hormone Receptor, *nhr-25*, Cause

Sensitivity to Distinct, but Overlapping Stresses. PLoS ONE. 2014;9: e92552.

doi:10.1371/journal.pone.0092552.s013