Specific Mitotic Events Drive Cytoskeletal Remodeling Required for Left-Right

Organizer Development

Yan Wu[1,2,&], Yiling Lan[1,2,&], Favour Ononiwu[1,2], Abigail Poole[3], Kirsten Rasmussen[1],

Jonah Da Silva[1], Abdalla Wael Shamil[1], Li-En Jao[4], and Heidi Hehnly[1,2,*]

[1]Department of Biology, Syracuse University, Syracuse, 13244 USA

[2]BioInspired Institute, Syracuse University, Syracuse, 13244 USA

[3]Worcester Polytechnic Institute, Worcester, 01609 MA

[4]Department of Cell Biology and Human Anatomy, University of California Davis

School of Medicine, Sacramento, 95817 USA

[&] Equal contribution

*Lead contact, correspondence: hhehnly@syr.edu, Twitter: @LovelessRadio

https://orcid.org/0000-0001-6660-5254

Summary

Cellular proliferation is essential for tissue development, including the Left-Right

Organizer (LRO), a transient organ vital for establishing the vertebrate LR body plan.

This study examines cell redistribution and progenitor cell roles in LRO formation,

focusing on cell lineage and behavior. Using zebrafish, we mapped mitotic events in

Kupffer's Vesicle (KV), revealing an FGF-dependent, anteriorly enriched mitotic

pattern. With a KV-specific fluorescent microtubule (MT) line, we observed mitotic

spindles aligning along KV's longest axis until the rosette stage, where post-spin

spindles are excluded. Aligned spindles form cytokinetic bridges directing MT

bundles to a junction where a rosette forms. After abscission, repurposed MT

bundles target the rosette center, aiding actin recruitment. Additional cells, both

cytokinetic and non-cytokinetic, are incorporated into the rosette, assembling MT

bundles before actin recruitment. These findings demonstrate the importance of

early divisions for rosette assembly, MT patterning, and actin remodeling during KV

development.

Keywords: Left-Right Organizer, Cell Patterning, Rosette, Lumen Formation, Mitosis,

Cytokinesis, Tight Junctions, Microtubules, Actin, Cilia.

1 **Introduction**

2   The Left-Right Organizer (LRO) is one of the earliest ciliated organs formed

3 during vertebrate development. It is a small and yet important structure—e.g., the

4 node in mice and Kupffer's vesicle (KV) in zebrafish—that only exists transiently

5 during early embryogenesis to help establish the left-right patterning of the body

6 plan. The LRO initiates the development of left-right asymmetry through its cilia-

7 driven, directional fluid flow that induces asymmetric gene expression (summarized

8 in (Grimes & Burdine, 2017; Okabe *et al*, 2008; Dasgupta & Amack, 2016; Forrest *et*

9 *al*, 2022)). Compromised formation of LRO or its ciliary motility (e.g., mutations in the

10 axonemal dynein motor (McGrath *et al*, 2003; Layton, 1976; Lee & Anderson, 2008))

11 would lead to LR patterning defects.

12   Previous studies using the mouse and zebrafish models (**Fig 1A**, reviewed for

13 fish in (Dasgupta & Amack, 2016; Grimes & Burdine, 2017), study in mice (Gredler &

14 Zallen, 2023)) have shown that LRO is formed through the organization of the LRO

15 progenitor cells (e.g., dorsal forerunner cells in zebrafish) into rosette-like structures

16 before displaying epithelial-like characteristics and initiating lumen formation. Using

17 in vitro three-dimensional (3D) cultured Madin-Darby Canine Kidney (MDCK) cells

18 suspended in extracellular matrix (Li *et al*, 2014; Mangan *et al*, 2016; Blasky *et al*,

19 2015; Klinkert *et al*, 2016), lumen formation has been further shown to be preceded

20 by cell division, followed by the establishment of apical polarity and cell-cell contacts

21 adjacent to the cytokinetic bridge. However, those in vitro 3D cell culture models do

22 not recapitulate the mesenchymal to epithelial transition, nor the dynamic changes in

23 cell shape and expression profiles during LRO formation in a living organism. In

24 addition, without the in vivo biological context, different in vitro 3D models often lead

25 to different conclusions. For example, the MDCK (Boehlke *et al*, 2013) and Inner

26 Medullary Collecting Duct (IMCD)(Mahjoub & Stearns, 2012) models showed a

27 different timing of cilia formation. Therefore, despite the insights gained from different

28 in vitro 3D cell culture models, the cellular mechanisms and signals underlying the

29 formation and maintenance of the LRO remain unresolved.

To understand how a functional LRO is formed in a living organism—from a small group of precursor cells to a fluid-filled sac lined with both motile and non-motile ciliated cells (Aljiboury *et al*, 2023; Tavares *et al*, 2017), we studied the development of the KV at single-cell resolution using a series of novel transgenic zebrafish lines and cell biological techniques, including high-resolution live microscopy, lineage tracing, laser ablation, and pharmacological treatments. Our study shows that FGF signaling-mediated early divisions of KV precursor cells are pivotal for proper KV development as the progeny of these early division events constitute a proportion of cells in the mature KV. Our data also suggest that apical microtubule (MT) bundles—originated from the daughter cells of the early KV precursors—facilitate actin recruitment that is crucial for KV development.

Results

Identification of an anteriorly positioned and pre-lumen enriched KV mitotic events and their post-mitotic distribution map.

To follow how the KV is formed at single-cell resolution in a living vertebrate, a Tg(Sox17:GFP-CAAX; h2afx:h2afv-mCherry) was used, where the KV progenitor cells, i.e., the dorsal forerunner cells (DFCs), were labeled by membrane-bound GFP while all histones were labeled by the red fluorescent protein mCherry. We followed the movement and division of DFCs by time-lapse microscopy starting at approximately 8 hours post fertilization (hpf) when 15–30 DFCs have formed (**Fig 1A, 1B, Video S1**). The KV transitions from approximately 15 DFCs that then expand in number and organize into rosette configurations before transitioning to a lumen (**Fig 1A**). Mitotic events predominantly occurred in the anterior region of the future KV before lumen opening (**Fig 1B-C, Fig S1A-B, Video S1**). The same conclusion was also reached when fixed Sox17:GFP-CAAX embryos were analyzed by mitotic cell marker, anti-phospho-histone 3 (PH3) staining at 8 hpf, where we

detected a significant fraction of mitotic events in the anterior zone (**Fig 1D, 1E**) with an enrichment to the left anterior quadrant (**Fig S1C**).

Next, we investigated the distribution of post-mitotic daughter cells from KV precursors in the fully formed KV. For this purpose, embryos were incubated with BrdU from 70% epiboly to 12 hpf to label cells that had undergone cell division before KV completion at 12 hpf. We observed that these BrdU-positive (post-mitotic) cells were distributed throughout the KV, with a significantly higher concentration in the posterior region (**Fig 1D-E, Fig S1D**).

To determine specifically how KV progenitor cells and their progeny reposition following cell division, we performed cell lineage tracing analysis of H2B-Dendra expressing, Sox17:GFP marked KV precursor cells (**Fig 1F-G**). In this experiment, individual H2B-Dendra-positive cells (green) were photoconverted at metaphase and the trajectories of the resulting daughter cells (red) were followed throughout the KV developmental stages until lumen expansion (**Fig 1A, Fig 1F-G**). We found that the daughter cells of a given KV precursor stay together and move within the KV boundary after division (representative events shown in **Fig 1F'** and **Fig S1E**). Among 8 photoconverted metaphase cells and their progeny were followed, all the daughter cells did not divide again during KV development, indicating that at least 50% of KV progenitor cells divide once and only once after 8 hpf (**Fig 1H**). We found that most of the post-mitotic cells were organized in the anterior region of the KV until the rosette stage at the 120 min time point, and then redistributed to the posterior region of the KV during lumen formation (180 min, **Fig 1H, 1I**). When we determined the ratio of anteriorly versus posteriorly positioned post-mitotic cells, this ratio started to approach 1 right before lumen opening and after the cell number in the KV had reached its maximum (**Fig 1I**). Together these results indicate that KV progenitor cells divide preferentially in the anterior region of the future KV, and the resulting post-mitotic cells were incorporated throughout the mature KV with more cells in the posterior than in the anterior regions.

An anterior/posterior FGF signaling gradient is required for anterior zones of

mitotic activity.

Since the early mitotic events of KV precursors took place predominantly

anterior to the location of the future KV (**Fig 1**), we hypothesized that an

anteroposterior signal(s) coordinates the cell division and the later LRO

development. To test this, we individually inhibited the major signaling pathways that

had been implicated in LRO development (Neugebauer *et al*, 2009; Meyers & Martin,

1999; Caron *et al*, 2012) using well-established pharmacological inhibitors, and then

determined whether KV development is affected (**Fig 2A, Fig S2A-E**). We found that

blocking fibroblast growth factor (FGF) signaling, but not Wnt or Sonic hedgehog

(Shh) signaling, attenuated the division of KV precursors through PH3 staining at 8

hpf (**Fig 2B, 2D**) and caused KV cells to not expand in number at pre-lumenal to

lumenal stages (**Fig S2E**). Inhibiting FGF signaling also disrupted spatially concen-

trated mitotic events in the anterior quadrant of the developing KV and caused

severe KV developmental defects (**Fig 2B, 2E-F, S2A-D**). Specifically, only FGF

inhibition demonstrated a significant defect in KV development measured by lumen

area (**Fig 2B, 2F, S2A-C**). These findings suggest that an anteroposterior FGF

signaling gradient is required for the mitotic activity in the anterior region of the

developing KV.

To determine whether loss of the anteriorly biased mitotic events is sufficient

to disrupt proper KV formation at the later stage, we disrupted mitotic events

independent of FGF signaling using pericentrin maternal-zygotic mutant embryos

(*pcnt*[-/-], (Sepulveda *et al*, 2018)) and assessed KV development. Pericentrin is a

centrosomal protein known to regulate spindle assembly, and loss of pericentrin can

result in a decrease in mitotic entry and failure to complete mitosis (Mikule *et al*,

2006; Purohit *et al*, 1999; Chen *et al*, 2014; Delaval & Doxsey, 2010; Zimmerman *et*

al, 2004). We found that the overall cell division (**Fig 2C-D**) and the anterior

enrichment of mitotic events (**Fig 2E**) of KV precursors decreased significantly in

maternal-zygotic *pcnt[-/-]* embryos. This decrease in anteriorly positioned mitotic

events was also associated with a disruption in KV development, as demonstrated

by a significant reduction in the lumen area at 12 hpf (**Fig 2C, 2F**). Together, these

findings suggest that potential anteroposterior FGF signaling regulates the early,

anteriorly biased KV precursor divisions, which are required for subsequent KV

formation.

Early KV mitotic events hold greater significance to KV development

compared to later KV mitotic events.

To determine the role and relevance of individual KV mitotic events, we

employed laser ablation to selectively remove mitotic events (**Fig 3A-B**). Temporal

characterization of mitotic events involved six ablation conditions, encompassing two

control scenarios—no ablation and ablation of non-mitotic cells (3 to 8 cells). The

first experimental ablation comparison was with ablating all mitotic events compared

to no ablation controls (**Fig 3C, 3E, 3G**). Then specific temporal ablation patterns

were performed and compared (**Fig 3D, 3F, 3H**): (1) ablating the first 4 mitotic events

when fewer than 20 KV precursor cells were present (Condition 1); (2) ablating 4

mitotic events when more than 20 KV precursor cells were present (Condition 2);

and (3) ablating all mitotic events when more than 20 KV precursor cells were

present (Condition 3). During video acquisition, mitotic cells were ablated by

positioning a region of interest (ROI) over the metaphase plate, distinguished by

h2afx:h2afv-mCherry, and delivering a 355 nm pulsed laser within the ROI (**Fig 3B**).

Monitoring encompassed all cells, including the cell with the ablated plate, and

revealed no evident apoptotic events in the non-ablated cells. The ablated cells

failed to complete mitosis and were ultimately extruded from the developing KV.

Our studies indicate that the first 4 KV divisions are crucial for KV

development, while later divisions near lumen formation are not essential (**Fig 3C-I**).

KVs did not significantly increase in cell number between 8 hpf and 12 hpf when all

mitotic events were ablated or when all mitotic events were ablated with more than

20 KV precursor cells present (**Fig 3E-F**), suggesting the importance of mitotic events in KV cell number expansion during development. Significant defects in lumen formation kinetics (**Fig 3C-D, 3G-H, Video S2**) and the final lumen area (**Fig 3I**) occurred when all mitotic events or only the first 4 events were ablated (Condition 1), but not when later mitotic events were ablated (Condition 2 and 3), compared to non-mitotic cell ablation control (**Fig 3I**). Embryos subjected to ablation and monitoring for lumen formation kinetics (**Fig 3C-D**) were fixed at 12 hpf and immunostained for cilia. Like lumen formation defects, significant reductions in the percentage of ciliated cells were observed only when ablating all mitotic events or the first 4 events (**Fig 3C-D, 3J**). These findings underscore the pivotal role of early KV mitotic events in orchestrating both KV development and overall KV ciliogenesis.

KV spindles stably align along KVs longest axis until the KV starts rounding, then spindles spin and are extruded.

Proper spindle orientation during mitosis is important for tissue morphogenesis (Bergstralh *et al*, 2017; Vertii *et al*, 2018). Given the significance of early mitotic events in KV development (**Fig 3**), we aimed to understand the relationship between spindle positioning and division timing by analyzing the spindle behaviors in the KV precursor cells over time. To specifically label MTs in KV precursors, we adapted a similar strategy we have used before (Rathbun *et al*, 2020) to generate a Tg(Sox17:EMTB-3xGFP) transgenic line, where the MTs in KV precursors were labeled by GFP tagging the MT-binding domain from the protein ensconsin. This transgenic line allowed us to discern changes in intracellular MT patterning during KV development in live embryos for the first time (**Fig 4A**). Through a series of time-lapse imaging analysis, we found that the spindles of the dividing KV precursors formed and positioned primarily along the long axis of the developing KV in the pre-lumenal stage (**Fig 4A**, 8 min, **Video S3**). As the KV precursors were incorporated to form the rosette structure, the MT bundles started to concentrate at cytokinetic bridges near the rosette center (**Fig 4A**, 97 min, **Video S3**). Strikingly, as

the KV structure went from a structure that had a long and short axis (**Fig 4A**, 8 min) to a structure that was more rounded (**Fig 4A**, 97 to 163 min), the spindles that formed started to rotate or "spin" and were subsequently extruded from the KV (**Fig 4A**, 163 min, **Fig 4B**, **Video S3**). These findings indicate that the rosette was formed by the daughter cells of the early-dividing KV precursors. In contrast, daughter cells from the later cell divisions were excluded from the KV. These results also explain why the later, but not the early, division events are dispensable for KV formation (**Fig 3**). To our knowledge, this is also the first evidence to show that cell extrusion is part of the developmental program of KV formation, akin to the cell extrusion process that has been documented in other developmental contexts such as the generation and maintenance of the tight epithelial barrier (Gudipaty & Rosenblatt, 2017; Marinari *et al*, 2012; Nanavati *et al*, 2020; Eisenhoffer & Rosenblatt, 2013).

To characterize the robustness of these observed events, the aspect ratio (A. R.), defined as the longest axis over the shortest axis of the KV (**Fig 4C**), was measured during KV development (**Fig 4D**) and compared to the degree of spindle angle changes per minute (**Fig 4C, 4E**). Time was normalized to lumen opening so that negative time values are pre-lumen formation and positive values are post-lumen formation (**Fig 4D-4F**). We identified that as the aspect ratio of the KV approached a mean value of less than 2 (dashed line, **Fig 4D**), that spindles started to spin which relates to an increased rate of rotation (angular velocity, **Fig 4E,** direct comparison between A.R. less than 2 and greater than 2 in **4E'**). A population of these mitotic events then become extruded once the lumen starts to open (**Fig 4F,** direct comparison between A.R. less than 2 and greater than 2 in **4F'**). When we ablated the first four mitotic events (condition 1) or ablated all mitotic events post-reaching 20 KV progenitor cells (condition 3, refer to **Fig 3A, 4G, Video S3 and S4**), we observed a loss of post-mitotic extrusion events. Interestingly, even when we only ablated the first 4 mitotic events, no later extrusion events occurred, suggesting that extrusion may be a response to cell packing. This suggests that if early events

are alleviated, later division events can still be incorporated into the KV, but the KV

doesn't develop appropriately (**Fig 3**).

KV progenitor cells position cytokinetic bridges towards tjp1 marked tight

junctions where they maintain cytokinetic bridge derived MT bundles post bridge

cleavage.

The findings in **Fig 4A** demonstrate that cytokinetic bridges are organized

toward the center of a KV cell rosette. However, the coordination of this process with

other temporally regulated cellular events, such as tight junction organization and

apical actin recruitment, remains unclear. To investigate this, we utilized TgKI

(tjp1atdTomato); Sox17:EMTB-3xGFP embryos to label the tight junction marker,

tjp1, at its endogenous locus (Levic *et al*, 2021) while simultaneously marking MTs in

KV cells. In these embryos, tjp1-labeled tight junctions were assembled prior to

division events. Subsequently, early KV divisions positioned their cytokinetic bridges

toward a tjp1-labeled junction (**Fig 5A, 5B**), where MT bundles derived from the

cytokinetic bridge were maintained post-bridge cleavage (**Fig 5C**, arrow at 2m

indicating midbody loss). This finding contrasts with studies in human cells grown in

culture, which do not assemble into 3D structures, where MT enrichment at bridges

diminishes as abscission completes (Steigemann & Gerlich, 2009; Steigemann *et al*,

2009; Guizetti *et al*, 2011; Addi *et al*, 2018). This suggests that the assembly of a 3D

tissue may impose additional requirements on these MT-based structures as

modeled in **Fig 5D**.

Converging cytokinetic cells initiate actin recruitment at forming rosette

centers.

To examine the temporal and spatial dynamics of actin in relation to MTs, we

injected Sox17:EMTB-3xGFP embryos with Lifeact-mRuby mRNA (**Fig 6A**). When

imaging KV conditions with fewer than 20 cells, the first cytokinetic bridge was found

to accumulate actin (magenta arrow, **Fig 6B**, **Video S5**) and serves as the initial site

for rosette formation. A second cytokinetic bridge, also recruiting actin, can move

towards the rosette, allowing the daughter cells connected by this bridge to integrate

into the developing structure (blue arrow, **Fig. 6B**). Surrounding cells, which are not

connected by cytokinetic bridges, can then assemble into the rosette, organizing MT

bundles toward its center (non-division-derived MT bundle, NDMT, **Fig 6B**, modeled

in **Fig 6C**).

To determine whether the initial mitotic events are required for actin

recruitment to rosette centers, we ablated the first four divisions (condition 1, **Fig 3A**)

and compared them to control, non-ablated conditions (**Fig 6C**). Under ablated

conditions, actin was unable to accumulate at rosette centers that had accumulated

non-division-derived MT bundles (NDMT, **Fig 6D**, right) compared to controls (**Fig**

6D, left). Actin recruitment to MT bundles across three conditions: 1) cytokinetic

bridge-derived MT bundles (CT-CB), 2) non-division-derived MT bundles (CT-

NDMT), and 3) NDMT from condition 1 ablated cells (Cond.1-NDMT). While non-

division-derived MT bundles (CT-NDMT) can recruit actin (**Fig 6F**), the recruitment is

significantly less compared to cytokinetic bridge-derived MT bundles over a 10-

minute period (CT-CB, **Fig 6F, 6G**). However, ablating the first four mitotic cells does

not significantly decrease the ability of non-division-derived MT bundles (Cond.1-

NDMT) to recruit actin (**Fig 6E-G**). This suggests that while the properties of

cytokinetic bridge-derived MT bundles are associated with increased actin

recruitment compared to non-division-associated MT bundles, post-mitotic cells do

not influence the ability of neighboring rosette cells with non-division-derived MT

bundles to assemble actin.

Cytokinetic bridge derived MT bundles are required for actin recruitment at

rosette during lumen formation.

To further investigate whether cytokinetic bridge-derived MT bundles are

necessary for actin recruitment during the rosette stage, we conducted targeted

ablations on cytokinetic bridges and non-division-derived MT (NDMT) bundles

prior to the expected time of actin recruitment, then measured actin accumulation

and MT recovery (**Fig 7A**). For the cytokinetic bridge ablations, we ablated one

side of the bridge adjacent to where the midbody should form (CB-A), while

leaving the other side unablated (CB-NA) (**Fig 7A, 7B**). We found that actin

recruitment on the ablated side was significantly diminished over a 10-minute

period, while the non-ablated side retained its ability to recruit actin (**Fig 7B, 7E,**

S3B). When comparing the ability of MTs to recover after cytokinetic bridge

ablation (CB-A) to ablated non-division-derived MT bundles (AB-NDMT), we found

that MTs were unable to recover under CB-A conditions, whereas they did recover

under AB-NDMT conditions (**Fig 7B-7D, 7F, S3C**). This suggests that the

properties of MT bundles in the rosette differ between those derived from

cytokinetic bridges and those originating from non-mitotic cells. When examining

whether ablated NDMT bundles could recruit actin after 10 minutes, we found they

were unable to do so compared to control non-ablated NDMT conditions (CT-

NDMT, **Fig 7G, S3D**). These findings suggest that while the ability of MTs to

recover varies between cytokinetic bridge-derived MT bundles and non-division-

derived MT bundles (**Fig 7F**), both MT bundled structures are necessary for actin

accumulation (**Fig 7E, 7G**), with cytokinetic bridge-derived bundles recruiting actin

to a greater extent (**Fig 6F, G**).

Discussion

Studies of LRO formation have elucidated the fundamental role of LRO

progenitors in forming the rosette-like structure with epithelial-like characteristics

before the initiation of lumen formation (Gredler & Zallen, 2023; Aljiboury *et al*, 2023;

Essner *et al*, 2005). However, the steps preceding lumen formation is not clear,

especially in the context of a living embryo. By studying KV formation in zebrafish

embryos with novel transgenic and cell biological tools, our results suggest a model

in which FGF-mediated signals induce the proliferation of a subset of early KV

precursor cells, predominantly located anteriorly to the developing KV. Their progeny

is then incorporated into the rosette before lumen formation, whereas the progeny

from the later cell divisions were excluded (**Fig 8**). How FGF signaling coordinates

the proliferation of the early KV precursors is an important question that awaits future

studies.

We find that the apical microtubule (MT) bundles, which originate from the

daughter cells of early Kupffer's Vesicle (KV) precursors, persist after cytokinesis

and are oriented towards regions marked by tjp1-labeled tight junctions (**Fig 8**).

These apical MT bundles are crucial for recruiting actin to these sites. Actin

recruitment has shown to be a process essential for apical membrane formation and

lumen development, not only in KV but also in other systems such as mouse node

formation (Gredler & Zallen, 2023) and pluripotent stem cell-derived epiblasts

(Indana *et al*, 2024). For instance, in pluripotent stem cell-derived epiblasts, apical

actin expansion is required for lumen formation, with subsequent lumen expansion

driven by osmotic pressure-induced fluid transfer (Indana *et al*, 2024). Under

conditions of early mitotic ablation in KV where there is a significant reduction of

actin to forming rosette structures, we observe that although a lumen forms, its size

is significantly reduced (**Fig 3**). One possible explanation is that defects in tight

junction assembly may lead to fluid leakage (Kuo *et al*, 2022; Odenwald *et al*, 2017;

Fanning *et al*, 2012; Indana *et al*, 2024; Mukenhirn *et al*, 2024). However, our

examination of lumen morphology in early mitotic-ablated KVs did not reveal the

folded apical membrane appearance characteristic of such defects (Mukenhirn *et al*,

2024). This suggests that tight junction protein recruitment at the apical membrane in

early mitotic-ablated KVs may be functional and formation occurs independently of

mitotic events during lumen development. This is consistent with our observation that

small tjp1-labeled tight junction structures are present before early mitotic events,

with cytokinetic bridges pinching towards these structures (**Fig 8**). Nonetheless, the

question remains whether osmolarity-driven fluid flow is sufficient to drive lumen

opening under actin-deficient conditions and how lumen formation is initiated in such

contexts.

1   In summary, our analysis of KV organization offers new insights into how

2   spindle orientation, followed by mitotic exit driving MT bundle formation, contributes

3   to actin recruitment during rosette formation, essential for KV development.

Acknowledgements

We thank SU's BioImaging Core for their assistance and In Vivo Biosystems for helping us create the Sox17:EMTB-3xGFP line. This work was supported by National Institutes of Health Grants no. R01GM-127621 (H.H.), no. R01GM-130874 (H.H.), and no. OD026946 (SU BioImaging Center) for the Zeiss LSM980 with Airyscan2.

Disclosure and competing interest statement

The authors declare no competing interests.

Supplemental Information:

Video S1. *Identification of an anteriorly positioned and pre-lumen enriched Kupffer's Vesicle (KV) mitotic events and their post-mitotic distribution map.* Related to Fig 1B.

Video S2. *Early KV mitotic events hold greater significance to KV development compared to later KV mitotic events.* Related to Fig 3C, D.

Video S3. *Spindles stably align along KVs longest axis until the KV starts rounding, then spindles spin and are extruded.* Related to Fig 4A, B.

Video S4. *KV spindles stably align along KVs longest axis until the KV starts rounding, then spindles spin and are extruded.* Related to Fig 4G.

Video S5. *Converging cytokinetic cells initiate actin recruitment at forming rosette centers.* Related to Fig 6A-6B.

Resource Availability

Lead contact: For further information or to request resources/reagents, contact Lead Contact, Dr. Heidi Hehnly (hhehnly@syr.edu)

Materials availability: New materials generated for this study are available for

distribution.

Data and code availability: All data sets analyzed for this study are displayed.

Experimental model and subject details

Fish Lines

Zebrafish lines were maintained in accordance with protocols approved by the

Institutional Animal Care Committee of Syracuse University (IACUC Protocol #18-

006). Embryos were raised at 28.5^{o}C and staged (as described in (Kimmel *et al*,

1995)). Wildtype and/or transgenic zebrafish lines used for live imaging and

immunohistochemistry are listed in Key Resource Table.

Method Details

Chemical inhibitors

Dechorionated live embryos were treated with 10 µM XAV939 (Sigma-Aldrich), 0.5 to

5 µM Infigratinib (MedChem Express), and 50 µM cyclopamine (Sigma-Aldrich) were

prepared. Tg(sox17:GFP-CAAX) transgenic zebrafish embryos were dechorionated

on 2% agarose pad dishes and at 50% epiboly incubated with vehicle control (0.4%

DMSO, 0.5 % Methanol) and in listed concentrations of inhibitor diluted in zebrafish

embryo water (0.03% sea salt, 1*10^{-5}% Methylene Blue, RO water) and also taking

into account the volume of agarose pad. Embryos were incubated to 8 hours post

fertilization (hpf) and 12 hpf and fixed in 4% paraformaldehyde (PFA) (refer to

immunofluorescence section).

BrdU Incorporation

Tg(sox17:GFP-CAAX) embryos were dechorionated at 50% epiboly, followed by

incubation in E3 buffer (5mM NaCl,0.17mM KCl, 0.33mM $CaCl_2$,0.33mM $MgSO_4$) for

15 minutes at room temperature. Embryos were then incubated in 10 mM BrdU in E3

buffer containing 12% DMSO and maintained at temperatures between 28.5 and

30°C until 10 hpf. Following serial dilution washes in E3 medium, the embryos were

incubated until 12 hpf and fixed in 4% PFA and overnight dehydration in cold

methanol at -20°C. Embryos then underwent a sequential rehydration process using

methanol to PBST mixture ratios of 3:1, 1:1, and 1:3 for 5 minutes each step.

Embryos were refixed for 15 minutes in 4% PFA with 0.5% Triton-X 100, followed by

acid treatment with 2N HCl for 1 hour and subsequent neutralization in 0.1M borate

buffer (0.1M Boric Acid, 0.076M NaOH) for 20 minutes. Then standard

immunofluorescent approaches were used to label for BrdU (anti-BrdU, mouse,

1:100, Sigma-Aldrich,11170376001: BMC9318). See immunofluorescence section

and Key Resource Table.

Plasmids and mRNA for injection experiments

Plasmids were generated using Gibson cloning methods (NEBuilder HiFi DNA

assembly Cloning Kit) and maxi-prepped before injection and/or transfection. mRNA

was made using mMESSAGE mMACHINE™SP6 transcription kit. See key resource

table for a list of plasmid constructs and mRNA used. Injections of 1 cell staged

embryos were performed as described in (Aljiboury *et al*, 2021).

Immunofluorescence

Immunostaining for phospho-histone H3 (PH3), acetylated tubulin and GFP,

Tg(sox17:GFP-CAAX) transgenic zebrafish embryos were fixed at 8 and 12 hours

post fertilization (hpf) using 4% PFA with 0.5% Triton X-100 overnight at 4°C.

Embryos were dechorionated after washing with PBST (0.1% Tween-20 in

Phosphate Buffered Saline (PBS)) 3 times. Embryos were blocked in wash solution

(1%DMSO,1% BSA,0.1%Triton-X) for 1 hour at room temperature with gentile

agitation. Primary antibody incubation (diluted in wash solution) occurs overnight at

4°C. Primary antibodies used include: anti-Phospho-Histone H3 (Rabbit) antibody

(1:200, Cell Signaling Technology, 9701S), Anti-GFP (Chicken) (1:300, GeneTex,

GTX13970: AB_371416) or Anti-Acetylated Tubulin (Mouse) (1:300, Sigma

Aldrich,T6793: RRID: AB_477585), Anti-GFP (Rabbit) (1:300, Molecular Probes, A-

11122: AB_221569), refer to key resource table. Embryos were then washed and

incubated with secondary antibodies for 2-4 hours at room temperatures or overnight

at 4°C. Secondary antibodies used include: Alexa Fluor Anti-Mouse 568 (1:300, Life

Technologies, A10037; RRID: AB_2534013), Alexa Fluor Anti-Chicken 488 (1:300,

Fisher scientific, A11039), or Alexa Fluor Anti-Mouse 647 (1:300, Life Technologies,

A31571; RRID: AB_162542), Alexa Fluor Anti-Rabbit 488 (1:300, Life Technologies,

A21206; RRID: AB_2535792). Embryos were stained with DAPI (1 µg/mL) to label

nuclei after 3 times washes with wash solution. Embryos were mounted by 2%

agarose after washing with PBS. Refer to Key resource table.

Imaging

Fixed or live dechorionated embryos are embedded in low-melting 1.5% agarose

(Key resource table) with the KV positioned at the bottom of a #1.5 glass bottom

MatTek plate (Key resource table) and imaged using a spinning disk confocal

microscope or laser scanning confocal microscope. Zebrafish embryos were imaged

using Leica DMi8 (Leica, Bannockburn, IL) equipped with a X-light V2 Confocal unit

spinning disk equipped with a Visitron VisiFRAP-DC photokinetics attached to 405

and 355nm lasers, a Leica SP8 (Leica, Bannockburn, IL) laser scanner confocal

microscope (LSCM) and/or a Zeiss LSCM 980 (Carl Zeiss, Germany) with an

Airyscan 2 detector. The Leica DMi8 is equipped with a Lumencore SPECTRA X

(Lumencore, Beaverton, OR), Photometrics Prime-95B sCMOS Camera, and 89

North-LDi laser launch. VisiView software was used to acquire images. Optics used

with this unit are HC PL APO x40/1.10W CORR CS2 0.65 water immersion

objective, HC PL APO x40/0.95 NA CORR dry and HCX PL APO x63/1.40-0.06 NA

oil objective. The SP8 laser scanning confocal microscope is equipped with HC PL

APO 20x/0.75 IMM CORR CS2 objective, HC PL APO 40x/1.10 W CORR CS2 0.65

water objective and HC PL APO x63/1.3 Glyc CORR CS2 glycerol objective. LAS-X

software was used to acquire images. The Zeiss LSM 980 is equipped with a T-PMT,

GaASP detector, MA-PMT, Airyscan 2 multiplex with 4Y and 8Y. Optics used with

this unit are PL APO x63/1.4 NA oil DIC. Zeiss Zen 3.2 was used to acquire the

images. A Leica M165 FC stereomicroscope equipped with DFC 9000 GT sCMOS

camera was used for staging and phenotypic analysis of zebrafish embryos.

Photoconversion

Tg(sox17:GFP) embryos were injected with 75ng of Dendra-H2B mRNA (see

methods in (Aljiboury *et al*, 2021)). The embryos were incubated at 30°C until 8 hpf,

dechorionated and embedded in 1.5% agarose. To photoconvert H2B-Dendra,

405 nm laser beam was applied to a Region of Interest (ROI) placed over a mitotic

plane. The 405 nm laser was used at 8 mW power with 2 FRAP cycles at 50 ms/pixel

dwell time. Time-lapse videos were captured using 470 nm and 555 nm lasers to

capture green and red emission from non-photoconverted and photoconverted

respectively. Z-stacks were acquired at z-stack with a 2 μm step size every 2 minutes

until 12 hpf.

Laser Ablation

Tg(sox17:GFP-CAAX; h2afx:h2afv-mCherry) or Tg(sox17:EMTB-3XGFP)

dechorionated live embryos were imaged on a spinning disk with VisiView kinetics

unit starting at 6 hpf.

Mitotic ablation: Time-lapse videos capturing GFP-CAAX and h2afx:h2afv-mCherry

used 470 nm and 555 nm lasers respectively across a z-stack with a 2 μm step size

every 2 minutes until 12 hpf. KV mitotic events were ablated using a 355 nm pulsed

laser operating at 65% power. The laser was applied within a ROI over mitotic cell

for 2 cycles of 50 ms/pixel. Non-mitotic ablation control involved ablating a region

containing 3-8 interphase cells of the KV. Ablated embryos were fixed at 1hpf and

stained for cilia (acetylated tubulin).

MT bundles ablation: Time-lapse videos capturing EMTB-3XGFP and Lifeact-mRuby

used 470 nm and 555 nm lasers, respectively, across a z-stack with a 2 μm step size

every 2 minutes until 12 hpf. MT bundles (cytokinetic bridge-derived and non-

division-derived bridge) were ablated using a 355 nm pulsed laser operating at 65%

power. The laser was applied within an ROI over the proximal end of the bundles for

1-3 cycles of 50 ms/pixel until a break in the bundle was observed live.

Image and data analysis

Images were processed using FIJI/ImageJ. Graphs and statistical analysis were

produced using Prism 9 software. Additional analysis of KV cells were performed

using Bitplane IMARIS software. Videos were created using FIJI/ImageJ or IMARIS.

All time-lapse video projections are registered using FIJI. For percentage of ciliated

KV cells, the number of cells with cilia was counted and represented as a percentage

over the total number of cells in the cyst forming tissue.

Mitotic index and Anterior vs Posterior mitotic event calculation. For mitotic index,

the number of mitotic cells (PH3 positive) in the KV was divided by the total number

of KV cells (DAPI and Sox17:GFP-CAAX positive), resulting in a percentage of

mitotic cells out of the entire population. To calculate fraction of mitotic events in the

anterior versus posterior regions, anterior and posterior KV mitotic cell number was

divided by the total number of mitotic cells in the KV.

KV Aspect Ratio. KV aspect ratio was calculated from a video projection of

Sox17:EMTB-3xGFP embryos from 7 to 12 hpf. Images were acquired at maximum

2-to-3-minute intervals. For each captured frame, the aspect ratio was calculated

using FIJI/ImageJ. Longest axis over shortest axis was determined giving the aspect

ratio over time. Data was presented by averaging aspect ratios across 10 bins for

the duration of KV development.

Angular Velocity of the Spindle. The metaphase derived spindle angle relative to the

KV's longest axis was measured over time. To do this, a line was positioned along

the longest KV axis in relation to a second line that passes through the two spindle

poles and the longest axis of a KV mitotic spindle. The ImageJ/FIJI angle tool

calculated the angle at which the two lines intersect providing a spindle angle

($\angle$Spindle). This spindle angle was calculated from metaphase to anaphase

completion. The angular velocity was measured by taking the difference of identified

spindle angle across consecutive time point 1 ($\angle$Spindle$_{T1}$) and time point 2

($\angle$Spindle$_{T2}$) and dividing the difference by time interval (T_1-T_2).

$$Angular\ Velocity = \left| \frac{\angle\text{Spindle}_{T1} - \angle\text{Spindle}_{T2}}{T_1 - T_2} \right|$$

Data was presented by averaging spindle angular velocity from each mitotic event

across 20 min bins for the duration of KV development between 7 to 12 hpf.

Lumen area.

A region was drawn around the lumen perimeter and area calculated using the

measure function. Where applicable, values were normalized to the control mean by

dividing each lumen area by the mean value of the control lumens.

Actin intensity measurement at nascent MT bundles.

KV cells of interest were masked across Z slices at five different time points using

Imaris Software. A square ROI with constant size across all samples was placed at a

newly formed MT bundle or MT bundle ablation site, labeled with EMTB-3xGFP, to

quantify actin (Lifeact-mRuby) from max-projected datasets over time using the ROI

manager in FIJI/ImageJ. Data normalization was performed by dividing the

fluorescence intensity at each time point by the mean fluorescence intensity at the

final time point for all measured bridges or intensity at the first time point after

subtracting background intensity.

Statistical Analysis

PRISM9 software was used for all graph preparations that include all individual data

points across embryos and clutches denoted by color and size of points respectively

and as noted in legends. These plots were presented as violin, floating bar, and a

modified scatter plot that is termed "super-plot" (Lord *et al*, 2020) to denote individual

embryos across clutches. Unpaired two-tailed t-tests for pairwise numerical

differences in mean and one way ANOVA were performed when comparing greater

than two groups using PRISM9 software. **** denotes a p-value<0.0001, *** p-

value<0.001, **p-value<0.01, *p-value<0.05, ns, not significant. For further

information on detailed statistical analysis see **Table S1**.

Key resources table

REAGENT or RESOURCE	SOURCE	IDENTIFIER
Antibodies		
Acetylated Tubulin	Sigma Aldrich	T6793: RRID: AB_477585
Anti-GFP (Chicken)	GeneTex	GTX13970: AB_371416
Anti-GFP (Rabbit)	Molecular Probes	A-11122: AB_221569
Anti-PH3 (Rabbit)	Cell Signaling Technology	9701S
Anti-BrdU (Mouse)	Sigma-Aldrich	11170376001: BMC9318
Alexa Fluor Anti-Rabbit 488	Life Technologies	A21206; RRID: AB_2535792
Alexa Fluor Anti-Rabbit 568	Life Technologies	A10042; RRID: AB_2534017
Alexa Fluor Anti-Chicken 488	Fisher scientific	A11039
Alexa Fluor Anti-Mouse 568	Life Technologies	A10037; RRID: AB_2534013

Alexa Fluor Anti-Mouse 647	Life Technologies	A31571; RRID: AB_162542
Alexa Fluor® 647 Phalloidin	Cell Signaling Technology	8940S
Bacterial and virus strains		
N/A		
Biological samples		
N/A		
Chemicals, peptides, and recombinant proteins		
DAPI	Sigma Aldrich	D9542-10mg
Agarose	Thermo Fisher	16520100
BSA	Fisher Scientific	BP1600-100
Dimethylsulphoxide	Fisher Scientific	BP231-100
Paraformaldehyde	Fisher Scientific	O4042-500
Phosphate Buffered Saline	Fisher Scientific	10010023
Triton X-100	Fisher Scientific	BP151500
Tween 20	Thermo Fisher	BP337500
Sodium Chloride	Fisher Scientific	BP358
5-Bromo-2'-deoxyuridine, ≥99% (HPLC)	Sigma-Aldrich	Product Number : B5002 CAS-No. : 59-14-3
XAV939	Sigma-Aldrich	Product Number : X3004 CAS-No. : 284028-89-3
Infigratinib	MedChem Express	Catalog No. : HY-13311 CAS No. : 872511-34-7
Cyclopamine	Sigma-Aldrich	Product Number : PHL82510 CAS-No. : 4449-51-8
Critical commercial assays		
BIO BASIC Maxi Prep Kit	BIO BASIC	9K-0060023
NEBuilder HiFi DNA assembly Cloning Kit	New England BioLabs	E5520S
mMESSAGE mMACHINETMSP6	Invitrogen	AM1340
Deposited data		
N/A		
Experimental models: Cell lines		
N/A		
Experimental models: Organisms/strains		

Zebrafish	Zebrafish International Resource Center	AB-Wildtype
Zebrafish	Heidi Hehnly Lab	Sox17:GFP-CAAX; h2afx:h2afv-mCherry
Zebrafish	Jeffrey Amack Lab	Tg (sox17:GFP-CAAX)sny101
Zebrafish	Li-En Jao Lab	PCNT (-/-)
Zebrafish	Zebrafish International Resource Center	Tg(sox17:GFP)
Zebrafish	Heidi Hehnly lab	Tg(sox17:EMTB-3×GFP)
Zebrafish	Heidi Hehnly lab	sox17:GFP-CAAX;PCNT(-/-)
Zebrafish	Heidi Hehnly lab Michel Bagnat lab	Sox17:GFP-CAAX; Tjp1atdTomato
Oligonucleotides		
H2B-Dendra (mRNA)	Heidi Hehnly lab	Plasmid: pCS2-H2B-Dendra
Lifeact-mRuby (mRNA)	Heidi Hehnly lab	Plasmid: pCS2-Lifeact-mRuby
mKate-MKLP1 (mRNA)	Heidi Hehnly lab	Plasmid: pCS2-mKate-MKLP1
Recombinant DNA		
Plasmid: pCS2-H2B-Dendra	Addgene	Plasmid #140559
Plasmid: pCS2-Lifeact-mRuby	Addgene	Plasmid #194339
Plasmid: pCS2-mKate-MKLP1	Addgene	Plasmid #140570
Software and algorithms		
ImageJ/FIJI	NIH and Laboratory for Optical and Computational Instrumentation	https://imagej.net/Fiji
IMARIS, Bitplane	Oxford Instruments	https://imaris.oxinst.com/
PRISM9	GraphPad	https://www.graphpad.com/scientific-software/prism/

LAS-X Software	Leica Microsystems	https://www.leica-microsystems.com/products/microscope-software/p/leica-las-x-ls/
VisiView	Visitron	https://www.visitron.de/products/visiviewr-software.html
Other		
N/A		

References Cited

Addi C, Bai J & Echard A (2018) Actin, microtubule, septin and ESCRT filament remodeling during late steps of cytokinesis. *Curr Opin Cell Biol* 50: 27–34

Aljiboury AA, Ingram E, Krishnan N, Ononiwu F, Pal D, Manikas J, Taveras C, Hall NA, Da Silva J, Freshour J, *et al* (2023) Rab8, Rab11, and Rab35 coordinate lumen and cilia formation during zebrafish left-right organizer development. *PLoS Genet* 19: e1010765

Aljiboury AA, Mujcic A, Cammerino T, Rathbun LI & Hehnly H (2021) Imaging the early zebrafish embryo centrosomes following injection of small-molecule inhibitors to understand spindle formation. *STAR Protoc* 2: 100293

Bergstralh DT, Dawney NS & St Johnston D (2017) Spindle orientation: a question of complex positioning. *Development* 144: 1137–1145

Blasky AJ, Mangan A & Prekeris R (2015) Polarized protein transport and lumen formation during epithelial tissue morphogenesis. *Annu Rev Cell Dev Biol* 31: 575–591

Boehlke C, Kotsis F, Buchholz B, Powelske C, Eckardt KU, Walz G, Nitschke R & Kuehn EW (2013) Kif3a guides microtubular dynamics, migration and lumen formation of MDCK cells. *PLoS One* 8

Caron A, Xu X & Lin X (2012) Wnt/β-catenin signaling directly regulates Foxj1 expression and ciliogenesis in zebrafish Kupffer's vesicle. *Development* 139: 514–524

Chen C-TT, Hehnly H, Yu Q, Farkas D, Zheng G, Redick SD, Hung H-FF, Samtani R, Jurczyk A, Akbarian S, *et al* (2014) A Unique Set of Centrosome Proteins Requires Pericentrin for Spindle-Pole Localization and Spindle Orientation. *Current Biology* 24: 2327–2334

Dasgupta A & Amack JD (2016) Cilia in vertebrate left-right patterning. *Philos Trans R Soc Lond B Biol Sci* 371

Delaval B & Doxsey SJ (2010) Pericentrin in cellular function and disease. *J Cell Biol* 188: 181–190

Eisenhoffer GT & Rosenblatt J (2013) Bringing balance by force: live cell extrusion controls epithelial cell numbers. *Trends Cell Biol* 23: 185–192

Essner JJ, Amack JD, Nyholm MK, Harris EB & Yost HJ (2005) Kupffer's vesicle is a ciliated organ of asymmetry in the zebrafish embryo that initiates left-right development of the brain, heart and gut. *Development* 132: 1247–1260

Fanning AS, Van Itallie CM & Anderson JM (2012) Zonula occludens-1 and -2 regulate apical cell structure and the zonula adherens cytoskeleton in polarized epithelia. *Mol Biol Cell* 23: 577–590

Forrest K, Barricella AC, Pohar SA, Hinman AM & Amack JD (2022) Understanding laterality disorders and the left-right organizer: Insights from zebrafish. *Front Cell Dev Biol* 10

Gredler ML & Zallen JA (2023) Multicellular rosettes link mesenchymal-epithelial
transition to radial intercalation in the mouse axial mesoderm. *Dev Cell* 58: 933-
950.e5
Grimes DT & Burdine RD (2017) Left–Right Patterning: Breaking Symmetry to
Asymmetric Morphogenesis. *Trends in Genetics* 33: 616–628
Gudipaty SA & Rosenblatt J (2017) Epithelial Cell Extrusion: pathways and
pathologies. *Semin Cell Dev Biol* 67: 132
Guizetti J, Schermelleh L, Mäntler J, Maar S, Poser I, Leonhardt H, Müller-Reichert T
& Gerlich DW (2011) Cortical constriction during abscission involves helices of
ESCRT-III-dependent filaments. *Science (1979)* 331: 1616–1620
Indana D, Zakharov A, Lim Y, Dunn AR, Bhutani N, Shenoy VB & Chaudhuri O
(2024) Lumen expansion is initially driven by apical actin polymerization
followed by osmotic pressure in a human epiblast model. *Cell Stem Cell* 31:
640-656.e8
Kimmel CB, Ballard WW, Kimmel SR, Ullmann B & Schilling TF (1995) Stages of
embryonic development of the zebrafish. *Developmental dynamics : an official
public* 203: 253–310
Klinkert K, Rocancourt M, Houdusse A & Echard A (2016) Rab35 GTPase couples
cell division with initiation of epithelial apico-basal polarity and lumen opening.
Nat Commun 7
Kuo WT, Odenwald MA, Turner JR & Zuo L (2022) Tight junction proteins occludin
and ZO-1 as regulators of epithelial proliferation and survival. *Ann N Y Acad Sci*
1514: 21–33
Layton WM (1976) Random determination of a developmental process: reversal of
normal visceral asymmetry in the mouse. *J Hered* 67: 336–338
Lee JD & Anderson K V. (2008) Morphogenesis of the node and notochord: the
cellular basis for the establishment and maintenance of left-right asymmetry in
the mouse. *Dev Dyn* 237: 3464–3476
Levic DS, Yamaguchi N, Wang S, Knaut H & Bagnat M (2021) Knock-in tagging in
zebrafish facilitated by insertion into non-coding regions. *Development* 148
Li D, Mangan A, Cicchini L, Margolis B & Prekeris R (2014) FIP5 phosphorylation
during mitosis regulates apical trafficking and lumenogenesis. *EMBO Rep* 15:
428–437
Lord SJ, Velle KB, Dyche Mullins R & Fritz-Laylin LK (2020) SuperPlots:
Communicating reproducibility and variability in cell biology. *Journal of Cell
Biology* 219
Mahjoub MR & Stearns T (2012) Supernumerary Centrosomes Nucleate Extra Cilia
and Compromise Primary Cilium Signaling. *Current Biology* 22: 1628–1634
Mangan AJ, Sietsema D V, Li D, Moore JK, Citi S & Prekeris R (2016) Cingulin and
actin mediate midbody-dependent apical lumen formation during polarization of
epithelial cells. *Nat Commun* 7: 12426

Marinari E, Mehonic A, Curran S, Gale J, Duke T & Baum B (2012) Live-cell
delamination counterbalances epithelial growth to limit tissue overcrowding.
Nature 484: 542–545
McGrath J, Somlo S, Makova S, Tian X & Brueckner M (2003) Two populations of
node monocilia initiate left-right asymmetry in the mouse. *Cell* 114: 61–73
Meyers EN & Martin GR (1999) Differences in left-right axis pathways in mouse and
chick: functions of FGF8 and SHH. *Science* 285: 403–406
Mikule K, Delaval B, Kaldis P, Jurcyzk A, Hergert P & Doxsey S (2006) Loss of
centrosome integrity induces p38—p53—p21-dependent G1—S arrest. *Nature*
Cell Biology 2006 9:2 9: 160–170
Mukenhirn M, Wang C-H, Guyomar T, Bovyn MJ, Staddon MF, van der Veen RE,
Maraspini R, Lu L, Martin-Lemaitre C, Sano M, *et al* (2024) Tight junctions
control lumen morphology via hydrostatic pressure and junctional tension. *Dev*
Cell
Nanavati BN, Yap AS & Teo JL (2020) Symmetry Breaking and Epithelial Cell
Extrusion. *Cells* 9
Neugebauer JM, Amack JD, Peterson AG, Bisgrove BW & Yost HJ (2009) FGF
signalling during embryo development regulates cilia length in diverse epithelia.
Nature 458: 651–654
Odenwald MA, Choi W, Buckley A, Shashikanth N, Joseph NE, Wang Y, Warren
MH, Buschmann MM, Pavlyuk R, Hildebrand J, *et al* (2017) ZO-1 interactions
with F-actin and occludin direct epithelial polarization and single lumen
specification in 3D culture. *J Cell Sci* 130: 243–259
Okabe N, Xu B & Burdine RD (2008) Fluid dynamics in zebrafish Kupffer's vesicle.
Dev Dyn 237: 3602–3612
Purohit A, Tynan SH, Valle R & Doxsey SJ (1999) Direct Interaction of Pericentrin
with Cytoplasmic Dynein Light Intermediate Chain Contributes to Mitotic Spindle
Organization. *J Cell Biol* 147: 481
Rathbun LI, Aljiboury AA, Bai X, Hall NA, Manikas J, Amack JD, Bembenek JN &
Hehnly H (2020) PLK1- and PLK4-Mediated Asymmetric Mitotic Centrosome
Size and Positioning in the Early Zebrafish Embryo. *Current Biology* 30: 4519-
4527.e3
Sepulveda G, Antkowiak M, Brust-Mascher I, Mahe K, Ou T, Castro NM,
Christensen LN, Cheung L, Yoon D, Huang B, *et al* (2018) Co-translational
protein targeting facilitates centrosomal recruitment of PCNT during centrosome
maturation. *Co-translational protein targeting facilitates centrosomal recruitment*
of PCNT during centrosome maturation in vertebrates 7: 241083
Steigemann P & Gerlich DW (2009) Cytokinetic abscission: cellular dynamics at the
midbody. *Trends Cell Biol* 19: 606–616
Steigemann P, Wurzenberger C, Schmitz MHA, Held M, Guizetti J, Maar S & Gerlich
DW (2009) Aurora B-mediated abscission checkpoint protects against
tetraploidization. *Cell* 136: 473–484

Tavares B, Jacinto R, Sampaio P, Pestana S, Pinto A, Vaz A, Roxo-Rosa M,
Gardner R, Lopes T, Schilling B, *et al* (2017) Notch/Her12 signalling modulates,
motile/immotile cilia ratio downstream of Foxj1a in zebrafish left-right organizer.
Elife 6
Vertii A, Kaufman PD, Hehnly H & Doxsey S (2018) New dimensions of asymmetric
division in vertebrates. *Cytoskeleton* 75: 87–102 doi:10.1002/cm.21434
[PREPRINT]
Zimmerman WC, Sillibourne J, Rosa J & Doxsey SJ (2004) Mitosis-specific
anchoring of γ tubulin complexes by pericentrin controls spindle organization
and mitotic entry. *Mol Biol Cell* 15: 3642–3657
11
12

Figure Legends

Figure 1. Identification of an anteriorly positioned and pre-lumen enriched Kupffer's

Vesicle (KV) mitotic events and their post-mitotic distribution map.

(A) Model depicting KV cells transitioning from a small population, likely proliferative,

then organize into rosette structures then transition to a cyst of ciliated (light blue)

cells surrounding a fluid filled lumen.

(B) Representative confocal volumetric projections from live KV development video

(refer to **Video S1**). Pre-KV lumen time point (-22 min before lumen opening) and

post-lumen (26 min past lumen opening) shown. KV cell membranes marked by

Sox17:GFP-CAAX (magenta) and nuclei with h2afx:h2afv-mCherry (non-mitotic,

cyan; mitotic, yellow). Scale bar, 10 μm.

(C) Percentage of KV cells that are mitotic across a 120 min video time course (n=3

embryos ±SEM).

(D) Confocal projections of mitotic (Phospho-Histone H3, PH3, magenta) and post-

mitotic (BrdU, magenta) events at pre lumen (8 hpf) and post lumen (12 hpf). KV cell

membranes marked with Sox17:GFP-CAAX (inverted gray). Scale bar, 10 μm.

(E) Fraction of PH3 and BrdU-positive cells in the anterior vs posterior regions of the

KV (n=4 clutches, 36 embryos; and n=3 clutches, n=17 embryos respectively). Each

point represents an embryo, the point colors represent clutch, n=3 clutches,

**p<0.01.

(F-G) Confocal projections of H2B-Dendra photoconversion (non-photoconverted

black nuclei, photo-converted in fire-LUT). Cyan and magenta outlines mark mitotic

photoconverted event at 10 min. A total of 8 mitotic events were photoconverted.

Daughter cell positioning at (**F**) 96 min and (**G**) 180 min shown. **(F')** Trajectory of

marked mitotic events from division completion to 96 min. KV cell boundary, green

outline. Scale bar, 10 μm.

(H) Temporal schematic of KV mitotic event for representative embryo in (**F**),

daughter cells did not divide again.

(I) KV cell number (blue) and ratio of post mitotic anterior positioned or posterior

positioned cells were calculated over time from acquisition in (**F-G**).

Statistical results detailed in **Table S1**.

Figure 2. An anterior/posterior FGF signaling gradient is required for anterior zones

of mitotic activity.

(**A**) Model illustrating experimental approach to identify whether a signal influences

mitotic events and how those events contribute to KV development.

(**B-C**) Confocal projections of pre-lumen (8hpf, top) and post-lumen (12 hpf, bottom)

KVs treated with DMSO, 10 µM XAV939 (Wnt inhibitor), and 1µM infigratinib (FGF

inhibitor) (**B**) or lacking Pericentrin (*pcnt*$^{/-}$) compared with wild-type control (*pcnt*$^{+/+}$)

(**C**). KV cell membranes marked with Sox17:GFP-CAAX (inverted gray), mitotic

events with PH3 (magenta). Lumens highlighted in gold. Scale bar, 10 µM.

(**D-F**) Percent of mitotic KV cells (**D**), anterior vs posterior positioned mitotic events

(**E**), and lumen area (**F**) were measured for n>33 embryos (small dots represent

embryos, **D-F**) across n≥3 clutches (large dots represent clutch, **D, F**) per condition.

Refer to Figure S2. Graphs are scatter plots with mean of clutches shown ±SEM (**D,**

F) or a violin plot (**E**). DMSO, 10 µM XAV939, 1 µM infigratinib, *pcnt*$^{+/+}$ and *pcnt*$^{/-}$

shown. *p<0.05, **p<0.01, ***p<0.005, ****p<0.0001, ns, not significant. Statistical

results detailed in **Table S1**.

Figure 3. *Early KV mitotic events hold greater significance to KV development*

compared to later KV mitotic events.

(**A**) Model illustrating laser ablation conditions during KV development.

(**B**) Representative images of KV ablation experiment showing a pre-ablation and

post-ablation cell. White arrow highlights KV cell ablated. KV cell membranes

marked by Sox17:GFP-CAAX (magenta) and nuclei with h2afx:h2afv-mCherry

(cyan). Scale bar, 10 µm.

(C, D) Confocal projections taken from a video acquisition (0-200m, left, Sox17:GFP-

CAAX, inverted gray; lumen, gold, Refer to **Video S2**). The embryo was then fixed at

6SS (12 hpf) and stained for cilia (acetylated tubulin, cyan, right; Sox17:GFP-CAAX

in magenta). Number of cilia in each representative image labeled. Scale bar, 10

μm.

(E, F) Truncated violin plot depicting the number of KV cells at 8 hpf and 12 hpf for

no ablation and ablating all mitotics (**E**) or ablation conditions 1, 2, or 3 (**F**) across

n≥3 embryos (each point represents a single embryo). Median denoted with black

line and early/late temporal zone threshold (20 cells) denoted with grey dashed line.

(**G, H**) Lumen area was measured over time for over 100 min for no ablation (control)

and ablating all mitotics (**G**) or ablation conditions 1, 2 or 3 (**H**). n≥3 ±SEM embryos

measured per condition. Unpaired two-tailed t-tests in (**G**), One way ANOVA with

Dunnett's multiple comparison in (**H**). $*p<0.05$, $**p<0.01$, $***p<0.005$, ns, not

significant.

(**I, J**) Truncated violin plots depicting KV lumen area (12 hpf, **I**) and percentage of

ciliated KV cells (**J**). n≥3 embryos at each condition. One way ANOVA with Dunnett's

multiple comparison to non-mitotic ablation controls. $*p<0.05$, $***p<0.005$, ns, not

significant.

Statistical results detailed in **Table S1**.

Figure 4. *KV spindles stably align along KVs longest axis until the KV starts*

rounding, then spindles spin and are extruded.

(**A, B**) Time-lapse video of MT organization during KV formation (Sox17: EMTB-

3xGFP, gray). KV are outlined (white dashed line). Highlighted are spindles (yellow

arrows) and cytokinetic bridges (pink arrows). The blue boxed region is magnified in

(**B**) showcasing a spinning mitotic spindle that becomes extruded. Mitotic spindle

poles noted with blue and pink arrowheads. Extruded cytokinetic KV cell post

spinning event outlined (orange dashed line). Scale bars, 10 μm. Refer to **Video S3**.

(**C**) Methods to measure KV aspect ratio (A.R.) and the angular velocity of the

spindle (ω∠a), illustrated in model.

(**D, E**) Floating bar graph depicting the maximum, minimum and mean of KV A.R. (**D**)

and mean angular velocity (**E**) over 20 min binned time periods relative to lumen

formation. The gray dashed line marks when A.R. equals 2. n=7 embryos measured

across 6 clutches.

(**E'**) Angular velocity were compared when A.R. was <2 or >2 from data sets in (**E**).

32 spindle poles from n=7 embryos across 6 clutches were monitored. Unpaired two-

tailed t-tests, **p<0.01.

(**F**) Floating bar graph depicting the maximum, minimum and mean of the

percentage of mitotic events with post mitotic cell extruded at various time bin (each

40 min) relative to lumen formation. n=7 embryos from 6 clutches. (**F'**) Percentage of

mitotic events with post mitotic cell extruded pre- versus post-lumen formation. n=7

embryos from 6 clutches. Unpaired two-tailed t-tests, *p<0.05,

(**G**) Truncated violin plot depicting the frequency of mitotic events with post mitotic

cell extruded in no ablation controls or mitotic ablated condition 1 and 3. n>6

embryos measured over 6 clutches. Unpaired two-tailed t-tests, *p<0.05. Refer to

Video S4.

Statistical results detailed in **Table S1**.

Figure 5: *KV progenitor cells position cytokinetic bridges towards tjp1 marked tight*

junctions where they maintain cytokinetic bridge derived MT bundles post bridge

cleavage.

(A-B) Time-lapse image of MT (Sox17:EMTB-3xGFP, gray) and tjp1 (TgKI

(tjp1atdTomato, magenta)) distribution in KV mitotic cells. Time-lapse image of

metaphase cell in (**A**) highlighted by orange box is magnified in (**B**). Scale bars, 5

μm.

(C) Time-lapse image of an abscission event (yellow arrow) with a cytokinetic bridge

and midbody labeled with mKate-MKLP1 (cyan) and MT (EMTB-3xGFP, magenta).

Insets show MKLP1 (cyan) decorating a midbody (0 min), bridge abscission (yellow

arrow, 2 min), and midbody loss at 8 min. Scale bar, 10 µm.

(D) Working model depicting mitotic events affecting midbody distribution and tight

junction formation during KV development.

Figure 6: *Converging cytokinetic cells initiate actin recruitment at forming rosette*

centers.

(A-B) Time-lapse video of actin (Lifeact-mRuby, FireLUT) and MTs (Sox17:EMTB-

3xGFP, inverted Gray). The region denoted by orange box in **(A)** is magnified 1.5X in

(B) first cytokinetic bridge is marked (magenta arrow) in relation to second

cytokinetic bridge (blue arrow)**.** Scale bar, 10 µm, calibration bar, Lifeact-mRuby

fluorescent intensity values. Refer to **Video S5.**

(C) Testable model for how KV forms MT bundles and actin enriches in no ablation

control compared to mitotic ablation conditions.

(D) Representative confocal volumetric projections of KVs depicting MTs

(Sox17:EMTB-3xGFP, inverted gray) and actin (Lifeact-mRuby, FireLUT) of no

ablation control (CT, Left) or mitotic ablated condition 1 (Cond.1, right) at rosette

stage. Scale bar, 10 µm.

(E) Time lapse images of non-division-derived MT bundle formation (Sox17:EMTB-

3xGFP, inverted gray) and actin (Lifeact-mRuby, FireLUT) recruitment at tips of the

bundles from no ablation control condition (CT-NDMT, Top) and mitotic ablated

condition 1 (Cond.1-NDMT, bottom). Scale bar, 10 µm. Calibration bar on the right

denotes Lifeact-mRuby intensity.

(F-G) Line chart depicting the relative actin intensity 10 min (**F**) or floating bar graph

depicting area under curve (**G**) for relative actin intensity over 10 min interval

featured in (**F**). One way ANOVA with Dunnett's multiple comparison, *$p<0.05$, ns,

not significant, n≥3 for each condition over ≥2 clutches.

Statistical results detailed in **Table S1**.

Figure 7. *Cytokinetic bridge MT bundles are required for actin recruitment at rosette during lumen formation.*

(A) Model depicting laser ablation conditions during KV development.

(B-D) Volumetric projections from video acquisition of an abscission event. Actin is labeled with Lifeact-mRuby (Fire-LUT), and MT are labeled with EMTB-3xGFP (Gray). Scale bar, 10 µm. Ablation conditions are noted above image and outlined in (**A**).

(E-G) Area under the curve of relative actin (**E,G**) or MT intensity (**F**) over 10 min from **Fig S3A-C**. One-way ANOVA with Dunnett's multiple comparisons, *p<0.05, n≥4 cell, n≥2 clutches per condition for (**E**). Unpaired t-test, *p<0.05, **p<0.01. n≥3 cell, n=2 clutches per condition for (**F, G**).

Statistical results detailed in **Table S1**.

Figure 8: *Proposed model for cell division contribution to lumen formation.*

Model depicting lumen formation through FGF mediated asymmetrical cell cycle entry which drives formation of cytokinetic bridge derived MT bundles that recruit apical actin patches at rosette centers. MT (cyan), actin (orange), post-mitotic nuclei (magenta), pre-mitotic nuclei (dark blue), tight junction protein 1 (tjp1, spring green), and midbody (MKLP1, purple) are shown.

Supplemental Table Legends:

Table S1. *Detailed statistical analysis of results reported in this study.* This table includes detailed result of all statistical analysis used. Figure where analysis applied, category of analysis, embryo sample number, number of clutches, statistic test applied, parameter, significant level, and specific p-values are shown.

Supplemental Figure Legends:

Figure S1. *Identification of an anteriorly positioned and pre-lumen enriched Kupffer's Vesicle (KV) mitotic events and their post-mitotic distribution map.*

*Related to **Fig 1**.*

(A) KV mitotic cell position was mapped during KV development from **Video S1**.

Different colors indicate different 20 min time bins pre and post lumen formation.

(B) Time bins from (**A**) and **Video S1** were mapped as percentage of mitotic KV cells

in relation to lumen formation.

(C, D) Bubble plot depicting the positioning of mitotic event (PH3, **C**) and post mitotic

cells (BrdU-positive, **D**) in KV at 8 and 12 hpf respectively. Large circle, represents >

7 events. Small circle represents 3 events. Tiny circle represents 1 event. n=26

embryos, across 3 clutches.

(E) 2 mitotic photoconversion events shown in **Fig 1F** highlighted. Scale bar, 10 µm.

Statistical results detailed in **Table S1**.

Figure S2. *An anterior/posterior FGF signaling gradient is required for anterior zones*

*of mitotic activity. Related to **Fig 2**.*

(A-C) Confocal projections of post lumen (12 hpf) embryos treated with methanol, 50

µM Cyclopamine (**A**), DMSO, 10 µM XAV939 (**B**), and 0.5µM, 1µM, 3µM, 5µM

infigratinib (**C**). KV cell membranes marked with Sox17:GFP-CAAX (inverted gray),

lumens highlighted in gold. Scale bar, 10 µm.

(D) KV morphologies characterized following infigratinib treatment with varying

concentrations at 12 hpf, n>25 embryos per condition across 2 clutches.

(E) KV cell number at pre-lumen (8hpf) and post-lumen (12 hpf) following DMSO, 10

µM XAV939, and 1 µM infigratinib treatment. KV cell number was compared

between $pcnt^{+/+}$ and $pcnt^{-/-}$ zebrafish. n>16 embryos (small dots represent embryos)

across n≥3 clutches (large dots represent clutch) per condition. Graphs are scatter

plots with mean of clutches ±SEM shown, n≥3 clutches. **p<0.01, ***p<0.005,

****p<0.0001, ns, not significant.

Statistical results detailed in **Table S1.**

Figure S3. *Cytokinetic bridge MT bundles are required for actin recruitment at*

*rosette during lumen formation. Related to **Fig 7**.*

(A) Model depicting laser ablation conditions during KV development.

(B-D) Actin intensity (Lifeact-mRuby, **S3B, D**) and MT (EMTB-3xGFP, **S3C**) was measured over 10 min for cytokinetic bridge ablation conditions (**S3B, C**) and non-division-derived MT bundle (**S3C, D**) ablation conditions outlined in (**S3A**) and images in **Fig 7B-D**. n≥3±SEM.

Video S1*. Identification of an anteriorly positioned and pre-lumen enriched Kupffer's Vesicle (KV) mitotic events and their post-mitotic distribution map. Related to **Fig 1B***. Live confocal registered video showing mitotic events within KV in an embryo from the outcross of Sox17:GFP-CAAX and H2afx:h2afv-mCherry transgenic lines. KV cell plasma membranes marked by magenta (Sox17: GFP-CAAX) and nuclei (cyan, H2afx: h2afv-mCherry). Scale bar, 20 μm.

Video S2. *Early KV mitotic events hold greater significance to KV development compared to later KV mitotic events. Related to **Fig 3C, D***. Live confocal video showing lumen formation from a Sox17:GFP-CAAX, H2afx:h2afv-mCherry embryos under different laser ablation conditions. Sox17:GFP-CAAX (inverted gray) shown. Scale bar, 10 μm.

Video S3: *KV spindles stably align along KVs longest axis until the KV starts rounding, then spindles spin and are extruded. Related to **Fig 4A, B***. Live confocal video showing MT (grey, Sox17:EMTB-3xGFP) in KV (left). Highlighted from KV video on left is a spinning spindle (middle) and cell extrusion event (right). Scale bar, 20 μm.

Video S4: *KV spindles stably align along KVs longest axis until the KV starts rounding, then spindles spin and are extruded. Related to **Fig 4G***. Time lapse data set of Sox17:EMTB-3xGFP (MTs, Fire LUT) embryos under control (no ablation) conditions compared to laser ablation conditions (cond.1, first four

mitotic events ablated; and cond. 3, all mitotic events ablated after KV reaches 20

cells). Scale bar, 20 µm.

Video S5: *Converging cytokinetic cells initiate actin recruitment at forming rosette*

centers. Related to ***Fig 6A-6B****.*

Time lapse of actin (Lifeact-mRuby, fire-LUT) and KV MTs (Sox17:EMTB-3xGFP,

shown in inverted gray) during KV rosette formation. Scale bar, 10 µm.

Fig 1. Identification of an anteriorly positioned and pre-lumen enriched KV mitotic events and their post-mitotic distribution map.

Fig S1. Identification of an anteriorly positioned and pre-lumen enriched KV mitotic events and their post-mitotic distribution map.

Fig 2. An anterior/posterior FGF signaling gradient is required for anterior zones of mitotic activity.

Fig S2. An anterior/posterior FGF signaling gradient is required for anterior zones of mitotic activity.

Fig 3. Early KV mitotic events hold greater significance to KV development compared to later KV mitotic events.

Fig 4. KV spindles stably align along KVs longest axis until the KV starts rounding, then spindles spin and are extruded.

Fig 5. KV progenitor cells position cytokinetic bridges towards tjp1-marked tight junctions where they maintain cytokinetic bridge derived MT bundles post bridge cleavage.

Fig 6. Converging cytokinetic cells initiate actin recruitment at forming rosette centers.

Fig 7. Cytokinetic bridge derived MT bundles are required for actin recruitment at rosette during lumen formation.

Fig S3. Cytokinetic bridge derived MT bundles are required for actin recruitment at rosette during lumen formation

Fig 8. Proposed model for cell division contribution to lumen formation.

Table S1. Detailed statistical analysis of results reported in this study.

Figure	Category	n embryo	n Clutch	Statistical Test	Parameters	Result	p-value
1C	%Mitotic KV cells	3	3	N/A	N/A	N/A	N/A
1E	pH3 positive cell Anterior	36	4	Unpaired t-test	t=5.409, df=6	N/A	N/A
	pH3 positive cell Posterior					**	0.0016
	BrdU positive cell Anterior	17	3	Unpaired t-test	t=4.627, df=4	N/A	N/A
	BrdU positive cell Posterior					**	0.0098
1I	Ratio of Postmitotic anterior/posterior cell	1	1	N/A	N/A	N/A	N/A
S1A	%Mitotic KV cells	1	1	N/A	N/A	N/A	N/A
S1B	Position of the PH3 positive cell	1	1	N/A	N/A	N/A	N/A
S1C	pH3 positive cell	26	3	N/A	N/A	N/A	N/A
S1D	BrdU positive cell	3	3	N/A	N/A	N/A	N/A
2D	DMSO	53	5	One Way ANOVA (multiple comparison)	F (2,1.51) =5.621	N/A	N/A
	XAV939	33	3			ns	0.4369
	Infigratinib	68	3			**	0.0022
	pcnt (+/+)	46	3	Unpaired t-test	t=2.638, df=89	N/A	N/A
	pcnt (-/-)	45	3			**	0.0098
2E	DMSO	34	5	Unpaired t-test	t=4.447, df=64	****	<0.0001
	XAV939	20	3	Unpaired t-test	t=2.262, df=38	*	0.0295
	Infigratinib	34	3	Unpaired t-test	t=0.03508, df=66	ns	0.9721
	pcnt (+/+)	31	3	Unpaired t-test	t=3.440, df=60	**	0.0011
	pcnt (-/-)	30	3	Unpaired t-test	t=2.199, df=58	*	0.0319
2F	DMSO	29	5	One Way ANOVA	F (2,71) = 76.33	N/A	N/A
	XAV939	16	3			ns	0.2

	Infigratinib	29	3	(multiple comparison)		****	<0.0001
	pcnt (+/+)	16	4	Unpaired t-test	t=6.018, df=57	N/A	N/A
	pcnt (-/-)	29	3			****	<0.0001
S2D	DMSO	39	2	N/A	N/A	N/A	N/A
	0.5 µM Infigratinib	34	2	N/A	N/A	N/A	N/A
	1 µM Infigratinib	39	2	N/A	N/A	N/A	N/A
	3 µM Infigratinib	25	2	N/A	N/A	N/A	N/A
	5 µM Infigratinib	57	2	N/A	N/A	N/A	N/A
S2E	DMSO pre-lumen stage	36	5	Unpaired t-test	t=5.693, df=71	N/A	N/A
	DMSO post-lumen stage	37	4			****	<0.0001
	XAV939 pre-lumen stage	22	3	Unpaired t-test	t=3.679, df=36	N/A	N/A
	XAV939 post-lumen stage	16	3			***	0.0008
	Infigratinib pre-lumen stage	32	3	Unpaired t-test	t=3.032, df=59	N/A	N/A
	Infigratinib post-lumen stage	29	3			**	0.0036
	pcnt (+/+) pre-lumen stage	54	3	Unpaired t-test	t=14.55, df=75	N/A	N/A
	pcnt (+/+) post-lumen stage	23	4			****	<0.0001
	pcnt (-/-) pre-lumen stage	35	3	Unpaired t-test	t=1.906, df=56	N/A	N/A
	pcnt (-/-) post-lumen stage	23	3			ns	0.0617
3E	No Ablation 8hpf	4	4	Unpaired t-test	t=3.586, df=7	**	0.0089
	No Ablation 12hpf	5	5				
	Ablate All Mitotics 8hpf	3	3	Unpaired t-test	t=1.413, df=4	ns	0.2306
	Ablate All Mitotics 12hpf	3	3				
3F	Cond. 1_8hpf	4	4	Unpaired t-test	t=2.782, df=5	*	0.0388
	Cond. 1_12hpf	3	3				

	Cond. 2_8hpf	3	3	Unpaired t-test	t=4.230, df=4	*	0.0134
	Cond. 2_12hpf	3	3				
	Cond. 3_8hpf	6	6	Unpaired t-test	t=1.040, df=9	ns	0.3257
	Cond. 3_12hpf	5	5				
3G	Lumen area: No ablation control 20min	3	3	Unpaired t-test	t=1.000, df=4	ns	0.3739
	Lumen area: All mitotic ablated 20min	3	3				
	Lumen area: No ablation control 30min	3	3	Unpaired t-test	t=1.231, df=4	ns	0.2856
	Lumen area: All mitotic ablated 30min	3	3				
	Lumen area: No ablation control 40min	3	3	Unpaired t-test	t=1.356, df=4	ns	0.2466
	Lumen area: All mitotic ablated 40min	3	3				
	Lumen area: No ablation control 50min	3	3	Unpaired t-test	t=2.049, df=4	ns	0.1098
	Lumen area: All mitotic ablated 50min	3	3				
	Lumen area: No ablation control 60min	3	3	Unpaired t-test	t=1.777, df=4	ns	0.1503
	Lumen area: All mitotic ablated 60min	3	3				
	Lumen area: No ablation control 70min	3	3	Unpaired t-test	t=2.000, df=4	ns	0.1161
	Lumen area: All mitotic ablated 70min	3	3				

	Lumen area: No ablation control 80min	3	3	Unpaired t-test	t=2.174, df=4	ns	0.0954
	Lumen area: All mitotic ablated 80min	3	3				
	Lumen area: No ablation control 90min	3	3	Unpaired t-test	t=3.096, df=4	*	0.0364
	Lumen area: All mitotic ablated 90min	3	3				
	Lumen area: No ablation control 100min	3	3	Unpaired t-test	t=3.298, df=4	*	0.0300
	Lumen area: All mitotic ablated 100min	3	3				
	Lumen area: No ablation control 110min	3	3	Unpaired t-test	t=4.597, df=4	*	0.0101
	Lumen area: All mitotic ablated 110min	3	3				
3H	Lumen area: Cond. 1_10min	3	3	One Way ANOVA (multiple comparison)	F (2, 6) = 0.5000	N/A	N/A
	Lumen area: Cond. 2_10min	3	3			ns	0.6180
	Lumen area: Cond. 3_10min	4	4			ns	0.6180
	Lumen area: Cond. 1_20min	3	3	One Way ANOVA (multiple comparison)	F (2, 6) = 0.5000	N/A	N/A
	Lumen area: Cond. 2_20min	3	3			ns	0.6180
	Lumen area: Cond. 3_20min	4	4			ns	0.6180
	Lumen area: Cond. 1_30min	3	3	One Way ANOVA (multiple comparison)	F (2, 6) = 0.7719	N/A	N/A
	Lumen area: Cond. 2_30min	3	3			ns	0.4955
	Lumen area: Cond. 3_30min	4	4			ns	0.4955
	Lumen area: Cond. 1_40min	3	3	One Way ANOVA	F (2, 6) = 2.493	N/A	N/A

Lumen area: Cond. 2_40min	3	3	(multiple comparison)		ns	0.1697
Lumen area: Cond. 3_40min	4	4			ns	0.1697
Lumen area: Cond. 1_50min	3	3	One Way ANOVA	F (2, 6) = 1.520	N/A	N/A
Lumen area: Cond. 2_50min	3	3	(multiple comparison)		ns	0.2945
Lumen area: Cond. 3_50min	4	4			ns	0.2945
Lumen area: Cond. 1_60min	3	3	One Way ANOVA	F (2, 6) = 8.375	N/A	N/A
Lumen area: Cond. 2_60min	3	3	(multiple comparison)		*	0.0215
Lumen area: Cond. 3_60min	4	4			*	0.0215
Lumen area: Cond. 1_70min	3	3	One Way ANOVA	F (2, 6) = 19.10	N/A	N/A
Lumen area: Cond. 2_70min	3	3	(multiple comparison)		**	0.0031
Lumen area: Cond. 3_70min	4	4			**	0.0031
Lumen area: Cond. 1_80min	3	3	One Way ANOVA	F (2, 6) = 48.23	N/A	N/A
Lumen area: Cond. 2_80min	3	3	(multiple comparison)		***	0.0003
Lumen area: Cond. 3_80min	4	4			***	0.0003
Lumen area: Cond. 1_90min	3	3	One Way ANOVA	F (2, 6) = 17.48	N/A	N/A
Lumen area: Cond. 2_90min	3	3	(multiple comparison)		**	0.0039
Lumen area: Cond. 3_90min	4	4			**	0.0039
Lumen area: Cond. 1_100min	3	3	One Way ANOVA	F (2, 6) = 35.51	N/A	N/A
Lumen area: Cond. 2_100min	3	3	(multiple comparison)		***	0.0006
Lumen area: Cond. 3_100min	4	4			***	0.0006
Lumen area: Cond. 1_110min	3	3	One Way ANOVA	F (2, 6) = 25.79	N/A	N/A
Lumen area: Cond. 2_110min	3	3	(multiple comparison)		**	0.0016

	Lumen area: Cond. 3_110min	4	4			**	0.0016
3I	Non-mitotic Control	10	6	One Way ANOVA (multiple comparison)	$F_{(3, 17)} = 3.302$	N/A	N/A
	Ablate Mitotic Cond. 1	3	3			*	0.0493
	Ablate Mitotic Cond.2	4	4			ns	0.7471
	Ablate Mitotic Cond. 3	5	5			ns	0.0893
3J	Non-mitotic Control	10	6	One Way ANOVA (multiple comparison)	$F_{(3, 18)} = 9.935$	N/A	N/A
	Ablate Mitotic Cond. 1	3	3			***	0.0003
	Ablate Mitotic Cond. 2	4	4			ns	0.6507
	Ablate Mitotic Cond. 3	5	5			ns	0.5365
4D	Aspect Ratio of KV	7	6	N/A	N/A	N/A	N/A
4E	Spindle Angular Velocity	43 Mitotic Cells	6	N/A	N/A	N/A	N/A
4E'	A.R.>2	12 cells (7 embryos)	6	Unpaired t-test	t=3.114, df=30	N/A	0.0040
	A.R.<2	20 cells (7 embryos)	6			**	0.6507
4F, F'	Pre-Lumen	7	6	Unpaired t-test	t=2.381, df=12	N/A	0.0347
	Post Lumen	7	6			*	
4G	No Ablation Control	7	6	Unpaired t-test	t=2.488, df=11	N/A	0.0301
	Ablate Mitotics Cond. 1 and 3	6	6			*	
6F	CT-CB	5	2	N/A	N/A	N/A	N/A

	CT-NDMT	5	2	N/A	N/A	N/A	N/A
	Cond.1-NDMT	3	1	N/A	N/A	N/A	N/A
6G	CT-CB	5	2	One-way ANOVA (Multiple comparison)	F(2,10)=8.989	*	0.0185
	CT-NDMT	5	2			N/A	
	Cond.1-NDMT	3	1			ns	0.4377
7E	CT-CB	6	3	One-way ANOVA (Multiple comparison)	F (2, 11) = 4.477	*	0.022
	CB-NA	4	3			*	0.0257
	CB-A	4	3			N/A	
7F	CB-A	4	2	Unpaired t-test	t=4.103, df=5	N/A	0.0093
	AB-NDMT	3	2			**	
7G	CT-NDMT	3	2	Unpaired t-test	t=2.623, df=5	N/A	0.00469
	AB-NDMT	4	2			*	
S3B	CT-CB	6	3	N/A	N/A	N/A	N/A
	CB-NA	4	3	N/A	N/A	N/A	N/A
	CB-A	4	2	N/A	N/A	N/A	N/A
S3C	CB-A	3	2	N/A	N/A	N/A	N/A
	AB-NDMT	4	2	N/A	N/A	N/A	N/A
S3D	CT-NDMT	3	2	N/A	N/A	N/A	N/A
	AB-NDMT	4	2	N/A	N/A	N/A	N/A