**A novel motif in calcimembrin/C16orf74 dictates multimeric dephosphorylation by calcineurin**

Devin A. Bradburn[1,7], Joana C. Reis[2,3,7], Shariq Qayyum[2,4,5], Thibault Viennet[2,3,6], Haribabu Arthanari[2,3], Martha S. Cyert[1]*

1. Department of Biology, Stanford University, Stanford, CA 94305, USA

2. Department of Cancer Biology, Dana-Farber Cancer Institute, Boston, Massachusetts, USA.

3. Department of Biological Chemistry and Molecular Pharmacology, Harvard Medical School, Boston, Massachusetts, USA.

4. Men's Health: Aging & Metabolism, Brigham and Women's Hospital, Harvard Medical School, Boston, MA, USA.

5. Cardiovascular Research Center, Department of Medicine, Massachusetts General Hospital and Harvard Medical School, Boston, MA, USA

6. Department of Chemistry and iNANO, Aarhus University, Aarhus, Denmark

7. Contributed equally to this work

*Corresponding author:
Department of Biology
410B Bass Biology Building
Stanford University
Stanford, CA 94305-5020
phone: 650 723-9970
fax:  650 724-9945
E-mail: mcyert@stanford.edu

**Abstract:**

Calcineurin, the $Ca^{2+}$/calmodulin-activated protein phosphatase, recognizes substrates and regulators via

short linear motifs, PxIxIT and LxVP, which dock to distinct sites on calcineurin and determine calcineurin

distribution and catalysis, respectively. Calcimembrin/C16orf74 (CLMB), an intrinsically disordered

microprotein whose expression correlates with poor cancer outcomes, targets calcineurin to membranes

where it may promote oncogenesis by shaping calcineurin signaling. We show that CLMB associates with

membranes via lipidation, i.e. N-myristoylation and reversible S-acylation. Furthermore, CLMB contains

an unusual composite 'LxVPxIxIT' motif, that binds the PxIxIT-docking site on calcineurin with

extraordinarily high affinity when phosphorylated, $^{33}$LxVPxIxITxx(p)T$^{44}$. Calcineurin dephosphorylates

CLMB to decrease this affinity, but Thr44 is protected from dephosphorylation when PxIxIT-bound. We

propose that CLMB is dephosphorylated in multimeric complexes, where one PxIxIT-bound CLMB recruits

calcineurin to membranes, allowing a second CLMB to engage via its LxVP motif to be dephosphorylated.

*In vivo* and *in vitro* data, including nuclear magnetic resonance (NMR) analyses of CLMB-calcineurin

complexes, supports this model. Thus, the CLMB composite motif imposes unique properties to calcineurin

signaling at membranes including sensitivity to CLMB:calcineurin ratios, CLMB phosphorylation and

dynamic S-acylation.

**Introduction**

Calcium ($Ca^{2+}$) ions are spatially and temporally controlled in cells to regulate a wide array of biological

processes (*1*). $Ca^{2+}$ signaling frequently occurs within membrane-associated microdomains that contain

both effector proteins and their substrates. Thus, subcellular localization of calcineurin, the only $Ca^{2+}$-

calmodulin (CaM) regulated serine/threonine protein phosphatase, dictates its signaling specificity(*2*).

Calcineurin (CNA/B) is an obligate heterodimer composed of catalytic (CNA) and regulatory (CNB)

subunits that is activated when $Ca^{2+}$/calmodulin binding relieves autoinhibition carried out by the CNA C-

terminal domain. Calcineurin interacts with partners via two short linear motifs (SLiMs), LxVP and PxIxIT,

which cooperate to promote dephosphorylation. PxIxIT binds to CNA independent of $Ca^{2+}$ and anchors

calcineurin to substrates and regulators. In contrast, LxVP binds to a site composed of residues in both CNA

and CNB and orients substrates for dephosphorylation at a separate catalytic site, marked by $Zn^{2+}$ and $Fe^{2+}$

ions (*3*). LxVP binding occurs only after activation by $Ca^{2+}$/CaM and is blocked by autoinhibitory sequences

in CNA as well as the immunosuppressant drugs, Cyclosporin A and FK506 (*4, 5*).

Systematic discovery of PxIxIT and LxVP SLiMs has significantly expanded our knowledge of calcineurin

signaling (*6-9*) and surprisingly identified at least 31 human proteins that harbor composite calcineurin-

binding motifs containing  contiguous LxVP and PxIxIT sequences that share a proline (LxVPxIxIT)(*9*).

Although each SLiM in a composite motif can bind to its cognate site, they are unlikely to do so

simultaneously, as their docking surfaces are ~60 Å apart on calcineurin. Thus, how or if these motifs

promote dephosphorylation is unclear.

Here we elucidate how C16orf74, a short unstructured protein associated with poor outcomes in melanoma,

pancreatic and head and neck cancer, is regulated by calcineurin via a composite motif  (*10, 11*). We name

C16orf74 calcimembrin (CLMB), because it associates with membranes via lipidation (N-myristoylation

and S-acylation) and recruits calcineurin to these sites. CLMB is phosphorylated at Thr 44 (pThr44), and

binds calcineurin with exceptionally high affinity because the flanking LxVP and pThr44 features

(LxVPxIxITxx(p)T) enhance PxIxT-mediated binding to CNA. Calcineurin also dephosphorylates pThr44.

but this residue is protected when CLMB is bound to the PxIxT-docking surface. Thus, we propose that

dephosphorylation occurs in multimers, with one CNA/B bound to two CLMB molecules at the PxIxIT and

LxVP-docking sites, respectively. NMR confirms formation of these complexes *in vitro*. Furthermore, this

multimeric model predicts that CLMB dephosphorylation is sensitive to the relative levels of CLMB to

calcineurin, which we demonstrate through in cell analyses and kinetic modeling of dephosphorylation *in*

*vitro*. In summary, this work reveals how CLMB can dynamically shape calcineurin signaling through a

mechanism dictated by its composite motif and reversible association with membranes.

## Results

**Calcimembrin is myristoylated and S-acylated.** CLMB lacks predicted transmembrane domains but may associate with membranes via lipidation (Figure 1A) (*10*). CLMB contains a glycine at position 2 (Gly2), a site frequently modified by myristate, a 14-carbon lipid whose co-translational addition can promote protein association with membranes (*12*). CLMB also has two cysteines near in its N-terminus, Cys7 and Cys14, one of which, Cys7, is predicted to be modified by S-acylation, the reversible addition of a fatty acid, usually a 16-carbon palmitate, that regulates membrane localization (*13*). To determine if CLMB is lipidated, we metabolically labeled HEK293 Flp-In T-REx cells inducibly expressing wildtype CLMB-FLAG (CLMB$_{WT}$) with YnMyr and 17-ODYA, alkyne-containing analogs of myristate and palmitate respectively (*14*). Following immunoprecipitation, analog incorporation was detected via addition of biotin using CLICK-chemistry followed by visualization with streptavidin (Figure 1B). CLMB$_{WT}$ incorporated YnMyr and 17-ODYA, indicating that the protein is both myristoylated and palmitoylated. To identify which residues were lipidated, we mutated proposed sites of S-acylation (CLMB$_{C7,14S}$), and myristoylation (CLMB$_{G2A}$ (Figure 1B). The mutants were expressed at low levels suggesting instability, which we countered by including a proteasome inhibitor (MG132) in the analyses. CLMB$_{C7,14S}$ showed no 17-ODYA incorporation but maintained YnMyr suggesting that Cys7 and/or Cys14 are S-acylated. By contrast, neither lipid modification was detected for CLMB$_{G2A}$, which confirms that Gly2 is myristoylated.

To determine if CLMB-FLAG is dually S-acylated at Cys7 and Cys14, we used an acyl-PEG exchange assay (APE) in which S-acylated groups are cleaved using hydroxylamine (NH$_2$OH) and exchanged for a PEGylated molecule that increases protein mass by 5 kDa for each S-acylated residue(*15*) (Figure 1C). Two slower migrating bands were observed for CLMB$_{WT}$. CLMB$_{C7S}$ lacked the upper band, while CLMB$_{C7,14S}$ was unmodified, indicating that CLMB$_{WT}$ contains two S-acylated residues, i.e. Cys7 and Cys14. CLMB$_{G2A}$ also displayed two sites of S-acylation, suggesting that metabolic labeling (Figure 1B) was not sufficiently sensitive to detect these modifications. Together these findings establish that CLMB is myristoylated at Gly2 and S-acylated at Cys7 and Cys14.

**Dynamic S-acylation of CLMB mediates membrane association.** To examine whether CLMB-FLAG S-

acylation is dynamic (*16*), we carried out a pair of pulse-chase analyses with two metabolic labels (Figure

1D and 1E). L-AHA, an azide-containing methionine analog detected CLMB-FLAG protein levels, while

in parallel, 17-ODYA detected S-acylation. CLMB-FLAG was immunopurified (IP) after labeling for one

hour followed by a 30-minute chase with DMSO or Palmostatin B (PalmB), a pan-inhibitor of thioesterases

that remove S-acylation (*17*). Quantifying the 17-ODYA to L-AHA ratio revealed that incorporation of

palmitate into CLMB increased ~3-fold when thioesterases were inhibited (PalmB vs DMSO). This

indicates that S-acylation of CLMB turns over rapidly relative to its rate of degradation.

To determine if lipidation mediates CLMB membrane recruitment, we analyzed the localization of

$CLMB_{WT}$ and lipidation defective mutants, $CLMB_{G2A}$ and $CLMB_{C7,14S}$. Detergent-assisted subcellular

fractionation showed that $CLMB_{WT}$ was primarily membrane associated, while both mutants were

predominantly cytosolic (Figure 1F and 1G). Therefore, N-myristoylation and dynamic S-acylation of

CLMB stabilize the protein and mediate its association with membranes.

**The composite SLiM in CLMB mediates binding to and dephosphorylation by calcineurin.** Next, we

examined how each part of CLMB's composite motif, LDVPDIIIT, contributes to calcineurin binding by

generating mutants, $CLMB_{IxITMut}$ (LDVPD**AIAA**) and $CLMB_{LxVMut}$, (**ADA**PDIIIT) (Figure 1A) that disrupt

each motif independently without altering the shared proline. To examine binding of these mutants to

calcineurin, the association of CLMB-FLAG with GFP-tagged calcineurin (CNA-GFP complexed with

endogenous CNB) was assessed following purification of calcineurin from lysates of HEK293 cells co-

transfected with these constructs (Figure 2A and 2B). $CLMB_{WT}$ but not $CLMB_{IxITMut}$ co-purified with

calcineurin, showing that this interaction was PxIxIT-dependent. Co-purification of $CLMB_{LxVMut}$ with

calcineurin was somewhat reduced relative to $CLMB_{WT}$, indicating a smaller contribution to calcineurin

binding. Furthermore, when expressed and analyzed by SDS-polyacrylamide gel electrophoresis (SDS-

PAGE), both mutants displayed two electrophoretic forms, with a slower migrating form also faintly

observed for $CLMB_{WT}$. These represent CLMB containing pThr44 and are absent in mutants lacking this

residue (CLMB_{T44A}, CLMB_{T44A+LxVMut}). These observations indicate that calcineurin binds to CLMB and

dephosphorylates pT44, which is disrupted when either the LxVP or PxIxIT portion of the composite motif

is compromised. Finally, we confirmed that pThr44 enhances binding to calcineurin, as the slower migrating

forms of CLMB_{WT} and CLMB_{LxVMut} predominantly co-purified with calcineurin (*10*). Indeed, CLMB_{T44A}

displayed reduced binding to calcineurin, which was further reduced by mutating LxV (CLMB_{T44A+LxVMut}).

The contributions of PxIxIT, LxVP, and pThr44 to calcineurin binding were also examined by measuring

the extent of CLMB co-localization with calcineurin (CNA-GFP complexed with endogenous CNB) via

immunofluorescence (IF). When co-transfected into MCF7 cells CLMB_{WT}-FLAG recruits calcineurin to

membranes (Figure 2C, 2D). In contrast, when co-expressed with CLMB_{IxITMut} calcineurin is predominantly

cytosolic. CLMB_{LxVMut} and CLMB_{T44A} also display reduced colocalization with calcineurin, mirroring the

trends observed in co-purification analyses.

Finally, the ability of calcineurin to dephosphorylate pThr44 in CLMB was demonstrated *in vitro* using

peptide substrates that encode residues 21-50 (Figure 1A and 2 E). The peptide from pCLMB_{WT} but not

pCLMB_{LxVMut} was dephosphorylated showing that, as for other calcineurin substrates, the LxVP motif is

required for catalysis (*18*). Together, these data show that both the LxVP and PxIxIT components of the

composite motif are functional, that CLMB is a calcineurin substrate, and that pThr44 enhances CLMB

binding to calcineurin.

**Phosphorylated Thr44 and the LxVP motif enhance CLMB binding to the PxIxIT docking site.** To

quantitatively assess CLMB-calcineurin interaction, we used isothermal titration calorimetry (ITC) and

fluorescence polarization anisotropy (FP) to measure binding of CLMB peptides containing the composite

SLiM (residues 21-50) to purified, catalytically inactive CNA/B containing CNA_{Trunc+H151Q}. This mutant

lacks the C-terminal autoinhibitory region of CNA, which allows LxVP motifs to bind to CNA/B in the

absence of $Ca^{2+}$/CaM(*19*). ITC revealed that the phosphorylated CLMB_{WT} peptide (pCLMB_{WT}) binds

calcineurin at a one-to-one ratio with the highest reported affinity for any substrate, ($K_D$= 8.29 nM), (Figure

3A). FP using FITC-labelled peptides revealed that phosphorylation of Thr44 enhances CLMB binding to

calcineurin ~30-fold ($K_D$ = 12.29nM vs 384.5 nM) (Figure 3B). The concordance between measurements

made with a tag-free CLMB peptide (ITC) vs. a peptide tagged with fluorescein isothiocyanate (FITC) (FP)

(8.29 vs 12.29 nM for pCLMB$_{WT}$), shows that the tag does not significantly affect binding affinity. Given

the high-throughput capability of FP, this method was used for the remaining comparisons. To test if the

high affinity pCLMB$_{WT}$-calcineurin binding observed was PxIxIT-mediated, we used FP to assess the ability

of CLMB peptides to compete for calcineurin binding with FITC-PVIVIT, a well characterized PxIxIT

peptide([20, 21]) (Figure 3C). pCLMB$_{WT}$ competed with PVIVIT most effectively, (Figure 3C, 3E) with

peptides lacking the LxVP motif (pCLMB$_{LxVMut}$), or pThr44 (CLMB$_{WT}$) showing reduced competition.

pCLMB$_{IxITMut}$ lacking the PxIxIT motif was unable to displace PVIVIT from CNA/B.  When these

experiments were repeated using FITC-pCLMB$_{WT}$ peptide in place of PVIVIT (Figure 3D, 3E), a similar

pattern was observed, although the absence of pThr44 showed a much greater (~30x) reduction in $K_i$.

Importantly, full length CLMB protein displayed a near equal ability to displace FITC-pCLMB$_{WT}$, showing

that CLMB peptides accurately mimic full length CLMB binding to calcineurin (Figure S1A and S1B).

Altogether, these data show that, like PVIVIT, pCLMB$_{WT}$ peptides bind to the PxIxIT docking groove on

CNA and that flanking residues, i.e. LxV and pThr44, enhance binding to form an extended

LxVPxIxITxxT(p) sequence with exceptionally high affinity for calcineurin.

**NMR analysis shows CLMB binding to two sites on calcineurin.** Next, to investigate the structure of

CLMB in the presence and absence of calcineurin, we used solution-state NMR. 88% of non-proline

backbone resonances were assigned using a standard set of triple-resonance experiments, utilizing full-

length dually [13]C- and [15]N-labeled CLMB (76 residues) that was expressed in *E.coli* (Figure 4A and S2A).

To assess CLMB secondary structure, we employed Talos-N, a computational method that predicts

backbone torsion angles from NMR chemical shifts. This revealed that CLMB is disordered overall, but

contains two regions that displayed a tendency to form a β-strands (Figure S2B):  Residues I38-I40 include

the PxIxIT, consistent with binding of this motif-type to CNA as a β-strand (*4, 22*). N-terminal residues,

Q12-V15, were also predisposed to localized folding suggesting possible involvement in other binding

interactions.

To map the interaction interface of calcineurin on CLMB, we focused first on PxIxIT-mediated interactions

by titrating $^{15}$N-labeled CLMB with varying molar ratios of monomeric recombinant CNA in the absence

of CNB and monitoring the interaction through $^{15}$N-$^1$H SOFAST-HMQC experiments. CNA can bind to

PxIxIT motifs but lacks the LxVP binding surface found at the interface of CNA and CNB(4). Due to the

size of the CLMB-CNA complex, we anticipated significant broadening of NMR signals at the interaction

interface. This broadening can also arise from the exchange kinetics between the observable CLMB and

the titrant, CNA, making it challenging to distinguish the two effects. As expected, at a molar ratio of 2:1

(CLMB: CNA), many of the CLMB signals undergo significant line broadening, which become more

pronounced at a molar ratio to 1:1, due to the majority of CLMB being bound to CNA. This is particularly

the case for signals corresponding to the PxIxIT and LxVP motifs, indicating, as observed in FP

experiments, that LxV residues participate in binding to the PxIxIT interface (Figure 4B and C, Figure

S2C). Line broadening across the majority of CLMB residues indicates that the interaction interface on

CLMB extends beyond the PxIxIT and LxVP motifs (Figure 4B and C). Interactions with the N- and C-

terminal residues of CLMB may be an avidity-driven effect, anchored by PxIxIT binding, as has been

reported for other calcineurin substrates (23).

To investigate binding that may occur at the LxVP docking site of calcineurin, we next titrated CLMB with

the heterodimeric CNA/B complex. Signal broadening at the N-terminus of CLMB was more pronounced

with CNA/B relative to CNA alone, suggesting possible interactions between CLMB and CNB (Figure 4B,

C vs 4D, E, and S2D). These include the Q9-V12 region that displayed partial folding in the absence of

calcineurin.  At a molar ratio of 2:1 (CLMB: CNA/B), more broadening is observed in LxVP but not PxIxIT

residues when compared to the interaction with CNA (Figure 4B vs 4D, and S2C). However, at a molar

ratio of 1:1, where essentially all CLMB should be PxIxIT-bound (given the higher $K_D$ for PxIxIT vs LxVP

and our working concentrations), the pattern of line broadening at the composite motif is similar when

complexed with either CNA or CNA/B (Figure 4C vs 4E). Together these findings show that when in excess

(2 CLMB: 1 CNA/B), CLMB engages calcineurin (CNA/B) through both its LxVP and PxIxIT motifs. To

investigate these interactions further, we carried out structural predictions for 2 CLMB molecules and

CNA/B using AlphaFold 3(*24*). Consistent with our NMR analyses, the models show two copies of CLMB

simultaneously bound to CNA/B, one each at the LxVP and PxIxIT docking sites. Predicted binding for

CLMB containing pThr44 to CNA/B was nearly identical, however with the LxVP-bound CLMB showing

an additional interaction of pThr44 with the active site as would occur during catalysis (Figure S3A-S3C).

Thus, these models suggest that formation of multimeric complexes (2 CLMB: 1 CNA/B) will facilitate

dephosphorylation.

**Calcineurin dephosphorylates CLMB by forming multimers.** Next, we sought to determine if, as

predicted, dephosphorylation of pT44 is promoted when CLMB is in excess (2 CLMB: 1 CNA/B) but is

inhibited by lower CLMB: CNA/B ratios which favor formation of PxIxIT-bound complexes that protect

pThr44 from dephosphorylation. To this end, we inducibly expressed CLMB$_{WT}$-FLAG in HEK293 Flp-In

T-REx cells and examined its phosphorylation status in cells where calcineurin is expressed at endogenous

levels or overexpressed (i.e. transfected with GFP-CNA, which complexes with endogenous CNB). In stark

contrast to its behavior with typical substrates, overexpression of calcineurin caused CLMB-FLAG

phosphorylation to increase (Figure 5A, B, and S4A). Intact PxIxIT binding rather than catalytic activity

was required for this effect, as overexpressing calcineurin that is catalytically inactive (containing GFP-

CNA$_{H151Q}$), but not calcineurin that is defective for PxIxIT binding (containing GFP-CNA$_{NIR}$), increased

CLMB-FLAG phosphorylation (Figure S4B)(*25*). We predicted that further increasing CLMB expression

in these calcineurin overexpressing cells would increase the CLMB:CNA/B ratio, leading to greater

dephosphorylation of CLMB-FLAG. Thus, we overexpressed V5-tagged CLMB in HEK293 Flp-In T-REx

cells inducibly expressing CLMB-FLAG with overexpressed calcineurin (GFP-CNA complexed with

endogenous CNB). As predicted, the amount of phosphorylated CLMB$_{WT}$-FLAG observed was reduced by

introducing additional CLMB$_{WT}$-V5 (Figure 5A-C). This effect was dependent on PxIxIT binding, as

introducing CLMB$_{IxITMut}$-V5 did not decrease CLMB$_{WT}$-FLAG phosphorylation, but CLMB$_{LxVMut}$-V5,

which contains an intact PxIxIT motif, did. We repeated these experiments using CLMB$_{IxITMut}$-FLAG,

which is partially phosphorylated regardless of calcineurin levels. Surprisingly, CLMB$_{IxITMut}$-FLAG was

predominantly dephosphorylated when co-expressed with CLMB$_{WT}$-V5 or CLMB$_{LxVMut}$-V5 but not

CLMB$_{IxITMut}$-V5 (Figure 5A and B). Thus, although the LxVP and PxIxIT components of the CLMB

composite motif are both required for efficient dephosphorylation, they do not need to be on the same

molecule. A single copy of CLMB must have an LxVP to mediate its own dephosphorylation but can utilize

the PxIxIT of neighboring copies to recruit calcineurin, promote LxVP binding, and be dephosphorylated

(Figure 5C).

To demonstrate more precisely how the ratio of CLMB to calcineurin affects dephosphorylation, we

conducted a series of *in vitro* dephosphorylation assays using a fixed substrate concentration (100nM

pCLMB$_{WT}$ or pCLMB$_{IxITMut}$ peptides) with varied concentrations of purified calcineurin (CNA/B) (20nM,

100nM and 200nM). For pCLMB$_{IxITMut,}$ dephosphorylation rate increased with the calcineurin

concentration. However, pCLMB$_{WT}$ showed no change under these same conditions (Figure 5D). We

propose that this unusual result reflects competition between the two different modes of pCLMB binding

to calcineurin: high affinity binding to the PxIxIT-docking site, which prevents T44 dephosphorylation, vs

low affinity binding to the LxVP-docking site which promotes dephosphorylation. With a substrate that

contains only an LxVP motif (i.e. pCLMB$_{IxITMut}$), increasing calcineurin concentration accelerates substrate

binding to the LxVP docking site and dephosphorylation. However, with a substrate that contains a

composite motif (i.e. pCLMB$_{WT}$), this increased LxVP-mediated binding is offset by an equivalent

reduction in the amount of pCLMB that is available to be dephosphorylated due to substrate binding via

the PxIxIT docking site and pThr44 protection. We used kinetic modeling to assess how well this proposed

mechanism explains the observed dynamics of pCLMB$_{WT}$ and pCLMB$_{IxITMut}$ dephosphorylation (Figure

S5A and B). When simultaneously fitting results with both substrates, the kinetic model showed remarkable

agreement with the data and estimated rate constants that approach values reported in the literature (Figure

5D and S5C)(*6, 18*). Together, these analyses provide evidence that the phosphorylated composite motif in

CLMB imposes unique attributes on its dephosphorylation. In addition to a requirement for $Ca^{2+}$ and CaM,

the presence of this composite motif dictates that substrate must be in excess of calcineurin so that

multimeric complexes can form and carry out dephosphorylation.

**Discussion**

In this study, we reveal how a composite LxVPxIxITxx(p)T motif functions in calcineurin binding and

dephosphorylation. For engaging calcineurin (CNA/B), we can break down the thermodynamic

contributions of the motif in CLMB into three parts: (i) the core PxIxIT motif, (ii) pThr44, and (iii) the

LxVP motif. Our FP analyses reveal that the PxIxIT motif provides the largest contribution to the binding

free energy, as mutating this motif (pCLMB$_{IxITMut}$) decreased binding by more than two orders of magnitude.

Phosphorylation acts as the second major driver of binding affinity, imparting a ~30-fold increase in binding

affinity, from $K_D$ ~400nM to 13 nM for the non-phosphorylated versus phosphorylated peptide (Figure 3).

Similarly, phosphorylation at this position was reported to enhance the affinity of the PxIxIT motif in

Cabin1, a calcineurin regulator that like CLMB, contains a PxIxIT followed by "PP(p)TP"(*26, 27*). In

contrast, mutating the LxVP motif only modestly reduces affinity for calcineurin (~5X), indicating a smaller

role in binding. These data suggest that the PxIxIT motif primarily anchors the peptide, while

phosphorylation significantly enhances the overall affinity.

Using full length recombinant CLMB, NMR studies reveal that the binding interface for calcineurin extends

beyond the composite motif, with notable involvement of the N-terminal residues in interactions. However,

this interaction may differ in a cellular environment, where the N-terminus is anchored to the membrane

via N-myristoylation and S-acylation. At higher molar ratios of CLMB to calcineurin (CNA/B), NMR data

suggest that a second CLMB molecule binds, by utilizing the LxVP motif to bind its cognate stie at the

CNA/B heterodimeric interface. These findings are supported by *in vivo* data, showing that increasing

calcineurin expression enhanced levels of phosphorylated CLMB, which then decreased when CLMB

expression was further increased.

Based on all our data, we present two models for how CLMB may shape calcineurin activity *in vivo* (Figure

6). Dynamic S-acylation promotes CLMB association with membranes (Figure 6A, Step 1), where, when

phosphorylated, it recruits a localized pool of calcineurin via high affinity PxIxIT-mediated binding (Figure

6A, Step 2). In the presence of a $Ca^{2+}$ signal, $Ca^{2+}$ and CaM activate calcineurin which exposes the LxVP

docking site(*28*).  If CLMB is in excess relative to calcineurin and PxIxIT binding is fully saturated, a

second CLMB molecule will bind via its LxVP motif which positions pThr44 for dephosphorylation at the

active site (Figure 6A, Step 3).  As the population of CLMB shifts to the dephosphorylated form, decreased

PxIxIT binding affinity (Figure 3) promotes calcineurin dissociation from CLMB, allowing the enzyme to

engage and dephosphorylate other nearby LxVP- and PxIxIT-containing substrates (Figure 6A, Step 4). A

similar requirement for 'low' PxIxIT affinity was demonstrated for the another signaling scaffold, AKAP5,

which binds and concentrates both calcineurin and NFAT near L-type $Ca^{2+}$ channels in neurons to promote

NFAT dephosphorylation. In that case, increasing the affinity of the native PxIxIT in AKAP5 for calcineurin

~two-fold significantly impaired NFAT activation due to increased sequestering of calcineurin(*29*).

In our second model (Figure 6B), active calcineurin-pCLMB dimers (1 CLMB: 1 CNA/B) directly engage

and dephosphorylate substrates at the membrane that contain only an LxVP motif (Figure 6B, step 3). Many

candidate calcineurin substrates are predicted to contain only an LxVP, but if and how the majority are

dephosphorylated by calcineurin has yet to be uncovered. The best characterized example is the cAMP-

dependent protein kinase type II-alpha regulatory subunit (PRKAR2A), which depends on the PxIxIT motif

within its binding partner, the AKAP5 scaffold, to recruit calcineurin for its dephosphorylation(*9, 30*). In

our second model (Figure 6B), active calcineurin-pCLMB dimers (1 CLMB: 1 CNA/B) directly engage

and dephosphorylate such PxIxIT-less substrates at the membrane (Figure 6B, step 3). Simply recruiting

calcineurin to membranes, where association kinetics are 20-30 fold faster than in the cytosol of eukaryotic

cells, may enhance LxVP binding causing their efficient dephosphorylation despite lacking their own

PxIxIT motif(*31*). Such division of the PxIxIT and LxVP motifs among two molecules may also allow for

the extremely high affinity PxIxIT observed in CLMB. In substrates that contain both LxVP and PxIxIT

motifs, stable PxIxIT binding may limit enzymatic efficiency by slowing product release. However, acting

only as a scaffold, CLMB can remain tightly associated with calcineurin via PxIxIT-binding while other

substrates cycle in and out of the LxVP-binding surface and are dephosphorylated. In this scenario (Figure

6B), calcineurin remains at the membrane either until CLMB levels increase sufficiently to allow CLMB

dephosphorylation and calcineurin dissociation (Figure 6B step 4) or until CLMB is de-acylated. In both

models, intracellular $Ca^{2+}$ dynamics, CLMB protein levels, and S-acylation/de-acylation activities combine

to modulate calcineurin signaling. Future studies aim to identify the palmitoyl acyltransferases (PATs) and

acyl protein thioesterases (APTs) that dynamically regulate CLMB S-acylation to better understand how

these critical regulators contribute to calcineurin-CLMB signaling *in vivo*.

Another missing player in our model is the kinase(s) that phosphorylates CLMB. One candidate is the

stress-activated p38 mitogen-activated protein kinase (MAPK), which phosphorylates the PP(p)TP site in

Cabin1 and is predicted to phosphorylate Thr44 of CLMB (*32*). Because phosphorylation increases CLMB-

calcineurin affinity, kinase activation is required to prime regulation of calcineurin signaling by CLMB.

However, phosphorylation also establishes a negative feedback mechanism, with dephosphorylation

decreasing calcineurin's interaction with its substrate. Thus, regulators that modify kinase activity may also

significantly affect CLMB-calcineurin signaling dynamics.

Because the sequence in CLMB is the first composite motif to be analyzed, further studies are needed to

establish whether dephosphorylation via multimeric complexes is a general property of such motifs. Several

predicted LxVPxIxIT motifs do occur in ion channels, such as TRPM6, that assemble into higher order

complexes(*9, 33*). One possibility is that such composite motifs, which are present in at least thirty

additional human proteins, may in some cases provide a safety mechanism, by ensuring that calcineurin

dephosphorylates only multimeric substrates(*9*).

The major challenge that lies ahead is to identify the molecular targets and physiological functions that are

regulated by calcineurin-CLMB. CLMB increases proliferation and migration of some pancreatic ductal

adenocarcinoma (PDAC) derived cell lines, suggesting that CLMB shapes calcineurin signaling to promote

oncogenesis(*10*). Indeed, high CLMB expression is associated with poor patient outcomes for PDAC,

melanoma and head and neck cancer(*10, 11*). Mechanisms underlying these activities have not yet been

identified, but these cancers may provide good starting points for examining how CLMB modifies

calcineurin signaling in a biologically and clinically relevant context(*34*).

**Materials and Methods**

**Cell culture and transfection:**

Cells were cultured at 37 °C in 5% $CO_2$. HEK293 Flp-In TReX and MCF7 cell lines were grown in standard

media containing Dulbecco's modified eagle's medium (DMEM) with 4.5 mg/mL glucose, L-glutamine,

and sodium pyruvate (Cytiva SH30243.FS) supplemented with 10% fetal bovine serum (Corning

MT35010CV). Transfections were done according to manufacturer instructions. HEK293 cells were

transfected using lipofectamine 2000 (Thermo Fisher Scientific 11668019). MCF7 cells were transfected

using Transfex ATCC transfection reagent (ATCC ACS-4005).

**Stable cell line generation:**

On day one, $9.6 \times 10^5$ HEK293 Flp-In TReX cells were transfected using lipofectamine 2000 with 1.5 µg

pOG44 and 0. 5 µg pcDNA5 FRT TO encoding proteins of interest in a 6 well dish (plasmid # 2801, 2852,

2860, 2861, 2862, 2863, 2864, and 2880). Transfections were done according to manufacturer instructions.

On day two, transfected cells were passaged onto 10cm plates. On day three, selection was started using

media containing 3 µg blasticidin and 200 ng/mL hygromycin. Selection media was replaced every 2-3

days until distinct colonies appeared. Colonies were selected using cloning rings and expanded before being

stored in liquid nitrogen.

**Metabolic labeling to detect CLMB lipidation:**

HEK293 Flp-In TReX lines stably expressing $CLMB_{WT}$, $CLMB_{C7,14S}$, or $CLMB_{G2A}$ were induced with 10

ng/mL doxycycline for 24 hours and split evenly among three 10 cm tissue culture dishes. The following

day, cells were rinsed with PBS and incubated in DMEM containing 4.5 mg/mL glucose, L-glutamine,

sodium pyruvate, 2% FBS, and either 20 µM YnMyr (Click Chemistry Tools 1164-5), 30 µM 17-ODYA

(Cayman Chemical Company 34450-18-5), or vehicle (ethanol and DMSO) overnight. Cells were then co-

incubated with 20 µM YnMyr, 30 µM 17-ODYA, or vehicle and 20 µM MG132 (EMD Millipore 474787-

10MG) for 4 hours before washed twice in PBS and frozen at -80 °C. Cells were lysed in 400 µL TEA lysis

buffer (50 mM TEA pH 7.4, 150 mM NaCl, 1% triton X-100, and 1 mM PMSF) by inverting for 15 minutes

at 4 °C. Lysates were passed five times through a 27-gauge needle, centrifuged at 16,000 x *g* for 15 minutes

at 4 °C and protein content determined by Pierce BCA protein assay kit (Thermo Fisher Scientific 23225).

25 µg total lysate was removed as input and 0.5 mg total protein was incubated with pre-washed magnetic

M2 anti-FLAG (Sigma-Aldrich M8823-5ML) beads at 4 °C for 1.5 hours while inverting. Beads were

washed 3X with modified RIPA (50 mM TEA pH 7.4, 150 mM NaCl, 1% Triton X-100, 0.1% SDS, and

1% sodium deoxycholate) and resuspended in 40 µL 1x PBS. 10 µL of 5x click reaction mix (1x PBS, 0.5

mM biotin azide, 5 mM TCEP, 0.5 mM TBTA, 5 mM $CuSO_4$) was added and reactions incubated for 1 hour

at 24 °C, shaking at 550 RPM. Beads were washed 3X in modified RIPA and eluted 2X using 30 µL of 1

mg/mL FLAG peptide in modified RIPA buffer. 12 µL of non-reducing 6x SDS sample buffer was added

to eluate before heating at 95 °C for 2 minutes and analysis by SDS-PAGE. Note that addition of reducing

agents to SDS sample buffer results in loss of 17-ODYA signal. Samples were transferred to nitrocellulose

membrane and lipidation was detected using fluorophore-tagged streptavidin, and total CLMB with anti-

FLAG. Immunoblots were imaged using a Licor Odyssey CLx and quantified with Image Studio software.

**Metabolic pulse-chase labeling assay to assess dynamic CLMB S-acylation:**

HEK293 Flp-In TReX cells stably expressing CLMB$_{WT}$-FLAG were induced with 10 ng/mL doxycycline

for 48 hours. Cells were rinsed 2X in PBS and incubated in methionine free DMEM supplemented with 5%

charcoal-stripped FBS, 1 mM L-glutamate, 1 mM sodium pyruvate, 0.4 mM cysteine and 10 ng/mL

doxycycline along with vehicle (ethanol), 30 μM 17-ODYA, 50 μM L-AHA, or 17-ODYA and L-AHA

together. After two hours, cells were washed 2X in PBS and either harvested immediately or incubated in

standard media supplemented with 10 μM palmitic acid and 10 ng/mL doxycycline, with or without 10 μM

palmostatin B (EMD Millipore 178501-5MG). Cells were rinsed in PBS and frozen at -80 °C. Metabolic

labeling was done as described in the previous section with the exception that two rounds of labeling were

done. First, Cy5-azide (Click Chemistry Tools AZ118-1) was linked to 17-ODYA. Then, cells were washed

and L-AHA was linked to AF-488 (Click Chemistry Tools 1277-1) alkyne before being washed three

additional times and eluted using FLAG peptide. Samples were resolved via SDS-PAGE and in-gel

fluorescence was read using the GE Amersham Typhoon imaging system. Images were quantified in Image

Studio. The proportion of S-acylated CLMB is calculated as Cy5/A-488 normalized to the no chase control.

**Acyl-PEG exchange:**

HEK293 Flp-In TReX lines stably expressing CLMB$_{WT}$, CLMB$_{C7S}$, CLMB$_{C7,14S}$, or CLMB$_{G2A}$ were induced

with 10 ng/mL doxycycline for 48 hours. Cells were rinsed 2X in PBS and stored at -80 °C. Cells were

lysed in 400 μL lysis buffer (50 mM TEA pH 7.3, 150 mM NaCl, 2.5% SDS, 1 mM PMSF, 100 units

benzonase) at 37 °C for 20 minutes, shaking at 800 RPM. Lysates were passed 10X through a 27-gauge

needle, centrifuged at 16,000 x $g$ for 5 minutes and protein quantified by BCA. Lysates were adjusted to 2

mg/mL and incubated with 10 mM TCEP for 30 minutes at room temperature. Cysteines were protected by

incubating with 25 mM NEM for 2 hours at 40 °C, shaking at 800RPM. Protein was precipitated over night

with four volumes of acetone at -20 °C, centrifuged at 16,000 x $g$ for 20 minutes at 4 °C and washed 4X in

70% acetone. Acetone was removed, pellets air-dried to remove all traces of acetone and resuspended in 75

μL Buffer A (50 mM TEA pH 7.3, 150 mM NaCl, 4% SDS, 0.2% triton X-100, 4 mM EDTA) for 1 hour at

40 °C, shaking at 800 RPM. Samples were centrifuged at 16,000 x $g$ for 2 minutes, 7 μL removed as input,

and the remainder split into +cleavage and -cleavage fractions. The +cleavage sample containing 1% SDS

was treated with 750 µM hydroxylamine (final concentration) for 1 hour at 30 °C, rotating end over end.

Following methanol-chloroform precipitation (ice cold, final concentrations 42% methanol, 16%

chloroform) samples were centrifuged at 16,000 x $g$ for 20 minutes at 4 °C, aqueous layer removed, washed

3X with 100% methanol and air dried. Pellets were resuspended as described in Buffer A and incubated in

1 mM mPEG-Mal (Methoxypolyethylene glycol maleimide, 5 kDa, 63187 Sigma-Aldrich) for 2 hours at

room temp, rotating end over end. Samples were methanol-chloroform precipitated as described,

resuspended in Lysis buffer with 1X SDS-PAGE sample buffer, resolved using SDS-PAGE and visualized

via immunoblotting with mouse anti-FLAG to detect CLMB-FLAG. Endogenous calnexin was detected

using anti-calnexin antibodies. Immunoblots were imaged using a Licor Odyssey CLx and quantified with

Image Studio software.

**Detergent-assisted subcellular fractionation:**

HEK293 Flp-In TReX lines stably expressing CLMB$_{WT}$, CLMB$_{C7,14S}$, or CLMB$_{G2A}$ were induced with 10

ng/mL doxycycline for 48 hours and 20 µM MG132 (EDM Millipore 474787-10MG) for 4 hours prior to

harvesting. Cells were washed 2X in PBS, harvested and immediately lysed in 600 µL of Cytoplasm Buffer

(10 mM HEPES pH 6.8, 100 mM NaCl, 300 mM sucrose, 3 mM MgCl$_2$, 5 mM EDTA, 0.015% digitonin,

1x halt protease inhibitor cocktail (Thermo Fischer scientific 78442)) at 4 °C for 15 minutes. 60 µL of total

lysate was removed as input and the remainder centrifuged at 1000 x $g$ for 10 min at 4 °C. The supernatant

was removed as cytoplasmic fraction and the pellet washed 2X in PBS and resuspended in 600 µL

membrane buffer (10 mM HEPES pH 7.5, 100 mM NaCl, 300 mM sucrose, 3 mM MgCl$_2$, 3 mM EDTA,

0.5% Triton X-100, 1x halt protease inhibitor cocktail) for 30 minutes at 4 °C turning end over end.

Insoluble material was pelleted by centrifugation at 5000 x $g$ for 10 minutes at 4 °C and supernatant

removed as membrane fraction. Both fractions were centrifuged at 12,000 x $g$ at 4 °C for 10 minutes.

Samples were heated in 1X SDS sample buffer at 95 °C for 5 minutes, resolved via SDS-PAGE,

immunoblotted and CLMB-FLAG detected using anti-FLAG antibodies; anti-GM130 and anti-α-tubulin

were used as membrane and cytosolic markers respectively. Immunoblots were imaged using a Licor

Odyssey CLx and quantified with Image Studio software. CLMB in cytosolic and membrane fractions were

quantified as $CLMB_{Cyto}/(CLMB_{Cyto}+ CLMB_{Mem})$ and $CLMB_{Mem}/(CLMB_{Cyto} + CLMB_{Mem})$, respectively.

**Immunofluorescence, microscopy, and image analysis:**

MCF7 cells were co-transfected (as described previously) with equal amounts of pcDNA5 encoding

$CLMB_{WT}$, $CLMB_{IxITMut}$, $CLMB_{LxVMut}$, $CLMB_{T44A}$ and pcDNA5 encoding CNA-GFP. The following day

cells were plated on poly-lysine coated 12 mm, #1.5H glass coverslips (ThorLabs). On day three, cells were

washed in PBS and fixed in 100% methanol for 10 minutes at -20 °C. Coverslips were washed 3 times in

1x PBS, blocked (1x PBS, 200 mM glycine, 2.5% FBS) for 30 minutes at room temperature, then incubated

in primary antibody diluted in blocking buffer for 1 hour before being washed 3 times in 1x PBS and

incubated for 1 hour in secondary antibody solution. Following three additional washes, coverslips were

mounted using Prolong Glass mountant with NucBlue stain. After sitting overnight at room temperature,

coverslips were imaged using a Lionheart FX automated widefield microscope with a 20X Plan Fluorite

WD 6.6 NP 0.45 objective. For each coverslip, a 7 by 7 grid of non-overlapping images was automatically

taken. All images were manually assessed and those with defects (debris, bubbles, out-of-focus, etc.) were

filtered out. Images were analyzed using an automated pipeline using CellProfiler(*35*). Briefly, cells were

identified by their nuclei and all those lacking CLMB-FLAG staining removed. CLMB-FLAG signal was

used to define the cell body and all cells containing one or more fully saturated pixels in the CLMB-FLAG

or CNA-GFP channels were removed. Pearson's correlation coefficient between CLMB-FLAG and CNA-

GFP was determined for each individual cell. >500 cells per condition were analyzed.

**GFP-Trap coimmunoprecipitation:**

HEK293 cells were co-transfected (as described previously) with pcDNA5 encoding CNA-GFP and

pcDNA5 encoding $CLMB_{WT}$, $CLMB_{IxITMut}$, $CLMB_{LxVMut}$, $CLMB_{T44A}$, or $CLMB_{LxVMut+T44A}$. 24 hours post-

transfection, cells were washed 2X in PBS and frozen at -80 °C. Cells were lysed in 500 µL lysis buffer (50

mM Tris pH 7.5, 150 mM NaCl, 1% NP-40), aspirated 5X with a 27-gauge needle, centrifuged at 16,000 x

$g$ for 20 min at 4 °C and protein quantified via BCA. 100 mg total protein was removed as input. Lysates

were adjusted to 0.5% NP-40 and 500 µg of protein incubated with pre-washed magnetic GFP-TRAP beads

(ChromoTek gtma-20) for 2 hours at 4 °C, turning end over end. Beads were magnetized, washed 3 times

and eluted with 50 µL of 2x SDS PAGE sample buffer. Samples were resolved via SDS-PAGE and

visualized via immunoblotting with anti-flag antibodies to detect CLMB-FLAG and anti-GFP to detect

CNA-GFP. Anti-GAPDH was used as a loading control for inputs. Immunoblots were imaged using a Licor

Odyssey CLx and CLMB-calcineurin binding was quantified with Image Studio software. Co-purified

CLMB was normalized using the following: $(CLMB_{IP}/CLMB_{Input})/GFP-CNA_{IP}$. Additionally, all values are

displayed relative to $CLMB_{WT}$ which is set at 1.

**Protein expression and purification:**

CNA catalytic dead mutant H151Q with an N-terminal $His_6$-CNA (residues 1–391) and CNB (residues 1–

170) subunits were expressed using *E. coli* BL21(DE3) cells in Lysogeny Broth (LB) supplemented with

50 µg/mL ampicillin. Protein expression was induced with 1 mM isopropyl β-d-1-thiogalactopyranoside

(IPTG) at 18 °C. Cells were lysed using a Fisherbrand™ Model 505 Sonic Dismembrator and lysates were

centrifuged at $46,000 \times g$ at 4 °C for 45 minutes. The protein was purified by incubation with nickel-

nitriloacetic acid (Ni-NTA) agarose resin (Qiagen) for 3h at 4 °C. The resin was washed with 50 mM Tris

pH 8.0, 250 mM NaCl, 20 mM imidazole, and 1 mM TCEP. A buffer containing 50 mM Tris pH 8.0, 250

mM NaCl, 300 mM imidazole, and 1 mM TCEP was used for elution. Calcineurin was further purified by

SEC (Superdex® 200 Increase 10/300 GL) using Phosphate-buffered saline (PBS) and $CaCl_2$ 1mM as the

final buffer.

**Fluorescence polarization binding assay:**

All FP measurements were performed on a multifunctional microplate reader (EnVision, Perkin Elmer) in

black 384-well microplates (Corning, Cat No 3575) with 30 µL of the assay solution per well.  480-nm

excitation and 535-nm emission filters were used for the FP measurements. In the FP saturation binding

experiments, 5 nM FITC labeled peptides were mixed with increasing concentrations of calcineurin in PBS

and 500 μM CaCl$_2$ buffer (from 0 to 25 μM). The FP values were plotted against the log of the protein

concentrations, and the dissociation constant (apparent $K_D$) was obtained from the resulting sigmoidal curve

as analyzed in GraphPad Prism 10.

**Flourescence polarization competition assay:**

For the competitive binding assay, a mixture containing 5 nM FITC-labeled CLMB peptides and CNA/B

(0.08 and 2 μM for CLMB vs PVIVIT experiments) was incubated with serial dilutions of unlabeled

peptides for 30 minutes at room temperature. The FP values were determined, and the IC$_{50}$ values, i.e., the

concentrations required for 50% displacement of the FITC labeled peptide, were calculated in GraphPad

Prism 10. The $K_i$ values of competitive inhibitors were calculated based on a previously reported

method(*36*), further adapted by(*37*). Briefly, the relation of $K_i$ and IC$_{50}$ in competitive inhibition as applied

to protein-ligand–inhibitor (P–L–I) interactions, can be summarized by the following equation:

471
$$K_i = \frac{IC_{50}}{(\frac{L}{K_d} + 1)}$$

Where L stands for the concentration of ligand, i.e., the FITC-labeled peptide used in the FP binding assay;

and $K_D$ corresponds to the dissociation constant of the same FITC-labeled peptide resulting from the

performed FP binding assay.

**ITC measurements:**

Isothermal titration calorimetry (ITC) experiments were performed using an Affinity ITC from TA

Instruments (New Castle, DE) equipped with an autosampler in a buffer containing 50 mM HEPES, pH

8.0, 150 mM NaCl, 1 mM TCEP, 500μM CaCl$_2$ at 25 °C. 40 μM protein solution in the calorimetric cell

was titrated with 100 μM peptide solution using 0.5 μL injection in 200-sec intervals using stirring speed

at 125rpm. The resulting isotherm was fitted with a single site model to yield thermodynamic parameters

of ΔH, ΔS, stoichiometry, and $K_D$ using NanoAnalyze software (TA instruments).

**NMR analyses:**

All NMR data were acquired using TopSpin3.4 (Bruker), processed with NMRPipe v.2020/2021 and

analyzed with CCPNmr v3(*38, 39*). Structural biology applications were compiled by the SBGrid

consortium. Three-dimensional (3D) NMR experiments were conducted on a Bruker spectrometer

operating at 600 MHz, equipped with TCI cryoprobes and z-shielded gradients. Samples of

approximately180-μM 2H-, $^{13}$C- and $^{15}$N-calcimembrin in 20 mM Hepes pH 6.9, 100 mM NaCl, 0.5 mM

EDTA, 1 mM TCEP and 5% v/v $^2$H2O were used. Temperature was set to 25˚C. Chemical shift assignments

are deposited in the Biological Magnetic Resonance Bank (BMRB) under accession code 52641.

The NMR titration samples were prepared by adding unlabeled calcineurin A/B complex or unlabeled

calcineurin subunit A into solutions of U-[$^{15}$N]-labeled calcimembrin (50 μM) to reach a ratio of 0.5:1 (25

μM unlabeled calcineurin) or 1:1 (50 μM unlabeled calcineurin). Two-dimensional (2D) spectra were

acquired using SOFAST-HMQC on a Bruker 800 MHz Avance III spectrometer equipped with a TXO

cryoprobe and z-shielded gradients(*40*). Data was processed using NmrPipe and analyzed with CCPNmr.

Secondary structure prediction was analyzed using Talos-N(*22*).

**AlphaFold 3 modeling:**

Calcineurin-CLMB complex modeling was done using AlphaFold 3 on AlphaFold server(*24*). Modeling

was done using truncated $CNA_{1-491}$, CNB, and two copies of CLMB phosphorylated at Thr44. $Zn^{2+}$ and $Fe^{3+}$

(found in the calcineurin active site) and four $Ca^{2+}$ ions (bound to CNB) were included in modeling. All

images were generated in UCSF ChimeraX 1.8(*41*).

*In vivo* **analysis of CLMB phosphorylation status:**

HEK293 FLIP-In TreX cells stably expressing CLMB$_{WT}$-FLAG, CLMB$_{IxITMut}$-FLAG, CLMB$_{LxVMut}$-FLAG, or CLMB$_{T44A}$-FLAG were transfected with empty vector or pcDNA5 FRT TO encoding CNA-GFP alongside empty vector or pcDNA5 FRT TO encoding CLMB$_{WT}$-V5, CLMB$_{IxITMut}$-V5, or CLMB$_{LxVMut}$-V5. Protein expression was induced for 24 hours with 10 ng/mL doxycycline, cells washed 2X in PBS and stored at -80 °C. Cells were lysed (50 mM Tris pH 8, 150 mM NaCl, 1% NP-40, 0.1% SDS, 0.5% sodium dodecyl sulfate, 1x Halt protease inhibitor cocktail) and resolved via SDS-PAGE and detected via immunoblotting.

***In vitro* CLMB phospho-peptide dephosphorylation:**

CLMB$_{WT}$, CLMB$_{IxITMut}$, and CLMB$_{LxVMut}$ phospho-peptides were synthesized by Vivitide, resuspended in 10 mM Tris pH 8.0 and dephosphorylated at 26 °C in reaction buffer (50 mM Tris pH 8.0, 100 mM NaCl, 6 mM MgCl$_2$, 2 mM CaCl$_2$, 1 mM EGTA, 0.1 mg/mL BSA, 1 mM DTT, 0.5 µM MDCC-PBP (Thermo Scientific PV4406), 1 µM CaM (Millipore Sigma 208694-1MG)). Reactions were initiated by addition of CLMB peptide to a final concentration of 200nM. Reactions took place in a total of 50 µL in a corning black 96 well vessel and were monitored continuously via fluorescence emitted by phosphate bound MDCC-PBP (EX:425/ EM:460). Phosphate standards were used to convert fluorescence intensity to phosphate concentration. Data were analyzed using Prism software and the rate of phosphate production was obtained by fitting data to a single exponential.

**Kinetic Modeling:**

Dephosphorylation data were fit by simulation and non-linear regression using Kintek Explorer™. Sigma values for each individual replicate were obtained using the double exponential afit function. All replicates were simultaneously fit to the models described in supplemental figures 2A and 2B. Curve offsets capped at a maximum offset of 50nM. Substrate concentration was allowed to vary by 15% during fitting. Offsets and substrate concentrations for each curve and fit quality metrics are displayed in Tables S1-S3. Final graphs display averaged (n $\geq$ 5) phosphate released at each timepoint with the corresponding fit.

**Statistical analysis:**

Statistical analysis was done using Graphpad Prism version 10.2.2. All data is displayed as representative

images or mean values with standard deviation error bars unless otherwise noted. All data represent three

independent experiments unless otherwise indicated in figure legends. Image analysis of CLMB-calcineurin

colocalization was conducted using the following number of cells: $CLMB_{WT} = 525$, $CLMB_{IxITMut} = 641$,

$CLMB_{LxVMut} = 500$, $CLMB_{T44A} = 517$. Statistical significance was calculated using one-way analysis of

variance (ANOVA) with corrections for multiple comparisons as indicated.

**Data availability:**

All original scans of immunoblots, immunofluorescence images, and AlphaFold predictions can be

accessed through Mendeley Data (to be made available upon publication). Plasmids and reagents will be

shared upon request. NMR chemical shift assignments are deposited in the Biological Magnetic Resonance

Bank (BMRB) under accession code 52641.

**Acknowledgements:**

We thank Norman Davey and members of the Cyert and Arthanari labs, in particular Angela Barth and

Sneha Roy, for helpful discussion. We are grateful to Callie Wigington and Makena Pule for their early

contributions to the project and to Rick Russell for feedback on the kinetic modeling. We thank Ilya

Bezprozvanny for suggesting the name, calcimembrin.

**Funding:**

MSC and DAB are supported by NIH grant R35 GM136243. DAB also acknowledges support from NIH

grant T32 GM007276. HA, JCR, SQ and TV acknowledge grant R01 GM136859 from NIH.

**Author Contributions:**

MSC, DAB, HA and JCR conceived the studies. SQ and TV provided early NMR analysis of CLMB-

calcineurin complexes. DAB, supervised by MSC, carried out experiments and analyzed data for Figures

1, 2, and 5, and prepared figures. JCR, supervised by HA, carried out experiments and analyzed data for

Figures 3 and 4. TV carried out secondary structure analyses. DAB and MSC wrote the manuscript in

consultation with HA and JCR.

**References:**

1.  M. D. Bootman, G. Bultynck, Fundamentals of Cellular Calcium Signaling: A Primer.
*Cold Spring Harb Perspect Biol* **12**, (2020).
2.  I. Ulengin-Talkish, M. S. Cyert, A cellular atlas of calcineurin signaling. *Biochim Biophys*
*Acta Mol Cell Res* **1870**, 119366 (2023).
3.  J. Aramburu, A. Rao, C. B. Klee, Calcineurin: from structure to function. *Curr Top Cell*
*Regul* **36**, 237-295 (2000).
4.  S. Grigoriu *et al.*, The molecular mechanism of substrate engagement and
immunosuppressant inhibition of calcineurin. *PLoS biology* **11**, e1001492 (2013).
5.  J. Roy, M. S. Cyert, Identifying New Substrates and Functions for an Old Enzyme:
Calcineurin. *Cold Spring Harb Perspect Biol* **12**, (2020).
6.  B. L. Brauer *et al.*, Leveraging New Definitions of the LxVP SLiM To Discover Novel
Calcineurin Regulators and Substrates. *ACS Chem Biol* **14**, 2672-2682 (2019).
7.  S. R. Sheftic, R. Page, W. Peti, Investigating the human Calcineurin Interaction Network
using the piLxVP SLiM. *Scientific reports* **6**, 38920 (2016).
8.  E. Tsekitsidou *et al.*, Calcineurin associates with centrosomes and regulates cilia length
maintenance. *J Cell Sci* **136**, (2023).
9.  C. P. Wigington *et al.*, Systematic Discovery of Short Linear Motifs Decodes Calcineurin
Phosphatase Signaling. *Mol Cell* **79**, 342-358 e312 (2020).
10. T. Nakamura *et al.*, Overexpression of C16orf74 is involved in aggressive pancreatic
cancers. *Oncotarget* **8**, 50460-50475 (2017).
11. M. Uhlen *et al.*, Towards a knowledge-based Human Protein Atlas. *Nat Biotechnol* **28**,
1248-1250 (2010).
12. T. Meinnel, C. Dian, C. Giglione, Myristoylation, an Ancient Protein Modification
Mirroring Eukaryogenesis and Evolution. *Trends Biochem Sci* **45**, 619-632 (2020).
13. M. Blanc, F. P. A. David, F. G. van der Goot, SwissPalm 2: Protein S-Palmitoylation
Database. *Methods Mol Biol* **2009**, 203-214 (2019).
14. M. C. Yap *et al.*, Rapid and selective detection of fatty acylated proteins using omega-
alkynyl-fatty acids and click chemistry. *Journal of lipid research* **51**, 1566-1580 (2010).
15. A. Percher *et al.*, Mass-tag labeling reveals site-specific and endogenous levels of protein
S-fatty acylation. *Proc Natl Acad Sci U S A* **113**, 4302-4307 (2016).
16. M. F.S. *et al.*, Mechanisms and functions of protein S-acylation. *Nat Rev Mol Cell Biol*,
(2024).
17. F. J. Dekker *et al.*, Small-molecule inhibition of APT1 affects Ras localization and
signaling. *Nat Chem Biol* **6**, 449-456 (2010).

18.     D. K. Blumenthal, K. Takio, R. S. Hansen, E. G. Krebs, Dephosphorylation of cAMP-
599             dependent protein kinase regulatory subunit (type II) by calmodulin-dependent protein
600             phosphatase. Determinants of substrate specificity. *J Biol Chem* **261**, 8140-8145 (1986).

601     19.     B. A. Perrino, L. Y. Ng, T. R. Soderling, Calcium regulation of calcineurin phosphatase
602             activity by its B subunit and calmodulin. Role of the autoinhibitory domain. *J Biol Chem*
603             **270**, 340-346 (1995).

604     20.     J. Aramburu *et al.*, Affinity-driven peptide selection of an NFAT inhibitor more selective
605             than cyclosporin A. *Science* **285**, 2129-2133 (1999).

606     21.     H. Li, L. Zhang, A. Rao, S. C. Harrison, P. G. Hogan, Structure of calcineurin in complex
607             with PVIVIT peptide: portrait of a low-affinity signalling interaction. *J Mol Biol* **369**,
608             1296-1306 (2007).

609     22.     Y. Shen, A. Bax, Protein structural information derived from NMR chemical shift with
610             the neural network program TALOS-N. *Methods Mol Biol* **1260**, 17-32 (2015).

611     23.     R. Hendus-Altenburger *et al.*, Molecular basis for the binding and selective
612             dephosphorylation of Na(+)/H(+) exchanger 1 by calcineurin. *Nat Commun* **10**, 3489
613             (2019).

614     24.     J. Abramson *et al.*, Accurate structure prediction of biomolecular interactions with
615             AlphaFold 3. *Nature* **630**, 493-500 (2024).

616     25.     H. Li, A. Rao, P. G. Hogan, Structural delineation of the calcineurin-NFAT interaction
617             and its parallels to PP1 targeting interactions. *J Mol Biol* **342**, 1659-1674 (2004).

618     26.     S. Lee *et al.*, The optimized core peptide derived from CABIN1 efficiently inhibits
619             calcineurin-mediated T-cell activation. *Exp Mol Med* **54**, 613-625 (2022).

620     27.     H. Q. Nguyen *et al.*, Quantitative mapping of protein-peptide affinity landscapes using
621             spectrally encoded beads. *Elife* **8**, (2019).

622     28.     S. J. Li *et al.*, Cooperative autoinhibition and multi-level activation mechanisms of
623             calcineurin. *Cell research* **26**, 336-349 (2016).

624     29.     H. Li *et al.*, Balanced interactions of calcineurin with AKAP79 regulate Ca2+-
625             calcineurin-NFAT signaling. *Nature structural & molecular biology* **19**, 337-345 (2012).

626     30.     T. W. Church *et al.*, AKAP79 enables calcineurin to directly suppress protein kinase A
627             activity. *Elife* **10**, (2021).

628     31.     W. Y. C. Huang, S. G. Boxer, J. E. Ferrell, Jr., Membrane localization accelerates
629             association under conditions relevant to cellular signaling. *Proc Natl Acad Sci U S A* **121**,
630             e2319491121 (2024).

631     32.     J. L. Johnson *et al.*, An atlas of substrate specificities for the human serine/threonine
632             kinome. *Nature* **613**, 759-766 (2023).

633     33.     C. H. Huang, Y. C. Chen, C. C. Chen, Physical interaction between calcineurin and
634             Cav3.2 T-type Ca2+ channel modulates their functions. *FEBS Lett* **587**, 1723-1730
635             (2013).

636     34.     A. S. Manalan, C. B. Klee, Activation of calcineurin by limited proteolysis. *Proc Natl*
637             *Acad Sci U S A* **80**, 4291-4295 (1983).

638     35.     D. R. Stirling *et al.*, CellProfiler 4: improvements in speed, utility and usability. *BMC*
639             *Bioinformatics* **22**, 433 (2021).

640     36.     Y. Cheng, W. H. Prusoff, Relationship between the inhibition constant (K1) and the
641             concentration of inhibitor which causes 50 per cent inhibition (I50) of an enzymatic
642             reaction. *Biochem Pharmacol* **22**, 3099-3108 (1973).

37.  R. Z. Cer, U. Mudunuri, R. Stephens, F. J. Lebeda, IC50-to-Ki: a web-based tool for
converting IC50 to Ki values for inhibitors of enzyme activity and ligand binding.
*Nucleic acids research* **37**, W441-445 (2009).
38.  F. Delaglio *et al.*, NMRPipe: a multidimensional spectral processing system based on
UNIX pipes. *J Biomol NMR* **6**, 277-293 (1995).
39.  W. F. Vranken *et al.*, The CCPN data model for NMR spectroscopy: development of a
software pipeline. *Proteins* **59**, 687-696 (2005).
40.  P. Schanda, E. Kupce, B. Brutscher, SOFAST-HMQC experiments for recording two-
dimensional heteronuclear correlation spectra of proteins within a few seconds. *J Biomol
NMR* **33**, 199-211 (2005).
41.  E. C. Meng *et al.*, UCSF ChimeraX: Tools for structure building and analysis. *Protein Sci*
**32**, e4792 (2023).

**Figure 1: Calcimembrin localizes to membranes via protein lipidation.**

**a** Schematic of CLMB. N-myristoylation (red); S-acylated cysteines (yellow); phospho-Thr44 (green); Calcineurin binding motifs LxVP (pink) and PxIxIT (purple). Mutant sequences are shown for CLMB$_{\text{LxVMut}}$ and CLMB$_{\text{IxITMut}}$.

**b** CLMB incorporates analogs of palmitate (17-ODYA) or myristate (YnMyr) as indicated. Cells were treated with 20 µM MG132 prior to harvesting. Representative immunoblot shown of FLAG tagged CLMB wild type (CLMB$_{\text{WT}}$), S-acylation defective (CLMB$_{\text{C7,14S}}$) or myristoylation defective (CLMB$_{\text{G2A}}$), immunoprecipitated and detected using CLICK chemistry with biotin-azide followed by detection with IRdye 800CW streptavidin (n = 3 independent experiments). Asterisks indicate CLMB dimers formed due to absence of reducing agents (see methods).

**c** Representative immunoblot of acyl-PEG exchange performed on lysates expressing CLMB$_{\text{WT}}$-FLAG or lipidation mutants (CLMB$_{\text{C7S}}$, CLMB$_{\text{C7,14S}}$, CLMB$_{\text{G2A}}$). PEGylation events, a proxy for S-acylation, are indicated by asterisks. For each sample, input, cleaved (NH$_2$OH) and PEGylated (mPEG), and non-cleaved PEGylated conditions are shown. Cells were treated with 20 µM MG132 for 4 hours prior to harvesting. Endogenous, dually S-acylated calnexin is a positive control (n = 3 independent experiments).

**d** Pulse chase analysis of CLMB-FLAG S-acylation turnover using analogs of palmitate (17-ODYA) or methionine (L-AHA). Incorporation into immunoprecipitated CLMB-FLAG in the absence of analog (Neg Ctrl), or after 2 hours of analog treatment followed by 0 or 30 minutes without analog including DMSO (control) or palmostatin B. 17-ODYA or L-AHA were detected using CLICK with Cy5 and AZDye 488, respectively. A representative in-gel fluorescence scan is shown (n = 3 independent experiments).

**e** Data from (**d**) showing mean ± SEM of 17-ODYA signal normalized to L-AHA. Indicated p-values calculated by two-way ANOVA corrected by Tukey's multiple comparison test. N=3 independent experiments.

**f** Sub-cellular fractionation of lysates containing CLMB$_{\text{WT}}$-FLAG or lipidation mutants (CLMB$_{\text{C7,14S}}$, CLMB$_{\text{G2A}}$). Cells were treated with 20 µM MG132 for 4 hours prior to harvesting. Representative immunoblot showing input, cytoplasmic (cyto), and membrane (mem) fractions. GM130 and α-tubulin mark membrane and cytosolic fractions respectively.

**g** Data from (**f**) showing mean ± SEM cytoplasmic or membrane signals normalized to total (cytoplasmic + membrane signals). Indicated p-values calculated by two-way ANOVA corrected by Tukey's multiple comparison test. N= 3 independent experiments.

**Figure 2: Composite SLiM in calcimembrin mediates binding to and dephosphorylation by calcineurin.**

**a** Representative immunoblot of anti-GFP purification from lysates of HEK293 cells co-transfected with CNA-GFP and CLMB-FLAG (CLMB$_{WT}$, CLMB$_{IxITMut}$, CLMB$_{LxVMut}$, CLMB$_{T44A}$, or CLMB$_{LxVMut+T44A}$).

**b** Data from (**a**) showing mean ± SEM CLMB-FLAG pulled down normalized to CLMB-FLAG input levels and CNA-GFP pulled down. Indicated p-values calculated by two-way ANOVA corrected by Dunnett's multiple comparison test (n = 3 independent experiments).

**c** Representative immunofluorescence images of fixed MCF7 cells co-transfected with CLMB-FLAG (CLMB$_{WT}$, CLMB$_{IxITMut}$, CLMB$_{LxVMut}$, or CLMB$_{T44A}$) and CNA-GFP. CLMB was visualized using anti-FLAG, calcineurin with anti-GFP, and nuclei with Hoechst 33342. Scale bar = 50 μm.

**d** Data from (**c**) showing median with 95% CI Pearson's correlation coefficient of CLMB-calcineurin colocalization. Each point represents a single cell (n ≥ 500 for each condition). Indicated p-values calculated by Kruskal-Wallis with Dunn's multiple comparisons correction.

**e** Plot of phosphate released from 200 nM pCLMB$_{WT}$ (blue circles) or pCLMB$_{LxVMut}$ (pink squares) phospho-peptides dephosphorylated by 20 nM CNA/B and 1 μM CaM. Displayed is the mean ± SEM of phosphate released for ≥ 4 replicates at each timepoint.

A

$K_D(M) = 8.29 \pm 5.71\ E\text{-}09$
$n = 0.96 \pm 0.02$

$\Delta H\ (kcal/mol) = -12.96 \pm 0.66$
$\Delta S\ (kcal/mol) = -6.48$

B

FITC-pCLMB$_{WT}$
FITC-CLMB$_{WT}$

C

FITC-VIVIT Peptide

D

FITC-pCLMB$_{WT}$ Peptide

pCLMB$_{WT}$   CLMB$_{WT}$
pCLMB$_{IxITMut}$   pCLMB$_{LxVMut}$

E

| Peptide Competitor | FITC-VIVIT | | FITC-pCLMB$_{WT}$ | |
|---|---|---|---|---|
| | IC$_{50}$(nM) | K$_i$(nM) | IC$_{50}$(nM) | K$_i$(nM) |
| pCLMB$_{WT}$ | 81.4 ± 16.2 | 80.9 ± 16.1 | 18.9 ± 1.2 | 13.4 ± 0.8 |
| pCLMB$_{IxITMut}$ | NA | NA | 6687 ± 878 | 4754 ± 624 |
| pCLMB$_{LxVMut}$ | 159 ± 30 | 158 ± 30 | 103 ± 9 | 73.2 ± 6.4 |
| CLMB$_{WT}$ | 267 ± 21 | 265 ± 21 | 585 ± 57 | 426 ± 40 |

**Figure 3: Phosphorylation and the LxVP motif contribute to PxIxIT binding.**

**a** Isothermal calorimetric enthalpy change displaying the thermometric evaluation of the affinity between CNA/B complex and unlabeled pWT peptide (top), and binding isotherm obtained by titration of unlabeled pWT into CNA/B complex (40 µM) (bottom). A dissociation constant $(K_D) = 8.3 \pm 5.7$ nM was obtained.

**b** Fluorescence polarization (FP) saturation binding curves of two different FITC-labeled peptide tracers (5nM), FITC-pCLMB$_{WT}$ and FITC-CLMB$_{WT}$ were incubated with serial dilutions of CNA/B (0 to 12.5µM). The FP values were recorded at room temperature after a 30 minute incubation time. FP data were fitted to a one-site binding model (mean $\pm$ s.d., n = 2 independent experiments) using GraphPad Prism 10.0.

**c** Competitive binding assay of unlabeled pCLMB$_{WT}$, CLMB$_{WT}$, pCLMB$_{IxITMut}$, and pCLMB$_{LxVMut}$ peptides. The unlabeled peptides were titrated against a constant concentration of FP probe (FITC-PVIVIT peptide, 5 nM) in the presence of 2 µM CNA/B. Data (mean $\pm$ s.d., n = 2 independent experiments) were analyzed in GraphPad Prism 10.0.

**d** Competitive binding assay of unlabeled pCLMB$_{WT}$, CLMB$_{WT}$, pCLMB$_{IxITMut}$, and pCLMB$_{LxVMut}$ peptides. The peptides were titrated against a constant concentration of FP probe (FITC-pCLMB$_{WT}$ peptide, 5 nM) in the presence of 0.08 µM CNA/B . Data (mean $\pm$ s.d., n = 2 independent experiments) were analyzed in GraphPad Prism 10.0.

**e** Table displaying the experimentally obtained IC$_{50}$ values and converted K$_i$ values resulting from the competitive binding assay of unlabeled pCLMB$_{WT}$, CLMB$_{WT}$, pCLMB$_{IxITMut}$, and pCLMB$_{LxVMut}$ peptides. The unlabeled peptides were titrated against a constant concentration of FP probe (FITC-pCLMB$_{WT}$ or FITC-PVIVIT, 5 nM) in the presence of 0.08 µM and 2µM CNA/B, respectively. Data (mean $\pm$ s.d., n = 2 independent experiments) were analyzed in GraphPad Prism 10.0.

A

B  2 CLMB: 1 CNA

C  1 CLMB: 1 CNA

D  2 CLMB: 1 CNA/B

E  1 CLMB: 1 CNA/B

**Figure 4**: **CLMB-CN binding extends beyond the LxVP and PxIxIT motifs.**

**a** $^{15}$N HSQC spectra of free CLMB with the resonance assignment (red).

**b** NMR spectral analysis of the interaction of CLMB with CNA at 2:1 (CLMB: CNA); **c** with CNA at 1:1 (CLMB: CNA); **d** with CNA/B complex at 2:1 (CLMB: CNA/B); **e** with CNA/B at 1:1 (CLMB: CNA/B). The graphs display the intensity ratios of the CLMB resonances with and without the titrant (CNA or CNA/B) at different molar ratios. The ratios corresponding to the LxV motif, represented in pink, and IxIT motif, in purple. Residues where we cannot access the intensity ratio are represented with negative numbers. Peaks resulting in complete broadening are shown as negative bars in green ($I/I_0 = -0.4$). Residues that were not assigned in the apo CLMB are shown in blue ($I/I_0 = -0.2$), prolines that are not represented in a $^{15}$N-HSQC are shown in grey ($I/I_0 = -0.2$), and residues where we cannot unambiguously assign the bound resonance is shown in red, ($I/I_0 = -0.2$). Residues that have an overlap in the resonance and hence when one of the resonances broadens results in $I/I_0$ higher than 1 is marked with a * sign.

**Figure 5: Calcineurin dephosphorylates calcimembrin by forming multimers.**

**a** Representative immunoblot of CLMB-FLAG phosphorylation-induced mobility shifts induced by calcineurin overexpression. Performed using lysates from Flp-In-T-REx HEK cells inducibly expressing CLMB$_{WT}$-FLAG, CLMB$_{IxITMut}$-FLAG, or CLMB$_{LxVMut}$-FLAG and transfected with empty vector, CNA-GFP, or CNA-GFP with CLMB-V5 (CLMB$_{WT}$, CLMB$_{IxITMut}$, CLMB$_{LxVMut}$). Asterisks indicate phosphorylated CLMB.

**b** Data from (**a**) showing mean ± SEM phospho-CLMB-FLAG normalized to total CLMB-FLAG. Displayed p-values calculated by two-way ANOVA corrected by Dunnett's multiple comparison test (n = 3 independent experiments).

**c** Cartoon representation of calcineurin-CLMB complex formation and CLMB dephosphorylation under the conditions used in (**a**), i.e. CLMB-FLAG (blue) overexpression, CLMB-FLAG (blue with "F") and calcineurin overexpression, or CLMB-FLAG, calcineurin, and CLMB-V5 (green with "V") overexpression. Cartoons depict complexes formed by CLMB$_{WT}$-FLAG, CLMB$_{IxITMut}$-FLAG, and CLMB$_{LxVMut}$-FLAG with red X's to indicate motif mutation. Black arrows depict CLMB dephosphorylation.

**d** (Right) Plots of phosphate released from 100 nM pCLMB$_{WT}$ (blue circles) or pCLMB$_{IxITMut}$ (purple squares) phosphopeptide dephosphorylated by 20 nM (left), 100 nM (middle) or 200 nM (right) CNA/B and 1 μM calmodulin. Displayed is the mean ± SEM of phosphate released for ≥ 5 replicates at each timepoint. (Left) Plots fit by simulation to the model shown in S5A and S5B using Kintek Explorer.

A

LDVPDIIIT  P

LDVPDIIIT  P

2  →

LDVPDIIIT  P

3  →

LDVP  P    LDVPDIIIT  P

4  →

LDVPDIIIT    PXIxIT  LxVP  P    P    P

Ca²⁺

B

LDVPDIIIT  P

LDVPDIIIT  P

2  →

LDVPDIIIT  P

3  →

LxVP  P    LDVPDIIIT  P

4  →

LDVP  P    LDVPDIIIT  P

5  →

LDVPDIIIT    LDVPDIIIT    P    P

Ca²⁺

Ca²⁺
Channel    S-Acylation    N-Myristoylation    Calcimembrin    Phosphate    Calcineurin    Calcium/
Calmodulin    Calcineurin
Substrate

P

**Figure 6: A model for how calcimembrin modulates calcineurin signaling.**

**a** (1) CLMB is dynamically acylated and localizes to membranes where it (2) recruits CN. When CLMB levels are low, calcineurin remains tightly bound to phosphorylated CLMB regardless of $Ca^{2+}$ signals. (3) As CLMB levels rise, $Ca^{2+}$ and CaM binding allow calcineurin-CLMB multimers to form and CLMB to be dephosphorylated. (4) Calcineurin is then freed to engage and dephosphorylate nearby substrates.

**b** (1) CLMB is dynamically acylated and localizes to membranes where it (2) recruits calcineurin. When CLMB levels are low, calcineurin remains tightly bound to phosphorylated CLMB. (3) $Ca^{2+}$ and CaM activate calcineurin, allowing CLMB to present it to nearby substrates for LxVP-mediated binding and dephosphorylation. (4) When CLMB accumulates at membranes, calcineurin-CLMB multimers form, and (5) dephosphorylation triggers calcineurin release to terminate signaling.