A novel motif in calcimembrin/C16orf74 dictates multimeric dephosphorylation by calcineurin

Devin A. Bradburn[1,7], Joana C. Reis[2,3,7], R.Yvette Moreno[1], Shariq Qayyum[2,4,5], Thibault Viennet[2,3,6], Haribabu Arthanari[2,3], Martha S. Cyert[1]*

1. Department of Biology, Stanford University, Stanford, CA 94305, USA

2. Department of Cancer Biology, Dana-Farber Cancer Institute, Boston, Massachusetts, USA.

3. Department of Biological Chemistry and Molecular Pharmacology, Harvard Medical School, Boston, Massachusetts, USA.

4. Men's Health: Aging & Metabolism, Brigham and Women's Hospital, Harvard Medical School, Boston, MA, USA.

5. Cardiovascular Research Center, Department of Medicine, Massachusetts General Hospital and Harvard Medical School, Boston, MA, USA

6. Department of Chemistry and iNANO, Aarhus University, Aarhus, Denmark

7. Contributed equally to this work

*Corresponding author:
Department of Biology
410B Bass Biology Building
Stanford University
Stanford, CA 94305-5020
phone: 650 723-9970
fax: 650 724-9945
E-mail: mcyert@stanford.edu

Abstract:

Calcineurin, the Ca^{2+}/calmodulin-activated protein phosphatase, recognizes substrates and regulators via

short linear motifs, PxIxIT and LxVP, which dock to distinct sites on calcineurin to determine enzyme

distribution and catalysis, respectively. Calcimembrin/C16orf74 (CLMB), an intrinsically disordered

microprotein whose expression correlates with poor cancer outcomes, targets calcineurin to membranes

where it may promote oncogenesis by shaping calcineurin signaling. We show that CLMB associates with

membranes via lipidation, i.e. N-myristoylation and reversible S-acylation. Furthermore, CLMB contains

an unusual composite 'LxVPxIxIT' motif, that binds the PxIxIT-docking site on calcineurin with

extraordinarily high affinity when phosphorylated, [33]LDVPDIIITPP(p)T[44]. Calcineurin dephosphorylates

CLMB to decrease this affinity, but Thr44 is protected from dephosphorylation when PxIxIT-bound. We

propose that CLMB is dephosphorylated in multimeric complexes, where one PxIxIT-bound CLMB recruits

calcineurin to membranes, allowing a second CLMB to engage via its LxVP motif to be dephosphorylated.

In vivo and *in vitro* data, including nuclear magnetic resonance (NMR) analyses of CLMB-calcineurin

complexes, support this model. Thus, CLMB with its composite motif imposes unique properties to

calcineurin signaling at membranes including sensitivity to CLMB:calcineurin ratios, CLMB

phosphorylation and dynamic S-acylation.

Introduction

Calcium (Ca^{2+}) ions are spatially and temporally controlled in cells to regulate a wide array of biological

processes [1]. Ca^{2+} signaling frequently occurs within membrane-associated microdomains that contain both

effector proteins and their substrates. Thus, subcellular localization of calcineurin, the only Ca^{2+}-calmodulin

(CaM) regulated serine/threonine protein phosphatase, dictates its signaling specificity[2]. Calcineurin

(CNA/B) is an obligate heterodimer composed of catalytic (CNA) and regulatory (CNB) subunits that is

activated when Ca^{2+}/calmodulin binding relieves autoinhibition mediated by the CNA C-terminal domain[3].

Calcineurin interacts with partners via two short linear motifs (SLiMs), LxVP and PxIxIT, which cooperate

to promote dephosphorylation. PxIxIT binds to CNA independent of Ca^{2+} and anchors calcineurin to

substrates and regulators. In contrast, LxVP binds to a site composed of residues in both CNA and CNB

and orients substrates for dephosphorylation at a separate catalytic site, marked by Zn^{2+} and Fe^{2+} ions[2, 3].

LxVP binding occurs only after activation by Ca^{2+}/CaM and is blocked by autoinhibitory sequences in CNA

as well as the immunosuppressant drugs, Cyclosporin A and FK506[4, 5].

Systematic discovery of PxIxIT and LxVP SLiMs has significantly expanded our knowledge of calcineurin

signaling[6, 7, 8, 9] and surprisingly identified at least 31 human proteins that harbor composite calcineurin-

binding motifs containing contiguous LxVP and PxIxIT sequences that share a proline (LxVPxIxIT)[9].

Although each SLiM in this composite motif can bind to its cognate site, they are unlikely to do so

simultaneously, as their docking surfaces are ~60 Å apart on calcineurin[4, 9]. Thus, how or if these motifs

promote dephosphorylation is unclear.

Here we elucidate how C16orf74, a short unstructured microprotein associated with poor outcomes in

melanoma, pancreatic and head and neck cancer, is regulated by calcineurin via a composite LxVPxIxIT

motif [9, 10]. We name C16orf74 calcimembrin (CLMB), because it associates with membranes via lipidation

(N-myristoylation and S-acylation) and recruits calcineurin to these sites. CLMB is phosphorylated at Thr

44 (pThr44) and binds calcineurin with exceptionally high affinity because the flanking LxVP and pThr44

features (LxVPxIxITxx(p)T) enhance PxIxT-mediated binding to CNA. Calcineurin also dephosphorylates

pThr44, but this residue is protected when CLMB is bound to the PxIxT-docking surface. Thus, we propose

that dephosphorylation occurs in multimers, with one CNA/B bound to two CLMB molecules at the PxIxIT

and LxVP-docking sites, respectively. NMR confirms formation of these complexes *in vitro*. Furthermore,

this multimeric model predicts that CLMB dephosphorylation varies depending on the amount of CLMB

relative to calcineurin, which we demonstrate via in cell analyses and kinetic modeling of

dephosphorylation *in vitro*. In summary, this work reveals how CLMB can dynamically shape calcineurin

signaling through a mechanism dictated by its composite motif and reversible association with membranes.

Results

Calcimembrin is myristoylated and S-acylated. CLMB lacks predicted transmembrane domains but may associate with membranes via lipidation (Fig. 1a)[10]. CLMB contains a glycine at position 2 (Gly2), a site frequently modified by myristate, a 14-carbon lipid whose co-translational addition can promote protein association with membranes[11]. CLMB also has two cysteines near in its N-terminus, Cys7 and Cys14, one of which, Cys7, is predicted to be modified by S-acylation, the reversible addition of a fatty acid, usually a 16-carbon palmitate, that regulates membrane localization[12]. To determine if CLMB is lipidated, we metabolically labeled HEK293 Flp-In T-REx cells inducibly expressing wildtype CLMB-FLAG (CLMB$_{WT}$) with YnMyr and 17-ODYA, alkyne-containing analogs of myristate and palmitate respectively[13]. Following immunoprecipitation, analog incorporation was detected via addition of biotin using CLICK-chemistry followed by visualization with streptavidin (Fig. 1b). CLMB$_{WT}$ incorporated YnMyr and 17-ODYA, indicating that the protein is both myristoylated and palmitoylated. To identify which residues were lipidated, we mutated proposed sites of S-acylation (CLMB$_{C7,14S}$), and myristoylation (CLMB$_{G2A}$ (Fig. 1b). The mutants were expressed at low levels suggesting instability, which we countered by including a proteasome inhibitor (MG132) in the analyses. CLMB$_{C7,14S}$ showed no 17-ODYA incorporation but maintained YnMyr suggesting that Cys7 and/or Cys14 are S-acylated. By contrast, neither lipid modification was detected for CLMB$_{G2A}$, which confirms that Gly2 is myristoylated. Because myristoylation can promote S-acylation by anchoring a protein to the membrane[14], we then examined CLMB$_{G2A}$ for S-acylation with increased sensitivity by including palmostatin B (PalmB), a pan-inhibitor of thioesterases that remove S-acylation[15]. In the presence of this inhibitor, 17-ODYA incorporation into both CLMB$_{WT}$ and CLMB$_{G2A}$ was observed (Supplementary Fig.1), indicating that CLMB$_{G2A}$ is S-acylated, but with reduced efficiency.

To determine if CLMB-FLAG is dually S-acylated at Cys7 and Cys14, we used acyl-PEG exchange (APE) in which S-acylated groups are cleaved using hydroxylamine (NH$_2$OH) and exchanged for a PEGylated molecule that increases protein mass by 5 kDa for each S-acylated residue[16] (Fig.1c). Endogenous calnexin,

a dually acylated protein, served as a positive control for each sample. For CLMB$_{WT}$, two slower migrating

bands were observed. CLMB$_{C7S}$ lacked the upper band, while CLMB$_{C7,14S}$ was unmodified, indicating that

Cys7 and Cys14 are S-acylated in CLMB$_{WT}$ (Fig. 1c). CLMB$_{G2A}$ displayed two sites of S-acylation, but

with reduced intensity relative to CLMB$_{WT}$, consistent with results from metabolic labeling (Fig. 1b).

Together these findings establish that CLMB is myristoylated at Gly2 and S-acylated at Cys7 and Cys14.

Dynamic S-acylation of CLMB mediates membrane association. To examine whether S-acylation of

CLMB is dynamic[17], we carried out a pair of pulse-chase analyses with two metabolic labels (Fig. 1d, e).

L-AHA, an azide-containing methionine analog detected CLMB-FLAG protein levels, while in parallel,

17-ODYA detected S-acylation. CLMB-FLAG was immunopurified (IP) after labeling for one hour

followed by a 30-minute chase with DMSO or PalmB. Quantifying the 17-ODYA to L-AHA ratio revealed

that palmitate incorporation into CLMB increased ~3-fold when thioesterases were inhibited (Fig, 1e,

PalmB vs DMSO). This indicates that S-acylation of CLMB turns over rapidly relative to its rate of

degradation.

To determine if lipidation mediates CLMB membrane recruitment, we analyzed the localization of

CLMB$_{WT}$ and lipidation defective mutants, CLMB$_{G2A}$ and CLMB$_{C7,14S}$. Detergent-assisted subcellular

fractionation showed that CLMB$_{WT}$ was primarily membrane associated, appearing in fractions marked by

GM130, a Golgi protein. By contrast, both mutants were predominantly cytosolic, appearing in fractions

marked by α-tubulin (Fig. 1 f, g). Together these analyses indicate that N-myristoylation, in combination

with dynamic dual S-acylation, promote the stable association of CLMB with membranes.

The composite SLiM in CLMB mediates binding to and dephosphorylation by calcineurin. Next, we

examined how each part of CLMB's composite motif, LDVPDIIIT, contributes to calcineurin binding by

generating mutants, CLMB$_{IxITMut}$ (LDVPDAIAA) and CLMB$_{LxVMut}$, (**ADAPDIIIT**) (Fig. 1a) that disrupt

each motif independently without altering the shared proline. To examine binding of these mutants to

calcineurin, the association of CLMB-FLAG with GFP-tagged calcineurin (CNA-GFP complexed with

endogenous CNB) was assessed following purification of calcineurin from lysates of HEK293 cells co-

transfected with these constructs (Fig. 2a, b). $CLMB_{WT}$ but not $CLMB_{IxITMut}$ co-purified with calcineurin,

showing that this interaction was PxIxIT-dependent. Less calcineurin co-purified with $CLMB_{LxVMut}$ vs.

$CLMB_{WT}$, showing that the LxVP also makes a modest contribution to calcineurin binding. Furthermore,

when expressed and analyzed by SDS-polyacrylamide gel electrophoresis (SDS-PAGE), both mutants

displayed two electrophoretic forms, with a slower migrating band also faintly observed for $CLMB_{WT}$ (Fig.

2a, input). Phosphorylation, which often reduces electrophoretic mobility, has been reported for CLMB at

Thr44 (pThr44) and less frequently at Thr46[18]. We confirmed that the slower migrating forms of CLMB

contained pThr44, as they were absent in mutants lacking this residue ($CLMB_{T44A}$, $CLMB_{T44A+LxVMut}$,

$CLMB_{T44A,T46A}$), but not in $CLMB_{T46A}$ (Fig. 2a, Supplemental Fig. 2a). These observations indicate that

calcineurin binds to CLMB and dephosphorylates pThr44, which is disrupted when either the LxVP or

PxIxIT portion of the composite motif is compromised.

Finally, we confirmed that pThr44 enhance binding to calcineurin, as the slower migrating (phosphorylated)

forms of $CLMB_{WT}$ and $CLMB_{LxVMut}$ predominantly co-purified with calcineurin (Fig. 2a, IP)[10]. Indeed,

$CLMB_{T44A}$ displayed reduced binding to calcineurin, which was further reduced by mutating LxV

($CLMB_{T44A+LxVMut}$) (Fig. 2a, b).

The contributions of PxIxIT, LxVP, and pThr44 to calcineurin binding were also examined by measuring

the extent of CLMB co-localization with calcineurin (CNA-GFP complexed with endogenous CNB) via

immunofluorescence (IF) (Fig. 2c, d). When co-transfected into MCF7 cells $CLMB_{WT}$-FLAG and

calcineurin colocalize at membranes (see regions marked by arrows in Fig. 2c). In contrast, when co-

expressed with $CLMB_{IxITMut}$, this colocalization was greatly reduced and calcineurin was predominantly

cytosolic. $CLMB_{LxVMut}$ and $CLMB_{T44A}$ also displayed reduced colocalization with calcineurin (Fig. 2c, d),

mirroring the trends observed in co-purification analyses (Fig. 2a, b). Thus, CLMB recruits calcineurin to

membranes primarily via PxIxT-mediated binding.

Finally, dephosphorylation of pThr44 in CLMB by calcineurin was demonstrated *in vitro* using

phosphorylated peptide substrates that encode residues 21-50 (Fig. 1a). The peptide from pCLMB$_{WT}$ was

robustly dephosphorylated compared to pCLMB$_{LxVMut}$ (Fig. 2e) showing that, as for other calcineurin

substrates, the LxVP motif promotes catalysis[19]. Finally, the presence of a functional LxVP motif in the

CLMB$_{WT}$ peptide was also confirmed by its ability to inhibit dephosphorylation of the RII peptide, an

LxVP-containing calcineurin substrate derived from the regulatory subunit of protein kinase A (RII

isoform)[19] (Supplemental Fig. 2b). Together, these data show that both the LxVP and PxIxIT components

of the composite motif are functional, that CLMB is a calcineurin substrate, and that pThr44 increases

CLMB binding to calcineurin.

Phosphorylated Thr44 and the LxVP motif enhance CLMB binding to the PxIxIT docking site. To

quantitatively assess CLMB-calcineurin interaction, we used isothermal titration calorimetry (ITC) and

fluorescence polarization anisotropy (FP) to measure binding of CLMB peptides containing the composite

SLiM (residues 21-50) to purified, catalytically inactive CNA/B containing CNA$_{Trunc+H151Q}$. This mutant

CNA lacks the C-terminal autoinhibitory region, which allows LxVP motifs to bind to CNA/B in the

absence of Ca^{2+}/CaM[20]. ITC revealed that the phosphorylated CLMB$_{WT}$ peptide (pCLMB$_{WT}$) binds

calcineurin at a one-to-one ratio with high affinity (K_D= 37 nM), (Fig. 3a). FP using FITC-labelled peptides

at pH 7.4 revealed that phosphorylation of Thr44 enhances CLMB binding to calcineurin ~20-fold (K_D =

14.1 nM vs 291 nM) (Fig. 3b). Furthermore, the affinity of pCLMB$_{WT}$ for calcineurin was similar over a

pH range (6.9-8.0) which should change the protonation state of histidine residues (Supplemental Fig. 3a,

b). This included the histidine immediately preceding the LxVP (H32) which is expected to engage with

calcineurin during LxVP docking[7]. The concordance between measurements made with CLMB peptide that

is tag-free (ITC) vs. tagged with fluorescein isothiocyanate (FITC) (FP) (37 vs 14.1 nM for pCLMB$_{WT}$)

(Fig. 3a, b), shows that the tag does not significantly affect binding affinity. Given the high-throughput

capability of FP, this method was used for the remaining comparisons. To test if the high affinity pCLMB$_{WT}$-

calcineurin binding observed was PxIxIT-mediated, we used FP to assess the ability of CLMB peptides to

compete with a well characterized PxIxIT peptide, FITC-PVIVIT [21, 22], for calcineurin binding (Fig. 3c, e).

pCLMB$_{WT}$ competed with PVIVIT most effectively, with peptides lacking the LxVP motif (pCLMB$_{LxVMut}$),

or pThr44 (CLMB$_{WT}$) showing reduced competition (Fig. 3c, e). pCLMB$_{IxITMut}$ lacking the PxIxIT motif

was unable to displace PVIVIT from CNA/B. When these experiments were repeated using FITC-

pCLMB$_{WT}$ peptide in place of FITC-PVIVIT (Fig. 3d, e), a similar pattern was observed, although

phosphorylation of T44 (pCLMB$_{WT}$ vs CLMB$_{WT}$) caused a much greater reduction in Ki (30X vs 3X for

PVIVIT competition). Importantly, CLMB peptides accurately mimic full length CLMB binding to

calcineurin as both caused equivalent displacement of FITC-pCLMB$_{WT}$ (Supplemental Fig. 3c,d).

Altogether, these data show that, like PVIVIT, pCLMB$_{WT}$ peptides bind to the PxIxIT docking groove on

CNA and that flanking residues, i.e. LxV and pThr44, enhance binding to form an extended

LxVPxIxITxxT(p) sequence with exceptionally high affinity for calcineurin.

NMR analysis shows CLMB binding to two sites on calcineurin. Next, to investigate the structure of

CLMB in the presence and absence of calcineurin, we used solution-state NMR. 88% of non-proline

backbone resonances were assigned using a standard set of triple-resonance experiments, utilizing full-

length dually ^{13}C- and ^{15}N-labeled CLMB (76 residues) that was expressed in *E.coli* (Fig. 4a, Supplemental

Fig. 4a). To assess CLMB secondary structure, we employed TALOS-N, a computational method that

predicts backbone torsion angles from NMR chemical shifts[23]. This revealed that CLMB is disordered

overall, but contains two regions with a tendency to form a β-strand (Supplemental Fig. 4b): Residues I38-

I40 include the PxIxIT, consistent with binding of this motif-type to CNA as a β-strand [22]. N-terminal

residues, Q12-V15, were also predisposed to localized folding suggesting possible involvement in other

binding interactions. Similar analyses carried out for CLMB using chemical shift secondary structure

population interference (CheSPI)[24] confirmed the absence of secondary structure (Supplemental Fig. 4c).

To map the interaction interface of calcineurin on CLMB, we focused first on PxIxIT-mediated interactions

by titrating ^{15}N-labeled CLMB with varying molar ratios of monomeric recombinant CNA in the absence

of CNB and monitoring the interaction through ^{15}N-^{1}H SOFAST-HMQC experiments (Fig.4). No additional

insights were gained from BEST-TROSY experiments[25]. CNA can bind to PxIxIT motifs but lacks the LxVP

binding surface found at the interface of CNA and CNB[4]. Due to the size of the CLMB-CNA complex, we

anticipated significant broadening of NMR signals at the interaction interface. This broadening can also

arise from exchange kinetics between the observable CLMB and the titrant, CNA, making it challenging to

distinguish the two effects. As expected, at a molar ratio of 2:1 (CLMB: CNA), many of the CLMB signals

undergo significant line broadening (Fig.4b), which become more pronounced at a molar ratio to 1:1, due

to the majority of CLMB being bound to CNA (Fig.4c). This is particularly true for signals corresponding

to the PxIxIT and LxVP motifs, indicating, as observed in FP experiments, that LxV residues participate in

binding to the PxIxIT interface (Fig. 4b, c, Supplemental Fig. 2d). Line broadening across the majority of

CLMB residues indicates that the interaction interface on CLMB extends beyond the PxIxIT and LxVP

motifs (Fig.4 b, c). Interactions with the N- and C-terminal residues of CLMB may be an avidity-driven

effect, anchored by PxIxIT binding, as has been reported for other calcineurin substrates [26].

To investigate binding that may occur at the LxVP docking site of calcineurin, we next titrated CLMB with

the heterodimeric CNA/B complex (Fig.4d, e). Signal broadening at the N-terminus of CLMB was more

pronounced with CNA/B relative to CNA alone, suggesting possible interactions between CLMB and CNB

(Fig. 4b, c vs 4d, e, and Supplemental Fig. 4d). These include the Q9-V12 region that displayed partial

folding in the absence of calcineurin. At a molar ratio of 2:1 (CLMB: CNA/B), more broadening is observed

in LxVP but not PxIxIT residues when compared to the interaction with CNA (Fig. 4d vs 4b, and

Supplemental Fig. 4c). This includes not only L33 and V35, but also H32, another residue expected to

participate in LxVP binding to CNA/B[7]. However, at a molar ratio of 1:1, where essentially all CLMB

should be PxIxIT-bound (given the higher K_D for PxIxIT vs LxVP and our working concentrations), the

pattern of line broadening at the composite motif is similar when complexed with either CNA or CNA/B

(Fig. 4c vs 4e). To better visualize LxVP-mediated binding of CLMB to CNA/B, NMR analyses were also

performed in the presence of pCLMB$_{LxVPmut}$ peptide (which was not isotopically labeled) to saturate the

PxIxIT binding site on CNA/B (Supplemental Fig. 5). While addition of this peptide on its own caused no

change in the NMR spectrum for ^{15}N-labeled CLMB (Supplemental Fig. 5a), in the presence of CNA/B

(2:1 or 1:1 CLMB: CNA) residues in the LxVP motif, particularly H32 and L33, displayed shifting and

broadening effects in addition to N-terminal residues, as previously observed (Supplemental Fig. 5b,c).

Together these findings show that when in excess (2 CLMB: 1 CNA/B), CLMB engages calcineurin

(CNA/B) through both its LxVP and PxIxIT motifs.

To investigate the CLMB-calcineurin interaction further, we carried out structural predictions using

AlphaFold 3 for 2 CLMB molecules and CNA/B containing truncated CNA lacking the C-terminal

calmodulin binding and regulatory domains (Supplemental Fig. 6a-c) [27]. Consistent with our NMR analyses,

the models show two copies of CLMB simultaneously bound to CNA/B, one each at the LxVP and PxIxIT

docking sites. Predicted interactions of pCLMB containing pThr44 with CNA/B was nearly identical,

however with the LxVP-bound CLMB molecule showing an additional interaction of pThr44 with the

active site, as would occur during catalysis (Supplemental Fig. 6a-c). Thus, these models suggest that

formation of multimeric complexes (2 CLMB: 1 CNA/B) will facilitate dephosphorylation of pCLMB.

Calcineurin dephosphorylates CLMB by forming multimers. Next, we sought to test the prediction that

pThr44 dephosphorylation is promoted when CLMB is in excess (2 CLMB: 1 CNA/B) but inhibited at

lower CLMB: CNA/B ratios where PxIxIT-bound complexes are favored and protect pThr44 from

dephosphorylation. To this end, we inducibly expressed CLMB$_{WT}$-FLAG in HEK293 Flp-In T-REx cells

and compared its phosphorylation status with endogenous vs overexpressed calcineurin (i.e. transfection of

GFP-CNA, which complexes with endogenous CNB). In stark contrast to a typical substrate, CLMB-FLAG

phosphorylation at T44 increased upon calcineurin overexpression (Fig. 5a,b, Supplemental Fig.7a). Intact

PxIxIT binding rather than catalytic activity was required for this effect, as overexpressing catalytically

inactive calcineurin (containing GFP-CNA$_{H151Q}$) [28], but not calcineurin that is defective for PxIxIT binding

(containing GFP-CNA$_{NIR}$) [29], increased CLMB-FLAG phosphorylation (Supplemental Fig. 7b). We

predicted that further increasing CLMB levels in these calcineurin overexpressing cells would promote

dephosphorylation of CLMB-FLAG by increasing the CLMB:CNA/B ratio. Thus, we overexpressed V5-

tagged CLMB in HEK293 Flp-In T-REx cells that inducibly expressed CLMB-FLAG and overexpressed

calcineurin (GFP-CNA complexed with endogenous CNB) (Fig. 5a-c). As predicted, phosphorylation of

$CLMB_{WT}$-FLAG decreased upon expression of $CLMB_{WT}$-V5. This effect was dependent on PxIxIT binding,

as introducing $CLMB_{IxITMut}$-V5 did not decrease $CLMB_{WT}$-FLAG phosphorylation, but $CLMB_{LxVMut}$-V5,

which contains an intact PxIxIT motif, did (Fig, 5a-c). for $CLMB_{WT}$, when $CLMB_{LxVMut}$ was expressed,

its phosphorylation was enhanced by overexpression of calcineurin and rescued by co-expression of

$CLMB_{WT}$-V5 or $CLMB_{LxVMut}$-V5 but not $CLMB_{IxITMut}$-V5. However, because dephosphorylation is

impaired by mutation of the LxV (Fig. 2e), $CLMB_{LxV}$-FLAG displayed increased levels of phosphorylation

relative to $CLMB_{WT}$ under all conditions (Fig.5a, b). Finally, we repeated these analyses using $CLMB_{IxITMut}$-

FLAG, which displayed partial phosphorylation regardless of calcineurin levels. Surprisingly, $CLMB_{IxITMut}$-

FLAG was predominantly dephosphorylated when co-expressed with $CLMB_{WT}$-V5 or $CLMB_{LxVMut}$-V5 but

not $CLMB_{IxITMut}$-V5 (Fig. 5a, b). Thus, although the LxVP and PxIxIT components of the CLMB composite

motif are both required for efficient dephosphorylation, they do not need to be on the same molecule. A

single copy of CLMB must have an LxVP to mediate its own dephosphorylation but can utilize the PxIxIT

of neighboring copies to recruit calcineurin, promote LxVP binding, and be dephosphorylated (Fig. 5c,

bottom row).

To demonstrate more precisely how the ratio of CLMB to calcineurin affects dephosphorylation, we

conducted a series of *in vitro* dephosphorylation assays using a fixed substrate concentration (100nM

$pCLMB_{WT}$ or $pCLMB_{IxITMut}$ peptides) with increasing concentrations of purified calcineurin (CNA/B) (20,

100 or 200nM) (Fig. 5d, top row). Under these *in vitro* conditions, $pCLMB_{IxITMut}$ was efficiently

dephosphorylated due to high enzyme and substrate concentrations and the absence of any competing

PxIxIT-containing proteins. For $pCLMB_{IxITMut}$, we observed that dephosphorylation rates increased

significantly with the calcineurin concentration (Fig. 5d, top row). In contrast, dephosphorylation of

$pCLMB_{WT}$ increased modestly as calcineurin increased from 20 vs 100 nM, and then decreased slightly at

200nM calcineurin, where enzyme concentration exceeded that of substrate (100 nM) (Fig. 5d, top row).

To better understand this unusual reaction behavior, we turned to kinetic modeling. First, we developed a

model describing the complex series of reactions we expect to occur, based on our NMR and in cell analyses

of CLMB-calcineurin interactions (Supplemental Fig. 8). For $pCLMB_{IxITMut}$, which contains only an LxVP

motif (Supplemental Fig. 8b), substrate binds to the LxVP-docking site of calcineurin (k_2, k_{-2}) and is

dephosphorylated (k_3) leading to product formation. Thus, dephosphorylation accelerates with increasing

enzyme concentration (Fig. 5d, top row). However, for dephosphorylation of $pCLMB_{WT}$ (Supplemental Fig

8a), we propose competition between two binding modes: high affinity binding to the PxIxIT-docking site

(k_1, k_{-1}), which prevents pT44 phosphorylation, or lower affinity LxVP-mediated binding (k_2, k_{-2}), which

leads to dephosphorylation (k_3). Furthermore, the model allows one enzyme molecule to bind substrate at

each of these sites simultaneously (ES1S2). For this complex reaction mechanism, increasing enzyme

concentration drives formation of more LxVP-bound substrate and dephosphorylation, but this is offset by

reduced substrate availability due to increased PxIxIT-mediated substrate binding and pThr44 protection.

Thus, dephosphorylation rates decrease as enzyme concentration exceeds that of substrate (Fig. 5d, top

row).

To further probe this proposed mechanism, we then used kinetic modeling software (KintekExploer[TM])

which fits experimental data directly through dynamic simulation and does not require simplifying

assumptions[30]. After providing initial values for rate constants consistent with published reports[6, 31], our

experimental data were fit to the proposed kinetic model (Fig. 5d bottom row and Supplemental Fig. 8).

The resulting rate parameters were well within expected values, and the predicted K_D for phosphorylated

PxIxIT binding to calcineurin was 7nM (Supplemental Fig. 8c), which agrees well with experimentally

determined values (Fig. 3a, c). However, we noted that some parameters, particularly on and off rates for

substrate binding were poorly constrained by the model. Therefore, we manually manipulated different

parameters to better understand how relationships between them influenced the fit to our data

(Supplemental Fig. 9). This revealed that the equilibrium binding constant for phospho-PxIxIT binding was

critical rather than specific on and off rates (k_1 and k_{-1}) (Supplemental Fig. 9a and b). Furthermore, the k_1

and k_{-1} values predicted by kinetic simulations represented lower limits for these rates as decreasing them

caused significant deviations from the data despite keeping the K_D constant (Supplemental Fig. 9b).

Similarly, k_{off} for LxVP binding (k_{-2}) and k_{cat} (k_3) could be increased proportionally without significantly

affecting the overall fit (Supplemental Fig. 9c), suggesting that the catalytic efficiency of LxVP-mediated

dephosphorylation is a key driver of the observed reaction kinetics. Confirming this, we found that

maintaining the second order rate constant was critical for good agreement between the data and the model

(Supplemental Fig. 9d). Thus overall, kinetic modeling of our *in vitro* dephosphorylation data supports the

hypothesis that phosphorylated CLMB is dephosphorylated via multimer formation. This novel mechanism,

which requires substrate levels to exceed that of calcineurin for efficient dephosphorylation, is dictated by

the composite LxVPxIxIT motif.

Discussion

Here we describe multivalent interactions between calcineurin and CLMB, an intrinsically disordered

microprotein and calcineurin substrate that dynamically associates with membranes via protein lipidation.

We elucidate roles for a novel composite motif in CLMB, LxVPxIxIT, that combines two SLiMs, PxIxIT

and LxVP, with distinct binding modes and functions in calcineurin signaling.

Calcineurin, the conserved and ubiquitously expressed calcium/calmodulin regulated protein phosphatase,

has been categorized as a dynamic or date hub and specifically as a linear motif binding hub (LMB-hub)

because it engages a diverse set of partners that bind competitively to the structurally plastic PxIxIT docking

groove[32]. PxIxIT peptides from natural substrates bind to calcineurin as a β strand, vary greatly in sequence

and typically have low affinities (~2.5 to 250 micomolar)[26, 33], while regulators and scaffolds, including

PVIVIT, the selected inhibitor peptide, have K_D's of ~0.5 micromolar [21, 34, 35]. By contrast, we show that

the [32]HLDVPDIIITPP(p)TP[45] sequence in CLMB binds calcineurin at the PxIxIT-docking groove with a

K_D of ~14 nM. This extraordinary affinity is mediated predominantly via binding of the core PxIxIT motif

which is enhanced ~20 fold by phosphorylation at the +3 position (pThr44), and ~5 fold by the N-terminal

LxV sequence (Fig. 3). Such high affinities have also been achieved by engineering the PVIVIT peptide to

include a hydrophobic residue at the -1 position and a phosphothreonine at the +3 position, although

phosphorylation does not universally enhance PxIxIT affinity[35]. A natural PxIxIT peptide derived from an

endogenous inhibitor of calcineurin (CABIN1),[2145]PPEITVTPP(p)TP[2156] also shows high affinity (~170

nM) due to phosphorylation of its C-terminal sequence, PPTP, which is identical to that found in CLMB[34].

This proline-rich sequence is predicted to form a polyproline helix that may function to extend the β-strand

interaction with calcineurin[36, 37]. Thus, the composite LxVPxIxIT motif in CLMB combines several features

that consistently stabilize PxIxT-docking. Because calcineurin is an LMB-hub, such high affinity PxIxIT

peptides typically inhibit signaling by preventing its interaction with substrates[35, 38]. It is surprising,

therefore that CLMB exhibits high affinity PxIxIT-mediated docking yet is also dephosphorylated by

calcineurin. As we demonstrate, this is only possible due to the LxVP portion of the composite SLiM in

CLMB, and dephosphorylation can only occur when excess CLMB levels allow lower affinity LxVP-

docking to occur on a molecule of calcineurin whose PxIxIT-docking site is saturated.

Based on our findings, we present two models for how CLMB may shape calcineurin activity *in vivo* (Figure

6). First, dynamic S-acylation promotes CLMB association with membranes (Fig. 6a, Step 1), where, when

phosphorylated, it recruits a localized pool of calcineurin via high affinity PxIxIT-mediated binding, (Fig.

6a, Step 2). This membrane sequestration may prevent calcineurin from dephosphorylating substrates in

other cellular locations. In the presence of a Ca^{2+} signal, Ca^{2+} and CaM then activate calcineurin, which

exposes the LxVP docking site[39]. If CLMB is in excess relative to calcineurin and PxIxIT binding is fully

saturated, a second CLMB molecule will bind via its LxVP motif which positions pThr44 for

dephosphorylation at the active site (Fig. 6a, Step 3). However, as the population of CLMB shifts to the

dephosphorylated form, PxIxIT binding affinity decreases (Fig. 3) which promotes calcineurin dissociation

from CLMB, allowing the enzyme to engage and dephosphorylate other nearby LxVP- and PxIxIT-

containing substrates (Fig. 6a, Step 4). This tuning of PxIxIT affinity is critical for signaling as demonstrated

for AKAP5, a scaffold that concentrates both calcineurin and NFAT near L-type Ca^{2+} channels in neurons

to promote NFAT dephosphorylation[40]. In that case, increasing the AKAP5 PxIxIT affinity for calcineurin

~two-fold significantly impaired NFAT activation due to increased sequestering of calcineurin[41].

In our second model (Fig. 6b), active calcineurin-pCLMB dimers (1 CLMB: 1 CNA/B) directly engage and

dephosphorylate substrates at the membrane that contain an LxVP, but no PxIxIT motif (Fig. 6b, step 3).

Such substrates are exemplified by PRKAR2A, the type II-alpha regulatory subunit of protein kinase A,

which contains an LxVP but depends on the PxIxIT motif within its binding partner, AKAP5, to recruit

calcineurin and promote dephosphorylation[9, 42]. Proteome-wide predictions suggest that many proteins may

similarly harbor an LxVP without an associated PxIxIT [9]. In our second model (Fig. 6b), active calcineurin-

pCLMB dimers (1 CLMB: 1 CNA/B) directly engage and dephosphorylate such PxIxIT-less substrates at

the membrane (Fig. 6b, step 3). Simply recruiting calcineurin to membranes, where association kinetics are

20-30 fold faster than in the cytosol of eukaryotic cells, may enhance LxVP binding causing their efficient

dephosphorylation in the absence of a PxIxIT motif [43]. Such division of the PxIxIT and LxVP motifs among

two molecules may also allow for the extremely high affinity PxIxIT observed in CLMB. In substrates that

contain both a LxVP and PxIxIT motif, stable PxIxIT binding would be expected to limit enzymatic

efficiency by slowing product release. However, acting only as a scaffold, CLMB can remain tightly

associated with calcineurin via PxIxIT-binding while other substrates cycle in and out of the LxVP-binding

surface to be dephosphorylated. In this scenario (Fig. 6b), calcineurin remains at the membrane either until

CLMB levels increase sufficiently to allow CLMB dephosphorylation and calcineurin dissociation (Fig. 6b

step 4) or until CLMB is de-acylated. In both models, intracellular Ca^{2+} dynamics, CLMB protein levels,

and S-acylation/de-acylation activities combine to modulate calcineurin signaling. Future studies aim to

identify the palmitoyl acyltransferases (PATs) and acyl protein thioesterases (APTs) that dynamically

regulate CLMB S-acylation to better understand how these critical regulators contribute to calcineurin-

CLMB signaling *in vivo*.

Another missing player in our model is the kinase(s) that phosphorylates CLMB at Thr44. One candidate

is the stress-activated p38 mitogen-activated protein kinase (MAPK), which phosphorylates the identical

PxIxIT-flanking sequence in Cabin1 (PP(p)TP)[44]. Because phosphorylation increases CLMB-calcineurin

affinity, kinase activation is required to prime regulation of calcineurin signaling by CLMB. However,

phosphorylation also establishes a negative feedback mechanism, with dephosphorylation decreasing

calcineurin's interaction with its substrate. Thus, regulators that modify kinase activity may also

significantly affect CLMB-calcineurin signaling dynamics.

This study highlights the critical role that SLiMs play in conferring flexibility, tunability and dynamism to

cell signaling at the molecular level. Further studies are needed to establish whether composite LxVPxIxT

motifs always limit substrate dephosphorylation to occurring in a multimeric complex. Of at least 30 human

proteins containing predicted LxVPxIxIT motifs, several are ion channels, such as TRPM6, that assemble

into higher order complexes[45], suggesting that these motifs could be a fidelity mechanism that limits

dephosphorylation to substrates that are properly assembled into complexes. Similarly, multivalency is a

characteristic exhibited by many SLiMs. Some composite SLiMs bind multiple partners in an exclusive

manner, for example requiring a kinase and a phosphatase to compete for access to a shared substrate[46, 47].

For the intermediate chain of dynein, multivalency regulates assembly of the dynein complex and its

interaction with distinct binding partners[48]. In the cell cycle regulator CDKN1A (p21), sequences that flank

the PCNA-binding SLiM also encode an essential nuclear localization sequence[49]. Thus, 'composite' SLiMs

that encode multiple interactions may be more common that currently appreciated. This complexity

emphasizes how critical SLiM-based interactions are for integrating and coordinating the plethora of

possible protein-protein interactions that dictate cellular responses.

A major challenge that lies ahead is to identify the molecular targets and physiological functions that are

regulated by calcineurin-CLMB. While this manuscript was being revised for publication, another study

showed that C16orf74/MICT1 is highly expressed in brown fat in mice and controls thermogenesis by

regulating the dephosphorylation of PKA RIIβ by calcineurin[50]. In some pancreatic ductal adenocarcinoma

(PDAC) derived cell lines, CLMB increases proliferation and migration suggesting that CLMB shapes

calcineurin signaling to promote oncogenesis[10]. Indeed, high CLMB expression is associated with poor

patient outcomes for PDAC, melanoma and head and neck cancer[10, 51]. Mechanisms underlying these

activities have not yet been identified, but these cancers may provide good starting points for examining

how CLMB modifies calcineurin signaling in a clinically relevant context.

Methods

Cell culture and transfection:

Cells were cultured at 37 °C in 5% CO_2. HEK293 Flp-In TReX and MCF7 cell lines were grown in standard

media containing Dulbecco's modified eagle's medium (DMEM) with 4.5 mg/mL glucose, L-glutamine,

and sodium pyruvate (Cytiva SH30243.FS) supplemented with 10% fetal bovine serum (Corning

MT35010CV). Transfections were done according to manufacturer instructions. HEK293 cells were

transfected using lipofectamine 2000 (Thermo Fisher Scientific 11668019). MCF7 cells were transfected

using Transfex ATCC transfection reagent (ATCC ACS-4005).

Stable cell line generation:

On day one, 9.6 x 10^5 HEK293 Flp-In TReX cells were transfected using lipofectamine 2000 with 1.5 μg

pOG44 and 0. 5 μg pcDNA5 FRT TO encoding proteins of interest in a 6 well dish (plasmid # 2801, 2852,

2860, 2861, 2862, 2863, 2864, and 2880). Transfections were done according to manufacturer instructions.

On day two, transfected cells were passaged onto 10cm plates. On day three, selection was started using

media containing 3 μg blasticidin and 200 ng/mL hygromycin. Selection media was replaced every 2-3

days until distinct colonies appeared. Colonies were selected using cloning rings and expanded before being

stored in liquid nitrogen.

Metabolic labeling to detect CLMB lipidation:

HEK293 Flp-In TReX lines stably expressing CLMB$_{WT}$, CLMB$_{C7,14S}$, or CLMB$_{G2A}$ were induced with 10

ng/mL doxycycline for 24 hours and split evenly among three 10 cm tissue culture dishes. The following

day, cells were rinsed with PBS and incubated in DMEM containing 4.5 mg/mL glucose, L-glutamine,

sodium pyruvate, 2% FBS, and either 20 µM YnMyr (Click Chemistry Tools 1164-5), 30 µM 17-ODYA

(Cayman Chemical Company 34450-18-5), or vehicle (ethanol and DMSO) overnight. Cells were then co-

incubated with 20 µM YnMyr, 30 µM 17-ODYA, or vehicle and 20 µM MG132 (EMD Millipore 474787-

10MG) for 4 hours before washed twice in PBS and frozen at -80 °C. Cells were lysed in 400 µL TEA lysis

buffer (50 mM TEA pH 7.4, 150 mM NaCl, 1% triton X-100, and 1 mM PMSF) by inverting for 15 minutes

at 4 °C. Lysates were passed five times through a 27-gauge needle, centrifuged at 16,000 x g for 15 minutes

at 4 °C and protein content determined by Pierce BCA protein assay kit (Thermo Fisher Scientific 23225).

25 µg total lysate was removed as input and 0.5 mg total protein was incubated with pre-washed magnetic

M2 anti-FLAG (Sigma-Aldrich M8823-5ML) beads at 4 °C for 1.5 hours while inverting. Beads were

washed 3X with modified RIPA (50 mM TEA pH 7.4, 150 mM NaCl, 1% Triton X-100, 0.1% SDS, and

1% sodium deoxycholate) and resuspended in 40 µL 1x PBS. 10 µL of 5x click reaction mix (1x PBS, 0.5

mM biotin azide, 5 mM TCEP, 0.5 mM TBTA, 5 mM $CuSO_4$) was added and reactions incubated for 1 hour

at 24 °C, shaking at 550 RPM. Beads were washed 3X in modified RIPA and eluted 2X using 30 µL of 1

mg/mL FLAG peptide in modified RIPA buffer. 12 µL of non-reducing 6x SDS sample buffer was added

to eluate before heating at 95 °C for 2 minutes and analysis by SDS-PAGE. Note that addition of reducing

agents to SDS sample buffer results in loss of 17-ODYA signal. Samples were transferred to nitrocellulose

membrane and lipidation was detected using fluorophore-tagged streptavidin, and total CLMB with anti-

FLAG. Immunoblots were imaged using a Licor Odyssey CLx and quantified with Image Studio software.

Metabolic pulse-chase labeling assay to assess dynamic CLMB S-acylation:

HEK293 Flp-In TReX cells stably expressing CLMB$_{WT}$-FLAG were induced with 10 ng/mL doxycycline

for 48 hours. Cells were rinsed 2X in PBS and incubated in methionine free DMEM supplemented with 5%

charcoal-stripped FBS, 1 mM L-glutamate, 1 mM sodium pyruvate, 0.4 mM cysteine and 10 ng/mL

doxycycline along with vehicle (ethanol), 30 µM 17-ODYA, 50 µM L-AHA, or 17-ODYA and L-AHA

together. After two hours, cells were washed 2X in PBS and either harvested immediately or incubated in

standard media supplemented with 10 µM palmitic acid and 10 ng/mL doxycycline, with or without 10 µM

palmostatin B (EMD Millipore 178501-5MG). Cells were rinsed in PBS and frozen at -80 °C. Metabolic

labeling was done as described in the previous section with the exception that two rounds of labeling were

done. First, Cy5-azide (Click Chemistry Tools AZ118-1) was linked to 17-ODYA. Then, cells were washed

and L-AHA was linked to AF-488 (Click Chemistry Tools 1277-1) alkyne before being washed three

additional times and eluted using FLAG peptide. Samples were resolved via SDS-PAGE and in-gel

fluorescence was read using the GE Amersham Typhoon imaging system. Images were quantified in Image

Studio. The proportion of S-acylated CLMB is calculated as Cy5/A-488 normalized to the no chase control.

Acyl-PEG exchange:

HEK293 Flp-In TReX lines stably expressing $CLMB_{WT}$, $CLMB_{C7S}$, $CLMB_{C7,14S}$, or $CLMB_{G2A}$ were induced

with 10 ng/mL doxycycline for 48 hours. Cells were rinsed 2X in PBS and stored at -80 °C. Cells were

lysed in 400 µL lysis buffer (50 mM TEA pH 7.3, 150 mM NaCl, 2.5% SDS, 1 mM PMSF, 100 units

benzonase) at 37 °C for 20 minutes, shaking at 800 RPM. Lysates were passed 10X through a 27-gauge

needle, centrifuged at 16,000 x g for 5 minutes and protein quantified by BCA. Lysates were adjusted to 2

mg/mL and incubated with 10 mM TCEP for 30 minutes at room temperature. Cysteines were protected by

incubating with 25 mM NEM for 2 hours at 40 °C, shaking at 800RPM. Protein was precipitated over night

with four volumes of acetone at -20 °C, centrifuged at 16,000 x g for 20 minutes at 4 °C and washed 4X in

70% acetone. Acetone was removed, pellets air-dried to remove all traces of acetone and resuspended in 75

µL Buffer A (50 mM TEA pH 7.3, 150 mM NaCl, 4% SDS, 0.2% triton X-100, 4 mM EDTA) for 1 hour at

40 °C, shaking at 800 RPM. Samples were centrifuged at 16,000 x g for 2 minutes, 7 µL removed as input,

and the remainder split into +cleavage and -cleavage fractions. The +cleavage sample containing 1% SDS

was treated with 750 µM hydroxylamine (final concentration) for 1 hour at 30 °C, rotating end over end.

Following methanol-chloroform precipitation (ice cold, final concentrations 42% methanol, 16%

chloroform) samples were centrifuged at 16,000 x g for 20 minutes at 4 °C, aqueous layer removed, washed

3X with 100% methanol and air dried. Pellets were resuspended as described in Buffer A and incubated in

1 mM mPEG-Mal (Methoxypolyethylene glycol maleimide, 5 kDa, 63187 Sigma-Aldrich) for 2 hours at

room temp, rotating end over end. Samples were methanol-chloroform precipitated as described,

resuspended in Lysis buffer with 1X SDS-PAGE sample buffer, resolved using SDS-PAGE and visualized

via immunoblotting with mouse anti-FLAG to detect CLMB-FLAG. Endogenous calnexin was detected

using anti-calnexin antibodies. Immunoblots were imaged using a Licor Odyssey CLx and quantified with

Image Studio software.

Detergent-assisted subcellular fractionation:

HEK293 Flp-In TReX lines stably expressing $CLMB_{WT}$, $CLMB_{C7,14S}$, or $CLMB_{G2A}$ were induced with 10

ng/mL doxycycline for 48 hours and 20 µM MG132 (EDM Millipore 474787-10MG) for 4 hours prior to

harvesting. Cells were washed 2X in PBS, harvested and immediately lysed in 600 µL of Cytoplasm Buffer

(10 mM HEPES pH 6.8, 100 mM NaCl, 300 mM sucrose, 3 mM $MgCl_2$, 5 mM EDTA, 0.015% digitonin,

1x halt protease inhibitor cocktail (Thermo Fischer scientific 78442)) at 4 °C for 15 minutes. 60 µL of total

lysate was removed as input and the remainder centrifuged at 1000 x g for 10 min at 4 °C. The supernatant

was removed as cytoplasmic fraction and the pellet washed 2X in PBS and resuspended in 600 µL

membrane buffer (10 mM HEPES pH 7.5, 100 mM NaCl, 300 mM sucrose, 3 mM $MgCl_2$, 3 mM EDTA,

0.5% Triton X-100, 1x halt protease inhibitor cocktail) for 30 minutes at 4 °C turning end over end.

Insoluble material was pelleted by centrifugation at 5000 x g for 10 minutes at 4 °C and supernatant

removed as membrane fraction. Both fractions were centrifuged at 12,000 x g at 4 °C for 10 minutes.

Samples were heated in 1X SDS sample buffer at 95 °C for 5 minutes, resolved via SDS-PAGE,

immunoblotted and CLMB-FLAG detected using anti-FLAG antibodies; anti-GM130 and anti-α-tubulin

were used as membrane and cytosolic markers respectively. Immunoblots were imaged using a Licor

Odyssey CLx and quantified with Image Studio software. CLMB in cytosolic and membrane fractions were

quantified as $CLMB_{Cyto}/(CLMB_{Cyto}+ CLMB_{Mem})$ and $CLMB_{Mem}/(CLMB_{Cyto} + CLMB_{Mem})$, respectively.

Immunofluorescence, microscopy, and image analysis:

MCF7 cells were co-transfected (as described previously) with equal amounts of pcDNA5 encoding

$CLMB_{WT}$, $CLMB_{IxITMut}$, $CLMB_{LxVMut}$, $CLMB_{T44A}$ and pcDNA5 encoding CNA-GFP. The following day

cells were plated on poly-lysine coated 12 mm, #1.5H glass coverslips (ThorLabs). On day three, cells were

washed in PBS and fixed in 100% methanol for 10 minutes at -20 °C. Coverslips were washed 3 times in

1x PBS, blocked (1x PBS, 200 mM glycine, 2.5% FBS) for 30 minutes at room temperature, then incubated

in primary antibody diluted in blocking buffer for 1 hour before being washed 3 times in 1x PBS and

incubated for 1 hour in secondary antibody solution. Following three additional washes, coverslips were

mounted using Prolong Glass mountant with NucBlue stain. After sitting overnight at room temperature,

coverslips were imaged using a Lionheart FX automated widefield microscope with a 20X Plan Fluorite

WD 6.6 NP 0.45 objective. For each coverslip, a 7 by 7 grid of non-overlapping images was automatically

taken. All images were manually assessed and those with defects (debris, bubbles, out-of-focus, etc.) were

filtered out. Images were analyzed using an automated pipeline using CellProfiler [52]. Briefly, cells were

identified by their nuclei and all those lacking CLMB-FLAG staining removed. CLMB-FLAG signal was

used to define the cell body and all cells containing one or more fully saturated pixels in the CLMB-FLAG

or CNA-GFP channels were removed. Pearson's correlation coefficient between CLMB-FLAG and CNA-

GFP was determined for each individual cell. >500 cells per condition were analyzed.

GFP-Trap coimmunoprecipitation:

HEK293 cells were co-transfected (as described previously) with pcDNA5 encoding CNA-GFP and

pcDNA5 encoding $CLMB_{WT}$, $CLMB_{IxITMut}$, $CLMB_{LxVMut}$, $CLMB_{T44A}$, or $CLMB_{LxVMut+T44A}$. 24 hours post-

transfection, cells were washed 2X in PBS and frozen at -80 °C. Cells were lysed in 500 µL lysis buffer (50

mM Tris pH 7.5, 150 mM NaCl, 1% NP-40), aspirated 5X with a 27-gauge needle, centrifuged at 16,000 x

g for 20 min at 4 °C and protein quantified via BCA. 100 mg total protein was removed as input. Lysates

were adjusted to 0.5% NP-40 and 500 µg of protein incubated with pre-washed magnetic GFP-TRAP beads

(ChromoTek gtma-20) for 2 hours at 4 °C, turning end over end. Beads were magnetized, washed 3 times

and eluted with 50 µL of 2x SDS PAGE sample buffer. Samples were resolved via SDS-PAGE and

visualized via immunoblotting with anti-flag antibodies to detect CLMB-FLAG and anti-GFP to detect

CNA-GFP. Anti-GAPDH was used as a loading control for inputs. Immunoblots were imaged using a Licor

Odyssey CLx and CLMB-calcineurin binding was quantified with Image Studio software. Co-purified

CLMB was normalized using the following: $(CLMB_{IP}/CLMB_{Input})/GFP\text{-}CNA_{IP}$. Additionally, all values are

displayed relative to $CLMB_{WT}$ which is set at 1.

Protein expression and purification:

CNA catalytic dead mutant H151Q with an N-terminal His_6-CNA (residues 1–391) and CNB (residues 1–

170) subunits were expressed using *E. coli* BL21(DE3) cells in Lysogeny Broth (LB) supplemented with

50 μg/mL ampicillin. Protein expression was induced with 1 mM isopropyl β-d-1-thiogalactopyranoside

(IPTG) at 18 °C. Cells were lysed using a Fisherbrand™ Model 505 Sonic Dismembrator and lysates were

centrifuged at $46,000 \times g$ at 4 °C for 45 minutes. The protein was purified by incubation with nickel-

nitriloacetic acid (Ni-NTA) agarose resin (Qiagen) for 3h at 4 °C. The resin was washed with 50 mM Tris

pH 8.0, 250 mM NaCl, 20 mM imidazole, and 1 mM TCEP. A buffer containing 50 mM Tris pH 8.0, 250

mM NaCl, 300 mM imidazole, and 1 mM TCEP was used for elution. Calcineurin was further purified by

SEC (Superdex® 200 Increase 10/300 GL) using Phosphate-buffered saline (PBS) and $CaCl_2$ 1mM as the

final buffer.

Fluorescence polarization binding assay:

All FP measurements were performed on a multifunctional microplate reader (EnVision, Perkin Elmer) in

black 384-well microplates (Corning, Cat No 3575) with 30 μL of the assay solution per well. 480-nm

excitation and 535-nm emission filters were used for the FP measurements. In the FP saturation binding

experiments, 5 nM FITC labeled peptides were mixed with increasing concentrations of calcineurin in PBS

buffer (pH 7.4) (commercially available tablets, Thermo Fisher #18912014) and 500 μM $CaCl_2$ buffer (from

0 to 25 μM). Where indicated (Supplemental Fig. 3), these analyses were carried out at different pHs using

a buffer with 20 mM Hepes pH 6.9, 100 mM NaCl, 0.5 mM EDTA, 1 mM TCEP or a buffer containing 50

mM HEPES, pH 8.0, 150 mM NaCl, 1 mM TCEP, 500µM $CaCl_2$. FP values were plotted against the log

of the protein concentrations, and the dissociation constant (apparent K_D) was obtained from the resulting

sigmoidal curve as analyzed in GraphPad Prism 10. Protein concentrations above 5µM resulted in an upturn

in fluorescence polarization signal, indicative of possible aggregation or artifacts; to maintain assay

integrity, the considered concentrations were limited to ≤5µM.

Fluorescence polarization competition assay:

For the competitive binding assay, a mixture containing 5 nM FITC-labeled CLMB peptides and CNA/B

(0.08 and 2 µM for CLMB vs PVIVIT experiments) was incubated with serial dilutions of unlabeled

peptides for 30 minutes at room temperature. The FP values were determined, and the IC_{50} values, i.e., the

concentrations required for 50% displacement of the FITC-labeled peptide, were calculated in GraphPad

Prism 10. The K_i values of competitive inhibitors were calculated based on a previously reported method[53],

further adapted by[54]. Briefly, the relation of K_i and IC_{50} in competitive inhibition as applied to protein-

ligand–inhibitor (P–L–I) interactions, can be summarized by the following equation:

545
$$K_i = \frac{IC_{50}}{(\frac{L}{K_d} + 1)}$$

Where L stands for the concentration of ligand, i.e., the FITC-labeled peptide used in the FP binding assay;

and K_D corresponds to the dissociation constant of the same FITC-labeled peptide resulting from the

performed FP binding assay.

ITC measurements:

Isothermal titration calorimetry (ITC) experiments were performed using an Affinity ITC from TA

Instruments (New Castle, DE) equipped with an autosampler in a buffer containing 50 mM HEPES, pH

8.0, 150 mM NaCl, 1 mM TCEP, 500µM $CaCl_2$ at 25 °C. 40 µM protein solution in the calorimetric cell

was titrated with 100 µM peptide solution using 0.5 µL injection in 200-sec intervals using stirring speed

at 125rpm. The resulting isotherm was fitted with a single site model to yield thermodynamic parameters

of ΔH, ΔS, stoichiometry, and K_D using NanoAnalyze software (TA instruments). All results are

representative of three independent experiments (n = 3).

NMR analyses:

All NMR data were acquired using TopSpin3.4 (Bruker), processed with NMRPipe v.2020/2021 and

analyzed with CCPNmr v3[55, 56]. Structural biology applications were compiled by the SBGrid consortium.

Three-dimensional (3D) NMR experiments were conducted on a Bruker spectrometer operating at 600

MHz, equipped with TCI cryoprobes and z-shielded gradients. Samples of approximately180-µM 2H-, ^{13}C-

and ^{15}N-calcimembrin in 20 mM Hepes pH 6.9, 100 mM NaCl, 0.5 mM EDTA, 1 mM TCEP and 5% v/v

^{2}H2O were used. Temperature was set to 25˚C. Chemical shift assignments are deposited in the Biological

Magnetic Resonance Bank (BMRB) under accession code 52641.

The NMR titration samples were prepared by adding unlabeled calcineurin A/B complex or unlabeled

calcineurin subunit A into solutions of U-[^{15}N]-labeled calcimembrin (50 µM) to reach a ratio of 0.5:1 (25

µM unlabeled calcineurin) or 1:1 (50 µM unlabeled calcineurin). Titration experiments featuring

pCLMB$_{LxVMut}$ (40µM) peptide with free CLMB (50µM), and with calcineurin:CLMB at 1:1 and 0.5:1

stoichiometries were also performed. Two-dimensional (2D) spectra were acquired using SOFAST-HMQC

on a Bruker 800 MHz Avance III spectrometer equipped with a TXO cryoprobe and z-shielded gradients[57].

Data was processed using NmrPipe and analyzed with CCPNmr. Secondary structure prediction was

analyzed using TALOS-N[58] and CheSPI[24].

AlphaFold 3 modeling:

Calcineurin-CLMB complex modeling was done using AlphaFold 3 on AlphaFold server[27]. Modeling was

done using truncated CNA$_{1-391}$, CNB, and two copies of CLMB, either unmodified or phosphorylated at

Thr44. Zn^{2+} and Fe^{3+} (found in the calcineurin active site) and four Ca^{2+} ions (bound to CNB) were included

in modeling. All images were generated in UCSF ChimeraX 1.8[59].

In vivo analysis of CLMB phosphorylation status:

HEK293 FLIP-In TreX cells stably expressing $CLMB_{WT}$-FLAG, $CLMB_{IxITMut}$-FLAG, $CLMB_{LxVMut}$-FLAG,

or $CLMB_{T44A}$-FLAG were transfected with empty vector or pcDNA5 FRT TO encoding CNA-GFP

alongside empty vector or pcDNA5 FRT TO encoding $CLMB_{WT}$-V5, $CLMB_{IxITMut}$-V5, or $CLMB_{LxVMut}$-V5.

Protein expression was induced for 24 hours with 10 ng/mL doxycycline, cells washed 2X in PBS and

stored at -80 °C. Cells were lysed (50 mM Tris pH 8, 150 mM NaCl, 1% NP-40, 0.1% SDS, 0.5% sodium

dodecyl sulfate, 1x Halt protease inhibitor cocktail) and resolved via SDS-PAGE and detected via

immunoblotting.

In vitro CLMB phosphopeptide dephosphorylation:

 $CLMB_{WT}$, $CLMB_{IxITMut}$, and $CLMB_{LxVMut}$ phospho-peptides were synthesized by Vivitide, resuspended in

10 mM Tris pH 8.0 and dephosphorylated at 26 °C in reaction buffer (50 mM Tris pH 8.0, 100 mM NaCl,

6 mM $MgCl_2$, 2 mM $CaCl_2$, 1 mM EGTA, 0.1 mg/mL BSA, 1 mM DTT, 0.5 µM MDCC-PBP (Thermo

Scientific PV4406), 1 µM CaM (Millipore Sigma 208694-1MG)). Reactions were initiated by addition of

CLMB peptide to a final concentration of 100nM. Reactions took place in a total of 50 µL in a corning

black 96 well vessel and were monitored continuously via fluorescence emitted by phosphate bound

MDCC-PBP (EX:425/ EM:460). Phosphate standards were used to convert fluorescence intensity to

phosphate concentration. Data were analyzed using Prism software. Because the experimental design does

not comply with assumptions of standard kinetic models, reaction rates were obtained by fitting data to a

single exponential model as an empirical description of the curves to facilitate comparison between

experimental conditions.

In vitro RII phospho-peptide dephosphorylation:

1 µM RII peptide was dephosphorylated at 26 °C in reaction buffer (50 mM Tris pH 8.0, 100 mM NaCl, 6

mM $MgCl_2$, 2 mM $CaCl_2$, 1 mM EGTA, 0.1 mg/mL BSA, 1 mM DTT, 0.5 µM MDCC-PBP (Thermo

Scientific PV4406), 100nM calmodulin, 10 nM calcineurin, and CLMB peptide at the indicated

concentrations (Millipore Sigma 208694-1MG)). Reactions were initiated by adding Ca^{2+} and CaM to their

final concentrations. Reactions took place in a total of 50 μL in a corning black 96 well vessel and were

monitored continuously via fluorescence emitted by phosphate bound MDCC-PBP (EX:400/ EM:460).

Phosphate standards were used to convert fluorescence intensity to phosphate concentration.

Kinetic Modeling:

Dephosphorylation data were fit by simulation and non-linear regression using Kintek Explorer™. Sigma

values for each individual replicate were obtained using the double exponential afit function. All replicates

were simultaneously fit to the models described in Supplemental Fig.s 8 (a,b). Curve offsets were capped

at a maximum offset of 50nM. Substrate concentration was allowed to vary by 15% during fitting. Initial

parameter values were chosen based on previously published rates, when available. The final offset and

initial substrate concentration values for each curve, along with fitting information, are shown in

Supplemental Tables 1-3. Final graphs display averaged ($n \geq 5$) phosphate released at each timepoint with

the corresponding fit. Simulated data were exported using 200 simulation steps and plotted using GraphPad

Prism.

Statistical analysis

Statistical analysis was done using Graphpad Prism version 10.2.2. All data is displayed as representative

images or mean values with standard deviation error bars unless otherwise noted. All data represent three

independent experiments unless otherwise indicated in figure legends. Image analysis of CLMB-calcineurin

colocalization (Fig. 2C) was conducted using the following number of cells: $CLMB_{WT}$ = 525, $CLMB_{IxITMut}$

= 641, $CLMB_{LxVMut}$ = 500, $CLMB_{T44A}$ = 517. Statistical significance was calculated using one-way analysis

of variance (ANOVA) with corrections for multiple comparisons as indicated.

Data availability:

All original scans of immunoblots, immunofluorescence images, and AlphaFold predictions can be
accessed through Mendeley Data (to be made available upon publication). Plasmids and reagents will be
shared upon request. NMR chemical shift assignments are deposited in the Biological Magnetic Resonance
Bank (BMRB) under accession code 52641.

Acknowledgements:

We thank Norman Davey and members of the Cyert and Arthanari labs, in particular Angela Barth and
Sneha Roy, for helpful discussion. We are especially grateful to Rick Russell (UT Austin) for feedback on
the kinetic modeling and thank Callie Wigington and Makena Pule for their early contributions to the
project. We thank Ilya Bezprozvanny (UT Southwestern) for suggesting the name, calcimembrin.

Funding:

MSC, DAB and RYM are supported by NIH grant R35 GM136243. DAB also acknowledges support from
NIH grant T32 GM007276. HA, JCR, SQ and TV acknowledge grant R01 GM136859 from NIH.

Author Contributions:

MSC, DAB, HA and JCR conceived the studies. SQ and TV provided early NMR analysis of CLMB-
calcineurin complexes. DAB, supervised by MSC, carried out experiments and analyzed data for Figures
1, 2, and 5, and prepared figures. RYM, supervised by MSC, carried out experiments and analyzed data for
Supplementary figures 1 and 2a. JCR, supervised by HA, carried out experiments and analyzed data for
Figures 3 and 4. TV carried out secondary structure analyses. DAB and MSC wrote the manuscript in
consultation with HA and JCR.

References

1.   Bootman MD, Bultynck G. Fundamentals of Cellular Calcium Signaling: A Primer. *Cold*
658   *Spring Harb Perspect Biol* **12**, (2020).

660   2.   Ulengin-Talkish I, Cyert MS. A cellular atlas of calcineurin signaling. *Biochim Biophys*
661   *Acta Mol Cell Res* **1870**, 119366 (2023).

662
663   3.   Aramburu J, Rao A, Klee CB. Calcineurin: from structure to function. *Curr Top Cell*
664   *Regul* **36**, 237-295 (2000).

665
666   4.   Grigoriu S*, et al.* The molecular mechanism of substrate engagement and
667   immunosuppressant inhibition of calcineurin. *PLoS Biol* **11**, e1001492 (2013).

668
669   5.   Roy J, Cyert MS. Identifying New Substrates and Functions for an Old Enzyme:
670   Calcineurin. *Cold Spring Harb Perspect Biol* **12**, (2020).

671
672   6.   Brauer BL*, et al.* Leveraging New Definitions of the LxVP SLiM To Discover Novel
673   Calcineurin Regulators and Substrates. *ACS Chem Biol* **14**, 2672-2682 (2019).

674
675   7.   Sheftic SR, Page R, Peti W. Investigating the human Calcineurin Interaction Network
676   using the piLxVP SLiM. *Sci Rep* **6**, 38920 (2016).

677
678   8.   Tsekitsidou E*, et al.* Calcineurin associates with centrosomes and regulates cilia length
679   maintenance. *J Cell Sci* **136**, (2023).

680
681   9.   Wigington CP*, et al.* Systematic Discovery of Short Linear Motifs Decodes Calcineurin
682   Phosphatase Signaling. *Mol Cell* **79**, 342-358 e312 (2020).

683
684   10.   Nakamura T*, et al.* Overexpression of C16orf74 is involved in aggressive pancreatic
685   cancers. *Oncotarget* **8**, 50460-50475 (2017).

686
687   11.   Meinnel T, Dian C, Giglione C. Myristoylation, an Ancient Protein Modification
688   Mirroring Eukaryogenesis and Evolution. *Trends Biochem Sci* **45**, 619-632 (2020).

689
690   12.   Blanc M, David FPA, van der Goot FG. SwissPalm 2: Protein S-Palmitoylation Database.
691   *Methods Mol Biol* **2009**, 203-214 (2019).

692
693   13.   Yap MC*, et al.* Rapid and selective detection of fatty acylated proteins using omega-
694   alkynyl-fatty acids and click chemistry. *J Lipid Res* **51**, 1566-1580 (2010).

695
14.   Wang Y, Windh RT, Chen CA, Manning DR. N-Myristoylation and betagamma play roles
beyond anchorage in the palmitoylation of the G protein alpha(o) subunit. *J Biol Chem*
274, 37435-37442 (1999).

15.   Dekker FJ, *et al.* Small-molecule inhibition of APT1 affects Ras localization and
signaling. *Nat Chem Biol* **6**, 449-456 (2010).

16.   Percher A, Ramakrishnan S, Thinon E, Yuan X, Yount JS, Hang HC. Mass-tag labeling
reveals site-specific and endogenous levels of protein S-fatty acylation. *Proc Natl Acad*
Sci U S A **113**, 4302-4307 (2016).

17.   F.S. M, Abrami L, Linder ME, Bamji SX, Dickinson BC, van der Goot FG. Mechanisms
and functions of protein S-acylation. *Nat Rev Mol Cell Biol*, (2024).

18.   Hornbeck PV, *et al.* 15 years of PhosphoSitePlus(R): integrating post-translationally
modified sites, disease variants and isoforms. *Nucleic Acids Res* **47**, D433-D441 (2019).

19.   Blumenthal DK, Takio K, Hansen RS, Krebs EG. Dephosphorylation of cAMP-
dependent protein kinase regulatory subunit (type II) by calmodulin-dependent protein
phosphatase. Determinants of substrate specificity. *J Biol Chem* **261**, 8140-8145 (1986).

20.   Perrino BA, Ng LY, Soderling TR. Calcium regulation of calcineurin phosphatase activity
by its B subunit and calmodulin. Role of the autoinhibitory domain. *J Biol Chem* **270**,
7012 (1995).

21.   Aramburu J, Yaffe MB, Lopez-Rodriguez C, Cantley LC, Hogan PG, Rao A. Affinity-
driven peptide selection of an NFAT inhibitor more selective than cyclosporin A. *Science*
285, 2129-2133 (1999).

22.   Li H, Zhang L, Rao A, Harrison SC, Hogan PG. Structure of calcineurin in complex with
PVIVIT peptide: portrait of a low-affinity signalling interaction. *J Mol Biol* **369**, 1296-
1306 (2007).

23.   Shen Y, Bax A. Protein backbone and sidechain torsion angles predicted from NMR
chemical shifts using artificial neural networks. *J Biomol NMR* **56**, 227-241 (2013).

24.   Nielsen JT, Mulder FAA. CheSPI: chemical shift secondary structure population
inference. *J Biomol NMR* **75**, 273-291 (2021).

25.   Solyom Z, Schwarten M, Geist L, Konrat R, Willbold D, Brutscher B. BEST-TROSY
experiments for time-efficient sequential resonance assignment of large disordered
proteins. *J Biomol NMR* **55**, 311-321 (2013).

26. Hendus-Altenburger R, *et al.* Molecular basis for the binding and selective
dephosphorylation of Na(+)/H(+) exchanger 1 by calcineurin. *Nat Commun* **10**, 3489
(2019).

742
743 27. Abramson J, *et al.* Accurate structure prediction of biomolecular interactions with
744   AlphaFold 3. *Nature* **630**, 493-500 (2024).

745
746 28. Shibasaki F, Price ER, Milan D, McKeon F. Role of kinases and the phosphatase
747   calcineurin in the nuclear shuttling of transcription factor NF-AT4. *Nature* **382**, 370-373
748   (1996).

749
750 29. Li H, Rao A, Hogan PG. Structural delineation of the calcineurin-NFAT interaction and
751   its parallels to PP1 targeting interactions. *J Mol Biol* **342**, 1659-1674 (2004).

752
753 30. Johnson KA. Fitting enzyme kinetic data with KinTek Global Kinetic Explorer. *Methods
754   Enzymol* **467**, 601-626 (2009).

755
756 31. Klee CB, Draetta GF, Hubbard MJ. Calcineurin. *Adv Enzym* **61**, 149-200 (1988).

757
758 32. Jespersen N, Barbar E. Emerging Features of Linear Motif-Binding Hub Proteins. *Trends
759   Biochem Sci* **45**, 375-384 (2020).

760
761 33. Li H, Rao A, Hogan PG. Interaction of calcineurin with substrates and targeting proteins.
762   *Trends Cell Biol* **21**, 91-103 (2011).

763
764 34. Lee S, *et al.* The optimized core peptide derived from CABIN1 efficiently inhibits
765   calcineurin-mediated T-cell activation. *Exp Mol Med* **54**, 613-625 (2022).

766
767 35. Nguyen HQ, *et al.* Quantitative mapping of protein-peptide affinity landscapes using
768   spectrally encoded beads. *Elife* **8**, (2019).

769
770 36. O'Brien KT, Mooney C, Lopez C, Pollastri G, Shields DC. Prediction of polyproline II
771   secondary structure propensity in proteins. *R Soc Open Sci* **7**, 191239 (2020).

772
773 37. Tubiana J, *et al.* Funneling modulatory peptide design with generative models: Discovery
774   and characterization of disruptors of calcineurin protein-protein interactions. *PLoS
775   Comput Biol* **19**, e1010874 (2023).

777   38.   Aramburu J, Garcia-Cozar F, Raghavan A, Okamura H, Rao A, Hogan PG. Selective
778   inhibition of NFAT activation by a peptide spanning the calcineurin targeting site of
779   NFAT. *Mol Cell* **1**, 627-637 (1998).

780
781   39.   Li SJ*, et al.* Cooperative autoinhibition and multi-level activation mechanisms of
782   calcineurin. *Cell Res* **26**, 336-349 (2016).

783
784   40.   Oliveria SF, Dell'Acqua ML, Sather WA. AKAP79/150 anchoring of calcineurin controls
785   neuronal L-type Ca2+ channel activity and nuclear signaling. *Neuron* **55**, 261- 275
786   (2007).

787
788   41.   Li H, Pink MD, Murphy JG, Stein A, Dell'Acqua ML, Hogan PG. Balanced interactions
789   of calcineurin with AKAP79 regulate Ca2+-calcineurin-NFAT signaling. *Nat Struct Mol*
790   *Biol* **19**, 337-345 (2012).

791
792   42.   Church TW*, et al.* AKAP79 enables calcineurin to directly suppress protein kinase A
793   activity. *Elife* **10**, (2021).

794
795   43.   Huang WYC, Boxer SG, Ferrell JE, Jr. Membrane localization accelerates association
796   under conditions relevant to cellular signaling. *Proc Natl Acad Sci U S A* **121**,
797   e2319491121 (2024).

798
799   44.   Johnson JL*, et al.* An atlas of substrate specificities for the human serine/threonine
800   kinome. *Nature* **613**, 759-766 (2023).

801
802   45.   Li M, Jiang J, Yue L. Functional characterization of homo- and heteromeric channel
803   kinases TRPM6 and TRPM7. *J Gen Physiol* **127**, 525-537 (2006).

804
805   46.   Goldman A*, et al.* The Calcineurin Signaling Network Evolves via Conserved Kinase-
806   Phosphatase Modules that Transcend Substrate Identity. *Mol Cell* **55**, 422-435 (2014).

807
808   47.   Hirschi A, Cecchini M, Steinhardt RC, Schamber MR, Dick FA, Rubin SM. An
809   overlapping kinase and phosphatase docking site regulates activity of the retinoblastoma
810   protein. *Nat Struct Mol Biol* **17**, 1051-1057 (2010).

811
812   48.   Jara KA*, et al.* Multivalency, autoinhibition, and protein disorder in the regulation of
813   interactions of dynein intermediate chain with dynactin and the nuclear distribution
814   protein. *Elife* **11**, (2022).

816   49.   Simonsen S*, et al.* Extreme multivalency and a composite short linear motif facilitate
817   PCNA-binding, localisation and abundance of p21 (CDKN1A). *FEBS J*, (2025).

818
819   50.   Dinh J*, et al.* The microprotein C16orf74/MICT1 promotes thermogenesis in brown
820   adipose tissue. *EMBO J*, (2025).

821
822   51.   Uhlen M*, et al.* Towards a knowledge-based Human Protein Atlas. *Nat Biotechnol* **28**,
823   1248-1250 (2010).

824
825   52.   Stirling DR, Swain-Bowden MJ, Lucas AM, Carpenter AE, Cimini BA, Goodman A.
826   CellProfiler 4: improvements in speed, utility and usability. *BMC Bioinformatics* **22**, 433
827   (2021).

828
829   53.   Cheng Y, Prusoff WH. Relationship between the inhibition constant (K1) and the
830   concentration of inhibitor which causes 50 per cent inhibition (I50) of an enzymatic
831   reaction. *Biochem Pharmacol* **22**, 3099-3108 (1973).

832
833   54.   Cer RZ, Mudunuri U, Stephens R, Lebeda FJ. IC50-to-Ki: a web-based tool for
834   converting IC50 to Ki values for inhibitors of enzyme activity and ligand binding.
835   *Nucleic Acids Res* **37**, W441-445 (2009).

836
837   55.   Delaglio F, Grzesiek S, Vuister GW, Zhu G, Pfeifer J, Bax A. NMRPipe: a
838   multidimensional spectral processing system based on UNIX pipes. *J Biomol NMR* **6**,
839   277-293 (1995).

840
841   56.   Vranken WF*, et al.* The CCPN data model for NMR spectroscopy: development of a
842   software pipeline. *Proteins* **59**, 687-696 (2005).

843
844   57.   Schanda P, Kupce E, Brutscher B. SOFAST-HMQC experiments for recording two-
845   dimensional heteronuclear correlation spectra of proteins within a few seconds. *J Biomol*
846   *NMR* **33**, 199-211 (2005).

847
848   58.   Shen Y, Bax A. Protein structural information derived from NMR chemical shift with the
849   neural network program TALOS-N. *Methods Mol Biol* **1260**, 17-32 (2015).

850
851   59.   Meng EC*, et al.* UCSF ChimeraX: Tools for structure building and analysis. *Protein Sci*
852   **32**, e4792 (2023).

Figure Legends

Fig. 1 Calcimembrin localizes to membranes via protein lipidation. **a.** CLMB schematic showing N-

myristoylation (red); S-acylation (yellow), phosphorylation (green), calcineurin-binding motifs LxVP

(pink) and PxIxIT (purple). Sequences of wild-type and mutated regions shown. **b.** Representative

immunoblot showing metabolic labeling of CLMB-Flag (wild-type or indicated mutants) with palmitate

(17-ODYA) or myristate (YnMyr) analogs, analyzed via CLICK chemistry with biotin-azide and

detection with IRdye 800CW streptavidin ("Biotin"). Inputs and immunoprecipitated (IP) samples are

shown. Asterisks indicate CLMB dimers. (n = 3 independent experiments). **c.** Representative immunoblot

showing S-acylation of CLMB-FLAG (wild-type or indicated mutants) and endogenous calnexin

(positive control) detected via Acyl-PEG exchange. Asterisks show PEGylation events. Input, cleaved

(NH_2OH) and PEGylated (mPEG), and non-cleaved PEGylated conditions are indicated (n = 3

independent experiments). **d.** Representative in-gel fluorescence scan showing pulse-chase analysis of

CLMB-FLAG using palmitate (17-ODYA) or methionine (L-AHA) analogs detected via CLICK

chemistry with Cy5 or AZDye 488, respectively. Incorporation into CLMB-FLAG with no analog (Neg

Ctrl), at 0 or 30 minutes after unlabeled chase with DMSO (control) or PalmB. shown (n = 3 independent

experiments). **e.** Data from (**d**) showing mean ± SEM of 17-ODYA signal normalized to L-AHA; * p <

0.05, T0 vs PalmB p = 0.0103, ** p < 0.01, DMSO vs PalmB p = 0.007**. f.** Representative immunoblot

showing sub-cellular fractionation of CLMB-FLAG (wild-type or indicated mutants) with input,

cytoplasmic (cyto), and membrane (mem) fractions. GM130 and α-tubulin mark mem and cyto fractions,

respectively (n= 3 independent experiments). **g**. Data from (**f**) showing mean ± SEM cyto or mem signals

normalized to total (cyto + mem); n.s. not significant, **** p < .0001. CLMBC$_{7,14S}$ vs CLMB$_{G2A}$:

cytoplasm, p = 0.238, membrane p = 0.238. For (**e, g**) p-values were calculated by two-way ANOVA

corrected by Tukey's multiple comparison.

Fig. 2 Composite LxVPxIxIT SLiM in calcimembrin mediates binding to and dephosphorylation by

calcineurin. a. Representative immunoblot showing co-purification of CLMB-FLAG (wild-type and

indicated mutants) with CNA-GFP. Inputs and purified (IP) samples shown (n = 3 independent

experiments). Glyceraldehyde-3-Phosphate Dehydrogenase (GAPDH) was used as a loading control. **b.**

Data from (**a**) showing mean ± SEM of co-purified CLMB-FLAG normalized to CLMB-FLAG input

levels and precipitated CNA-GFP. Indicated p-values calculated by two-way ANOVA corrected by

Dunnett's multiple comparison test. **** p < 0.0001, *** p<0.001, * p<0.05; $CLMB_{WT}$ vs $CLMB_{LxVmut}$ p

= 0.0325, $CLMB_{WT}$ vs $CLMB_{T44A}$ p = 0.0002. **c.** Representative images of MCF7 cells co-transfected

with CLMB-FLAG (wild-type or indicated mutants) and CNA-GFP. Fixed cells immunostained with

anti-FLAG (magenta), anti-GFP (yellow), and nuclei (DNA) with Hoechst 33342. Scale bar = 50 μm.

Arrows indicate some examples of CLMB and CNA co-localization to plasma membrane. Data from (**c**)

showing median with 95% CI Pearson's correlation coefficient of CLMB-calcineurin colocalization. Each

point represents a single cell (n ≥ 500 for each condition). p-values calculated by Kruskal-Wallis with

Dunn's multiple comparisons correction. * p < 0.05 (p = 0.0116), ** p < 0.01 (p = 0.0092), **** p <

0.0001. **e** Dephosphorylation of $pCLMB_{WT}$ (blue) or $pCLMB_{LxVMut}$ (pink) phosphopeptides with 20 nM

CNA/B and 1 μM CaM. Mean ± SEM of phosphate released displayed for ≥ 4 replicates at each timepoint

with reaction rate as shown.

Fig. 3 Thr44 Phosphorylation and the LxVP motif contribute to PxIxIT binding. a. Determination

of the affinity between CNA/B and unlabeled $pCLMB_{WT}$ peptide using isothermal calorimetry.

Dissociation constant (K_D) = 37 ± 1.22 nM, reaction stoichiometry (n value) = 1.00 ± 0.02,

enthalpy (ΔH) = -3.97 ± 0.76, and entropy (TΔS) = 20.8 ± 3.01 are shown. Data are mean ± s.d.

for *n* = 3 independent experiments. **b.** Binding isotherm obtained by titration of unlabeled

$pCLMB_{WT}$ peptide into CNA/B (40 μM). **c.** Fluorescence polarization (FP) saturation binding

curves of FITC-$pCLMB_{WT}$ and FITC-$CLMB_{WT}$ (5nM) incubated with serial dilutions of CNA/B

(0 to 5µM). FP data were fitted to a one-site binding model (mean ± s.d., n = 2 independent

experiments). **d**. Competitive binding assay of unlabeled pCLMB$_{WT}$, CLMB$_{WT}$, pCLMB$_{IxITMut}$,

and pCLMB$_{LxVMut}$ peptides titrated against FITC-PVIVIT peptide (5 nM) in the presence of 2

µM CNA/B. Data are mean ± s.d. of n = 2 independent experiment. **e.** Competitive binding assay

of unlabeled pCLMB$_{WT}$, CLMB$_{WT}$, pCLMB$_{IxITMut}$, and pCLMB$_{LxVMut}$ peptides titrated against

FITC-pCLMB$_{WT}$ peptide, (5 nM) in the presence of 0.08 µM CNA/B. Data are mean ± s.d. for n

= 2 independent experiments. **f.** Table displaying the experimentally obtained IC$_{50}$ values and

converted K$_i$ values from data in (**d, e**).

Fig. 4 CLMB-calcineurin binding extends beyond the LxVP and PxIxIT motifs. a. ^{15}N HSQC spectra

of free CLMB with the resonance assignment. **b.** NMR spectral analysis of CLMB interaction with CNA

at 2:1 (CLMB: CNA); **c**. with CNA at 1:1(CLMB: CNA); **d.** with CNA/B at 2:1 (CLMB: CNA/B); **e.**

with CNA/B at 1:1 (CLMB: CNA/B). **b-e.** Graphs display the intensity ratios of CLMB resonances with

and without the titrant (CNA or CNA/B) at different molar ratios. The ratios corresponding to the LxV

motif (pink) and IxIT (purple) are highlighted. Residues with no access to the intensity ratio are

represented with negative numbers, and peaks resulting in complete broadening are shown in green (I/I0

= -0.4). Unassigned residues in apo CLMB are blue (I/I0 = -0.2), prolines (which are not represented in

^{15}N-HSQC are grey (I/I0 = -0.2), and residues for which the bound resonance cannot be unambiguously

assigned are red, (I/I0 = -0.2). * Residues with I/I0 higher than 1 due to overlapping resonances.

Fig. 5 Calcineurin dephosphorylates calcimembrin by forming multimers. a. Representative

immunoblot showing phosphorylation-induced mobility shifts in CLMB-FLAG (wild-type or indicated

mutants) when co-expressed with GFP-CNA and CLMB-V5 (wild-type or indicated mutants). GAPDH

was used as a loading control **b.** Data from (**a**) showing mean ± SEM phospho-CLMB-FLAG normalized

to total CLMB-FLAG. p-values calculated by two-way ANOVA corrected by Dunnett's multiple

comparison test (n = 3 independent experiments). n.s., not significant; without vs with CLMB$_{IxIxITmut}$-V5

expression: p = 0.995, 0.748, 0.999 from left to right. CLMB$_{IxITmut}$ without vs with CNA-GFP p = 0.626.

**** p < 0.0001, *** p < 0.001, p = 0.0005. **c.** Cartoon representation of calcineurin (turquoise)-CLMB

complex formation and CLMB dephosphorylation (removal of pink "P") under conditions in (**a**), i.e.

CLMB-FLAG (blue with "F") overexpression with or without co-expression of Calcineurin-GFP and

CLMB-V5 (green with "V") as indicated. Top row shows CLMB$_{WT}$-FLAG, bottom row shows

CLMB$_{IXITMut}$-FLAG (labelled LDVPxxxx). **d**. (Top) Experimental data showing phosphate released from

100 nM pCLMB$_{WT}$ (blue) or pCLMB$_{IxITMut}$ (purple) phosphopeptide dephosphorylated by 20 nM (left),

100 nM (middle) or 200 nM (right) CNA/B and 1 μM calmodulin. Each dephosphorylation rate is

displayed and data is mean ± SEM for ≥ 5 replicates at each timepoint. (Bottom) Overlay of experimental

data (circles) with results of kinetic simulations (solid lines) using the model shown in Fig. S5a, b for

each calcineurin concentration.

Fig. 6 Models explaining how calcimembrin may modulate calcineurin signaling. a (1) CLMB

localizes to membranes via dynamic acylation where it (2) recruits calcineurin (turquoise). At low CLMB

levels, calcineurin remains tightly bound to phosphorylated CLMB (purple) regardless of Ca^{2+} signals. (3)

At higher CLMB levels, CLMB (purple and green) forms multimers with calcineurin and is

dephosphorylated upon calcineurin activation by Ca^{2+} and CaM. (4) Calcineurin is then released and can

engage and dephosphorylate membrane-associated substrates (red) that contain PxIxT and LxVP SLiMs.

b (1) CLMB localizes to membranes via dynamic acylation where it (2) recruits calcineurin (turquoise).

At low CLMB levels, calcineurin remains tightly bound to phosphorylated CLMB (purple). (3) Ca^{2+} and

CaM activate calcineurin, allowing it to engage and dephosphorylate LxVP-containing substrates (red).

(4) At high CLMB levels (purple and green) calcineurin-CLMB multimers form. (5) This promotes

CLMB dephosphorylation and triggers calcineurin release to terminate signaling.

a

1- MGLKMSCLKGFQMCVSSSSSSHDEAPVLNDKHLDVPDIIITPPTPTGMMLPRDLGSTVWLDETGSCPDDGEIDPEA -76

G2   C7   C14   33- LxVP -36   T44
                                            36- PxIxIT -41

Construct Name	Sequence
CLMB_WT	33-LDVPDIIITPPTP-45
CLMB_LxVMut	33-ADAPDIIITPPTP-45
CLMB_IxITMut	33-LDVPDAIAAPPTP-45

b

IP: FLAG
    Biotin
    CLMB-FLAG
Input CLMB-FLAG

c

CLMB-FLAG
Calnexin

d

17-ODYA (Cy5)
L-AHA (AZDye 488)

e

f

CLMB-FLAG
GM130
α-Tubulin

g

Peptide Competitor	FITC-PVIVIT		FITC-pCLMB$_{WT}$	
	IC$_{50}$(nM)	K$_i$ (nM)	IC$_{50}$(nM)	K$_i$ (nM)
pCLMB$_{WT}$	81.4 ± 16.2	80.9 ± 16.1	18.9 ± 1.2	13.4 ± 0.8
pCLMB$_{IxITMut}$	NA	NA	6687 ± 878	4754 ± 624
pCLMB$_{LxVMut}$	159 ± 30	158± 30	103 ± 9	73.2 ± 6.4
CLMB$_{WT}$	267 ± 21	265 ± 21	585 ± 57	426 ± 40

a

b

Ca²⁺
Channel

S-Acylation

N-Myristoylation

Calcimembrin

Phosphate

P

Calcineurin

Calcium/
Calmodulin

Calcineurin
Substrate