Functional genomics reveals strain-specific genetic requirements conferring hypoxic

growth in *Mycobacterium intracellulare*

Yoshitaka Tateishi[1,2*], Yuriko Ozeki[1], Akihito Nishiyama[1], Yuta Morishige[3], Yusuke

Minato[4,5], Anthony D. Baughn[5], Sohkichi Matsumoto[1,6,7,8]

1. Department of Bacteriology, Graduate School of Medicine and Dental Sciences,

Niigata University, 1-757, Asahimachi-dori, Chuo-ku, Niigata 951-8510, Japan.

2. Department of Microbiology, Fukushima Medical University, 1, Hikariga-oka,

Fukushima City, Fukushima, 960-1295, Japan.

3. Department of Mycobacterium Reference and Research, Research Institute of

Tuberculosis, Japan Anti-Tuberculosis Association, Kiyose, Tokyo, 2048533, Japan.

4. Department of Microbiology, Fujita Health University School of Medicine, Toyoake,

Aichi, Japan.

5. Department of Microbiology and Immunology, University of Minnesota Medical

School, 689 23[rd] Avenue S.E. Microbiology Research Facility, Minneapolis, 55455,

MN, USA.

6. Laboratory of Tuberculosis, Institute of Tropical Disease, Universitas Airlangga,

Kampus C JI. Mulyorejo, Surabaya, 60113, Indonesia

7. Department of Bacteriology, Osaka Metropolitan University Graduate School of

Medicine, 1-4-3 Asahi-machi, Abeno-ku, Osaka 545-8586, Japan

8. Division of Research Aids, Hokkaido University Institute for Vaccine Research &

Development, Kita 20, Nishi 10, Kita-Ku, Sapporo, Hokkaido 001-0020, Japan

e-mail: y-tateishi@med.niigata-u.ac.jp, y-tate25@fmu.ac.jp

Abstract

Mycobacterium intracellulare is a major etiological agent of the recently expanding

Mycobacterium avium–intracellulare complex pulmonary disease (MAC-PD).

Therapeutic regimens that include a combination of macrolides and antituberculous

drugs have been implemented with limited success. To identify novel targets for drug

development that accommodate the genomic diversity of *M. avium–intracellulare*, we

subjected eight clinical MAC-PD isolates and the type strain ATCC13950 to

genome-wide profiling to comprehensively identify universally essential functions by

transposon sequencing (TnSeq). Among these strains, we identified 131 shared essential

or growth-defect-associated genes by TnSeq. Unlike the type strain, the clinical strains

showed increased requirements for genes involved in gluconeogenesis and the type VII

secretion system under standard growth conditions, the same genes required for hypoxic

pellicle-type biofilm formation in ATCC13950. Consistent with the central role of

hypoxia in the evolution of *M. intracellulare*, the clinical MAC-PD strains showed more

rapid adaptation to hypoxic growth than the type strain. Importantly, the increased

requirements of hypoxic fitness genes were confirmed in a mouse lung infection model.

These findings confirm the concordant genetic requirements under hypoxic conditions

in vitro and hypoxia-related conditions *in vivo*, and highlight the importance of using

clinical strains and host-relevant growth conditions to identify high-value targets for

drug development.

Introduction

In contrast to the ongoing decline in the rate of tuberculosis, *Mycobacterium*

avium–intracellulare complex pulmonary disease (MAC-PD) is increasing in many

parts of the world[1,2]. The standard therapy for MAC-PD involves multidrug

chemotherapy that includes clarithromycin and several antituberculous drugs.

Unfortunately, treatment failure is common, resulting in relapse, disease progression,

and death[3]. Therefore, it is critical to investigate the bacterial physiological

characteristics that underlie the ability of MAC-PD strains to establish and maintain

chronic drug-recalcitrant infections.

Recent advances in comparative genomics have revealed major differences in

the genomic features of reference type strains and circulating pathogenic clinical

isolates. Extensive genomic diversity has been recognized across clinical isolates of

nontuberculous mycobacteria (NTM). *Mycobacterium avium* has been divided into four

subspecies including subspecies, *avium*, *hominissuis*, *silvaticum*, and *paratuberculosis*[4].

Whereas *M. intracellulare* has been divided into two subgroups, typical *M.*

intracellulare (TMI) and *M. paraintracellulare–M. indicus pranii* (MP-MIP), with

currently two additional distinguished subspecies, *yongonense*, and *chimaera*[5,6]. Despite

the increasing number of sequenced MAC-PD strains, the molecular genetic

mechanisms that underlie their virulence and pathogenesis are still poorly understood.

The burgeoning field of functional genomics has allowed the rapid assessment

of molecular aspects of organisms on a genome-wide scale. Transposon (Tn) sequencing

(TnSeq) is a functional genomic approach that combines saturation-level

transposon-insertion mutagenesis with next-generation sequencing (NGS) to

comprehensively evaluate the fitness costs associated with gene inactivation in bacteria.

Over the last decade, TnSeq has been applied to numerous bacterial species using

various approaches[7]. Whereas initial studies applied TnSeq to reference strains, recent

approaches have included the comparative assessment of representative clinical strains

to better understand the fundamental aspects of pathogen biology[8,9]. We have previously

used TnSeq to globally characterize the genes that are essential for growth versus those

that are essential for hypoxic pellicle formation in the *M. intracellulare* type strain

ATCC13950[10]. However, given the recent expansion of MAC-PD and the

well-recognized high levels of genomic diversity in this complex of bacteria, data

obtained from the study of a decades-old reference strain is not expected to represent the

diversity of extant clinical isolates. ATCC13950 was originally isolated from the

abdominal lymph node of the 34-month-old female[11]. Even though the etiological

bacterial species are the same, the clinical manifestation of infant lymphadenitis differs

markedly from MAC-PD, which often occurs in middle-aged and elderly female

patients with no predisposing immunological disorder. So far, there are a total of two

reports on TnSeq that compare genetic requirements between clinical mycobacterial

strains and the type strains in *M. tuberculosis*[8] and *M. abscessus*[9]. They reported the

diversity of genetic requirements among strains including the type strain such as *M.*

tuberculosis H37Rv and *M. abscessus* ATCC19977. Therefore, it is essential to

investigate a collection of strains that captures the diversity of recent clinical MAC-PD

strains in order to define key molecular aspects of their pathogenesis and virulence.

In this study, we used TnSeq to analyze a set of recently isolated clinical *M.*

intracellulare strains with diverse genotypes. We integrated the gene essentiality data

for these strains and the type strain to identify the common essential and

growth-defect-associated genes that represent the genomic diversity of this group of

pathogens. We also identified the greater requirements of genes involved in

gluconeogenesis, the type VII secretion system, and cysteine desulfurase in the clinical

MAC-PD strains than in the type strain. Furthermore, the requirement for these genes

was confirmed in a mouse lung infection model. The profiles of genetic requirements in

clinical MAC-PD strains suggest the mechanism of hypoxic adaptation in NTM in

terms of promising drug targets, especially for strains that cause clinical MAC-PD.

Results

Common essential and growth-defect-associated genes representing the genomic

diversity of *M. intracellulare* strains

To obtain comprehensive information on the genome-wide gene essentiality of *M.*

intracellulare, we included nine representative strains of *M. intracellulare* with diverse

genotypes in this study (Fig. 1a, b, Supplementary Fig. 1). We used TRANSIT[12] to

obtain 2–6 million reads of Tn insertions. The average number of Tn insertion reads was

> 50 at Tn-inserted TA sites, which confirmed the TnSeq data (Supplementary Table 1).

The numbers of essential genes in the clinical MAC-PD strains, calculated with the

hidden Markov model (HMM; a transition probability model), tended to be smaller than

that in ATCC13950 (Supplementary Table 2). In total, 131 genes were identified as

essential or growth-defect-associated with the HMM analysis across all *M.*

intracellulare strains, including the eight MAC-PD clinical strains and ATCC13950 (Fig.

2a, Supplementary Table 3). The genes identified as universal essential or

growth-defect-associated were involved in fundamental functions, such as glyoxylate

metabolism, purine/pyrimidine metabolism, amino acid synthesis, lipid biosynthesis,

arabinogalactan/peptidoglycan biosynthesis, porphyrin metabolism, DNA replication,

transcription, amino acid tRNA ligases, ribosomal proteins, general secretion proteins,

components of iron–sulfur cluster assembly, some ABC transporters, and the type VII

secretion system. The genes identified corresponded to the genes that encode

conventional and under-development drug targets, such as those encoding gyrase,

arabinosyltransferase, enoyl-(acyl-carrier-protein) reductase, alanine racemase,

DNA-directed RNA polymerase, malate synthase[13] and F_0F_1 ATP synthase (Fig. 2b).

These data suggest that the set of universal essential or growth-defect-associated genes

identified with the HMM analysis of *M. intracellulare* strains with diverse genotypes

has potential utility as antibiotic drug targets.

By comparative genomic analysis, we have revealed 3,153 core genes among

55 *M. intracellulare* strains including those used in this study[5] (Supplementary Fig. 1).

Of these core genes, 439 genes were classified as strain-dependent essential or

growth-defect-associated with the HMM analysis (Supplementary Table 4). And among

5,824 accessory genes, 222 genes were classified as essential or

growth-defect-associated with the HMM analysis. The proportion of these

strain-dependent and accessory essential or growth-defect-associated genes was

comparable to the similar analysis of *Streptococcus pneumoniae*[14]. Of note, 17 (3.8%)

of strain-dependent essential or growth-defect-associated genes and 14 (6.3%) of

accessory essential or growth-defect-associated genes were also included in the 175

genes required for hypoxic pellicle formation in ATCC13950 reported in our previous

study[10] (Supplementary Table 5). Gene set enrichment analysis (GSEA)[15] showed that

9.1% (16 out of 175) genes were hit as core enrichment (Supplementary Table 5). Of

them, 4 genes were hit commonly as genes showing increased genetic requirements

analyzed by resampling plus HMM analyses including genes of phosphoenolpyruvate

carboxykinase *pckA* (OCU_RS48660), type VII secretion-associated serine protease

mycP5 (OCU_RS38275), type VII secretion protein eccC5 (OCU_RS38345) and

glycine cleavage system amino-methyltransferase *gcvT* (OCU_RS35955).

Partial overlap of the genes showing increased genetic requirements in clinical

MAC-PD strains with those required for hypoxic pellicle formation in ATCC13950

The profiles of genetic requirements in each bacterial strain reflect the adaptation to

the environment in which each strain lives. When the strains are placed in a special

environment, they can adapt to the situation by altering the profiles of genetic

requirements, resulting in the remodeling of metabolic pathways. The sharing of

strain-dependent and accessory essential or growth-defect-associated genes with genes

required for hypoxic pellicle formation in ATCC13950 prompts us to consider that the

profiles of genetic requirements in clinical MAC-PD strains may be associated with the

genes required for hypoxic pellicle formation in ATCC13950. To clarify this, we

compared the genetic requirements of the clinical MAC-PD strains with that in

ATCC13950 by integrating the HMM analysis with the resampling analysis, a method

of calculating the number of significantly altered Tn insertion reads (a gene-based

permutation model). With this integrating analysis, 121 genes in the clinical MAC-PD

strains were shown to have significantly fewer Tn insertions than ATCC13950 (Fig. 3,

Supplementary Tables 6, 7). Of these, nine genes were identified in 3–5 clinical

MAC-PD strains. These included genes encoding phosphatases (OCU_RS30185,

OCU_RS35795), cysteine desulfurase (*csd* [OCU_RS40305]), methyl transferase

(OCU_RS41540), a hypothetical protein (OCU_RS47290), type VII secretion

components (*mycP5* [OCU_RS38275], *eccC5* [OCU_RS38345]), a glycine cleavage

system aminomethyltransferase (*gcvT* [OCU_RS35955]), and phosphoenolpyruvate

carboxykinase (*pckA* [OCU_RS48660]) (Figs 3, 4). The genes involved in

gluconeogenesis (such as *pckA*), the glycine cleavage system, and the type VII secretion

system were also identified as required for hypoxic pellicle formation in ATCC13950 in

our previous study[9]. The Tn insertion reads were significantly reduced in the pathways

of gluconeogenesis, including in fructose-1,6-bisphosphatase (*glpX*),

phosphoenolpyruvate carboxykinase (*pckA*), and pyruvate carboxylase (*pca*), but were

significantly increased in those of glycolysis, including in pyruvate kinase (*pykA*) and

pyruvate dehydrogenase (*aceE*, *lpdC*), suggesting that carbohydrate metabolism was

metabolically remodeled in the clinical MAC-PD strains, which mimics the metabolism

observed during hypoxic pellicle formation in ATCC13950 (Fig. 5).

On the other hand, 162 genes were detected with significantly more Tn insertions in

the clinical MAC-PD strains than in ATCC13950 (Fig. 3). Of these, 12 genes were

identified in more than seven strains. The increased Tn insertion reads in the genes

encoding superoxide dismutase (OCU_RS25945) and pyruvate dehydrogenase (*aceE*

[OCU_RS35720]) in the clinical MAC-PD strains suggest the inessentiality of the

detoxification of oxygen radicals and glycolysis metabolism in these strains (Figs 3–5).

These genes also included those encoding amidohydrolases (OCU_RS34400,

OCU_RS34425, OCU_RS34445), some transcription factors (OCU_RS28440,

OCU_RS30230), a CoA transferase (OCU_RS34390), and some predicted transporters

(OCU_RS28435, OCU_RS39950, OCU_RS39955, OCU_RS44685). Furthermore, the

increase in Tn insertion reads in predicted multidrug-resistant genes, such as those

encoding the DrrAB-ABC transporter complex (OCU_RS39950, OCU_RS39955)[16] and

a putative lipoprotein LpqB (OCU_RS44685), encoded by the two-component system

mtrAB operon[17-18], in the clinical MAC-PD strains were found.

In order to investigate the effect of gene duplication on the change of genetic

requirements between strains, we conducted blast search for the 162 genes showing

increased Tn insertion reads in the clinical MAC-PD strains compared to ATCC13950.

We found that M019 has duplicate genes of OCU_RS44705 coding

adenosylhomocysteinase (LOCUS_42940: ahcY_1, LOCUS_21000: ahcY_2). However,

there were no duplicate genes found in the remaining 161 genes. These data suggest that

gene duplication has minor effects on the change of genetic requirements between

strains. Rather, sequence differences and accessory genes may play a key role in

determining the difference of genetic requirements.

Identification of genes in the clinical MAC-PD strains required for mouse lung

infection

Hypoxia is the major environmental condition in natural water[19], biofilms[20], and

infections in humans and animals characterized by caseous granuloma lesions[21]. The

impact of hypoxia on mycobacteria under various ecological circumstances implies that

the genes required for pathogenesis of MAC-PD may be in some degrees, overlapped

with the genes with increased requirements in the clinical MAC-PD strains compared to

ATCC13950, and also with the genes required for hypoxic pellicle formation in

ATCC13950. To identify genes required for *in vivo* infection of clinical MAC-PD

strains, we performed TnSeq in mice infected intratracheally with the Tn mutant library

strains of M.i.198 or M.i.27, which we had demonstrated to be virulent in mice (Fig. 1b,

Supplementary Fig. 3)[22]. The time course of the changes in the bacterial burden showed

a pattern similar to those of the wild-type strains M.i.198 and M.i.27, respectively,

except that it was not possible to harvest sufficient colonies (as few as 10^4/mouse) in the

few mice infected with the M.i.27 Tn mutant strain in week 8 and week 16

(Supplementary Fig. 3b, Supplementary Table 8). With a resampling analysis, 629 and

512 genes were shown to be required for *in vivo* infection with transposon mutant

library strains M.i.27 and M.i.198, respectively, at least at one time point between day 1

and week 16 of infection (Fig. 6a, Supplementary Fig. 3c, d, Supplementary Tables 9,

10). Of these genes, 72 (11.9%) and 50 (9.76%) in M.i.27 and M.i.198, respectively,

were also required for hypoxic pellicle formation in ATCC13950. Forty-one genes also

required for hypoxic pellicle formation in ATCC13950 were identified in both M.i.27

and M.i.198, and included genes associated with succinate production (multifunctional

2-oxoglutarate metabolism enzyme) and the glyoxylate cycle (isocitrate lyase), cysteine

and methionine synthesis (thiosulfate sulfurtransferase [CysA],

5-methyltetrahydrofolate-homocysteine methyltransferase [MetH]), the

serine/threonine-protein kinase PknG system, proteasome subunits (PrcA, PrcB), DNA

repair UvrC, secretion protein EspR, cell-wall synthesis proteins (such as

phosphatidylinositol [polyprenol-phosphate-mannose-dependent

alpha-[1-2]-phosphatidylinositol pentamannoside mannosyltransferase], peptidoglycan

(penicillin-binding protein A pbpA), and L,D-transpeptidase 2), putative membrane

proteins and transporters (such as the MmpL family, transport accessory protein MmpS3,

transport protein MmpL11, vitamin B_{12} transport ATP-binding protein BacA, ABC

transporter substrate-binding protein, and ABC transporter permease), and mammalian

cell entry (Mce) proteins (OCU_RS48855–48875) (Supplementary Table 11).

At all timepoints from Day1 to Week 16, there were 74 and 99 genes were

shown to be required for lung infection with M.i.27 and M.i.198, respectively. Of them,

21 (28.3%) and 12 (12.1%) genes belonged to the genes involved in hypoxic pellicle

formation in the type strain. At timepoints throughout Week 4 to Week 16 except for

Day1, there were 121 and 172 genes were shown to be required for lung infection with

M.i.27 and M.i.198, respectively. Of them, 21 (23.1%) and 30 (18.0%) genes belonged

to genes involved in hypoxic pellicle formation in the type strain. These hypoxic

pellicle-associated genes detected both in M.i.27 and M.i.198 encoded methionine

synthesis (MetH), acyl-CoA dehydrogenase, isocitrate lyase, MMPL family transporter

(MmpL11) (from Day1 to Week 16), additionally multifunctional oxoglutarate

decarboxylase/dehydrogenase, proteasome subunits (PrcA, PrcB), ABC transporter

ATP-binding protein/permease, lipase chaperone (from Week 4 to Week 16) (new

Supplementary Tables 12 and 13).

The genes only identified in M.i.27 that were also required for hypoxic pellicle

formation in ATCC13950 included genes associated with gluconeogenesis (*pckA*, *glpX*),

the glycine cleavage system (*gcvT*), the type VII secretion system (*eccD5, espG5,*

eccC5 [OCU_RS38280-OCU_RS38285-OCU_RS38345], *eccC3* [OCU_RS48145]),

and D-inositol 3-phosphate glycosyltransferase (Supplementary Table 11). These three

systems, gluconeogenesis, the glycine cleavage system, and the type VII secretion

system, also showed increased genetic requirements in 3–5 clinical MAC-PD strains

than in ATCC13950 (Figs 3, 6). Although M.i.27 was excluded as a relevant strain

showing increased genetic requirements of these three metabolic systems, the increase

in genetic requirements for infection *in vivo* suggests a positive relationship between

hypoxic growth *in vitro* and bacterial growth *in vivo*. In contrast, M.i.198 already

showed increased genetic requirements in these three metabolic systems under aerobic

conditions *in vitro*, so the genes associated with these three systems were not identified

as having fewer Tn insertion reads during infection *in vivo*. Genes only identified in

M.i.198 and required for hypoxic pellicle formation in ATCC13950 included genes

associated with succinate synthesis (2-oxoglutarate oxidoreductase subunit KorA) or

encoding the HTH-type transcriptional repressor KstR and menaquinone reductase

(Supplementary Table 11). As increased genetic requirements in 3–5 clinical strains

(excluding M.i.198) compared with ATCC13950, an increased requirement for the RNA

methyltransferase OCU_RS41540 was identified in M.i.198 during infection *in vivo*.

These *in vivo* TnSeq data suggest that, despite the differences in the profiles of the

genes required for *in vivo* infection between strains, increase of genetic requirements for

hypoxic growth in part contribute to the pathogenesis *in vivo*.

We have performed a statistical enrichment analysis of gene sets by GSEA[15] to

investigate whether the genes required for growth of M.i.27 and M.i.198 in mouse lungs

overlap the gene sets required for hypoxic pellicle formation in ATCC13950 as well as

the gene sets showing increased genetic requirements observed in the clinical MAC-PD

strains listed in Fig 3. Of these gene sets including a total of 180 genes, 54% (98 genes)

and 40% (73 genes) were listed as enriched in genes required for mouse infection in

M.i.27 and M.i.198, respectively ($P < 0.01$ in M.i.27 and $P = 0.016$ in M.i.198)

(Supplementary Fig. 4, Supplementary Tables 14). The enriched genes both in M.i.27

and M.i.198 included genes encoding fructose -1,6-bisphosphatase (*glpX*), a type VII

secretion protein (*eccC5*), glycine cleavage system aminomethyltransferase (*gcvT*),

some phosphatase (OCU_RS30185) and a hypothetical protein (OCU_RS47290) (Fig.

6). The enriched genes only in M.i.27 included genes of phosphoenolpyruvate

carboxylinase (*pckA*), cysteine desulfurase (*csd*) and the ESX-5 type VII secretion

component serine protease (*mycP5*). The enriched genes only in M.i.198 included genes

encoding some phosphatase (OCU_RS35975) and class I SAM-dependent RNA

methyltransferase (OCU_RS41540). Similarly, 40-50% of the genes required for

ATCC13950's hypoxic pellicle formation and strain-dependent/accessory essential

genes are significantly enriched individually with the genes required for *in vivo* bacterial

growth (Supplementary Figs. 5, 6, Supplementary Tables 15, 16). These data suggest the

sharing of the hypoxia-adaptation pathways between *in vitro* hypoxic pellicle formation

and *in vivo* bacterial growth.

Preferential hypoxic adaptation of clinical MAC-PD strains evaluated with

bacterial growth kinetics

The metabolic remodeling, such as the increased genetic requirements of

gluconeogenesis and the type VII secretion system, and the overlap of the genes

required for mouse lung infection and those required for hypoxic pellicle formation

involved by these metabolic pathways suggest the preferential adaptation of the clinical

MAC-PD strains to hypoxic conditions. To confirm this, we investigated the growth

kinetics of each strain under aerobic and hypoxic conditions. In terms of growth kinetics,

most of the clinical MAC-PD strains (except M001) entered logarithmic (log) phase

earlier than ATCC13950 under 5% oxygen (Fig. 7a, 7b). However, under aerobic

conditions, the clinical MAC-PD strains entered log phase at a similar time to

ATCC13950. The growth rate at mid-log point (midpoint) was significantly reduced

under hypoxic conditions in a total of 5 clinical MAC-PD strains (M.i.198, M.i.27,

M003, M019 and M021) compared to aerobic conditions (Fig. 7a, 7c). The growth rate

at midpoint under hypoxic conditions was significantly lower in these 5 clinical

MAC-PD strains than in ATCC13950. By contrast, the growth rate at midpoint was

significantly increased under hypoxic conditions compared to aerobic conditions in

ATCC13950. And there was no significant change in the growth rate between aerobic

and hypoxic conditions in the rest 3 clinical MAC-PD strains (M018, M001 and

MOTT64). Despite a longer lag phase compared to other strains under hypoxic

conditions observed in ATCC13950 and M018, additional two strains (M005, M016)

phylogenetically more closely related to ATCC13950 showed different growth pattern

under hypoxia; M005 reached midpoint later than ATCC13950, by contrast M016

reached midpoint three-quarters earlier than ATCC13950 (Supplementary Fig. 7). These

data suggest that genotypes seem to have some impact on bacterial growth pattern under

hypoxia; however, since there was a significant difference in the timing of hypoxic

adaptation among ATCC13950 and its neighbor strains, bacterial growth pattern under

hypoxia seems to be determined by multiple factors other than genotypes such as

strain-specific profiles of genetic requirements and unproven regulatory systems. In

summary, the data of more rapid adaptation to hypoxia followed by reduced growth rate

once adapted indicate the greater metabolic advantage on continuous logarithmic

bacterial growth under hypoxic conditions in the clinical MAC-PD strains than in

ATCC13950.

Effects of knockdown of universal and accessory/strain-dependent essential or

growth-defect-associate genes in clinical MAC-PD strains

To facilitate the functional analysis of the TnSeq-hit genes, it is necessary to construct

the genetically-engineered strains. Still now, there are very few reports of gene

engineering of MAC strains and none of them has been accompanied by the follow-up

studies[23-28]. Moreover, no attempts have been reported to construct the knockdown

strains in MAC strains. With an intention to evaluate the effect of suppressing TnSeq-hit

genes on bacterial growth, we used the pRH2052/pRH2521 clustered regularly

interspaced short palindromic repeats interference (CRISPR-i) plasmids to construct

knockdown strains where the expression of the target gene is suppressed[29-31]. Of the

nine *M. intracellulare* strains analyzed in this study, seven could be transformed with

the vectors for single guide RNA (sgRNA) expression and dCas9 expression. However,

the other two strains, M001 and MOTT64, could not be transformed with these

CRISPR-i vectors. To confirm the gene essentiality detected with the HMM analysis,

we evaluated the consequent growth inhibition in the knockdown strains of

representative universal essential or growth-defect-associated genes, including *glcB*,

inhA, *gyrB*, and *embB*. The knockdown strains showed growth suppression with

comparative growth rates of knockdown strains to vector control strains that do not

express single sgRNA reaching 10^{-1} to 10^{-4} (Fig. 8a).

As representative accessory and strain-dependent essential or

growth-defect-associated genes, we evaluated the 9 genes showing reduced Tn insertion

reads in the 3–5 clinical MAC-PD strains as listed in Fig. 3, as well as *glpX*, a gene of

gluconeogenesis where four MAC-PD strains showed reduced Tn insertion reads in a

resampling analysis (Fig. 5b). The growth was suppressed in the *pckA*-knockdown

strains in M.i.198 and M003, in the *glpX*-knockdown strain in M003, and in the cysteine

desulfurase gene (*csd*)-knockdown strains in M.i.198 and M003, with comparative

growth rates of knockdown strains to vector control strains reaching 10^{-2} to 10^{-4} (Fig.

8b). The growth tended to be suppressed in the type VII secretion protein gene

(*eccC5*)-knockdown strain in M003 and in the type VII secretion-associated serine

protease mycosin gene (*mycP5*)-knockdown strain in M003, with comparative growth

rates of knockdown strains to vector control strains reaching less than 10^{-1}. By contrast,

the growth was not suppressed in these knockdown strains in ATCC13950, with

comparative growth rates of knockdown strains to vector control strains not reaching

10^{-1}, which was consistent to the inessentiality of these genes under aerobic conditions

in ATCC13950. As for M.i.198 and M003, the consistency between TnSeq data and

CRISPR-i data was supported by a correlation between the growth suppression of the

knockdown strains and the fitness costs evaluated by a resampling analysis

(Supplementary Table 17).

In contrast to the cases of M.i.198 and M003, the growth of knockdown strains

of the accessory and strain-dependent essential genes was not effectively suppressed in

the other MAC-PD strains despite being positively detected in TnSeq (Fig. 8b,

Supplementary Fig. 8). Although the gene expression levels were not suppressed to <

1% as reported in *M. tuberculosis* (Mtb)[32], the suppression of gene expression was

confirmed to approximately 20-70% in the knockdown strains of these accessory and

strain-dependent essential or growth-defect-associate genes, as well as universal

essential or growth-defect-associated genes such as *glcB*, *inhA*, *gyrB*, and *embB*

(Supplementary Fig. 9). The possible reason may be explained by the recently revealed

bypass mechanism of gene essentiality in strain-dependent/accessory essential or

growth-defect-associated genes, where gene essentiality can be bypassed by several

mechanisms including the composition of the accessory genome and pathway

rewiring[14].

To be summarized, we could validate the universal essential or

growth-defect-associated genes by the CRISPR-i system in the *M. intracellulare* strains

used in this study. And, although not successful with all strains, we could validate the

accessory and strain-dependent essential or growth-defect-associated genes, especially

genes of gluconeogenesis, type VII secretion components and iron-sulfur cluster

assembly in several MAC-PD strains like M.i.198 and M003.

Discussion

In this study, using TnSeq of nine genetically diverse *M. intracellulare* strains, we have

identified 131 shared essential and growth-defect-associated genes with an HMM

analysis. The genes identified are promising drug targets, representing the genomic

diversity of the clinical MAC-PD strains and reflecting the actual clinical situation. The

strategy of narrowing the essential genes of bacteria is a useful approach for refining the

database of drug targets. We have also demonstrated an overview of the differences in

genetic requirements for hypoxic growth between the ATCC type strain and the clinical

M. intracellulare strains responsible for MAC-PD, with the evidence of earlier entry to

log phase followed by slower growth rate observed in most of these clinical strains

compared to ATCC13950. This is the first study to demonstrate the positive relationship

between the profiles of increased genetic requirements for hypoxic metabolic

remodeling and the phenotypes advantageous for bacterial survival under hypoxic

conditions.

In this study, 121 genes were detected as significantly reduced Tn insertion in the

clinical MAC-PD strains compared to ATCC13950, which was higher than the case of

Mtb showing 10–50 genes with altered Tn insertion reads detected by a resampling

analysis[8]. This may reflect the genomic diversity of *M. intracellulare* compared with

that of Mtb. The proportion of accessory genes is larger in NTM than in Mtb[33]. In a

clinical trial of phage therapy for *M. abscessus* infection, the number of accessory genes

was shown to decrease with the loss of virulence in *M. abscessus* during the course of

therapy[34], suggesting some role for accessory genes for its clinical pathogenesis. Our

data of the overlap of the accessory essential or growth-defect-associated genes such as

pckA and *mycP5* (Supplementary Table 6) with genes required for mouse lung infection

(Fig. 6b, Supplementary Fig. 3) supports the idea that accessory genes (non-core genes)

may encode functions required for bacterial survival, especially under nonoptimal

conditions, such as hypoxia and during the infection of human and animal bodies.

Our TnSeq data are consistent with metabolomic data that have shown the

importance of the gluconeogenic carbon flow of TCA cycle intermediates for bacterial

growth under hypoxic conditions in terms of the essentiality of PckA and the reverse

TCA cycle[35–38]. Gluconeogenesis is essential for bacterial growth when fatty acids are

used as a major nutrient[37]. PckA catalyzes the conversion of the metabolites derived

from the TCA cycle to the metabolites of gluconeogenesis[36]. The reaction catalyzed by

PckA is essential for bacterial growth when the available nutrients are predominantly

fatty acids[39,40]. The main nutrients for mycobacteria inside hypoxic granulomas in an

infected human or mouse body are lipids, such as fatty acids and cholesterol, rather than

carbohydrates[37,41]. In the present study, *M. intracellulare* grew under 5% oxygen,

whereas Mtb stops growing under 1% oxygen and enters a dormant state[36]. The

depletion of phosphoenolpyruvate in the *pckA* deletion mutant of Mtb reduced the

phosphoenolpyruvate–carbon flux essential for bacterial growth and drug sensitivity[36].

The present study supports the idea that PckA is essential for the microaerobic growth

of mycobacteria. Our TnSeq data imply the preferential adaptation of the clinical

MAC-PD strains for circumstances *in vivo*, which are characterized by hypoxia.

On the other hand, such metabolic shift towards gluconeogenesis and reverse TCA

cycle can also be observed under reactive nitrogen species and *in vivo* infection as well.

This suggests that the metabolic shift observed under hypoxia is a general stress

response rather than being specific to hypoxia due to the nature of the

provided/available carbon source[42]. Several factors other than hypoxia can be estimated

to be related to the change of the genetic requirements in the clinical MAC-PD strains

such as host-related factors and general stress conditions (i.e. nutrition of abundant fatty

acids and less carbohydrates, low pH circumstances)[37,39-41]. Except for our data of the

profile of genes required for hypoxic pellicle formation in *M. intracellulare*[10], no

fundamental genome-wide data are available on the change of the genetic requirements

in response to each stress factor in non-tuberculous mycobacteria. Further studies are

needed to discriminate which factors are specifically involved in the increased

requirements of each gene. Although of interest, it is beyond the scope of the current

study.

About 10% of the genes required for mouse infection are also required for hypoxic

pellicle formation. The genes showing increased genetic requirements in the clinical

MAC-PD strains, such as *pckA* and genes encoding the ESX-5 type VII secretion

system, including *eccC5* and *mycP5*, also showed increased genetic requirements in

mouse infections. In a TnSeq study of mice infected intravenously with *M. avium*, the

genes of the ESX-5 type VII secretion system were required for infection of the liver

and spleen[43]. The ESX-5 type VII secretion system is involved in the transport of fatty

acids and the virulence of mycobacteria[44,45]. These similarities in the profiles of genetic

requirements prompt us to propose some role of biofilm formation in mycobacterial

pathogenesis in animals and humans. In the present study, we shed light on the impact

of hypoxia on biofilm formation, the acquisition of virulence, and the *in vivo*

pathogenesis of NTM in terms of global genetic requirements. Although the

morphologies of the pellicle and macroscopic lesions (including cavitation and nodular

bronchiectasis) are not similar, monitoring the expression of the genes identified in this

study may provide clues to the mechanism of biofilm formation *in vivo*.

In this study, we revealed the pattern of hypoxic adaptation characteristic to the

clinical MAC-PD strains by the growth curve analysis: an early entry to log phase

followed by a reduced growth rate (Fig. 7). This pattern implicates the continuous

impact on the infected hosts during logarithmic bacterial growth, which may be

involved in the persistent and steadily progressive illness for years in humans[3]. The

advantage of slow growth in mycobacteria may be inferred by the role of a histone-like

nucleoid protein, mycobacterial DNA-binding protein 1 (MDP1), also designated

HupB[46-48]. The expression of MDP1 is enhanced in the stationary phase, resulting in

reduced growth speed and long-term survival with increased resistance to reactive

oxygen species (ROS)[46-48]. The reduced growth rate after entry to log phase may allow

long-term survival of mycobacteria under ROS-rich circumstances inside host cells. The

pattern of hypoxic adaptation as observed in the clinical MAC-PD strains may enhance

pathogenesis by extending the period of the repeated process of bacterial invasion and

replication in infected macrophages.

Contrary to our expectation, not all clinical strains showed rapid adaptation from

aerobic to hypoxic conditions relative to ATCC13950. These data suggest that some, but

not a major proportion of, clinical *M. intracellulare* strains do not have an advantageous

phenotype for continuous logarithmic growth under hypoxia. In Mtb, the adaptation to

hypoxia is considered to be necessary for clinical pathogenesis because the conditions

in vivo, such as inside macrophages or granulomas, are hypoxic[42,49]. However, our strain

M001 entered log phase later than ATCC13950 under 5% oxygen. Our inability to

construct knockdown strains in M001 prevented us from clarifying its clinical

pathogenesis without preferential hypoxic adaptation. However, because we have shown

that most of the *M. intracellulare* strains with different genotypes showed preferential

adaptation to hypoxia in this study, our results largely reflect the clinical situation of the

etiological agents of MAC-PD. A possible strategy for identifying factors that confer

clinical pathogenesis will be to focus on the major clinical strains that show preferential

hypoxic adaptation.

The data of validation of the universal essential or growth-defect associated genes

by the CRISPR-i system was overall acceptable in the clinically MAC-PD strains whose

knockdown strains could be constructed (Fig. 8a). However, we have encountered the

difficulties of validating the TnSeq-hit genes by CRISPR-i experiment, especially

accessory and strain-dependent essential or growth-defect-associated genes in the

MAC-PD strains other than M.i.198 or M003 (Fig. 8b, Supplementary Fig. 8). We found

the lower efficiency of the suppression of gene expression by CRISPR-i in *M.*

intracellulre than in Mtb that is reported to be suppressed to 6% under 10 ng/ml

anhydrotetracycline (aTc) and by < 1% under 100 ng/ml aTc in the knockdown strains,

compared to strains expressing scrambled single guide RNAs (sgRNA)[32]. However,

because the levels of knockdown efficiency were comparable between

strain-dependent/accessory essential genes and universally essential genes

(Supplementary Fig. 9), the low efficiency of knockdown does not seem to explain the

reason of the discrepancy between TnSeq and CRISPR-i results. Although it is difficult

to find the clear definitive reason for this phenomenon only from the current study, the

possible reason may be speculated by the bypass mechanism of gene essentiality which

is characteristically observed in strain-dependent/accessory essential or

growth-defect-associated genes. According to the publication by Rosconi[14] reporting the

'forced-evolution experiments' of 36 clinical *Streptococcus pneumoniae* strains., gene

essentiality can be bypassed by several mechanisms including the composition of the

accessory genome and pathway rewiring, They recovered successfully knockout

mutants from transformation experiments in strain-dependent/accessory essential genes

such as cytidine monophosphate kinase, a folate pathway enzyme formate

tetrahydrofolate ligase and an undecaprenyl phosphate-biosynthesis pathway enzyme

farnesyl-diphosphate synthase. The bypassing of gene essentiality could be suggested

by observing suppressor mutations and synthetic lethality in knockout strains. By

contrast, universal essential genes were reported to fulfill the three categories including

high levels of conservation within and often across species, limited genetic diversity,

and high and stable expression levels. Consequently, universal essential genes were

estimated to be rigid, largely immutable key components to an organism's survival. The

knockout recovery of the universal essential genes was reported to fail as shown by no

colonies or only appearance of merodiploids. Taking into consideration such bypass

mechanism of gene essentiality, the inconsistent effect of gene silencing of

strain-dependent/accessory essential genes on bacterial growth may reflect some kinds

of pathway rewiring that helps the bacteria grow under suppression of the target gene

expression. Nevertheless, our study has opened the avenue for functional analysis of

NTM that leads to the development of novel drugs targeting universal and

accessory/strain-dependent essential metabolic pathways, especially by making use of

M.i.198 and M003 as optimal NTM strains for gene manipulation.

As in the *glpX*-knockdown strains, *csd* knockdown caused growth suppression in

several MAC-PD strains. This type II Csd is involved in the biosynthesis of the

iron–sulfur clusters of the SUF machinery. Compared with the type I cysteine

desulfurase IscS, the SUF machinery is active under oxidative-stress and iron-limiting

conditions[50]. Representatives of iron–sulfur cluster proteins include components of the

electron-transport system, such as NADH dehydrogenase I (Nuo) and succinate

dehydrogenase (Sdh), which are components of respiratory complexes I and II,

respectively. Under hypoxic conditions, the non-proton-translocating type II NADH

dehydrogenase Ndh is upregulated, and cytochrome *bd* oxidase (CydAB) and the

proton-pumping type I NADH dehydrogenase (Nuo) are downregulated[42]. In the reverse

TCA cycle on the same reaction pathway as Sdh, fumarate reductase (Fnr) is also an

iron–sulfur cluster protein that produces succinate, and is a proposed alternative electron

acceptor under hypoxic conditions[38,51]. The increased essentiality of the *csd* gene

observed in the clinical MAC-PD strains seems to reflect the switching of the

electron-transport system to less-energy-efficient respiration to support preferential

hypoxic survival, including *in vivo*.

We tried to use inhibitors of phosphoenolpyruvate carboxykinase to confirm the

increased essentiality of the *pckA* gene. However, we detected no positive effect because

millimolar 3-mercaptopicolinic acid is required to show the effect of inhibition, even in

eukaryotic cells, which have no cell wall[52]. Moreover, 3-alkyl-1,8-dibenzylxanthine[53]

also only effectively inhibits PckA in eukaryotic cells. However, PckA has been

proposed as a drug target for Mtb[54], so our TnSeq data seem to be consistent with

studies directed towards antimycobacterial drug discovery.

We found the difference in capacity of transformation between strains belonging to

the same genomic *M. paraintracellulare-M. indicus pranii* (MP-MIP) subgroup.

Although the direct mechanism determining the competency for foreign DNA has not

been elucidated in *M. intracellulare* and other NTM species, several studies on general

bacteria suggest the difficulties of introducing foreign DNA into clinical strains

compared to the laboratory strains. As suggested in *Staphylococcus aureus*[55], some

clinical strains develop elimination systems of foreign nucleic acids such as a type

III-like restriction endonuclease. As suggested in gran-negative bacteria[56], there may be

some difference in cell surface structures between strains, resulting in the necessity of

polymyxin B nonapeptide targeting cell membrane for transforming clinical strains. The

efficiency of eliminating foreign DNA may be attributed to various kinds of

strain-specific factors including restriction endonuclease, natural CRISPR-interference

system and cell wall structures rather than a simple genotype factor.

There may be a claim that the saturation of Tn insertion in most of each replicate

were 50-60% in our libraries. However, our method of constructing Tn mutant libraries

is in accordance with the previous studies of "high-density" transposon libraries[9]

because the saturation of the Tn mutant libraries by combining replicates are 62-79% as

follows: ATCC13950: 67.6%, M001: 72.9%, M003: 63.0%, M018: 62.4%, M019:

74.5%, M.i.27: 76.6%, M.i.198: 68.0%, MOTT64: 77.6%, M021: 79.9%. That is, we

predicted gene essentiality from the Tn mutant libraries with 62-79% saturation in each

strain. However, there is non-permissive sequence pattern in mariner transposon

mutagenesis and using more than 10 replicates of Tn mutant libraries is reported to be a

more accurate method[57,58]. The issue of high probability of judging nonessential region

as mistakenly as "essential" are mainly found in small ORFs (2-9 TA sits) rather than

larger ORFs (TA sites $> = 10$))[57]. There are 4136 (6.43% of total ORFs) nonpermissive

TA sites in ATCC13950, which is less than in *M. tuberculosis* H37Rv (9% of total

ORFs)[57] and in *M. abscessus* ATCC19977 (8.1% of total ORFs)[58]. As for small ORFs

(2-9 TA sites), there are nonpermissive TA sites in 41 genes (ORFs) of common

essential or growth-defect- associated (1.35% of a total of 3021 larger ORFs in

ATCC13950). With respect to the covering of non-permissive sites, our TnSeq data may

need to be improved to classify quite accurately the gene essentiality particularly on the

nonstructural small regulatory genes including small RNAs, although the study by

DeJesus shows the comparable number of essential genes in large genes possessing

more than 10 TA sites between 2 and 14 TnSeq datasets, most of which seem to be

structural genes[57].

In summary, we have identified 131 universal essential and

growth-defect-associated genes that cover the genomic diversity of *M. intracellulare*

strains responsible for MAC-PD, which narrowed down the drug targets against

MAC-PD. Furthermore, we have demonstrated preferential adaptation to hypoxia in

clinical NTM strains compared with the type strain in terms of the increased genetic

requirements for hypoxic growth, such as those encoding phosphoenolpyruvate

carboxykinase, the ESX-5 type VII secretion system, and cysteine desulfurase. These

genes were shown to have a significant impact on infection *in vivo* by increasing their

genetic requirements. Our data highlight the importance of using clinical strains and

hypoxic conditions in experimental research, and provide fundamental resources for

developing novel drugs directed towards nontuberculous mycobacterial strains that

cause severe clinical diseases.

Methods

Bacterial strains and phages used to prepare Tn mutants

A total of nine strains were used in this study, including seven clinical strains isolated

from patients with MAC-PD in Japan (M.i.198, M.i.27, M018, M003, M019, M021,

and M001)[22], *M. intracellulare* genovar *paraintracellulare* type strain MOTT64 isolated

from MAC-PD in South Korea[59], and the *M. intracellulare* type strain ATCC13950 (Fig.

1). The Tn mutant libraries were constructed using glucose as a carbon source under

aerobic conditions according to the transduction method used for ATCC13950[10], except

that a higher concentration of kanamycin (100 μg/ml) was used to select the transposon

mutants than was used for ATCC13950 (25 μg/ml). In brief, a hundred milliliter of the

bacteria were cultured aerobically in Middlebrook 7H9 medium (Difco Laboratories,

Detroit, MI, USA) containing 0.2% glycerol, 0.1% Tween 80 (MP Biochemicals,

Illkrich, France), and 10% Middlebrook albumin–dextrose–catalase (7H9/ADC/Tween

80) until mid-log phase (optical density at a wavelength of 600 nm [OD_{600}] of

approximately 0.4–0.6). After harvesting the bacteria, 10 mL of high-titer

mycobacteriophage phAE180[60] was transduced into the bacteria 1-day at 37°C for

inserting Tn and plated on Middlebrook 7H10 solid medium supplemented with 10%

oleic acid-albumin-dextrose-catalase (Becton Dickinson and Co., Sparks, MD)

(7H10/OADC) agar plates containing kanamycin. After 2 or 3 week-cultivation at 37°C

aerobically, the colonies were harvested *en masse* and used as Tn mutant libraries. The

kanamycin-resistant colonies ($> 1 \times 10^5$) were harvested and evenly resuspended in

7H9/ADC medium containing 20% glycerol, aliquoted, and stored at −80 °C until

further use.

Next-generation sequencing and TnSeq analysis

DNA libraries were prepared as reported previously[61]. The TnSeq libraries were

sequenced with the NextSeq 500 System, 150-bp single-end run (Illumina, San Diego,

CA, USA). The sequence reads were analyzed as previously described[10]. Genetic

requirements were calculated, and gene essentiality was predicted by resampling and

HMM analyses using TRANSIT, respectively[12]. First, an HMM analysis was conducted

to calculate the fluctuation of the number of Tn insertion reads individually for each

strain. HMM method can categorize the gene essentiality throughout the genome

including "Essential", "Growth-defect", "Non-essential" and "Growth-advantage".

"Essential" genes are defined as no insertions in all or most of their TA sites.

"Non-essential" genes are defined as regions that have usual read counts.

"Growth-defect" genes are defined as regions that have unusually low read counts.

"Growth-advantage" genes are defined as regions that have unusually high low read

counts. Universal essential and growth-defect-associated genes were identified as the

genes classified as "Essential" and "Growth-defect" with an HMM analysis in all 9

subject strains (Supplementary Table 3). Next, a resampling analysis was conducted to

calculate the difference of the number of Tn insertion reads between each clinical strain

and ATCC13950 (Supplementary Table 7). By comparing the data obtained by HMM

and resampling analyses, we summarized the difference of genetic requirements

between the clinical MAC-PD strains and ATCC13950. To compare the genetic

requirements of the clinical MAC-PD strains and ATCC13950, the annotation for the

clinical MAC-PD strains was adapted from that of ATCC13950 by adjusting the START

and END coordinates of each open reading frame (ORF) in the clinical MAC-PD strains

according to their alignment with the corresponding ORFs of ATCC13950. On

TRANSIT software, we set the replicate option as Sum to combine read counts. And we

used Beta-Geometric correction (BGC) to normalize the datasets to fit an "ideal"

geometric distribution with a variable probability parameter ρ[12]. The function of

TnSeq-hit genes was categorized according to the Kyoto Encyclopedia of Genes and

Genomes (KEGG) database.

Comparison of growth curves under aerobic and hypoxic conditions

First, we precultured the bacteria in 7H9/ADC broth until the bacteria reached early

log-phase (OD_{600} approximately 0.1–0.3). Next, the bacterial cultures were diluted with

new 7H9/ADC broth (final OD_{600} 0.003) for the aerobic and hypoxic growth assay.

Bacterial cells were grown statically in 96-well plates containing 250 μl of 7H9/ADC

broth in each well under aerobic conditions or 5% oxygen at 37 °C with humidification

(APM-30D incubator, Astec, Tokyo, Japan). The culture was sampled every 1–3 days.

At sampling, the culture was mixed by pipetting 50 times. An aliquot (4 μl) of the

culture and its diluents were streaked onto 7H10/OADC agar and the CFUs counted.

The growth curve was fitted to the logistic 4P model with the JMP software (SAS

Institute Inc., Cary, NC, USA) to calculate the inflection point (midpoint) and growth

rate at midpoint in the log growth phase. Entry into the log phase was evaluated from

the timing of the inflection point on the growth curve.

Animals

Female C57BL/6JJcl mice were purchased from CLEA Japan (Tokyo, Japan).

Six-week-old mice were used for the infection experiment (Supplementary Fig. 3). All

the mice were kept under specific-pathogen-free conditions in the animal facility of

Niigata University Graduate School of Medicine, according to the institutional

guidelines for animal experiments. The mice were housed (four per cage) under a

12-h/12-h light/dark cycle. Food and water were available *ad libitum.* Kanamycin was

added to the drinking water to a final concentration of 100 µg/ml to maintain the mice's

resistance to the Tn mutant strains. Euthanasia was conducted with an intraperitoneal

injection of a combined anesthetic (15 µg of medetomidine, 80 µg of midazolam, and

100 µg of butorphanol), followed by cervical dislocation.

Mouse TnSeq experiment

The protocol was based on our previous mouse lung infection experiment[22]. Twenty

mice each were used for the Tn mutant strains M.i.198 and M.i.27. The mice were

injected intratracheally with 3.25×10^8 and 5.93×10^7 CFUs of the Tn mutant strains,

respectively. The infected lungs were expected to be harvested from five mice at day 1,

week 4, week 8, and week 16 after infection. However, the lungs were harvested from

only four mice infected with the M.i.27 Tn mutant strains in week 8 because we

experienced difficulties with sampling, and the lungs were harvested from only three,

one, and two mice infected with M.i.198 Tn mutant strains in week 8, week 12, and

week 16, respectively, because morbidity was high in the mice infected with M.i.198.

The harvested lungs were homogenized in 4.5 ml of distilled water with the

gentleMACS™ Tissue Dissociator (Miltenyi Biotec, Bergish Gladbach, Germany) and

the homogenates were plated on 7H10/OADC agar plates and cultured for 3 weeks. The

colonies were harvested *en masse* for TnSeq.

Gene set enrichment analysis

To investigate the overlap of the gene sets required for growth in mouse lungs with

those required for hypoxic pellicle formation observed in ATCC13950 and those

showing increased genetic requirements observed in the clinical MAC-PD strains, a

statistical enrichment of gene sets was performed by GSEA 4.3.2[15]. Parameters set for

GSEA were as follows: permutations = 1,000, permutation type: gene_set, enrichment

statistic: weighted, metric for ranking genes: Signal2Noise, max size: 500, min size: 15.

Construction and growth assay of knockdown strains

Knockdown strains were constructed with the CRISPR-dCas9-mediated RNA

interference system, as reported previously[29]. Gene-specific 20-nt sequences were

selected as protospacers for incorporation into the sgRNAs (Supplementary Table 18).

pRH2502 (a vector expressing an inactivated version of the *cas9* DNA sequence) and

pRH2521 (an sgRNA-expressing vector whose expression is regulated by the

TetR-regulated smyc promotor P_{myc1}tetO) were introduced into the bacteria to obtain

colonies of the knockdown strains. For selection, 25 µg/ml kanamycin and 50 µg/ml

hygromycin were used for ATCC13950, and 100 µg/ml kanamycin and 50 µg/ml

hygromycin were used for the clinical strains. The growth assay of the knockdown

strains was performed in 7H9/ADC broth containing the above-mentioned

concentrations of kanamycin and hygromycin under aerobic conditions at 37 °C with

humidification (APM-30D incubator, Astec, Tokyo, Japan). During the assay, the

cultures were supplemented with anhydrotetracycline (aTc) to a final concentration of

50 ng/ml (as a concentration that is not toxic to ATCC13950 and M.i.27) or 200 ng/ml

(for the other clinical strains). aTc was added repeatedly every 48 h to maintain the

induction of dCas9 and sgRNAs in experiments that extended beyond 48 h[29]. An aliquot

(4 μl) of the culture and its diluents were streaked onto 7H10/OADC agar plates

containing kanamycin and hygromycin, and the CFUs were counted. Growth inhibition

was evaluated as the ratio of the CFUs of knockdown strains compared with those of the

vector control strains lacking the guide RNA sequence. To confirm the downregulation

of gene expression in the knockdown strains, quantitative reverse transcription PCR

(qRT-PCR) was performed. Bacteria were cultured in 10 ml of 7H9/ADC/Tween 80

containing kanamycin and hygromycin until mid-log phase (OD_{600} approximately

0.4–0.6). The cultures were supplemented with aTc every 48 h, as in the growth assay

for the knockdown strains. One day after the second addition of aTc, RNA was extracted

by bead-beating six times at 5000 rpm for 30 s with cooling (Micro Smash MS 100R

Cell Disruptor; Tomy Seiko, Tokyo, Japan) and cDNA was synthesized with the

Direct-zol™ RNA Miniprep (Zymo Research, Orange, CA, USA) and ReverTra Ace

qPCR RT Master Mix with gDNA Remover (Toyobo, Osaka, Japan). qRT-PCR was

performed with the CFX Connect Real-time PCR Detection System (Bio-Rad, Hercules,

CA, USA), according to the manufacturer's instructions. The data were normalized to

the expression of *sigA*.

Statistical analyses

The cut-off of fold-change ($\log_2$FC) to compare genetic requirements by TnSeq

resampling analysis was set as adjusted *P*-values less than 0.05. Pearson's correlation

test was performed between the fitness costs ($\log_2$FC calculated by the resampling

assay), efficiency of knockdown (assessed by qRT-PCR) and growth suppression in the

knockdown strains. Statistical analyses of the growth curves were performed using a

4-parameter logistic model (logistic 4P model), Wilcoxon two-sample test, and a

one-way analysis of variance (ANOVA) followed by Steel's multiple comparisons test

using the JMP software program (SAS Institute Inc., Cary, NC, USA). Data are

presented as the mean ± SD. Statistical significance was set as *P* values less than 0.05.

[References]

1.   Adjemian, J., Olivier, K.N., Seitz, A.E., Holland, S.M. & Prevots, D.R.

740   Prevalence of nontuberculous mycobacterial lung disease in U.S. Medicare

741   beneficiaries. *Am J Respir Crit Care Med* **185**, 881-886 (2012).

742   2.   Namkoong, H. *et al.* Epidemiology of pulmonary nontuberculous mycobacterial

743   disease, Japan. *Emerg Infect Dis* **22**, 1116-1117 (2016).

744   3.   Daley, C.L. *et al.* Treatment of nontuberculous mycobacterial pulmonary

745   disease: an official ATS/ERS/ESCMID/IDSA clinical practice guideline. *Eur*

746   *Respir J* **56**, 2000535 (2020).

747   4.   Uchiya, K.I. *et al.* Comparative genome analyses of *Mycobacterium avium*

748   reveal genomic features of its subspecies and strains that cause progression of

749   pulmonary disease. *Sci Rep* **7**, 39750 (2017).

750   5.   Tateishi, Y. *et al.* Comparative genomic analysis of *Mycobacterium*

751   *intracellulare*: implications for clinical taxonomic classification in pulmonary

752   *Mycobacterium avium-intracellulare* complex disease. *BMC Microbiol* **21**, 103

753   (2021).

754   6.   Tortoli, E. *et al.* Genome-based taxonomic revision detects a number of

755   synonymous taxa in the genus *Mycobacterium*. *Infect Genet Evol* **75**, 103983

756   (2019).

7.   van Opijnen, T. & Camilli, A. Transposon insertion sequencing: a new tool for

systems-level analysis of microorganisms. *Nat Rev Microbiol* **11**, 435-442

(2013).

8.   Carey, A.F. *et al.* TnSeq of *Mycobacterium tuberculosis* clinical isolates reveals

strain-specific antibiotic liabilities. *PLoS Pathog* **14**, e1006939 (2018).

9.   Akusobi, C. *et al..* Transposon-sequencing across multiple *Mycobacterium*

abscessus isolates reveals significant functional genomic diversity among strains.

mBio **6**, e0337624 (2025).

10.   Tateishi, Y. *et al.* Genome-wide identification of essential genes in

Mycobacterium intracellulare by transposon sequencing - Implication for

metabolic remodeling. *Sci Rep* **10**, 5449 (2020).

11.   Cuttino, J.T. & Mc, C.A. Pure granulomatous nocardiosis, a new fungus disease

distinguished by intracellular parasitism; a description of a new disease in man

due to a hitherto undescribed organism, *Nocardia intracellularis*, n. sp.,

including a study of the biologic and pathogenic properties of this species. *Am J*

Pathol **25**, 1-47 (1949).

12.   DeJesus, M.A., Ambadipudi, C., Baker, R., Sassetti, C. & Ioerger, T.R.

TRANSIT--A Software Tool for Himar1 TnSeq Analysis. *PLoS Comput Biol* **11**,

e1004401 (2015).

13.   Krieger, I.V. *et al.* Structure-guided discovery of phenyl-diketo acids as potent

inhibitors of *M. tuberculosis* malate synthase. *Chem Biol* **19**, 1556-1567 (2012).

14.   Rosconi, F. *et al.* A bacterial pan-genome makes gene essentiality

strain-dependent and evolvable. *Nat Microbiol* **7**, 1580–1592 (2022).

15.   Subramanian, A. *et al.* Gene set enrichment analysis: A knowledge-based

approach for interpreting genome-wide expression profiles. *Proc Natl Acad Sci*

U S A **102,** 15545–15550 (2005).

16.   Coll, F. *et al.* Genome-wide analysis of multi- and extensively drug-resistant

Mycobacterium tuberculosis. *Nat Genet* **50**, 307-316 (2018).

17.   Nguyen, H.T., Wolff, K.A., Cartabuke, R.H., Ogwang, S. & Nguyen, L. A

lipoprotein modulates activity of the MtrAB two-component system to provide

intrinsic multidrug resistance, cytokinetic control and cell wall homeostasis in

Mycobacterium. *Mol Microbiol* **76**, 348-364 (2010).

18.   Li, S. *et al.* CRISPRi chemical genetics and comparative genomics identify

genes mediating drug potency in *Mycobacterium tuberculosis*. *Nat Microbiol* **7**,

766–779 (2022).

19.   Kirschner, R.A., Jr., Parker, B.C. & Falkinham, J.O., 3rd. Epidemiology of

infection by nontuberculous mycobacteria. *Mycobacterium avium,*

Mycobacterium intracellulare, and *Mycobacterium scrofulaceum* in acid,

brown-water swamps of the southeastern United States and their association

with environmental variables. *Am Rev Respir Dis* **145**, 271-275 (1992).

20. Werner, E. *et al.* Stratified growth in *Pseudomonas aeruginosa* biofilms. *Appl*

Environ Microbiol **70**, 6188-6196 (2004).

21. Via, L.E. *et al.* Tuberculous granulomas are hypoxic in guinea pigs, rabbits, and

nonhuman primates. *Infect Immun* **76**, 2333-2340 (2008).

22. Tateishi, Y. *et al.* Virulence of *Mycobacterium intracellulare* clinical strains in a

mouse model of lung infection - role of neutrophilic inflammation in disease

severity. *BMC Microbiol* **23**, 94 (2023).

23. Kryzywinska, E., Bhatnagar, S., Sweet, L., Chatterjee, D. & Schorey, J.S.

Mycobacterium avium 104 deleted of the methyltransferase D gene by allelic

replacement lacks serotype-specific glycopeptidolipids and shows attenuated

virulence in mice. *Mol Microbiol* **56**, 1262-1273 (2005).

24. Irani, V.R., Lee, S.-H., Eckstein, T.M., Inamine, J.M., Belisle, J.T. & Maslow,

J.N. Utilization of a *ts-sacB* selection system for the generation of a

Mycobacterium avium serovar-8 specific glycopeptidolipid allelic exchange

811    mutant. A*nnals Clin Microbiol Antimicrob* **3**, 18 (2004).

812 25. Marklund, B.I., Speert, D.P. & Stokes, R.W. Gene replacement through

813    homologous recombination in *Mycobacterium intracellulare*. *J Bacteriol* **177**,

814    6100-6105 (1995).

815 26. Marklund, B.I., Mahenthiralingam, E. & Stokes, R.W. Site-directed mutagenesis

816    and virulence assessment of the *katG* gene of *Mycobacterium intracellulare*. *Mol*

817    *Microbiol* **29**, 999-1008 (1998).

818 27. Mahenthiralingam, E., Marklund, B.I., Brooks, L.A., Smith, D.A., Bancroft, G.J.

819    & Stokes, R.W. Site-directed mutagenesis of the 19-kilodalton lipoprotein

820    antigen reveals No essential role for the protein in the growth and virulence of

821    *Mycobacterium intracellulare*. *Infect Immun.* **66**, 3626-3634 (1998).

822 28. Hinds, J. *et al.* Enhanced gene replacement in mycobacteria. *Microbiology* **145**,

823    519-527 (1999).

824 29. Singh, A.K. *et al.* Investigating essential gene function in *Mycobacterium*

825    *tuberculosis* using an efficient CRISPR interference system. *Nucleic Acids Res*

826    **44**, e143 (2016).

827 30. Choudhary, E., Thakur, P., Pareek, M. & Agarwal, N. Gene silencing by CRISPR

828    interference in mycobacteria. *Nat Commun* **6**, 6267 (2015).

31.   Rock, J.M. *et al.* Programmable transcriptional repression in mycobacteria using

an orthogonal CRISPR interference platform. *Nat Microbiol* **2,** 16274 (2017).

32.   McNeil, M.B. & Cook, G.M. Utilization of CRISPR interference to validate

MmpL3 as a drug target in *Mycobacterium tuberculosis*. *Antimicrob Agents*

Chemother **63,** e00629-19 (2019).

33.   Yang, T. *et al.* Pan-genomic study of *Mycobacterium tuberculosis* reflecting the

primary/secondary genes, generality/individuality, and the interconversion

through copy number variations. *Front Microbiol* **9**, 1886 (2018).

34.   Nick, J.A. *et al.* Host and pathogen response to bacteriophage engineered against

Mycobacterium abscessus lung infection. *Cell* **185**, 1860-1874.e12 (2022).

35.   Eoh, H. *et al.* Metabolic anticipation in *Mycobacterium tuberculosis*. *Nat*

Microbiol **2**, 17084 (2017).

36.   Lim, J. *et al.* Phosphoenolpyruvate depletion mediates both growth arrest and

drug tolerance of *Mycobacterium tuberculosis* in hypoxia. *Proc Natl Acad Sci U*

S A **118,** e2105800118 (2021).

37.   Marrero, J., Rhee, K.Y., Schnappinger, D., Pethe, K. & Ehrt, S. Gluconeogenic

carbon flow of tricarboxylic acid cycle intermediates is critical for

Mycobacterium tuberculosis to establish and maintain infection. *Proc Natl Acad*

Sci U S A **107**, 9819-9824 (2010).

38.   Watanabe, S. *et al.* Fumarate reductase activity maintains an energized

membrane in anaerobic *Mycobacterium tuberculosis*. *PLoS Pathog* **7**, e1002287

(2011).

39.   Gouzy, A., Healy, C., Black, K.A., Rhee, K.Y. & Ehrt, S. Growth of

Mycobacterium tuberculosis at acidic pH depends on lipid assimilation and is

accompanied by reduced GAPDH activity. *Proc Natl Acad Sci U S A* **118,**

e2024571118 (2021).

40.   Quinonez, C.G. *et al.* The role of fatty acid metabolism in drug tolerance of

Mycobacterium tuberculosis. *mBio* **13**, e0355921 (2022).

41.   Pandey, A.K. & Sassetti, C.M. Mycobacterial persistence requires the utilization

of host cholesterol. *Proc Natl Acad Sci U S A* **105**, 4376-4380 (2008).

42.   Prosser, G. *et al.* The bacillary and macrophage response to hypoxia in

tuberculosis and the consequences for T cell antigen recognition. *Microbes*

Infect **19**, 177-192 (2017).

43.   Dragset, M.S. *et al.* Global assessment of *Mycobacterium avium* subsp.

hominissuis genetic requirement for growth and virulence. *mSystems* **4,**

e00402-19 (2019).

44.   Ates, L.S. *et al.* Essential role of the ESX-5 secretion system in outer membrane

permeability of pathogenic mycobacteria. *PLoS Genet* **11**, e1005190 (2015).

45.   Ates, L.S. *et al.* The ESX-5 system of pathogenic mycobacteria is nvolved in

capsule integrity and virulence through its substrate PPE10. *PLoS Pathog* **12**,

e1005696 (2016).

46.   Savitskaya, A. *et al.* C-terminal intrinsically disordered region-dependent

organization of the mycobacterial genome by a histonelike protein. *Sci Rep* 8,

1–15 (2018).

47.   Enany, S. *et al.* Mycobacterial DNA-binding protein 1 is critical for long term

survival of *Mycobacterium smegmatis* and simultaneously coordinates cellular

functions. *Sci Rep* 7, 1–11 (2017).

48.   Shaban, A.K. *et al.* Mycobacterial DNA□binding protein 1 is critical for BCG

survival in stressful environments and simultaneously regulates gene expression.

Sci Rep 13, 14157 (2023).

49.   Rustad, T.R., Sherrid, A.M., Minch, K.J. & Sherman, D.R. Hypoxia: a window

into *Mycobacterium tuberculosis* latency. *Cell Microbiol* **11**, 1151-1159 (2009).

50.   Takahashi, Y. & Tokumoto, U. A third bacterial system for the assembly of

iron-sulfur clusters with homologs in archaea and plastids. *J Biol Chem* **277**,

883  28380-28383 (2002).

884 51. Cecchini, G., Schröder, I., Gunsalus, R.P. & Maklashina, E. Succinate

885  dehydrogenase and fumarate reductase from *Escherichia coli*. *Biochim Biophys*

886  *Acta* **1553**, 140-157 (2002).

887 52. Brearley, M.C., Daniel, Z., Loughna, P.T., Parr, T. & Brameld, J.M. The

888  phosphoenolpyruvate carboxykinase (PEPCK) inhibitor, 3-mercaptopicolinic

889  acid (3-MPA), induces myogenic differentiation in C2C12 cells. *Sci Rep* **10**,

890  22177 (2020).

891 53. Montal, E.D. *et al.* Inhibition of phosphoenolpyruvate carboxykinase blocks

892  lactate utilization and impairs tumor growth in colorectal cancer. *Cancer Metab*

893  **7**, 8 (2019).

894 54. McLeod, M.J., Krismanich, A.P., Assoud, A., Dmitrienko, G.I. & Holyoak, T.

895  Characterization of 3-[(carboxymethyl)thio]picolinic acid: a novel inhibitor of

896  phosphoenolpyruvate carboxykinase. *Biochemistry* **58**, 3918-3926 (2019).

897 55. Corvaglia, A.R., François, P., Hernandez, D., Perron, K., Linder, P. & Schrenzel,

898  J. A type III-like restriction endonuclease functions as a major barrier to

899  horizontal gene transfer in clinical *Staphylococcus aureus* strains. *Proc Natl*

900  *Acad Sci U S A* **107,** 11954-11958 (2010).

56. Qin, J., Hong, Y., Pullela, K., Morona, R., Henderson, I.R. & Totsika, M. A

method for increasing electroporation competence of Gram-negative clinical

isolates by polymyxin B nonapeptide. *Sci Rep* **12,**:11629 (2022).

57. DeJesus, M.A. *et al.* Comprehensive essentiality analysis of the *Mycobacterium*

tuberculosis genome via saturating transposon mutagenesis. *mBio* **8,** e02133-16

(2017).

58. Rifat, D., Chen L., Kreiswirth, B.N. & Nuermberger, E.L.. Genome-wide

essentiality analysis of *Mycobacterium abscessus* by aaturated rransposon

mutagenesis and deep sequencing. *mBio* **12,** e0104921 (2021).

59. Kim, B.J. *et al.* Complete genome sequence of *Mycobacterium intracellulare*

clinical strain MOTT-64, belonging to the INT1 genotype. *J Bacteriol* **194**, 3268

(2012).

60. Kriakov, J., Lee, S. & Jacobs, W.R., Jr. Identification of a regulated alkaline

phosphatase, a cell surface-associated lipoprotein, in *Mycobacterium smegmatis*.

J Bacteriol **185**, 4983-4991 (2003).

61. Minato, Y. *et al.* Genomewide assessment of *Mycobacterium tuberculosis*

conditionally essential metabolic pathways. *mSystems* **4,** e00070-19 (2019).

62. Shannon, P. *et al.* Cytoscape: a software environment for integrated models of

biomolecular interaction networks. *Genome Res* **13**, 2498-2504 (2003).

Acknowledgments

The authors thank Dr. Torin Weisbrod and Dr. William R. Jacobs, Jr. (Albert Einstein

College of Medicine, Bronx, NY, USA) for kindly providing mycobacterial phage

phAE180, and Dr. Norikazu Hara and Dr. Takeshi Ikeuchi (Department of Molecular

Genetics, Bioresource Science, Brain Center for Bioresources, Brain Research Institute,

Niigata University) for next-generation sequencing with the NextSeq 500 System. The

authors thank Mr. Naomi Yoshihara and Ms. Chisa Kumagai for constructing Tn mutant

library strains and hypoxic growth experiments, respectively. This work was supported

by Grants-in-Aid for Scientific Research (grant numbers 20KK0216, 21K19637, and

22H03115 to Yoshitaka Tateishi) from the Ministry of Health, Labour, and Welfare, of

Japan This work was also supported by grants from the Japanese Ministry of Education,

Culture, Sports, Science, and Technology, the Ministry of Health, and the Research

Program on Emerging Infectious Disease from the Japan Agency for Medical Research

and Development (23fk0108673h0501, AMED-CREST 25gm1610009h0004, and

JP223fa627005).

Author contributions

Y.T. conceived, designed and performed the experiments. Y.O., A.N., and S.M.

contributed reagents or materials. Y. Morishige., Y. Minato, and A.D.B. contributed

analytical tools. Y. Morishige., Y. Minato, A.D.B., and S.M. provided critical

suggestions throughout the preparation of the manuscript. Y.T. wrote the manuscript. All

authors read and approved the final manuscript.

Availability of data and materials

The datasets used or analyzed in this study are available from the corresponding author

upon reasonable request.

Declarations

Ethics approval and consent to participate

All experimental protocols were approved by the Institutional Review Board of Niigata

University (approval numbers: 2019-0020 for the use of human tissue samples and

SA00506 for experiments on live vertebrates). All experiments on humans and human

samples were conducted in accordance with relevant guidelines and regulations.

Informed consent was obtained from all patients. All experiments on animals were

conducted in accordance with the relevant guidelines and regulations, and the study

complied with the ARRIVE guidelines.

Consent for publication

According to the protocols approved by the Institutional Review Board of Niigata

University mentioned above, consent for publication was obtained from the patients

enrolled in this study.

Competing interests

The authors declare that they have no competing interests.

Fig. 1 Phylogenetic tree of the *M. intracellulare* strains and strategy of the TnSeq

analyses in this study. a Phylogenetic tree of the *M. intracellulare* strains used in this

study. The tree was generated based on average nucleotide identity (ANI) with the

neighbor-joining method. TMI: typical *M. intracellulare*; MP-MIP: *M.*

paraintracellulare–M. indicus pranii. The detail of phylogenetic tree is shown in

Supplementary Fig. 1. **b** Strategy of the procedure of TnSeq analyses.

Fig. 2 Identification of the essential and growth-defect-associated genes across all

nine *M. intracellulare* strains used in this study. a Functional categories of 131 genes

identified as universal essential or growth-defect-associated with an HMM analysis. The

genes were categorized according to the Kyoto Encyclopedia of Genes and Genomes

(KEGG) database. **b** Universal essential or growth-defect-associated genes

corresponding to the genes of existing antituberculous drug targets. **c** The number of the

essential or growth-defect-associated genes in the accessory genes, and the number of

strain-dependent essential or growth-defect-associated genes in the core genes.

Fig. 3 Detection of genes showing increased or reduced genetic requirements in the

clinical *M. intracellulare* strains. Left panel shows the genes identified as having

fewer Tn insertion reads than the type strain ATCC13950. The fold changes in the

number of Tn insertion reads calculated by a resampling analysis are represented by the

color scale. Red squares indicate genes required for hypoxic pellicle formation in

ATCC13950[10]. *Genes identified with a combination of resampling and HMM analyses.

†Genes identified only with resampling analysis. Right panel shows the genes identified

as having more Tn insertion reads than the type strain ATCC13950. NH: no homolog

with ATCC13950, NS: no significant increase in Tn insertion reads by a resampling

analysis (adjusted $P < 0.05$).

Fig. 4 Overview of the differences in genetic requirements between the clinical *M.*

***intracellulare* strains and ATCC13950, drawn with Cytoscape[62].** Each central node

represents the functional category of genes, assigned with a KEGG pathway analysis.

Each peripheral node represents the genes showing significant changes in the number of

Tn insertion reads. The size of each node represents the number of strains identified.

Each edge represents a strain identified as showing significant changes in the number of

Tn insertion reads. The thickness of each edge represents the adjusted *P* value. The

color scale of each edge represents the fold change in the number of Tn insertion reads

calculated by a resampling analysis. The graphical organization was referenced to the

previous publication by Carey, *et al*[8].

1001

Fig. 5 Preferential genetic requirements for gluconeogenesis in the clinical *M.*

***intracellulare* strains inferred from the TnSeq results. a** Carbohydrate pathway

showing the changes in Tn insertion reads. Genes identified with the resampling

analysis are framed by squares. The essentiality of gluconeogenesis-related genes (*pckA,*

glpX) was higher and that of glycolysis-related genes (*aceE, lpdC*) was lower in the

clinical *M. intracellulare* strains compared with those in ATCC13950. **b** Data on genetic

requirements calculated with a resampling analysis in the clinical *M. intracellulare*

strains compared with ATCC13950. Red indicates a reduction in Tn insertion reads in

the clinical *M. intracellulare* strains. Blue indicates an increase of Tn insertion reads in

the clinical *M. intracellulare* strains. Asterisks indicate statistical significance of $\log_2$

fold changes ($\log_2$FCs). The graphical organization was referenced to the previous

publication by Carey, *et al*[8].

1014

Fig. 6 Detection of genes required for mouse lung infection in the clinical *M.*

***intracellulare* strains. a** Consistency between the genes required for infection in mouse

lungs and the 175 genes required for hypoxic pellicle formation in ATCC13950. **b**

Changes in the requirements for genes with increased genetic requirements observed in

the clinical *M. intracellulare* strains after the infection of mouse lungs. Highlighted

genes are also required for hypoxic pellicle formation by ATCC13950.

Genetic requirements of the strain itself: genetic requirements compared with that of

ATCC13950. Genetic requirements in mouse lungs: changes in the genetic requirements

in the infected mouse lungs compared with those before infection. Gene set enrichment

analysis: The genes listed as core enrichment are shown as "Yes" and the genes not

listed as core enrichment are shown as "No."

1026

Fig. 7 Comparison of the timing of entry into logarithmic growth and logarithmic

growth rate by the clinical *M. intracellulare* strains and ATCC13950. a

Representative data on the growth curves of each strain under aerobic and 5% oxygen

conditions. The assay was performed three times in 96-well plates containing 250 µl of

broth medium per well, in triplicate. Data are represented as CFUs in 4 µl sample at

each timepoint. Data on the growth curves are the means of triplicate experiments. Data

from one experiment representative of three independent experiments ($N = 3$) are shown.

b Time at the inflection point (midpoint) on the sigmoid growth curve. *significantly

earlier than an aerobic culture of ATCC13950; #significantly earlier than a hypoxic

culture of ATCC13950; †significantly later than an aerobic culture of ATCC13950;

‡significantly later than a hypoxic culture of ATCC13950. **c** Growth rate at midpoint of

the growth curve in each strain. #significantly slower than a hypoxic culture of

ATCC13950; †significantly slower than an aerobic culture of the corresponding strain;

‡significantly faster than an aerobic culture of ATCC13950. Open bars: aerobic; closed

bars: 5% O_2. Data are shown as the means ± SD of triplicate experiments. Data from

one experiment representative of three independent experiments ($N = 3$) are shown.

1043

Fig. 8 Evaluation of the effect of the suppression of TnSeq-hit genes on the

bacterial growth using the CRISPR-i system. Open bar: day 3; closed bar: day 7. **a**

Comparative growth rates of the knockdown strains relative to those of the vector

control strains in the representative universal essential or growth-defect-associated

genes: *glcB, inhA*, *gyrB* and *embB*. **b** Comparative growth rates of the knockdown

strains relative to those of the vector control strains in the representative accessory and

strain-dependent essential or growth-defect-associated genes: *pckA*, *glpX*, *csd* and

ESX-5 type VII secretion components. Data are shown as the means ± SD of triplicate

experiments. Data from one experiment representative of two independent experiments

($N = 2$) are shown.

1054

a

b

1. Detection of shared essential (ES) or growth-defect –associated (GD) genes among study strains by the Hidden Markov Model (HMM) analysis

2. Detection of altered genetic requirements in clinical MAC-PD strains compared to ATCC13950 by the resampling analysis

3. Detection of genes required for mouse lung infection

a

Not included in pathway (3 genes)

Signaling and cellular processes (10 genes)

Genetic information processing (39 genes)

Metabolism (79 genes)

Metabolism: glyoxylate, purine/pyrimidine, amino acid, lipid, arabinogalactan/peptidoglycan, porphyrin

Genetic information processing: DNA replication, transcription, amino acid tRNA ligases, ribosomal proteins, general Sec proteins

Signaling and cellular processes: ABC transporter, type VII secretion system, iron-sulfur cluster assembly

Not Included in Pathway: hypothetical proteins

b

Gene locus	gene	function	target drug
OCU_RS25020, OCU_RS25025	*gyrB, gyrA*	DNA topoisomerase subunit B, intein-containing DNA gyrase subunit A	fluoroquinolones
OCU_RS26150	*embB*	arabinosyltransferase	ethambutol
OCU_RS40185	*inhA*	enoyl[acyl-carrier-protein] reductase FabI	isoniazid/ethionamid
OCU_RS45570	*alr*	alanine racemase	cycloserine
OCU_RS46225	*rpoC*	DNA-directed RNA polymerase subunit beta'	rifampicin
OCU_RS38100	*glcB*	malate synthase G	malate synthase inhibitors
OCU_RS32840, OCU_RS32845, OCU_RS32850	*atpA, atpG, atpB*	F_0F_1 ATP synthase subunit alpha, gamma, beta	bedaquiline

c

- Universal essential or growth-defect-associated genes
- Strain-dependent essential or growth-defect-associated genes
- Accessory essential or growth-defect-associated genes
- Non-essential genes

Fig. 2

Tn decrease in clinical strains

A total of 121 genes were detected as significantly reduced Tn in clinical MAC-PD strains compared to ATCC13950. Of those, 9 genes were commonly hit in 3 to 5 clinical strains.

OCU_RS30185 Ppx/GppA family phosphatase

OCU_RS35795 Bifunctional RNaseH/acid phosphatase

OCU_RS35955 Glycine cleavage system amino-methyltransferase *gcvT*

OCU_RS38275 Type VII secretion-associated serine protease mycosin *mycP5*

OCU_RS38345 Type VII secretion protein *eccC5*

OCU_RS40305 Cysteine desulfurase *csd*

OCU_RS41540 Class I SAM-dependent RNA methyltransferase

OCU_RS47290 Hypothetical protein

OCU_RS48660 Phosphoenolpyruvate carboxykinase *pckA*

log₂ (fold-change)

-6.42   0   +11.14

Fig. 3

Tn increase in clinical strains

A total of 162 genes were detected as significantly increased Tn insertion compared to ATCC13950. Of those, 12 genes were commonly hit in more than 7 clinical MAC-PD strains.

OCU_RS25945 Superoxide dismutase [Mn]

OCU_RS28435 RND family transporter

OCU_RS28440 TetR/AcrR transcriptional regulator

OCU_RS30230 Response regulator transcription factor

OCU_RS34390 CoA transferase

OCU_RS34400 amidohydrolase

OCU_RS34425 amidohydrolase

OCU_RS34445 amidohydrolase

OCU_RS35720 Pyruvate dehydrogenase (acetyl-transferring)

OCU_RS39950 daunorubicin resistance protein DrrA family ABC transporter ATP-binding protein

OCU_RS39955 ABC transporter

OCU_RS44685 MtrAB system accessory protein *lpqB*

Fig. 4

Fig 5.

a

M.i.27
629 required genes in mouse lungs

M.i.198
512 required genes in mouse lungs

72 genes (11.4%) corresponding to the 175 genes required for hypoxic pellicle formation

50 genes (9.76%) corresponding to the 175 genes required for hypoxic pellicle formation

72 genes

50 genes

31 genes

41 genes

9 genes

41 genes commonly detected in M.i.27 and M.i.198 as corresponding genes required for hypoxic pellicle formation

b

Locus_tag	Gene annotation	Genetic require-ments of strain itself	M.i.27 Genetic require-ments in mouse lungs	Gene set enrich-ment analysis	Genetic require-ments of strain itself	M.i.198 Genetic require-ments in mouse lungs	Gene set enrich-ment analysis
OCU_RS48660	Phoshoenolpyruvate carboxykinase PckA	NC	UP	Yes	UP	NC	No
OCU_RS30540	Fructose-1,6-bisphosphatase GlpX	NC	UP	Yes	NC	NC	Yes
OCU_RS40305	Cysteine desulfurase Csd	NC	UP	Yes	UP	NC	No
OCU_RS38345	TypeVII secretion protein EccC5	NC	UP	Yes	UP	NC	Yes
OCU_RS38275	TypeVII serine protease MycP5	NC	UP	Yes	UP	NC	No
OCU_RS35955	Glycine cleavage system aminomethyltransferase GcvT	NC	UP	Yes	UP	NC	Yes
OCU_RS30185	Ppx/GppA family phosphatase	NC	NC	Yes	NC	NC	Yes
OCU_RS35795	Bifunctional RNaseH/acid phosphatase	NC	NC	No	NC	NC	Yes
OCU_RS41540	Class I SAM-dependent RNA methyltransferase	UP	NC	No	NC	UP	Yes
OCU_RS47290	Hypothetical protein	UP	NC	Yes	NC	NC	Yes

Fig. 6

Fig. 7

Fig. 8