Functional genomics reveals the mechanism of hypoxic adaptation in nontuberculous

mycobacteria

Yoshitaka Tateishi[1*], Yuriko Ozeki[1], Akihito Nishiyama[1], Yuta Morishige[2], Yusuke

Minato[3,4], Anthony D. Baughn[4], Sohkichi Matsumoto[1,5]

1. Department of Bacteriology, Graduate School of Medicine and Dental Sciences,

Niigata University, 1-757, Asahimachi-dori, Chuo-ku, Niigata 951-8510, Japan.

2. Department of Mycobacterium Reference and Research, Research Institute of

Tuberculosis, Japan Anti-Tuberculosis Association, Kiyose, Tokyo, 2048533, Japan.

3. Department of Microbiology, Fujita Health University School of Medicine, Toyoake,

Aichi, Japan.

4. Department of Microbiology and Immunology, University of Minnesota Medical

School, 689 23[rd] Avenue S.E. Microbiology Research Facility, Minneapolis, 55455,

MN, USA.

5. Laboratory of Tuberculosis, Institute of Tropical Disease, Universitas Airlangga,

Surabaya, East Java, Indonesia.

e-mail: y-tateishi@med.niigata-u.ac.jp

Abstract

Mycobacterium intracellulare is a major etiological agent of the recently expanding

Mycobacterium avium–intracellulare complex pulmonary disease (MAC-PD).

Therapeutic regimens that include a combination of macrolides and antituberculous

drugs have been implemented with limited success. To identify novel targets for drug

development that accommodate the genomic diversity of *M. avium–intracellulare*, we

subjected eight clinical MAC-PD isolates and the type strain ATCC13950 to genome-

wide profiling to comprehensively identify universally essential functions. Among these

strains, we identified 131 shared essential or growth-defect-associated genes. Unlike the

type strain, the clinical strains showed an increased requirement for genes involved in

gluconeogenesis and the type VII secretion system under standard growth conditions,

the same genes required for hypoxic pellicle-type biofilm formation in ATCC13950.

Consistent with the central role of hypoxia in the evolution of *M. intracellulare*, the

clinical MAC-PD strains showed more rapid adaptation to hypoxic growth than the type

strain. Importantly, the increased essentiality of hypoxic fitness genes was confirmed in

a mouse lung infection model. These findings confirm the concordant gene essentiality

under hypoxic conditions *in vitro* and hypoxia-related conditions *in vivo*, and highlight

the importance of using clinical strains and host-relevant growth conditions to identify

high-value targets for drug development.

Introduction

In contrast to the ongoing decline in the rate of tuberculosis, *Mycobacterium avium–*

intracellulare complex pulmonary disease (MAC-PD) is increasing in many parts of the

world[1,2]. The standard therapy for MAC-PD involves multidrug chemotherapy that

includes clarithromycin and several antituberculous drugs. Unfortunately, treatment

failure is common, resulting in relapse, disease progression, and death[3]. Therefore, it is

critical to investigate the bacterial physiological characteristics that underlie the ability

of MAC-PD strains to establish and maintain chronic drug-recalcitrant infections.

Recent advances in comparative genomics have revealed major differences in

the genomic features of reference type strains and circulating pathogenic clinical

isolates. Extensive genomic diversity has been recognized across clinical isolates of

nontuberculous mycobacteria (NTM). *Mycobacterium avium* has been divided into four

subspecies including subspecies, *avium*, *hominissuis*, *silvaticum*, and *paratuberculosis*[4].

Whereas *M. intracellulare* has been divided into two subgroups, typical *M.*

intracellulare (TMI) and *M. paraintracellulare–M. indicus pranii* (MP-MIP), with

currently two additional distinguished subspecies, *yongonense*, and *chimaera*[5,6]. Despite

the increasing number of sequenced MAC-PD strains, the molecular genetic

mechanisms that underlie their virulence and pathogenesis are still poorly understood.

The burgeoning field of functional genomics has allowed the rapid assessment

of molecular aspects of organisms on a genome-wide scale. Transposon (Tn) sequencing

(TnSeq) is a functional genomic approach that combines saturation-level transposon-

insertion mutagenesis with next-generation sequencing (NGS) to comprehensively

evaluate the fitness costs associated with gene inactivation in bacteria. Over the last

decade, TnSeq has been applied to numerous bacterial species using various

approaches[7]. Whereas initial studies applied TnSeq to reference strains, recent

approaches have included the comparative assessment of representative clinical strains

to better understand the fundamental aspects of pathogen biology[8]. We have previously

used TnSeq to globally characterize the genes that are essential for growth versus those

that are essential for hypoxic pellicle formation in the *M. intracellulare* type strain

ATCC13950[9]. However, given the recent expansion of MAC-PD and the well-

recognized high levels of genomic diversity in this complex of bacteria, data obtained

from the study of a decades-old reference strain is not expected to represent the

diversity of extant clinical isolates. ATCC13950 was originally isolated from the

abdominal lymph node of the 34-month-old female[10]. Even though the etiological

bacterial species are the same, the clinical manifestation of infant lymphadenitis differs

markedly from MAC-PD, which often occurs in middle-aged and elderly female

patients with no predisposing immunological disorder. Therefore, it is essential to

investigate a collection of strains that captures the diversity of recent clinical MAC-PD

strains in order to define key molecular aspects of their pathogenesis and virulence.

In this study, we used TnSeq to analyze a set of recently isolated clinical *M.*

intracellulare strains with diverse genotypes. We integrated the gene essentiality data

for these strains and the type strain to identify the common essential and growth-defect-

associated genes that represent the genomic diversity of this group of pathogens. We

also identified the greater essentiality of genes involved in gluconeogenesis, the type

VII secretion system, and cysteine desulfurase in the clinical MAC-PD strains than in

the type strain. Furthermore, the requirement for these genes was confirmed in a mouse

lung infection model. The profiles of gene essentiality in clinical MAC-PD strains

suggest the mechanism of hypoxic adaptation in NTM in terms of promising drug

targets, especially for strains that cause clinical MAC-PD.

Results

Common essential and growth-defect-associated genes representing the genomic

diversity of *M. intracellulare* strains

To obtain comprehensive information on the genome-wide gene essentiality of *M.*

intracellulare, we included nine representative strains of *M. intracellulare* with diverse

genotypes in this study (Fig. 1a, b). We used TRANSIT[11] to obtain 2–6 million reads of

Tn insertions. The average number of Tn insertion reads was > 50 at Tn-inserted TA

sites, which confirmed the TnSeq data (Supplementary Table 1). The numbers of

essential genes in the clinical MAC-PD strains, calculated with the hidden Markov

model (HMM; a transition probability model), tended to be smaller than that in

ATCC13950 (Supplementary Table 2). In total, 131 genes were identified as essential or

growth-defect-associated with the HMM analysis across all *M. intracellulare* strains,

including the eight MAC-PD clinical strains and ATCC13950 (Fig. 2a, Supplementary

Table 3). The genes identified as universal essential or growth-defect-associated were

involved in fundamental functions, such as glyoxylate metabolism, purine/pyrimidine

metabolism, amino acid synthesis, lipid biosynthesis, arabinogalactan/peptidoglycan

biosynthesis, porphyrin metabolism, DNA replication, transcription, amino acid tRNA

ligases, ribosomal proteins, general secretion proteins, components of iron–sulfur

cluster assembly, some ABC transporters, and the type VII secretion system. The genes

identified corresponded to the genes that encode conventional and under-development

drug targets, such as those encoding gyrase, arabinosyltransferase, enoyl-(acyl-carrier-

protein) reductase, alanine racemase, DNA-directed RNA polymerase, malate

synthase[12] and F_0F_1 ATP synthase (Fig. 2b). These data suggest that the set of universal

essential or growth-defect-associated genes identified with the HMM analysis of *M.*

intracellulare strains with diverse genotypes has potential utility as antibiotic drug

targets.

116   By comparative genomic analysis, we have revealed 3,153 core genes among

55 *M. intracellulare* strains including those used in this study[5]. Of these core genes, 439

genes were classified as strain-dependent essential or growth-defect-associated with the

HMM analysis (Supplementary Table 4). And among 5,824 accessory genes, 222 genes

were classified as essential or growth-defect-associated with the HMM analysis. The

proportion of these strain-dependent and accessory essential or growth-defect-associated

genes was comparable to the similar analysis of *Streptococcus pneumoniae*[13]. Of note,

17 (3.8%) of strain-dependent essential or growth-defect-associated genes and 14

(6.3%) of accessory essential or growth-defect-associated genes were also included in

the 175 genes required for hypoxic pellicle formation in ATCC13950 reported in our

previous study[9] (Supplementary Table 4).

Genes with increased gene essentiality in clinical MAC-PD strains are also

required for hypoxic pellicle formation in the type strain

The sharing of strain-dependent and accessory essential or growth-defect-associated

genes with genes required for hypoxic pellicle formation in ATCC13950 prompts us to

consider the possibility that the clinical MAC-PD strains increase the gene essentiality

of the genes involved in hypoxic growth compared to ATCC13950. To clarify this, we

compared the gene essentiality of the clinical MAC-PD strains with that in ATCC13950

by integrating the HMM analysis with the resampling analysis, a method of calculating

the number of significantly altered Tn insertion reads (a gene-based permutation

model). With this integrating analysis, 121 genes in the clinical MAC-PD strains were

shown to have significantly fewer Tn insertions than ATCC13950 (Fig. 3,

Supplementary Tables 5, 6). Of these, nine genes were identified in 3–5 clinical MAC-

PD strains. These included genes encoding phosphatases (OCU_RS30185,

OCU_RS35795), cysteine desulfurase (*csd* [OCU_RS40305]), methyl transferase

(OCU_RS41540), a hypothetical protein (OCU_RS47290), type VII secretion

components (*mycP5* [OCU_RS38275], *eccC5* [OCU_RS38345]), a glycine cleavage

system aminomethyltransferase (*gcvT* [OCU_RS35955]), and phosphoenolpyruvate

carboxykinase (*pckA* [OCU_RS48660]) (Figs 3, 4). The genes involved in

gluconeogenesis (such as *pckA*), the glycine cleavage system, and the type VII secretion

system were also identified as required for hypoxic pellicle formation in ATCC13950 in

our previous study[9]. The Tn insertion reads were significantly reduced in the pathways

of gluconeogenesis, including in fructose-1,6-bisphosphatase (*glpX*),

phosphoenolpyruvate carboxykinase (*pckA*), and pyruvate carboxylase (*pca*), but were

significantly increased in those of glycolysis, including in pyruvate kinase (*pykA*) and

pyruvate dehydrogenase (*aceE*, *lpdC*), suggesting that carbohydrate metabolism was

metabolically remodeled in the clinical MAC-PD strains, which mimics the metabolism

observed during hypoxic pellicle formation in ATCC13950 (Fig. 5).

On the other hand, 162 genes were detected with significantly more Tn insertions in

the clinical MAC-PD strains than in ATCC13950 (Fig. 3). Of these, 12 genes were

identified in more than seven strains. The increased Tn insertion reads in the genes

encoding superoxide dismutase (OCU_RS25945) and pyruvate dehydrogenase (*aceE*

[OCU_RS35720]) in the clinical MAC-PD strains suggest the inessentiality of the

detoxification of oxygen radicals and glycolysis metabolism in these strains (Figs 3–5).

These genes also included those encoding amidohydrolases (OCU_RS34400,

OCU_RS34425, OCU_RS34445), some transcription factors (OCU_RS28440,

OCU_RS30230), a CoA transferase (OCU_RS34390), and some predicted transporters

(OCU_RS28435, OCU_RS39950, OCU_RS39955, OCU_RS44685). Furthermore, the

increase in Tn insertion reads in predicted multidrug-resistant genes, such as those

encoding the DrrAB-ABC transporter complex (OCU_RS39950, OCU_RS39955)[14] and

a putative lipoprotein LpqB (OCU_RS44685), encoded by the two-component system

mtrAB operon[15-16], in the clinical MAC-PD strains were found.

Requirement of genes with increased gene essentiality in the clinical MAC-PD

strains for mouse lung infection

Hypoxia is the major environmental condition in natural water[17], biofilms[18], and

infections in humans and animals characterized by caseous granuloma lesions[19]. The

impact of hypoxia on mycobacteria under various ecological circumstances implies that

the genes with increased essentiality in the clinical MAC-PD strains compared to

ATCC13950 and the genes required for hypoxic pellicle formation in ATCC13950 may

be required for infection *in vivo*. To examine whether the set of genes with increased

essentiality in the clinical MAC-PD strains also shows increased essentiality during

infection *in vivo*, we performed TnSeq in mice infected intratracheally with the Tn

mutant library strains of M.i.198 or M.i.27, which we had demonstrated to be virulent in

mice (Supplementary Fig. 1a). The time course of the changes in the bacterial burden

showed a pattern similar to those of the wild-type strains M.i.198 and M.i.27,

respectively, except that it was not possible to harvest sufficient colonies (as few as

10^4/mouse) in the few mice infected with the M.i.27 Tn mutant strain in week 8 and

week 16 (Supplementary Fig. 1b, Supplementary Table 7). With a resampling analysis,

629 and 512 genes were shown to be required for *in vivo* infection with transposon

mutant library strains M.i.27 and M.i.198, respectively, between day 1 and week 16 of

infection (Fig. 6a, Supplementary Fig. 1c, d, Supplementary Tables 8, 9). Of these

genes, 72 (11.9%) and 50 (9.76%) in M.i.27 and M.i.198, respectively, were also

required for hypoxic pellicle formation in ATCC13950. Forty-one genes also required

for hypoxic pellicle formation in ATCC13950 were identified in both M.i.27 and

M.i.198, and included genes associated with succinate production (multifunctional 2-

oxoglutarate metabolism enzyme) and the glyoxylate cycle (isocitrate lyase), cysteine

and methionine synthesis (thiosulfate sulfurtransferase [CysA], 5-

methyltetrahydrofolate-homocysteine methyltransferase [MetH]), the serine/threonine-

protein kinase PknG system, proteasome subunits (PrcA, PrcB), DNA repair UvrC,

secretion protein EspR, cell-wall synthesis proteins (such as phosphatidylinositol

[polyprenol-phosphate-mannose-dependent alpha-[1-2]-phosphatidylinositol

pentamannoside mannosyltransferase], peptidoglycan (penicillin-binding protein A

pbpA), and L,D-transpeptidase 2), putative membrane proteins and transporters (such as

the MmpL family, transport accessory protein MmpS3, transport protein MmpL11,

vitamin B_{12} transport ATP-binding protein BacA, ABC transporter substrate-binding

protein, and ABC transporter permease), and mammalian cell entry (Mce) proteins

(OCU_RS48855–48875) (Supplementary Table 10).

The genes only identified in M.i.27 that were also required for hypoxic pellicle

formation in ATCC13950 included genes associated with gluconeogenesis (*pckA, glpX*),

the glycine cleavage system (*gcvT*), the type VII secretion system (*eccD5, espG5,*

eccC5 [OCU_RS38280-OCU_RS38285-OCU_RS38345], *eccC3* [OCU_RS48145]),

and D-inositol 3-phosphate glycosyltransferase (Supplementary Table 10). These three

systems, gluconeogenesis, the glycine cleavage system, and the type VII secretion

system, also showed increased gene essentiality in 3–5 clinical MAC-PD strains than in

ATCC13950 (Figs 3, 6). Although M.i.27 was excluded as a relevant strain showing

increased gene essentiality of these three metabolic systems, the increase in gene

requirements for infection *in vivo* suggests a positive relationship between hypoxic

growth *in vitro* and bacterial growth *in vivo*. In contrast, M.i.198 already showed

increased gene essentiality in these three metabolic systems under aerobic conditions *in*

vitro, so the genes associated with these three systems were not identified as having

fewer Tn insertion reads during infection *in vivo*. Genes only identified in M.i.198 and

required for hypoxic pellicle formation in ATCC13950 included genes associated with

succinate synthesis (2-oxoglutarate oxidoreductase subunit KorA) or encoding the

HTH-type transcriptional repressor KstR and menaquinone reductase (Supplementary

Table 10). As increased gene essentiality in 3–5 clinical strains (excluding M.i.198)

compared with ATCC13950, an increased requirement for the RNA methyltransferase

OCU_RS41540 was identified in M.i.198 during infection *in vivo*. Despite the

differences in the profiles of the genes required for *in vivo* infection between strains,

these data suggest that increased gene essentiality for hypoxic growth confers

advantages for pathogenesis *in vivo*.

We have performed a statistical enrichment analysis of gene sets by GSEA[20] to

investigate whether the genes required for growth of M.i.27 and M.i.198 in mouse lungs

overlap the gene sets required for hypoxic pellicle formation in ATCC13950 as well as

the gene sets showing increased gene essentiality observed in the clinical MAC-PD

strains listed in Fig 3. Of these gene sets including a total of 180 genes, 54% (98 genes)

and 40% (73 genes) were listed as enriched in genes required for mouse infection in

M.i.27 and M.i.198, respectively ($P < 0.01$ in M.i.27 and $P = 0.016$ in M.i.198)

(Supplementary Fig. 2, Supplementary Table 11). The enriched genes both in M.i.27 and

M.i.198 included genes encoding fructose -1,6-bisphosphatase (*glpX*), a type VII

secretion protein (*eccC5*), glycine cleavage system aminomethyltransferase (*gcvT*),

some phosphatase (OCU_RS30185) and a hypothetical protein (OCU_RS47290) (Fig.

6). The enriched genes only in M.i.27 included genes of phosphoenolpyruvate

carboxylinase (*pckA*), cysteine desulfurase (*csd*) and the ESX-5 type VII secretion

component serine protease (*mycP5*). The enriched genes only in M.i.198 included genes

encoding some phosphatase (OCU_RS35975) and class I SAM-dependent RNA

methyltransferase (OCU_RS41540). These data suggest the sharing of the hypoxia-

adaptation pathways between *in vitro* hypoxic pellicle formation and *in vivo* bacterial

growth.

Preferential hypoxic adaptation of clinical MAC-PD strains evaluated with

bacterial growth kinetics

The metabolic remodeling, such as the increased gene essentiality of gluconeogenesis

and the type VII secretion system, and the overlap of the genes required for mouse lung

infection and those required for hypoxic pellicle formation involved by confer these

metabolic pathways suggests the preferential adaptation of the clinical MAC-PD strains

to hypoxic conditions. To confirm this, we investigated the growth kinetics of each

strain under aerobic and hypoxic conditions. In terms of growth kinetics, most of the

clinical MAC-PD strains (except M001) entered logarithmic (log) phase earlier than

ATCC13950 under 5% oxygen (Fig. 7a, Supplementary Fig. 3). However, under aerobic

conditions, the clinical MAC-PD strains entered log phase at a similar time to

ATCC13950. The growth rate at mid-log point (midpoint) was significantly reduced

under hypoxic conditions in a total of 5 clinical MAC-PD strains (M.i.198, M.i.27,

M003, M019 and M021) compared to aerobic conditions (Fig. 7b, Supplementary Fig.

3). The growth rate at midpoint under hypoxic conditions was significantly lower in

these 5 clinical MAC-PD strains than in ATCC13950. By contrast, the growth rate at

midpoint was significantly increased under hypoxic conditions compared to aerobic

conditions in ATCC13950. And there was no significant change in the growth rate

between aerobic and hypoxic conditions in the rest 3 clinical MAC-PD strains (M018,

M001 and MOTT64). The data of more rapid adaptation to hypoxia followed by

reduced growth rate once adapted indicate the greater metabolic advantage on

continuous logarithmic bacterial growth under hypoxic conditions in the clinical MAC-

PD strains than in ATCC13950.

Effects of knockdown of universal and accessory/strain-dependent essential or

growth-defect-associate genes in clinical MAC-PD strains

To facilitate the functional analysis of the TnSeq-hit genes, it is necessary to construct

the genetically-engineered strains. Still now, there are very few reports of gene

engineering of MAC strains and none of them has been accompanied by the follow-up

studies[21-26]. Moreover, no attempts have been reported to construct the knockdown

strains in MAC strains. With an intention to break through this current situation, we

used the pRH2052/pRH2521 clustered regularly interspaced short palindromic repeats

interference (CRISPR-i) system to construct knockdown strains where the expression of

the target gene is suppressed[27]. Of the nine *M. intracellulare* strains analyzed in this

study, seven could be transformed with the vectors for single guide RNA (sgRNA)

expression and dCas9 expression. However, the other two strains, M001 and MOTT64,

could not be transformed with these CRISPR-i vectors. To confirm the gene essentiality

detected with the HMM analysis, we evaluated the consequent growth inhibition in the

knockdown strains of representative universal essential or growth-defect-associated

genes, including *glcB*, *inhA*, *gyrB*, and *embB*. The knockdown strains showed growth

suppression with comparative growth rates of knockdown strains to vector control

strains that do not express single sgRNA reaching 10^{-1} to 10^{-4} (Fig. 8a).

As representative accessory and strain-dependent essential or growth-defect-

associated genes, we evaluated the 9 genes showing reduced Tn insertion reads in the 3–

5 clinical MAC-PD strains as listed in Fig. 3, as well as *glpX*, a gene of gluconeogenesis

where four MAC-PD strains showed reduced Tn insertion reads in a resampling analysis

(Fig. 5b). The growth was suppressed in the *pckA*-knockdown strains in M.i.198 and

M003, in the *glpX*-knockdown strain in M003, and in the cysteine desulfurase gene

(*csd*)-knockdown strains in M.i.198 and M003, with comparative growth rates of

knockdown strains to vector control strains reaching 10^{-2} to 10^{-4} (Fig. 8b). The growth

tended to be suppressed in the type VII secretion protein gene (*eccC5*)-knockdown

strain in M003 and in the type VII secretion-associated serine protease mycosin gene

(*mycP5*)-knockdown strain in M003, with comparative growth rates of knockdown

strains to vector control strains reaching less than 10^{-1}. By contrast, the growth was not

suppressed in these knockdowns trains in ATCC13950, with comparative growth rates

of knockdown strains to vector control strains not reaching 10^{-1}, which was consistent to

the inessentiality of these genes under aerobic conditions in ATCC13950. As for

M.i.198 and M003, the consistency between TnSeq data and CRISPR-i data was

supported by a correlation between the growth suppression of the knockdown strains

and the fitness costs evaluated by a resampling analysis (Supplementary Table 12).

307   In contrast to the cases of M.i.198 and M003, the growth of knockdown strains

308 of the accessory and strain-dependent essential genes was not effectively suppressed in

309 the other MAC-PD strains despite positively detected in TnSeq (Fig. 5b, Supplementary

310 Fig. 4). These data suggest the difficulties in the effective operation of genetically

311 modified system using foreign genes from different bacterial species in MAC-PD

312 strains. Although the suppression of gene expression was confirmed in the knockdown

313 strains of these accessory and strain-dependent essential or growth-defect-associate

314 genes, the gene expression levels were not suppressed to < 1% as reported in *M.*

315 *tuberculosis* (Mtb)[28] (Supplementary Fig. 5). The differences between strains as well as

316 between genes in the degree of suppression of gene expression required to inhibit

317 bacterial growth may explain such inconsistencies between the TnSeq and CRISPR-i

318 data.

319   To be summarized, we could validate the universal essential or growth-defect-

320 associated genes by the CRISPR-i system in the *M. intracellulare* strains used in this

321 study. And, although not successful with all strains, we could validate the accessory and

322 strain-dependent essential or growth-defect-associated genes, especially genes of

323 gluconeogenesis, type VII secretion components and iron-sulfur cluster assembly in

324 several MAC-PD strains like M.i.198 and M003.

Discussion

In this study, using TnSeq of nine genetically diverse *M. intracellulare* strains, we have

identified 131 shared essential and growth-defect-associated genes with an HMM

analysis. The genes identified are promising drug targets, representing the genomic

diversity of the clinical MAC-PD strains and reflecting the actual clinical situation. The

strategy of narrowing the essential genes of bacteria is a useful approach for refining the

database of drug targets. We have also demonstrated an overview of the differences in

gene essentiality required for hypoxic growth between the ATCC type strain and the

clinical *M. intracellulare* strains responsible for MAC-PD, with the evidence of earlier

entry to log phase followed by slower growth rate observed in most of these clinical

strains compared to ATCC13950. This is the first study to demonstrate the positive

relationship between the profiles of increased gene essentiality required for hypoxic

metabolic remodeling and the phenotypes advantageous for bacterial survival under

hypoxic conditions.

In this study, 121 genes were detected as significantly reduced Tn insertion in the

clinical MAC-PD strains compared to ATCC13950, which was higher than the case of

Mtb showing 10–50 genes with altered Tn insertion reads detected by a resampling

analysis[8]. This may reflect the genomic diversity of *M. intracellulare* compared with

that of Mtb. The proportion of accessory genes is larger in NTM than in Mtb[29]. In a

clinical trial of phage therapy for *M. abscessus* infection, the number of accessory genes

was shown to decrease with the loss of virulence in *M. abscessus* during the course of

therapy[30], suggesting some role for accessory genes for its clinical pathogenesis. Our

data of the overlap of the accessory essential or growth-defect-associated genes such as

pckA and *mycP5* (Supplementary Table 5) with genes required for mouse lung infection

(Fig. 6b, Supplementary Fig. 2) supports the idea that accessory genes (non-core genes)

may encode functions required for bacterial survival, especially under nonoptimal

conditions, such as hypoxia and during the infection of human and animal bodies.

Our TnSeq data are consistent with metabolomic data that have shown the

importance of the gluconeogenic carbon flow of TCA cycle intermediates for bacterial

growth under hypoxic conditions in terms of the essentiality of PckA and the reverse

TCA cycle[31–34]. Gluconeogenesis is essential for bacterial growth when fatty acids are

used as a major nutrient[33]. PckA catalyzes the conversion of the metabolites derived

from the TCA cycle to the metabolites of gluconeogenesis[32]. The reaction catalyzed by

PckA is essential for bacterial growth when the available nutrients are predominantly

fatty acids[35,36]. The main nutrients for mycobacteria inside hypoxic granulomas in an

infected human or mouse body are lipids, such as fatty acids and cholesterol, rather than

carbohydrates[33,37]. In the present study, *M. intracellulare* grew under 5% oxygen,

whereas Mtb stops growing under 1% oxygen and enters a dormant state[32]. The

depletion of phosphoenolpyruvate in the *pckA* deletion mutant of Mtb reduced the

phosphoenolpyruvate–carbon flux essential for bacterial growth and drug sensitivity[32].

The present study supports the idea that PckA is essential for the microaerobic growth

of mycobacteria. Our TnSeq data imply the preferential adaptation of the clinical MAC-

PD strains for circumstances *in vivo*, which are characterized by hypoxia.

On the other hand, such metabolic shift towards gluconeogenesis and reverse TCA

cycle can also be observed under reactive nitrogen specie and *in vivo* infection as well.

This suggests that the metabolic shift observed under hypoxia is a general stress

response rather than being specific to hypoxia due to the nature of the

provided/available carbon source[38]. Several factors other than hypoxia can be estimated

to be related to the change of the gene essentiality in the clinical MAC-PD strains such

as host-related factors and general stress conditions (i.e. nutrition of abundant fatty

acids and less carbohydrates, low pH circumstances)[33,35-37]. Except for our data of the

profile of genes required for hypoxic pellicle formation in *M. intracellulare*[9], no

fundamental genome-wide data are available on the change of the gene essentiality in

response to each stress factors in non-tuberculous mycobacteria. Further studies are

needed to discriminate which factors are specifically involved in the increased

essentiality of each gene. Although of interest, it is beyond the scope of the current

study.

About 10% of the genes required for mouse infection are also required for hypoxic

pellicle formation. The genes showing increased gene essentiality in the clinical MAC-

PD strains, such as *pckA* and genes encoding the ESX-5 type VII secretion system,

including *eccC5* and *mycP5*, also showed increased gene requirements in mouse

infections. In a TnSeq study of mice infected intravenously with *M. avium*, the genes of

the ESX-5 type VII secretion system were required for infection of the liver and

spleen[39]. The ESX-5 type VII secretion system is involved in the transport of fatty acids

and the virulence of mycobacteria[40,41]. These similarities in the profiles of gene

essentiality prompt us to propose some role of biofilm formation in mycobacterial

pathogenesis in animals and humans. In the present study, we shed light on the impact

of hypoxia on biofilm formation, the acquisition of virulence, and the *in vivo*

pathogenesis of NTM in terms of global gene essentiality. Although the morphologies of

the pellicle and macroscopic lesions (including cavitation and nodular bronchiectasis)

are not similar, monitoring the expression of the genes identified in this study may

provide clues to the mechanism of biofilm formation *in vivo*.

In this study, we revealed the pattern of hypoxic adaptation characteristic to the

clinical MAC-PD strains by the growth curve analysis: an early entry to log phase

followed by a reduced growth rate (Fig. 7). This pattern implicates the continuous impact

on the infected hosts during logarithmic bacterial growth, which may be involved in the

persistent and steadily progressive illness for years in humans[3]. The advantage of slow

growth in mycobacteria may be inferred by the role of a histone-like nucleoid protein,

Mycobacterial DNA-binding protein 1 (MDP1), also designated HupB[42-44]. The

expression of MDP1 is enhanced in the stationary phase, resulting in reduced growth

speed and long-term survival with increased resistance to reactive oxygen species

(ROS)[42-44]. The reduced growth rate after entry to log phase may allow long-term survival

of mycobacteria under ROS-rich circumstances inside host cells. The pattern of hypoxic

adaptation as observed in the clinical MAC-PD strains may enhance the pathogenesis by

extending the period of the repeated process of bacterial invasion and replication in

infected macrophages.

Contrary to our expectation, not all clinical strains showed rapid adaptation from

aerobic to hypoxic conditions relative to ATCC13950. These data suggest that some, but

not a major proportion of, clinical *M. intracellulare* strains do not have an advantageous

phenotype for continuous logarithmic growth under hypoxia. In Mtb, the adaptation to

hypoxia is considered to be necessary for clinical pathogenesis because the conditions

in vivo, such as inside macrophages or granulomas, are hypoxic[38,45]. However, our strain

M001 entered log phase later than ATCC13950 under 5% oxygen. Our inability to

construct knockdown strains in M001 prevented us from clarifying its clinical

pathogenesis without preferential hypoxic adaptation. However, because we have shown

that most of the *M. intracellulare* strains with different genotypes showed preferential

adaptation to hypoxia in this study, our results largely reflect the clinical situation of the

etiological agents of MAC-PD. A possible strategy for identifying factors that confer

clinical pathogenesis will be to focus on the major clinical strains that show preferential

hypoxic adaption.

The data of validation of the universal essential or growth-defect associated

genes by the CRISPR-i system was overall acceptable in the clinically MAC-PD strains

whose knockdown strains could be constructed (Fig. 8a). However, we have

encountered the difficulties of validating the TnSeq-hit genes by CRISPR-i experiment,

especially accessory and strain-dependent essential or growth-defect-associated genes in

the MAC-PD strains other than M.i.198 or M003 (Fig. 8b, Supplementary Fig. 4). As a

possible reason, the low efficiency of the CRISPR-i system may obscure the increase in

gene essentiality suggested by the TnSeq data (Supplementary Fig. 5). The efficiency of

the suppression of gene expression by CRISPR-i in accessory and strain-dependent

essential or growth-defect-associated genes was lower in our *M. intracellulare* strains

than in Mtb[28]. In Mtb, the expression of *mmpL3*, a gene encoding a mycolic acid

transporter, is reported to be suppressed to 6% under 10 ng/ml anhydrotetracycline (aTc)

and by < 1% under 100 ng/ml aTc in the knockdown strains, compared to strains

expressing scrambled single guide RNAs (sgRNA)[28]. The gene expression levels were

suppressed in our knockdown strains by around 10%–40% under 50–200 ng/ml aTc

relative to the levels found in strains not expressing sgRNA (Supplementary Fig. 5). The

growth inhibition effect is theoretically dependent upon the degree to which gene

expression is suppressed. On the other hand, the inconsistent growth phenotypes of

knockdown strains of accessory and strain-dependent essential or growth-defect-

associated genes between MAC-PD strains implies some differences in the degree of

suppression of gene expression required to inhibit bacterial growth between genes as

well as between bacterial strains. Despite the issue of the lower efficiency of the

CRISPR-I system in *M. intracellulare* than in Mtb, this study has opened the avenue for

functional analysis of NTM that leads to the development of novel drugs that target

universal and accessory/strain-dependent essential metabolic pathways that has

significant impact on bacterial growth, especially by making use of M.i.198 and M003

as optimal NTM strains for gene manipulation.

As in the *glpX*-knockdown strains, *csd* knockdown caused growth suppression in

several MAC-PD strains. This type II Csd is involved in the biosynthesis of the iron–

sulfur clusters of the SUF machinery. Compared with the type I cysteine desulfurase

IscS, the SUF machinery is active under oxidative-stress and iron-limiting conditions[46].

Representatives of iron–sulfur cluster proteins include components of the electron-

transport system, such as NADH dehydrogenase I (Nuo) and succinate dehydrogenase

(Sdh), which are components of respiratory complexes I and II, respectively. Under

hypoxic conditions, the non-proton-translocating type II NADH dehydrogenase Ndh is

upregulated, and cytochrome *bd* oxidase (CydAB) and the proton-pumping type I

NADH dehydrogenase (Nuo) are downregulated[38]. In the reverse TCA cycle on the

same reaction pathway as Sdh, fumarate reductase (Fnr) is also an iron–sulfur cluster

protein that produces succinate, and is a proposed alternative electron acceptor under

hypoxic conditions[34,47]. The increased essentiality of the *csd* gene observed in the

clinical MAC-PD strains seems to reflect the switching of the electron-transport system

to less-energy-efficient respiration to support preferential hypoxic survival, including *in*

vivo.

We tried to use inhibitors of phosphoenolpyruvate carboxykinase to confirm the

increased essentiality of the *pckA* gene. However, we detected no positive effect because

millimolar 3-mercaptopicolinic acid is required to show the effect of inhibition, even in

eukaryotic cells, which have no cell wall[48]. Moreover, 3-alkyl-1,8-dibenzylxanthine[49]

also only effectively inhibits PckA in eukaryotic cells. However, PckA has been

proposed as a drug target for Mtb[50], so our TnSeq data seem to be consistent with

studies directed towards antimycobacterial drug discovery.

In summary, we have identified 131 universal essential and growth-defect-

associated genes that covers the genomic diversity of *M. intracellulare* strains

responsible for MAC-PD, which narrowed down the drug targets against MAC-PD.

Furthermore, we have demonstrated a preferential adaptation to hypoxia in clinical

NTM strains compared with the type strain in terms of the increased essentiality of

genes required for hypoxic growth, such as those encoding phosphoenolpyruvate

carboxykinase, the ESX-5 type VII secretion system, and cysteine desulfurase. These

genes were shown to have a significant impact on infection *in vivo* by increasing their

gene requirements. Our data highlight the importance of using clinical strains and

hypoxic conditions in experimental research, and provide fundamental resources for

developing novel drugs directed towards nontuberculous mycobacterial strains that

cause severe clinical diseases.

Methods

Bacterial strains and phages used to prepare Tn mutants

A total of nine strains were used in this study, including seven clinical strains isolated

from patients with MAC-PD in Japan (M.i.198, M.i.27, M018, M003, M019, M021,

and M001)[51], *M. intracellulare* genovar *paraintracellulare* type strain MOTT64 isolated

from MAC-PD in South Korea[52], and the *M. intracellulare* type strain ATCC13950 (Fig.

1). The Tn mutant libraries were constructed using glucose as a carbon source under

aerobic conditions according to the transduction method used for ATCC13950[9], except

that a higher concentration of kanamycin (100 μg/ml) was used to select the transposon

mutants than was used for ATCC13950 (25 μg/ml). In brief, a hundred milliliter of the

bacteria were cultured aerobically in 7H9/ADC broth until mid-log phase (optical

density at a wavelength of 600 nm [OD_{600}] of approximately 0.4–0.6). After harvesting

the bacteria, 10 mL of high-titer mycobacteriophage phAE180[53] was transduced into the

bacteria 1-day at 37°C for inserting Tn and plated on 7H10/OADC agar plates

containing kanamycin. After 2 or 3 week-cultivation at 37°C aerobically, the colonies

were harvested *en masse* and used as Tn mutant libraries. The kanamycin-resistant

colonies ($> 1 \times 10^5$) were harvested and evenly resuspended in 7H9/ADC medium

containing 20% glycerol, aliquoted, and stored at -80 °C until further use.

Next-generation sequencing and TnSeq analysis

DNA libraries were prepared as reported previously[54]. The TnSeq libraries were

sequenced with the NextSeq 500 System, 150-bp single-end run (Illumina, San Diego,

CA, USA). The sequence reads were analyzed as previously described[9]. Gene

essentiality was calculated with resampling and HMM analyses using TRANSIT[11].

First, an HMM analysis was conducted to calculate the fluctuation of the number of Tn

insertion reads individually for each strain. Next, a resampling analysis was conducted

to calculate the difference of the number of Tn insertion reads between each clinical

strain and ATCC13950 (Supplementary Table 6). By comparing the data obtained by

HMM and resampling analyses, we summarized the difference of gene essentiality

between the clinical MAC-PD strains and ATCC13950. To compare the gene

essentiality of the clinical MAC-PD strains and ATCC13950, the annotation for the

clinical MAC-PD strains was adapted from that of ATCC13950 by adjusting the START

and END coordinates of each open reading frame (ORF) in the clinical MAC-PD strains

according to their alignment with the corresponding ORFs of ATCC13950. The function

of TnSeq-hit genes was categorized according to the Kyoto Encyclopedia of Genes and

Genomes (KEGG) database.

Comparison of growth curves under aerobic and hypoxic conditions

First, we precultured the bacteria in 7H9/ADC broth until the bacteria reached early log-

phase (OD_{600} approximately 0.1–0.3). Next, the bacterial cultures were diluted with new

7H9/ADC broth (final OD_{600} 0.003) for the aerobic and hypoxic growth assay. Bacterial

cells were grown statically in 96-well plates containing 250 µl of 7H9/ADC broth in

each well under aerobic conditions or 5% oxygen at 37 °C with humidification (APM-

30D incubator, Astec, Tokyo, Japan). The culture was sampled every 1–3 days. At

sampling, the culture was mixed by pipetting 50 times. An aliquot (4 µl) of the culture

and its diluents were streaked onto 7H10/OADC agar and the CFUs counted. The

growth curve was fitted to the logistic 4P model with the JMP software (SAS Institute

Inc., Cary, NC, USA) to calculate the inflection point (midpoint) and growth rate at

midpoint in the log growth phase. Entry into the log phase was evaluated from the

timing of the inflection point on the growth curve.

Animals

Female C57BL/6JJcl mice were purchased from CLEA Japan (Tokyo, Japan). Six-

week-old mice were used for the infection experiment (Supplementary Fig. 1). All the

mice were kept under specific-pathogen-free conditions in the animal facility of Niigata

University Graduate School of Medicine, according to the institutional guidelines for

animal experiments. The mice were housed (four per cage) under a 12-h/12-h light/dark

cycle. Food and water were available *ad libitum*. Kanamycin was added to the drinking

water to a final concentration of 100 μg/ml to maintain the mice's resistance to the Tn

mutant strains. Euthanasia was conducted with an intraperitoneal injection of a

combined anesthetic (15 μg of medetomidine, 80 μg of midazolam, and 100 μg of

butorphanol), followed by cervical dislocation.

Mouse TnSeq experiment

The protocol was based on our previous mouse lung infection experiment[51]. Twenty

mice each were used for the Tn mutant strains M.i.198 and M.i.27. The mice were

injected intratracheally with 3.25×10^8 and 5.93×10^7 CFUs of the Tn mutant strains,

respectively. The infected lungs were expected to be harvested from five mice at day 1,

week 4, week 8, and week 16 after infection. However, the lungs were harvested from

only four mice infected with the M.i.27 Tn mutant strains in week 8 because we

experienced difficulties with sampling, and the lungs were harvested from only three,

one, and two mice infected with M.i.198 Tn mutant strains in week 8, week 12, and

week 16, respectively, because morbidity was high in the mice infected with M.i.198.

The harvested lungs were homogenized in 4.5 ml of distilled water with the

gentleMACS™ Tissue Dissociator (Miltenyi Biotec, Bergish Gladbach, Germany) and

the homogenates were plated on Middlebrook 7H10–oleic acid ADC (7H10-OADC)

agar plates and cultured for 3 weeks. The colonies were harvested *en masse* for TnSeq.

Gene set enrichment analysis

To investigate the overlap of the gene sets required for growth in mouse lungs with

those required for hypoxic pellicle formation observed in ATCC13950 and those

showing increased gene essentiality observed in the clinical MAC-PD strains, a

statistical enrichment of gene sets was performed by GSEA 4.3.2[20]. Parameters set for

GSEA were as follows: permutations = 1,000, permutation type: gene_set, enrichment

statistic: weighted, metric for ranking genes: Signal2Noise, max size: 500, min size: 15.

Construction and growth assay of knockdown strains

Knockdown strains were constructed with the CRISPR-dCas9-mediated RNA

interference system, as reported previously[27]. Gene-specific 20-nt sequences were

selected as protospacers for incorporation into the sgRNAs (Supplementary Table 13).

pRH2502 (a vector expressing an inactivated version of the *cas9* DNA sequence) and

pRH2521 (an sgRNA-expressing vector whose expression is regulated by the TetR-

regulated smyc promotor $P_{myc1}tetO$) were introduced into the bacteria to obtain colonies

of the knockdown strains. For selection, 25 μg/ml kanamycin and 50 μg/ml hygromycin

were used for ATCC13950, and 100 μg/ml kanamycin and 50 μg/ml hygromycin were

used for the clinical strains. The growth assay of the knockdown strains was performed

in 7H9/ADC broth containing the above-mentioned concentrations of kanamycin and

hygromycin under aerobic conditions at 37 °C with humidification (APM-30D

incubator, Astec, Tokyo, Japan). During the assay, the cultures were supplemented with

anhydrotetracycline (aTc) to a final concentration of 50 ng/ml (as a concentration that is

not toxic to ATCC13950 and M.i.27) or 200 ng/ml (for the other clinical strains). aTc

was added repeatedly every 48 h to maintain the induction of dCas9 and sgRNAs in

experiments that extended beyond 48 h[27]. An aliquot (4 μl) of the culture and its diluents

were streaked onto 7H10/OADC agar plates containing kanamycin and hygromycin,

and the CFUs were counted. Growth inhibition was evaluated as the ratio of the CFUs

of knockdown strains compared with those of the vector control strains lacking the

guide RNA sequence. To confirm the downregulation of gene expression in the

knockdown strains, quantitative reverse transcription PCR (qRT-PCR) was performed.

Bacteria were cultured in 10 ml of Middlebrook 7H9 medium (Difco Laboratories,

Detroit, MI, USA) containing 0.2% glycerol, 0.1% Tween 80 (MP Biochemicals,

Illkrich, France), and 10% Middlebrook albumin–dextrose–catalase (7H9/ADC/Tween

80), kanamycin, and hygromycin until mid-log phase (OD_{600} approximately 0.4–0.6).

The cultures were supplemented with aTc every 48 h, as in the growth assay for the

knockdown strains. One day after the second addition of aTc, RNA was extracted by

bead-beating six times at 5000 rpm for 30 s with cooling (Micro Smash MS 100R Cell

Disruptor; Tomy Seiko, Tokyo, Japan) and cDNA was synthesized with the Direct-zol™

RNA Miniprep (Zymo Research, Orange, CA, USA) and ReverTra Ace qPCR RT

Master Mix with gDNA Remover (Toyobo, Osaka, Japan). qRT-PCR was performed

with the CFX Connect Real-time PCR Detection System (Bio-Rad, Hercules, CA,

USA), according to the manufacturer's instructions. The data were normalized to the

expression of *sigA*.

Statistical analyses

Pearson's correlation test was performed between the fitness costs ($\log_2$FC calculated by

the resampling assay), efficiency of knockdown (assessed by qRT-PCR) and growth

suppression in the knockdown strains. Statistical analyses of the growth curves were

performed using a 4-parameter logistic model (logistic 4P model), Wilcoxon two-

sample test, and a one-way analysis of variance (ANOVA) followed by Steel's multiple

comparisons test using the JMP software program (SAS Institute Inc., Cary, NC, USA).

Data are presented as the mean ± SD. Statistical significance was set as P values less

than 0.05.

[References]

1.   Adjemian, J., Olivier, K.N., Seitz, A.E., Holland, S.M. & Prevots, D.R.

616   Prevalence of nontuberculous mycobacterial lung disease in U.S. Medicare

617   beneficiaries. *Am J Respir Crit Care Med* **185**, 881-886 (2012).

618   2.   Namkoong, H. *et al.* Epidemiology of pulmonary nontuberculous mycobacterial

619   disease, Japan. *Emerg Infect Dis* **22**, 1116-1117 (2016).

620   3.   Daley, C.L. *et al.* Treatment of nontuberculous mycobacterial pulmonary

621   disease: an official ATS/ERS/ESCMID/IDSA clinical practice guideline. *Eur*

622   *Respir J* **56**, 2000535 (2020).

623   4.   Uchiya, K.I. *et al.* Comparative genome analyses of *Mycobacterium avium*

624   reveal genomic features of its subspecies and strains that cause progression of

625   pulmonary disease. *Sci Rep* **7**, 39750 (2017).

626   5.   Tateishi, Y. *et al.* Comparative genomic analysis of *Mycobacterium*

627   *intracellulare*: implications for clinical taxonomic classification in pulmonary

628   *Mycobacterium avium-intracellulare* complex disease. *BMC Microbiol* **21**, 103

629   (2021).

630   6.   Tortoli, E. *et al.* Genome-based taxonomic revision detects a number of

631   synonymous taxa in the genus *Mycobacterium*. *Infect Genet Evol* **75**, 103983

632   (2019).

7.   van Opijnen, T. & Camilli, A. Transposon insertion sequencing: a new tool for

systems-level analysis of microorganisms. *Nat Rev Microbiol* **11**, 435-442

(2013).

8.   Carey, A.F. *et al.* TnSeq of *Mycobacterium tuberculosis* clinical isolates reveals

strain-specific antibiotic liabilities. *PLoS Pathog* **14**, e1006939 (2018).

9.   Tateishi, Y. *et al.* Genome-wide identification of essential genes in

Mycobacterium intracellulare by transposon sequencing - Implication for

metabolic remodeling. *Sci Rep* **10**, 5449 (2020).

10. Cuttino, J.T. & Mc, C.A. Pure granulomatous nocardiosis, a new fungus disease

distinguished by intracellular parasitism; a description of a new disease in man

due to a hitherto undescribed organism, *Nocardia intracellularis*, n. sp.,

including a study of the biologic and pathogenic properties of this species. *Am J*

Pathol **25**, 1-47 (1949).

11. DeJesus, M.A., Ambadipudi, C., Baker, R., Sassetti, C. & Ioerger, T.R.

TRANSIT--A Software Tool for Himar1 TnSeq Analysis. *PLoS Comput Biol* **11**,

e1004401 (2015).

12. Krieger, I.V. *et al.* Structure-guided discovery of phenyl-diketo acids as potent

inhibitors of *M. tuberculosis* malate synthase. *Chem Biol* **19**, 1556-1567 (2012).

13. Rosconi, F. *et al.* A bacterial pan-genome makes gene essentiality strain-

dependent and evolvable. *Nat Microbiol* **7**, 1580–1592 (2022).

14. Coll, F. *et al.* Genome-wide analysis of multi- and extensively drug-resistant

Mycobacterium tuberculosis. *Nat Genet* **50**, 307-316 (2018).

15. Nguyen, H.T., Wolff, K.A., Cartabuke, R.H., Ogwang, S. & Nguyen, L. A

lipoprotein modulates activity of the MtrAB two-component system to provide

intrinsic multidrug resistance, cytokinetic control and cell wall homeostasis in

Mycobacterium. *Mol Microbiol* **76**, 348-364 (2010).

16. Li, S. *et al.* CRISPRi chemical genetics and comparative genomics identify

genes mediating drug potency in *Mycobacterium tuberculosis*. *Nat Microbiol.* **7**,

766–779 (2022).

17. Kirschner, R.A., Jr., Parker, B.C. & Falkinham, J.O., 3rd. Epidemiology of

infection by nontuberculous mycobacteria. *Mycobacterium avium,*

Mycobacterium intracellulare, and *Mycobacterium scrofulaceum* in acid, brown-

water swamps of the southeastern United States and their association with

environmental variables. *Am Rev Respir Dis* **145**, 271-275 (1992).

18. Werner, E. *et al.* Stratified growth in *Pseudomonas aeruginosa* biofilms. *Appl*

Environ Microbiol **70**, 6188-6196 (2004).

19.   Via, L.E. *et al.* Tuberculous granulomas are hypoxic in guinea pigs, rabbits, and

670   nonhuman primates. *Infect Immun* **76**, 2333-2340 (2008).

671   20.   Subramanian, A. *et al.* Gene set enrichment analysis: A knowledge-based

672   approach for interpreting genome-wide expression profiles. *Proc Natl Acad Sci*

673   *U S A* **102,** 15545–15550 (2005).

674   21.   Kryzywinska, E., Bhatnagar, S., Sweet, L., Chatterjee, D. & Schorey, J.S.

675   *Mycobacterium avium* 104 deleted of the methyltransferase D gene by allelic

676   replacement lacks serotype-specific glycopeptidolipids and shows attenuated

677   virulence in mice. *Mol Microbiol* **56**, 1262-1273 (2005).

678   22.   Irani, V.R., Lee, S.-H., Eckstein, T.M., Inamine, J.M., Belisle, J.T. & Maslow,

679   J.N. Utilization of a ts-sacB selection system for the generation of a

680   *Mycobacterium avium* serovar-8 specific glycopeptidolipid allelic exchange

681   mutant. A*nnals Clin Microbiol Antimicrob* **3**, 18 **(**2004).

682   23.   Marklund, B.I., Speert, D.P. & Stokes, R.W. Gene replacement through

683   homologous recombination in *Mycobacterium intracellulare*. *J Bacteriol* **177**,

684   6100-6105 (1995).

685   24.   Marklund, B.I., Mahenthiralingam, E. & Stokes, R.W. Site-directed mutagenesis

686   and virulence assessment of the *katG* gene of *Mycobacterium intracellulare*. *Mol*

687  *Microbiol* **29**, 999-1008 (1998).

688 25. Mahenthiralingam, E., Marklund, B.I., Brooks, L.A., Smith, D.A., Bancroft, G.J.

689   & Stokes, R.W. Site-directed mutagenesis of the 19-kilodalton lipoprotein

690   antigen reveals No essential role for the protein in the growth and virulence of

691   *Mycobacterium intracellulare*. *Infect Immun*. **66**, 3626-3634 (1998).

692 26. Hinds, J. *et al.* Enhanced gene replacement in mycobacteria. *Microbiology* **145**,

693   519-527 (1999).

694 27. Singh, A.K. *et al.* Investigating essential gene function in *Mycobacterium*

695   *tuberculosis* using an efficient CRISPR interference system. *Nucleic Acids Res*

696   **44**, e143 (2016).

697 28. McNeil, M.B. & Cook, G.M. Utilization of CRISPR interference to validate

698   MmpL3 as a drug target in *Mycobacterium tuberculosis*. *Antimicrob Agents*

699   *Chemother* **63,** e00629-19 (2019).

700 29. Yang, T. *et al.* Pan-genomic study of *Mycobacterium tuberculosis* reflecting the

701   primary/secondary genes, generality/individuality, and the interconversion

702   through copy number variations. *Front Microbiol* **9**, 1886 (2018).

703 30. Nick, J.A. *et al.* Host and pathogen response to bacteriophage engineered against

704   *Mycobacterium abscessus* lung infection. *Cell* **185**, 1860-1874.e12 (2022).

705   31.   Eoh, H. *et al.* Metabolic anticipation in *Mycobacterium tuberculosis*. *Nat*

706   *Microbiol* **2**, 17084 (2017).

707   32.   Lim, J. *et al.* Phosphoenolpyruvate depletion mediates both growth arrest and

708   drug tolerance of *Mycobacterium tuberculosis* in hypoxia. *Proc Natl Acad Sci U*

709   *S A* **118,** e2105800118 (2021).

710   33.   Marrero, J., Rhee, K.Y., Schnappinger, D., Pethe, K. & Ehrt, S. Gluconeogenic

711   carbon flow of tricarboxylic acid cycle intermediates is critical for

712   *Mycobacterium tuberculosis* to establish and maintain infection. *Proc Natl Acad*

713   *Sci U S A* **107**, 9819-9824 (2010).

714   34.   Watanabe, S. *et al.* Fumarate reductase activity maintains an energized

715   membrane in anaerobic *Mycobacterium tuberculosis*. *PLoS Pathog* **7**, e1002287

716   (2011).

717   35.   Gouzy, A., Healy, C., Black, K.A., Rhee, K.Y. & Ehrt, S. Growth of

718   *Mycobacterium tuberculosis* at acidic pH depends on lipid assimilation and is

719   accompanied by reduced GAPDH activity. *Proc Natl Acad Sci U S A* **118,**

720   e2024571118 (2021).

721   36.   Quinonez, C.G. *et al.* The role of fatty acid metabolism in drug tolerance of

722   *Mycobacterium tuberculosis*. *mBio* **13**, e0355921 (2022).

37.   Pandey, A.K. & Sassetti, C.M. Mycobacterial persistence requires the utilization

of host cholesterol. *Proc Natl Acad Sci U S A* **105**, 4376-4380 (2008).

38.   Prosser, G. *et al.* The bacillary and macrophage response to hypoxia in

tuberculosis and the consequences for T cell antigen recognition. *Microbes*

Infect **19**, 177-192 (2017).

39.   Dragset, M.S. *et al.* Global assessment of *Mycobacterium avium* subsp.

hominissuis genetic requirement for growth and virulence. *mSystems* **4,** e00402-

19 (2019).

40.   Ates, L.S. *et al.* Essential role of the ESX-5 secretion system in outer membrane

permeability of pathogenic mycobacteria. *PLoS Genet* **11**, e1005190 (2015).

41.   Ates, L.S. *et al.* The ESX-5 system of pathogenic mycobacteria is nvolved in

capsule integrity and virulence through its substrate PPE10. *PLoS Pathog* **12**,

e1005696 (2016).

42.   Savitskaya, A. *et al.* C-terminal intrinsically disordered region-dependent

organization of the mycobacterial genome by a histonelike protein. *Sci Rep* 8, 1–

15 (2018).

43.   Enany, S. *et al.* Mycobacterial DNA-binding protein 1 is critical for long term

survival of Mycobacterium smegmatis and simultaneously coordinates cellular

functions. *Sci Rep* 7, 1–11 (2017).

44.   Shaban, A.K. *et al.* Mycobacterial DNA-binding protein 1 is critical for BCG

survival in stressful environments and simultaneously regulates gene expression.

Sci Rep 13, 14157 (2023).

45.   Rustad, T.R., Sherrid, A.M., Minch, K.J. & Sherman, D.R. Hypoxia: a window

into *Mycobacterium tuberculosis* latency. *Cell Microbiol* **11**, 1151-1159 (2009).

46.   Takahashi, Y. & Tokumoto, U. A third bacterial system for the assembly of iron-

sulfur clusters with homologs in archaea and plastids. *J Biol Chem* **277**, 28380-

28383 (2002).

47.   Cecchini, G., Schröder, I., Gunsalus, R.P. & Maklashina, E. Succinate

dehydrogenase and fumarate reductase from *Escherichia coli*. *Biochim Biophys*

Acta **1553**, 140-57 (2002).

48.   Brearley, M.C., Daniel, Z., Loughna, P.T., Parr, T. & Brameld, J.M. The

phosphoenolpyruvate carboxykinase (PEPCK) inhibitor, 3-mercaptopicolinic

acid (3-MPA), induces myogenic differentiation in C2C12 cells. *Sci Rep* **10**,

22177 (2020).

49.   Montal, E.D. *et al.* Inhibition of phosphoenolpyruvate carboxykinase blocks

lactate utilization and impairs tumor growth in colorectal cancer. *Cancer Metab*

7, 8 (2019).

50.   McLeod, M.J., Krismanich, A.P., Assoud, A., Dmitrienko, G.I. & Holyoak, T.

Characterization of 3-[(carboxymethyl)thio]picolinic acid: a novel inhibitor of

phosphoenolpyruvate carboxykinase. *Biochemistry* **58**, 3918-3926 (2019).

51.   Tateishi, Y. *et al.* Virulence of *Mycobacterium intracellulare* clinical strains in a

mouse model of lung infection - role of neutrophilic inflammation in disease

severity. *BMC Microbiol* **23**, 94 (2023).

52.   Kim, B.J. *et al.* Complete genome sequence of *Mycobacterium intracellulare*

clinical strain MOTT-64, belonging to the INT1 genotype. *J Bacteriol* **194**, 3268

(2012).

53.   Kriakov, J., Lee, S. & Jacobs, W.R., Jr. Identification of a regulated alkaline

phosphatase, a cell surface-associated lipoprotein, in *Mycobacterium smegmatis*.

J Bacteriol **185**, 4983-4991 (2003).

54.   Minato, Y. *et al.* Genomewide assessment of *Mycobacterium tuberculosis*

conditionally essential metabolic pathways. *mSystems* **4,** e00070-19 (2019).

55.   Shannon, P. *et al.* Cytoscape: a software environment for integrated models of

biomolecular interaction networks. *Genome Res* **13**, 2498-2504 (2003).

Acknowledgments

The authors thank Dr. Torin Weisbrod and Dr. William R. Jacobs, Jr. (Albert Einstein College of Medicine, Bronx, NY, USA) for kindly providing mycobacterial phage phAE180, and Dr. Norikazu Hara and Dr. Takeshi Ikeuchi (Department of Molecular Genetics, Bioresource Science, Brain Center for Bioresources, Brain Research Institute, Niigata University) for next-generation sequencing with the NextSeq 500 System. This work was supported by Grants-in-Aid for Scientific Research (grant numbers 20KK0216, 21K19637, and 22H03115 to Yoshitaka Tateishi) from the Ministry of Health, Labour, and Welfare, of Japan This work was also supported by grants from the Japanese Ministry of Education, Culture, Sports, Science, and Technology, the Ministry of Health, and the Research Program on Emerging Infectious Disease from the Japan Agency for Medical Research and Development (23fk0108673h0501).

Author contributions

Y.T. conceived, designed and performed the experiments. Y.O., A.N., and S.M. contributed reagents or materials. Y. Morishige., Y. Minato, and A.D.B. contributed analytical tools. Y. Morishige., Y. Minato, A.D.B., and S.M. provided critical suggestions throughout the preparation of the manuscript. Y.T. wrote the manuscript. All authors read

and approved the final manuscript.

Availability of data and materials

The datasets used or analyzed in this study are available from the corresponding author

upon reasonable request.

Declarations

Ethics approval and consent to participate

All experimental protocols were approved by the Institutional Review Board of Niigata

University (approval numbers: 2019-0020 for the use of human tissue samples and

SA00506 for experiments on live vertebrates). All experiments on humans and human

samples were conducted in accordance with relevant guidelines and regulations. Informed

consent was obtained from all patients. All experiments on animals were conducted in

accordance with the relevant guidelines and regulations, and the study complied with the

ARRIVE guidelines.

Consent for publication

According to the protocols approved by the Institutional Review Board of Niigata

University mentioned above, consent for publication was obtained from the patients

enrolled in this study.

Competing interests

The authors declare that they have no competing interests.

Fig. 1 Phylogenetic tree of the *M. intracellulare* strains and strategy of the

experiment in this study. a Phylogenetic tree of the *M. intracellulare* strains used in this

study. The tree was generated based on average nucleotide identity (ANI) with the

neighbor-joining method. TMI: typical *M. intracellulare*; MP-MIP: *M.*

paraintracellulare–M. indicus pranii. **b** Strategy of the experimental procedure.

Fig. 2 Identification of the essential and growth-defect-associated genes across all

nine *M. intracellulare* strains used in this study. a Functional categories of 131 genes

identified as universal essential or growth-defect-associated with an HMM analysis. The

genes were categorized according to the Kyoto Encyclopedia of Genes and Genomes

(KEGG) database. **b** Universal essential or growth-defect-associated genes corresponding

to the genes of existing antituberculous drug targets. **c** The number of the essential or

growth-defect-associated genes in the accessory genome, and the number of strain-

dependent essential or growth-defect-associated genes in the core genome.

Fig. 3 Detection of genes showing increased or reduced gene essentiality in the

clinical *M. intracellulare* strains. Left panel shows the genes identified as having fewer

Tn insertion reads than the type strain ATCC13950. The fold changes in the number of

Tn insertion reads calculated by a resampling analysis are represented by the color scale.

Red squares indicate genes required for hypoxic pellicle formation in ATCC13950[9].

*Genes identified with a combination of resampling and HMM analyses. †Genes

identified only with resampling analysis. Right panel shows the genes identified as having

more Tn insertion reads than type strain ATCC13950. NH: no homolog with ATCC13950,

NS: no significant increase in Tn insertion reads by a resampling analysis (adjusted $P <$

0.05).

Fig. 4 Overview of the differences in gene essentiality between the clinical _M._

intracellulare strains and ATCC13950, drawn with Cytoscape[55]. Each central node

represents the functional category of genes, assigned with a KEGG pathway analysis.

Each peripheral node represents the genes showing significant changes in the number of

Tn insertion reads. The size of each node represents the number of strains identified. Each

edge represents a strain identified as showing significant changes in the number of Tn

insertion reads. The thickness of each edge represents the adjusted P value. The color

scale of each edge represents the fold change in the number of Tn insertion reads

calculated by a resampling analysis. The graphical organization was referenced to the

previous publication by Carey, _et al_[8].

Fig. 5 Preferential gene requirements for gluconeogenesis in the clinical *M.*

intracellulare **strains inferred from the TnSeq results. a** Carbohydrate pathway

showing the changes in Tn insertion reads. Genes identified with the resampling

analysis are framed by squares. The essentiality of gluconeogenesis-related genes (*pckA,*

glpX) was higher and that of glycolysis-related genes (*aceE, lpdC*) was lower in the

clinical *M. intracellulare* strains compared with those in ATCC13950. **b** Data on gene

essentiality calculated with a resampling analysis in the clinical *M. intracellulare* strains

compared with ATCC13950. Red indicates a reduction in Tn insertion reads in the

clinical *M. intracellulare* strains. Blue indicates an increase of Tn insertion reads in the

clinical *M. intracellulare* strains. Asterisks indicate statistical significance of $\log_2$ fold

changes ($\log_2$FCs). The graphical organization was referenced to the previous

publication by Carey, *et al*[8].

Fig. 6 Significance of increased gene essentiality profiles *in vitro* on the

pathogenesis of mouse lung infection. a Consistency between the genes required for

infection in mouse lungs and the 175 genes required for hypoxic pellicle formation in

ATCC13950. **b** Changes in the requirements for genes with increased essentiality

observed in the clinical *M. intracellulare* strains after the infection of mouse lungs.

Highlighted genes are also required for hypoxic pellicle formation by ATCC13950.

Gene essentiality of the strain itself: gene essentiality compared with that of

ATCC13950. Gene requirements in mouse lungs: changes in the gene requirements in

the infected mouse lungs compared with those before infection. Gene set enrichment

analysis: The genes listed as core enrichment are shown as "Yes" and the genes not

listed as core enrichment are shown as "No."

Fig. 7 Comparison of the timing of entry into logarithmic growth and logarithmic

growth rate by the clinical *M. intracellulare* strains and ATCC13950. a Time at the

inflection point (midpoint) on the sigmoid growth curve. *significantly earlier than an

aerobic culture of ATCC13950; #significantly earlier than a hypoxic culture of

ATCC13950; †significantly later than an aerobic culture of ATCC13950; ‡significantly

later than a hypoxic culture of ATCC13950. **b** Growth rate at midpoint of the growth

curve in each strain. #significantly slower than a hypoxic culture of ATCC13950;

†significantly slower than an aerobic culture of the corresponding strain; ‡significantly

faster than an aerobic culture of ATCC13950. Open bars: aerobic; closed bars: 5% O_2.

Data are shown as the means ± SD of triplicate experiments. Data from one experiment

representative of three independent experiments ($N = 3$) are shown.

Fig. 8 Validation of the inhibition of bacterial growth caused by the suppression of

the expression of TnSeq-hit genes using the CRISPR-i system. Open bar: day 3;

closed bar: day 7. **a** Comparative growth rates of the knockdown strains relative to those

of the vector control strains in the representative universal essential or growth-defect-

associated genes: *glcB, inhA*, *gyrB* and *embB*. **b** Comparative growth rates of the

knockdown strains relative to those of the vector control strains in the representative

accessory and strain-dependent essential or growth-defect-associated genes: *pckA*, *glpX*,

csd and ESX-5 type VII secretion components. Data are shown as the means ± SD of

triplicate experiments. Data from one experiment representative of two independent

experiments ($N = 2$) are shown.

a

| | TMI group |
| MP-MIP group |

0.300

b

Prepare Tn mutant libraries

Clinical MAC-PD strains	Reference strain
M.i.198 M001 M021 M.i.27 M003 MOTT64 M018 M019	ATCC13950

TnSeq

Detect universal and accessory/strain-dependent essential or growth- defect-associated genes among clinical MAC-PD strains and ATCC13950 by an Hidden Markov Model (HMM) analysis

Compare gene essentiality between clinical MAC-PD strains and ATCC13950 by HMM and resampling analyses (The annotation for the clinical strains has been adapted from the ATCC13950.)

Detect genes showing significantly reduced or increased Tn insertion reads

Detect genes required for mouse lung infection and evaluate the impact of genes showing increased gene essentiality in clinical MAC-PD strains compared to ATCC13950 on mouse lung infection

Validate the gene essentiality by CRISPR-i for the representative universal and accessory/strain-dependent shared essential or growth- defect-associated genes and for the genes showing significantly reduced Tn insertion reads in clinical MAC-PD strains compared to ATCC13950

Fig. 1

a

Not Included in Pathway (3 genes)

Signaling and cellular processes (10 genes)

Genetic information processing (39 genes)

Metabolism (79 genes)

Metabolism: glyoxylate, purine/pyrimidine, amino acid, lipid, arabinogalactan/peptidoglycan, porphyrin

Genetic information processing: DNA replication, transcription, amino acid tRNA ligases, ribosomal proteins, general Sec proteins

Signaling and cellular processes: ABC transporter, type VII secretion system, iron-sulfur cluster assembly

Not Included in Pathway: hypothetical proteins

b

Gene locus	gene	function	target drug
OCU_RS25020, OCU_RS25025	*gyrB, gyrA*	DNA topoisomerase subunit B, intein-containing DNA gyrase subunit A	fluoroquinolones
OCU_RS26150	*embB*	arabinosyltransferase	ethambutol
OCU_RS40185	*inhA*	enoyl[acyl-carrier-protein] reductase FabI	isoniazid/ethionamid
OCU_RS45570	*alr*	alanine racemase	cycloserine
OCU_RS46225	*rpoC*	DNA-directed RNA polymerase subunit beta'	rifampicin
OCU_RS38100	*glcB*	malate synthase G	malate synthase inhibitors
OCU_RS32840, OCU_RS32845, OCU_RS32850	*atpA, atpG, atpB*	F_0F_1 ATP synthase subunit alpha, gamma, beta	bedaquiline

c

Number of genes

Universal essential or growth-defect-associated genes

Strain-dependent essential or growth-defect-associated genes

Accessory essential or growth-defect-associated genes

Non-essential genes

Accessory genome 222

Core genome 131 / 439

Fig. 2

Tn decrease in clinical strains

A total of 121 genes were detected as significantly reduced Tn in clinical MAC-PD strains compared to ATCC13950. Of those, 9 genes were commonly hit in 3 to 5 clinical strains.

Tn increase in clinical strains

A total of 162 genes were detected as significantly increased Tn insertion compared to ATCC13950. Of those, 12 genes were commonly hit in more than 7 clinical MAC-PD strains.

Fig. 3

Fig. 4

a

b

Log$_2$ (fold-change)

-6.42   +4.67

Fig 5.

a

b

Locus_tag	Gene annotation	Gene essentia-lity of strain itself	M.i.27 Gene require-ments in mouse lungs	Gene set enrichi-ment analysis	Gene essentia-lity of strain itself	M.i.198 Gene require-ments in mouse lungs	Gene set enrichi-ment analysis
OCU_RS48660	Phoshoenolpyruvate carboxykinase PckA	NC	UP	Yes	UP	NC	No
OCU_RS30540	Fructose-1,6-bisphosphatase GlpX	NC	UP	Yes	NC	NC	Yes
OCU_RS40305	Cysteine desulfurase Csd	NC	UP	Yes	UP	NC	No
OCU_RS38345	TypeVII secretion protein EccC5	NC	UP	Yes	UP	NC	Yes
OCU_RS38275	TypeVII serine protease MycP5	NC	UP	Yes	UP	NC	No
OCU_RS35955	Glycine cleavage system aminomethyltransferase GcvT	NC	UP	Yes	UP	NC	Yes
OCU_RS30185	Ppx/GppA family phosphatase	NC	NC	Yes	NC	NC	Yes
OCU_RS35795	Bifunctional RNaseH/acid phosphatase	NC	NC	No	NC	NC	Yes
OCU_RS41540	Class I SAM-dependent RNA methyltransferase	UP	NC	No	NC	UP	Yes
OCU_RS47290	Hypothetical protein	UP	NC	Yes	NC	NC	Yes

Fig. 6

Fig. 7

Fig. 8