Passive accumulation of alkaloids in putatively non-toxic frogs challenges paradigms of the origins of acquired chemical defenses

Rebecca D. Tarvin[1,*], Jeffrey L. Coleman[2,3], David A. Donoso[4,5], Mileidy Betancourth-Cundar[6], Karem López-Hervas[7], Kimberly S. Gleason[8], J. Ryan Sanders[8], Jacqueline M. Smith[8], Santiago R. Ron[9], Juan C. Santos[10], Brian E. Sedio[2,3], David C. Cannatella[2,*], and Richard Fitch[8,*]

Affiliations

[1] Museum of Vertebrate Zoology and Department of Integrative Biology, University of California, Berkeley, Berkeley, CA 94720 USA

[2] Department of Integrative Biology and Biodiversity Collections, University of Texas at Austin, Austin, TX

[3] Smithsonian Tropical Research Institute, Apartado 0843–03092, Balboa, Ancón, Republic of Panama

[4] Grupo de Investigación en Ecología Evolutiva en los Trópicos (EETROP), Universidad de las Américas, Quito, Ecuador

[5] Ecological Networks Lab, Technische Universität Darmstadt, Darmstadt, Germany

[6] Departamento de Ciencias Biológicas, Universidad de los Andes, Bogotá, Colombia, 111711

[7] Max Planck Institute for Evolutionary Biology, Plön, Germany 24306

[8] Department of Chemistry and Physics, Indiana State University, Terre Haute, IN 47809, USA

[9] Museo de Zoología, Escuela de Biología, Facultad de Ciencias Exactas y Naturales, Pontificia Universidad Católica del Ecuador, Quito, Ecuador

[10] Department of Biological Sciences, St John's University, NY, USA 11439

*Corresponding authors: rdtarvin@berkeley.edu, catfish@utexas.edu, Richard.Fitch@indstate.edu

Abstract

Understanding the origins of novel, complex phenotypes is a major goal in evolutionary biology. Poison frogs of the family Dendrobatidae have evolved the novel ability to acquire alkaloids from their diet for chemical defense at least three times. However, taxon sampling for alkaloids has been biased towards colorful species, without similar attention paid to inconspicuous ones that are often assumed to be undefended. As a result, our understanding of how chemical defense evolved in this group is incomplete. Here we provide new data showing that, in contrast to previous studies, species from each undefended poison frog clade have measurable yet low amounts of alkaloids. We confirm that undefended dendrobatids regularly consume mites and ants, which are known sources of alkaloids. Further, we confirm the presence of alkaloids in two putatively non-toxic frogs from other families. Our data suggest the existence of a phenotypic intermediate between toxin consumption and sequestration—passive accumulation—that differs from active sequestration in that it involves no derived forms of transport and storage mechanisms yet results in low levels of toxin accumulation. We discuss the concept of passive accumulation and its potential role in the origin of chemical defenses in poison frogs and other toxin-sequestering organisms.

Keywords: toxin sequestration, toxin resistance, bioaccumulation, novelty, adaptive landscape, toxicokinetics

1. Introduction

Complex phenotypes can evolve by leveraging phenotypic plasticity in existing traits with concerted change in developmental modules [1]. However, the evolutionary trajectory that animals take to traverse an adaptive landscape from one phenotype to another may be difficult to reconstruct given that they often must cross or avoid adaptive valleys, which include phenotypes that are not always readily observed in populations. Nevertheless, phenotype diversity can help us unravel origins of novel traits and reveal the physiological trade-offs associated with their evolutionary trajectory.

Acquired chemical defenses, or the ability to sequester and use chemicals from the environment against predators or parasites, is one complex phenotype whose evolutionary history has proved difficult to characterize [2,3]. How is it that animals transition from consuming to sequestering toxins? The following phases are likely to occur: 1) consistent exposure to a toxic compound; 2) prior existence or evolution of some resistance to the toxin; 3) change in the elimination rate of the compound that may lead to its prolonged retention, hereafter "passive accumulation" (see Fig. 1); 4) co-option of molecular pathways to transport and store the compound in a specific location, hereafter "active sequestration", which may in turn select for enhanced resistance. Note that while we focus on the processes underlying toxin resistance and sequestration, other phenotypes and selection pressures such as conspicuous coloration or predators may influence these patterns [4]. In the following text we use the terms alkaloid and toxin interchangeably, although the toxicity of each compound is not always known or very straightforward [5]. Similarly, for simplicity we broadly bin species into putatively aposematic (high alkaloid content and conspicuous coloration) and putatively undefended (low or zero alkaloid content and usually lacking conspicuous coloration) categories.

Figure 1. Major processes involved in the transition from the undefended to defended phenotype: 1) toxin intake, here visualized with two discrete points representing low and high rates; 2) toxin elimination rate (Elim. Rate), e.g., via toxin metabolism; 3) toxin sequestration rate (Seq. Rate), i.e., the active transport of toxins for storage in a specific location such as the skin; and 4) toxin accumulation rate (Acc. Rate), or the rate at which toxins are accumulated in the animal. Defense phenotypes are ultimately a result of how these processes interact over time, here arbitrarily from 0 h, immediately after toxin ingestion, to 24 h following ingestion. Although toxin intake influences the total possible amount of toxin accumulation, it cannot fully explain the defensive phenotype. We hypothesize that the "no accumulation" phenotype is characterized by the absence of any ability to sequester toxins in combination with a high rate of elimination, resulting in 0 toxin accumulation (solid grey line); this phenotype is a likely ancestral state for many animals. In contrast, we hypothesize that "passive accumulation" is characterized by lower elimination rates than the no accumulation phenotype, leading to a low amount of toxin accumulation (solid yellow line); however, some mechanisms of toxin transport could also exist, in which case a low sequestration rate could result in a passive accumulation phenotype when elimination rate is high (dashed yellow line). We hypothesize that the "active sequestration" phenotype evolves from an intermediate passive accumulation phenotype through the addition of novel sequestration mechanisms that result in high sequestration rates (solid purple line). However, elimination rates could still modulate the amount of toxins ultimately accumulated, with lower elimination rates resulting in a higher proportion of toxin accumulation overall (dashed purple line).

| | **Process** | | | |
Accumulation Phenotype	Elim. Rate	Seq. Rate	Acc. Rate	Line Type
No accumulation	+++	0	0	—
Passive accumulation	++	0	+	— (yellow)
	+++	+	+	--- (yellow)
Active sequestration	++	+++	++	— (purple)
	+	+++	+++	--- (purple)

We use new data from poison frogs (Anura: Dendrobatidae) to shed some light on this complex
transition. Approximately 100 of the ~340 dendrobatid poison frog species [6] fall into three
conspicuously colored and alkaloid-sequestering (aposematic) clades: *Ameerega, Epipedobates,* and
Dendrobatinae; the other ~240 species compose several other primarily inconspicuously colored clades
that for the most part have been assumed to lack alkaloid defenses: i.e., all Aromobatinae (e.g.,
Allobates, Rheobates, and *Aromobates*), *Colostethus, Silverstoneia, Leucostethus,* and all Hyloxalinae
(*Hyloxalus*) (Fig. 2). According to the phylogenetic placement of these traits, poison frogs have evolved
sequestration of lipophilic alkaloids from consumed arthropods at least three times [7], making them an
ideal group to study complex phenotypic transitions. Much of the research on poison frogs has focused
on changes in diet (toxin intake) in the origins of chemical defenses [8–11] without much focus on rate
of toxin elimination versus accumulation (Fig. 2; but see [12,13]). However, rates of intake,
sequestration, and elimination all shape the ability of an animal to accumulate a compound (Fig. 1).
Thus, characterizing the metabolism and sequestration of alkaloids in defended and undefended
dendrobatid lineages is essential to understand the origins of chemical defense [14]. We propose that
changes in toxin metabolism through selection on mechanisms of toxin resistance likely play a major
role in the evolution of toxicity.

Figure 2. Evolutionary models of toxin sequestration in Dendrobatidae have changed over time. A) When several
species of aposematic dendrobatids (purple lines) were found to have narrower dietary niches than undefended
dendrobatids and other frogs [10,15,16], researchers hypothesized that diet specialization may have driven the
radiation of aposematic dendrobatids [17]. **B)** Chemists hypothesized that aposematic dendrobatids sequester
dietary alkaloids via an alkaloid uptake system [11]. Daly [18] postulated that an alkaloid uptake system was
present in the ancestor of Dendrobatidae (here denoted as passive accumulation) and that it is "overexpressed" in
aposematic dendrobatids (here denoted as active sequestration). **C)** A phylogenetic analysis of Dendrobatidae
revealed that aposematism and diet specialization evolved independently several times [9]. The new information

helped generate the diet-toxicity hypothesis, which posits that shifts from a generalist to a specialist diet drove the
multiple origins of alkaloid uptake through enhanced resistance and/or more efficient sequestration systems [4,8].
D) Here we propose a combination of these hypotheses, i.e., that passive accumulation, alkaloid consumption, and
some level of alkaloid resistance was present in an early dendrobatid lineage; enhanced resistance and active
sequestration mechanisms then arose later, resulting in the chemical defense phenotype. This hypothesis places
less emphasis on dietary changes and more strongly emphasizes novel molecular mechanisms (e.g., binding
proteins and target-site insensitivity [19–21]). Phylogenies in each subpanel highlight how increasing resolution
impacted our understanding of phenotypic diversification in Dendrobatidae. All images of frogs were taken by RDT.

2. Results and Discussion

(a) Phases 1 and 2: Consistent exposure to toxins may select for resistance and sequestration

Many animals occasionally or frequently consume toxins, and a multitude have evolved toxin resistance.
Some invertebrate pests resist pesticides [22–25], many insect herbivores resist plant toxins [26,27],
some predators resist toxic prey [28], many animals resist environmental pollutants [29], and toxic
organisms usually evolve resistance to their own defenses ("autoresistance") [3,30].

The general mechanisms of toxin resistance are toxin avoidance, toxin metabolism, and target
modification [31]. If an animal does not or cannot *avoid* toxin exposure, it will need to *survive* exposure
using toxin metabolism or target modification mechanisms such as biotransformation, elimination,
alternative targets, and target-site resistance (see [31] for more details). Toxin metabolism, also known
as toxicokinetics [32], is a set of mechanisms based on detoxification pathways than may provide toxin
resistance. These pathways are common to all animals and were likely used by the ancestors of most if
not all animals that eventually evolved toxin sequestration (Fig. 1).

The type of toxin resistance present in an animal may eventually affect that animal's ability or
propensity to sequester toxins. For example, animals that possess target-site resistance may be more
likely to evolve toxin sequestration than animals that avoid toxins [31]. Although one might expect that
toxin metabolism may also prevent toxin sequestration, the ability to metabolize toxins can in some
cases augment toxin defenses [33], increase the toxicity of a compound (e.g., pumiliotoxin to
allopumiliotoxin in the poison frogs *Adelphobates galactonotus*, *A. castaneoticus*, *Dendrobates auratus*,
and *D. tinctorius* [34,35]), or result in some amount of passive accumulation through increased toxin
exposure [33,36]. In general, toxin-sequestering animals often have specialized mechanisms of toxin
resistance when compared to non-toxic relatives [31]. For example, three amino acid replacements in
the ATPα protein evolved in association with cardenolide sequestration in Danainae butterflies [36,37]
and predatory fireflies that sequester lucibufagins have ATPα gene duplications that enhance lucibufagin
resistance [38].

In dendrobatids, mechanisms of toxin resistance are still understudied [39]. Target-site resistance to
some alkaloids appears to have evolved in several toxic clades and in some non-toxic species [20,21].
Some toxic species also appear to have alternative target mechanisms including binding proteins like
alpha-binding globulin [19] and saxiphillin [40] that might prevent alkaloids from accessing their
molecular targets. Accumulation of alkaloids in skin glands could help to prevent alkaloids from reaching
their targets. Although direct evidence is lacking, some poison frogs may biotransform alkaloids into less
toxic forms until they can be eliminated from the body, e.g., using cytochrome p450s [41].

(b) Phases 3 and 4: Passive accumulation and active sequestration in poison frogs

Although the inconspicuously colored clades of poison frogs have long been considered to lack chemical

defenses (i.e., they are undefended), many species have not been comprehensively assessed. By
reviewing existing data from inconspicuous poison frog species, we found that of the 245 inconspicuous
species described to date [6], only 30 have been assessed for toxicity, sometimes using methods that
would not necessarily detect lipophilic alkaloids (Table S1). Available data suggest that at least nine of
these species might have alkaloids: *Allobates femoralis*, *Al. kingsburyi*, *Al. zaparo*, *Hyloxalus maculosus*,
H. nexipus, H. vertebralis, Leucostethus fugax, Paruwrobates erythromos, and *Silverstoneia punctiventris*
[4,8,42,43]. However, evolutionary studies have not fully incorporated these data (Fig. 1, Table S1, and
see below).

We tested for possible alkaloid presence in a broad selection of inconspicuously colored poison frog
lineages using GCMS. In total, we surveyed 89 animals representing 30 species of Neotropical frogs
including 27 dendrobatid poison frogs and representatives from most of the major undefended clades in
Dendrobatidae (Table 1). As far as we are aware, we provide alkaloid data for the first time for seven
undefended species (*Rheobates palmatus, Allobates juanii, H. shuar, H.* sp. Agual Azul*, H. toachi, S.* aff.
gutturalis, and *S. erasmios*) and one defended species (*Epipedobates* sp. 1). Overall, we detected
alkaloids in skins from 12 of 13 undefended species included in our study, although often with less
diversity and relatively lower quantities than in aposematic lineages (Fig. 3, Table 1, Table S2, Table S3).
We find substantially higher diversities of alkaloids in aposematic dendrobatid species than previously
reported [5,42,44–46], and expand knowledge on major classes of alkaloids within genera. Because
chemical standards for most poison frog alkaloids do not exist, it is not possible to provide absolute
quantification of alkaloids. Reported values are in units of integrated area, which do not directly
correspond to alkaloid quantity because of differences in ion yield. Nevertheless, qualitative
comparisons of integrated areas can provide insight into how species differ in degrees of magnitude.

Table 1. Range and median of alkaloid quantity (estimated by the sum of integrated areas) and alkaloid diversity
(number of different compounds) by species. The presumed chemical defense phenotype for poison frogs is given
according to Santos and Cannatella [4]. Purple rows highlight toxic species.

Family	Subfamily	Species	Phenotype	Sample Size (frogs)	Log (Total Integrated Area)		Alkaloid Number	
					Range	Median	Range	Median
Dendrobatidae	Aromobatinae	*Rheobates palmatus*	undefended	4	13.07 – 14.24	14.00	1 – 4	1.5
Dendrobatidae	Aromobatinae	*Allobates insperatus*	undefended	8	13.47 – 15.44	14.99	1 – 9	5.0
Dendrobatidae	Aromobatinae	*Allobates juanii*	undefended	1	14.10	14.10	1	1.0
Dendrobatidae	Aromobatinae	*Allobates kingsburyi*	undefended	1	13.63	13.63	2	2.0
Dendrobatidae	Aromobatinae	*Allobates talamancae*	undefended	3	14.89 – 16.27	15.09	2 – 4	3.0
Dendrobatidae	Aromobatinae	*Allobates zaparo*	undefended	1	16.78	16.78	8	8.0
Dendrobatidae	Colostethinae	*Leucostethus fugax*	undefended	8	12.57 – 15.33	14.00	3 – 8	4.5
Dendrobatidae	Colostethinae	*Ameerega bilinguis*	aposematic	1	21.97	21.97	133	133.0
Dendrobatidae	Colostethinae	*Ameerega hahneli*	aposematic	4	20.21 – 22.29	21.68	85 – 140	128.5
Dendrobatidae	Colostethinae	*Silverstoneia* aff. *gutturalis*	undefended	9	11.80 – 17.33	15.40	1 – 10	3.0
Dendrobatidae	Colostethinae	*Silverstoneia erasmios*	undefended	2	14.70 – 16.11	15.41	15 – 15	15.0
Dendrobatidae	Colostethinae	*Epipedobates* aff. *espinosai*	aposematic	2	18.44 – 20.20	19.32	83 – 131	107.0
Dendrobatidae	Colostethinae	*Epipedobates anthonyi*	aposematic	1	20.54	20.54	127	127.0
Dendrobatidae	Colostethinae	*Epipedobates boulengeri*	aposematic	2	18.87 – 19.39	19.13	77 – 94	85.5
Dendrobatidae	Colostethinae	*Epipedobates* sp. 1	aposematic	2	19.49 – 19.68	19.59	99 – 105	102.5
Dendrobatidae	Colostethinae	*Epipedobates espinosai*	aposematic	2	18.82 – 21.33	20.08	85 – 146	115.5
Dendrobatidae	Colostethinae	*Epipedobates machalilla*	aposematic	2	12.98 – 15.67	14.32	8 – 38	23.0
Dendrobatidae	Colostethinae	*Epipedobates tricolor*	aposematic	2	18.36 – 19.07	18.72	91 – 114	102.5
Dendrobatidae	Hyloxalinae	*Hyloxalus awa*	undefended	7	0.00 – 16.05	13.58	0 – 12	3.0
Dendrobatidae	Hyloxalinae	*Hyloxalus shuar*	undefended	1	14.92	14.92	5	5.0
Dendrobatidae	Hyloxalinae	*Hyloxalus* sp. Agua Azul	undefended	1	14.30	14.30	8	8.0
Dendrobatidae	Hyloxalinae	*Hyloxalus toachi*	undefended	2	0.00 – 0.00	0.00	0 – 0	0.0
Dendrobatidae	Dendrobatinae	*Phyllobates aurotaenia*	aposematic	4	17.72 – 21.08	18.88	48 – 118	67.5

Dendrobatidae	Dendrobatinae	*Dendrobates truncatus*	aposematic	3	20.05 – 23.95	20.42	111 – 172	115.0
Dendrobatidae	Dendrobatinae	*Oophaga sylvatica*	aposematic	5	22.86 – 24.85	23.76	152 – 189	175.0
Dendrobatidae	Dendrobatinae	*Andinobates fulguritus*	aposematic	2	20.09 – 20.51	20.30	80 – 85	82.5
Dendrobatidae	Dendrobatinae	*Andinobates minutus*	aposematic	4	16.57 – 18.77	18.07	34 – 80	66.0
Bufonidae		*Amazophrynella siona*	NA	2	14.12 – 14.40	14.26	1 – 1	1.0
Bufonidae		*Atelopus* aff. *spurrelli*	NA	1	11.58	11.58	4	4.0
Leptodactylidae	Leptodactylinae	*Lithodytes lineatus*	NA	2	0.00 – 0.00	0.00	0 – 0	0.0

For Aromobatinae, we surveyed the undefended genera *Rheobates* and *Allobates.* Alkaloids were
detected in all four *R. palmatus* individuals sampled, with one individual having at least four classes of
compounds represented (4,6-disubstituted quinolizidines, 3,5-disubstituted indolizidines, 3,5-
disubstituted pyrrolizidines, and unclassified). We found that five species of *Allobates* all had detectable
levels of alkaloids. *Allobates insperatus* had a relatively high level of alkaloid diversity, with at least
eighteen alkaloids from nine classes detected, and at least one class found in each of the eight sampled
individuals. In contrast, only one unclassified alkaloid was identified in a single individual of *Allobates*
juanii while two were found in one individual of *Allobates kingsburyi*. At least two alkaloids were
identified in each of the three sampled individuals of *Allobates talamancae* (including the lehmizidine
277A and five new alkaloids). Eight alkaloids were identified in the single surveyed *Allobates zaparo*
individual (including the spiropyrrolizidines **222-1** and **222-2** as well as six unclassified alkaloids). Prior
assessments using thin-layer chromatography suggested the presence of alkaloids in three *Al. kingsburyi*
[4], but none in twelve *Al. insperatus* [8]. Four studies (Table S1) failed to identify any alkaloids in
Allobates talamancae. Allobates zaparo was shown to possibly have trace alkaloids, although the
interpretation of these data was absence of alkaloids [8]. There are no known aposematic species from
this subfamily, although we note conflicting evidence on the presence of alkaloids in *Allobates femoralis*
[42,47–49] (Table S1).

Within Colostethinae, we surveyed individuals from two undefended clades, *Leucostethus* and
Silverstoneia, and from two aposematic clades, *Epipedobates* and *Ameerega.* From *Leucostethus fugax*,
we identified a total of twelve 5-substituted indolizidine, 5,6-disubstituted indolizidine, pyrrolidine,
spiropyrrolizidine, and unclassified alkaloids (**196A**, **225C**, **222-1**, **222-2**, and eight new alkaloids), with
three to eight unique compounds detected in each of the eight sampled individuals. Our data are
consistent with prior thin-layer chromatography data showing that *Leucostethus fugax* tested positive
for skin compounds [4], though prior interpretation of these data were different (Table S1). We also
surveyed two species of *Silverstoneia*. We found alkaloids in all nine *S.* aff. *gutturalis*, with a total of
fourteen alkaloids identified across seven classes (**196A**, **223I**, **233A**, **235B**, **237U**, three isomers of
239AB, two isomers of **239CD**, and four new alkaloids). In just two individuals of *S. erasmios,* we
detected a total of 26 alkaloids, including some pumiliotoxins (**325B**, **323B**) and pyrrolizidines (**225C**).
While *S. erasmios* and *S.* aff. *gutturalis* had not been surveyed for alkaloids previously, a study detected
13 alkaloids in *S. punctiventris* [43]. In addition, we conducted a highly sensitive, untargeted analysis
(UHPLC-HESI-MS/MS) of *S. flotator*, which revealed that alkaloids were present in all 12 sampled
individuals (>99% alkaloid pathway probability; Table S6). At this probability level, we found 67 alkaloids
including one quinolizidine, two pyridines, and an analog of epibatidine (Tables S5 and S6).

In terms of the aposematic clades of Colostethinae that we sampled, most of the individual skins of
Epipedobates and *Ameerega* contained dozens to more than one hundred unique alkaloids (see Table S3
for full details). For *Ameerega*, we surveyed 5 individuals representing 2 species, all of which had
integrated areas that were more than 75,000x compared to individuals of its sister clade, *Leucostethus*
fugax (Table 1). Similarly, alkaloid diversity was 10–20x greater in *Ameerega* than in *Leucostethus.*
Histrionicotoxins and decahydroquinolines were considered previously to be the dominant alkaloid
classes in genus *Ameerega* [50]; here we also found high levels of indolizidines (Fig. 3). Patterns for

Epipedobates as compared to sister genus *Silverstoneia* were similar, although less extreme. We
surveyed 13 individuals representing 7 species in *Epipedobates* and identified at least 370 alkaloids,
which contrasts with studies using a less sensitive method (thin-layer chromatography) that found
mixed evidence for the presence of alkaloids in *E.* aff. *espinosai* (then referred to as *E. boulengeri*) and *E.*
machalilla [4,8]. However, the quantity and diversity of alkaloids in *E. machalilla* was substantially lower
than in other *Epipedobates* species, occurring at levels similar to *Silverstoneia* spp. (Table 1, Fig. 3).
Except for *E. machalilla*, each *Epipedobates* species had about 10x higher quantities and diversities of
alkaloids compared to members of *Silverstoneia.* We found trace levels of epibatidine in *Epipedobates*
anthonyi but not in other *Epipedobates* species. Epibatidine and its analogs have also been detected in
E. espinosai, Ameerega silverstonei, *S. flotator* [51], this study], and *Ameerega petersi* or a closely
related, undescribed species (reported as *Dendrobates pictus* from Loreto, Peru in [42], but see
taxonomic revision by [52])*.

Within Hyloxaline, a generally undefended clade, we surveyed four species of *Hyloxalus*, three of which
had detectable levels of alkaloids. We identified seventeen different alkaloids in *H. awa* (**197D**, **197H**,
199B, **217B**, **221P**, **223AB**, **231A**, **231C**, **247E**, and eight previously undescribed alkaloids), with the seven
sampled individuals having zero to twelve alkaloids each. We detected five alkaloids in a single individual
of *H. shuar* (**197D**, **199B**, **237G**, and two isomers of **239K**) and eight alkaloids in a single individual of *H.*
sp. Agua Azul (**195C**, **197D**, **199B**, **251K**, and four new alkaloids). Our detection of low levels of alkaloids
in *H. awa* are consistent with the observations that avian predators consume *H. awa* [53]. No alkaloids
were detected in two individuals of *H. toachi*, the only undefended species from which we failed to
detect alkaloids*.

According to the most recent phylogenetic reconstructions [7], the sister clade to Hyloxalinae is
Dendrobatinae. Dendrobatinae contains exclusively (or near exclusively) toxic species. From this
subfamily, we surveyed 18 individuals representing 5 species. We identified a total of 187 unique
alkaloids from four *Phyllobates aurotaenia*, 316 alkaloids from five *Oophaga sylvatica*, and 213 alkaloids
from three *Dendrobates truncatus*. These three species are all relatively large poison frogs (snout-vent
lengths 20–35 mm; Table S2), which may in part explain their high alkaloid diversities and quantities. In
Andinobates minutus and *Andinobates fulguritus*, which are members of the same subfamily but are
much smaller in size (11–15 mm; Table S2), we detected 129 and 109 alkaloids, respectively. Three of
the *Andinobates minutus* individuals were juveniles. The total alkaloid quantities (integrated areas) in *D.*
truncatus and *O. sylvatica* were comparable to those of *Ameerega* but were higher than quantities
detected in *Epipedobates*. We also report for the first time, to the best of our knowledge, the
occurrence of N-Methyldecahydroquinolines outside the genus *Ameerega* (in *E.* aff. *espinosai*, *E.* sp. 1, *S.*
aff. *gutturalis*, *Andinobates minutus*, *Andinobates fulguritus*, *D. truncatus*, *O. sylvatica*, *P. aurotaenia*,
and *Allobates insperatus*) [50]. The ability to N-methylate DHQ (demonstrated experimentally in
Adelphobates galactonotus [12]) may thus be conserved in dendrobatids, or, non-exclusively, arthropod
sources of the alkaloid class (likely myrmicine ants [54]) are widespread.

Figure 3. From left to right: an ultrametric tree showing phylogenetic relationships inferred previously [59] among sampled species with the three aposematic poison frog clades highlighted in purple, the undefended clades in dark gray, and non-dendrobatids in light gray (Bufonidae: *Amazophrynella siona* and *Atelopus* aff. *spurrelli*; Leptodactylidae: *Lithodytes lineatus*). Tile color indicates the log of the total quantity of alkaloids in each class as measured by the sum of integrated areas of alkaloids of that class from GCMS data per individual. The number in each tile indicates the number of alkaloids (including isomers) detected in each individual for each class. On the right are prey items recovered from the stomach of each individual, colored by arthropod group and scaled to 1

(total number of prey identified are shown under N). Note the large proportion of ants (Formicidae, dark purple)
and mites (Acari, light purple) in many of the individuals compared to other prey types. See Table S3 for alkaloid-
level data and Table S4 for raw diet data. Poison frog genera names are abbreviated as follows: *All., Allobates*;
Ame., Ameerega; *And., Andinobates*; *D., Dendrobates*; *E., Epipedobates*; *H., Hyloxalus*; *Le., Leucostethus*; *O.,*
Oophaga; *P., Phyllobates*; *R., Rheobates*; *S., Silverstoneia*; Alkaloid class abbreviations are based on [50,60] and are
as follows: HTX, histrionicotoxins; PTX, pumiliotoxins; PTXB, Pumiliotoxin B; aPTX, allopumiliotoxins; DeoxyPTX,
deoxypumiliotoxins; hPTX, homopumiliotoxins; deoxy-hPTX, deoxy-homopumiliotoxins; DHQ,
decahydroquinolines; NMeDHQ, N-Methyldecahydroquinolines; HO-DHQ, hydroxy-decahydroquinolines; 3,5-P,
3,5-disubstituted pyrrolizidines; HO-3,5-P, hydroxy-3,5-disubstituted pyrrolizidines; 5-I, 5-substituted indolizidines;
3,5-I, 3,5-disubstituted indolizidines; 5,6-I, 5,6-disubstituted indolizidines; 5,8-I, 5,8-disubstituted indolizidines;
Dehydro-5,8-I, Dehydro-5,8-Indolizidines; 5,6,8-I, 5,6,8-trisubstituted indolizidines; HO-5,6,8-I, Hydroxy-5,6,8-
trisubstituted indolizidines; 1,4-Q, 1,4-disubstituted quinolizidines; 4,6-Q, 4,6-disubstituted quinolizidines; 3,5-Q,
3,5-disubstituted quinolizidines; 1,3,4-Q, 1,3,4-trisubstituted quinolizidines; Lehm, lehmizidines; Epiquinamide,
epiquinamide; 2-Pyr, 2-substituted pyrrolidine; 3-Pyr, 3-substituted pyrrolidine; 2,5-Pyr, 2,5-disubstituted
pyrrolidines; Pyr, pyrrolizidine of indeterminate substitution; 2,6-Pip, 2,6-disubstituted piperidines; Pip, other
piperidines; Pyri, pyridines (including epibatidine); GTX, gephyrotoxins; Tricyclic, coccinelline-like tricyclics; SpiroP,
spiropyrrolizidines; Necine, unspecified necine base; Unclass, unclassified alkaloids without known structures.

Outside of Dendrobatidae, we detected a new unclassified alkaloid, New159, in each of two
Amazophrynella siona (Bufonidae) and four alkaloids in one individual of *Atelopus* aff. *spurrelli* (Anura:
Bufonidae; 3,5-disubstituted pyrrolizidine **237R**-1, decahydroquinoline **243A**-3, 5,8-disubstituted
indolizidine **251B**-2, and an unclassified alkaloid, New267-2). To the best of our knowledge, the
detection of a decahydroquinoline and a 3,5-disubstituted pyrrolizidine in a bufonid frog other than
Melanophryniscus [55] is novel and may provide useful context for understanding the evolution of
chemical defense in the family. We detected no alkaloids in two *Lithodytes lineatus* (Leptodactylidae)
individuals, which is surprising because *Lithodytes lineatus* has been hypothesized to be a Müllerian
mimic of poison frogs, though the composition of its chemical defenses may be primarily proteinaceous
[56]. These findings are also interesting in light of the fact that *Lithodytes lineatus* live and breed in ant
colonies using chemical signals that provide camouflage [57]. In addition, we identified 55 alkaloids in 3
sampled individuals of the frog *Eleutherodactylus cystignathoides* (Eleutherodactylidae), 40 of which
were likely identical to compounds identified in *S. flotator* according to our analyses (Tables S5, S6). A
few other species of *Eleutherodactylus* frogs from Cuba are also known to have alkaloids [58]. Thus,
these patterns suggest that some alkaloids may be widespread byproducts of frog metabolism or that
passive accumulation may occur more generally in frogs. Our data do not allow us to discriminate
between these possibilities.

Dietary data from these same specimens point to the ubiquity of mites and ants in dendrobatid diets,
and possibly more generally in other leaf-litter dwelling frogs (Fig. 3; see below). This finding in concert
with the detection of low levels of alkaloids in the lineages that putatively lack chemical defenses leads
us to hypothesize that dietary shifts are not sufficient to explain the presence or absence of the
chemical defense phenotype within Dendrobatidae or possibly in other families (Bufonidae,
Eleutherodactylidae). The total amount of alkaloids accumulated is a result of multiple rates including
toxin intake, toxin elimination, and toxin sequestration (Fig. 1) – not just intake alone. Answers to the
following questions would help further evaluate the relative roles of diet versus sequestration
mechanisms in the evolution of toxin sequestration in frogs.

1) Is total alkaloid intake lower in undefended lineages? If so, this would imply that behavioral or
environmental changes affect diet and impact the defensive phenotype. Several of the lipophilic
alkaloids found in dendrobatid frogs have been traced to arthropod sources, specifically mites [61], ants
[62], and beetles [63], but the amount and diversity of alkaloids in each arthropod prey type is generally
unknown. Shifts in diet content towards a higher proportion of ants and mites have been hypothesized

to play an important role in the origin of chemical defense in poison frogs [4,8]. We quantified gut
contents for the same individuals that we analyzed by GCMS and found that both undefended and
aposematic species consume a large proportion of ants and mites (Fig. 3; Table S4). Although the
aposematic clades tend to consume proportionally more ants and mites, as found in other studies, the
undefended lineages do consume a high proportion of ants and mites. Other data support this general
pattern: ants and mites constituted up to 51% and 60% of the stomach contents of the undefended
species *Allobates talamancae* [64] and *H. sauli* [8]*, respectively. Ants and mites compose nearly 50% of
the arthropods (36 and 10%, respectively) found in the *S. flotator* stomachs we analyzed (Table S4).
Sympatric populations of the undefended *H. awa* and aposematic *E. espinosai* (formerly *E.*
darwinwallacei [65]) are both diet specialized, with the former consuming mostly ants and beetles and
the latter consuming mostly mites and springtails [66]. In a lab experiment, the aposematic species *D.*
tinctorius was shown to prefer fruit fly larvae over ants when given the choice [67], suggesting that even
in aposematic species, consumption of possible alkaloid-containing prey is not necessarily a preference.
One study found that *O. sylvatica* alkaloid quantity is inversely correlated with ant and mite stomach
contents; however, this species consumed more mites and ants than sympatric *H. infraguttatus* [46].
Although in many cases aposematic species consume more mites and ants than undefended species, the
undefended lineages clearly consume toxic prey items, and some of the consumed alkaloids reach the
skin (Fig. 3). In sum, the available data do not strongly support that changes in diet alone are sufficient
to explain differences in alkaloid skin quantities.

2) Is the rate of toxin elimination faster in undefended lineages? Faster elimination would imply that
toxin metabolism impacts the defensive phenotype. Only a few studies have reviewed toxin metabolism
and elimination in dendrobatids, and none provided data for non-toxic species. Nevertheless, the
available data from aposematic species show species-level variation and plasticity in the metabolism and
elimination of alkaloids. *Epipedobates anthonyi, R. variabilis,* and *R. imitator* accumulate more than
twice as much ingested epibatidine compared to *P. vittatus* and *D. tinctorius* [68]. *Oophaga sylvatica* and
D. tinctorius upregulate detoxification genes such as cytochrome p450s upon alkaloid consumption
[35,41]. *Adelphobates galactonotus* sequesters the alkaloids HTX and DHQ less efficiently at higher
doses [12]. Some species metabolically alter the structure of alkaloids: *A. galactonotus*, *A. castaneoticus*,
D. auratus, and *D. tinctorius* can hydroxylate pumiliotoxin **251D** [34,35], making it more toxic (to mice);
A. galactonotus can also N-methylate DHQ [12]. These studies indicate that alkaloid elimination rate and
metabolism varies among aposematic species, but not enough information exists to infer much about
elimination rates in undefended lineages. Given that undefended lineages consume alkaloids yet show
much lower levels of alkaloids in the skin (Fig. 2), we hypothesize that their toxin elimination rates are
faster than in aposematic lineages (Fig. 1).

3) Are active sequestration mechanisms (Fig. 1) unique to chemically defended species, or can they also
be found in undefended ones? This would imply that the efficiency of sequestration mechanisms
impacts the defensive phenotype. Little is known regarding the mechanisms of toxin sequestration in
poison frogs or in other toxin-sequestering animals. An alkaloid-binding globulin was recently
characterized in the poison frog *O. sylvatica* [19]. While plasma assays demonstrated that the
aposematic species *O. sylvatica*, *E. tricolor*, and *D. tinctorius* can bind and sequester a PTX-like
photoprobe, plasma from the undefended *Allobates femoralis* showed no binding activity. In addition,
the evolutionarily distant mantellid species *Mantella aurantiaca*, which also sequesters alkaloids, did not
show binding activity. These data hint at variation in molecular mechanisms for alkaloid uptake across
lineages [19], which may be tuned to availability of specific alkaloids in each species' diet. One GCMS
analysis did not detect alkaloids in the skins of *Allobates talamancae* and *C. panamansis* after they
consumed fruit flies dusted with 5,8-disubstituted indolizidine **209B**, decahydroquinoline **195A**, and
histrionicotoxin **285C** for five weeks [11]. Other unpublished data suggest an inability by brightly colored

H. azureiventris to accumulate alkaloids (identities not reported) from fruit flies, though the frogs
apparently accumulated alkaloids dissolved in a methanol-saline solution [69]. Sparteine, a quinolizidine
structurally similar to epibatidine, was detected in *Allobates femoralis* skin after being fed sparteine-
dusted fruit flies for over a month [47].

Additional data on potential uptake mechanisms in dendrobatids exist for benzocaine, a synthetic
lipophilic compound that is used for anesthesia and euthanasia in amphibians and has a structure similar
to poison frog alkaloids. Benzocaine is readily taken up orally to the skin in the aposematic poison frog
D. auratus, the non-toxic ranid frog (*Lithobates clamitans*), and the alkaloid-sequestering bufonid toad
Melanophryniscus moreirae [48]. Although the same amount of benzocaine was injected into each frog,
twice as much benzocaine was recovered from *D. auratus* than *L. clamitans* and three times as much
was recovered from *M. moreirae* (see their Fig. 2)*,* suggesting that lipophilic compound uptake occurs
without specialized mechanisms of sequestration in *L. clamitans* (e.g., possibly "passive sequestration")
but that *D. auratus* and *M. moreirae* likely have distinct active sequestration mechanisms that result in
much higher levels of benzocaine retention.

Although more data will be necessary to evaluate phylogenetic patterns of active sequestration
mechanisms, these data suggest that active sequestration mechanisms might be absent in undefended
lineages, and that sequestering species differ substantially in their ability to actively transport and store
specific compounds (Fig. 1).

(c) Predictions arising from the passive accumulation hypothesis

Data from this and other studies indicate that nearly all dendrobatids consume alkaloid-containing prey
and species vary in their ability to clear alkaloids. Some species appear to lack specific transport and
storage mechanisms for consumed alkaloids ("active sequestration"), yet they have detectable levels of
alkaloids in their skin; we refer to this phenotype as "passive accumulation" and suggest that it is an
evolutionary intermediate between toxin consumption (with no sequestration) and sequestration (Fig.
1). We predict that the ancestral state of poison frogs (and potentially other clades with alkaloid-
sequestering species, such as *Melanophryniscus* and *Eleutherodactylus*) is alkaloid consumption and low
levels of alkaloid resistance, accompanied by passive alkaloid accumulation. Interestingly, we also
detected small amounts of alkaloids in two species of bufonid toads and one eleutherodactylid (but not
in a leptodactylid), suggesting that passive accumulation may be present in an even older ancestor.
Importantly, our concept of passive accumulation requires no major evolutionary innovations, only the
tweaking of metabolic efficiency and/or toxin intake, along with the ability to survive consuming certain
toxins (Fig. 2). Passive accumulation would also be expected to result in the diffusion of alkaloids across
many tissues, rather than concentration of alkaloids within a specific tissue. Desorption electrospray
ionization mass spectrometry imaging data indicate that alkaloids diffuse across various tissues in the
aposematic dendrobatid *Dendrobates tinctorius* immediately following intake, possibly an evolutionary
trace of the low elimination rates that may have initial evolved in an ancestor with the passive
accumulation phenotype [13].

Toxin resistance is associated with toxin sequestration in dendrobatid poison frogs [20,21]. Although
available data supports the presence of target-site resistance in some but not all poison frogs, we
anticipate that some alkaloid resistance evolved in the ancestor of Dendrobatidae or in an even older
ancestor, but is yet to be described [3,8] (Fig. 1D). Such resistance may be difficult to characterize using
the comparative method if it involves mutations of small effect [70] or pleiotropic processes. Regardless,
it is clear that all or nearly all dendrobatid poison frogs consume some amount of alkaloid-containing
arthropods [4,8,10] (Fig. 3) and they do not appear to suffer substantially from doing so (e.g. [35,47,68]).
While little to no adaptation appears necessary to passively accumulate lipophilic alkaloids, additional

adaptive changes are likely necessary to more efficiently clear or accumulate alkaloids. New research is
beginning to identify major molecular players involved in this process [19].

Passive accumulation of toxins is not a novel concept, as it has been discussed previously in terms of
self-medication [71,72] and bioaccumulation (e.g., of environmental pollutants [32]), and it is also
conceptually analogous to some medical treatments in humans (e.g., chemotherapy). Any organism that
consumes something toxic might simultaneously suffer from toxin exposure yet benefit from the
compound's effect on disease, infection, parasites, or predators. For example, in the presence of
parasitoids, *Drosophila suzukii* flies preferentially lay their eggs on the insecticide atropine, which
protects them from being parasitized but prolongs development [73]. Mechanisms that likely underlie
passive accumulation may also be analogous to key organismal functions. For example, humans
accumulate vitamin E in the liver [74] and use a transfer protein abundant in liver cells to shuttle the
vitamin into the plasma where it becomes bioavailable [75]. The transition from passive accumulation to
active sequestration in poison frogs may also rely on overexpression of genes whose encoded proteins
bind to and transport alkaloids [41] (Fig. 1B). Because most poison-frog alkaloids are fat-soluble, the
passive diffusion of alkaloids, perhaps using fat-storage mechanisms, could have evolved with few
changes to the ancestral physiological machinery.

In sum, for toxin-resistant organisms, there is little cost to accumulating a toxin, yet there may be
benefits in doing so. If toxin accumulation is both low-cost and beneficial, then slow toxin elimination
rates could evolve quite readily, resulting in passive accumulation. Two recent studies support the idea
that some toxin resistance permits toxin intake and results in passive accumulation. In one, nicotine-
resistant *Drosophila melanogaster* fruit flies that were fed nicotine accumulated measurable amounts of
the toxin in their bodies, more than nicotine-sensitive flies [33]. In another study, ouabain-resistant *D.
melanogaster* flies that were fed ouabain accumulated measurable amounts of ouabain in their bodies,
more than ouabain-sensitive flies [36]. In a more extreme scenario, cardenolide defense in milkweed
butterflies may not rely on any active forms of toxin transport or storage, but rather is simply a result of
a high rate of toxin intake relative to toxin clearance [76]. Two cardenolide-resistant species (*Danaus
plexippus* and *D. gilippus*) accumulate the cardenolides in the midgut and store the compounds in the
hemolymph as well as integument and adhering tissues. In contrast, a susceptible, non-sequestering
species (*Euploea core*) appears to degrade and clear cardenolides. In another case, the aphid *Athalia
rosae* shows constant turnover of its glucosinolate toxins, suggesting that they cannot effectively store
glucosinolates, yet their metabolic clearing is inefficient enough that they still maintain a high level of
toxins in the hemolymph [77]. It is conceivable, then, that in some cases, accumulation of defensive
chemicals results from a mechanism that enables high net toxin intake, followed by passive entry into
the bloodstream and long-term storage in tissues.

Are these cases of active sequestration? Under our definition they are not, given that these species do
not actively transport and store these compounds, as far as we know. Rather, these organisms merely
fail to efficiently metabolize and eliminate these compounds, leading to their temporary diffusion in
certain tissues that provides a transient benefit against parasites or predators. Evidence for this "passive
accumulation" phenotype as an intermediate stage on the path towards toxin sequestration is scarce,
but passive accumulation is a pervasive pattern in studies of ecological toxicology and may be more
common in toxin-sequestering lineages than we currently know.

(d) Limitations

Our study presents a novel alkaloid dataset for dendrobatid frogs and some relatives, yet it is limited in
the following ways. For some species we only sampled one or two individuals, which may paint an
incomplete picture of toxin diversity and quantity in the group. Poison frogs vary substantially over time

and seasons in their alkaloid profiles [78], yet we did not conduct serial sampling. Standards are
unavailable for most frog alkaloids and thus we could not measure absolute quantity. Relative
quantitation was performed based on integration of the extracted ion chromatogram of the base peak
for each alkaloid for maximum sensitivity and selectivity. The nature of these data mean that qualitative
comparisons may be meaningful but quantitative comparisons across alkaloid structures could be
misleading, especially given our small sample sizes for some species. Therefore, we refrained from
conducting additional quantitative analyses of integrated area data. Finally, batrachotoxin and
tetrodotoxin are too heavy to study using GCMS; we cannot exclude the possibility that they occur in the
sampled species.

3. Conclusion

The large-scale evolutionary transition from consuming to sequestering toxins has occurred in a plethora
of invertebrates [79] and vertebrates [30]. Here we provide new evidence showing that undefended
poison frogs and frogs in closely related families (Bufonidae, Eleutherodactylidae) contain measurable
amounts of alkaloids. We confirm that they consume some amount of toxic arthropod prey. We propose
that passive accumulation of consumed alkaloids is an ancestral state in the group, and possibly in
related taxa, and that selection acted on the efficiency of toxin elimination and sequestration to result in
chemical defense. Future studies of the kinetics of alkaloids in different tissues of both aposematic and
undefended poison frogs will provide further insight into these putative intermediate evolutionary
steps.

4. Methods

(a) Field collection

In the case of *Silverstoneia flotator* and *Eleutherodactylus cystignathoides*, animals were collected and
euthanized with benzocaine in 2022 in Gamboa, Panama (9.136, -79.723) and in 2024 in Austin, Texas,
USA (30.285, -97.736). Dorsal and ventral skins were removed and placed separately in ~1-mL MeOH in
1-dram glass vials for UHPLC-HESI-MS/MS analyses (see below). For all other species, animals were
collected in 2014 and euthanized with an overdose of lidocaine. Whole skins were removed and placed
in ~1-mL MeOH in glass vials with PTFE-lined caps. Stomachs of all species were removed and placed in
95% ethanol.

Instruments and dissection surfaces were cleaned with 95% ethanol between dissections. Although
contamination across samples is possible, it is unlikely to invalidate the identification of alkaloids in
undefended species based on the following patterns. 1) At several sites, we only sampled undefended
species, and these individuals were found to contain alkaloids (e.g., Las Brisas – *Rheobates palmatus*, El
Valle – *Silverstoneia* aff. *gutturalis*, and Santa Maria – *Hyloxalus* sp. Agua Azul); i.e. these cannot possibly
have come from contamination by aposematic species. 2) In one site where we collected both
undefended and aposematic species, the undefended species show no alkaloids (Lita – *H. toachi*); i.e.,
the preparation of both types does not imply cross-contamination of samples. 3) At two sites where the
undefended species were prepared on a different day from the aposematic species (Valle Hermoso – *H.
awa* and *E. boulengeri*; Canelos – *L. fugax* and *A. hahneli*) and could not have been cross-contaminated,
the undefended species still show evidence of alkaloids. 4) All chromatograms in the sequence and
integration data were inspected manually. Peaks with low areas or following samples with high areas
and subject to carryover were excluded from further analysis. 5) Data from Panama collected by a
different team using different methods also identify alkaloids in an undefended dendrobatid (*S.*

flotator).

(b) Alkaloid identification and quantification

(b) Alkaloid identification and quantification

For samples from Ecuador and Colombia, a 100-μL aliquot of the MeOH was sampled from each vial and
transferred to a 200-μL limited volume insert and analyzed directly by GC-MS. The system used was a
Thermo AS-3000 autosampler interfaced to a Trace GC Ultra interfaced to a iTQ 1100 ion trap mass
spectrometer autotuned with FC-43 (PFTBA) operating in positive ion mode. AS conditions were as
follows: 2 pre-wash cycles of 5 μL MeOH, then 3 plunger strokes and withdrawal of 1.00 μL sample with
1-μL air gap, injection with no pre- or post-injection dwell followed by 3 post wash cycles of 5 μL MeOH.
GC conditions were as follows: splitless injection, splitless time 1.00 min with surge (200 kPa for 0.70
min, to sharpen early peaks), split flow 50 mL/min; injector temperature 250C, oven temperature
program 100C for one minute, then ramped at 10C/min to 280C and held 10 min; transfer line
temperature 300C. MS conditions were as follows: for electron ionization (EI), collection mode profile, 1
microscan, 25 μsec max ion time, range 35–650 μ, source temperature 250 C, solvent delay 3.00 min,
source voltage 70 eV; for chemical ionization (CI), reagent gas NH3 (1.8 mL/min). Samples for CI were
run in ddMS2 mode (3 precursor ions) with 1 microscan, 50 ms max ion time, 0.450 μ precursor width
and dynamic exclusion duration 0.2 min.

EI spectra were compared with published data [51,60,80] to identify class and likely ID. A set of known
standards was run to give accurate retention times across the range of alkaloids and normalized to
literature data using linear regression. Sample retention times were then normalized, and molecular
weights were obtained from CI MS1 spectra. These were then directly compared to archival Daly GC-MS
data where possible. CI MS2 spectra were also used where possible to confirm functional groups such as
alcohols by loss of water, etc. Kovats retention indexes (semi-standard nonpolar) are also provided
based on retention times and published indexes for background silicone impurities. Accuracy of index
assignments were confirmed based on fatty acid methyl esters from skin lipids present in extracts.
Epibatidine coelutes with the lipid methyl palmitoleate and the latter caused a number of false positives
in the GC-MS data. We thus reviewed LC-HRMS data at the known elution time relative to a known
standard. Epibatidine was only found in one sample in trace quantities and is marked as such.

Samples from Panama and Texas were extracted on separate occasions, then filtered and run in tandem
with ultra-high-performance liquid-chromatography heated-electrospray-ionization tandem mass
spectrometry (UHPLC-HESI-MS/MS), following an untargeted metabolomics profile, with conditions
optimized specifically for retention and subsequent identification of alkaloids [81]. Briefly, for extraction,
methanol was evaporated and skins were homogenized with stainless steel beads in a TissueLyser
QIAGEN^{TM} and resuspended in 1800 μL of extraction solvent (9:1 MeOH: pH 5 water). Samples were
then extracted for 3 hr at 4˚C in a ThermoMixer (Eppendorf US, Enfield, CT, USA), followed by
evaporation of the methanol component with a speedvac concentrator (Thermo Fisher Scientific,
Waltham, MA, USA). Next, samples were freeze-dried with a lyophilizer overnight and resuspended in
500 μL extraction solvent. Resuspended samples were then filtered, diluted 1:7 in 100% MeOH, and
analyzed using UHPLC-HESI-MS/MS on a Thermo Vanquish LC and QExactive quadrupole-orbitrap MS.
Instrumental methods were identical to those described by [81]. A positive reference of 1 μg/μL ≥98%
(±)-epibatidine dihydrochloride hydrate (Sigma-Aldrich, St. Louis, MO, USA) was included in the run.

Following UHPLC-HESI-MS/MS, chromatographic data were processed using MZmine 3 (v3.9.0) [82],
applying a stringent MS1 noise threshold parameter >100000, as used by other workers (e.g., [81]). We
did not use a gap filling algorithm, a step often used in analysis of chemically homogeneous datasets to
backfill overlooked metabolite occurrences, so as to avoid the creation of false positive metabolite
observations. MZmine 3 assigns chromatographic features to putative compounds based on molecular

mass and retention time. MZmine 3 feature tables and MS2 data were then uploaded to the Global
Natural Products Social Molecular Networking (GNPS) platform [83] for Feature-Based Molecular
Networking [84]. We used SIRIUS [85] and CSI:FingerID [86] to infer molecular formulae and predict
structures including the elements H, C, N, O, P, and S. CANOPUS was used to classify metabolites [87],
following the ClassyFire [88] and NPClassifier molecular taxonomies [89]. Only compounds assigned to
the alkaloid pathway with an NPClassifier pathway probability score >99% were retained in the feature
table; epibatidine (the positive reference) was among the compounds recovered at this confidence level.
This untargeted metabolomics approach yielded correct annotations for epibatidine at the levels of
most specific class ("epibatidine analogues": ClassyFire) and class and superclass ("pyridine alkaloids"
and "nicotinic alkaloids": NPClassifier). As expected, the compound was detected only in the positive
reference sample.

(c) Diet identification

556 Stomach content was inspected under a stereoscope and all prey items identified to order (or family, in
557 the case of Formicidae). Given the low sample sizes in many individuals, we did not conduct statistical
558 comparisons of diet composition across species.

(d) Analyses

We summarized and plotted data from Ecuadorian and Colombian samples in R v4.3.1 [90] using the
packages *ggplot2* [91], *cowplot* v1.1.1 [92], and *dplyr* v1.1.2 [93]. Samples from Panama and Texas were
analyzed using a different instrument that has higher sensitivities to detect more diverse compounds
but lower retention-time resolution, as well as untargeted analytical methods, reducing confidence in
structural inferences. Therefore, data are not directly comparable, and they could not be included in
Figure 3. Phylogenies were subsetted from [59] using *ape* v5.7.1 [94] and *phytools* v1.9.16 [95]. Any
compounds known to co-elute with other compounds were unable to be identified, so we averaged
their quantities across the co-eluting compounds. Corrections for mass were not included; we instead
opted to provide data from full skins.

Acknowledgements. We thank Fray Arriaga, Josué Collins (STRI, Panama), Cristian Florez-Pai (FELCA,
Colombia), Valentina Gómez-Bahamón, Camilo Isaza (Cafam, Colombia), Roberto Márquez, Daniel
Nastacuaz, Pablo Palacios-Rodriguez, Andrea Paz, Santiago Vega, and many others for their assistance in
the field. We thank the communities of Laguna de Cube (Esmeraldas) and Laguna de San Pedro
(Orellana) in Ecuador for their support and efforts towards conserving local ecosystems. We also thank
Kameron T. Bell, Nicholas R. Andreasen, and Megan M. Reid for their help acquiring and processing
GCMS data. Finally, JLC would like to thank Raineldo Urriola, Isis Ochoa, Lil Marie Camacho, Zurenayka
Alain, Roberto Ibáñez, Roberto Cambra, Roberto Borrell, Rivieth De Liones, Félix Rodriguez, and Orelis
Arosemena for their warmth, generosity, and logistical support during his time at STRI.

Funding. RDT was supported by an NIH MIRA (1R35GM150574), start-up funding from University of
California Berkeley, and grants from the Society of Systematic Biologists, North Carolina Herpetological
Society, Society for the Study of Reptiles and Amphibians, Chicago Herpetological Society, Texas
Herpetological Society, the EEB Program at University of Texas at Austin, National Science Foundation
Graduate Research Fellowship Program Graduate Research Opportunities Worldwide (in partnership
with USAID), and a 2014 National Geographic Young Explorer Grant. Additional support to DCC, RWF,
JLC, and RDT was provided by NSF DBI-1556967, and BES received support from Stengl-Wyer
Endowment Grant SWG-22-01. RWF and students KSG, JMS, and JRS were supported by NSF DUE-
0942345, NSF CHE-1531972, and NSF IOS-1556982.

Author contributions. RDT: conceptualization, data curation, formal analysis, funding acquisition, investigation, methodology, project administration, resources, supervision, visualization, writing-original draft, writing-review & editing; JLC: data curation, formal analysis, funding acquisition, investigation, resources, writing-original draft, writing-review & editing; SRR, MBC, KLH, JCS: investigation, resources, writing-review & editing; DD: data curation, investigation, resources, writing-review & editing; KSG, JMS, JRS: data curation, investigation, writing-review & editing; BES: formal analysis, funding acquisition, investigation, resources; DCC: funding acquisition, investigation, project administration, resources, supervision, writing-original draft, writing-review & editing; RWF: data curation, formal analysis, funding acquisition, investigation, methodology, project administration, resources, supervision, visualization, writing-original draft, writing-review & editing. All authors gave final approval for publication and agreed to be held accountable for the work described herein.

Ethics. Collection was performed under permits (COL: Res. 1177 at Universidad de los Andes) and Contrato Marco Acceso a los Recursos Genéticos Nro. 005-14 IC-FAU-DNB/MA (Ecuador) as well as Ministerio de Ambiente Permiso de Colecta Científica No. ARBG-0038-2022 and Permiso de Transferencia de Material Genético y/o Biológico No. PA-01-ARG-096-2022 (Panama). The animal use protocols were approved by the University of Texas at Austin (IACUC AUP-2012-00032) and the Smithsonian Tropical Research Institute (SI-22017). Voucher specimens are deposited in the Museo de Zoología (QCAZ) de Pontificia Universidad Católica del Ecuador (PUCE) and the Museo de Historia Natural C.J. Marinkelle (ANDES) at the Universidad de los Andes in Bogotá, Colombia.

Use of Artificial Intelligence (AI) and AI-assisted technologies. No AI or AI-assisted technologies were used in the preparation of this manuscript.

Data accessibility. The datasets supporting this article have been uploaded as part of the supplementary material. GCMS and LCMS data are available at the Global Natural Product Social Molecular Networking (GNPS) (accession numbers pending). Other raw data are available here as supplementary tables.

Supplementary Information

Table S1. A summary of data available on alkaloid detection in "non-toxic" lineages of poison frogs.

Table S2. Collection localities, specimen numbers, size, sex, and summary of alkaloid quantities and diversity for each individual.

Table S3. Alkaloid-level data for every individual analyzed by GCMS.

Table S4. Stomach content data for every individual.

Table S5. List of the subset of classes and most specific classes of compounds in *Silverstoneia flotator* annotated as alkaloids ("Alkaloid Pathway" of NPClassifier) at >99% probability and data on presence/absence of each in *Eleutherodactylus cystignathoides*.

Table S6. Feature table with details about determined chemical properties, annotations, and presence/absence information for a) *Silverstoneia flotator* skin alkaloids, and b) identifying information for run numbers listed in Table S6a columns.

References

1. West-Eberhard MJ. 2003 *Developmental plasticity and evolution*. 1st edn. New York: Oxford
University Press.

2. Berenbaum MR. 1995 The chemistry of defense: theory and practice. *Proc. Natl. Acad. Sci. U. S. A.*
92, 2–8.

3. Santos JC, Tarvin RD, O'Connell LA. 2016 A review of chemical defense in poison frogs
(Dendrobatidae): ecology, pharmacokinetics, and autoresistance. In *Chemical Signals in Vertebrates*
13 (eds BA Schulte, TE Goodwin, MH Ferkin), pp. 305–337. Switzerland: Springer International
Publishing.

4. Santos JC, Cannatella DC. 2011 Phenotypic integration emerges from aposematism and scale in
poison frogs. *Proc. Natl. Acad. Sci. U. S. A.* **108**, 6175–6180.

5. Lawrence JP, Rojas B, Blanchette A, Saporito RA, Mappes J, Fouquet A, Noonan BP. 2023 Linking
predator responses to alkaloid variability in poison frogs. *J. Chem. Ecol.* **49**, 195–204.

6. AmphibiaWeb. 2024 https://amphibiaweb.org/ University of California, Berkeley, CA, USA.
Accessed 14 March 2024.

7. Santos JC, Baquero M, Barrio-Amorós C, Coloma LA, Erdtmann LK, Lima AP, Cannatella DC. 2014
Aposematism increases acoustic diversification and speciation in poison frogs. *Proc. R. Soc. Lond. B*
Biol. Sci. **281**, 20141761.

8. Darst CR, Menéndez-Guerrero PA, Coloma LA, Cannatella DC. 2005 Evolution of dietary
specialization and chemical defense in poison frogs (Dendrobatidae): A comparative analysis. *Am.*
Nat. **165**, 56–69.

9. Santos JC, Coloma LA, Cannatella DC. 2003 Multiple, recurring origins of aposematism and diet
specialization in poison frogs. *Proc. Natl. Acad. Sci. U. S. A.* **100**, 12792–12797.

10. Toft CA. 1995 Evolution of diet specialization in poison-dart frogs (Dendrobatidae). *Herpetologica*
51, 202–216.

11. Daly JW, Secunda SI, Garraffo HM, Spande TF, Wisnieski A, Cover JF Jr. 1994 An uptake system for
dietary alkaloids in poison frogs (Dendrobatidae). *Toxicon* **32**, 657–663.

12. Jeckel AM *et al.* 2022 Dose-dependent alkaloid sequestration and N-methylation of
decahydroquinoline in poison frogs. *J. Exp. Zool. A Ecol. Integr. Physiol.* **337**, 537–546.

13. Jeckel AM, Matsumura K, Nishikawa K, Morimoto Y, Saporito RA, Grant T, Ifa DR. 2020 Use of
whole-body cryosectioning and desorption electrospray ionization mass spectrometry imaging to
visualize alkaloid distribution in poison frogs. *J. Mass Spectrom.* **55**, 1–6.

14. Gonzalez M, Carazzone C. 2023 Eco-Metabolomics Applied to the Chemical Ecology of Poison Frogs
(Dendrobatoidea). *J. Chem. Ecol.* (doi:10.1007/s10886-023-01443-0)

15. Toft CA. 1980 Feeding ecology of thirteen syntopic species of anurans in a seasonal tropical
environment. *Oecologia* **45**, 131–141.

16. Toft CA. 1981 Feeding ecology of Panamanian litter anurans: Patterns in diet and foraging mode. *J.*
Herpetol. **15**, 139–144.

17. Caldwell JP. 1996 The evolution of myrmecophagy and its correlates in poison frogs (Family
Dendrobatidae). *J. Zool.* **240**, 75–101.

18. Daly JW. 1998 Thirty years of discovering arthropod alkaloids in amphibian skins. *J. Nat. Prod.* **61**, 162–172.

19. Alvarez-Buylla A *et al.* 2023 Binding and sequestration of poison frog alkaloids by a plasma globulin. *Elife* **12**, e85096.

20. Tarvin RD, Santos JC, O'Connell LA, Zakon HH, Cannatella DC. 2016 Convergent substitutions in a sodium channel suggest multiple origins of toxin resistance in poison frogs. *Mol. Biol. Evol.* **33**, 1068–1081.

21. Tarvin RD, Borghese CM, Sachs W, Santos JC, Lu Y, O'Connell LA, Cannatella DC, Harris RA, Zakon HH. 2017 Interacting amino acid replacements allow poison frogs to evolve epibatidine resistance. *Science* **357**, 1261–1266.

22. Andreev D, Kreitman M, Phillips TW, Beeman RW, ffrench-Constant RH. 1999 Multiple origins of cyclodiene insecticide resistance in *Tribolium castaneum* (Coleoptera: Tenebrionidae). *J. Mol. Evol.* **48**, 615–624.

23. Daborn PJ *et al.* 2002 A single p450 allele associated with insecticide resistance in *Drosophila*. *Science* **297**, 2253–2256.

24. Chiu T-L, Wen Z, Rupasinghe SG, Schuler MA. 2008 Comparative molecular modeling of *Anopheles gambiae* CYP6Z1, a mosquito P450 capable of metabolizing DDT. *Proc. Natl. Acad. Sci. U. S. A.* **105**, 8855–8860.

25. Richard H, Carroll L. 2013 The molecular genetics of insecticide resistance. *Genetics* **194**, 807–815.

26. Dobler S, Petschenka G, Pankoke H. 2011 Coping with toxic plant compounds - The insect's perspective on iridoid glycosides and cardenolides. *Phytochemistry* **72**, 1593–1604.

27. Agrawal AA, Petschenka G, Bingham RA, Weber MG, Rasmann S. 2012 Toxic cardenolides: Chemical ecology and coevolution of specialized plant-herbivore interactions. *New Phytol.* **194**, 28–45.

28. Arbuckle K, Rodríguez de la Vega RC, Casewell NR. 2017 Coevolution takes the sting out of it: Evolutionary biology and mechanisms of toxin resistance in animals. *Toxicon* **140**, 118–131.

29. Whitehead A, Clark BW, Reid NM, Hahn ME, Nacci D. 2017 When evolution is the solution to pollution: Key principles, and lessons from rapid repeated adaptation of killifish (*Fundulus heteroclitus*) populations. *Evol. Appl.* **10**, 762–783.

30. Savitzky AH, Mori A, Hutchinson DA, Saporito RA, Burghardt GM, Lillywhite HB, Meinwald J. 2012 Sequestered defensive toxins in tetrapod vertebrates: Principles, patterns, and prospects for future studies. *Chemoecology* **22**, 141–158.

31. Tarvin RD, Pearson KC, Douglas TE, Ramírez-Castañeda V, Navarrete MJ. 2023 The diverse mechanisms that animals use to resist toxins. *Annu. Rev. Ecol. Evol. Syst.* **54**, 283–306.

32. Spurgeon D, Lahive E, Robinson A, Short S, Kille P. 2020 Species sensitivity to toxic substances: evolution, ecology and applications. *Front. Environ. Sci. Eng. China* **8**. (doi:10.3389/fenvs.2020.588380)

33. Douglas TE, Beskid SG, Gernand CE, Nirtaut BE, Tamsil KE, Fitch RW, Tarvin RD. 2022 Trade-offs between cost of ingestion and rate of intake drive defensive toxin use. *Biol. Lett.* **18**, 20210579.

34. Daly JW, Garraffo HM, Spande TF, Clark VC, Ma J, Ziffer H, Cover JF Jr. 2003 Evidence for an enantioselective pumiliotoxin 7-hydroxylase in dendrobatid poison frogs of the genus *Dendrobates*. *Proc. Natl. Acad. Sci. U. S. A.* **100**, 11092–11097.

35. Alvarez-Buylla A, Payne CY, Vidoudez C, Trauger SA, O'Connell LA. 2022 Molecular physiology of
pumiliotoxin sequestration in a poison frog. *PLoS One* **17**, e0264540.

36. Karageorgi M *et al.* 2019 Genome editing retraces the evolution of toxin resistance in the monarch
butterfly. *Nature* **574**. (doi:10.1038/s41586-019-1610-8)

37. Petschenka G, Fandrich S, Sander N, Wagschal V, Boppré M, Dobler S. 2013 Stepwise evolution of
resistance to toxic cardenolides via genetic substitutions in the Na+/K+-ATPase of milkweed
butterflies (Lepidoptera: Danaini). *Evolution* **67**, 2753–2761.

38. Yang L *et al.* 2023 Predatory fireflies and their toxic firefly prey have evolved distinct toxin
resistance strategies. *Curr. Biol.* **33**, 5160-5168.e7.

39. Coleman JL, Cannatella DC. 2023 The molecular basis and evolution of toxin resistance in poison
frogs. *Evol. Ecol.* (doi:10.1007/s10682-023-10258-0)

40. Abderemane-Ali F *et al.* 2021 Evidence that toxin resistance in poison birds and frogs is not rooted
in sodium channel mutations and may rely on "toxin sponge" proteins. *J. Gen. Physiol.* **153**,
e202112872.

41. Caty SN *et al.* 2019 Molecular physiology of chemical defenses in a poison frog. *J. Exp. Biol.* **222**,
jeb204149.

42. Daly JW, Myers CW, Whittaker N. 1987 Further classification of skin alkaloids from neotropical
poison frogs (Dendrobatidae), with a general survey of toxic/noxious substances in the Amphibia.
Toxicon **25**, 1023–1095.

43. Gonzalez M, Palacios-Rodriguez P, Hernandez-Restrepo J, González-Santoro M, Amézquita A,
Brunetti AE, Carazzone C. 2021 First characterization of toxic alkaloids and volatile organic
compounds (VOCs) in the cryptic dendrobatid *Silverstoneia punctiventris*. *Front. Zool.* **18**, 39.

44. Cipriani I, Rivera M. 2009 Detección de alcaloides en la piel de cuatro especies de anfibios
ecuatorianos (Anura: Dendrobatidae). *Rev. Ecuat. Med. Cienc. Biol.* **30**, 42–49.

45. Moskowitz NA *et al.* 2020 Land use impacts poison frog chemical defenses through changes in leaf
litter ant communities. *Neotropical Biodiversity* **6**, 75–87.

46. Moskowitz NA, Alvarez-Buylla A, Morrison CR, Chamba A, Rentería J, Tapia EE, Coloma LA, Donoso
DA, O'Connell LA. 2022 Poison frog diet and chemical defense are influenced by availability and
selectivity for ants. *bioRxiv.* , 2022.06.14.495949. (doi:10.1101/2022.06.14.495949)

47. Sanchez E *et al.* 2019 Transcriptomic signatures of experimental alkaloid consumption in a poison
frog. *Genes* **10**, 733.

48. Saporito RA, Grant T. 2018 Comment on Amézquita et al. (2017) "Conspicuousness, color
resemblance, and toxicity in geographically diverging mimicry: The pan-Amazonian frog *Allobates
femoralis* ." *Evolution* , 1009–1014.

49. Amézquita A, Ramos Ó, González MC, Rodríguez C, Medina I, Simões PI, Lima AP. 2017
Conspicuousness, color resemblance, and toxicity in geographically diverging mimicry: The pan-
Amazonian frog *Allobates femoralis*. *Evolution* **71**, 1039–1050.

50. Daly JW, Ware N, Saporito RA, Spande TF, Garraffo HM. 2009 N-methyldecahydroquinolines: An
unexpected class of alkaloids from Amazonian poison frogs (Dendrobatidae). *J. Nat. Prod.* **72**, 1110–
1114.

51. Daly JW, Garraffo HM, Spande TF. 1999 Alkaloids from amphibian skins. In *Alkaloids: Chemical and*

Biological Perspectives (ed SW Pelletier), pp. 1–161. New York: Pergamon.

52.   Guillory WX *et al.* 2020 Phylogenetic relationships and systematics of the Amazonian poison frog
750   genus *Ameerega* using ultraconserved genomic elements. *Mol. Phylogenet. Evol.* **142**, 106638.

751   53.   Darst CR, Cummings ME. 2006 Predator learning favours mimicry of a less-toxic model in poison
752   frogs. *Nature* **440**, 208–211.

753   54.   Jones TH, Gorman JST, Snelling RR, Delabie JHC, Blum MS, Garraffo HM, Jain P, Daly JW, Spande TF.
754   1999 Further alkaloids common to ants and frogs: Decahydroquinolines and a quinolizidine. *J.*
755   *Chem. Ecol.* **25**, 1179–1193.

756   55.   Rodríguez C, Rollins-Smith L, Ibáñez R, Durant-Archibold AA, Gutiérrez M. 2017 Toxins and
757   pharmacologically active compounds from species of the family Bufonidae (Amphibia, Anura). *J.*
758   *Ethnopharmacol.* **198**, 235–254.

759   56.   Prates I, Antoniazzi MM, Sciani JM, Pimenta DC, Toledo F, Haddad FB, Jared C. 2011 Skin glands,
760   poison and mimicry in dendrobatid and leptodactylid amphibians. *Journal of Morphology* **273**, 279–
761   290.

762   57.   de Lima Barros A, López-Lozano JL, Lima AP. 2016 The frog *Lithodytes lineatus* (Anura:
763   Leptodactylidae) uses chemical recognition to live in colonies of leaf-cutting ants of the genus *Atta*
764   (Hymenoptera: Formicidae). *Behav. Ecol. Sociobiol.* **70**, 2195–2201.

765   58.   Rodríguez A, Poth D, Schulz S, Gehara M, Vences M. 2013 Genetic diversity, phylogeny and
766   evolution of alkaloid sequestering in Cuban miniaturized frogs of the *Eleutherodactylus limbatus*
767   group. *Mol. Phylogenet. Evol.* **68**, 541–554.

768   59.   Wan YC *et al.* 2023 Selection on visual opsin genes in diurnal Neotropical frogs and loss of the SWS2
769   opsin in poison frogs. *Mol. Biol. Evol.* **40**. (doi:10.1093/molbev/msad206)

770   60.   Daly JW, Spande TF, Garraffo HM. 2005 Alkaloids from amphibian skin: A tabulation of over eight-
771   hundred compounds. *J. Nat. Prod.* **68**, 1556–1575.

772   61.   Saporito RA, Donnelly MA, Norton R a., Garraffo HM, Spande TF, Daly JW. 2007 Oribatid mites as a
773   major dietary source for alkaloids in poison frogs. *Proc. Natl. Acad. Sci. U. S. A.* **104**, 8885–8890.

774   62.   Saporito RA, Garraffo HM, Donnelly MA, Edwards AL, Longino JT, Daly JW. 2004 Formicine ants: An
775   arthropod source for the pumiliotoxin alkaloids of dendrobatid poison frogs. *Proc. Natl. Acad. Sci.*
776   *U. S. A.* **101**, 8045–8050.

777   63.   Dumbacher JP, Wako A, Derrickson SR, Samuelson A, Spande TF, Daly JW. 2004 Melyrid beetles
778   (*Choresine*): a putative source for the batrachotoxin alkaloids found in poison-dart frogs and toxic
779   passerine birds. *Proc. Natl. Acad. Sci. U. S. A.* **101**, 15857–15860.

780   64.   Mebs D, Yotsu-Yamashita M, Pogoda W, Vargas Alvarez J, Ernst R, Köhler G, Toennes SW. 2018 Lack
781   of alkaloids and tetrodotoxin in the neotropical frogs *Allobates* spp. (Aromobatidae) and
782   *Silverstoneia flotator* (Dendrobatidae). *Toxicon* **152**. (doi:10.1016/j.toxicon.2018.07.027)

783   65.   López-Hervas K, Santos JC, Ron SR, Betancourth-Cundar M, Cannatella DC, Tarvin RD. 2024 Deep
784   divergences among inconspicuously colored clades of Epipedobates poison frogs. *Mol. Phylogenet.*
785   *Evol.* , 108065.

786   66.   Sánchez-Loja S, Donoso DA, Páez-Vacas MI. 2023 Conspicuous and cryptic poison frogs are picky
787   and prefer different meals in syntopy. *Evol. Ecol.* (doi:10.1007/s10682-023-10282-0)

788   67.   Moskowitz NA, D'Agui R, Alvarez-Buylla A, Fiocca K, O'Connell LA. 2022 Poison frog dietary

789  preference depends on prey type and alkaloid load. *PLOS ONE* **17**, e0276331.

790 68. Waters KR, Dugas MB, Grant T, Saporito RA. 2023 The ability to sequester the alkaloid epibatidine is
791  widespread among dendrobatid poison frogs. *Evol. Ecol.* (doi:10.1007/s10682-023-10260-6)

792 69. Saporito RA, Spande TF, Garraffo HM, Donnelly MA. 2009 Arthropod alkaloids in poison frogs: A
793  review of the 'Dietary Hypothesis.' *Heterocycles* **79**, 277–297.

794 70. ffrench-Constant RH, Daborn PJ, Goff GL. 2004 The genetics and genomics of insecticide resistance.
795  *Trends Genet.* **20**, 163–170.

796 71. Clayton DH, Wolfe ND. 1993 The adaptive significance of self-medication. *Trends Ecol. Evol.* **8**, 60–
797  63.

798 72. Singer MS, Mace KC, Bernays EA. 2009 Self-medication as adaptive plasticity: increased ingestion of
799  plant toxins by parasitized caterpillars. *PLoS One* **4**, e4796.

800 73. Poyet M, Eslin P, Chabrerie O, Prud'homme SM, Desouhant E, Gibert P. 2017 The invasive pest
801  *Drosophila suzukii* uses trans-generational medication to resist parasitoid attack. *Sci. Rep.* **7**, 43696.

802 74. Violet P-C *et al.* 2020 Vitamin E sequestration by liver fat in humans. *JCI Insight* **5**.
803  (doi:10.1172/jci.insight.133309)

804 75. Arita M, Sato Y, Miyata A, Tanabe T, Takahashi E, Kayden HJ, Arai H, Inoue K. 1995 Human alpha-
805  tocopherol transfer protein: cDNA cloning, expression and chromosomal localization. *Biochem. J*
806  **306 (Pt 2)**, 437–443.

807 76. Petschenka G, Agrawal AA. 2015 Milkweed butterfly resistance to plant toxins is linked to
808  sequestration, not coping with a toxic diet. *Proc. Biol. Sci.* **282**, 20151865.

809 77. Müller C, Wittstock U. 2005 Uptake and turn-over of glucosinolates sequestered in the sawfly
810  *Athalia rosae*. *Insect Biochem. Mol. Biol.* **35**, 1189–1198.

811 78. Saporito RA, Donnelly MA, Jain P, Martin Garraffo H, Spande TF, Daly JW. 2007 Spatial and
812  temporal patterns of alkaloid variation in the poison frog *Oophaga pumilio* in Costa Rica and
813  Panama over 30 years. *Toxicon* **50**, 757–778.

814 79. Duffey SS. 1980 Sequestration of plant natural products by insects. *Annu. Rev. Entomol.* **25**, 447–
815  477.

816 80. Daly JW, Brown GB, Mensah-Dwumah M, Myers CW. 1978 Classification of skin alkaloids from
817  neotropical poison-dart frogs (Dendrobatidae). *Toxicon* **16**, 163–188.

818 81. Sedio BE *et al.* 2021 Chemical similarity of co-occurring trees decreases with precipitation and
819  temperature in North American forests. *Frontiers in Ecology and Evolution* **9**.
820  (doi:10.3389/fevo.2021.679638)

821 82. Schmid R *et al.* 2023 Integrative analysis of multimodal mass spectrometry data in MZmine 3. *Nat.*
822  *Biotechnol.* **41**, 447–449.

823 83. Wang M *et al.* 2016 Sharing and community curation of mass spectrometry data with Global
824  Natural Products Social Molecular Networking. *Nat. Biotechnol.* **34**, 828–837.

825 84. Nothias L-F *et al.* 2020 Feature-based molecular networking in the GNPS analysis environment. *Nat.*
826  *Methods* **17**, 905–908.

827 85. Dührkop K, Fleischauer M, Ludwig M, Aksenov AA, Melnik AV, Meusel M, Dorrestein PC, Rousu J,
828  Böcker S. 2019 SIRIUS 4: a rapid tool for turning tandem mass spectra into metabolite structure

information. *Nat. Methods* **16**, 299–302.

86. Dührkop K, Shen H, Meusel M, Rousu J, Böcker S. 2015 Searching molecular structure databases
with tandem mass spectra using CSI:FingerID. *Proc. Natl. Acad. Sci. U. S. A.* **112**, 12580–12585.

87. Dührkop K *et al.* 2021 Systematic classification of unknown metabolites using high-resolution
fragmentation mass spectra. *Nat. Biotechnol.* **39**, 462–471.

88. Djoumbou Feunang Y *et al.* 2016 ClassyFire: automated chemical classification with a
comprehensive, computable taxonomy. *J. Cheminform.* **8**, 61.

89. Kim HW *et al.* 2021 NPClassifier: A Deep Neural Network-Based Structural Classification Tool for
Natural Products. *J. Nat. Prod.* **84**, 2795–2807.

90. R Core Team. 2023 *R: A Language and Environment for Statistical Computing*. Vienna, Austria: R
Foundation for Statistical Computing. See https://www.R-project.org/.

91. Wickham H. 2016 ggplot2: Elegant Graphics for Data Analysis.

92. Wilke CO. 2020 *cowplot: Streamlined plot theme and plot annotations for "ggplot2."* See
https://cran.r-project.org/package=cowplot.

93. Wickham H, François R, Henry L, Müller K, Vaughan D. 2023 *dplyr: A grammar of data*
manipulation. See https://cran.r-project.org/package=dplyr.

94. Paradis E, Schliep K. 2019 ape 5.0: an environment for modern phylogenetics and evolutionary
analyses in R. *Bioinformatics*. **35**, 526–528. (doi:10.1093/bioinformatics/bty633)

95. Revell LJ. 2012 phytools: An R package for phylogenetic comparative biology (and other things).
Methods in Ecology and Evolution. **3**, 217–223. (doi:10.1111/j.2041-210X.2011.00169.x)