Title: Y chromosome linked UTY modulates sex differences in valvular fibroblast methylation in response to nanoscale extracellular matrix cues

Short title: UTY alters methylation states in male valve cells

Authors: Rayyan M. Gorashi[1,2], Talia Baddour[1,2], Sarah J. Chittle[1,2], Nicole E. Félix Vélez[1,2], Michaela A. Wenning[3,4], Kristi S. Anseth[5,6], Luisa Mestroni[7,8], Brisa Peña[7,8,9], Peng Guo[10], and Brian A. Aguado[1,2]*

Affiliations:

[1] Shu Chien-Gene Lay Department of Bioengineering, University of California San Diego, La Jolla, CA 92093, USA.

[2] Sanford Consortium for Regenerative Medicine, La Jolla, CA 92037, USA.

[3] Department of Chemical Engineering, Massachusetts Institute of Technology, Cambridge, MA 02139, USA.

[4] Harvard-MIT Program in Health Sciences and Technology, Massachusetts Institute of Technology, Cambridge, MA 02139, USA.

[5] Department of Chemical and Biological Engineering, University of Colorado Boulder, CO 80303, USA.

[6] BioFrontiers Institute, University of Colorado Boulder, CO 80309, USA.

[7] Cardiovascular Institute, University of Colorado Denver Anschutz Medical Campus, School of Medicine, Division of Cardiology, at 12700 E.19th Avenue, Bldg. P15, Aurora, Colorado, 80045, United States.

[8] Department of Medicine, Division of Cardiology, University of Colorado Anschutz Medical Campus, at 12700 E.19th Avenue, Bldg. P15, Aurora, Colorado, 80045, United States.

23   [9] Bioengineering Department, University of Colorado Denver Anschutz Medical Campus, at

24   Bioscience 2 1270 E. Montview Avenue, Suite 100, Aurora, Colorado, 80045, United States.

25   [10] Nikon Imaging Center, Department of Cellular and Molecular Medicine, University of

26   California San Diego, La Jolla, CA 92093, USA.

28   *Correspondence to Brian A. Aguado, PhD. Email: baguado@ucsd.edu.

Abstract

Aortic valve stenosis (AVS) is a progressive disease wherein males develop valve

calcification relative to females that develop valve fibrosis. Valvular interstitial cells (VICs)

aberrantly activate to myofibroblasts during AVS, driving the fibrotic valve phenotype in females.

Myofibroblasts further differentiate into osteoblast-like cells and secrete calcium nanoparticles,

driving valve calcification in males. We hypothesized the lysine demethylase UTY (ubiquitously

transcribed tetratricopeptide repeat containing, Y-linked) decreases methylation uniquely in

response to nanoparticle cues in the valve extracellular matrix to promote an osteoblast-like

phenotype. Here, we describe a bioinspired hydrogel cell culture platform to interrogate how

nanoscale cues modulate sex-specific methylation states in VICs activating to myofibroblasts and

osteoblast-like cells. We found UTY (ubiquitously transcribed tetratricopeptide repeat containing,

Y-linked) modulates VIC phenotypes in response to nanoscale cues uniquely in males. Overall, we

reveal a novel role of UTY in the regulation of calcification processes in males during AVS

progression.

Introduction

Aortic valve stenosis (AVS) is sexually dimorphic disease where aortic valve tissue progressively undergoes fibro-calcification[1]. AVS impacts approximately 13% of the population aged over 75, and gender-based disparities exist with clinical management of AVS, given retrospective studies suggesting women are under recommended for valve replacements. Given that valve replacement patients are also at risk for increased co-morbidities after the procedure, a significant need exists to identify non-surgical based strategies to delay the onset of AVS[2]. Clinical research further suggests the degree of fibro-calcification in the valve depends on biological sex, where males exhibit increased calcification relative to females[3,4]. Given the sex dimorphisms observed in valve tissue during AVS onset, biological sex must be considered when identifying paths toward non-surgical pharmacologic treatments for AVS.

Our collective understanding of the sex-specific fibro-calcification processes in valve tissue during AVS remain limited. Valvular interstitial cells (VICs) are the resident fibroblast-like cells in valve tissue that are known to regulate valve fibro-calcification[3]. In early stages of AVS, VICs chronically activate to myofibroblasts and express alpha smooth muscle actin (aSMA) stress fibers[5]. In moderate to late stages of AVS, myofibroblasts further differentiate into osteoblast-like valve cells[6,7]. Osteoblast-like valve cells are characterized by the nuclear localization of runt-related transcription factor (RUNX2) and contribute to the accumulation of stiff, spherical calcium phosphate nanoparticles in valve tissue[8,9]. Over time, nanoparticles increase in size and abundance and further exacerbate valve fibro-calcification[3]. We posit the initiation of sex specific AVS progression likely depends on VIC interactions with the valve extracellular microenvironment.

Calcific lesions observed in healthy and late stage AVS tissue contain dense, mineralized spherical nanoparticles (NPs) of varying size in the valve extracellular microenvironment[10]. Prior

studies have shown that osteoblast-like cells experience altered cytoskeletal dynamics after culture

on substrates containing stiff NPs[11]. In particular, the NPs affected actin organization, disrupted

microtubule networks, and reduced overall cell spreading. VICs interact with their surrounding

microenvironment via transmembrane integrin receptors[12,13]. Integrins are directly linked to the

actin cytoskeleton and mediate cell response to external stimuli through alterations to the nuclear

chromatin structure[14–16]. Prior studies have also identified differences in chromatin structure of

VICs from both healthy and diseased patients[17,18]. As such, chromatin re-arrangement in male and

female VICs in response to nanoscale cues in the extracellular matrix is hypothesized to modulate

sex-specific cellular phenotypes during valve fibro-calcification.

Furthermore, cellular responses to the extracellular microenvironment also depend on

intracellular sex chromosome effects[19–22]. Sex differences in AVS progression have been attributed

to both patient-specific sex hormone levels and sex chromosome activity[23–25]. To detangle the

effects of sex hormones from sex chromosome activity, approaches using hydrogel biomaterials to

recapitulate the valve extracellular matrix have been used to show female VICs have increased

myofibroblast activation due to genes that escape X chromosome inactivation[26,27]. Outside of

VICs, previous work has also shown hydrogels to be critical tools to understand sex-specific

mechanotransduction in endothelial cells cultured *in vitro*[28,29]. Our understanding of how VICs

respond to nanoscale cues in the extracellular matrix will continue to harness *in vitro* culture tools

to probe sex-specific cell-nanomaterial interactions and subsequent sex-specific effects on cell

phenotypes.

Here, we engineered a bioinspired hydrogel cell culture platform to interrogate how

nanoscale stiffness cues modulate the VIC-to-myofibroblast-to-osteoblast transition in males and

females. Hydrogel biomaterials have been used previously to culture primary VICs isolated from

aortic valve leaflets as a strategy to determine how sex chromosome linked genes modulate sex

differences in VICs phenotypes[26,27]. In our approach, we incorporated stiff, polystyrene

nanoparticles (PSNPs) into a water-swollen crosslinked poly(ethylene glycol) (PEG) norbornene

hydrogel. After culture on our PSNP hydrogel, our results collectively show male-specific

responses to nanoscale stiffness cues through Y-chromosome linked histone demethylase,

Ubiquitously Transcribed Tetratricopeptide Repeat Containing, Y-linked, UTY. After UTY

demethylase knockdown, we show increased methylation as well as upregulation of Transforming

Growth Factor Beta Receptor 1 (TGFBR1), a key promoter of calcification processes in VICs.

Overall, we have potentiated a role of the Y-linked UTY lysine demethylase in causing sex-specific

effects that guide male VICs towards calcification.

Results

Hydrogels containing polystyrene nanoparticles mimic the valve microenvironment.

We engineered a polystyrene nanoparticle (PS-NPs) hydrogel that recapitulates the valve

matrix that contains nano-scale calcium phosphate particles. We obtained age-matched male and

female aortic valve tissue (UCSD IRB 804209) from two healthy patients and two patients

diagnosed with AVS. Using scanning electron microscopy (SEM), we visualized the accumulation

of nano-scale calcium phosphate particles in aortic valve tissue from male and female AVS patients

relative to healthy patients (Figure 1A). After quantifying particle diameter, we observe

significantly larger particle sizes in males relative to females (Supplementary Figure 1). To capture

the range of observed particle sizes in the tissue microenvironment, we incorporated 200 or 2000

nm diameter PS-NPs into our polyethylene(glycol) norbornene (PEG-Nb) hydrogel network to

create a matrix containing nano-scale particles (Figure 1B). We quantified bulk stiffness using

shear rheology (Figure 1C, 1D) and surface stiffness using atomic force microscopy

(Supplementary Figure 2). We determined that the elastic modulus (E) of our soft, blank hydrogel

matrix formulation ($E = 2.48 \pm 0.05$ kPa) does not significantly change with the incorporation of

200 nm PS-NPs ($E_{200 \text{ nm PS-NPs, low}} = 2.37 \pm 0.12$ kPa and $E_{200 \text{ nm PS-NPs, high}} = 2.98 \pm 0.19$ kPa) and

2000 nm PS-NPs ($E_{2000 \text{ nm PS-NPs, low}} = 3.36 \pm 0.09$ kPa and $E_{2000 \text{ nm PS-NPs, high}} = 2.55 \pm 0.02$ kPa),

relative to our stiff hydrogel formulation ($E = 21.80 \pm 0.36$ kPa). We also utilized SEM to confirm

PS-NP presence on the surface of the hydrogel matrix (Figure 1E).

Sex-specific myofibroblast activation and osteoblast-like differentiation on nanoparticle

hydrogels.

To test the effect of PS-NP nanoparticles on sex-specific valve cell phenotypes, we isolated

valvular interstitial cells (VICs) from porcine aortic valve tissue for culture on our PS-NP

hydrogels (Figure 2A). First, using alpha smooth muscle actin (αSMA) stress fibers as a marker

of myofibroblast activation, we observed female VICs having significantly higher levels of

myofibroblast activation on the control, blank hydrogel without particles and PS-NP hydrogel

conditions, compared to male VICs (Figures 2B, 2C). Second, we evaluated the nuclear

localization of the transcription factor runt-related transcription factor 2 (RUNX2) as a marker of

osteoblast-like differentiation. Male VICs displayed significantly higher levels of RUNX2 nuclear

localization on the control hydrogel and most PS-NP hydrogel conditions, compared to female

VICs (Figures 2D, 2E). Notably, we did not observe dose-dependent or PS-NP size-dependent

trends in the phenotypes. Collectively, our results provide evidence that male and female VICs

exhibit sex-specific differences in myofibroblast and osteoblast-like phenotypes in response to PS-

NPs in culture. Though we detected sex-specific myofibroblast and osteoblast-like VIC

phenotypes in response to PS-NP hydrogels, we did not observe consistent dose-dependent or PS-

NP size-dependent trends in the phenotypes.

Sex-specific methylation and acetylation states on PS-NP hydrogels.

Recognizing that epigenetic modifications drive cellular phenotype plasticity via

alterations to methylation and acetylation states in the nucleus, we sought to characterize sex-

specific epigenetic states in VICs on our PS-NP gels (Figure 3A). First, to understand potential

cell-matrix interactions that modulate transcriptional activity, we sought to investigate methylation

states in VICs cultured on PS-NP hydrogels. Male VICs displayed reduced tri-methylation across

most hydrogel conditions relative to female VICs (Figure 3B, 3C). Interestingly, we did observe

dose-dependent and PS-NP size-dependent changes in methylation after culture on PS-NP

hydrogels. Next, we determined acetylation states in VICs cultured on PS-NP gels, with the

hypothesis that male VICs would have increased acetylation to drive expression of osteoblast-

associated genes. Interestingly, we did observe a unique trend in the males, where only larger

particles affected the acetylation state (Figure 3D, 3E). Male VICs also displayed significantly

higher acetylation states (Figure 3D) in most of our blank and PS-NP hydrogel conditions, when

compared to female VICs (Figure 3E). Our evidence suggests that male-specific epigenetic

modifications may regulate male-specific osteoblast-like differentiation on PS-NP hydrogels.

UTY modulates male-specific methylation on PS-NP hydrogels.

To investigate how epigenetic alterations regulate male-specific osteoblast-like phenotypes

on PS-NP hydrogels, we next hypothesized epigenetic modifiers coded on the Y-chromosome may

contribute to male-specific phenotypes. Specifically, we posit that a Y-linked demethylase may

play a key role in regulating VIC response to nanoscale cues in the extracellular matrix. We

interrogated the role of the Y-linked global demethylase Ubiquitously Transcribed

Tetratricopeptide Repeat Containing, Y-linked (UTY) in regulating VIC phenotype and variable

methylation on PS-NP hydrogels. We first validated the male-specific mRNA expression of *UTY*

using real time quantitative polymerase chain reaction (RT-qPCR) (Figure 4A). Next, we employed

a small interfering RNA (siRNA) targeting the *UTY* gene to transcriptionally silence *UTY* while

VICs are cultured on PS-NP hydrogels. We utilized RT-qPCR to validate that we significantly

reduced relative *UTY* gene expression compared to our Silencer control uniquely in male VICs

(Figure 4B). Female VICs did not show significant *UTY* expression in all our conditions, further

validating a male-specific reduction in *UTY* expression. As another means of validation, we next

used immunofluorescence to detect global methylation in male and female VIC nuclei (Figure 4C).

As hypothesized, UTY knockdown in male VICs lead to a significant increase in methylation on

blank hydrogels (Figure 4D). Interestingly, UTY knockdown in female VICs also increased

methylation on blank gels, although the effect was more pronounced in male VICs. On PS-NP

gels, UTY knockdown did not impact methylation in male or female VICs. Together, we suggest

UTY modulates male-specific methylation states in VICs cultured on our engineered hydrogels.

Bulk sequencing and gene ontology unveils upregulation of pro-osteogenic differentiation

pathways after *UTY* siRNA.

Given the sex-specific differences in myofibroblast and osteoblast-like phenotypes, in

addition to sex-specific methylation and acetylation states, we performed bulk mRNA sequencing

to understand the sex-specific transcriptomic responses to *UTY* siRNA treatment. In both blank

and PS-NP gels, male VICs displayed a larger number of differentially expressed genes (DEGs)

compared to female VICs (Figure 5A-B, Supplemental Figure 3A-3B). Gene ontology (GO)

enrichment analysis of male VICs with *UTY* knockdown reveal upregulated DEGs associated with

organelle assembly, cell morphogenesis, responses to mechanical stimulus, cilium organization,

and Transforming Growth Factor Beta Receptor (TGFBR) signaling, among others (Figure 5C).

After *UTY* knockdown, downregulated DEGs in male VICs associated with GO terms including

GTP binding, guanyl ribonucleotide binding, and protein kinase binding (Figure 5D). In females,

many upregulated and downregulated DEGs after siRNA treatment were associated with pathways

involving cell metabolism and cell cycle regulations (Supplemental Figure 3C-3D). We speculate

that these pathways were differentially expressed because of the internalization of the siRNA

delivery vehicle, rather than the presence of the *UTY* siRNA itself. Collectively, our RNAseq

data suggest *UTY* modulates pathway activity uniquely in male VICs responding to PS-NP gels.

Discussion

Here, we highlight that the Y-chromosome modulates male-specific VIC phenotypes to

nanoscale cues in the extracellular matrix relative to female VICs (Figure 6). Using PEG hydrogels

with polystyrene nanoparticles as a platform for cell culture, we show the protein-coding gene

UTY may modulate male-specific demethylation in VICs during calcification in AVS. Recent

studies have described the significant role of the mosaic loss of the Y-chromosome (mLOY) in

male-specific cardiovascular disease progression[30,31,32]. mLOY occurs as an aging-related

mutation in males, though the heterogeneity in mLOY in varying cell types and relation disease

progression is not well understood. In the context of AVS, significant positive associations between

transcatheter aortic valve replacement (TAVR) patient mortality and presence of mLOY in

circulating monocytes have been reported[30]. Transcriptomic studies on mLOY monocytes revealed

a profibrotic gene signature, suggesting their role in driving restenosis post-TAVR. Furthermore,

UTY is also known to protect against proinflammatory phenotypes in lung tissue during pulmonary

hypertension. Specifically, *UTY* knockdown resulted in significant increases in the production of

proinflammatory cytokines CXCL9 and CXCL10, recruiting macrophages and promoting a

fibrotic phenotype in the lung tissue[33]. Our results here further corroborate prior efforts implicating

Y chromosome linked genes, including *UTY*, as key male-specific modulators of male-specific cell

phenotypes[33,30,31]. Together, our work here provides a hydrogel cell culture platform that enables

the interrogation of Y chromosome linked genes and their impact on male-specific cellular

phenotypes.

To the best of our knowledge, our study is the first to directly show *UTY* influences male-

specific VIC myofibroblast activation and osteoblast-like differentiation in response to

extracellular mechanical cues. Primary VICs from porcine sources have long been utilized to

evaluate myofibroblast activation and osteogenic differentiation processes in AVS in response to

different bulk substrate stiffnesses[28,34–39]. For example, in response to tissue culture plastic (elastic

modulus, E ~ 1 GPa), VICs automatically activate to myofibroblasts through the expression of α-

SMA stress fibers[26]. As an alternative approach, hydrogel biomaterial platforms composed of PEG

and a variety of crosslinking chemistries have been used to study VIC mechanosensitivity and sex-

specific modulation of myofibroblast activation and osteoblast differentiation states. For example,

previous work has shown that VICs cultured in 2D or 3D PEG-norbornene hydrogels in the ~1

kPa to ~40 kPa stiffness regime recapitulates sex-specific myofibroblast phenotypes observed in

valve tissue[26,27]. PEG hydrogel cell culture platforms also showed X chromosome inactivation

genes and the p38 MAPK signaling pathway partially regulate female-specific myofibroblast

activation processes[27]. Using a similar hydrogel system, we implemented a bio-inspired approach

where we incorporated nanoparticles into the hydrogel matrix to mimic nanoscale features in valve

tissue, as apparent in our SEM images and previous work. Inflammatory cytokines were found to

direct myofibroblasts towards the osteogenic pathway toward calcification[40]. Our work suggests

that nuclear localization of runt-related transcription factor 2 (RUNX2) in VICs responding to

nanoscale features in tissue may modulate a male-specific bias toward the initiation of calcification

processes. One of the most upregulated GO terms in males after *UTY* siRNA on PS-NP hydrogels

included Transforming Growth Factor Beta Receptor (TGFBR) signaling. Prior studies have

identified TGFBR1 as a key mediator in myofibroblast and osteogenic differentiation processes,

indicating a potential role for *UTY* in regulating pathways associated with fibro-calcification[41–43].

Enabled with our hydrogel platform, our work provides evidence for a novel mechanism

describing UTY as a candidate regulator of myofibroblast and osteoblast-like phenotypes in male

VICs, furthering our knowledge on how sex chromosomes may broadly contribute to somatic cell

phenotypes.

Our study also showcased the role of *UTY* in modulating the global transcriptional

landscape in VICs responding to hydrogel microenvironments via modulations in tri-methylation

and acetylation markers. Many studies have linked activity of mechanosensitive cell surface

receptors in modulating epigenetic states in the nucleus[17,18]. Epigenetic modifications can occur in

response to an external mechanical stimulus, which alters the conformation of the cell's surface

receptor, altering the structure of the connected intracellular cytoskeletal network tied to the

nucleus[22]. Diseased myofibroblasts have also been shown to maintain distinct chromatin structures

in response to bulk stiffness cues in PEG-acrylate gels, compared to healthy myofibroblasts. The

altered mechanical tension on the nucleus influences epigenetic processes that can transcriptionally

activate (acetylation) or silence (methylation) different regions of chromatin. A prior study using

VICs showed that stiff hydrogels promoted increased acetylation in male VICs compared to soft

hydrogels, however, the specific epigenetic alterations and the mechanisms by which they occur

are still poorly understood. Building upon this work, we show that male specific methylation and

acetylation states may be modulated via *UTY* demethylase activity. Prior studies have investigated

various phenotypic changes of VICs in response to a variety of mechanical cues. When evaluating

methylation in males, we observed a consistent trend in decreased methylation correlating with

particle size and density, indicating male myofibroblasts uniquely interact with nanoscale features

in an extracellular microenvironment. Interestingly, female VICs did not show a decrease in

methylation when interacting with nanoscale cues, but did reduce methylation with larger

nanoparticles, indicating other demethylases may participate in female-specific cell-particle

interactions. A previous study showed that female human embryonic stem cells cultured with silver

nanoparticles have compromised differentiation processes via disordered X chromosome

inactivation, although female-specific effects of nanoparticles on pluripotent vs somatic cell

cultures have yet to be compared[44]. Future efforts will need to assess the role of other demethylases

in common between XY and XX cells, or potential activity of X-linked demethylases (e.g. *UTX,*

KDM5C, etc.) to supplement our observations in male VICs.

There are several limitations of the PSNP hydrogel platform described in our study. First,

our study did not decouple the effects of sex hormones from sex chromosomes. For future work,

a gonadectomized animal model can probe the effects of the sex chromosome on VIC phenotype.

Sex hormones could also be dosed into the hydrogel platform, further interrogating the intersection

between sex hormone and sex chromosome effects. Secondly, our study sought to characterize the

VIC-nanoparticle interactions on a soft hydrogel matrix. In the diseased AV environment, VICs

interact with a variety of mechanical and biochemical cues. Our study did not include biochemical

factors such as secreted inflammatory cytokines, which are present in the diseased AV due to

immune cell infiltration. Future cell culture platforms can be adapted to include inflammatory

factors secreted from macrophages to evaluate epigenetic modifications in response to

inflammatory cytokines and nanoscale stiffness cues. Third, our study also evaluated cell

phenotypes after two days in culture. Acknowledging that AVS is a time-dependent progressive

disease, we suggest future studies where longer timepoints are used to evaluate sex-specific

chromatin structures (e.g. after 7 days in culture) as VICs acquire a more permanent myofibroblast

phenotype reflective of diseased valve tissue. Fourth, future efforts to use spherical calcium

phosphate particles instead of PS-NPs may be used to further enhance the physiological relevance

of the *in vitro* culture platform and evaluate effects of UTY and other Y-linked genes on

calcification[45].

Taken together, our study implicates the importance of sex chromosome-linked genes in

the progression of disease-driving phenotypes in AVS. Our bioinspired hydrogel platform provides

evidence that nanoscale cues in the extracellular matrix influence sex-specific phenotypes in VICs,

and broadly suggests biomaterials are useful tools to answer research questions related to sex-

specific biology[46,47]. We posit that Y-linked epigenetic regulators, such as UTY, regulate male-

specific osteoblast-like phenotypes, contributing to earlier onset of valvular calcification in males.

Future studies will explore the interplay between mechanical stimuli and sex-specific epigenetic

regulation in males and females.

Materials and Methods

Nanoparticle hydrogel fabrication

8-arm, 40 kDa polyethylene(glycol)-norbornene (PEG-Nb) was synthesized as previously described[48]. Hydrogel precursor solutions were made mixing together 4% (w/v) PEG-Nb, 5 kDa PEG-dithiol crosslinker (2.96 mM, JenKem), CRGDS peptide (2 mM, Bachem), phosphate buffered saline (PBS, Gibco, 14-190-250), and photoinitiator lithium phenyl-2,4,6-trimethyl-benzoylphosphinate (1.7 mM, Sigma, 900889). To incorporate nanoparticles into hydrogel precursor solutions, polystyrene nanoparticle solutions, we utilized 1% wt/vol aqueous solutions of 200 nm (Nanocs, PS01-200) or 2000 nm (Nanocs, PS01-2u) polystyrene nanoparticles (PS-NPs) instead of PBS in the hydrogel precursor formulation. All hydrogels were made utilizing a 0.99:1 thiol-to-ene ratio. Vapor deposition was used to functionalize 12 mm glass coverslips (Mercedes Scientific, MER0012) and 25 mm glass coverslips (Mercedes Scientific, MER0025) with thiol groups to allow for hydrogel adhesion. Briefly, coverslips were placed in an autoclave jar along with 100 µL of mercaptopropyltrimethoxysilane (MPTS, Sigma-Aldrich, X) in a 60°C oven overnight for thiol functionalization. Glass slides were coated with Sigmacote (Sigma-Aldrich, 175617) to create a hydrophobic glass surface during hydrogel fabrication. Either 11.3 µL or 65 µL of hydrogel precursor solution was pipetted onto the Sigmacote glass slide and covered with a 12 mm or 25 mm coverslip, respectively. The hydrogels were photopolymerized at 4 mW/cm^2, removed with a razor blade, sterilized with 5% isopropyl alcohol/PBS solution, and rinsed three times with PBS. After the final PBS rinse, cell culture media containing Media 199 (Gibco, 11043023) supplemented with 1% Fetal Bovine Serum (Gibco, 16000069), and 50 units penicillin, and 0.05 mg/ml streptomycin (Sigma Aldrich, P4458) was added to the hydrogels for overnight swelling prior to cell culture.

Rheology

Hydrogels containing nanoparticles were prepared as described above for rheological measurements. Storage (G') and loss (G'') moduli were measured using a DHR-3 rheometer (TA Instruments) with an 8 mm parallel plate geometry. Oscillatory shear rheology was used with an amplitude of 1% and frequency of 1 Hz[26]. To obtain the elastic modulus (E), the following formula was used: $E = 2 * G' (1 + v)$ where v is Poisson's ratio. We utilized a value of $v = 0.5$, assuming G' >>> G'' for viscoelastic hydrogels.

Scanning Electron Microscopy (SEM) on Hydrogels and Human Valve Tissue and quantification of particle size distribution

For scanning electron microscopy (SEM) of hydrogels, samples were prepared on coverslips, frozen in liquid nitrogen, and lyophilized. SEM was performed using a JSM-6010LA instrument (JEOL Ltd, CU Anschutz) using manufacturer settings. For SEM on human valve tissue, samples were fixed, paraffin embedded, and sectioned onto glass slides. Samples were sputter coated with iridium for 30 seconds at 75% power and imaged using a FEI Quanta FEG 250 (FEI) using manufacturer settings. Average calcium particle size was determined from two biological replicates for each sex in the disease condition using ImageJ.

Atomic Force Microscopy (AFM)

For atomic force microscopy, hydrogels mounted on coverslips were immersed in PBS and analyzed using a JPK NanoWizard 4a instrument (JPK Instruments). Hydrogels were probed using CB3 cantilevers with triangular tips (Nanosensors, qp-BioAC-CI) until an initial force of 1 nN was reached. Calibration of the cantilever was made applying the thermal noise method. Force

spectroscopy was performed at each node of a square grid distributed over a 20 µm by 20 µm grid.

Force–displacement (F–d) data were collected using a peak load of 5 nN at a displacement rate of

10 µm/s. Force–deformation (F–δ) data were calculated using F–d curves by subtracting the initial

cantilever deflection. To determine Young's modulus E at each node, the loading portion of each

F–δ curve was fit to an analytical model for a rigid conical tip in contact with an elastic half-space

(38), i.e., $F=(2/\pi)(E/1-v^2)(\tan\alpha)\delta^2$, using angle α determined from SEM images, an assumed

Poisson's ratio for the hydrogel ($v = 0.5$), and E as the sole fitting parameter.

Porcine valvular interstitial cell isolation and culture

Aortic valves (AVs) were dissected from 5-8-month-old male or female pigs (Midwest

Research Swine) to obtain valvular interstitial cells (VICs). For each isolation, four to eight porcine

hearts were separated by sex and pooled. Each batch of VICs obtained from an isolation is

considered as a biological replicate, in which 2-3 biological replicates were used for Figures 2-6.

After dissection, AV leaflets were submerged and rinsed in Earle's Balanced Salt Solution (EBSS,

Sigma Aldrich, E2888) containing 50 units penicillin and 0.05 mg/ml streptomycin. Next, leaflets

were digested in EBSS containing 250 units of Collagenase type 2 (Worthington Biochemical,

LS004176) for 30 minutes at 37°C under constant agitation at 100 rpm. After, collagenase solution

was aspirated from the AV leaflets to remove endothelial cells. A secondary digestion with fresh

collagenase solution was performed for 60 minutes under agitation at 100 rpm. To collect VICs,

the remaining tissue was scraped against a 100 µm cell strainer, washed with VIC media, then

centrifuged at 400 g for 10 minutes. The supernatant was aspirated and replaced with fresh VIC

expansion media, comprised of Media 199 supplemented with 15% FBS, and 50 units penicillin

and 0.05 mg/ml streptomycin. Freshly isolated VICs were cultured in VIC expansion media with

media changes every other day until confluency. VICs were frozen down at passage 1. All

experiments utilized passage 2 VICs at 35,000 cells/cm^2. To reduce proliferation during

experimentation, passage 2 VICs were cultured in VIC media containing 1% FBS.

Small-interfering RNA (siRNA) Gene Knockdown

Gene knockdowns in the porcine VICs were conducted using small interfering RNAs

(siRNAs). We utilized Ubiquitously Transcribed Tetratricopeptide Repeats, Y-linked (*UTY*) siRNA

(Ambion, 4392421 ID: s14740). On Day 0, cells were seeded at 42,000 cells/cm^2 coverslips in 1%

VIC media, without antibiotics or antifungal supplements, and cultured overnight. We increased

cell seeding density per manufacturer recommendations for siRNA experiments. On Day 1, *UTY*

siRNAs and Silencer negative control (Ambion, AM4635) were incubated with Lipofectamine

3000 Transfection Reagent (Thermo Scientific, L3000008) for 15 minutes at room temperature

before adding it to fresh 1% VIC media without antibiotics. The final siRNA concentration for the

12 mm and 25 mm hydrogels was 20 pmol and 100 pmol, respectively. On Day 2, the samples

were monitored without media change. On Day 3, the samples were fixed for immunofluorescence

and lysed for RNA collection. In total, *UTY* siRNA treatment lasted 48 hours.

Immunofluorescence, Imaging, and Analysis

Cells were cultured on NP hydrogels for 48 hours before fixing with 4% (w/v)

paraformaldehyde (Electron Microscopy Sciences, 15710) for 20 minutes. Afterwards, cells were

permeabilized with 0.1% (w/v) Triton-X-100 (Sigma Aldrich, 93443) for one hour.

Permeabilization buffer was removed and added 5% (w/v) bovine serum albumin (BSA, Sigma

Aldrich, A8327) as blocking buffer overnight at 4°C. The following primary antibodies were

diluted in blocking buffer and incubated for one hour at room temperature: α-smooth muscle actin

(aSMA, 1:300, Abcam, ab7817), runt-related transcription factor 2 (RUNX2, 1:300, Abcam,

ab23981), Methyl-Lysine (MeK, 1:350, Novus Biologicals, NB600824), and acetylated lysine

(AcK, 1:350, Abcam, ab190479). Samples were rinsed with 0.05% (v/v) Tween-20 (Sigma

Aldrich, P1379) for 5 minutes. The following secondary antibodies were diluted in blocking buffer

and incubated for one hour at room temperature protected from light: AlexaFluor 488 goat anti-

mouse (1:200, Thermo Scientific, A11001), AlexaFluor 647 goat anti-rabbit (1:200, Thermo

Scientific, A21245), or AlexaFluor 647 donkey anti-rabbit (1:200, Thermo Scientific, A32790).

Nuclei were stained with DAPI (1:500, Roche, 10236276001) and cytoplasm were stained with

CellMask Orange (1:5000, Fisher Scientific, H32713). Sample coverslips were rinsed with PBS

and transferred to a glass-bottom well plate (CellVis, P24-1.5H-N) for imaging on a Nikon Eclipse

Ti2-E. Fluorescence was quantified after modifying a freely available MATLAB code[27], and

fluorescence channel exposure was adjusted consistently in all conditions for representative image

clarity. Alpha smooth muscle actin intensity ratios were calculated by normalizing α-SMA

intensity with CellMask cytoplasmic intensity. Nuclear RUNX2 intensity ratios were calculated

by normalizing RUNX2 nuclear intensity to cytoplasmic intensity. Nuclear methylated lysine

ratios were calculated by normalizing MeK nuclear intensity to cytoplasmic intensity. Nuclear

acetylated lysine ratios were calculated by normalizing AcK nuclear intensity to DAPI intensity.

RNA Isolation and RT-qPCR

RNA was collected at specified timepoints using the RNeasy Micro Kit (Qiagen, 74004)

following the manufacturer's protocol. Gels on 25 mm coverslips were inverted onto lysis buffer

for 3 minutes then rinsed with 70% (v/v) ethanol to collect lysed cellular material. After following

manufacturer's protocol for RNA isolation, samples were analyzed for mRNA concentration using

a NanoDrop 2000 spectrophotometer (Thermo Scientific). Sample concentrations were normalized

to 1 ng/µL during cDNA preparation using the iScript Synthesis Kit (Bio-Rad, 1708841), following

manufacturer's protocol. To measure relative gene expression, iQ SYBR Green Supermix (Bio-

Rad, 1708882) was mixed with primer sequences found in Supplemental Table 1. Cq values were

determined using a CFX384 iCycler (Bio-Rad). All gene expression calculations were normalized

to *RPL30* expression.

Bulk RNA Sequencing and Analysis

Prior to bulk paired-end RNA sequencing, RNA was isolated as described above. RNA quality

was assessed by Agilent TapeStation 4200 and RNA quantity was determined by QuBit 3.0

Fluorometer. Approximately 200 ng of RNA per sample was used for sequencing. The Sanford

Consortium for Regenerative Medicine's Genomics Core conducted cDNA library preparation and

indexing using the Illumina Stranded mRNA kit and IDT Illumina RNA UD Indexes, respectively.

Sequences were checked for quality control using fastqc (v0.12.1) and subsequently removed

adapter sequences using trimmomatic (v0.39) if adapter sequence content exceeded the threshold

identified by fastqc. Alignment to the porcine genome (Sscrofa11.1) was conducted using HISAT2

2.2.1. Subsequent differential gene expression analysis was conducted in R (4.3.2) and R Studio

(2023.12.1) using DESeq2 (Release 3.19) before we filtered for genes with a false discovery rate

above 0.1. We utilized gene ontology analysis features in Ingenuity Pathway Analysis (Qiagen) for

gene enrichment analyses of upregulated and downregulated genes in the PS-NP samples.

 Statistical Analysis

For immunofluorescence studies, one-way ANOVA with Tukey's posttests were

conducted. We set a threshold of significance cutoff at $P < 0.0001$. Due to the experimental sample

size (N > 700 cells), we sought to reduce statistical bias by conducting a Cohen's d-value test to

determine significance independent of sample size. Thresholds of significance were represented

by the following key: *** = d > 0.8, ** = d > 0.5, * = d > 0.2 for between sex significance; ### =

d > 0.8, ## = d > 0.5, # = d > 0.2 for male group significance; and \$\$\$ = d > 0.8, \$\$ = d > 0.5, \$ =

d > 0.2 for female group significance. For RT-qPCR studies, one-way ANOVA with Tukey's

posttests were conducted. For Figure 1, data is shown as the mean $\pm$ standard deviation, as

calculated in GraphPad PRISM.

Figure Captions

Figure 1. PS-NP hydrogels recapitulate nano-scale features in aortic valve tissue ECM.

(A) Scanning electron microscopy (SEM) images of male healthy, diseased, and female diseased

human aortic valve leaflets. Scale bar = 5 μm. (B) Schematic of process to incorporate polystyrene

nanoparticles (PS-NPs) into PEG-Norbornene hydrogel precursor solutions to create PS-NP

hydrogels adhered to glass coverslips via UV polymerization. (C) Rheological characterization of

UV-mediated polymerization of soft, stiff, and PS-NP hydrogels made with 200 nm and 2000 nm

nanoparticles (n=3 representative measurements) at low and high concentrations (0.02 mg/ml and

0.7 mg/ml, respectively). (D) Young's modulus (E) values for soft, stiff, and PS-NP hydrogels

(N=3 gels). Significance determined via one-way ANOVA (****p<0.0001 denoting statistical

significance between hydrogel formulations). Mean ± SD shown. (E) SEM of blank and PS-NP

hydrogels (200 nm, 2000 nm). Yellow arrows denote PS-NPs. Scale bar = 5 μm.

Figure 2. Male and female VICs exhibit sex-specific myofibroblast and osteoblast-like

phenotypes.

(A) Schematic describing experimental workflow for probing myofibroblast activation with α-

smooth muscle actin (α-SMA) and osteoblast-like differentiation with runt related (RUNX2).

Figure drawn with BioRender. (B) Representative confocal microscopy images of alpha smooth

muscle stress fibers (green) and nuclei (blue). Scale bar = 50 μm. (C) Normalized α-SMA

expression ratios for male and female VICs cultured on control hydrogels with no PS-NPs, 200

nm PS-NPs, or 2000 nm PS-NPs at low and high concentrations (0.02 mg/ml and 0.7 mg/ml,

respectively). Significance determined via Cohen's d-test (*d < 0.2, **d < 0.5, ***d<0.8,

****d<1.4 denoting significance between biological sex in the provided table, red asterisk color

indicates significant increase in female groups, [#]d < 0.2, [##]d < 0.5 denoting significance relative to

male control with no particles, [$$]d < 0.5, [$$$]d < 0.8 denoting significance relative to female control

with no particles). (D) Representative confocal microscopy images of RUNX2 expression (red)

and nuclei (blue). Scale bar = 50 μm. (E) RUNX2 nuclear to cytoplasmic intensity ratios of male

and female VICs cultured on control hydrogels with no PS-NPs, 200 nm PS-NPs, or 2000 nm PS-

NPs at low and high concentrations (0.02 mg/ml and 0.7 mg/ml, respectively). Significance

determined via Cohen's d-test (*d < 0.2, **d < 0.5, ***d<0.8, ****d<1.4 denoting significance

between biological sex in the provided table, blue asterisk color indicates significant increase in

male groups, [#]d < 0.2, [##]d < 0.5 denoting significance relative to male control with no particles,

[$$]d < 0.5, [$$$]d < 0.8 denoting significance relative to female control with no particles.

Figure 3. Male and female VICs exhibit sex-specific changes in global methylation and

acetylation states.

(A) Schematic describing experimental workflow for assessing sex-specific methylation and

acetylation states in VICs cultured on PS-NP gels. Figure drawn with BioRender. (B)

Representative confocal microscopy images of methylated lysine (yellow) and nuclei (blue). Scale

bar = 50 μm. (C) Normalized nuclear intensity ratios of methylated lysine to DAPI in male and

female VICs cultured on control hydrogels with no PS-NPs, 200 nm PS-NPs, or 2000 nm PS-NPs

at low and high concentrations (0.02 mg/ml and 0.7 mg/ml, respectively) . Significance determined

via Cohen's d-test (**d < 0.5, ***d<0.8, ****d<1.4 denoting significance between biological sex

in the provided table, red asterisk color indicates significant increase in female groups, [#]d < 0.2,

[##]d < 0.5, [###]d<0.8, [####]d<1.4 denoting significance relative to male control with no particles, [$]d <

0.2 denoting significance relative to female control with no particles). (D) Representative confocal

microscopy images of acetylated lysine (yellow) and nuclei (blue). Scale bar = 50 μm. (E)

Normalized nuclear intensity ratios of acetylated lysine to DAPI in male and female VICs cultured

on control hydrogels with no PS-NPs, 200 nm PS-NPs, or 2000 nm PS-NPs at low and high

concentrations (0.02 mg/ml and 0.7 mg/ml, respectively). Significance determined via Cohen's d-

test (*d<0.2, **d < 0.5 denoting significance between biological sex in the provided table, blue

asterisk color indicates significant increase in male groups, [#]d < 0.2, [###]d<0.8 denoting significance

relative to male control with no particles, [$]d < 0.2, [$$]d < 0.5 denoting significance relative to female

control with no particles).

Figure 4. Male-specific methylation is uniquely increased in male VICs by siRNA-targeting

of Y-linked demethylase, UTY.

(A) Real time quantitative polymerase chain reaction (RT-qPCR) for relative expression of *UTY*

in male and female VICs cultured on blank, 200 nm, and 2000 nm PS-NP hydrogels. (B) RT-qPCR

to validate partial silencing of *UTY* in male VICs after introduction of siRNA vector. (C)

Representative confocal microscopy images of methyl lysine (red) and nuclei (blue). Scale bar =

50 μm. (D) Normalized nuclear intensity ratios of methylated lysine in male and female VICs

cultured on control hydrogels with no PS-NPs and 2000 nm PS-NPs. Significance determined via

Cohen's d-test (*d<0.2, **d < 0.5 denoting significance between biological sex in the provided

table, blue asterisk color indicates significant increase in male groups, [#]d < 0.2, [###]d<0.8 denoting

significance relative to male control with no particles, [$]d < 0.2, [$$]d < 0.5 denoting significance

relative to female control with no particles).

Figure 5. Bulk sequencing and gene ontology enrichment analysis in male VICs unveils

upregulated and downregulated GO terms.

(A) Bulk mRNA sequencing shows differentially expressed genes in male VICs (blue =

downregulated, red = upregulated, gray = not significant) in control and PS-NP hydrogels using

DESeq2 (n=2 biological replicates). (B) Upregulated and (C) downregulated gene ontology-based

pathways that are enriched for genes differentially expressed in control and PS-NP hydrogels after

UTY siRNA treatment.

Figure 6. Proposed mechanism revealing UTY as a candidate mediator of male-specific demethylation.

(A) Male VICs have increased nuclear localization on blank hydrogels, and nuclear localization of RUNX2 increases when male VICs respond to PS-NPs embedded in the matrix. UTY reduces methylation in male VICs responding to PS-NPs, and methylation is restored after UTY knockdown. (B) Female VICs have increased α-SMA stress fiber formation relative to male VICs on blank hydrogels, with lower nuclear localization of RUNX2 and equivalent methylation and acetylation. Female VICs responding to PS-NPs have uniquely reduced α-SMA stress fiber formation with increases in acetylation.

Acknowledgements

R.M.G. acknowledges support from a National Institutes of Health Research Supplement to Promote Diversity in Health-Related Research (R00 HL148542) and the San Diego ARCS Foundation. T.B. acknowledges support from the San Diego Match Fellowship. N.E.F. acknowledges support from a National Institutes of Health Research Supplement to Promote Diversity in Health-Related Research (R00 HL148542) and support from the National Science Foundation Graduate Research Fellowship Program. B.P and L.M acknowledge funding from the National Institutes of Health (NIH) National Heart, Lung, and Blood Institute (NHLBI): grants K25HL148386 (B.P.), R01HL169578 (B.P.), and R01HL147064 (L.M.), the NIH National Institute of Aging (NIA) R21AG080257 (BP), the Ludeman Family Center for Women's Health Research seed grant (B.P.), the American Heart Association (AHA): 23CDA1052411 (B.P.). This study was also supported by generous grants of the John Patrick Albright (L.M. and B.P.). B.A.A. acknowledges support from the National Institutes of Health (R00 HL148542), the NIH Director's

New Innovator Award (DP2 HL173948), the Chan Zuckerberg Initiative Science Diversity Leadership Award, and the American Heart Association (22CDA942253). We acknowledge the help and resources of Dr. Guillaume Castillon at the Electron Microscopy Core and Dr. Zhicheng Long at the Nanofabrication core at the University of California San Diego.

References

1. Carabello BA, Paulus WJ. Aortic stenosis. *The Lancet*. 2009;373(9667):956-966. doi:10.1016/S0140-6736(09)60211-7

2. Myasoedova VA, Ravani AL, Frigerio B, et al. Novel pharmacological targets for calcific aortic valve disease: Prevention and treatments. *Pharmacol Res*. 2018;136:74-82. doi:10.1016/j.phrs.2018.08.020

3. Simard L, Côté N, Dagenais F, et al. Sex-Related Discordance Between Aortic Valve Calcification and Hemodynamic Severity of Aortic Stenosis. *Circ. Res.* 2017;120(4):681-691. doi: 10.1161/CIRCRESAHA.116.309306.

4. Gourgas O, Khan K, Schwertani A, Cerruti M. Differences in mineral composition and morphology between men and women in aortic valve calcification. *Acta Biomater*. 2020;106:342-350. doi:10.1016/j.actbio.2020.02.030

5. Ayoub S, Lee CH, Driesbaugh KH, et al. Regulation of valve interstitial cell homeostasis by mechanical deformation: implications for heart valve disease and surgical repair. *J R Soc Interface*. 2017;14(135):20170580. doi:10.1098/rsif.2017.0580

6. Goody PR, Hosen MR, Christmann D, et al. Aortic Valve Stenosis. *Arterioscler Thromb Vasc Biol*. 2020;40(4):885-900. doi:10.1161/ATVBAHA.119.313067

7. Blaser MC, Kraler S, Lüscher TF, Aikawa E. Multi-Omics Approaches to Define Calcific Aortic Valve Disease Pathogenesis. *Circ Res*. 2021;128(9):1371-1397. doi:10.1161/CIRCRESAHA.120.317979

8. Nagy E, Eriksson P, Yousry M, et al. Valvular osteoclasts in calcification and aortic valve

stenosis severity. *Int J Cardiol*. 2013;168(3):2264-2271. doi:10.1016/j.ijcard.2013.01.207

9. Yu C, Li L, Xie F, et al. LncRNA TUG1 sponges miR-204-5p to promote osteoblast

differentiation through upregulating Runx2 in aortic valve calcification. *Cardiovasc Res*.

2018;114(1):168-179. doi:10.1093/cvr/cvx180

10.   Bertazzo S, Gentleman E. Aortic valve calcification: a bone of contention. *Eur Heart J*.

2017;38(16):1189-1193. doi:10.1093/eurheartj/ehw071

11.   Ibrahim M, Schoelermann J, Mustafa K, Cimpan MR. TiO2 nanoparticles disrupt cell

adhesion and the architecture of cytoskeletal networks of human osteoblast-like cells in a size

dependent manner. *J Biomed Mater Res A*. 2018;106(10):2582-2593.

doi:10.1002/jbm.a.36448

12.   Chen JH, Simmons CA, Towler DA. Cell–Matrix Interactions in the Pathobiology of

Calcific Aortic Valve Disease. *Circ Res*. 2011;108(12):1510-1524.

doi:10.1161/CIRCRESAHA.110.234237

13.   Kodigepalli KM, Thatcher K, West T, et al. Biology and Biomechanics of the Heart Valve

Extracellular Matrix. *J Cardiovasc Dev Dis*. 2020;7(4):57. doi:10.3390/jcdd7040057

14.   Barczyk M, Carracedo S, Gullberg D. Integrins. *Cell Tissue Res*. 2010;339(1):269-280.

doi:10.1007/s00441-009-0834-6

15.   Takada Y, Ye X, Simon S. The integrins. *Genome Biol*. 2007;8(5):215. doi:10.1186/gb-

2007-8-5-215

16.   van der Flier A, Sonnenberg A. Function and interactions of integrins. *Cell Tissue Res*.

2001;305(3):285-298. doi:10.1007/s004410100417

17.   Walker CJ, Crocini C, Ramirez D, et al. Nuclear mechanosensing drives chromatin

remodelling in persistently activated fibroblasts. *Nat Biomed Eng*. 2021;5(12):1485-1499.

doi:10.1038/s41551-021-00709-w

18.   Walker CJ, Batan D, Bishop CT, et al. Extracellular matrix stiffness controls cardiac

valve myofibroblast activation through epigenetic remodeling. *Bioeng Transl Med*.

2022;7(3):e10394. doi:10.1002/btm2.10394

19.   Walker CJ, Schroeder ME, Aguado BA, Anseth KS, Leinwand LA. Matters of the heart:

Cellular sex differences. *J Mol Cell Cardiol*. 2021;160:42-55.

doi:10.1016/j.yjmcc.2021.04.010

20.   Hartman RJG, Huisman SE, den Ruijter HM. Sex differences in cardiovascular

epigenetics—a systematic review. *Biol Sex Differ*. 2018;9(1):19. doi:10.1186/s13293-018-

0180-z

21.   Deegan DF, Nigam P, Engel N. Sexual Dimorphism of the Heart: Genetics, Epigenetics,

and Development. *Front Cardiovasc Med*. 2021;8. doi:10.3389/fcvm.2021.668252

22.   Wijchers PJ, Festenstein RJ. Epigenetic regulation of autosomal gene expression by sex

chromosomes. *Trends Genet*. 2011;27(4):132-140. doi:10.1016/j.tig.2011.01.004

23.   Dobson LE, Fairbairn TA, Plein S, Greenwood JP. Sex Differences in Aortic Stenosis and

Outcome Following Surgical and Transcatheter Aortic Valve Replacement. *J Womens Health*.

2015;24(12):986-995. doi:10.1089/jwh.2014.5158

24.   Côté N, Clavel MA. Sex Differences in the Pathophysiology, Diagnosis, and

Management of Aortic Stenosis. *Cardiol Clin*. 2020;38(1):129-138.

doi:10.1016/j.ccl.2019.09.008

25.   Fleury MA, Annabi MS, Voisine M, et al. Impact of sex and sex hormones on

pathophysiology and progression of aortic stenosis in a murine model. *Physiol Rep*. 2022;

10(16): e15433. doi: 10.14814/phy2.15433

26.   Aguado BA, Walker CJ, Grim JC, et al. Genes That Escape X Chromosome Inactivation

Modulate Sex Differences in Valve Myofibroblasts. *Circulation*. 2022;145(7):513-530.

doi:10.1161/CIRCULATIONAHA.121.054108

27.   Vogt BJ, Peters DK, Anseth KS, Aguado BA. Inflammatory serum factors from aortic

valve stenosis patients modulate sex differences in valvular myofibroblast activation and

osteoblast-like differentiation. *Biomater Sci*. 2022;10(22):6341-6353.

doi:10.1039/D2BM00844K

28.   James BD, Allen JB. Sex-Specific Response to Combinations of Shear Stress and

Substrate Stiffness by Endothelial Cells In Vitro. *Adv Healthc Mater*. 2021;10(18):2100735.

doi:10.1002/adhm.202100735

29.   Sánchez Marrero G, Villa-Roel N, Li F, et al. Single-Cell RNA sequencing investigation

of female-male differences under PAD conditions. *Front Cardiovasc Med*. 2023;10.

doi:10.3389/fcvm.2023.1251141

30.   Mas-Peiro S, Abplanalp WT, Rasper T, et al. Mosaic loss of Y chromosome in monocytes

is associated with lower survival after transcatheter aortic valve replacement. *Eur Heart J*.

2023;44(21):1943-1952. doi:10.1093/eurheartj/ehad093

31.   Sano S, Horitani K, Ogawa H, et al. Hematopoietic loss of Y chromosome leads to

cardiac fibrosis and heart failure mortality. *Science*. 2022;377(6603):292-297.

doi:10.1126/science.abn3100

32.   Thompson DJ, Genovese G, Halvardson J, et al. Genetic predisposition to mosaic Y

chromosome loss in blood. *Nature*. 2019;575(7784):652-657. doi:10.1038/s41586-019-1765-3

33.   Y-Chromosome Gene, Uty, Protects Against Pulmonary Hypertension by Reducing

Proinflammatory Chemokines | American Journal of Respiratory and Critical Care Medicine.

Accessed May 13, 2024. https://www.atsjournals.org/doi/full/10.1164/rccm.202110-2309OC

34.   Puperi DS, Balaoing LR, O'Connell RW, West JL, Grande-Allen KJ. 3-Dimensional

spatially organized PEG-based hydrogels for an aortic valve co-culture model. *Biomaterials*.

2015;67:354-364. doi:10.1016/j.biomaterials.2015.07.039

35.   Puperi DS, Kishan A, Punske ZE, et al. Electrospun Polyurethane and Hydrogel

Composite Scaffolds as Biomechanical Mimics for Aortic Valve Tissue Engineering. *ACS*

Biomater Sci Eng. 2016;2(9):1546-1558. doi:10.1021/acsbiomaterials.6b00309

36.   Porras AM, Westlund JA, Evans AD, Masters KS. Creation of disease-inspired

biomaterial environments to mimic pathological events in early calcific aortic valve disease.

Proc Natl Acad Sci. 2018;115(3):E363-E371. doi:10.1073/pnas.1704637115

37.   Scott AJ, Simon LR, Hutson HN, Porras AM, Masters KS. Engineering the aortic valve

extracellular matrix through stages of development, aging, and disease. *J Mol Cell Cardiol*.

2021;161:1-8. doi:10.1016/j.yjmcc.2021.07.009

38.   Floy ME, Shabnam F, Givens SE, et al. Identifying molecular and functional similarities

and differences between human primary cardiac valve interstitial cells and ventricular

fibroblasts. *Front Bioeng Biotechnol*. 2023;11. Accessed July 2, 2023.

https://www.frontiersin.org/articles/10.3389/fbioe.2023.1102487

39.   Driscoll K, Cruz AD, Butcher JT. Inflammatory and Biomechanical Drivers of

Endothelial-Interstitial Interactions in Calcific Aortic Valve Disease. *Circ Res*.

2021;128(9):1344-1370. doi:10.1161/CIRCRESAHA.121.318011

40.   Grim JC, Aguado BA, Vogt BJ, et. al. Secreted Factors From Proinflammatory

Macrophages Promote an Osteoblast-Like Phenotype in Valvular Interstitial Cells.

Arterioscler. Thromb. Vasc. Biol. 2020; ;40(11):e296-e308. doi:

10.1161/ATVBAHA.120.315261

41.   Chen G, Deng C, Li YP. TGF-β and BMP Signaling in Osteoblast Differentiation and

Bone Formation. *Int J Biol Sci*. 2012;8(2):272-288. doi:10.7150/ijbs.2929

42.   Vander Ark A, Cao J, Li X. TGF-β receptors: In and beyond TGF-β signaling. *Cell

Signal*. 2018;52:112-120. doi:10.1016/j.cellsig.2018.09.002

43. Yang P, Troncone L, Augur ZM, Kim SSJ, McNeil ME, Yu PB. The role of bone

morphogenetic protein signaling in vascular calcification. *Bone*. 2020;141:115542.

doi:10.1016/j.bone.2020.115542

44. Zhang J, Chen Y, Gao M, et. al. Silver Nanoparticles Compromise Female Embryonic

Stem Cell Differentiation through Disturbing X Chromosome Inactivation. *ACS Nano*. 2019;

13, 2, 2050–2061. doi:10.1021/acsnano.8b08604

45. Richards JM, Kunitake JAMR, Hunt HB, et al. Crystallinity of hydroxyapatite drives

myofibroblastic activation and calcification in aortic valves. *Acta Biomater*. 2018;71:24-36.

doi:10.1016/j.actbio.2018.02.024

46. Félix Vélez NE, Gorashi RM, and Aguado BA. Chemical and molecular tools to probe

biological sex differences at multiple length scales. J. Mater. Chem. B. 2022;10, 7089-7098.

doi: 10.1039/D2TB00871H

47. Fogg K, Tseng NH, Peyton SR, et. al. Roadmap on biomaterials for women's health. J.

Phys. Mater. 2023;6 012501. doi: 10.1088/2515-7639/ac90ee

48. Aguado BA, Schuetze KB, Grim JC, et al. Transcatheter aortic valve replacements alter

circulating serum factors to mediate myofibroblast deactivation. Sci Trans Med,

2019;11(509):eaav3233. doi: 10.1126/scitranslmed.aav3233

Figure 1

759 **Figure 2**

760

781

Figure 3

Figure 4

Figure 5

851   **Figure 6**

852