1
2
3
4   ***Cone bipolar cell synapses generate transient versus sustained signals in parallel ON***
5   ***pathways of the mouse retina***
6
7
8
9

10 Sidney P. Kuo[1,2], Wan-Qing Yu[3], Prerna Srivastava[4], Haruhisa Okawa[3], Luca Della Santina[3],
11 David M. Berson[5], Gautam B. Awatramani[4], Rachel O.L. Wong[3], Fred Rieke[1]
12
13 [1]Department of Physiology and Biophysics, University of Washington, Seattle WA
14 [2]Department of Neuroscience, University of Minnesota, Minneapolis MN
15 [3]Department of Biological Structure, University of Washington, Seattle WA
16 [4]Department of Biology, University of Victoria, Victoria BC
17 [5]Department of Neuroscience, Brown University, Providence RI
18
19 Correspondence: Fred Rieke, rieke@uw.edu

20   *Abstract*
21
Parallel processing is a fundamental organizing principle in the nervous system, and
understanding how parallel neural circuits generate distinct outputs from common inputs is a
key goal of neuroscience. In the mammalian retina, divergence of cone signals into multiple
feed-forward bipolar cell pathways forms the initial basis for parallel retinal circuits dedicated to
specific visual functions. Here, we used patch-clamp electrophysiology, electron microscopy
and two photon imaging of a fluorescent glutamate sensor to examine how kinetically distinct
responses arise in transient versus sustained ON alpha RGCs (ON-T and ON-S RGCs) of the
mouse retina. We directly compared the visual response properties of these RGCs with their
presynaptic bipolar cell partners, which we identified using 3D electron microscopy
reconstruction. Different ON bipolar cell subtypes (type 5i, type 6 and type 7) had
indistinguishable light-driven responses whereas extracellular glutamate signals around RGC
dendrites and postsynaptic excitatory currents measured in ON-T and ON-S RGCs in response
to the identical stimuli used to probe bipolar cells were kinetically distinct. Anatomical
examination of the bipolar cell axon terminals presynaptic to ON-T and ON-S RGCs suggests
bipolar subtype-specific differences in the size of synaptic ribbon-associated vesicle pools may
contribute to transient versus sustained kinetics. Our findings indicate bipolar cell synapses are
a primary point of divergence in kinetically distinct visual pathways.

39
40   *Introduction*
41
Divergence into parallel pathways allows neural circuits to generate diverse outputs from a
common set of inputs. This functional diversity can be quite striking; in vision, for example, light
inputs initially encoded by the same collection of cone photoreceptors mediate sensitivity to
several kinds of motion, to color and to spatial structure, all in the same region of space. These
diverse functional sensitivities arise through parallel processing of light inputs, first in the retina
and subsequently in downstream visual areas. Here we focus on the mechanisms producing
transient and sustained responses – a key differentiating characteristic of retinal output cells
across species (Baden et al., 2016; Bae et al., 2018; Cleland et al., 1973; Ikeda and Wright,
1972; Roska and Werblin, 2001; Werblin, 1970).
51
The retina provides a unique opportunity to study the mechanisms that differentiate signals in
parallel pathways. We have a near-complete list of retinal cell types and substantial information
about which of these cell types participate in specific parallel pathways (Helmstaedter et al.,
2013; Masland, 2012). Under daylight conditions, retinal processing starts with cone
photoreceptors transducing light inputs and providing divergent output to ~13 types of bipolar
cell (Euler et al., 2014; Greene et al., 2016; Helmstaedter et al., 2013; Shekhar et al., 2016;
Tsukamoto and Omi, 2017). The resulting bipolar cell signals are transmitted to ~20-40 retinal
ganglion cell (RGC) types (Baden et al., 2016; Bae et al., 2018; Goetz et al., 2022). RGC inputs
are strongly shaped by the activity of a large and diverse collection of amacrine cells that
receive input from bipolar cells and provide presynaptic inhibition to bipolar cells and
postsynaptic inhibition to RGCs (Diamond, 2017).
63
This circuitry produces highly distinct responses across RGC types - with cells that sample
inputs on a variety of spatial and temporal scales, cells with distinct chromatic properties, cells
that sense the direction of moving objects, and cells that separate local from global motion
(Gollisch and Meister, 2010; Kerschensteiner, 2022). Several mechanisms contribute to these
functional differences, starting with the division of signals into ON and OFF pathways at the
cone-to-bipolar synapse (Euler et al., 2014; Slaughter and Miller, 1981). Temporal properties of
responses of different bipolar cells can also differ for some stimuli (Awatramani and Slaughter,

2000; Baden et al., 2013; Euler et al., 2014; Euler and Masland, 2000; Ichinose et al., 2014;
Ichinose and Hellmer, 2016; Wu et al., 2000). In the inner retina, integration of excitatory and
inhibitory synaptic inputs produces directionally-selective responses in specific RGC types
(Taylor and Vaney, 2002; Vaney et al., 2012) and differences in spatial weighting functions in
other RGCs (Cafaro and Rieke, 2013; Roska and Werblin, 2001; Zhang et al., 2012). Properties
of dendritic integration and RGC spike generation also differentiate responses of some RGCs
(Margolis et al., 2010; Ran et al., 2020; Wienbar and Schwartz, 2022).
78
A fundamental difference among RGCs is whether their responses are sustained or transient
(Baden et al., 2016; Bae et al., 2018; Cleland et al., 1973; Ikeda and Wright, 1972; Roska and
Werblin, 2001). Cone photoreceptors respond to changes in light intensity with graded
potentials that have both a transient and sustained component (Angueyra and Rieke, 2013;
Cangiano et al., 2012; Szikra et al., 2014). Both kinetic components of the cone response are
maintained in the firing rates of sustained cells, which track the prevailing contrast. However,
only the initial component of the cone response is preserved in the spiking response of transient
cells, which therefore respond primarily to changes in contrast. Where and how these
differences originate is not clear. Here, we show that the distinct kinetics of responses of ON-
sustained alpha and ON-transient alpha RGCs (1) persist for stimuli that elicit near-identical
responses among the relevant bipolar cells, (2) do not depend on amacrine cell-mediated
inhibition and, (3) are correlated with anatomical differences in output synapses of bipolar cell
subtypes that are presynaptic to these RGCs. Together, these findings indicate that bipolar cell
synapses are an important point of divergence in kinetically distinct visual pathways.
93
94 ***Results***
95
We focused on two ON RGC cell types, ON-transient alpha (ON-T) and ON-sustained alpha
(ON-S) RGCs, that could be readily identified in a flat-mount preparation of the isolated mouse
retina (Fig. 1A) by their characteristic spiking response to a step increment in light intensity (Fig.
1B) (Bae et al., 2018; Goetz et al., 2022; Krieger et al., 2017; Kuo et al., 2016). Based on their
light response, dendritic field size and stratification pattern in the inner plexiform layer (IPL) (see
below; Fig. S4), the ON-T RGCs we studied here likely correspond to the 6sw RGC type of Bae
et al. (2018) and the medium-field ON-transient RGC of Goetz et al. (2022).
103
We characterized the temporal spiking response properties of ON-T and ON-S RGCs by using
an extracellular cell-attached patch electrode to measure action potential firing from either cell
type while presenting a spatially uniform, randomly fluctuating light stimulus (Gaussian noise)
that allowed us to efficiently probe a variety of stimulus contrasts and temporal frequencies.
These and all subsequent physiological recordings were conducted using a photopic mean
background light level (~3300 R*/rod/s; ~2000 P*/Scone/s) to suppress rod photoreceptor
responses and therefore focus on cone-mediated signaling (Grimes et al., 2018; Ke et al.,
2014). Figure 1C shows the spiking response of example ON-T and ON-S RGCs when we
presented multiple repetitions of the same random stimulus contrast sequence; these RGC
types clearly responded differently to temporally modulated light input. In particular, noise-
evoked ON-T RGC spike responses were comprised primarily of large amplitude, less frequent
spike bursts whereas ON-S RGC responses consisted of a more even distribution of small to
large amplitude events (Fig. 1C and Fig S1A).
117

118
Figure 1*. Distinct temporal spiking response properties of ON-transient versus ON-sustained*
RGCs. (A) Maximum intensity projection z-stack fluorescence images of ON-T (left, cyan) and
ON-S (right, magenta) RGCs dialyzed with Alexa Fluor 594 during whole-cell patch-clamp
recording. Scale bar 50 µm. (B) Characteristic spiking responses to light step increment of
example ON-T and ON-S RGCs. (C) Example spiking responses to repeated trials of spatially
uniform (300 µm spot diameter) Gaussian noise stimulus (top black trace) presented with a
mean luminance of ~2400 P*/Scone/s (~3000 R*/rod/s). ON-T (cyan) and ON-S (magenta)
RGCs were recorded from sequentially in the same piece of retina. PSTHs (bottom) calculated
from 20 repetitions of noise stimulus. (D) ON-T and ON-S STAs derived from responses to non-
repeated noise stimuli. Colored lines and shaded areas show mean ± SEM STA waveforms
from ON-T (n= 15) and ON-S (n= 13) RGCs. Note time axis is reversed to facilitate comparison
with linear filters in Figures 2 and 4. (E) Quantification of STA kinetics in different RGCs. Open
circles show measurements from individual cells and filled circles with error bars show mean ±
SEM values. Significance values from Wilcoxan rank sum tests.
133

We summarized the response of each ON-T and ON-S RGC by computing the average stimulus
preceding each recorded spike (the spike-triggered average or STA stimulus) when presenting
a non-repeated Gaussian noise stimulus (Fig. 1D) (Chichilnisky, 2001) and we quantified
response kinetics from these STA waveforms by measuring the time of zero-crossing (between
trough and peak) as well as the ratio of amplitudes of the trough and peak (biphasic index). ON-
T RGC STAs were significantly more biphasic (biphasic index = 0.63 ± 0.07; mean ± S.D.; n =
15) than those measured for ON-S RGCs (biphasic index = 0.43 ± 0.11; mean ± S.D.; n = 13)
(Fig. 1E, right), consistent with their more transient response kinetics that are apparent with step
stimuli (Fig. 1B). ON-T RGC responses were also slightly delayed (zero-cross time = - 0.080 ±
0.0063 sec; mean ± S.D.; n = 15) relative to those of ON-S RGCs (zero-cross time = -0.072 ±
0.0052 sec; mean ± S.D.; n = 13) (Fig. 1C, bottom and 1E, left). We used the STA to quantify
spiking behavior because it provides a convenient model of RGC temporal response properties.
However, it should be noted that the STA alone does not capture all differences between the
RGC spike responses – e.g. the discreteness of the ON-T responses in Figure 1C.
148
149 *Bipolar cell to RGC synapses are an important site of signal diversification in ON pathways*
150
A variety of mechanisms could potentially contribute to the distinct temporal response
characteristics of ON-T versus ON-S RGCs, including differences in excitatory and/or inhibitory
input conveyed by presynaptic circuits. Parallel visual channels are initially established by the
divergence of cone signals to different cone bipolar cell subtypes, which have axons that
terminate at distinct depths within the IPL and can correspondingly contact distinct RGC types
(Euler et al., 2014; Wässle et al., 2009). We therefore sought to understand to what extent kinetic
differences in bipolar cell-mediated excitatory synaptic input could account for transient versus
sustained responses in these RGCs.
159
To do this, we used somatic whole-cell voltage-clamp recordings to measure excitatory synaptic
currents from both RGC types while stimulating the retina with the same spatially uniform
Gaussian noise stimulus we used when measuring RGC spike output (Fig. 2A). We used these
measurements to construct linear-nonlinear (LN) models comprised of a linear temporal filter (Fig.
2B and Fig. S2A) and a time-invariant (static) nonlinearity (Fig. S2B) that provide a simple
estimate of the transformation between visual input and excitatory synaptic input to each RGC
type (Kim and Rieke, 2001). LN models for each RGC type accurately predicted measured
responses to noise stimuli that were not used to construct the models, although LN models of
ON-S RGC responses performed better than those for ON-T RGCs (correlation coefficient
between average response to repeated noise stimulus and model prediction: 0.61 ± 0.057 (ON-
S, n = 9) and 0.50 ± 0.13 (ON-T, n =11), mean ± S.D.; p = 0.028 Wilcoxan rank-sum test). This
may in part be because ON-T stimulus-evoked excitatory currents were biased towards large
amplitude events compared to those of ON-S RGCs (Fig. 2A, Fig. S1B), similar to the distinct
amplitude distributions of spike events in these cell types noted above (Fig. 1C and Fig. S1A).
The linear filter captures the kinetics of the RGC response and is analogous to the time-reversed
STA derived from spike responses (Chichilnisky, 2001). As with the STAs in Figure 1D, excitatory
linear filters were slower and more biphasic in ON-T (zero-cross time = 0.10 ± 0.013 s; biphasic
ind. = 0.45 ± 0.090; mean ± S.D.; n = 13) compared to ON-S RGCs (zero-cross time = 0.083 ±
0.0074 s; biphasic ind. = 0.31 ± 0.10; mean ± S.D.; n = 15) (Fig. 2C). This latter difference is
consistent with more transient light-evoked excitatory synaptic input to ON-T RGCs. Indeed, light
step-evoked excitatory currents to ON-T RGCs decayed rapidly to baseline levels within a few
hundred milliseconds, even in the continued presence of light input, whereas excitatory inputs to
ON-S RGCs had a sustained component that persisted for the duration of the light step (Fig. 2D-
E).
184

These results show that transient versus sustained kinetics are already present in the excitatory
pathways that provide input to ON-T and ON-S RGCs. These findings do not rule out additional
contributions to the distinct spiking behavior of these RGC types from amacrine cell-mediated
inhibitory synaptic input to the RGCs or from different intrinsic spiking properties. However, the
clear differences we observed in excitatory synaptic input between ON-T and ON-S RGCs
indicates that bipolar cell pathways have a central role in establishing transient versus sustained
kinetics.

192
Figure 2. *Distinct excitatory synaptic input to ON-T and ON-S RGCs. (A) Excitatory synaptic*
currents measured in example ON-T (cyan) and ON-S (pink) RGCs in response to same
Gaussian noise stimulus as in Fig. 1. Traces show average currents from ten repetitions of the
same stimulus and are baseline subtracted to facilitate comparison. (B) Linear filters computed
from responses to non-repeated noise stimulus presentation. Lines and shaded areas show
mean ± SEM of responses from ON-T (n= 13) and ON-S (n= 15) RGCs. (C) Quantification of
filter kinetics in different RGCs. (D) Excitatory currents measured from example ON-T and ON-S
RGCs in response to a 0.5 second step increase in light intensity (50% contrast; mean response
to five repetitions of stimulus). (E) Quantification of step response kinetics (n= 19 ON-T RGCs;
n= 13 ON-S RGCs). In (C) and (E) open circles show measurements from individual cells and
filled circles with error bars show mean ± SEM values. Significance values from Wilcoxan rank
sum tests.
205
206
207
208

ON-T and ON-S RGCs receive input from distinct bipolar types

In general, retinal circuits dedicated to transient versus sustained signaling are located at the center and borders of the IPL, respectively (Baden et al., 2016; Bae et al., 2018; Franke et al., 2017; Roska and Werblin, 2001). We compared stratification patterns of ON-T and ON-S RGCs in a subset of whole-cell recordings by filling cells with a fluorescent dye and imaging the dendrites of physiologically-characterized RGCs using two-photon microscopy. We assessed IPL depth in live tissue by using retinas from a transgenic mouse line in which type 5 bipolar cells strongly express EGFP (*Gjd2-EGFP* mice). These EGFP+ bipolar cells have axons that occupy a narrow band just above and overlapping with the dendrites of ON starburst amacrine cells (Fig. S3A) and express calcium binding protein 5 (CaBP5) (Fig. S3B-D) (Ghosh et al., 2004). Based on the regular spacing of their somas, these cells likely correspond to one of the three subtypes of type 5 bipolar cells previously identified in mouse retina (Greene et al., 2016; Rodieck, 1991) (Fig. S3E-F). Imaging of RGCs together with EGFP+ axon terminals in *Gjd2-EGFP* mice revealed that ON-T and ON-S RGC dendrites occupy distinct, mostly non-overlapping regions of the IPL (Fig. S4).

The distinct IPL stratification of ON-T and ON-S RGCs indicates they receive input from different bipolar cell populations. A recent connectomic analysis that used a published electron microscopy data set from mouse retina (Ding et al., 2016) found that type 6 and type 7 bipolar cells account for ~50% and 46% of the excitatory synapses formed upon ON-S RGC dendrites, respectively (Sabbah et al., 2018). However, the bipolar cell subtype(s) presynaptic to ON-T RGCs have not previously been identified. Our observation that ON-T RGC dendrites occupy a narrow band in the IPL that very closely matches the stratification pattern of EGFP+ type 5 bipolar cell axons in *Gjd2-EGFP* mice (Fig. S5) suggests that these bipolar cells likely provide a substantial fraction of the excitatory input to ON-T RGCs. To test this possibility, we transfected retinas from *Gjd2-EGFP* mice with a plasmid encoding PSD-95 fused to the red fluorescent reporter mCherry to label postsynaptic sites of excitatory input (Fig. S5). We quantified the fraction of putative synapses, defined as sites with overlapping PSD-95 and EGFP fluorescent signals, that arose from *Gjd2-EGFP+* axon terminals using confocal microscopy (Schwartz et al., 2012). From this analysis, we concluded that the labeled type 5 bipolar cells in *Gjd2-EGFP* mice provide ~40% of the ribbon synaptic input to ON-T RGCs (Fig. S5D).

We next performed correlative fluorescence imaging and serial block-face scanning electron microscopy (SBFSEM) to identify which type 5 bipolar subtype is labeled in Gjd2-EGFP mice and to further characterize what other bipolar cell types provide the remaining input to ON-T RGCs. We first used 2P microscopy to image the dendritic morphology of two dye-filled ON-T RGCs that were identified from their light-evoked spike responses in flat-mounted retina from *Gjd2-EGFP* transgenic mice (Fig. 3A). After 2P imaging, we fixed the retina and used the near infrared branding (NIRB) technique to create fiducial marks in the tissue around the dye-filled ON-T RGCs (Bishop et al., 2011). After acquiring the SBFSEM volume, we located the somas of the two ON-T RGCs using these fiducial marks and reconstructed the dendritic arbors of both cells (e.g. Fig. 3B). The dendritic morphologies from our EM reconstruction matched those from the light microscopy images. This suggests that we successfully identified the two ON-T RGCs recorded physiologically in our EM volume. We chose the two longest dendritic arbors from the RGC soma to the dendritic tip and identified all the synapses from bipolar cells along these dendrites (Fig. 3C). We then reconstructed all the presynaptic bipolar cells and classified them according to their axon morphology and stratification (Fig. 3D). Careful examination of axonal tiling among type 5i, 5o and 5t subtypes provided additional confirmation of the classification of these type 5 bipolar cells (Fig. S6A) (Greene et al., 2016). When comparing the EM images with light microscopy images, it is clear that *GJD2-GFP+* bipolar cells are type 5i bipolar cells (Fig. S6B).

261
Figure 3. *Identification of bipolar cell subtypes presynaptic to ON-T RGCs. (A) Maximum-projections z-stacks of two dye-filled ON-T RGCs (green) in wholemount retina from Gjd2-EGFP (red) transgenic mice. (B) Traced skeletons of dendritic arbors of the two ON-T RGCs from EM volume. (C) An EM micrograph showing a ribbon synapse (arrowhead) on a BC axonal terminal onto the dendrite of a ON-T RGC. (D) An example skeleton for each presynaptic BC type (types 5i, 5t, 5o, X, 6 and 7) of ON-T RGCs. Top: whole-mount view, Bottom: side view. (E) Bar plot showing the proportion of input synapses to ON-T RGCs from each BC type. Mean ± SEM (n=3 RGCs, 2 from this study and one from Ding et al. (2016)).*

Results from the SBFSEM reconstructions agreed closely with those from the light microscopy approach described above. Type 5i bipolar cell axon terminals provided the largest fraction of synapses onto the two ON-T RGCs within our SBFSEM volume (~30-35%). Other inputs arose from type 5o (~10%), type 5t (~5-15%), XBC (~15-30%), and, somewhat surprisingly given the differences in stratification, from type 6 (~20%) bipolar cells (Fig. 3E). We confirmed this connectivity pattern by additionally analyzing bipolar cell synaptic inputs to a putative ON-T RGC

277 (RGC7089) in the same publicly available SBFSEM volume (Ding et al., 2016) that was used by
278 Sabbah et al., (2018) to map inputs to ON-S RGCs (Fig. 3E).
279

280
Figure 4. *Bipolar cell subtypes presynaptic to ON-T and ON-S RGCs have indistinguishable*
temporal response properties. (A) Maximum-projection z-stacks of fluorescent dye-filled bipolar
cells targeted for electrophysiological recording in retinal slices from Gjd2-EGFP (left), Grm6-
tdTomato (middle), and Gus8.4-EGFP (right) transgenic mice. (B) Responses to spatially
uniform Gaussian noise stimulation (same as in Figs. 1-2) in example type 5i (cyan), type 6
(magenta) and type 7 (green) bipolar cells. Voltage traces are average responses to ten
repetitions of the noise stimulus. (C) Linear filters computed from responses to non-repeated
noise stimulus presentation. Lines and shaded areas show mean ± SEM of responses (type 5i,
n= 11; type 6, n= 7; type 7, n= 6). (D) Quantification of response kinetics from linear filters. No
significant differences across types (p = 0.19 zero-cross times; p= 0.89 biphasic index; Kruskal-
Wallis test). (E) Responses to step light increment. Lines and shaded areas show peak-
normalized mean ± SEM responses across type 5i bipolar cells (n= 8) and type 6 bipolar cells
(n= 5). (F) Quantification of step response kinetics. Time to peak and steady-state response
amplitude were not different between type 5i and type 6 bipolar cells (p= 0.72 and p= 0.35,
respectively; Wilcoxan rank sum test).
296
297
298

299 *Bipolar cell light responses differ minimally across types*
300
Are transient versus sustained excitatory inputs inherited from the response properties of the
different bipolar cells presynaptic to ON-T and ON-S RGCs? Or is the distinct excitatory
synaptic input these RGCs receive primarily due to differences at the bipolar cell to RGC
synapse? To answer these questions, we compared the light-driven response properties of the
bipolar cell subtypes that comprise most of the excitatory input to ON-T or ON-S RGCs. We
made targeted electrophysiological recordings from type 5i, type 6 and type 7 bipolar cells in
retinal slices from three transgenic mouse lines in which these different bipolar cell subtypes
express fluorescent proteins (see Methods). In these experiments, we used somatic whole-cell
perforated-patch recordings to limit perturbation of the G-protein signaling cascade that
mediates ON bipolar cell light responses.
311
Somatic resting membrane potentials did not differ significantly among these bipolar cell types
(mean ± S.D. V_{rest}: -38 ± 6.3 mV in type 5i (n= 11); -33.7 ± 6.3 mV in type 6 (n= 7); - 36.5 ± 4.7
mV in type 7 (n=6); Kruskal-Wallis test, p= 0.35). We characterized light responses using the
same spatially uniform Gaussian noise stimulus (Fig. 4B) and linear-nonlinear model analysis
(Fig. 4C and Fig. S7) as in RGC recordings to facilitate direct comparison of results from bipolar
cells and RGCs (Kim and Rieke, 2001; Rieke, 2001). LN models of bipolar cell responses
accurately predicted responses to noise stimuli (correlation coefficient between average
response to repeated noise stimulus and LN model prediction = 0.75 ± 0.15 for n = 18 bipolar
cells (n = 13 type 5; n = 3 type 6; n = 2 type 7)). Unlike for ON-T and ON-S RGCs, temporal
characteristics of type 5i, type 6 and type 7 bipolar cell linear filters were indistinguishable from
each other under our recording conditions (Fig 4B-D). Although there was some variability in the
temporal response properties across the population of bipolar cells we recorded, these
differences were not associated with morphologically and molecularly defined bipolar cell types.
Rather, we observed variability in response kinetics within each bipolar cell subtype and
temporal responses thus overlapped substantially across different type 5i, type 6 and type 7
bipolar cells (Fig. 4 and Fig. S7).
328
Responses to light steps measured in a subset of type 5i and type 6 bipolar cell recordings
confirmed our conclusion from Gaussian noise stimuli that different bipolar cell types exhibit
nearly identical light response kinetics under our recording conditions (Fig. 4E-F). Unlike the
clear differences in light step-evoked excitatory currents we measured in ON-T and ON-S RGCs
(Fig. 2D-E), type 5i and type 6 bipolar cells both had sustained responses to light steps. These
findings indicate that mechanisms upstream of the generation of the cone bipolar cell voltage
response, such as synaptic transmission from cones to cone bipolar cells and intrinsic response
properties of the cone bipolar cells, cannot explain the kinetically distinct responses in parallel
ON channels mediated by type 5i, type 6 or type 7 bipolar cells.
338
339 *Presynaptic inhibition does not establish kinetic differences in bipolar cell-mediated excitation*
340
Bipolar cells participate in diverse inhibitory circuits with amacrine cells, which are comprised of
>40 morphologically and functionally distinct cell types (Diamond, 2017; Helmstaedter et al.,
2013). Recent work suggests that amacrine cell-mediated presynaptic inhibition is critical to
bipolar cell subtype-specific patterns of light-evoked glutamate release (Franke et al., 2017).
Our bipolar cell voltage recordings, which should reflect a combination of light-evoked excitatory
and inhibitory input, argue against the involvement of presynaptic inhibition in generating
differential bipolar cell synaptic output between type 5i, type 6 and type 7 bipolar cells. However,
the slicing procedure we used to prepare tissue for bipolar cell recordings may damage some
inhibitory connections, especially long-range amacrine cell inputs. Distance-dependent

attenuation of voltage signals between bipolar cell axon terminals and the somatic recording site
could also limit our ability to accurately measure the influence of presynaptic inhibition in these
recordings. We therefore performed additional anatomical and pharmacological experiments to
investigate a possible role for presynaptic inhibition in generating kinetically distinct excitation to
ON-T and ON-S RGCs.

355
Figure 5. Presynaptic inhibition does not establish kinetic differences between bipolar cell
inputs to ON-T versus ON-S RGCs. (A) Example EM micrograph showing a ribbon dyad (red
arrowhead) between a BC and an ON-T RGC and an amacrine cell, with an adjacent feedback
inhibitory synapse (yellow asterisk).　(B) Bar plots showing the percentage of RGC/AC ribbon
dyads with feedback inhibition. Mean ± SEM (n=2 ON-S RGCs, 2 ON-T RGCs) (C) Excitatory
currents measured in response to light step increment in control conditions (cyan) and in the
presence of GABA receptor antagonists (10 µM SR95531 and 50 µM TPMPA; dashed gray) in
an example ON-T RGC. Traces are mean response from six repeats of stimulus presentation.
(D) Summary of steady-state amplitudes measured from step responses in control or with
GABA receptor antagonists. No significant difference between conditions (p= 0.063; Wilcoxan
signed rank test) (E) Control excitatory currents (cyan) or with addition of strychnine (1 µM;
dashed gray) to bath in an example ON-T RGC. (F) Summary of strychnine experiments. No
significant difference between conditions (p= 0.438; Wilcoxan signed rank test). In (D) and (F),
open circles with lines show measurements from individual ON-T RGCs. Filled circles with error
bars show mean ± SEM.
371

Many bipolar cell output synapses are organized into a "dyad" arrangement in which a bipolar
cell ribbon synapse releases glutamate onto the processes of two different postsynaptic cells. At
dyads, bipolar cell output synapses can be presynaptic to the processes of two amacrine cells,
two RGCs, or an amacrine cell and a RGC (Cohen and Sterling, 1990; Dowling and Boycott,
1966; McGuire et al., 1984; Yu et al., 2023). Dyad synapses comprised of an amacrine cell and
a RGC may shape bipolar cell glutamate release dynamics onto the RGC partner because
glutamatergic excitation of the postsynaptic amacrine cell can elicit release of inhibitory
neurotransmitters (GABA or glycine) from the amacrine cell processes that feed back onto the
bipolar cell axon terminal. This form of inhibitory feedback control of bipolar cell output, termed
reciprocal inhibition, could potentially contribute to differences in glutamate release onto ON-T
and ON-S RGCs.
383
We analyzed bipolar cell dyads with postsynaptic amacrine cell processes and either ON-T or
ON-S RGC dendrites in SBFSEM volumes to examine whether reciprocal inhibition is likely to
generate more transient glutamate release onto ON-T RGCs. We defined putative reciprocal
synapses as bipolar cell/amacrine cell/RGC dyads in which we observed clustered vesicles
within the amacrine cell process that were within close proximity of the bipolar cell ribbon
synapse (Fig. 5A). Our ultrastructural analysis revealed that the prevalence of reciprocal
synapses was similar at dyad synapses with postsynaptic ON-T versus ON-S RGCs (Fig. 5A-B).
This finding is not consistent with an important role for feedback/reciprocal inhibition in
generating differential bipolar cell output onto the different RGC types.
393
We also performed pharmacological experiments to further examine what role, if any, inhibition
may have in establishing transient glutamate release onto ON-T RGC dendrites. We used
ACSF in place of Ames solution for the dissection and maintenance of tissue for these
experiments. ACSF is often used in retina experiments, including the glutamate release
measurements describe below, and we wanted to check that our results generalized across
conditions. Light step-evoked currents in both ON-T and ON-S RGCs were more sustained than
responses measured using Ames' solution (Fig. S8A-B). Nonetheless, the relative differences
between RGC types appear to be maintained across different experimental conditions.
402
Blockade of $GABA_A$ and $GABA_C$ receptors via bath co-application of SR95531 and TPMPA
enhanced both the initial transient component and the sustained component of light step-evoked
excitatory synaptic currents in ON-T RGCs (Fig. 5C). The strong enhancement of the initial
phase of the response we observed is consistent with prior work showing that $GABA_C$ receptor-
mediated inhibition limits multivesicular release from bipolar cells (Sagdullaev et al., 2006). On
average, block of GABA receptors did not significantly change in the ratio of sustained versus
transient components of excitatory inputs to ON-T RGCs compared to control conditions (Fig
5D). If anything, these drugs tended to make excitatory currents more transient, which is the
opposite effect expected if presynaptic GABA receptors contribute to generating transient
responses in these cells. Thus, GABAergic inhibition appears to control response gain but not
kinetics of bipolar cell synaptic output to ON-T RGCs. Similarly, we found that glycinergic
inhibition is not likely to contribute importantly to more transient glutamate release from bipolar
cells presynaptic to ON-T RGCs; bath application of the glycine receptor antagonist strychnine
did not significantly alter the kinetics of light step-evoked excitatory currents in these cells (Fig.
5E-F). Taken together with our ultrastructural analysis, these results show that amacrine cell-
mediated inhibition of bipolar cell axon terminals does not contribute to kinetic differences in
excitatory synaptic input to ON-T and ON-S RGCs.
420
421
422

423
Figure 6. *Distinct glutamate input kinetics at ON-S vs ON-T RGC dendrites. (A) ON-S RGC*
dendrites expressing iGluSnFR in a KCNG-Cre mouse were filled with sulforhodamine 101 (left)
after measuring their spiking response to a 200 µm spot extracellularly (Right). Yellow bands
indicate stimulus duration. (B) iGluSnFR expressing ON-S and ON-T dendrites at different depths
of the inner-plexiform layer (Left) were imaged using two-photon microscopy. Example iGluSnFR
signals extracted from small regions of interest (ROIs 1-8) from ON-S and ON-T RGC dendrites
illustrate the kinetic differences in their respective inputs. Black, mean responses; gray, individual
trials. (C) Distribution of steady-state response amplitudes of the individual (gray) and average
(black) ROIs from different recordings done from ON-S and ON-T dendritic field of views evoked
by a 200 µm static spot (n = 7 FOVs, 6 retinas for ON-S and 5 FOVs, 5 retinas for ON-T RGC
dendrites, *p<0.001; t-test).

Bipolar cell glutamate release differs across types
436
Taken together, the bipolar cell and RGC recordings in Figures 2 and 4 indicate that bipolar cell
synapses have a key role in the separation of visual signals into transient or sustained channels
within ON pathways in the retina. Presynaptic and/or postsynaptic mechanisms could contribute
to the distinct kinetics of bipolar cell-mediated excitatory input to ON-T and ON-S RGCs. To
investigate whether presynaptic differences in glutamate release from bipolar cell axon
terminals shape RGC response kinetics, we expressed the fluorescent glutamate reporter
iGluSnFR on the dendrites of ON-T and ON-S RGCs. Using two-photon glutamate sensor
imaging, we found that light-evoked glutamate signals had significantly faster kinetics at ON-T
RGC dendrites (steady-state amplitude as % of peak amplitude = 21 ± 13; mean ± S.D.; n = 224
ROIs) compared to those associated with ON-S RGC dendrites (steady-state amp. (% of peak
amp.) = 44 ± 12; mean ± S.D.; n = 307 ROIs) (Fig. 6). These results show that differences in
glutamate release from presynaptic bipolar cells contribute to transient versus sustained ON
RGC responses. However, these experiments do not rule out the possibility that additional
postsynaptic mechanisms, for example differential glutamate receptor expression, may also
contribute to kinetic differences between RGC types.
452
Although whole-cell excitatory currents in ON-T and ON-S RGCs (Fig. 2, Fig. S8) and iGluSnFR
signals measured from ON-T and ON-S RGC dendrites (Fig. 6) both exhibited clear kinetic
differences between the RGC types, there were quantitative differences in the response to light
steps between the different measurement approaches. As described above, these differences
appear to arise from differences in the external solution (Ames vs ACSF). The iGluSnFR
experiments used an ACSF solution, like the patch-clamp experiments in Figs. 5 and S8. Both
iGluSnFR responses and excitatory synaptic inputs measured in ACSF showed a larger
sustained component of the step response at both ON-T and ON-S dendrites compared to
currents measured in Ames (compare Fig. 2D-E, Fig. S8A-B and Fig. 6B-C). This observation
highlights the importance of comparing responses between different RGC types, as well as
between RGCs and bipolar cells, under identical recording conditions.
464
465   Importantly, the kinetics of glutamate release measured using iGluSnFR are much more similar
466   to the excitatory synaptic inputs to ON-T and ON-S RGCs than they are to the relevant bipolar
467   voltage responses (Fig 4). This indicates that the difference between the ON-T and ON-S
468   responses originates at least in part in the conversion of bipolar voltage to glutamate release.
469
470   *Ribbon size and vesicle replenishment*
471
Prior work has shown that bipolar cells can exhibit profound short-term depression of glutamate
release due to use-dependent depletion of the readily-releasable pool of synaptic vesicles
(Burrone and Lagnado, 2000; Singer and Diamond, 2006, 2003; von Gersdorff and Matthews,
1997). Strong vesicle depletion at the cone bipolar cell synapses onto ON-T RGCs could
generate more transient responses in these cells compared to ON-S RGCs. To examine this
possibility, we used a paired flash stimulus paradigm, in which two brief light flashes separated
by varying time intervals were presented while measuring excitatory synaptic currents in ON-T
or ON-S RGCs. As in Figures 1, 2 and 4, we used Ames' medium for tissue preparation and
maintenance in these experiments.
481
Flash-evoked excitatory synaptic currents were suppressed in both RGC types at short inter-
flash intervals, consistent with incomplete recovery from vesicle depletion at the time of the
second flash (Fig. 7A top/middle, B). However, the extent of suppression was much stronger for
currents measured in ON-T RGCs compared to ON-S RGCs. We quantified the extent of

suppression using a paired flash ratio, which was calculated as the peak amplitude of the
response to the second flash divided by the peak amplitude of the response to the first flash. At
the shortest interval we tested (50 ms), the response to a second flash was almost completely
suppressed in ON-T RGCs (paired flash ratio = 0.04 ± 0.01; mean ± SEM) whereas the
response to a second flash in ON-S was suppressed considerably less (paired flash ratio = 0.29
± 0.04). To rule out the possibility that a reduction in bipolar cell voltage response to the second
flash could account for the strong paired flash suppression of excitatory input to RGCs, we also
measured type 5i bipolar cell responses to the paired flash stimuli. The amplitude of flash-
evoked voltage responses in type 5i bipolar cells was unchanged by a preceding flash at all time
intervals (Fig. 7A bottom, C), which argues that paired flash suppression arises at the bipolar
cell synaptic terminal. These results are consistent with the idea that differences in the extent of
stimulus-evoked vesicle pool depletion in bipolar cells could underlie transient versus sustained
excitatory input to ON-T and ON-S RGCs.
499

500
501
Figure 7. *Stronger flash-evoked suppression of excitatory currents in ON-T versus ON-S RGCs.*
(A) Paired flash-evoked excitatory currents measured at different inter-flash intervals (50, 100,
200, 400 ms) in an example ON-T (top; cyan) and an ON-S (middle, magenta) RGC. Bottom
black traces show voltage responses to same stimulus measured in an example type 5i bipolar
cell. Arrows at top indicate timing of first flash (black) and second flashes (gray). Flashes were
400% contrast and 10 ms duration. Traces show mean responses to five repeats of each paired
flash interval. (B) Filled circles with error bars show mean ± SEM paired-flash ratios at different
inter-flash intervals across all ON-T RGCs (cyan; n= 6), ON-S RGCs (magenta; n= 5) and type
5i bipolar cells (black; n= 6).
511
512

513
514
Figure 8. *Bipolar cell subtype-specific differences in ribbon size. (A) 3D reconstructions*
showing segmented axon stalk and axonal terminal (gray) and all the synaptic ribbons within the
axonal terminal a type 5i (cyan) and a type 6 (magenta) BC. (B) Violin plot showing distribution
of ribbon volumes (each data point is a ribbon) for the type 5i and type 6 BC in (A). The boxes
indicate the interquartile range (thick lines, 25%-75%), median value (white dot), and 1.5 times
interquartile range (thin line). Shaded gray areas represent rotated kernel density plots. (C) 3D
reconstructions showing segmented dendritic arbor (gray) and all the synaptic ribbons
presynaptic to these segments of ON-T (cyan) or ON-S (magenta) RGC dendrites. (D) Violin
plot with scattered dots showing ribbon volumes for each ribbon presynaptic to a stretch of
reconstructed ON-T or ON-S RGC dendrite. Each ribbon (dot) is colored differently for each BC
type. (E) An EM micrograph (left) and 3D reconstruction (right) of a ribbon and its adjacent
vesicles. The ribbon is indicated by the red arrow in the EM image. In the 3D reconstruction,
yellow dots indicate the adjacent vesicles surrounding the presynaptic ribbon (red) in the BC
terminals (gray). Postsynaptic ON-T RGC is in blue. (F) Scatter plot showing ribbon size in
volume in relationship with the number of adjacent vesicles. Gray line represents the linear
regression model fitted to the data. Linear regression: $\beta = 6.98e{-}06$, $p < 0.001$, $r^2 = 0.66$.

We next looked for ultrastructural correlates of differential synapse function in bipolar cell axon
terminals reconstructed from SBFSEM volumes. Interestingly, type 6 bipolar cell synaptic
ribbons were ~2-3 fold larger than those of types 5i bipolar cells (Fig. 8A-B). These differences
did not depend on the identity of the postsynaptic RGC partner (Fig. 8D). Ribbons in XBCs, type
5o and type 5t bipolar cells were also smaller than type 6 bipolar cell ribbons (Fig. 8D). This
bipolar cell subtype-specific difference in ribbon size and the difference in ratio of type 6 to type
5i input between RGC types meant that ribbons were on average smaller at synapses made on
ON-T compared to ON-S dendrites (Fig. 8C-D).
539
We found that ribbon size of several ON bipolar cell types is strongly correlated with the total
number of vesicles that are in close contact with the synaptic ribbon across different bipolar cell
types (Fig. 8E-F). Ribbon-associated vesicles are thought to correspond to the releasable pool
of vesicles, comprised of both a readily-releasable pool that are docked at the presynaptic
active zone as well as a reserve pool of vesicles that are bound to the ribbon at more distal sites
(Lagnado and Schmitz, 2015). These two vesicle pools exhibit distinct release kinetics, with the
readily-releasable pool corresponding to the fastest phase of glutamate release and the reserve
pool contributing to slower glutamate release that occurs with prolonged stimuli (Mennerick and
Matthews, 1996; Neves and Lagnado, 1999). Our observations that the average excitatory
synaptic input to ON-T RGCs is smaller and that most excitatory input to ON-T cells originates
from bipolar synapses with fewer ribbon-associated vesicles are consistent with a role for
vesicle depletion in establishing transient glutamate release onto these RGCs. The larger pool
of ribbon-associated vesicles in type 6 bipolar cell terminals, which comprise ~ 55% of the
synaptic input to ON-S RGCs, may support sustained synaptic transmission onto these RGCs.
554
555 ***Discussion***
556
In this study, we sought to understand how kinetically distinct visual pathways are established in
the retina by directly comparing the visual response properties of transient and sustained ON
RGCs with their presynaptic bipolar cells. Our results show that bipolar cell synapses have a
key role in establishing different temporal channels in ON circuits of the retina. Further,
amacrine cell-mediated presynaptic inhibition does not appear to account for transient versus
sustained bipolar cell synaptic output. Rather, bipolar cell subtype-specific differences in
synaptic release properties likely generate kinetically distinct output to postsynaptic targets. Our
anatomical analysis suggests that these differences may at least in part be due to differences in
synaptic vesicle pool size.
566
A key aspect of our work is that we measured RGC and bipolar cell responses using identical
stimuli and under the same experimental recording conditions. To our knowledge, a direct
comparison of RGC responses with those of their presynaptic bipolar cells has not previously
been reported. The importance of measuring RGC and bipolar cell responses under the same
conditions is highlighted by our observation that excitatory synaptic currents measured in ON
alpha RGCs exhibited quantitatively different kinetics, although relative differences between
types were preserved, when measurements were made from retinas maintained in ACSF
versus Ames' bath solution (compare Fig. 2 (Ames') and Figs. 6 and S8 (ACSF)).
575
We found that the bipolar cell subtypes that provide the major input to ON-T (type 5i) or ON-S
RGCs (types 6 and 7) had indistinguishable light-evoked voltage responses (Figure 4). This was
not entirely surprising given that most or all ON cone bipolar cell subtypes are indirectly coupled
to other nearby ON cone bipolar cells via electrical synapses between bipolar cell axon
terminals and the distal processes of AII amacrine cells. These electrical synapses form the
basis for the primary rod pathway that mediates vision under dim light conditions, but lateral

signals can also flow among neighboring bipolar cells under conditions where cone signaling
dominates (Demb and Singer, 2012). Indeed, a substantial fraction (~30-50%) of light-evoked
responses in the bipolar cell types we investigated here can be attributed to lateral signals that
arrive via electrical synapses at photopic light levels (Kuo et al., 2016). Each AII amacrine cell is
electrically coupled to multiple ON cone bipolar subtypes as well as to other AII amacrine cells
(Marc et al., 2014; Tsukamoto and Omi, 2017; Veruki and Hartveit, 2002a, 2002b). Different ON
cone bipolar cell subtypes may also be directly coupled via gap junctions to each other
(Sigulinsky et al., 2020; Tsukamoto and Omi, 2017).
590
Direct or indirect (via AII) coupling may limit the extent to which different bipolar cells can
generate subtype-specific voltage responses. However, our findings differ from a previous study
that found some differences among mouse ON cone bipolar cell subtypes. Ichinose et al.,
(2014) reported that a subset of type 5 bipolar cells, which the authors termed type 5f, and type
7 bipolar cells had more transient responses than type 6 bipolar cells. Methodological
differences may account for the discrepancy between our results and the previous work by
Ichinose et al., (2014). For example, the previous study measured bipolar cell responses using
whole-cell recordings and included inhibitory receptor antagonists in their experimental solutions
whereas we used gramicidin perforated-patch recordings and left inhibitory signaling intact.
Regardless of the reasons for the different findings, our observation that bipolar cell subtypes
exhibited similar responses under the same experimental conditions for which we observed
obvious differences among postsynaptic RGCs argues that mechanisms upstream of bipolar
cell glutamate release (e.g. cone to bipolar cell synapses or intrinsic voltage response
properties of bipolar cells) are unlikely to have a major role in shaping RGC type-specific
responses for the stimuli we used here.
606
A limitation of our work is that we did not record from every one of the bipolar cell subtypes that
are presynaptic to ON-T RGCs. Instead, we measured responses from type 5i and type 6
bipolar cells, which together comprise >~50% of the synaptic input to ON-T RGC dendrites. It is
possible that the other presynaptic bipolar cell types (type 5o, type 5t and XBC) may have more
transient light-evoked voltage responses that could contribute to the transient excitation
measured in ON-T RGCs. In fact, a previous study in mouse retina found that XBCs and a
subset of type 5 bipolar cells express voltage-gated Na^+ channels (Hellmer et al., 2016). These
channels may enhance response transience or might even contribute to the generation of light-
driven all-or-none spiking responses in these bipolar cells (Ichinose et al., 2014; Puthussery et
al., 2013; Saszik and DeVries, 2012). Despite the possibility that intrinsic response properties of
the bipolar cell types we did not record from could contribute to the generation of transient
responses in ON-T RGCS, we consider it unlikely that this alone could not account for the
kinetics of the excitatory synaptic currents we measured in these RGCs. Type 5i and type 6
bipolar cells have a significant sustained component of their light step-evoked voltage response
(Fig. 4E-F), whereas excitatory currents measured for the same stimulus and under the same
conditions in ON-T RGCs had little to no sustained component (Fig. 2D-E). Even if the synapses
formed by type 5o, type 5t and XBC provide a major fraction of the functional synaptic input to
ON-T RGCs (despite our anatomical estimate that together they comprise <50% of the total
ribbon input), we would still expect to observe some sustained excitatory input if type 5i and
type 6 bipolar cell synaptic output follows the voltage response of these cells. Thus, a marked
transformation of the sustained bipolar cell membrane voltage response to a transient synaptic
current must take place. Our measurements of extracellular glutamate at ON-T and ON-S
dendrites (Fig. 5) suggest much of this transformation occurs at the level of bipolar cell
glutamate release. It is interesting to note in this context that ON-T and ON-S RGCs both
receive input from type 6 bipolar cells (comprising ~20% and ~50% of total synaptic input,
respectively). Due to differences in dendrite stratification, type 6 bipolar cell synapses onto ON-

T RGCs arise from a region of the axon that is located more proximal to the bipolar cell soma
than the synapses formed onto ON-S RGCs, which are found within the distal axon lobules.
However, we did not observe any differences in ribbon size at distal versus proximal locations in
type 6 bipolar cell axons (not shown). Whether postsynaptic target-specific synapse function, as
has previously been reported in salamander retina (Asari and Meister, 2014, 2012), contributes
to signal diversification in circuits involving type 6 bipolar cells will require further investigation.
639
Our results indicate that amacrine cell-mediated inhibition at bipolar cell synaptic terminals is not
necessary for generating transient glutamate release kinetics onto ON-T RGC dendrites. This is
seemingly at odds with prior work that has noted an essential role for surround inhibition in
establishing type-specific bipolar cell glutamate release dynamics (Franke et al., 2017).
However, we note that our analysis here focused only on whether presynaptic inhibition shapes
the kinetics of excitatory inputs to ON-T RGCs. Our anatomical and pharmacological
experiments together argue that presynaptic inhibition is not an important factor in the
transformation from sustained bipolar cell voltage responses to transient excitatory synaptic
currents in ON-T RGCs. This is not to say that amacrine cell-mediated inhibition does not have
a role in generating subtype-specific response properties for other aspects of bipolar cell
function. Indeed, inhibition does have in important role in establishing differential spatial tuning
of excitatory inputs to the RGC types we studied; excitatory synaptic currents in ON-T RGCs
exhibit stronger surround inhibition than ON-S RGCs, and this surround inhibition is reduced by
bath application of GABA receptor antagonists (Fig. S8C).
654
We conclude that transient versus sustained excitatory inputs to ON-T and ON-S RGCs most
likely arise from bipolar cell subtype-specific differences in synaptic release properties.
Excitatory inputs to ON-T RGCs exhibited stronger paired flash depression than those to ON-S
RGCs (Fig. 7), which indicates distinct vesicle cycling dynamics at the bipolar cell axon
terminals presynaptic to the different RGC types. Our ultrastructural analysis suggests that
bipolar subtype-dependent differences in ribbon-associated vesicle pool size could be a factor in
establishing transient versus sustained glutamate release kinetics (Fig. 8). This finding is
consistent with the idea that sustained transmission from ribbon synapses is determined by the
rate of replenishment of the readily-releasable pool of synaptic vesicles following exocytosis
(Lagnado and Schmitz, 2015). It is important to note, however, that type 7 bipolar cells (~45% of
excitatory input to ON-S RGCs) have small synaptic ribbons, yet iGluSnFR imaging of type 7
bipolar cell axon terminals found that these cells exhibit sustained glutamate release (Srivastava
et al., 2022). Additional related mechanisms that we did not directly investigate here may
contribute as well. For example, presynaptic resting potential has a strong effect on steady-state
vesicle release and thus stimulus-evoked release dynamics. Exocytosis that occurs when
bipolar cells rest at voltages near release threshold can lead to a depletion of readily-releasable
vesicles that limits the amount of release that occurs upon further light-induced depolarizations
(Grimes et al., 2014; Jarsky et al., 2011; Oesch and Diamond, 2011). Thus, an initial phasic
component of release can be reduced by depolarized resting potentials whereas hyperpolarized
potentials can facilitate greater occupancy of vesicles in the releasable pool, which can enhance
response transience. Bipolar subtype-dependent differences in release threshold due to
differential Ca^{2+} channel expression/regulation (Pangrsic et al., 2018) and/or coupling between
Ca^{2+} channel activity and vesicle release (Burrone et al., 2002; Burrone and Lagnado, 2000)
could also impact release dynamics. Several aspects of excitatory synaptic currents in the two
RGC types we studied here are consistent with a situation in which bipolar cell inputs to ON-T
RGCs have a higher threshold for glutamate release compared to those at ON-S RGCs; ON-T
RGCs receive less spontaneous excitatory synaptic input than ON-S RGCs (Kuo et al., 2016)
and stimulus-evoked EPSCs are slower, sparser and more biased towards larger amplitude
events in ON-T RGCs compared to those in ON-S RGCs (Fig. 2, Fig. S1B). Further

investigation will be required to determine whether this is due to differences in axon terminal
resting potential and/or other presynaptic factors. We did not find evidence for subtype-specific
differences in resting membrane potential among type 5i, type 6 and type 7 bipolar cells, but our
somatic voltage recordings likely have limited ability to resolve small voltage differences at the
bipolar cell axon terminals.
689
Our results do not rule out additional roles for postsynaptic mechanisms in establishing transient
versus sustained RGC spike output. For example, delayed amacrine cell-mediated inhibitory
synaptic input to RGC dendrites could truncate spiking responses (Nirenberg and Meister,
1997). RGC cell-intrinsic mechanisms, such as RGC subtype-specific differences in voltage-
gated Na^+ channel properties and/or expression pattern could also contribute to distinct
response kinetics (Brombas et al., 2022; Chang et al., 2023; Werginz et al., 2020). While a
combination of pre- and postsynaptic factors may ultimately act in concert to generate transient
versus sustained RGC spike output (Warwick et al., 2018; Werginz et al., 2020), our results
establish a fundamental role for bipolar cell synapses in the segregation of visual input into
kinetically distinct channels in ON pathways of the retina.
700
701 ***Materials and Methods***
702
703 *Animals*
704
Experiments used 5- to 15-week-old wild-type, *Grm6-tdTomato* (Morgan et al., 2011), *Gus8.4-*
EGFP (Huang et al., 1999), *Gjd2-EGFP*, or Kcng4-Cre (Duan et al., 2014) transgenic mice.
Transgenic mouse strains were backcrossed into the same genetic background as wild-type
mice (C57BL/6J; Jackson Laboratory). Animal care and handling followed procedures approved
by the Institutional Animal Care and Use Committee of the University of Washington or the
University of Victoria Animal Care Committee.
711
712 *Tissue preparation*
713
Retinal tissue was prepared for physiological recordings as previously described (Kuo et al.,
2016; Srivastava et al., 2022). Briefly, mice were killed by cervical dislocation or decapitation
and retinas were dissected from both eyes in oxygenated (95% O_2, 5% CO_2), bicarbonate-
buffered Ames' (Sigma) or artificial cerebrospinal fluid (ACSF) solution and either flat-mounted
with the vitreal surface facing up onto a poly-D-lysine coated glass coverslip (Corning) (RGC
patch-clamp recordings and iGluSnFR imaging) or sliced in an orientation perpendicular to the
flat-mount plane using a vibratome (200 µm slice thickness; bipolar cell recordings) before being
transferred to an upright 2P fluorescence microscope. ACSF contained (in mM): 125 NaCl, 2.5
KCl, 1.2 $CaCl_2$, 1.5 $MgCl_2$, 1.25 NaH_2PO_4, 20 glucose, 0.5 L-glutamine, 0.1 sodium ascorbate,
and 26 $NaHCO_3$ (pH 7.4, 310mOsm/L). All experiments exclusively used tissue from ventral
retina.
725
726 *Electrophysiology*
727
During recordings, retinal tissue was continuously perfused (~6-8 mL/min) with oxygenated,
bicarbonate-buffered Ames' solution maintained at 30-34 ºC. In a subset of electrophysiological
recordings (Figs. 6C-F and S8), tissue was perfused with ACSF rather than Ames' solution.
Electrical signals were acquired at 10 kHz and low-pass filtered at 3 kHz using a Multiclamp
700B or Multiclamp 700A amplifier (Molecular Devices) and Symphony Data Acquisition
software (http://symphony-das.github.io). The somas of different ON bipolar cell subtypes were
targeted for gramicidin perforated-patch current-clamp recordings using 2P laser scanning

excitation (960nm) in retinal slices prepared from transgenic mouse lines in which bipolar cell
subtypes expressed fluorescent proteins. Tissue from *Grm6-tdTomato* mice, in which a sparse
subset of ON bipolar cell types (type 6, 7 and 8) express tdTomato was used to target
recordings to type 6 bipolar cells (Morgan et al., 2011; Schwartz et al., 2012). Type 7 bipolar
cells were targeted for recording in *Gus8.4-EGFP* mice, which express EGFP in a type 7 bipolar
cells and rod bipolar cells (Huang et al., 2003). Type 5i bipolar cells were targeted in retinas
from *Gjd2-EGFP* mice, which we found express EGFP selectively in these bipolar cells (Fig. S6)
as well as less robustly in an uncharacterized amacrine cell type (not shown). For bipolar cell
recordings, we minimized light exposure from the 2P laser by restricting scan regions to the
inner retina, away from photoreceptors, and by keeping laser exposure brief (<~30 sec/scan
region; <2 mW post-objective laser power) (Kuo et al., 2016). After identification of fluorescent
cell bodies, we acquired recordings using infrared differential interference contrast microscopy.
Bipolar recordings used an internal solution containing 123 mM K-aspartate, 10 mM KCl, 10 mM
HEPES, 1 mM $MgCl_2$, 1 mM $CaCl_2$, 2 mM EGTA, 4 mM Mg-ATP, 0.5 mM Tris-GTP, 100 µM
Alexa Fluor 594-hydrazide and 25-30 µg/mL gramicidin D (280 mOsm/L; pH 7.2). Pipettes (~7-9
Ω tip resistance) were front-filled with gramicidin-free internal solution to facilitate formation of
high resistance (>1 GΩ) seals between patch pipettes and the cell membrane. After seal
formation, series resistance (R_s) was monitored in voltage-clamp mode by applying a 10 mV
voltage step across the pipette tip. Experiments were initiated once R_s declined to <50 MΩ,
which typically took 20-30 min. During recordings, we occasionally confirmed the integrity of the
cell membrane under the recording electrode by monitoring Alexa Fluor 594 fluorescence using
2P excitation; if any dye was visible in the cell body, we abandoned the recording. No holding
current was applied during current-clamp recordings from bipolar cells. At the conclusion of
each bipolar cell recording, the membrane under the patch pipette was intentionally ruptured
and fluorescence images of Alexa Fluor 594-filled cells were acquired using 2P microscopy to
confirm bipolar cell subtype identities (Wässle et al., 2009).
761
RGC recordings were performed in a flat-mount preparation of the retina from wild-type mice or
Gjd2-EGFP transgenic mice following previously described methods (Kuo et al., 2016)
Excitatory synaptic currents were isolated in whole-cell voltage-clamp recordings by holding
RGCs near the reversal potential for Cl⁻- mediated conductances (-68.5 mV). Patch pipettes
(2.5-4 MΩ) were filled with an internal solution containing (in mM): 105 Cs-methanesulfonate, 10
Tetraethylammonium-Cl, 20 HEPES, 10 EGTA, 2 QX-314, 5 Mg-ATP, 0.5 Tris-GTP (~280
mOsm; pH 7.2). Series resistance (<15 MΩ) was compensated by 70% using the amplifier
circuitry. Inhibitory receptor antagonists (SR-95531, TPMPA, strychnine) were obtained from
Tocris or Sigma Aldrich and added to the bath solution from concentrated aqueous stock
solutions.
772
773 *iGluSnFR expression and two-photon imaging*
774
Intravitreal injection of a cre-dependent viral plasmid (pAAV.hSyn.Flex.iGluSnFR.WPRE.SV40)
was performed into Kcng4-cre mice to selectively express the glutamate sensor, iGluSnFR, in
alpha RGCs, including ON-T and ON-S RGCs (Krieger et al., 2017). All injection procedures,
tissue preparations, and subsequent two-photon imaging were performed as previously
described in Srivastava et al., 2022. Briefly, for iGluSnFR imaging, the laser was tuned to 920
nm and all recordings were acquired at a frame rate of 58.25 Hz (256 x 256 pixels). In KCNG4-
Cre mice, ON-S alpha RGCs were first identified by their spiking properties using extracellular
cell-attached recordings. To identify the dendrites of ON-S RGCs, cells were filled with
sulphorhodamine 101. ON-T RGC dendrites were then recorded at the same location but
deeper into the retina (Krieger et al., 2017).
785

Visual stimulation

Visual stimuli were presented from a photopic mean background illumination (~3,300 rhodopsin
isomerizations per rod per second (R*/rod/s)) to focus specifically on cone-mediated signaling
(Grimes et al., 2018). Visual stimuli for electrophysiological recordings were generated using
Stage Visual Stimulation Software (http://stage-vss.github.io). For all electrophysiological
recordings except those in Figs. 6C-F and S8, light stimuli consisted of a spatially uniform
circular spot (520 µm diameter) delivered from a LED (405 nm peak output; Hosfelt) that was
focused through the microscope substage condenser (Kuo et al., 2016). In the experiments
shown in Figs. 6C-F and S8, a Digital Light Processing (DLP) projector driven by a 405 nm LED
(Wintech Digital) was used to project spots of various sizes through the microscope objective
(Kuo et al., 2020). For iGluSnFR imaging (Fig. 5), background illuminance was kept around
1000 photons/µm^2/s. For experiments, static spots of different sizes were generated and
presented using StimGen software (https://github.com/benmurphybaum/StimGen). Stimuli were
presented via a DLP through the substage condenser focused onto the photoreceptor layer of
the retina. All visual stimuli were presented 4 s after the start of the recording for a duration of 2
s.

Data analysis and statistics

Spike triggered average (STA) waveforms (Fig. 1) and linear-nonlinear (LN) models of light-
driven RGC (Fig. 2) and bipolar cell (Fig.4) responses were derived from responses to randomly
fluctuating light stimuli (Gaussian distribution of light intensities with standard deviation = 50% of
mean intensity; 0-60 Hz bandwidth) as described previously (Chichilnisky, 2001; Kim and Rieke,
2001; Rieke, 2001). For LN model construction, the linear filter was calculated by computing the
cross-correlation of the light stimulus with the measured voltage (bipolar cell) or current (RGC)
response and dividing the result by the power spectrum of the stimulus. The static nonlinearity
was constructed from a point-by-point comparison of the predicted filter output with the
measured response and binning the result along the prediction axis. In almost all cases, the
linear filter had a biphasic shape with initial and secondary lobes that are opposite polarity. A
small number of bipolar cells had triphasic filters (Fig. S7). We quantified the kinetics of each
STA or linear filter by measuring the time at which the STA/filter waveform crossed zero ("zero-
cross time"; e.g. Fig. 1C, left) between the first and second lobes of the filter (e.g. between peak
and trough of STA waveforms) and by calculating a biphasic index = |A| / |B|, where |B| is the
absolute peak amplitude of the initial lobe of the filter and |A| is the amplitude of the peak
amplitude of the secondary lobe (e.g. Fig. 2C, right). We quantified the kinetics of responses to
light step stimuli by calculating the time between light step onset and the peak amplitude
response ("time to peak"; e.g. Fig. 2E, left) and dividing the mean steady-state response
(calculated from the last 100 ms light step increment) by the peak amplitude initial response at
step onset (and multiplying by 100) (e.g. Fig. 2E, right). We note that alternative measures of
step response or linear filter kinetics than those presented in Fig. 4 were also not statistically
different between bipolar cell subtypes (not shown). Statistical tests were performed using
MATLAB (Mathworks) or Igor Pro (WaveMetrics) software.

Biolistic transfection

DNA plasmids encoding mTFP1-myc (24 µg) and PSD95-mCherry (12 µg) under the control of
the cytomegalovirus (CMV) promoter were coated onto gold particles (1.6 µm diameter, 12.5
mg, Bio-Rad). Using a Helios gene gun (40 psi, Bio-Rad), these particles were then delivered to
retinal ganglion cells in whole-mount retinas. Subsequently, the transfected retinas were
incubated overnight in mACSF within a humid, oxygenated chamber maintained at 33°C.

Immunohistochemistry
838
Retinal sections (60 μm thick) were prepared using a vibratome (Leica) from retinal whole
mount retinas of *Gjd2-EGFP* mice. Both whole mount retinas and retinal sections were fixed in
4% PFA for 30 minutes. Retinal sections were embedded in low melting point agarose (Sigma-
Aldrich). Retinal tissues were first incubated with blocking solution (5% normal donkey serum,
0.5% Triton-X100 in PBS, Sigma-Aldrich) for 2 hours at 4°C, then incubated overnight at 4°C
with primary antibodies dissolved in blocking solution. Excess primary antibody was rinsed 3x10
minutes with PBS, then sections were incubated for 2 hours with secondary antibodies
dissolved in PBS. Excess secondary antibody was rinsed 3x10 minutes with PBS and sections
were mounted on standard microscopy glass slides using Vectashield mounting medium
(Vector). Primary antibody used in the study were: goat anti-ChAT diluted 1:1000 Chemicon
AB144P, mouse anti-CaBP5 diluted 1:1000 courtesy of Dr. Haeseleer, mouse anti-myc diluted
at 1:1000, chicken anti-GFP diluted at 1:1000. Secondary antibody were donkey anti-goat IgG
Dylight-568 diluted 1:1000, donkey anti-mouse Dylight-647 diluted 1:1000, donkey anti-mouse
Dylight-405 diluted at 1:1000 , donkey anti-chicken Alexa 488, the nuclear dye TOPRO3 (diluted
1:200, Thermo-Fisher).
854
855 *Near-infrared branding (NIRB) and serial block face scanning electron microscopy (SBFSEM)*
856
Two neighboring ON-T RGCs were physiologically identified in *Gjd2-EGFP* mice. After
identifying ON-T RGCs via their characteristic step-evoked extracellular spike response (e.g.
Figure 1B), they were filled with Alexa 488 or Alexa 555. Dendritic arbors and T5i-BC axonal
arbors were imaged using a confocal microscope. The retina was then fixed with 4%
glutaraldehyde for 30 minutes at room temperature (RT). The retina was washed with 0.1M Na+
Cacodylate buffer for three times and flat-mounted with 0.1M Na+ Cacodylate buffer. To
relocate the two RGCs under electron microscopy, we applied the near-infrared branding
(NIRB) technique previously published to burn fiducial markers in the retina (Bishop et al.). After
NIRBing, the retina tissue was unmounted and fixed in 4% glutaraldehyde overnight. The retina
was processed for SBFSEM subsequently according to the protocol described previously (Glumi
paper). Briefly, the tissue was initially treated with reduced osmium for 1 hour in the refrigerator.
Subsequently, after washing with double-distilled H2O (ddH2O), the tissue was placed in a
freshly prepared thiocarbohydrazide solution for 20 minutes at RT. Following another round of
ddH2O rinsing, the tissue was incubated in 2% osmium tetroxide for 30 minutes at RT. After
rinsing, the sample was incubated in 1% uranyl acetate at 4°C overnight. Post-wash, the tissue
was stained with Walton's lead aspartate for 30 minutes at 60°C. Following another wash, the
retinal piece underwent dehydration using a graded alcohol series (20%, 50%, 70%, 90%,
100%, 100%), each step lasting 5 minutes, followed by two changes of 100% propylene oxide at
RT for 10 minutes each. Finally, the tissue was embedded in Epon. The resulting block was
then trimmed and mounted in the SBEM microscope (GATAN/Zeiss SIGMA, 3View), and image
stacks were acquired at a voxel size of 5 x 5 x 50 nm.
878
879
880

References

Angueyra JM, Rieke F. 2013. Origin and effect of phototransduction noise in primate cone
photoreceptors. *Nat Neurosci* **16**:1692–1700. doi:10.1038/nn.3534
Asari H, Meister M. 2014. The projective field of retinal bipolar cells and its modulation by
visual context. *Neuron* **81**:641–652. doi:10.1016/j.neuron.2013.11.029
Asari H, Meister M. 2012. Divergence of visual channels in the inner retina. *Nat Neurosci*
15:1581–1589. doi:10.1038/nn.3241
Awatramani GB, Slaughter MM. 2000. Origin of transient and sustained responses in ganglion
cells of the retina. *J Neurosci* **20**:7087–7095. doi:10.1523/JNEUROSCI.20-18-07087.2000
Baden T, Berens P, Bethge M, Euler T. 2013. Spikes in mammalian bipolar cells support
temporal layering of the inner retina. *Curr Biol* **23**:48–52. doi:10.1016/j.cub.2012.11.006
Baden T, Berens P, Franke K, Román Rosón M, Bethge M, Euler T. 2016. The functional diversity
of retinal ganglion cells in the mouse. *Nature* **529**:345–350. doi:10.1038/nature16468
Bae JA, Mu S, Kim JS, Turner NL, Tartavull I, Kemnitz N, Jordan CS, Norton AD, Silversmith WM,
Prentki R, Sorek M, David C, Jones DL, Bland D, Sterling ALR, Park J, Briggman KL, Seung
HS, Eyewirers. 2018. Digital Museum of Retinal Ganglion Cells with Dense Anatomy and
Physiology. *Cell* **173**:1293-1306.e19. doi:10.1016/j.cell.2018.04.040
Bishop D, Nikić I, Brinkoetter M, Knecht S, Potz S, Kerschensteiner M, Misgeld T. 2011. Near-
infrared branding efficiently correlates light and electron microscopy. *Nat Methods*
8:568–570. doi:10.1038/nmeth.1622
Brombas A, Zhou X, Williams SR. 2022. Light-evoked dendritic spikes in sustained but not
transient rabbit retinal ganglion cells. *Neuron* **110**:2802-2814.e3.
doi:10.1016/j.neuron.2022.06.009
Burrone J, Lagnado L. 2000. Synaptic depression and the kinetics of exocytosis in retinal bipolar
cells. *J Neurosci* **20**:568–578. doi:10.1523/JNEUROSCI.20-02-00568.2000
Burrone J, Neves G, Gomis A, Cooke A, Lagnado L. 2002. Endogenous calcium buffers regulate
fast exocytosis in the synaptic terminal of retinal bipolar cells. *Neuron* **33**:101–112.
doi:10.1016/s0896-6273(01)00565-7
Cafaro J, Rieke F. 2013. Regulation of spatial selectivity by crossover inhibition. *J Neurosci*
33:6310–6320. doi:10.1523/JNEUROSCI.4964-12.2013
Cangiano L, Asteriti S, Cervetto L, Gargini C. 2012. The photovoltage of rods and cones in the
dark-adapted mouse retina. *J Physiol* **590**:3841–3855. doi:10.1113/jphysiol.2011.226878
Chang L, Ran Y, Yang M, Auferkorte O, Butz E, Hüser L, Haverkamp S, Euler T, Schubert T. 2023.
Spike desensitisation as a mechanism for high-contrast selectivity in retinal ganglion
cells. *Front Cell Neurosci* **17**:1337768. doi:10.3389/fncel.2023.1337768
Chichilnisky EJ. 2001. A simple white noise analysis of neuronal light responses. *Network*
12:199–213.
Cleland BG, Levick WR, Sanderson KJ. 1973. Properties of sustained and transient ganglion cells
in the cat retina. *J Physiol* **228**:649–680. doi:10.1113/jphysiol.1973.sp010105
Cohen E, Sterling P. 1990. Demonstration of cell types among cone bipolar neurons of cat
retina. *Philos Trans R Soc Lond B Biol Sci* **330**:305–321. doi:10.1098/rstb.1990.0201
Demb JB, Singer JH. 2012. Intrinsic properties and functional circuitry of the AII amacrine cell.
Vis Neurosci **29**:51–60. doi:10.1017/S0952523811000368

Diamond JS. 2017. Inhibitory Interneurons in the Retina: Types, Circuitry, and Function. *Annu
Rev Vis Sci* **3**:1–24. doi:10.1146/annurev-vision-102016-061345
Ding H, Smith RG, Poleg-Polsky A, Diamond JS, Briggman KL. 2016. Species-specific wiring for
direction selectivity in the mammalian retina. *Nature* **535**:105–110.
doi:10.1038/nature18609
Dowling JE, Boycott BB. 1966. Organization of the primate retina: electron microscopy. *Proc R
Soc Lond B Biol Sci* **166**:80–111. doi:10.1098/rspb.1966.0086
Duan X, Krishnaswamy A, De la Huerta I, Sanes JR. 2014. Type II cadherins guide assembly of a
direction-selective retinal circuit. *Cell* **158**:793–807. doi:10.1016/j.cell.2014.06.047
Euler T, Haverkamp S, Schubert T, Baden T. 2014. Retinal bipolar cells: elementary building
blocks of vision. *Nat Rev Neurosci* **15**:507–519. doi:10.1038/nrn3783
Euler T, Masland RH. 2000. Light-evoked responses of bipolar cells in a mammalian retina. *J
Neurophysiol* **83**:1817–1829. doi:10.1152/jn.2000.83.4.1817
Franke K, Berens P, Schubert T, Bethge M, Euler T, Baden T. 2017. Inhibition decorrelates visual
feature representations in the inner retina. *Nature* **542**:439–444.
doi:10.1038/nature21394
Ghosh KK, Bujan S, Haverkamp S, Feigenspan A, Wässle H. 2004. Types of bipolar cells in the
mouse retina. *J Comp Neurol* **469**:70–82. doi:10.1002/cne.10985
Goetz J, Jessen ZF, Jacobi A, Mani A, Cooler S, Greer D, Kadri S, Segal J, Shekhar K, Sanes JR,
Schwartz GW. 2022. Unified classification of mouse retinal ganglion cells using function,
morphology, and gene expression. *Cell Rep* **40**:111040.
doi:10.1016/j.celrep.2022.111040
Gollisch T, Meister M. 2010. Eye smarter than scientists believed: neural computations in
circuits of the retina. *Neuron* **65**:150–164. doi:10.1016/j.neuron.2009.12.009
Greene MJ, Kim JS, Seung HS, EyeWirers. 2016. Analogous Convergence of Sustained and
Transient Inputs in Parallel On and Off Pathways for Retinal Motion Computation. *Cell
Rep* **14**:1892–1900. doi:10.1016/j.celrep.2016.02.001
Grimes WN, Baudin J, Azevedo AW, Rieke F. 2018. Range, routing and kinetics of rod signaling in
primate retina. *Elife* **7**:e38281. doi:10.7554/eLife.38281
Grimes WN, Schwartz GW, Rieke F. 2014. The synaptic and circuit mechanisms underlying a
change in spatial encoding in the retina. *Neuron* **82**:460–473.
doi:10.1016/j.neuron.2014.02.037
Hellmer CB, Zhou Y, Fyk-Kolodziej B, Hu Z, Ichinose T. 2016. Morphological and physiological
analysis of type-5 and other bipolar cells in the Mouse Retina. *Neuroscience* **315**:246–
258. doi:10.1016/j.neuroscience.2015.12.016
Helmstaedter M, Briggman KL, Turaga SC, Jain V, Seung HS, Denk W. 2013. Connectomic
reconstruction of the inner plexiform layer in the mouse retina. *Nature* **500**:168–174.
doi:10.1038/nature12346
Huang L, Max M, Margolskee RF, Su H, Masland RH, Euler T. 2003. G protein subunit G gamma
13 is coexpressed with G alpha o, G beta 3, and G beta 4 in retinal ON bipolar cells. *J
Comp Neurol* **455**:1–10. doi:10.1002/cne.10396
Huang L, Shanker YG, Dubauskaite J, Zheng JZ, Yan W, Rosenzweig S, Spielman AI, Max M,
Margolskee RF. 1999. Ggamma13 colocalizes with gustducin in taste receptor cells and

mediates IP3 responses to bitter denatonium. *Nat Neurosci* **2**:1055–1062.
doi:10.1038/15981

Ichinose T, Fyk-Kolodziej B, Cohn J. 2014. Roles of ON cone bipolar cell subtypes in temporal
coding in the mouse retina. *J Neurosci* **34**:8761–8771. doi:10.1523/JNEUROSCI.3965-
13.2014

Ichinose T, Hellmer CB. 2016. Differential signalling and glutamate receptor compositions in the
OFF bipolar cell types in the mouse retina. *J Physiol* **594**:883–894. doi:10.1113/JP271458

Ikeda H, Wright MJ. 1972. Receptive field organization of "sustained" and "transient" retinal
ganglion cells which subserve different function roles. *J Physiol* **227**:769–800.
doi:10.1113/jphysiol.1972.sp010058

Jarsky T, Cembrowski M, Logan SM, Kath WL, Riecke H, Demb JB, Singer JH. 2011. A Synaptic
Mechanism for Retinal Adaptation to Luminance and Contrast. *J Neurosci* **31**:11003–
11015. doi:10.1523/JNEUROSCI.2631-11.2011

Ke J-B, Wang YV, Borghuis BG, Cembrowski MS, Riecke H, Kath WL, Demb JB, Singer JH. 2014.
Adaptation to background light enables contrast coding at rod bipolar cell synapses.
Neuron **81**:388–401. doi:10.1016/j.neuron.2013.10.054

Kerschensteiner D. 2022. Feature Detection by Retinal Ganglion Cells. *Annu Rev Vis Sci* **8**:135–
169. doi:10.1146/annurev-vision-100419-112009

Kim KJ, Rieke F. 2001. Temporal contrast adaptation in the input and output signals of
salamander retinal ganglion cells. *J Neurosci* **21**:287–299. doi:10.1523/JNEUROSCI.21-
01-00287.2001

Krieger B, Qiao M, Rousso DL, Sanes JR, Meister M. 2017. Four alpha ganglion cell types in
mouse retina: Function, structure, and molecular signatures. *PLoS One* **12**:e0180091.
doi:10.1371/journal.pone.0180091

Kuo SP, Chiang P-P, Nippert AR, Newman EA. 2020. Spatial Organization and Dynamics of the
Extracellular Space in the Mouse Retina. *J Neurosci* **40**:7785–7794.
doi:10.1523/JNEUROSCI.1717-20.2020

Kuo SP, Schwartz GW, Rieke F. 2016. Nonlinear Spatiotemporal Integration by Electrical and
Chemical Synapses in the Retina. *Neuron* **90**:320–332.
doi:10.1016/j.neuron.2016.03.012

Lagnado L, Schmitz F. 2015. Ribbon Synapses and Visual Processing in the Retina. *Annu Rev Vis*
Sci **1**:235–262. doi:10.1146/annurev-vision-082114-035709

Marc RE, Anderson JR, Jones BW, Sigulinsky CL, Lauritzen JS. 2014. The AII amacrine cell
connectome: a dense network hub. *Front Neural Circuits* **8**:104.
doi:10.3389/fncir.2014.00104

Margolis DJ, Gartland AJ, Euler T, Detwiler PB. 2010. Dendritic Calcium Signaling in ON and OFF
Mouse Retinal Ganglion Cells. *J Neurosci* **30**:7127–7138. doi:10.1523/JNEUROSCI.5694-
09.2010

Masland RH. 2012. The Neuronal Organization of the Retina. *Neuron* **76**:266–280.
doi:10.1016/j.neuron.2012.10.002

McGuire BA, Stevens JK, Sterling P. 1984. Microcircuitry of bipolar cells in cat retina. *J Neurosci*
4:2920–2938. doi:10.1523/JNEUROSCI.04-12-02920.1984

Mennerick S, Matthews G. 1996. Ultrafast exocytosis elicited by calcium current in synaptic
terminals of retinal bipolar neurons. *Neuron* **17**:1241–1249. doi:10.1016/s0896-
6273(00)80254-8
Morgan JL, Soto F, Wong ROL, Kerschensteiner D. 2011. Development of cell type-specific
connectivity patterns of converging excitatory axons in the retina. *Neuron* **71**:1014–
1021. doi:10.1016/j.neuron.2011.08.025
Neves G, Lagnado L. 1999. The kinetics of exocytosis and endocytosis in the synaptic terminal of
goldfish retinal bipolar cells. *J Physiol* **515 (Pt 1)**:181–202. doi:10.1111/j.1469-
7793.1999.181ad.x
Nirenberg S, Meister M. 1997. The light response of retinal ganglion cells is truncated by a
displaced amacrine circuit. *Neuron* **18**:637–650. doi:10.1016/s0896-6273(00)80304-9
Oesch NW, Diamond JS. 2011. Ribbon synapses compute temporal contrast and encode
luminance in retinal rod bipolar cells. *Nat Neurosci* **14**:1555–1561. doi:10.1038/nn.2945
Pangrsic T, Singer JH, Koschak A. 2018. Voltage-Gated Calcium Channels: Key Players in Sensory
Coding in the Retina and the Inner Ear. *Physiol Rev* **98**:2063–2096.
doi:10.1152/physrev.00030.2017
Puthussery T, Venkataramani S, Gayet-Primo J, Smith RG, Taylor WR. 2013. NaV1.1 channels in
axon initial segments of bipolar cells augment input to magnocellular visual pathways in
the primate retina. *J Neurosci* **33**:16045–16059. doi:10.1523/JNEUROSCI.1249-13.2013
Ran Y, Huang Z, Baden T, Schubert T, Baayen H, Berens P, Franke K, Euler T. 2020. Type-specific
dendritic integration in mouse retinal ganglion cells. *Nat Commun* **11**:2101.
doi:10.1038/s41467-020-15867-9
Rieke F. 2001. Temporal contrast adaptation in salamander bipolar cells. *J Neurosci* **21**:9445–
9454.
Rodieck RW. 1991. The density recovery profile: a method for the analysis of points in the plane
applicable to retinal studies. *Vis Neurosci* **6**:95–111. doi:10.1017/s095252380001049x
Roska B, Werblin F. 2001. Vertical interactions across ten parallel, stacked representations in
the mammalian retina. *Nature* **410**:583–587. doi:10.1038/35069068
Sabbah S, Papendorp C, Koplas E, Beltoja M, Etebari C, Gunesch AN, Carrete L, Kim MT, Manoff
G, Bhatia-Lin A, Zhao T, Schreck D, Dowling H, Briggman KL, Berson DM. 2018. Synaptic
circuits for irradiance coding by intrinsically photosensitive retinal ganglion cells
(preprint). Neuroscience. doi:10.1101/442954
Sagdullaev BT, McCall MA, Lukasiewicz PD. 2006. Presynaptic inhibition modulates spillover,
creating distinct dynamic response ranges of sensory output. *Neuron* **50**:923–935.
doi:10.1016/j.neuron.2006.05.015
Saszik S, DeVries SH. 2012. A mammalian retinal bipolar cell uses both graded changes in
membrane voltage and all-or-nothing Na+ spikes to encode light. *J Neurosci* **32**:297–307.
doi:10.1523/JNEUROSCI.2739-08.2012
Schwartz GW, Okawa H, Dunn FA, Morgan JL, Kerschensteiner D, Wong RO, Rieke F. 2012. The
spatial structure of a nonlinear receptive field. *Nat Neurosci* **15**:1572–1580.
doi:10.1038/nn.3225
Shekhar K, Lapan SW, Whitney IE, Tran NM, Macosko EZ, Kowalczyk M, Adiconis X, Levin JZ,
Nemesh J, Goldman M, McCarroll SA, Cepko CL, Regev A, Sanes JR. 2016.

Comprehensive Classification of Retinal Bipolar Neurons by Single-Cell Transcriptomics.
Cell **166**:1308-1323.e30. doi:10.1016/j.cell.2016.07.054

Sigulinsky CL, Anderson JR, Kerzner E, Rapp CN, Pfeiffer RL, Rodman TM, Emrich DP, Rapp KD,
Nelson NT, Lauritzen JS, Meyer M, Marc RE, Jones BW. 2020. Network Architecture of
Gap Junctional Coupling among Parallel Processing Channels in the Mammalian Retina. *J*
Neurosci **40**:4483–4511. doi:10.1523/JNEUROSCI.1810-19.2020

Singer JH, Diamond JS. 2006. Vesicle depletion and synaptic depression at a mammalian ribbon
synapse. *J Neurophysiol* **95**:3191–3198. doi:10.1152/jn.01309.2005

Singer JH, Diamond JS. 2003. Sustained Ca2+ entry elicits transient postsynaptic currents at a
retinal ribbon synapse. *J Neurosci* **23**:10923–10933. doi:10.1523/JNEUROSCI.23-34-
10923.2003

Slaughter MM, Miller RF. 1981. 2-amino-4-phosphonobutyric acid: a new pharmacological tool
for retina research. *Science* **211**:182–185. doi:10.1126/science.6255566

Srivastava P, de Rosenroll G, Matsumoto A, Michaels T, Turple Z, Jain V, Sethuramanujam S,
Murphy-Baum BL, Yonehara K, Awatramani GB. 2022. Spatiotemporal properties of
glutamate input support direction selectivity in the dendrites of retinal starburst
amacrine cells. *Elife* **11**:e81533. doi:10.7554/eLife.81533

Szikra T, Trenholm S, Drinnenberg A, Jüttner J, Raics Z, Farrow K, Biel M, Awatramani G, Clark
DA, Sahel J-A, da Silveira RA, Roska B. 2014. Rods in daylight act as relay cells for cone-
driven horizontal cell-mediated surround inhibition. *Nat Neurosci* **17**:1728–1735.
doi:10.1038/nn.3852

Taylor WR, Vaney DI. 2002. Diverse synaptic mechanisms generate direction selectivity in the
rabbit retina. *J Neurosci* **22**:7712–7720.

Tsukamoto Y, Omi N. 2017. Classification of Mouse Retinal Bipolar Cells: Type-Specific
Connectivity with Special Reference to Rod-Driven AII Amacrine Pathways. *Front*
Neuroanat **11**:92. doi:10.3389/fnana.2017.00092

Vaney DI, Sivyer B, Taylor WR. 2012. Direction selectivity in the retina: symmetry and
asymmetry in structure and function. *Nat Rev Neurosci* **13**:194–208.
doi:10.1038/nrn3165

Veruki ML, Hartveit E. 2002a. Electrical synapses mediate signal transmission in the rod
pathway of the mammalian retina. *J Neurosci* **22**:10558–10566.
doi:10.1523/JNEUROSCI.22-24-10558.2002

Veruki ML, Hartveit E. 2002b. AII (Rod) amacrine cells form a network of electrically coupled
interneurons in the mammalian retina. *Neuron* **33**:935–946. doi:10.1016/s0896-
6273(02)00609-8

von Gersdorff H, Matthews G. 1997. Depletion and replenishment of vesicle pools at a ribbon-
type synaptic terminal. *J Neurosci* **17**:1919–1927. doi:10.1523/JNEUROSCI.17-06-
01919.1997

Warwick RA, Kaushansky N, Sarid N, Golan A, Rivlin-Etzion M. 2018. Inhomogeneous Encoding
of the Visual Field in the Mouse Retina. *Curr Biol* **28**:655-665.e3.
doi:10.1016/j.cub.2018.01.016

Wässle H, Puller C, Müller F, Haverkamp S. 2009. Cone contacts, mosaics, and territories of
bipolar cells in the mouse retina. *J Neurosci* **29**:106–117. doi:10.1523/JNEUROSCI.4442-
08.2009

Werblin FS. 1970. Response of retinal cells to moving spots: intracellular recording in Necturus
maculosus. *J Neurophysiol* **33**:342–350. doi:10.1152/jn.1970.33.3.342
Werginz P, Raghuram V, Fried SI. 2020. Tailoring of the axon initial segment shapes the
conversion of synaptic inputs into spiking output in OFF-α T retinal ganglion cells. *Sci*
Adv **6**:eabb6642. doi:10.1126/sciadv.abb6642
Wienbar S, Schwartz GW. 2022. Differences in spike generation instead of synaptic inputs
determine the feature selectivity of two retinal cell types. *Neuron* **110**:2110-2123.e4.
doi:10.1016/j.neuron.2022.04.012
Wu SM, Gao F, Maple BR. 2000. Functional architecture of synapses in the inner retina:
segregation of visual signals by stratification of bipolar cell axon terminals. *J Neurosci*
20:4462–4470. doi:10.1523/JNEUROSCI.20-12-04462.2000
Yu W-Q, Swanstrom R, Sigulinsky CL, Ahlquist RM, Knecht S, Jones BW, Berson DM, Wong RO.
2023. Distinctive synaptic structural motifs link excitatory retinal interneurons to diverse
postsynaptic partner types. *Cell Rep* **42**:112006. doi:10.1016/j.celrep.2023.112006
Zhang Y, Kim I-J, Sanes JR, Meister M. 2012. The most numerous ganglion cell type of the
mouse retina is a selective feature detector. *Proc Natl Acad Sci U S A* **109**:E2391-2398.
doi:10.1073/pnas.1211547109
1114
1115

1116 *Supplemental Information*
1117
1118

1119
1120
Figure S1. Noise-evoked responses are biased towards large amplitude events in ON-T
compared to ON-S RGCs. (A) Spike event peak spike rate cumulative distributions for ON-T
(magenta; n= 6 cells; 131 spike events) and ON-S RGC (cyan; n= 7 cells; 300 spike events).
Firing rates from average responses to a repeated noise stimulus (see Fig. 1B-C) were
normalized to the maximal firing rate in each cell before combining events across cells. (B)
EPSC amplitude cumulative distributions for ON-T (n= 13 cells; 275 EPSCs) and ON-S (n= 10
cells; 365 EPSCs) RGCs. Same stimulus and normalization procedure as in (A).
1128

1129
1130

1131
1132
Figure S2. *Linear-nonlinear models of excitatory synaptic input to ON-T and ON-S RGCs. (A)*
Linear filters computed from responses to non-repeated noise stimulus presentation. Lines
show filters from individual ON-T (cyan; n= 13) and ON-S (magenta; n= 15) RGCs. Filters were
normalized to the peak amplitude of the primary inward component to facilitate visual
comparison of temporal responses. Same cells as shown in average filters in Fig. 2B. (B)
Nonlinearities derived by plotting the predicted response (convolution of linear filter and
stimulus) against the measured excitatory currents for each RGC shown in (A). Linear
predictions were calculated using non-normalized filters.
1141

1142
1143
Figure S3. Characterization of Gjd2-EGFP+ bipolar cells. (A) Vertical section of the Gjd2-EGFP
mouse line showing EGFP positive bipolar cells axons (green) stratifying in the ON sublamina of
the IPL, co-labeled with cholinergic amacrine cell (magenta, ChAT antibody). Transmitted light
image of the retina in blue. (B) Vertical section of the Gjd2-EGFP mouse line showing EGFP
positive bipolar cells (green) colocalized with the T5 marker CaBP5 (magenta). (C) En-face
image of a whole mount retina from the GJD2 mouse line showing EGFP positive bipolar cells
bodies (green) co-labeled with the T5 bipolar cell marker CaBP5 (magenta). (D) Same field as
in (C) with cell bodies of EGFP positive bipolar cells outlined (green) by application of a digital
edge outlining filter in ImageJ. (E) Position of cell body centroids for EGFP positive bipolar cells
in a larger field of view (20X objective) outlined by green circles. (F) Density recovery profile
analysis showing the density profile as a function of distance from the reference neuron for
EGFP positive bipolar cells (green histograms) and for an equal amount of neurons randomly
positioned in the same space (grey histograms) showing the presence of a characteristic
exclusion distance for EGFP positive bipolar cells, hallmark of a retinal mosaic.

1158

1159

Figure S4. *ON-T and ON-S RGC dendrites ramify in distinct IPL sublaminae. (A) Side*

projections from 2P z-stack images of dye-filled RGCs (ON-T: top, cyan; ON-S: bottom,

magenta) and EGFP signal (yellow) in retinas from Gjd2-EGFP mice. RGCs are same cells as

shown in en face view in Fig. 1A (B) Quantification of fluorescence signals from dye filled RGCs

in Gjd2-EGFP mouse retina as in (A). Colored lines and shaded areas show mean ± SEM

fluorescence intensity with respect to distance from peak EGFP (yellow) fluorescence intensity

(n= 6 ON-T, cyan and n= 4 ON-S, magenta, RGCs).

1167

1168

1169
1170
Figure S5. *Gjd2-EGFP+ type 5 bipolar cells provide ~40% of total synapses onto ON-T RGCs.*
(A) An example ON-T RGC biolistically labeled by Cerulean *Fluorescent Protein (blue) and*
PSD95-mCherry (green) in retina from a Gjd2-EGFP (red) transgenic mouse. Top: en face view,
Bottom: side views. (B) Top: Dot map of PSD95-mCherry apposed (magenta) or not apposed
(green) to GFP+ BC axon terminals. Bottom: Magnified view of the raw images and the dot map
for a segment of dendritic arbor in the while box. (C) Quantification of total excitatory synapse
density across the dendritic arbor of ON-T RGCs. (D) Quantification of excitatory synapses in
ON-T dendrites that are adjacent to GFP+ axon terminals.
1179

1180

1181
1182
Figure S6. *Classification of three subtypes of type 5 bipolar cells. (A) Wholemount views of the*
axon terminals of type 5i, type 5t and type 5o bipolar cells in the SBFSEM volume in Fig. 3. The
tiling of the space validates the categorization of the three subtypes of type 5 bipolar cells. (B)
Identification of Gjd2-EGFP+ bipolar cells as type 5i bipolar cells, supported by correlated light
(left) and electron (right) microscopy images. The reconstructed skeleton of the type 5i bipolar
cell in (B) is delineated by dashed lines in (A).
1189

1190
1191
Figure S7. *Linear-nonlinear models of bipolar cell voltage responses. (A) Individual linear filters*
for all type 5i (cyan; n= 11), type 6 (magenta; n= 7), and type 7 (green; n=6) bipolar cells
computed from responses to non-repeated noise stimulus presentation. Filters were normalized
to the peak amplitude of the primary inward component to facilitate visual comparison of
temporal responses. Same cells as shown in average filters in Fig. 4B. (B) Nonlinearities
derived by plotting the predicted response (convolution of linear filter and stimulus) against the
measured voltage response for each bipolar cell shown in (A). Linear predictions were
calculated using non-normalized filters.
1200

1201
1202
Figure S8. *Presynaptic inhibition does not affect response kinetics but does affect spatial*
integration for excitatory inputs to ON-T RGCs (A) Step-evoked (100% contrast, 300 µm spot
diameter, 500 ms duration) excitatory synaptic currents in example ON-T (cyan) and ON-S
RGCs measured in ACSF. Traces show mean responses to ten trials and are baseline
subtracted to facilitate comparison. (B) Summary of light step-evoked steady-state response
amplitude for different stimulus sizes measured in ACSF. Note that excitatory inputs in ON-T are
more transient than those in ON-S RGCs across all stimulus sizes, but responses in both RGC
types are more sustained than when measurements were performed using Ames' solution
(compare to Fig. 2D-E). (C) Summary of total response, measured as the integrated excitatory
current response during the light step, versus stimulus spot size. Responses were normalized to
the maximal response in each cell prior to averaging. Excitation in ON-T RGCs is tuned to
smaller spot sizes than in ON-S RGCs and this tuning relies in part on GABAergic inhibition.
Symbols with error bars in (B) and (C) show mean ± SEM (ON-T control n = 12; ON-S control n
= 6; ON-T GABA-R block n = 5; ON-T Gly-R block n = 5).
1217