1
2
3
4   ***Cone bipolar cell synapses generate transient versus sustained signals in parallel ON***
5   ***pathways of the mouse retina***
6
7
8
9

10   Sidney P. Kuo[1,2], Wan-Qing Yu[1], Prerna Srivastava[3], Haruhisa Okawa[1], Luca Della Santina[1],
11   David M. Berson[4], Gautam B. Awatramani[3], Rachel O.L. Wong[1], Fred Rieke[1]
12
13   [1]Department of Neurobiology and Biophysics, University of Washington, Seattle WA
14   [2]Department of Neuroscience, University of Minnesota, Minneapolis MN
15   [3]Department of Biology, University of Victoria, Victoria BC
16   [4]Department of Neuroscience, Brown University, Providence RI
17
18   Correspondence: Fred Rieke, rieke@uw.edu

Abstract

Parallel processing is a fundamental organizing principle in the nervous system and understanding how parallel neural circuits generate distinct outputs from common inputs is a key goal of neuroscience. In the mammalian retina, divergence of cone signals into multiple feedforward bipolar cell pathways forms the initial basis for parallel retinal circuits dedicated to specific visual functions. Here, we used patch-clamp electrophysiology, electron microscopy and two photon imaging of a fluorescent glutamate sensor to examine how kinetically-distinct responses arise in transient versus sustained ON alpha RGCs (ON-T and ON-S RGCs) of the mouse retina. We directly compared the visual response properties of these RGCs with their presynaptic bipolar cell partners, which we identified using 3D electron microscopy reconstruction. Different ON bipolar cell subtypes (type 5i, type 6 and type 7) had indistinguishable light-driven responses whereas extracellular glutamate signals around RGC dendrites and postsynaptic excitatory currents measured in ON-T and ON-S RGCs in response to the identical stimuli used to probe bipolar cells were kinetically distinct. Anatomical examination of the bipolar cell axon terminals presynaptic to ON-T and ON-S RGCs suggests that bipolar subtype-specific differences in the size of synaptic ribbon-associated vesicle pools may contribute to transient versus sustained kinetics. Our findings indicate that feedforward bipolar cell synapses are a primary point of divergence in kinetically distinct visual pathways.

Introduction

Divergence into parallel pathways allows neural circuits to generate diverse outputs from a common set of inputs. This functional diversity can be quite striking; in vision, for example, light inputs initially encoded by the same collection of cone photoreceptors mediate sensitivity to several kinds of motion, to color and to spatial structure, all in the same region of space. These diverse functional sensitivities arise through parallel processing of light inputs, first in the retina and subsequently in downstream visual areas. Here we focus on the origin of transient and sustained responses – a key differentiating characteristic of retinal output cells across species (Awatramani and Slaughter, 2000; Baden et al., 2016; Bae et al., 2018; Cleland et al., 1973; Ikeda and Wright, 1972; Roska and Werblin, 2001; Werblin, 1970).

The retina provides a unique opportunity to study the mechanisms that differentiate signals in parallel pathways. We have a near-complete list of retinal cell types and substantial information about which of these cell types participate in specific parallel pathways (Helmstaedter et al., 2013; Masland, 2012). Under daylight conditions, retinal processing starts with cone photoreceptors transducing light inputs and providing divergent output to ~13 types of bipolar cell (Euler et al., 2014; Greene et al., 2016; Helmstaedter et al., 2013; Shekhar et al., 2016; Tsukamoto and Omi, 2017). The resulting bipolar cell signals are transmitted to ~20-40 retinal ganglion cell (RGC) types (Baden et al., 2016; Bae et al., 2018; Goetz et al., 2022). RGC responses are also strongly shaped by the activity of a large and diverse collection of amacrine cells that receive input from bipolar cells and provide presynaptic inhibition to bipolar cells and postsynaptic inhibition to RGCs (Diamond, 2017; Jadzinsky and Baccus, 2013). Finally, RGC responses are shaped by feedforward synaptic properties, particularly synaptic depression and facilitation (DeVries, 2000; Grabner et al., 2016; Jarsky et al., 2011; Lagnado and Schmitz, 2015; Nikolaev et al., 2013; Oesch and Diamond, 2011).

These mechanisms collectively produce highly distinct responses across RGC types - with cells that sample inputs on a variety of spatial and temporal scales, cells with distinct chromatic properties, cells that sense the direction of moving objects, and cells that separate local from

global motion (reviewed by Gollisch and Meister, 2010; Kerschensteiner, 2022). Several
mechanisms contribute to these functional differences, starting with the division of signals into
ON and OFF pathways at the cone-to-bipolar synapse (Euler et al., 2014; Slaughter and Miller,
1981). Temporal properties of responses of different bipolar cells can also differ for some stimuli
(Awatramani and Slaughter, 2000; Baden et al., 2013; Euler et al., 2014; Euler and Masland,
2000; Ichinose et al., 2014; Ichinose and Hellmer, 2016; Wu et al., 2000). In the inner retina,
integration of excitatory and inhibitory synaptic inputs produces directionally-selective responses
in specific RGC types (Taylor and Vaney, 2002; Vaney et al., 2012) and differences in spatial
weighting functions in other RGCs (Cafaro and Rieke, 2013; Roska and Werblin, 2001; Zhang
et al., 2012). Properties of dendritic integration and RGC spike generation also differentiate
responses of some RGCs (Margolis et al., 2010; Ran et al., 2020; Wienbar and Schwartz,
2022).
81
A fundamental difference among RGCs is whether their responses are sustained or transient
(Awatramani and Slaughter, 2000; Baden et al., 2016; Bae et al., 2018; Cleland et al., 1973;
Ikeda and Wright, 1972; Roska and Werblin, 2001). Cone photoreceptors respond to changes in
light intensity with graded potentials that have both a transient and sustained component
(Angueyra and Rieke, 2013; Cangiano et al., 2012; Szikra et al., 2014). Both kinetic
components of the cone response are maintained in the firing rates of sustained cells, which
track the prevailing contrast. However, only the initial component of the cone response is
preserved in the spiking response of transient cells, which therefore respond primarily to
changes in contrast. Where and how these differences originate is not clear. Here, we show that
the distinct kinetics of responses of ON-sustained alpha and ON-transient alpha RGCs (1)
persist for stimuli that elicit near-identical responses among the relevant bipolar cells, (2) persist
without amacrine cell-mediated inhibition and, (3) are correlated with anatomical differences in
output synapses of bipolar cell subtypes that are presynaptic to these RGCs. These findings
highlight the importance of feedforward bipolar cell output synapses in producing kinetically
distinct retinal outputs.
97
98 ***Results***
99
We focused on two ON RGC cell types, ON-transient alpha (ON-T) and ON-sustained alpha
(ON-S) RGCs, that could be readily identified in a flat-mount preparation of the isolated mouse
retina (Fig. 1A) by their characteristic spiking response to a step increment in light intensity (Fig.
1B) (Bae et al., 2018; Farrow et al., 2013; Goetz et al., 2022; Krieger et al., 2017; Kuo et al.,
2016). Based on their light response, dendritic field size and stratification pattern in the inner
plexiform layer (IPL) (see below; Fig. S4), the ON-T RGCs we studied here likely correspond to
the PV2 RGC type of Farrow et al. (2013), 6sw RGC type of Bae et al. (2018) and the medium-
field ON-transient RGC of Goetz et al. (2022). ON-S RGCs correspond to the PV1 RGC type of
Farrow et al. (2013) and 8w RGC type of Bae et al. (2018).
109
110

Figure 1. *Distinct temporal spiking response properties of ON-transient (ON-T) versus ON-sustained (ON-S) RGCs. (A) Maximum intensity projection z-stack fluorescence images of ON-T (left, cyan) and ON-S (right, magenta) RGCs dialyzed with Alexa Fluor 594 during whole-cell patch-clamp recording. Scale bar 50 μm. (B) Characteristic spiking responses to a light increment for example ON-T and ON-S RGCs. (C) Example spiking responses to repeated trials of spatially uniform (520 μm spot diameter) Gaussian noise stimulus (top black trace) presented with a mean luminance of ~2000 P*/S-cone/s (~3300 R*/rod/s; ~1650 P*/M-cone/s). ON-T (cyan) and ON-S (magenta) RGCs were recorded sequentially in the same piece of retina. Average spike responses (bottom) calculated from 20 repetitions of the noise stimulus. (D) ON-T and ON-S STAs derived from responses to non-repeated noise stimuli. Colored lines and shaded areas show mean ± SEM STA waveforms from ON-T (n= 15) and ON-S (n= 13) RGCs. The STA time axis is reversed to facilitate comparison with linear filters in Figures 2 and 4. The mean line obscures the SEM shading for much of the filter in both cell types. (E) Quantification of STA kinetics in different RGCs. Open circles show measurements from individual cells and filled circles with error bars show mean ± SEM values. Significance values from Wilcoxon rank sum tests.*

We characterized the temporal spiking response properties of ON-T and ON-S RGCs by using
an extracellular cell-attached patch electrode to measure action potential firing from either cell
type while presenting a spatially uniform, randomly fluctuating light stimulus (Gaussian noise)
that allowed us to efficiently probe a variety of stimulus contrasts and temporal frequencies.
These and all subsequent physiological recordings were conducted using a photopic mean
background light level (~3300 R*/rod/s; ~2000 P*/S-cone/s; ~1650 P*/M-cone/s) to suppress rod
photoreceptor responses and focus on cone-mediated signaling (Grimes et al., 2018; Ke et al.,
2014). Figure 1C shows the spiking response of example ON-T and ON-S RGCs to multiple
repetitions of the same random stimulus contrast sequence; these RGC types clearly responded
differently to temporally modulated light input. In particular, noise-evoked ON-T RGC spike
responses were comprised primarily of large amplitude, less frequent spike bursts whereas ON-
S RGC responses were less discrete and consisted of events with a wider range of amplitudes
(Fig. 1C and Fig S1A).

We summarized the response of each ON-T and ON-S RGC by computing the average stimulus
preceding each recorded spike (the spike-triggered average or STA stimulus) when presenting
a non-repeated Gaussian noise stimulus (Fig. 1D) (Chichilnisky, 2001). W   e quantified
response kinetics from these STA waveforms by measuring the time of zero-crossing (between
trough and peak) as well as the ratio of amplitudes of the trough and peak (biphasic index). ON-
T RGC STAs were significantly more biphasic (biphasic index = 0.63 ± 0.07; mean ± S.D.; n =
15) than ON-S RGC STAs (biphasic index = 0.43 ± 0.11; mean ± S.D.; n = 13) (Fig. 1E, right),
consistent with their more transient responses to step stimuli (Fig. 1B). ON-T RGC responses
were also slightly delayed (zero-cross time = - 0.080 ± 0.0063 sec; mean ± S.D.; n = 15) relative
to those of ON-S RGCs (zero-cross time = -0.072 ± 0.0052 sec; mean ± S.D.; n = 13) (Fig. 1C,
bottom and 1E, left). The STA provides a convenient summary of RGC temporal response
properties, but it does not capture all differences between the RGC spike responses – e.g. the
discreteness of the ON-T responses in Figure 1C. Nonetheless, the combination of direct
observation of the spike responses and STA kinetics illustrate how ON-S and ON-T responses
differ.

Bipolar cell to RGC synapses are an important site of signal diversification in ON pathways

A variety of mechanisms could contribute to the distinct temporal response characteristics of ON-
T versus ON-S RGCs, including differences in excitatory and/or inhibitory input conveyed by
presynaptic circuits. Parallel visual channels are initially established by the divergence of cone
signals to different cone bipolar cell subtypes, which have axons that terminate at distinct depths
within the IPL and can correspondingly contact distinct RGC types (Euler et al., 2014; Wässle et
al., 2009). Prior studies have demonstrated IPL layer-specific light-evoked glutamate release
kinetics from different bipolar cell synaptic terminals (Borghuis et al., 2013a; Franke et al., 2017).
We therefore sought to understand to what extent kinetic differences in bipolar cell-mediated
excitatory synaptic input could account for the transient versus sustained responses of these
RGCs.

To do this, we used somatic whole-cell voltage-clamp recordings to measure excitatory synaptic
currents from both RGC types while stimulating the retina with the same spatially uniform
Gaussian noise stimulus that we used to characterize RGC spike output (Fig. 2A). We used these
measurements to construct linear-nonlinear (LN) models comprised of a linear temporal filter (Fig.
2B and Fig. S2A) and a time-invariant (static) nonlinearity (Fig. S2B) that together provide a simple
estimate of the transformation between visual input and excitatory synaptic input to each RGC
type (Kim and Rieke, 2001). LN models for each RGC type accurately predicted measured

responses to noise stimuli that were not used to construct the models, although LN models of
ON-S RGC responses performed better than those for ON-T RGCs (correlation coefficient
between average response to repeated noise stimulus and model prediction: 0.61 ± 0.06 (ON-S,
n = 9) and 0.50 ± 0.13 (ON-T, n =11), mean ± S.D.; p = 0.028 Wilcoxon rank-sum test). ON-T
stimulus-evoked excitatory currents were biased towards large amplitude events compared to
those of ON-S RGCs (Fig. 2A, Fig. S1B), like the distinct amplitude distributions of spike events
in these cell types noted above (Fig. 1C and Fig. S1A).

The linear filter captures the kinetics of the RGC response and is analogous to the time-reversed
STA derived from spike responses (Chichilnisky, 2001). As with the STAs in Figure 1D, excitatory
linear filters were slower and more biphasic in ON-T (zero-cross time = 0.10 ± 0.013 s; biphasic
ind. = 0.45 ± 0.090; mean ± S.D.; n = 13) compared to ON-S RGCs (zero-cross time = 0.083 ±
0.0074 s; biphasic ind. = 0.31 ± 0.10; mean ± S.D.; n = 15) (Fig. 2C). This latter difference is
consistent with more transient light-evoked excitatory synaptic input to ON-T RGCs. Indeed, light
step-evoked excitatory currents to ON-T RGCs decayed to baseline levels within a few hundred
milliseconds, even in the continued presence of light input, whereas excitatory inputs to ON-S
RGCs had a sustained component that persisted for the duration of the light step (Fig. 2D-E).

These results show that transient versus sustained kinetics are already present in the excitatory
pathways that provide input to ON-T and ON-S RGCs. However, these findings do not rule out
additional contributions to the distinct spiking behavior of these RGC types from amacrine cell-
mediated inhibitory synaptic input to the RGCs or from different intrinsic spiking properties. To
examine whether postsynaptic inhibition contributes importantly to kinetic differences between
ON-T and ON-S RGCs, we measured both excitatory and inhibitory synaptic currents in response
to Gaussian noise stimuli in a subset of RGC recordings (Fig. S3). Like excitatory currents, light-
evoked inhibitory currents were more transient in ON-T RGCs (linear filter biphasic ind. = 0.34 ±
0.12; mean ± S.D.; n= 8) compared to those in ON-S RGCs (biphasic ind. = 0.16 ± 0.09; mean ±
S.D.; n= 8) (Fig. S3B-C; p= 0.01, Wilcoxon rank sum test). We also quantified the relative strength
of excitatory and inhibitory synaptic inputs by calculating the ratio between the largest noise-
evoked excitatory and inhibitory conductances (G_{exc} and G_{inh}; derived from the measured
currents) in response to a repeated noise stimulus in each RGC (Fig. S3A, D-E). This ratio did
not differ significantly between the two cell types (ON-T RGCs: peak G_{exc} / peak G_{inh} = 1.46 ±
0.65; mean ± S.D.; n = 6; ON-S RGCs: peak G_{exc} / peak G_{inh} = 4.04 ± 2.48; mean ± S.D.; n = 7;
p=0.10, Wilcoxon rank sum test). The relatively small inhibitory inputs together with the clear
differences we observed in excitatory synaptic input between ON-T and ON-S RGCs indicate that
bipolar cell pathways have a central role in establishing transient versus sustained kinetics.

215
Figure 2*. Distinct excitatory synaptic input to ON-T and ON-S RGCs. (A) Excitatory synaptic*
currents measured in example ON-T (cyan) and ON-S (pink) RGCs in response to the same
Gaussian noise stimulus used   in Fig. 1. Traces show average currents from ten repetitions of
the same stimulus and are baseline subtracted to facilitate comparison. (B) Linear filters
computed from responses to non-repeated noise stimulus presentation. Lines and shaded areas
show mean ± SEM of responses from ON-T (n= 13) and ON-S (n= 15) RGCs. The mean line
obscures the SEM shading for much of the filter in both cell types. (C) Quantification of filter
kinetics in different RGCs. (D) Excitatory currents measured from example ON-T and ON-S
RGCs in response to a 0.5 second step increase in light intensity (50% contrast; mean response
to five repetitions of stimulus). (E) Quantification of step response kinetics (n= 19 ON-T RGCs;
n= 13 ON-S RGCs). In (C) and (E) open circles show measurements from individual cells and
filled circles with error bars show mean ± SEM values. Significance values from Wilcoxon rank
sum tests.

ON-T and ON-S RGCs receive input from distinct bipolar types

Retinal circuits dedicated to transient versus sustained signaling are generally located at the
center and borders of the IPL (Baden et al., 2016; Bae et al., 2018; Beaudoin et al., 2019; Borghuis
et al., 2013b; Franke et al., 2017; Roska and Werblin, 2001). We compared stratification patterns
of ON-T and ON-S RGCs in a subset of whole-cell recordings by filling cells with a fluorescent

dye and imaging the dendrites of physiologically-characterized RGCs using two-photon
microscopy. We assessed IPL depth in live tissue by using retinas from a transgenic mouse line
in which type 5 bipolar cells strongly express EGFP (*Gjd2-EGFP* mice). These EGFP+ bipolar
cells express calcium binding protein 5 (CaBP5; Fig. S3B-D) and have axons that occupy a narrow
band just above and overlapping with the dendrites of ON starburst amacrine cells (Fig. S3A)
(Ghosh et al., 2004). Based on the regular spacing of their somas, these cells likely correspond
to one of the three subtypes of type 5 bipolar cells previously identified in mouse retina (Greene
et al., 2016; Rodieck, 1991) (Fig. S3E-F). Imaging of RGCs together with EGFP+ axon terminals
in *Gjd2-EGFP* mice revealed that ON-T and ON-S RGC dendrites occupy distinct, mostly non-
overlapping regions of the IPL (Fig. S4).
246
The distinct IPL stratification of ON-T and ON-S RGCs indicates they receive input from different
bipolar cell populations. Recent serial block-face electron microscopy (SBFSEM) connectomic
analyses found that type 6 and type 7 bipolar cells account for ~50-52% and 46% of the excitatory
synapses formed upon ON-S RGC dendrites (Sabbah et al., 2018; Swygart et al., 2024).
However, the bipolar cell subtype(s) presynaptic to ON-T RGCs have not previously been
identified. Our observation that ON-T RGC dendrites occupy a narrow band in the IPL that
closely matches the stratification pattern of EGFP+ type 5 bipolar cell axons in *Gjd2-EGFP* mice
(Fig. S5) suggests that these bipolar cells likely provide a substantial fraction of the excitatory
input to ON-T RGCs. To test this possibility, we transfected retinas from *Gjd2-EGFP* mice with a
plasmid encoding PSD-95 fused to the red fluorescent reporter mCherry to label postsynaptic
sites of excitatory input (Fig. S5). We used confocal microscopy to quantify   the fraction of
putative synapses, defined as sites with overlapping PSD-95 and EGFP fluorescent signals, that
arose from *Gjd2-EGFP*+ axon terminals (Schwartz et al., 2012). From this analysis, we concluded
that the labeled type 5 bipolar cells in *Gjd2-EGFP* mice provide ~40% of the ribbon synaptic input
to ON-T RGCs (Fig. S5D).
262
We next performed correlative fluorescence imaging and SBFSEM to identify which type 5 bipolar
subtype is labeled in Gjd2-EGFP mice and to further characterize what other bipolar cell types
provide the remaining input to ON-T RGCs. We first used 2P microscopy to image the dendritic
morphology of two dye-filled ON-T RGCs that were identified from their light-evoked spike
responses in flat-mounted retina from *Gjd2-EGFP* transgenic mice (Fig. 3A). After 2P imaging,
we fixed the retina and used the near infrared branding (NIRB) technique to create fiducial marks
in the tissue around the dye-filled ON-T RGCs (Bishop et al., 2011). After acquiring the SBFSEM
volume, we located the somas of the two ON-T RGCs using these fiducial marks and
reconstructed the dendritic arbors of both cells (e.g. Fig. 3B). The dendritic morphologies from
our EM reconstruction matched those from the light microscopy images. This suggests that we
successfully identified the two ON-T RGCs recorded physiologically in our EM volume. We chose
the two longest dendritic arbors from the RGC soma to the dendritic tip and identified all the
synapses from bipolar cells along these dendrites (Fig. 3C). We then reconstructed all the
presynaptic bipolar cells and classified them according to their axon morphology and stratification
(Fig. 3D). Careful examination of axonal tiling among type 5i, 5o and 5t subtypes provided
additional confirmation of the classification of these type 5 bipolar cells (Fig. S7A) (Greene et al.,
2016). When comparing the EM images with light microscopy images, it is clear that *GJD2-GFP*+
bipolar cells are type 5i bipolar cells (Fig. S7B).
281

Figure 3. *Identification of bipolar cell subtypes presynaptic to ON-T RGCs. (A) Maximum-projection   z-stacks of two dye-filled ON-T RGCs (green) in wholemount retina from Gjd2-EGFP (red) transgenic mice. (B) Traced skeletons of dendritic arbors of the two ON-T RGCs from EM volume. (C) An EM micrograph showing a ribbon synapse (arrowhead) on a BC axonal terminal onto the dendrite of a ON-T RGC. (D) An example skeleton for each presynaptic BC type (types 5i, 5t, 5o, X, 6 and 7) of ON-T RGCs. Top: whole-mount view, Bottom: side view. (E) Bar plot showing the proportion of input synapses to ON-T RGCs from each BC type. Mean ± SEM (n=3 RGCs, 2 from this study and one from Ding et al. (2016)).*

Results from the SBFSEM reconstructions agreed closely with those from the light microscopy approach described above. Type 5i bipolar cell axon terminals provided the largest fraction of synapses onto the two ON-T RGCs within our SBFSEM volume (~30-35%). Other inputs arose from type 5o (~10%), type 5t (~5-15%), XBC (~15-30%), and, somewhat surprisingly given the differences in stratification, from type 6 (~20%) bipolar cells (Fig. 3E). We confirmed this connectivity pattern by additionally analyzing bipolar cell synaptic inputs to a putative ON-T RGC (RGC7089) in the same publicly available SBFSEM volume (Ding et al., 2016) that was used by Sabbah et al., (2018) to map inputs to ON-S RGCs (Fig. 3E).

Figure 4. *Bipolar cell subtypes presynaptic to ON-T and ON-S RGCs have indistinguishable*
temporal response properties. (A) Maximum-projection z-stacks of fluorescent dye-filled bipolar
cells targeted for electrophysiological recording in retinal slices from Gjd2-EGFP (left), Grm6-
tdTomato (middle), and Gus8.4-EGFP (right) transgenic mice. (B) Responses to spatially
uniform Gaussian noise stimulation (same as in Figs. 1-2) in example type 5i (cyan), type 6
(magenta) and type 7 (green) bipolar cells. Voltage traces are average responses to ten
repetitions of the noise stimulus. Scale bar = 5 mV. (C) Linear filters computed from responses
to non-repeated noise stimulus presentation. Lines and shaded areas show mean ± SEM of
responses (type 5i, n= 11; type 6, n= 7; type 7, n= 6). (D) Quantification of response kinetics
from linear filters. No significant differences across types (p = 0.19 zero-cross times; p= 0.89
biphasic index; Kruskal-Wallis test). (E) Responses to step light increment. Lines and shaded
areas show peak-normalized mean ± SEM responses across type 5i bipolar cells (n= 8) and
type 6 bipolar cells (n= 5). (F) Quantification of step response kinetics. Time to peak and
steady-state response amplitude were not different between type 5i and type 6 bipolar cells (p=
0.72 and p= 0.35, respectively; Wilcoxon rank sum test).

Bipolar cell light responses differ minimally across types

Are transient versus sustained signals already differentiated in the voltage responses of the different bipolar cells presynaptic to ON-T and ON-S RGCs? Or is the distinct excitatory synaptic input that these RGCs receive primarily due to differences at the bipolar cell to RGC synapse? To answer these questions, we compared the light-driven response properties of the bipolar cell subtypes that comprise most of the excitatory input to ON-T or ON-S RGCs. We made targeted electrophysiological recordings from type 5i, type 6 and type 7 bipolar cells in retinal slices from three transgenic mouse lines in which these different bipolar cell subtypes express fluorescent proteins (see Methods). In these experiments, we used somatic whole-cell perforated-patch recordings to limit perturbation of the G-protein signaling cascade that mediates ON bipolar cell light responses.

Somatic resting membrane potentials did not differ significantly among these bipolar cell types (mean ± S.D. V_{rest}: -38 ± 6.3 mV in type 5i (n= 11); -33.7 ± 6.3 mV in type 6 (n= 7); - 36.5 ± 4.7 mV in type 7 (n=6); Kruskal-Wallis test, p= 0.35; values are not corrected for the liquid junction potential since that is hard to estimate in perforated-patch recordings). We characterized light responses using the same spatially-uniform Gaussian noise stimulus (Fig. 4B) and linear-nonlinear model analysis (Fig. 4C and Fig. S7) as in RGC recordings to facilitate direct comparison of results from bipolar cells and RGCs (Kim and Rieke, 2001; Rieke, 2001). LN models of bipolar cell responses accurately predicted responses to noise stimuli (correlation coefficient between average response to repeated noise stimulus and LN model prediction = 0.75 ± 0.15 for n = 18 bipolar cells (n = 13 type 5; n = 3 type 6; n = 2 type 7)). Unlike for ON-T and ON-S RGCs, temporal characteristics of type 5i, type 6 and type 7 bipolar cell linear filters were indistinguishable from each other under our recording conditions (Fig 4B-D). Although there was some variability in the temporal response properties across the population of bipolar cells we recorded, these differences were not associated with morphologically and molecularly defined bipolar cell types. Rather, we observed variability in response kinetics within each bipolar cell subtype and temporal responses thus overlapped substantially across different type 5i, type 6 and type 7 bipolar cells (Fig. 4 and Fig. S7).

Responses to light steps measured in a subset of type 5i and type 6 bipolar cell recordings confirmed our conclusion from Gaussian noise stimuli that different bipolar cell types exhibit nearly identical light response kinetics under our recording conditions (Fig. 4E-F). Unlike the clear differences in light step-evoked excitatory currents we measured in ON-T and ON-S RGCs (Fig. 2D-E), type 5i and type 6 bipolar cells both had sustained responses to light steps. These findings indicate that mechanisms upstream of the generation of the cone bipolar cell voltage response, such as synaptic transmission from cones to cone bipolar cells and intrinsic response properties of the cone bipolar cells, cannot explain the kinetically distinct responses in parallel ON channels mediated by type 5i, type 6 or type 7 bipolar cells. Consistent with this conclusion, ON-T and ON-S RGCs exhibit kinetically distinct light step-evoked spiking responses under scotopic conditions (not shown), where much of the RGC response originates from signals carried from outer to inner retina via the rod bipolar cell pathway (Fig. 1B) (Grimes et al., 2014; Ke et al., 2014). Because AII amacrine cells distribute rod bipolar cell signals directly to cone bipolar cell axon terminals, differences observed under scotopic conditions likely arise at and/or downstream of cone bipolar cell synaptic output.

Figure 5. *Presynaptic inhibition   is not required for kinetic differences between bipolar cell inputs to ON-T versus ON-S RGCs. (A) Example EM micrograph showing a ribbon dyad (red arrowhead) between a BC and an ON-T RGC and an amacrine cell, with an adjacent feedback inhibitory synapse (yellow asterisk). (B) Bar plots showing the percentage of RGC/AC ribbon dyads with feedback inhibition. Mean ± SD (n=2 ON-S RGCs 85 dyads analyzed, 2 ON-T RGCs with 83 dyads analyzed) (C) Excitatory currents measured in response to light step increment in control conditions (cyan) and in the presence of GABA receptor antagonists (10 µM SR95531 and 50 µM TPMPA; dashed gray) in an example ON-T RGC. Traces are mean response from six repeats of stimulus presentation. (D) Summary of steady-state amplitudes measured from step responses in control or with GABA receptor antagonists. Step-evoked currents were significantly more transient when GABA$_{A/C}$ receptors were blocked (p= 0.031; Wilcoxon signed rank test; n= 6 RGCs) (E) Control excitatory currents (cyan) or with addition of strychnine (1 µM; dashed gray) to bath in an example ON-T RGC. (F) Summary of strychnine experiments. No significant difference between conditions (p= 0.44; Wilcoxon signed rank test; n= 5 RGCs). In (D) and (F), open circles with lines show measurements from individual ON-T RGCs. Filled circles with error bars show mean ± SEM.*

Presynaptic inhibition does not establish kinetic differences in bipolar cell-mediated excitation

Bipolar cells participate in diverse inhibitory circuits with amacrine cells, which are comprised of >40 morphologically and functionally distinct cell types (Diamond, 2017; Helmstaedter et al., 2013). Past work suggests that amacrine cell-mediated presynaptic inhibition is critical to bipolar cell subtype-specific patterns of light-evoked glutamate release (Franke et al., 2017). Our bipolar cell voltage recordings, which should reflect a combination of light-evoked excitatory and inhibitory input, argue that presynaptic inhibition is not required for generating differential synaptic output between type 5i, type 6 and type 7 bipolar cells. However, the slicing procedure we used to prepare tissue for bipolar cell recordings may damage some inhibitory connections, especially long-range amacrine cell inputs. Distance-dependent attenuation of voltage signals between bipolar cell axon terminals and the somatic recording site could also limit our ability to accurately measure the influence of presynaptic inhibition in these recordings. We therefore performed additional anatomical and pharmacological experiments to investigate a possible role for presynaptic inhibition in generating kinetically distinct excitation to ON-T and ON-S RGCs. Many bipolar cell output synapses are organized into a "dyad" arrangement in which a bipolar cell ribbon synapse releases glutamate onto the processes of two different postsynaptic cells. At dyads, bipolar cell output synapses can be presynaptic to the processes of two amacrine cells, two RGCs, or an amacrine cell and a RGC (Cohen and Sterling, 1990; Dowling and Boycott, 1966; McGuire et al., 1984; Yu et al., 2023). Dyad synapses comprised of an amacrine cell and a RGC may shape bipolar cell glutamate release dynamics onto the RGC partner because glutamatergic excitation of the postsynaptic amacrine cell can elicit release of inhibitory neurotransmitters (GABA or glycine) from the amacrine cell processes that feed back onto the bipolar cell axon terminal. This form of inhibitory feedback control of bipolar cell output, termed reciprocal inhibition, could potentially contribute to differences in glutamate release onto ON-T and ON-S RGCs.

We analyzed bipolar cell dyads with postsynaptic amacrine cell processes and either ON-T or ON-S RGC dendrites in SBFSEM volumes to examine whether reciprocal inhibition is likely to generate more transient glutamate release onto ON-T RGCs. We defined putative reciprocal synapses as bipolar cell/amacrine cell/RGC dyads in which we observed clustered vesicles within the amacrine cell process that were in close proximity to the bipolar cell ribbon synapse (Fig. 5A). Our ultrastructural analysis revealed that the prevalence of reciprocal synapses was similar at dyad synapses with postsynaptic ON-T versus ON-S RGCs (Fig. 5A-B). This finding is not consistent with an important role for feedback/reciprocal inhibition in generating differential bipolar cell output onto the different RGC types.

We also performed pharmacological experiments to further examine what role, if any, inhibition may have in establishing transient glutamate release onto ON-T RGC dendrites. We used ACSF in place of Ames solution for the dissection and maintenance of tissue for these experiments. ACSF is often used in retina experiments, including the glutamate release measurements describe below, and we wanted to check that our results generalized across conditions. Light step-evoked currents in both ON-T and ON-S RGCs were more sustained than responses measured using Ames' solution (Fig. S9A-B). Nonetheless, the difference between sustained and transient RGC types was maintained across different experimental conditions.

In ON-T RGCs, blockade of $GABA_A$ and $GABA_C$ receptors via bath co-application of SR95531 and TPMPA enhanced the peak amplitude of the initial transient component of light-step evoked excitatory synaptic currents (-2023 ± pA 372.7; mean ± S.D.) compared to control (-833.6 ± 103.2 pA) (p= 0.031, Wilcoxon sign rank test; n= 6) whereas the sustained component at the

end of the light step was not significantly affected (control: -200.7 ± 14.71 pA; GABA-R block: -
290.9 ± 43.69; p = 0.16, Wilcoxon sign rank test; n= 6) (Fig. 5C). The strong enhancement of
the initial phase of the response we observed is consistent with prior work showing that $GABA_C$
receptor-mediated inhibition limits multivesicular release from bipolar cells (Sagdullaev et al.,
2006). Block of GABA receptors thus made excitatory currents in ON-T RGCs more transient
compared to control conditions (Fig 5D). Note that this is opposite the expected effect if
presynaptic GABA receptors contribute to generating transient responses in these cells; GABA
receptor antagonists should have made excitatory currents more sustained if presynaptic
GABAergic inhibition was generating transient bipolar cell input to ON-T RGCs. Thus,
GABAergic inhibition appears to control response gain and time course of bipolar cell synaptic
output to ON-T RGCs but is not responsible for the transient kinetics of excitatory input to these
cells.

We also found that glycinergic inhibition is not likely to contribute importantly to the more
transient glutamate release from bipolar cells presynaptic to ON-T RGCs; bath application of the
glycine receptor antagonist strychnine did not significantly alter the kinetics of light step-evoked
excitatory currents in these cells (Fig. 5E-F). Taken together with our ultrastructural analysis,
these results show that amacrine cell-mediated GABAergic or glycinergic inhibition of bipolar
cell axon terminals does not explain kinetic differences in excitatory synaptic input to ON-T and
ON-S RGCs.

Bipolar cell glutamate release differs across types

Taken together, the bipolar cell and RGC recordings in Figures 2 and 4 indicate that bipolar cell
synapses have a key role in the separation of visual signals into transient or sustained channels
within ON pathways in the retina. Presynaptic and/or postsynaptic mechanisms could contribute
to the distinct kinetics of bipolar cell-mediated excitatory input to ON-T and ON-S RGCs. To
investigate whether presynaptic differences in glutamate release from bipolar cell axon
terminals shape RGC response kinetics, we expressed the fluorescent glutamate reporter
iGluSnFR on the dendrites of ON-T and ON-S RGCs (see Methods). Using two-photon
glutamate sensor imaging, we found that light-evoked glutamate signals had significantly faster
kinetics at ON-T RGC dendrites (steady-state amplitude as % of peak amplitude = 21 ± 13;
mean ± S.D.; n = 224 ROIs) compared to   ON-S RGC dendrites (steady-state amp. as % of
peak amp. = 44 ± 12; mean ± S.D.; n = 307 ROIs) (Fig. 6).

The experiments summarized in Figure 5 suggest that presynaptic inhibitory input to bipolar
axon terminals is not required for the difference between ON-T and ON-S responses. The
glutamate release measurements provided a further test. We compared the kinetics of release
before and after suppressing spiking amacrine cell activity (with bath application of the voltage-
gated Na^+ channel blocker TTX) or all amacrine cells (with bath application of the glutamate
receptor antagonist NBQX). This approach avoided the confounding changes in excitatory
signals we encountered with inhibitory receptor blockers (e.g. the increase in excitatory input to
ON-T RGCs in Figure 5C and oscillatory bursts of excitation in ON-S RGCs (not shown)). The
more transient glutamate release kinetics at ON-T vs ON-S dendrites remained even when
amacrine cell responses were   suppressed (control: steady-state amp. as % of peak amp. 13
±10; mean ± S.D.; n = 189 ROIs/4 FOVs for ON-T dendrites vs. 40 ± 12; mean ± S.D.; n = 287
ROIs/8 FOVs for ON-S dendrites; TTX: 16 ± 9; mean ± S.D.; n = 50 ROIs/3 FOVs for ON-T
dendrites vs. 35 ± 11; mean ± S.D.; n = 146 ROIs/6 FOVs for ON-S dendrites; *p<0.001; t-test);
NBQX: 7 ± 3; mean ± S.D.; n = 112 ROIs/1 FOV for ON-T dendrites vs 24 ± 9; mean ± S.D.; n =
97 ROIs/2 FOVs for ON-S dendrites; *p<;0.001; t-test; Figure 7).

492
Figure 6. *Distinct glutamate input kinetics at ON-S vs ON-T RGC dendrites. (A) ON-S RGC*
dendrites expressing iGluSnFR in a Kcng4-Cre mouse were filled with sulforhodamine 101 (left)
after measuring their spiking response to a 200 μm spot extracellularly (Right). Yellow bands
indicate stimulus duration. (B) iGluSnFR expressing ON-S and ON-T dendrites at different depths
of the inner-plexiform layer (Left) were imaged using two-photon microscopy. Example iGluSnFR
signals extracted from small regions of interest (ROIs 1-8) from ON-S and ON-T RGC dendrites
illustrate the kinetic differences in their respective inputs. Black, mean responses; gray, individual
trials. (C) Distribution of steady-state response amplitudes (mean fluorescence during last 1 s of
stimulus) of the individual (gray) and average (black) ROIs from   ON-S and ON-T dendrites
evoked by a 200 μm static spot (n = 7 FOVs, 6 retinas for ON-S and 5 FOVs, 5 retinas for ON-T
*RGC dendrites, *p<0.001; t-test beween ROIs).*

Figure 7. *Distinction between sustained and transient responses remains when amacrine cell*
activity is blocked.   A. iGluSnFR measurements of glutamate release in control conditions and
in the presence of TTX (which blocks spike-dependent amacrine cell output); or in the presence
of NBQX, which blocks AMPA receptor mediated input to all amacrine cells (as well as
horizontal cells). B. Ratio of steady state to transient response across ROIs before (top) and
during (bottom) application of TTX/NBQX.

513
514 These results show that differences in glutamate release from presynaptic bipolar cells
515 contribute to transient versus sustained ON RGC responses and that the differences in release
516 are retained without amacrine feedback to bipolar cells. These experiments do not, however,
517 rule out the possibility that additional postsynaptic mechanisms, for example differential
518 glutamate receptor expression, contribute to kinetic differences between RGC types.

519
Although whole-cell excitatory currents in ON-T and ON-S RGCs (Fig. 2, Fig. S8) and iGluSnFR
signals measured from ON-T and ON-S RGC dendrites (Fig. 6) both exhibited clear kinetic
differences between the RGC types, there were quantitative differences in the response to light
steps between the different measurement approaches. As described above, these differences
appear to arise from differences in the external solution (Ames vs ACSF). The iGluSnFR
experiments used an ACSF solution, like the patch-clamp experiments in Figs. 5 and S8. Both
iGluSnFR responses and excitatory synaptic inputs measured in ACSF showed a larger
sustained component of the step response at both ON-T and ON-S dendrites compared to
currents measured in Ames (compare Fig. 2D-E, Fig. S8A-B and Fig. 6B-C). This observation
highlights the importance of comparing responses between different RGC types, as well as
between RGCs and bipolar cells, under identical recording conditions.

531
532 Importantly, the kinetics of glutamate release measured using iGluSnFR resemble the excitatory
533 synaptic inputs to ON-T and ON-S RGCs more closely than the relevant bipolar voltage
534 responses (Fig. 4). This indicates that the difference between the ON-T and ON-S responses
535 originates at least in part in the conversion of bipolar voltage to glutamate release.
536

537   *Ribbon size and vesicle replenishment*
538
Prior work has shown that bipolar cells can exhibit profound short-term depression of glutamate
release due to use-dependent depletion of the readily-releasable pool of synaptic vesicles
(Burrone and Lagnado, 2000; Singer and Diamond, 2006, 2003; von Gersdorff and Matthews,
1997). Strong vesicle depletion at the cone bipolar cell synapses onto ON-T RGCs could
generate more transient responses in these cells compared to ON-S RGCs. To examine this
possibility, we measured excitatory synaptic inputs to ON-T and ON-S RGCs in response to a
paired flash stimulus in which two brief light flashes were separated by varying time intervals. As
in Figures 1, 2 and 4, we used Ames' medium for tissue preparation and maintenance in these
experiments. These experiments used the same photopic background light level (~3300
R*/rod/s; ~2000 P*/Scone/s; ~1650 P*/Mcone/s) as used for Gaussian noise and step stimuli.
549
Flash-evoked excitatory synaptic currents were suppressed in both RGC types at short inter-
flash intervals, consistent with incomplete recovery from vesicle depletion at the time of the
second flash (Fig. 8A top/middle, B). This suppression was much stronger for currents
measured in ON-T RGCs compared to ON-S RGCs. We quantified the extent of suppression
using a paired flash ratio, which was calculated as the peak amplitude of the response to the
second flash divided by the peak amplitude of the response to the first flash. At the shortest
interval we tested (50 ms), the response to a second flash was almost completely suppressed in
ON-T RGCs (paired flash ratio = 0.04 ± 0.03; mean ± S.D.; n = 6) whereas the response to a
second flash in ON-S RGCs was suppressed considerably less (paired flash ratio = 0.29 ± 0.08;
n = 5). To rule out the possibility that a reduction in bipolar cell voltage response to the second
flash could account for the strong paired flash suppression of excitatory input to RGCs, we also
measured type 5i bipolar cell responses to the paired flash stimuli. The amplitude of flash-
evoked voltage responses in type 5i bipolar cells was unchanged by a preceding flash at all time
intervals (Fig. 8A bottom, C), which argues that paired flash suppression arises at the bipolar
output synapse. These results are consistent with the idea that differences in the extent of
stimulus-evoked vesicle pool depletion in bipolar cells could underlie transient versus sustained
excitatory input to ON-T and ON-S RGCs.
567

Figure 8. _Stronger flash-evoked suppression of excitatory currents in ON-T versus ON-S RGCs._
(A) Paired flash-evoked excitatory currents measured at different inter-flash intervals (50, 100,
200, 400 ms) in an example ON-T (top; cyan) and an ON-S (middle, magenta) RGC. Bottom
black traces show voltage responses to same stimulus measured in an example type 5i bipolar
cell. Arrows at top indicate timing of first flash (black) and second flashes (gray). Flashes were
400% contrast and 10 ms duration. Traces show mean responses to five repeats of each paired
flash interval. (B) Filled circles with error bars show mean ± SEM paired-flash ratios at different
inter-flash intervals across all ON-T RGCs (cyan; n= 6), ON-S RGCs (magenta; n= 5) and type
5i bipolar cells (black; n= 6).

Figure 9. *Bipolar cell subtype-specific differences in ribbon size (volume). (A) 3D*
reconstructions showing segmented axon stalk and axonal terminal (gray) and all the synaptic
ribbons within the axonal terminal for a type 5i (cyan) and a type 6 (magenta) BC. (B) Violin plot
showing distribution of ribbon volumes (each data point is a ribbon) for the type 5i and type 6
BC in (A). The boxes indicate the interquartile range (thick lines, 25%-75%), median value
(white dot), and 1.5 times interquartile range (thin line). Shaded gray areas represent rotated
kernel density plots. (C) 3D reconstructions showing segmented dendritic arbor (gray) and all
the synaptic ribbons presynaptic to these segments of ON-T (cyan) or ON-S (magenta) RGC
dendrites. (D) Violin plot with scattered dots showing ribbon volumes for each ribbon
presynaptic to a stretch of reconstructed ON-T or ON-S RGC dendrite. Each ribbon (dot) is
colored differently for each BC type. (E) An EM micrograph (left) and 3D reconstruction (right) of
a ribbon and its adjacent vesicles. The ribbon is indicated by the red arrow in the EM image. In
the 3D reconstruction, yellow dots indicate the adjacent vesicles surrounding the presynaptic
ribbon (red) in the BC terminals (gray). Postsynaptic ON-T RGC is in blue. (F) Scatter plot
showing ribbon size in volume in relationship with the number of adjacent vesicles. Gray line
represents the linear regression model fitted to the data. Linear regression: $\beta = 6.98e{-}06$, p <
0.001, $r^2 = 0.66$.

We next looked for ultrastructural correlates of differential synapse function in bipolar cell axon
terminals reconstructed from SBFSEM volumes. Interestingly, type 6 bipolar cell synaptic
ribbons were ~2-3 fold larger than those of types 5i bipolar cells (Fig. 9A-B, Fig. S10A). These
differences did not depend on the identity of the postsynaptic RGC partner (Fig. 9D, Fig. S10B).
Ribbons in XBCs, type 5o and type 5t bipolar cells were also smaller than type 6 bipolar cell
ribbons (Fig. 9D, Fig. S10B). This bipolar cell subtype-specific difference in ribbon size and the
difference in ratio of type 6 to type 5i input between RGC types meant that ribbons were on
average smaller at synapses made on ON-T compared to ON-S dendrites (Fig. 9C-D, Fig.
S10B).
606
607   ***Discussion***
608
We sought to understand how kinetically distinct visual pathways are established in the retina by
directly comparing the visual response properties of transient and sustained ON RGCs with their
presynaptic bipolar cells. Our results show that bipolar cell synapses have a key role in
establishing different temporal channels in ON circuits of the retina. Further, amacrine cell-
mediated presynaptic inhibition does not appear to account for transient versus sustained
bipolar cell synaptic output. Rather, bipolar cell subtype-specific differences in synaptic release
properties likely generate kinetically distinct output to postsynaptic targets. Our anatomical
analysis suggests that these differences may at least in part be due to differences in synaptic
vesicle pool size.
618
A key aspect of our work is that we measured RGC and bipolar cell responses using identical
stimuli and under the same experimental recording conditions. To our knowledge, a direct
comparison of RGC responses with those of their presynaptic bipolar cells has not previously
been reported. The importance of measuring RGC and bipolar cell responses under the same
conditions is highlighted by our observation that excitatory synaptic currents measured in ON
alpha RGCs exhibited quantitatively different kinetics, although relative differences between
types were preserved, when measurements were made from retinas maintained in ACSF
versus Ames' bath solution (compare Fig. 2 (Ames') and Figs. 5 and S9 (ACSF)).
627
We found that the bipolar cell subtypes that provide the major input to ON-T (type 5i) or ON-S
RGCs (types 6 and 7) had indistinguishable light-evoked voltage responses (Figure 4). This was
not entirely surprising given that most or all ON cone bipolar cell subtypes are indirectly coupled
to other nearby ON cone bipolar cells via electrical synapses between bipolar cell axon
terminals and the distal processes of AII amacrine cells. These electrical synapses form the
basis for the primary rod pathway that mediates vision under dim light conditions, but lateral
signals can also flow among neighboring bipolar cells under conditions where cone signaling
dominates (Demb and Singer, 2012). Indeed, a substantial fraction (~30-50%) of light-evoked
responses in the bipolar cell types we investigated here can be attributed to lateral signals that
arrive via electrical synapses at photopic light levels (Kuo et al., 2016). Each AII amacrine cell is
electrically coupled to multiple ON cone bipolar subtypes as well as to other AII amacrine cells
(Marc et al., 2014; Tsukamoto and Omi, 2017; Veruki and Hartveit, 2002a, 2002b). Different ON
cone bipolar cell subtypes may also be directly coupled via gap junctions to each other
(Sigulinsky et al., 2020; Tsukamoto and Omi, 2017).
642
Direct or indirect (via AII) coupling may limit the extent to which different bipolar cells can
generate subtype-specific voltage responses. These findings differ from a previous study that
found some differences among mouse ON cone bipolar cell subtypes. Ichinose et al., (2014)
reported that a subset of type 5 bipolar cells, which the authors termed type 5f, and type 7

bipolar cells had more transient responses than type 6 bipolar cells. Methodological differences
may account for the discrepancy between our results and the previous work by Ichinose et al.,
(2014). For example, Ichinose et al. included inhibitory receptor antagonists in their
experimental solutions and used 100 μm diameter spots to characterize bipolar cell responses
whereas we left inhibitory signaling intact and   used larger (520 μm diameter) spots to match
the stimuli we used for RGC recordings. Interestingly, a recent investigation of spatial and
temporal response properties of light-evoked intracellular calcium signals across bipolar cell
types suggests that ON bipolar cells can exhibit type-specific response kinetics for small stimuli
(<~200 μm diameter) but not larger spot sizes, presumably due to surround inhibition (Hsiang et
al., 2024). Regardless of the reasons for the different findings, our observation that bipolar cell
subtypes exhibited similar responses under the same recording and light stimulus conditions for
which we observed obvious differences among postsynaptic RGCs argues that mechanisms
upstream of bipolar cell glutamate release (e.g. cone to bipolar cell synapses or intrinsic voltage
response properties of bipolar cells) are unlikely to have a major role in shaping RGC type-
specific responses for the stimuli we used here.
662
A limitation of our work is that we did not record from every one of the bipolar cell subtypes that
are presynaptic to ON-T RGCs. Instead, we measured responses from type 5i and type 6
bipolar cells, which together comprise >~50% of the synaptic input to ON-T RGC dendrites. It is
possible that the other presynaptic bipolar cell types (type 5o, type 5t and XBC) may have more
transient light-evoked voltage responses that could contribute to the transient excitation
measured in ON-T RGCs. In fact, a previous study in mouse retina found that XBCs and a
subset of type 5 bipolar cells express voltage-gated Na$^+$ channels (Hellmer et al., 2016). These
channels may enhance response transience or might even contribute to the generation of light-
driven all-or-none spiking responses in these bipolar cells (Ichinose et al., 2014; Puthussery et
al., 2013; Saszik and DeVries, 2012).
673
Even if transient intrinsic response properties of type 5o, 5t and XBC bipolar cells contribute to
the generation of transient responses in ON-T RGCS,   this alone is unlikely to   account for
the kinetics of the excitatory synaptic currents we measured in these RGCs. Type 5i and type 6
bipolar cells have a significant sustained component of their light step-evoked voltage response
(Fig. 4E-F), whereas excitatory currents measured for the same stimulus and under the same
conditions in ON-T RGCs had little to no sustained component (Fig. 2D-E). We would expect to
observe some sustained excitatory input to ON-T RGCs if type 5i and type 6 bipolar cell
synaptic output follows the voltage response of these bipolar cells. Thus, a marked
transformation of the sustained bipolar cell membrane voltage response to a transient synaptic
current must take place. Our measurements of extracellular glutamate at ON-T and ON-S
dendrites (Fig. 5) suggest that much of this transformation occurs at the level of bipolar cell
glutamate release. It is interesting in this context that ON-T and ON-S RGCs both receive input
from type 6 bipolar cells (comprising ~20% and ~50% of total synaptic input, respectively). Due
to differences in dendrite stratification, type 6 bipolar cell synapses onto ON-T RGCs arise from
a region of the axon that is located more proximal to the bipolar cell soma than the synapses
formed onto ON-S RGCs, which are found within the distal axon lobules. However, we did not
observe any differences in ribbon size at distal versus proximal locations in type 6 bipolar cell
axons (Figure 9D). Whether postsynaptic target-specific synapse function, as has previously
been reported in salamander retinal bipolar cells (Asari and Meister, 2014, 2012) and mouse
starburst amacrine cells (Pottackal et al., 2021), contributes to signal diversification in circuits
involving type 6 bipolar cells will require further investigation.
695

Our electrophysiological and glutamate imaging results indicate that amacrine cell-mediated
inhibition at bipolar cell synaptic terminals is not necessary for generating transient glutamate
release kinetics onto ON-T RGC dendrites. This is seemingly at odds with prior work that has
indicated an essential role for surround inhibition in establishing type-specific bipolar cell
glutamate release dynamics (Franke et al., 2017). However, we note that our analysis here
focused only on whether presynaptic inhibition shapes the kinetics of excitatory inputs to ON-T
RGCs. Our anatomical and pharmacological experiments together argue that presynaptic
inhibition is not an important factor in the transformation from sustained bipolar cell voltage
responses to transient excitatory synaptic currents in ON-T RGCs. This is not to say that
amacrine cell-mediated inhibition does not have a role in generating subtype-specific response
properties for other aspects of bipolar cell function. Indeed, inhibition does have an important
role in establishing differential spatial tuning of excitatory inputs to the RGC types we studied;
excitatory synaptic currents in ON-T RGCs exhibit stronger surround inhibition than ON-S
RGCs, and this surround inhibition is reduced by bath application of GABA receptor antagonists
(Fig. S9C). Interestingly, recent work revealed postsynaptic target-specific surround inhibition at
neighboring type 6 bipolar cell output synapses within the same axon terminal (Swygart et al.,
2024). Whether target-specific presynaptic inhibition of type 6 bipolar cells also contributes to
differential surround inhibition between ON-T and ON-S RGCs or is primarily due to differences
across bipolar cell types (e.g. type 5i/5t/XBC vs. type 6/7), remains to be determined.
715
We conclude that transient versus sustained excitatory inputs to ON-T and ON-S RGCs most
likely arise from bipolar cell subtype-specific differences in synaptic release properties.
Excitatory inputs to ON-T RGCs exhibited stronger paired flash depression than those to ON-S
RGCs (Fig. 8), which indicates distinct vesicle cycling dynamics at the bipolar cell axon
terminals presynaptic to the different RGC types. Our ultrastructural analysis suggests that
bipolar subtype-dependent differences in ribbon size may be a factor in establishing transient
versus sustained glutamate release kinetics (Fig. 9). It is important to note, however, that type 7
bipolar cells (~45% of excitatory input to ON-S RGCs) have small synaptic ribbons, yet
iGluSnFR imaging of type 7 bipolar cell axon terminals found that these cells exhibit sustained
glutamate release (Srivastava et al., 2022).
726
We found that ribbon size is strongly correlated with the total number of vesicles that are in
close contact with the synaptic ribbon across different ON bipolar cell types (Fig. 9E-F, Fig.
S10C). Thus, there are fewer ribbon-associated vesicles at glutamatergic synapses formed on
ON-T RGC dendrites compared to those at ON-S RGCs. Ribbon-associated vesicles are
thought to correspond to the releasable pool of vesicles, comprised of both a readily-releasable
pool that are docked at the base of the ribbon, at the presynaptic active zone, and a reserve
(alternatively referred to as 'releaseable') pool of vesicles that are bound to the ribbon at more
distal sites (Grabner et al., 2023; Lagnado and Schmitz, 2015; von Gersdorff et al., 1996; Wan
and Heidelberger, 2011). The number of ribbon-associated vesicles, as well as the rate at which
the readily-releasable and reserve vesicle pools are replenished following exocytosis, together
define the amplitude and time course of stimulus-evoked vesicle release (Jackman et al., 2009;
Lagnado and Schmitz, 2015; Neves and Lagnado, 1999; Oesch and Diamond, 2011; Singer
and Diamond, 2006; von Gersdorff and Matthews, 1997). For example, a small pool of readily-
releasable vesicles and relatively slow replenishment in rod bipolar cell axon terminals
contributes to transient release kinetics from these cells (Singer and Diamond, 2006). Our
observation that the average excitatory synaptic input to ON-T RGCs is smaller and that most
excitatory input to ON-T cells originates from bipolar synapses with fewer ribbon-associated
vesicles are consistent with a role for vesicle depletion in establishing transient glutamate
release onto these RGCs. The larger pool of ribbon-associated vesicles in type 6 bipolar cell

terminals, which comprise ~ 55% of the synaptic input to ON-S RGCs, may support sustained
synaptic transmission onto these RGCs. However, additional mechanisms, such as bipolar cell
type-specific differences in the rate of refilling of the ribbon-associated vesicles could also
contribute to sustained versus transient glutamate release.
750
Direct measurements of exocytosis at ribbon synapses show that release rates increase with
increasing ribbon size and the number of vesicles bound to the ribbon. Evidence for this
dependence includes a tight correspondence between functional measures of the releasable
vesicle pool and anatomical measurements of the number of vesicles bound to the ribbon
(Grabner et al., 2023; von Gersdorff et al., 1996). Ribbon size and the number of ribbon-
associated vesicles vary across the day-night cycle; these morphological changes are
correlated with diurnal changes in maintained vesicle fusion rates (Babai et al., 2016; Dembla et
al., 2020; Hull et al., 2006). Finally, release rates decrease when ribbon size is reduced (Mehta
et al., 2013; Wakeham et al., 2023), or when ribbons are ablated either abruptly or genetically
(Grabner and Moser, 2021; Maxeiner et al., 2016; Mehta et al., 2013; Snellman et al., 2011).
These biophysical studies are consistent with the correlation that we observe here between
ribbon size and the extent to which transmitter release is transient vs. sustained. A note of
caution in making this comparison is that exocytosis is sensitive to the voltage trajectory
(Snellman et al., 2009), and the biophysical studies often use more rapid and larger voltage
changes than those elicited by physiological light inputs. In fact, genetic deletion of ribbons does
not strongly alter light-evoked excitatory currents in ON-S RGCs (Okawa et al., 2019). Why
ribbon deletion has relatively little effect on light-evoked responses in ON-S RGCs remains to
be determined. It is interesting to note that ~30% of synapses formed by type 6 bipolar cells
onto ON-S RGC dendrites lack ribbons even in wildtype retina (Okawa et al., 2019). Future
studies will be needed to investigate to what extent such ribbonless synapses could contribute
to bipolar cell type-specific differences in glutamate release such as we observed here.
772
The dynamics of transmitter release are also impacted by vesicle depletion and the time
required for refilling of releasable vesicle pools following strong stimulation. In particular, the
sustained transmitter release rate should be equal to the sustained refilling rate. Direct
measurements of refilling at the cone output synapse following strong activation show that it is
quite slow – with a time constant near 1 sec (Grabner et al., 2016; Jackman et al., 2009; Rabl et
al., 2006). This is far too slow to explain the recovery from paired-flash depression that we
observe here. Several factors may contribute to this difference. First, postsynaptic receptor
saturation can contribute to paired-pulse depression, and it can recover from such depression
more quickly than presynaptic release (Grabner et al., 2016). Second, the paired-flash
depression that we measure in RGC synaptic inputs is largely absent in the bipolar voltage
responses. This suggests that it is dominated by a history-dependence at the cone bipolar
output synapses and refilling of vesicle pools may proceed more quickly at bipolar synapses
than cone synapses.
786
Mechanisms that we did not directly investigate here may also contribute to the difference
between transient and sustained responses. For example, presynaptic resting potential has a
strong effect on steady-state vesicle release and thus stimulus-evoked release dynamics.
Exocytosis that occurs when bipolar cells rest at voltages near release threshold can lead to a
depletion of readily-releasable vesicles that limits the amount of release that occurs upon further
light-induced depolarizations (Grimes et al., 2014; Jarsky et al., 2011; Oesch and Diamond,
2011). Thus, an initial phasic component of release can be reduced by depolarized resting
potentials whereas hyperpolarized potentials can facilitate greater occupancy of vesicles in the
releasable pool, which can enhance response transience. Bipolar subtype-dependent

differences in release threshold due to differential Ca^{2+} channel expression/regulation (Pangrsic
et al., 2018) and/or coupling between Ca^{2+} channel activity and vesicle release (Burrone et al.,
2002; Burrone and Lagnado, 2000) could also impact release dynamics. Several aspects of
excitatory synaptic currents in the two RGC types we studied here are consistent with a
situation in which bipolar cell inputs to ON-T RGCs have a higher threshold for glutamate
release compared to those at ON-S RGCs; ON-T RGCs receive less spontaneous excitatory
synaptic input than ON-S RGCs (Kuo et al., 2016) and stimulus-evoked EPSCs are slower,
sparser and more biased towards larger amplitude events in ON-T RGCs compared to those in
ON-S RGCs (Fig. 2, Fig. S1B). Further investigation will be required to determine whether this is
due to differences in axon terminal resting potential and/or other presynaptic factors. We did not
find evidence for subtype-specific differences in resting membrane potential among type 5i, type
6 and type 7 bipolar cells, but our somatic voltage recordings likely have limited ability to resolve
small voltage differences at the bipolar cell axon terminals.
809
Our results do not rule out additional roles for postsynaptic mechanisms in establishing transient
versus sustained RGC spike output. For example, delayed amacrine cell-mediated inhibitory
synaptic input to RGC dendrites could truncate spiking responses (Nirenberg and Meister,
1997). We did not observe a significant difference in the light-evoked excitation/inhibition
amplitude ratio between ON-S and ON-T RGCs (Fig. S3E), but our data does not   exclude a
role for postsynaptic inhibition in generating transient responses. RGC cell-intrinsic
mechanisms, such as RGC subtype-specific differences in voltage-gated Na^+ channel
properties and/or expression pattern could also contribute to distinct response kinetics
(Brombas et al., 2022; Chang et al., 2023; Werginz et al., 2020). While a combination of pre-
and postsynaptic factors may ultimately act in concert to generate transient versus sustained
RGC spike output (Warwick et al., 2018; Werginz et al., 2020), our results establish a
fundamental role for bipolar cell synapses in the segregation of visual input into kinetically
distinct channels in ON pathways of the retina.
823
824 **_Materials and Methods_**
825
826 *Animals*
827
Experiments used 5- to 15-week-old wild-type, *Grm6-tdTomato* (Morgan et al., 2011), *Gus8.4-*
EGFP (Huang et al., 1999), *Gjd2-EGFP*, or Kcng4-Cre (Duan et al., 2014) transgenic mice.
Transgenic mouse strains were backcrossed into the same genetic background as wild-type
mice (C57BL/6J; Jackson Laboratory). Animal care and handling followed procedures approved
by the Institutional Animal Care and Use Committee of the University of Washington or the
University of Victoria Animal Care Committee.
834
835 *Tissue preparation*
836
Retinal tissue was prepared for physiological recordings as previously described (Kuo et al.,
2016; Srivastava et al., 2022). Briefly, mice were killed by cervical dislocation or decapitation
and retinas were dissected from both eyes in oxygenated (95% O_2, 5% CO_2), bicarbonate-
buffered Ames' (Sigma) or artificial cerebrospinal fluid (ACSF) solution and either flat-mounted
with the vitreal surface facing up onto a poly-D-lysine coated glass coverslip (Corning) (RGC
patch-clamp recordings and iGluSnFR imaging) or sliced in an orientation perpendicular to the
flat-mount plane using a vibratome (200 μm slice thickness; bipolar cell recordings) before being
transferred to an upright 2P fluorescence microscope. ACSF contained (in mM): 125 NaCl, 2.5
KCl, 1.2 $CaCl_2$, 1.5 $MgCl_2$, 1.25 NaH_2PO_4, 20 glucose, 0.5 L-glutamine, 0.1 sodium ascorbate,
and 26 $NaHCO_3$ (pH 7.4, 310mOsm/L). For preparation of retinal slices, isolated retina was

surrounded with melted agarose (maintained at ~40ºC) followed by rapid cooling using an ice
block. The embedded retina was then positioned in front of the vibratome blade in an orientation
orthogonal to the flat mount plane for sectioning. All experiments exclusively used tissue from
ventral retina.
851
852 *Electrophysiology*
853
During recordings, retinal tissue was continuously perfused (~6-8 mL/min) with oxygenated,
bicarbonate-buffered Ames' solution maintained at 30-34 ºC. In a subset of electrophysiological
recordings (Figs. 6C-F and S8), tissue was perfused with ACSF rather than Ames' solution.
Electrical signals were acquired at 10 kHz and low-pass filtered at 3 kHz using a Multiclamp
700B or Multiclamp 700A amplifier (Molecular Devices) and Symphony Data Acquisition
software (http://symphony-das.github.io). The somas of different ON bipolar cell subtypes were
targeted for gramicidin perforated-patch current-clamp recordings using 2P laser scanning
excitation (960nm) in retinal slices prepared from transgenic mouse lines in which bipolar cell
subtypes expressed fluorescent proteins. Tissue from *Grm6-tdTomato* mice, in which a sparse
subset of ON bipolar cell types (type 6, 7 and 8) express tdTomato was used to target
recordings to type 6 bipolar cells (Morgan et al., 2011; Schwartz et al., 2012). Type 7 bipolar
cells were targeted for recording in *Gus8.4-EGFP* mice, which express EGFP in a type 7 bipolar
cells and rod bipolar cells (Huang et al., 2003). Type 5i bipolar cells were targeted in retinas
from *Gjd2-EGFP* mice, which we found express EGFP selectively in these bipolar cells (Fig. S6)
as well as less robustly in an uncharacterized amacrine cell type (not shown). For bipolar cell
recordings, we minimized light exposure from the 2P laser by restricting scan regions to the
inner retina, away from photoreceptors, and by keeping laser exposure brief (<~30 sec/scan
region; <2 mW post-objective laser power) (Kuo et al., 2016). After identification of fluorescent
cell bodies, we acquired recordings using infrared differential interference contrast microscopy.
Bipolar recordings used an internal solution containing 123 mM K-aspartate, 10 mM KCl, 10 mM
HEPES, 1 mM $MgCl_2$, 1 mM $CaCl_2$, 2 mM EGTA, 4 mM Mg-ATP, 0.5 mM Tris-GTP, 100 µM
Alexa Fluor 594-hydrazide and 25-30 µg/mL gramicidin D (280 mOsm/L; pH 7.2). Pipettes (~7-9
Ω tip resistance) were front-filled with gramicidin-free internal solution to facilitate formation of
high resistance (>1 GΩ) seals between patch pipettes and the cell membrane. After seal
formation, series resistance (R_s) was monitored in voltage-clamp mode by applying a 10 mV
voltage step across the pipette tip. Experiments were initiated once R_s declined to <50 MΩ,
which typically took 20-30 min. During recordings, we occasionally confirmed the integrity of the
cell membrane under the recording electrode by monitoring Alexa Fluor 594 fluorescence using
2P excitation; if any dye was visible in the cell body, we abandoned the recording. No holding
current was applied during current-clamp recordings from bipolar cells. At the conclusion of
each bipolar cell recording, the membrane under the patch pipette was intentionally ruptured
and fluorescence images of Alexa Fluor 594-filled cells were acquired using 2P microscopy to
confirm bipolar cell subtype identities (Wässle et al., 2009).
887
RGC recordings were performed in a flat-mount preparation of the retina from wild-type mice or
Gjd2-EGFP transgenic mice following previously described methods (Kuo et al., 2016).
Excitatory synaptic currents were isolated in whole-cell voltage-clamp recordings by holding
RGCs near the reversal potential for Cl- mediated conductances (-68.5 mV). Patch pipettes
(2.5-4 MΩ) were filled with an internal solution containing (in mM): 105 Cs-methanesulfonate, 10
Tetraethylammonium-Cl, 20 HEPES, 10 EGTA, 2 QX-314, 5 Mg-ATP, 0.5 Tris-GTP (~280
mOsm; pH 7.2). Series resistance (<15 MΩ) was compensated by 70% using the amplifier
circuitry. Inhibitory receptor antagonists (SR-95531, TPMPA, strychnine) were obtained from
Tocris or Sigma Aldrich and added to the bath solution from concentrated aqueous stock
solutions.

iGluSnFR expression and two-photon imaging

Intravitreal injection of a cre-dependent viral vector (pAAV.hSyn.Flex.iGluSnFR.WPRE.SV40;
Addgene Catalog #98929-AAV1) was performed into Kcng4-cre mice to selectively express the
glutamate sensor, iGluSnFR, in alpha RGCs, including ON-T and ON-S RGCs (Krieger et al.,
2017). All injection procedures, tissue preparations, and subsequent two-photon imaging were
performed as previously described in Srivastava et al., 2022. Briefly, for iGluSnFR imaging, the
laser was tuned to 920 nm and all recordings were acquired at a frame rate of 58.25 Hz (256 x
256 pixels). In KCNG4-Cre mice, ON-S alpha RGCs were first identified by their spiking
properties using extracellular cell-attached recordings. To identify the dendrites of ON-S RGCs,
cells were filled with sulphorhodamine 101. ON-T RGC dendrites were then recorded at the
same location but deeper into the retina (Krieger et al., 2017).

Visual stimulation

Visual stimuli were presented from a photopic mean background illumination (~3,300 rhodopsin
isomerizations per rod per second (R*/rod/s)) to focus specifically on cone-mediated signaling
(Grimes et al., 2018). Visual stimuli for electrophysiological recordings were generated using
Stage Visual Stimulation Software (http://stage-vss.github.io). For all electrophysiological
recordings except those in Figs. 6C-F and S8, light stimuli consisted of a spatially uniform
circular spot (520 μm diameter) delivered from a LED (405 nm peak output; Hosfelt) that was
focused through the microscope substage condenser (Kuo et al., 2016). In the experiments
shown in Figs. 6C-F and S8, a Digital Light Processing (DLP) projector driven by a 405 nm LED
(Wintech Digital) was used to project spots of various sizes through the microscope objective
(Kuo et al., 2020). For iGluSnFR imaging (Fig. 5), background illuminance was kept around
1000 photons/μm^2/s. For experiments, static spots of different sizes were generated and
presented using StimGen software (https://github.com/benmurphybaum/StimGen). Stimuli were
presented via a DLP through the substage condenser focused onto the photoreceptor layer of
the retina. All visual stimuli were presented 4 s after the start of the recording for a duration of 2
s.

Data analysis and statistics

Spike triggered average (STA) waveforms (Fig. 1) and linear-nonlinear (LN) models of light-
driven RGC (Fig. 2) and bipolar cell (Fig.4) responses were derived from responses to randomly
fluctuating light stimuli (Gaussian distribution of light intensities with standard deviation = 50% of
mean intensity; 0-60 Hz bandwidth) as described previously (Chichilnisky, 2001; Kim and Rieke,
2001; Rieke, 2001). For LN model construction, the linear filter was calculated by computing the
cross-correlation of the light stimulus with the measured voltage (bipolar cell) or current (RGC)
response and dividing the result by the power spectrum of the stimulus. The static nonlinearity
was constructed from a point-by-point comparison of the predicted filter output with the
measured response and binning the result along the prediction axis. In almost all cases, the
linear filter had a biphasic shape with initial and secondary lobes that are opposite polarity. A
small number of bipolar cells had triphasic filters (Fig. S7). We quantified the kinetics of each
STA or linear filter by measuring the time at which the STA/filter waveform crossed zero ("zero-
cross time"; e.g. Fig. 1C, left) between the first and second lobes of the filter (e.g. between peak
and trough of STA waveforms) and by calculating a biphasic index = |A| / |B|, where |B| is the
absolute peak amplitude of the initial lobe of the filter and |A| is the amplitude of the peak
amplitude of the secondary lobe (e.g. Fig. 2C, right). We quantified the kinetics of responses to
light step stimuli by calculating the time between light step onset and the peak amplitude

response ("time to peak"; e.g. Fig. 2E, left) and dividing the mean steady-state response
(calculated from the last 100 ms light step increment) by the peak amplitude initial response at
step onset (and multiplying by 100) (e.g. Fig. 2E, right). We note that alternative measures of
step response or linear filter kinetics than those presented in Fig. 4 were also not statistically
different between bipolar cell subtypes (not shown). For light step-evoked responses, steady-
state measurements were obtained from the mean current/voltage over the last 100 ms of the
500 ms light steps (electrophysiology) or the mean fluorescence over the last 1 sec of the 2 sec
light step (iGluSnFR). Statistical tests were performed using MATLAB (Mathworks) or Igor Pro
(WaveMetrics) software.
958
959 *Biolistic transfection*
960
DNA plasmids encoding mTFP1-myc (24 µg) and PSD95-mCherry (12 µg) under the control of
the cytomegalovirus (CMV) promoter were coated onto gold particles (1.6 µm diameter, 12.5
mg, Bio-Rad). Using a Helios gene gun (40 psi, Bio-Rad), these particles were then delivered to
retinal ganglion cells in whole-mount retinas. Subsequently, the transfected retinas were
incubated overnight in mACSF within a humid, oxygenated chamber maintained at 33°C.
Immunohistochemistry
967
Retinal sections (60 µm thick) were prepared using a vibratome (Leica) from retinal whole
mount retinas of *Gjd2-EGFP* mice. Both whole mount retinas and retinal sections were fixed in
4% PFA for 30 minutes. Retinal sections were embedded in low melting point agarose (Sigma-
Aldrich). Retinal tissues were first incubated with blocking solution (5% normal donkey serum,
0.5% Triton-X100 in PBS, Sigma-Aldrich) for 2 hours at 4°C, then incubated overnight at 4°C
with primary antibodies dissolved in blocking solution. Excess primary antibody was rinsed 3x10
minutes with PBS, then sections were incubated for 2 hours with secondary antibodies
dissolved in PBS. Excess secondary antibody was rinsed 3x10 minutes with PBS and sections
were mounted on standard microscopy glass slides using Vectashield mounting medium
(Vector). Primary antibody used in the study were: goat anti-ChAT diluted 1:1000 Chemicon
AB144P, mouse anti-CaBP5 diluted 1:1000 courtesy of Dr. Haeseleer, mouse anti-myc diluted
at 1:1000, chicken anti-GFP diluted at 1:1000. Secondary antibody were donkey anti-goat IgG
Dylight-568 diluted 1:1000, donkey anti-mouse Dylight-647 diluted 1:1000, donkey anti-mouse
Dylight-405 diluted at 1:1000 , donkey anti-chicken Alexa 488, the nuclear dye TOPRO3 (diluted
1:200, Thermo-Fisher).
983
984 *Near-infrared branding (NIRB) and serial block face scanning electron microscopy (SBFSEM)*
985
Two neighboring ON-T RGCs were physiologically identified in *Gjd2-EGFP* mice. After
identifying ON-T RGCs via their characteristic step-evoked extracellular spike response (e.g.
Figure 1B), they were filled with Alexa 488 or Alexa 555. Dendritic arbors and T5i-BC axonal
arbors were imaged using a confocal microscope. The retina was then fixed with 4%
glutaraldehyde for 30 minutes at room temperature (RT). The retina was washed with 0.1M Na+
Cacodylate buffer for three times and flat-mounted with 0.1M Na+ Cacodylate buffer. To
relocate the two RGCs under electron microscopy, we applied the near-infrared branding
(NIRB) technique previously published to burn fiducial markers in the retina (Bishop et al.). After
NIRBing, the retina tissue was unmounted and fixed in 4% glutaraldehyde overnight. The retina
was processed for SBFSEM subsequently according to the protocol described previously (Della
Santina et al., 2016). Briefly, the tissue was initially treated with reduced osmium for 1 hour in
the refrigerator. Subsequently, after washing with double-distilled H_2O (ddH2O), the tissue was
placed in a freshly prepared thiocarbohydrazide solution for 20 minutes at RT. Following
another round of ddH2O rinsing, the tissue was incubated in 2% osmium tetroxide for 30

minutes at RT. After rinsing, the sample was incubated in 1% uranyl acetate at 4°C overnight.
Post-wash, the tissue was stained with Walton's lead aspartate for 30 minutes at 60°C.
Following another wash, the retinal piece underwent dehydration using a graded alcohol series
(20%, 50%, 70%, 90%, 100%, 100%), each step lasting 5 minutes, followed by two changes of
100% propylene oxide at RT for 10 minutes each. Finally, the tissue was embedded in Epon.
The resulting block was then trimmed and mounted in the SBEM microscope (GATAN/Zeiss
SIGMA, 3View), and image stacks were acquired at a voxel size of 5 x 5 x 50 nm.
1007
1008
1009

References

Angueyra JM, Rieke F. 2013. Origin and effect of phototransduction noise in primate cone photoreceptors. *Nat Neurosci* **16**:1692–1700. doi:10.1038/nn.3534

Asari H, Meister M. 2014. The projective field of retinal bipolar cells and its modulation by visual context. *Neuron* **81**:641–652. doi:10.1016/j.neuron.2013.11.029

Asari H, Meister M. 2012. Divergence of visual channels in the inner retina. *Nat Neurosci* **15**:1581–1589. doi:10.1038/nn.3241

Awatramani GB, Slaughter MM. 2000. Origin of transient and sustained responses in ganglion cells of the retina. *J Neurosci* **20**:7087–7095. doi:10.1523/JNEUROSCI.20-18-07087.2000

Babai N, Sendelbeck A, Regus-Leidig H, Fuchs M, Mertins J, Reim K, Brose N, Feigenspan A, Brandstätter JH. 2016. Functional Roles of Complexin 3 and Complexin 4 at Mouse Photoreceptor Ribbon Synapses. *J Neurosci* **36**:6651–6667. doi:10.1523/JNEUROSCI.4335-15.2016

Baden T, Berens P, Bethge M, Euler T. 2013. Spikes in mammalian bipolar cells support temporal layering of the inner retina. *Curr Biol* **23**:48–52. doi:10.1016/j.cub.2012.11.006

Baden T, Berens P, Franke K, Román Rosón M, Bethge M, Euler T. 2016. The functional diversity of retinal ganglion cells in the mouse. *Nature* **529**:345–350. doi:10.1038/nature16468

Bae JA, Mu S, Kim JS, Turner NL, Tartavull I, Kemnitz N, Jordan CS, Norton AD, Silversmith WM, Prentki R, Sorek M, David C, Jones DL, Bland D, Sterling ALR, Park J, Briggman KL, Seung HS, Eyewirers. 2018. Digital Museum of Retinal Ganglion Cells with Dense Anatomy and Physiology. *Cell* **173**:1293-1306.e19. doi:10.1016/j.cell.2018.04.040

Beaudoin DL, Kupershtok M, Demb JB. 2019. Selective synaptic connections in the retinal pathway for night vision. *J Comp Neurol* **527**:117–132. doi:10.1002/cne.24313

Bishop D, Nikić I, Brinkoetter M, Knecht S, Potz S, Kerschensteiner M, Misgeld T. 2011. Near-infrared branding efficiently correlates light and electron microscopy. *Nat Methods* **8**:568–570. doi:10.1038/nmeth.1622

Borghuis BG, Marvin JS, Looger LL, Demb JB. 2013a. Two-photon imaging of nonlinear glutamate release dynamics at bipolar cell synapses in the mouse retina. *J Neurosci* **33**:10972–10985. doi:10.1523/JNEUROSCI.1241-13.2013

Borghuis BG, Marvin JS, Looger LL, Demb JB. 2013b. Two-photon imaging of nonlinear glutamate release dynamics at bipolar cell synapses in the mouse retina. *J Neurosci* **33**:10972–10985. doi:10.1523/JNEUROSCI.1241-13.2013

Brombas A, Zhou X, Williams SR. 2022. Light-evoked dendritic spikes in sustained but not transient rabbit retinal ganglion cells. *Neuron* **110**:2802-2814.e3. doi:10.1016/j.neuron.2022.06.009

Burrone J, Lagnado L. 2000. Synaptic depression and the kinetics of exocytosis in retinal bipolar cells. *J Neurosci* **20**:568–578. doi:10.1523/JNEUROSCI.20-02-00568.2000

Burrone J, Neves G, Gomis A, Cooke A, Lagnado L. 2002. Endogenous calcium buffers regulate fast exocytosis in the synaptic terminal of retinal bipolar cells. *Neuron* **33**:101–112. doi:10.1016/s0896-6273(01)00565-7

Cafaro J, Rieke F. 2013. Regulation of spatial selectivity by crossover inhibition. *J Neurosci* **33**:6310–6320. doi:10.1523/JNEUROSCI.4964-12.2013

Cangiano L, Asteriti S, Cervetto L, Gargini C. 2012. The photovoltage of rods and cones in the
dark-adapted mouse retina. *J Physiol* **590**:3841–3855. doi:10.1113/jphysiol.2011.226878
Chang L, Ran Y, Yang M, Auferkorte O, Butz E, Hüser L, Haverkamp S, Euler T, Schubert T. 2023.
Spike desensitisation as a mechanism for high-contrast selectivity in retinal ganglion
cells. *Front Cell Neurosci* **17**:1337768. doi:10.3389/fncel.2023.1337768
Chichilnisky EJ. 2001. A simple white noise analysis of neuronal light responses. *Network*
12:199–213.
Cleland BG, Levick WR, Sanderson KJ. 1973. Properties of sustained and transient ganglion cells
in the cat retina. *J Physiol* **228**:649–680. doi:10.1113/jphysiol.1973.sp010105
Cohen E, Sterling P. 1990. Demonstration of cell types among cone bipolar neurons of cat
retina. *Philos Trans R Soc Lond B Biol Sci* **330**:305–321. doi:10.1098/rstb.1990.0201
Demb JB, Singer JH. 2012. Intrinsic properties and functional circuitry of the AII amacrine cell.
Vis Neurosci **29**:51–60. doi:10.1017/S0952523811000368
Dembla E, Dembla M, Maxeiner S, Schmitz F. 2020. Synaptic ribbons foster active zone stability
and illumination-dependent active zone enrichment of RIM2 and Cav1.4 in
photoreceptor synapses. *Sci Rep* **10**:5957. doi:10.1038/s41598-020-62734-0
DeVries SH. 2000. Bipolar cells use kainate and AMPA receptors to filter visual information into
separate channels. *Neuron* **28**:847–856. doi:10.1016/s0896-6273(00)00158-6
Diamond JS. 2017. Inhibitory Interneurons in the Retina: Types, Circuitry, and Function. *Annu*
Rev Vis Sci **3**:1–24. doi:10.1146/annurev-vision-102016-061345
Ding H, Smith RG, Poleg-Polsky A, Diamond JS, Briggman KL. 2016. Species-specific wiring for
direction selectivity in the mammalian retina. *Nature* **535**:105–110.
doi:10.1038/nature18609
Dowling JE, Boycott BB. 1966. Organization of the primate retina: electron microscopy. *Proc R*
Soc Lond B Biol Sci **166**:80–111. doi:10.1098/rspb.1966.0086
Duan X, Krishnaswamy A, De la Huerta I, Sanes JR. 2014. Type II cadherins guide assembly of a
direction-selective retinal circuit. *Cell* **158**:793–807. doi:10.1016/j.cell.2014.06.047
Euler T, Haverkamp S, Schubert T, Baden T. 2014. Retinal bipolar cells: elementary building
blocks of vision. *Nat Rev Neurosci* **15**:507–519. doi:10.1038/nrn3783
Euler T, Masland RH. 2000. Light-evoked responses of bipolar cells in a mammalian retina. *J*
Neurophysiol **83**:1817–1829. doi:10.1152/jn.2000.83.4.1817
Farrow K, Teixeira M, Szikra T, Viney TJ, Balint K, Yonehara K, Roska B. 2013. Ambient
illumination toggles a neuronal circuit switch in the retina and visual perception at cone
threshold. *Neuron* **78**:325–338. doi:10.1016/j.neuron.2013.02.014
Franke K, Berens P, Schubert T, Bethge M, Euler T, Baden T. 2017. Inhibition decorrelates visual
feature representations in the inner retina. *Nature* **542**:439–444.
doi:10.1038/nature21394
Ghosh KK, Bujan S, Haverkamp S, Feigenspan A, Wässle H. 2004. Types of bipolar cells in the
mouse retina. *J Comp Neurol* **469**:70–82. doi:10.1002/cne.10985
Goetz J, Jessen ZF, Jacobi A, Mani A, Cooler S, Greer D, Kadri S, Segal J, Shekhar K, Sanes JR,
Schwartz GW. 2022. Unified classification of mouse retinal ganglion cells using function,
morphology, and gene expression. *Cell Rep* **40**:111040.
doi:10.1016/j.celrep.2022.111040

Gollisch T, Meister M. 2010. Eye smarter than scientists believed: neural computations in
circuits of the retina. *Neuron* **65**:150–164. doi:10.1016/j.neuron.2009.12.009
Grabner CP, Futagi D, Shi J, Bindokas V, Kitano K, Schwartz EA, DeVries SH. 2023. Mechanisms
of simultaneous linear and nonlinear computations at the mammalian cone
photoreceptor synapse. *Nat Commun* **14**:3486. doi:10.1038/s41467-023-38943-2
Grabner CP, Moser T. 2021. The mammalian rod synaptic ribbon is essential for Cav channel
facilitation and ultrafast synaptic vesicle fusion. *Elife* **10**:e63844.
doi:10.7554/eLife.63844
Grabner CP, Ratliff CP, Light AC, DeVries SH. 2016. Mechanism of High-Frequency Signaling at a
Depressing Ribbon Synapse. *Neuron* **91**:133–145. doi:10.1016/j.neuron.2016.05.019
Greene MJ, Kim JS, Seung HS, EyeWirers. 2016. Analogous Convergence of Sustained and
Transient Inputs in Parallel On and Off Pathways for Retinal Motion Computation. *Cell
Rep* **14**:1892–1900. doi:10.1016/j.celrep.2016.02.001
Grimes WN, Baudin J, Azevedo AW, Rieke F. 2018. Range, routing and kinetics of rod signaling in
primate retina. *Elife* **7**:e38281. doi:10.7554/eLife.38281
Grimes WN, Schwartz GW, Rieke F. 2014. The synaptic and circuit mechanisms underlying a
change in spatial encoding in the retina. *Neuron* **82**:460–473.
doi:10.1016/j.neuron.2014.02.037
Hellmer CB, Zhou Y, Fyk-Kolodziej B, Hu Z, Ichinose T. 2016. Morphological and physiological
analysis of type-5 and other bipolar cells in the Mouse Retina. *Neuroscience* **315**:246–
258. doi:10.1016/j.neuroscience.2015.12.016
Helmstaedter M, Briggman KL, Turaga SC, Jain V, Seung HS, Denk W. 2013. Connectomic
reconstruction of the inner plexiform layer in the mouse retina. *Nature* **500**:168–174.
doi:10.1038/nature12346
Hsiang J-C, Shen N, Soto F, Kerschensteiner D. 2024. Distributed feature representations of
natural stimuli across parallel retinal pathways. *Nat Commun* **15**:1920.
doi:10.1038/s41467-024-46348-y
Huang L, Max M, Margolskee RF, Su H, Masland RH, Euler T. 2003. G protein subunit G gamma
13 is coexpressed with G alpha o, G beta 3, and G beta 4 in retinal ON bipolar cells. *J
Comp Neurol* **455**:1–10. doi:10.1002/cne.10396
Huang L, Shanker YG, Dubauskaite J, Zheng JZ, Yan W, Rosenzweig S, Spielman AI, Max M,
Margolskee RF. 1999. Ggamma13 colocalizes with gustducin in taste receptor cells and
mediates IP3 responses to bitter denatonium. *Nat Neurosci* **2**:1055–1062.
doi:10.1038/15981
Hull C, Studholme K, Yazulla S, von Gersdorff H. 2006. Diurnal changes in exocytosis and the
number of synaptic ribbons at active zones of an ON-type bipolar cell terminal. *J
Neurophysiol* **96**:2025–2033. doi:10.1152/jn.00364.2006
Ichinose T, Fyk-Kolodziej B, Cohn J. 2014. Roles of ON cone bipolar cell subtypes in temporal
coding in the mouse retina. *J Neurosci* **34**:8761–8771. doi:10.1523/JNEUROSCI.3965-
13.2014
Ichinose T, Hellmer CB. 2016. Differential signalling and glutamate receptor compositions in the
OFF bipolar cell types in the mouse retina. *J Physiol* **594**:883–894. doi:10.1113/JP271458

Ikeda H, Wright MJ. 1972. Receptive field organization of "sustained" and "transient" retinal
1139   ganglion cells which subserve different function roles. *J Physiol* **227**:769–800.
1140   doi:10.1113/jphysiol.1972.sp010058
1141   Jackman SL, Choi S-Y, Thoreson WB, Rabl K, Bartoletti TM, Kramer RH. 2009. Role of the
1142   synaptic ribbon in transmitting the cone light response. *Nat Neurosci* **12**:303–310.
1143   doi:10.1038/nn.2267
1144   Jadzinsky PD, Baccus SA. 2013. Transformation of visual signals by inhibitory interneurons in
1145   retinal circuits. *Annu Rev Neurosci* **36**:403–428. doi:10.1146/annurev-neuro-062012-
1146   170315
1147   Jarsky T, Cembrowski M, Logan SM, Kath WL, Riecke H, Demb JB, Singer JH. 2011. A Synaptic
1148   Mechanism for Retinal Adaptation to Luminance and Contrast. *J Neurosci* **31**:11003–
1149   11015. doi:10.1523/JNEUROSCI.2631-11.2011
1150   Ke J-B, Wang YV, Borghuis BG, Cembrowski MS, Riecke H, Kath WL, Demb JB, Singer JH. 2014.
1151   Adaptation to background light enables contrast coding at rod bipolar cell synapses.
1152   *Neuron* **81**:388–401. doi:10.1016/j.neuron.2013.10.054
1153   Kerschensteiner D. 2022. Feature Detection by Retinal Ganglion Cells. *Annu Rev Vis Sci* **8**:135–
1154   169. doi:10.1146/annurev-vision-100419-112009
1155   Kim KJ, Rieke F. 2001. Temporal contrast adaptation in the input and output signals of
1156   salamander retinal ganglion cells. *J Neurosci* **21**:287–299. doi:10.1523/JNEUROSCI.21-
1157   01-00287.2001
1158   Krieger B, Qiao M, Rousso DL, Sanes JR, Meister M. 2017. Four alpha ganglion cell types in
1159   mouse retina: Function, structure, and molecular signatures. *PLoS One* **12**:e0180091.
1160   doi:10.1371/journal.pone.0180091
1161   Kuo SP, Chiang P-P, Nippert AR, Newman EA. 2020. Spatial Organization and Dynamics of the
1162   Extracellular Space in the Mouse Retina. *J Neurosci* **40**:7785–7794.
1163   doi:10.1523/JNEUROSCI.1717-20.2020
1164   Kuo SP, Schwartz GW, Rieke F. 2016. Nonlinear Spatiotemporal Integration by Electrical and
1165   Chemical Synapses in the Retina. *Neuron* **90**:320–332.
1166   doi:10.1016/j.neuron.2016.03.012
1167   Lagnado L, Schmitz F. 2015. Ribbon Synapses and Visual Processing in the Retina. *Annu Rev Vis*
1168   *Sci* **1**:235–262. doi:10.1146/annurev-vision-082114-035709
1169   Marc RE, Anderson JR, Jones BW, Sigulinsky CL, Lauritzen JS. 2014. The AII amacrine cell
1170   connectome: a dense network hub. *Front Neural Circuits* **8**:104.
1171   doi:10.3389/fncir.2014.00104
1172   Margolis DJ, Gartland AJ, Euler T, Detwiler PB. 2010. Dendritic Calcium Signaling in ON and OFF
1173   Mouse Retinal Ganglion Cells. *J Neurosci* **30**:7127–7138. doi:10.1523/JNEUROSCI.5694-
1174   09.2010
1175   Masland RH. 2012. The Neuronal Organization of the Retina. *Neuron* **76**:266–280.
1176   doi:10.1016/j.neuron.2012.10.002
1177   Maxeiner S, Luo F, Tan A, Schmitz F, Südhof TC. 2016. How to make a synaptic ribbon: RIBEYE
1178   deletion abolishes ribbons in retinal synapses and disrupts neurotransmitter release.
1179   *EMBO J* **35**:1098–1114. doi:10.15252/embj.201592701
1180   McGuire BA, Stevens JK, Sterling P. 1984. Microcircuitry of bipolar cells in cat retina. *J Neurosci*
1181   **4**:2920–2938. doi:10.1523/JNEUROSCI.04-12-02920.1984

Mehta B, Snellman J, Chen S, Li W, Zenisek D. 2013. Synaptic ribbons influence the size and frequency of miniature-like evoked postsynaptic currents. *Neuron* **77**:516–527. doi:10.1016/j.neuron.2012.11.024

Morgan JL, Soto F, Wong ROL, Kerschensteiner D. 2011. Development of cell type-specific connectivity patterns of converging excitatory axons in the retina. *Neuron* **71**:1014–1021. doi:10.1016/j.neuron.2011.08.025

Neves G, Lagnado L. 1999. The kinetics of exocytosis and endocytosis in the synaptic terminal of goldfish retinal bipolar cells. *J Physiol* **515 (Pt 1)**:181–202. doi:10.1111/j.1469-7793.1999.181ad.x

Nikolaev A, Leung K-M, Odermatt B, Lagnado L. 2013. Synaptic mechanisms of adaptation and sensitization in the retina. *Nat Neurosci* **16**:934–941. doi:10.1038/nn.3408

Nirenberg S, Meister M. 1997. The light response of retinal ganglion cells is truncated by a displaced amacrine circuit. *Neuron* **18**:637–650. doi:10.1016/s0896-6273(00)80304-9

Oesch NW, Diamond JS. 2011. Ribbon synapses compute temporal contrast and encode luminance in retinal rod bipolar cells. *Nat Neurosci* **14**:1555–1561. doi:10.1038/nn.2945

Okawa H, Yu W-Q, Matti U, Schwarz K, Odermatt B, Zhong H, Tsukamoto Y, Lagnado L, Rieke F, Schmitz F, Wong ROL. 2019. Dynamic assembly of ribbon synapses and circuit maintenance in a vertebrate sensory system. *Nat Commun* **10**:2167. doi:10.1038/s41467-019-10123-1

Pangrsic T, Singer JH, Koschak A. 2018. Voltage-Gated Calcium Channels: Key Players in Sensory Coding in the Retina and the Inner Ear. *Physiol Rev* **98**:2063–2096. doi:10.1152/physrev.00030.2017

Pottackal J, Singer JH, Demb JB. 2021. Computational and Molecular Properties of Starburst Amacrine Cell Synapses Differ With Postsynaptic Cell Type. *Front Cell Neurosci* **15**:660773. doi:10.3389/fncel.2021.660773

Puthussery T, Venkataramani S, Gayet-Primo J, Smith RG, Taylor WR. 2013. NaV1.1 channels in axon initial segments of bipolar cells augment input to magnocellular visual pathways in the primate retina. *J Neurosci* **33**:16045–16059. doi:10.1523/JNEUROSCI.1249-13.2013

Rabl K, Cadetti L, Thoreson WB. 2006. Paired-pulse depression at photoreceptor synapses. *J Neurosci* **26**:2555–2563. doi:10.1523/JNEUROSCI.3667-05.2006

Ran Y, Huang Z, Baden T, Schubert T, Baayen H, Berens P, Franke K, Euler T. 2020. Type-specific dendritic integration in mouse retinal ganglion cells. *Nat Commun* **11**:2101. doi:10.1038/s41467-020-15867-9

Rieke F. 2001. Temporal contrast adaptation in salamander bipolar cells. *J Neurosci* **21**:9445–9454.

Rodieck RW. 1991. The density recovery profile: a method for the analysis of points in the plane applicable to retinal studies. *Vis Neurosci* **6**:95–111. doi:10.1017/s095252380001049x

Roska B, Werblin F. 2001. Vertical interactions across ten parallel, stacked representations in the mammalian retina. *Nature* **410**:583–587. doi:10.1038/35069068

Sabbah S, Papendorp C, Koplas E, Beltoja M, Etebari C, Gunesch AN, Carrete L, Kim MT, Manoff G, Bhatia-Lin A, Zhao T, Schreck D, Dowling H, Briggman KL, Berson DM. 2018. Synaptic circuits for irradiance coding by intrinsically photosensitive retinal ganglion cells (preprint). Neuroscience. doi:10.1101/442954

Sagdullaev BT, McCall MA, Lukasiewicz PD. 2006. Presynaptic inhibition modulates spillover,
creating distinct dynamic response ranges of sensory output. *Neuron* **50**:923–935.
doi:10.1016/j.neuron.2006.05.015
Saszik S, DeVries SH. 2012. A mammalian retinal bipolar cell uses both graded changes in
membrane voltage and all-or-nothing Na+ spikes to encode light. *J Neurosci* **32**:297–307.
doi:10.1523/JNEUROSCI.2739-08.2012
Schwartz GW, Okawa H, Dunn FA, Morgan JL, Kerschensteiner D, Wong RO, Rieke F. 2012. The
spatial structure of a nonlinear receptive field. *Nat Neurosci* **15**:1572–1580.
doi:10.1038/nn.3225
Shekhar K, Lapan SW, Whitney IE, Tran NM, Macosko EZ, Kowalczyk M, Adiconis X, Levin JZ,
Nemesh J, Goldman M, McCarroll SA, Cepko CL, Regev A, Sanes JR. 2016.
Comprehensive Classification of Retinal Bipolar Neurons by Single-Cell Transcriptomics.
Cell **166**:1308-1323.e30. doi:10.1016/j.cell.2016.07.054
Sigulinsky CL, Anderson JR, Kerzner E, Rapp CN, Pfeiffer RL, Rodman TM, Emrich DP, Rapp KD,
Nelson NT, Lauritzen JS, Meyer M, Marc RE, Jones BW. 2020. Network Architecture of
Gap Junctional Coupling among Parallel Processing Channels in the Mammalian Retina. *J*
Neurosci **40**:4483–4511. doi:10.1523/JNEUROSCI.1810-19.2020
Singer JH, Diamond JS. 2006. Vesicle depletion and synaptic depression at a mammalian ribbon
synapse. *J Neurophysiol* **95**:3191–3198. doi:10.1152/jn.01309.2005
Singer JH, Diamond JS. 2003. Sustained Ca2+ entry elicits transient postsynaptic currents at a
retinal ribbon synapse. *J Neurosci* **23**:10923–10933. doi:10.1523/JNEUROSCI.23-34-
10923.2003
Slaughter MM, Miller RF. 1981. 2-amino-4-phosphonobutyric acid: a new pharmacological tool
for retina research. *Science* **211**:182–185. doi:10.1126/science.6255566
Snellman J, Mehta B, Babai N, Bartoletti TM, Akmentin W, Francis A, Matthews G, Thoreson W,
Zenisek D. 2011. Acute destruction of the synaptic ribbon reveals a role for the ribbon in
vesicle priming. *Nat Neurosci* **14**:1135–1141. doi:10.1038/nn.2870
Snellman J, Zenisek D, Nawy S. 2009. Switching between transient and sustained signalling at
the rod bipolar-AII amacrine cell synapse of the mouse retina. *J Physiol* **587**:2443–2455.
doi:10.1113/jphysiol.2008.165241
Srivastava P, de Rosenroll G, Matsumoto A, Michaels T, Turple Z, Jain V, Sethuramanujam S,
Murphy-Baum BL, Yonehara K, Awatramani GB. 2022. Spatiotemporal properties of
glutamate input support direction selectivity in the dendrites of retinal starburst
amacrine cells. *Elife* **11**:e81533. doi:10.7554/eLife.81533
Swygart D, Yu W-Q, Takeuchi S, Wong ROL, Schwartz GW. 2024. A presynaptic source drives
differing levels of surround suppression in two mouse retinal ganglion cell types. *Nat*
Commun **15**:599. doi:10.1038/s41467-024-44851-w
Szikra T, Trenholm S, Drinnenberg A, Jüttner J, Raics Z, Farrow K, Biel M, Awatramani G, Clark
DA, Sahel J-A, da Silveira RA, Roska B. 2014. Rods in daylight act as relay cells for cone-
driven horizontal cell-mediated surround inhibition. *Nat Neurosci* **17**:1728–1735.
doi:10.1038/nn.3852
Taylor WR, Vaney DI. 2002. Diverse synaptic mechanisms generate direction selectivity in the
rabbit retina. *J Neurosci* **22**:7712–7720.

Tsukamoto Y, Omi N. 2017. Classification of Mouse Retinal Bipolar Cells: Type-Specific
Connectivity with Special Reference to Rod-Driven AII Amacrine Pathways. *Front*
Neuroanat **11**:92. doi:10.3389/fnana.2017.00092
Vaney DI, Sivyer B, Taylor WR. 2012. Direction selectivity in the retina: symmetry and
asymmetry in structure and function. *Nat Rev Neurosci* **13**:194–208.
doi:10.1038/nrn3165
Veruki ML, Hartveit E. 2002a. Electrical synapses mediate signal transmission in the rod
pathway of the mammalian retina. *J Neurosci* **22**:10558–10566.
doi:10.1523/JNEUROSCI.22-24-10558.2002
Veruki ML, Hartveit E. 2002b. AII (Rod) amacrine cells form a network of electrically coupled
interneurons in the mammalian retina. *Neuron* **33**:935–946. doi:10.1016/s0896-
6273(02)00609-8
von Gersdorff H, Matthews G. 1997. Depletion and replenishment of vesicle pools at a ribbon-
type synaptic terminal. *J Neurosci* **17**:1919–1927. doi:10.1523/JNEUROSCI.17-06-
01919.1997
von Gersdorff H, Vardi E, Matthews G, Sterling P. 1996. Evidence that vesicles on the synaptic
ribbon of retinal bipolar neurons can be rapidly released. *Neuron* **16**:1221–1227.
doi:10.1016/s0896-6273(00)80148-8
Wakeham CM, Shi Q, Ren G, Haley TL, Duvoisin RM, von Gersdorff H, Morgans CW. 2023.
Trophoblast glycoprotein is required for efficient synaptic vesicle exocytosis from retinal
rod bipolar cells. *Front Cell Neurosci* **17**:1306006. doi:10.3389/fncel.2023.1306006
Wan Q-F, Heidelberger R. 2011. Synaptic release at mammalian bipolar cell terminals. *Vis*
Neurosci **28**:109–119. doi:10.1017/S0952523810000453
Warwick RA, Kaushansky N, Sarid N, Golan A, Rivlin-Etzion M. 2018. Inhomogeneous Encoding
of the Visual Field in the Mouse Retina. *Curr Biol* **28**:655-665.e3.
doi:10.1016/j.cub.2018.01.016
Wässle H, Puller C, Müller F, Haverkamp S. 2009. Cone contacts, mosaics, and territories of
bipolar cells in the mouse retina. *J Neurosci* **29**:106–117. doi:10.1523/JNEUROSCI.4442-
08.2009
Werblin FS. 1970. Response of retinal cells to moving spots: intracellular recording in Necturus
maculosus. *J Neurophysiol* **33**:342–350. doi:10.1152/jn.1970.33.3.342
Werginz P, Raghuram V, Fried SI. 2020. Tailoring of the axon initial segment shapes the
conversion of synaptic inputs into spiking output in OFF-α T retinal ganglion cells. *Sci*
Adv **6**:eabb6642. doi:10.1126/sciadv.abb6642
Wienbar S, Schwartz GW. 2022. Differences in spike generation instead of synaptic inputs
determine the feature selectivity of two retinal cell types. *Neuron* **110**:2110-2123.e4.
doi:10.1016/j.neuron.2022.04.012
Wu SM, Gao F, Maple BR. 2000. Functional architecture of synapses in the inner retina:
segregation of visual signals by stratification of bipolar cell axon terminals. *J Neurosci*
20:4462–4470. doi:10.1523/JNEUROSCI.20-12-04462.2000
Yu W-Q, Swanstrom R, Sigulinsky CL, Ahlquist RM, Knecht S, Jones BW, Berson DM, Wong RO.
2023. Distinctive synaptic structural motifs link excitatory retinal interneurons to diverse
postsynaptic partner types. *Cell Rep* **42**:112006. doi:10.1016/j.celrep.2023.112006

Zhang Y, Kim I-J, Sanes JR, Meister M. 2012. The most numerous ganglion cell type of the
mouse retina is a selective feature detector. *Proc Natl Acad Sci U S A* **109**:E2391-2398.
doi:10.1073/pnas.1211547109
1314
1315

1316　　***Supplemental Information***
1317
1318

1319
1320
Figure S1. *Noise-evoked responses are biased towards large amplitude events in ON-T*
compared to ON-S RGCs. (A) Spike event peak spike rate cumulative distributions for ON-T
(cyan; n= 6 cells; 131 spike events) and ON-S RGC (magenta; n= 7 cells; 300 spike events).
Firing rates from average responses to a repeated noise stimulus (see Fig. 1B-C) were
normalized to the maximal firing rate in each cell before combining events across cells. (B)
EPSC amplitude cumulative distributions for ON-T (n= 13 cells; 275 EPSCs) and ON-S (n= 10
cells; 365 EPSCs) RGCs. Same stimulus and normalization procedure as in (A).
1328

1329
1330

1331
1332
Figure S2. *Linear-nonlinear models of excitatory synaptic input to ON-T and ON-S RGCs. (A)*
Linear filters computed from responses to non-repeated noise stimulus presentation. Lines
show filters from individual ON-T (cyan; n= 13) and ON-S (magenta; n= 15) RGCs. Filters were
normalized to the peak amplitude of the primary inward component to facilitate visual
comparison of temporal responses. Same cells as shown in average filters in Fig. 2B. (B)
Nonlinearities derived by plotting the predicted response (convolution of linear filter and
stimulus) against the measured excitatory currents for each RGC shown in (A). Linear
predictions were calculated using non-normalized filters.
1341

1342
Figure S3. *Inhibitory synaptic input to ON-T and ON-S RGCs. (A) Inhibitory synaptic currents*
measured at V_{hold} = +10 mV (reversal potential for excitation) in example ON-T (cyan) and ON-S
(pink) RGCs in response to same Gaussian noise stimulus as in Figs. 1-2. Traces show
average currents from ten repetitions of the same stimulus and are baseline subtracted to
facilitate comparison. (B) Linear filters computed from responses to non-repeated noise stimulus
presentation. Lines and shaded areas show mean ± SEM of responses from ON-T (n= 8) and
ON-S (n= 8) RGCs. (C) Quantification of filter kinetics in different RGCs. Zero-cross times were
not significantly different (p= 0.57). Significance values from Wilcoxon rank sum test. (D) Peak
amplitudes of excitatory and inhibitory conductances (G_{exc} and G_{inh}, respectively) calculated
from average current responses to repeated noise stimuli such as shown in Figs. 2A and S3A.
Lines connect peak G_{exc} and G_{inh} values measured from the same RGCs. Circles with error bars
show mean ± SEM across ON-T RGCs (n= 6) and ON-S RGCs (n= 7) (E) Summary of peak
excitatory/inhibitory conductance ratios. Open circles show measurements from individual cells
and filled circles with error bars show mean ± SEM values. No significant difference between
ON-T and ON-S RGCs; p= 0.10, Wilcoxon rank sum test.
1358

1359

1360
1361
Figure S4. *Characterization of Gjd2-EGFP+ bipolar cells. (A) Vertical section of the Gjd2-EGFP*
mouse line showing EGFP positive bipolar cells axons (green) stratifying in the ON sublamina of
the IPL, co-labeled with cholinergic amacrine cell (magenta, ChAT antibody). Transmitted light
image of the retina in blue. (B) Vertical section of the Gjd2-EGFP mouse line showing EGFP
positive bipolar cells (green) colocalized with the T5 marker CaBP5 (magenta). (C) En-face
image of a whole mount retina from the GJD2 mouse line showing EGFP positive bipolar cells
bodies (green) co-labeled with the T5 bipolar cell marker CaBP5 (magenta). (D) Same field as
in (C) with cell bodies of EGFP positive bipolar cells outlined (green) by application of a digital
edge outlining filter in ImageJ. (E) Position of cell body centroids for EGFP positive bipolar cells
in a larger field of view (20X objective) outlined by green circles. (F) Density recovery profile
analysis showing the density profile as a function of distance from the reference neuron for
EGFP positive bipolar cells (green histograms) and for an equal amount of neurons randomly
positioned in the same space (grey histograms) showing the presence of a characteristic
exclusion distance for EGFP positive bipolar cells, hallmark of a retinal mosaic.

1376
1377
Figure S5. *ON-T and ON-S RGC dendrites ramify in distinct IPL sublaminae. (A) Side*
projections from 2P z-stack images of dye-filled RGCs (ON-T: top, cyan; ON-S: bottom,
magenta) and EGFP signal (yellow) in retinas from Gjd2-EGFP mice. RGCs are same cells as
shown in en face view in Fig. 1A (B) Quantification of fluorescence signals from dye filled RGCs
in Gjd2-EGFP mouse retina as in (A). Colored lines and shaded areas show mean ± SEM
fluorescence intensity with respect to distance from peak EGFP (yellow) fluorescence intensity
(n= 6 ON-T, cyan and n= 4 ON-S, magenta, RGCs).
1385

1386

1387
1388
Figure S6. *Gjd2-EGFP+ type 5 bipolar cells provide ~40% of total synapses onto ON-T RGCs.*
(A) An example ON-T RGC biolistically labeled by Cerulean *Fluorescent Protein (blue) and*
PSD95-mCherry (green) in retina from a Gjd2-EGFP (red) transgenic mouse. Top: en face view,
Bottom: side views. (B) Top: Dot map of PSD95-mCherry apposed (magenta) or not apposed
(green) to GFP+ BC axon terminals. Bottom: Magnified view of the raw images and the dot map
for a segment of dendritic arbor in the while box. (C) Quantification of total excitatory synapse
density across the dendritic arbor of ON-T RGCs. (D) Quantification of excitatory synapses in
ON-T dendrites that are adjacent to GFP+ axon terminals.
1397

1398

1399
1400
Figure S7. *Classification of three subtypes of type 5 bipolar cells. (A) Wholemount views of the*
axon terminals of type 5i, type 5t and type 5o bipolar cells in the SBFSEM volume in Fig. 3. The
tiling of the space validates the categorization of the three subtypes of type 5 bipolar cells. (B)
Identification of Gjd2-EGFP+ bipolar cells as type 5i bipolar cells, supported by correlated light
(left) and electron (right) microscopy images. The reconstructed skeleton of the type 5i bipolar
cell in (B) is delineated by dashed lines in (A).
1407
1408

1409
Figure S8. *Linear-nonlinear models of bipolar cell voltage responses. (A) Individual linear filters*
for all type 5i (cyan; n= 11), type 6 (magenta; n= 7), and type 7 (green; n=6) bipolar cells
computed from responses to non-repeated noise stimulus presentation. Filters were normalized
to the peak amplitude of the primary inward component to facilitate visual comparison of
temporal responses. Same cells as shown in average filters in Fig. 4B. (B) Nonlinearities
derived by plotting the predicted response (convolution of linear filter and stimulus) against the
measured voltage response for each bipolar cell shown in (A). Linear predictions were
calculated using non-normalized filters.
1418

1419
1420
Figure S9*. Presynaptic inhibition does not affect response kinetics but does affect spatial*
integration for excitatory inputs to ON-T RGCs (A) Step-evoked (100% contrast, 52   0 μm spot
diameter, 500 ms duration) excitatory synaptic currents in example ON-T (cyan) and ON-S
RGCs measured in ACSF. Traces show mean responses to ten trials and are baseline
subtracted to facilitate comparison. (B) Summary of light step-evoked steady-state response
amplitude for different stimulus sizes measured in ACSF. Note that excitatory inputs in ON-T are
more transient than those in ON-S RGCs across all stimulus sizes, but responses in both RGC
types are more sustained than when measurements were performed using Ames' solution
(compare to Fig. 2D-E). (C) Summary of total response, measured as the integrated excitatory
current response during the light step, versus stimulus spot size. Responses were normalized to
the maximal response in each cell prior to averaging. Excitation in ON-T RGCs is tuned to
smaller spot sizes than in ON-S RGCs and this tuning relies in part on GABAergic inhibition.
Symbols with error bars in (B) and (C) show mean ± SEM (ON-T control n = 12; ON-S control n
= 6; ON-T GABA-R block n = 5; ON-T Gly-R block n = 5).

1435
Figure S10. Bipolar cell subtype-specific differences in ribbon surface areas. (A) Violin plot
showing distribution of ribbon surface areas (each data point is a ribbon) for the type 5i and type
6 BC in (Fig 9A). (B) Violin plot with scattered dots showing ribbon surface areas for each ribbon
presynaptic to a stretch of reconstructed ON-T or ON-S RGC dendrite. (C) Scatter plot showing
ribbon size in surface area in relationship with the number of adjacent vesicles. Gray line
represents the linear regression model fitted to the data. Linear regression: $\beta = 0.0003$, $p <$
0.001, $r^2 = 0.69$.
1443