# Mucin-2 ER-to-Golgi transport mechanism identifies source of ER stress
in inflammatory bowel disease

Margaretha A.J. Morsink[1,†,#], Lena S. Koch[1,†], Shixian Hu[2,3], Rinse K. Weersma[2], Harry van
Goor[4], Arno R. Bourgonje[2,+], Kerensa Broersen[1,*]

[1] Department of Applied Stem Cell Technologies, TechMed Center, University of Twente,
Enschede, The Netherlands

[2] Department of Gastroenterology and Hepatology, University of Groningen, University
Medical Center Groningen, Groningen, the Netherlands

[3] Institute of Precision Medicine, The First Affiliated Hospital, Sun Yat-Sen University,
Guangzhou, Guangdong, China

[4] Department of Pathology and Medical Biology, University of Groningen, University Medical
Center Groningen, Groningen, the Netherlands

[†] These authors share first authorship.

[#] Current affiliation: Laboratory for Stem Cells and Tissue Engineering, Columbia University,
New York, NY, United States of America

[+] Current affiliation: The Henry D. Janowitz Division of Gastroenterology, Department of
Medicine, Icahn School of Medicine at Mount Sinai, New York, NY, United States of America

[*] Corresponding author Kerensa Broersen

**Email:** k.broersen@utwente.nl

**Running Title:** MUCIN-2 ER-TO-GOLGI TRANSPORT

**Keywords:** mucin-2/ COPII-enlargement/ inflammatory bowel disease/ ER-to-Golgi
trafficking/ TGF-β

**Acknowledgments**
We thank Dr. Michael Rape (University of California, Berkeley, CA, USA) for kindly gifting the KLHL12-FLAG in pcDNA5 construct and Dr. Vivek Malhotra (CRG-Centre de Regulacio Genomica, Barcelona, Spain) for kindly providing the TANGO1-FLAG plasmid in pcDNA3.1.

**Disclosure and competing Interest Statement:** RKW acted as consultant for Takeda, received unrestricted research grants from Takeda, Johnson & Johnson, Tramedico and Ferring and received speaker fees from MSD, AbbVie, and Janssen Pharmaceuticals. ARB received speaker fees from AbbVie. KB has a part-time affiliation at NIZO food research, Ede, the Netherlands. All other authors have no competing interests to declare.

**Author contributions:**

Margaretha A.J. Morsink – Conceptualization; Data curation; Formal analysis; Validation; Investigation; Visualization; Methodology; Carried out the research and collected and analyzed the data; Writing—original draft;

Lena S. Koch – Conceptualization; Resources; Data curation; Validation; Visualization; Investigation; Formal analysis; Methodology; Supervision; Writing – review and editing; Writing – final draft

Shixian Hu – Clinical validation: Conceptualization; Data curation; Formal analysis; Validation; Investigation; Visualization; Methodology;

Rinse K. Weersma – Clinical validation: Conceptualization; Resources; Supervision; Funding acquisition; Data curation; Project administration;

Harry van Goor – Clinical validation: Clinical validation: Conceptualization; Data curation; Formal analysis; Validation; Investigation; Visualization; Methodology;

Arno R. Bourgonje – Clinical validation: Clinical validation: Conceptualization; Data curation; Formal analysis; Validation; Investigation; Visualization; Methodology;

Kerensa Broersen - Conceptualization; Resources; Supervision; Funding acquisition; Data curation; Project administration; Writing – original draft; Writing—review and editing

**Funding**: This work was supported by the Netherlands Organ-on-Chip Initiative, an NWO Gravitation project (024.003.001) and an NWO Vici project (19424), both funded by the Ministry of Education, Culture and Science of the government of the Netherlands. RKW is supported by the Seerave Foundation and the Netherlands Organization for Scientific Research. ARB is supported by a Rubicon fellowship from NWO (452022317). The funders had no role in study design, data collection and analysis, preparation of the manuscript or decision to publish.

# The Paper Explained

## Problem

Inflammatory bowel disease (IBD) is characterized by deficits in the mucous layer covering
our intestines. While the secretion process of mucin-2, the protein that makes up the mucous
layer, is largely well-documented, transport between the ER and the Golgi apparatus
remains enigmatic and poses additional challenges due to the size discrepancy of mucin-2
and regular ER-to-Golgi transport vesicles. We hypothesize that defective mucin-2 ER-to-
Golgi transport may contribute to the observed ER stress observed in IBD patients.

## Results

We discovered that mucin-2 is transported from ER-to-Golgi involving coat protein complex II
(COPII) vesicles facilitated by proteins such as TANGO1 and cTAGE5. KLHL12 promotes
the vesicle enlargement needed to accommodate the large mucin-2 protein. We found that
TANGO1 expression is regulated by TGF-β. Aberrant TGF-β levels are observed in IBD. Our
findings imply that the impaired formation of COPII vesicles and the reduced secretion of
mucin-2 contribute to the increased ER stress seen in patients suffering from IBD. An
analysis of the gene expression in colonic tissues of IBD patients has provided further
confirmation of elevated ER stress, along with alterations in the expression levels of TGF-β,
mucin-2 and various proteins involved in COPII-mediated transport.

## Impact

Our study is the first to describe the intracellular ER-to-Golgi trafficking of mucin-2, while
elucidating ER stress origins in IBD by describing the relationship between COPII-
enlargement, TGF-β, and mucin-2. This mechanism has clinical implications for IBD and can
thereby provide different treatment strategies.

27
28

**Abstract**

The intestinal mucous layer relies on mucin-2 secretion. While the mucin-2 secretory
pathway has been studied, endoplasmic reticulum (ER) to Golgi apparatus (Golgi) trafficking
remains poorly understood. The size of mucin-2 exceeds the capacity of regular coat protein
complex II (COPII) vesicles, responsible for ER-to-Golgi transport. After confirming
conventional secretion of mucin-2, we showed that COPII vesicle enlargement is facilitated
by TANGO1 and cTAGE5, and promoted by KLHL12. Inflammatory bowel disease (IBD) is
characterized by a compromised mucous layer, altered activity of Transforming Growth
Factor β (TGF-β), and increased ER stress. Using a cell culture, we showed that TGF-β
inhibition induces TANGO1-mediated ER stress. Mucosal gene expression analysis in IBD
patients confirmed elevated ER stress and validated concomitantly altered mRNA levels of
TGF-β with mucin-2 and transport proteins TANGO1 and cTAGE5. In conclusion, we
propose that the unsuccessful formation of enlarged COPII vesicles could be a source of ER
stress in IBD, because of lowered TANGO1 protein expression, subsequently leading to
decreased mucin-2 secretion.

1 **Introduction**
2
The mucous membrane covers the entire epithelial lining of the human body facing the
exterior, including the respiratory, digestive, and urinary tracts (Medicine & Virus, 1999). In
the intestines, mucin-2 is the predominant mucous protein, which is continuously secreted by
intestinal goblet cells (Tytgat *et al*, 1996). Besides acting as the first line of defense against
invading bacteria and microorganisms, the mucous layer provides a nutritional adherence
site for commensal gut bacteria (Arike & Hansson, 2016). Moreover, the mucous layer
prevents dehydration, damage from gastric juices, and mechanical damage due to the
peristaltic movement of the intestines (Johansson *et al*, 2008; Linden *et al*, 2008).
Impairment of this layer has detrimental effects, as can be observed in patients suffering
from inflammatory bowel disease (IBD) (Baumgart & Carding, 2007; Podolsky, 1991), and
has been linked to possible defects in post-transcriptional processing and secretion of
mucin-2 (Larsson *et al*, 2011). After post-translational dimerization and N-glycosylation
within the endoplasmic reticulum (ER) (Garbes *et al*, 2015; Johansson *et al*, 2013; Lidell *et*
*al*, 2003; Rodriguez-Pineiro *et al*, 2013), mucin-2 is extensively O-glycosylated in the Golgi
apparatus. It is then oligomerized and subsequently packaged into goblet cell-residing
secretory granules (Ambort *et al,* 2012a; Forstner, 1995). Following secretion, mucin-2
absorbs water to form a gel-like sheet with a protective function. The formation of secretory
granules and post-Golgi transport mechanisms of mucin-2 has been well described (Ambort
*et al*, 2012b; Birchenough *et al*, 2015; Perez-Vilar, 2008; Perez-Vilar, 2007). However, the
ER-to-Golgi transport mechanism of mucin-2 remains largely unknown.
23
Transport of proteins out of the ER can be described by a conventional pathway (reviewed
by (Lee *et al*, 2004)) which enables ER-to-Golgi transport, or through unconventional
secretion pathways (reviewed by (Rabouille, 2017)) where proteins bypass the Golgi
altogether. Conventional protein secretion requires a 60-80 nm large coat protein complex 2
(COPII)-coated vesicle for transport, which consists of inner COPII proteins SEC23 and
SEC24, and outer COPII proteins SEC13 and SEC31 (reviewed by (Jensen & Schekman,
2011)). Brefeldin A (BFA), which inhibits COPII-mediated ER-to-Golgi transport (Fujiwara *et*
*al*, 1988), and Monensin, which inhibits trans-Golgi transport (Mollenhauer *et al*, 1990), are
the most widely used compounds to inhibit conventional protein secretion (Nylander &
Kalies, 1999). Early research has shown inhibited secretion of mucin-2 upon the addition of
BFA and Monensin, indicating mucin-2 is conventionally secreted involving COPII vesicle
transport from ER to Golgi (Dekker & Strous, 1990; van Klinken *et al*, 1998). However,
mucin-2 exhibits a hydrodynamic radius of 176 nm upon exiting the ER, exceeding the
dimensions of regular COPII vesicles (Perez-Vilar, 2008; Perez-Vilar, 2007). Therefore,
additional mechanisms must be involved to accommodate the large cargo in the vesicles.
39
Pro-collagen I (PC1), an essential extracellular matrix protein, is a 300-400 nm fibril upon
exiting the ER. Its COPII-mediated ER-to-Golgi transport has been well examined by various
researchers. Electron microscopy revealed the formation of enlarged COPII vesicles around
the PC1 fibril (Gorur *et al*, 2017) and various proteins facilitating this mechanism have been
established. Cullin-3 ubiquitin ligase complex containing Kelch like protein 12 (Cul3-KLHL12)
was identified as a key contributor for the enlarged COPII vesicle for PC1 trafficking (Jin *et*
*al*, 2012). It mono-ubiquitylates outer COPII protein SEC31, which drives the formation of an
enlarged COPII vesicle. In addition, overexpression of *KLHL12* induced accelerated
secretion of PC1 (Gorur *et al.*, 2017; Yuan *et al*, 2017; Yuan *et al*, 2018). Moreover,
transport and Golgi organization protein 1 (TANGO1) has been shown to enable transport of
bulky cargo (Saito *et al*, 2009). TANGO1 interacts with cutaneous T-cell lymphoma-
associated antigen 5 (cTAGE5) at ER exit sites (ERES) mediating the formation of enlarged
COPII (Saito *et al*, 2011). The complementary role of TANGO1 and cTAGE5 has been
described to facilitate transport of PC1, albeit the exact mechanisms are incompletely
understood (Gorur *et al.*, 2017; Raote *et al*, 2018; Yuan *et al.*, 2018). Mucin variant MUC26B
is trafficked through a TANGO1-mediated enlarged COPII structure in *Drosophila* (Reynolds

*et al*, 2019). Therefore, we hypothesized a similar mechanism may enable the COPII-
dependent transport of mucin-2. We further hypothesized that pathologies characterized by
mucus layer deficiencies such as IBD, encompassing ulcerative colitis (UC) and Crohn's
disease (CD), are associated with differential regulation of ER-to-Golgi trafficking of mucin-2.
This could in turn account for increased ER stress levels observed during IBD (Kaser &
Blumberg, 2010).
7
In this study, we reveal that mucin-2 ER-to-Golgi trafficking follows a process analogous to
PC1, occurring through COPII vesicles and facilitated by TANGO1 and cTAGE5.
Additionally, KLHL12 shows a promotive function in expanding COPII vesicles, aiding in the
trafficking of the bulky mucin-2 from the ER to the Golgi apparatus. Furthermore, we identify
a regulatory role of TGF-β for TANGO1. This discovery suggests that impaired COPII vesicle
formation and deficient mucin-2 secretion may contribute to the elevated ER stress observed
in IBD, a condition also characterized by altered levels of TGF-β. Analysis of mucosal gene
expression of colonic tissue from IBD patients further confirmed high ER stress associated
with altered expression levels of TGF-β1, mucin-2 and COPII transport proteins.
17
18
19

1  **Results**
2
*Mucin-2 is Transported in Large COPII Vesicles Facilitated by TANGO1 and*
*cTAGE5, and Promoted by KLHL12*
To confirm previously published data suggesting COPII involvement in mucin-2 trafficking
(Dekker & Strous, 1990; van Klinken *et al.*, 1998), we studied post-Golgi mucin-2 levels in
the intestinal goblet cell representing cell line HT29-MTX. Exposure to Brefeldin A (BFA),
and Monensin both reduced post-Golgi mucin-2 levels, without affecting cell viability
**(Supplementary Figure 1, Supplementary Figure 2 A-E)**, with BFA reducing post-Golgi
mucin-2 levels in a dose-dependent manner. Additional observations, such as the partial
localization of mucin-2 with the outer COPII coat protein SEC31A and the decrease in post-
Golgi mucin-2 upon silencing of SEC31A **(Supplementary Figure 3, Supplementary**
**Figure 2 F-H)**, further provided support for the involvement of COPII in ER-to-Golgi mucin-2
transport.

Nevertheless, it remains enigmatic how the large dimensions of mucin-2 (diameter ~350 nm)
are accommodated by the limited size of COPII vesicles (60-80 nm). Previously, large PC1
was shown to be trafficked to the Golgi in enlarged COPII vesicles facilitated by a set of
identified proteins (Gorur *et al.*, 2017). We aimed to establish whether mucin-2 ER-to-Golgi
trafficking involved a similar enlargement mechanism. After analyzing the size of the
SEC31A punctae **(Supplementary Figure 2H)**, we determined the degree of colocalization
of SEC31A punctae larger than 300 nm with mucin-2. Approximately 80% (27 out of 34
punctae) colocalized suggesting trafficking of mucin-2 in large COPII vesicles analogous to
PC1.

Previously, TANGO1, cTAGE5, and KLHL12 were shown to facilitate the enlargement of
COPII vesicles (**Figure 1A**) (Gorur *et al.*, 2017; Jin *et al.*, 2012; Raote *et al.*, 2018) while
silencing of these chaperones resulted in decreased secretion of PC1 (Maiers *et al*, 2017).
To investigate the involvement of these proteins for mucin-2 ER-to-Golgi trafficking,
TANGO1, cTAGE5, and KLHL12 were targeted for silencing using siRNA (**Supplementary**
**Figure 4**) and levels of post-Golgi mucin-2 and SEC31A protein expression were measured
using Western Blot. A significant decrease in post-Golgi mucin-2 levels was observed
following the decrease in expression of the COPII-enlarging proteins, indicating their
involvement in mucin-2 trafficking (**Figure 1B-C**) matching previous reports for PC1 (Maiers
*et al.*, 2017). Moreover, a decrease in the level of expressed SEC31A was observed by
silencing TANGO1 and KLHL12, but not for cTAGE5 (**Figure 1D-E**), consistent with
published data (Saito *et al.*, 2011). Given the requirement of TANGO1 and KLHL12 for
mucin-2 trafficking, we hypothesized that overexpression of TANGO1 and KLHL12, using a
TANGO1-FLAG and KLHL12-FLAG plasmid (**Supplementary Figure 5**), would result in an
increased level of post-Golgi mucin-2. While overexpression of KLHL12 indeed induced an
increase in post-Golgi mucin-2 levels, interestingly, TANGO1 overexpression had no effect
(**Figure 1F-G**). The combination of TANGO1 and KLHL12 overexpression contributed to an
increased post-Golgi mucin-2 level, although not significantly (p=0.501). The involvement of
TANGO1 and KLHL12 in mucin-2 trafficking was further verified using immunofluorescence.
Partial colocalization of TANGO1 and mucin-2 was observed (**Figure 1H**). Overexpression
of KLHL12-FLAG and its detection by means of a FLAG-antibody showed colocalization with
mucin-2 as well (**Figure 1I**). These results were analogous to immunofluorescence results
obtained for PC1 (Gorur *et al.*, 2017), suggesting trafficking in large COPII vesicles.
Altogether, these results suggest that mucin-2 is trafficked in enlarged COPII vesicles
facilitated by TANGO1 and cTAGE5, and promoted by KLHL12.

*TGF-β regulates TANGO1 protein expression*
Recently, TGF-β was reported to upregulate TANGO1 protein expression to, in turn,
facilitate PC1 secretion (Maiers *et al.*, 2017). As TANGO1 was found to facilitate trafficking
to the Golgi for mucin-2, we predicted that exposure of HT29-MTX cells to TGF-β would

mediate TANGO1-facilitated mucin-2 trafficking. We first investigated the effect of TGF-β1,
as the most abundant isoform of the TGF-β family (Kubiczkova *et al*, 2012), on TANGO1
expression levels using Western Blot. In line with our expectations, TGF-β1 induced an
upregulation of TANGO1 protein expression (**Figure 2B-C**) without affecting TANGO1
mRNA levels (**Figure 2N**), similar to results reported by Maiers and colleagues (Maiers *et*
*al.*, 2017). Next, we sought if mucin-2 trafficking would be facilitated in response to TGF-β1-
mediated TANGO1 upregulation. However, contradicting our expectations, we observed a
significant decrease in mucin-2 post-Golgi levels (**Figure 2D-E**). *MUC2* transcription has
been shown to converge with inflammatory pathways (Van Seuningen *et al*, 2001). The
transcription factor NF-kB, a key regulator in inflammatory gene expression (Verstrepen *et*
*al*, 2008), is known to also play a positive role in *MUC2* transcription. IL-1β has been shown
to induce NF-kB activity (Scholz *et al*, 2013) while TGF-β1 is known as an inhibitor
(Baumgart & Carding, 2007; Sedda *et al*, 2015). In light of this, it is possible that mucin-2
protein levels were inadvertently decreased as a function of TGF-β1 exposure (Ahn *et al*,
2005). Using RT-qPCR we confirmed that *MUC2* mRNA levels were indeed severely
downregulated by TGF-β1 (**Figure 2J**). We anticipated that these decreased *MUC2* mRNA
levels upon exposure to TGF-β1 would be the result of TGF-β1-induced NF-kB
downregulation. However, NF-κB levels were not altered by TGF-β1 (**Figure 3L**). Therefore,
our attempts to rescue *MUC2* transcription using IL-1β (**Figure 2A, 2J**) were idle. Due to the
low mRNA levels of *MUC2* following TGF-β1 stimulation, we anticipate reduced trafficking of
mucin-2 from ER-to-Golgi, given there was less cargo to transport.

As our experiments indicated that TGF-β1 plays a role in mucin-2 trafficking, we decided to
inhibit TGF-β1 using the small molecules Galunisertib, SB431542, and their combination.
Galunisertib is a small molecule inhibitor of the TGF-β1 receptor I, specifically
downregulating the phosphorylation of Smad2, inhibiting activation of the canonical pathway
of TGF-β (Holmgaard *et al*, 2018). SB431542 also inhibits TGF-β1 receptor I, but, instead,
by blocking phosphorylation of ALK4, ALK5, and ALK7 (activin receptor-like kinase)
receptors (Chambers *et al*, 2009; Halder *et al*, 2005). In line with our expectations, TANGO1
protein expression was decreased significantly by inhibition of TGF-β1 (**Figure 2F-G**). In a
response to the decreased TANGO1 protein levels, post-Golgi mucin-2 levels were also
attenuated as measured by Western Blot (**Figure 2H-I**). This suggests the facilitating role of
TGF-β1 in mucin-2 trafficking. However, we observed a decrease in *MUC2* mRNA levels as
well (**Figure 2K**), unaffected by NF-κB mRNA levels (**Figure 2M**). Although *MUC2* mRNA
levels were downregulated by Galunisertib, the post-Golgi mucin-2 levels were not,
demonstrating low MUC2 mRNA levels still facilitate mucin-2 translation sufficient to study
mucin-2 trafficking.

Due to the discrepancy between observed TANGO1 mRNA and protein levels in response to
TGF-β1 stimulation, we investigated the TANGO1 mRNA levels in response to TGF-β1
inhibition (**Figure 2O**). We noted an upregulation of TANGO1 mRNA by inhibiting TGF-β1,
directly contrasting the observed TANGO1 protein levels in response to TGF-β1 inhibition.
The discrepancy between TANGO1 mRNA and protein levels suggests TANGO1 expression
is mediated by post-transcriptional regulation (Maeda *et al*, 2020; Zhang *et al*, 2014).
Collectively, TGF-β1 plays an undeniable role in TANGO1 protein expression, which has
been shown to facilitate mucin-2 secretion. However, due to the intricate role of TGF-β1 in
various signaling pathways and cellular processes, we were not able to unambiguously
report the effect on mucin-2 trafficking. The lack of TANGO1, as a result of TGF-β1
inhibition, severely impeded mucin-2 secretion, thereby suggesting the facilitating role of
TGF-β1 in mucin-2 secretion.

*Inhibition of TGF-β leads to IBD-Associated ER Stress*
Given the facilitating role of TGF-β1 in large COPII formation for mucin-2 trafficking, we
hypothesized that the inhibition of TGF-β1 could be responsible for ER stress in goblet cells
as has been observed in mucus-deficiency pathologies like IBD (Kaser & Blumberg, 2010),

due to compromised generation of large COPII vesicles. We examined several different
genes in the Unfolded Protein Response (UPR) pathways through RT-qPCR (**Figure 3A**),
previously described as a proxy for measuring ER stress (Oslowski & Urano, 2011).
Thapsigargin stimulation served as a positive control (Lindner *et al*, 2020). We showed
elevated mRNA levels of spliced X-box Protein 1 (sXBP1), C/EBP-homologous protein
(CHOP), immunoglobin binding protein (BiP), Activating Transcription Factor 4 (ATF4), and
ER-degradation-enhancing-α-mannidose-like protein (EDEM) upon Thapsigargin stimulation
(**Supplementary Figure 7**). To determine whether ER stress could be caused by the
unsuccessful generation of large COPII vesicles, we silenced COPII protein SEC31A and
COPII enlarging proteins TANGO1, KLHL12, and cTAGE5. Indeed, we observed increased
ER stress levels upon inhibiting large COPII formation (**Figure 3B**), in line with published
reports (Bisnett *et al*, 2021; Fang *et al*, 2015; Liu *et al*, 2019). Moreover, overexpression of
TANGO1 and KLHL12, which facilitates trafficking in large COPII vesicles, did not elevate
ER stress levels (**Figure 3C**). Collectively, these results elucidate the potential importance of
the generation of large COPII vesicles to preclude ER stress.

Next, we investigated the effects of TGF-β1 and its inhibition on ER stress. Due to the
positive effect of TGF-β1 on TANGO1 protein levels, therefore facilitating large COPII
formation, we anticipated TGF-β1 to not induce ER stress. Indeed, no ER stress was
observed upon stimulating the HT29-MTX cells with TGF-β1 (**Figure 3D**). Concomitantly, as
the inhibition of TGF-β1 impedes large COPII formation, we predicted it would induce ER
stress. In line with this expectation, elevated mRNA levels of all UPR targets were observed
upon TGF-β1 inhibition (**Figure 3E**) consistent with previous reports (Kassan *et al*, 2012;
Okumura *et al*, 2017). These results demonstrate the importance of TGF-β1-mediated large
COPII formation to inhibit generation of ER stress.

*TANGO1 is a Target of the Unfolded Protein Response*
The observation that TANGO1 mRNA levels in response to TGF-β1 activation and inhibition
(**Figure 2**) correlate with ER stress levels, prompted us to investigate TANGO1 mRNA levels
of other ER stress situations. We hypothesized a UPR-mediated upregulation of TANGO1
mRNA levels as a result of ER stress. At the same time, we anticipated decreased TANGO1
protein levels through incomplete post-transcriptional regulation by low levels of TGF-β1. To
address this hypothesis, we first confirmed increased TANGO1 transcription upon
stimulation with the known ER stressor Thapsigargin in the goblet cell model HT29-MTX
(**Figure 3F**). Moreover, silencing of COPII (enlarging) proteins SEC31A, KLHL12 and
cTAGE5, which are known provokers of ER stress, showed a significant increase in
TANGO1 mRNA levels as well (**Figure 3G**). Interestingly, silencing of TANGO1 somehow
still upregulated TANGO1 mRNA levels (**Figure 3G**), but not protein levels (**Supplementary
Figure 4**). Collectively, these findings suggest TANGO1 transcription is a target of the UPR.
Further investigation of TANGO1 post-transcriptional regulation and clinical verification of
TANGO1 mRNA and protein levels in IBD divulge new avenues for future research.

*Intestinal tissue from patients with IBD shows elevated ER stress levels and*
*increased expression of MUC-2 trafficking genes*
To validate our findings from the *in vitro* goblet cell model HT29-MTX, we investigated
intestinal biopsy data from a cohort comprising individuals diagnosed with either UC or CD.
We studied the expression profiles of genes involved in the UPR pathway, as well as
components of the proposed ER-to-Golgi trafficking pathway in inflamed and non-inflamed
regions of both the ileum and colon. In line with our *in vitro* findings, patients with UC and CD
exhibited higher levels of ER stress compared to healthy controls with significantly increased
mRNA levels of *XBP1*, *CHOP*, *BiP*, *ATF4* and *EDEM1* (**Figure 4A-E**)**.** There was no
difference in gene expression among the two disease states, however, ER stress was shown
to be higher during active inflammation. We then evaluated the mucosal expression of
mucin-2 and genes involved in its ER-to-Golgi trafficking. Compared to healthy controls, in
patients with UC and CD, we observed a higher mRNA expression level of mucin-2 (**Figure

**5A**). This was even more prominently increased in inflamed regions. Genes involved in
mucin-2 ER-to-Golgi trafficking, such as SEC31a, TANGO1 and cTAGE5, were significantly
increased in both UC and CD with a higher degree of expression during inflammation
(**Figure 5B-D**). Against our expectations, *KLHL12* expression levels were not altered in both
IBD subtypes compared to controls (**Figure 5E**). As a major negative regulatory element of
mucosal inflammation, TGF-β1 levels (**Figure 5F**) were significantly increased in patients
with IBD. The mucosal gene expression analysis of tissues from patients with IBD confirm
that ER stress levels are increased in IBD and support our findings of differential expression
of mucin-2 and its associated trafficking components.

**Discussion**

This study reports on ER-to-Golgi trafficking of the mucin-2 protein, which is an important
physiological component of the mucus layer lining the human intestines. Our results show
that mucin-2 trafficking is COPII-dependent and is facilitated by proteins that have previously
been documented to facilitate enlargement of COPII vesicles. We further show that TGF-β1
regulates TANGO1 protein expression and thereby facilitates COPII enlargement. We
demonstrate that TGF-β1 inhibition is responsible for ER stress, in analogy to the ER stress
as has been shown in IBD. Lastly, we validate our findings with IBD patient data which
substantiate high ER stress levels and show high *MUC2* and associated COPII transport
gene expression in association with upregulated TGF-β1 mRNA expression. Collectively,
these findings indicate that …

*Mutations in COPII-Encoding Genes in Intestinal Diseases Illustrate COPII-*
*Dependency of Mucin-2*

Using BFA and Monensin, we demonstrated conventional trafficking of mucin-2 consistent
with previous reports (Dekker & Strous, 1990; Okada *et al*, 2011; van Klinken *et al.*, 1998).
Silencing of SEC31A further corroborated the COPII-dependency, which had not been
shown for mucin-2 previously. Mutations in COPII genes have been linked to various
pathophysiological defects in pro-collagen I (Boyadjiev *et al*, 2006; Garbes *et al.*, 2015).
However, no such direct links about COPII defects in mucin-2 related diseases are yet
known, however, loss of TANGO1 protein levels is found in colon carcinomas, which are, in
turn, associated with loss of mucous layer (Arndt & Bosserhoff, 2007; Zhang *et al.*, 2014).
miRNA-429 has been indicated to downregulate inner COPII protein SEC23A and is
associated with decreased mucin-2 secretion as well in a murine model of colitis (Mo *et al*,
2016), endorsing our findings of a COPII-mediated mucin-2 ER-to-Golgi transport
mechanism. Interestingly, a clinical cohort study revealed single nucleotide polymorphisms
(SNPs) in the SEC24B and SEC24B-antisense RNA 1 genes are associated with mucinous
colorectal cancer (Penney *et al*, 2018) – a carcinoma which volume consists of at least 50%
mucins (Nozoe *et al*, 2000). However, there is no evidence that proves that a defect in
SEC24B affects mucin-2 secretion. Lastly, YIPF6 has been implicated in COPII vesicle
formation, through interaction with inner COPII proteins SEC23 and SEC24 (Kranjc *et al*,
2017; Ran *et al*, 2019), particularly facilitating the secretion of antimicrobial proteins in the
intestines to maintain intestinal homeostasis. Mutations in YIPF6 cause spontaneous
intestinal inflammation in mice and associated loss of mucin-2 was observed (Brandl *et al*,
2012; Moresco & Brandl, 2013). Although the exact mechanism through which YIPF6
induces colitis remains unknown, it is suggestive of mucin-2's COPII dependency and COPII
defects leading to pathophysiological disruptions in mucin-2 secretion.

Mutations in COP-II enlarging genes were furthermore implicated in various other diseases
spanning cardiac (Buniello *et al*, 2019; Lekszas *et al*, 2020), neurological (Lemos *et al*,
2011; Zhang *et al*, 2018) and liver (Gupta & Beggs, 2014; Norman *et al*, 2015) defects,
illustrating the consequences of disrupted ER-to-Golgi trafficking of various proteins.
Therefore, future efforts should investigate the role and association of mutations in the
COPII-enlarging proteins in intestinal diseases.

*Trafficking of Mucin-2 in Enlarged COPII Vesicles is Reminiscent of Transport of*
*Other Large Proteins Including Pro-Collagen I*

TANGO1 deficiency or knockout models have been associated with defects in secretory
pathway genes and protein expression, as well as disrupted secretion of small and large
cargo proteins including but not limited to PC1 (McCaughey *et al*, 2021b). In vivo, TANGO1
knockdown in Drosophila decreased mucin-26B by disrupting COPII formation (Reynolds *et*
*al.*, 2019). TANGO1 and cTAGE5 have been implicated in transport of bulky chylomicrons in

large COPII vesicles as well, and knockout of TANGO1 led to retained apoB within the cell
(Santos *et al*, 2016). Moreover, KLHL12 was shown to colocalize with apoB and its silencing
attenuated apoB secretion, corresponding to the results obtained for mucin-2 (Butkinaree *et*
*al*, 2014). Overexpressing TANGO and KLHL12 confirmed their promoting role for mucin-2
trafficking as has been observed for PC1 (Jin *et al.*, 2012), collagen IV (Kawaguchi *et al*,
2018) and apoB (Butkinaree *et al.*, 2014) trafficking, through the mono-ubiquitylation of
SEC31.

TANGO1's facilitating role in formation of large COPII vesicles is unequivocal (McCaughey
*et al*, 2021a; Saito *et al.*, 2009), yet there are reports of its ambiguous mechanism in the
generation of large COPII vesicles. Schekman and colleagues suggest a co-packaging of
TANGO1 with the cargo (Yuan *et al*, 2018), whereas Malhotra and colleagues demonstrated
that TANGO1 does not leave the ER, but forms a ring around the COPII vesicle (Raote *et*
*al.*, 2018). The distanced colocalized punctae of TANGO1 and mucin-2 (**Figure 1B**, yellow
arrow) suggest transport of TANGO1 away from the ER, which has been validated in recent
results that show TANGO1 colocalizing with Sec16A indicating an ERES localization
(McCaughey *et al.*, 2021b). In addition, TANGO1 might play another role in the mucin-2
secretory pathway occurring in the Golgi apparatus, as it has been shown to aid in O-
glycosylation for MUC26B in *Drosophila* (Reynolds *et al.*, 2019).
19

*TGF-β Facilitates Mucin-2 Trafficking Through TANGO1 Protein Expression,*
*However, Molecular Mechanism of TGF-β on MUC2 Transcription Remains*
*Enigmatic*
TGF-β has been implicated in IBD as lower active levels are observed (Lee *et al*, 2018), due
to increased expression of TGF-β inhibitor Smad7 in the intestinal mucosa (Monteleone *et*
*al*, 2023; Troncone *et al*, 2018), prompting us to investigate the effect of TGF-β on mucin-2
trafficking. Consistent with previous research (Connolly *et al*, 2023; Maiers *et al.*, 2017),
TGF-β induced expression of TANGO1. The effect of inhibiting TGF-β on TANGO1 has not
been reported previously, however, the decreased expression is in line as an antagonistic
response to TGF-β stimulation. Decreased post-Golgi mucin-2 levels resulting from TGF-β
inhibition also endorse a facilitating role of TGF-β in TANGO1 and mucin-2 regulation. The
specific role of TGF-β in mucin regulation remains ambivalent, however. In line with our
findings on MUC2 mRNA levels, TGF-β was found to induce MUC5AC mRNA expression
(Lee *et al*, 2021). At the same time, however, TGF-β was implicated in downregulation of
MUC5AC expression (Jonckheere *et al*, 2004). Moreover, a functional cooperation of TGF-β
upregulation and NF-κB signaling was found to elevate MUC2 mRNA levels in response to
the nontypeable Haemophilus influenzae bacteria (Jono *et al*, 2002). Others suggest an
inhibitory effect of TGF-β to NF-κB (Baumgart & Carding, 2007; Sedda *et al.*, 2015), yet our
findings show no effect of TGF-β on NF-κB. IL-1β has been shown to activate transcription
factor NF-κB (Scholz *et al.*, 2013), corresponding to our results. However, MUC2
transcription did not seem altered by NF-κB, contradicting numerous findings on MUC2
transcription (Ahn *et al.*, 2005; Kim *et al*, 2002; Mejías-Luque *et al*, 2010; Mesquita *et al*,
2003; Rex *et al*, 2019; Wu *et al*, 2008; Yamamoto *et al*, 2003). Mucin-2 protein levels do not
always correlate to MUC2 mRNA levels due to extensive post-translational modifications
(Cullen, 2011), as observed by the aberration in response to Galunisertib. Ultimately, the
molecular mechanism through which MUC2 mRNA levels and mucin-2 trafficking are
decreased in response to TGF-β exposure remains unknown.

The discrepancy between TANGO1 mRNA and protein levels suggests post-transcriptional
regulation. Based on our findings related to TGF-β and its inhibition, we suspect it plays a
role in TANGO1 post-transcriptional regulation, however we lack the evidence to support this
statement. In general, the transcription and translation of TANGO1 have been poorly
investigated (Luo *et al*, 2017). We have shown TANGO1 transcription is a target of the UPR,
consistent with what is known about TANGO1 and other COPII proteins (Amodio *et al*, 2013;
Maiers *et al.*, 2017). Other post-transcriptional and post-translational modifications of

TANGO1 that have been reported including O-glycosylation (Zhang *et al.*, 2014) and
phosphorylation (Maeda *et al.*, 2020), both impacting COPII formation. Various other
phosphorylation sites of TANGO1 have been described, yet their exact mechanisms have
not been investigated (Bisnett *et al.*, 2021). Generally, post-transcriptional and post-
translational modifications play an important role in the formation of large COPII vesicles
(McCaughey & Stephens, 2018). Recently, it was shown that TANGO1 expression was
dependent on inflammasome activation of caspase-1, as well as signaling from IL-1 and
TGF-β (Connolly *et al.*, 2023). Elucidating the post-transcriptional mechanism of TANGO1,
possibly mediated by TGF-β, could identify therapeutic targets for IBD and fibrosis.
10
11

*TGF-β Inhibition Generates ER Stress in Intestinal Diseases Due to Unsuccessful*
*Formation of Large COPII Vesicles*

In addition to investigating the effects of TGF-β on mucin-2 trafficking, we examined its effect
on other hallmarks of IBD pathogenesis and pathology, namely increased levels of ER stress
in the goblet cells (Ma *et al*, 2017). There are several factors that could induce ER stress in
IBD, although the exact mechanisms are not fully understood (Sheng *et al*, 2012).
Environmental factors, such as the interactions with the microbiota, are thought to contribute
to ER stress (Kaser & Blumberg, 2010). Moreover, mutations in UPR genes, such as XBP1,
have been implicated in early onset Crohn's disease (Redhai & Boutros, 2021), and
mutations in the MUC2 gene, leading to unsuccessful O-glycosylation in the ER, have been
shown to cause IBD-related ER stress (Ma *et al.*, 2017; Ramanan & Cadwell, 2016).
Increased transcription of MUC2 has been shown to induce ER stress as well, due to the
complex folding and post-translation processing of mucin-2 in the ER (Tawiah *et al*, 2018).
Inflammatory states in the intestines induce mucin transcription and translation, thereby
exerting a sudden increase in workload of the ER, leading to ER stress (Dohrman *et al*,
1998). Therefore, proper UPR signaling is essential to maintain ER homeostasis and cell
functionality, as it attempts to restore balance. However, if protein accumulation at the ER
lumen is beyond restoration, CHOP signaling is activated and induces apoptosis (Oslowski &
Urano, 2011). Alterations in COPII genes have been shown to increase ER stress and
activate UPR pathways in many cases (Bisnett *et al.*, 2021; Fang *et al.*, 2015; Liu *et al.*,
2019), endorsing our findings. TANGO1 and KLHL12 overexpression were not shown to
induce ER stress, elucidating their importance for ER homeostasis. TGF-β, and the
subsequent increased TANGO1 protein expression, indeed did not affect ER stress levels, in
contrast to findings of Maiers and colleagues (Maiers *et al.*, 2017), however in support of our
theory that TGF-β enables mucin-2 transport. Concomitantly, TGF-β inhibition, and the
decreased TANGO1 protein levels, did activate the UPR signaling pathways, supporting our
theory.
Impaired TGF-β signaling is also found in Celiac Disease, also known as gluten intolerance,
associated with elevated ER stress levels in the intestines and an altered mucosal layer
(Benahmed *et al*, 2007; Ferrari *et al*, 2021; Schumann *et al*, 2017), further suggesting the
importance of TGF-β in maintaining ER homeostasis. Lastly, increased availability of IL-1β
has been implicated in ER stress due to aberrant mucin formation, explaining our observed
increased BiP levels (Heazlewood *et al*, 2008). Collectively, we predict the increased Smad7
signaling inhibiting TGF-β in IBD contributes to the increased ER stress, due to a lack of
TANGO1 protein availability, ultimately leading to decreased mucin-2 secretion.
47

*Patient Data Provide a Complex View on Regulation of Mucin-2 and Associated ER*
*Stress*

IBD is a complex and multifactorial disease caused by genetic, environmental, and microbial
imbalances (Kaser & Blumberg, 2010). The two major forms, ulcerative colitis (UC) and
Crohn's disease (CD) are both marked by a loss of mucus layer functionality (Moehle *et al*
2006). Corroborating our in vitro findings using the goblet cell line HT29-mtx, the mucosal
gene expression analysis of intestinal tissue from patients with UC and CD showed

significantly elevated ER stress levels in intestinal biopsies. Several studies have reported
on increased ER stress in UC and CD intestinal tissues in connection with impaired mucin
formation (Heazlewood *et al.*, 2008; Kaser *et al*, 2008; Shkoda *et al*, 2007). Our patient data
revealed significantly increased mucin-2 mRNA and its associated ER-to-Golgi trafficking
genes. Mucin-2 mRNA and protein levels during IBD have been reported to vary. This was
attributed to different origins of the disease, be it genetic or environmental, analysis methods
and even patient cohort demographics (as reviewed in (Bankole *et al*, 2021)). However,
multiple studies agreed on an increase in MUC2 mRNA in UC and CD which was
exacerbated in inflamed regions matching our findings (Ahn *et al.*, 2005; Cari *et al*, 2023;
Gersemann *et al*, 2009; Hanski *et al*, 1999; Heazlewood *et al.*, 2008). Furthermore, this
human data confirms our earlier conclusions that increased expression levels of mucin-2 are
concomitant with increased levels of SEC31A, cTAGE5 and TANGO1, signifying their
association. In opposition to our discoveries, TGF-β1 mRNA expression levels were
significantly elevated in UC and CD intestinal samples, now showing a positive relationship
between TGF-β and TANGO1 mRNA levels paralleling the TANGO1 protein regulation by
TGF-β we had established earlier. Naturally, patient data present a more holistic view on the
molecular processes during IBD compared to our data derived from the in vitro goblet cell
model HT29-mtx, also taking into account for complex immune system interactions during
chronic and acute inflammation. While we resolved the pathway of ER-to-Golgi mucin-2
trafficking in an isolated goblet cell model during health and intestinal inflammation, the in
vivo patient data shows a differential gene expression. We mimicked short term inflammation
to reveal the role of TGF-β mediated mucin trafficking. The lack of an integrated immune
component denotes one limitation of the in vitro study. Integrating LPMCs, PBMCs or
dendritic cells could help elucidate the role of TGF-β mediated TANGO1 and subsequent
mucin-2 regulation as they have been shown to be crucial for TGF- β signaling in the
intestinal mucosa in inflammatory states (Di Sabatino *et al*, 2008; Ihara *et al,* 2016;
Monteleone *et al*, 2004).

Similar to mucin-2, TGF-β expression has been shown to vary in patients with IBD with a
discrepancy of mRNA and protein levels (as reviewed in (Ihara *et al*, 2017; Naghdalipour *et*
*al*, 2022)). The increased mRNA levels could be an indicator of its regulatory role in mucosal
immunity to actively reduce inflammatory responses (Hatamzade Esfahani & Day, 2023;
Sedda *et al*., 2015). We propose that increased TGF-β1 concurrent with high MUC2 mRNA
expression and trafficking signaled by SEC31A, cTAGE5 and TANGO1 levels could reflect
efforts to start reparative processes to restore the broken mucosal barrier (Cari *et al.*, 2023;
Ihara *et al.*, 2016).
36

*Conclusion*
This study is the first to describe that mucin-2 ER-to-Golgi transport occurs in large COPII
vesicles, similar to the transport of other large proteins, such as PC1 and chylomicrons. The
enlargement of the COPII vesicles for mucin-2 transport is facilitated by cTAGE5 and
TANGO1 and promoted by Cullin-3 adaptor KLHL12. Successful generation of these large
COPII vesicles is essential to maintain ER homeostasis and cellular function. IBD, an
intestinal disease associated with loss of mucous layer and decreased TGF-β protein levels,
presents increased ER stress levels of unknown cause. We demonstrated that TGF-β
inhibition led to increased ER stress levels, as a result of decreased TANGO1 protein
expression. Interestingly, TGF-β inhibition upregulated TANGO1 mRNA levels, which we
hypothesize to be a mechanism of the UPR to ameliorate ER stress. Moreover, the
discrepancy between TANGO1 mRNA and protein levels is suggestive of post-transcriptional
regulation, possibly by TGF-β, divulging a promising avenue for future research. Mucosal
gene expression data from intestinal tissue of patients with IBD substantiated high ER stress
levels and showed high MUC2 and associated COPII transport gene expression, however, in
association with upregulated TGF-β1 mRNA expression. In conclusion, we propose that the
unsuccessful formation of large COPII vesicles, as a result of the observed TGF-β inhibition

by Smad7, could be one of the sources of ER stress in IBD, because of lowered TANGO1
protein expression, subsequently leading to decreased mucin-2 secretion.
3

**Materials and Methods**

*Cell Culture, Transfections, and Plasmids*

HT29-MTX cells were cultured in Dulbecco's Minimum Eagle Media F-12 (DMEM, Gibco,
Cat no. 11320033) containing 2 mM GlutaMAX, supplemented with 10% Fetal Bovine Serum
(FBS, Gibco, Cat no. 10270106), 100 units/mL of penicillin (Gibco, Cat no. 15140130), and
100 µg/mL of streptomycin (Gibco, Cat no. 15140130), and 1% non-essential amino acids
(NEAA, Gibco, Cat no. 11140-050). Cells were maintained at 37 ºC, 100% humidity, and 5%
$CO_2$ and were passaged twice per week using DBPS (Gibco, Cat no. 14190-094) and
Trypsin-EDTA (Gibco, Cat no. 15400054). Starvation cell culture medium contained DMEM
F-12 supplemented with 2 mM GlutaMAX, 100 units/mL of penicillin, 100 µg/mL of
streptomycin, and 1% NEAA.
Cells were seeded at 15 000 cells/cm$^2$ in a 6 wells plate (Greiner Bio-One, Cellstar, Cat no.
657 160) and cultured for 2 days. Four hours prior to transfection, the medium was changed
to starvation medium. Subsequently, silencing RNAs (siRNA, Ambion Life Technologies, 25
pmol per well) were diluted in starvation medium. DharmaFect 1 (DharmaCon, 7.5 µL per
well, Cat no. T-2001-02) was diluted in starvation medium as well and combined in a 1:1
ratio with the siRNA. The mixture was incubated for 20 minutes at room temperature before
addition to the cells. Cells were transfected for approximately 40 hours after samples were
collected for further analysis. The Silencer Pre-Designed siRNAs (Ambion) used were
SEC31A (147540), TANGO1 (UNQ6077, 124540), cTAGE5 (144132), and KLHL12
(133405). A non-targeting siRNA (ASO2CMLU) was used as negative control for the
silencing experiments.
The KLHL12-FLAG in pcDNA5 construct was a kind gift from Dr. Michael Rape (University of
California, Berkeley, CA, USA) and the TANGO1-FLAG plasmid in pcDNA3.1 was kindly
provided by Dr. Vivek Malhotra (CRG-Centre de Regulacio Genomica, Barcelona, Spain). E.
Coli bacteria containing KLHL12-FLAG plasmid were cultured on a LB Agar plate (Fisher
BioReagents, Cat no. BP1425-500) at 37 ºC and the plasmid was isolated from single
colonies using the PureYield Plasmid Midiprep kit (Promega, Cat no. A2495), according to
the manufacturer's instructions. Slbtl2 bacteria containing TANGO1-FLAG plasmid were
handled similarly yet cultured at 30 ºC. Plasmid concentrations were measured using the
ND-1000 Spectrophotometer (NanoDrop, Unity Lab Services, Thermo Fisher Scientific) and
the sequences were partially validated using Sanger Sequencing (Eurofins Genomics). Prior
to transfection, 15 000 cells/cm$^2$ were plated in a 6-wells plate and after one day transfected
using Lipofectamine 2000 (Thermofisher, Cat no. 11668030) in OptiMEM (Gibco, Cat no.
31985-070) for 40 hours. The TANGO1-FLAG plasmid was transfected at 1 µg/mL, whereas
the KLHL12-FLAG plasmid was transfected at 100 ng/mL, due to the formation of
aggregates at higher concentrations. Transfection efficiency was determined by quantifying
immunofluorescence using a FLAG antibody and DAPI counterstaining in ImageJ (v1.52p
NIH).
It should be noted that the transfection efficiency of KLHL12-FLAG was lower than
TANGO1-FLAG, due to using a ten times diluted plasmid and transfection agent
(**Supplementary Figure 5**). However, the use of 1 µg of plasmid resulted in toxic
aggregation of KLHL12, as observed by immunofluorescence of the FLAG protein
(**Supplementary Figure 6A**). TANGO, a computational tool to predict beta aggregation of
proteins, predicted beta aggregation for the KLHL12 protein (**Supplementary Figure 6B**)
(Fernandez-Escamilla *et al*, 2004; Linding *et al*, 2004; Rousseau *et al*, 2006). Therefore,
smaller amounts (0.1 µg) of KLHL12-FLAG were used to overexpress KLHL12 in HT29-MTX
cells.

*Treatments*

Brefeldin A (BFA) stock solution was prepared in DMSO at 500 µg/mL and Monensin stock
solution was prepared in 70% ethanol at 1 mM. The cells were stimulated with BFA at 1

µg/mL, 10 µg/mL, or 100 µg/mL and Monensin at 1 µM for 45 minutes. Subsequently, the
medium was refreshed to starvation medium for 24 hours after which samples were
collected. Other treatments were incubated with the cells for 24 hours in starvation medium
before sample collection. All treatments are summarized in **Table 1**.
5

*Antibodies*
Commercially available antibodies used for immunofluorescence (IF) and Western Blotting
(WB) were as follows: mouse anti-mucin-2 (Santa Cruz, sc-7314, Lot #A3019, 1:200 for IF
and 1:500 in TBST-XL for WB), rabbit anti-SEC31A (Cell Signaling Technologies, E1J5M,
Lot #0, 1:800 for IF and 1:1000 in 5% Bovine Serum Albumin (Sigma, Cat no. A9418-50G)
for WB), rabbit anti-TANGO1 (Sigma, Sigma, Cat no. HPA055922, Lot #000008249, 1:200
for IF, 1:1000 in 5% milk for WB), mouse anti-actin (Abcam, ab8224, 1:2000 in 5% milk for
WB), mouse anti-FLAG (F1804-50UG, IF 1:400), rabbit anti-FLAG (Sigma, Cat no. F7425,
Lot #06M4757, IF 1:100).
15

*Post-Golgi Mucin-2 Isolation for Western Blot*
HT29-MTX cells were cultured in medium deprived of FBS 24h in 6-wells plates prior to
mucus collection, as this avoided noisy Western Blot results and increased mucin-2
secretion, which was also observed by Cantero-Recasens et al. in HT29 cells (Cantero-
Recasens *et al*, 2019), without decreasing cell viability (**Supplementary Figure 7**). After
aspiration of medium, the cell layers were washed with 200 µL of 6 M Guadinium
Hydrochloride (Gdn-HCl, Sigma, Cat no. G3272-25G), 5 mM EDTA (Fluka, Cat no. 03690-
100ML), and 10 mM $NaH_2PO_4$ (Sigma, Cat no. 331988) at pH 6.5, and aspirated after 1
minute, loosely based on the method of Cantero-Recasens *et al*. (Cantero-Recasens *et al*.,
2019). Subsequently, 1 mL of DBPS was added and incubated for 30 minutes at 37 ⁰C,
which was then carefully collected and precipitated using methanol/chloroform protein
precipitation. Briefly, 4 mL of methanol (Sigma, Cat no. 322415-2L), 1 mL of chloroform
(Sigma, Cat no. 288306), and 3 mL of ultrapure $H_2O$ (MilliQ, Salmenkipp, Purelab Flex,
Elga) were added in that order and each to mix vortexed for 2 seconds. Subsequently, the
samples were centrifuged for 5 minutes at 4000 rpm and the top aqueous layer was
removed, as the protein was residing between the layers. Next, 4 mL of methanol was
added, vortexed, and centrifuged for 14 minutes at 4000 rpm. The methanol was removed,
and the protein pellet was dried using $N_2$ air flow and dissolved in 12 µL MilliQ and 8 µL 5x
Laemmli Buffer, consisting of 1.5 M Tris (Thermo Fisher, Cat no. 1.08382), 10% SDS
(Sigma-Aldrich, L3771-500G), 0.05% Bromophenol Blue (Bio-Rad, Cat no. 1610-404), 50%
glycerol (CalBiochem, Cat no. 356350-1000ML), and 1.43 M β-mercaptoethanol (Thermo
Fisher, Cat no. 31350010). Samples were used directly for SDS-PAGE and Western Blot
purposes or stored at -80 ⁰C until further use.
39

*Cell Lysate Sample Preparation*
Cell lysates were obtained from different serum-deprived 6 wells plates, as the Gdn-HCl
mucin-2 extraction interfered with obtaining proper cell lysates. Samples were washed twice
with ice cold DBPS and subsequently lysed in 200 µL RIPA buffer (50 mM Tris, 1% Triton X-
100, 12 mM sodium deoxycholate (Sigma, Cat no. 30970-25G), 0.1% (w/v) SDS, 150 mM
NaCl (Sigma, 746398-500G) in MilliQ), containing 1X Proteinase Inhibitor Cocktail
(Calbiochem, Cat no. 539134). Subsequently, the cell lysate was collected and agitated for
30 min at 800 rpm at 4 ⁰C in a Thermomixer (Eppendorf). Afterwards, the lysate was
sonicated (42 kHz, Branson 1510, Bransonic) in an ice bath intermittently 3 times for 10
seconds each, maintaining 1 minute at 4 ⁰C in between sonication pulses. Lastly, the sample
was centrifuged for 20 minutes at 12000 rpm and 100 µL of the supernatant was used or

stored at -80 ºC until further use. Protein concentrations were measured using the DC
Protein Assay (Bio-Rad, Cat no. 5000112) according to the manufacturer's instructions.
3

*Western Blot*

Mucin-2 precipitated samples could directly be processed, whereas 17.5 µg protein of cell
lysates were combined with 8 µL of 5x Laemmli buffer. Protein samples were denatured at
100 ºC for 5 minutes and briefly centrifuged before loading onto a 4-12% Tris- Glycine
Novex Wedgewell gel (Fisher Scientific, Cat no. XP04125BOX). SDS-PAGE was performed
using standard running buffer in Xcell Surelock Electrophoresis Cell (Invitrogen) at 150V for
75 minutes. Protein transfer was completed in a transfer buffer, containing 20% Emsure
methanol, 0.1% (w/v) SDS, 25 mM TRIS-Base (Sigma, Cat no.), and 19.2 mM Glycine
(Sigma, Cat no. G7126) in MilliQ for 100 minutes at 100 V onto a 0.45 µm PVDF membrane
(Immobilon). Protein transfer was confirmed using a Coommassie Blue stain (Fisher
Scientific, Cat no. 23236). The membrane was incubated in blocking buffer (5% non-fat dry
milk (Carlroth, Cat no. T145.1) in TBST containing 0.005% Tween-20 (Sigma, Cat no.
P7949-500ML), or 0.02% Tween-20 (TBST-XL)), for 1 hour at room temperature on an
orbital shaker at 50 rpm. Primary antibodies were incubated overnight at 4 ºC on an orbital
shaker, as mentioned in section 2.3. The next morning, membranes were washed 3 times for
5 minutes in TBST or TBST-XL for mucin-2 and incubated with anti-mouse IgG HRP
Conjugate (1:10000, Promega, Cat no. W4021) or anti-rabbit IgG HRP Conjugate (1:10000,
Promega, Cat no. W4011) in blocking buffer and developed using the SuperSignal West
Femto Maximum Sensitivity Substrate kit (Thermofisher Scientific, Cat no. 34095).
Membranes were imaged using FluorChem M Imager (ProteinSimple) and analyzed in
ImageJ (v1.52p, NIH).
25

*Immunofluorescence*

Cells were seeded and cultured at 15 000 cells/cm2 in a 24 wells plate onto a #1.5 coverslip
with a diameter of 12 mm (Thermo Scientific, Cat no. 1217N79) and deprived of serum for
24 hours before fixation. After 3 days, the cells were fixated using 4% formaldehyde (Sigma,
Cat no. 252549) in DPBS for 15 minutes at room temperature and subsequently washed
three times for 5 minutes with DPBS. Afterwards, samples were permeabilized in 0.1% (v/v)
Triton X-100 (Sigma-Aldrich, Cat no. T8787-250ML) for 10 minutes at room temperature and
blocked using the blocking buffer consisting of 5% (v/v) FBS in DBPS for 1 hour at room
temperature under gentle shaking. The primary antibodies were diluted in blocking buffer
and incubated overnight at 4 ºC. The next day, samples were washed three times for 5
minutes in DPBS. The secondary antibodies (Goat anti-Mouse IgG Alexa Fluor 488,
Invitrogen; Goat anti-Rabbit IgG Alexa Fluor 647, Invitrogen) were diluted 1:500 in blocking
solution and incubated for 60 minutes at room temperature in the dark under gentle shaking.
Subsequently, three washes of 5 minutes in DPBS were performed. The nuclei were stained
using DAPI (1:4000 in DPBS, Invitrogen, Cat no. D1306) and the samples were mounted
with DAKO mounting medium (Agilent, Cat no. S302380-2) onto a glass slide (Thermofisher,
Cat no. 28908). Samples were imaged using the Fluorescent EVOS Microscope (Invitrogen
EVOS FL Imaging System, ThermoFisher Scientific) or the Zeiss LSM 880 Confocal
Microscope (Zeiss). F-actin immunostaining was performed by incubating fixated,
permeabilized, and blocked cells with Phalloidin Alexa Fluor 633 (1:40 in DBPS, Invitrogen,
Cat no. A22284) for 30 minutes at room temperature under gentle shaking. Subsequently,
cells were washed 3 times with DPBS, counterstained with DAPI, and imaged using a
fluorescent microscope.
49

*RT-qPCR*

mRNA was isolated from (treated) cells in a 12 wells plate (Greiner Bio-One, Cellstar, Cat
no. 665 180) using the NucleoSpin RNA kit (Macherey-Nagel, Cat no. 740955.250)
according to the manufacturer's instructions. RNA quality was ensured with 260/230 ratio
between 1.8 and 2.2 using a ND-1000 spectrophotometer. cDNA samples were obtained
from 1 μg RNA using the iScript$^{TM}$ cDNA synthesis kit (Bio-Rad, Cat no. 170-8891) according
to the manufacturer's instructions. Reverse transcriptase quantitative polymerase chain
reaction (RT-qPCR) was performed with 5 ng cDNA per sample, 10 μM of forward and
reverse primer, and 2x SensiMix$^{TM}$ SYBR & Fluorescein (Bioline, Cat no. QT615-05) in a 384
wells plate (Bio-Rad, Cat no. HSP3805). Thermal cycling was performed in a Bio-Rad
CFX384 thermocycler using 40 cycles of 95 ºC melting for 15 seconds, 60 ºC annealing for
15 seconds, and 72 ºC extension for 15 seconds, prior to an initial hot start at 95 ºC for 10
minutes (see **Supplementary Table 1** for primer sequences).
13

*Cell Viability*

Cell viability was measured by measuring ATP production using Cell Titer Glo assay 2.0
(Promega, Cat no. G9241). Cells were seeded at a density of 15 000 cells/cm$^2$ in an opaque
96 well plate (Greiner Bio-one, Cat no. 655098) in 100 μL cell culture medium. After 3 days
of culture and relevant cell treatments, a volume of 100 μL of reagent was added to the cell
culture medium and allowed to react for 10 minutes at room temperature, under slow orbital
shaking (50 rpm, KS250 Basic, Ika Labortechnik). Staurosporin (10 μM, Millipore Sigma, Cat
no. 569397-100UG) was used as a negative control. Luminescence was measured using
Victor3 1420 Multilabel Counter (PerkinElmer) for 1 second per sample.
23

*Clinical validation in a cohort of individuals with inflammatory bowel disease (IBD)*

A human validation of the *in vitro* results was performed using intestinal biopsy samples from
individuals diagnosed with inflammatory bowel disease (IBD). Participants were recruited
from the outpatient clinic of the University Medical Center Groningen (UMCG) as part of the
1000IBD project, which involved gathering comprehensive phenotypic information and
molecular profiles (Imhann *et al*, 2019). Patients that were included were >18 years of age,
enrolled in the period of 2003 until 2019, and were diagnosed with Crohn's disease (CD) or
ulcerative colitis (UC). The diagnosis of IBD was based on various criteria, including clinical
symptoms, laboratory tests, endoscopic examination, and histopathological analysis of
biopsies. Additionally, biopsy samples from 16 healthy individuals without IBD were included
for comparison. Their biopsies were obtained during endoscopies conducted due to
suspected intestinal disease or as part of colon cancer screening, all of which yielded
negative results. Prior to sample collection, written informed consent was obtained from all
participants. The study was approved by the Institutional Review Board (IRB) of the UMCG
in Groningen, the Netherlands (in Dutch: 'Medisch Ethische Toetsingscommissie', METc;
IRB nos. 2008/338 and 2016/424), and adhered to the principles outlined in the Declaration
of Helsinki (2013).
41

*Collection of intestinal biopsies*

A total of 711 intestinal biopsies were obtained from 420 patients diagnosed with IBD along
with 52 intestinal biopsies from 16 healthy non-IBD controls (Hu *et al*, 2024). During an
endoscopic procedure, biopsies were immediately frozen in liquid nitrogen by either an
endoscopy nurse or a research technician. Biopsies were taken from both inflamed and non-
inflamed regions in adjacent areas, and their inflammatory status was assessed
histologically by certified pathologists. For patients with CD and UC, biopsies were taken
from both the ileal and colonic tissue. However, inflamed ileal biopsies from UC patients
(n=3) were excluded from the analysis, likely due to backwash ileitis. The biopsies were
stored at -80ºC until further processing.

*RNA isolation, sequencing, and data processing*

RNA isolation was conducted using the AllPrep DNA/RNA mini kit (Qiagen, ref no: 80204) following the manufacturer's instructions. To homogenize the intestinal biopsies, the Qiagen Tissue Lyser with stainless steel beads (diameter 5 mm, reference number: 69989) was utilized in RLT lysis buffer containing β-mercaptoethanol. In the first batch of samples, the BioScientific NEXTflexTM Rapid Directional RNA-Seq Kit (Perkin-Elmer) was employed for sample preparation. Paired-end RNA sequencing was performed on the Illumina NextSeq500 sequencer (Illumina). For the second batch of samples, the Eukaryotic Transcriptome Library (Novogene) was constructed for sample preparation, followed by paired-end RNA sequencing on the Illumina HiSeq PE250 platform. Sequencing was carried out in two separate batches, and pseudo-randomization was employed across plates to mitigate potential batch effects, taking into account the type of IBD diagnosis, biopsy location, and disease activity. Batch effects were considered in all the analyses. On average, approximately 25 million reads were generated per sample. The quality of raw reads was assessed using FastQC with default parameters (ref v.0.11.7). Adapters identified by FastQC were removed using Cutadapt (ref v1.1) with default settings. Sickle (ref v1.200) was used to trim low-quality ends from the reads (length <25 nucleotides, quality <20). Alignment of the reads to the human genome (human_glk_v37) was performed using HISAT (ref v0.1.6) with a maximum allowance of two mismatches, and read sorting was done using SAMtools (ref v0.1.19). SAMtools flagstat and Picard tools (ref v2.9.0) were utilized to obtain mapping statistics. Six samples with a low percentage of read alignment (<90%) were excluded from the analysis. Gene expression was estimated using HTSeq (ref v0.9.1) based on Ensemble version 75 annotation, resulting in an RNA expression dataset comprising 15,934 genes. Gene level expression data were normalized using a trimmed mean of M values, and clr transformation was applied, resulting in a set of 826 mucosal RNA-seq samples.

*Statistical Analysis*

Evaluation of statistical significance was determined using a one- or two-way ANOVA followed by Bonferroni's post-hoc test using GraphPad Prism 8.0.1 (GraphPad Software, La Jolla, CA). A p-value <0.05 was considered to indicate statistical significance. All experiments were conducted in triplicates (n = 3) unless otherwise stated. For human intestinal mucosal RNA-sequencing analysis, gene expressions were compared between group of individuals with CD, UC and controls using Mann-Whitney *U*-tests followed by post-hoc Bonferroni correction for multiple testing. Analyses and visualization of gene expressions were performed using the Python programming language (v.3.9.0; Python Software Foundation; https://python.org) using the *pandas* (v.1.4.2), *numpy* (v.1.21.0), *matplotlib* (v.3.5.1), and *seaborn* (v.0.12.1) packages.

**References**

Ahn DH, Crawley SC, Hokari R, Kato S, Yang SC, Li JD, Kim YS (2005) TNF-alpha Activates MUC2 Transcription via NF-kappaB but Inhibits via JNK Activation. *Cellular Physiology and Biochemistry* 15: 029-040

Ambort D, Johansson ME, Gustafsson JK, Nilsson HE, Ermund A, Johansson BR, Koeck PJ, Hebert H, Hansson GC (2012a) Calcium and pH-dependent packing and release of the gel-forming MUC2 mucin. *Proceedings of the National Academy of Sciences of the United States of America* 109: 5645-5650

Ambort D, Johansson MEV, Gustafsson JK, Nilsson HE, Ermund A, Johansson BR, Koeck PJB, Hebert H, Hansson GC (2012b) Calcium and pH-dependent packing and release of the gel-forming MUC2 mucin. *Proceedings of the National Academy of Sciences* 109: 5645-5650

Amodio G, Venditti R, De Matteis MA, Moltedo O, Pignataro P, Remondelli P (2013) Endoplasmic reticulum stress reduces COPII vesicle formation and modifies Sec23a cycling at ERESs. *FEBS Lett* 587: 3261-3266

Arike L, Hansson GC (2016) The Densely O-Glycosylated MUC2 Mucin Protects the Intestine and Provides Food for the Commensal Bacteria. *Journal of Molecular Biology* 428: 3221-3229

Arndt S, Bosserhoff AK (2007) Reduced expression of TANGO in colon and hepatocellular carcinomas. *Oncology reports* 18: 885-891

Bankole E, Read E, Curtis MA, Neves JF, Garnett JA (2021) The Relationship between Mucins and Ulcerative Colitis: A Systematic Review. *J Clin Med* 10

Baumgart DC, Carding SR (2007) Inflammatory bowel disease: cause and immunobiology. *The Lancet* 369: 1627-1640

Benahmed M, Meresse B, Arnulf B, Barbe U, Mention JJ, Verkarre V, Allez M, Cellier C, Hermine O, Cerf-Bensussan N (2007) Inhibition of TGF-beta signaling by IL-15: a new role for IL-15 in the loss of immune homeostasis in celiac disease. *Gastroenterology* 132: 994-1008

Birchenough GMH, Johansson ME, Gustafsson JK, Bergström JH, Hansson GC (2015) New developments in goblet cell mucus secretion and function. *Mucosal Immunology* 8: 712-719

Bisnett BJ, Condon BM, Lamb CH, Georgiou GR, Boyce M (2021) Export Control: Post-transcriptional Regulation of the COPII Trafficking Pathway. *Front Cell Dev Biol* 8: 618652-618652

Boyadjiev SA, Fromme JC, Ben J, Chong SS, Nauta C, Hur DJ, Zhang G, Hamamoto S, Schekman R, Ravazzola M *et al* (2006) Cranio-lenticulo-sutural dysplasia is caused by a SEC23A mutation leading to abnormal endoplasmic-reticulum-to-Golgi trafficking. *Nature genetics* 38: 1192-1197

Brandl K, Tomisato W, Li X, Neppl C, Pirie E, Falk W, Xia Y, Moresco EMY, Baccala R, Theofilopoulos AN *et al* (2012) Yip1 domain family, member 6 (Yipf6) mutation induces spontaneous intestinal inflammation in mice. *Proceedings of the National Academy of Sciences* 109: 12650-12655

Buniello A, MacArthur JAL, Cerezo M, Harris LW, Hayhurst J, Malangone C, McMahon A, Morales J, Mountjoy E, Sollis E *et al* (2019) The NHGRI-EBI GWAS Catalog of published genome-wide association studies, targeted arrays and summary statistics 2019. *Nucleic Acids Res* 47: D1005-D1012

Butkinaree C, Guo L, Ramkhelawon B, Wanschel A, Brodsky JL, Moore KJ, Fisher EA (2014) A regulator of secretory vesicle size, Kelch-like protein 12, facilitates the secretion of apolipoprotein B100 and very-low-density lipoproteins--brief report. *Arteriosclerosis, thrombosis, and vascular biology* 34: 251-254

Cantero-Recasens G, Butnaru CM, Brouwers N, Mitrovic S, Valverde MA, Malhotra V (2019) Sodium channel TRPM4 and sodium/calcium exchangers (NCX) cooperate in the control of Ca2+-induced mucin secretion from goblet cells. *Journal of Biological Chemistry* 294: 816-826

Cari L, Rosati L, Leoncini G, Lusenti E, Gentili M, Nocentini G, Riccardi C, Migliorati G, Ronchetti S (2023) Association of GILZ with MUC2, TLR2, and TLR4 in Inflammatory Bowel Disease. *Int J Mol Sci* 24

Chambers SM, Fasano CA, Papapetrou EP, Tomishima M, Sadelain M, Studer L (2009) Highly efficient neural conversion of human ES and iPS cells by dual inhibition of SMAD signaling. *Nat Biotechnol* 27: 275-280

Connolly LM, McFalls CM, McMahon IG, Bhat AM, Artlett CM (2023) Caspase 1 Enhances Transport and Golgi Organization Protein 1 Expression to Promote Procollagen Export From the Endoplasmic Reticulum in Systemic Sclerosis Contributing to Fibrosis. *Arthritis Rheumatol* 75: 1831-1841

Cullen PJ (2011) Post-translational regulation of signaling mucins. *Curr Opin Struct Biol* 21: 590-596

Dekker J, Strous GJ (1990) Covalent oligomerization of rat gastric mucin occurs in the rough endoplasmic reticulum, is N-glycosylation-dependent, and precedes initial O-glycosylation. *The Journal of biological chemistry* 265: 18116-18122

Di Sabatino A, Pickard KM, Rampton D, Kruidenier L, Rovedatti L, Leakey NA, Corazza GR, Monteleone G, MacDonald TT (2008) Blockade of transforming growth factor beta upregulates T-box transcription factor T-bet, and increases T helper cell type 1 cytokine and matrix metalloproteinase-3 production in the human gut mucosa. *Gut* 57: 605-612

Dohrman A, Miyata S, Gallup M, Li JD, Chapelin C, Coste A, Escudier E, Nadel J, Basbaum C (1998) Mucin gene (MUC 2 and MUC 5AC) upregulation by Gram-positive and Gram-negative bacteria. *Biochimica et biophysica acta* 1406: 251-259

Domińska K, Kowalska K, Matysiak ZE, Płuciennik E, Ochędalski T, Piastowska-Ciesielska AW (2017) Regulation of mRNA gene expression of members of the NF-κB transcription factor gene family by angiotensin II and relaxin 2 in normal and cancer prostate cell lines. *Mol Med Rep* 15: 4352-4359

Fang J, Liu M, Zhang X, Sakamoto T, Taatjes DJ, Jena BP, Sun F, Woods J, Bryson T, Kowluru A *et al* (2015) COPII-Dependent ER Export: A Critical Component of Insulin Biogenesis and β-Cell ER Homeostasis. *Molecular endocrinology (Baltimore, Md)* 29: 1156-1169

Fernandez-Escamilla AM, Rousseau F, Schymkowitz J, Serrano L (2004) Prediction of sequence-dependent and mutational effects on the aggregation of peptides and proteins. *Nat Biotechnol* 22: 1302-1306

Ferrari E, Monzani R, Saverio V, Gagliardi M, Pańczyszyn E, Raia V, Villella VR, Bona G, Pane M, Amoruso A *et al* (2021) Probiotics Supplements Reduce ER Stress and Gut Inflammation Associated with Gliadin Intake in a Mouse Model of Gluten Sensitivity. *Nutrients* 13: 1221

Forstner G (1995) Signal transduction, packaging and secretion of mucins. *Annu Rev Physiol* 57: 585-605

Fujiwara T, Oda K, Yokota S, Takatsuki A, Ikehara Y (1988) Brefeldin A causes disassembly of the Golgi complex and accumulation of secretory proteins in the endoplasmic reticulum. *Journal of Biological Chemistry* 263: 18545-18552

Garbes L, Kim K, Rieß A, Hoyer-Kuhn H, Beleggia F, Bevot A, Kim MJ, Huh YH, Kweon HS, Savarirayan R *et al* (2015) Mutations in SEC24D, encoding a component of the COPII machinery, cause a syndromic form of osteogenesis imperfecta. *American journal of human genetics* 96: 432-439

Gersemann M, Becker S, Kubler I, Koslowski M, Wang G, Herrlinger KR, Griger J, Fritz P, Fellermann K, Schwab M *et al* (2009) Differences in goblet cell differentiation between Crohn's disease and ulcerative colitis. *Differentiation* 77: 84-94

Gorur A, Yuan L, Kenny SJ, Baba S, Xu K, Schekman R (2017) COPII-coated membranes function as transport carriers of intracellular procollagen I. *Journal of Cell Biology* 216: 1745-1759

Gupta VA, Beggs AH (2014) Kelch proteins: emerging roles in skeletal muscle development and diseases. *Skelet Muscle* 4: 11

Halder SK, Beauchamp RD, Datta PK (2005) A specific inhibitor of TGF-beta receptor kinase, SB-431542, as a potent antitumor agent for human cancers. *Neoplasia* 7: 509-521

Hanski C, Born M, Foss HD, Marowski B, Mansmann U, Arasteh K, Bachler B, Papenfuss M, Niedobitek F (1999) Defective post-transcriptional processing of MUC2 mucin in ulcerative colitis and in Crohn's disease increases detectability of the MUC2 protein core. *J Pathol* 188: 304-311

Hatamzade Esfahani N, Day AS (2023) The Role of TGF-β, Activin and Follistatin in Inflammatory Bowel Disease. *Gastrointestinal Disorders* 5: 167-186

Heazlewood CK, Cook MC, Eri R, Price GR, Tauro SB, Taupin D, Thornton DJ, Png CW, Crockford TL, Cornall RJ *et al* (2008) Aberrant Mucin Assembly in Mice Causes Endoplasmic Reticulum Stress and Spontaneous Inflammation Resembling Ulcerative Colitis. *PLOS Medicine* 5: e54

Holmgaard RB, Schaer DA, Li Y, Castaneda SP, Murphy MY, Xu X, Inigo I, Dobkin J, Manro JR, Iversen PW *et al* (2018) Targeting the TGFβ pathway with galunisertib, a TGFβRI small molecule inhibitor, promotes anti-tumor immunity leading to durable, complete responses, as monotherapy and in combination with checkpoint blockade. *Journal for ImmunoTherapy of Cancer* 6: 47

Hu S, Bourgonje AR, Gacesa R, Jansen BH, Bjork JR, Bangma A, Hidding IJ, van Dullemen HM, Visschedijk MC, Faber KN *et al* (2024) Mucosal host-microbe interactions associate with clinical phenotypes in inflammatory bowel disease. *Nat Commun* 15: 1470

Huang X, Gao Y, Li S, Wu C, Wang J, Zheng N (2019) Modulation of Mucin (MUC2, MUC5AC and MUC5B) mRNA Expression and Protein Production and Secretion in Caco-2/HT29-MTX Co-Cultures Following Exposure to Individual and Combined Aflatoxin M1 and Ochratoxin A. *Toxins (Basel)* 11: 132

Ihara S, Hirata Y, Koike K (2017) TGF-beta in inflammatory bowel disease: a key regulator of immune cells, epithelium, and the intestinal microbiota. *J Gastroenterol* 52: 777-787

Ihara S, Hirata Y, Serizawa T, Suzuki N, Sakitani K, Kinoshita H, Hayakawa Y, Nakagawa H, Ijichi H, Tateishi K *et al* (2016) TGF-beta Signaling in Dendritic Cells Governs Colonic Homeostasis by Controlling Epithelial Differentiation and the Luminal Microbiota. *Journal of immunology (Baltimore, Md : 1950)* 196: 4603-4613

Imhann F, Van der Velde KJ, Barbieri R, Alberts R, Voskuil MD, Vich Vila A, Collij V, Spekhorst LM, Van der Sloot KWJ, Peters V *et al* (2019) The 1000IBD project: multi-omics data of 1000 inflammatory bowel disease patients; data release 1. *BMC Gastroenterol* 19: 5

Jensen D, Schekman R (2011) COPII-mediated vesicle formation at a glance. *Journal of Cell Science* 124: 1-4

Jin L, Pahuja KB, Wickliffe KE, Gorur A, Baumgärtel C, Schekman R, Rape M (2012) Ubiquitin-dependent regulation of COPII coat size and function. *Nature* 482: 495-500

Johansson ME, Sjovall H, Hansson GC (2013) The gastrointestinal mucus system in health and disease. *Nat Rev Gastroenterol Hepatol* 10: 352-361

Johansson MEV, Phillipson M, Petersson J, Velcich A, Holm L, Hansson GC (2008) The inner of the two Muc2 mucin-dependent mucus layers in colon is devoid of bacteria. *Proceedings of the National Academy of Sciences* 105: 15064-15069

Jonckheere N, Van Der Sluis M, Velghe A, Buisine M-P, Sutmuller M, Ducourouble M-P,
Pigny P, Büller HA, Aubert J-P, Einerhand AWC *et al* (2004) Transcriptional activation of
the murine Muc5ac mucin gene in epithelial cancer cells by TGF-beta/Smad4 signalling
pathway is potentiated by Sp1. *Biochem J* 377: 797-808
Jono H, Shuto T, Xu H, Kai H, Lim DJ, Gum JR, Kim YS, Yamaoka S, Feng X-H, Li J-D
(2002) Transforming growth factor-beta -Smad signaling pathway cooperates with NF-kappa
B to mediate nontypeable Haemophilus influenzae-induced MUC2 mucin transcription. *The
Journal of biological chemistry* 277: 45547-45557
Kaser A, Blumberg RS (2010) Endoplasmic reticulum stress and intestinal inflammation.
*Mucosal Immunol* 3: 11-16
Kaser A, Lee AH, Franke A, Glickman JN, Zeissig S, Tilg H, Nieuwenhuis EE, Higgins DE,
Schreiber S, Glimcher LH *et al* (2008) XBP1 links ER stress to intestinal inflammation and
confers genetic risk for human inflammatory bowel disease. *Cell* 134: 743-756
Kassan M, Galan M, Partyka M, Saifudeen Z, Henrion D, Trebak M, Matrougui K (2012)
Endoplasmic reticulum stress is involved in cardiac damage and vascular endothelial
dysfunction in hypertensive mice. *Arteriosclerosis, thrombosis, and vascular biology* 32:
1652-1661
Kawaguchi K, Endo A, Fukushima T, Madoka Y, Tanaka T, Komada M (2018) Ubiquitin-
specific protease 8 deubiquitinates Sec31A and decreases large COPII carriers and collagen
IV secretion. *Biochem Biophys Res Commun* 499: 635-641
Kim S, Domon-Dell C, Wang Q, Chung DH, Di Cristofano A, Pandolfi PP, Freund JN, Evers
BM (2002) PTEN and TNF-alpha regulation of the intestinal-specific Cdx-2 homeobox gene
through a PI3K, PKB/Akt, and NF-kappaB-dependent pathway. *Gastroenterology* 123: 1163-
1178
Kranjc T, Dempsey E, Cagney G, Nakamura N, Shields DC, Simpson JC (2017) Functional
characterisation of the YIPF protein family in mammalian cells. *Histochemistry and Cell
Biology* 147: 439-451
Kubiczkova L, Sedlarikova L, Hajek R, Sevcikova S (2012) TGF-β – an excellent servant but
a bad master. *Journal of Translational Medicine* 10: 183
Larsson JM, Karlsson H, Crespo JG, Johansson ME, Eklund L, Sjovall H, Hansson GC
(2011) Altered O-glycosylation profile of MUC2 mucin occurs in active ulcerative colitis and
is associated with increased inflammation. *Inflamm Bowel Dis* 17: 2299-2307
Lee MC, Miller EA, Goldberg J, Orci L, Schekman R (2004) Bi-directional protein transport
between the ER and Golgi. *Annual review of cell and developmental biology* 20: 87-123
Lee SH, Kwon Je, Cho M-L (2018) Immunological pathogenesis of inflammatory bowel
disease. *Intest Res* 16: 26-42
Lee SU, Kim MO, Kang MJ, Oh ES, Ro H, Lee RW, Song YN, Jung S, Lee JW, Lee SY *et al*
(2021) Transforming Growth Factor beta Inhibits MUC5AC Expression by Smad3/HDAC2
Complex Formation and NF-kappaB Deacetylation at K310 in NCI-H292 Cells. *Mol Cells*
44: 38-49
Lekszas C, Foresti O, Raote I, Liedtke D, Konig EM, Nanda I, Vona B, De Coster P,
Cauwels R, Malhotra V *et al* (2020) Biallelic TANGO1 mutations cause a novel syndromal
disease due to hampered cellular collagen secretion. *Elife* 9
Lemos RR, Oliveira DF, Zatz M, Oliveira JR (2011) Population and computational analysis
of the MGEA6 P521A variation as a risk factor for familial idiopathic basal ganglia
calcification (Fahr's disease). *J Mol Neurosci* 43: 333-336
Lidell ME, Johansson ME, Morgelin M, Asker N, Gum JR, Jr., Kim YS, Hansson GC (2003)
The recombinant C-terminus of the human MUC2 mucin forms dimers in Chinese-hamster
ovary cells and heterodimers with full-length MUC2 in LS 174T cells. *Biochem J* 372: 335-
345

Linden SK, Sutton P, Karlsson NG, Korolik V, McGuckin MA (2008) Mucins in the mucosal barrier to infection. *Mucosal Immunology* 1: 183-197

Linding R, Schymkowitz J, Rousseau F, Diella F, Serrano L (2004) A comparative study of the relationship between protein structure and beta-aggregation in globular and intrinsically disordered proteins. *J Mol Biol* 342: 345-353

Lindner P, Christensen SB, Nissen P, Moller JV, Engedal N (2020) Cell death induced by the ER stressor thapsigargin involves death receptor 5, a non-autophagic function of MAP1LC3B, and distinct contributions from unfolded protein response components. *Cell Commun Signal* 18: 12

Liu L, Cai J, Wang H, Liang X, Zhou Q, Ding C, Zhu Y, Fu T, Guo Q, Xu Z *et al* (2019) Coupling of COPII vesicle trafficking to nutrient availability by the IRE1α-XBP1s axis. *Proceedings of the National Academy of Sciences of the United States of America* 116: 11776-11785

Luo C, Wang F, Ren X, Ke T, Xu C, Tang B, Qin S, Yao Y, Chen Q, Wang QK (2017) Identification of a molecular signaling gene-gene regulatory network between GWAS susceptibility genes ADTRP and MIA3/TANGO1 for coronary artery disease. *Biochim Biophys Acta Mol Basis Dis* 1863: 1640-1653

Ma X, Dai Z, Sun K, Zhang Y, Chen J, Yang Y, Tso P, Wu G, Wu Z (2017) Intestinal Epithelial Cell Endoplasmic Reticulum Stress and Inflammatory Bowel Disease Pathogenesis: An Update Review. *Front Immunol* 8: 1271

Maeda M, Komatsu Y, Saito K (2020) Mitotic ER Exit Site Disassembly and Reassembly Are Regulated by the Phosphorylation Status of TANGO1. *Developmental Cell* 55: 237-250.e235

Maiers JL, Kostallari E, Mushref M, deAssuncao TM, Li H, Jalan-Sakrikar N, Huebert RC, Cao S, Malhi H, Shah VH (2017) The unfolded protein response mediates fibrogenesis and collagen I secretion through regulating TANGO1 in mice. *Hepatology* 65: 983-998

McCaughey J, Mantell JM, Neal CR, Heesom K, Stephens DJ (2021a) MIA3/TANGO1 enables efficient secretion by constraining COPII vesicle budding. *bioRxiv*: 2021.2002.2024.432632

McCaughey J, Stephens DJ (2018) COPII-dependent ER export in animal cells: adaptation and control for diverse cargo. *Histochemistry and Cell Biology* 150: 119-131

McCaughey J, Stevenson NL, Mantell JM, Neal CR, Paterson A, Heesom K, Stephens DJ (2021b) A general role for TANGO1, encoded by MIA3, in secretory pathway organization and function. *J Cell Sci* 134

Medicine I, Virus CAFSNV (1999) *Assessment of Future Scientific Needs for Live Variola Virus*. National Academies Press

Mejías-Luque R, Lindén SK, Garrido M, Tye H, Najdovska M, Jenkins BJ, Iglesias M, Ernst M, de Bolós C (2010) Inflammation modulates the expression of the intestinal mucins MUC2 and MUC4 in gastric tumors. *Oncogene* 29: 1753-1762

Mesquita P, Peixoto AJ, Seruca R, Hanski C, Almeida R, Silva F, Reis C, David L (2003) Role of site-specific promoter hypomethylation in aberrant MUC2 mucin expression in mucinous gastric carcinomas. *Cancer Lett* 189: 129-136

Mo J-S, Alam KJ, Kim H-S, Lee Y-M, Yun K-J, Chae S-C (2016) MicroRNA 429 Regulates Mucin Gene Expression and Secretion in Murine Model of Colitis. *Journal of Crohn's and Colitis* 10: 837-849

Moehle C, Ackermann N, Langmann T, Aslanidis C, Kel A, Kel-Margoulis O, Schmitz-Madry A, Zahn A, Stremmel W, Schmitz G (2006) Aberrant intestinal expression and allelic variants of mucin genes associated with inflammatory bowel disease. *J Mol Med (Berl)* 84: 1055-1066

Mollenhauer HH, James Morré D, Rowe LD (1990) Alteration of intracellular traffic by monensin; mechanism, specificity and relationship to toxicity. *Biochimica et Biophysica Acta (BBA) - Reviews on Biomembranes* 1031: 225-246

Monteleone G, Laudisi F, Stolfi C (2023) Smad7 as a positive regulator of intestinal inflammatory diseases. *Curr Res Immunol* 4: 100055

Monteleone G, Mann J, Monteleone I, Vavassori P, Bremner R, Fantini M, Del Vecchio Blanco G, Tersigni R, Alessandroni L, Mann D *et al* (2004) A failure of transforming growth factor-beta1 negative regulation maintains sustained NF-kappaB activation in gut inflammation. *The Journal of biological chemistry* 279: 3925-3932

Moresco EMY, Brandl K (2013) Linking membrane trafficking and intestinal homeostasis. *Tissue Barriers* 1: e23119

Naghdalipour M, Moradi N, Fadaei R, Rezghi Barez S, Sayyahfar S, Mokhtare M, Fard TK, Fallah S, Esteghamati A (2022) Alteration of miR-21, miR-433 and miR-590 tissue expression related to the TGF-beta signaling pathway in ulcerative colitis patients. *Arch Physiol Biochem* 128: 1170-1174

Norman GL, Yang CY, Ostendorff HP, Shums Z, Lim MJ, Wang J, Awad A, Hirschfield GM, Milkiewicz P, Bloch DB *et al* (2015) Anti-kelch-like 12 and anti-hexokinase 1: novel autoantibodies in primary biliary cirrhosis. *Liver Int* 35: 642-651

Nozoe T, Anai H, Nasu S, Sugimachi K (2000) Clinicopathological characteristics of mucinous carcinoma of the colon and rectum. *Journal of surgical oncology* 75: 103-107

Nylander S, Kalies I (1999) Brefeldin A, but not monensin, completely blocks CD69 expression on mouse lymphocytes:: efficacy of inhibitors of protein secretion in protocols for intracellular cytokine staining by flow cytometry. *Journal of Immunological Methods* 224: 69-76

Okada SF, Zhang L, Kreda SM, Abdullah LH, Davis CW, Pickles RJ, Lazarowski ER, Boucher RC (2011) Coupled nucleotide and mucin hypersecretion from goblet-cell metaplastic human airway epithelium. *Am J Respir Cell Mol Biol* 45: 253-260

Okumura N, Hashimoto K, Kitahara M, Okuda H, Ueda E, Watanabe K, Nakahara M, Sato T, Kinoshita S, Tourtas T *et al* (2017) Activation of TGF-beta signaling induces cell death via the unfolded protein response in Fuchs endothelial corneal dystrophy. *Sci Rep* 7: 6801

Oslowski CM, Urano F (2011) Measuring ER stress and the unfolded protein response using mammalian tissue culture system. *Methods in enzymology* 490: 71-92

Penney ME, Parfrey PS, Savas S, Yilmaz YE (2018) Associations of single nucleotide polymorphisms with mucinous colorectal cancer: genome-wide common variant and gene-based rare variant analyses. *Biomark Res* 6: 17-17

Perez-Vilar (2008) Formation of mucin granules. In: *The Golgi Apparatus*, Mironov A.A. P.M. (ed.)Springer: Vienna

Perez-Vilar J (2007) Mucin granule intraluminal organization. *Am J Respir Cell Mol Biol* 36: 183-190

Podolsky DK (1991) Inflammatory Bowel Disease. *New England Journal of Medicine* 325: 928-937

Rabouille C (2017) Pathways of Unconventional Protein Secretion. *Trends in Cell Biology* 27: 230-240

Ramanan D, Cadwell K (2016) Intrinsic Defense Mechanisms of the Intestinal Epithelium. *Cell Host & Microbe* 19: 434-441

Ran Y, Xiong MG, Xu ZS, Luo WW, Wang SY, Wang YY (2019) YIPF5 Is Essential for Innate Immunity to DNA Virus and Facilitates COPII-Dependent STING Trafficking. *Journal of immunology (Baltimore, Md : 1950)* 203: 1560-1570

Raote I, Ortega-Bellido M, Santos AJM, Foresti O, Zhang C, Garcia-Parajo MF, Campelo F,
Malhotra V (2018) TANGO1 builds a machine for collagen export by recruiting and spatially
organizing COPII, tethers and membranes. *eLife* 7: e32723
Redhai S, Boutros M (2021) The Role of Organelles in Intestinal Function, Physiology, and
Disease. *Trends in Cell Biology* 31: 485-499
Rex J, Lutz A, Faletti LE, Albrecht U, Thomas M, Bode JG, Borner C, Sawodny O, Merfort I
(2019) IL-1β and TNFα Differentially Influence NF-κB Activity and FasL-Induced
Apoptosis in Primary Murine Hepatocytes During LPS-Induced Inflammation. *Frontiers in*
*Physiology* 10
Reynolds HM, Zhang L, Tran DT, Ten Hagen KG (2019) Tango1 coordinates the formation
of endoplasmic reticulum/Golgi docking sites to mediate secretory granule formation.
*Journal of Biological Chemistry* 294: 19498-19510
Rodriguez-Pineiro AM, Bergstrom JH, Ermund A, Gustafsson JK, Schutte A, Johansson ME,
Hansson GC (2013) Studies of mucus in mouse stomach, small intestine, and colon. II.
Gastrointestinal mucus proteome reveals Muc2 and Muc5ac accompanied by a set of core
proteins. *Am J Physiol Gastrointest Liver Physiol* 305: G348-356
Rousseau F, Schymkowitz J, Serrano L (2006) Protein aggregation and amyloidosis:
confusion of the kinds? *Curr Opin Struct Biol* 16: 118-126
Saito K, Chen M, Bard F, Chen S, Zhou H, Woodley D, Polischuk R, Schekman R, Malhotra
V (2009) TANGO1 Facilitates Cargo Loading at Endoplasmic Reticulum Exit Sites. *Cell*
136: 891-902
Saito K, Yamashiro K, Ichikawa Y, Erlmann P, Kontani K, Malhotra V, Katada T (2011)
cTAGE5 mediates collagen secretion through interaction with TANGO1 at endoplasmic
reticulum exit sites. *Mol Biol Cell* 22: 2301-2308
Santos AJ, Nogueira C, Ortega-Bellido M, Malhotra V (2016) TANGO1 and Mia2/cTAGE5
(TALI) cooperate to export bulky pre-chylomicrons/VLDLs from the endoplasmic reticulum.
*The Journal of cell biology* 213: 343-354
Scholz CC, Cavadas MAS, Tambuwala MM, Hams E, Rodríguez J, Kriegsheim Av, Cotter P,
Bruning U, Fallon PG, Cheong A *et al* (2013) Regulation of IL-1β–induced NF-κB by
hydroxylases links key hypoxic and inflammatory signaling pathways. *Proceedings of the*
*National Academy of Sciences* 110: 18490-18495
Schumann M, Siegmund B, Schulzke JD, Fromm M (2017) Celiac Disease: Role of the
Epithelial Barrier. *Cellular and Molecular Gastroenterology and Hepatology* 3: 150-162
Sedda S, Marafini I, Dinallo V, Di Fusco D, Monteleone G (2015) The TGF-β/Smad System
in IBD Pathogenesis. *Inflammatory Bowel Diseases* 21: 2921-2925
Sheng YH, Hasnain SZ, Florin THJ, McGuckin MA (2012) Mucins in inflammatory bowel
diseases and colorectal cancer. *Journal of Gastroenterology and Hepatology* 27: 28-38
Shkoda A, Ruiz PA, Daniel H, Kim SC, Rogler G, Sartor RB, Haller D (2007) Interleukin-10
blocked endoplasmic reticulum stress in intestinal epithelial cells: impact on chronic
inflammation. *Gastroenterology* 132: 190-207
Tawiah A, Cornick S, Moreau F, Gorman H, Kumar M, Tiwari S, Chadee K (2018) High
MUC2 Mucin Expression and Misfolding Induce Cellular Stress, Reactive Oxygen
Production, and Apoptosis in Goblet Cells. *The American journal of pathology* 188: 1354-
1373
Troncone E, Marafini I, Stolfi C, Monteleone G (2018) Transforming Growth Factor-
β1/Smad7 in Intestinal Immunity, Inflammation, and Cancer. *Frontiers in Immunology* 9
Tytgat KM, Opdam FJ, Einerhand AW, Büller HA, Dekker J (1996) MUC2 is the prominent
colonic mucin expressed in ulcerative colitis. *Gut* 38: 554-563
van Klinken BJ-W, Einerhand AWC, Büller HA, Dekker J (1998) The oligomerization of a
family of four genetically clustered human gastrointestinal mucins. *Glycobiology* 8: 67-75

Van Seuningen I, Pigny P, Perrais M, Porchet N, Aubert JP (2001) Transcriptional regulation
of the 11p15 mucin genes. Towards new biological tools in human therapy, in inflammatory
diseases and cancer? *Front Biosci* 6: D1216-1234
Verstrepen L, Bekaert T, Chau TL, Tavernier J, Chariot A, Beyaert R (2008) TLR-4, IL-1R
and TNF-R signaling to NF-kappaB: variations on a common theme. *Cell Mol Life Sci* 65:
2964-2978
Wu J, Gong J, Geng J, Song Y (2008) Deoxycholic acid induces the overexpression of
intestinal mucin, MUC2, via NF-kB signaling pathway in human esophageal adenocarcinoma
cells. *BMC Cancer* 8: 333
Yamamoto H, Bai YQ, Yuasa Y (2003) Homeodomain protein CDX2 regulates goblet-
specific MUC2 gene expression. *Biochem Biophys Res Commun* 300: 813-818
Yuan L, Baba S, Bajaj K, Schekman R (2017) Cell-free Generation of COPII-coated
Procollagen I Carriers. *Bio-protocol* 7: e2450
Yuan L, Kenny SJ, Hemmati J, Xu K, Schekman R (2018) TANGO1 and SEC12 are
copackaged with procollagen I to facilitate the generation of large COPII carriers.
*Proceedings of the National Academy of Sciences* 115: E12255-E12264
Zhang F, Wang Y, Wang T, Yao L, Lam SM, Huang X, Fan J, Wang Q, Liu L, Jiang Y *et al*
(2018) cTAGE5/MEA6 plays a critical role in neuronal cellular components trafficking and
brain development. *Proceedings of the National Academy of Sciences of the United States of*
*America* 115: E9449-E9458
Zhang L, Syed ZA, van Dijk Härd I, Lim J-M, Wells L, Ten Hagen KG (2014) O-
Glycosylation regulates polarized secretion by modulating Tango1 stability. *Proceedings of*
*the National Academy of Sciences* 111: 7296-7301
24
25
26

1  **Tables**

3  **Table 1**: Treatments, concentrations, manufacturer, catalogue number, and lot number
4

| Treatment | Concentration | Duration | Manufacturer, catalogue number, lot number |
|---|---|---|---|
| **Brefeldin A** | 1, 10, 100 µg/mL | 45 minutes | Sigma, Cat no. B6542-5MG, Lot # 029M4075V |
| **Monensin** | 1 mM | 45 minutes | Sigma, Cat no. M5273-500MG, Lot # SLBK4090V |
| **TGF-β1** | 10 ng/mL | 24 hours | PeproTech, Cat no. 100-21, Lot #1218209 |
| **IL-1β** | 10 ng/mL | 24 hours | BioLegend, Cat no. 00000575102, Lot #B262053 |
| **Galunisertib** | 10 µM | 24 hours | MedChemExpress, Cat no. HY-13226, Batch #81834 |
| **SB431542** | 10 µM | 24 hours | R&D Systems, Cat no. 1614/50, Lot #14A1236410 |
| **Thapsigargin** | 1 µM | 24 hours | Invitrogen, Cat no. T7458, Lot # 2186060 |

**Figures**

**Figure 1: Mucin-2 transport in large COPII vesicles is promoted by KLHL12 and facilitated by**
**TANGO1.** A) A schematic representation of COPII formation for regular (top) and large (bottom) sized
vesicles, showing the involvement of TANGO1, cTAGE5, and KLHL12. Created with BioRender.com.
B) Western blot of post-Golgi mucin-2 after silencing of TANGO1, cTAGE5, and KLHL12. C) Semi-
quantification of the Western Blot using ImageJ, comparing the signal intensity of mucin-2 to the
signal intensity of β-actin, indicating a statistically significant decrease of post-Golgi mucin-2 levels
upon silencing of TANGO1, cTAGE5, and KLHL12. D) Western blot of SEC31A after silencing of
TANGO1, cTAGE5, and KLHL12. E) Semi-quantification of the Western Blot using ImageJ,
comparing the signal intensity of SEC31A to the signal intensity of β-actin, indicating a statistically
significant decrease of SEC31A expression upon silencing of TANGO1 and KLHL12, and similar
expression after silencing cTAGE5. F) Western blot of mucin-2 upon overexpression of TANGO1 and
KLHL12. G) Semi-quantification of the Western Blot using ImageJ, comparing the signal intensity of

mucin-2 to the signal intensity of β-actin, indicating a statistically significant increase of post-Golgi
mucin-2 levels upon overexpression of KLHL12 and a non-significant increase by overexpression of
the combination of KLHL12 and TANGO1. H) Immunofluorescent staining of mucin-2 (green),
TANGO1 (red), and nuclei (blue) showing partial colocalization of mucin-2 and TANGO1, as indicated
by the white arrows. The yellow arrow indicates mucin-2 and TANGO1 colocalization further from the
ER. Scale bar represents 10 µm. I) Immunofluorescence of mucin-2 (green), KLHL12-FLAG (red),
and nuclei (blue), showing colocalization of mucin-2 and KLHL12-FLAG as indicated by the arrows.
Scale bar represents 10 µm. Data was analyzed using one-way ANOVA followed by Bonferroni's test.
Significance was indicated as * ($p < 0.05$); ** ($p < 0.01$); *** ($p < 0.001$) with respect to Control.

**Figure 2: TGF-β facilitates mucin-2 trafficking through TANGO1.** A) Schematic representation of the interaction of TGF-β1 with TANGO1, COPII, IL-1β, NF-κB, and MUC2. Created with BioRender.com. B) Western blot of TANGO1 after stimulation with TGF-β1 and IL-1β. C) Semi-quantification of the Western Blot using ImageJ, comparing the signal intensity of TANGO1 to the signal intensity of β-actin, indicating a statistically significant increase in TANGO1 expression upon addition of TGF-β1, and the combination of TGF-β1 and IL-1β. D) Western blot of post-Golgi mucin-2 after stimulation with TGF-β1 and IL-1β. E) Semi-quantification of the Western Blot using ImageJ, comparing the signal intensity of mucin-2 to the signal intensity of β-actin, indicating a statistically significant decrease of post-Golgi mucin-2 levels upon addition of TGF-β1, and the combination of TGF-β1 and IL-1β. F) Western blot of TANGO1 upon inhibiting TGF-β1 using Galunisertib and SB431542. G) Semi-quantification of the Western Blot using ImageJ, comparing the signal intensity of TANGO1 to the signal intensity of β-actin, indicating a statistically significant decrease of TANGO1 expression upon inhibition of TGF-β1 using Galunisertib and SB431542. H) Western blot of post-Golgi mucin-2 after inhibition of TGF-β1 using Galunisertib and SB431542. I) Semi-quantification of the Western Blot using ImageJ, comparing the signal intensity of mucin-2 to the signal intensity of β-actin, indicating a statistically significant decrease of post-Golgi mucin-2 levels upon inhibition of TGF-β1 using Galunisertib and SB431542. J-O) mRNA levels of MUC2, TANGO1, and NF-κB, normalized to GAPDH and control group using the $2^{-\Delta\Delta Ct}$ method determined with RT-qPCR. J) Decreased MUC2 mRNA levels upon addition of TGF-β1, and the combination of TGF-β1 and IL-1β. K) Decreased MUC2 mRNA levels upon inhibition of TGF-β1 using Galunisertib and SB431542. L) Increased NF-κB mRNA levels upon addition of IL-1β and TGF-β1 and IL-1β. M) Increased NF-κB mRNA levels upon inhibition of TGF-β1 through Galunisertib. N) TANGO1 mRNA levels after stimulation with TGF-β1 and IL-1β show no significant difference. O) TANGO1 mRNA levels upon inhibition of TGF-β1,

1    showing significantly increased TANGO1 expression for the combination of Galunisertib and
2    SB431542. Data was analyzed using one-way ANOVA followed by Bonferroni's test. Significance was
3    indicated as * ($p < 0.05$); ** ($p < 0.01$); *** ($p < 0.001$) with respect to Control.

**Figure 3: ER stress is increased by inhibiting the formation of large COPII vesicles.** A) Schematic representation of the Unfolded Protein Response as measure of ER stress. The UPR consist of three pathways, namely ATF6, PERK, and IRE1. From each pathway, at least one target has been examined on mRNA level to indicate ER stress. Created with BioRender.com. B) ER stress is induced through silencing the formation of large COPII vesicles results in increased levels of ER stress in the CHOP, BiP, ATF4, and EDEM mRNA levels, but not by splicing of XBP1. C) ER stress is not induced by overexpression of TANGO1 and KLHL12 in sXBP1, ATF4, and EDEM. Elevated levels of CHOP and BiP are observed for the co-transfection of TANGO1 and KLHL12. D) ER stress as a result of exposure to TGF-β1 and IL-1β only shows elevated levels of BiP, yet other pathways are not altered as expected. E) ER stress is elevated upon inhibition of TGF-β1 by Galunisertib and SB431542. F) Thapsigargin upregulates TANGO1 mRNA levels. G) Silencing of SEC31A, TANGO1, KLHL12, and cTAGE5 upregulates TANGO1 mRNA levels. Data was analyzed using two-way ANOVA followed by Bonferroni's test. Significance was indicated as * (p < 0.05); ** (p < 0.01); *** (p <0.001) with respect to Control of each specific gene tested.

**Figure 4: ER stress gene expression is increased in intestinal tissue from patients with**
**ulcerative colitis (UC) and Crohn's disease (CD).** Intestinal biopsies from patients with IBD
diagnosed with either UC or CD were sequenced and gene expression levels of ER stress markers
XBP1, CHOP, BiP, ATF4 and EDEM1 (A-E) were determined. Gene expressions were compared
between group of individuals with CD, UC and healthy controls using Mann-Whitney U-tests followed
by post-hoc Bonferroni correction for multiple testing. Significance was indicated as * ($p < 0.05$); ** ($p
< 0.01$); *** ($p < 0.001$) with respect to healthy controls.

**Figure 5: Gene expression data from patients with IBD possibly indicates an anti-inflammatory role of TGF-β and its regulatory effect on mucin-2 transcription and transport.** Intestinal biopsies from an IBD patient cohort diagnosed with either UC or CD were sequenced and gene expression levels of mucin-2 and associated ER-to-Golgi transport factors in connection to TGF-β1 were evaluated. Patient data revealed elevated levels of MUC2 (A) along with genes involved in COPII-facilitated transport SEC31A (B), cTAGE5 (C) and TANGO1 (D). KLHL12 showed expression comparable to healthy controls. Upregulation of TGF-β1 (F) during UC and CD might divulge (reveal?) its role in promoting mucosal barrier function through amplified mucin-2 transcription and trafficking. Gene expressions were compared between group of individuals with CD, UC and healthy controls using Mann-Whitney U-tests followed by post-hoc Bonferroni correction for multiple testing. Significance was indicated as * (p < 0.05); ** (p < 0.01); *** (p <0.001) with respect to healthy controls.

Supplementary Figures

**Supplementary Figure 1:** ATP production as measure of viability of HT29-MTX cells upon addition of
BFA and Monensin. A) Concentrations of BFA and Monensin used to determine conventional
secretion of MUC2 (10 µg/mL of BFA and 1 µM of Monensin). B) Concentrations of BFA used to
determine BFA-dose dependency (25 µg/mL; 50 µg/mL; 100 µg/mL). Staurosporin (10 µM) was used
as a negative control. Data was analyzed using one-way ANOVA followed by Bonferroni's test.
Significance was indicated as * ($p < 0.05$); ** ($p < 0.01$); *** ($p < 0.001$) with respect to 0% FBS.

**Supplementary Figure 2: Mucin-2 is trafficked in a COPII-dependent manner.** A) A schematic representation of the effect of Brefeldin A (BFA) and Monensin (Mon) on conventional protein secretion. Created with BioRender.com. B) Western Blot of post-Golgi mucin-2 as a result of BFA and Mon exposure for 45 minutes. C) Semi-quantification of the Western Blot using ImageJ, comparing the signal intensity of mucin-2 to the signal intensity of β-actin, indicating a statistically significant decrease of post-Golgi mucin-2 upon addition of BFA and Mon. D) Western Blot showing BFA dose dependency on post-Golgi mucin-2 levels. E) Semi-quantification of the Western Blot using ImageJ, comparing the signal intensity of mucin-2 to the signal intensity of β-actin, indicating a statistically significant decrease of post-Golgi mucin-2 secretion upon addition of different BFA concentrations. F) Western blot of silencing of COPII protein SEC31A showing decreased post-Golgi mucin-2 levels compared to controls, including non-targeting siRNA and transfection agent DharmaFect. G) Semi-quantification of the Western Blot using ImageJ, comparing the signal intensity of mucin-2 to the signal intensity of β-actin, indicating a statistically significant decrease of post-Golgi mucin-2 levels upon silencing of outer COPII protein SEC31A. H) Immunofluorescence of mucin-2 (green), SEC31A (red), and nuclei (blue), showing a partial colocalization of mucin-2 and SEC31A. Scale bar represents 10 µm. Data was analyzed using one-way ANOVA followed by Bonferroni's test. Significance was indicated as * (p < 0.05); ** (p < 0.01); *** (p <0.001) with respect to Control (C and G) or 0 µM BFA (E).

**Supplementary Figure 3:** Silencing efficiency of SEC31A was determined at 39.72 ± 0.01882%. A) Western Blot data of SEC31A and β-actin as loading control for the control cells, the effect of DharmaFect, the effect of non-targeting siRNA, and the effect of SEC31A siRNA. B) Quantification of SEC31A expression using ImageJ, normalized to β-actin and control. Data was analyzed using one-way ANOVA followed by Bonferroni's test. Significance was indicated as * ($p < 0.05$); ** ($p < 0.01$); *** ($p < 0.001$) with respect to Control.

**Supplementary Figure 4:** Silencing efficiency of TANGO1 was determined at 44.84 ± 0.085%. A) Western Blot data of TANGO1 and β-actin as loading control for the control cells, the effect of DharmaFect, the effect of non-targeting siRNA, and the effect of TANGO1 siRNA. B) Quantification of TANGO1 expression using ImageJ, normalized to β-actin and control. Data was analyzed using one-way ANOVA followed by Bonferroni's test. Significance was indicated as * (p < 0.05); ** (p < 0.01); *** (p <0.001) with respect to Control.

3 **Supplementary Figure 5**: Transfection efficiency of TANGO1-FLAG and KLHL12-FLAG. A-B)
4 Immunofluorescence of F-actin (Magenta), and nuclei (blue) and A) TANGO1-FLAG (green) and B)
5 KLHL12-FLAG (green). Scale bar represents 50 µm. C) Transfection efficiency (%) of 80% for
6 TANGO1-FLAG and 35% for KLHL12-FLAG. D) TANGO1 mRNA levels in transfected conditions.

2 **Supplementary Figure 6**: Toxic aggregation of KLHL12-FLAG by transfecting 1 µg. A)
3 Immunofluorescent imaging of KLHL12-FLAG (Magenta) and nuclei (blue), showing large aggregates.
4 Scale bar 100 µm. B) TANGO predicted beta sheet aggregation propensity indicating the aggregation
5 of KLHL12 (Fernandez-Escamilla *et al.*, 2004; Linding *et al.*, 2004; Rousseau *et al.*, 2006).

**Supplementary Figure 7**: Response of HT29-MTX to different serum conditions. A-C) Immunofluorescence staining of HT29-MTX cells with MUC2 (green), F-actin (magenta), and nuclei (blue) for 10%, 0%, and 0.5% FBS for 24 hours respectively. Scale bar 50 μm. D) ATP production of HT29-MTX upon 10% FBS as positive control, and 0% and 0.5% FBS for 24h and 48h. Staurosporin (10μM) is used as negative control. Data was analyzed using one-way ANOVA followed by Bonferroni's test. Significance was indicated as * (p < 0.05); ** (p < 0.01); *** (p <0.001) with respect to Control.

2  **Supplementary Figure 8:** ER stress is measured by quantifying mRNA levels of sXPB1, CHOP, BiP,
3  ATF4, and EDEM, as shown by stimulation with ER stressor Thapsigargin. Data was analyzed using
4  one-way ANOVA followed by Bonferroni's test. Significance was indicated as * ($p < 0.05$); ** ($p <$
5  $0.01$); *** ($p < 0.001$) with respect to Control.

1 **Supplementary Table 1**: Primer sets for RT-qPCR

| Gene of interest | Primer sequence (5' → 3') | Reference |
|---|---|---|
| **GAPDH fwd**<br>**GAPDH rev** | CTCTGCTCCTCCTGTTCGAC<br>TTAAAAGCAGCCCTGGTGAC | (Maiers *et al.*, 2017) |
| **NFKB fwd**<br>**NFKB rev** | GTGGTGCCTCACTGCTAACT<br>GGATGCACTTCAGCTTCTGT | (Domińska *et al*, 2017) |
| **MUC2 fwd**<br>**MUC2 rev** | AAGACGGCACCTACCTCG<br>TTGGAGGAATAAACTGGAGAACC | (Huang *et al*, 2019) |
| **TANGO1 fwd**<br>**TANGO1 rev** | GAACCTGAACCTGAACCAGTAG<br>GCTCTCCCTGAGCATGATTT | (Maiers *et al.*, 2017) |
| **sXBP1 fwd**<br>**sXBP1 rev** | CTGAGTCCGAATCAGGTGCAG<br>ATCCATGGGGAGATGTTCTGG | (Oslowski & Urano, 2011) |
| **ATF4 fwd**<br>**ATF4 rev** | GTTCTCCAGCGACAAGGCTA<br>ATCCTGCTTGCTGTTGTTGG | (Oslowski & Urano, 2011) |
| **CHOP fwd**<br>**CHOP rev** | AGAACCAGGAAACGGAAACAGA<br>TCTCCTTCATGCGCTGCTTT | (Oslowski & Urano, 2011) |
| **BiP fwd**<br>**BiP rev** | TGTTCAACCAATTATCAGCAAACTC<br>TTCTGCTGTATCCTCTTCACCAGT | (Oslowski & Urano, 2011) |
| **EDEM fwd**<br>**EDEM rev** | CAAGTGTGGGTACGCCACG<br>AAAGAAGCTCTCCATCCGGTC | (Oslowski & Urano, 2011) |