TITLE

Kynurenine monooxygenase blockade reduces endometriosis-like lesions, improves visceral hyperalgesia, and rescues mice from a negative behavioural phenotype in experimental endometriosis.

AUTHORS

Ben Higgins[1], Ioannis Simitsidellis[1,2,3], Xiaozhong Zheng[1], Frances Collins[1,2,3], Natalie Z.M. Homer[4, 8], Scott G. Denham[4], Joanna P. Simpson[4], Mike Millar[5], Lyndsey Boswell[5], Hee Y. Lee[6], Yeon G. Kim[6], Kyung H. Park[6], Larry C. Park[6], Patrick J. Sweeney[6], Gerard Feraille[7], Alessandro Taddei[7], David Chagras[7], Thierry Alvarez[7], Scott P. Webster[8], Andrew Horne[†2,3], Philippa T.K. Saunders[†2,3], and Damian J. Mole[†*1,9].

AFFILIATIONS

[1] Centre for Inflammation Research, Institute for Regeneration and Repair, Edinburgh Bioquarter, The University of Edinburgh, UK.

[2] EXPPECT Edinburgh, Institute for Regeneration and Repair, Edinburgh Bioquarter, The University of Edinburgh, UK.

[3] MRC Centre for Reproductive Health, Institute for Regeneration and Repair, Edinburgh Bioquarter, The University of Edinburgh, UK.

[4] Mass Spectrometry Core, Edinburgh Clinical Research Facility, The University of Edinburgh, UK.

[5] SuRF Molecular Histology Facility, Queen's Medical Research Institute, The University of Edinburgh, UK.

[6] Naason Science, Inc., Republic of Korea.

[7] Syneos Health France, Les Templiers, 2400 route des Colles, 06410 Biot, Sophia-Antipolis, France.

[8] Centre for Cardiovascular Science, Queen's Medical Research Institute, The University of Edinburgh, UK

[9] Clinical Surgery, The University of Edinburgh, UK

*Corresponding Author and Lead Contact: Damian J Mole. ORCID 0000-0001-6884-7302; Email: damian.mole@ed.ac.uk

†Indicates senior authors.

SUMMARY

Endometriosis is a common and debilitating neuro-inflammatory disorder that is associated with chronic pain. Definitive diagnosis is based on the presence of endometrial-like tissue (lesions) in sites outside the uterus. Kynurenine monooxygenase (KMO) is a mitochondrial enzyme of tryptophan metabolism that regulates inflammation and immunity. Here, we show that KMO is expressed in epithelial cells in human endometriosis tissue lesions and in corresponding lesions in a mouse model of endometriosis. In mice, oral treatment with the potent KMO inhibitor KNS898 induced a biochemical state of KMO blockade with accumulation of kynurenine, diversion to kynurenic acid and ablation of 3-hydroxykynurenine production. In the mouse model of endometriosis, KMO inhibition improved histological outcomes and endometriosis pain-like behaviours, even when KNS898 treatment commenced one week after initiation of lesions. Taken together, these results suggest that KMO blockade is a promising new non-hormonal therapeutic modality for endometriosis.

143 words.

KEYWORDS

Kynurenine monooxygenase; Endometriosis; Drug discovery; Non-hormonal therapies; Visceral hyperalgesia

INTRODUCTION

Endometriosis is a life-altering condition that affects approximately 10% of females. It is an oestrogen-dependent neuroinflammatory disorder associated with debilitating pelvic pain, excessive fatigue, gastrointestinal and urinary symptoms, and infertility[1]. Worldwide, 200 million prevalent cases are forecast by 2026. Endometriosis is defined by the presence of endometrial-like tissue ('lesions') outside the uterus. Physiological hormonal fluctuations in women induce cyclical episodes of cell proliferation, inflammation, injury, and repair within lesions that favour fibroblast to myofibroblast differentiation and fibrosis[1]. We and others have identified metabolic dysfunction in cells associated with development of endometriosis lesions[2]. At present, therapeutic options are largely limited to surgery (that often needs to be repeated) or medical therapies that target hormonal activity with resultant side effects (block conception, menopausal symptoms)[1]. Patient surveys consistently show frustration with the lack of available treatments that can give long term relief from symptoms including pain, low mood and bloating[3]. Analysis of recent clinical trials directed at endometriosis[1] has highlighted an unmet need for new, non-hormonal approaches to symptom relief. The studies in the current paper have addressed this need by focusing on an enzyme that is known to play a key role in inflammatory processes that are implicated in the aetiology of endometriosis, but which has not previously been investigated as a target.

Our proposed solution to this unmet medical need is by targeting the enzyme kynurenine 3-monooxygenase (KMO). KMO is a critical regulator of inflammation at multiple organ sites that acts by altering metabolic flux through the kynurenine pathway of tryptophan metabolism[4]. KMO is known to be expressed in non-pathological endometrium[5], but whether KMO is over-expressed in endometriosis lesions and linked to the severity of inflammation remains to be determined. KMO has been identified as a critical step in converting kynurenine to the cytotoxic metabolite, 3-hydroxykynurenine (3HK), that is an oxidative stressor, causes protein cross-linking, and regulates the immune-metabolic interface[4]. Although there is no specific information about a direct role of KMO in endometriosis, there is evidence of dysregulated tryptophan metabolism in a recent study using a preclinical non-human primate model of endometriosis[6], and increased kynurenine pathway flux at the immune-metabolic interface between stromal cells and NK immune cells in endometriosis lesions[7].

At present, there is a scientific rationale for KMO inhibition in endometriosis, but it remains to be shown in preclinical experiments whether KMO inhibition is efficacious in decreasing lesion volume or behavioural symptoms which are used as a surrogate for pain responses in model systems. KNS898 is a highly specific small molecule KMO inhibitor with potential for use by women with endometriosis, based on favourable characteristics for oral development in terms of bioavailability and predicted half-life[8,9]. KNS898 is a competitive inhibitor of kynurenine substrate at the active site of KMO with a pIC50 of 8.8[8-10]. We propose that KMO inhibition is a novel therapeutic strategy for endometriosis and, if successful, we will make a significant positive impact for women with this painful, disabling condition. The aim of this project was to obtain proof-of-concept for KMO inhibition as a novel therapy for endometriosis. Specifically, we sought to explore the expression of KMO in biobanked human endometrial and endometriosis lesion tissues, confirm target inhibition of KMO by KNS898 in mice, and define the preclinical efficacy of KNS898 in improving clinical features of disease (specifically hyperalgesia and altered cage behaviour) and reducing endometriosis lesion volume in an experimental mouse model of endometriosis.

RESULTS

KMO is expressed in human eutopic endometrium and human endometriosis tissue lesions. To explore whether we could detect variations between expression of KMO in endometrium (eutopic) within the uterus and a variety of lesions obtained from patients, we conducted detailed immunohistochemistry with a primary antibody specific for KMO. On fixed tissue sections of normal human endometrium KMO expression was most striking in epithelial cells lining the glands (Figure 1a, insert B) with lower levels in the luminal layer (insert C). Notably expression in the glands was not uniform (**Figure 1a**). KMO was also strongly immunopositive in human peritoneal endometriosis lesions (**Fig. 1d to Fig. 1g**), and evidently mostly localised to the epithelial tissues surrounding the distended endometrial gland-like structures (DEGLS) (**Fig. 1e and Fig. 1g**). Expression in the stromal compartment appeared variable. In human ovarian endometriosis lesions, KMO was present at low expression levels in the mesothelial layers (**Fig. 1h and Fig. 1i**). Duplex immunohistochemistry with cell phenotype markers CD68 (macrophages) did not show KMO colocalising with these immune cells (data not shown)

Oral KNS898 inhibits KMO in mice. Next, we established that oral dosing of KNS898 by gavage in mice resulted in inhibition of KMO. Using n=3 mice per group, we administered KNS898 at 0.01 mg/kg, 5 mg/kg, and 25 mg/kg twice daily (b.d.) in vehicle for seven days, as described. Plasma drug levels and metabolite concentrations are shown in **Figure 2.** KNS898 dosed at 0.01 mg/kg b.d. resulted in a mean ($\pm$ S.E.M.) plasma drug level of 0.18 ± 0.01 ng/mL, 5 mg/kg resulted in 88.8 ± 22.6 µg/mL, and 25 mg/kg gave 483.9 ± 84.0 µg/mL. The difference between groups was statistically significant by one-way ANOVA with post hoc Tukey's test ($P = 0.001$) (**Fig.2a**). KMO blockade with KNS898 was clearly measurable. A backlog in the KMO substrate KYN was evident: KNS898 dosed at 0.01 mg/kg b.d. resulted in a mean ($\pm$ S.E.M.) plasma level of KYN of 339 ± 39 ng/mL, 5 mg/kg resulted in 4940 ± 483 ng/mL, and 25 mg/kg gave 3682 ± 634 ng/mL. The difference between groups was statistically significant by one-way ANOVA with post hoc Tukey's test ($P = 0.001$). The increase in KYN at maximal inhibition was approximately 14-fold compared to the level seen after KNS898 0.01 mg/kg (**Fig.2b**). Excess KYN was metabolised to KA by kynurenine aminotransferase: KNS898 dosed at 0.01 mg/kg b.d. resulted in a mean ($\pm$ S.E.M.) plasma level of KA of 629 ± 103 ng/mL, 5 mg/kg resulted in 14399 ± 3394 ng/mL, and 25 mg/kg gave 15965 ± 789 ng/mL. The difference between groups was statistically significant by one-way ANOVA with post hoc Tukey's test ($P = 0.001$). The fold increase in KA at maximal inhibition was approximately 25-fold compared to the level seen after KNS898 0.01 mg/kg (**Fig.2c**). KMO blockade resulted in a statistically-significant reduction of 3HK in plasma: KNS898 dosed at 0.01 mg/kg b.d. resulted in a mean ($\pm$ S.E.M.) plasma level of 3HK of 27.7 ± 7.2 ng/mL, 5 mg/kg resulted in 4.2 ± 0.3 ng/mL, and 25 mg/kg gave 0.9 ± 0.4 ng/mL. The difference between groups was statistically significant by one-way ANOVA with post hoc Tukey's test ($P = 0.001$). The fold decrease in 3HK at maximal inhibition was approximately 30-fold compared to the level seen after KNS898 0.01 mg/kg (**Fig.2d**). Overall, there was a clear dose response to KNS898 administration leading to maximal KMO blockade at 25 mg/kg b.d. This dose was therefore selected for efficacy experiments going forward. A diagrammatic representation of the kynurenine pathway is shown as Figure 2e for reference.

KMO blockade reduces endometrial gland-like lesion burden in experimental endometriosis in mice. The
experimental design for the mouse model of endometriosis is shown in **Figure 3a.** The pharmacological effect
of KNS898 therapy showed appropriate levels of KNS898 detected in plasma (**Fig 3b**), with accumulation of
kynurenine (**Fig. 3c**), inhibition of 3HK production (**Fig. 3d**) and diverted metabolism of accumulated
kynurenine to kynurenic acid (**Fig. 3e**). All recipient mice inoculated with donor tissue (groups G3, G4, and G5)
developed distended endometrial gland-like structures (DEGLS). The incidence of DEGLS formation was
enumerated at autopsy, and the axial length of each DEGLS was measured after excision from the surrounding
tissue. In G3 (endometriosis + vehicle), 8 of 15 (53%) of the inoculated animals had developed DEGLS. In
KNS898-treated group G4 (endometriosis + treatment from Day 19), DEGLS formed in 4 of 15 mice (26.7%)
and in G5 (Endo + treatment start on Day 26) in 6 of 15 mice (40%). As expected, no DEGLS were formed in
the non-inoculated control and sham groups. The total number of DEGLS per animal in each group was highest
in G3 with an average of 4.0 per animal with DEGLS (total = 32 DEGLS in 8 mice in G3). Mice with
endometriosis receiving KNS898 from the time of inoculation (G4) had an average of 2.0 DEGLS per animal
with DEGLS (total = 8 DEGLS in 4 mice in G4) and those receiving KNS898 1 week after inoculation (G5) had
an average of 1.8 DEGLS per animal (total = 11 DEGLS in 6 mice in G5) (**Figs. 3f and 3g**). Statistical analysis
by ANOVA showed a significant difference in endometriosis DEGLS burden between groups (P = 0.0295 for
DEGLS per animal; P = 0.004 for DEGLS per group). DEGLS axial length and derived volume did not differ
between groups (**Supplementary Fig. 1a and b**). All recipient mice inoculated with donor tissue lost body
weight following inoculation which then gradually recovered. After recovery, body weight of all three
inoculated groups was lower compared to the control groups for the duration of the study. Overall, there was no
significant difference in body weight between G3 and the KNS898 treatment groups G4 and G5 (endometriosis
+ treatment from Day 26) (**Suppl. Fig. 1c**).

KMO is expressed in experimental endometriosis in mice. Histological examination of DEGLS identified
them as containing cystic structures lined with epithelial layers identifiable as columnar epithelium,
pseudostratified epithelium, squamous epithelium, and cuboidal epithelium, with goblet cells. These DEGLS
were considered to represent endometriosis-like lesions derived from the implanted basal
endometrial/myoepithelial layers of the donor mice uteri (**Suppl. Fig 2**). Immunohistochemistry using an
antibody to KMO showed KMO protein expression localised mainly to the epithelial cells lining of the DEGLS,
with a lesser degree of KMO positive staining in the closest surrounding connective tissue, in keeping with the
previously observed KMO expression pattern in human endometriosis lesion tissue (**Fig. 4a and 4d and
Supplementary Fig S3**). The thickness (area divided by length) of the KMO positive epithelial layer was
quantified for each DEGLS section using QuPath and there was no difference between groups G3, G4 and G5
(**Fig. 4b and 4e**). However, quantification of KMO expression confirmed the high intensity of KMO staining in
the epithelial lining layers (**Fig. 4c and 4f**), but also showed a clear and statistically significant reduction in
KMO expression intensity in those areas in DEGLS removed from mice treated with the KMO inhibitor
KNS898 (**Fig. 4g**; P = 0.008). Representative micrographs from each of groups G3, G4 and G5 are presented in
Figures 4h, 4i and 4j.

KMO inhibition reduces mechanical allodynia in experimental endometriosis. Clinical endometriosis is

associated with visceral hyperalgesia and central sensitisation to pain[11]. Visceral and central hyperalgesia may

be tested in rodents using the Von Frey filament test[12]. Baseline reaction values for hind paw and bladder Von

Frey tests showed no significant difference in mechanical allodynia before inoculation. In established

endometriosis without treatment (group G3), the mechanical allodynia threshold in the hind paw was

statistically significantly lower compared to baseline for the group. When compared to the control groups at the

corresponding time point beginning 1 week after inoculation and continuing until the end of the study. Day 26

KNS898-treated group (G5) showed a statistically-significant improvement in mechanical allodynia in the hind

paw using the Von Frey test compared to mice in G3 with untreated endometriosis given vehicle control (Two-

way ANOVA, Group effect P = 0.003, time effect P < 0.0001; Dunnett's multiple comparison test G5 vs G3

P=0.001) (**Fig. 5a**). The mechanical allodynia threshold for the bladder reflex also was lower in mice with

endometriosis compared to baseline throughout the study, and KNS898 treatment commencing at D26 (G5) was

associated with a statistically significant improvement in bladder mechanical allodynia threshold at D42

compared to mice with untreated endometriosis given vehicle control (G3)(Two-way ANOVA, Group effect P =

0.038, time effect P < 0.001; Dunnett's multiple comparison test G5 vs G3 P=0.021)(**Fig. 5b**).

KMO inhibition rescues impaired cage exploration behaviour and mobility in mice with endometriosis.

HCA peripheral moving speed, time at cage edge, and illness behaviour, including temperature, motility and

cage exploration behaviour was quantified using Home Cage Analysis (HCA). Baseline HCA was recorded

before inoculation and at the end of the experiment. Mice with endometriosis without treatment showed an

overall reduction in activity in moving distance and moving speed relative to baseline, and compared to sham-

operated control groups, indicating a negative effect on behaviour due to endometriosis. Importantly, mice with

endometriosis treated with KNS898 showed marked improvement in motility and cage exploration behaviour

compared to untreated endometriosis mice, and although this difference between groups was statistically

significant by Welch's one-way ANOVA, post hoc testing (Dunnet's T3) was not significant between groups.

This qualitative difference of time spent exploring the periphery of the cage being lower for mice with

endometriosis treated with vehicle control was seen in both the day and night phases (**Figs. 5c and 5d**). A

similar improvement with KNS898 treatment compared to vehicle control mice with endometriosis was seen for

total moving distance, total moving speed, and peripheral distance, but not for total moving or climbing time,

for which no difference between groups was detected. Together, these data indicate that KMO inhibition with

KNS898 results in an improvement in well-being evidenced by improved cage exploration behaviour in addition

to improved objective histological measures of endometriosis disease burden.

DISCUSSION

In this study, we set out to investigate the potential for KMO inhibition as a non-hormonal therapy for

endometriosis. First, we confirmed that KMO was expressed in human endometrium by immunohistochemistry,

and then showed that KMO was clearly expressed in the epithelial cells in human endometriosis lesions. Next,

we demonstrated that the highly specific KMO inhibitor KNS898 was orally bioavailable when given twice

daily by gavage in mice, and clearly blocked KMO activity in a dose-dependent manner at a dose of 25mg/kg.

We therefore used that dose to test the efficacy of KMO blockade with KNS898 in mice with experimentally-

induced endometriosis. One important finding of this project is that KMO blockade resulted in a reduction in
endometriosis severity compared to untreated mice with endometriosis, specifically in terms of reducing i) the
number of mice that developed endometriosis tissue lesions, and ii) the number of lesions per mouse in those
that did develop lesions. The histopathology of the experimental endometriosis lesions was sufficiently similar
macroscopically to that seen in the examined human DEGLS, and KMO was evidently highly expressed in the
same tissue distribution in model lesions compared to human disease. KMO blockade also decreased lesion
KMO expression. Critically, and importantly from a translational perspective, therapeutic blockade of KMO
improved visceral hyperalgesia measured by reduced mechanical allodynia and restored normal cage
exploration behaviour and mobility in treated mice compared to untreated mice with endometriosis. Together,
these data show that KMO is expressed in human and mouse endometriosis tissue lesions and that therapeutic
KMO blockade reduces the number of endometriosis lesions and improves holistic metrics of disease behaviour
in mice.

The model of endometriosis, using inoculation of endometrial tissue of ovariectomized donor mice,
reliably induced the pathophysiological symptoms indicative of endometriosis in recipient mice. Test groups
inoculated with endometrial tissue (G3-G5) showed significant growth of ectopic endometrial tissue. Groups
treated with test article experienced significantly less DEGLS development (significantly fewer DEGLS were
noted in treated groups when compared to vehicle treated groups). Disease burden in the treatment group that
had treatment starting immediately after inoculation was lower than that of the vehicle-only treated group. It is
not clear why mean cystic size and cystic volume in treated animals was not smaller in treated animals. We can
only speculate that KMO blockade may potentiate rapid involution of cysts, but this cannot be proven
mechanistically here.

Mice that received inoculated endometrial tissue showed a measurable and increased visceral
hyperalgic pain response (lower mechanical threshold) as measured by bladder response to von Frey filament
testing, and improvement in a surrogate marker of central sensitisation to pain measured by hind paw Von Frey
filament testing when compared to control mice. The mechanical threshold of both treatment groups trended
higher compared to the vehicle treated group. One interpretation of these data is that KMO inhibition reduced
responses to pain caused by the presence of endometriosis, i.e. improving visceral hyperalgesia.

Home cage behaviour using HCA indicated a reduced overall activity in endo-inoculated mice when
compared to control mice. It should also be noted that test article-treated animals in both groups showed more
locomotor behaviour and a seemingly better quality of life within the home cage environment when the home-
cage dynamics were monitored. This supports the notion that treated mice exhibit less propensity for behaviours
that are, at times, typical of depressive and anxiety-like behaviour in home-cage, group housed conditions.

Under the experimental conditions imposed, treatment with KNS898 on two dosing schedules provided
a significant reduction in DEGLS formation within the inoculated mice, as well as a seemingly higher pain
threshold. This attenuation of pain response was coupled with increased activity levels in the home cage. Taken
together, these data suggest a therapeutic effect to alleviation of certain salient symptoms of endometriosis, as
well as a reduction in the number and size of DEGLS.

Non-pathological endometrium is a site of high KMO expression. Because endometriosis lesions in
women are 'endometrial-like' tissue rather than normal endometrium, we tested, and demonstrated expression of
KMO in the epithelial layers of endometriosis lesions sampled from women undergoing surgery for

endometriosis. First-line medical treatment for endometriosis is the contraceptive pill or other ovarian steroid
hormone suppressive drugs. Treatment failures are frequent, side effects are common, all approaches are
contraceptive. Many women opt for invasive surgery to remove or ablate the endometriosis lesions.

In conclusion, KMO is expressed in human endometriosis tissue lesions and in a mouse model of
endometriosis in the epithelial layers of distended endometrial gland-like structures. Oral KNS898 reliably
induced a biochemical state of KMO blockade with accumulation of kynurenine, diversion to kynurenic acid
and ablation of 3-hydroxykynurenine production. KMO blockade improved histological and symptomatic
behavioural endometriosis disease features with an overall benefit, even when treatment commenced one week
after establishment of the disease. KMO blockade is therefore a promising avenue for a new non-hormonal
therapeutic modality for endometriosis.

MATERIALS AND METHODS

Ethical approvals and permissions. The human tissue samples were obtained from participants who had given
fully informed written consent under ethical approval granted by Lothian Research Ethics Committee (LREC
11/AL/0376). Human tissue samples were obtained with ethical approval and fully informed consent from
individuals attending the Royal Infirmary of Edinburgh as described below. Animal experiments conducted by
NAASON Inc were carried out according to the National Institute of Health (NIH) & National Institutes of
Health Korea (NIHK) guidelines for the care and use of laboratory animals and approved by Naason Science in
accordance with all applicable FELASA, IACUC and AAALAC guidelines. Animal experiments outsourced to
Syneos Health were conducted with institutional ethical approval.

Human Patients and Samples. Tissue samples were collected from patients undergoing a diagnostic
laparoscopy for suspected endometriosis following Endometriosis Phenome and Biobanking Harmonisation
Project (EPHect) guidelines[13]. Patient summary characteristics are presented in **Supplementary Table S1**. Note
there was a range of disease stages assigned at time of surgery according to American Fertility Society (AFS)
criteria[14]. Cycle stage was determined by measuring hormones in blood according to standard protocols and
assessment of eutopic endometrial tissue histology when such samples were available[15]. Lesions were recovered
from 17 patients, of these n=10 were recovered from the peritoneal side wall consistent with classification as
superficial peritoneal endometriosis lesions and n=5 from the cysts of ovarian disease (endometrioma) (n=2 on
hormones). Eutopic endometrium was from 4 patients n=3 of which had no lesions at time of surgery (noted as
stage 0). General histology of samples was assessed using H&E staining.

Immunohistochemistry (human endometrium and human endometriosis tissue lesions). 5 µm sections of
formalin-fixed paraffin-embedded tissue blocks were mounted on SuperFrost Plus adhesion slides (Thermo
Fisher Scientific). Sections were deparaffined with xylene and rehydrated prior to heat-induced antigen retrieval
using Instant Pot: Tris-EDTA pH9[16]. Sections were washed with tap water and incubated in phosphate buffered
saline (PBS) for 5 minutes. Endogenous peroxidase was blocked with 0.3% hydrogen peroxide in 70% v/v
methanol for 30 mins at room temperature then washed in PBS prior to blocking in Normal Goat Serum
(NGS)/PBS/bovine serum albumin (BSA)(5%) for 30 mins and streptavidin for 15 mins. Sections were washed

twice in PBS and then blocked with biotin for 15 mins and washed in PBS. The primary antibody to KMO

(KMO Rabbit polyclonal, Proteintech, Catalog Number:10698-1-AP)[17] was diluted to a final concentration of

1:1000 in NGS/PBS/BSA and incubated overnight at 4°C in a humidity chamber. Sections were washed twice

with PBS/Tween 0.05% (1ml Tween in 2L PBS) for 5 mins. The secondary detection antibody Goat Anti-Rabbit

Biotinylated (Vector Cat number: BA-1000) was diluted in NGS/PBS/BSA (1:500) and incubated for 30 mins,

prior to washing twice in PBS/Tween 0.05%, for 5 mins before adding the detection system reagent

Streptavidin-HRP (DAKO Cat Number P0397) 1:500 in PBS for 30 min, washed and stained with DAB (DAKO

Cat Number K3468) as per manufacturer's directions and incubated for 5 mins before a final wash with tap

water. Sections were counterstained with Haematoxylin, dehydrated through graded ethanol and mounted.

Sections of stained slides were scanned on a Zeiss Axioscan Z1 slide scanner and exported as TIFF files: images

were evaluated for stromal, epithelial and immune cell content.

KNS898 preparation for oral administration. KNS898 powder was weighed and dissolved at the required

concentrations in a final vehicle of 2% DMSO, 20% PEG200, 78% 0.15M NaCl by volume. Brief sonication on

ice was done to facilitate disolution.

In vivo confirmation of KMO inhibition by KNS898 in mice. This experiment was outsourced to Syneos

Health (Les Templiers, 2400 route des Colles, 06410 Biot, Sophia-Antipolis, France). A formal

pharmacokinetic/pharmacodynamic study was not required at this stage. Female C5Bl/6J mice aged 12 weeks

were purchased from Charles River Laboratories, maintained on standard 12 hour light-dark cycle, given free

access to water and standard chow before being randomised to one of three dose levels of KNS898 (n=3 per

group, total n=9 mice). Dose levels tested were 0.01mg/kg, 5mg/kg and 25mg/kg. Mice were gavaged with 0.5

mL of drug in vehicle twice daily for 7 days before euthanasia and plasma sampling.

Plasma samples. Blood was sampled into Sarstedt Microvette CB K2EDTA 300 μL tubes and centrifuged at

5,000 rpm (2380 RCF) for 3 mins. Plasma was aliquoted, frozen on dry ice and transferred to storage at -80°C

prior to temperature-controlled shipping.

LC-MS/MS analysis of plasma drug levels and kynurenine metabolites. Plasma samples (100 μL) were

diluted at a 1:1 ratio with 4% phosphoric acid and enriched with 50 ng $^{13}C_6$-kynurenine, $^{13}C_6$-3-

hydroxykynurenine (Sigma Aldrich, custom synthesis) and d5-kynurenic acid (CDN isotopes). 12-point

calibration standards (0.1 to 100 ng) were prepared for KNS898, kynurenine (KYN), kynurenic acid (KA), and

3-hydroxykynurenine (3HK) and extracted alongside samples using solid phase extraction plates (Waters Oasis

HLB, 10 mg sorbent, 30 μm particle size). Extracts were dried down under nitrogen and reconstituted in LC-MS

grade water (100 μL). 10 μL was injected onto a column (Ace C18-PFP column; 100 x 2.1 mm internal diameter

1.7 μm; HiChrom (VWR, Lutterworth)) using an Acquity I-Class UPLC liquid chromatography system (Waters)

linked to a QTRAP 6500+ mass spectrometer (AB Sciex)[10]. The flow rate was set at 0.4 mL/min with a column

temperature of 40°C. Separation was carried out using a gradient mobile phase system of A – 0.1% aqueous

formic acid and B – 0.1% formic acid in methanol, starting at 15%B, rising to 85%B over 6 mins and returning

to 15%B by 9 mins. Mass spectrometry settings were for positive mode electrospray (5.5 kV, 700°C) and

multiple reaction monitoring m/z 209.0 → 192.2 for KYN, m/z 189.9 → 144.1 for KA m/z 225.0 → 208.0 for

3HK and m/z 361.1 → 120.1 for KNS898 and for internal standards were m/z 231.0 → 214.0 for $^{13}C_6$-3HK, m/z

195.1 → 177.2 for d5-KA and m/z 215.0 → 197.8 for $^{13}C_6$-kynurenine. Retention times for KYN, KA, 3HK and

KNS898 were 1.8, 3.2, 1.2 and 6.5 mins, respectively and 3.2 mins for d5KA, 1.8 mins for $^{13}C_6$-3HK and 1.2

mins for $^{13}C_6$-KYN. Data were acquired by Analyst 1.7.1 software (AB Sciex) and linear regression analysis

was carried out on MultiQuant 3.0.3 software (AB Sciex) where peak integrations and amounts of each

kynurenine metabolite and KNS898 were calculated using the peak area ratio of compound/internal standard,

with data further handled in Microsoft Excel 2016 as described[18].

Experimental mouse model of endometriosis. This experiment was outsourced to Naason Science Inc.,

Osong, Korea (KBIO New Drug Development Center #506, Chungbuk, Korea, 28160) using protocols

originally developed by the Saunders team in Edinburgh[19,20]. Experimental design and groups are shown in

Figure 3a and **Supplementary Table S2**. There were 5 groups of mice with n=10-15/group):25 mg/kg KNS898

was administered twice a day via oral gavage in two of the groups of mice. Group 4 received KNS898 from the

time of endometrial tissue inoculation (Day 19; G4); group 5 commenced dosing 1 week after inoculation (Day

26; G5). Group 3 received vehicle (2% DMSO, 20% PEG200, 78% 0.15M NaCl) in the same regimen.

To perform the mouse model of endometriosis, donor female C57Bl/6 mice aged 6 weeks were acclimatized for

2 weeks prior to surgery. Ovariectomy (Day 0) was performed at 8 weeks of age under general anaesthesia with

monitoring, with analgesia that extended to the post-operative period with buprenorphine (0.03 ml) (Veterges

ic® 3 mg/ml, Ceva Inc., Korea) subcutaneously. To prepare donor tissue that would best replicate menstrual-like

tissue in women, ovariectomized (OVX) mice were primed with daily s.c. injections of 100 ng 17β-estradiol

(E2) on days 7, 8 and 9. On days 13 – 19 a silastic progesterone (P4) pellet was implanted subcutaneously.

These animals were injected once daily with E2 (5 ng in sesame oil) on days 13, 14 and 15. Decidualization was

induced in one uterine horn with an injection of 20 µl sesame oil 4 hours after the last E2 injection. On day 19

(4 days after induction of decidual response), donor mice were killed 4 hours after removal of the P4 pellet.

Endometrial tissue was then scraped from the myometrial layer of the decidualized uterine horn, suspended in

500 µl PBS and injected via a XG needle sprayed into the lower abdominal cavity of the recipient mouse under

general anaesthesia with monitoring and post-operative analgesia as described. The ratio of donor to recipient

mouse was 1:1 (from one donor to one recipient). Recipient mice had intact ovaries to ensure ongoing hormonal

stimulation of the injected tissue: group allocations are shown in **Supplementary Table S2**.

Mechanical allodynia test by the Von Frey method. Abdominal and hind paw Von Frey tests were performed

in the recipient animals before inoculation (baseline), and 1, 2, and 3 weeks after inoculation. Mechanical

threshold was measured using Von Frey filaments. For the hind-paw, 15, 8, 6, 4, 2, 1.4, 0.6, 0.4 g filaments were

used, and for the bladder reflex to filament application to the lower abdomen, 60, 26, 10, 8, 6, 4, 2, 1 g filaments

were used. The experimenter was blind to the group allocation in order to reduce bias.

Cage exploration and behavioural assay using Home Cage Analysis. Home Cage Analysis (HCA) was

performed in the recipient animals before inoculation (baseline), and at the late stage of treatment. Recipient

mice were randomly housed using a Monte Carlo randomization. All animals had a micro-chip (BioMark, USA)

inserted to the abdomen prior to being placed in the home-cage. This procedure does not cause undue

discomfort or hamper, in any way, animal movement. Total moving distance, total moving time, moving speed,

isolation/separation distance, isolated time, peripheral time, peripheral distance, in centre zones time, in centre

zones distance, climbing time, and body temperature were tracked automatically by an ActualHCA™ Home

Cage Analyzer (ActualAnalytics Ltd., Edinburgh, UK) and processed with proprietary machine learning and

artificial intelligence algorithms.

Endpoint tissue and plasma sampling. On experimental Day 40, all recipient mice were euthanized, and blood

collected via cardiac puncture. Whole blood was collected into heparinised tubes, and plasma was separated by

centrifugation (3000 rpm for 15 min) at 4°C. Separated plasma was collected in Eppendorf microtubes, frozen

on dry ice and stored at -80°C. Photographs of the abdominal cavity were obtained. Lesions from the abdominal

cavity were harvested. DEGLS were dissected from the surrounding abdominal tissue and measured for size and

volume. DEGLS volume was measured using the following formula[21]: Volume = long diameter × (short

diameter/2)2 × π. DEGLS were fixed in a 4% paraformaldehyde solution and prepared for standard H&E. If

more than one DEGLS was present in an animal, DEGLS that were not used for H&E were snap-frozen in

liquid nitrogen and stored at -80oC.

Endometriosis lesion histology. Endometriosis lesion tissue blocks were sectioned at a uniform thickness of 5

μm and were mounted onto a microscope slide. The slide then underwent deparaffination and hydration.

Paraffin was removed from the slide using xylene, then hydration through graded ethanol and washing were

performed. Slides were then stained with Harris haematoxylin and alcoholic eosin Y and mounted after

dehydration and clearing with xylene. The H&E-stained images were visualized using a Slide Scanner

(Panoramic scan, 3D HISTECH).

Immunohistochemistry (mouse DEGLS). Immunohistochemistry to detect KMO in mouse DEGLS tissue was

performed on a Leica Bond III automated immunostaining robot. 5 μm thick sections obtained from FFPE

(formalin Fixed Paraffin Embedded) samples mounted on superfrost plus slides were stained as follows. Heat

induced epitope retrieval (HIER) was performed using Epitope Retrieval Solution 1 (Leica, ER1 pH 6.0 citrate

based solution) for 20 minutes at 99^0C. Tissue sections were then incubated for 10 minutes in hydrogen peroxide

to block endogenous hydrogen peroxidase activity followed by 10 minute blocking with normal goat serum. The

primary antibody against KMO (Proteintech 10698-1-AP @1:1000 Rabbit)[17] was incubated for 1 hour, then

incubated with a goat anti-rabbit peroxidase conjugated secondary antibody for 30 minutes prior to visualisation

with diaminobenzoate (DAB) using standard protocols.

Digital slide scanning. Whole sections were scanned using a Zeiss Axioscan Z1 whole slide scanner. The image

files (.czi) were batch converted to Tif format for image analysis and quantification, acquisition and batch export

used Carl Zeiss Zen v2.5 software.

Quantitative image analysis. Image analysis was carried out using QuPath v0.4.2. Analysis was standardised

by using multiple regions of interest from individual slides and collating them into a training image. To improve

stain contrast, overlapping DAB and haematoxylin staining was deconvoluted by manual optimisation of the

stain vectors. Individual slide epithelia were annotated, with relative DAB optical densities and annotation shape

measurements taken. Epithelial thickness and area were calculated to enable correlation with KMO intensity. To

generate the heat map of cell KMO expression, a cell detection was carried out on haematoxylin staining using

standard parameters, a 5-100um2 area range, cell expansion of 1um and a threshold of 0.14.

Statistical analysis. Power calculations were performed using G*Power (v3.1.9.4) software. Input parameters

were used: 2-tailed, α-error probability = 0.05, and power (1-β error probability) = 0.80. Continuous variable

data were tested for Normality of distribution with a one sample Kolmogorov-Smirnov test. Normally

distributed data were analysed by one-way ANOVA with post-hoc Dunnett's T3 for multiple groups. Data

comparing treatment group effects at multiple time-points were analysed by two-way ANOVA with multiple

comparison testing by Dunnett's method . Data comparing multiple groups with unequal variances were

analyzed with Welch's ANOVA. Data not following the Normal distribution were analysed with non-parametric

Kruskal-Wallis test. Categorical and proportions data were analysed by Fisher's exact test. Data were visualised

with GraphPad Prism.

LIST OF SUPPLEMENTARY MATERIALS

Fig. S1 and Fig. S2.

Table S1 and Table S2.

REFERENCES

1. Saunders, P.T.K., and Horne, A.W. (2021). Endometriosis: Etiology, pathobiology, and therapeutic prospects. Cell *184*, 2807-2824. 10.1016/j.cell.2021.04.041.

2. Dorning, A., Dhami, P., Panir, K., Hogg, C., Park, E., Ferguson, G.D., Hargrove, D., Karras, J., Horne, A.W., and Greaves, E. (2021). Bioluminescent imaging in induced mouse models of endometriosis reveals differences in four model variations. Dis Model Mech *14*. 10.1242/dmm.049070.

3. Horne, A.W., Saunders, P.T.K., Abokhrais, I.M., Hogg, L., and (appendix), E.P.S.P.S.G. (2017). Top ten endometriosis research priorities in the UK and Ireland. Lancet *389*, 2191-2192. 10.1016/S0140-6736(17)31344-2.

4. Mole, D.J., Webster, S.P., Uings, I., Zheng, X., Binnie, M., Wilson, K., Hutchinson, J.P., Mirguet, O., Walker, A., Beaufils, B., et al. (2016). Kynurenine-3-monooxygenase inhibition prevents multiple organ failure in rodent models of acute pancreatitis. Nat Med *22*, 202-209. 10.1038/nm.4020.

5. Thul, P.J., and Lindskog, C. (2018). The human protein atlas: A spatial map of the human proteome. Protein Sci *27*, 233-244. 10.1002/pro.3307.

6. Atkins, H.M., Bharadwaj, M.S., O'Brien Cox, A., Furdui, C.M., Appt, S.E., and Caudell, D.L. (2019). Endometrium and endometriosis tissue mitochondrial energy metabolism in a nonhuman primate model. Reprod Biol Endocrinol *17*, 70. 10.1186/s12958-019-0513-8.

7. Liu, X.T., Sun, H.T., Zhang, Z.F., Shi, R.X., Liu, L.B., Yu, J.J., Zhou, W.J., Gu, C.J., Yang, S.L., Liu, Y.K., et al. (2018). Indoleamine 2,3-dioxygenase suppresses the cytotoxicity of 1 NK cells in response to ectopic endometrial stromal cells in endometriosis. Reproduction *156*, 397-404. 10.1530/REP-18-0112.

8. Liddle, J., Beaufils, B., Binnie, M., Bouillot, A., Denis, A.A., Hann, M.M., Haslam, C.P., Holmes, D.S., Hutchinson, J.P., Kranz, M., et al. (2017). The discovery of potent and selective kynurenine 3-monooxygenase inhibitors for the treatment of acute pancreatitis. Bioorg Med Chem Lett *27*, 2023-2028. 10.1016/j.bmcl.2017.02.078.

9. Walker, A.L., Ancellin, N., Beaufils, B., Bergeal, M., Binnie, M., Bouillot, A., Clapham, D., Denis, A., Haslam, C.P., Holmes, D.S., et al. (2017). Development of a Series of Kynurenine 3-Monooxygenase Inhibitors Leading to a Clinical Candidate for the Treatment of Acute Pancreatitis. J Med Chem *60*, 3383-3404. 10.1021/acs.jmedchem.7b00055.

10. Hayes, A.J., Zheng, X., O'Kelly, J., Neyton, L.P.A., Bochkina, N.A., Uings, I., Liddle, J., Baillie, J.K., Just, G., Binnie, M., et al. (2023). Kynurenine monooxygenase regulates inflammation during critical illness and recovery in experimental acute pancreatitis. Cell Rep *42*, 112763. 10.1016/j.celrep.2023.112763.

11. Ryan, A., Healey, M., Cheng, C., Dior, U., and Reddington, C. (2022). Central sensitisation in pelvic pain: A cohort study. Aust N Z J Obstet Gynaecol *62*, 868-874. 10.1111/ajo.13596.

12. Deuis, J.R., Dvorakova, L.S., and Vetter, I. (2017). Methods Used to Evaluate Pain Behaviors in Rodents. Front Mol Neurosci *10*, 284. 10.3389/fnmol.2017.00284.

13. Rahmioglu, N., Fassbender, A., Vitonis, A.F., Tworoger, S.S., Hummelshoj, L., D'Hooghe, T.M., Adamson, G.D., Giudice, L.C., Becker, C.M., Zondervan, K.T., et al. (2014). World Endometriosis Research Foundation Endometriosis Phenome and

Biobanking Harmonization Project: III. Fluid biospecimen collection, processing, and
storage in endometriosis research. Fertil Steril *102*, 1233-1243.
10.1016/j.fertnstert.2014.07.1208.

14.   Revised American Society for Reproductive Medicine classification of endometriosis:
1996. (1997). Fertil Steril *67*, 817-821. 10.1016/s0015-0282(97)81391-x.

15.   Noyes, R.W., Hertig, A.T., and Rock, J. (1975). Dating the endometrial biopsy. Am J
Obstet Gynecol *122*, 262-263. 10.1016/s0002-9378(16)33500-1.

16.   Kearns, N., Norville, K., and Frelinger, J.G. (2023). Instant Pot for antigen retrieval: a
simple, safe and economical method for use in immunohistochemistry.
Biotechniques *74*, 237-241. 10.2144/btn-2022-0043.

17.   Lai, M.H., Liao, C.H., Tsai, N.M., Chang, K.F., Liu, C.C., Chiu, Y.H., Huang, K.C., and Lin,
C.S. (2021). Surface Expression of Kynurenine 3-Monooxygenase Promotes
Proliferation and Metastasis in Triple-Negative Breast Cancers. Cancer Control *28*,
10732748211009245. 10.1177/10732748211009245.

18.   Homer, N.Z.M. Using MultiQuant and Excel software to evaluate and report multi-
analyte targeted LC-MS/MS data. https://www.protocols.io/view/using-multiquant-
and-excel-software-to-evaluate-an-cxmrxk56.

19.   Greaves, E., Cousins, F.L., Murray, A., Esnal-Zufiaurre, A., Fassbender, A., Horne, A.W.,
and Saunders, P.T. (2014). A novel mouse model of endometriosis mimics human
phenotype and reveals insights into the inflammatory contribution of shed
endometrium. Am J Pathol *184*, 1930-1939. 10.1016/j.ajpath.2014.03.011.

20.   Horne, A.W., Ahmad, S.F., Carter, R., Simitsidellis, I., Greaves, E., Hogg, C., Morton,
N.M., and Saunders, P.T.K. (2019). Repurposing dichloroacetate for the treatment of
women with endometriosis. Proc Natl Acad Sci U S A *116*, 25389-25391.
10.1073/pnas.1916144116.

21.   Yun, C.W., Yun, S., Lee, J.H., Han, Y.S., Yoon, Y.M., An, D., and Lee, S.H. (2016).
Silencing Prion Protein in HT29 Human Colorectal Cancer Cells Enhances Anticancer
Response to Fucoidan. Anticancer Res *36*, 4449-4458. 10.21873/anticanres.10989.

ACKNOWLEDGEMENTS

We would like to thank the University of Edinburgh MRC Confidence in Concept award team: Andrew McBride, Lorraine Jackson. We thank Susan Bodie, Dave Pritchard from Edinburgh Innovations. We thank all staff and support team members at Syneos Health and NAASON Science Inc.

FUNDING

UKRI Medical Research Council Confidence-in-Concept grant MRC/CIC8/73 (DJM, SPW, PTKS, AH)

UKRI Medical Research Council Senior Clinical Fellowship MR/P008887/1 (DJM)

AUTHOR CONTRIBUTIONS

Conceptualization: DJM, SPW, PTKS, AH

Methodology: BH, IS, NZMH, SGD, JPS, MM, LB, LCP, PJS, AT, SPW, AH, PTKS, DJM

Investigation: BH, IS, XZ, FC, SGP, JPS, LB, HYL, YGK, KHP, LCP GF, AT, DC, TA

Visualization: BH, IS, KHP, LCP, DJM

Funding acquisition: DJM, SPW, PTKS, AH.

Project administration: DJM, XZ, LCP, PES, AT, PTKS, AH, NZMH

Supervision: MM, LCP, AT, PTKS, DJM

Writing – original draft: DJM, LCP

Writing – review & editing: BH, IS, XZ, FC, NZMH, SGD, JPS, MM, LB, LCP, PJS, AT, SPW, AH, PTKS, DJM.

DECLARATION OF INTERESTS

The following authors have interests to declare: S.P.W., D.J.M. are co-founders of Kynos Therapeutics Ltd.. D.J.M. is a Board Member of Kynos. The University of Edinburgh controls Patents WO2015/091647, WO2016/097144, WO2016/188827 that relate to inhibitors of KMO inhibitors, and include the compound used in this paper. The remaining authors declare no competing interests.

DATA AND MATERIALS AVAILABILITY

All data are available in the main text or the supplementary materials.

KNS898 availability is restricted under a Material Transfer Agreement. Please contact the corresponding author in the first instance.

 FIGURES

 Fig. 1

Figure 1. Immunohistochemistry of KMO expression in human endometrium and distended endometriosis gland-like lesions. Fixed tissue sections were stained with anti-KMO antibody (1:500 dilution) and visualized with DAB as described in the Methods section. **Panel A.** Normal human endometrium (200 X magnification); **B**[1] and **C**[1] insets denote areas shown in panels **B** and **C** at higher magnification. KMO expression is demonstrated as dark brown DAB-positive staining and is most intense in the epithelial cells lining the glands. **Panels D through G.** Human peritoneal endometriosis tissue lesions stained with anti-KMO antibody visualized with DAB. D/**E** PIN3652, Stage II, F/**G** PIN3306 Stage I. Note intense staining of cells lining glandular structures. **Panel H.** Ovarian-type endometriosis tissue lesion with higher magnification inset

($\mathbf{I^1}$) showing KMO expression present but at lower intensity in the mesothelial tissue surface (this sample does

not have epithelial cells in the lesion).

Fig. 2

Figure 2. KNS898 plasma levels and pharmacodynamic effect of KMO blockade. Mice (n=3 per group,

individual data shown) were given KNS898 twice daily by gavage at the doses shown. After 7 days, blood was

sampled at euthanasia and KNS898 levels and kynurenine pathway metabolite levels measured by LC-MS/MS.

A. KNS898 drug levels. **B.** Kynurenine. **C.** Kynurenic acid. **D**. 3-hydroxykynurenine (logarithmic scale).

Comparison between groups by one way ANOVA with post hoc Tukey's test. *P <0.05, **P<0.01, ***P<0.001,

n.s. not statistically significant. **E.** A diagram of the kynurenine pathway showing the key step catalyzed by
KMO.

Figure 3. Therapeutic effect of KNS898 in an experimental mouse model of endometriosis. A.

Experimental design. Ovariectomized (OVX) donor mice were hormonally stimulated as shown (E2 estradiol,

P4 progesterone). At Day 19, donor mouse endometrial fragments were inoculated into recipient mice in a 1:1 ratio. KNS898 treatment 25 mg/kg twice daily by oral gavage was commenced at Day 19 or after a 1 week interval on Day 26 and in both cases continued for 2 weeks. Groups were G1: n=10, control mice; G2: n=10, sham-operated control mice; G3: n=15, endometriosis + vehicle; G4: n=15, endometriosis with KNS898 commenced at Day 19; G5: n=15, endometriosis with KNS898 commenced at Day 26. **B.** KNS898 drug levels (relative concentrations). **C.** Kynurenine. **D.** 3-hydroxykynurenine. **E**. Kynurenic acid. Individual data are shown in panels B through E . **Panels F and G.** Enumerated distended endometriosis gland-like structures (DEGLS) in recipient mice by treatment group. **F**. Total number of DEGLS per group. **G**. Total number of DEGLS per animal for all animals in the group; bars show mean with s.e.m. (**G**). Comparison between groups by one way ANOVA with post hoc Tukey's test. *P <0.05, **P<0.01, ***P<0.001, ****P<0.0001, n.s. not statistically significant.

Figure 4. Quantification of KMO expression in mouse model distended endometriosis gland-like structure
(DEGLS) lesions. Sections were visualized at 200 X magnification (**A, B, C**) with higher magnification insets
shown (**D, E, F**). **Panel A and D.** Fixed tissue sections were stained with anti-KMO antibody (1:500 dilution)
and visualized with DAB as described in the Methods section. **B and E.** QuPath was used to identify the
epithelial tissue layers (yellow arrows denote the boundary) which were quantified by thickness. **C and F.**
KMO expression intensity quantified and heat map expression values are overlayed. **G.** KMO expression
staining intensity per unit area of endometriosis DEGLS epithelium, categorized by treatment group (G3
endometriosis + vehicle; G4 endometriosis + KNS898 from D19; G5 endometriosis + KNS898 from D26.
Individual data points shown. Comparison between groups by one way ANOVA with post hoc Tukey's test. *P
<0.05. . **H, I and J.** Representative micrographs from G3 (**H**), G4 (**I**) and G5 (**J**) showing KMO expression in
the epithelial cells lining each DEGLS.

 Fig. 5

Figure 5. Effect of KNS898 on mechanical allodynia and illness behaviour in an experimental mouse
model of endometriosis. A. Hind paw Von Frey filament test showing effect of endometriosis and KNS898
treatment in groups G3, G4 and G5 **B.** Bladder Von Frey filament test **C.** Home Cage Analysis of motility
showing a daytime motility deficit in mice with endometriosis compared to control mice, and clear restitution of
normal motility in KNS898 treated groups. **D**. Nighttime home cage motility analysis showing the benefit of
KNS898 treatment on normalizing the motility deficit seen in mice with endometriosis. Data are mean with
s.e.m. For A and B, statistical comparison between groups was by two-way ANOVA to compare Group effect
and Time effect, with multiple group comparison using Dunnett's T3 test. Asterisks represent treatment group
effect statistical significance of the Dunnett's T3 test comparing treatment group G5 to G3 vehicle control *P
<0.05, **P <0.01. For C and D, Welch's ANOVA with multiple group comparison using Dunnett's T3 test was
used. Although the ANOVA was statistically significant, the post hoc Dunnett's T3 was not, therefore no
asterisks are marked.