Comparative Analysis of Small Secreted Peptide Signaling during Defense Response: Insights from Vascular and Non-Vascular Plants

Irina Lyapina[1], Daria Ganaeva[1], Eugene A. Rogozhin[1,2], Ekaterina V. Ryabukhina[1], Dmitry Yu. Ryazantsev[1], Anna Mamaeva[1] and Igor Fesenko[1*]

[1]Shemyakin-Ovchinnikov Institute of Bioorganic Chemistry, Miklukho-Maklaya 16/10, 117997 Moscow, Russia

[2]All-Russian Institute for Plant Protection, Pushkin 196608, Russia

*Corresponding author

Abstract

Peptides play a crucial role in regulating plant responses to stress conditions. Small secreted peptides (SSPs) are known to be involved in immune response and are found in all land plants. However, the evolution of stress peptide signaling remains poorly understood. Here, we compared the pathogen-induced transcriptomes of the moss *Physcomitrium patens*, *Zea mays*, *Brassica napus*, and *Solanum tuberosum* to gain insight into the role of established SSP families from vascular and non-vascular plants in immune response. We identified 50, 76, 135, and 309 known SSP genes, differentially expressed during infection, in *P. patens*, *S. tuberosum*, *Z. mays,* and *B. napus*, respectively. In addition, a novel family of bryophyte-specific PSY-like peptides that are differentially regulated under infection was identified. The similarity of SSPs across the examined plant species was validated through a comparative analysis of peptide motifs and predicted three-dimensional structures. However, despite the similar patterns of antimicrobial peptide expression, only the RALF peptide family was differentially regulated under infection in both vascular and non-vascular plants. We also found that EPFL peptides, which are involved in growth and development processes in angiosperms, were differentially regulated in *P. patens* in response to pathogen infection. Thus, our study showed the complex nature of SSP signaling and shed light on the regulation of SSPs in different plant lineages during infection.

Introduction

Peptides regulate different processes, including the response to stress conditions (Hou et al., 2021; Lyapina et al., 2019; Olsson et al., 2019). Like animal cytokines (J.-M. Zhang & An, 2007), a group of plant peptides modulating immune response was named "phytocytokines" (Hou et al., 2021). The most characterized phytocytokines include PLANT ELICITOR PEPTIDES (PEP), PAMP-INDUCED PEPTIDES (PIPs), RAPID ALKALINIZATION FACTOR (RALF), PHYTOSULFOKINES (PSKs), PLANT PEPTIDE CONTAINING SULFATED TYROSINE1 (PSY1), SERINE-RICH ENDOGENOUS PEPTIDES (SCOOPs), GOLVENs (GLVs), CLAVATA3/EMBRYO SURROUNDING REGION (CLV3/CLE), CLAVATA3/EMBRYO SURROUNDING REGION (CLV3/CLE) and INFLORESCENCE DEFICIENT IN ABSCISSION (IDA) families (Rzemieniewski & Stegmann, 2022). In addition to the stress response, phytocytokines are shown to be involved in the regulation of plant growth and development (Hou et al., 2021).

Besides phytocytokines, antimicrobial peptides (AMPs) play an important role in plant defense, triggering pathogen cell lysis due to lipid membrane disruption (Moravej et al., 2018). Plant AMPs are small, cysteine-rich proteins (Kulaeva et al., 2020; Tam et al., 2015) that have diverse structures, characteristics, and action mechanisms. The classification of plant AMPs is mostly based on sequence similarity and structures. The most characterized AMPs include thionins, defensins, hevein-like peptides, knottins, stable-like peptides, lipid transfer proteins, snakins, and cyclotides (Lima et al., 2022). Antimicrobial peptides were found in most plant taxa.

The phytocytokines, antimicrobial peptides, and plant peptide hormones that control growth and development are often referred to as Small Secreted Peptides (SSPs). The plant SSPs are involved in cell-to-cell communications and share common features: 1) the active peptide sequences are embedded at the C-terminus of 100–250 aa functional or non-functional protein precursors; 2) protein precursors contain a hydrophobic N-terminal secretion signal; 3) peptides are released by specific proteases in certain conditions (de Bang et al., 2017; Hu et al., 2021). Nevertheless, precursors of widely recognized phytocytokines, including systemin (SYS) and PEP precursors, do not possess an export signal, which highlights the shortcomings

of the SSP classification (Hu et al., 2021; Tavormina et al., 2015). The group of SSPs consists

of post-translationally modified (PTM) peptides, cysteine-rich peptides (CRP), and non-Cys-

rich and non-PTM peptides. The families of PTM peptides such as PSK, PSY,

RGF/GLV/CLEL, CLE, and IDA are widely conserved and were presented in the most recent

common ancestor of angiosperms and bryophytes. The cysteine-rich peptides, such as RALF,

EPIDERMAL PATTERNING FACTOR (EPF), and EPF-LIKE (EPFL), are shown to be

involved in the regulation of many aspects of plant growth and development (Tian et al., 2022).

These families were also present in the most recent common ancestor of angiosperms and

bryophytes (Furumizu & Shinohara, 2024). The families of systemin (SYS) and PEP peptides

are the most well-known phytocytokines, not belonging to the aforementioned two categories.

During the past decade, thousands of SSPs belonging to more than 70 families have

been identified in different plant genomes (Boschiero et al., 2020). In *Triticum aestivum*, 4,981

putative SSPs were identified, among which 1,790 SSPs were grouped into 38 known SSP

families (Tian et al., 2022). In *Oryza sativa* 236 SSPs, including 52 previously unannotated,

were shown to be involved in the immune response (P. Wang et al., 2020). In *Medicago*

trancatula, 4439 genes were predicted to encode SSPs (Boschiero et al., 2020). The

bioinformatic search in three bryophyte lineages — mosses, liverworts, and hornworts

identified hundreds of SSPs, confirming the important role of peptide signaling in all plant

lineages (Lyapina et al., 2021).

The databases of plant SSPs, such as PlantSSP (Ghorbani et al., 2015), OrysSSP (Pan

et al., 2013), and MtSSPdb (Boschiero et al., 2020) have been published in recent years.

Nevertheless, the catalogue of plant SSPs is far from complete, and new peptide families are

still being discovered. Recently, the new family of plant phytocytokines, SCREW/CTNIP, was

identified in Arabidopsis based on analysis of stress-induced transcriptomes (Z. Liu et al.,

2022; Rhodes et al., 2022). Moreover, plant genomes contain tens of thousands of small open

reading frames (smORFs, < 100 aa) that can encode unannotated secreted peptides or be a raw

material for the evolution of novel SSPs (Fesenko et al., 2021; Hanada et al., 2013). Such

translatable smORFs can be located on transcripts that were annotated as long non-coding

RNAs (lncRNAs) (Hsu & Benfey, 2018).

The evolution and role of SSPs in land colonization by plants is currently being

discussed (Bowman, 2022; Bowman et al., 2017; Ghorbani et al., 2015). The analysis of

phytocytokines in different plant groups showed that most of them are family- or species-
specific. That probably reflects their fast evolution, which is driven by arms races between
plants and microbes. For example, a SCOOP family is specific to *Brassicaceae* family (Yang
et al., 2023). Systemin and hydroxyproline-rich systemin were identified only in *Solanaceae*
family (Ryan & Pearce, 2003). The PEP family is present in many angiosperm families, but
their sequences are highly divergent (Lori et al., 2015; Tanaka & Heil, 2021). A species-
specific *Z. mays* immune signaling peptide 1 (Zip1) was found in *Zea mays* (Ziemann et al.,
2018). Nevertheless, such PTM phytocytokine families as PSK, PSY, RGF/GLV/CLEL, CLE,
and IDA were probably present in the most recent common ancestor of angiosperms and
bryophytes (Furumizu & Shinohara, 2024). The most phytocytokines were identified in
angiosperms. However, the role of RALF peptides in the immune response was recently
demonstrated in the moss *P. patens* (Mamaeva et al., 2023). Nevertheless, it is currently unclear
if plant lineages such as bryophytes, lycophytes, ferns, and gymnosperms contain unannotated
peptide families regulating response to stress conditions. Moreover, the interplay of different
families of phytocytokines in different plant lineages is poorly understood. For example, the
understanding of the interaction between several families of phytocytokines in the immune
response in diverse plant lineages is currently incomplete.

In this study, we aimed to compare patterns of SSP expression under pathogen invasion
in vascular and non-vascular plants. We performed comparative analysis of pathogen-induced
transcriptomes of three angiosperm species - *Zea mays*, *Brassica napus*, and *Solanum*
tuberosum, and one bryophyte — the moss *Physcomitrium patens*. Our analysis revealed
different patterns of SSP transcription between vascular plants as well as between vascular and
non-vascular plants. In addition, our bioinformatic search identified two novel putative peptide
families, specific to bryophytes.

Materials and Methods

Pre-processing

Raw RNA-sequencing data of the following plants under biotic stress were analyzed: model organism moss *Physcomitrium patens*, and angiosperm plants: *Zea mays*, *Brassica napus*, *Solanum tuberosum* (accessions: SRP274010, SRP390856, SRP053361, SRP066006). Three biological replicates for mock and infection samples were sequenced using Illumina HiSeq2000. The following infection time points were chosen: 4, 8 and 24 hours post inoculation (hpi) for moss *P. patens* infected by *Botrytis cinerea*, for angiosperms – 24, 40, 60 hpi for *Z. mays* infected by *Colletotrichum graminicola;* 24, 48, 96 hpi for *B. napus* infected by *Sclerotinia sclerotiorum*; 12, 24, 72 hpi for *S. tuberosum* infected by *Pectobacterium carotovorum.*

FastQC (Andrews, 2010) was used for quality control, Trimmomatic (Bolger et al., 2014) was used to clip adapters and nucleotides of low quality. Genomes and genome annotations were accessed via Phytozome (https://phytozome-next.jgi.doe.gov/, Goodstein et al., 2012) and Ensembl Plants (https://plants.ensembl.org/). *P. patens* v3.3, *Z. mays* Ref_Gen_V4, *B. napus* AST_PRJEB5043_v1, and *S. tuberosum* v6.1 versions were selected.

Assembly and ORF prediction

Genomes were indexed using hisat2-build, reads were mapped to the reference genome using HISAT2 (Kim et al., 2015). Samtools (H. Li et al., 2009) was used for collecting mapping statistics.

De novo transcripts assembly was done using Stringtie (Pertea et al., 2015), Stringtie -e mode was applied for quantification. Prediction of open reading frames (ORFs) from AUG start codon for assembled transcripts with further automatic translation to amino acid sequences of length over 30 aa was done using orfipy (Singh & Wurtele, 2021). The longest ORF per transcript was selected, considering that the longest ORF represents actual coding sequence.

Functional annotation

Interproscan (Quevillon et al., 2005) was used to predict protein domains. Analysis of homology with known families of small secreted peptides (SSPs) was done using the SSP

141 prediction tool (https://mtsspdb.zhaolab.org/prediction/, Boschiero et al., 2020). Plant long non-
142 coding RNAs were accessed via PLncDB (https://www.tobaccodb.org/plncdb/).

DE analysis and filtration

Differential gene expression analysis was done using the edgeR package (Robinson et al.,
2010). Transcripts with 1 > logFC < -1 and FDR < 0.05 were considered differentially expressed.

Translated ORFs of length less than 200 amino acids were considered small ORFs
(smORFs). ORFs predicted as known SSPs by SSP prediction tool were assigned to the
corresponding SSP family based on homology to SSPs from other plants. Search for potential
novel candidates of signaling peptides was performed in smORFs corresponding to DE transcripts,
with predicted signal sequence, no transmembrane or known protein domains, and with no
homology to known SSPs from other plants.

Sequence alignment and structure prediction

Multiple sequence alignments (MSAs) were created using the high precision L-INS-I
algorithm of MAFFT v7 (Katoh & Standley, 2013). Alignments were visualized using JalView
v2.11.2.2, Clustalx color (Waterhouse et al., 2009).

Prediction of three-dimensional structures was done using AlphaFold2 (Jumper et al.,
2021). Structural comparison was performed using TM align (Y. Zhang & Skolnick, 2005). Such
metrics as RMSD and TM-score were considered. Structural search against PDB and AlphaFold
database (AFDB) was performed using Foldseek (van Kempen et al., 2024).

Search for homologs

Search for paralogs of novel immune peptide candidates was performed using BLASTP
against all major ORFs assigned for each transcript. Hits with E-value < 0.00001 were considered
significant. Search for orthologs and paralogs for novel candidates in other databases was done
using NCBI BLASTP against non-redundant protein sequences database, organism: *Viridiplantae.*
Sequences with no matches or with matches to unannotated proteins ("putative", "unknown",
"uncharacterized", etc) were selected. TBLASTN against 1000 plant transcriptomes (1KP,
https://db.cngb.org/datamart/plant/DATApla4/) was run to search for homologs of novel
candidates among translated nucleotide sequences of other plants published in 1KP. The following

default parameters were customized: 250 max. target sequences, word size 3. Hits with E-value <

0.01 were considered significant.

Cloning and recombinant protein expression

The pET32b+ vector ("Novagen", USA) was used for recombinant plasmids construction.

When the obtained recombinant plasmid is expressed, a hybrid protein is created, that contains

thioredoxin, polyhistidine domain, enteropeptidase cleavage site (DDDDR) and the peptide

sequences. Previously an assembly of synthetic genes encoded MEG and EPFL peptides was

carried by PCR method. This procedure was made according to Shevchuk et al. (2004): the long

oligonucleotides (Supplementary Table 1) (100 ng of each) and all required components were

added to PCR mixture (the Phusion HS II DNA polymerase was used). The PCR amplification

was carried out in using with the following program: 94 ∘C – 60 s; 94 ∘C – 20 s, 60 ∘C – 20 s, 72

∘C – 20 s – 10 cycles; 72 ∘C – 60 s. Then 2.5 µl from the obtained mixture was added to a novel

amplification mixture containing flanking for the corresponding peptides oligonucleotides for (0.5

µM of each), and the amplification was carried out according to the same program (20 cycles).

The PCR product was purified from the agarose gel and cloned into the pET32b+ vector by using

HihdIII/KpnI restriction endonucleases. The obtained construction was transformed into *E. coli*

strain Origami. The induction and purification of the recombinant protein was carried out under

native conditions according to the QIAexpressionist manual ("Qiagen", Germany). For the

purpose of fusion protein purification from bacterial cells 1000 ml LB growth medium was

inoculated by 25 ml of overnight *E. coli* culture. The induction was carried out for 4 h in the

presence of 1 mM IPTG, then the cells were centrifuged, and the precipitate was resuspended in

50 ml of buffer A (50 mM NaH_2PO_4, 300 mM NaCl, 10 mM imidazole, pH 8.0). The *E. coli* cells

disruption was realized for 30 min at ice with addition of lysozyme (1 mg/ml) followed by

sonication. The clarified cell lysate was applied to Ni-NTA agarose column (10 ml) that was

preliminarily equilibrated by buffer A, then the column was washed by buffer A containing 20

mM imidazole. The fusion protein was eluted by buffer B (50 mM NaH_2PO_4, 300 mM NaCl, 250

mM imidazole, pH 8.0). The purity and size of recombinant hybrid proteins was evaluated by

SDS-PAGE electrophoresis.

Plant growth

Protonemata of *Physcomitrium patens* subsp. patens Gransden 2004 (Freiburg, Germany)
were grown in 100 mL liquid Knop medium containing 500 mg/L ammonium tartrate under white
light with a photon flux of 61 μmol/m2 s under a 16-hour photoperiod at 24°C. Seven-day-old
protonema was used for analysis.

ROS detection

For intracellular ROS detection we used the fluorescent dye 2′,7′-Dichlorofluorescin
Diacetate (DCFH-DA, Sigma-Aldrich, USA). *P. patens* protonemata was treated with 5 μM
synthesized PpEPFL or PpMEG for 10 min. Samples were incubated with 10 μM DCFH-DA for
15 min. The detection was performed on the fluorescent microscope Axio Imager M2 (Zeiss) with
an AxioCam 506 mono digital camera and filter units. The No. 44 filter (λex BP 475 nm/40 nm;
λem BP 530 nm/50 nm) was used for DCFH-DA fluorescence detection. The exposure was set to
300 msec, FITC and Brightlight channels were used. Data on the fluorescence intensity were
obtained from the related Zeiss software Zen.

Results

Identification of SSPs in pathogen-induced transcriptomes

In order to analyze expression patterns of known and previously unannotated SSPs during immune response, we re-analyzed four published transcriptomes of phylogenetically distant species: the moss *Physcomitrium patens* (Physcomitrella), *Zea mays*, *Brassica napus*, and *Solanum tuberosum* (Kwenda et al., 2016; Reboledo et al., 2021; D. Wang et al., 2022; Xia et al., 2023; Figure 1A). These transcriptomes were selected based on the following criteria: 1) the plants were infected with either necrotrophic or hemibiotrophic pathogens; 2) the quality of raw data (see Methods); and 3) RNA-seq analysis was performed at different time points after pathogen inoculation.

The RNA-seq reads were re-mapped to the corresponding genomes with the following alignment rate: 91% for *P. patens*; 96% for *Z. mays*; 83% for *B. napus*; and 85% for *S. tuberosum*. Overall, we identified from 28,234 to 67,590 transcripts at different hpi (hours post infection) across all selected species (Figure 1B). The following number of transcripts were assigned to annotated genes: 23,843 for *P. patens*; 15,269 for *S. tuberosum*; 15,138 for *Z. mays* and 19,875 for *B. napus*. To annotate all assembled transcripts, open reading frames (ORFs) larger than 30 amino acids (aa) that start with AUG codons were predicted using orfipy tool (Singh & Wurtele, 2021). For annotation, the major ORF was selected for each transcript, resulting in approximately 45,000 predicted proteins in *P. patens* and *S. tuberosum*; approximately 40,000 and 60,500 predicted proteins in *Z. mays* and *B. napus*, respectively.

	Physcomitrium patens			Zea mays			Solanum tuberosum			Brassica napus		
Plant												
Pathogen	B.cinerea			C. graminicola			P. carotovorum			S. sclerotiorum		
Time Point, h	4	8	24	24	40	60	12	24	72	24	48	96
Transcripts	43780	43861	43978	42208	41277	28234	48590	49887	49434	65613	66116	67590
DE transcripts (% lncRNA)	251 (0.4)	6256 (0.21)	13190 (0.17)	18129 (0.41)	8572 (0.36)	9685 (0.41)	2266 (0.75)	704 (0.85)	1900 (1.05)	7938 (0.38)	10523 (0.29)	15143 (0.32)
DE short ORF + SP (% lncRNA)	2 (0)	172 (0.6)	348 (0.57)	1184 (0.34)	698 (0.14)	987 (0.61)	296 (1.35)	109 (3.7)	240 (1.25)	633 (0.32)	1121 (0.27)	1535 (0.52)
DE SSP	2	27	38	53	35	48	42	14	20	70	99	140
All SSP	129	129	129	89	94	92	174	174	157	314	289	250

Figure 1. A - General scheme of work pipeline; **B -** transcripts assembly statistics.

The predicted proteins were then annotated using the InterProScan tool (Quevillon et al., 2005). This resulted in assigning to known protein domains about 57% of assembled transcripts in *P. patens* and *B. napus*, 53% in *S. tuberosum* and 70% in *Z. may*. Among the

most represented domains, such as chloroplast- or ribosome-related, a significant part of
predicted ORFs belonged to uncharacterized domains or were unannotated (Supplementary
Table 2).

We next used the online SSP prediction tool (https://mtsspdb.zhaolab.org/prediction/,
Boschiero et al., 2020) to determine the transcripts encoding precursors of known SSP families
(Figure 1B; Supplementary Table 3). Overall, 129, 92, 168, and 284 transcripts encoding
potential SSP precursors were identified in the assembled transcriptomes of *P. patens*, *Z. mays*,
S. tuberosum, and *B. napus*, respectively (Figure 1B; Supplementary Table 3).

For each species, we next identified differentially expressed (DE) transcripts (-1 ≤ log2
fold change ≥ 1, *P*adj < 0.05) at each time point (Table 1B; Supplementary Table 2). In *P.*
patens and *Z. mays*, the highest number of DE transcripts was found at 24 hpi - 30% and 43%
of all transcripts, respectively (Figure 1B; Supplementary Table 2). For *S. tuberosum* the largest
number of DE transcripts was found at 12 hpi (4.7 % of overall transcripts), and for *B. napus* -
at 96 hpi (22.4% of overall transcripts) (Figure 1B; Supplementary Table 2).

We additionally identified lncRNAs in assembled transcriptomes that could potentially
encode for unannotated peptide precursors (Hsu & Benfey, 2018). To estimate the proportion
of lncRNAs in differentially regulated transcripts, we selected the assembled transcripts that 1)
do not code for annotated proteins; 2) do not assign to known SSP families; 3) contain smORFs
below 100 codons; and 4) do not overlap known genes on the same strand. We annotated such
transcripts using the lncRNA database PLncDB (https://www.tobaccodb.org/plncdb/). At each
time point, lncRNA transcripts accounted for approximately 0.17–1.05% of differentially
regulated transcripts in each species, with *S. tuberosum* having a higher percentage and *P.*
patens having a lower one (Figure 1B; Supplementary Table 2). We suggest that it can be the
result of both biological and technical reasons.

For further analysis, we selected 50, 76, 135, and 309 SSP genes, differentially
expressed during infection, in *P. patens*, *S. tuberosum, Z. mays* and *B. napus*, respectively
(Figure 1B; Supplementary Table 3). Moreover, we additionally selected DE transcripts, which
code for unannotated proteins ≤ 200 aa, containing N-terminal signals (Figure 1B;
Supplementary Table 2). The corresponding predicted proteins were used for the identification
of novel, previously unannotated peptide families. Some of them were classified as long
noncoding RNAs (Figure 1B; Supplementary Table 2).

Transcriptional patterns of SSP genes in different plant lineages

271   We next analyzed the transcriptional patterns of different groups of SSPs in response
272 to phytopathogen infection. The first group consists of selected known phytocytokines. The
273 second group included "peptide hormones," which have been demonstrated to control growth
274 and development but not stress responses. The third group consists of known families of AMPs.

The transcriptional regulation of phytocytokines

The most common differentially regulated phytocytokines in all examined species were
from conserved phytocytokine families such as PSK, CLE, PSY, and RALF. The transcripts
that encode RALF peptides were identified in all examined species at three time points. The
members of this family were differentially regulated under infection in *P. patens*, *B. napus* and
Z. mays. In *P. patens*, the PpRALF1 (Pp3c3_15280) gene was significantly up-regulated under
infection (Figure 2; Supplementary Table 3). Some members of the RALF family were
significantly down-regulated under infection in *B. napus* and *Z. mays* (e.g., CDX78035 in *B.*
napus and Zm00001d033941_T001 in *Z. mays*; Figure 2).

Figure 2. Groups of differentially expressed gene families predicted as known SSPs involved
in immunity. Red arrows indicate up-regulated transcripts, blue arrows - down-regulated.

Previously, PSK peptides were shown to regulate the immune response to biotrophic
and necrotrophic pathogens in an antagonistic manner in Arabidopsis (Y. Li et al., 2024). The
genes belonging to the PSK family were differentially regulated in vascular plants but not in
P. patens. In Z. *mays*, PHYTOSULFOKINES 6-RELATED gene (Zm00001d023385_T001)
was up-regulated during infection. In addition, another phytosulfokine gene
(Zm00001d023384_T001) was up- and down-regulated at different hpi, and
Zm00001d048006_T001 was down-regulated. In *B. napus*, we identified 22 up-regulated
transcripts of PSK genes, and two genes, CDX68551 and CDY36518, were down-regulated.
One PSK gene (Soltu.DM.02G027440.2) was upregulated in *S. tuberosum* (Figure 2;
Supplementary Table 3).

The PSY family was recently shown to play a role in regulating the trade-off between
plant growth and stress response in vascular plants (Ogawa-Ohnishi et al., 2022). PSY
receptors (PSYRs) activate the expression of various genes encoding stress response
transcription factors upon detection of the ligands (Ogawa-Ohnishi et al., 2022). We found that
PSY transcripts were significantly down-regulated in all vascular plants under infection:
MSTRG.18847.1 in *S. tuberosum*; CDY71288, CDX80347 and MSTRG.42712.1 in *B. napus*;
14 transcripts in *Z. mays* (Figure 2; Supplementary Table 3). Some PSY genes in *Z. mays* and
B. napus were also up-regulated: Zm00001d038523_T001, Zm00001d020443_T001,
MSTRG.18595.2 and MSTRG.17209.2 in *Z. mays*; CDY64688, CDY35659, CDY69338 and
MSTRG.43937.2 in *B. napus* (Figure 2; Supplementary Table 3). However, our analysis did
not detect the transcript of the PSY gene in P. *patens*.

The analysis of phytocytokines specific to vascular plants showed that PIP (7 up-
regulated transcripts, 3 annotated: two isoforms of CDY38932 and CDX96041), CEP (12 up-
regulated transcripts, 3 annotated: CDY50603, CDX93232 and CDY63206; one down
MSTRG.44689.1), IDA (CDY08563), and SCOOP (three up-regulated: MSTRG.37091.1,
MSTRG.4039.1 and MSTRG.4077.1; one down-regulated MSTRG.30356.2) peptides were
significantly up-regulated, and some down-regulated, in *B. napus* (Figure 2; Supplementary
Table 3).

Transcripts of IDA peptide precursors were detected in *S. tuberosum* plants, but their
transcription did not show significant changes under infection.

Transcripts of PEP precursors (PROPEPs) were not detected in our analysis across the
examined plant species, likely due to their low transcription levels.

Thus, our analysis showed that only the RALF peptide family was differentially
regulated under infection in vascular and non-vascular plants.

The transcriptional patterns of known peptide growth hormones

The number of SSP families that were shown to regulate only growth and development
processes is limited. For example, EPIDERMAL PATTERNING FACTOR (EPF) and EPF-
LIKE (EPFL) family play a role in regulating plant growth and development (Y. He et al.,
2023; Xiong et al., 2022). Recently, the EPFL family was found to be significantly down-
regulated upon *F. oxisporum* infection and elicitor treatments in tomato (Slezina et al., 2021).
Our analysis revealed transcripts encoding members of the EPFL family in all angiosperms,
although their transcription level remained unchanged during infection. However, we found
that some members of EPFL family were significantly down-regulated in the moss *P. patens*.
According to our analysis, 14 transcripts belonging to the EPFL peptide family were identified
in *P. patens*. From them, 9 transcripts were significantly down-regulated (six annotated genes:
Pp3c6_12270, Pp3c23_5720, Pp3c24_9860, Pp3c23_11350, Pp3c1_26030, Pp3c6_27020;
Figure 2; Supplementary Table 3). The function of this peptide family in bryophytes is
currently unknown. To this end, we measured the accumulation of intracellular reactive oxygen
species (ROS) in moss protonemata after treatment with 5 μM synthetic EPFL (Pp3c6_12270)
for 10 min. The treatment resulted in a significant accumulation of ROS compared to the
control (paired t-test: *P-value* < 0.0005; Supplementary Figure 1A). Thus, these results suggest
that moss EPFL peptides may be involved in the regulation of the immune response, as has
been recently shown in tomato (Slezina et al., 2021).

The genes encoding members of the cysteine-rich TAX (TAXIMIN) family peptide
were significantly up-regulated in *Z. mays* (four transcripts, three isoforms of
Zm00001d008923_T001 and one not annotated MSTRG.18593.1), *B. napus* (seven transcripts,
two annotated: CDY31955 and CDY19064), *P. patens* (Pp3c4_17790) and down-regulated in

S. tuberosum (MSTRG.21268.1; Figure 2 Supplementary Table 3). This family was suggested

to play a role in nutrient-status-related signaling (de Bang et al., 2017).

Another peptide family of cysteine-rich peptides, MEG (Maternally Expressed Gene),

was previously shown to be down-regulated in tomato plants under infection (Slezina et al.,

2021). We found that members of this family were significantly down-regulated in *P. patens*

(two isoforms of Pp3c5_6640) and *B. napus* (MSTRG.22520.1) and some members were

significantly up- and down-regulated in *Z. mays* (three up-regulated transcripts: two isoforms

of Zm00001d005037_T001 and Zm00001d052375_T002; two down-regulated transcripts:

Zm00001d052375_T002 and Zm00001d042365_T001; Figure 2 Supplementary Table 3).

Since the functions of this peptide in bryophytes are also unknown, we synthesized MEG from

moss *P. patens* and measured the accumulation of ROS molecules in moss protonema cells

after treatment with 5 µM PpMEG, although no significant differences from the control were

observed (Supplementary Figure 1B).

TPD 1 (TAPETUM DETERMINANT 1) is a secreted cysteine-rich small protein that

was shown to control anthers cell differentiation in Arabidopsis (Huang et al., 2016). We

identified two TPD transcripts in the moss *P. patens* (STRG.20746.1, STRG.20746.2) and

three of the five transcripts encoded TPD-like peptides in *Z. mays* (two of them already

annotated (Zm00001d049823_T001, Zm00001d006677_T001) were significantly up-

regulated (Supplementary Table 3).

Thus, our analysis shows that cysteine-rich peptides, which were not shown to play a

role in regulating immune responses, were actively regulated during infection in all plant

lineages. Our findings show that the widely conserved RALF and EPFL peptide families may

play a role in the immune response in both vascular and non-vascular plants.

The transcriptional regulation of antimicrobial SSPs

We identified several different SSPs classified as antimicrobial and protease inhibitors

in the analyzed species, such as Gibberellic acid-stimulated Arabidopsis (GASA), ARACIN,

Potato type II proteinase inhibitor, SubIn and others. The most known antimicrobial peptides,

hevein and thionin, were identified in the transcriptomes of all examined species

(Supplementary Table 3). However, hevein transcripts were significantly up-regulated during

infection only in vascular plants (Figure 2; Supplementary Table 3). Thionins were

significantly up-regulated in *B. napus* and *S. tuberosum*, but we did not identify differential
expression in other species.

The Non-specific lipid transfer proteins (nsLTP) are a plant-specific superfamily of
cysteine-rich AMPs (Santos-Silva et al., 2023). The transcripts of nsLTP genes were identified
in all analyzed plant species and were the most highly represented group of transcripts, both
up- and down-regulated (Supplementary Table 3). Our findings showed that nsLTP have
similar patterns of regulation in the response to pathogen attack in vascular and non-vascular
plants (Figure 2).

Comparative analysis of SSP motifs

We next compared the peptide motifs of bryophyte and angiosperm SSPs, differentially
regulated in the transcriptomes of the examined species. The SSPs were selected based on the
following criteria: 1) differentially expressed in *P. patens* and in at least one angiosperm
species; or 2) differentially expressed in *P. patens*. We used only functional peptide sequences
for this analysis (Figure 3A; Supplementary Figure 2).

The EPFL peptides identified in *P. patens*, *S. tuberosum*, and *B. napus* revealed a
typical cysteine pattern as well as a conserved GS region (Figure 3A, Supplementary Figure
2A), which is similar to the previous studies (Takata et al., 2013; Xiong et al., 2022).

The YISY motif, which is shown to play a role in receptor binding (R. Zhang et al., 2023), was
identified in most RALF peptides from *P. patens*, *S. tuberosum,* and *B. napus*. However, *Z.*
mays peptides contain an ISY motif (Figure 3A, Supplementary Figure 2B). The RGC region
was found in all studied plants, suggesting that RALF peptides share known motifs in all
examined species (R. Zhang et al., 2023).

Figure 3. A - Sequence similarity analysis for peptide families in *Physcomitrium patens* and angiosperms; **B -** Structural comparison of angiosperm and *Physcomitrium patens* peptide predicted structures.

Previously, MEG peptides with 8 and 12 cysteine residues were identified in angiosperms (Gutiérrez-Marcos et al., 2004; Slezina et al., 2021). We found that MEG peptides from *P. patens* and *Z. mays*, *S. tuberosum,* and *B. napus* contain 12 conserved cysteine residues,

including two CC regions (Figure 3A, Supplementary Figure 2C). In contrast to the cysteine
patterns, the MEG peptide sequences showed significant variation among the examined
sequences.

Thus, our analysis showed that EPFL peptides are the most conserved among all the
analyzed plants. The largest difference in peptide motifs was observed in MEG peptides,
including the number of cysteine residues and overall amino acid composition.

Structure-based comparison of the identified SSPs

The prediction of peptide structures using AlphaFold2

The accuracy of sequence-based methods for predicting peptide structures has been
demonstrated in recent years (McDonald et al., 2023). Moreover, current methods for structure
comparison allow the detection of distant orthologs (Monzon et al., 2022). Therefore, we next
compared structures of EPFL, RALF, CLE, MEG, nsLTP, CAPE, TPD, and TAX peptides
from different species. For this, 135 peptide structures were predicted by AlphaFold2 (Jumper
et al., 2021), and structures with an average pLDDT > 70 were selected for further analysis.
Initially, the peptide structures of *P. patens* were compared with those of angiosperms.
Structural alignment was performed using the TM-align tool, and metrics including sequence
identity (Seq_Id), Root Mean Square Deviation (RMSD), and TM-score were calculated
(Supplementary Table 4). Because clear secondary structures were not predicted for RALF and
CLE peptide, they were excluded from further analysis. Overall, 21 peptide structures of MEG,
EPFL, CAPE, TPD, TAX, and nsLTP peptides with the highest average pLDDT per family for
each plant species were selected for structural comparisons (Supplementary Table 4,
Supplementary Materials).

The predicted structures of the MEG peptides consist of multiple beta-sheets. The
highest similarity to *P. patens* MEG structures was observed in the case of *Z. mays* MEG (TM-
score 0.80; Figure 3B) which is in line with our motif comparison. The lowest structural
similarity was found between *P. patens* and *S. tuberosum* MEG structures (TM-score 0.44,
Supplementary Table 4). The MEG of *Z. mays* contains an additional loop composed of alpha-
helices, which is absent in the predicted MEG structure of *P. patens*.

The structure of EPFL peptides comprises two anti-parallel beta-sheets connected by a
loop. The structures of EPFL peptides from *Z. mays* and *P. patens* showed the highest similarity
(TM-score 0.68; Figure 3A) compared to other analyzed plants (TM-scores 0.55 for *B. napus*
and *S. tuberosum* (Supplementary Table 4).

The predicted structure of non-specific lipid transport proteins (nsLTPs) contains four
alpha-helices. *B. napus* nsLTP peptide structure revealed the highest similarity to *P. patens*
ndLTP structures (Figure 3B). TM-scores for *Z. mays* and *S. tuberosum* structures compared
to *P. patens* were 0.50 and 0.56, respectively (Supplementary Table 4).

The predicted structures of the TPD and TAX peptides showed high structural
similarity across all examined species (Supplementary Table 4). The highest similarity between
the predicted structures of *P. patens* and *B. napus* peptides was observed (Figure 3B).

The predicted structures of the CAPE peptides contain two beta-sheets and an alpha-
helix. The highest similarity to *P. patens* CAPE was observed between P. *patens* and *B. napus*
structures (Figure 3B).

Therefore, the cysteine-rich peptides (MEG, EPFL) from *P. patens* showed the highest
similarity to *Z. mays* homologs. The structures of CAPE, nsLTP, TAX, and TPD peptides from
P. patens were the most similar to the predicted structures from *B. napus*. The TPD peptide has
the most conserved structure among *P. patens, Z. mays,* and *B. napus* species.

The search for structural homologs of identified SSPs

Then, we searched for structural homologs of *P. patens* peptide families MEG, EPFL,
CAPE, nsLTP, TAX, and TPD against AlphaFold Protein Structure Database
(https://alphafold.ebi.ac.uk) and PDB (https://www.rcsb.org) databases by Foldseek (van
Kempen et al., 2024).

The analysis of MEG peptides yielded three significant hits (TM-score > 0.5,
Probability > 0.8) against AlphaFold Database (AFDB) with CRP7-Cysteine-rich family
protein from *Zea mays* (Supplementary Figure 3).

For EPFL peptides, structural similarity to ~70 EPFL proteins from *Oryza sativa,*
Glycine max, Arabidopsis thaliana, Nicotiana tabacum, etc. was identified. Hits to structures
from the PDB database were also obtained, such as EPFL and plant receptor ERL1-TMM in
complex with EPF1 from *Arabidopsis thaliana* (Accessions in PDB database: 2liy_A, 5xkj_F,

5xjo_E; Supplementary Figure 3). We also found hits to the structures of nsLTP, TPD, TAX,
and CAPE peptides in AlphaFold and PDB databases (Supplementary Figure 3).

Thus, our analysis revealed high structural similarity among MEG, EPFL, CAPE,
nsLTP, TAX, and TPD peptides from different plant taxonomic groups. For CAPE and EPFL
precursors, hits to PDB structures were observed. Therefore, our results confirmed that
searches by Foldseek (van Kempen et al., 2024) can be used to find peptide homologs at the
structural level.

Identification of novel families of SSPs in the moss *P. patens*

Novel peptide families have emerged during the evolution of land plants (Furumizu &
Shinohara, 2024). However, the identification of taxon-specific peptides and novel peptide
families is a challenging task. Moreover, lineage-specific peptide families have been identified
only in angiosperms to date. Therefore, we next sought to search for novel SSPs involved in
immune response in the non-vascular plant, *P. patens,* using our custom pipeline.

At first, we selected 356 ORFs below 200 aa located on DE transcripts that contained a
predicted N-terminus signal sequence but did not contain a transmembrane domain or known
peptide motif (Figure 1). We additionally filtered out "candidate" SSPs that have protein
annotation or have hits in the NCBI non-redundant protein sequence database (BLASTP, E-
value < 0.01, organism: *Viridiplantae*). We next searched for possible orthologs of 58
"candidate" SSPs in the 1000 plant transcriptome (1KP) database
(https://db.cngb.org/datamart/plant/DATApla4/) using TBLASTN (E-value < 0.01). Then, we
manually checked the presence of possible peptide motifs based on multiple sequence
alignments (MSAs).

Based on the analysis of possible paralogs in the *P. patens* genome, we identified two
families of putative novel SSPs in *P. patens*. The consensus sequences of identified motifs are
presented in Figure 4A (Supplementary figure 4). Family 1 comprises three *P. patens* genes
(*Pp3c2_4250*, *Pp3c14_17750*, and *Pp3c21_20040*) that were not functionally annotated yet.
One of these genes, *Pp3c21_20040*, was differentially regulated upon *B. cinerea* infection
(Figure 4A, Supplementary figure 4A). The members of Family 1 are cysteine-rich and
comprise 8 conservative cysteine residues as well as conserved "NATH" and "SCGF" motifs
(Figure 4A). The TBLASTN search in the 1KP database identified possible orthologs in 49

species, including mosses, liverworts, and lycophytes (Figure 4A, Supplementary Figure 5).
The median length of predicted ORFs was 119 aa.

Figure 4. **A -** Alignment logos of novel candidate families found in *P. patens* (**Family 1:**
STRG.19963.1_ORF.1; STRG.1695.1_ORF.2; STRG.14124.1_ORF.4; alignment logo with
1KP; **Family 2:** STRG.4116.1_ORF.1; STRG.4116.2_ORF.1; STRG.4476.2_ORF.1;
STRG.4476.3_ORF.1; STRG.4473.1_ORF.1; alignment logo with 1KP); **B -** Alignment
logos of novel peptide candidates corresponding to unique *P. patens* ORFs with 1KP
(STRG.12453.1_ORF.5; STRG.18601.1_ORF.1; STRG.17637.1_ORF.1).

The putative peptide precursors of Family 2 shared two similar peptide motifs:
DY*D***N**H**P and DY*D***N***H**P (Figure 4A, Supplementary Figure 4B). These
motifs are similar to the known PSY peptide motif DY*(D)(P)**N**H*P (Furumizu et al.,
2021), but differ in the position of conserved residues H and P. Previously, one *P. patens* gene
(Pp3c8_3200) was identified as a precursor of PSY peptide (Furumizu et al., 2021; Tost et al.,
2021). Two predicted protein precursors from Family 2, STRG.4116.1_ORF.1 (117 aa) and
STRG.4116.2_ORF.1 (91aa), shared the DY*D***N**H**P motif at the C-end and came
from the *Pp3c3_33940V3.3* gene that was erroneously annotated as alternative splicing factor
SRp20/9G8 in Phytozome 13. The STRG.4116.1 transcript was differentially up-regulated
upon infection with *B. cinerea*. The other three predicted precursors contain the motif

DY*D***N***H**P. Two of them, STRG.4476.2_ORF.1 (114 aa) and STRG.4476.3_ORF.1

(118 aa) belonged to the gene *Pp3c4_6140*. The assembled transcript STRG.4473.1 intersects

the *P. patens* gene *Pp3c4_6040* and encodes 118 aa precursor STRG.4473.1_ORF.1. Thus, we

identified a novel family of PSY-like peptides that are encoded by three *P. patens* loci. The

TBLASTN analysis against 1000 Plant transcriptomes identified 20 possible orthologs of

Family 2 in mosses and one liverwort species (Supplementary Figure 6). The median length of

predicted ORFs was 110 aa which is similar to known PSY genes.

Because SSP can be presented by only one copy in the genomes of non-vascular plants,

we additionally analyzed putative SSP precursors without possible paralogs in *P. patens*. We

searched for their possible homologs against transcripts from 1KP database. The resulting

alignments of three most interesting up-regulated peptide candidates are presented in Figure

4B. Full alignments can be found in Supplementary Figure 7. These putative peptide precursors

contain motifs that are conserved among different plant species. In the case of

STRG.12453.1_ORF.5 (140aa), P***TPSPGHN motif is observed, which similar to CLE and

IDA peptide motifs (Furumizu et al., 2021). This candidate was found in four moss species

(Figure 4B, Supplementary Figure 7A). Candidate STRG.18601.1_ORF.1 (115aa) contains a

conserved LDLVE motif, which is observed in 13 other species, including mosses and

liverworts (Figure 4A, Supplementary Figure 7B). Finally, *P. patens* candidate

STRG.17637.1_ORF.1 (121aa) is cysteine-rich and, notably, it contains a conserved DYY

region. Hits to four other species were found, including moss, liverwort, and hornwort plants

(Figure 4B, Supplementary Figure 7C).

Thus, we propose an approach to identify novel, not annotated, plant peptide families. This

approach focuses on motif analysis of ORFs from DE transcripts, both with the presence of

SSP features (proper length, N-terminal signal sequence, no TM-domain) and the absence of

annotated protein domains, homology to known proteins. We identified 5 potential candidates

for novel signaling peptides in *P. patens*. Two of them comprise families based on sequence

similarity, and some of them are unique ORFs predicted in *P. patens*. Nevertheless, all potential

novel candidates are found in other plant species belonging to non-vascular plants. For two

candidates, distant homology to known peptide families (PSY, IDA, and CLE) is suspected.

Discussion

Recent studies showed that peptides derived from protein precursors play an important role in plant immunity, modulating the defense response. The plant cell-surface receptors that recognize these peptide ligands contain Leucine-Rich Repeat (LRR) or Malectin ectodomains and have an ancient origin (Ngou et al., 2024). Contrary to immune receptors, the origin and evolution of immune peptides are still poorly understood (Furumizu et al., 2021). One of the reasons is that, due to rapid changes, they appear to quickly enter the "twilight zone", where sequence similarity is no longer detectable. Nevertheless, the several databases that include plant Small Secreted Peptides (SSPs) were recently published (Boschiero et al., 2020; Ghorbani et al., 2015; Pan et al., 2013). Sequence similarity, however, does not necessarily mean functional similarity across different plant phyla. In this study, we compared pathogen-induced transcriptomes in vascular and non-vascular plants to understand how SSP families from different plant lineages respond to infection. Along with conserved SSPs associated with immune responses that were found in the transcriptomes of both vascular and non-vascular plants, we observed species-specific regulation of certain SSP families.

Our analysis shows that cysteine-rich peptides were actively regulated during infection in all plant lineages. The EPFL, RALF, TAX, and MEG were differentially regulated in *P. patens* and angiosperm species. The widely conserved family of EPIDERMAL PATTERNING FACTOR (EPF) peptides is involved in regulating plant growth or development in vascular and non-vascular land plants (Caine et al., 2016; Y. He et al., 2023). The nine EPFL genes that are upregulated in the developing sporophyte have been identified in previous bioinformatics studies of the *P. patens* genome (Caine et al., 2016; Lyapina et al., 2021; Takata et al., 2013), but their functions remain unknown. However, the role of EPF and EPF-LIKE (EPFL) peptides as positive or negative regulators of immune response has not been shown to date. Nevertheless, several genes of the EPFL family were shown to be significantly down-regulated upon pathogen infection in tomato (Slezina et al., 2021). In this study, we found that the transcriptional level of six EPFL genes was significantly down-regulated in *P. patens*. This result can be explained by a specific response to pathogens or by the inhibition of growth processes under stress conditions.

Our treatment of moss protonemata with synthetic EPFL peptide resulted in an increase in intracellular reactive oxygen species (ROS) levels, suggesting the activation of downstream

signaling. In Arabidopsis, the receptor-like kinase (RLK) ERECTA and its two closely related

homologs, ERECTA-LIKE 1 (ERL1) and ERL2 are responsible for perceiving the EPF/EPFL

family peptides (Y. He et al., 2023; Tameshige et al., 2017). In the genome of *P. patens,* six

potential orthologues of the ERECTA genes were found (Villagarcia et al., 2012). Thus, the

EPFL family might be a target for future work on understanding the evolution of

phytocytokines in land plants.

The RALF peptides were shown to be involved in immune response across different

plant lineages (Y.-H. He et al., 2023; Mamaeva et al., 2023; Slezina et al., 2021; Wolf & Höfte,

2014). We found both up-regulated and down-regulated RALF genes in pathogen-induced

transcriptomes, which is consistent with previous results showing different roles of different

members of RALFs in modulating plant immunity (Y.-H. He et al., 2023; L. Liu et al., 2024;

Mamaeva et al., 2023; Stegmann et al., 2017).

The poorly characterized cysteine-rich peptides, such as TAX and MEG were

differentially regulated in both vascular and non-vascular plant species. However, how TAX

peptides are involved in immune response is currently unknown. Previously, the tomato MEGs

were found to be down-regulated in response to infection (Slezina et al., 2021). This is similar

to our findings and suggests the conservative role of this SSP family in vascular and non-

vascular land plants. Nevertheless, the role of MEG peptides in plant growth and defense is

poorly understood.

In addition to common approaches for the identification of SSPs based on sequence

similarity (Boschiero et al., 2020), we developed a new pipeline to identify novel lineage-

specific secreted peptides. Our pipeline resulted in the identification of two putative SSP

families in *P. patens*. One of these families (Family 2) consists of PSY-like peptides, which

are similar to the known phytocytokine PSY family in angiosperms. The PSY peptide family

was present in the most recent common ancestor of vascular plants and bryophytes (Furumizu

& Shinohara, 2024). Previously, one gene potentially encoding the PSY peptide and another

encoding the cognate PSYR receptor were identified in *P. patens* (Furumizu et al., 2021;

Furumizu & Aalen, 2023). Thus, we significantly expand the family of PSY-like peptides in

bryophytes. The identification of previously unknown PSY-like peptides in bryophyte genomes

could expand our understanding of the peptide evolution in land plants.

In conclusion, our comprehensive analysis of small secreted peptides in the pathogen-induced transcriptomes of four plant species showed the conservation of cysteine-rich peptides across different plant lineages. We found high motif and structural conservation of these peptides in vascular and non-vascular plants. In addition, we identified previously uncharacterized putative SSP families in bryophytes. One of these bryophyte-specific families shows similarity to previously characterized PSY peptides. Thus, our study provides a new avenue to explore the evolution of known and previously uncharacterized peptide families.

Author contributions

IF supervised the study. IL, DG conducted all experiments and performed all bioinformatic analyses. AM performed ROS measurement. EAR, DYR and EVR synthesized recombinant moss *P. patens* peptides. IL, DG and IF wrote the manuscript. All authors contributed to the article and approved the submitted version.

Funding

This work was supported by Russian Science Foundation (project no. 23-74-10048).

Data availability statement

The data that support the findings were derived from the NCBI Sequence Reads Archive (SRA) available in the public domain with following accession numbers: SRP274010, SRP390856, SRP053361, SRP066006.

References

Andrews, S. (2010). *FastQC: A Quality Control tool for High Throughput Sequence Data.* https://www.bioinformatics.babraham.ac.uk/projects/fastqc/

Bolger, A. M., Lohse, M., & Usadel, B. (2014). Trimmomatic: A flexible trimmer for Illumina sequence data. *Bioinformatics (Oxford, England)*, *30*(15), 2114–2120. https://doi.org/10.1093/bioinformatics/btu170

Boschiero, C., Dai, X., Lundquist, P. K., Roy, S., Bang, T. C. de, Zhang, S., Zhuang, Z., Torres-

Jerez, I., Udvardi, M. K., Scheible, W.-R., & Zhao, P. X. (2020). MtSSPdb: The Medicago
truncatula Small Secreted Peptide Database. *Plant Physiology*, *183*(1), 399–413.
https://doi.org/10.1104/pp.19.01088

Bowman, J. L. (2022). The origin of a land flora. *Nature Plants*, *8*(12), 1352–1369.
https://doi.org/10.1038/s41477-022-01283-y

Bowman, J. L., Kohchi, T., Yamato, K. T., Jenkins, J., Shu, S., Ishizaki, K., Yamaoka, S.,
Nishihama, R., Nakamura, Y., Berger, F., Adam, C., Aki, S. S., Althoff, F., Araki, T., Arteaga-
Vazquez, M. A., Balasubrmanian, S., Barry, K., Bauer, D., Boehm, C. R., … Schmutz, J.
(2017). Insights into Land Plant Evolution Garnered from the Marchantia polymorpha
Genome. *Cell*, *171*(2), 287-304.e15. https://doi.org/10.1016/j.cell.2017.09.030

Caine, R. S., Chater, C. C., Kamisugi, Y., Cuming, A. C., Beerling, D. J., Gray, J. E., &
Fleming, A. J. (2016). An ancestral stomatal patterning module revealed in the non-vascular
land plant Physcomitrella patens. *Development*, *143*(18), 3306–3314.
https://doi.org/10.1242/dev.135038

de Bang, T. C., Lay, K. S., Scheible, W.-R., & Takahashi, H. (2017). Small peptide signaling
pathways modulating macronutrient utilization in plants. *Current Opinion in Plant Biology*,
39, 31–39. https://doi.org/10.1016/j.pbi.2017.05.005

Fesenko, I., Shabalina, S. A., Mamaeva, A., Knyazev, A., Glushkevich, A., Lyapina, I.,
Ziganshin, R., Kovalchuk, S., Kharlampieva, D., Lazarev, V., Taliansky, M., & Koonin, E. V.
(2021). A vast pool of lineage-specific microproteins encoded by long non-coding RNAs in
plants. *Nucleic Acids Research*, *49*(18), 10328–10346. https://doi.org/10.1093/nar/gkab816

Furumizu, C., & Aalen, R. B. (2023). Peptide signaling through leucine-rich repeat receptor
kinases: Insight into land plant evolution. *New Phytologist*, *238*(3), 977–982.
https://doi.org/10.1111/nph.18827

Furumizu, C., Krabberød, A. K., Hammerstad, M., Alling, R. M., Wildhagen, M., Sawa, S., &
Aalen, R. B. (2021). The sequenced genomes of nonflowering land plants reveal the innovative
evolutionary history of peptide signaling. *The Plant Cell*, *33*(9), 2915–2934.
https://doi.org/10.1093/plcell/koab173

Furumizu, C., & Shinohara, H. (2024). Land plant peptide signaling: What we know—and
don't know—about its evolution. *Physiologia Plantarum*, *176*(1), e14172.
https://doi.org/10.1111/ppl.14172

Ghorbani, S., Lin, Y.-C., Parizot, B., Fernandez, A., Njo, M. F., Van de Peer, Y., Beeckman,
T., & Hilson, P. (2015). Expanding the repertoire of secretory peptides controlling root
development with comparative genome analysis and functional assays. *Journal of
Experimental Botany*, *66*(17), 5257–5269. https://doi.org/10.1093/jxb/erv346

Goodstein, D. M., Shu, S., Howson, R., Neupane, R., Hayes, R. D., Fazo, J., Mitros, T., Dirks,
W., Hellsten, U., Putnam, N., & Rokhsar, D. S. (2012). Phytozome: A comparative platform
for green plant genomics. *Nucleic Acids Research*, *40*(Database issue), D1178-1186.
https://doi.org/10.1093/nar/gkr944

Gutiérrez-Marcos, J. F., Costa, L. M., Biderre-Petit, C., Khbaya, B., O'Sullivan, D. M.,

Wormald, M., Perez, P., & Dickinson, H. G. (2004). Maternally expressed gene1 Is a novel maize endosperm transfer cell-specific gene with a maternal parent-of-origin pattern of expression. *The Plant Cell*, *16*(5), 1288–1301. https://doi.org/10.1105/tpc.019778

Hanada, K., Higuchi-Takeuchi, M., Okamoto, M., Yoshizumi, T., Shimizu, M., Nakaminami, K., Nishi, R., Ohashi, C., Iida, K., Tanaka, M., Horii, Y., Kawashima, M., Matsui, K., Toyoda, T., Shinozaki, K., Seki, M., & Matsui, M. (2013). Small open reading frames associated with morphogenesis are hidden in plant genomes. *Proceedings of the National Academy of Sciences*, *110*(6), 2395–2400. https://doi.org/10.1073/pnas.1213958110

He, Y., He, X., Wang, X., Hao, M., Gao, J., Wang, Y., Yang, Z.-N., & Meng, X. (2023). An EPFL peptide signaling pathway promotes stamen elongation via enhancing filament cell proliferation to ensure successful self-pollination in Arabidopsis thaliana. *New Phytologist*, *238*(3), 1045–1058. https://doi.org/10.1111/nph.18806

He, Y.-H., Chen, S.-Y., Chen, X.-Y., Xu, Y.-P., Liang, Y., & Cai, X.-Z. (2023). RALF22 promotes plant immunity and amplifies the Pep3 immune signal. *Journal of Integrative Plant Biology*, *65*(11), 2519–2534. https://doi.org/10.1111/jipb.13566

Hou, S., Liu, D., & He, P. (2021). Phytocytokines function as immunological modulators of plant immunity. *Stress Biology*, *1*(1), 8. https://doi.org/10.1007/s44154-021-00009-y

Hsu, P. Y., & Benfey, P. N. (2018). Small but Mighty: Functional Peptides Encoded by Small ORFs in Plants. *PROTEOMICS*, *18*(10), 1700038. https://doi.org/10.1002/pmic.201700038

Hu, X.-L., Lu, H., Hassan, M. M., Zhang, J., Yuan, G., Abraham, P. E., Shrestha, H. K., Villalobos Solis, M. I., Chen, J.-G., Tschaplinski, T. J., Doktycz, M. J., Tuskan, G. A., Cheng, Z.-M. (Max), & Yang, X. (2021). Advances and perspectives in discovery and functional analysis of small secreted proteins in plants. *Horticulture Research*, *8*(1), 1–14. https://doi.org/10.1038/s41438-021-00570-7

Huang, J., Zhang, T., Linstroth, L., Tillman, Z., Otegui, M. S., Owen, H. A., & Zhao, D. (2016). Control of Anther Cell Differentiation by the Small Protein Ligand TPD1 and Its Receptor EMS1 in Arabidopsis. *PLOS Genetics*, *12*(8), e1006147. https://doi.org/10.1371/journal.pgen.1006147

Jumper, J., Evans, R., Pritzel, A., Green, T., Figurnov, M., Ronneberger, O., Tunyasuvunakool, K., Bates, R., Žídek, A., Potapenko, A., Bridgland, A., Meyer, C., Kohl, S. A. A., Ballard, A. J., Cowie, A., Romera-Paredes, B., Nikolov, S., Jain, R., Adler, J., … Hassabis, D. (2021). Highly accurate protein structure prediction with AlphaFold. *Nature*, *596*(7873), 583–589. https://doi.org/10.1038/s41586-021-03819-2

Katoh, K., & Standley, D. M. (2013). MAFFT Multiple Sequence Alignment Software Version 7: Improvements in Performance and Usability. *Molecular Biology and Evolution*, *30*(4), 772–780. https://doi.org/10.1093/molbev/mst010

Kim, D., Langmead, B., & Salzberg, S. L. (2015). HISAT: A fast spliced aligner with low memory requirements. *Nature Methods*, *12*(4), 357–360. https://doi.org/10.1038/nmeth.3317

Kulaeva, O., Kliukova, M., Afonin, A., Sulima, A., Zhukov, V., & Tikhonovich, I. (2020). The role of plant antimicrobial peptides (AMPs) in response to biotic and abiotic environmental

factors. *Biological Communications*, *65*(2), Article 2. https://doi.org/10.21638/spbu03.2020.205

Kwenda, S., Birch, P. R. J., & Moleleki, L. N. (2016). Genome-wide identification of potato long intergenic noncoding RNAs responsive to Pectobacterium carotovorum subspecies brasiliense infection. *BMC Genomics*, *17*(1), 614. https://doi.org/10.1186/s12864-016-2967-9

Li, H., Handsaker, B., Wysoker, A., Fennell, T., Ruan, J., Homer, N., Marth, G., Abecasis, G., Durbin, R., & 1000 Genome Project Data Processing Subgroup. (2009). The Sequence Alignment/Map format and SAMtools. *Bioinformatics (Oxford, England)*, *25*(16), 2078–2079. https://doi.org/10.1093/bioinformatics/btp352

Li, Y., Di, Q., Luo, L., & Yu, L. (2024). Phytosulfokine peptides, their receptors, and functions. *Frontiers in Plant Science*, *14*. https://doi.org/10.3389/fpls.2023.1326964

Lima, A. M., Azevedo, M. I. G., Sousa, L. M., Oliveira, N. S., Andrade, C. R., Freitas, C. D. T., & Souza, P. F. N. (2022). Plant antimicrobial peptides: An overview about classification, toxicity and clinical applications. *International Journal of Biological Macromolecules*, *214*, 10–21. https://doi.org/10.1016/j.ijbiomac.2022.06.043

Liu, L., Liu, X., Bai, Z., Tanveer, M., Zhang, Y., Chen, W., Shabala, S., & Huang, L. (2024). Small but powerful: RALF peptides in plant adaptive and developmental responses. *Plant Science*, *343*, 112085. https://doi.org/10.1016/j.plantsci.2024.112085

Liu, Z., Hou, S., Rodrigues, O., Wang, P., Luo, D., Munemasa, S., Lei, J., Liu, J., Ortiz-Morea, F. A., Wang, X., Nomura, K., Yin, C., Wang, H., Zhang, W., Zhu-Salzman, K., He, S. Y., He, P., & Shan, L. (2022). Phytocytokine signalling reopens stomata in plant immunity and water loss. *Nature*, *605*(7909), 332–339. https://doi.org/10.1038/s41586-022-04684-3

Lori, M., van Verk, M. C., Hander, T., Schatowitz, H., Klauser, D., Flury, P., Gehring, C. A., Boller, T., & Bartels, S. (2015). Evolutionary divergence of the plant elicitor peptides (Peps) and their receptors: Interfamily incompatibility of perception but compatibility of downstream signalling. *Journal of Experimental Botany*, *66*(17), 5315–5325. https://doi.org/10.1093/jxb/erv236

Lyapina, I., Filippova, A., & Fesenko, I. (2019). The Role of Peptide Signals Hidden in the Structure of Functional Proteins in Plant Immune Responses. *International Journal of Molecular Sciences*, *20*(18), 4343. https://doi.org/10.3390/ijms20184343

Lyapina, I., Filippova, A., Kovalchuk, S., Ziganshin, R., Mamaeva, A., Lazarev, V., Latsis, I., Mikhalchik, E., Panasenko, O., Ivanov, O., Ivanov, V., & Fesenko, I. (2021). Possible role of small secreted peptides (SSPs) in immune signaling in bryophytes. *Plant Molecular Biology*, *106*(1–2), 123–143. https://doi.org/10.1007/s11103-021-01133-z

Mamaeva, A., Lyapina, I., Knyazev, A., Golub, N., Mollaev, T., Chudinova, E., Elansky, S., Babenko, V. V., Veselovsky, V. A., Klimina, K. M., Gribova, T., Kharlampieva, D., Lazarev, V., & Fesenko, I. (2023). RALF peptides modulate immune response in the moss Physcomitrium patens. *Frontiers in Plant Science*, *14*. https://doi.org/10.3389/fpls.2023.1077301

McDonald, E. F., Jones, T., Plate, L., Meiler, J., & Gulsevin, A. (2023). Benchmarking

AlphaFold2 on Peptide Structure Prediction. *Structure (London, England : 1993)*, *31*(1), 111–
119.e2. https://doi.org/10.1016/j.str.2022.11.012

Monzon, V., Paysan-Lafosse, T., Wood, V., & Bateman, A. (2022). Reciprocal best structure
hits: Using AlphaFold models to discover distant homologues. *Bioinformatics Advances*, *2*(1),
vbac072. https://doi.org/10.1093/bioadv/vbac072

Moravej, H., Moravej, Z., Yazdanparast, M., Heiat, M., Mirhosseini, A., Moosazadeh
Moghaddam, M., & Mirnejad, R. (2018). Antimicrobial Peptides: Features, Action, and Their
Resistance Mechanisms in Bacteria. *Microbial Drug Resistance*, *24*(6), 747–767.
https://doi.org/10.1089/mdr.2017.0392

Ngou, B. P. M., Wyler, M., Schmid, M. W., Kadota, Y., & Shirasu, K. (2024). Evolutionary
trajectory of pattern recognition receptors in plants. *Nature Communications*, *15*(1), 308.
https://doi.org/10.1038/s41467-023-44408-3

Ogawa-Ohnishi, M., Yamashita, T., Kakita, M., Nakayama, T., Ohkubo, Y., Hayashi, Y.,
Yamashita, Y., Nomura, T., Noda, S., Shinohara, H., & Matsubayashi, Y. (2022). Peptide
ligand-mediated trade-off between plant growth and stress response. *Science (New York, N.Y.)*,
378(6616), 175–180. https://doi.org/10.1126/science.abq5735

Olsson, V., Joos, L., Zhu, S., Gevaert, K., Butenko, M. A., & De Smet, I. (2019). Look Closely,
the Beautiful May Be Small: Precursor-Derived Peptides in Plants. *Annual Review of Plant
Biology*, *70*, 153–186. https://doi.org/10.1146/annurev-arplant-042817-040413

Pan, B., Sheng, J., Sun, W., Zhao, Y., Hao, P., & Li, X. (2013). OrysPSSP: A comparative
Platform for Small Secreted Proteins from rice and other plants. *Nucleic Acids Research*,
41(D1), D1192–D1198. https://doi.org/10.1093/nar/gks1090

Pertea, M., Pertea, G. M., Antonescu, C. M., Chang, T.-C., Mendell, J. T., & Salzberg, S. L.
(2015). StringTie enables improved reconstruction of a transcriptome from RNA-seq reads.
Nature Biotechnology, *33*(3), 290–295. https://doi.org/10.1038/nbt.3122

Quevillon, E., Silventoinen, V., Pillai, S., Harte, N., Mulder, N., Apweiler, R., & Lopez, R.
(2005). InterProScan: Protein domains identifier. *Nucleic Acids Research*, *33*(Web Server
issue), W116–W120. https://doi.org/10.1093/nar/gki442

Reboledo, G., Agorio, A. D., Vignale, L., Batista-García, R. A., & Ponce De León, I. (2021).
Transcriptional profiling reveals conserved and species-specific plant defense responses during
the interaction of Physcomitrium patens with Botrytis cinerea. *Plant Molecular Biology*,
107(4–5), 365–385. https://doi.org/10.1007/s11103-021-01116-0

Rhodes, J., Roman, A.-O., Bjornson, M., Brandt, B., Derbyshire, P., Wyler, M., Schmid, M.
W., Menke, F. L., Santiago, J., & Zipfel, C. (2022). Perception of a conserved family of plant
signalling peptides by the receptor kinase HSL3. *eLife*, *11*, e74687.
https://doi.org/10.7554/eLife.74687

Robinson, M. D., McCarthy, D. J., & Smyth, G. K. (2010). edgeR: A Bioconductor package
for differential expression analysis of digital gene expression data. *Bioinformatics (Oxford,
England)*, *26*(1), 139–140. https://doi.org/10.1093/bioinformatics/btp616

Ryan, C. A., & Pearce, G. (2003). Systemins: A functionally defined family of peptide signals
that regulate defensive genes in Solanaceae species. *Proceedings of the National Academy of*
Sciences, *100*(suppl_2), 14577–14580. https://doi.org/10.1073/pnas.1934788100

Rzemieniewski, J., & Stegmann, M. (2022). Regulation of pattern-triggered immunity and
growth by phytocytokines. *Current Opinion in Plant Biology*, *68*, 102230.
https://doi.org/10.1016/j.pbi.2022.102230

Santos-Silva, C. A. dos, Ferreira-Neto, J. R. C., Amador, V. C., Bezerra-Neto, J. P., Vilela, L.
M. B., Binneck, E., Rêgo, M. de S., da Silva, M. D., Mangueira de Melo, A. L. T., da Silva, R.
H., & Benko-Iseppon, A. M. (2023). From Gene to Transcript and Peptide: A Deep Overview
on Non-Specific Lipid Transfer Proteins (nsLTPs). *Antibiotics*, *12*(5), Article 5.
https://doi.org/10.3390/antibiotics12050939

Shevchuk, N. A., Bryksin, A. V., Nusinovich, Y. A., Cabello, F. C., Sutherland, M., & Ladisch,
S. (2004). Construction of long DNA molecules using long PCR-based fusion of several
fragments   simultaneously.   *Nucleic   Acids   Research*,   *32*(2),   e19.
https://doi.org/10.1093/nar/gnh014

Singh, U., & Wurtele, E. S. (2021). orfipy: A fast and flexible tool for extracting ORFs.
Bioinformatics, *37*(18), 3019–3020. https://doi.org/10.1093/bioinformatics/btab090

Slezina, M. P., Istomina, E. A., Korostyleva, T. V., Kovtun, A. S., Kasianov, A. S., Konopkin,
A. A., Shcherbakova, L. A., & Odintsova, T. I. (2021). Molecular Insights into the Role of
Cysteine-Rich Peptides in Induced Resistance to Fusarium oxysporum Infection in Tomato
Based on Transcriptome Profiling. *International Journal of Molecular Sciences*, *22*(11), 5741.
https://doi.org/10.3390/ijms22115741

Stegmann, M., Monaghan, J., Smakowska-Luzan, E., Rovenich, H., Lehner, A., Holton, N.,
Belkhadir, Y., & Zipfel, C. (2017). The receptor kinase FER is a RALF-regulated scaffold
controlling   plant   immune   signaling.   *Science*,   *355*(6322),   287–289.
https://doi.org/10.1126/science.aal2541

Takata, N., Yokota, K., Ohki, S., Mori, M., Taniguchi, T., & Kurita, M. (2013). Evolutionary
Relationship and Structural Characterization of the EPF/EPFL Gene Family. *PLOS ONE*, *8*(6),
e65183. https://doi.org/10.1371/journal.pone.0065183

Tam, J. P., Wang, S., Wong, K. H., & Tan, W. L. (2015). Antimicrobial Peptides from Plants.
Pharmaceuticals, *8*(4), 711–757. https://doi.org/10.3390/ph8040711

Tameshige, T., Ikematsu, S., Torii, K. U., & Uchida, N. (2017). Stem development through
vascular tissues: EPFL-ERECTA family signaling that bounces in and out of phloem. *Journal*
of Experimental Botany, *68*(1), 45–53. https://doi.org/10.1093/jxb/erw447

Tanaka, K., & Heil, M. (2021). Damage-Associated Molecular Patterns (DAMPs) in Plant
Innate Immunity: Applying the Danger Model and Evolutionary Perspectives. *Annual Review*
of Phytopathology, *59*(Volume 59, 2021), 53–75. https://doi.org/10.1146/annurev-phyto-
082718-100146

Tavormina, P., De Coninck, B., Nikonorova, N., De Smet, I., & Cammue, B. P. A. (2015). The
Plant Peptidome: An Expanding Repertoire of Structural Features and Biological Functions.

The Plant Cell, *27*(8), 2095–2118. https://doi.org/10.1105/tpc.15.00440

Tian, D., Xie, Q., Deng, Z., Xue, J., Li, W., Zhang, Z., Dai, Y., Zheng, B., Lu, T., De Smet, I.,
& Guo, Y. (2022). Small secreted peptides encoded on the wheat (triticum aestivum L.) genome
and their potential roles in stress responses. *Frontiers in Plant Science*, *13*, 1000297.
https://doi.org/10.3389/fpls.2022.1000297

Tost, A. S., Kristensen, A., Olsen, L. I., Axelsen, K. B., & Fuglsang, A. T. (2021). The PSY
Peptide Family—Expression, Modification and Physiological Implications. *Genes*, *12*(2),
Article 2. https://doi.org/10.3390/genes12020218

van Kempen, M., Kim, S. S., Tumescheit, C., Mirdita, M., Lee, J., Gilchrist, C. L. M., Söding,
J., & Steinegger, M. (2024). Fast and accurate protein structure search with Foldseek. *Nature*
Biotechnology, *42*(2), 243–246. https://doi.org/10.1038/s41587-023-01773-0

Villagarcia, H., Morin, A.-C., Shpak, E. D., & Khodakovskaya, M. V. (2012). Modification of
tomato growth by expression of truncated ERECTA protein from Arabidopsis thaliana. *Journal*
of Experimental Botany, *63*(18), 6493–6504. https://doi.org/10.1093/jxb/ers305

Wang, D., Jin, S., Chen, Z., Shan, Y., & Li, L. (2022). Genome-wide identification of the pectin
methylesterase inhibitor genes in Brassica napus and expression analysis of selected members.
Frontiers in Plant Science, *13*, 940284. https://doi.org/10.3389/fpls.2022.940284

Wang, P., Yao, S., Kosami, K., Guo, T., Li, J., Zhang, Y., Fukao, Y., Kaneko-Kawano, T.,
Zhang, H., She, Y.-M., Wang, P., Xing, W., Hanada, K., Liu, R., & Kawano, Y. (2020).
Identification of endogenous small peptides involved in rice immunity through
transcriptomics- and proteomics-based screening. *Plant Biotechnology Journal*, *18*(2), 415–
428. https://doi.org/10.1111/pbi.13208

Waterhouse, A. M., Procter, J. B., Martin, D. M. A., Clamp, M., & Barton, G. J. (2009). Jalview
Version 2—A multiple sequence alignment editor and analysis workbench. *Bioinformatics*,
25(9), 1189–1191. https://doi.org/10.1093/bioinformatics/btp033

Wolf, S., & Höfte, H. (2014). Growth Control: A Saga of Cell Walls, ROS, and Peptide
Receptors. *The Plant Cell*, *26*(5), 1848–1856. https://doi.org/10.1105/tpc.114.125518

Xia, X., Wei, Q., Xiao, C., Ye, Y., Li, Z., Marivingt-Mounir, C., Chollet, J.-F., Liu, W., & Wu,
H. (2023). Genomic survey of NPF and NRT2 transporter gene families in five inbred maize
lines and their responses to pathogens infection. *Genomics*, *115*(2), 110555.
https://doi.org/10.1016/j.ygeno.2022.110555

Xiong, L., Huang, Y., Liu, Z., Li, C., Yu, H., Shahid, M. Q., Lin, Y., Qiao, X., Xiao, J., Gray,
J. E., & Jin, J. (2022). Small EPIDERMAL PATTERNING FACTOR-LIKE2 peptides regulate
awn development in rice. *Plant Physiology*, *190*(1), 516–531.
https://doi.org/10.1093/plphys/kiac278

Yang, H., Kim, X., Skłenar, J., Aubourg, S., Sancho-Andrés, G., Stahl, E., Guillou, M.-C.,
Gigli-Bisceglia, N., Tran Van Canh, L., Bender, K. W., Stintzi, A., Reymond, P., Sánchez-
Rodríguez, C., Testerink, C., Renou, J.-P., Menke, F. L. H., Schaller, A., Rhodes, J., & Zipfel,
C. (2023). Subtilase-mediated biogenesis of the expanded family of SERINE RICH
ENDOGENOUS PEPTIDES. *Nature Plants*, *9*(12), 2085–2094.

https://doi.org/10.1038/s41477-023-01583-x

Zhang, J.-M., & An, J. (2007). Cytokines, inflammation, and pain. *International*
Anesthesiology Clinics, *45*(2), 27–37. https://doi.org/10.1097/AIA.0b013e318034194e

Zhang, R., Shi, P.-T., Zhou, M., Liu, H.-Z., Xu, X.-J., Liu, W.-T., & Chen, K.-M. (2023). Rapid
alkalinization factor: Function, regulation, and potential applications in agriculture. *Stress*
Biology, *3*(1), 16. https://doi.org/10.1007/s44154-023-00093-2

Zhang, Y., & Skolnick, J. (2005). TM-align: A protein structure alignment algorithm based on
the TM-score. *Nucleic Acids Research*, *33*(7), 2302–2309. https://doi.org/10.1093/nar/gki524

Ziemann, S., van der Linde, K., Lahrmann, U., Acar, B., Kaschani, F., Colby, T., Kaiser, M.,
Ding, Y., Schmelz, E., Huffaker, A., Holton, N., Zipfel, C., & Doehlemann, G. (2018). An
apoplastic peptide activates salicylic acid signalling in maize. *Nature Plants*, *4*(3), Article 3.
https://doi.org/10.1038/s41477-018-0116-y

878