Hypocotyl Development in Arabidopsis and other Brassicaceae Displays Evidence of Photoperiodic Memory

James Ronald[1], Sarah C.L Lock[1], Will Claydon[1], Zihao Zhu[2], Kayla McCarthy[1], Elizabeth Pendlington[1], Ethan J. Redmond[1], Gina Y.W. Vong[1], Sanoj P. Stanislas[1], Seth J. Davis[1], Marcel Quint[2], Daphne Ezer[1]*.

[1]Department of Biology, University of York, Wentworth Way, Heslington, York YO10 5DD, UK.
[2]Institute of Agricultural and Nutritional Sciences, Martin Luther University Halle-Wittenberg, Halle (Saale), Germany.

*Corresponding authors:
Daphne Ezer (daphne.ezer@york.ac.uk)

Abstract

Sensing and responding to photoperiod changes is essential for plants to adapt to seasonal

progression. Most of our understanding of how plants sense photoperiodic changes is through

studies on flowering time [1]. However, other aspects of plant development are regulated by the

photoperiod, including hypocotyl elongation [2, 3]. Unlike flowering, hypocotyl elongation

displays a greater plasticity to changes in the photoperiod with increases in daylength causing

greater inhibition of growth until a threshold is met [3]. Although we have a good understanding

of how the photoperiod regulates hypocotyl elongation [2, 3], previous investigations have only

looked at hypocotyl development under a constant photoperiod. It is unknown if changes in the

photoperiod during development influence hypocotyl elongation and whether the initial

photoperiod exposure continues to impact subsequent development. Here, we developed a

physiological assay to investigate this question. We have discovered that hypocotyl elongation is

influenced by a memory of past photoperiod exposure in Arabidopsis and Brassicaceae cultivars

used for microgreen agriculture. Photoperiodic memory persisted for multiple days, although it

weakened over time, and the strength of the memory was dependent on the genetic background.

We identified that phyB and ELF3, key regulators of hypocotyl development [4, 5], were required

for photoperiodic memory. Finally, we identified that the circadian clock is unlikely to function as

a repository for photoperiodic memory as circadian rhythms quickly re-aligned with the new

photoperiod. In summary, our work highlights for the first-time evidence of a photoperiodic

memory that can control plant development.

Results and Discussion

Designing a photoperiodic memory experimental assay

To understand whether hypocotyl development in *Arabidopsis thaliana* (Arabidopsis hereafter)

is influenced by prior exposure to a given different photoperiod (Figure 1A) we designed the

following assay. Here, stratified seeds were firstly germinated under constant white light for 24

hours to synchronize germination before seedlings were then provided two days of

photoperiodic information, either short-days (8/16 light/dark) or long-days (16/8 light/dark).

On day 3, seedlings were then either transferred to the reciprocal photoperiod or kept under the

existing photoperiod. Hypocotyl measurements were then taken on day 5 and day 7 to

understand the immediate and long-term impact of the shift, respectively. We transferred

seedlings on day 3 as hypocotyl development in de-etiolated Arabidopsis seedlings primarily

occurs between three- and four-days post-germination [6]. Supporting this, we found minimal

difference in median hypocotyl length of Ws-2 seedlings grown under SD and LD on day 3 (Figure

1B). To compare the relative amount of photoperiodic memory between Arabidopsis ecotypes

and mutants with different hypocotyl lengths, we implemented a photoperiodic memory index

(PMI, Figure 1C). The PMI scales the hypocotyl length relative to the median hypocotyl length of

growth under LD and SD conditions. Thus, seedlings with a PMI closer to 1 have a hypocotyl

length similar to those grown under constant SD, while a score closer to 0 indicates development

similar to that of constant LD.

Arabidopsis seedlings display photoperiodic memory

We firstly analyzed the photoperiodic memory of wild-type Arabidopsis Ws-2 seedlings. As

discussed above, Ws-2 seedlings were exposed to the photoperiodic shift on day 3 (Figure 2A).

By day 5, hypocotyl development in the Ws-2 seedlings that underwent either photoperiod shift

had a PMI of ~0.45 (Figure 2B, Supplementary Figure 1), indicating that exposure to the initial

photoperiod does influence hypocotyl development. By day 7, hypocotyl development in the

seedlings that underwent the photoperiod shift had diverged from each other and were more

closely resembling those under constant conditions (Figure 2C). However, for neither

photoperiod shift did hypocotyl elongation mirror that of seedlings grown under a constant

photoperiod (Figure 2C). Together, these results suggest that hypocotyl development displays

evidence of a lasting photoperiodic memory.

Next, we investigated whether other Arabidopsis ecotypes also displayed photoperiodic memory

and the heterogeneity in response between these ecotypes. Ecotypes were picked to cover a range

of latitudes from the equator to northernly latitudes (Supplementary Table 1). For all ecotypes,

hypocotyl elongation again displayed evidence of photoperiodic memory and this memory

continued to persist on day 7 (Figure 2B-C, Supplementary Figure 1). However, for the Sha, Cvi

and Bay-0 ecotypes the memory of the initial photoperiod was weaker than Ws-2 for both

photoperiod shifts. We also observed that certain ecotypes displayed stronger memory of a

particular photoperiod shift. For example, Tanzania (Tnz) had a stronger memory of the LD-SD

than SD-LD transitions on day 5, while Sapporo (Sap) displayed the opposite behavior (Figure

2B). However, neither effect persisted after an additional two days in the final photoperiod

(Figure 2C). Importantly, for all ecotypes tested except Ws-2, there was a difference in hypocotyl

length between the different photoperiod shifts on day 5 even though the seedlings were exposed

to the same amount of total light (Supplementary Figure 2-3). Therefore, it is the order of the

photoperiod exposure that is important in determining hypocotyl elongation.

phyB and ELF3 regulate the rate that Ws-2 responds to changes in day length

To understand the genetic pathways underpinning the response to the photoperiodic shifts, we

characterized different mutants in the Ws-2 ecotype background (wild type, WT), as this ecotype

had the strongest photoperiodic memory (Figure 2). We firstly focused on phyB and ELF3 as they

are key regulators of hypocotyl elongation [4, 5]. On day 5, the *elf3* mutant had reduced
photoperiodic memory of both photoperiod shifts relative to Ws-2 (Figure 2D, Supplementary
Figure 4-5). On day 7, hypocotyl elongation in the *elf3* mutant remained more sensitive to the SD-
LD shift but had a similar response to WT for the LD-SD shift (Figure 2E). For the *phyB* mutant,
seedlings phenocopied the WT response to the LD-SD on day 5 but then displayed reduced growth
by day 7 (Figure 2D-E). Under the SD-LD shift, *phyB* mutants had a similar PMI to those kept under
constant LD (Figure 2D), indicating a weak memory of the initial SD photoperiod. This phenotype
became further enhanced by day 7, with *phyB* mutants moved from SD-LD now shorter than those
kept under continuous LD (Figure 2E). We also analyzed the *elf3/phyB* mutant phenotype to
understand the genetic relationship between ELF3 and phyB. For the LD-SD shift, the *elf3/phyB*
mutant phenocopied the WT response at both timepoints (Figure 2D-E). Similarly, for the SD-LD
shift the *elf3/phyB* mutant had a similar response to WT at the first timepoint (Figure 2D).
However, at the second timepoint the *elf3/phyB* mutant had reduced photoperiodic memory than
WT (Figure 2E). Together, our results highlight a complex requirement of phyB and ELF3 in
regulating photoperiodic memory: ELF3 is necessary to regulate a plant's initial sensitivity to
changes in the photoperiod, while phyB prevents over-compensation for the SD-LD shift and
sensitizes plants to the shift to SD.
90
The impact of photoperiodic memory is dependent on developmental age
Hypocotyl elongation is dependent on the accumulation of PIFs at specific time of day. Maximal
hypocotyl elongation occurs under SD photoperiods as PIF accumulation coincides with the late
evening, where PIF activity displays minimal inhibition [3, 7]. This led us to investigate how the
dynamics of hypocotyl elongation changes across time in seedlings that underwent the
photoperiod shift. For this experiment, seeds were stratified in water rather than on the surface
of the agar-media for 3 days. This small change in the protocol delayed the onset of hypocotyl
emergence until day 3, the point at which the photoperiod shift initiates. Thus, the perception of
the initial photoperiod under these conditions occurred exclusively in the germinating seed. This
allowed us to subsequently investigate how photoperiodic perception in seeds influences
seedling development alongside the effects of the photoperiod shifts on the timing of hypocotyl
elongation.
103
Intriguingly, WT seedlings whose germinating seeds were exposed to LD, but then shifted to SD
had extremely rapid hypocotyl elongation, compared to all other conditions (Figure 3A). The
timing of growth in seeds that were shifted from LD-SD was consistent with those grown under
constant SD conditions (Figure 3B). However, the maximal growth rate in the seeds that were
shifted from LD-SD was much higher on the first night and remained elevated on the second night.

Thus, the shift from LD-SD caused a temporary acceleration of growth. For the SD-LD shift, Ws-2
displayed a less pronounced memory of the initial SD exposure, with overall hypocotyl elongation
(Figure 3A) and the timing of growth (Figure 3B) closely mirroring seedlings that were only
grown under constant LD. In summary, Ws-2 seeds display photoperiodic memory of prior LD
exposure, and this results in sustained levels of increased hypocotyl elongation when shifted to
SD.
115
As ELF3 and phyB were required for photoperiodic memory in seedlings, we investigated
whether they contributed to photoperiodic memory in seeds. phyB is functional in seeds [8, 9]
and the circadian clock is also active in imbibed seeds [10]. The role of ELF3 in seeds remains to
be determined. As in WT, the *elf3, phyB* and *elf3/phyB* mutants' overall growth and the timing of
growth showed weak memory of the SD-LD shift (Figure 3A-B). For the LD-SD shift, the *elf3*
mutant did not display enhanced growth on the first night (Figure 3B). Instead, the enhanced
growth was delayed until the second night. Furthermore, the peak timing of this growth was
shifted from late evening to dusk, where ELF3 functions within the evening complex to repress
PIF4/5 expression [11]. As with WT, *elf3* mutants displayed two evenings of accelerated growth
following the LD-SD shift before the rate of hypocotyl elongation became more consistent with
growth under constant SD (Figure 3B). In the *phyB* mutant, there was no evidence of accelerated
growth following from the shift from LD-SD. Instead, growth displayed two smaller peaks at dusk
and in the late evening. These two smaller peaks of growth likely compensate for the absence of
one predominant period of growth. The *elf3/phyB* mutant displayed similar phenotypes to the
respective single mutants for the LD-SD shift, with both a delay in the adjustment to the new
photoperiod and an absence of a major peak of growth (Figure 3B). Thus, phyB and ELF3
seemingly work independently to control the acceleration of growth and adjustment to the new
photoperiod, respectively, following the LD-SD shift.
134
Circadian clocks adapt to photoperiod shifts quickly in Arabidopsis seedlings
Although our data suggests that seeds and seedlings have a memory of the prior photoperiod, it
is unclear how this photoperiodic information is stored. One possible mechanism is through the
circadian clock. The circadian clock has a critical role in regulating photoperiodic responses in
plants [1-3]. Furthermore, the circadian clock is active in imbibed seeds [10] and the expression
of *PIF4* and *PIF5* display photoperiod-dependent changes in the phase of their expression [3].
Thus, we hypothesized that the circadian clock may function as a repository for photoperiodic
information.
143

To test this possibility, we analyzed the circadian waveform of the *CCR2::LUC* reporter in the Ws-
2 background. Here, seedlings were entrained for ten days under SD (8/16 light/dark) and then
shifted to LD for three days. As a control, seedlings were entrained and then kept under SD or LD
for the length of the experiment. For seedlings grown under either constant SD or LD conditions,
CCR2::LUC was phased to the early evening of each respective photoperiod as reported previously
[12, 13]. In seedlings that underwent the photoperiod shift, the phase of *CCR2::LUC* was initially
between the phase of *CCR2::LUC* in either constant SD or constant LD seedlings (Figure 4A).
However, by the second day post-shift, the phase of the *CCR2::LUC* reporter became aligned with
seedlings that were grown under constant LD. As hypocotyl elongation displayed memory of the
photoperiod information for at least four days post-shift (Figure 2-3), it is unlikely photoperiodic
information stored by the circadian clock can sufficiently explain the developmental responses
we have observed.

Although it is unlikely that circadian clock provides photoperiodic memory, we were interested
in understanding the heterogeneity of the response to the SD-LD shift. To investigate this, we
visualized the individual responses of each plant that underwent the shift using a functional
Principle Component Analysis (PCA) (see methods). We included three points for each individual
plant measured from the day before the shift, the day of the shift, and the day after the shift. Each
point represents the shape of the circadian rhythm across the respective day. This plot can be
interpreted in the same way as a traditional PCA plot, so days with similarly phased circadian
rhythms should group together. We found that there were individual plants that had rhythms that
resembled those under LD prior to the SD to LD shift and conversely there were individual plants
that had rhythms resembling SD plants two days after the shift (Figure 4B). However, this was a
very small proportion of the total plants evaluated. Therefore, the re-adjustment of circadian
phase following the photoperiod shift is very robust and uniform. In summary, hypocotyl
photoperiodic memory is unlikely to be directly controlled by the circadian clock.

Other Brassicaceae display strong photoperiodic memory.
Next, we investigated whether photoperiodic memory was present in other plants. Here, we
focused on microgreens that are grown in vertical farms as the hypocotyl is the primary
commercial product [14]. Microgreens grown under LD photoperiods have enhanced nutritional
quality, but the longer photoperiods are associated with higher operational energy costs.
Alongside reducing energy consumption, shorter photoperiods will also induce hypocotyl
elongation, increasing the quantity of yield. We hypothesized that photoperiodic memory may
enable us to combine the benefit of growth under both photoperiods. To investigate this
possibility, we firstly focused on kale and radish crops that are grown as microgreens within

vertical farms. As with Arabidopsis, kale and radish displayed evidence of photoperiodic memory
(Figure 4C, Supplementary Figure 6). However, all three cultivars that were tested displayed
stronger photoperiodic memory than Arabidopsis. Seedlings shifted from SD-LD had a PMI closer
to those grown under constant SD, while those shifted from LD-SD had a PMI similar to those
grown under constant LD. Moreover, the plants that experienced a SD to LD shift were also less
etiolated and had greener coloration, which is an agriculturally advantageous trait (Figure 4C).
These results suggest that photoperiod shifts may have immediate uses in optimizing growth
conditions in vertical farms.
188
Concluding statement
In summary, our work has a previously unknown mechanism of photoperiodic memory that
controls hypocotyl development. Distinct pathways seem to operate to control the response to
different directions of photoperiodic shifts, and the impact of the memory is dependent on the
developmental age of the plant. The genetic pathway that regulates photoperiodic memory is
unclear, but it seems to occur independently of transcriptional circadian rhythms. Photoperiodic
memory may have immediate direct agricultural implications by allowing the optimization of
nutritional quality, yield, and operational costs of vertical farms. It will be important to test the
presence of photoperiodic memory outside of Brassicaceae to understand the wider potential
benefits of photoperiodic memory in other agricultural settings.
199
Methods
<u>Plant lines and husbandry.</u>
The following *Arabidopsis thaliana* ecotypes were used in this work: Bay-0, Cape Verde Island
(Cvi), Sapporo (Sap), Shakdara (Sha), Tanzania-1 (Tnz-1) and Wassilewskija-2 (Ws-2). All
ecotypes harbored the *CCR2::LUC* reporter gene. Bay-0 *CCR2::LUC*, Cvi *CCR2::LUC*, Sha *CCR2::LUC*
and Ws-2 *CCR2::LUC* have been described previously [15-17]. For the Sap and Tnz *CCR2::LUC*
lines, the respective parent background were transformed with the *CCR2::LUC* reporter by floral
dip [18]. A luminesce transgenic plant was then identified and backcrossed to the respective
parental line. This backcrossed transgenic was then self-fertilized before a homozygous, single
transgene segregant was identified and bulked in the following generation. The following
Arabidopsis mutants in the Ws-2 background were also used: *elf3-4 CCR2::LUC*, *phyB-10*
CCR2::LUC, *elf3-4/phyB-10 CCR2::LUC*. These mutants have been previously described in earlier
work [4, 19, 20]. Prior to experiments, all lines were propagated on soil in environmentally
controlled growth chambers with a long-day photoperiod (16/8 L/D) at a light intensity of 100
$\mu mol/m^{-2}/s^{-1}$ and a set temperature of 22°C.
215

<u>Hypocotyl measurements.</u>

For the collection of data presented in figure 1, figure 2 and supplementary figure 1-4, all seeds were surface sterilized before being plated onto 1x Murashige and Skoog (MS) media with 1.5% w/v phytoagar (Duchefa), 1% w/v sucrose (Sigma) and 0.5 g/L MES (Sigma). Seeds were then stratified for 48-96 hours. After stratification, germination was induced by exposure to 75 μmol/m^{-2}/s^{-1} of white light for 24 hours and a constant temperature of 19.5°C. After 24 hours, germinating seeds were provided photoperiod information. They were either grown under short-days (SD, 6/18) or long-days (LD, 16/8) photoperiods with a constant temperature of 19.5°C and a light intensity of 75 μmol/m^{-2}/s^{-1}. After 48 hours, the photoperiod shifts were initiated. Seedlings were either kept under the initial photoperiod as a control or transferred to the reciprocal photoperiod. All photoperiod shifts occurred at ZT6, before lights off for the SD growth conditions. Measurements were taken on day 5 (timepoint 1) or day 7 (timepoint 2) at ZT6. All experiments were repeated twice with a similar result observed across the two experimental repeats. The presented data is a combination of the two experimental repeats, with each individual experiment including at least fifteen seedlings. Hypocotyl images were measured using imageJ [21]. For figures 1 and 2, hypocotyl measurements were scaled so that the median LD hypocotyl length was 0 and the median SD hypocotyl length was 1, using the formula: *Scaled length = (length – median length LD)/(median length SD – median length LD).* Adjusted p-values were calculated using TukeyHSD tests. We have included all statistical and raw hypocotyl measurements as a supplementary dataset (Supplementary File).

<u>Luciferase assay.</u>

Arabidopsis Ws-2 *CCR2::LUC* WT seeds were surface sterilized and plated on 1x MS medium containing 3% w/v sucrose (Sigma), 1.5% w/v phytoagar (Duchefa), 0.5 g/L MES (Sigma) and 15 μg/ml Hygromycin-B. After 5 days of stratification at 4°C, seeds were transferred to the specified photoperiod for 6 days. For all photoperiods, a constant temperature of 20°C was used and a light intensity of between 80 - 100 μmol/m^{-2}/s^{-1}. On day 6, seedlings were transferred into a 96 well plate containing the same media as described above but additionally 15 μl of 1 mM Luciferin. Plates were then placed back in the growth cabinet until they were then transferred to the TOPCOUNT (TC) at dawn on day 7. Plates in the TC were grown under the stated photoperiod with red plus blue light (2.4-3.6 μmol/m^{-2}/s^{-1} red light and 0.8-1.4 μmol/m^{-2}/s^{-1} blue light) and a constant temperature of 20°C for a further 4 days. For samples undergoing a photoperiod shift, this shift occurred on day 11, exactly 96 hours after being placed in the TC. The fda package in R was used to smooth curves using B-splines and perform functional principal component analysis.

<u>Continuous monitoring of hypocotyl elongation.</u>

Sterilized Arabidopsis seeds were stratified in darkness at 4°C for 72 hours. To avoid contact of seedlings with the media surface during hypocotyl elongation, two strips were cut and removed from the cultivation medium. Solid *A. thaliana* solution (ATS) nutrient medium [22] with 1% w/v sucrose was used as the cultivation media for these experiments. Seeds were sown in small indentations on the surface of the cut to prevent excessive movement as described previously [23]. Plates were placed vertically at 21°C and constant light (LL, 40 µmol m^{-2}s^{-1}) for one day to allow uniform germination of seeds, before being transferred to the initial photoperiod: either LD (16/8 light/dark) or SD (8/16 light/dark) conditions. After two days, plates were transferred to the alternative photoperiod (SD or LD, dawn-aligned), or were kept under the initial photoperiod. The analysis of hypocotyl elongation then started 72 hours after sowing (ZT0). Seedlings were illuminated by two infrared light emitting diodes and were imaged hourly by a Panasonic G5 camera modified with a long pass filter (830 nm). To detect minor changes between time points, image stacks were imported as "Image Sequence" in ImageJ. Each seedling (n≳6) was analyzed separately, and the hypocotyl length was measured using the "Segmented Line" tool, by slightly adjusting the control nodes between time points.

<u>Hypocotyl Elongation in Microgreens</u>

Microgreens (Supplementary Table 2) seeds were sown into trays containing a compost mix comprised of equal parts F2+S (Levington), coir and vermiculite supplemented with 0.7 g/L osmocote fertiliser. Seeds were sown approximately 1cm below the surface of the soil. Microgreens were then grown under either a long-day (16/8) or short-day photoperiod (8/16) photoperiod with a constant temperature of 19°C, 60% humidity and 70 µmol/m^{-2}/s^{-1} of light. Trays were inspected daily to determine when individual hypocotyls had emerged from the soil. Once 50% of the respective microgreen cultivar had a visible hypocotyl, seedlings were grown for three days under the initial photoperiod. This period was determined as the initial photoperiod exposure. After this period, the microgreens were then either switched to the reciprocal photoperiod or kept under the same photoperiod as a control for a further 3 days. Trays were switched at ZT5 to ensure that most of the plants that underwent photoperiod shifts had equal exposure to each photoperiod. Each experiment was carried out three times with 40 individual plants imaged per variety per experiment. To image the microgreen hypocotyls, leaves were firstly dissected before the hypocotyls were removed from the soil and laid flat in an Epson scanner. High-resolution scans were then taken. Images were imported into ImageJ for measurement. All imaging took place at ZT5.

 Figure Legends.

Figure 1 – Designing an assay to test for photoperiodic memory. (A) In the context of this

manuscript, we define a 'memory-less system' as one whose growth only depends on the current

photoperiod. Conversely, 'memory' refers to growth that also depends on prior environmental

conditions. (B) Hypocotyl growth in Arabidopsis seedlings occurs predominantly during the

period after the photoperiod shift has occurred. (C) A hypocotyl memory index was used to

compare photoperiodic memory and the sensitivity to photoperiodic shifts between ecotypes and

mutants that have large magnitudes differences in hypocotyl elongation. Hypocotyl lengths were

scaled so that the median hypocotyl length at LD (16/8hr light/dark) is 0 and the median

hypocotyl length at SD (8hr/16hr light/dark) is 1.

Figure 2 – Hypocotyl development in Arabidopsis displays evidence of photoperiodic memory.
(A) Diagram explaining the four different photoperiodic growth conditions, the day phase is
white, and the night phase is black. Different Arabidopsis wild type accessions were either
measured (B) two or (C) four days after the respective photoperiod shift or were kept under the
same photoperiod for the duration of the experiment. Photoperiodic memory response in *elf3-4*,
phyB-10 and *elf3-4/phyB-10* mutants after either (D) two days or (E) four days following the
photoperiod shift. Letters signify a significant adjusted p-value of $p < 0.01$, determined by
TukeyHSD.

Figure 3 – Photoperiodic memory response is dependent on the developmental age of the plant.

(A) Hypocotyl length in Arabidopsis Ws-2 wild-type, *elf3-4*, *phyB-10* and *elf3-4/phyB-10* seedlings

under different photoperiodic conditions across time. (B) Hypocotyl growth rate (mm/hr) in the

respective genotypes. Highlighted above the plots are the respective photoperiods, long-days in

green and short-days in blue. The color boxes highlight the dark phase under the respective

photoperiod. All measurements were taken after the photoperiod shift had occurred at ZT72.

Note that the seeds germinated later in this experiment than in the experiment depicted in Figure

2, with hypocotyl emergence occurring primarily on day 3.

Figure 4 – The circadian clock is unlikely to store photoperiodic memory in Arabidopsis. (A)

Waveform of *CCR2::LUC* bioluminescence in Ws-2 wild-type seedlings grown under continuous

SD (blue) or LD (green) or in seedlings shifted from SD to LD (yellow). (B) A functional PCA (fPCA)

plot, where each point represents a different bioluminescence curve for an individual plant, color-

coded by their photoperiod condition that plant was exposed to, using a color-scheme consistent

with (A). fPCA is interpreted in the same way as a standard PCA, as a projection of data (in this

case, coefficients of a complex B-spline function) to a lower dimensional space. (C) Kale (CN

seeds), Kale (Tozer seeds) and radish plants used in microgreen agriculture displayed evidence

of photoperiodic memory. Letters signify a significant adjusted p-value of $p < 0.01$, determined

by TukeyHSD.

Supplementary Figure 1 – Representative scans of seedlings from wild-type Arabidopsis

ecotypes used in this work. Images are from day 5, two days after the photoperiod shift. Scale bar

is 5 mm.

Supplementary Figure 2 – Hypocotyl length (mm) of wild-type accessions sorted by total light exposure (hours) on day 5. Dark blue is constant SD, purple are seedlings shifted from LD-SD, yellow is seedlings shifted from LD-SD and green is constant LD.

Supplementary Figure 3 - Hypocotyl length (mm) of wild-type accessions sorted by total light exposure (hours) on day 7. Dark blue is constant SD, purple are seedlings shifted from LD-SD, yellow is seedlings shifted from LD-SD and green is constant LD.

Supplementary Figure 4 – Representative scans of seedlings from wild type Arabidopsis Ws-2, *elf3-4*, *phyB-10* and *elf3-4/phyB-10.* Images are from day 5, the first measurement point. Scale bar is 5 mm.

Supplementary Figure 5 - Hypocotyl length (mm) of wild-type Ws-2, *elf3-4*, *phyB-10* and *elf3-4/phyB-10* sorted by total light exposure (hours) on (A) day 5 and (B) day 7. Dark blue is constant SD, purple are seedlings shifted from LD-SD, yellow is seedlings shifted from LD-SD and green is constant LD.

Supplementary Figure 6 – Representative scans of microgreen seedlings. Microgreens are approximately 11 days old at the end of the experiment Scale bar is 15 mm.

Supplementary File – Raw data and statistical tests associated with the figures of this manuscript.

Acknowledgements

We would like to thank Maarten Koornneef for providing the Tnz-1 seeds. We would also like to acknowledge A. L. Tozer Limited for providing us with radish and kale seeds and the York Biology horticulture staff for their assistance and support with this work.

Conflicts of Interest

D.E. and W.C. are currently part of a Better Food for All Innovate UK grant in collaboration with Farm Urban LTD, A.L. Tozer Limited, Vertically Urban Limited, Nutritank Community Interest Company and Alder Hey Living Hospital Limited.

Grant Funding

We would like to acknowledge the following funding sources: the Royal Society (RGS\R2\212345: D.E.), Biotechnology and Biological Sciences Research Council (Responsive Mode) (BB/V006665/1: S.L, K.M., D.E. and S.J.D.), the Biotechnology and Biological Sciences Research Council (White Rose Doctoral Training Partnership) (BB/T007222/1: E.J.R. and G.Y.W.V.), and GenerationResearch (W.C.).

Supplementary Table 1 – Arabidopsis ecotypes used in this work.

Ecotype	Latitude	Longitude	Country
Wassilewskija-2 (Ws-2)	52.3	30.0	Belarus
Tanzania (Tnz)	-2.9	36.2	Tanzania
Bay-0	49.0	11.0	Germany
Shakdara (Sha)	38.4	68.5	Tajikistan
Cape Verde Island (Cvi)	15.1	-23.6	Cape Verde Island
Sapporo (Sap)	43.1	141.3	Japan

Supplementary Table 2 – Microgreen cultivars used in this work

Microgreen	Species	Variety
Kale (Tozer Seeds)	*Brassica oleracea*	Dwarf Green Curled
Kale (CN Seeds)	*Brassica oleracea*	Dwarf Blue
Radish	*Raphanus raphanistrum*	Sangrin

References

1. Song, Y.H., et al., *Photoperiodic flowering: time measurement mechanisms in leaves.* Annu Rev Plant Biol, 2015. **66**: p. 441-64.

2. Nozue, K., et al., *Rhythmic growth explained by coincidence between internal and external cues.* Nature, 2007. **448**(7151): p. 358-61.

3. Niwa, Y., T. Yamashino, and T. Mizuno, *The circadian clock regulates the photoperiodic response of hypocotyl elongation through a coincidence mechanism in Arabidopsis thaliana.* Plant Cell Physiol, 2009. **50**(4): p. 838-54.

4. Reed, J.W., et al., *Mutations in the gene for the red/far-red light receptor phytochrome B alter cell elongation and physiological responses throughout Arabidopsis development.* The Plant Cell, 1993. **5**(2): p. 147-157.

5. Zagotta, M.T., et al., *The Arabidopsis ELF3 gene regulates vegetative photomorphogenesis and the photoperiodic induction of flowering.* Plant J, 1996. **10**(4): p. 691-702.

6. Gendreau, E., et al., *Cellular Basis of Hypocotyl Growth in Arabidopsis thaliana.* Plant Physiology, 1997.

7. Leivar, P. and E. Monte, *PIFs: systems integrators in plant development.* Plant Cell, 2014. **26**(1): p. 56-78.

8. Lee, K.P., et al., *Spatially and genetically distinct control of seed germination by phytochromes A and B.* Genes & Development, 2012. **26**(17): p. 1984-1996.

9. Piskurewicz, U., et al., *The Arabidopsis endosperm is a temperature-sensing tissue that implements seed thermoinhibition through phyB.* Nature Communications, 2023. **14**(1): p. 1202.

10. Zhong, H.H., et al., *Imbibition, but not release from stratification, sets the circadian clock in Arabidopsis seedlings.* The Plant Cell, 1998. **10**(12): p. 2005-2017.

11. Nusinow, D.A., et al., *The ELF4-ELF3-LUX complex links the circadian clock to diurnal control of hypocotyl growth.* Nature, 2011. **475**(7356): p. 398-402.

12. Strayer, C., et al., *Cloning of the Arabidopsis Clock Gene TOC1, an Autoregulatory Response Regulator Homolog.* Science, 2000. **289**(5480): p. 768-771.

13. Schultz, T.F., et al., *A role for LKP2 in the circadian clock of Arabidopsis.* Plant Cell, 2001. **13**(12): p. 2659-70.

14. Bhaswant, M., et al., *Microgreens-A Comprehensive Review of Bioactive Molecules and Health Benefits.* Molecules, 2023. **28**(2).

15. Doyle, M.R., et al., *The ELF4 gene controls circadian rhythms and flowering time in Arabidopsis thaliana.* Nature, 2002. **419**(6902): p. 74-7.

16. Boikoglou, E., et al., *Environmental memory from a circadian oscillator: the Arabidopsis thaliana clock differentially integrates perception of photic vs. thermal entrainment.* Genetics, 2011. **189**(2): p. 655-64.

17. Anwer, M.U., et al., *Natural variation reveals that intracellular distribution of ELF3 protein is associated with function in the circadian clock.* eLife, 2014. **3**: p. e02206.

18. Davis, A.M., et al., *Protocol: Streamlined sub-protocols for floral-dip transformation and selection of transformants in Arabidopsis thaliana.* Plant Methods, 2009. **5**: p. 3.

19. Hicks, K.A., et al., *Conditional circadian dysfunction of the Arabidopsis early-flowering 3 mutant.* Science, 1996. **274**(5288): p. 790-792.

20. Ronald, J., et al., *Cellular localization of Arabidopsis EARLY FLOWERING3 is*
responsive to light quality. Plant Physiology, 2022.
21. Schneider, C.A., W.S. Rasband, and K.W. Eliceiri, *NIH Image to ImageJ: 25 years*
of image analysis. Nature Methods, 2012. **9**(7): p. 671-675.
22. Lincoln, C., J.H. Britton, and M. Estelle, *Growth and development of the axr1*
mutants of Arabidopsis. Plant Cell, 1990. **2**(11): p. 1071-80.
23. Janitza, P., et al., *A Guide to Quantify Arabidopsis Seedling*
Thermomorphogenesis at Single Timepoints and by Interval Monitoring, in
Thermomorphogenesis: Methods and Protocols, M. Chen, Editor. 2024, Springer
US: New York, NY. p. 3-16.