STEMorph: A Set of Morphed Emotional Face Stimuli from Angry to Happy Derived

from *NimStim*

Mohammad Ebrahim Katebi[1,2*], Mohammad Hossein Ghafari[3*], and Tara Ghafari[4]

[1] School of Cognitive Sciences, Institute for Research in Fundamental Sciences (IPM), Tehran,

Iran

[2] School of Medicine, Tehran University of Medical Sciences, Tehran, Iran

[3] School of Medicine, Shahid Beheshti University of Medical Sciences, Tehran, Iran

[4] Centre for Human Brain Health, School of Psychology, University of Birmingham, UK

16   **Author Note**

17   [*] These authors contributed equally.

18   Correspondence concerning this article should be addressed to Tara Ghafari, Centre for

19   Human Brain Health, School of Psychology, University of Birmingham, Birmingham, UK, B15 2TT. Email:

20   t.ghafari@bham.ac.uk

21   **Abstract**

Emotion recognition through facial expressions is crucial for interpreting social cues. However,

it is often influenced by biases, i.e, systematic recognition advantages for particular emotions.

Nevertheless, these biases are inconsistently reported across studies, likely due to methodological

variations which underline the necessity for a standardized approach. Traditional face morphing

methods can create unnatural-looking stimuli, and may confound the interpretation of emotions. To

address this issue, we here introduce *STEMorph*, a validated stimulus set based on the NimStim set.

We employed neutral-anchored morphing and neural-network-generated masks to ensure the natural

appearance and integrity of the depicted emotions. We validated our stimulus set by having

participants rate each face on a 9-point scale ranging from angry to happy, assessing the perceived

emotional intensity. *STEMorph's* validity was confirmed through linear regression analysis, showing a

strong correlation between subjective ratings and targeted emotional states. Moreover, aligning with

previous research highlighting gender as a key factor in emotion recognition, *STEMorph* also exhibits

variations in ratings across gender subgroups. *STEMorph's* reliability was confirmed through a two-

week follow-up rating session with a subgroup of the same participants. By introducing a controlled,

validated, and ecologically valid stimulus set of emotional faces, *STEMorph* paves the way for further

investigations into facial emotion recognition, deepening our understanding of this vital aspect of

human interaction.

Keywords: *STEMorph*, Stimulus Set, Facial Emotional Expressions, Morphed Stimuli, *NimStim*

41   **Introduction**

Emotion recognition is a fundamental aspect of human social interaction, allowing individuals

to interpret and respond appropriately to the emotional states of others. Facial expressions serve as a

primary modality for conveying emotions, providing valuable cues about an individual's internal states

(Russell, 1994). However, despite its significance, the process of emotion recognition is not devoid of

biases. These biases manifest as systematic recognition advantages for particular emotions and have

significant implications for both clinical and non-clinical populations (Silvia et al., 2006).

Recognition biases can vary across different conditions. Individuals with major depressive

disorder (MDD) tend to interpret neutral or ambiguous faces as sad or threatening, reflecting a

negativity bias. This aligns with findings that people with MDD are more attuned to negative emotions

and less responsive to positive ones, which impairs their ability to interpret emotional cues

(Leppänen, 2006; Penton-Voak et al., 2017). Such biases may reinforce mood-congruent cognitive

distortions, contributing to the maintenance and exacerbation of depressive symptoms (Harmer et al.,

2009; Penton-Voak et al., 2012). Similarly, individuals with generalized anxiety disorder (GAD) and

social anxiety disorder (SAD) exhibit hypervigilance toward negative emotional cues, such as anger or

fear, which can lead to misinterpretations of facial expressions and reinforce maladaptive emotional

responses (Surguladze et al., 2004; Waters et al., 2008). Therefore, these biases can significantly

impact social interactions and clinical outcomes; emphasizing the importance of their precise

measurement for both research and clinical applications.

Studies often report inconsistent results regarding the direction and magnitude of emotional

recognition biases (Münkler et al., 2015). For instance, one study found that healthy individuals did

not demonstrate a bias toward happiness or sadness when presented with neutral facial expressions

(Leppänen et al., 2004). While another study showed that healthy participants tended to perceive

neutral faces as expressing happiness, whereas depressed individuals interpreted them as neutral

(Surguladze et al., 2004). It is uncertain whether these mixed findings reflect genuine individual

differences in emotion perception or are due to methodological variations between the studies

(Vikhanova et al., 2022). Particularly, the quality and ecological validity of facial stimuli can significantly

impact participants' responses. While standardized photographic stimuli are commonly used for their

control over visual properties, they may not fully capture the complexities of real-world facial

expressions, which can lead to discrepancies in findings. Research indicates that more naturalistic

stimuli, such as dynamic or spontaneous facial expressions, enhance ecological validity but introduce

variability that complicates experimental control (Dawel et al., 2022; Fernando, 2015). Consequently,

A key challenge lies in creating stimuli that maintain natural appearance while allowing precise control

over emotional expressions (Dawel et al., 2022; Fernando, 2015).

One widely used approach to generating a continuum of emotions is through face morphing

techniques. Face morphing is an image processing technique employed to achieve transformation

from one face to another and facilitates a gradual transition between two facial images. By morphing

facial features between two extreme emotional expressions, researchers generate a spectrum of face

images with ambiguous emotional expressions for participants to interpret (Joormann & Gotlib, 2006;

Paiva-Silva et al., 2016). However, traditional face morphing presents challenges, particularly

concerning the formation of unrealistic and unnatural facial stimuli (Paiva-Silva et al., 2016) which may

confound interpretations and obscure real emotion recognition processes.

To maintain naturalism, proper masking of the face images (extracting the face area while

removing the background) is required, as morphing introduces artifacts in non-face regions. Using no

mask (full head) or simple oval/square masks which are employed by many studies (Dawel et al., 2022)

may cause distortions that risk emotional content loss, since various facial features hold different

degrees of significance depending on the type of expression (Elfenbein & Ambady, 2002; Schurgin et

al., 2014). For instance, the mouth region is particularly salient for recognizing happiness, while the

eyes are more critical for identifying fear or sadness (Guarnera et al., 2018). These regional differences

can lead to biases in emotion perception, as individuals may focus on different facial features based on

cultural or individual preferences (Elfenbein & Ambady, 2002). These underscore the necessity for

stimuli preserving anatomical fidelity through methods like neutral-anchored morphing (Vikhanova et

al., 2022) and standardized masking. However, there are no existing publicly available stimulus set

without these methodological confounds (Dawel et al., 2022; Paiva-Silva et al., 2016; Testerman,

2023). While sets like the Karolinska Directed Emotional Faces (KDEF) (Goeleven et al., 2008) and the

Pictures of Facial Affect (POFA) (Ekman, 1976) provide valuable resources, they often rely on basic

morphing techniques that can produce artifacts, particularly around facial features crucial for emotion

recognition such as the eyes and mouth. Furthermore, many existing sets use simple geometric masks

(oval or rectangular) that can create artificial boundaries and reduce ecological validity (Dawel et al.,

2022; Paiva-Silva et al., 2016). To address this gap, we developed and validated a new stimulus set

utilizing neutral-anchored morphing alongside neural-network-generated masks, to optimize

naturalism and coherence.

We derived the stimuli from the *NimStim set of facial expressions* (Tottenham et al., 2009)

which is a comprehensive collection of facial images portraying a spectrum of emotions including

happy, angry, and neutral expressions. The selection of anger and happiness as the target emotions for

this study was based on their distinct valence and arousal profiles, which make them suitable for

exploring biases in emotion recognition. Additionally, these emotions are commonly used in emotion

research (Silvia et al., 2006; Vikhanova et al., 2022), facilitating comparisons with existing literature.

Investigating the transition between these emotions allows us to examine biases in emotional

perception across a clear positive-negative emotional gradient, which is particularly relevant for

studies on mood disorders and social cognition (Silvia et al., 2006). We adopt a nuanced approach to

morphing emotional faces by generating distinct naturalistic face masks for each image using neural

networks. Furthermore, we utilize the neutral facial expression as a central anchor point to enhance

the naturalistic quality of the stimuli. We then evaluate the validity and reliability of *STEMorph* by

asking a group of participants to provide ratings of the emotional content of these stimuli.

116   **Method**

117   **Participants**

118   A total of 50 participants (age range: 18-26, mean age: 21.5, 22 females) were recruited for

the validity testing and post hoc power analysis confirmed that the sample size was sufficient.

Participants were medical students from Tehran University of Medical Sciences (TUMS) and Shahid

Beheshti University of Medical Sciences (SBUMS). All participants reported normal or corrected-to-

normal vision as well as no history of diagnosed neurological or psychiatric disorders., Informed

consent was obtained from all participants, and they were compensated for their time with money or

course credit. This study was conducted with ethical approval from the ethics committee of Shahid

Beheshti University of Medical Sciences (IR.SBMU.MSP.REC.1396.757).

126   **Experiment design & Face Generation**

127   We used face morphing to produce a linear transition of emotions from anger to happiness

using neutral expression as an intermediary point. The stimuli were derived from the *NimStim set of*

facial expressions (Tottenham et al., 2009). Each emotional expression has an open-mouth and closed-

mouth variation in *NimStim*, from which we selected closed-mouth variations. *Abrosoft Fantamorph*

(Abrosoft Fantamorph Deluxe 5.4.8) software was used for morphing, involving nine steps to transition

from the angry face to neutrality and subsequently to the happy face. We used the original facial

expressions of anger, happiness, and neutral as reference points and generated three intermediate

emotional states between each original face.

135   We selected 22 individuals (11 males and 11 females) from *NimStim*, prioritizing those whose

facial expressions facilitated smooth and natural morphing while avoiding individuals with extreme

facial expressions (highly exaggerated or intense displays of emotions). The final set is composed of

198 distinct faces, representing 9 different emotional steps for each of the 22 individuals (Figure 1).

Figure 1

Example Stimuli from the Set.

Note. Top row illustrates the two ends of the spectrum (happy and angry) along with the middle step

(neutral). Middle row shows the intermediary morphing stages without masking. Bottom row shows the final

outcome with corresponding neural-network-generated masks applied to each morphing step.

To address artifacts introduced in the ears, hair, and other non-landmark regions of the face

during conventional face morphing, we generated distinct naturalistic face masks for each image using

the *FaceSwap (Faceswap)* software, which utilizes the *Bisenet-FP (face-parsing.PyTorch)* neural

network. *BiSeNet-FP* is a modified lightweight neural network designed for face parsing. The model

was set to not include hair or ears and extract only the face region. Manual editing was performed to

correct for any artifacts arising from stray hair (e.g., a strand of hair overlapping the face resulted in a

split in the face) or other extraneous features.

Experimental Setup

Participants were asked to rate the emotional state of each presented stimulus on a 9-point

scale from angry to happy. The task was developed and implemented using *Psychtoolbox (Brainard,*

1997) within *MATLAB R2022a* (Figure 2). Participants were comfortably seated in a dimly lit room

facing a laptop monitor. The resolution of monitor screen was 1920 × 1080 and it was positioned ~54

cm from the participant. Participants were instructed to freely observe the images without fixation

requirements.

Figure 2

Schematic presentation of the experimental design.

Note. Stimuli appeared at the center of the screen, and participants rated the emotional intensity of

each face within a 4000 ms viewing window without fixation restrictions. Each stimulus was followed by a

randomized inter-trial interval (ITI) of 1000 to 2000 ms, during which a noise mask was displayed.

The task started with pressing a key, after which participants were asked to rate the emotional

state of each presented face based on a 9-point scale where 1 indicated the angriest emotional state

and 9 indicated the happiest emotional state. The exact instructions were: 'Please rate the emotional

state of each face on a scale from 1 to 9, where 1 represents the angriest expression and 9 represents

the happiest expression.' To control for motor and numerical biases, the scale was reversed for half of

the participants. For these individuals, the instructions were adjusted accordingly, and their responses

were rescaled during analysis to align with the original scale (1 = angry, 9 = happy). Each participant

had a 4-second window to provide their rating; in the absence of a response within this timeframe,

the task automatically proceeded to the subsequent face.

Each of the 198 stimuli was presented twice in a randomized order, resulting in a total of 396

trials per participant, interspersed by two 2-minute breaks. Following each trial, a randomized inter-

trial interval (ITI) of 1000 to 2000 ms ensued, during which a noise mask was displayed at the center

of the screen to prevent from interference between trials and minimize priming effects (Aguado et al.,

2015; Sweeny et al., 2009). To generate the noise mask the stimulus presented in the previous trial

was scrambled into 4-pixel squares.

Reliability Assessment

To assess reliability, the same behavioral task was readministered to a subgroup of 23

participants two weeks following their initial participation. This subgroup consisted of participants

who agreed to return for retesting (age range: 18-26, mean age: 22, 5 females).

Data Analysis

Validity

To investigate the relationship between morph levels and subjective ratings while accounting

for gender effects, we conducted an Ordinary Least Squares (OLS) linear regression analysis using

statsmodels (Seabold & Perktold, 2010). The model included morph step (1-9), face gender

(male/female), participant gender (male/female), and all two-way interactions as predictors of

participants' emotional ratings as below:

Equation (1):

$$\textbf{\textit{Participans' Rating}} = \boldsymbol{\beta_0} + \boldsymbol{\beta_1 MS} + \boldsymbol{\beta_2 FG} + \boldsymbol{\beta_3 PG} + \boldsymbol{\beta_4(MS \times FG)} + \boldsymbol{\beta_5(MS \times PG)} + \boldsymbol{\beta_6(FG \times PG)} + \boldsymbol{\varepsilon}$$

where MS corresponds to Morph Step (1-9), FG to Face Gender (0=female, 1=male), PG to

Participant Gender (0=female, 1=male), $\boldsymbol{\beta}$ to coefficients of each regressor, and ε to error term. Given

that this is an effort to validate a stimulus set, all collected participants were included in analyses, with

no outlier removal being performed. We then calculated partial residuals for morph steps by removing

variance explained by gender-related terms to isolate the morph-rating relationship.

To quantify evidence strength against null models, we computed Bayes Factors using the

BayesFactor (Morey & Rouder, 2023) R package. Statistical power (1- β error) was determined through

*G*Power* (Faul et al., 2009) with α = 0.05 relative to a non-correlation model ($\rho\ H_0 = 0$). To visualize

partial regression plots we generated violin plots using matplotlib (Hunter, 2007) and seaborn

(Waskom, 2021).

Gender Subgroup Analysis

To examine gender-specific effects on emotion perception, we extended our primary

regression framework (Equation 1) through partial regression analysis of participant gender and face

gender subgroups. We computed condition-specific partial residuals from the full model (Equation 1)

to isolate morph-rating relationships within gender subgroups.

For participant gender subgroups, we calculated partial residuals by retaining morph step,

participant gender (Female/Male), and their interaction (Morph Step × Participant Gender), while

regressing out variance associated with face gender effects (Face Gender + Morph Step × Face Gender

+ Face Gender × Participant Gender) (Equation 2). Additionally, in face gender subgroup analysis,

morph step, face gender (Female/Male), and their interaction (Morph Step × Face Gender) were

preserved while variance attributable to participant gender effects (Participant Gender + Morph Step

× Participant Gender + Face Gender × Participant Gender) were removed (Equation 3). This approach

enabled examination of gender-specific emotion perception while controlling for cross-gender

interactions.

Equation (2)

$$\textbf{\textit{Participant Gender Partial Rating}} = \textbf{\textit{Particiapans' Rating}} - (\boldsymbol{\beta_2}\textbf{\textit{FG}} + \boldsymbol{\beta_4}(\textbf{\textit{MS}} \times \textbf{\textit{FG}}) + \boldsymbol{\beta_6}(\textbf{\textit{FG}} \times \textbf{\textit{PG}}))$$

Equation (3)

$$\textbf{\textit{Face Gender Partial Rating}} = \textbf{\textit{Particiapans' Rating}} - (\boldsymbol{\beta_3}\textbf{\textit{PG}} + \boldsymbol{\beta_5}(\textbf{\textit{MS}} \times \textbf{\textit{PG}}) + \boldsymbol{\beta_6}(\textbf{\textit{FG}} \times \textbf{\textit{PG}}))$$

Where MS corresponds to Morph Step (1-9), FG to Face Gender (0=female, 1=male), PG to Participant

Gender (0=female, 1=male), $\boldsymbol{\beta}$ to coefficients of each regressor and ε to error term.

Reliability

To investigate reliability, we analyzed data from participants who engaged in the retesting (n =

23) and compared it with their validity data. We exactly matched trials from the two tasks based on

the presented face and performed OLS linear regression between participants' ratings in the validity

task and the retest conducted two weeks later as below:

Equation (4):

$$\boldsymbol{Particiapans'\ Retest\ Rating\ =\ \beta_0\ +\ \beta_1 VR\ +\ \varepsilon}$$

where VR corresponds to Validity Ratings (1-9), $\boldsymbol{\beta}$ to coefficient of VR, and ε to error term. We

then calculated the Bayes Factor and Statistical power using the methods previously described.

234   **Results**

The aim of this study was to develop and validate a new stimulus set for emotion recognition

utilizing neutral-anchored face morphing and neural-network-generated masks to optimize naturalism

and coherence. A total of 50 participants were recruited to rate the emotional content of the face

stimuli on a 9-point scale ranging from angry to happy with a subset of 23 participants returning for a

reliability assessment two weeks later.

Validity

We employed linear regression to examine the emotional ratings provided by participants

(mean = 4.85, SD = 2.14) against the emotional morph level of the stimulus set, while accounting for

face gender, participant gender, and their interactions (Table 1). The model revealed a strong positive

association between morph step and emotional ratings, with a coefficient estimate of 0.735 (95% CI

[0.724, 0.746], p < 0.001). The model explained a substantial proportion of variance in ratings (R^2 =

0.741, p < 0.001; Figure 3). The Bayes factor comparing the model to the null hypothesis (R^2 = 0)

approached infinity, indicating strong evidence in favor of the model. Furthermore, the statistical

power (1 − β error) relative to a non-correlation model (ρ H_o = 0) was 0.999, demonstrating robust

sensitivity to detect the observed effect.

Figure 3

Validity of emotional ratings across morphing steps.

.

Note. Morphing steps range from angry (1) to happy (9) on the x-axis. The y-axis presents participants'

emotional ratings as partial residuals after removing variance explained by gender-related terms (face gender,

participant gender, and interactions). Violin plots show kernel density estimates of rating distributions at each

morph step. Extreme outliers (±2SD from morph-level means) were excluded only for visualization clarity

purposes (all data retained in statistical models). Small grey dots depict raw residuals, large black dots indicate

mean ratings, and the dashed line shows the regression slope. The equation reflecting the full mode is:

Participants' Rating = 0.735 × MS + -0.248 × FG + 0.326 × PG + 0.026 × (MS×FG) + -0.065 × (MS×PG) + -0.008 ×

(FG×PG) + 1.236 where MS corresponds to Morph Step (1-9), FG to Face Gender (0=female, 1=male) and PG to

Participant Gender (0=female, 1=male). Colors correspond to morph steps.

Figure 4

Regression coefficients for the model.

.

Note. Beta estimates (95% CIs) from the OLS regression model predicting emotional ratings. Predictors

include Morph Step (MS; 1–9), Face Gender (FG; 0 = female, 1 = male), Participant Gender (PG; 0 = female, 1 =

male), and their two-way interactions. Asterisks denote terms significant at α = 0.05.

Table 1

Ordinary Least Squares Regression Results.

Predictor	β	SE	t	P > |t|	[0.025	0.975]
const	1.236	0.032	38.715	**<0.001**	1.174	1.299
Morph Step	0.735	0.005	134.882	**<0.001**	0.724	0.746
Face_Gender	-0.248	0.038	-6.485	**<0.001**	-0.323	-0.173
Participant_Gender	0.326	0.038	8.665	**<0.001**	0.252	0.400
MS_FG	0.026	0.006	4.324	**<0.001**	0.014	0.038
MS_PG	-0.065	0.006	-10.669	**<0.001**	-0.077	-0.053
FG_PG	-0.008	0.031	-0.252	0.801	-0.069	0.054

Note. OLS regression results from the full interactive model (Equation 1). The model explains 74.1% of

variance in emotional ratings (R² = 0.741, N = 19,596 observations). Morph Step (MS) represents emotion

intensity progression (1: angry to 9: happy). Face/Participant Gender coded 0 = Female, 1 = Male. Significant

terms (α = 0.05) marked with asterisks. The non-significant Face × Participant Gender interaction (p = 0.801)

indicates additive rather than multiplicative gender effects.

Gender Subgroup Validity Analysis

To assess the robustness of validity across gender dimensions, we included gender of face

stimuli and gender of participant in our model as described above (Equation 1). Significant effects

were observed for face gender (β =-0.248, p < 0.001), participant gender (β = 0.326, p < 0.001), and

the interactions between morph step and face gender (β = 0.026, p < 0.001) as well as morph step and

participant gender (β =-0.065, p < 0.001; see Figure 4). The interaction between face gender and

participant gender was not significant (β =-0.008, p = 0.801). In other words, female faces received

higher ratings than male faces, whereas male participants provided higher ratings than females.

To evaluate the validity of participants' ratings in gender subgroups, we analyzed partial

residual for each subgroup. Strong predictive validity across subgroups were observed. Morph step

slopes for face gender subgroups (Figure 5a) were nearly identical (R^2 female faces: 0.759; R^2 male

faces: 0.759), with slopes and interactions as below:

Female Faces: 1.24 + 0.74 × Morph Step; Male Faces: 0.99 + 0.76 × Morph Step

Participant gender subgroups exhibited greater divergence where female raters showed

steeper slopes (R^2 = 0.748) than male raters (R^2 = 0.721, Figure 5b), suggesting gender-related

differences in emotional rating:

Female Participants: 1.24 + 0.74 × Morph Step; Male Participants: 1.56 + 0.67 × Morph Step

The strong R^2 of the partial regression analysis across subgroups confirm robust validity across

gender dimensions while highlighting nuanced differences in how participant gender relates to

emotion perception accuracy.

Figure 5

Gender-based subgroup analysis of the validity data.

Note. A) Face gender subgroups: Female faces (blue) and male faces (orange) show comparable morph-

rating slopes derived from partial regression analysis. B) Participant gender subgroups: Female participants

(blue) demonstrate steeper slope than male participants (orange) in partial residuals. Violin plots display

distributions of gender-specific partial residuals at each morph step (1: angry to 9: happy), with dashed lines

showing ordinary least squares fits to residualized data. Large markers indicate mean ratings per morph step.

Extreme outliers (±2SD from morph-level means) were excluded only for visualization clarity purposes (all data

retained in statistical models). Trendline equations reflect slope estimates from residualized relationships, with

shared R² value from the full interactive model.

Reliability

To evaluate reliability, we examined the relationship between the validity and the retest

ratings using a linear regression model. This model had a coefficient of determination (R^2) of 0.736 (p-

value < 0.001) (Figure 6) and the coefficient estimate for the validity answers was 0.84, with an

intercept of 0.814. The high R^2 value indicates strong test-retest reliability, suggesting minimal

variability in interpretations of the stimulus set across multiple exposures. The Bayes factor, comparing

the calculated regression to the null hypothesis ($R^2 = 0$), was infinite, showing strong evidence in favor

of H1. Moreover, the calculated statistical power ($1 - \beta$ error) in comparison with the non-correlation

model (Correlation ρ H0 = 0) was 0.999, demonstrating robust sensitivity to detect the observed

consistency in participants' ratings.

Figure 6

Reliability of participants' ratings between the initial test and the retest session.

Note. The x-axis denotes the ratings from the first task (validity ratings), and the y-axis corresponds to

ratings from the retest. The violin plots depict the distribution and probability density of ratings at each step, with

individual trials displayed by small grey dots. The means of the retest ratings are marked with black dots, and the

regression line is presented by a dashed line. The regression outcome and R^2 values are displayed in the figure.

The colors are used for visualization purposes only.

Discussion

In this paper, we introduced *STEMorph*, a newly developed stimulus set based on the NimStim

dataset, featuring facial expressions transitioning from anger to happiness . We employed neutral-

anchored morphing and neural-network-generated masking and evaluated the validity and reliability

of this dataset.

Participants' subjective ratings showed a robust positive correlation with the intended

emotional states, indicating that our morphing methodology successfully created a naturalistic

continuum of emotional transformations. This finding aligns with prior research emphasizing the

importance of realistic stimuli in emotion recognition studies (Paiva-Silva et al., 2016). Moreover,

STEMorph addresses limitations in earlier datasets, such as artifacts caused by basic morphing

techniques (Dawel et al., 2022), thereby enhancing ecological validity.

Furthermore, STEMorph's ability to capture gender differences in emotion recognition

provides a valuable tool for addressesing gaps in gender-specific emotion perception research. These

results align with previous studies suggesting that facial emotion recognition is influenced by the

gender of the stimuli (Harris et al., 2016) and the perceiver (Cavieres et al., 2021; Forni-Santos &

Osório, 2015; Wingenbach et al., 2018). Prior research have consistently reported gender-based

variations in emotion recognition (Elfenbein & Ambady, 2002) highlighting the importance of using

stimuli that account for such differences. Additionally, the high test-retest reliability observed in our

study suggests minimal variability in interpretations across multiple exposures. This finding

underscores the importance of well-designed stimuli in yielding consistent results in emotion

perception research.

Our methodological choices also reflect advancements over traditional approaches.

Specifically, utilizing a neutral expression as an intermediary anchor point minimized bias at the

central position and facilitated smoother transitions between angry and happy expressions. This

approach aligns with findings from earlier studies exploring emotional gradients (Silvia et al., 2006)

and builds on work demonstrating that neutral anchoring enhances naturalism in morphed stimuli

(Vikhanova et al., 2022). Additionally, by generating distinct neural-network-based masks for each

image, we mitigated issues like distortions in non-facial regions (e.g., hair or ears), which are common

in datasets relying on simpler masking techniques (Schurgin et al., 2014). Moreover, while removing

the background may reduce the real-world context of the stimuli, this step eliminates distractions,

focusing solely on facial features for emotion recognition. These innovations ensure that *STEMorph*

preserves anatomical fidelity while maintaining focus on facial features critical for emotion

recognition.

361   Importantly, by making the *STEMorph* stimulus set publicly available along with all associated

362   data and codes, we facilitate broader adoption and encourage further research in this critical domain

363   of human interaction. This transparency aligns with calls within the literature for accessible resources

364   to advance emotion recognition studies (Münkler et al., 2015). Furthermore, *STEMorph's* design

365   principles could inform future efforts to create ecologically valid stimuli for diverse applications,

366   including clinical diagnostics and machine learning algorithms for facial expression analysis.

367   Despite these strengths, this study has limitations that warrant consideration. While our

strong linear relationships provide a promising foundation for future research, further examination

across diverse samples is essential to assess the generalizability and external validity of our findings.

For instance, expanding participant demographics beyond young adults of similar ethnic backgrounds

could reveal how age or cultural differences influence emotion perception biases. Additionally,

restricting participant evaluations to a scale from anger to happiness may simplify responses.

Nontheless, we believe that focusing on the anger-happiness continuum provides a foundation for

future studies exploring other emotional transitions with more flexible response formats. Emotions

such as sadness or fear may exhibit unique perceptual biases requiring tailored stimulus sets.

Therefore, developing complementary transitions between various emotions could enrich our

understanding of emotion recognition processes across different contexts.

In conclusion, our results underscore the importance of methodological rigor in stimulus

design within the field of emotion recognition. The incorporation of neutral-anchored morphing and

neural-network-generated masks effectively addresses limitations observed in earlier studies regarding

the unnatural appearance of stimuli. By enhancing the naturalism and coherence of facial stimuli, our

work significantly contributes to the ongoing efforts to improve methodologies in studying facial

expressions. Furthermore, variations in emotional ratings across gender subgroups highlight the

relevance of considering both participant and facial expressor characteristics in emotion recognition

research. We believe that *STEMorph* can serve as a valuable resource for future studies exploring the

dynamics of emotion perception across diverse populations.

Declarations

Funding

389   No funding was received for conducting this study.

Conflicts of interest

391   The authors have no competing interests to declare that are relevant to the content of this

392 article.

Ethics approval

394   Approval was obtained from the ethics committee of Shahid Beheshti University of Medical

395 Sciences (IR.SBMU.MSP.REC.1396.757) on 22/01/2018. The procedures used in this study adhere to

396 the tenets of the Declaration of Helsinki.

Consent to participate

398   Informed consent was obtained from all individual participants included in the study.

Availability of data and materials

The data acquired and source codes for the psychophysics task and data analysis performed in

this study can be found in the following GitHub repository: https://github.com/Mekatebi/STEMorph

To use the stimulus set, visit the *NimStim Set of Facial Expressions* website and follow the

instructions to request access: https://danlab.psychology.columbia.edu/content/nimstim-set-facial-

expressions

The stimulus set is also available from the corresponding author on reasonable request.

Use of generative AI and AI-assisted technologies in the writing process

During the preparation of this work the authors used *Claude AI* in order to improve the

language and readability. After using this tool/service, the authors reviewed and edited the content as

needed and take full responsibility for the content of the published article.

410   **References**

Abrosoft Fantamorph Deluxe 5.4.8. https://www.fantamorph.com/index.html

Aguado, L., Pedraza, I. S., & Gutiérrez, A. G. (2015). A comparison of backward masking of faces in

expression and gender identification. *Psicologica*, *35*, 171-194.

Brainard,   D.   H.   (1997).   The   Psychophysics   Toolbox.   *Spat   Vis*,   *10*(4),   433-436.

https://www.ncbi.nlm.nih.gov/pubmed/9176952

Cavieres, A., Maldonado, R., Bland, A., & Elliott, R. (2021). Relationship Between Gender and

Performance on Emotion Perception Tasks in a Latino Population. *Int J Psychol Res (Medellin)*,

14(1), 106-114. https://doi.org/10.21500/20112084.5032

Dawel, A., Miller, E. J., Horsburgh, A., & Ford, P. (2022). A systematic survey of face stimuli used in

psychological   research   2000–2020.   *Behavior   research   methods*,   *54*(4),   1889-1901.

https://doi.org/10.3758/s13428-021-01705-3

Ekman, P. F., Wallace V. (1976). *Pictures of Facial Affect*. Consulting Psychologists Press.

Elfenbein, H. A., & Ambady, N. (2002). On the universality and cultural specificity of emotion recognition:

a meta-analysis. *Psychological bulletin*, *128*(2), 203.

face-parsing.PyTorch. https://github.com/zllrunning/face-parsing.PyTorch

Faceswap. https://faceswap.dev/

Faul, F., Erdfelder, E., Buchner, A., & Lang, A.-G. (2009). Statistical Power Analyses Using G*Power 3.1:

Tests for Correlation and Regression Analyses. *Behavior research methods*, *41*, 1149-1160.

https://doi.org/10.3758/BRM.41.4.1149

Fernando, F.-S. (2015). Facial Emotion Processing in the Laboratory (and elsewhere): Tradeoffs between

Stimulus Control and Ecological Validity. *AIMS Neuroscience*, *2*(4), 236-239.

https://doi.org/10.3934/Neuroscience.2015.4.236

Forni-Santos, L., & Osório, F. L. (2015). Influence of gender in the recognition of basic facial expressions:

A critical literature review. *World J Psychiatry*, *5*(3), 342-351.

https://doi.org/10.5498/wjp.v5.i3.342

Goeleven, E., De Raedt, R., Leyman, L., & Verschuere, B. (2008). The Karolinska Directed Emotional

Faces: A validation study. *Cognition and Emotion*, *22*(6), 1094-1118.

https://doi.org/10.1080/02699930701626582

Guarnera, M., Magnano, P., Pellerone, M., Cascio, M. I., Squatrito, V., & Buccheri, S. L. (2018). Facial

expressions and the ability to recognize emotions from the eyes or mouth: A comparison among

old adults, young adults, and children. *J Genet Psychol*, *179*(5), 297-310.

https://doi.org/10.1080/00221325.2018.1509200

Harmer, C. J., Goodwin, G. M., & Cowen, P. J. (2009). Why do antidepressants take so long to work? A

cognitive neuropsychological model of antidepressant drug action. *British Journal of Psychiatry*,

195(2), 102-108. https://doi.org/10.1192/bjp.bp.108.051193

Harris, D. A., Hayes-Skelton, S. A., & Ciaramitaro, V. M. (2016). What's in a Face? How Face Gender and

Current Affect Influence Perceived Emotion. *Front Psychol*, *7*, 1468.

https://doi.org/10.3389/fpsyg.2016.01468

Hunter, J. D. (2007). Matplotlib: A 2D graphics environment. *Computing in Science & Engineering*, *9*(3),

90-95. https://doi.org/10.1109/MCSE.2007.55

Joormann, J., & Gotlib, I. H. (2006). Is this happiness I see? Biases in the identification of emotional facial

expressions in depression and social phobia. *J Abnorm Psychol*, *115*(4), 705-714.

https://doi.org/10.1037/0021-843X.115.4.705

Leppänen, J., Milders, M., Bell, J., Terriere, E., & Hietanen, J. (2004). Depression biases the recognition

of emotionally neutral faces. *Psychiatry Research*, *128*, 123-133.

https://doi.org/10.1016/j.psychres.2004.05.020

Leppänen, J. M. (2006). Emotional information processing in mood disorders: a review of behavioral

and neuroimaging findings. *Curr Opin Psychiatry*, *19*(1), 34-39.

https://doi.org/10.1097/01.yco.0000191500.46411.00

Morey, R. D., & Rouder, J. N. (2023). BayesFactor: Computation of Bayes Factors for Common Designs.

https://CRAN.R-project.org/package=BayesFactor

Münkler, P., Rothkirch, M., Dalati, Y., Schmack, K., & Sterzer, P. (2015). Biased Recognition of Facial Affect

in Patients with Major Depressive Disorder Reflects Clinical State. *PLOS ONE*, *10*(6), e0129863.

https://doi.org/10.1371/journal.pone.0129863

Paiva-Silva, A. I. d., Pontes, M. K., Aguiar, J. S. R., & de Souza, W. C. (2016). How do we evaluate facial

emotion recognition? *Psychology & neuroscience*, *9*(2), 153.

Penton-Voak, I. S., Bate, H., Lewis, G., & Munafò, M. R. (2012). Effects of emotion perception training on

mood in undergraduate students: randomised controlled trial. *Br J Psychiatry*, *201*(1), 71-72.

https://doi.org/10.1192/bjp.bp.111.107086

Penton-Voak, I. S., Munafò, M. R., & Looi, C. Y. (2017). Biased facial-emotion perception in mental health

disorders: A possible target for psychological intervention? *Current Directions in Psychological*

Science, *26*(3), 294-301. https://doi.org/10.1177/0963721417704405

Russell, J. A. (1994). Is there universal recognition of emotion from facial expression? A review of the

cross-cultural studies. *Psychol Bull*, *115*(1), 102-141. https://doi.org/10.1037/0033-

2909.115.1.102

Schurgin, M. W., Nelson, J., Iida, S., Ohira, H., Chiao, J. Y., & Franconeri, S. L. (2014). Eye movements

during emotion recognition in faces. *Journal of Vision*, *14*(13), 14-14.

https://doi.org/10.1167/14.13.14

Seabold, S., & Perktold, J. (2010, 2010). *statsmodels: Econometric and statistical modeling with python*

9th Python in Science Conference,

Silvia, P. J., Allan, W. D., Beauchamp, D. L., Maschauer, E. L., & Workman, J. O. (2006). Biased Recognition

of Happy Facial Expressions in Social Anxiety. *Journal of Social and Clinical Psychology*, *25*(6),

585-602. https://doi.org/10.1521/jscp.2006.25.6.585

Surguladze, S. A., Young, A. W., Senior, C., Brebion, G., Travis, M. J., & Phillips, M. L. (2004). Recognition

accuracy and response bias to happy and sad facial expressions in patients with major

depression. *Neuropsychology*, *18*(2), 212-218. https://doi.org/10.1037/0894-4105.18.2.212

Sweeny, T. D., Grabowecky, M., Suzuki, S., & Paller, K. A. (2009). Long-lasting effects of subliminal

affective priming from facial expressions. *Consciousness and Cognition*, *18*(4), 929-938.

https://doi.org/https://doi.org/10.1016/j.concog.2009.07.011

Testerman, M. (2023). *Face Image Databases: An A-Z directory of databases of face stimulus images for*

use in behavioral research. Retrieved 2024/02/07 from

https://libguides.princeton.edu/facedatabases

Tottenham, N., Tanaka, J. W., Leon, A. C., McCarry, T., Nurse, M., Hare, T. A., Marcus, D. J., Westerlund,

A., Casey, B. J., & Nelson, C. (2009). The NimStim set of facial expressions: Judgments from

untrained research participants. *Psychiatry Research*, *168*(3), 242-249.

https://doi.org/https://doi.org/10.1016/j.psychres.2008.05.006

Vikhanova, A., Mareschal, I., & Tibber, M. (2022). Emotion recognition bias depends on stimulus

morphing strategy. *Attention, Perception, & Psychophysics*, *84*(6), 2051-2059.

https://doi.org/10.3758/s13414-022-02532-0

Waskom, M. L. (2021). seaborn: statistical data visualization. *Journal of Open Source Software*, *6*(60),

3021. https://doi.org/10.21105/joss.03021

Waters, A. M., Mogg, K., Bradley, B. P., & Pine, D. S. (2008). Attentional bias for emotional faces in

children with generalized anxiety disorder. *J Am Acad Child Adolesc Psychiatry*, *47*(4), 435-442.

https://doi.org/10.1097/CHI.0b013e3181642992

Wingenbach, T. S. H., Ashwin, C., & Brosnan, M. (2018). Sex differences in facial emotion recognition

across varying expression intensity levels from videos. *PLOS ONE*, *13*(1), e0190634.

https://doi.org/10.1371/journal.pone.0190634

Goeleven, E., R. De Raedt, L. Leyman and B. Verschuere (2008). "The Karolinska Directed

Emotional Faces: A validation study." Cognition and Emotion **22**(6): 1094-1118.

Ekman, P., & Friesen, W. V. (1976). *Pictures of Facial Affect*. Retrieved 2025/02/05

from https://books.google.com/books/about/Pictures_of_Facial_Affect.html?id=gbfMSgAACAAJ

Guarnera M, Magnano P, Pellerone M, Cascio MI, Squatrito V, Buccheri SL. Facial expressions

and the ability to recognize emotions from the eyes or mouth: A comparison among old adults, young

adults, and children. J Genet Psychol. 2018 Sep-Oct;179(5):297-310. doi:

10.1080/00221325.2018.1509200. Epub 2018 Oct 22. PMID: 30346916.

Fernando, F.-S. (2015). "Facial Emotion Processing in the Laboratory (and elsewhere):

Tradeoffs between Stimulus Control and Ecological Validity." AIMS Neuroscience 2(4): 236-239.

Harmer, C. J., G. M. Goodwin and P. J. Cowen (2009). "Why do antidepressants take so long to

work? A cognitive neuropsychological model of antidepressant drug action." British Journal of

Psychiatry 195(2): 102-108.

Leppänen, J. M. (2006). "Emotional information processing in mood disorders: a review of

behavioral and neuroimaging findings." Curr Opin Psychiatry 19(1): 34-39.

Penton-Voak, I. S., M. R. Munafò and C. Y. Looi (2017). "Biased facial-emotion perception in

mental health disorders: A possible target for psychological intervention?" Current Directions in

Psychological Science 26(3): 294-301.

Penton-Voak, I. S., H. Bate, G. Lewis and M. R. Munafò (2012). "Effects of emotion perception

training on mood in undergraduate students: randomised controlled trial." <u>Br J Psychiatry</u> 201(1): 71-

72.

Waters, A. M., K. Mogg, B. P. Bradley and D. S. Pine (2008). "Attentional bias for emotional

faces in children with generalized anxiety disorder." <u>J Am Acad Child Adolesc Psychiatry</u> 47(4): 435-

442.

532