Multi-omics phenotyping characterizes molecular divergence underlying different clinical scenarios of inflammatory bowel disease

Nguyen Tran Nam Tien[1,#], Eun Jeong Choi[2,#], Nguyen Quang Thu[1], Seung Jung Yu[2], Duc Ninh Nguyen[3], Dong Hyun Kim[1], Nguyen Phuoc Long[1,*], Hong Sub Lee[2,*]

[1]Department of Pharmacology and PharmacoGenomics Research Center, Inje University College of Medicine, Busan 47392, Republic of Korea

[2]Department of Internal Medicine, Busan Paik Hospital, Inje University College of Medicine, Busan, 47392, Republic of Korea

[3]Section for Comparative Pediatrics and Nutrition, Department of Veterinary and Animal Sciences, University of Copenhagen, Frederiksberg 1870, Denmark

[#]These authors contributed equally to this work

*Corresponding authors: N.P.L. (email: phuoclong@inje.ac.kr) and H.S.L. (email: hslee@paik.ac.kr)

ABSTRACT

Clinically heterogeneous spectrum and molecular phenotypes of inflammatory bowel disease (IBD) remain to be comprehensively elucidated. This study set out to explore the serum molecular profiles (I) of IBD subtypes; in association with (II) elevated fecal calprotectin and (III) disease activity states; (IV) upon treatment escalation; and (V) in patients who needed treatment escalation. The serum proteome, metabolome, and lipidome of 75 treated IBD patients were profiled. Single- and multi-omic data analysis was performed to determine differential analytes and integrative biosignatures. (I) Chronic inflammation, and phosphatidylcholine and bile acid homeostasis disturbances underlined the differences between Crohn's disease (CD) and ulcerative colitis. (II) Elevated calprotectin was associated with higher levels of inflammatory proteins and sphingomyelins (SM) and lower levels of bile acids, amino acids, and triacylglycerols (TG). Relative to patient remission, active disease state (III) was characterized by decreased SMs and increased inflammatory proteins and TGs. (IV) Treatment escalation was associated with augmented levels of inflammatory response-related proteins and reduced levels of amino acids. Most TG species increased in the post-treatment escalation. Moreover, needed-treatment-escalation patients had significantly lower levels of TGs (V). They also showed increased SMs and decreased signaling receptor binding proteins. Multi-omics analysis revealed biosignatures that captured the differences between groups of each scenario. Eight analytes, including NFASC, ANGPTL4, and chenodeoxycholate, were found in at least three biosignatures. Collectively, disturbances in immune response, bile acid homeostasis, amino acids, and lipids alteration potentially underlie the clinically heterogeneous spectrum of IBD.

Keywords: IBD; deep phenotyping; molecular phenotype; multi-omics; DIABLO

INTRODUCTION

Inflammatory bowel disease (IBD) poses a significant health burden worldwide, impacting millions of individuals [1,2]. The disease emerges from a multifaceted interplay involving genetic predisposition, the gut microbiota, diverse environmental exposures, and immunological disturbances [3,4]. Upon diagnosis, IBD is typically classified as Crohn's disease (CD) and ulcerative colitis (UC) [5]. Treatment plans are tailored based on disease activity, aiming to mitigate inflammation, alleviate symptoms, and enhance the patient's quality of life [6]. Multiple evaluation methods are needed to achieve this goal.

Biomarkers including fecal calprotectin serve as a surrogate endpoint of gastrointestinal tract inflammation and clinicians implement symptomology to manage the patients [6]. Failure of standard therapy requires treatment escalation where biologics target key immunoregulatory-related processes to reduce inflammation, improve symptoms, and promote healing for IBD patients [7,8]. However, disease heterogeneity, inadequate drug concentration, and immunogenicity development can compromise biologics's efficacy [4]. Therefore, a deeper understanding of molecular profiles associated with the disease pathogenesis, inflammation level, disease activity state, and molecular profiles upon treatment escalation is essential [5].

Deep molecular phenotyping represents a paradigm shift in disease characterization and refers to a comprehensive and highly detailed approach beyond solely relying on clinical, biochemical, and imaging findings [9,10]. Multi-omics stands out as a promising approach for comprehensively characterizing complex diseases like IBD. Indeed, the approach has gradually become a cornerstone of IBD research [11-19]. Several molecular perturbations driving disease pathophysiology, including barrier dysfunction, chronic immunological response, microbial dysbiosis, and consequences alteration in tricarboxylic acid cycle metabolites, amino acid, lipid

metabolism, and oxidative pathways, have been elucidated thanks to the omics approach.

However, gaps persist in current IBD omics-based research, e.g., limited independent validation

and a small number of multi-omic studies [18]. Additionally, most (multi)-omics studies focused on

molecular profiling for IBD subtypes, with a limited number of studies for other clinical scenarios,

including disease activity, inflammation level, and treatment response.

This study aimed to explore the differences in serum molecular profiles (proteome,

metabolome, and lipidome) underpinning different clinical scenarios of the under-treatment IBD

population. Specifically, we characterized the single- and multi-ome differences among patients

diagnosed with either CD or UC, those with elevated or normal fecal calprotectin levels, and

individuals who presented an active or remission disease state. Moreover, we investigated the

molecular perturbations in patients who underwent treatment escalation with samples collected

before and after the treatment initiation, as well as those who needed treatment escalation and those

who did not. The study provided comprehensive insights into the functional multi-ome profiles of

IBD patients across five distinct clinical scenarios and laid a foundation for subsequent studies on

improving disease characterization and management.

METHODS

Ethical statement and study population

The study protocol was reviewed and approved by the Institutional Review Board of Busan Paik Hospital (No. BPIRB 2024-02-046). The study was conducted following the Helsinki Declaration. Before enrollment, all subjects provided written informed consent. Previously diagnosed IBD patients who underwent extensive treatment were recruited in the current study. Participants were excluded if they had severe comorbidities or extraintestinal manifestations. Endoscopy was performed to classify patients into UC and CD subtypes at enrolment (**Table 1**). Fecal calprotectin was used to categorize participants into elevated or normal groups using a 150 µg/g cut-off [20,21]. Median imputation was used in 2 cases with missing calprotectin levels. The C-reactive proteins were significantly higher in the CD than in the UC group and in the elevated fecal calprotectin group compared to the normal counterpart (**Table 1**, **Supplementary Table S1**). The MAYO score cut-off of 1 and CD activity index (CDAI) score cut-off of 150 concerning CD and UC were used to reflect the disease status as either active state (i.e., mild, moderate, or severe conditions) or remission. Remission-state patients tended to have a more prolonged illness duration (**Supplementary Table S2**). Patients with samples collected before and after the treatment escalation were assigned to the pseudo-pre- and post-treatment escalation design as an unpaired comparison (**Supplementary Table S3**). The former group (i.e., pre-treatment escalation) also served as patients who needed treatment escalation, compared to those who did not need it (**Supplementary Table S4**). These constituted five clinically meaningful scenarios, which are schematized in **Fig. 1**.

Metabolomics and lipidomics sample preparation and data acquisition

The metabolomics and lipidomics experiments followed our previously established procedure [22-

26]. Detailed plasma sample preparation, sample treatment, and instrumental analysis of

metabolomics and lipidomics were described in the **supplementary methods**.

Proteomics sample preparation and data acquisition

For proteomics analysis, Olink® panels of Explore 384 Inflammation and Target 96

Cardiovascular III (Olink, Uppsala, Sweden) were used. Sample treatment for protein extraction

followed Olink's established procedure. This method utilizes a highly specific antibody probe

labeled with two oligonucleotides to bind the target protein. The real-time PCR amplification of

the oligonucleotide sequence then allows for quantitative measurement of the resulting DNA

sequence. The experiment was conducted by DNALink (Seoul, Republic of Korea).

Data preprocessing, data treatment, and normalization

MS-DIAL (version 4.9.0) was used for MS data processing [27]. In-house libraries of authentic

standards, MS-DIAL in-built libraries, and public libraries were used to annotate metabolites and

lipids [23,24]. Only metabolites and lipids with reliable annotations were subjected to subsequent

analyses. Next, the aligned data was imported to the *MetaboAnalystR* package version 4.0.0 for

further data treatment [28]. The normalized protein expression values were used for proteomics data,

and the values recorded below the limit of detection were considered missing. Data points from

the Olink® Target 96 Cardiovascular III were used for downstream analyses when proteins were

quantified in 2 panels.

Analytes (metabolites, lipids, and proteins) with a missing rate $\geq 50\%$ were excluded, whereas

the retained missing values were inputted via the k-nearest neighbors algorithm. For metabolomics

and lipidomics, analytes whose relative standard deviation was > 25% in the pooled QCs samples

were also filtered out.

Single-omics differential analysis

For differential analysis, the *limma* method was used. The analysis was performed in the

MetaboAnalystR package version 4.0.0 for each omics data separately [28]. Pre-treated and

normalized data without scaling was used to obtain differential analytes (DAs) of metabolites,

lipids, and proteins between groups. A P-value of 0.05 was considered the putative significant cut-

off. The heatmap of DAs of each omics was visualized via the *ComplexHeatmap* version 2.14.0 in

R 4.2.3 [29].

Integrative multi-omics analysis and biosignature identification

Regarding integrative multi-omics analysis, we employed the data integration analysis for

biomarker discovery using the latent components (DIABLO) method, which was performed using

the *mixOmics* package version 6.23.4 [30,31]. Normalized data of each omics data was used as input

data. The scaling was conducted internally by DIABLO. For model tuning, we first explored the

weight and chose the value of 0.5 to satisfy a trade-off between correlation and discriminability.

Next, the number of components and the distance method were selected via a 10-fold cross-

validation (CV). Afterward, the number of features within each data layer was determined. The

performance of selected biosignatures to discriminate groups of interest was obtained separately.

The area under the receiver operating characteristic curve (AUC) was calculated by averaging

sequentially over 10-fold cross-validation, 10-repeated, and three data blocks [30,31].

RESULTS

To investigate the molecular perturbation underlying different clinical scenarios of IBD, we stratified the patient cohort using their diagnostic results, fecal calprotectin level, disease activity score, and treatment escalation information (**Fig. 1**).

Treated UC and CD patients presented divergence in molecular perturbations linked to chronic inflammation, phosphatidylcholines, and bile acid homeostasis

UC and CD are traditionally the two main subtypes of IBD, which are macro-phenotyped based on endoscopy results and clinical manifestations (**Fig. 2a**). Herein, we determined molecules had differential abundance between UC *vs.* CD. Employing an age- and sex-adjusted linear model, we found 44, 11, and 29 differential proteins, metabolites, and lipids, respectively. These differential analytes (DAs) showed relatively straightforward patterns of differences across three data modalities, especially lipids, and proteins (**Fig. 2b-d**). Most differential proteins (e.g., IFNG, IL1R1, IL12RB1, CCL20, AGER) were linked to inflammation and immune signaling pathways. We also observed a consistent up-regulation of cell adhesion molecules, including ITGB2, ITGB6, ITGA11, LAMA4, and ADA, among others (**Fig. 2b**). Regarding metabolomics modality, changes in bile acid biosynthesis were observed where levels of three conjugated bile acids (glycocholate, taurocholate, and taurodeoxycholate) increased 2-3.5 times in the UC compared to the CD group. Changes of two amino acids (dihydroorotate and 5-hydroxylysine) were also observed (**Fig. 2c**). Lipidomics analyses showed that phosphatidylcholines (PC) and its derivatives of lysoPC (LPC) and ether-linked PC [PC(O-)] and LPC(O-) were the major altered species. Abundances of LPC, lysophosphatidylethanolamines (LPE), ceramides (Cer), and triacylglycerols (TG) were decreased, whereas levels ether-linked phosphatidylethanolamine [PE(O-)] were augmented in the UC group. PCs showed up- and down-regulation patterns (**Fig. 2d**).

Utilizing an integrative omics approach, we evaluated the correlation of the data layers and

determined the highly correlated and discriminant features across the three data blocks. The

proteomics modality correlated well with two others (**Fig. 2e**, **Supplementary Fig. S1a**). Among

16 DIABLO-identified proteins, most of them showed significantly higher abundance in UC,

except an 8-fold lower level of CLIP2. Proteins (especially differential proteins of ITGA11,

ITGB2, and IL32) showed strong negative correlation with lipids, particularly LPC(O-18:1) and

LPC(16:0-SN1). These lipids correlated negatively with 2,3-dihydroxybenzoate. Other highly

correlated and discriminant metabolites (e.g., taurodeoxycholate, taurochenodeoxycholate) and

lipids (e.g., Cer(18:1;2O/24:1)) were also statistically different.

Elevated fecal calprotectin associated with increased inflammation-related proteins, SMs;

decreased bile acids, amino acids, and TGs

Fecal calprotectin has emerged as a valuable tool in managing IBD, including screening,

diagnosis, disease activity and severity assessment, treatment monitoring, and relapse prediction

of IBD patients [6,32]. We explored the molecular perturbations of two IBD subpopulations, i.e.,

elevated and normal fecal calprotectin levels (**Fig. 3a**). We found 34 differentially expressed

proteins between the two groups. Most of them (32 proteins) were up-regulated in patients with

elevated fecal calprotectin levels (**Fig. 3b**). These differential proteins belonged to some of the

innate and adaptive immune responses, such as TNF, MAPK, IL-17, and NF-κB signaling

pathways. Change in proteins involved in the sphingolipid signaling (TNFRSF1A, TRAF2) and

PI3K-Akt signaling (CSF1, HGF, OSM, TGFA) was also observed. Besides, 15 differential

metabolites were found (**Fig. 3c**). Abundances of chenodeoxycholate and conjugated bile acids

(i.e., glycocholate, taurochenodeoxycholate, and glycochenodeoxycholate) decreased (about 2.5-

fold), except glycohyodeoxycholate increased slightly in the elevated calprotectin group. Higher

levels of two amino acids (creatinine, glutamate) and two DNA metabolism-related metabolites

(cytidine, deoxyguanosine-monophosphate) were also observed. Lipids, especially lysoPC, were

also found in the metabolomics modality. The lipid modality contributed the most significant

number of DAs, with 47 differential lipids (**Fig. 3d**). They presented a clear contrast mostly linked

to the low abundance of TG and high abundance of SM. Levels of other lipid subclasses [i.e., PC,

PC(O-), LPC, LPE, Cer, and cholesteryl esters (CE)] were also increased.

Integrative omics analysis revealed a good correlation between different data blocks, in which

negative correlations were generally observable between proteomics and two other data layers

(**Fig. 3e**, **Supplementary Fig. S1b**). However, a strong correlation was the positive correlation of

glutamate and creatine with Cer species. Proteins served as the major modality to distinguish 2

groups. Thirty multi-omics-identified proteins were related to immune activation. Only 5 of them

were differential analytes. Multi-ome biosignatures also included five metabolites, such as 2 amino

acids (creatinine and glutamate; also were differential analytes), potential bacterial-source

metabolites (pyrocatechol), and chenodeoxycholate. Eight selected lipids belonged to the Cer, PC,

and LPC species, seven of which were differential lipids.

Active IBD disease state was characterized by an upregulation of inflammation-related

proteins, a decrease in SM, and an increase in TG levels

The molecular response underlying patients in active or remission states was investigated (**Fig.**

4a). While lipids showed a significant contribution with 57 differential analytes, 18 and 7

differential proteins and metabolites were observed. Overall, patients with active disease were

characterized by the upregulation of all inflammatory-related proteins, a decrease in SM, and an

increase in TG levels compared to the remission group (**Fig. 4b-d**). The significantly altered

proteins, such as CXCL9, CXCL1, IL17A, TNFSF11, ITGB2, and ADGRE2, were prominently

related to activated immune response (**Fig. 4b**). Regarding differential metabolites, muricholate

and taurocholate were elevated (2-fold), whereas coprocholate was slightly declined (**Fig. 4c**).

Levels of lactate, proline, and acetylcarnitine were decreased. We observed that patients who did

not need treatment escalation contributed mainly to the augmented levels of TGs in the active

group, whereas abundances of TGs were reduced in the remission group regardless of biologics

status (**Fig. 4d**). Lipidomics analysis also revealed a lower abundance of PCs, LPCs, and CEs in

the active group.

In multi-omics analysis, a pair-wise correlation from DIABLO showed a strong correlation

between omics layers (**Supplementary Fig. S1c**, **Fig. 4e**). Metabolites and lipids were major

drivers for differentiating active and remission disease states. The biosignature comprised 8

proteins, 25 metabolites, and 30 lipids. None of the identified proteins were differentially

expressed (**Fig. 4e**). Acylcarnitines were detected primarily within the metabolomics data, yet their

abundance between the 2 groups was subtle, except for acetylcarnitine. Two bile acids

(muricholate and taurocholate) were also at the top, indicating differences between groups. Within

the lipidomics layer, 4 TGs and 9 SMs showed the highest differences between the active and

remission groups. Besides, fatty acyls were identified in the multi-omics analysis. They showed a

strong positive correlation with ANGPTL4 and acylcarnitines.

Treatment escalation promoted upregulation of inflammatory response-related proteins,

TGs, and decreased amino acids, purine metabolites

We explored molecular perturbations between patients who received treatment escalation with

samples collected before and after the therapeutic initiation (**Fig. 5a**). Treatment escalation

triggered significant alterations in proteome, metabolome, and lipidome response in IBD patients

(**Fig. 5b-d**). We identified 16, 24, and 20 differential proteins, metabolites, and lipids. The majority

of altered proteins, which were up-regulated, were linked to the perturbations of the cellular

receptor signaling pathway (e.g., COL1A1, FLT3LG, ITGB6, LAMA4), PI3K-Akt signaling

pathway, and focal adhesion (**Fig. 5b**). A consistent decrease in levels of differential metabolites

was observed (**Fig. 5c**). They were linked to lipid metabolism [e.g., LPC(O-16:0), LPC(O-16:1),

LPE(O-18:1), and CAR(12:0;O)], arginine biosynthesis (arginine, aspartate, ornithine, and

glutamine), pentose phosphate pathway (hexose-monophosphate, gluconate, and glycerate), purine

metabolism (hypoxanthine, glutamine, and deoxyguanosine monophosphate), and amino acid

alterations (e.g., creatinine, DOPA, and oxoproline). Lipidome alterations were mainly

characterized by a higher level of TGs with a 2-3-fold increase (**Fig. 5d**).

A high correlation of metabolites and proteins, along with good separation between the two

groups were observed, indicated by the distinction in the centroids of each group, yet with an

observable moderate amount of overlap between each sample (**Supplementary Fig. S1d**, **Fig. 5e**).

Most metabolites (4/5, e.g., hypoxanthine) and lipids (5/6, mainly TGs) in the multi-ome

biosignatures were also differential analytes, whereas 11/30 proteins had differential abundance.

Lower abundance of hypoxanthine and glycate were correlated strongly with decreased

abundances of TGs and increased levels of proteins such as CCL13, ITGB6, and COLEC12.

Patients who needed to escalate their treatment showed downregulation of signaling receptor

binding proteins, lower levels of TGs, higher levels of SMs and PCs

The multi-ome profiles of patients who needed treatment escalation and those who did not

were compared (**Fig. 6a**). The adjusted linear model revealed 25 proteins, mostly linked to

signaling receptor binding, differentially expressed (11 up- and 14 down-regulated) between two

groups (**Fig. 6b**). Proteins linked to cytokine-cytokine receptor interaction were down-regulated

(2 to 10-fold; CXCL6, CD40LG, CCL13, TNFRSF14, and TNFSF13B) in patients who needed

treatment escalation, except CXCL9 increased 6-fold. Four proteins involved in MAPK and PI3K-Akt signaling pathways (i.e., FLT3LG, EGF, EGFR, and PDGFA) were also consistently down-regulated. Changes in the proteins mentioned above might also indicate the alteration of NF-κB and JAK-STAT signaling pathways. For metabolomics data, we determined 10 differential metabolites (**Fig. 6c**). A marginal increase in the abundance of three amino acids (DOPA, arginine, and ornithine) was observed, which might imply a change in arginine metabolism. Regarding lipidome alteration, 31 lipids were statistically significant (**Fig. 6d**). TG species showed a lower level (2-3-fold) in patients who required treatment escalation, whereas other lipid subclasses, such as PC(O-), PC, and Cer had increased abundance.

A good correlation between proteome, metabolome, and lipidome profiles was observed within two groups (**Supplementary Fig. S1e**, **Fig. 6e**). The negative correlation pattern was mainly observed, such as between NFASC and TREM2 with SM(34:1;2O). Proteins identified by multi-omics analysis, which mainly down-regulated, were linked to the immune response. While metabolites range from acylcarnitines to bile acids (chenodeoxycholate) and bacterial-sourced metabolites (pyrocatechol), selected lipids were mainly SM and PC(O-) species, whose higher abundance in patients needed treatment escalation.

Integrative biosignatures captured differences between groups of five scenarios

The predictive performance of selected multi-ome biosignatures is shown in **Table 2**. Differences between CD and UC groups, as well as between individuals with elevated and normal fecal calprotectin levels, could be captured by the selected multi-ome biosignatures (AUC = 0.80, P-value = 0.008; AUC = 0.73, P-value = 0.03; respectively). However, the model failed to differentiate active from remission disease state groups (AUC = 0.56, P-value = 0.42). Regarding treatment escalation, the biosignatures exhibited an excellent performance in distinguishing post-

from pre-treatment escalation groups (AUC = 0.90, P-value = 0.01). Additionally, a good

capability was observed in distinguishing patients needing to escalate their treatment from those

not (AUC = 0.77, P-value = 0.02). Among these five integrative biosignatures, 8 analytes were

detected in at least 3 of 5 scenarios (**Supplementary Fig. S2**). They included NFASC, CD83,

chenodeoxycholate, ANGPTL4, pyrocatechol, Cer(18:1;2O/24:1), ISM1, and EPHA1. Their

abundances across 5 scenarios are shown in **Supplementary Figs. S3-10**.

DISCUSSION

Owing to the heterogeneous nature of IBD, the current disease classification could preclude the endeavors to understand the pathophysiology of the disease. A classification comprising multiple biomarkers that reflect the IBD spectrum holds promise to account for diverse manifestations of IBD [5]. Hence, using multi-ome molecular profiles may be a better approach. Our study set out to investigate five commonly seen clinical scenarios and provide the evidence about the molecular differences of the comparable groups at the functional multi-ome data. It pinpointed the serum multi-ome profiles underlying the disease and associated with either the fecal calprotectin levels or disease activity states, as well as molecular profiles upon treatment escalation.

We first aimed to elucidate the molecular differences between UC and CD. Barrier dysfunction is known to be the primary mechanism underlying IBD pathophysiology [4]. In UC, the injury only extends to the mucosal layer, whereas transmural inflammation occurs in CD [12]. In our study, cell adhesion molecules were consistently higher in UC compared to CD, suggesting the importance and potential utility of these proteins. Among them, the increased expression of ITGA11, ITGB2, and IL32 demonstrated strong negative correlation with LPC(O-18:1) and LPC(16:0-SN1). The alterations of PC species and their derivatives suggests the disruption of the gut mucus layer. Previous studies found that PCs and LPCs species decreased in UC versus CD, whereas we observed a consistent decrease of LPC species and PCs showed species-dependent patterns [16]. The shift of phospholipid composition in intestinal mucus could lead to an increase in the permeability of the intestinal mucus, promoting mucosal inflammation [33]. Bile acids are mainly re-absorbed via the ileum, a well-known location for CD compared to UC. In our study, changes in bile acid

homeostasis indicated by altered levels of bile acids and FGF19 expression between two disease

subtypes also might induce chronic inflammation, resulting in an impaired mucosal barrier [34,35].

Fecal calprotectin could be used as a surrogate endpoint presenting gastrointestinal tract

inflammation in IBD. Mechanistically, non-apoptotic epithelial cell death, oxidative stress, and

homeostasis pathway dysregulation result in an excessive release of calprotectin, promoting

inflammation [36]. Establishing the association between serum multi-ome profiles and fecal

calprotectin to shed light on the systemic molecular alterations upon inflammation. In our study,

patients with elevated fecal calprotectin presented up-regulation of proteins involved in

inflammation-promoting signaling pathways. Inflammation and microbiome dysbiosis might lead

to malabsorption and disrupt bile acid function in the gastrointestinal tract. This might result in

systemic changes in amino acids and DNA-related metabolites, as observed in our study [14,15].

Besides, determining the relationship between serum multi-omic profiles and disease activity

could illuminate systemic molecular alterations associated with active disease manifestations. We

found that active disease was characterized by up-regulation of Th1 and Th17 immune activation-

related proteins such as CXCL9, CXCL1, ITGB2, and IL17A. Between these two scenarios, we

observed differences, yet significance in the direction of change of TGs and SMs. In the gut, the

immunomodulatory role of SMs, either of host- or microbiome-origin, has been recognized as

contributing to cell growth, differentiation, apoptosis, and membrane structure integrity [16,37,38].

Additionally, intestinal spontaneous inflammations could result from reduced bacterial

sphingolipids. IBD patients suffer from lipid abnormalities that correlate with disease activity [39].

We observed consistently increased levels of TGs in the active disease state, which may be helpful

biomarkers and associated with the pathophysiology of the active disease state.

We also designed the pre- and post-treatment scenarios to investigate the molecular

perturbations upon treatment escalation. Treatment escalation appeared to be associated with an

increase of inflammatory response-related proteins and TGs; and a decrease of amino acids and

purine metabolites. The high abundance of TGs after treatment escalation could be linked to the

reduced abundance of TGs in the elevated fecal calprotectin group as a response to inflammation

alleviation. Moreover, needed-treatment-escalation patients had a lower level of TGs compared to

patients who did not require treatment escalation. It should also be noted that increased TGs in the

active-disease-state patients were attributed predominantly to individuals who did not need

treatment escalation. Therefore, TGs might have the potential for treatment monitoring and

complication detection. However, a more sophisticated analysis in the context of improved study

design and sample size is warranted to establish the association [40]. Patients required to escalate

their treatment tend to suffer more severity of disease and thus need to escalate their treatment.

Nevertheless, disease severity assessment is a hard-to-implement process requiring multiple

criteria [32]. Therefore, fast and convenient biosignatures might open the door for early treatment

intervention. We observed the decreased levels of signaling receptor binding proteins and higher

levels of SMs and PCs. However, CXCL9 increased 6-fold in needed-treatment-escalation

patients. The abundance of this protein was also significantly increased in patients with active

disease states and elevated calprotectin. Notably, CXCL9 significantly increased the risk of disease

development in CD patients [41]. Therefore, pre-clinical to clinical activation of CXCL9 might be of

potential for clinical utilization.

Integrative biosignature holds the potential to account for diverse manifestations and capture

the holistic change of IBD. We evaluated predictive performance to give an idea about the potential

of the proposed biosignature. This *priori* information might be of potential use for further studies.

Some common analytes detected might be potential candidates for IBD management. For example,

neurofascin (NFASC), which was only detected by integrative analysis, was a biomarker candidate

among 3 of 5 investigated scenarios with consistently lower levels in elevated-calprotectin, active-

disease-state, and needed-treatment-escalation patients. Also, the NFASC levels were lower in CD

patients and tended to be normalized in the post-treatment escalation group. Neurofascin is known

for its mechanistic and clinical role in peripheral neuropathy, which is one of the most common

yet unrecognized neurological disorders in the IBD population, especially in CD patients [42-46]. A

previous study showed that aged monkey brain tissues presented increased interaction between

neurofascin and mutant α-synuclein, and transfected neurofascin could ameliorate its neuritic

toxicity in neuronal cells. Alpha-synuclein is known to cause Parkinson's disease, which is co-

occurrence and increased risk in IBD patients [47]. Additionally, early anti-inflammatory treatment,

especially anti-TNF-α agents, has been associated with decreased Parkinson's disease risk in the

IBD population [48,49]. Another protein, ANGPTL4, known for its regulatory role in inflammatory

responses across various diseases [50], was augmented in patients with elevated calprotectin and in

patients who needed treatment escaltion, yet decreased in active cases. Studies showed elevated

plasma levels of ANGPTL4 in mild/moderate IBD patients, returning to normal levels in severe

cases [51]. Increased ANGPTL4 expression in the inflamed colons correlated with the reduced

infiltration of CD8+ cells, suggesting its potential role in modulating gut immune responses [51].

Some limitations in our study should be noted. First, the genetic background of the patients

was unavailable. This hampered the exploration of the genetics and molecular profiles. Second,

the limited number of participants might restrict the generalizability of some findings and make it

hard to detect subtle but potentially important differences in the multi-ome profiles. Besides, the

inclusion of treated patients due to the convenient sampling setting prevented us from exploring

the effects of the underlying disease itself and the effects of the treatment. Although the molecular

profiles reported in our study were comprehensive, the cross-sectional study design made it

necessary to improve the study design to draw the causative relationship.

CONCLUSION

This study conducted a functional multi-omics approach toward deep molecular phenotyping

of IBD. The comprehensive analysis revealed distinct molecular biosignatures across five typical

IBD scenarios. Our findings provide deeper insights into serum multi-ome profiles underlying the

disease and in association with either the fecal calprotectin levels or disease activity states, and

molecular profiles upon treatment escalation. These biosignatures pointed towards disturbances in

key pathophysiological processes like immune response, bile acid homeostasis, amino acids, and

lipids alteration, potentially underlying the heterogeneous spectrum of IBD. Some potential

candidates, such as NFASC, ANGPTL4, chenodeoxycholate, and TGs hold promise as potential

candidates for IBD management. Future validation in more extensive and longitudinal cohorts and

translation into clinical tools can pave the way for a more precise approach to IBD diagnosis,

stratification, and treatment.

ACKNOWLEDGEMENTS

This work was supported by Eisai Korea Inc. The funder was not involved in the study design, data acquisition, data analysis, data interpretation, manuscript writing and editing, or publication.

The biospecimens and data used in this study were provided by the Biobank of Inje University Busan Paik Hospital, member of Korea Biobank Network (KBN4_A08).

AUTHOR CONTRIBUTIONS STATEMENT

N.T.N.T.: Data curation, Methodology, Investigation, Formal analysis, Validation, Writing – Original Draft, Writing – Review & Editing. **E.J.C.**: Data curation, Methodology, Investigation, Formal analysis, Visualization, Writing – Review & Editing. **N.Q.T.**: Data curation, Methodology, Software, Validation, Writing – Review & Editing. **S.J.Y.**: Data curation, Investigation, Validation, Writing – Review & Editing. **D.N.N.**: Methodology, Supervision, Validation, Writing – Review & Editing. **D.H.K.**: Conceptualization, Investigation, Supervision, Resources, Writing – Review & Editing, Funding acquisition. **N.P.L.**: Conceptualization, Methodology, Investigation, Formal analysis, Validation, Supervision, Resources, Writing – Original draft, Writing – Review & Editing, Funding acquisition. **H.S.L.**: Conceptualization, Methodology, Investigation, Formal analysis, Validation, Resources, Supervision, Writing – Original draft, Writing – Review & Editing, Funding acquisition.

DATA AVAILABILITY STATEMENT

The data supporting this study's findings are available upon reasonable request.

ADDITIONAL INFORMATION

Competing Interests Statement

407 The authors declare that they have no known competing financial interests or personal

408 relationships that could have appeared to influence the work reported in this paper.

409 **SUPPLEMENTARY INFORMATION**

410 Supplementary materials for this article can be found online.

REFERENCES

Alatab, S. *et al.* The global, regional, and national burden of inflammatory bowel disease in 195 countries and territories, 1990-2017: a systematic analysis for the Global Burden of Disease Study 2017. *Lancet Gastroenterol. Hepatol.* **5**, 17–30 (2020).

Kaplan, G. G. & Windsor, J. W. The four epidemiological stages in the global evolution of inflammatory bowel disease. *Nat. Rev. Gastroenterol. Hepatol.* **18**, 56–66 (2021).

Ananthakrishnan, A. N. Epidemiology and risk factors for IBD. *Nat. Rev. Gastroenterol. Hepatol.* **12**, 205–217 (2015).

Chang, J. T. Pathophysiology of Inflammatory Bowel Diseases. *New Engl. J. Med.* **383**, 2652–2664 (2020).

Verstockt, B., Bressler, B., Martinez-Lozano, H., McGovern, D. & Silverberg, M. S. Time to Revisit Disease Classification in Inflammatory Bowel Disease: Is the Current Classification of Inflammatory Bowel Disease Good Enough for Optimal Clinical Management? *Gastroenterology* **162**, 1370–1382 (2022).

Walsh, A. J., Bryant, R. V. & Travis, S. P. L. Current best practice for disease activity assessment in IBD. *Nat. Rev. Gastroenterol. Hepatol.* **13**, 567–579 (2016).

Danese, S., Vuitton, L. & Peyrin-Biroulet, L. Biologic agents for IBD: practical insights. *Nat. Rev. Gastroenterol. Hepatol.* **12**, 537–545 (2015).

Juillerat, P. *et al.* Positioning biologics in the treatment of IBD: A practical guide – Which mechanism of action for whom? *Curr. Res. Pharmacol. Drug. Discov.* **3**, 100104 (2022).

Robinson, P. N. Deep phenotyping for precision medicine. *Hum. Mutat.* **33**, 777–780 (2012).

Delude, C. M. Deep phenotyping: The details of disease. *Nature* **527**, S14–S15 (2015).

Gisbert, J. P. & Chaparro, M. Clinical Usefulness of Proteomics in Inflammatory Bowel Disease: A Comprehensive Review. *J. Crohns Colitis* **13**, 374–384 (2018).

Titz, B. *et al.* Proteomics and Lipidomics in Inflammatory Bowel Disease Research: From Mechanistic Insights to Biomarker Identification. *Int. J. Mol. Sci.* **19**, 2775 (2018).

Bourgonje, A. R. *et al.* Oxidative Stress and Redox-Modulating Therapeutics in Inflammatory Bowel Disease. *Trends Mol. Med.* **26**, 1034–1046 (2020).

Gallagher, K., Catesson, A., Griffin, J. L., Holmes, E. & Williams, H. R. T. Metabolomic Analysis in Inflammatory Bowel Disease: A Systematic Review. *J. Crohns Colitis* **15**, 813–826 (2021).

Aldars-García, L., Gisbert, J. P. & Chaparro, M. Metabolomics Insights into Inflammatory Bowel Disease: A Comprehensive Review. *Pharmaceuticals* **14**, 1190 (2021).

Lee, E. G. *et al.* Systematic Review of Recent Lipidomics Approaches Toward Inflammatory Bowel Disease. *Biomol. Ther. (Seoul)* **29**, 582–595 (2021).

Agrawal, M., Allin, K. H., Petralia, F., Colombel, J.-F. & Jess, T. Multiomics to elucidate inflammatory bowel disease risk factors and pathways. *Nat. Rev. Gastroenterol. Hepatol.* **19**, 399–409 (2022).

Sudhakar, P. *et al.* Tailoring Multi-omics to Inflammatory Bowel Diseases: All for One and One for All. *J. Crohns Colitis* **16**, 1306–1320 (2022).

Fiocchi, C. Omics and Multi-Omics in IBD: No Integration, No Breakthroughs. *Int. J. Mol. Sci.* **24**, 14912 (2023).

Costa, F. *et al.* Calprotectin is a stronger predictive marker of relapse in ulcerative colitis than in Crohn's disease. *Gut* **54**, 364 (2005).

21   Gisbert, J. P. *et al.* Fecal Calprotectin and Lactoferrin for the Prediction of Inflammatory
Bowel Disease Relapse. *Inflamm. Bowel Dis.* **15**, 1190-1198 (2009).

22   Long, N. P. *et al.* Comprehensive lipid and lipid-related gene investigations of host immune
responses to characterize metabolism-centric biomarkers for pulmonary tuberculosis. *Sci.*
Rep. **12**, 13395 (2022).

23   Yen, N. T. H. *et al.* Multimodal plasma metabolomics and lipidomics in elucidating
metabolic perturbations in tuberculosis patients with concurrent type 2 diabetes. *Biochimie*
211, 153–163 (2023).

24   Anh, N. K. *et al.* Metabolic phenotyping and global functional analysis facilitate metabolic
signature discovery for tuberculosis treatment monitoring. *Biochim. Biophys. Acta. Mol.*
Basis. Dis. **1870**, 167064 (2024).

25   Cajka, T. *et al.* Optimization of Mobile Phase Modifiers for Fast LC-MS-Based Untargeted
Metabolomics and Lipidomics. *Int. J. Mol. Sci.* **24**, 1987 (2023).

26   Nguyen, B. T. *et al.* Omics analysis unveils changes in the metabolome and lipidome of
Caenorhabditis elegans upon polydopamine exposure. *J. Pharm. Biomed. Anal.* **244**,
116126 (2024).

27   Tsugawa, H. *et al.* MS-DIAL: data-independent MS/MS deconvolution for comprehensive
metabolome analysis. *Nat. Methods* **12**, 523–526 (2015).

28   Pang, Z. *et al.* MetaboAnalyst 5.0: narrowing the gap between raw spectra and functional
insights. *Nucleic Acids Res.* **49**, W388–W396 (2021).

29   Gu, Z., Eils, R. & Schlesner, M. Complex heatmaps reveal patterns and correlations in
multidimensional genomic data. *Bioinformatics* **32**, 2847–2849 (2016).

30   Rohart, F., Gautier, B., Singh, A. & KA, L. C. mixOmics: An R package for 'omics feature
selection and multiple data integration. *PLoS Comput. Biol.* **13**, e1005752 (2017).

31   Singh, A. *et al.* DIABLO: an integrative approach for identifying key molecular drivers
from multi-omics assays. *Bioinformatics* **35**, 3055–3062 (2019).

32   Peyrin-Biroulet, L. *et al.* Defining Disease Severity in Inflammatory Bowel Diseases:
Current and Future Directions. *Clin. Gastroenterol. Hepatol.* **14**, 348–354. (2016).

33   Bischoff, S. C. *et al.* Intestinal permeability – a new target for disease prevention and
therapy. *BMC Gastroenterol.* **14**, 189 (2014).

34   Gadaleta, R. M. *et al.* Fibroblast Growth Factor 19 modulates intestinal microbiota and
inflammation in presence of Farnesoid X Receptor. *EBioMedicine* **54**, 102719 (2020).

35   Di Ciaula, A., Bonfrate, L., Khalil, M. & Portincasa, P. The interaction of bile acids and
gut inflammation influences the pathogenesis of inflammatory bowel disease. *Intern.*
Emerg. Med. **18**, 2181–2197 (2023).

36   Boyapati, R. K., Rossi, A. G., Satsangi, J. & Ho, G. T. Gut mucosal DAMPs in IBD: from
mechanisms to therapeutic implications. *Mucosal Immunol.* **9**, 567-582 (2016).

37   Boldyreva, L. V., Morozova, M. V., Saydakova, S. S. & Kozhevnikova, E. N. Fat of the
Gut: Epithelial Phospholipids in Inflammatory Bowel Diseases. *Int. J. Mol. Sci.* **22**, 11682
(2021).

38   Kayama, H. & Takeda, K. Emerging roles of host and microbial bioactive lipids in
inflammatory bowel diseases. *Eur. J. Immunol.* **53**, 2249866 (2023).

39   Tews, H. C. *et al.* Unique Metabolomic and Lipidomic Profile in Serum From Patients
With Crohn's Disease and Ulcerative Colitis Compared With Healthy Control Individuals.
Inflamm. Bowel Dis. (2023).

40 Cainzos-Achirica, M. *et al.* Inflammatory Bowel Disease and Atherosclerotic
Cardiovascular Disease: JACC Review Topic of the Week. *J. Am. Coll. Cardiol.* **76**, 2895-
2905 (2020).
41 Leibovitzh, H. *et al.* Immune response and barrier dysfunction-related proteomic
signatures in preclinical phase of Crohn's disease highlight earliest events of pathogenesis.
Gut **72**, 1462 (2023).
42 Ng, J. K. M. *et al.* Neurofascin as a target for autoantibodies in peripheral neuropathies.
Neurology **79**, 2241–2248 (2012).
43 Koike, H. & Katsuno, M. Pathophysiology of Chronic Inflammatory Demyelinating
Polyneuropathy: Insights into Classification and Therapeutic Strategy. *Neurol. The.* **9**,
213–227 (2020).
44 Gao, Y., Kong, L., Liu, S., Liu, K. & Zhu, J. Impact of Neurofascin on Chronic
Inflammatory Demyelinating Polyneuropathy via Changing the Node of Ranvier Function:
A Review. *Front. Mol. Neurosci.* **14** (2021).
45 Helmar Christoph, L., David, B. & Satoshi, K. Chronic inflammatory demyelinating
polyneuropathy: update on diagnosis, immunopathogenesis and treatment. *J. Neurol.*
Neurosurg. Psychiatry **90**, 981 (2019).
46 García-Cabo, C. & Morís, G. Peripheral neuropathy: An underreported neurologic
manifestation of inflammatory bowel disease. *Eur. J. Intern. Med.* **26**, 468–475 (2015).
47 Yang, W. *et al.* Mutant alpha-synuclein causes age-dependent neuropathology in monkey
brain. *J. Neurosci.* **35**, 8345–8358 (2015).
48 Peter, I. *et al.* Anti–Tumor Necrosis Factor Therapy and Incidence of Parkinson Disease
Among Patients With Inflammatory Bowel Disease. *JAMA Neurol.* **75**, 939-946 (2018).
49 Zhu, Y. *et al.* Association between inflammatory bowel diseases and Parkinson's disease:
systematic review and meta-analysis. *Neural Regen. Res.* **17**, 344–353 (2022).
50 Zuo, Y., He, Z., Chen, Y. & Dai, L. Dual role of ANGPTL4 in inflammation. *Inflammation*
Res. **72**, 1303–1313 (2023).
51 Zhang, X. *et al.* Increased angiopoietin-like 4 expression ameliorates inflammatory bowel
diseases via suppressing CD8+ T cell activities. *Biochem. Biophys. Res. Commun.* **612**, 37–
43 (2022).

    FIGURE LEGENDS

Fig. 1. Overall study workflow spanning from patient recruitment, sampling, multi-ome profiling, and data integration, analysis, and interpretation of five clinically meaningful scenarios of treated inflammatory bowel disease patients.

Fig. 2. Single and integrative omics analysis presenting molecular perturbances between treated UC and CD patients. (a) Sample collection and grouping. Heatmap of differential **(b)** proteins, **(c)** metabolites, and **(d)** lipids. **(e)** DIABLO-selected biosignatures.

Abbreviations: IBD: inflammatory bowel disease, UC: ulcerative colitis, CD: Crohn's disease. * and bold: analytes detected in both

single- and multi-omics data analysis.

Fig. 3. Single and integrative omics analysis presenting molecular perturbances associated with fecal calprotectin levels. (a)

Sample collection and grouping. Heatmap of differential **(b)** proteins, **(c)** metabolites, and **(d)** lipids. **(e)** DIABLO-selected biosignatures.

Abbreviations: IBD: inflammatory bowel disease, UC: ulcerative colitis, CD: Crohn's disease. * and bold: analytes detected in both

single- and multi-omics data analysis.

Fig. 4. Single and integrative omics analysis presenting molecular perturbances associated with disease activity states. (a) Sample

collection and grouping. Heatmap of differential **(b)** proteins, **(c)** metabolites, and **(d)** lipids. **(e)** DIABLO-selected biosignatures.

Abbreviations: IBD: inflammatory bowel disease, UC: ulcerative colitis, CD: Crohn's disease. * and bold: analytes detected in both

single- and multi-omics data analysis.

Fig. 5. Single and integrative omics analysis presenting molecular perturbances for the pseudo pre- and post-treatment escalation design. (a) Sample collection and grouping. Heatmap of differential **(b)** proteins, **(c)** metabolites, and **(d)** lipids. **(e)** DIABLO-selected biosignatures. Abbreviations: IBD: inflammatory bowel disease, UC: ulcerative colitis, CD: Crohn's disease. * and bold: analytes detected in both single- and multi-omics data analysis.

Fig. 6. Single and integrative omics analysis presenting molecular perturbances between IBD patients who needed treatment

escalation and those who did not. (a) Sample collection and grouping. Heatmap of differential **(b)** proteins, **(c)** metabolites, and **(d)**

559   lipids. **(e)** DIABLO-selected biosignatures. Abbreviations: IBD: inflammatory bowel diseases, Tx escalation: treatment escalation, UC:

560   ulcerative colitis, CD: Crohn's disease. * and bold: analytes detected in both single- and multi-omics data analysis.

TABLES

Table 1. Clinical characteristics by disease subtypes.

	CD, N = 18[1]	UC, N = 57[1]
Age at recruitment [years]	30 (25, 53)	41 (27, 55)
BMI [kg.m^{-2}]	23.6 (20.8, 25.4)	22.0 (20.7, 23.8)
Sex, females	5 (28%)	22 (39%)
Time since diagnosis [years]	1.08 (0.32, 2.07)	1.48 (0.50, 2.71)
Smokers		
-	11 (61%)	41 (72%)
No	6 (33%)	12 (21%)
Yes	1 (6%)	4 (7%)
Disease state[2]		
Active	7 (39%)	38 (67%)
Remission	11 (61%)	19 (33%)
Fecal calprotectin[3]		
Elevated	11 (61%)	19 (33%)
Normal	7 (39%)	38 (67%)
CRP [mg/L]	0.28 (0.08, 1.01)	0.08 (0.04, 0.19)*
5-aminosalicylic acid, Yes	17 (94%)	56 (98%)
Steroids, Yes	3 (17%)	15 (26%)
Azathioprine, Yes	12 (67%)	18 (32%)*
Biologics, Yes	8 (44%)	15 (26%)

* P-value < 0.05. Wilcoxon rank-sum test and Fisher's exact test were used for continuous and categorical variables.

[1] Data are presented as median (interquartile range) or n (%).

[2] The MAYO score cut-off of 1 and CD activity index (CDAI) score cut-off of 150 concerning CD and UC were used to reflect the disease activity in either active (i.e., mild, moderate, or severe conditions) or remission.

[3] A 150 µg/g cut-off was used to categorize participants into elevated or normal groups.

564 **Table 2. Performance regarding the classification of the biosignatures in five scenarios.**

	AUC[1]	P-value
IBD subtypes, CD *vs.* UC	0.80	0.008
Fecal calprotectin levels, elevated *vs.* normal	0.73	0.03
Disease state, active *vs.* remission	0.56	0.42
Post- *vs.* pre-treatment escalation	0.90	0.01
Needed *vs.* not needed treatment escalation	0.77	0.02
[1] Average AUC via cross-validation procedure.		