**Differential effects of prolonged post-fixation on immunohistochemical and histochemical staining for postmortem human brains**

Weiya Ma[1], Yashar Zeighami[1,2*] and Mahsa Dadar[1,2*]

[1]Cerebral Imaging Centre, Douglas Mental Health University Institute and [2]Department of Psychiatry, McGill University, Montréal, Québec, Canada    H4H 1R3

[*] Equal contribution

Corresponding author:

Mahsa Dadar, PhD
Weiya Ma, PhD
Douglas Mental Health University Institute and Department of Psychiatry
McGill University,
6875 LaSalle Boulevard, Verdun, Montreal,
Quebec, Canada, H4H 1R3
email: mahsa.dadar@mcgill.ca

Running title*: Effect of post-fixation on human brain staining*

Number of Pages:  46
Number of figures: 9
Number of Tables: 4

Category: Original research article

**Abstract**

Immunohistochemical (IHC) and histochemical (HC) staining is widely used for human brains postfixed in formalin for years and decades. Understanding the effect of prolonged post-fixation, postmortem interval (PMI) and age on these staining procedures is important for interpreting their outcomes, thus improving diagnosis and research of brain disorders. In this study, we performed both IHC and HC staining for prefrontal cortex (PFC) of postmortem human brains postfixed for 1, 5, 10, 15 and 20 years. A negative correlation was detected between the intensity of neuron marker NeuN, microglia marker Iba1, cresyl violet and Luxol fast blue staining versus post-fixation times. By contrast, a positive correlation was seen between the intensity of astrocyte marker GFAP and H&E staining versus post-fixation times. No correlation was found between most of these parameters versus age or PMI. These data suggest that prolonged post-fixation exerts both positive and negative effects, but age and PMI have limited effects, on these IHC and HC parameters. Hence these differential changes need to be considered in interpretation of the results when using tissues with prolonged post-fixation. It is recommended to perform IHC and HC staining for human brains with similar post-fixation time to offset its impact on downstream analyses.

Keywords: prefrontal cortex, immunohistochemistry, histochemistry, NeuN, GFAP, Iba1, hematoxylin, myelin, postmortem interval.

1. Introduction

Immunohistochemistry (IHC) and histochemistry (HC) are widely used for diagnostic and research purposes. Both approaches are essential to brain research exploring the mechanisms underlying brain functions as well as neurological and psychiatric disorders. Postmortem human brains are important research materials to uncover the underpinning mechanisms. In brain banks worldwide, it is a common practice to postfix the donated postmortem human brains in 10% neutral-buffered formalin, i.e., 4% formaldehyde solution buffered to a neutral pH until future use. As such, most postmortem brains have usually been preserved in neutral formalin for months, years and decades. An important concern in using these tissues is whether prolonged formalin post-fixation of postmortem brains affects the quality of subsequent IHC and HC staining. Although neutral formalin provides excellent tissue preservation to prevent degradation in IHC (Paavilainen et al., 2010), at the same time it leads to cross-linking of proteins and nucleic acids in tissues (Helander, 1994), thus masking antigen epitopes to reduce immunoreactivity in IHC (Arnold et al., 1996) and mRNA signals in in situ hybridization (Mostegl et al., 2011). Hence prolonged formalin post-fixation is generally believed to prevent the efficient use of postmortem human brains archived for years and decades. To improve IHC staining, some procedures including enzyme-based and heat-based antigen retrieval (HIAR) have been applied to break formalin-induced crosslinks in postfixed tissues (Shi et al., 2011). Of these approaches, citrate buffer-based HIAR has widely been used for IHC staining on postfixed human brains (Shi et al., 1998; Jiao et al., 1999; Ramos-Vara and Beissenherz, 2000; Leong et al., 2010). Using HIAR, the effect of prolonged post-fixation on IHC and/or HC staining for paraffin embedded human brain sections were explored (Lyck et al., 2008; Webster et al., 2009;

Alrafiah and Alshali, 2019; Wu et al., 2022). These studies showed that the influence of prolonged post-fixation on IHC and/or HC staining was either negative or unaltered. However, the time spans of post-fixation for the examined group versus the control group from these prior studies were either very short (weeks or months) (Webster et al., 2009; Pikkarainen et al., 2010; Wu et al., 2022) or rather wide (a group with mixed post-fixation times from 1 to 20 years) (Alrafiah and Alshali, 2019). Furthermore, the detailed information on post-fixation times for presented data were missing and no quantitative analysis was performed (Alrafiah and Alshali, 2019), so it was not possible to systematically correlate the qualitative data with the post-fixation times.

In this study, we aimed to determine the effect of formalin post-fixation for 1, 5, 10, 15 and 20 years on IHC and HC staining for postmortem human brains. IHC staining for neuron nuclear specific marker (NeuN), astrocyte marker glial fibrillary acidic protein (GFAP) and microglia marker ionized calcium-binding adaptor molecule 1 (Iba1) are commonly used to evaluate the three major cell types in human brain tissues. NeuN, GFAP and Iba1 are important cell type specific markers involved in various brain functions and in pathogenesis of numerous brain disorders and widely used in brain research (Reddy and Abeygunaratne, 2022). Therefore, it is important to examine whether prolonged post-fixation exerts an effect on IHC staining of the three markers in human brains. Hematoxylin (H) and eosin Y (E) staining as well as cresyl violet (CV) staining are very common HC staining methods extensively used in clinical diagnostic and brain research. The two staining techniques help to identify different types of cells and tissues in the human brain and provide important information regarding the pattern, shape, and structure of cells. Hence it is also necessary to determine the effect of prolonged post-fixation on H&E- and CV-based HC staining for human brains.

Axonal myelination in both central and peripheral nervous system is essential for a fast and efficient transmission of electrical signals along neurons (Nave and Werner, 2014). Demyelination or myelin damage leads to numerous neurological conditions (Stadelmann et al., 2019), including multiple sclerosis (Staugaitis et al., 2012) and Alzheimer's disease (Papuc and Rejdak, 2020). Myelin research on postmortem human brains is essential to understand the underlying mechanisms of these disorders. Luxol fast blue (LFB) staining is a most common HC technique to illustrate normal myelin as well as to assess Wallerian degeneration in diagnosis and research of these diseases. Hence it is important to evaluate the effect of prolonged post-fixation duration on LFB myelin staining for human brains.

Postmortem interval (PMI) was defined as the time elapsed after death until the samples were fixed in formalin. It is generally believed that a shorter PMI could alleviate degradation of proteins and nucleic acids to preserve tissue and cell structures. Prior studies found that PMI over days did not seem to significantly affect IHC staining (Blair et al., 2016; Mizee et al., 2017; Krassner et al., 2023). Regarding the role of the donor age in IHC staining, mixed outcomes were reported (Gonzalez-Maeso et al., 2002; Mizee et al., 2017; Krassner et al., 2023).  In this study, we also examined whether PMI and donor age play a role in IHC and HC staining for human brains.

To address the above-mentioned issues, in the current study we performed NeuN, GFAP and Iba1 IHC staining as well as H&E, CV and LFB HC staining for the prefrontal cortex (PFC) of human brains postfixed for 1, 5, 10, 15 and 20 years. We also performed a series of linear regression analysis to examine the relationships between IHC and HC staining versus post-fixation time, PMI and donor age.

2.  Materials and methods

2.1  Postmortem human brain acquisition

All experiments were performed with the approval of Human Research Ethics Committees of

McGill University and Douglas Mental Health University Institute. After reception of the whole

brain at the Douglas-Bell Canada Brain Bank (DBCBB; https://douglasbrainbank.ca),

postmortem human brains were collected and hemispheres were immediately separated by a

sagittal cut in the middle of the brain, brainstem, and cerebellum. One unsliced hemisphere (right

or left, in alternation), 1⁄2 brainstem, and 1⁄2 cerebellum were fixed in 10% neutral buffered

formalin. For the current study, we requested PFC blocks (n=20) from human brains which have

been postfixed for 1, 5, 10, 15 and 20 years (n=4 per group, in total, n=20). All specimens were

similarly processed and postfixed in the same solution (i.e. 10% formalin), with the only

difference in post-fixation times. All brains were donated to the DBCBB by familial consent

through the Quebec Coroner's Office. Blood toxicology assessment was performed and

individuals with evidence of drugs or psychotropic medications were excluded. Individuals with

a known history of neurological disorders or head injury were also excluded from this study.

Demographic characteristics associated with each sample are listed in Table 1. Groups were

matched as closely as possible for sex, age, race and PMI.

2.2  Cryostat sectioning of PFC samples

After reception from the DBCBB, 20 formalin-fixed PFC containing blocks from human brains

post-fixed for 1, 5, 10, 15 and 20 years were transferred to 30% sucrose in 1X phosphate

buffered saline (PBS) for cryoprotection for at least one week. PFC blocks were embedded in

M1 embedding medium (Fisher Scientific, Saint-Laurent, Quebec, Canada) and cut into 10- and 50-µm-thick sections on a cryostat (Leica, Feasterville, Pennsylvania, USA). The 10-µm-thick sections were cut from the middle part of the blocks and sequentially mounted on gelatin pre-coated slides and stored in -80°C freezer until they were used for HC such as H&E and CV staining. The 50-µm-thick sections were sequentially collected in cryoprotectant contained in 24-well culture plates and kept in -20°C until being used for IHC and HC such as LBF staining.

2.3 NeuN, GFAP and Iba1 immunohistochemistry

On the day of free-floating immunostaining, PFC sections were taken out of the freezer and washed in 1XPBS, then transferred into Eppendorf tubes containing the antigen retrieval buffer (10mM sodium citrate, 0.05% Tween 20, pH 6.0). The Eppendorf tubes were heated on boiling water (100 to 110°C) in a 1-liter beaker for 20 min. Then the tubes were cooled down on ice and sections were rinsed with cold double distilled (dd) $H_2O$ for 10 min. Sections were transferred into 1X PBS containing 0.05% Triton-100 (PBS+T) in wells of a 24-well culture plate. Sections were quenched in 0.03% $H_2O_2$ for 15 min to abolish the endogenous peroxidase in the tissue and next incubated in 50% ethanol for 1 hour to increase the tissue penetration by antisera. Sections were incubated in 10% normal goat serum (NGS) and 1% Bovine albumin diluted in PBS+T for 2 hours to block the endogenous non-specific antigenic sites. Finally, sections were incubated, respectively, in a mouse monoclonal antibody raised against a neuronal marker NeuN antibody (1:5000, Abcam, Cat. ab10424, Toronto, Ontario, Canada), in a rabbit monoclonal antibody raised against a neuron marker (1:5000, Abcam, ab177487), in a polyclonal rabbit antiserum raised against a astrocyte marker GFAP (1: 5000, Novus Biologicals, Cat. NB300-141, Centennial, Colorado, USA) or in a rat monoclonal antibody raised against microglia marker

Iba1 (1:2000, Abcam, Cat. ab283346) for 18 hours at room temperature. Sections were next

incubated in a biotinylated goat anti-mouse, in a biotinylated goat anti-rabbit IgG or in a

biotinylated goat anti-rat IgG (1:200, MJS Biolynx Inc., Brockville, Ontario, Canada) for 1 hours

and further processed by using an Elite ABC kit (MJS Biolynx Inc.) for another 1 hours

according to the manufacturer instruction. Finally, the immunoprecipitates were developed using

3,3-diaminobenzidine (DAB, Sigma-Aldrich, Oakville, Ontario, Canada) as the chromogen and

enhanced by the glucose oxidase-nickel-DAB method (Shu et al., 1988). Between incubations,

PFC sections were washed thoroughly in PBS + T twice. Finally, sections were mounted on the

pre-cleaned super plus glass slides (Fisher Scientific) or gelatin-coated slides, dehydrated in

ascending ethanol solutions and cleared in xylene, and cover-slipped with xylene-based

mounting medium (Micromount, Leica). After air drying overnight, sections were observed

under the bright field of a microscope (Olympus Microscopes, 1X73, PIF, Montreal, Canada)

and the digital images of immunoreactive cells were captured using the CellSens software

(Olympus). Omission of primary antisera in all tested slides resulted in negative

immunostaining.

2.4    Hemetoxylin and eosin Y staining and Cresyl violet staining

On the day of HC staining, slides mounted with the 10-µm-thick PFC sections from the 5 groups

were taken out from -80$^o$C freezer and air dried for about 30 min to completely remove the

moisture.

For H&E staining, sections were directly stained with 0.1% Mayers Hematoxylin solution

(Sigma, Cat. MHS-16) for 10 min before bluing in warm running tape water (pH 7.8) for 5 min.

Sections were stained with 0.5% ethanol eosin Y solution (Sigma, Cat. HT110116) for 30 sec.

Sections were then dipped in ddH$_2$O, 50% and 70% ethanol briefly, then dehydrated in 95%

ethanol and 100% ethanol (2X, 5 min each) and cleared in xylene (2X, 5 min each). Finally,

slides were cover-slipped, aired dried, and observed under the microscope as mentioned above.

The nuclei of stained cells appeared as dark blue, the cytoplasm as pink while the erythrocytes

were stained as bright red. The capillary and small blood vessels were stained as bright red

which can be easily identified in the neuropils.

For CV staining, sections were incubated in 100% ethanol for 5 min and in 95% ethanol

for 3 min. Then sections were rehydrated in 70% ethanol for 30 sec and in ddH$_2$O for 30 sec.

Sections were stained in 0.5% CV solution for 4 min. Sections were rinsed in ddH$_2$O briefly

before being differentiated in 70% acetified ethanol for 30 sec. Sections were then dehydrated in

90% and 100% ethanol (2X, 2 min each). Finally, sections were cleared in xylene (2X, 5 min

each). Slides were cover-slipped, air dried and observed under the microscope as mentioned

above. CV stained Nissl substances in the cytoplasm of neurons appeared as purple.

2.5 Luxol fast blue-Eosin Y- Cresyl violet staining

On the day of LFB-EY-CV staining, 50µm-thick human PFC sections in cryoprotactant were

taken out from -20$^o$C freezer, rinsed in PBS, mounted on the gelatin-coated slides, and air dried

for about 2 hours to completely remove the moisture. Sections were soaked in xylene (2X, 15

min each), 100% and 95% ethanol (2X, 5 min each), to de-fat the myelin in PFC sections.

Sections were then incubated in 1% LFB solution (Luxol fast blue stain kit, Abcam, Cat.

Ab150675) at 45$^o$C overnight. On the second day, sections were briefly rinsed in 95% ethanol

and ddH$_2$O (5 sec each), differentiated in 0.05% lithium carbonate solution, 70% (2X, 30 sec

each), and rinsed in ddH$_2$O briefly. The differentiating procedures were repeated until a sharp

contrast between the dark blue white matter and the colorless gray matter was observed. Briefly rinsed in 70% ethanol, sections were stained in 0.05% acidified ethanol EY for 30 sec. After a brief wash in ddH$_2$O, sections were stained in 0.5% CV solution for 1 min. Then sections were rinsed in ddH$_2$O briefly, dehydrated through 95% and 100% ethanol and cleared in xylene (2X, 5 min each). Section mounted slides were cover-slipped, air dried and observed under microscope as mentioned above. Myelin in the white matter of PFC sections were stained as dark blue or blue by LFB, but sparse or absent in the gray matter, particularly absent in the superficial layers. Neuropil or the cytoplasm of cells were stained as pink while the red blood cells in the blood vessels were stained as bright red by EY. The cytoplasm of neurons was specifically stained by CV as purple.

2.6  Image capture, quantification and statistical analysis for IHC and HC stained sections

In addition to the same staining procedures for all sections from the 5 groups, the setup for image capturing was consistent for all PFC sections from the 5 groups. Bright field images of IHC or HC stained profiles were captured at the same magnification (20X), with the same threshold of illumination or exposure time on the Olympus microscope (1X73 PIF). Each captured field results in an image with a dimension of 650µm x 475µm. The average optical intensity for each image was measured automatically using ImageJ (ImageJ ver.1.53, Softonic, Barcelona, Spain). The values of average optical intensity measured fall into the 1 to 256 range of the grayscale.

The intensity from the measurements from 5 images were averaged for each brain sample. The mean intensities from all groups (1, 5, 10, 15 and 20 years of post-fixation) were statistically compared by using one-way ANOVA with Student-Newman-Keuls multiple comparison tests using SigmaPlot V14 (Grafiti, St Palo Alto, California, USA). Linear regression

analysis was performed to examine the relationships between the intensity of each staining and the post-fixation years, donor age and PMI for all samples from the 5 groups (n=20) using the software of SigmaPlot v14 (Grafiti). P and R values were determined by the software automatically. For all analyses, the significance level was set as p<0.05.

3. Results

3.1 Effect of prolonged post-fixation on NeuN, GFAP and Iba1 IHC staining

In the current study, we used two monoclonal antibodies raised against NeuN provided by Abcam (see Table 2). Using the mouse monoclonal anti-NeuN antibody for the 50-µm-thick PFC section from the 5 post-fixation groups, we observed that abundant MsxNeuN-IR neurons were present in the gray matter of human brains postfixed from 1 to 20 years (Fig 1A to 1E). MsxNeuN-IR neurons exhibited typical and clear neuron morphology with IR mainly localized in the nuclei, but also seen in the cytoplasm. From quantitative analysis, the MsxNeuN-IR intensity from the 10, 15 and 20-year-groups, but not from the 1-year-group, was significantly lower than from the 5-year-group (Fig. 1F, p>0.05). No significant difference was found among the 1, 10, 15 and 20-year-groups. Linear regression analysis showed a significant negative correlation between the MsxNeuN-IR neurons and the postfixation years (Fig. 2A, p<0.03, R= -0.486).

We also used another rabbit monoclonal anti-NeuN antiserum from Abcam (Table 2) for the 50-µm-thick PFC section of the 5 groups. It was the same antibody used by a prior study showing a negative correlation of NeuN-IR and post-fixation time (Wu et al., 2022). Abundant RbxNeuN-IR neurons were only found in the gray matter of PFC sections from the 1- year-group (Fig 3A). Reliable RbxNeuN-IR neurons were invisible in the 5, 10, 15 and 20 year-groups (Fig

3B to 3E). Quantitatively, the intensity of RbxNeuN-IR neurons was significantly lower in PFC

of the 5, 10, 15, and 20-year-groups than the 1-year-group (Fig. 3F, p<0.001). Linear regression

analysis showed a significant negative correlation between the intensity of RbxNeuN-IR neurons

and the post-fixation years (Fig. 2B, p<0.001, R=-0.711). Together, the above observations

suggest that prolonged post-fixation reduces NeuN IHC staining for human brains.

Using a rabbit anti-GFAP antibody from Novus Biological (Table 2), we found that

GFAP-IR astrocytes were abundantly distributed throughout the gray and white matter of PFC

section from human brains postfixed for 1, 5, 10, 15 and 20 years (Fig 4A to 4E). GFAP-IR

profiles exhibited typical characteristics of astrocytes, i.e., stellate soma with numerous rather

long processes and branches. Quantitatively, the intensity of GFAP-IR astrocytes from the 1, 5,

10, 15-year-groups was significantly lower than from the 20-year-group (Fig. 4F, p<0.05 to

0.001). No significant difference was detected among the 1, 5, 10 and 15-year-groups. Linear

regression analysis disclosed a significant positive correlation between the intensity of GFAP-IR

astrocytes and the post-fixation times (Fig 2C, p<0.014, R=0.542). These data suggest that

prolonged post-fixation enhances GFAP IHC staining for human brains.

In the current study, we used a rat anti-Iba1 monoclonal antibody (Table 2) for IHC

staining on the 50-µm-thick PFC sections of human brains postfixed for 1, 5, 10, 15 and 20

years. We found that Iba1-IR microglia were abundantly distributed throughout the gray and

white matter of the PFC of human brain postfixed for 1 (Fig 5A) and 5 (Fig 5B) years. Iba1-IR

microglia exhibited typical characteristics of microglia, with small irregular soma and multiple

short processes. Numerous Iba1-IR microglia were also present in PFC of human brains

postfixed for 10 (Fig 5C) and 15 (Fig 5D) years, but were much fewer in PFC of human brain

postfixed for 20 years (Fig 5E). Quantitatively, the intensity of Iba1-IR microglia from the 5, 10,

15 and 20-year-groups was very significantly reduced compared to the 1-year-group (Fig 5F, p<0.001). No significant difference was detected among the 5, 10, 15 and 20-year-groups. Linear regression analysis revealed a significant negative correlation between the intensity of Iba1-IR microglia and the post-fixation years (Fig. 2D, p<0.006, R=-0.596). These observations suggest that prolonged post-fixation reduces Iba1 IHC staining on human brains.

3.2 Effect of prolonged post-fixation on hemetoxylin & eosin Y staining

In this study, we examined H&E staining for the 10-µm-thick PFC sections of human brains post-fixed for 1, 5, 10, 15 and 20 years. The nuclei of H&E stained cells were dark blue while the cytoplasm was pink (Fig. 6A to 6E). In contrast, red blood cells in the small vessels (Fig 6A to 6E, arrows) and capillaries (Fig 6D and 6E, arrowheads) were stained as bright red in the neuropils. Quantitatively, the intensity of H&E staining from the 5 and 15-year-groups was significantly higher than from the 1-year-group (Fig. 6F, p<0.05 and p<0.001, respectively). No significant difference was detected between any other two groups (Fig. 6F). Linear regression analysis showed a significant positive correlation between the intensity of H&E staining and the post-fixation years (Fig. 7A, p<0.011, R=0.553). These data suggest that prolonged post-fixation enhances H&E staining for human brains.

3.3 Effect of prolonged post-fixation on cresyl violet staining

We performed CV staining for the 10-µm-thick PFC sections from human brains postfixed for 1, 5, 10, 15 and 20 years. CV stained neurons were observed in the gray matter of PFC sections from all groups (Fig. 8A to 8E). Nissl substances in the cytoplasm of neurons was stained as purple or dark purple. No apparent differences in the abundance and intensity of CV stained neurons among the 1 and 5-year-groups (Fig 8A and 8B). However, the abundance and intensity

of CV stained neurons in PFC of the 10, 15 and 20-year-group were evidently reduced (Fig 8C to 8E). Quantitatively, the intensity of CV staining from the 10, 15 and 20-year-groups was significantly lower than from the 1 (Fig. 8F, $p<0.05$) and 5-year-groups (Fig. 8F, $p<0.05$). No significant difference was detected among the 10, 15 and 20-year-groups. linear regression analysis revealed a very significant negative correlation between the intensity of CV staining and the post-fixation years (Fig. 7C, $p<0.011$, $R=-0.558$). These data suggest that prolonged post-fixation reduces CV staining for human brains.

3.4 Effect of prolonged post-fixation on myelin staining

In this study, using LFB-EY-CV staining protocol, we performed myelin HC staining for the 50-µm-thick PFC sections from human brains postfixed for 1, 5, 10, 15 and 20 years. We observed that the myelin enriched white matter of PFC from all brains of the 5 groups were stained as dark blue or blue by LFB while the neuropils and blood vessels were stained as pink and red by EY (Fig 9A to 9E). In the gray matter (data not shown), where myelin was absent or sparse, neurons were stained by CV as purple. Qualitatively, no obvious difference in the abundance and intensity of stained myelin could be found among the 5 groups (Fig 9A to 9E). However, stained myelin bundles in PFC of the 10, 15 and 20-year groups (Fig 9C to 9E) were apparently more diffuse in the margin and the myelin neurofilaments were not as clear as those in the 1 and 5-year groups (Fig.9A and 9B). Quantitatively, the myelin intensity from the 15 and 20-year groups appeared lower than those from the 1, 5 and 10-year groups, but the difference was not statistically significant (Fig. 9F). However, linear regression analysis revealed a significant negative correlation between the myelin intensity and the post-fixation years (Fig. 7C, $R=-0.461$, $p<0.041$,). These data suggest that prolonged post-fixation reduces myelin staining of human brains.

3.5 Effects of donor age and postmortem interval on IHC and HC staining

In the current study, to examine the possible effect of donor age and postmortem interval on IHC and HC staining we performed linear regression analysis (Tables 3 and 4). There was a significant negative correlation between donor age and the intensity of GFAP-IR astrocytes in PFC of human brains (Table 3, R=-0.483, p<0.031). However, there was no significant correlation between donor age and the intensity of any other IHC or HC staining (Table 3). Moreover, we did not observe any significant correlation between PMI and the intensity of any IHC and HC staining (Table 4). These data suggest that age and PMI exert limited effect on IHC and HC staining.

4. Discussion

4.1 Summary of major findings

In the current study, we examined NeuN, GFAP and Iba1 IHC staining as well as H&E, CV and LFB HC staining on PFC sections from human brains postfixed for 1, 5, 10, 15 and 20 years. A negative correlation was detected between the IHC staining of NeuN and Iba1and the HC staining of CV and LFB versus the post-fixation times. By contrast, a positive correlation was found between GFAP IHC staining and H&E HC staining versus the post-fixation times. These data suggest that prolonged post-fixation exerts differentiating facilitating and inhibiting effects on IHC and HC staining. Furthermore, except for a negative correlation between age and GFAP IHC staining, no significant correlation was detected between age or PMI versus any IHC and HC staining. These data suggest that unlike post-fixation time, age and PMI play limited roles in IHC and HC staining of postmortem human brains.

4.2 Effects of prolonged post-fixation on NeuN, GFAP and Iba1 IHC staining for human brains

Prior studies showed a negative correlation between NeuN-IR intensity and post-fixation times ranging from immediate to 3 years (Wu et al., 2022) or from post-fixation durations of 24 hours, 4 months and 10 years (Lyck et al., 2008). Another prior IHC study on postfixed swine brain showed that NeuN expression was reduced in swine brain postfixed for 2 months, but completely disappeared after 3-month-postfixation (Lundstrom et al., 2019). These prior findings suggest that prolonged post-fixation even within a couple of months or years suppresses NeuN expression in IHC staining. In this study, we greatly expanded the time span of examination for human brain tissues. Consistent with these prior studies, we detected a negative correlation between NeuN-IR intensity and post-fixation time on human brains by using two anti-NeuN antisera from Abcam. However, the two antisera resulted in rather different IHC staining patterns although the trend of post-fixation exerted effect was similar. Reliable NeuN-IR neurons stained with a mouse monoclonal antibody appeared in all groups, even visible in brains from the 20-year-group. However, NeuN-IR neurons stained with a rabbit monoclonal antibody, the same antibody used in the study by (Wu et al., 2022), only existed in brains from the 1-year-group. Our data suggest that the effect of prolonged post-fixation might also depend on used antibodies or relevant molecular changes in tissues. A suitable NeuN antibody could reveal target protein even in brain samples postfixed for two decades even though the staining intensity was significantly reduced compared to samples from early years of post-fixation. As such, it is possible to compensate for the prolonged post-fixation reduced immunoreactivity with a more effective antibody. It is recommended to assess more available antibodies and find the most optimal antibody to perform IHC staining for postmortem brains, thus offsetting the negative effect of prolonged post-fixation on IHC staining.

GFAP is an important biomarker widely used in diagnostic and research of neurological disorders such as Alzheimer disease (Abdelhak et al., 2022; Kim et al., 2023). Astrocytosis could be an alternative pathway for the pathogenesis of AD (Pereira et al., 2021). To understand the mechanisms underlying neurological disorders, it is important to examine GFAP expression in archived human AD brain postfixed for years and decades. Previously, it was shown that GFAP levels was reduced in human brains with post-fixation ranging from 24 hours to 4 months and up to 10 years by immunoarray, but GFAP-IR astrocytes were also sufficiently seen in tissue fixed for 10 years when applying HIER in IHC (Lyck et al., 2008). Another study showed that GFAP-IR intensity remained unaltered in human brains postfixed ranging for months up to 3 years by IHC staining (Wu et al., 2022). Contrary to these studies, for the first time we observed here a positive correlation between GFAP-IR intensity and post-fixation time ranging over 1, 5, 10, 15 and 20 years. We further observed that the mean GFAP-IR intensity from the 20-year-group was significantly higher than other 4 groups, but no significant difference was detected among other 4 groups. Indeed, we also observed abundant strongly stained GFAP-IR astrocytes in PFC and hippocampus of human brain postfixed for 25 years (n=1, supplemental Fig. 1). Our data strongly suggest that prolonged post-fixation exerts a positive or facilitating effect on GFAP IHC staining. Of course, more GFAP antisera available from other commercial sources are required to confirm this conclusion. Detection of GFAP in human brains postfixed for >20 years definitively paves a novel way for the use of large quantities of donated and archived AD brains to uncover the underlying mechanisms of this deteriorating disease.

Increased biomarker for microglia is a consistent feature of AD brains (Hopperton et al., 2018), suggesting a crucial role of microglia activation in AD pathogenesis. Thus it is important to determine the effect of post-fixation on microglia marker such as Iba1 in IHC studies. Prior

studies using immunoassay showed that microglia marker (CD45+ and CD68) in postmortem

human brains was negatively correlated with post-fixation time (from 24hr to 4 months and up to

10 years) (Lyck et al., 2008). However, a subsequent study showed no alteration in Iba1 IHC

staining detected in human brains postfixed from weeks up to 3 years (Wu et al., 2022).

Consistent with the study by Lyck et al (Lyck et al., 2008), in the current study, we found a very

significant negative correlation between Iba1-IR intensity and post-fixation time. The mean

intensity of Iba1-IR microglia was significantly reduced in human brains postfixed for 5, 10, 15

and 20 years compared to the 1-year-group. These data suggest that extended post-fixation

duration reduces Iba1 IHC expression in human brains.

It remains unknown why prolonged formalin post-fixation exerts differential positive

and negative effects on IHC staining. For the negative effect, it was assumed that prolonged

post-fixation progressively enhances the crosslinking between proteins to mask most antigen

epitopes, thus leading to the reduced immunoreactivity. On the contrary, formalin induced

protein crosslinking in human tissues might change protein conformation to expose some

antigens. However, in our pioneer study (Supplemental Fig. 1), we observed that citrate buffer

(pH6.0) based HIAR, which breaks up formalin induced crosslinks, enhanced GFAP IHC

staining for brains postfixed for 25 years. This observation indicates that formalin induced

crosslinks is unlikely to be the reason for the positive effect of prolonged post-fixation on GFAP

IHC staining. Moreover, we still observed that prolonged post-fixation also exerts differential

positive and negative effects on HC staining (see below), which further support the assumption

that more antigen exposures by formalin induced crosslinks is unlikely the reason for the positive

effect.

Prior studies suggest that molecular-dependent changes occurring in prolonged formalin-fixed brain tissues are likely associated with the effect of post-fixation on IHC (Lyck et al., 2008; Wu et al., 2022). Indeed, it was shown that the effects of post-fixation delays >12 hour specifically relied on each antigen since IR-cells could be artificially increased, decreased, or unaffected depending upon the targeted antigen (Lyck et al., 2008; Engel and Moore, 2011). Based on these findings, the differential positive and negative effect of prolonged post-fixation on IHC staining might be the reflection of the molecular changes in brain tissues.

4.3  Effects of prolonged formalin post-fixation on H&E, CV and LFB HC staining for postmortem human brains

Hematoxylin and eosin (H&E) staining is the standard histological protocol as it produces a good overview of the tissue and the cellular components to clearly display different types of structures. In the current study, for the first time we demonstrated a positive correlation between H&E staining intensity and post-fixation time of 1, 5, 10, 15 and 20 years. This observation suggest that post-fixation improves H&E HC staining.

Cresyl violet (CV) staining is a fundamental method to visualize the rough endoplasmic reticulum and ribosomes, the Nissl substance, along the course of dendrites or in neuronal perikaryons. Regarding the effect of prolonged post-fixation on CV staining, it was previously shown that intensity of CV stained neurons was negatively correlated with post-fixation time (from weeks to 3 years) (Wu et al., 2022). Consistently, we found a negative correlation between CV staining intensity and post-fixation time ranging over 1, 5, 10, 15 and 20 years.

In addition to magnetic resonance imaging (MRI) and IHC staining for myelin basic protein (MBP), Luxol fast blue (LFB) staining is a most common HC technique to examine

myelin and Wallerian degeneration in diagnostic and research. Prior studies showed that formalin post-fixation of postmortem human brains for 43 and 64 days altered MRI indices of myelin images (Schmierer et al., 2010; Shatil et al., 2018). Using tissue immunoassay, it was reported that MBP-IR was reduced in human brains postfixed from 4 months to 10 years (Lyck et al., 2008). Another more recent qualitative study showed that prolonged post-fixation of human brains (a group of 1 to 20 years) reduced myelin intensity on LFB staining compared to the control group (< 1 year) (Alrafiah and Alshali, 2019). Consistent with these prior studies, we found a negative correlation between myelin intensity and post-fixation time (1, 5, 10, 15, and 20 years) using linear regression analysis. However, using one-way ANOVA with multiple comparison test, no significant difference in the myelin intensity was detected among the 5 groups, although myelin intensity of the 15- and 20-year-groups were evidently lower than these of the 1, 5 and 10 year-groups. These data suggest that prolonged post-fixation reduces myelin staining, but this reduction in the first 10 years was not evident.

Taking above mentioned data into consideration, a question was raised as to why prolonged post-fixation exerts differential positive and negative effects on HC staining. Formalin induced protein crosslinks seem suitable to explain a stronger H&E staining as post-fixation prolongs. But this presumption fails to interpret the negative effect of prolonged post-fixation on CV and LFB staining. Alternative theories must be explored such as degradation of certain proteins such as myelin or structures as well as Nissl substances. But the loss of NeuN stained neurons and Iba1 stained microglia is surpassed by the gain of GFAP stained astrocytes or H&E stained other cell types so that an enhanced H&E staining could be noticed. Further studies are needed to address these issues.

It is worth mentioning that most prior studies exploring the effect of prolonged post-fixation on IHC and HC staining were performed on paraffin-embedded human brain sections. The current study for the first time examined this issue on thin and thick cryosections of postmortem human brains. Although paraffin embedding provides better tissue preservation, its process also prevents the penetration of antibodies (Ramos-Vara et al., 2014). Cryosectioning is a fast and convenient technique to obtain large quantities of thin (10 to 20 µm thick) and thick (40 to 50µm) sections in a short time. IHC for mounted thin sections or for free-floating thick sections makes staining more versatile and flexible. Following HIAR, the quality of IHC staining was comparable for both paraffin embedded sections and cryosections (Shi et al., 2011). It was also shown that formalin post-fixation and subsequent cryosectioning did not affect the size and shape of ocular tissues (Tran et al., 2017). Our data support the reliability of using cryosections from human brains postfixed for years and decades in research of neurological and psychiatric disorders.

4.4  Effects of age and postmortem interval on IHC and HC staining

In this study, except for a negative correlation between age and GFAP IHC staining, no significant correlation between age or PMI versus IHC and HC staining was observed. Our data were consistent with previous studies showing no correlation between PMI and IHC. For example, it was shown that immunostaining profiles for most proteins remained unchanged even after PMI of over 50 h (Blair et al., 2016) and cellular features appeared to remain intact for more extended PMI (Krassner et al., 2023). Even when RNA was degraded, the protein levels in postmortem human remained stable (Stan et al., 2006). All these data suggest that PMI up to 3 days likely exerts no major effects on IHC and HC staining of postmortem brains. A recent

review mentioned that the evidence for the interaction of age with PMI effects is mixed and may depend on the structural feature considered (Krassner et al., 2023). Isolation of primary microglia from the human post-mortem brain was not affected by donor age and PMI (Mizee et al., 2017). However, a negative correlation was reported between age and immunoreactivity for G-protein subunit $G_{\alpha i1/2}$-, but not for subunit $G_{\alpha i3}$-, $G_{\alpha o}$-, and $G_{\alpha s}$-proteins (Gonzalez-Maeso et al., 2002). In this study, we only detect a negative correlation between age and GFAP-IR, but not other IHC and HC staining. GFAP expression was known to be increased in aging human brains (Wruck and Adjaye, 2020; Chatterjee et al., 2021). Elevated GFAP expression is likely involved in the pathogenesis of neurodegenerative diseases such as Alzheimer's disease, Parkinson's disease and amyotrophic lateral sclerosis (Palmer and Ousman, 2018). However, in our brain samples without any neurological disorders, we found that GFAP IHC staining was negatively affected by donor age, but positively affected by post-fixation time. Hence, to achieve consistent results, it is recommended to perform GFAP IHC staining on postmortem brain within similar ranges of age and post-fixation time.

4.5  Significance of the findings from the current study

In the current study, we demonstrated a positive effect of prolonged post-fixation on GFAP IHC staining and H&E HC staining as well as a negative effect on NeuN, Iba1 IHC staining and CV and IFB HC staining for PFC from postmortem human brains. The differential positive and negative effects could result from the molecular conformational changes associated with post-fixation times. Moreover, we found no correlation between the donor age or PMI with IHC and HC staining for most target proteins or structures, except for GFAP, Thus, successful IHC and HC staining depends on post-fixation times, target molecules, available antibodies and staining

procedures. Therefore, it is recommended to perform IHC and HC staining for postmortem human brains postfixed at the same times, to use the most optimal antibodies and staining procedures. Since prolonged post-fixation exerts both positive and negative effects on IHC and HC staining, depending on the target molecules, it is necessary or unnecessary to perform these procedures as soon as possible after post-fixation is initiated. Since all IHC and HC staining was performed on thin or thick cryosections and reliable staining results were achieved, this study also supports the application of cryosectioning in morphological studies of postmortem brains.

The current study expands the time span of possible application of archived human brains postfixed for years and decades for clinical diagnosis and brain research.  Hence the data collected from the current study could be used to model and adjust for the impact of post-fixation on some of IHC and HC staining measures, if not all. Exploration of these long archived and postfixed human brains will advance our understanding the mechanisms underlying neurological and psychiatric disorders.

5.  Acknowledgements

We are grateful to Dr. Eve-Marie Frigon from Université du Québec à Trois-Rivières for her technical help.

6. Competing interests

The authors have no competing interest.

7. Funding sources

The current study was supported by research funds from the Healthy Brains, Healthy Lives (HBHL, Ref# 2b-NISU-18), Natural Sciences and Engineering Research Council of Canada (NSERC, Ref # RGPIN-2023-04038), and Fonds de Recherche du Québec - Santé (FRQS, Ref # CB - 330750) of Dr. Mahsa Dadar.

8. Author contributions

WM performed immunohistochemical and histochemical staining, analyzed all data and drafted the manuscript. MD and YZ designed the concept of this study and participated in the revision of this manuscript. All authors have read and approved the final manuscript.

Table 1. Complete demographic for human brain cohort

| Groups | Genders | Ages | Races | Causes of death | PMI (hours) | Post-fixation (years) |
|--------|---------|------|-------|-----------------|-------------|-----------------------|
| 1yr-1 | M | 50 | Caucasian | Natural | 54.50 | 1 |
| 1yr-2 | F | 41 | Caucasian | Natural | 51.50 | 1 |
| 1yr-3 | M | 19 | Caucasian | Suicide | 46.00 | 1 |
| 1yr-4 | M | 33 | Caucasian | Accidental | 70.50 | 1 |
| 5yr-1 | M | 53 | Caucasian | Accidental | 49.50 | 5 |
| 5yr-2 | M | 59 | Caucasian | Suicide | 47.50 | 5 |
| 5yr-3 | M | 55 | Caucasian | Accidental | 51.00 | 5 |
| 5yr-4 | M | 59 | Caucasian | Natural | 46.42 | 5 |
| 10yr-1 | M | 75 | Caucasian | Natural | 15.88 | 10 |
| 10yr-2 | M | 91 | Caucasian | Natural | 72.68 | 10 |
| 10yr-3 | M | 80 | Caucasian | Natural | 42.58 | 10 |
| 10yr-4 | F | 84 | Caucasian | Natural | 10.72 | 10 |
| 15yr-1 | F | 49 | Caucasian | Suicide | 59.50 | 15 |
| 15yr-2 | M | 81 | Caucasian | Accidental | 98.75 | 15 |
| 15yr-3 | M | 30 | Caucasian | Suicide | 51.50 | 15 |
| 15yr-4 | M | 36 | Haitian | Natural | 55.25 | 15 |
| 20yr-1 | F | 25 | Caucasian | Suicide | 20.00 | 20 |
| 20yr-2 | M | 22 | Caucasian | Suicide | 24.00 | 20 |
| 20yr-3 | F | 55 | Caucasian | Suicide | 36.00 | 20 |
| 20yr-4 | F | 44 | Caucasian | Suicide | 29.50 | 20 |

Footnotes: Age of the individuals (years): 19-91, average: 52.05; Postmortem interval (hours): 10.72-98.75, average: 46.66.

Table 2.  Summary of primary antisera used in this study

| Name | Source | Cat # | species | Dilution | Datasheet website |
|------|--------|-------|---------|----------|-------------------|
| Anti-NeuN | Abcam | ab104224 | Mouse monoclonal | 1:5000 | https://www.abcam.com/products/primary-antibodies/neun-antibody-1b7-neuronal-marker-ab104224.html |
| Anti-NeuN | Abcam | ab177487 | Rabbit monoclonal | 1:5000 | https://www.abcam.com/products/primary-antibodies/neun-antibody-1b7-neuronal-marker-ab104224.html |
| Anti-GFAP | Novus Bilogical | NB300-141 | Rabbit polyclonal | 1:5000 | https://www.novusbio.com/products/gfap-antibody_nb300-141 |
| Anti-Iba1 | Abcam | Ab263346 | Rat monoclonal | 1:2500 | https://www.abcam.com/products/primary-antibodies/iba1-antibody-epr16589-rat-igg2a-chimeric-ab283346.html |

Table 3.  Summary of linear regression analysis between ages and intensity of IHC and HC

| Intensity | Age | Size | R | P |
|---|---|---|---|---|
| MsxNeuN-IR | 19-91 | 20 | 0.0914 | 0.702 |
| RbxNeuN-IR | 19-91 | 20 | 0.390 | 0.089 |
| GFAP-IR | 19-91 | 20 | -0.483 | 0.031 |
| Iba1-IR | 19-91 | 20 | 0.179 | 0.449 |
| | | | | |
| H&E staining | 19-91 | 20 | 0.139 | 0.559 |
| CV staining | 19-91 | 20 | 0.269 | 0.251 |
| LFB staining | 19-91 | 20 | 0.198 | 0.403 |

Footnotes: A significant negative correlation was detected between GFAP-IR intensity and age (R=-0.483, p<0.031). There was no significant correlation between intensity of any other IHC and HC staining versus age.

Table 4.  Summary of linear regression analysis between PMI and intensity of IHC and HC

| Intensity | PMI (hours) | Size | R | P |
|---|---|---|---|---|
| MsxNeuN-IR | 10.72 - 98.75 | 20 | 0.0678 | 0.777 |
| RbxNeuN-IR | 10.72 - 98.75 | 20 | 0.217 | 0.358 |
| GFAP-IR | 10.72 - 98.75 | 20 | 0.257 | 0.275 |
| Iba1-IR | 10.72 - 98.75 | 20 | 0.153 | 0.519 |
| | | | | |
| H&E staining | 10.72 - 98.75 | 20 | 0.193 | 0.416 |
| CV staining | 10.72 - 98.75 | 20 | 0.0583 | 0.807 |
| LFB staining | 4.49 - 98.75 | 20 | 0.0630 | 0.792 |

Footnotes: There was no significant correlation between intensity of any IHC and HC staining versus PMI.

**Figure 1** Expression of MsxNeuN-IR neurons in the gray matter of 50-µm-thick PFC sections of human brains post-fixed for 1, 5, 10, 15 and 20 years. Numerous NeuN-IR neuronal profiles were distributed throughout all layers in the gray matter of PFC sections of human brains postfixed from 1 to 20 years (**A** to **E**). Immunoprecipitates were not only localized in the nuclei, but also in the cytoplasm of stained neurons. Quantitatively, the NeuN-IR intensity from the 10, 15 and 20-year-groups, but not from the 1-year-group, was significantly lower than from the 5-year-group (F, * p>0.05). No significant difference was found among the 1, 10, 15 and 20-year-groups. Mean±SEM, n=4, one-way ANOVA with Student-Newman-Keuls multiple comparison test. The significance level was set at p<0.05. Scale bar = 100µm.

**Figure 2** Linear regression analysis of the correlation between the intensity of MxNeuN-IR neurons (**A**), RbxNeuN-IR neurons (**B**), GFAP-IR astrocytes (**C**) and Iba1-IR microglia (**D**) in PFC of human brains verse the post-fixation times. A significant negative correlation was found between the intensity of MsxNeuN-IR neurons (**A**, p<0.03, R=-0.486), the intensity of RbxNeuN-IR neurons (**B**, p<0.001, R=-0.711), and the intensity of Iba1-IR microglia (**D**, p<0.006, R=-0.596) versus the post-fixation years. However, a significant positive correlation was detected between the intensity of GFAP-IR astrocytes and the post-fixation years (**C**, p<0.014, R=0.542). Linear regression analysis, n=20. Significance level was set as p< 0.05.

**Figure 3.** Expression of RbxNeuN-IR neurons in the gray matter of 50-µm-thick PFC sectopms of human brains post-fixed for 1, 5, 10, 15 and 20 years. Abundant strongly stained NeuN-IR neurons were only observed in PFC of the 1-year-group (**A**). However, reliable NeuN-IR neurons were almost invisible in PFC of all other groups (**B** to **E**). Quantitatively, the intensity of NeuN-IR neurons from the 5, 10, 15 and 20-year-groups was very significantly lower than from the 1-year-group (F, $p<0.001$). Mean±SEM, n=5 sections, one-way ANOVA with Student-Newman-Keuls multiple comparison test. The significance level was set at $p<0.05$. Scale bar = 100µm.

**Figure 4** Distribution of GFAP-IR astrocytes in the 50-µm-thick PFC sections of human brains post-fixed for 1, 5, 10, 15 and 20 years. GFAP-IR astrocytes were abundantly distributed throughout the gray and white matter of PFC from human brain postfixed for 1, 5, 10, 15 and 20 years (**A** to **E**). GFAP-IR astrocytes exhibited typical characteristics of astrocytes, a stellate soma with rather long and branched processes. Quantitative analysis revealed that the intensity of GFAP-IR astrocytes from the 1, 5, 10, 15-year-groups was significantly lower than from the 20-year-group (F, $p < 0.05$ to 0.001). Mean±SEM, n=4, one-way ANOVA with Student-Newman-Keuls multiple comparison test. The significance level was set at $p < 0.05$. Scale bar = 100µm.

**Figure 5** Distribution of Iba1-IR microglia in the 50-µm-thick PFC sections of human brains post-fixed for 1, 5, 10, 15 and 20 years. Iba1-IR microglia were abundantly distributed throughout the gray and white matter of PFC of human brain postfixed for 1 (**A**) and 5 (**B**) years. Numerous Iba1-IR microglia were seen in PFC of human brains postfixed for 10 (**C**) and 15 (**D**) years. However, Iba1-IR microglia were much less abundant in PFC of the 20-year-group (**E**). Quantitatively, the intensity of Iba1-IR microglia from the 5, 10, 15 and 20-years groups was very significantly reduced compared to that from the 1-year-group (**F**, p<0.001). No significant difference was detected among the 5, 10, 15 and 20-years groups. Mean±SEM, n=4, one-way ANOVA with Student-Newman-Keuls multiple comparison test. The significance level was set at p<0.05. Scale bar = 100µm.

**Figure 6** Hematoxylins (H) and Eosin Y (E) staining for 10-µm-thick PFC sections of human brains post-fixed for 1, 5, 10, 15 and 20 years. In H&E-stained PFC sections (**A** to **E**), the nuclei of cells were stained as dark blue by H while the cytoplasm was stained as pink by E. In contrast, the red blood cells in the small vessels (**A** to **E**, arrows) and capillaries (**D** and **E**, arrowheads) were stained by E as bright red, which is easy to identify in the neuropils. Quantitatively, the intensity of H&E staining from the 5 and 15-year-groups was significantly higher than from the 1-year-group (**F,** * to ***, $p<0.05$ to 0.001). There was no significant difference between any other two groups (**F**). Mean±SEM, n=4, one-way ANOVA with Student-Newman-Keuls multiple comparison test. The significance level was set at $p<0.05$. Scale bar = 100µm.

**Figure 7** Linear regression analysis of the correlation between the intensity of H&E staining (**A**), CV staining (**B**) or LFB staining (**C**) in PFC of human brains and the post-fixation years. A significant positive correlation was detected between the intensity of H&E staining and the post-fixation years (**A,** p<0.011, R=0.553). A significant negative correlation between the intensity of CV staining (**B,** p<0.11, R=0.558) or LFB staining (**C,** p<0.041, R=-0.461) and the post-fixation times. Linear regression analysis, n=20. Significance level was set as p< 0.05.

**Figure 8** Cresyl violet staining of the 10µm-thick PFC sections of human brain post-fixed for 1, 5, 10, 15 and 20 years. In CV-stained PFC sections (**A** to **E**), the cytoplasm of neurons was stained as purple. Quantitatively, the intensity of CV staining from the 10, 15 and 20-year-groups was significantly lower than from the 1 (**F**, *, p<0.05) and 5-year-groups (**F**, #, p<0.05). No significant difference was detected among the 10, 15 and 20-year-groups. Mean±SEM, n=4, one-way ANOVA with Student-Newman-Keuls multiple comparison test. The significance level was set at p<0.05. Scale bar = 100µm.

**Figure 9** Luxol fast blue (LFB)-Eosin Y(EY)-cresyl violet (CV) staining of the 50µm-thick PFC sections from human brain post-fixed for 1, 5, 10, 15 and 20 years. In the myelin enriched white matter of PFC (**A** to **E**), heavily LFB-stained myelin profiles were observed. The neuropil and blood vessels were counterstained by EY as pink and bright red, respectively. Neurons in the myelin lacking gray matter was stained as purple (not shown). Quantitatively, the myelin intensity in PFC from the 15 and 20-year groups were lower than those from the 1, 5 and 10-year groups (**F**). However, the difference was not statistically significant (**F**). Mean±SEM, n=4, one-way ANOVA with Student-Newman-Keuls multiple comparison test. The significance level was set at p<0.05. Scale bar = 100µm.

## References

Abdelhak A, Foschi M, Abu-Rumeileh S, Yue JK, D'Anna L, Huss A, Oeckl P, Ludolph AC, Kuhle J, Petzold A, Manley GT, Green AJ, Otto M, and Tumani H. Blood GFAP as an emerging biomarker in brain and spinal cord disorders. Nat Rev Neurol. 2022 18:158-172.

Alrafiah A, and Alshali R. The effect of prolonged formalin fixation on the staining characteristics of archival human brain tissue. Folia Morphol (Warsz). 2019 78:230-236.

Arnold MM, Srivastava S, Fredenburgh J, Stockard CR, Myers RB, and Grizzle WE. Effects of fixation and tissue processing on immunohistochemical demonstration of specific antigens. Biotech Histochem. 1996 71:224-230.

Blair JA, Wang C, Hernandez D, Siedlak SL, Rodgers MS, Achar RK, Fahmy LM, Torres SL, Petersen RB, Zhu X, Casadesus G, and Lee HG. Individual Case Analysis of Postmortem Interval Time on Brain Tissue Preservation. PLoS One. 2016 11:e0151615.

Chatterjee P, Pedrini S, Stoops E, Goozee K, Villemagne VL, Asih PR, Verberk IMW, Dave P, Taddei K, Sohrabi HR, Zetterberg H, Blennow K, Teunissen CE, Vanderstichele HM, and Martins RN. Plasma glial fibrillary acidic protein is elevated in cognitively normal older adults at risk of Alzheimer's disease. Transl Psychiatry. 2021 11:27.

Engel KB, and Moore HM. Effects of preanalytical variables on the detection of proteins by immunohistochemistry in formalin-fixed, paraffin-embedded tissue. Arch Pathol Lab Med. 2011 135:537-543.

Gonzalez-Maeso J, Torre I, Rodriguez-Puertas R, Garcia-Sevilla JA, Guimon J, and Meana JJ. Effects of age, postmortem delay and storage time on receptor-mediated activation of G-proteins in human brain. Neuropsychopharmacology. 2002 26:468-478.

Helander KG. Kinetic studies of formaldehyde binding in tissue. Biotech Histochem. 1994 69:177-179.

Hopperton KE, Mohammad D, Trepanier MO, Giuliano V, and Bazinet RP. Markers of microglia in post-mortem brain samples from patients with Alzheimer's disease: a systematic review. Mol Psychiatry. 2018  23:177-198.

Jiao Y, Sun Z, Lee T, Fusco FR, Kimble TD, Meade CA, Cuthbertson S, and Reiner A. A simple and sensitive antigen retrieval method for free-floating and slide-mounted tissue sections. J Neurosci Methods. 1999  93:149-162.

Kim KY, Shin KY, and Chang KA. GFAP as a Potential Biomarker for Alzheimer's Disease: A Systematic Review and Meta-Analysis. Cells. 2023  12.

Krassner MM, Kauffman J, Sowa A, Cialowicz K, Walsh S, Farrell K, Crary JF, and McKenzie AT. Postmortem changes in brain cell structure: a review. Free Neuropathol. 2023  4.

Leong TY, Cooper K, and Leong AS. Immunohistology--past, present, and future. Adv Anat Pathol. 2010  17:404-418.

Lundstrom Y, Lundstrom P, Popova SN, Lindblom RPF, and Alafuzoff I. Detection of Changes in Immunohistochemical Stains Caused by Postmortem Delay and Fixation Time. Appl Immunohistochem Mol Morphol. 2019  27:238-245.

Lyck L, Dalmau I, Chemnitz J, Finsen B, and Schroder HD. Immunohistochemical markers for quantitative studies of neurons and glia in human neocortex. J Histochem Cytochem. 2008 56:201-221.

Mizee MR, Miedema SS, van der Poel M, Adelia, Schuurman KG, van Strien ME, Melief J, Smolders J, Hendrickx DA, Heutinck KM, Hamann J, and Huitinga I. Isolation of primary

microglia from the human post-mortem brain: effects of ante- and post-mortem variables. Acta Neuropathol Commun. 2017  5:16.

Mostegl MM, Richter B, Dinhopl N, and Weissenbock H. Influence of prolonged formalin fixation of tissue samples on the sensitivity of chromogenic in situ hybridization. J Vet Diagn Invest. 2011  23:1212-1216.

Nave KA, and Werner HB. Myelination of the nervous system: mechanisms and functions. Annu Rev Cell Dev Biol. 2014  30:503-533.

Paavilainen L, Edvinsson A, Asplund A, Hober S, Kampf C, Ponten F, and Wester K. The impact of tissue fixatives on morphology and antibody-based protein profiling in tissues and cells. J Histochem Cytochem. 2010  58:237-246.

Palmer AL, and Ousman SS. Astrocytes and Aging. Front Aging Neurosci. 2018  10:337.

Papuc E, and Rejdak K. The role of myelin damage in Alzheimer's disease pathology. Arch Med Sci. 2020  16:345-351.

Pereira JB, Janelidze S, Smith R, Mattsson-Carlgren N, Palmqvist S, Teunissen CE, Zetterberg H, Stomrud E, Ashton NJ, Blennow K, and Hansson O. Plasma GFAP is an early marker of amyloid-beta but not tau pathology in Alzheimer's disease. Brain. 2021  144:3505-3516.

Pikkarainen M, Martikainen P, and Alafuzoff I. The effect of prolonged fixation time on immunohistochemical staining of common neurodegenerative disease markers. J Neuropathol Exp Neurol. 2010  69:40-52.

Ramos-Vara JA, and Beissenherz ME. Optimization of immunohistochemical methods using two different antigen retrieval methods on formalin-fixed paraffin-embedded tissues: experience with 63 markers. J Vet Diagn Invest. 2000  12:307-311.

Ramos-Vara JA, Webster JD, DuSold D, and Miller MA. Immunohistochemical evaluation of the effects of paraffin section storage on biomarker stability. Vet Pathol. 2014  51:102-109.

Reddy DS, and Abeygunaratne HN. Experimental and Clinical Biomarkers for Progressive Evaluation of Neuropathology and Therapeutic Interventions for Acute and Chronic Neurological Disorders. Int J Mol Sci. 2022  23.

Schmierer K, Thavarajah JR, An SF, Brandner S, Miller DH, and Tozer DJ. Effects of formalin fixation on magnetic resonance indices in multiple sclerosis cortical gray matter. J Magn Reson Imaging. 2010  32:1054-1060.

Shatil AS, Uddin MN, Matsuda KM, and Figley CR. Quantitative Ex Vivo MRI Changes due to Progressive Formalin Fixation in Whole Human Brain Specimens: Longitudinal Characterization of Diffusion, Relaxometry, and Myelin Water Fraction Measurements at 3T. Front Med (Lausanne). 2018  5:31.

Shi SR, Cote RJ, and Taylor CR. Antigen retrieval immunohistochemistry used for routinely processed celloidin-embedded human temporal bone sections: standardization and development. Auris Nasus Larynx. 1998  25:425-443.

Shi SR, Shi Y, and Taylor CR. Antigen retrieval immunohistochemistry: review and future prospects in research and diagnosis over two decades. J Histochem Cytochem. 2011  59:13-32.

Shu SY, Ju G, and Fan LZ. The glucose oxidase-DAB-nickel method in peroxidase histochemistry of the nervous system. Neurosci Lett. 1988  85:169-171.

Stadelmann C, Timmler S, Barrantes-Freer A, and Simons M. Myelin in the Central Nervous System: Structure, Function, and Pathology. Physiol Rev. 2019  99:1381-1431.

Stan AD, Ghose S, Gao XM, Roberts RC, Lewis-Amezcua K, Hatanpaa KJ, and Tamminga CA. Human postmortem tissue: what quality markers matter? Brain Res. 2006  1123:1-11.

Staugaitis SM, Chang A, and Trapp BD. Cortical pathology in multiple sclerosis: experimental approaches to studies on the mechanisms of demyelination and remyelination. Acta Neurol Scand Suppl. 2012:97-102.

Tran H, Jan NJ, Hu D, Voorhees A, Schuman JS, Smith MA, Wollstein G, and Sigal IA. Formalin Fixation and Cryosectioning Cause Only Minimal Changes in Shape or Size of Ocular Tissues. Sci Rep. 2017 7:12065.

Webster JD, Miller MA, Dusold D, and Ramos-Vara J. Effects of prolonged formalin fixation on diagnostic immunohistochemistry in domestic animals. J Histochem Cytochem. 2009 57:753-761.

Wruck W, and Adjaye J. Meta-analysis of human prefrontal cortex reveals activation of GFAP and decline of synaptic transmission in the aging brain. Acta Neuropathol Commun. 2020 8:26.

Wu X, Deng C, Su Y, Zhang C, Chen M, Tian K, Wu H, and Xu S. The effect of prolonged formalin fixation on the expression of proteins in human brain tissues. Acta Histochem. 2022 124:151879.