A direct neural signature of serial dependence in working memory

Cora Fischer[1,2], Jochen Kaiser[1,2], Christoph Bledowski[1,2*]

Affiliations

[1] Goethe University Frankfurt, Institute of Medical Psychology, Heinrich-Hoffmann-

Strasse 10, 60528 Frankfurt am Main, Germany.

[2] Goethe University Frankfurt, Brain Imaging Center, Medical Faculty, 60528 Frankfurt am

Main, Germany

* Corresponding authors:

Christoph Bledowski, Goethe University Frankfurt, Institute of Medical Psychology,

Heinrich-Hoffmann-Strasse 10, 60528 Frankfurt am Main, Germany.

E-Mail: bledowski@em.uni-frankfurt.de

Keywords

working memory | serial dependence | neural representation | multivariate analysis |

magnetoencephalography

Running Title

A direct neural signature of serial dependence

Conflicts of interest

None.

Significance Statement

Our perception is biased towards sensory input from the recent past. While this "serial

dependence" has been well established in behavior, a direct neural correlate of the

phenomenon has not been determined. Applying multivariate analyses to

magnetoencephalographic data during a visual working memory task, we found an

attraction of neural representations to previous stimuli, which was consistent with

behavioral responses. This is the first evidence for a direct neural signature of serial

dependence. In addition, the present task, together with the high temporal resolution of

magnetoencephalography, allowed us to identify the timing of serial dependence. In line

with current theoretical models, we found serial dependence to emerge at late, post-

encoding stages of information processing in working memory.

Abstract

Serial dependence describes the phenomenon that current object representations are

attracted to previously encoded and reported representations. While attractive biases

have been observed reliably and across various modalities in behavioral reports, a direct

neural correlate has not been established. Previous studies have either shown a

reactivation of past information without observing a neural signal related to the bias of

the current information, or a repulsive distortion of current neural representations

contrasting the behavioral bias. The present study recorded neural signals with

magnetoencephalography during a working memory task to identify neural correlates of

serial dependence. Participants encoded and memorized two sequentially presented

motion directions per trial, one of which was later retro-cued for report. Multivariate

analyses provided reliable reconstructions of both motion directions. Importantly, the

reconstructed directions in the current trial were attractively shifted towards the target

direction of the previous trial. This neural bias mirrored the behavioral attractive bias, thus

reflecting a direct neural signature of serial dependence. The use of a retro-cue task in

combination with magnetoencephalography allowed us to determine that this neural bias

emerged at later, post-encoding time points. This timing suggests that serial dependence

in working memory affects memorized information during read-out and reactivation

processes that happen after the initial encoding. Taken together, we identified a direct

neural signature of serial dependence, which occurs during later processing stages of

working memory representations.

A direct neural signature of serial dependence in working memory

Our visual input can change quickly from moment to moment. Nevertheless, we represent

our environment as remarkably stable and coherent across short periods of time. This

coherence is thought to be supported by serial dependence (*1*), i.e., the phenomenon that

a current visual object is reported as more similar to a previously encountered visual object

than it actually was. This attractive bias acts most strongly between current and previous

objects that are close in time and space, that are attended and that are similar to each

other both with regard to the visual feature that has to be recalled and with regard to

accompanying context features (*1–3*). Furthermore, serial dependence occurs at different

levels of object processing including encoding (*4*), memorization (*5*) and read-out or

decision (*6, 7*).

While the cognitive aspects of serial dependence have been intensively studied (for

reviews see (*8*) or (*9*), its neural mechanisms are still largely unclear and a matter of active

research. One important aspect that has been investigated on the neural level is the nature

of the memory traces that impact current processing and the timing of their reactivation.

Single-unit recordings from monkey prefrontal cortex combined with

electroencephalography (EEG) and transcranial magnetic stimulation in humans have

demonstrated that information about a previous stimulus was reactivated shortly before

a new stimulus was encoded (*10*). This study found that the strength of this pre-encoding

reactivation in the prefrontal cortex predicted the attractive bias of the behavioral

response to the new stimulus. Moreover, subjects showed an enhanced serial bias after

single-pulse transcranial magnetic stimulation of their prefrontal cortex, which further

supported a causal link between the observed reactivation and serial dependence.

Another study (*11*) showed that memory traces from previous trials recorded from the

posterior parietal cortex (PPC) in rats reappeared during the inter-trial interval of a working

memory task, were present throughout the next trial and predicted the behavioral bias.

The observed results indicate that frontal and parietal regions are involved in the storage

of memory traces and their transfer into current processing.

Rather than looking for reactivation of past information, recent studies have investigated

the neural activity elicited by the current stimulus itself. This approach provides a more

direct measure of how the current neural representation is influenced by the recent past.

Neural activity in the visual cortex measured with functional magnetic resonance imaging

(fMRI) was influenced by orientations presented on a previous trial during the encoding

and maintenance of a current orientation (*12*). However, as the current and previous

orientations were either identical or orthogonal to each other, it remained unclear whether

this neural bias reflected an attraction or repulsion in relation to the past. In a more recent

fMRI study (*13*), the current neural representation of an orientation in the visual cortex was

repulsed from an orientation remembered in the previous trial, which was in contrast to

an observed behavioral attraction. In line with this finding, a recent

magnetoencephalography (MEG) study (*14*) observed a neural repulsion of a current

representation during encoding, in contrast to a behavioral attractive bias towards past

information. The authors of both studies proposed that the attractive behavioral bias is

driven by post-encoding processes. This view is supported by another computational

model (*15*), which states that early visual representations are repulsed and the behavioral

attraction emerges during post-encoding read-out of memorized information.

Until now, a direct reflection of the attractive bias of current information on the neural

level has not been observed. The current study applied an inverted encoding model

analysis to neural signals measured with magnetoencephalography (MEG) to reconstruct[1]

the information about currently remembered stimuli. We expected that a direct neural

signature of serial dependence should mirror the attractive bias in behavior by showing

---

[1] The term 'reconstruction' or 'reconstructed representation' refers to the reconstructed channel response, i.e. the output of the IEM analysis. It is not supposed to relate to the reconstruction of single-unit responses. It is used throughout the manuscript for brevity and readability.

an attractive shift of the reconstructed information towards the target information in the

previous trial. The aforementioned models (*13*, *15*) have assumed that neural

representations become attracted towards the recent past after encoding. We employed

a task that allowed us to differentiate between early processes including encoding and

maintenance and later, post-encoding read-out processes to inform these models and

determine not only if, but also during which processing stage the neural representation

becomes attracted towards the recent past. More specifically, we asked subjects to encode

two motion directions and memorize them for a short delay, followed by a retro-cue that

indicated which direction served as the target for an upcoming report. This retro-cue

design, in combination with the high temporal resolution of MEG, enabled us to track the

current representations across the whole trial to determine not only if, but also during

which processing stage ~~when~~ the neural representations become attractively biased

towards the direction reported in the previous trial.

Results

<u>Behavioral Results</u>

Ten human participants (seven females) memorized two sequentially presented visual

objects per trial (S1 and S2), while their neural signals were measured via MEG. Each

participant completed two sessions with a total of 1022 trials. The objects were colored

dot fields and participants had to remember the motion direction of the dots. After a short

delay, one of the motion directions was cued via the dot color (red or green, Fig. 1a). As

expected, we observed a serial dependence towards the target of the previous trial, as the

reported target motion direction of the current trial was systematically attracted to the

retro-cued motion direction of the previous trial. This attraction (Fig. 1b) followed a

derivative-of-Gaussian (DoG)-shaped curve with an amplitude parameter of 3.51°

(bootstrapped SD: 0.479°, lower 95% of permutations between -3.51° and 2.03°, $p < .001$,

R^2 = .236) and a w parameter of 0.024, equaling a width of 47.37° (full width at half

maximum, FWHM). We ran one-sided permutation tests (n = 10) to assess whether this

attraction differed from zero. In contrast, we did not observe a significant serial

dependence towards the non-target of the previous trial (amplitude: -1.02°, bootstrapped

SD: 0.788°, middle 95% of permutations between -1.63 and 1.49, $p = .206$, R^2 = .014; w

parameter: 0.046, equaling 24.53° FWHM), which replicated our previous observations (*2,*

3).

Figure 1

Fig. 1. Experimental paradigm and behavioral results. (A) Participants had to remember two motion directions (S1 and S2) per trial (motion directions are indicated here by gray arrows for illustration only). The stimuli were either red or green. After a short delay, a retro-cue indicated which of the two motion directions (red or green) they had to recall by adjusting a randomly oriented line. (B) Response errors in the current trial were biased towards the target direction of the previous trial (pink curve), but not towards the previous non-target direction (grey curve). The x-axis shows the motion direction difference between the previous and the current target, with positive values indicating that the previous target direction was more clockwise than the current target direction. The response error on the current trial is displayed on the y-axis, with positive values indicating a clockwise deviation from the true target direction. Therefore, shifts of the response errors to the lower left and upper right quadrant indicate an attraction whereas shifts to the upper left and lower right quadrant indicate a repulsion. The asterisk marks significant attractive serial dependence towards the previous target direction.

We also estimated serial dependence between the current and the previously cued object with regard to their color congruence. Serial dependence was modulated by task-relevant color. It was stronger when the current item had the same color as the cued item of the previous trial (amplitude = 4.02°, SD = 0. 528°, lower 95% of permutations between -3.70° and 2.41°, p < .001, R^2 = .167) than when their colors differed (amplitude = 2.99°, SD = 0. 538°, lower 95% of permutations between -2.99° and 1.79°, p = .001, R^2 = .106). The amplitude difference amounted to 1.02°, p = .038. This replicated the context effect found in our previous study (2). The behavioral results thus showed that our two-item working memory task, performed during a MEG recording, produced reliably an attractive bias between target items across trials.

<u>MEG Results</u>

We applied an inverted encoding model (IEM) (*16–18*) to the MEG data to obtain

reconstructed representations of the remembered (and currently relevant) motion

directions as well as the no longer relevant target motion direction from the previous trial.

If the MEG signal at a given time point carried direction-specific information, the

reconstruction should show a graded response function with a peak at the given direction.

If there was no direction information present, the reconstruction should be (near to) flat.

To evaluate whether the reconstruction contained direction information, we aligned all

reconstructed response functions to a common center direction (set to 180°), computed

its fidelity and tested it at each time point (in the resolution of 100 ms) against 0 (no

information present), using a cluster-based permutation test (exact test, 1-tailed) (see

Methods for details).

Reconstruction of the remembered directions per epoch

First, we estimated how well we could reconstruct the motion direction of both current

items throughout the current trial. For this aim we divided the current trial into three

epochs including the encoding and maintenance epochs of S1 and S2 (termed S1 and S2

epochs, respectively) and the retro-cue epoch. Specifically, during the S1 epoch, we

reconstructed the direction of S1, during the S2 epoch we reconstructed the direction of

S2 and during the retro-cue epoch, we reconstructed the motion direction of the currently

retro-cued S1 or S2 (current target). For all three epochs, we observed successful

reconstructions of the corresponding motion directions with a fidelity significantly greater

than zero (S1 epoch: $p < .002$, cluster extent: 300 to 1200ms, mean fidelity = 0.139; S2

epoch: $p < .002$, cluster extent: 0 to 1300ms, mean fidelity = 0.120; retro-cue epoch: $p <$

$.002$, cluster extent: 300 to 1500ms, mean fidelity = 0.115) (Fig. 2a-b). This shows that the

information about the motion direction of S1, S2 and the retro-cued target was
represented in the neural signals during the respective epochs.

For completeness, we also reconstructed the direction of S1 during the S2 epoch, as the
direction of S1 was still maintained in working memory in parallel to the concurrent
encoding and maintenance of the S2 direction (Fig. 3 a-b). We observed that during the
S2 epoch, the direction of S1 could be successfully reconstructed with a fidelity
significantly greater than zero (S1 during S2 epoch: $p = .005$, $p_{Bonferroni} = .015$, cluster
extent: 100 to 700ms, mean fidelity = 0.065). However, its mean fidelity within the
significant cluster was significantly smaller than the fidelities of the currently relevant
direction during all other epochs, i.e. S1 during S1 epoch, S2 during S2 epoch and target
during the retro-cue epoch (all p-values below .005).

Furthermore, we tried to reconstruct the motion direction that was not retro-cued during
the retro-cue epoch, i.e., the non-target of a given trial that became task-irrelevant after
the retro-cue. As expected, the information about the non-target item could not be
reconstructed in the retro-cue phase (all cluster p-values > .05).

Figure 2

Fig. 2. Reconstruction, fidelity and systematic shifts of currently relevant motion direction. (A) The

reconstructed direction of the currently relevant motion direction across the course of a trial. The left panel

shows the reconstruction of the S1 direction, the middle panel the reconstruction of the S2 direction and

the right panel the reconstruction of the cued direction (either S1 or S2). Higher reconstruction strength is

depicted in yellow and lower strength in dark blue. Time points (x-axis) with direction information show a

pronounced yellow area around the common center set to zero (y-axis) and darker areas with increasing

distance from this center. (B) Fidelity of reconstructed direction of the currently relevant motion direction

across the course of a trial. The left panel shows the fidelity of the S1 direction, the middle panel depicts the

fidelity of the S2 direction and the right panel shows the fidelity of the cued direction (either S1 or S2). Small

circles show the fidelity of each participant. Colored circles indicate time points with a fidelity significantly

greater than zero (cluster-based permutation test within each epoch). (C) The reconstructed direction during

all time points with a significant reconstruction fidelity was averaged within each epoch and the maximum

of the reconstructed direction was compared to the common center set to zero. A negative shift of the

reconstructed mean indicates a repulsion from the target direction of the previous trial whereas a positive

shift indicates an attraction. For a more detailed visualization, the range of direction shifts from -40° to +80°

is magnified and reconstructed shift values for each participant are indicated by dots. The upper row shows

the bootstrapping distributions (see section on MEG analysis below for details). Significant shifts are marked

with an asterisk.

Attractive shift of motion direction reconstructions as a direct signature of serial

dependence

The main analysis of our study aimed to identify the neural signature of serial dependence.

For this aim we looked for an attractive bias of the reconstructed neural representations

that mirrored the attractive behavioral bias by testing whether the mean of the significant

reconstructions within the S1, S2 or retro-cue epochs showed positive shifts of their

maxima from the common center of 180° towards the previous target (Fig. 2c; see

Methods for details). We found that the shift of the mean reconstruction of the first and

second item did not differ from 0° for the S1 epoch (-2.83°, *p = . 743*, bootstrapping test,

p_{FDR} = .870, 95% of permutations between -11.60° and 6.00°) or for the S2 epoch (-2.50°,

p = .870, bootstrapping test, *p_{FDR} = .870*, 95% of permutations between -7.57° and 2.11°),

respectively. In contrast, for the retro-cue epoch we found that the mean reconstruction

of the current target was shifted towards the previous target (10.22°, *p = .009*,

bootstrapping test, *p_{FDR} = .026*, 95% of permutations between 1.64° and 21.97°; Fig. 2c).

In addition, there was an attractive shift of the S1 direction towards the target of the

previous trial during the S2 epoch (11.69°, *p < .001*, bootstrapping test, 95% of

permutations between 4.96° and 31.06°; Fig. 3c). Thus, our MEG data provided evidence

for an attractive distortion of current neural representations toward the target information

in the previous trial, thereby revealing a direct neural signature of serial dependence.

Fig. 3. Reconstruction, fidelity and systematic shifts of S1 direction during the S2 epoch. (A) The reconstructed direction of the S1 motion direction during the S2 epoch. (B) Fidelity of reconstructed S1 motion direction across the course of the S2 epoch. Small circles show the fidelity values for each participant. Colored circles indicate time points with a fidelity significantly greater than zero. (C) The reconstructed direction for all time points with a significant reconstruction fidelity was averaged within the S2 epoch to indicate attractive and repulsive shifts in relation to the target of the previous trials. Significant shifts are marked with an asterisk. For details, refer to Figure 2.

Correlation between neural representation and behavioral response

To assess whether the observed item reconstructions from the MEG signals were behaviorally relevant, we correlated them with the subjects' behavioral responses.

Specifically, we calculated the circular correlation between the maximum of the stimulus

direction reconstructions and the response error for each time point and epoch separately

across *single trials*, i.e., we tested whether the shift/deviation of the neural response from

the true value, e.g., in the clockwise direction, was accompanied by a corresponding

shift/deviation of the behavioral response.

Indeed, Fig. 4 shows that-the reconstructed shift of the retro-cued target item during the

retro-cue phase predicted the upcoming behavior between 400 to 800ms after the onset

of the retro-cue ($p = .003$, permutation test, $p_{Bonferroni} = .009$, mean r = .030). In contrast,

there were no significant correlations for any other time points during the S1 and S2

epochs (cluster-based permutation test (exact test, 1-tailed). While the maximum of the

S1 reconstruction also showed a significant shift towards the previous target, there were

no significant correlations of the S1 reconstruction with the response error during the S2

epoch (no clusters with a p-value < .05).

Figure 4

Fig. 4. Correlation between single-trial shift of reconstruction and upcoming response. For each time

point, the shift of the maximum of the reconstructed direction in a single trial was correlated with the

response deviation from the target direction. Small circles show the r-values of each participant. Time points

where the shift of the reconstructed direction significantly correlated with the upcoming response deviation

are indicated by colored circles (cluster-based permutation test within each epoch).

The presented single trial-wise correlation between the neural shift and behavioral shift

did not directly incorporate the behavioral bias towards the previous trial. Thus, in order

to relate the neural attractive shift and the behavioral indicators of serial dependence more

directly, we performed an additional correlation analysis on aggregated data on the

between-subject level. In detail, we correlated the individual neuronal shift during the

retro-cue epoch with the two individual parameter fits of the behavior shift, i.e., amplitude

(a) and tuning width (w). In line with the correlation analysis on single-trial level, the

analysis on the *between-subject* level also revealed a significant correlation between the

w parameter of serial dependence and the neural shifts for the retro-cue epoch. Details of

this correlation analysis are displayed in the supporting information "Correlation between

the neural shift and individual serial dependence parameters" and Fig. S1.

Reactivation of the previous target direction

Finally, we used our IEM analysis approach to replicate previous results that have shown a

reactivation of memory traces from the previous trial before the stimulus presentation of

the current trial (*10*). Specifically, we reconstructed the motion direction of the target from

the previous trial during the one second of the inter-trial interval (ITI) immediately

preceding the current trial (Fig. 5). Consistent with previous findings from monkey single

cell and human EEG recordings (*10*), we observed that the information about the previous

target was reactivated just prior to the onset of the next trial. Direction reconstructions

showed a fidelity significantly greater than zero during a time period at the end of the ITI

($p = .019$, cluster extent: -200 to 100ms (relative to S1 onset), mean fidelity = 0.069).

Importantly, we also tracked the information of the previous target throughout the whole

current trial to test whether it was reactivated also during the processing of the current

items. Interestingly, we found that the target from the previous trial was reactivated once

more, but at a later time period during the current trial, i.e., when the retro-cue indicated

which item of the current trial was the target for the upcoming response ($p = .040$, cluster

extent: 2700 to 2900 ms relative to trial onset, mean fidelity = 0.057). In contrast, we found

no evidence of previous target reactivations during the encoding or maintenance of

current items (no clusters with a p-value < .05).

Fig. 5. Reconstruction and fidelity of previous target direction. (A) The target direction of the previous trial was reconstructed throughout the last second of the inter-trial interval (ITI) and the whole following trial. (B) Fidelity of reconstructed direction of the previous target. Small circles show the fidelity of each participant. Colored circles indicate time points with a fidelity significantly greater than zero. For details, refer to Figure 2.

Additionally, in order to test whether a re-activation of the previous target could explain the neural attractive shift observed in the current trial, we reconstructed the motion direction of the previous target from MEG data recorded during the different epochs of the current trial. Specifically, we trained the encoding model either on the presented stimuli in the S1 and S2 epochs or on the currently relevant item during the retro-cue epoch, respectively, and tried to reconstruct the target direction of the previous trial within each epoch. We observed that there was no cross-reconstruction between the current and the previous representation during the retro-cue epoch, which indicates that the observed attractive shift of the current target reconstruction following the retro-cue was not driven by a reactivation of the previous target in the same neural code. However, during the beginning of the S2 epoch, we observed a significant cross-reconstruction, i.e. previous target reactivation co-occurred with the reconstruction of S2 while it was presented to the subjects. The complete analysis and results are described in the supporting information

"Cross-validation reveals different codes for reactivation of previous information and

reconstructed biases of current information" and Fig. S2.

Eye movement control

To control for influences of eye movements on the MEG data, we recorded horizontal and

vertical electrooculograms to identify blinks and eye-movement-related independent

components of the MEG data that were removed during our MEG preprocessing (see

Methods). In addition, we also recorded continuous gaze position with a higher spatial

resolution using an eye-tracker. We used the eye-tracking data to estimate whether there

was a systematic relationship between eye-movements and the motion direction of

currently processed items in the S1, S2 or retro-cue-phases. We found that gaze directions

were systematically related to presentation and memorization of the stimulus direction

(S1 and S2 epoch) and its cue-based retrieval (retro-cue-phase). They also varied

considerably between subjects. Most importantly, gaze directions were not systematically

related to the MEG data, thus ruling out that they had driven our MEG results. The details

of the eye movement control analysis are displayed in the supporting information "Eye

movement control" and Fig. S3.

Discussion

The current study aimed to identify a direct neural signature of serial dependence.

Previous studies searching for neural markers of this attractive bias have investigated

either the reactivation of information from the past trial in the current trial, or have tried

to directly capture biases of current item representations by information from the past.

The present study found evidence for both types of neural correlates. We replicated

reactivations of target representations of the previous trial. More importantly, our study

provides the first evidence for an attractive bias of neural representations in the current

trial towards information from the previous trial, which mirrors the behavioral effects of

serial dependence and as such represents a neural signature of serial dependence.

The reactivation of past information in the current trial is crucial because an attractive bias

can only occur when information from the past has left a memory trace that interacts with

the processing of the current object. Barbosa, Stein and colleagues (*10*) have shown that

the working memory trace of an object representation from a previous trial, measured via

decoding of both neural activity in monkey prefrontal cortex and human EEG signals, was

reactivated just before new visual input was presented in the current trial. This was in line

with modulations of early visual evoked potentials in human EEG by past stimulus

information found in numerosity tasks (*19*, *20*). Using human EEG, Bae and Luck (*21*)

observed the reactivation of past stimulus information also during encoding and

maintenance phases in the current trial of an orientation working memory task. In our

study, we applied a similar analysis approach and tracked the reconstructed motion

direction of the target item from the previous trial both during the inter-trial interval, i.e.,

prior to the encoding of the new stimulus, and throughout the whole current trial. Similar

to the previous studies, we also found that the memory trace of the previous target was

reactivated just prior to the onset of the current trial and later in the current trial after the

retro-cue indicated the current target.

Reactivations of information from the previous trial may be seen as crucial but indirect

neural correlates of serial dependence. A memory trace of a previously relevant

representation is an important prerequisite for the emergence of serial dependence, since

a current representation can only be biased by integrating the information from the past

into the present trial. However, our main goal was to observe the direct effect of this past

information on current representations. More specifically, we asked whether neural

representations of current information are biased towards the recent past in the same way

as the behavioral response. Previous studies that aimed to measure neural correlates of

serial dependence directly found repulsive rather than attractive distortions of neural

representations in relation to the recent past, which were in contrast to the attractive bias

in the behavioral response (*13*, *14*). The repulsive neural biases were attributed to visual

adaptation during early perceptual processes. In contrast to these previous reports, the

present study found attractive biases of representations in the present trial towards

information from the previous trial that were consistent across the neural and behavioral

level.

We propose that our combination of a retro-cue paradigm with the high temporal

resolution of MEG accounts for the successful demonstration of serial dependence at the

neural level in the present study. Sheehan and Serences (*13*) recorded fMRI while

participants made a delayed report about a single memorized grating orientation. In

contrary to the attractive behavioral bias, the orientation reconstructed from the fMRI

signals measured in the visual cortex was repulsed away from the orientation presented

in the previous trial. The authors proposed a computational model in which early

encoding-related signals are repulsed from the recent past. The attraction observed in

behavior was attributed to a later decoding process that overcomes the repulsion of

sensory signals and leads to serial dependence but is not directly detectable in

hemodynamic response patterns. This is consistent with the notion of an ideal observer

model that combines two steps of object processing including an early encoding and a

later decoding of current information to explain serial dependence (*15*). According to this

model, a repulsive bias occurs early during encoding, whereas an attractive bias is due to

the read-out of the current object representation in a Bayesian history-dependent manner.

Previous behavioral studies have already identified read-out as a crucial processing stage

for serial dependence. Fritsche and colleagues (*6*), for example, asked participants either

to remember and report the orientation of a grating after a short delay (working memory

task) or to compare two simultaneously presented orientations directly (perception task).

They found a repulsive bias in the perception task, when the grating was still visually

present and compared to another grating, whereas in the working memory task, the

current target orientation was attracted towards the orientation reported in the preceding

trial. This supported an emergence of serial dependence at a late, post-perceptual,

processing stage.

Similarly, Hajonides and colleagues (*14*) observed a repulsion of currently remembered

orientations as decoded from MEG signals, while the behavioral recall was attracted

towards the past. Participants performed a working memory task with two sequentially

presented sample stimuli similar to our study. However, instead of using a temporally

separated retro-cue, Hajonides and colleagues (*14*) presented a cue that indicated the

target for later recall simultaneously with the second sample stimulus. As participants

knew which item would be the response target as soon as the second item was presented,

the read-out process for a specific item could not be disentangled from the encoding or

maintenance stages. Multivariate analysis of MEG signals revealed that the reconstructed

orientation information during and shortly after stimulus presentation was repulsed from

the orientation recalled in the previous trial. In line with Sheehan and Serences (*13*), the

authors assumed that the observed repulsion at the neural level reflected visual

adaptation, whereas that the attraction of the neural representation may have occurred at

later processing stages, which were not accessible by their analysis. Hajonides and

colleagues (*14*) speculated that the integration of past memory traces, putatively stored

in parietal areas (*11*), only happened during the behavioral response. This would mean

that the neural representation of the current representation is not affected by serial

dependence, but that this bias emerges as a result of integration processes during the

response.

The present study provided support for the notion that the neural markers of serial

dependence occur during post-perceptual phases of a working memory task, which is in

agreement with existing models and behavioral findings (*13–15*). The use of a retro-cue

task, which facilitates the segregation of earlier and later processing stages, in

combination with motion direction reconstructions from MEG signals, revealed shifts of

the current representation towards the motion direction of the target in the previous trial.

We observed such an attractive bias only at post-encoding processing stages in two time

windows: First, for S1 during and after the presentation of S2 and, second, for the cued

target stimulus after the presentation of the retro-cue. This demonstrates that serial

dependence can be observed on the neural level as a distortion of current neural

representations that mirrors the attractive bias in behavior. This direct neural signature of

serial dependence emerges after encoding but prior to the response, thus speaking

against the proposal that serial dependence is attributable to response-related processes

(*14*).

To further underscore the relevance of the observed neural distortions, we examined

whether the reconstructed neural representation predicted the behavioral response

deviation. Indeed we observed a positive correlation between the deviations of the neural

reconstruction and response errors on a single-trial level. Interestingly, this correlation was

found during the presentation of the retro-cue, coinciding with the time point at which

the neural representation of the current target stimulus was biased towards the recent

past. This observation was further supported by a positive correlation between the

attractive shift of the neural reconstruction during the retro-cue epoch and a behavioral

measure of serial dependence on the between-subject level. In addition, we observed a

reactivation of previous target information during the retro-cue presentation. The

distortion of the current neural representation could thus be the result of an integration

of past information. Taking these findings together, the retro-cue period emerges at the

time point when different neural markers of serial dependence appear together. This

suggests that the selective read-out of memorized information for an upcoming report

might be a crucial process for serial dependence during working memory.

The neural representation of S1 already became attracted towards the target of the

previous trial when S2 was presented. Notably, the reconstruction of the S1 direction had

dropped to baseline and reappeared during the S2 epoch. This fits well with the

observation that memory content that has become neurally silent can be successfully

reconstructed from the impulse response elicited by a "ping" or probe stimulus (*22, 23*).

This impulse-driven reconstruction is thought to reflect a "hidden state" of working

memory, in which unattended information is stored. Therefore, the observed S1

reconstruction elicited by S2 presentation in our study likely reflects the same type of

memory read-out, which might have driven the observed neural distortion. On the other

hand, S1 had been maintained for a certain time when S2 was presented. As Bliss and

colleagues (*5*) observed, serial dependence in behavior increases as a function of delay

duration, so the distortion of the neural representation of S1 during the S2 period could

also reflect an increasing bias during working memory maintenance.

Several recent studies have shown that serial dependence operates already during

perception (*4, 24*). In the current study, the neural representation of the current

information became biased only after encoding. However, we did observe a reactivation

of previous information as an indirect marker of SD before and during the onset of S1,

replicating previous results (*10*). This early reactivation of past information could reflect a

comparison of prior information with new visual input. Cicchini and colleagues (*25*)

proposed a predictive coding framework, in which priors informed by past visual input are

compared against incoming visual information during early encoding. This comparison

results in a prediction error, which determines whether a current stimulus will be biased

towards the past. In combination with the observed distortion of the neural

representation, this could indicate a two-step process: New visual information is compared

against past information during early encoding processes and the integration of past and

current information happens during later processing stages, before the actual report is

given.

How can these findings of serial dependence operating already during perception (*4, 24*)

be reconciled with other behavioral studies and our findings that pinpoint the emergence

of serial dependence at post-perceptual processing stages (*5, 6*)? One possibility is that

serial dependence does not operate at a single but rather at different levels of object

processing including early perceptual and later, memory-related processes (*26*). Moreover,

perception and working memory might not be as different as they appear: both functions

build mental representations from sensory input, involving feedforward and feedback

processing. Hence, encoding and decoding of information, including the neural

computations in lower- and higher-level cortical areas, represent integral parts of both

perception and working memory. Thus, in a perceptual task that requires an immediate

response when visual input is still physically present, read-out of the noisy information via

decoding is closely intertwined with its encoding. In working memory tasks that require

encoding and memorization of several items, the read-out of a single item is typically

postponed: It starts when a retro-cue or probe indicates which memorized item is task-

relevant. Serial dependence that operates during the read-out stage of object processing

appears to happen both early in perception and later in working memory tasks. However,

as shown in this study, in order to identify the neural basis of serial dependence, a multi-

item working memory task offers the advantage of separating different processing stages

to isolate the time point when an object representation is actually distorted. Furthermore,

while the present results indicate a particular importance of post-encoding processes for

serial dependence in working memory, they do not suggest that serial dependence could

not emerge during encoding. First, we cannot exclude that our data analysis was not

sensitive enough to detect small shifts in the neural signals during encoding, especially as

our analysis involved averaging across time points to increase the signal-to-noise ratio.

Second, with the used motion direction stimuli, reconstruction strength built up slowly

during the stimulus presentation, reaching its maximum only after stimulus offset.

Possibly, different stimulus material with faster build-ups might help to track the neural

representation during early encoding stages.

In summary, our study provides a direct neural signature of serial dependence that mirrors

the attractive behavioral bias. This demonstrates that serial dependence in working

memory directly acts on memory representations and biases them towards the past. The

attractive distortion of the current representation happened after object encoding,

suggesting that read-out processes during object decoding are important for the

emergence of serial dependence.

Materials and Methods

<u>Experimental Design</u>

Subjects

Thirteen participants were recruited from the Goethe-University Frankfurt and the

Fresenius University of Applied Sciences Frankfurt. All subjects reported normal or

corrected-to-normal vision and were screened for red-green-blindness. One subject

participated as a pilot subject for only one MEG session, one subject was excluded after

the first behavioral training session because of dependency on glasses and one subject

dropped out after the first MEG session due to exhaustion because of the MEG setting.

This resulted in a sample of 10 subjects (3 male), aged between 22 and 30 years (mean:

25.2 years). All subjects gave informed consent and received either a compensation of

€10/h or course credit. The study was approved by the Ethics Committee of the Medical

Faculty of the Goethe-University Frankfurt am Main.

Experimental Paradigm

Stimuli. Random dot patterns (RDP) were presented centrally on the screen and consisted

of 200 dots colored in red (RGB: 255, 0, 0) or green (RGB: 0, 86.4105, 0) on a black

background. The green was adjusted to match the luminance of the red in DKL color space

and then transferred back to RGB space. The dots were presented within an invisible

circular aperture which had a radius of 7.5° of visual angle. The dots had a diameter of

0.15° of visual angle and were placed randomly within the circular aperture of the RDP at

stimulus onset. The dots moved with a velocity of 3.5°/s and fully coherent in a direction

randomly drawn from a pool of directions between 5° and 355° spaced 10° from one

another, therefore avoiding cardinal directions. Dots reaching the edge of the aperture

were repositioned randomly on the edge of the opposing side of the aperture, so that dot

density was kept constant throughout the presentation. Throughout the whole experiment

a white fixation square with a diagonal of 0.15° of visual angle was presented centrally on

the screen, except for the cue presentation, when the fixation square changed its color to

red or green to cue which item should be reported. The item was reported by adjusting a

randomly oriented line to match the reported direction. The response line was white, with

a width of 0.6° and a length equaling the dot field radius. The starting point of the line

was the fixation square and the end point could be altered so that the line could point in

all possible directions.

Procedure. The experiment consisted of a delayed-estimation task. Specifically, each trial

began with the presentation of the first stimulus (S1) for 200 ms followed by a noise mask

for 100 ms consisting of dots moving with 0% coherence (i.e., randomly) and of the same

color as the preceding RDP. After a 1000 ms interval (ISI), the second stimulus (S2) and its

noise mask were presented for 200 ms and 100 ms, respectively (Figure 1). Subjects were

asked to memorized the motion direction of both RDP for a delay of 1000 ms. Then the

fixation square changed its color to red or green for 1500 ms, thereby cueing which motion

direction became a target item and had to be reported. Immediately after cue offset, a

randomly oriented line was presented. Subjects had to report the target motion direction

by rotating the line by moving an MEG-compatible track ball horizontally. No time limit

was given for the response and subjects were encouraged to work as precisely as possible.

After adjusting the line direction, subjects had to confirm their response by pressing the

left track ball button. At the end of each trial, a fixation screen of 2000 to 2500 ms (jittered

in steps of 10 ms, randomized) was presented. Subjects were instructed to fixate the

fixation square throughout the whole experiment.

Between trials, 36 possible direction differences (-170°:10°:180°) were balanced so that

every possible direction difference between the items of the current trial and the previous

target occurred equally often. This was done with the restriction that the items of the

current trial were not allowed to have the same motion direction difference to the previous

target item, i.e. cued for report in the previous trial. Thereby it was ensured that the items

within a trial could not have the same motion direction. Furthermore, the congruence of

color and temporal position between target items of the previous and current trial was

balanced, which automatically lead to a balancing for the non-target item of the current

trial, too and ensured that in each trial the two items had different colors from one another.

Every subject completed 1022 trials in two sessions on different days. Each session lasted

approximately 90 minutes and was divided into 7 blocks of 73 trials with breaks in

between. After the completion of each block, subjects received feedback ('You deviated

by more than 30° in XX% of trials.' – German: 'Du bist in XX% der Durchgänge um mehr

als 30° abgewichen.') about their performance in the previous block, displayed on the

screen. Subjects were seated at a viewing distance of approximately 50 cm from the

display. MATLAB software with the Psychophysics Toolbox extensions (Brainard, 1997;

Pelli, 1997) was used for stimulus generation and presentation. A PROPixx projector (VPixx

Technologies Inc.) with a resolution of 1920 x 1080 and running with 120 Hz refresh rate

was used.

Prior to the two MEG sessions, each subject was invited for a behavioral practice session.

This session consisted of 8 blocks of the task and was therefore slightly longer than the

MEG experiment. We screened the participants for red-green-blindness, checked MEG and

MRI contraindications, and instructed them in the behavioral task before participants

started to practice the task. In addition to the feedback presented at the end of each block

as in the MEG part of the study, subjects received a trialwise feedback in this first session.

If their response deviated more than 30° from the cued motion direction, we presented

feedback ('>30°!'). Thereby they could practice to be as precise as possible for the MEG

sessions.

<u>MEG recoding and preprocessing</u>

Data acquisition

We recorded MEG with a whole-head MEG system (Omega 2005; VSM MedTech Ltd., Port

Coquitlam, Canada) with 275 axial gradiometers (271 active) at a sampling rate of 1200 Hz

and without online filtering. Additionally, we recorded electrocardiogram (ECG), vertical

and horizontal electrooculograms (EOG) as well as continuous gaze position and pupil

dilation of the right eye by an MEG compatible eye-tracker (Eyelink CL 1000, SR Research

Ltd.). Head position was continuously recorded via head localization coils placed at the

nasion and above both ear canal entrances using ear-plugs. The initial head position was

saved at the beginning of the first MEG session and presented to the participants for

repositioning at the beginning of the second session as well as throughout each session

if repositioning was necessary. Behavioral responses were recorded with an MEG-

compatible trackball (Current Designs Inc.), enabling subjects to continuously recall

motion directions.

Preprocessing

MEG data were preprocessed using the python-based M/EEG analysis toolbox MNE

(version 0.22.0; (*27*)). The MEG signal was first notch-filtered at 50Hz and up to 250Hz in

steps of 50Hz to remove line noise and then cut into trials from 1000 ms before S1 onset

until 500 ms after response onset. Trials containing SQUID jump artifacts were manually

identified and excluded. The remaining trials were then highpass-filtered (1Hz) and

decomposed via independent component analysis (ICA) with the fastica method

implemented in MNE. The decomposition was then applied to the notch-filtered

continuous data and ICs representing cardiac and ocular activity were identified via

correlation with ECG and EOG signals. The notch-filtered continuous data were then

lowpass-filtered at 25Hz and cut for the respective analysis. To analyze reactivations of the

previous target direction, the data were cut into one long epoch (last second of the ITI up

until the end of the retro-cue delay). For the analysis regarding the directions of the

current trial, we used three epochs per trial: S1 encoding and maintenance (called S1

epoch), S2 encoding and maintenance (called S2 epoch) and retro-cue presentation and

delay (called retro-cue epoch). The epochs started 200 ms before stimulus/retro-cue onset

and lasted until 1500 ms after stimulus/ retro-cue onset. For the S1/S2 epochs, only 1300

ms after stimulus onset were included in the analysis as those reflected the presentation

+ delay time. Those epochs were baseline-corrected (-200 ms to 0 ms), the previously

identified ICs reflecting cardiac and ocular activity were removed and the data resampled

at 300Hz.

<u>Statistical Analysis</u>

Behavioral Analysis

First, we excluded trials in which the response error was at least 3 SDs higher than the

subject's mean response error, or in which the response time exceeded 20 s, indicating

potential attentional lapses. We also excluded the first trial of each block, as serial

dependence cannot occur on those trials, and demeaned the response errors by

subtracting the overall mean response error of a participant from each individual response

error to remove general individual response biases independent of serial dependence.

The evaluation of serial dependence was based on individual response errors, defined as

the deviation between presented and entered direction. The errors were sorted regarding

the difference between the target stimulus of the current trial and the target stimulus from

the previous trial as well as the relation of difference between the current item and the

item of the previous trial (clockwise or counter-clockwise). The difference was computed

by subtracting the direction of the current item from the direction of the item of the

previous trial. Therefore, when the current item was oriented more clockwise or more

counter-clockwise, this resulted in a negatively or positively signed distance, respectively.

A mean response error for a signed distance (distance*relation) deviating from 0 indicated

a systematic response bias. When the sign of this systematic bias matched the sign of the

distance between the directions, it indicated an attractive response bias. Conversely, an

opposite sign of the systematic bias compared to the signed distance indicated a repulsive

response bias. In addition, the signed response errors were sorted according to the color

congruence between the two items (same vs different color). The individual mean

response biases were used to evaluate the serial dependence per color congruence level.

We fitted the first derivative of a Gaussian curve (DoG; e.g. (*1*), a model which is usually

used to describe serial dependence. The DoG, given by

$$y = xawce^{-(wx)^2}$$

(1)

was fitted to the pooled mean response biases of all subjects (similar to the procedure by

(*6*) per factor level, i.e. one data point per subject and distance for the respective factor

level. In the DoG, *x* is the relative direction difference of two stimuli, *a* is the amplitude of

the curve peak, *w* scales the curve width, and *c* is the constant $y\sqrt{2}/e^{-0.5}$. The w parameter

was constrained to a value range of .01 to .1. We optimized the log likelihoods of our curve

fitting using Bayesian adaptive direct search (BADS; (*28*)). BADS alternates between a series

of fast, local Bayesian optimization steps and a systematic, slower exploration of a mesh

grid. To estimate the variability of the parameters *a* and *w*, we bootstrapped the DoG curve

fit 1,000 times, sampling the data with replacement on each iteration, and computed the

standard deviation of the resulting bootstrapped distributions of *a* (see (*2*) for a similar

procedure). To assess the significance of serial dependence on a group level, we used

permutation tests, i.e. we randomly inverted the signs of each participant's mean response

error, fitted a new DoG model to the pooled group data and collected the resulting

amplitude parameters a in a permutation distribution. We repeated this permutation

procedure 1,000 times and reported the percentage of permutations that led to equal or

higher values for *a* than the one estimated for the empirical data as p-values. The

significance level was set to α = 0.05 (one-sided permutation test).

MEG Analysis

Inverted encoding model analysis. To reconstruct motion directions from neural signals,

we used the so-called inverted encoding model technique (*29*). We built an encoding

model characterizing motion direction-selective responses in the MEG signal with the

underlying assumption that the MEG signal in each sensor reflects a weighted sum of

neural activity revealed by different motion directions. Our model consisted of a basis set

of 18 channels spanning the room of 0-360° with centers in steps of 20°. The channels had

the shape of sinusoids raised to the 18th power (adapted from (*30*) and (*31*)). This model

was applied to the MEG data, which was separated in a training and a test data set. For

each subject, we trained the model on one MEG session and tested it on the other one in

two iterations, so that each MEG session was once training and once test data and training

and test data were independent. To analyze the currently relevant motion directions, we

applied this reconstruction approach to the three types of epochs (S1 epoch, S2 epoch

and post-cue epoch) separately and in each epoch the currently relevant direction was the

aim of the reconstruction (S1 epoch: S1 direction, S2 epoch: S2 direction, post-cue epoch:

cued target direction). To increase the signal-to-noise ratio, the epoched data were

averaged across time bins of 100 ms, centered on 50 ms, 150 ms, etc. after epoch onset.

This led to 13 analysis time points for the S1 and S2 epoch and 15 analysis time points for

the post-cue epoch. For each epoch, the directions were reconstructed for each time point

separately in a time point by time point fashion. To check for reactivations of the previous

target direction, we applied the same analysis on the last second of the ITI and the whole

time course of the current trial, resulting in 51 analysis time points. This time, the aim of

the reconstruction was the previous target direction, therefore the data were trained and

tested on this direction.

In the first step, we modeled the activation of each sensor as a weighted sum of filter

responses:

$$B_1 = C_1 W$$

(2)

B_1 is the MEG signal in each sensor at a given time point (n trials * n sensors), C_1 the

modeled response of each channel function give the presented direction in each trial (n

trials * n channels) and W a weight matrix representing the contribution of each channel

to the measured signal in each sensor (n channels * n sensors). W was estimated by using

ordinary least-squares regression:

718                              $$\widehat{W} = (C_1^T C_1)^{-1} C_1^T B_1$$
719   (3)

720   In the second step, we inverted W and applied to the independent test MEG data B_2 (n

721   sensors * n trials) to compute the estimated channel responses $\hat{C}_2$ (n channels * n trials):

722                              $$\hat{C}_2 = (\widehat{W}^T \widehat{W})^{-1} \widehat{W}^T B_2$$
723   (4)

The estimated channel responses $\hat{C}_2$ were transformed from the channel space back to the

direction space by weighting $\hat{C}_2$ with each channel's sensitivity from the basis set. Those

reconstructed channel responses were then circularly shifted to a common center (180°).

This whole reconstruction procedure was done separately per subject, session, epoch and

time point and this was iterated 20 times. Across those iterations, the centers of the basis

set were shifted by 1° so that the channel centers were evenly spaced in steps of 1° across

the whole direction space as a result of those iterations. The single trial reconstructions

were then analyzed in two different ways, described in the following paragraphs.

Shifts of the reconstructed direction. The main goal of our analysis was to examine

potential shifts of the neural representation in relation to the target item of the previous

trial. Therefore, the single-trial reconstructions were ordered in two groups, those with a

previous target that was clockwise (CW) oriented in relation to the currently relevant item

and those with a previous target that was counter-clockwise (CCW) oriented. The CCW

reconstructions were flipped along the direction space. Thereby, for both CW and CCW

trials, a negative deviation of the maximum of the reconstruction from 180° indicated an

attraction toward the previous target, whereas a positive deviation indicated a repulsion.

Those reconstruction were then first averaged within each possible motion direction and

then across them to account for different presentation numbers of the directions. This

averaged reconstruction was obtained for each iteration (see above) and then later

averaged across sessions. Thereby, the analysis resulted in a reconstruction per participant,

epoch and time point. In the next step, the fidelity of those reconstructions was assessed

per epoch and time point to identify time points with significant direction information in

the MEG signal. A fidelity above 0 indicates a meaningful direction signal and was obtained

by projecting the polar vector of the reconstruction at every direction angle (in steps of

1°) onto the common center (180°). The length of this projected vector indicates the

information strengths of the reconstruction at each angle. The fidelity F was obtained by

averaging those information strengths:

$$F = mean(r(\theta)\, cos\,(\theta)\,)$$

(5)

We identified the time points in each epoch with significant direction reconstruction

fidelity by computing a one-sided cluster-based permutation test (as implemented in

MNE) against zero. As the effect of the neural shift was expected to be very small, we

averaged the reconstructions in each epoch only over those time points that showed a

significant reconstruction fidelity and thus interpretable data. To examine systematic

shifts, we then tested if the maximum of the reconstruction was systematically different

from the common center (180°). For display purposes, we subtracted the reconstructed

maximum from 180° to compute the direction shifts. A positive shift thus reflected

attraction and a negative shift reflected repulsion.

Statistical inference of shifts. To assess the significance of potential shifts, we used a

bootstrapping procedure similar to (*32*). To this aim, we randomly selected reconstructions

from 10 participants (with replacement), then computed the mean reconstruction over the

resampled participants and collected the shift. This was done on 10,000 iterations. The

distribution of shifts was then used to examine if the observed shift was significantly larger

than zero. A shift was significant when 95% of the resampling iterations led to a shift equal

or higher than 0°, i.e., we conducted a one-sided bootstrapping test as we expected an

attraction, which was indicated by a shift larger than 0°. To obtain an indicator for the

variability of the shifts, the middle 95% of shifts resulting from the described resampling

procedure are reported.

Correlation between neural representation and behavioral response. In addition, we

correlated the maximum of single trial reconstructions with the upcoming behavioral

response to test whether shifts in behavior and in the neural signal on a single-trial level

were systematically related to each other. Both behavioral and neural shifts in any direction

(clockwise or counter-clockwise) in a single-trial response can originate from or be

modulated by different sources like, e.g., noise in the motor response, but they can also

result from attractive serial dependence that can be observed by averaging over trials. We

collected the reconstructions before flipping them with regard to the previous target and

removed trials where the behavioral response deviated more than three SDs from the

individual response error. Then we computed the circular correlation between the

deviation of the reconstructed maximum from the presented direction and the response

error for each participant, session, epoch and time point. For S1 and S2 epochs, only

epochs from trials where the respective item was cued as the response target were

included.

Statistical inference of correlations. The obtained correlation coefficients were averaged

across sessions and then time points for each epoch with a correlation significantly

differing from zero were identified by computing a one-sided cluster-based permutation

test (as implemented in MNE) against zero.

Reconstruction of previous target direction. We also examined whether the neural signals

of the current trial contained memory traces of the previous target direction. Therefore,

another IEM analysis was employed, but this time trained and tested on the target

direction of the previous trial throughout the last second of the ITI and the current trial.

Time points that contained significant direction information from the previous target were

identified by computing the fidelity of the reconstructions as described above and tested

against zero with a one-sided cluster-based permutation test.

Multiple comparison correction. In order to correct for multiple comparisons, we applied

an FDR correction (Benjamini-Hochberg) to those analyses that resulted in one p-value

per epoch, i.e., the bootstrapping tests of the shifts against zero, and reported these

corrected values alongside the original ones. Furthermore, for cluster-based permutation

tests that were applied to each epoch individually, we multiplied p-values higher than the

minimally possible (p<.002) for significant clusters by the number of epochs (3), thus

resulting in a Bonferroni correction.

<u>Software</u>

All behavioral analysis were performed with MATLAB 2019a (*33*) and the following

toolboxes/functions: Circular Statistics Toolbox (*34*), BADS (*28*) and EzyFit v2.44 (*35*). The

MEG analysis were performed with Python 3.8 in the Anaconda distribution v.2020.11 (*36*)

using spyder v.4.1.5 (*37*) and the following toolboxes: autoreject v0.2.2 (*38*), MNE v0.22.0

(*27*), numpy v1.19.2 (*39*), pandas v1.1.3 (*40, 41*) and scipy v1.5.2 (*42*).

References

1. J. Fischer, D. Whitney, Serial dependence in visual perception. *Nature Neuroscience*. **17**, 738–743 (2014).

2. C. Fischer, S. Czoschke, B. Peters, B. Rahm, J. Kaiser, C. Bledowski, Context information supports serial dependence of multiple visual objects across memory episodes. *Nat Commun*. **11**, 1932 (2020).

3. S. Czoschke, C. Fischer, J. Beitner, J. Kaiser, C. Bledowski, Two types of serial dependence in visual working memory. *British Journal of Psychology*. **110**, 256–267 (2019).

4. Y. Murai, D. Whitney, Serial dependence revealed in history-dependent perceptual templates. *Current Biology*. **31**, 3185-3191.e3 (2021).

5. D. P. Bliss, J. J. Sun, M. D'Esposito, Serial dependence is absent at the time of perception but increases in visual working memory. *Scientific Reports*. **7** (2017), doi:10.1038/s41598-017-15199-7.

6. M. Fritsche, P. Mostert, F. P. de Lange, Opposite Effects of Recent History on Perception and Decision. *Current Biology*. **27**, 590–595 (2017).

7. D. Pascucci, G. Mancuso, E. Santandrea, C. D. Libera, G. Plomp, L. Chelazzi, Laws of concatenated perception: Vision goes for novelty, decisions for perseverance. *PLOS Biology*. **17**, e3000144 (2019).

8. A. Kiyonaga, J. M. Scimeca, D. P. Bliss, D. Whitney, Serial Dependence across Perception, Attention, and Memory. *Trends in Cognitive Sciences*. **21**, 493–497 (2017).

9. D. Pascucci, Ö. D. Tanrikulu, A. Ozkirli, C. Houborg, G. Ceylan, P. Zerr, M. Rafiei, Á. Kristjánsson, Serial dependence in visual perception: A review. *Journal of Vision*. **23**, 9 (2023).

10. J. Barbosa, H. Stein, R. L. Martinez, A. Galan-Gadea, S. Li, J. Dalmau, K. C. S. Adam, J. Valls-Solé, C. Constantinidis, A. Compte, Interplay between persistent activity and activity-silent dynamics in the prefrontal cortex underlies serial biases in working memory. *Nat Neurosci*. **23**, 1016–1024 (2020).

11. A. Akrami, C. D. Kopec, M. E. Diamond, C. D. Brody, Posterior parietal cortex represents sensory history and mediates its effects on behaviour. *Nature*. **554**, 368–372 (2018).

12. E. St. John-Saaltink, P. Kok, H. C. Lau, F. P. de Lange, Serial Dependence in Perceptual
Decisions Is Reflected in Activity Patterns in Primary Visual Cortex. *Journal of*
Neuroscience. **36**, 6186–6192 (2016).

13. T. C. Sheehan, J. T. Serences, Attractive serial dependence overcomes repulsive
neuronal adaptation. *PLOS Biology*. **20**, e3001711 (2022).

14. J. E. Hajonides, F. Van Ede, M. G. Stokes, A. C. Nobre, N. E. Myers, Multiple and
Dissociable Effects of Sensory History on Working-Memory Performance. *J. Neurosci*.
43, 2730–2740 (2023).

15. M. Fritsche, E. Spaak, F. P. de Lange, A Bayesian and efficient observer model explains
concurrent attractive and repulsive history biases in visual perception. *eLife*. **9**, e55389
(2020).

16. G. J. Brouwer, D. J. Heeger, Cross-orientation suppression in human visual cortex.
Journal of Neurophysiology. **106**, 2108–2119 (2011).

17. G. J. Brouwer, D. J. Heeger, Decoding and Reconstructing Color from Responses in
Human Visual Cortex. *Journal of Neuroscience.* **29**, 13992–14003 (2009).

18. T. C. Sprague, J. T. Serences, Attention modulates spatial priority maps in the human
occipital, parietal and frontal cortices. *Nature Neuroscience*. **16**, 1879–1887 (2013).

19. M. Fornaciai, J. Park, Attractive Serial Dependence in the Absence of an Explicit Task.
Psychological Science, 095679761773738 (2018).

20. M. Fornaciai, J. Park, Neural Dynamics of Serial Dependence in Numerosity Perception.
J Cogn Neurosci. **32**, 141–154 (2020).

21. G.-Y. Bae, S. J. Luck, Reactivation of Previous Experiences in a Working Memory Task.
Psychol Sci. **30**, 587–595 (2019).

22. M. J. Wolff, J. Jochim, E. G. Akyürek, M. G. Stokes, Dynamic hidden states underlying
working-memory-guided behavior. *Nature Neuroscience*. **20**, 864–871 (2017).

23. M. J. Wolff, J. Ding, N. E. Myers, M. G. Stokes, Revealing hidden states in visual working
memory using electroencephalography. *Frontiers in Systems Neuroscience*. **9** (2015),
doi:10.3389/fnsys.2015.00123.

24. M. Manassi, A. Liberman, A. Kosovicheva, K. Zhang, D. Whitney, Serial dependence in
position occurs at the time of perception. *Psychonomic Bulletin & Review* (2018),
doi:10.3758/s13423-018-1454-5.

25. G. M. Cicchini, A. Benedetto, D. C. Burr, Perceptual history propagates down to early
levels of sensory analysis. *Current Biology*. **31**, 1245-1250.e2 (2021).

26. D. Whitney, M. Manassi, Y. Murai, Searching for serial dependencies in the brain. *PLoS*
Biol. **20**, e3001788 (2022).

27. A. Gramfort, M. Luessi, E. Larson, D. Engemann, D. Strohmeier, C. Brodbeck, R. Goj, M.
Jas, T. Brooks, L. Parkkonen, M. Hämäläinen, MEG and EEG data analysis with MNE-
Python. *Frontiers in Neuroscience*. **7** (2013) (available at
https://www.frontiersin.org/articles/10.3389/fnins.2013.00267).

28. L. Acerbi, W. J. Ma, "Practical Bayesian Optimization for Model Fitting with Bayesian
Adaptive Direct Search" in *Advances in Neural Information Processing Systems 30*, I.
Guyon, U. V. Luxburg, S. Bengio, H. Wallach, R. Fergus, S. Vishwanathan, R. Garnett,
Eds. (2017; http://papers.nips.cc/paper/6780-practical-bayesian-optimization-for-
model-fitting-with-bayesian-adaptive-direct-search.pdf), pp. 1836–1846.

29. T. C. Sprague, K. C. S. Adam, J. J. Foster, M. Rahmati, D. W. Sutterer, V. A. Vo. *Eneuro,*
5(3).

30. E. F. Ester, T. C. Sprague, J. T. Serences, Parietal and Frontal Cortex Encode Stimulus-
Specific Mnemonic Representations during Visual Working Memory. *Neuron*. **87**, 893–
905 (2015).

31. T. C. Sprague, IEM-tutorial. *GitHub repository* (2016), (available at
https://github.com/tommysprague/IEM-tutorial).

32. E. F. Ester, T. C. Sprague, J. T. Serences, Categorical biases in human occipitoparietal
cortex. *Journal of Neuroscience*. **40.4**, 917–931 (2020).

33. MATLAB, *R2019a* (The MathWorks Inc., Natick, Massachusetts, 2019).

34. P. Berens, CircStat : A MATLAB Toolbox for Circular Statistics. *J. Stat. Soft*. **31** (2009),
doi:10.18637/jss.v031.i10.

35. F. Moisy, EzyFit 2.44 (https://www.mathworks.com/matlabcentral/fileexchange/10176-
ezyfit-2-44), MATLAB Central File Exchange. Retrieved April 6, 2018. (2016).

36. Anaconda Software Distribution. *Anaconda Documentation* (2020), (available at
https://docs.anaconda.com/).

37. P. Raybaut, Spyder-documentation. *Available online at: pythonhosted. org* (2009).

38. M. Jas, D. A. Engemann, Y. Bekhti, F. Raimondo, A. Gramfort, Autoreject: Automated
artifact rejection for MEG and EEG data (2016), doi:10.48550/ARXIV.1612.08194.

39. C. R. Harris, K. J. Millman, S. J. van der Walt, R. Gommers, P. Virtanen, D. Cournapeau,
E. Wieser, J. Taylor, S. Berg, N. J. Smith, R. Kern, M. Picus, S. Hoyer, M. H. van Kerkwijk,
M. Brett, A. Haldane, J. F. del Río, M. Wiebe, P. Peterson, P. Gérard-Marchant, K.
Sheppard, T. Reddy, W. Weckesser, H. Abbasi, C. Gohlke, T. E. Oliphant, Array
programming with NumPy. *Nature*. **585**, 357–362 (2020).

40. W. McKinney, others, "Data structures for statistical computing in python" in
Proceedings of the 9th Python in Science Conference (Austin, TX, 2010), vol. 445, pp.
51–56.

41. T. pandas development team, pandas-dev/pandas: Pandas (2020), ,
doi:10.5281/zenodo.3509134.

42. P. Virtanen, R. Gommers, T. E. Oliphant, M. Haberland, T. Reddy, D. Cournapeau, E.
Burovski, P. Peterson, W. Weckesser, J. Bright, S. J. van der Walt, M. Brett, J. Wilson, K.
J. Millman, N. Mayorov, A. R. J. Nelson, E. Jones, R. Kern, E. Larson, C. J. Carey, İ. Polat,
Y. Feng, E. W. Moore, J. VanderPlas, D. Laxalde, J. Perktold, R. Cimrman, I. Henriksen, E.
A. Quintero, C. R. Harris, A. M. Archibald, A. H. Ribeiro, F. Pedregosa, P. van Mulbregt,
SciPy 1.0 Contributors, SciPy 1.0: Fundamental Algorithms for Scientific Computing in
Python. *Nature Methods*. **17**, 261–272 (2020).

Acknowledgments:

We thank Magdalena Feldmann, Nicole Huizinga, Philipp Deutsch and Susanna Wenzel

for their help in data collection and Benjamin Rahm for helpful discussions.

Funding:

PhD scholarship by the German Academic Scholarship Foundation awarded to CF

Author contributions:

Conceptualization: CF, JK, CB

Methodology: CF, JK, CB

Investigation: CF

Formal analysis: CF

Visualization: CF, CB

Supervision: JK, CB

Writing—original draft: CF, CB

Writing—review & editing: CF, JK, CB

Competing interests:

Authors declare that they have no competing interests.

Data and code availability:

All code and preprocessed data used for the presented analyses are available on [to be

inserted upon publication].

Supporting Information

Correlation between the neural shift and individual serial dependence parameters. In addition to the single trial-based correlation between neural and behavioral shifts reported in the main manuscript, we calculated correlations between the neural shift and individual a and w parameter obtained from DoG serial dependence fits on a between-subject level. This analysis revealed no significant correlations during the S1 and S2 epochs for neither a nor w parameter (a parameter, S1: r = .01, p = .9767; S2: r = .08, p = .8114; w parameter: S1: r = .36, p = .3096; S2: r = .30, p = .4054). While during the retro-cue epoch, there was no significant correlation with the a parameter either (r = -.35, p = .3258), we did observe a significant correlation between individual neural shifts and w parameters (r = -.70, p = .0246). This means that smaller individual w parameters indicating a broader tuning of serial dependence went along with larger neural shifts. This further demonstrates that the neural shift during the retro-cue epoch reflected behavioral serial dependence. These results are displayed in the **Figure S1**.

Figure S1. Relation between the individual a (upper panels) and w (lower panels) parameters of a DoG serial dependence fit (x-axis) and the neural reconstruction shift (y-axis). Colored circles show individual values for the S1 (left panels), S2 (middle panels) and retro-cue epochs (right panels). Solid lines depict significant correlations, dashed grey lines mark a reconstruction shift of 0°.

Cross-validation reveals different codes for reactivation of previous information and reconstructed biases of current information. In the retro-cue epoch, we observed both an attractive bias towards the target of the previous trial and a transient reactivation of the previous target itself. Therefore, one alternative explanation of the observed attractive bias could be a reactivation of the previous target in the same neural code as the current target that in sum lead to an attractive shift of the reconstruction. To rule out this explanation, we ran a cross-validation analysis. Therefore, we trained the encoding model on the currently relevant item (S1 during the S1 epoch, S2 during the S2 epoch and the cued item during the retro-cue epoch) and aimed to reconstruct the motion direction of the previous trial.

This analysis revealed that during the retro-cue epoch, where we observed the neural shift toward the past, as well as during the S1 epoch, there was no reactivation in the same neural code as the current representation. In contrast, a reactivation of target information from the previous trial using the same neural code could be observed during stimulus presentation in the S2 epoch. This indicated that during the S2 phase, information from the current S2 and the previous target were reconstructed in the same neural code. These results are displayed in the **Figure S2**.

Fig. S2. Cross-validated reconstruction fidelity of previous direction, model trained on currently relevant direction. Fidelity of reconstructed direction of the previous target using a model that was trained on the currently relevant direction. The left panel shows the fidelity of the previous target during the S1 epoch, the middle panel depicts the fidelity during the S2 epoch and the right panel shows the fidelity of the cued direction (either S1 or S2). Small circles show the fidelity of each participant. Colored circles indicate time points with a fidelity significantly greater than zero (cluster-based permutation test within each epoch).

Eye Movement control. Participants were instructed to maintain fixation on a small square at the center of the screen throughout each trial and independent component analysis (ICA) was used during preprocessing to remove any eye movement artifacts. However, recent studies showed that systematic eye movements can potentially confound decoding results (e.g. 1, 2). In the present study, participants viewed dynamic stimuli (moving dot patterns) and had to remember a spatial feature (motion direction), which made systematic eye movement behavior more likely. Therefore, we thoroughly tested a) whether eye movements were associated with the presented/remembered motion direction and if so, b) whether eye movements could have driven our MEG results.

As a first step of our eye movement control analysis, we tested how much of the variance of gaze direction could be explained by the stimulus direction that was either currently presented or had to be maintained at a given moment in time. More specifically, we first transformed the x and y positions measured by the eye tracker (in relation to a baseline from 200ms until trial onset) into polar coordinates to obtain a gaze direction in degrees. This transformation was performed on the same time points as the MEG analysis, i.e., gaze directions were averaged within bins of 100ms. Then we computed the circular variance of these gaze directions separately for each specific stimulus direction and time bin per subject. This circular variance was compared to a random distribution of gaze direction variance. To obtain this distribution, we shuffled the motion direction labels randomly over trials and then computed the circular variance for each pseudo direction, which was unrelated to the direction that was actually presented in the trial. The empirical and shuffled variances were averaged across directions and then a ratio was computed between them. The resulting value reflects the extent to which the variance of the stimulus-related gaze directions was reduced compared with random gaze direction patterns. This procedure was able to reveal not only systematic gaze directions that were in accordance with the stimulus direction or the opposite direction, but also picked up all stimulus-related gaze directions, even if the relation differed across participants or time. We identified time points with a significant amount of stimulus-related gaze directions in each analyzed epoch with cluster-based permutation tests. **Fig. S3a** shows that gaze directions were systematically related to the presented or remembered motion direction at different time points during each epoch. However, stimulus-related gaze directions differed considerably between participants as indicated by the standard errors.

In the second step, we plotted the time courses of stimulus-related gaze direction separately for each subject. This is shown in **Fig. S3b**. This revealed that while some participants showed a high amount of stimulus-related gaze directions, others showed very little or no stimulus-related gaze directions at all. In **Fig. S3b,** we marked the time courses in dark red for subjects with the highest amount of direction-related gaze positions ("high") and dark green for those with the lowest amount ("low"), with gradients in between.

In the third step, we addressed the question of how the stimulus-related gaze directions related to our MEG results. To this aim, we plotted the MEG-based reconstruction fidelity of presented and remembered motion direction separately for each subject and epoch and, crucially, coded them by the amount of stimulus-related gaze directions shown by each subject. This is shown in **Fig. S3c**. Comparing the stimulus reconstruction fidelity time courses with the individual color codes derived from the gaze-direction analyses revealed that the reconstruction fidelities of each participant did not correspond to their amount of stimulus-related gaze behavior (compare **Fig. S3b** with **Fig. S3c**). This suggests that there was no systematic relationship at any time point between the fidelity measure and the amount of direction-related gaze positions.

To support this observation statistically, we calculated Spearman rank correlations between the amount of stimulus-related gaze directions and reconstruction fidelities individually for each time point of each epoch. The resulting p-values are shown in **Fig. S3d**. Even before correcting for multiple comparisons, the p-values were consistently above 0.05, except for one single time point in the retro-cue epoch. Moreover, the p-values within each epoch did not show any consistent time profile but varied considerably between time points, supporting our interpretation that MEG-based reconstruction fidelity was not driven by the amount of stimulus-related gaze directions.

In consequence, we also did not find any relationship between the attractive neural shift that was based on these reconstructed MEG signals (i.e., the main finding of our study) and stimulus-related gaze directions. Specifically, there was no correlation between each subject's rank of the strength of neural shift towards the previous target and the subject's rank of the amount of stimulus-related gaze directions in any of the epochs (S1: p = .986, S2: p = .260, retro-cue: p = .244).

We conclude that our MEG results were unlikely to be explained by eye movements.

Fig. S3. Systematic eye movements and reconstruction time courses were uncorrelated. (**a**) Amount of stimulus-related gaze direction (indicated by variance reduction) during each epoch as a function of time with error bars indicating standard errors and colored circles indicating time points with a variance reduction significantly greater than zero (cluster-based permutation test within each epoch; p<.05). (**b**) Time courses of individual stimulus-related gaze directions. Dark red indicates the subject with the highest average variance reduction in a given epoch while green indicates the one with the smallest average variance reduction, with color gradients in between according to the individual rank of average variance reduction. (**c**) Time courses of individual reconstruction fidelities. Lines are colored as in (**b**), reflecting the stimulus-related gaze direction rank of a subject in a given epoch. (**d**) Time courses of p-values (uncorrected for multiple comparisons) of time point-

by-time point Spearman rank correlations between the systematic gaze directions as shown in (**b**) and the reconstruction fidelity as shown in (**c**).

SI References

1. Mostert, P., Albers, A. M., Brinkman, L., Todorova, L., Kok, P., & De Lange, F. P. Eye movement-related confounds in neural decoding of visual working memory representations. *ENeuro*, **5**(4) (2018).
2. Quax, S. C., Dijkstra, N., van Staveren, M. J., Bosch, S. E., & van Gerven, M. A. Eye movements explain decodability during perception and cued attention in MEG. *Neuroimage*, **195**, 444-453 (2019).