Title: Metabolic rewiring in fat-depleted *Drosophila* reveals triglyceride:glycogen crosstalk and

identifies cDIP as a new regulator of energy metabolism

Authors: Rupali Ugrankar-Banerjee[1], Son Tran[1], Jade Bowerman[3], Blessy Paul[1], Lauren G.

Zacharias[2], Thomas P. Mathews[2], Ralph J. DeBerardinis[2,4], W Mike Henne[1,*]

Affiliations:

[1]Department of Cell Biology, UT Southwestern Medical Center, Dallas, TX 75390

[2]Children's Research Institute (CRI), UT Southwestern Medical Center, Dallas, TX 75390

[3]Department of Molecular Medicine, Cornell University, Ithaca, NY 14853

[4]Howard Hughes Medical Institute, UT Southwestern Medical Center, Dallas, TX 75390

*for correspondence: mike.henne@utsouthwestern.edu

Abstract:

Tissues store excess nutrients as triglyceride (TG) or glycogen, but how these reserves are

sensed and communicate remains poorly understood. Here we identify molecular players

orchestrating this metabolic balance, and uncover a signaling molecule that mediates energy

homeostasis during fat depletion. We show tissue-specific depletion of fatty acyl-CoA synthase

FASN1 in the *Drosophila* fat body (FB) causes near complete fat loss and metabolic remodeling

that dramatically elevates carbohydrate metabolism, leading to a striking ~20-fold increase in

glycogen storage. Proteomics and metabolomics identify key factors necessary for rewiring

including glycolysis enzymes and target-of-brain-insulin (tobi). FASN1-deficient flies are viable

but starvation sensitive, oxidatively stressed, and infertile. We also identify CG10824/cDIP as

upregulated in FASN1-depleted *Drosophila*. cDIP is a leucine-rich-repeat protein with homology

to secreted adipokines that fine-tune energy signaling. We find FB-specific cDIP loss promotes

Drosophila adult obesity, and is required for development in the absence of FASN1-dependent

lipid storage. Collectively, we show fat-depleted *Drosophila* rewire their metabolism to complete

development, and identify cDIP as a putative new cytokine that signals fat insufficiency and may

regulate energy homeostasis.

Key words:

Fatty acid synthase (FASN); glycolysis; glycogen; triglyceride (TG); nicotinamide adenine

dinucleotide phosphate (NADPH)

Introduction:

Organisms must sense and allocate nutrients to metabolism, development, and reproduction to

maintain homeostasis and populations[1,2]. Dietary carbohydrates are mobilized to glucose and

catabolized through glycolysis, the tricarboxylic acid (TCA) cycle, and oxidative phosphorylation

to yield carbon dioxide and ATP, the cellular energy currency[3–5]. Excess glucose is utilized for

the *de novo* synthesis of fatty acids and triglycerides (TG) and incorporated into lipid droplets

(LD) in the adipose tissue, or condensed to glycogen, a polymeric form of glucose stored

preferentially in liver and muscle[3,4,6–8]. Perturbations in energy storage or utilization are linked to

metabolic disorders such as obesity, Type 2 diabetes, cardiovascular disease, and cancers[4,5,9–

11]. However, the metabolic decision-making that dictates how tissues invest in and balance fat

and glycogen storage, and how these energetic reserves are sensed and communicate to

ensure homeostasis and development, are not well understood[1].

Drosophila melanogaster is a powerful model to interrogate metabolic balance and human

metabolic diseases. Not only does *Drosophila* carry functional homologs to ~75% of human

disease genes, but key metabolic pathways such as insulin and TOR signaling are evolutionarily

conserved in flies[10,12,13]. Like in humans, *Drosophila* can become obese, lipodystrophic, or

diabetic in response to caloric cues or genetic mutations[14–16]. The primary nutrient storage

tissue in the fly is the fat body (FB) which shares features of the mammalian adipose tissue and

liver[17,18]. The embryonic FB is comprised of a continuous monolayer of ~2200 polygonal cells

whose numbers remain unchanged but increase in size during larval development[19–22]. In

feeding larvae, most dietary carbohydrates are converted to TG and assimilated into LDs within

the FB. Glycogen is also detected in the FB, albeit at significantly lower concentrations than

lipids. Critically, the role of glycogen in fly adipose tissue is not clear, however recent studies

indicate that glycogen may be important for maintaining hemolymph trehalose, the predominant

circulating sugar in *Drosophila*, present at ~100-times the level of glucose[12,16,23].

Commitment to pupariation and metamorphosis requires that feeding larvae reach a biomass

checkpoint called 'critical weight'. Much of this body mass is contributed by TG-rich LDs in the

FB[24–26]. During metamorphosis juvenile larvae are transformed into adult flies. This non-feeding

developmental stage is energetically expensive and requires large-scale tissue remodeling,

imaginal cell differentiation, and growth[27–29]. However, there is no consensus as to whether FB

TG is necessary for pupal development. For example, lipodystrophic *Drosophila* models

encoding mutations in fly homologs of *Lipin*, a phosphatidic phosphatase for making the TG

precursor DAG, and *Seipin*, required for normal LD biogenesis, still achieve critical mass

despite having smaller adipose tissue size and/or greatly reduced TG content. In fact, ~60% of

dLipin mutants survive to the late pharate adult stage, and ~40% became adults. Similarly, no

lethality is observed for *dSeipin* mutants and adults were viable and fertile[15,30,31]. Recent studies

instead point to the blood-circulating sugar trehalose as an essential energy source for

metamorphosis[25,32,33]. These observations suggest FB TG stores may not be critical for fly

development, but how they specifically contribute to metamorphosis and development, and

whether other reserves provide additional energy or biomass remains poorly understood[25,32,33].

TG in the FB is produced from two lipid pools: absorbed lipids from the blood by lipophorin

trafficking or via *de novo* lipogenesis that requires fatty acid synthase (FASN)[9,34]. FASN is

evolutionarily conserved and catalyzes the synthesis of long chain fatty acids from

carbohydrate-derived acetyl-CoA, malonyl CoA, and the electron donor nicotinamide adenine

dinucleotide phosphate (NADPH)[11,35]. *Drosophila* encode three *FASN* genes with distinct tissue

expressions. *FASN1/CG3523* is ubiquitously expressed with highest expression in the FB,

whereas *FASN2/CG3524* and *FASN3/CG17374* are expressed primarily in the body wall,

oenocytes, and hemocytes (FASN2), or oenocytes and reproductive system (FASN3)[4,28,36,37].

Drosophila hypomorphs for *FASN1 and FASN2* exhibit reduced larval and adult TG, while null

mutants of *FASN1* and *FASN3* display embryonic or early larval developmental arrest[9,28]. Fatty

acid synthesis is thus crucial for the generation of energy reserves, cellular membranes, and

protection from sugar toxicity[4,35]. However, how FASN-driven lipid synthesis is coordinated with

carbohydrate metabolism to enable homeostasis and development remains incompletely

understood.

Here we identify molecular players that orchestrate the metabolic balance between lipid and

carbohydrate utilization, and dissect how TG and glycogen stores contribute to *Drosophila*

development. We find that most TG in the larval FB is generated via FASN1-mediated *de novo*

lipogenesis, but surprisingly both FASN1 and its TG products in the FB are dispensable for

larval development and metamorphosis. FB-specific *FASN1* RNAi knockdown *(*defined here as

FASN1^{FB-RNAi}) leads to fully developed larvae that lack >85% of TG lipid biomass, but

remarkably still complete their developmental cycle. We find this is achieved through metabolic

re-wiring that drastically elevates FB glycogen storage (with an excess of ~20-fold increase in

glycogen compared to controls) and carbohydrate metabolism through glycolysis and trehalose

export. *FASN1^{FB-RNAi}* adults are viable but lack detectable TG stores, and females are infertile

due to defects in ovarian differentiation and oocyte development. FASN1-depleted adults are

also sensitive to nutrient deprivation, exhibit shortened median lifespan, and larvae display

signatures of redox imbalance including lipid peroxidation. Through proteomics we uncover a

putative secreted cytokine, Common Dpr-interacting protein (cDIP)/CG10824 that is required for

larval development when fat is depleted. cDIP is highly upregulated in *FASN1*[FB-RNAi] FBs and

shows homology to secreted leucine-rich repeat proteins that regulate vertebrate insulin/IGF-like

signaling. In line with this, we find cDIP FB-specific knockdown elevates fat storage, and is

required for development when fat body lipid stores are depleted. Collectively, we hypothesize

that *FASN1*[FB-RNAi] *Drosophila* signal fat deficiency by secreting cDIP to coordinate fat-to-

carbohydrate metabolic homeostasis, ensuring metabolic adaptation and animal development.

Results:

Fat body specific-loss of FASN1 leads to triglyceride-depleted but viable *Drosophila*

The fat body (FB) is the central metabolic organ in *Drosophila,* and represents much of the

larval biomass necessary to achieve the critical weight for metamorphosis[18,24–26]. Most of the FB

lipid mass is composed of lipid droplet (LD) triglycerides (TGs) produced from *de novo*

lipogenesis in the FB[9,34,36]. To genetically manipulate these fat stores, we utilized the FB tissue-

specific driver *Dcg-Gal4*[16,38] to selectively RNAi-deplete the major *de novo* fatty acyl-CoA

synthase *FASN1*, which we previously identified as the major FASN isoform in FB TG

synthesis[34] (**Fig 1A**). FB-specific *FASN1* knockdown (*Dcg-Gal4>FASN1*[RNAi], denoted *FASN1*[FB-

RNAi]) led to visible reduction of larval fat deposits, still *FASN1*[FB-RNAi] larvae developed in tandem

with age-matched controls (*Dcg-Gal4*) and grew to full size. However, the adipose tissue in L3

pre-wandering larvae (~100 to 108 h AEL, After Egg Laying) appeared translucent instead of

the opaque white color typical of fat-engorged tissue (**Fig 1B, red arrows**). In line with this,

isolated FBs from *FASN1*[FB-RNAi] larvae now sank in PBS buffer, indicating they lost the buoyancy

provided by their fat stores (**Fig 1C, white arrows**). Of note, the amount of FB tissue itself was

not reduced and contained normal sized adipocytes, despite general loss of fat content,

indicating tissue development was not altered by FASN1 loss (**SFig 1A,B**). Perturbations in FB

fat storage is known to cause ectopic accumulation of lipids in other tissues such as the mid-

gut[39], however, gut TG levels were significantly reduced in *FASN1$^{FB-RNAi}$* larvae relative to

controls (**SFig 1C**).

To determine how FASN1 depletion impacted LDs in the FB, we conducted z-section confocal

imaging. *FASN1$^{FB-RNAi}$* larvae displayed a near complete lack of large LDs in the tissue mid-

plane (mid-plane LDs, mLDs) within FB cells. However, *FASN1$^{FB-RNAi}$* larvae still exhibited small

peripheral LDs (pLDs) near the tissue edges exposed to the animal hemolymph (**Fig 1D,E**). This

is consistent with previous work indicating these pLDs are not dependent on *de novo*

lipogenesis within the FB, but rather potentially derived from dietary lipids via lipophorin-

mediated lipid trafficking from other tissues[34]. pLDs near the cell surface were also observed by

transmission electron microscopy (TEM), whereas most of the *FASN1$^{FB-RNAi}$* FB cell cytoplasm

looked vacant, except for small "islands" of organelles (**Fig 1F**). As expected, total larval TG

was significantly impacted as FB-specific FASN1-depletion led to loss of ≥ 85% larval TGs

compared to control larvae (**Fig 1G**). To determine whether LD depletion in *FASN1$^{FB-RNAi}$* FB

cells occurred in a cell autonomous manner, we also knocked down *FASN1* using the

ubiquitous but mosaic driver *da-Gal4*, which selectively induces RNAi in a subset of FB cells.

Indeed, *da-Gal4>FASN1^{RNAi}* larval FB tissues exhibited a mosaic pattern of LD-depleted FB

cells (also positive for PM marker mcd8-GFP) adjacent to unaffected cells (no mcd8-GFP) with

typical intracellular LD populations (**SFig 1D**). Importantly, LD-depleted cells still contained

pLDs near the tissue surface, but lacked the larger mLDs near the tissue mid-plane, consistent

with FASN1 loss primarily impacting the large LD population. Collectively, this indicates that

FASN1 loss in FB cells causes cell autonomous blockage of *de novo* lipogenesis and abolishes

TG storage in mLDs.

Despite their dramatic reduction of TG, FASN1[FB-RNAi] larvae surprisingly pupated with no notable

developmental delay, completing metamorphosis and emerging as normal sized and apparently

healthy flies (**Fig 1H**). Collectively, this indicates that FASN1 is not necessary in the FB for

animal development, and that *Drosophila* larvae can reach critical weight and successfully

complete development despite the loss of >85% of their FB TG stores.

FASN1 is the primary FASN isoform for TG storage in the FB

Drosophila FB lipids are proposed to play important roles in animal development including as an

energy source for metamorphosis, biomaterials for tissue development, as well as a metabolic

buffer to counter caloric overload[10,26,28]. Since TG-depleted *FASN1[FB-RNAi]* FBs still supported fly

development, we investigated what factors and pathways were necessary for metamorphosis

into adults in the absence of FASN1 in the FB. We conducted unbiased global proteomics of

isolated FBs from control and *FASN1[FB-RNAi]* pre-wandering L3 larva using liquid-chromatography

coupled with tandem mass spectrometry (LC-MS/MS). Principal component analysis (PCA)

revealed control and *FASN1[FB-RNAi]* FBs exhibited significant differences in the relative

abundances of many proteins (**SFig 2A**). Proteomics also confirmed efficient knockdown of

FASN1 as it was among the most decreased of all detected proteins in its abundance in

FASN1[FB-RNAi] larval FBs (**Fig 2A**). Similarly, we noted significant decreases in the abundance of

other proteins important for TG synthesis or storage, such as the glycerol kinase Gk1 (important

for making glycerol-3-phosphate and TG), and the perilipins Lsd1 and Lsd2, as well as proteins

involved in TG breakdown such as Bmm (mammalian ATGL homolog), and pdgy and CPT2

(involved in fatty acid β-oxidation) (**Fig 2A**). GO term analysis revealed that lipid metabolic

processes, particularly those related to fatty acid and TG biosynthesis, as well as fatty acid

catabolism and oxidation were among the most significantly decreased pathways in *FASN1[FB-*

RNAi] FBs (**Fig 2B**). Notably, FASN2 protein levels were moderately increased in *FASN1[FB-RNAi]*

FBs (~3-fold), but FASN3 was undetected in both control and *FASN1[FB-RNAi]* FBs likely due to low

expression in this tissue (**Fig 2A**). Quantitative mRNA measurements by qPCR of extracted FBs

further confirmed successful *FASN1* targeting by RNAi, and mild upregulation of *FASN2*

transcripts in FASN1-depleted fat tissue (**SFig 2B**). This suggests that in the absence of

FASN1, the FB still expresses FASN2 and can produce fatty acyl-CoAs, likely to supply lipids

for membrane and organelle biosynthesis. In line with this, GO analysis indicated that

phospholipid biosynthesis and metabolism pathways were upregulated in *FASN1^FB-RNAi^* FBs

relative to controls (**SFig 2C**).

To determine whether loss of FASN2 or FASN3 would similarly deplete TG FB stores like

FASN1 loss, we knocked down *FASN3* in the FB using a *UAS-RNAi TRiP* line. Since a *UAS-*

RNAi line for *FASN2* was unavailable, we instead examined the FB of an established *FASN2*

mutant. Indeed, global loss FASN2 only mildly altered LD storage in the FB relative to *FASN1^FB-*

^RNAi, while *FASN3^FB-RNAi^* had no obvious LD defects in the FB and resembled control tissue

(**SFig 2D**). This implies that FASN1 may generate fatty acyl CoAs that are channeled primarily

into LD-TG in the larval FB.

FASN1-depleted FBs exhibit elevated glycogen storage and require target-of-brain-

insulin (tobi) for development

FB proteomics revealed that many proteins were significantly elevated in *FASN1^FB-RNAi^*

compared to controls, particularly those involved in carbohydrate metabolism. GO term analysis

indicated that carbohydrate catabolic processes, and glycolytic processes were among the top

scoring pathways increased in *FASN1^FB-RNAi^* FBs (**Fig 2C**). Among these up-regulated

carbohydrate-related proteins was target-of-brain-insulin (tobi), which was only detected in the

proteomics of *FASN1^FB-RNAi^* FBs (**Fig 2A**), and whose mRNA level increased ~8-fold in

*FASN1^FB-RNAi^* FBs compared to controls (**Fig 2D**). The *tobi* gene encodes a conserved alpha-

glucosidase expressed primarily in FB and gut, and is activated during sugar mobilization in the

FB[40,41]. Since alpha-glucosidases mediate glycogen breakdown to glucose, we examined how

FB glycogen levels compared between control and *FASN1[FB-RNAi]* FBs. Strikingly, periodic acid-

Schiff (PAS) staining revealed significantly concentrated glycogen deposits in *FASN1[FB-RNAi]* FBs

compared to controls, and biochemical measurements confirmed a remarkable >20-fold

glycogen increase in *FASN1[FB-RNAi]* FBs (**Fig 2E,F**). Increased glycogen granules were also

detected in *FASN1[FB-RNAi]* FB by imaging *Glycogenin-YFP,* as glycogenin functions as a primer

for glycogen polymerization[42] (**SFig 2E**). QPCR analyses showed elevated expression of other

key enzymes involved in glycogen synthesis (glycogenesis) and glycogen breakdown

(glycogenolysis) (**SFig 2F**). This suggested that, in the absence of TG stores, larvae may shunt

dietary nutrients into glycogen for energy storage and utilization in the FB. From this, we

hypothesized that *FASN1[FB-RNAi]* larvae relied on carbohydrate metabolism for development.

To test this, we RNAi depleted the glycogen synthesis enzyme GlyS, or the glycogen

mobilization enzyme tobi, either alone or in combination with FASN1 specifically in the larval FB.

Whereas RNAi loss of GlyS or tobi on their own did not impact larval development into flies,

depletion of either in combination with *FASN1[FB-RNAi]* led to developmental arrest during the

pupal stage (**Fig 2G,H**). Next, we determined whether FASN1-depleted larvae utilized glycogen

during pupal development to complete metamorphosis. We monitored TG and glycogen levels

through distinct developmental stages of larval growth, pupation, and fly emergence. Indeed, we

observed that during the transition from the pupal to pharate adult (late pupal) stage, glycogen

levels in *FASN1[FB-RNAi]* animals (but not controls) declined rapidly (**Fig 2I**). In fact, newly hatched

FASN1[FB-RNAi] adults contained less than 10% of their initial glycogen stores, consistent with

glycogen being heavily utilized to support fly development. In contrast, glycogen amounts in

controls remained steady through metamorphosis and into adulthood (**Fig 2I**). In contrast,

control animals appeared to mobilize FB TG stores during development, particularly during the

pupal period when major tissue remodeling and pharate adult development occurs[25] (**Fig 2J**). A

small decrease in TG levels of *FASN1*[FB-RNAi] was also observed, however, the energy available

through TG lipolysis in this fat-deficient animal is likely negligible (**Fig 2J**). Collectively, this

indicates that TG-depleted *FASN1*[FB-RNAi] larvae exhibit significantly elevated carbohydrate

storage in the form of FB glycogen, and functionally require glycogen synthesis and breakdown

for development into flies.

FASN1[FB-RNAi] larvae require glycolysis for development

Since *FASN1*[FB-RNAi] larvae elevate glycogen stores, which upon mobilization release glucose,

we queried whether they exhibited enhanced glucose catabolism, beginning with glycolysis. The

first step of glycolysis produces glucose-6-phosphate which can be channeled into glycogenesis

and the pentose phosphate pathway (PPP), or broken down into pyruvate via a multi-step

pathway driven by various enzymes[5,7] (**Fig 3A**). Proteomics profiling of *FASN1*[FB-RNAi] larval FBs

revealed that enzymes involved in glycolysis including Hex-A, Pgi, Pfk, Gapdh2, Eno, and Pyk

(pyruvate kinase) were significantly more abundant compared to controls (**Fig 3A,B**). Transcript

analysis also showed increased mRNA levels for Pyk which produces the end product pyruvate

(**SFig 3A**). To confirm that pyruvate itself was elevated in *FASN1*[FB-RNAi] larval FBs, we

expressed the fluorescent pyruvate biosensor *UAS-PyronicSF-GFP* in the FB using the *Dcg-*

Gal4 driver[43]. Indeed, PyronicSF-GFP fluorescence was substantially elevated in *FASN1*[FB-RNAi]

FBs compared to controls, indicating pyruvate accumulation (**Fig 3C**). Next, we queried whether

Pyk was required for *Drosophila* development in *FASN1*[FB-RNAi] animals by RNAi-depleting PyK

either alone or in combination with FASN1 in the FB. Strikingly, whereas PyK loss alone did not

impact *Drosophila* development into flies, FASN1 and PyK dual loss led to ~75% pupal arrest

and subsequent death (**Fig 3D**).

Since sustained glycolysis may also require elevated gluconeogenesis (in addition to

glycogenolysis), we also queried whether key gluconeogenic enzymes were necessary for fly

development when FASN1 was depleted. We knocked down Pepck (phosphoenolpyruvate

carboxykinase), which catalyzes the rate limiting step of gluconeogenesis, and G6P (glucose-6-

phosphatase), which converts glucose-6-P to terminal glucose. Indeed, development was not

impacted when either was targeted alone, but concurrent loss of Pepck2 or G6P with FASN1 in

the FB led to ~87% and ~95% pupal lethality, respectively (**Fig 3E**). Collectively, this indicates

that glycolysis is essential for *FASN1*$^{FB-RNAi}$ FB metabolism in animal development.

Pyruvate can be converted into acetyl-CoA and catabolized by the mitochondrial tricarboxylic

acid (TCA) cycle and electron transport chain (ETC) to generate ATP energy. During hypoxia,

pyruvate is converted to lactate in the cytoplasm and ATP is produced through anaerobic

glycolysis only, a less efficient process. This pathway is often upregulated in cancer cells, even

in the presence of oxygen, in the so-called Warburg effect[2,44,45]. To determine whether

Drosophila lacking FB FASN1 relied on anaerobic glycolysis or aerobic respiration for ATP, we

first measured mRNA levels of enzymes that catalyze production of lactate and acetyl-CoA. We

found that transcripts of lactate dehydrogenase (Ldh), which converts pyruvate to lactate, were

not elevated with FASN1 loss (**SFig 3A**). To functionally access if lactate production was

necessary to fuel *FASN1*$^{FB-RNAi}$ development, we knocked down *Ldh* in the FB either alone or in

combination with *FASN1*. Indeed, in both cases *Drosophila* development into flies was not

substantially impacted, indicating lactate production was not functionally required for

development even in the absence of TG stores (**SFig 3B**). Collectively, this indicates that

FASN1$^{FB-RNAi}$ larvae accumulate glycolytic enzymes and may functionally require glucose

catabolism for development. Pyruvate levels are elevated with FASN1 depletion, but lactate

metabolism is not necessary.

Proteomic and metabolic profiling indicate FASN1-KD larvae functionally down-regulate

mitochondrial TCA, OXPHOS, and FAO metabolism

Acetyl-CoA derived from pyruvate can be partitioned to *de novo* lipogenesis or to the TCA cycle,

which generates metabolic intermediates for energy[4,5]. However, in the absence of the key fatty

acid synthesis enzyme *FASN1*, acetyl-CoA should be channeled into the TCA cycle, resulting in

increased TCA activity. Surprisingly, proteomics profiling indicated that most TCA enzymes

were slightly *decreased* in abundance in *FASN1[FB-RNAi]* larval FBs compared to controls (**Fig 3F**).

In contrast, ATP citrate lyase (ATPCL), which converts the first TCA cycle metabolite, citrate,

into acetyl-CoA and oxaloacetate, was markedly elevated in *FASN1[FB-RNAi]* FBs (**Fig 3F**). ATPCL

represents a cataplerosis enzyme enabling TCA intermediates to leave the cycle[43] (**Fig 3A**). In

line with this, proteomics indicated decreased levels of mitochondrial ETC proteins involved in

oxidative phosphorylation (OXPHOS) in *FASN1[FB-RNAi]* FBs compared to controls, suggesting

mitochondrial ETC metabolism may be dampened in FASN1-depleted FBs (**SFig 3C**).

These observations suggested that, in contrast to glycolysis, TCA and ETC pathways were not

elevated in *FASN1[FB-RNAi]* FBs. We tested how depletion of enzymes that channel carbon into the

TCA cycle impacted *Drosophila* development. First, we RNAi depleted MPC1 that transports

pyruvate from the cytosol to mitochondria (the site of TCA metabolism), as well as Pdha, which

generates acetyl-CoA, the entry substrate of the TCA cycle. FB-specific MPC1 depletion in

FASN1[FB-RNAi] did result in ~50% reduction in fly development, but this appeared less severe

than perturbing glycolysis (**SFig 3D**). Furthermore, FB-specific depletion of Pdha in combination

with *FASN1[FB-RNAi]* only mildly impacted larval development into flies (**SFig 3E**). We also

monitored TCA cycle intermediate metabolites using LC-MS/MS metabolomics. TCA

metabolites including malate, fumarate, citrate/isocitrate, succinate, and alpha-ketoglutarate

were elevated in FASN1-depleted FBs, suggesting some alteration of the TCA cycle (**Fig 3G**).

Collectively, this supports a model where *FASN1[FB-RNAi]* animals primarily rely on glycolysis to

fuel their developmental life cycle, and may be less reliant on pyruvate oxidation.

Since tissues also generate energy through fatty acid oxidation (FAO)[46,47], we monitored

mitochondrial and peroxisomal FAO enzyme levels by proteomics. Mitochondrial and

peroxisomal FAO enzymes were both significantly decreased in $FASN1^{FB-RNAi}$ FBs, suggesting

FAO may be blunted with FASN1 loss (**SFig 3F**). To evaluate whether mitochondrial FAO was

suppressed in FASN1-depleted FBs, we monitored levels of acylcarnitines and other

bioenergetic metabolites via LC-MS/MS metabolomics. Indeed, $FASN1^{FB-RNAi}$ FBs exhibited

reduced levels of several long-chain acylcarnitines compared to control FBs, suggesting

potentially decreased lipid channeling into FAO[48] (**Fig 3G**). In contrast, $FASN1^{FB-RNAi}$ FBs

displayed elevated intermediate and end metabolites in glycolysis such as fructose-6-

phosphate, dihydroxyacetone phosphate, and acetyl-CoA, consistent with elevated glycolysis.

We also noted elevated metabolites of the pentose phosphate pathway (PPP), likely due to

elevated carbohydrate metabolism in FASN1-depleted FBs (**Fig 3A,G**).

Finally, we interrogated mitochondrial morphology in L3 larval FBs by using fluorescence

confocal microscopy. Whereas control FBs displayed numerous tubular mitochondria

throughout the cytoplasm, $FASN1^{FB-RNAi}$ FBs displayed fewer and fragmented mitochondria (**Fig**

3H). As fragmented mitochondria can be correlated with reduced mitochondria metabolic

activity, this supports a model where loss of FASN1 in the FB reduces mitochondrial

energetics[49]. Collectively this indicates that $FASN1^{FB-RNAi}$ larval FBs require glycolysis but may

remodel or blunt mitochondrial metabolism during development.

FASN1[FB-RNAi] flies are short-lived, starvation sensitive, and require dietary sugar and

trehalose for development

Given that FB depletion of FASN1 led to metabolic remodeling that reduced mitochondrial

bioenergetics and elevated glycogen storage and glycolysis, we next investigated how this

influenced organismal lifespan and stress responses. We first examined how $FASN1^{FB-RNAi}$ male

and female adult lifespans compared to controls, discovering that $FASN1^{FB-RNAi}$ flies of both

sexes displayed significantly shortened median lifespan on standard food (**Fig 4A**). We also

found $FASN1^{FB-RNAi}$ flies were starvation sensitive when maintained on water but no food

source, and controls survived approximately twice as long under these fasting conditions (**Fig**

4B). This was consistent with biochemical analysis indicating that $FASN1^{FB-RNAi}$ adults displayed

significantly lower TG and glycogen levels than control animals, and again indicates that the

elevated glycogen in FASN1-depleted larvae is consumed during metamorphosis (**Fig 4C,D**).

Since $FASN1^{FB-RNAi}$ larvae exhibited elevated glycolysis and glycogen stores, we monitored

whether they required dietary sugar for development. We created a minimal defined food by

mixing agar and yeast powder supplemented with varying amounts of sucrose (0% = no sugar

or 0S, 5% = standard sugar or 5S). We measured TG and glycogen of control and $FASN1^{FB-RNAi}$

larvae cultured on 0S and 5S food. In control animals, TG stores were boosted with the

consumption of 5S versus 0S, but FB glycogen remained unchanged between larvae eating

these diets (**Fig 4E,F**). Conversely, in FASN1-deficient larvae TG levels were similar between

larvae cultured on 0S or 5S food (and were ~80% lower than controls on 5S food), consistent

with a blockage of TG synthesis even in the presence of additional dietary sugars. Furthermore,

FBs of 0S-fed $FASN1^{FB-RNAi}$ larvae stored ~65% less glycogen than those fed 5S food, indicating

their glycogen stores were largely maintained by dietary sugars (**Fig 4F**). Though FB glycogen

in $FASN1^{FB-RNAi}$ larvae fed 0S food were ~4-fold elevated relative to controls on 0S, successful

development of $FASN1^{FB-RNAi}$ larvae to adults was significantly diminished on both 0S or 5S food

compared to controls (**Fig 4G**). Collectively, this indicates that $FASN1^{FB-RNAi}$ *Drosophila* rely on

dietary sugars to store the nutrient reserves for development into flies, and in the absence of

sufficient dietary sugar fail to store sufficient nutrients to complete metamorphosis**.**

Since *Drosophila* energy homeostasis is influenced by sugars in the animal hemolymph, we

also examined how the disruption of dietary carbohydrate-to-fat conversion and enhanced

glycogen storage impacted circulating glucose and trehalose. We measured these in control

and *FASN1^FB-RNAi* pre-wandering L3 larvae. Notably, glucose levels were not altered but

trehalose levels were substantially elevated in *FASN1^FB-RNAi* larvae, suggesting that glycogen in

FASN1^FB-RNAi FBs increased trehalose concentrations in circulation to aid animal development

(**Fig 4H,I**). To test this, we RNAi depleted the trehalose synthesis enzyme Tps1 in the FBs of

control and *FASN1^FB-RNAi* larvae. Strikingly, whereas Tps1 loss alone in the FB did not impact

development, dual depletion of Tps1 and FASN1 led to significant developmental arrest (**SFig**

4A). Collectively, this indicates that metabolic re-wiring in *FASN1^FB-RNAi* *Drosophila* creates a

dietary sugar dependency, and results in shortened lifespan and starvation sensitivity. It also

reveals that *FASN1^FB-RNAi* larvae require glycogen reserves and trehalose to support

development.

FB-specific FASN1 loss causes female infertility and reproductive arrest

As animal nutrient reserves are tightly linked with reproductive success[50,51], we next examined

how loss of FASN1 in the FB impacted fecundity by monitoring whether *FASN1^FB-RNAi* female

and male flies could successfully generate progeny. Indeed, eggs deposited by control (*Dcg-*

Gal4) females mated with *FASN1^FB-RNAi* males showed slightly decreased viability relative to

those from *Dcg-Gal4* fathers, suggesting loss of FASN1 in the FB of males mildly perturbed

their fecundity (**Fig 5A**). Strikingly, *FASN1^FB-RNAi* females mated with either control or *FASN1^FB-*

^RNAi males failed to lay eggs, indicating *FASN1^FB-RNAi* females exhibited substantial fertility

defects (**Fig 5A**).

The adult *Drosophila* female reproductive system consists of two ovaries, each with 20-30

ovarioles. An ovariole is organized into a procession of egg chambers (**Fig 5B**). Early egg

chambers are made of 16 sister cells enclosed by a monolayer of somatic follicle cells. A single

nurse cell is fated to become an oocyte, while the remaining 15 nurse cells supply the growing

oocyte with nutrients, proteins, and RNA. Oocyte maturation within the egg chamber occurs in

14 morphologically distinct stages[52,53]. Dissection of the internal reproductive organs of

FASN1[FB-RNAi] females revealed shrunken and underdeveloped ovaries which did not produce

mature oocytes (**Fig 5B,C, white arrows**). Upon close examination, *FASN1*[FB-RNAi] egg

chambers appeared to arrest at stage 9-10, when normally the follicle cells migrate to surround

the oocyte (**Fig 5D, follicle cells denoted by yellow arrows**). This also represents a critical

oogenesis checkpoint when extracellular diacylglycerol (DG), sourced primarily from the FB, are

absorbed by nurse cells via a lipophorin receptor (LpR2) dependent pathway. DG is acylated to

TG and assimilated in LDs, which accumulate in nurse cells and are transported through

connecting ring canals to the developing oocyte to drive its maturation and later fuel

embryogenesis[52–55]. In contrast to controls, *FASN1*[FB-RNAi] ovaries were largely devoid of LDs

(**Fig 5E, orange arrows**). This suggests they may have failed to clear this key metabolic

checkpoint, resulting in degeneration of the egg chambers as no discernable cells with nuclear

DAPI staining were visible in the interior of *FASN1*[FB-RNAi] ovaries (**Fig 5F, blue arrows).**

Collectively, we conclude that FB-specific FASN1 loss leads to female *Drosophila* infertility due

to defects in lipid delivery to ovaries for oogenesis.

FASN1-depleted FBs exhibit elevated oxidative stress signatures and rely on

antioxidants for development

Our metabolic and proteomic profiling revealed that in addition to increased glycolysis,

FASN1[FB-RNAi] FBs elevated pentose phosphate pathway (PPP) metabolism, which generates the

redox metabolite nicotinamide adenine dinucleotide (NADPH) as a bi-product. NADPH is an

electron donor and key metabolite in redox homeostasis[5,23]. Importantly, NADPH is required for

FASN-dependent *de novo* lipogenesis, and inhibiting FASN activity can elevate cellular NADPH

pools which perturb redox balance and drive oxidative stress through NAPDH Oxidase (NOX)-

mediated reactive oxygen species (ROS) generation[47,56]. For example, cancer cells typically

elevate FASN-dependent lipogenesis to support anabolic metabolism and cell growth.

Pharmacological FASN inhibition is an anti-cancer strategy that can drive NADPH accumulation

in cancer cells, promoting ROS signaling and cell death via NOX activity[4,57].

Given that FASN1 FB-depleted animals displayed shortened lifespans, we investigated whether

they exhibited signatures of altered redox homeostasis. Indeed, metabolomic profiling indicated

FASN1[FB-RNAi] FBs contained elevated concentrations of oxidized species of several redox

related metabolites including NADP+, NAD+, cystine, and glutathione disulfide (GSSG), the

oxidized form of glutathione (GSH)[24,58,59] (**Fig 6A**). In line with this, GSSG:GSH ratios were

significantly higher in *FASN1*[FB-RNAi] larvae compared to controls (**Fig 6B**). Proteomics also

indicated *FASN1*[FB-RNAi] FBs had elevated glutathione S-transferases (GST) and related

enzymes, suggesting FASN1 loss elevated oxidative toxins that required GST activity (**Fig 6C**).

Given that *FASN1*[FB-RNAi] larvae displayed signatures of oxidative stress, we interrogated whether

depletion of major cellular antioxidants negatively impacted larva-to-adult development. We

targeted three major antioxidants in the FB: superoxide dismutase 1 (SOD1), SOD2, and

catalase. We RNAi depleted these factors either alone or in combination with FASN1

specifically in the FB using the *Dcg-Gal4* driver. Notably, whereas RNAi-depletion of SOD1 or

SOD2 alone had no impact on fly development, their depletion together with FASN1 led to a ~

29% and ~70% reduction in fly eclosion, respectively (**Fig 6D, SFig 6A**). Similarly, RNAi

depletion of peroxisome-localized catalase did not affect *Drosophila* development, but its co-

depletion with FASN1 led to ~75% reduction in adult development (**Fig 6D**). In line with this,

FASN1[FB-RNAi] larval FBs displayed significantly more peroxisomes compared to controls,

suggesting peroxisomes may be elevated in FASN1-depleted FBs to support homeostasis

(**SFig 6B**). In support of this, RNAi depletion of peroxisomal biogenesis factor Pex5 in

conjunction with FASN1 reduced larva-to-fly development (**SFig 6C**). Collectively, this indicates

that FASN1 depleted FBs display signatures of defective redox homeostasis and functionally

require antioxidant enzymes and peroxisomes for developmental integrity.

cDIP is required for development of FASN1-depleted larvae and influences FB energy

and redox homeostasis

Given that *FASN1^FB-RNAi^ Drosophila* are fat-deficient, reliant on carbohydrate metabolism for

development, and displayed signatures of oxidative stress, we reasoned they may elevate the

expression of signaling molecules that influenced energy homeostasis of the FB. To potentially

uncover such new signaling molecules, we examined our FB proteomics for uncharacterized

proteins with signatures of signaling functions (**Fig 2A**). We noted that among the most up-

regulated proteins (~70-fold) in the *FASN1^FB-RNAi^* proteomics was Common Dpr-interacting

Protein (cDIP)/CG10824, a poorly characterized leucine rich-repeat (LRR) protein. Panther and

Phylome protein phylogeny databases indicate cDIP has low sequence homology to several

LRR proteins that function as adipokines or insulin-like growth factor binding proteins (IGFBPs),

secreted factors that fine-tune endocrine signaling by directly binding insulin or IGF-like peptides

to regulate their serum lifetime or receptor bioavailability[60]. For example, cDIP displays limited

homology (27% identical, 41% similar amino acids) to vertebrate IGF-binding protein acid-labile

subunit (IGFALS), a secreted factor that binds circulating insulin growth factor-1 (IGF-1) to

influence growth and energy storage[61,62]. cDIP also displays some similarity (33% identical, 23%

similar amino acids) to Leucine-rich alpha-2-glycoprotein 1 (LRG1) a secreted obesity-linked

cytokine that regulates insulin signaling[63,64]. Indeed, cDIP's domain architecture closely

resembles these factors with a predicted N-terminal signal peptide and LRR domain (**Fig 6E**).

Furthermore, AlphaFold2 structural predictions suggest structural similarities between cDIP,

LRG1, and IGFALS, implying potential functional similarities in energy signaling (**Fig 6F**). Based

on this, and since *Drosophila* feature a simplified insulin/IGF-like signaling (IIS) system that

governs energy storage, we hypothesized cDIP may function in some aspect of energy

homeostasis.

To probe into the functional relevance of cDIP, we RNAi-depleted *cDIP* in the FB (*Dcg-*

Gal4>cDIP^(RNAi)), and monitored whole body TG levels. cDIP loss did not alter TG in larvae but

led to fly obesity with a striking >2-fold TG increase in adult male and female flies compared to

controls (**Fig 6G**). Since obesity often correlates with elevated oxidative stress[65], we also

queried whether cDIP loss impacted FB redox homeostasis in the presence and absence of

FASN1. We stained larval FBs with the lipid peroxidation biosensor BODIPY-C11. Reduced

BODIPY-C11 exhibits a red fluorescence emission of ~590nm but shifts to green (~510nm)

emission when oxidized[66]. In control larval FBs, we noted red (reduced) BODIPY-C11 (BODIPY-

C11^{Red}) which was incorporated in LDs (**SFig 6D**). This is consistent with previous studies

indicating that LDs serve as lipo-protective depots that reduce lipid oxidation[28]. Control FBs

exhibited a diffuse dim green (oxidized) BODIPY-C11 (BODIPY-C11^{Oxid}), presumably decorating

cellular membranes, and a few green foci were observed (**Fig 6H**). In contrast, *FASN1^(FB-RNAi)*

FBs displayed significantly brighter BODIPY-C11^{Oxid} signal and more foci, consistent with

oxidative stress (**Fig 6H,I**). As expected, there was significantly less BODIPY-C11^{Red} associated

with LDs in *FASN1^(FB-RNAi)* tissues since they were largely depleted of fat, consistent with the

concept that loss of LDs promotes lipid peroxidation (**SFig 6D**). While loss of cDIP alone did not

elevate BODIPY-C11^{Oxid} intensity in FB cells, its dual depletion with FASN1 led to significantly

elevated BODIPY-C11^{Oxid} signal and foci, suggesting cDIP loss perturbed some aspect of redox

homeostasis (**Fig 6H,I**).

Given that cDIP appeared to influence energy storage and redox homeostasis in the FB, we

queried whether *Drosophila* required cDIP for development. Strikingly, for two independent

cDIP[RNAi] lines we noted that whereas FB-specific depletion of cDIP alone did not impact

Drosophila development, its joint loss with FASN1 in the FB resulted in a near complete loss of

fly eclosion, and animals died as pupae (**Fig 6J**). Collectively, this indicates that cDIP influences

some aspect of energy homeostasis and is essential for *Drosophila* development in the absence

of fat stores. We hypothesize that cDIP is secreted from FB cells in response to depletion of fat

stores and facilitates metabolic remodeling by blunting energy signaling to the FB, possibly by

regulating IIS activity. As such, we propose cDIP loss may perturb the ability of FASN1-depleted

larvae to rebalance their metabolism towards glycolysis and catabolic metabolism. Future

studies will dissect how cDIP modulates energy and redox homeostasis.

Discussion:

Energy balance is largely reliant on glucose, which drives glycolysis, the TCA cycle, and

OXPHOS, as well as adipose tissue metabolism[3,4,67]. Dietary sugars stimulate expression of *de*

novo lipogenic enzymes including acetyl CoA carboxylase (ACC) and FASN that convert acetyl

CoA derived from pyruvate to fatty acids that are esterified to TG and assimilated into LDs[5,18]. In

Drosophila, the FB which is the central lipid depot, performs liver-like functions and stores both

TG and glycogen[18,67]. We capitalized on this TG:glycogen dual utilization to identify factors

mediating energy balance and adaptation in animal development. We find that FB-specific loss

of FASN1 leads to a near complete loss of TG, and rebalancing to carbohydrate-oriented

metabolism. FASN1-depleted larvae maintain normal FB tissue size and complete their

developmental life cycle via elevated glycogen storage and glycolysis, but manifest adult female

infertility and shortened lifespans. FASN1-depleted FBs also display signatures of oxidative

stress and lipid peroxidation. We also identify cDIP, a putative cytokine-like factor with

homology to secreted IIS factors IGFALS and LRG1, as critical to FB energy homeostasis when

FASN1 is lost. Our work suggests cDIP may act as a 'sensor' of fat depletion in the FB, and

coordinator of metabolic remodeling that rewires FB metabolism away from nutrient storage and

towards catabolic metabolism.

Drosophila must reach critical weight to achieve pupation and metamorphosis, and the biomass for this is generally thought to come from FB lipids[24–26]. We find *FASN1[FB-RNAi]* larvae lack lipid stores yet grow to full size and pupate without developmental delay. We hypothesize that the ~20-fold concentrated glycogen deposits in *FASN1[FB-RNAi]* FBs may increase body mass by elevating FB cell water content. Glycogen absorbs 3-5 times its weight in water which may help compensate for lipid loss and help *FASN1[FB-RNAi]* larvae to reach critical weight[68]. Additionally, our data indicate that glycogen reserves, previously deemed dispensable for metamorphosis[6,8], are essential for *FASN1[FB-RNAi]* animals during pupal to pharate adult transition, while control animals rely mostly on TG. This metabolic rewiring reveals a previously unappreciated versatility to *Drosophila* development.

Transcriptomics and proteomics identified tobi, an alpha-glucosidase, as essential to bypass TG loss in FB tissue. Similarly, GO term analysis indicated significant reprogramming of glucose metabolism, whereby energy was channeled back into carbohydrates via glycogenesis and possibly gluconeogenesis, and metamorphosis was primarily fueled by aerobic glycolysis (but without lactate production, as in the Warburg effect). In line with this, loss of tobi, essential for glycogenolysis, or enzymes involved in gluconeogenesis, or glycolysis is fatal in *FASN1[FB-RNAi]* animals, but not in controls. It is known that FB glycogen, generally labelled a 'metabolic safeguard', can be mobilized to replenish trehalose in circulation during nutrient shortages[69]. Trehalose may be directly synthesized from glucose by trehalose-6-phosphate synthase (Tps1) exclusively in the FB[70]. Interestingly, recent studies show that 90% of hemolymph trehalose is utilized in early metamorphosis and *Tps1* null mutants do not complete metamorphosis[32,33]. We found that while FB-specific *Tps1-RNAi* larvae developed into adults, loss of *Tps1* and *FASN1* together resulted in a high degree of pupal lethality, suggesting that both FB glycogen and hemolymph trehalose support fly development in the absence of TG. Indeed, we detected

higher amounts trehalose in *FASN1*^FB-RNAi larval hemolymph, probably owing to enhanced

glycogen breakdown and gluconeogenesis in these animals.

Fat depleted *Drosophila* maintain their developmental integrity through rebalancing

fat:carbohydrate metabolism, but does this expose new vulnerabilities? While glycogen

metabolism facilitated *Drosophila* development, fertility was adversely affected in *FASN1*^FB-RNAi

females. *FASN1*^FB-RNA ovaries appeared morphologically small and underdeveloped. This may

be a consequence of a phenomenon called "organ sparing". When energy supply is limited,

animals preferentially support development of certain organs at the cost of others[51]. In *FASN1*^FB-

^RNAi female flies, we noted that egg chamber oocyte selection from nurse cells, and wrapping by

follicle cells, both occurred improperly, stalling oocyte maturation around stage 9-10 of

oogenesis. Since maternal TG deposition is a critical checkpoint for oocyte maturation and

embryo survival, failure of oogenesis in *FASN1*^FB-RNAi females likely results from low level of LDs

in nurse cells. TG stored in LDs of nurse cells is synthesized from DAG, 80% of which is

exogenously resourced from free-floating cells of the dissociated larval FB that persist in the

newly hatched fly[28,55]. We reason that lipophorin-mediated DAG delivery from *FASN1*^FB-RNAi lipid-

deficient fat cells to the ovaries was severely limited, resulting in organ developmental arrest. In

line with this, mid-oogenesis arrest and subsequent apoptosis of egg chambers due to low lipid

content has been reported by other studies[71,72].

Though *FASN1*^FB-RNAi larvae were able to complete metamorphosis and become adults, flies

harbored signatures of metabolic and redox stress. Adult flies exhibited barely detectable TG

stores and half the glycogen reserves of controls; unsurprisingly this rendered them starvation

sensitive. *FASN1*^FB-RNAi male and females displayed shorter median lifespans relative to

controls. We hypothesize this may be, in part, due to elevated ROS production in these animals,

as ROS has been linked to accelerated aging[73]. Metabolomic profiling revealed significant

elevations in the ratio of oxidized glutathione GSSG:GSH, and factors involved GSH redox

cycling in $FASN1^{FB-RNAi}$ FBs. Furthermore, $FASN1^{FB-RNAi}$ animals required SOD and catalase

antioxidants for development. Of note, cDIP depletion in the FB elevated not only nutrient

storage but also redox stress as indicated by BODIPY-C11 staining. The source of ROS

production is unclear but may be linked to elevated oxidative pathways coupled to energy

storage in the FB. Additionally, FASN1 depletion may elevate NADPH, a substrate of *de novo*

lipogenesis and bi-product of the pentose phosphate pathway, which is also upregulated in

$FASN1^{FB-RNAi}$ animals. NADPH accumulation can drive oxidative stress via NAPDH-dependent

oxidase (NOX) ROS production. Alternatively, elevated metabolism in the FASN1-depleted

background may drive ROS signaling from several pathways. Further studies will dissect how

FASN1 loss elevates oxidative stress, and how cDIP both coordinates energy metabolism and

redox homeostasis at the FB.

MATERIALS AND METHODS

Fly cultures and strains

All *Drosophila* animal research was performed in accordance with NIH recommended policies.

Drosophila L3 late feeding/pre-wandering larvae were used for fat tissue microscopy and

biochemical assays, unless otherwise stated. For lifespan and starvation assays, adult male and

female flies were used. Flies were reared and maintained on standard fly food containing

cornmeal, yeast, molasses (~5% sugars), and agar. Basic media for chronic sugar feeding

experiments was prepared with only yeast (10%) and agar (1.5%), supplemented with varying

concentrations of sucrose (0% or 5%). The *Dcg-Gal4* fat body (FB)-specific driver[38] was

provided by Jonathan M. Graff (former UTSW). *UAS-FASN1^{RNAi}* and all other *TRiP RNAi* lines,

PyronicSF-GFP, protein/organelle marker strains and other transgenic stocks used in this study

were obtained from the Bloomington Stock Center (Bloomington, IN). All Gal4/UAS crosses

were cultured at 25°C to maintain uniform transgene expression.

Larval fat body/ gut dissection and staining

Late feeding larvae were gently removed from inside the food media using a paintbrush and

rinsed in water to remove food particles. Larval fat bodies (FBs) or guts were dissected in PBS

using Dumont #5 forceps (Electron Microscopy Sciences). All tissues were fixed in 4%

paraformaldehyde (PFA) for 20 min at room temperature and rinsed briefly in PBS prior to

staining. Lipid droplets in FBs were stained by incubating FBs in 100 µM monodansylpentane

(MDH) LD stain (Abgent) for 20 min at room temperature, in the dark. LDs in guts were stained

with 1 µM Nile Red for 30 min at room temperature, in the dark. Next, FBs and guts were rinsed

in PBS and mounted on slides in SlowFade Gold antifade reagent with nuclear DAPI

(Invitrogen) for subsequent imaging.

Female ovary dissection and staining

Ovaries from 7 to 10-day old adult females were dissected in PBS, fixed in 4% PFA for 20 min

at room temperature. Following a brief wash in PBS, ovaries were incubated in CellMask Actin

stain (1:1000) or Phalloidin (1:400 in PBS-T) for staining of F-actin structures, according to

manufacturer's instructions (Invitrogen). LDs in ovaries were stained with BODIPY 493/503 C12

(1:500) for 30 mins in the dark. Excess stain from ovaries was removed by washing in PBS,

then mounted in SlowFade Gold antifade reagent containing DAPI (Invitrogen) for imaging.

Confocal fluorescence microscopy

Prepared slides were imaged using Zeiss LSM880 inverted laser scanning confocal microscope.

Tissues stained with organelle dyes (LD: MDH/ BODIPY C12/Nile Red; F-actin: phalloidin) and

those expressing transgenic fluorescent-tagged proteins (e.g., UAS-GFP; ER marker *UAS-*

Sec61B-TdT; PM marker *UAS-mcd8-GFP*; pyruvate sensor *PyronicSF-GFP;* mitochondrial

marker *UAS-MitoTimer*) were imaged using the appropriate channel filter DAPI/GFP/RFP. Most

FBs were imaged using a 40X oil immersion objective, and adult female ovaries with 10X or 20X

objective.

Lipid peroxidation assay

FBs dissected from late feeding larvae were incubated in 2 µM of BODIPY 581/591 (Invitrogen,

D3861), a fluorescent lipid peroxidation sensor, according to manufacturer's instructions.

Tissues were rinsed in PBS, fixed in 4% PFA for 20 min (in the dark) and washed briefly again

in PBS. FBs were mounted in SlowFade Gold antifade reagent, and GFP and RFP channel

filters were used to image green (oxidized) and red (reduced) fluorescence in FB cells. GFP and

RFP FB intensities were quantified using Fiji ImageJ and oxidized:reduced ratios were

calculated.

Light and other fluorescence microscopy

Whole 3L larvae without or with GFP-expressing (*Dcg-Gal4>UAS-GFP*) FBs were imaged using

the Leica S8AP0 stereoscope and EVOS FL Cell Imaging System (ThermoFisher), respectively.

Adult male and female flies were also imaged on Leica S8AP0 stereoscope.

Transmission electron microscopy

FB tissue was removed from *Drosophila* larvae, fixed in 2.5% (v/v) glutaraldehyde in 0.1 M

sodium cacodylate buffer paraformaldehyde, and processed in the UT Southwestern Electron

Microscopy Core Facility. They were post-fixed in 1% osmium tetroxide and 0.8% K3[Fe(CN6)]

in 0.1 M sodium cacodylate buffer for 1 hour at room temperature. Cells were rinsed with water

and en bloc stained with 2% aqueous uranyl acetate overnight. Next, they were rinsed in buffer

and dehydrated with increasing concentration of ethanol, infiltrated with Embed-812 resin and

polymerized in a 60°C oven overnight. Blocks were sectioned with a diamond knife (Diatome)

on a Leica Ultracut UCT (7) ultramicrotome (Leica Microsystems) and collected onto copper

grids, post stained with 2% aqueous uranyl acetate and lead citrate. Images were acquired on a

JOEL 1400 Plus transmission electron microscope using a voltage of 120 kV.

Quantitative PCR

Whole RNA was extracted from FB tissues using Trizol reagent (Ambion Life Technologies)

according to manufacturer's instructions. RNA was quantified by Denovix spectrophotometer,

and 1 μg of RNA was reverse transcribed to cDNA using iScript cDNA Synthesis Kit (BioRad).

QPCR was performed with cDNA using the SsoAdvanced Universal SYBR Green Supermix

(BioRad) on the BioRad CFX96 Real-Time System; mRNA expression data were normalized to

that of the fly housekeeping gene rp49. Primer sequences for all transcripts amplified are

available upon request.

Glucose and Trehalose assays

Feeding larvae washed out from culture vials were collected in 630 μm mesh-fitted baskets

(Genesee) and rinsed to get rid of adherent food particles. Larvae were dried and divided onto

into piles (10–12 larvae each) on a strip of parafilm. Larvae were bled by tearing the cuticle with

Dumont 5 forceps (Electron Microscopy Sciences). Two μl of colorless hemolymph was

aspirated from each pile and separately transferred to 96-well plates (Thermo-Scientific)

containing 0.1% N-Phenylthiourea (Sigma-Aldrich) in 50 μl PBS. 150 μl of Autokit Glucose

reagent (Wako) was added to each well and incubated at room temperature for 20 min before

measuring absorbance at 505 nm. Glucose concentration was calculated from a standard

curve generated with manufacturer's glucose standards. For trehalose assays, 8 μl of dilute

hemolymph was treated with 5 μl of (diluted 8X) porcine kidney trehalase (Sigma) overnight at

37□°C. Ten μl of treated sample was assayed for trehalose as described for glucose. Trehalose

amounts were calculated from standard glucose, as for glucose.

Glycogen assay

Whole larvae (10 larvae per sample, in triplicate) or dissected FBs (20-40 larvae per sample, in

triplicate) were homogenized in 300 μl ice-cold PBS using a pestel or sonication and syringe

(29G1/2 needle), respectively. After reserving 20 μl of the homogenate for protein Bradford

quantification, the rest of the homogenate was heat inactivated at 70°C for 10 min. Homogenate

was then centrifuged at maximum speed at 4°C for 10 min and supernatant was collected in a

new tube. In a 96-well plate, 30 μl of each sample was loaded in duplicate rows.   Then, 100 μl

of Autokit Glucose reagent + amyloglucosidase (2 μl amyloglucosidase {Sigma A1602; 25mg}

per 1 ml of Glucose reagent) was added to one row of samples, and 100 μl of Glucose reagent

alone (without amyloglucosidase) was added to the duplicate row of samples, and to the

glucose standards. The plate was incubated at 37°C for 30 min, after which color intensity was

measured using a microplate reader at 505 nm. Free glucose concentration in treated and

untreated samples was calculated based on the glucose standard curve. Glycogen

concentration (as free glucose) was determined by subtracting glucose in the untreated samples

from those treated with amyloglucosidase. Finally, total glycogen was normalized to protein

measured by the Bradford assay.

Periodic Acid-Schiff stain

Glycogen in FB tissues was stained using the Periodic Acid-Schiff (PAS) kit (Sigma). Larval FBs

were dissected in 1% BSA in PBS, fixed with 4% paraformaldehyde for 20 min, and washed

twice with 1% BSA in PBS. FBs were then incubated in 0.5% period acid solution for 5 min,

washed twice with 1% BSA in PBS, then stained with Schiff's reagent for 15 min, washed again,

and mounted on slides. PAS-stained FB tissues were imaged using the Leica DM6B light

microscope.

Thin layer chromatography

Larvae were weighed, then homogenized in 2:1:0.8 of methanol:chloroform:water. Samples

were incubated in a 37°C water bath for 1 hour. Chloroform and 1M KCl (1:1) were added to the

sample, centrifuged at 3000 rpm for 2 min, and the bottom layer containing lipids was aspirated

using a syringe. Lipids were dried using argon gas and resuspended in chloroform (100 µl of

chloroform/7mg of fly weight). Extracted lipids alongside serially diluted standard neutral lipids of

known concentrations were separated on TLC plates using hexane:diethyl ether:acetic acid

solvent (80:20:1, v/v/v). TLC plate was air dried for 10 min, spray stained with 3% copper (II)

acetate in 8% phosphoric acid and incubated at 145°C in the oven for 30 min to 1 hour to allow

bands to develop for scanning and imaging. Neutral lipid (TAG) band intensity was quantified

using Fiji ImageJ software, and lipid concentrations were calculated from the standard curve

generated with standard mixture.

Adult starvation assay

Freshly emerging adult flies were collected over a period of 1 to 3 days and transferred to fresh

standard food bottles. Flies were aged for 7 to 10 days, then male and female flies were

separately moved to food-free vials (10 flies/vial, in triplicate) containing a cotton plug at the

bottom soaked with 2 ml of tap water. Flies were kept at 25°C and vials were examined at least

3X daily to record number of dead flies. Surviving flies were moved to fresh vials with water-

soaked plug every 24 hours for optimal hydration and to avoid fungal growth.

Adult lifespan assay

Flies hatching between 0 to 3 hours were pooled, and 100 males and 100 females were

separately placed in fresh standard food vials (20 flies/vial). Vials were maintained at 25°C, and

dead flies were scored daily while moving surviving flies to fresh food.

Adult survivorship assay

Double FB knockdowns of FASN1 and candidate genes were generated by crossing 5 to 7 *Dcg-*

Gal4>UAS-FASN1-RNAi males to ~20 females from candidate *UAS-RNAi* line in standard food

vials cultured at 25°C. Mated females were allowed to lay eggs for 2 days, then parents were

removed to fresh vials. Larvae carrying *Dcg-Gal4* driver + both *UAS-RNAi* transgenes were

identified and moved to a new vial for developmental monitoring. Animals successfully

completing metamorphosis and hatching into adult flies were scored. This count was compared

to adult survivorship of *Dcg-Gal4 control*, *Dcg-Gal4>UAS-FASN1-RNAi*, *Dcg-Gal4>cand UAS-*

RNAi animals.

Fertility assays

All parents used were 7 to 10 days old. 10 virgin females were crossed with 10 males (controls)

in fresh food vials streaked with yeast paste and kept overnight at 25°C. The following morning,

flies were flipped to new food at 1 hour intervals (3X) to empty females of older eggs/embryos.

Flies were then allowed to lay on fresh plates supplemented with a drop of yeast paste for 4-6

hours, following which all parents were removed from bottles and number of eggs laid was

counted. The hatching of eggs on the food surface and development of progeny was closely

monitored and recorded.

Sample preparation for proteomic profiling

Fat bodies were extracted from *Drosophila* late feeding larvae in chilled Lysis Buffer + protease

inhibitor cocktail. Tissues were homogenized by sonication and supernatant (lysate) was

collected after centrifugation for 30 sec at 1000xg at 4°C. Centrifugation was repeated at

16,000xg for 10 min at 4°C, and supernatant transferred to fresh Eppendorf tube. Sample for

loading was prepared by mixing protein sample with 2X LDS Sample Buffer and Lysis Buffer,

and boiling at 95°C for 5 min. Sample was run on a 10% pre-cast polyacrylamide gel (BioRad)

at 100 V for ~25 min until the dye front was past the ladder. Gel was rinsed in MQ-water, then

stained with Coomassie BB R250 in 10% acetic acid, 50% methanol, 40% water for 20 min at

room temperature with rocking. Subsequently, gel was de-stained for 1 hour in 10% acetic acid,

50% methanol, 40% water with 4 solvent changes. When gel background appeared clear, de-

stain was exchanged for water. Protein gel bands were cut on clean glass surface of gel imager

with new razor, diced into 1 mm cubes, and moved to sterile Eppendorf tubes on ice for delivery

to Proteomics Core Facility at UTSW for protein identification by mass spectrometry.

Sample preparation for metabolomic profiling

Fat tissues dissected from late feeding larvae were homogenized in ice-cold methanol/water

80:20 (vol/vol) using a sonicator. Lysate was subjected to three freeze-thaw cycles between

liquid nitrogen and 37 °C, vortexed vigorously, and then centrifuged at ~20,160xg for 15 min at

4°C. Protein in the supernatant was measured using the Pierce BCA kit, and volume containing

10 µg protein was dried in a SpeedVac. Sample was re-suspended in 100 uL acetonitrile/water

80:20 (vol/vol), vortexed for 1 min, and centrifuged at ~20,160xg for 15 min at 4°C. Metabolite-

containing supernatant was transferred to a fresh Eppendorf tube and submitted to the CRI

Metabolomics Facility at UTSW for LC-MS metabolite analyses. Metabolites were quantified as

described previously [74,75].

Figure Legends:

Figure 1: Loss of FASN1 in *Drosophila* fat body produces virtually fat less flies

773  A) Cartoon of RNA interference-mediated knockdown using the UAS/Gal4 system; parental
774    cross between females carrying tissue (fat body)-specific *Dcg-Gal4* driver and males
775    containing the *UAS-FASN1^{RNAi}* transgene generate progeny lacking *FASN1* in the FB.
776  B) Control (*Dcg-Gal4*) and *FASN1$^{FB-RNAi}$* (*Dcg-Gal4>UAS-FASN1^{RNAi}*) age-matched late
777    feeding/pre-wandering third instar larvae (L3). Scale bar 1 mm.
778  C) Buoyancy-based test of fat content in extracted FBs from control and *FASN1$^{FB-RNAi}$*
779    larvae.
780  D) Confocal images of control and *FASN1$^{FB-RNAi}$* larval (L3) FB periphery and mid-plane
781    sections stained with LD stain monodansylpentane (MDH). Scale bar 25 µm.
782  E) Confocal images of control and *FASN1$^{FB-RNAi}$* FBs at their mid-plane and periphery from
783    larvae expressing ER marker (*Dcg-Gal4>UAS-Sec61B-tDTomato*) and stained with
784    MDH. *FASN1* loss results in adipocytes empty of LDs.
785  F) TEM micrographs of control and *FASN1$^{FB-RNAi}$* FB cells showing peripheral LDs (pLDs,
786    red arrows). *FASN1$^{FB-RNAi}$* cell appears vacant of medial LDs (mLDs) and contains little
787    'islands' of ER, LDs, and other organelles (green arrows).
788  G) Total triglycerides (TG) measured by thin layer chromatography (TLC) in whole L3 pre-
789    wandering control and *FASN1$^{FB-RNAi}$* larvae.
790  H) Size comparison of control and *FASN1$^{FB-RNAi}$* female (left) and male (right) adult flies.
791    Scale bar 2 mm.

793  **Figure 2: *FASN1*-deficient larvae channel dietary glucose to glycogen as energy reserves**
794  A) Volcano plot of proteins detected by LC-MS/MS analyses performed in FBs isolated from
795    control and *FASN1$^{FB-RNAi}$* L3 larvae.
796  B) Lipid metabolism processes were revealed to be amongst the most downregulated in
797    *FASN1$^{FB-RNAi}$* FBs. GO enrichment analysis of proteomics data was performed using
798    Pangea.
799  C) Carbohydrates metabolism processes downregulated in *FASN1$^{FB-RNAi}$* FBs (Pangea)
800  D) Relative expression of *tobi* mRNA in isolated FBs from control and *FASN1$^{FB-RNAi}$* larvae,
801    measured by QPCR.
802  E) Periodic acid-Schiff (PAS) glycogen stain of larval FBs from control and *FASN1$^{FB-RNAi}$*
803    larvae.
804  F) Quantification of glycogen extracted from control and *FASN1$^{FB-RNAi}$* larval FBs.
805  G) Percentage of L3 larvae (control, *FASN1$^{FB-RNAi}$*, *GlyS$^{FB-RNAi}$*, *FASN1$^{FB-RNA\,i}$+ GlyS$^{FB-RNAi}$*)
806    successfully completing metamorphosis to yield adult flies.

H) Percentage of L3 larvae (control, *FASN1*[FB-RNAi], *tobi*[FB-RNAi], *FASN1*[FB-RNA i]+ *tobi*[FB-RNAi])

successfully completing metamorphosis to adult flies.

I) Relative glycogen levels measured in control and *FASN1*[FB-RNAi] animals during distinct

developmental stages: Feeding Larvae: F Larvae (~100-108 h AEL), Wandering Larvae:

W Larvae (non-feeding, ~108-120 h AEL), Pupae (~132-140 h AEL), Pharate Adults

(~205-215 AEL), Virgin Flies (male + female, 0 to 2 h post-eclosion). AEL = After Egg

Laying. *$P<0.05$: FASN1 FL, WL, P, PA, VF relative to Dcg FL, *$P<0.05$: Dcg WL, P,

PA, VF relative to Dcg FL, *$P<0.05$ FASN1 WL, P, PA, VF relative to FASN1 FL.

J) Relative triglyceride levels measured in control and *FASN1*[FB-RNAi] animals during distinct

developmental stages: Feeding Larvae: F Larvae (~100-108 h AEL), Wandering Larvae:

W Larvae (non- feeding, ~108-120 h AEL), Pupae (~132-140 h AEL), Pharate Adults

(~205-215 AEL), Virgin Flies (male + female, 0 to 2 h post-eclosion). AEL = After Egg

Laying. *$P<0.05$: FASN1 FL, WL, P, PA, VF relative to Dcg FL, *$P<0.05$: Dcg WL, P,

PA, VF relative to Dcg FL, *$P<0.05$ FASN1 WL, P, PA, VF relative to FASN1 FL.

822 Fig 3: Larvae lacking FASN1 elevate glycolysis and PPP while down-regulating

823 mitochondrial TCA and ETC

A) Schematic representing the fate of glucose and role of FASN in *de novo* lipogenesis.

B) Relative increases (delta fold) in protein levels of glycolysis enzymes in *FASN1*[FB-RNAi]

FBs revealed by proteomics profiling.

C) Expression of fluorescent pyruvate biosensor *UAS-PyronicSF-GFP* in control and

FASN1[FB-RNAi] in larval FBs: Fire-LUT pseudo-colored (Fiji). Scale Bar 50 µm.

D) Percentage of L3 larvae (control, *FASN1*[FB-RNAi], *Pyk*[FB-RNAi], *FASN1*[FB-RNAi]+ *Pyk*[FB-RNAi])

successfully completing metamorphosis to adult flies.

E) Percentage of control and *FASN1*[FB-RNAi] larvae, and *G6P*[FB-RNAi] or *Pepck2*[FB-RNAi] larvae

with or without *FASN1* successfully completing metamorphosis to adult flies.

F) Relative changes (delta fold) in tricarboxylic acid cycle (TCA) enzymes in *FASN1*[FB-RNAi]

FBs detected by global proteomics.

G) Heat map of LC-MS/MS metabolomics showing raw abundances (colored by row) of up-

regulated and down-regulated proteins in the FBs of *FASN1*[FB-RNAi] larvae relative to *Dcg-*

Gal4 controls.

H) Confocal images of control and *FASN1*[FB-RNAi] FBs expressing mitochondrial fluorescent

marker *UAS-MitoTimer* to detect changes organelle density and morphology. Scale bar

25 µm, Scale bar (inset) 5 µm.

Fig 4: *FASN1^FB-RNAi* development to adults requires diet-dependent glycogen storage and trehalose production in the FB

    A) Longevity assay performed for control (*Dcg-Gal4*) and *FASN1^FB-RNAi* adult females and males (n = 100, each).

    B) Starvation assay using 7-10 days old control and *FASN1^FB-RNAi* female and male flies (n = 30, each).

    C) TG measured in control and *FASN1^FB-RNAi* male and female flies.

    D) Relative glycogen measured in control and *FASN1^FB-RNAi* male and female flies.

    E) TG measured in control and *FASN1^FB-RNAi* larvae reared on 0% and 5%sucrose (basic media) diet.

    F) Glycogen measured in control and *FASN1^FB-RNAi* larvae reared on 0% and 5% sucrose (basic media) diet.

    G) Percentage of control and *FASN1^FB-RNAi* eggs developing to adults on 0% and 5% sucrose diet.

    H) Control and *FASN1^FB-RNAi* larval hemolymph glucose levels (mg/DL).

    I) Control and *FASN1^FB-RNAi* larval hemolymph trehalose levels (mg/DL).

Fig 5: *FASN1^FB-RNAi* females display arrested oogenesis and are infertile

    A) Fertility assay performed to test viability of eggs produced by control (*Dcg-Gal4*) or *FASN1^FB-RNAi* females mated with either control or *FASN1^FB-RNAi* males.

    B) Confocal images of ovaries from 7–10-day old control and *FASN1^FB-RNAi* adult females stained with CellMask Actin stain. Entire *FASN1^FB-RNAi* ovary (right) is visible at 10X magnification. Parts of the control ovary are labelled (left). Scale bar 100 μm.

    C) Confocal images of phalloidin-stained ovaries from control and *FASN1^FB-RNAi* females. White arrows point to mature oocyte in control ovary and arrested oocyte in *FASN1^FB-RNAi*. Scale bar (20X) 100 μm, Scale bar (20X) 50 μm.

    D) Confocal images of ovaries from control and *FASN1^FB-RNAi* females stained with CellMask Actin stain. Yellow arrows point to follicle cells of stage 9/10 egg chambers. White arrows point to oocyte; oocyte maturation fails in *FASN1^FB-RNAi* ovaries. Scale bar 50 μm.

    E) Confocal images of LDs in control and *FASN1^FB-RNAi* ovaries stained with BODIPY. Orange arrow in control ovary indicates LDs accumulated in nurse cells and purple arrow indicates ring canal where movement of LDs from nurse cells to the developing

oocyte is occurring. There also appear to be plentiful LDs in the mature oocyte*. LDs

are sparse in *FASN1^FB-RNAi* egg chambers. Scale bar (20X) 100 µm, Scale bar (20X) 50

µm.

F) Confocal images of ovaries from control and *FASN1^FB-RNAi* females stained with nuclear

stain DAPI. Blue arrow in control ovary point to distinct nuclei in mature oocytes, absent

in *FASN1^FB-RNAi* ovary. Scale bar (20X) 100 µm, Scale bar (20X) 50 µm.

Fig 6: *FASN1^FB-RNAi* animals suffer from oxidative stress and require antioxidants to

complete development

A) Heat map of oxidation products of metabolites maintaining redox balance in the FBs of

FASN1^FB-RNAi larvae relative to *Dcg-Gal4* controls (raw abundances).

B) Ratios of glutathione disulfide (oxidized) to glutathione (reduced) in control and

FASN1^FB-RNAi FBs.

C) Relative changes (delta fold) in proteins involved in glutathione metabolism in *FASN1^FB-*

RNAi FBs relative to control.

D) Percentage of control and *FASN1^FB-RNAi* larvae, as well as those deficient of antioxidants

Sod2 or *catalase,* alone or also without *FASN1,* successfully developing into adult flies.

E) Domain architecture of *Drosophila melanogaster* (*D.m.*) cDIP, *Homo sapiens* (*H.s.*)

LRG1, and *H.s.* IGFALS.

F) AlphaFold2 structural predictions of *D.m.* cDIP, *H.s.* LRG1, and *H.s.* IGFALS.

Confidence score is colored blue-to-red (high-to-low) according to AlphaFold2.

G) Triglyceride (TG) levels for male (M) and female (F) adult control (*Dcg-Gal4*) and *Dcg-*

Gal4>cDIP^RNAi flies.

H) Confocal images of BODIPY-C11 stained larval FBs from control (*Dcg-Gal4*), or *Dcg-*

Gal4-driven *cDIP^RNAi*, *FASN1^RNAi*, or *FASN1^RNAi*+ *cDIP^RNAi* dual depletion.

I) Quantification of BODIPY-C11^Oxid puncta and signal intensity in control, *FASN1^FB-RNAi*,

cDIP^FB-RNAi, and *FASN1^FB-RNAi*+ *cDIP^FB-RNAi* FBs. Also, quantification of BODIPY-C11^Oxid

foci (#foci/100 square microns) and total signal (IntDen/square micron) for experiments

in Panel H.

J) Percentage of L3 larvae (control, *FASN1^FB-RNAi*, *cDIP^FB-RNAi*, *FASN1^FB-RNAi*+ *cDIP^FB-RNAi*)

developing to adult flies. Two independent *UAS-cDIP^RNAi* lines were tested.

Fig 7: Working model for how triglyceride: carbohydrate cross-talk and metabolic re-wiring occurs in a 'fat-less' *Drosophila* to promote growth and development

Supplemental Figures:

SFig 1:

A) Fluorescence micrograph of age-matched L3 control (*Dcg-Gal4*) and *FASN1^FB-RNAi* pre-wandering larvae expressing soluble GFP in their FBs (*Dcg-Gal4>UAS-GFP*).

B) Confocal mid-plane sections of FBs from control and *FASN1^FB-RNAi* age-matched L3 larvae expressing plasma membrane (PM) marker mcd8-GFP (*Dcg-Gal4>UAS-mcd8-GFP*).

C) Relative triglycerides (TG) in isolated larval guts from control and *FASN1^FB-RNAi* L3 larvae.

D) Confocal section images of FB periphery (top) and mid-plane (bottom) stained with LD stain MDH. FBs from the ubiquitous but mosaic driver *da-Gal4* (control) and *da-Gal4>UAS-FASN1^RNAi* L3 larvae also express PM marker mcd8-GFP; green indicates where *da-gal4* is expressed, and where *FASN1*-knockdown also occurs (right only).

SFig 2:

A) Principal Component Analysis (PCA) for global proteomics performed in FBs isolated from control and *FASN1^FB-RNAi* L3 larvae.

B) Relative (delta fold) changes in FASN isoform mRNA and protein expression detected in *FASN1^FB-RNAi* FBs by QPCR and proteomics analysis, respectively.

C) Phospholipid metabolism processes upregulated in *FASN1^FB-RNAi* FBs, determined by GO enrichment analysis of proteomics data using Pangea.

D) Confocal images of FB periphery and mid-plane sections from control (*Dcg-Gal4*) alongside FASN loss-of-functions isoforms (*FASN1^FB-RNAi*, *FASN2* global mutant, *FASN3^FB-RNAi*), stained with LD stain monodansylpentane (MDH). Scale bar 50 µm.

E) Confocal images of control and *FASN1^FB-RNAi* larval FB periphery and mid-plane sections expressing Glycogenin-YFP and stained with nuclear DAPI. Scale bar 50 µm.

F) mRNA levels of glycogen synthesis and degradation enzymes measured by QPCR in *FASN1^FB-RNAi* FBs relative to control.

SFig 3:

A) Gene expression (delta fold) changes in pyruvate metabolism enzymes in $FASN1^{FB-RNAi}$ FBs relative to controls, measured by QPCR.

B) Percentage of L3 larvae (control, $FASN1^{FB-RNAi}$, $Ldh^{FB-RNAi}$, $FASN1^{FB-RNA\,i}$+ $Ldh^{FB-RNAi}$) completing development to adult flies.

C) Relative changes (delta fold) in electron transport chain (ETC) enzymes in $FASN1^{FB-RNAi}$ FBs detected by global proteomics.

D) Percentage of L3 larvae (control, $FASN1^{FB-RNAi}$, $Mpc^{FB-RNAi}$, $FASN1^{FB-RNA\,i}$+ $Mpc^{FB-RNAi}$) completing development to adult flies.

E) Percentage of L3 larvae (control, $FASN1^{FB-RNAi}$, $Pdha^{FB-RNAi}$, $FASN1^{FB-RNA\,i}$+ $Pdha^{FB-RNAi}$) completing development to adult flies.

F) Relative changes (delta fold) in mitochondrial and peroxisomal fatty acid oxidation (FAO) enzymes in $FASN1^{FB-RNAi}$ FBs detected by global proteomics.

SFig 4:

A) Percentage of L3 larvae (control, $FASN1^{FB-RNAi}$, $Mpc^{FB-RNAi}$, $Tps1^{FB-RNA\,i}$+ $Tps1^{FB-RNAi}$) completing metamorphosis to adults.

SFig 6:

A) Percentage of L3 larvae (control, $FASN1^{FB-RNAi}$, $Sod1^{FB-RNAi}$, $FASN1^{FB-RNAi}$+ $Sod1^{FB-RNAi}$) developing to adult flies.

B) Confocal images of control and $FASN1^{FB-RNAi}$ larval FB cells expressing peroxisomal marker Pex3-SKL-GFP. Scale bar 50 μm.

C) Percentage of L3 larvae (control, $FASN1^{FB-RNAi}$, $Pex5^{FB-RNAi}$, $FASN1^{FB-RNAi}$+ $Pex5^{FB-RNAi}$) developing to adult flies.

D) Confocal images of FBs from control, $FASN1^{FB-RNAi}$, $cDIP^{FB-RNAi}$, and $FASN1^{FB-RNAi}$+ $cDIP^{FB-RNAi}$ larvae stained with lipid peroxidation biosensor BODIPY-C11. Reduced BODIPY-C11 (red) is shown. Scale bar 25 μm.

References:

1. Schönborn, J. W., Jehrke, L., Mettler-Altmann, T. & Beller, M. FlySilico: Flux balance modeling of Drosophila larval growth and resource allocation. *Sci. Rep.* **9**, 17156 (2019).

2. Wong, K. K. L., Liao, J. Z. & Verheyen, E. M. A positive feedback loop between Myc and

aerobic glycolysis sustains tumor growth in a Drosophila tumor model. *eLife* **8**, e46315

(2019).

3. Chatterjee, N. & Perrimon, N. What fuels the fly: Energy metabolism in Drosophila and its

application to the study of obesity and diabetes. *Sci. Adv.* **7**, eabg4336 (2021).

4. Garrido, D. *et al.* Fatty acid synthase cooperates with glyoxalase 1 to protect against sugar

toxicity. *PLoS Genet.* **11**, e1004995 (2015).

5. Mattila, J. & Hietakangas, V. Regulation of Carbohydrate Energy Metabolism in Drosophila

melanogaster. *Genetics* **207**, 1231–1253 (2017).

6. Gáliková, M. & Klepsatel, P. Endocrine control of glycogen and triacylglycerol breakdown in

the fly model. *Semin. Cell Dev. Biol.* **138**, 104–116 (2023).

7. Meschi, E. & Delanoue, R. Adipokine and fat body in flies: Connecting organs. *Mol. Cell.*

Endocrinol. **533**, 111339 (2021).

8. Gáliková, M. *et al.* Energy Homeostasis Control in Drosophila Adipokinetic Hormone

Mutants. *Genetics* **201**, 665–683 (2015).

9. Miao, T. *et al.* Acetyl-CoA-mediated autoacetylation of fatty acid synthase as a metabolic

switch of de novo lipogenesis in Drosophila. *Proc. Natl. Acad. Sci. U. S. A.* **119**, e2212220119

(2022).

10.   Musselman, L. P. & Kühnlein, R. P. Drosophila as a model to study obesity and metabolic

disease. *J. Exp. Biol.* **221**, jeb163881 (2018).

11.   Walls, S. M. *et al.* Ceramide-Protein Interactions Modulate Ceramide-Associated

Lipotoxic Cardiomyopathy. *Cell Rep.* **22**, 2702–2715 (2018).

12.   Baker, K. D. & Thummel, C. S. Diabetic larvae and obese flies-emerging studies of

metabolism in Drosophila. *Cell Metab.* **6**, 257–266 (2007).

13.   Verheyen, E. M. The power of Drosophila in modeling human disease mechanisms. *Dis.*

Model. Mech. **15**, dmm049549 (2022).

14.   Musselman, L. P. *et al.* A high-sugar diet produces obesity and insulin resistance in wild-

type Drosophila. *Dis. Model. Mech.* **4**, 842–849 (2011).

15.   Ugrankar, R., Liu, Y., Provaznik, J., Schmitt, S. & Lehmann, M. Lipin is a central regulator

of adipose tissue development and function in Drosophila melanogaster. *Mol. Cell. Biol.* **31**,

1646–1656 (2011).

16.   Ugrankar, R. *et al.* Drosophila glucome screening identifies Ck1alpha as a regulator of

mammalian glucose metabolism. *Nat. Commun.* **6**, 7102 (2015).

17.   Arrese, E. L. & Soulages, J. L. Insect fat body: energy, metabolism, and regulation. *Annu.*

Rev. Entomol. **55**, 207–225 (2010).

18.   Musselman, L. P. *et al.* Role of fat body lipogenesis in protection against the effects of

caloric overload in Drosophila. *J. Biol. Chem.* **288**, 8028–8042 (2013).

19.   Hoshizaki, D. K. *et al.* Embryonic fat-cell lineage in Drosophila melanogaster. *Dev. Camb.*

Engl. **120**, 2489–2499 (1994).

20.   Nelliot, A., Bond, N. & Hoshizaki, D. K. Fat-body remodeling in Drosophila melanogaster.

Genes. N. Y. N 2000 **44**, 396–400 (2006).

21.   Yongmei Xi, Y. Z. Fat Body Development and its Function in Energy Storage and Nutrient

Sensing in Drosophila melanogaster. *J. Tissue Sci. Eng.* **06**, (2015).

22.   Zheng, H., Yang, X. & Xi, Y. Fat body remodeling and homeostasis control in Drosophila.

Life Sci. **167**, 22–31 (2016).

23.   Chng, W.-B. A., Hietakangas, V. & Lemaitre, B. Physiological Adaptations to Sugar Intake:

New Paradigms from Drosophila melanogaster. *Trends Endocrinol. Metab. TEM* **28**, 131–142

(2017).

24.   Jacobs, H. T., George, J. & Kemppainen, E. Regulation of growth in Drosophila

melanogaster: the roles of mitochondrial metabolism. *J. Biochem. (Tokyo)* **167**, 267–277

(2020).

25.   Nishimura, T. Feedforward Regulation of Glucose Metabolism by Steroid Hormones

Drives a Developmental Transition in Drosophila. *Curr. Biol. CB* **30**, 3624-3632.e5 (2020).

26.   Tennessen, J. M. & Thummel, C. S. Coordinating growth and maturation - insights from

Drosophila. *Curr. Biol. CB* **21**, R750-757 (2011).

27.   Aguila, J. R., Suszko, J., Gibbs, A. G. & Hoshizaki, D. K. The role of larval fat cells in adult

Drosophila melanogaster. *J. Exp. Biol.* **210**, 956–963 (2007).

28.   Heier, C., Klishch, S., Stilbytska, O., Semaniuk, U. & Lushchak, O. The Drosophila model

to interrogate triacylglycerol biology. *Biochim. Biophys. Acta Mol. Cell Biol. Lipids* **1866**,

158924 (2021).

29.   Merkey, A. B., Wong, C. K., Hoshizaki, D. K. & Gibbs, A. G. Energetics of metamorphosis

in Drosophila melanogaster. *J. Insect Physiol.* **57**, 1437–1445 (2011).

30.   Cartwright, B. R. *et al.* Seipin performs dissectible functions in promoting lipid droplet

biogenesis and regulating droplet morphology. *Mol. Biol. Cell* **26**, 726–739 (2015).

31. Tian, Y. *et al.* Tissue-autonomous function of Drosophila seipin in preventing ectopic

lipid droplet formation. *PLoS Genet.* **7**, e1001364 (2011).

32. Yamada, T., Hironaka, K.-I., Habara, O., Morishita, Y. & Nishimura, T. A developmental

checkpoint directs metabolic remodelling as a strategy against starvation in Drosophila. *Nat.*

Metab. **2**, 1096–1112 (2020).

33. Yamada, T. *et al.* The role of glycogen in development and adult fitness in Drosophila.

Dev. Camb. Engl. **146**, dev176149 (2019).

34. Ugrankar, R. *et al.* Drosophila Snazarus Regulates a Lipid Droplet Population at Plasma

Membrane-Droplet Contacts in Adipocytes. *Dev. Cell* **50**, 557-572.e5 (2019).

35. D'Erchia, A. M., Tullo, A., Lefkimmiatis, K., Saccone, C. & Sbisà, E. The fatty acid synthase

gene is a conserved p53 family target from worm to human. *Cell Cycle Georget. Tex* **5**, 750–

758 (2006).

36. Heier, C. & Kühnlein, R. P. Triacylglycerol Metabolism in Drosophila melanogaster.

Genetics **210**, 1163–1184 (2018).

37. Hood, S. E. *et al.* Nuclear translocation ability of Lipin differentially affects gene

expression and survival in fed and fasting Drosophila. *J. Lipid Res.* **61**, 1720–1732 (2020).

38. Suh, J. M. *et al.* Hedgehog signaling plays a conserved role in inhibiting fat formation.

Cell Metab. **3**, 25–34 (2006).

39. Kühnlein, R. P. Thematic review series: Lipid droplet synthesis and metabolism: from

yeast to man. Lipid droplet-based storage fat metabolism in Drosophila. *J. Lipid Res.* **53**,

1430–1436 (2012).

40.   Buch, S., Melcher, C., Bauer, M., Katzenberger, J. & Pankratz, M. J. Opposing effects of

dietary protein and sugar regulate a transcriptional target of Drosophila insulin-like peptide

signaling. *Cell Metab.* **7**, 321–332 (2008).

41.   Rulifson, E. Fly-let biology and the high protein/low carb diet. *Cell Metab.* **7**, 281–283

(2008).

42.   Kilwein, M. D., Johnson, M. R., Thomalla, J. M., Mahowald, A. P. & Welte, M. A.

Drosophila embryos spatially sort their nutrient stores to facilitate their utilization. *Dev.*

Camb. Engl. **150**, dev201423 (2023).

43.   Arce-Molina, R. *et al.* A highly responsive pyruvate sensor reveals pathway-regulatory

role of the mitochondrial pyruvate carrier MPC. *eLife* **9**, e53917 (2020).

44.   Lee, J. E., Oney, M., Frizzell, K., Phadnis, N. & Hollien, J. Drosophila melanogaster

activating transcription factor 4 regulates glycolysis during endoplasmic reticulum stress. *G3*

Bethesda Md **5**, 667–675 (2015).

45.   Mengmeng Liu *et al.* Wingless signaling promotes lipid mobilization through signal-

induced transcriptional repression. *bioRxiv* 2023.01.25.525602 (2023)

doi:10.1101/2023.01.25.525602.

46.   Faust, J. E., Verma, A., Peng, C. & McNew, J. A. An inventory of peroxisomal proteins and

pathways in Drosophila melanogaster. *Traffic Cph. Den.* **13**, 1378–1392 (2012).

47.   Xu, M. *et al.* NAD kinase sustains lipogenesis and mitochondrial metabolismthrough

fatty acid synthesis. *Cell Rep.* **37**, 110157 (2021).

48.   Pachnis, P. *et al.* In vivo isotope tracing reveals a requirement for the electron transport

chain in glucose and glutamine metabolism by tumors. *Sci. Adv.* **8**, eabn9550 (2022).

49.   Chen, W., Zhao, H. & Li, Y. Mitochondrial dynamics in health and disease: mechanisms

and potential targets. *Signal Transduct. Target. Ther.* **8**, 333 (2023).

50.   Klepsatel, P., Knoblochová, D., Girish, T. N., Dircksen, H. & Gáliková, M. The influence of

developmental diet on reproduction and metabolism in Drosophila. *BMC Evol. Biol.* **20**, 93

(2020).

51.   Lanet, E. & Maurange, C. Building a brain under nutritional restriction: insights on

sparing and plasticity from Drosophila studies. *Front. Physiol.* **5**, 117 (2014).

52.   Avilés-Pagán, E. E. & Orr-Weaver, T. L. Activating embryonic development in Drosophila.

Semin. Cell Dev. Biol. **84**, 100–110 (2018).

53.   Welte, M. A. As the fat flies: The dynamic lipid droplets of Drosophila embryos. *Biochim.*

Biophys. Acta **1851**, 1156–1185 (2015).

54.   Sieber, M. H. & Spradling, A. C. Steroid Signaling Establishes a Female Metabolic State

and Regulates SREBP to Control Oocyte Lipid Accumulation. *Curr. Biol. CB* **25**, 993–1004

(2015).

55.   Ziegler, R. & Van Antwerpen, R. Lipid uptake by insect oocytes. *Insect Biochem. Mol.*

Biol. **36**, 264–272 (2006).

56.   Bernard, K. E., Parkes, T. L. & Merritt, T. J. S. A model of oxidative stress management:

moderation of carbohydrate metabolizing enzymes in SOD1-null Drosophila melanogaster.

PloS One **6**, e24518 (2011).

57.   Cui, Y. *et al.* NADPH accumulation is responsible for apoptosis in breast cancer cells

induced by fatty acid synthase inhibition. *Oncotarget* **8**, 32576–32585 (2017).

58.   Gao, L., Laude, K. & Cai, H. Mitochondrial pathophysiology, reactive oxygen species, and

1100   cardiovascular diseases. *Vet. Clin. North Am. Small Anim. Pract.* **38**, 137–155, vi (2008).

1101   59.   Strilbytska, O. *et al.* Dietary Sucrose Determines Stress Resistance, Oxidative Damages,

1102   and Antioxidant Defense System in Drosophila. *Scientifica* **2022**, 7262342 (2022).

1103   60.   Allard, J. B. & Duan, C. IGF-Binding Proteins: Why Do They Exist and Why Are There So

1104   Many? *Front. Endocrinol.* **9**, 117 (2018).

1105   61.   Baxter, R. C. Endocrine and cellular physiology and pathology of the insulin-like growth

1106   factor acid-labile subunit. *Nat. Rev. Endocrinol.* (2024) doi:10.1038/s41574-024-00970-4.

1107   62.   Camilli, C., Hoeh, A. E., De Rossi, G., Moss, S. E. & Greenwood, J. LRG1: an emerging

1108   player in disease pathogenesis. *J. Biomed. Sci.* **29**, 6 (2022).

1109   63.   He, S. *et al.* LRG1 is an adipokine that mediates obesity-induced hepatosteatosis and

1110   insulin resistance. *J. Clin. Invest.* **131**, e148545 (2021).

1111   64.   Choi, C. H. J. *et al.* LRG1 is an adipokine that promotes insulin sensitivity and suppresses

1112   inflammation. *eLife* **11**, e81559 (2022).

1113   65.   Manna, P. & Jain, S. K. Obesity, Oxidative Stress, Adipose Tissue Dysfunction, and the

1114   Associated Health Risks: Causes and Therapeutic Strategies. *Metab. Syndr. Relat. Disord.* **13**,

1115   423–444 (2015).

1116   66.   Lubojemska, A. *et al.* Adipose triglyceride lipase protects renal cell endocytosis in a

1117   Drosophila dietary model of chronic kidney disease. *PLoS Biol.* **19**, e3001230 (2021).

1118   67.   Ceperuelo-Mallafré, V. *et al.* Adipose tissue glycogen accumulation is associated with

1119   obesity-linked inflammation in humans. *Mol. Metab.* **5**, 5–18 (2016).

68.   Gefen, E., Marlon, A. J. & Gibbs, A. G. Selection for desiccation resistance in adult

Drosophila melanogaster affects larval development and metabolite accumulation. *J. Exp.*

Biol. **209**, 3293–3300 (2006).

69.   Yamada, T., Habara, O., Kubo, H. & Nishimura, T. Fat body glycogen serves as a

metabolic safeguard for the maintenance of sugar levels in Drosophila. *Dev. Camb. Engl.* **145**,

dev158865 (2018).

70.   Matsuda, H., Yamada, T., Yoshida, M. & Nishimura, T. Flies without trehalose. *J. Biol.*

Chem. **290**, 1244–1255 (2015).

71.   Parra-Peralbo, E. & Culi, J. Drosophila lipophorin receptors mediate the uptake of

neutral lipids in oocytes and imaginal disc cells by an endocytosis-independent mechanism.

PLoS Genet. **7**, e1001297 (2011).

72.   Buszczak, M., Lu, X., Segraves, W. A., Chang, T. Y. & Cooley, L. Mutations in the midway

gene disrupt a Drosophila acyl coenzyme A: diacylglycerol acyltransferase. *Genetics* **160**,

1511–1518 (2002).

73.   Lennicke, C. & Cochemé, H. M. Redox signalling and ageing: insights from Drosophila.

Biochem. Soc. Trans. **48**, 367–377 (2020).

74.   Aurora, A. B. *et al.* Loss of glucose 6-phosphate dehydrogenase function increases

oxidative stress and glutaminolysis in metastasizing melanoma cells. *Proc. Natl. Acad. Sci. U.*

S. A. **119**, e2120617119 (2022).

75.   Tasdogan, A. *et al.* Metabolic heterogeneity confers differences in melanoma metastatic

potential. *Nature* **577**, 115–120 (2020).

1141

1142

1143 **Figures:**

Fig. 1

1144

Fig. 2

1145

Fig. 3

1146

Fig. 4

Fig. 5

Fig. 6

Fig. 7

1150

Supp Fig. 1

A

Scores Plot

○ Ctrl
○ FASN1^FB-RNAi

PC 2 (21.8 %)

PC 1 (54.8 %)

B

Relative Expression in *FASN1^FB-RNAi*

All *P < 0.05

mRNA   Protein

Delta Fold change

ND

FASN1 FASN2 FASN3 FASN1 FASN2 FASN3

C

Phospholipid Metabolism (upregulated)

	P-value
phosphatidylcholine biosynthetic process	0.00037
phosphatidylcholine metabolic process	0.00109
phosphatidylethanolamine biosynthetic process	0.00356
phosphatidylethanolamine metabolic process	0.00527
phospholipid biosynthetic process	1.32E-05
glycerophospholipid biosynthetic process	0.00092
phospholipid metabolic process	0.00261
glycerophospholipid metabolic process	0.01957

Fold Enrichment

D

Control   *FASN1^FB-RNAi*   *FASN2 mutant*   *FASN3^FB-RNAi*

LD (MDH)

E

Control   *FASN1^FB-RNAi*

Glycogenin-YFP nuclear DAPI

F

Relative mRNA Expression in
FASN1^FB-RNAi

All *P < 0.05

Glycogenesis   Glycogenolysis

Fold change

Pgm GlyS AGBE   GlyP AGL

1152

Supp Fig. 2

Supp Fig. 3

A

Larvae to Adult Survivorship

1154

Supp Fig. 4

A

Larvae to Adult Survivorship

B

Control   FASN1^FB-RNAi

Pex3-SKL-GFP

C

Larvae to Adult Survivorship

D

Dcg-Gal4   FASN1^FB-RNAi

cDIP^FB-RNAi   FASN1^FBRNAi +cDIP^FB-RNAi

BODIPY C11^Red

1155

Supp Fig. 6