# Residues Neighboring an SH3-Binding Motif Participate in the Interaction *In Vivo*

David F. Jordan[1,2,3,4,*], Alexandre K. Dubé[1,2,3,4,5], Ugo Dionne[2,3,4,6,7,8], David Bradley[1,2,3,4,5] and Christian R. Landry [1,2,3,4,5,*]

1. Département de Biochimie, microbiologie et bio-informatique, Université Laval, 1045 Avenue de la Médecine, Québec, QC, Canada, G1V 0A6
2. Institut de Biologie Intégrative et des Systèmes (IBIS), 1030, Avenue de la Médecine, Université Laval, Québec, QC, Canada, G1V 0A6
3. PROTEO-Regroupement Québécois de Recherche sur la Fonction, l'Ingénierie et les Applications des Protéines, Université Laval, 1030, Avenue de la Médecine Québec, QC, Canada, G1V 0A6
4. Centre de Recherche en Données Massives de l'Université Laval, Université Laval, 1065, Avenue de la Médecine, Québec, QC, Canada, G1V 0A6
5. Département de Biologie, Université Laval, 1045 Avenue de la Médecine, Québec, QC, Canada, G1V 0A6
6. Centre de Recherche du Centre Hospitalier Universitaire (CHU) de Québec-Université Laval, 11, côte du Palais, Québec, QC, Canada, G1R 2J6
7. Centre de Recherche sur le Cancer de l'Université Laval, 9, rue McMahon, Québec, QC, Canada, G1R 3S3
8. Present address: Lunenfeld-Tanenbaum Research Institute, Sinai Health, 600 University Avenue, Toronto, ON, Canada, M5G 1X5

* Correspondance: 1030, Avenue de la Médecine, Université Laval, Québec, QC, Canada, G1V 0A6 Email: david.jordan.2@ulaval.ca. 1030, Avenue de la Médecine, Université Laval, Québec, QC, Canada, G1V 0A6 Email: christian.landry@bio.ulaval.ca

**Running head:** Residues neighboring motif interaction
**Keywords:** Short Linear motif, Src Homology 3 (SH3) domain, Deep mutational scanning, Binding affinity, Structure prediction

# Abstract

In signaling networks, protein-protein interactions are often mediated by modular domains that bind short linear motifs. The motifs' sequences affect many factors, among them affinity and specificity, or the ability to bind strongly and to the appropriate partners. Using Deep Mutational Scanning to create a mutant library, and protein complementation assays to measure protein-protein interactions, we determined the *in vivo* binding strength of a library of mutants of a binding motif on the MAP kinase kinase Pbs2, which binds the SH3 domain of the osmosensor protein Sho1 in *Saccharomyces cerevisiae*. These measurements were made using the full-length endogenous proteins, in their native cellular environment. We find that along with residues within the canonical motif, many mutations in the residues neighboring the motif also modulate binding strength. Interestingly, all Pbs2 mutations which increase binding are situated outside of the Pbs2 region that interacts with the canonical SH3 binding pocket, suggesting that other surfaces on Sho1 contribute to binding. We use predicted structures and mutations to propose a model of binding which involves residues neighboring the canonical Pbs2 motif binding outside of the canonical SH3 binding pocket. We compared this predicted structure with known structures of SH3 domains binding peptides through residues outside of the motif, and put forth possible mechanisms through which Pbs2 can bind specifically to Sho1. We propose that for certain SH3 domain-motif pairs, affinity and specificity are determined by a broader range of sequences than what has previously been considered, potentially allowing easier differentiation between otherwise similar partners.

# Summary

Protein-protein interactions are often mediated by a binding domain on one protein and a short disordered binding motif on another protein. We measured the binding strength of a mutant library of a binding motif situated in the yeast protein Pbs2 to the SH3 domain of Sho1. Many mutations in the residues neighboring the motif affect binding. A protein structure prediction of the interaction partners shows that residues neighboring the motif bind residues outside the known binding pocket on the SH3 domain. The Sho1-Pbs2 interaction differs enough from other known SH3-motif pairs to allow specific binding.

# Introduction

Cells possess complex and robust signaling networks that detect stimuli and trigger responses. These signaling networks are composed of a series of protein-protein interactions, which are often mediated by modular interaction domains. Many classes of protein interaction domains are shared across pathways and species, yet fulfill different roles and functions (Pawson *et al.* 2002). The different constraints placed on interaction domains by their distinct roles contribute to explaining the divergence in sequence between homologs of the same domain (Ernst *et al.* 2010; Dionne *et al.* 2022). Therefore, to understand how domain sequences have evolved and continue to evolve, the phenotypic consequences of mutations must be understood. Two of the major phenotypes for protein-protein interaction domains are affinity, that is, the strength of binding to the partner protein, and specificity, that is, the ability to bind to the appropriate partner proteins, and to not form spurious interactions with other proteins in the cellular environment (Ivarsson and Jemth 2019).

Many interaction domains bind short, intrinsically disordered stretches of their interaction partners, known as short linear motifs (SLiMs) or simply binding motifs (Gouw *et al.* 2018). Binding motifs are involved in a wide swath of cell signaling pathways, and underlie many important mechanisms in human and other cells (Kumar *et al.* 2024). Uncovering the determinants of binding can elucidate the functioning of human cells and certain diseases that affect them, such as various cancers in which motif-mediated signaling plays a part (Van Roey *et al.* 2014). Additionally, viruses also express their own binding motifs to disrupt the signaling machinery of their hosts (Davey *et al.* 2011). Understanding domain-motif binding can also allow them to be used as tools, for example by attaching therapeutic proteins to domains and motifs to hydrogels, to ensure the gradual release of the therapeutic proteins in human tissues (Delplace *et al.* 2019).

Interactions mediated by domain-motif associations are relatively weak, with dissociation constants ($K_D$) in the micromolar range, while domain-domain binding typically results in dissociation constants values orders of magnitude smaller, indicating much stronger binding (Van Roey *et al.* 2014). While such low affinity could simply be caused by physical limitations, such as the smaller interface, adaptive hypotheses have been put forward which suggest that an overly strong affinity can have deleterious effects on the cell, for instance by compromising specificity (Haslam and Shields 2012; Karlsson *et al.* 2016). Alternatively, this low intrinsic affinity can be compensated by other factors. For example, certain protein-protein interactions depend on the simultaneous binding of many domain-motif pairs (Liao *et al.* 2020). In other cases, motifs bind outside of the canonical binding pockets of their partner domains, although this does not always lead to increased affinity (Douangamath *et al.* 2002). The regions surrounding a domain can also have an effect on motif binding preference, as inserting domains into new protein backgrounds can change the motifs that the domain binds to (Dionne *et al.* 2021, 2022).

Binding affinity and specificity have previously been explored in the context of the interaction between binding domains and motifs, by measuring the binding of mutants *in vitro* (Yu *et al.* 1994; Zarrinpar *et al.* 2003; Tonikian *et al.* 2009; Vincentelli *et al.* 2015; Kazlauskas *et al.* 2016). However, the interaction of proteins in the cellular environment is a more complex situation, and

many interactions detected *in vivo* are not detected *in vitro,* and vice versa (Kelil *et al.* 2017).
Many factors can modulate binding in the cell, including colocalization of partners, expression in
the same cell cycle phases, and contributions from sequences outside of the immediate binding
domain and motif, as well as other proteins that can interact with one or both of the partners
(Ivarsson and Jemth 2019; Dionne *et al.* 2022). Earlier studies have measured the effects of
limited numbers of domain and motif mutants *in vivo* (Zarrinpar *et al.* 2003; Marles *et al.* 2004).
More recently, techniques such as deep mutational scanning (DMS) have been used to study the
impact of large libraries of mutations on protein stability and function *in vivo* (Fowler and Fields
2014). Protein complementation assays (Tarassov *et al.* 2008; Michnick *et al.* 2016) have also
successfully been used the binding strength of mutant libraries of artificially expressed constructs
(Diss and Lehner 2018; Faure *et al.* 2022; Robles *et al.* 2023; Bendel *et al.* 2024). Other studies
have even used genome editing in yeast to endogenously tag interaction partners, allowing the
measurement of the *in vivo* binding strength of libraries of mutants (Dionne *et al.* 2021;
Dibyachintan *et al.* 2025). A recent study showed the effect on binding of combining mutations in
a PDZ domain and its binding motif (Zarin and Lehner 2024). These *in vivo* techniques could be
used to determine how a large library of mutations can affect domain-motif interaction affinity.
137
One powerful model to study domain-motif interactions is SH3 domains. These globular domains
are composed of around 60 residues, and bind different proline-rich motifs (Kaneko *et al.* 2008).
The model yeast *Saccharomyces cerevisiae*'s proteome contains 27 different SH3 domains
(Dionne *et al.* 2022), including one in the High Osmolarity Glycerol (HOG) signaling pathway
protein Sho1 (Saito and Posas 2012). The SH3 domain of Sho1 is known to interact with a motif
on the MAPKK Pbs2, and this interaction can be strengthened by exposing cells to osmotic stress
(Maeda *et al.* 1995; Posas and Saito 1997; Saito and Posas 2012) (Figure 1a). The Sho1-Pbs2
interaction is thus a powerful model to study domain-motif interaction strength, due to the potential
for increasing or decreasing affinity through mutations and the simple method of inducing an
increase in the interaction strength using osmotic stress.
148
To better understand the impact of mutations on affinity, we measured the binding strength of a
nearly complete library of mutations in the Pbs2 binding motif *in vivo* using a DHFR Protein-
fragment Complementation Assay (PCA) (Tarassov *et al.* 2008; Michnick *et al.* 2016). We
measured the binding strength between single residue Pbs2 motif mutants and Sho1, the
canonical partner of Pbs2. We find that all single mutations that increase the interaction strength
are found outside the canonical binding motif, and that certain residues outside the binding motif
interact with Sho1 outside of the canonical binding pocket, and even outside of the SH3 domain.
We use this finding to propose a model of Sho1-Pbs2 binding, where residues outside the
canonical Pbs2 motif bind an additional binding pocket on Sho1. We also compare this proposed
model to other known SH3-motif interactions involving binding outside the canonical motif.

# Material and methods

## Growth conditions

*Escherichia coli* cells were grown in 2YT medium with shaking at 37 °C for liquid cultures. *S. cerevisiae* cells were grown in YPD medium, synthetic complete (SC) medium buffered to a pH of 6.0 or PCA medium (with or without 200 μg/ml methotrexate (MTX, Bioshop Canada) diluted in dimethyl sulfoxide (DMSO, Bioshop Canada), and 2.5% noble agar for solid plates), as specified. When needed, 200 μg/mL of G418 (Bioshop Canada), 100 μg/mL of nourseothricin (Nat, Jena Bioscience) or 250 μg/mL of hygromycin B (Hyg, Bioshop Canada) were used as selection agents. Also, when specified, 1 M of sorbitol (Bioshop Canada) was added to either liquid or solid PCA medium. Liquid cultures were grown at 30 °C with agitation at 250 rpm. See Table S4 in File S2 for complete composition of all growth media used.

## Cloning

All plasmids were constructed by Gibson assembly (New England Biolabs, Inc. 2020). As template for the *PBS2* DMS library construction, a pUC19 plasmid (Addgene #50005), was amplified and digested using SacI (New England Biolabs). This backbone was used for Gibson assembly along with an insert composed of a 248 base pair region of *PBS2* amplified from genomic DNA (pUC19-Pbs2). For CRISPR-Cas9 based genome editing, the pCAS plasmid (Addgene #60847) was modified as previously described (Ryan *et al.* 2016) to change the sgRNA to target either the Sho1 binding motif of Pbs2 (pCAS-Pbs2), the SH3 domain of Sho1 (pCAS-Sho1), or the stuffer sequence which was inserted into *PBS2* (pCAS-stuffer) (Dionne *et al.* 2021). The plasmids expressing fragments of *PBS2* were based on the pGD110 plasmid (Faure *et al.* 2022). For the two shorter *PBS2* fragments, primers were used to amplify part of the plasmid, with overhangs adding the *PBS2* fragment and homology to the plasmid backbone. For the two longer *PBS2* fragments, the fragments were amplified from genomic DNA, with primers adding homology to pGD110. In all cases, the pGD110 backbone was digested with HindIII (New England Biolabs), and Gibson assembly was used to add the Pbs2 fragments in 3' of an open reading frame containing DHFR F[3] and a linker, followed by the CYC terminator. Proper plasmid assembly was confirmed using PCR and Sanger sequencing. See Table S5 in File S2 for all primers used.

## Strain construction

For all strains used see Table S6 in File S2. In all cases, proper strain construction was verified by PCR and Sanger sequencing. DHFR fragment strains were either taken from the Yeast Protein Interactome Collection (Horizon Discovery) (Tarassov *et al.* 2008), from a previously published work (Dionne *et al.* 2021), or constructed for this project using the same methods. Further deletions were done in DHFR fragment fused strains in order to knock out either *PBS2* or *SHO1*, using the *LEU2* cassette from pUG37 (Gueldener *et al.* 2002), through homologous recombination. This produced the following strains: Hog1-DHFR F[1,2]/*pbs2::LEU2*, Sho1-DHFR F[1,2]/*pbs2::LEU2, sho1::LEU2*/Pbs2-DHFR F[3] and *sho1::LEU2*/Ybt1-DHFR F[3].
This approach prevents the additional wild-type allele from influencing the measurements of interaction strength of the variants.

The Pbs2 mutants of interest were constructed from the Pbs2-DHFR F[3] fusion strain. First, the
codons 71 to 126 were replaced with a stuffer sequence (GGCGGAAGTTCTGGAGGTGGTGGT)
that translates into a flexible linker sequence (GGSSGGGG), by co-transformation of the pCAS-
Pbs2 plasmid, and of a repair template consisting of the stuffer sequence flanked by homology
arms following a previously published protocol (Ryan *et al.* 2016). This produced the Pbs2-stuffer-
DHFR F[3] strain, which was used for the construction of the individually reconstructed mutants,
for the construction of the DMS libraries (see below), and also as a control to measure Pbs2
binding to SH3-containing yeast proteins. The mutants were individually constructed by co-
transforming the Pbs2-stuffer-DHFR F[3] strain with the pCAS-stuffer plasmid targeting the
aforementioned stuffer, and synthesized oligonucleotide sequences corresponding to the desired
mutation (Integrated DNA Technologies) (Table S7 in File S2).

Three additional haploid strains were built for the extended motif DMS library (positions 85-100)
and the individually reconstructed mutants used in the validation competition assays. The first two
strains were Hog1-DHFR F[1,2]/Pbs2-stuffer-DHFR F[3] and Sho1-DHFR F[1,2]/Pbs2-stuffer-
DHFR F[3]. These were built by adding the DHFR F[1,2] to either *HOG1* or *SHO1* in the MATα
Pbs2-stuffer-DHFR F[3] strain previously constructed through homologous recombination. The
DHFR F[1,2] moieties were amplified from plasmid pAG25-DHFR F[1,2]-linker-FLAG plasmid
(Dionne *et al.* 2021). The third strain was Pbs2-stuffer, which was used to measure the
proliferation of mutants without the effect of the DHFR fragment. This was constructed in the
haploid strain BY4742, using the same strategy to insert the stuffer into *PBS2* as detailed above.
The selected Pbs2 mutants for the validation assay were constructed in these three same strains
by transformation with repair templates containing the mutations of interest, generated through
fusion PCR. Briefly, forward and reverse primers were designed for each mutant to be
constructed, which contained the mutated sequence instead of the wild-type sequence. These
primers were used for separate PCR amplification of the *PBS2* sequence along with either
common forward or reverse primers to create two overlapping fragments. These were combined
in a second PCR which fused the overlapping sequences together, to create a single fragment of
338 base pairs, which was identical to *PBS2* except for the mutated codon of interest. These
fragments were transformed into the two DHFR F[1,2]/Pbs2-stuffer-DHFR F[3] strains and Pbs2-
stuffer using the CRISPR-Cas9 strategy described above. Most mutations were constructed in all
three strains, but a certain number could only be built in one or two of the strains. The successfully
constructed mutants are listed in Table S6 in File S2. The Sho1 point mutants were constructed
similarly to the Pbs2 mutants. First, a strain was constructed where *SHO1* was replaced by the
*HPHNT1* cassette from pFA6-HPHNT1 (Janke *et al.* 2004) through homologous recombination.
In parallel, mutant *SHO1* repair templates were constructed using fusion PCR with mutated
primers, following the same strategy as the *PBS2* point mutations. This fusion PCR was amplified
from genomic DNA where the *SHO1* locus was fused with DHFR F[1,2], and so the repair
templates contained the DHFR F[1,2]. These repair templates were transformed into the
*sho1*Δ::*HPHNT1* strain along with a pCas plasmid targeting the *HPHNT1* cassette (Ryan *et al.*
2016; Dibyachintan *et al.* 2025).

For the DHFR-PCA screen on solid media, the previously constructed wild-type Pbs2-DHFR F[3]
and Pbs2-stuffer-DHFR F[3] were used. The SH3-containing proteins DHFR F[1,2] strains were
obtained from the Yeast Protein Interactome Collection (Horizon Discovery). A version of Sho1
was also built where the entire SH3 domain was replaced by the same flexible stuffer used in the
Pbs2-stuffer strain, using the same method as *PBS2*, but using the pCas-Sho1 plasmid.
256
The inducible SH3 expression strains for the growth curves with the Pbs2 fragments were
constructed for this project following a previously published protocol (Lemieux *et al.* 2024). Using
a previously constructed strain, AKD0678, which contains the GEM expression system, where
expression of the *GAL1* promoter is driven by a β-estradiol inducible artificial transcription factor
(Aranda-Díaz *et al.* 2017), and where the coding sequence of *GAL1* is replaced by the stuffer
sequence. The DHFR F[1,2] was also added after the stuffer sequence. The SH3 domains of
*SHO1*, *ABP1* and *NBP2* were amplified and transformed into the stuffer, using the same strategy
as for the *PBS2* mutants.
265
<u>Construction of DMS libraries</u>
The initial DMS library covering the surrounding region of Pbs2 (positions 71-126) was
constructed as previously reported (Dionne *et al.* 2021). A region of 56 codons (168 nucleotides)
were targeted, which correspond to codons 71 to 126 of the *YJL128C*/*PBS2* ORF. A degenerate
oligonucleotide was designed for each codon to be mutated, with the three nucleotides of the
mutated codon replaced by an NNN codon (see Table S5 in File S2 for all degenerate
oligonucleotides used). This series of 56 oligonucleotides was used to amplify pUC19-Pbs2, in a
two-step PCR procedure (Miyazaki 2011). The mutated but unligated plasmids were digested with
DpnI (New England Biolabs) and then transformed into chemocompetent *E. coli* cells.
Transformant colonies were resuspended in 5 mL of 2YT, and mutant plasmids were extracted
using a miniprep plasmid extraction kit (Presto Mini Plasmid Kit, Geneaid) to obtain a plasmid
library containing all the DMS mutants. Prepared libraries were sequenced (see DNA sequencing
section), to verify that the desired codon diversity was present. Once diversity was confirmed, the
libraries were amplified from the plasmids, and transformed into the Pbs2-stuffer-DHFR F[3] yeast
strain using the CRISPR-Cas9 strategy described in the strain construction section (Ryan *et al.*
2016). Mutants for all codons were mixed together in a masterpool, with 5 optical density units
(OD) added for each codon. Frozen glycerol stocks were prepared for the individual suspensions
of each codon, and multiple stocks were prepared for the masterpool. Genomic DNA was
extracted from the masterpool using phenol/chloroform DNA extraction (Amberg *et al.* 2005), and
as before, the library was sequenced to verify diversity. Transformed yeast libraries were stored
at -80 °C. See File S3 for complete details on library construction.
287
A second DMS library was constructed for the extended motif of *PBS2*, comprising codons 85 to
100. An oligonucleotide pool (Integrated Data Technologies, see Table S8 in File S2) was
synthesized containing all possible NNK codons (K indicating either a G or T nucleotide) for the
16 codon region of interest. This pool was integrated into 2 different yeast strains: Sho1-DHFR
F[1,2]/Pbs2-stuffer-DHFR F[3] and Hog1-DHFR F[1,2]/Pbs2-stuffer-DHFR F[3], by replacing the
Pbs2 stuffer using the CRISPR-Cas9 strategy described in the strain construction section. As with

the first library, the integrated libraries were sequenced to verify diversity. Transformed yeast
libraries were stored at -80 °C.
296
<u>DHFR-PCA competition assays</u>
The PCA selection followed a previously published protocol (Dubé *et al.* 2022). Three liquid
cultures were started, each from 100 µL of the masterpool containing all mutated codon positions,
in SC complete medium. In the screen for the DMS library of the entire surrounding region
(positions 71-126), these cultures were mated with either Sho1-DHFR F[1,2] or Hog1-DHFR
F[1,2], and diploids were selected using nourseothricin and hygromycin B. A volume equivalent
to 5 OD was spun down and the supernatant was removed, to form a cellular pellet, which was
stored at -80 °C. This is the initial time point used for sequencing. The selected diploid cultures
were diluted to 0.1 OD/mL in 15 mL of liquid PCA medium with either 200 µg/mL of methotrexate
with 1 M of sorbitol, the same concentration of methotrexate without sorbitol, or a control without
methotrexate but with 1 M sorbitol. This PCA selection was grown at 30 °C for 96 hours, except
for the control cultures which were saturated after 24 hours. After this first growth cycle, the
cultures were diluted to 0.1 OD/mL into 15 mL of fresh liquid PCA medium, and a 5 OD pellet was
spun down and stored at -80 °C. The second PCA cycle lasted the same amount of time, and at
the end, as many 5 OD pellets as possible were prepared from each culture. Genomic DNA was
extracted from the pellets of the second PCA selection culture using phenol/chloroform DNA
extraction (Amberg *et al.* 2005). The Pbs2 locus was amplified and sequenced, as described in
the DNA sequencing section.
315
316 The liquid PCA selection for the extended motif DMS library (positions 85-100) was done following
317 the same protocol as the first DHFR-PCA selection, only ignoring the mating step, as the haploid
318 strains already contained both DHFR fragments. However, a fourth control condition of PCA
319 medium without methotrexate and without sorbitol was added. Three replicate pools were done
320 for each growth condition. Frozen pellets were prepared as in the first DHFR-PCA selection.
321
The individually reconstructed validation strains were pooled by combining 100 µL of saturated
overnight culture of each mutant in YPD. 6 pools were made for each condition, three pools each
from two individual reconstructed colonies of each mutant. The same conditions were used as
the second liquid PCA selection. For the cell proliferation measurements, SC medium with and
without 1 M of sorbitol was used. Two cycles of selection were done, with 96 hour cycles for the
conditions with methotrexate and 24 hour cycles for the conditions without methotrexate. Frozen
pellets were prepared as for the previous liquid PCA selections.
329
<u>DNA sequencing and analysis</u>
DNA sequencing was done in generally the same manner for all experiments, with minor changes
in certain cases, as detailed below. DNA was extracted from libraries and frozen pellets using a
standard phenol/chloroform genomic DNA extraction protocol (Amberg *et al.* 2005). The extracted
DNA was amplified and barcoded using a Row-Column DNA barcoding strategy as previously
described (Dubé *et al.* 2022). A summary of this strategy is included in File S3. These barcoded
pools were purified on magnetic beads, and sent for sequencing on Illumina NGS machines.
Unique combinations of barcodes were used to identify the DNA extracted from each pellet in

each replicate of each condition. After preliminary analysis of the sequencing of the DNA from the
pellets of the DHFR-PCA on the extended motif DMS library (positions 85-100), it was determined
that more reads would be needed. Additional sequencing libraries were prepared fresh from the
original DNA extractions. Instead of using a row-column barcoding approach, the *PBS2* region
was amplified using primers allowing the addition of Illumina Nextera barcodes (Illumina). These
allow automated demultiplexing by the Illumina MiSeq instrument. All other steps were as
described above.
345
The variant frequency in DMS libraries during construction and in the DHFR-PCA and proliferation
screens was evaluated using custom scripts based on (Després *et al.* 2022). Python libraries
pandas 1.5.2 (The pandas development team 2023), matplotlib 3.6.2 (Hunter 2007) and numpy
1.24.1 (Harris *et al.* 2020) were used for data manipulation and visualization. Quality was
assessed using FastQC 0.12.1 (Andrews 2010). Reads were trimmed using Trimmomatic 0.39
(Bolger *et al.* 2014), then demultiplexed using bowtie 1.3.1 for the sequencing of the surrounding
region DMS library (positions 71-126) (Langmead *et al.* 2009), Interstellar 1.0 following the RCP-
PCR configuration for the first sequencing run of the extended motif DMS library (positions 85-
100) (Kijima *et al.* 2023) and cutadapt 4.7 for the validation competition assay sequencing (Martin
2011). Forward and reverse reads were then merged using the PANDAseq software (Masella *et*
*al.* 2012). Next, identical reads were grouped using vsearch (Rognes *et al.* 2016), and aligned to
the wild-type *PBS2* sequence using the needle function from the EMBOSS software (Rice *et al.*
2000). From this, the frequency of each variant was obtained.
359
From the frequency of each variant, the interaction scores and selection coefficients of each
variant were calculated using custom R 4.3.1 (R Core Team 2023) scripts. The interaction score
was calculated as the log-2-fold-change of the frequency of each variant at the end of the second
cycle of methotrexate selection and the frequency of each variant after the selection for diploids
(so immediately preceding the methotrexate selection). These log-2-fold-change scores were
rescaled for each library, so that the median of nonsense codons was equivalent to -1 and the
value of wild-type variants was 0. Variants were then filtered to keep only those that had 20 or
more reads detected after diploid selection, in order to remove bias caused by low read counts
leading to inflated scores. A unique score was calculated for each variant by taking the median
score of all codons coding for the same amino acid variant in all replicates of the same condition.
Scores were only kept for amino acid variants that had at least 3 replicates from any combination
of codons not filtered out. For the competition assay of cell proliferation, a selection coefficient
was calculated, as in (McDonald 2019), with the generations calculated as the log-2-fold change
in optical density for each growth cycle.
374
We statistically verified which nonsense and silent mutants were abnormal using the R package
rstatix (Kassambara 2021), by comparing every nonsense or silent mutation to all other nonsense
or silent mutations (Two-sided Mann-Whitney U test, $p < 0.05$). The mutants which were
significantly different were removed from the interaction score dataset. Similarly, we verified which
mutants had defects in the control condition by comparing each missense mutant to the nonsense
mutants (Two-sided Mann-Whitney U test with false discovery rate corrected p-value > 0.05). The
missense mutants which were not significantly different from the nonsense mutants were also

removed from the interaction score dataset. We also removed from the interaction score dataset
the mutants for which an abundance change was detected through gain or loss of interaction
strength between Pbs2 and Hog1, as explained in the results section (Two-sided Mann-Whitney
U test with false discovery rate corrected p-value < 0.05). For details on the visualization of these
results, see File S3.
387
<u>DHFR-PCA growth curves</u>
Growth curves were measured for individually reconstructed *PBS2* mutants in PCA medium, to
validate Pbs2 interaction strength measurements. DHFR F[3] fused strains with *PBS2* mutations
were mated with Sho1-DHFR F[1,2]/*pbs2::LEU2*, in order to produce diploids with complementary
DHFR fusions. To mate these strains, 50 µL of precultures of each strain were mixed into 900 µL
of YPD and incubated at 30 °C overnight without agitation. 2 µL of each mated culture was spotted
on solid YPD+Nat+Hyg, to select for diploids. Exponential phase precultures of diploids were
diluted to 0.1 OD/mL in 80 µL of PCA medium, in a 384 well plate, in 4 replicates, at 30 °C.
Cultures were grown either with and without 200 µg/mL of methotrexate, and with and without 1
M sorbitol. The growth curves were kept for 72 hours at 30°C. The optical density of each well
was measured every 15 minutes in a Tecan Spark plate reader (Tecan). The growth rates for
*SHO1* mutants were measured in the same way, except that the mutant strains were mated with
either Pbs2-DHFR F[3]/*sho1Δ::LEU2* or Ybt1-DHFR F[3]/*sho1Δ::LEU2*. The latter strain is used
to measure Sho1 stability and abundance, as Ybt1 interacts constitutively with Sho1. In cases
where the mutation destabilizes Sho1, the Ybt1 interaction will be affected, analogously to how
Pbs2 destabilizing mutations affect the Pbs2-Hog1 interaction. The optical density was measured
in 200 µL in a 96 well plate using an Agilent Biotek Epoch 2 plate reader (Agilent). To measure
the interaction of the *PBS2* fragments, the plasmids containing the *PBS2* fragments with the
DHFR F[3] were transformed into 3 strains containing a DHFR F[1,2] at the *GAL1* locus: Sho1-
SH3-DHFR F[1,2], Abp1-SH3-DHFR F[1,2] Nbp2-SH3-DHFR F[1,2] as well as a YFP-DHFR
F[1,2] and a control wild-type strain. Three individual colonies were taken from each
transformation as independent replicates, and grown to exponential phase. The expression of the
DHFR F[1,2] fused fragments was induced using 20 nM of estradiol, which drives expression of
the *GAL1* promoter by the GEM artificial transcription factor, and should result in the same
expression level of each construct (Aranda-Díaz *et al.* 2017). The growth curves were measured
in the same manner as the *SHO1* mutants.
414
Growth curves data was analyzed using a custom script written in R (R Core Team 2023). The
maximal growth rate was calculated using a custom function, by measuring growth over 5 time
points, and taking the 98th percentile to avoid outliers. Pbs2 mutant P94R was disregarded, as
growth of all replicates in the control condition in the absence of methotrexate was much lower
than all other samples, suggesting a problem with strain construction. For the *PBS2* fragments,
an interaction score was calculated by normalizing the growth rate using the growth rate of the
strain measuring the interaction of the same SH3 domain with a DHFR F[3] fused to no Pbs2
fragment.
423

Structure predictions and visualization
Structure predictions were done for wild-type Pbs2 and Pbs2 mutants in complex with Sho1 using
AlphaFold 2.3.2 and the AlphaFold-Multimer implementation (Jumper *et al.* 2021; Evans *et al.*
2022). Default options were used, and only the top scoring model for each mutant was relaxed.
For Sho1, the sequence of the entire protein was used for prediction (Uniprot entry P40073). For
Pbs2, residues 71 to 126 were used for the prediction (Uniprot entry P08018). Predicted
structures were visualized using ChimeraX 1.8 (Pettersen *et al.* 2021). Contacts were predicted
using the ChimeraX "contacts" tool. See File S3 for details of contact analysis.
432
DHFR-PCA on solid media against SH3 domain-containing proteins
The DHFR-PCA on solid media was based on previous work (Tarassov *et al.* 2008; Rochette *et*
*al.* 2015). All colony manipulation was done using a robotic pin tool platform (BM5-SC1, S&P
Robotics Inc.). The strains containing DHFR F[1,2] fused SH3-containing proteins were taken
from the Yeast Protein Interactome Collection (Horizon Discovery) or built for this project.
Colonies were cherry picked from the 96-well plates, and arrayed onto YPD+Nat plates for the
DHFR F[1,2] strains or YPD+Hyg plates for the DHFR F[3] strains, in a 384 colony array. The
DHFR F[1,2] strains were organized into a randomized pattern to avoid any effects caused by
neighboring colonies. The 384 colony arrays were condensed into 1536 colony arrays on
YPD+Nat or YPD+Hyg, with the randomized DHFR F[1,2] pattern repeated four times. The outer
two rows and columns of the 1536 arrays were composed of LSM8-DHFR F[1,2] and CDC39-
DHFR F[3], which grow well on PCA medium and serve to avoid any measurement bias from
being on the edge of the array, for the interactions of interest. The 1536 colony arrays were mated
together by pinning the repeated DHFR F[1,2] array onto YPD plates, and then pinning one of the
DHFR F[3] arrays onto each plate. In this collection of arrays, each interaction between an SH3
containing protein and a Pbs2 variant was measured in 5 or 6 replicates. After 48 hours of mating
at 30 °C, the mated colonies were pinned onto YPD+Nat+Hyg plates to select for diploid cells.
Two growth cycles of 48 hours at 30 °C on YPD+Nat+Hyg were done. After diploid selection, the
plates were photographed using an EOS Rebel T5i camera (Canon), to verify that all colonies
grew correctly. The mated arrays were then pinned onto solid PCA medium plates, either with or
without methotrexate, and with 1 M of sorbitol. The arrays were grown for two cycles of 48 hours
at 30 °C in a custom growth and imaging platform (S&P Robotics Inc.), which incubated the plates
and took a picture of each plate every 2 hours.
456
These plate pictures were analyzed to measure the size of every colony. First, the ImageMagick
command line tool (ImageMagick Studio LLC 2023) was used to crop the images of the selection
plates as well as change them to grayscale and invert the colors. Colony sizes were then
quantified using the Python package Pyphe (Kamrad *et al.* 2020). See File S3 for complete details
of command lines used. The colony sizes were analyzed and visualized using a custom script
written in R 4.3.1 (R Core Team 2023), based on a previous analysis (Dionne *et al.* 2021). Colony
sizes were filtered, and colonies which had not grown at the end of the diploid selection were
removed from the PCA results. Colony sizes were log2 transformed and then normalized to avoid
bias from a position in the array or from the plate they grew on. Scores were rescaled, so that the
median of the scores of the Sho1 and wild-type Pbs2 interaction was equivalent to 1 and the
median of scores of the Sho1 and Pbs2-stuffer interaction was equivalent to 0.

# Results

<u>Deep Mutational Scan of the region surrounding the Pbs2 binding motif reveals that few mutations modify binding to Sho1</u>

The interaction between Sho1 and Pbs2 is modulated by an SH3 domain on Sho1 and the canonical binding motif on Pbs2, situated in positions 93-99, and which has the sequence KPLPPLP (Maeda *et al.* 1995). Previous computational work has suggested that the region surrounding binding motifs could play an important role in modulating binding, including in SH3 interactions and in particular in the Sho1-Pbs2 interaction (Stein and Aloy 2008; Kelil *et al.* 2016). To determine which Pbs2 residues play a role in binding, we undertook a Deep Mutational Scan (DMS) on 56 codons in positions 71-126, comprising the binding motif and its surrounding region. We generated a DNA library containing all possible single codon mutations within this region and used CRISPR-Cas9 to insert it into the *PBS2* locus of *S. cerevisiae*, replacing the wild-type sequence (Dionne *et al.* 2021). The *PBS2* gene was fused with a Dihydrofolate Reductase Protein-Fragment 3 (DHFR F[3]), and these cells were mated with cells containing a DHFR F[1,2] fusion with *SHO1*, creating diploid strains with complementary DHFR fragments on both interaction partners. Sho1-Pbs2 interaction reconstitutes a functional DHFR, which enables cell division in the presence of methotrexate (MTX) (Figure 1b). Growth in the presence of MTX in a strain with both DHFR fragments thus serves as a proxy for the amount of Sho1-Pbs2 complexes forming, which depends on both the binding affinity and the local abundance of the interacting proteins (Tarassov *et al.* 2008; Freschi *et al.* 2013). Since the DHFR fusions are C-terminal, nonsense mutations in the library result in the absence of DHFR fragment expression.

To measure the interaction strength of the various mutants in the DMS library, the mutants were pooled and DHFR-PCA competition assays were done in media containing both MTX and 1 M of sorbitol, in order to induce the HOG pathway (Ferrigno *et al.* 1998). The interaction score for each variant was calculated by normalizing the log2-fold change in variant frequencies before and after selection in MTX, determined by targeted sequencing of the *PBS2* locus. Within each replicate, scores were normalized such that wild-type Pbs2 had a score of 0, and the median of nonsense mutants had a score of -1. Mutants which possess a higher interaction strength than wild-type Pbs2 therefore have a positive interaction score, while mutants which possess a lower interaction score than wild-type Pbs2 have a negative interaction score.

While many mutations within the canonical motif altered interaction strength, most mutations in the surrounding region (positions 71-126) did not affect Sho1 binding, except for a short section neighboring the canonical motif itself (Figure 2a, Figure S1 and Figure S2 in File S1, Table S1 in File S2). This finding supports the critical role of the motif and validates computational predictions that extended the binding interface to include nearby residues (Stein and Aloy 2008; Kelil *et al.* 2016). Since the effect on binding was strongly position dependent, we categorized different sections of Pbs2 as follows: the canonical binding motif is the conserved type I binding motif sequence situated in positions 93 to 99, the extended motif is composed of the canonical motif and the neighboring residues which have a strong impact on binding, from positions 85 to 99.

DHFR-PCA signal reflects both binding affinity and protein abundance. To distinguish how mutations can impact these, we assessed the effect of Pbs2 mutants on the Hog1-Pbs2 interaction, which is independent of the SH3-binding motif (Murakami *et al.* 2008), thus serving as a control for Pbs2 local abundance. Therefore, mutations solely affecting Sho1 binding should not impact the Hog1-Pbs2 DHFR-PCA, whereas mutations affecting Pbs2 stability or local abundance will influence both the Sho1-Pbs2 and Hog1-Pbs2 assays.

We found that most mutations in the surrounding region of the Pbs2 motif had little effect on Hog1 binding (Figure 2a). Furthermore, mutants that had a negative effect on Hog1-Pbs2 binding also had a negative effect on Sho1-Pbs2 binding, suggesting that the reduction in signal results from a loss of local abundance of Pbs2 (Figure S3 in File S1). However, most mutations that affected Sho1 binding did not impact the Hog1 interaction, and therefore did not change the local abundance of Pbs2.

Extended motif DMS library identifies mutations modulating interaction strength

Based on the initial results, we focused our subsequent analysis on the extended Pbs2 motif (codons 85-99), including position 100 to measure potential effects immediately downstream of the canonical motif. The initial DHFR-PCA in a diploid strain led to incomplete measurement of Sho1-Pbs2 interaction, as some Pbs2 mutants presumably interacted with the Sho1 copy not fused with DHFR F[1,2]. Consequently, we built a new haploid strain containing both DHFR F[1,2] on Sho1 and DHFR F[3] on Pbs2. We also constructed a Hog1-DHFR F[1,2]/Pbs2-DHFR F[3] haploid strain to control for Pbs2 local abundance. These new strains capture all Pbs2 binding to Sho1 or Hog1, therefore allowing for more sensitive measurements. Using the same CRISPR-Cas9 approach as before, we introduced a second Pbs2 DMS library, comprising all possible single mutations in Pbs2 positions 85 to 100, into the two haploid DHFR strains. Mutant interaction strengths were measured via DHFR-PCA as previously described (Figures S4 and S5 in File S1). We then excluded 12 Pbs2 mutants showing significantly altered Hog1 interaction scores compared to wild-type and synonymous Pbs2 mutants (two-sided Mann-Whitney U test, FDR-corrected p-value < 0.05; Figure S6 in File S1). This filtering step eliminated mutants affecting interaction through changes in Pbs2 abundance or stability, which influence both Sho1 and Hog1 interactions.

The impacts on the Sho1-Pbs2 interaction were varied (Figure 2b, Figure S7 in File S1, Table S2 in File S2). Only a small fraction of mutants exhibited interaction scores comparable to nonsense mutants, indicating that single mutations rarely abolish Sho1-Pbs2 binding completely. Comparison of missense mutant interaction scores to wild-type and synonymous variants revealed that many mutants displayed significantly stronger or weaker interactions (two-sided Mann-Whitney U test, FDR-corrected p < 0.05). Specifically, 18 mutants (6.5% of missense) showed significantly increased interaction, while 173 mutants (62.7% of missense) showed significantly decreased interaction compared to wild-type Pbs2. The impact of mutants that reduced the interaction scores was generally greater.

Given that osmotic stress enhances the Sho1-Pbs2 interaction (Maeda *et al.* 1995; Posas and
Saito 1997; Saito and Posas 2012), we compared interaction strengths of Pbs2 mutants with and
without 1 M sorbitol (Figure S8 in File S1). A strong correlation was observed between interaction
scores under both conditions (Spearman's rho 0.98, p-value < $2.2 \times 10^{-16}$). As expected due to
HOG pathway induction, the effects on binding were stronger in the presence of sorbitol. Under
osmotic stress, strongly interacting Pbs2 mutants are even more often in contact with Sho1,
increasing the number of reconstituted DHFR complexes, and further outcompeting weakly
interacting mutants. Nevertheless, the high correlation suggests that Pbs2 mutants exhibit similar
relative binding changes regardless of osmotic stress, and the fundamental impact of mutations
is independent of HOG pathway activation.

To validate the pooled measurements, we individually reconstructed 24 mutants spanning the
range of measured interaction scores in Pbs2-DHFR F[3] strains and mated them with the Sho1-
DHFR F[1,2] strain. Growth rates of these diploids were then measured in DHFR-PCA medium.
As with the competition assay, the growth rate is a proxy for interaction strength, which reflects
affinity in the absence of a change in abundance. The growth curve results correlate strongly with
the interaction scores (Spearman's rho 0.86, p-value $2.4 \times 10^{-6}$ with 1 M sorbitol, spearman's rho
of 0.8, p-value $4.3 \times 10^{-6}$ without sorbitol), confirming the competition assay results (Figure S9a in
File S1). To further validate the pooled data, we performed a smaller-scale competition DHFR-
PCA, using 67 individually reconstructed mutants with diverse interaction scores (Figure S10 in
File S1, Table S3 in File S2).This validation assay showed a strong correlation with the DMS
library measurements (Spearman's rho 0.98, p-value < $2.2 \times 10^{-16}$ with 1 M sorbitol, spearman's
rho of 0.98, p-value < $2.2 \times 10^{-16}$ without sorbitol), thus validating the DMS library results (Figure
S9b in File S1). To assess the direct impact of mutations on cell proliferation, we constructed
strains with the same 67 mutations but lacking DHFR fragments. Cell growth and division thus
only depend on the effect of the mutations, and not interaction strength. We then performed a
pooled competitive growth assay in SC medium with 1 M sorbitol to measure the proliferation of
each mutant. With the exception of nonsense mutations, no significant effect on cell proliferation
was observed (Welch's t-test false discovery rate corrected p-value < 0.05) (Figure S11 in File
S1, Table S3 in File S2). Notably, the strongest interacting mutant, I87W, shows a detectable,
though not statistically significant, deleterious effect on cell proliferation.

<u>Structure prediction and Sho1 mutations suggest a secondary contact between extended motif</u>
<u>and Sho1</u>

While the crystal structure of the Sho1 SH3 domain bound to a 9-residue Pbs2 segment shows
the canonical motif (positions 93-99) occupying the entire canonical binding pocket (Kursula *et al.*
2008), our data reveals that many mutations outside this motif strongly affect binding as well. We
hypothesized that residues in the extended Pbs2 motif (positions 85-99) might interact with Sho1
outside its canonical SH3 domain pocket. This is supported by NMR structures and studies of
other SH3 domains showing that residues outside the canonical motif can modulate interaction
strength and specificity (Rickles *et al.* 1995; Feng *et al.* 1995; Stollar *et al.* 2009; Takaku *et al.*
2010; Gorelik and Davidson 2012). Given the absence of experimental structures for full-length
Sho1 or Pbs2, we used AlphaFold-Multimer predictions (Jumper *et al.* 2021; Evans *et al.* 2022)

to explore potential interactions beyond the canonical binding site (Figure S12 in File S1).
Modeling the entire Sho1 protein along with the surrounding region of the Pbs2 motif (positions
71-126), revealed potential interactions outside of the canonical binding pocket of Sho1, mediated
by residues neighboring the canonical binding motif of Pbs2 (Figure 3a). The predicted structure
shows a hydrophobic pocket formed by the Sho1 SH3 domain and a nearby loop, interacting with
an alpha helix formed by Pbs2 residues 84-92 (Figure 3b). Interestingly, all but one of the mutants
in the extended motif which significantly increase interaction strength are located within this
predicted Pbs2 helix. Although the model also predicted proximity of some C-terminal Pbs2
residues to Sho1, mutations in these positions had minimal impact on binding (Figure 2a), leading
us to focus on residues within and near the extended motif.
607
To understand how specific mutations alter binding, we predicted the structures of the five Pbs2
variants exhibiting the strongest Sho1 interaction, using the same method as for wild-type Pbs2
(Figure S12 in File S1). In particular, the strongest interacting mutant, I87W, is predicted to
position its tryptophan side chain in close proximity (1.77 Å) to the non-canonical hydrophobic
pocket, specifically within a sub-pocket formed by both SH3 and non-SH3 Sho1 residues.
Furthermore, it is predicted to form a hydrogen bond with D333 of Sho1 (Figure 3c). These two
factors potentially explain its strong interaction strength relative to all other Pbs2 variants.
Similarly, the strongly interacting mutants H86W, V91L, and V91M are all predicted to place
hydrophobic side chains within the non-canonical hydrophobic pocket, potentially contributing to
stronger interactions. For the remaining mutant, Q89D, the hydrophilic aspartate side chain is
predicted to face away from the hydrophobic pocket, thus not disrupting these interactions.
However, the mechanism of its increased binding strength remains unclear (Figure S13 in File
S1).
621
We hypothesized that replacing hydrophobic Pbs2 residues predicted to contact the hydrophobic
non-canonical pocket with non-hydrophobic ones would disrupt this interaction and weaken
binding. Consistent with this, analysis of mutations within the predicted alpha helix shows that
replacing the initially hydrophobic residues at positions 87, 90, and 91 with non-hydrophobic
residues reduces binding strength. Generally, substituting these positions with other hydrophobic
residues had a less detrimental effect, and for V91, many hydrophobic substitutions even
increased interaction strength (Figure 3d). These results support the prediction that the side
chains of residues at positions 87, 90, and 91 are located within the non-canonical hydrophobic
binding pocket. This finding helps explain the strong interactions strength of Pbs2 mutant I87W,
by showing the important role of hydrophobic interactions in position 87.
632
We further explored the presence of contacts between Pbs2 and surfaces outside the Sho1
canonical binding pocket. We introduced a series of Sho1 mutations in residues predicted to be
in proximity to Pbs2, in strains containing the DHFR fragments, to measure whether these
mutations would disrupt binding. We also built a Sho1 variant where the SH3 domain was
replaced by a flexible, neutral "stuffer" sequence (GGSSGGGG), to measure the interaction
strength without the contribution of the SH3 domain. We then assessed the effect of the mutations
on interaction strength using growth assays. To control for Sho1 stability and abundance, similar
to our approach with Hog1 for Pbs2 mutants, we measured the interaction of the Sho1 mutants

with Ybt1, a protein previously found to interact with Sho1 independently from the SH3 domain
(Dionne *et al.* 2021). We compared mutant growth rates to wild-type controls, looking for Sho1
variants that diminished Pbs2 binding but not Ybt1 binding (One-sided Student's t-test, FDR-
corrected p-value < 0.05, Figure S14 in File S1).
645
Contrary to our expectation, Ybt1 is not an ideal control, as the Sho1-Ybt1 interaction is weakened
by the absence of the SH3 domain, contrary to a previous report (Dionne *et al.* 2021). Also, certain
mutations seemingly increase the interaction strength between Sho1 and Ybt1, which could be
due to stabilization of Sho1, an increase in affinity with Ybt1 or a change in binding equilibrium
with Ybt1 caused by reduced binding to Pbs2. Still, in cases where the interaction of a Sho1
variant with Ybt1 is not significantly weaker than the interaction of wild-type Sho1 with Ybt1, we
assumed that the local abundance of Sho1 was not diminished. Two mutations show a clear
decrease in Pbs2 interaction, but no decrease in Ybt1 interaction: G287A and I350W (Figure 3e).
The first mutation is outside the SH3 domain, while the second is in the domain, but not in the
canonical binding site. Both mutated residues are predicted to be within 5 Å of the Pbs2 extended
motif. The G287 backbone is notably predicted to form a hydrogen bond with Pbs2 residue Q88,
located in the predicted alpha helix. Together, our structural predictions and interaction
measurements indicate that residues outside the canonical Pbs2 motif could directly interact with
residues both within and without the Sho1 SH3 domain, thereby modulating binding affinity.
660
661    <u>Comparison of Sho1-Pbs2 binding to other SH3-motif pairs</u>
662
Given the proposed role of the extended Pbs2 motif and the additional Sho1 binding pocket,
composed of the SH3 domain and a nearby loop, in modulating interaction strength, we
investigated if these features also contribute to the specificity of Pbs2 binding to Sho1. We
reasoned that if this additional pocket were unique among yeast SH3 domains, it could enhance
binding specificity, as the Pbs2 extended motif might be specifically adapted to this feature. First,
we performed a multiple sequence alignment of yeast SH3 domains, including 25 flanking
residues on each side, which encompasses all Sho1 residues predicted to be within 5 Å of Pbs2.
Apart from key conserved residues within the SH3 domain itself, this analysis revealed no
significant conservation of residues forming the additional Sho1 pocket (Figure S15a in File S1).
To assess potential structural conservation despite sequence divergence, we conducted a
multiple structural alignment using MUSTANG (Konagurthu *et al.* 2006) on AlphaFold-predicted
structures (Varadi *et al.* 2022) of yeast SH3 domains, with 25 neighboring residues. This analysis
showed structural conservation only within the SH3 domains, with no significant structural
conservation in the flanking regions, including those forming the additional Sho1 binding pocket
(Figure S15b in File S1). Therefore, we conclude that the additional binding pocket on Sho1
represents a unique structural feature among yeast SH3 domains.
679
Despite the lack of conserved structure of non-SH3 sequences, there are reports of three other
yeast SH3 domains binding extended motifs: Abp1, the second SH3 domain of Bem1 (referred to
as Bem1-2) and Nbp2 (Stollar *et al.* 2009; Takaku *et al.* 2010; Gorelik and Davidson 2012). We
investigated whether these SH3 domains interact with the extended Pbs2 motif *in vivo* using a
DHFR-PCA assay that measures colony growth on solid medium (Tarassov *et al.* 2008).The SH3-

containing proteins were fused with DHFR F[1,2] and Pbs2 with DHFR F[3]. As a control, we used
a Pbs2 variant where the extended motif (positions 85-99) and surrounding region (codons 71-
126, the same positions as the initial DMS library) were replaced by the neutral, flexible "stuffer"
peptide which was previously used to replace the Sho1 SH3 domain (GGSSGGGG). The
interaction strength of the SH3 domains was also measured against this Pbs2-stuffer variant.
690
Abp1 and Nbp2 interacted with Pbs2 at a similar strength to Sho1, while Bem1-2 showed no
detectable interaction (Figure 4a). However, Abp1 and Nbp2 also interacted with Pbs2-stuffer at
comparable levels to wild-type Pbs2, indicating that their binding is independent of the Sho1-
binding motif. Nbp2 is known to bind a second, distal motif on Pbs2 through its SH3 domain
(Mapes and Ota 2004), while Abp1 likely binds Pbs2 through an alternative, unknown site.
Screening all other known yeast SH3-containing proteins revealed significantly weaker
interactions with both Pbs2 variants compared to wild-type Sho1 (one-sided Mann-Whitney U test,
FDR-corrected p-value < 0.05; Figure S16 in File S1). Furthermore, wild-type Pbs2 showed a
stronger interaction with a Sho1 variant where the SH3 domain was replaced by the
aforementioned stuffer (Sho1-stuffer), than Pbs2-stuffer with Sho1-stuffer, confirming the role of
the non-SH3 portion of Sho1 in binding the extended motif of Pbs2 (Figure 4a). Given that Sho1-
stuffer abundance is similar to wild-type Sho1 (Dionne et al. 2021), these results demonstrate that
the extended Pbs2 motif specifically binds to Sho1 through both its SH3 and non-SH3 regions.
704
To further test whether the Pbs2 motif and surrounding region can bind other SH3 domains, we
used DHFR-PCA to measure the interaction of Pbs2 fragments of different lengths with the SH3
domains of Abp1, Nbp2 and Sho1. We expressed the SH3 domains at the same level to remove
bias resulting from the different cellular abundances of Abp1, Nbp2 and Sho1. We also measured
the interaction of the same Pbs2 fragments against a constitutively expressed YFP-DHFR F[1,2]
to control for proper expression of the Pbs2 fragments (Lemieux *et al.* 2025). The canonical Pbs2
motif (positions 93-99) showed a significantly stronger interaction (One-sided Student's t-test,
FDR-corrected p-value < 0.05) with the Sho1 SH3 domain compared to the Nbp2 SH3 domain
(Figure S17 in File S1). This preference for the Sho1 SH3 domain was maintained and the
interaction strength increased with two longer Pbs2 fragments: positions 85-100 (the region of the
second DMS screen) and positions 69-124 (corresponding roughly to the Pbs2-stuffer
replacement), which also showed significantly stronger binding to the Sho1 SH3 domain
compared to the Abp1 and Nbp2 SH3 domains. While interaction scores with the non-Sho1 SH3
domains increased with Pbs2 fragment length, they were always weaker than the Sho1
interaction. This confirms the role of the extended motif and residues even more distant from the
canonical site in maintaining binding specificity to the Sho1 SH3 domain.
721
To understand the basis of Pbs2 binding specificity to Sho1, we compared the predicted Sho1-
Pbs2 structure with NMR structures of Abp1-Ark1, Bem1-2-Ste20, and Nbp2-Ste20, all involving
peptides of similar length to the Pbs2 extended motif. The structures of the four SH3 domains and
their bound peptides showed a conserved binding mode, with each peptide occupying the
canonical SH3 domain pocket and extending towards the region corresponding to the additional
Sho1 binding pocket (Figure 4b, Figure S18 in File S1). However, Abp1 and Nbp2 appear to
interact with their respective extended motifs solely through the SH3 domain surface. In contrast,

Bem1-2 utilizes a second, non-SH3 interface that forms a pocket with its SH3 domain,
accommodating the extended motif, similar to Sho1. This Bem1-2 interface, identified as the
Cdc42-interacting (CI) region (Takaku *et al.* 2010), exhibits no structural or sequence similarity to
the corresponding region in Sho1. Furthermore, unlike Sho1, this additional binding region in
Bem1-2 is located C-terminal to its SH3 domain.
734
Beyond the conserved canonical motif positions, the extended motifs of Pbs2, Ark1, and Ste20
show limited sequence conservation. These extended motifs can be aligned using a previously
developed standard numbering system where the first conserved proline is assigned position 0,
preceding residues are numbered negatively, and following residues positively (Figure 4c) (Lim
*et al.* 1994). The binding specificity exhibited by Pbs2 may depend on the possible contacts
between the extended motif and the different SH3 domains, as well as the non-SH3 surfaces in
the cases of Sho1 and Bem1-2. By examining the interactions of these extended motifs with their
respective partners, particularly focusing on residues within the extended motif but outside the
canonical motif (positions 85-92), we aimed to identify determinants of Pbs2 binding specificity.
744
Three scenarios can differentiate Pbs2-Sho1 binding from the binding of other extended motifs to
their partners: either contacts are present in Pbs2-Sho1 but absent in other pairs, contacts are
absent in Pbs2-Sho1 but present in other pairs, or equivalent residues in extended motifs contact
different residues of their interaction partner. We find examples of all three scenarios. For
example, Pbs2 is predicted to contact Sho1 through a histidine residue in position -10, while Ark1,
for example, does not have contacts so far from the canonical motif (Figure S19a in File S1). Pbs2
is also missing certain interactions that are present in other pairs. For instance, in both Ste20 and
Ark1, the residue at position -7 is crucial for binding, as in both cases, mutation of the residue to
an alanine reduced binding affinity by at least an order magnitude, and this mutation had the
largest impact of all alanine substitutions outside the canonical motif (Stollar *et al.* 2009; Gorelik
and Davidson 2012). In contrast, position -7 in Pbs2 is not predicted to interact with Sho1, though
mutation to a hydrophilic residue increased interaction strength (Figure S19b in File S1, Table S2
in File S2). Finally, the same position in different extended motifs can form distinct interactions
with their partners. In position -6, the isoleucine in Pbs2 is predicted to contact Y319, W338 and
I350 on the SH3 domain. In contrast, the histidine in position -6 of Ark1 forms a salt bridge with
an aspartic acid on a loop of the Abp1 SH3 domain (Figure S19c in File S1). The structural
alignment of SH3 domains indicates that Sho1 has a gap at the position of the aspartic acid that
forms the salt bridge (Figure S15b in File S1, position 112). Even beyond these limited examples,
the four extended motifs show differences in how they bind their partners. This suggests that the
non-conserved regions within extended motifs enable a much greater degree of binding
differentiation than the conserved canonical motif alone.

# Discussion

Binding motifs face the double challenge of binding their partner with appropriate strength while avoiding spurious interactions with non-partner proteins. To understand this balance in the Sho1-Pbs2 interaction, we used deep mutational scanning of the Pbs2 motif and its flanking regions to quantify the impact of single residue mutations on Sho1 binding. Using structure predictions and mutations in both Pbs2 and Sho1, we conclude that binding occurs between Sho1 and Pbs2 outside of the canonical motif and the SH3 domain. We then compare the predicted Sho1-Pbs2 complex to existing structures of other SH3 domain-peptide complexes. Based on these comparisons, we propose several mechanisms by which the Pbs2 extended motif achieves specific binding to Sho1 over other SH3 domains.

One important consideration in domain-peptide interactions is the potential trade-off between interaction affinity and specificity. Intuitively, a domain that can strongly bind its target motif might also exhibit increased unwanted interactions with similar motifs within the same family, due to their physical and chemical resemblance. This idea is supported by previous studies on SH2 and PDZ domains, which, like SH3 domains, bind short motifs. These studies have suggested that mutants exhibiting higher affinity for their target tend to display lower specificity, implying a potential incompatibility between high affinity and high specificity (Ernst *et al.* 2010; Haslam and Shields 2012; Kaneko *et al.* 2012; Karlsson *et al.* 2016). However, a computational study of binding motifs, including SH3 binding motifs, proposed that increased affinity can correlate with increased specificity (Kelil *et al.* 2016). Other works have emphasized the importance of the residues surrounding the canonical binding motif in determining both affinity and specificity (Li 2005; Ivarsson and Jemth 2019). In fact, one study computed that nearly 30% of binding energy from the binding motif was contributed by residues outside the canonical motif, and suggested that the residues outside the canonical motif mostly play a role in determining specificity (Stein and Aloy 2008). Notably, our deep mutational scanning revealed that all mutations which significantly increase the interaction strength between Sho1 and Pbs2 are situated outside of the canonical motif. This suggests that the wild-type canonical motif might already be optimized for maximal binding strength. Therefore, we propose a model where the canonical motif residues in positions 93 to 99 are optimized and thus structurally constrained for affinity, while the neighboring residues are less constrained and able to modulate affinity and specificity when mutated.

We show that the modified binding strengths from mutations outside the Pbs2 canonical motif arise from interactions with surfaces beyond the SH3 domain's canonical pocket, even involving non-SH3 surfaces on Sho1. Non-binding pocket interactions have previously been observed in multiple SH3-mediated interactions (Lee *et al.* 1996; Dalgarno *et al.* 1997; Li 2005; Stollar *et al.* 2009; Takaku *et al.* 2010; Gorelik and Davidson 2012; Gaussmann *et al.* 2024), and we find that at least one mutation outside the Sho1 SH3 domain can disrupt binding, suggesting that the mutated residue plays a role in the interaction. Our AlphaFold-Multimer predictions support this, revealing a hydrophobic pocket formed by the Sho1 SH3 domain and a non-SH3 loop, which residues of the Pbs2 extended motif can occupy. While the disordered nature of the Pbs2 flanking region suggests a dynamic interaction, these models provide a plausible mechanism for non-

canonical motif-mediated binding. By removing the need for affinity to be dependent only on
canonical binding motifs, the residues that surround motifs could bind to protein surfaces which
are less conserved among SH3 domains, and therefore easier to differentiate. The presence of
less conserved components, such as extended motifs or non-canonical binding pockets could
thus contribute to specificity among domains of the same family. The Sho1 hydrophobic pocket,
formed by both SH3 and non-SH3 elements, exemplifies a unique structural feature that might
differentiate Sho1, enabling Pbs2 to achieve higher affinity without necessarily increasing binding
to other SH3 domains. This binding model, involving a larger number of residues in tuning affinity
and specificity, could potentially uncouple affinity and specificity, avoiding a trade-off between the
two. While affinity and specificity might be intrinsically coupled within the highly conserved
canonical binding interface, the less constrained surrounding residues offer a greater capacity for
independent variation of these traits.
826
SH3 domain-mediated interactions are known to be context-dependent, influenced not solely by
the domain itself but also by the surrounding protein environment. This context includes the non-
SH3 regions of the protein, interactions with other proteins, and cellular localization (Dionne *et al.*
2022). Consequently, an SH3 domain's interaction profile is not solely determined by its amino
acid sequence. Indeed, studies show that relocating an SH3 domain to a different protein or
expressing it in isolation significantly alters its interaction profile (Dionne *et al.* 2021; Lemieux *et*
*al.* 2024; Dibyachintan *et al.* 2025). The observed effects of Pbs2 mutations on binding strength
may be mediated not only through direct SH3 domain interactions but also within the broader
Sho1 protein context. The interaction of the Pbs2 extended motif with non-SH3 Sho1 residues
likely contributes to this context-dependent modulation. Other causes could include
stereochemical adjustments of SH3 binding loops or changes affecting Sho1's interactions with
other binding partners. Other proteins involved in the osmotic stress response, namely Ste11 and
Ste50, have been shown to interact with the Sho1 SH3 domain outside the canonical binding
pocket (Zarrinpar *et al.* 2004; Tatebayashi *et al.* 2006). Mutations in the Pbs2 peptide could
potentially influence the stability of these interactions as well.
842
Previous studies have investigated the binding of Pbs2 motif variants (Zarrinpar *et al.* 2003), and
our findings corroborate some of their observations. For instance, we also find that mutations near
the canonical Pbs2 motif can either increase or decrease binding strength. This study also
reported reduced fitness for a promiscuous Pbs2 double mutant (Zarrinpar *et al.* 2003). In
contrast, our analysis revealed no substantial fitness loss for single mutants, except for the I87W
variant, which exhibited a noticeable, though not statistically significant, decrease in fitness
(Figure S11 in File S1). Interestingly, I87W is also the mutant showing the greatest increase in
interaction strength, potentially explaining its slight fitness cost. It has previously been suggested
that affinity increasing mutants may have deleterious effects on cell homeostasis, by reducing the
dynamicity of the pathway, and impeding dissociation of the partners (Wang *et al.* 2018).
Consistent with this, prior work on the Sho1-Pbs2 interaction demonstrated that a Sho1 SH3
domain mutation that increased affinity also led to a minor reduction in cell growth (Marles *et al.*
2004), indicating that the wild-type affinity may be already optimized, and that stronger binding
may disrupt signaling.
857

In conclusion, we show that mutations in positions neighboring the canonical binding motif of Pbs2 can both increase and decrease binding strength with Sho1. We propose that these non-canonical Pbs2 residues can modulate binding affinity through interactions outside the conserved SH3 binding pocket. We hypothesize that Sho1 is similar to certain other yeast SH3 domains which bind longer motifs through additional binding pockets (Stollar *et al.* 2009; Takaku *et al.* 2010; Gorelik and Davidson 2012), and propose mechanisms through which an extended binding motif could better differentiate binding to different SH3 domains. Moving forward, studies of SH3-motif interactions should encompass a larger range of residues, considering both the SH3 domain surface beyond the canonical pocket and the regions surrounding the motif. Our findings, along with previous research, highlight that *in vitro* studies using isolated domains and minimal peptides may not fully represent the complexity of domain-motif interactions within their native protein context. By understanding and harnessing the potential of extended motif binding, researchers may be better able to understand the impact of motifs in cell signaling and disease, and better able to use motif binding in technical and medical applications.

# Data availability

Strains and plasmids are available upon request. All analysis scripts and visualization scripts are available on Github at the following address: https://github.com/Landrylab/Jordan_et_al_2024. All data, including demultiplexed sequencing data, results from the DHFR-PCA experiments, growth curve data, predicted structures, protein alignments, all files and information necessary to run the scripts, and supplementary Files S1 to S3 are available in the following Dryad repository: https://doi.org/10.5061/dryad.79cnp5j3z. The Supplementary Files are also available in the following Zenodo repository : https://doi.org/10.5281/zenodo.15721636. Results from the DHFR-PCA experiments on mutant libraries are also available as Tables S1 to S3 in File S2. File S1 contains Supplementary Figures S1 to S19, File S2 contains Supplementary Tables S1 to S10, and File S3 contains the Supplementary Methods.

# Acknowledgments

The authors thank Philippe C. Després, Moïra Dion and Dan Evans-Yamamoto for help with sequencing data analysis. We also thank all members of the LandryLab team for thoughtful comments on the experiments and on figure conception. All molecular graphics were performed using UCSF ChimeraX 1.8. The protein structure predictions were enabled by computing resources provided by the Digital Research Alliance of Canada.

# Funding

This work was supported by grants from the Canadian Institutes of Health Research numbers 387697 and 525827, and Human Frontier Science Program research grant RGP34/2018 to C.R.L. C.R.L. holds the Canada Research Chair in Cellular Systems and Synthetic Biology. D.J. was supported by the Canada Graduate Scholarship - Master's program from NSERC, a graduate scholarship from PROTEO, a graduate scholarship from the NSERC CREATE program EvoFunPath, a Postgraduate Scholarship - Doctoral from NSERC and a Doctoral scholarship

899 from FRQS. D.B. was funded by the EMBO Long-Term Fellowship (LTF) (ALTF 1069-
900 2019). U.D. was funded by an NSERC graduate fellowship.

# Author contributions

902 D.J., A.K.D., U.G. , D.B., and C.R.L. conceived and designed the study. C.R.L. supervised the
903 work and obtained funding. D.J. and A.K.D. built the strains and performed the experiments. D.J.
904 performed the computational analyses. D.J. wrote the first draft of the manuscript. D.J., A.K.D.,
905 U.G. , D.B., and C.R.L. contributed to revising the final version of the manuscript.
906

# Literature cited

908
Amberg D., D. C. Burke, and J. N. Strathern, 2005 *Methods in yeast genetics: a Cold Spring*
*Harbor Laboratory course manual.* Cold Spring Harbor Laboratory Press, Cold Spring
Harbor, N.Y.

Andrews S., 2010 FastQC: a quality control tool for high throughput sequence data

Aranda-Díaz A., K. Mace, I. Zuleta, P. Harrigan, and H. El-Samad, 2017 Robust synthetic
circuits for two-dimensional control of gene expression in yeast. ACS Synth. Biol. 6:
545–554.

Bendel A. M., A. J. Faure, D. Klein, K. Shimada, R. Lyautey, *et al.*, 2024 The genetic
architecture of protein interaction affinity and specificity. Nat. Commun. 15: 8868.

Bolger A. M., M. Lohse, and B. Usadel, 2014 Trimmomatic: a flexible trimmer for Illumina
sequence data. Bioinformatics 30: 2114–2120.

Dalgarno D. C., M. C. Botfield, and R. J. Rickles, 1997 SH3 domains and drug design: Ligands,
structure, and biological function. Pept. Sci. 43: 383–400.

Davey N. E., G. Travé, and T. J. Gibson, 2011 How viruses hijack cell regulation. Trends
Biochem. Sci. 36: 159–169.

Delplace V., A. Ortin-Martinez, E. L. S. Tsai, A. N. Amin, V. Wallace, *et al.*, 2019 Controlled
release strategy designed for intravitreal protein delivery to the retina. J. Control.
Release 293: 10–20.

Després P. C., A. F. Cisneros, E. M. M. Alexander, R. Sonigara, C. Gagné-Thivierge, *et al.*,
2022 Asymmetrical dose responses shape the evolutionary trade-off between antifungal
resistance and nutrient use. Nat Ecol Evol 6: 1501–1515.

Dibyachintan S., A. K. Dubé, D. Bradley, P. Lemieux, U. Dionne, *et al.*, 2025 Cryptic genetic
variation shapes the fate of gene duplicates in a protein interaction network. Nat.
Commun. 16: 1530.

Dionne U., É. Bourgault, A. K. Dubé, D. Bradley, F. J. M. Chartier, *et al.*, 2021 Protein context
shapes the specificity of SH3 domain-mediated interactions in vivo. Nat. Commun. 12:
1597.

Dionne U., L. J. Percival, F. J. M. Chartier, C. R. Landry, and N. Bisson, 2022 SRC homology 3

domains: multifaceted binding modules. Trends Biochem. Sci. 47: 772–784.

Diss G., and B. Lehner, 2018 The genetic landscape of a physical interaction. Elife 7.
https://doi.org/10.7554/eLife.32472

Douangamath A., F. V. Filipp, A. T. J. Klein, P. Barnett, P. Zou, *et al.*, 2002 Topography for
Independent Binding of α-Helical and PPII-Helical Ligands to a Peroxisomal SH3
Domain. Mol. Cell 10: 1007–1017.

Dubé A. K., R. Dandage, S. Dibyachintan, U. Dionne, P. C. Després, *et al.*, 2022 Deep
Mutational Scanning of Protein–Protein Interactions Between Partners Expressed from
Their Endogenous Loci In Vivo, pp. 237–259 in *Yeast Functional Genomics: Methods*
*and Protocols*, edited by Devaux F. Springer US, New York, NY.

Ernst A., D. Gfeller, Z. Kan, S. Seshagiri, P. M. Kim, *et al.*, 2010 Coevolution of PDZ domain-
ligand interactions analyzed by high-throughput phage display and deep sequencing.
Mol. Biosyst. 6: 1782–1790.

Evans R., M. O'Neill, A. Pritzel, N. Antropova, A. Senior, *et al.*, 2022 Protein complex prediction
with AlphaFold-Multimer. bioRxiv 2021.10.04.463034.

Faure A. J., J. Domingo, J. M. Schmiedel, C. Hidalgo-Carcedo, G. Diss, *et al.*, 2022 Mapping
the energetic and allosteric landscapes of protein binding domains. Nature 604: 175–
183.

Feng S., C. Kasahara, R. J. Rickles, and S. L. Schreiber, 1995 Specific interactions outside the
proline-rich core of two classes of Src homology 3 ligands. Proc. Natl. Acad. Sci. U. S. A.
92: 12408–12415.

Ferrigno P., F. Posas, D. Koepp, H. Saito, and P. A. Silver, 1998 Regulated nucleo/cytoplasmic
exchange of HOG1 MAPK requires the importin beta homologs NMD5 and XPO1.
EMBO J. 17: 5606–5614.

Fowler D. M., and S. Fields, 2014 Deep mutational scanning: a new style of protein science.
Nat. Methods 11: 801–807.

Freschi L., F. Torres-Quiroz, A. K. Dubé, and C. R. Landry, 2013 qPCA: a scalable assay to
measure the perturbation of protein-protein interactions in living cells. Mol. Biosyst. 9:
36–43.

Gaussmann S., R. Peschel, J. Ott, K. M. Zak, J. Sastre, *et al.*, 2024 Modulation of peroxisomal
import by the PEX13 SH3 domain and a proximal FxxxF binding motif. Nat. Commun.
15: 3317.

Gorelik M., and A. R. Davidson, 2012 Distinct peptide binding specificities of Src homology 3
(SH3) protein domains can be determined by modulation of local energetics across the
binding interface. J. Biol. Chem. 287: 9168–9177.

Gouw M., S. Michael, H. Sámano-Sánchez, M. Kumar, A. Zeke, *et al.*, 2018 The eukaryotic
linear motif resource – 2018 update. Nucleic Acids Res. 46: D428–D434.

Grant B. J., A. P. C. Rodrigues, K. M. ElSawy, J. A. McCammon, and L. S. D. Caves, 2006
Bio3d: an R package for the comparative analysis of protein structures. Bioinformatics

22: 2695–2696.

Gueldener U., J. Heinisch, G. J. Koehler, D. Voss, and J. H. Hegemann, 2002 A second set of
loxP marker cassettes for Cre-mediated multiple gene knockouts in budding yeast.
Nucleic Acids Res. 30: e23.

Harris C. R., K. J. Millman, S. J. van der Walt, R. Gommers, P. Virtanen, *et al.*, 2020 Array
programming with NumPy. Nature 585: 357–362.

Haslam N. J., and D. C. Shields, 2012 Peptide-Binding Domains: Are Limp Handshakes Safest?
Sci. Signal. 5: e40-pe40.

Hunter J. D., 2007 Matplotlib: A 2D Graphics Environment. Comput. Sci. Eng. 9: 90–95.

ImageMagick Studio LLC, 2023 *ImageMagick*.

Ivarsson Y., and P. Jemth, 2019 Affinity and specificity of motif-based protein-protein
interactions. Curr. Opin. Struct. Biol. 54: 26–33.

Janke C., M. M. Magiera, N. Rathfelder, C. Taxis, S. Reber, *et al.*, 2004 A versatile toolbox for
PCR-based tagging of yeast genes: new fluorescent proteins, more markers and
promoter substitution cassettes. Yeast 21: 947–962.

Jumper J., R. Evans, A. Pritzel, T. Green, M. Figurnov, *et al.*, 2021 Highly accurate protein
structure prediction with AlphaFold. Nature 596: 583–589.

Kamrad S., M. Rodríguez-López, C. Cotobal, C. Correia-Melo, M. Ralser, *et al.*, 2020 Pyphe, a
python toolbox for assessing microbial growth and cell viability in high-throughput colony
screens. Elife 9. https://doi.org/10.7554/eLife.55160

Kaneko T., L. Li, and S. S.-C. Li, 2008 The SH3 domain--a family of versatile peptide- and
protein-recognition module. Front. Biosci. 13: 4938–4952.

Kaneko T., H. Huang, X. Cao, X. Li, C. Li, *et al.*, 2012 Superbinder SH2 Domains Act as
Antagonists of Cell Signaling. Sci. Signal. 5: ra68–ra68.

Karlsson O. A., G. N. Sundell, E. Andersson, Y. Ivarsson, and P. Jemth, 2016 Improved affinity
at the cost of decreased specificity: a recurring theme in PDZ-peptide interactions. Sci.
Rep. 6: 34269.

Kassambara A., 2021 rstatix: Pipe-Friendly Framework for Basic Statistical Tests

Kazlauskas A., C. Schmotz, T. Kesti, J. Hepojoki, I. Kleino, *et al.*, 2016 Large-Scale Screening
of Preferred Interactions of Human Src Homology-3 (SH3) Domains Using Native Target
Proteins as Affinity Ligands. Mol. Cell. Proteomics 15: 3270–3281.

Kelil A., E. D. Levy, and S. W. Michnick, 2016 Evolution of domain–peptide interactions to
coadapt specificity and affinity to functional diversity. Proc. Natl. Acad. Sci. U. S. A. 113:
E3862–E3871.

Kelil A., B. Dubreuil, E. D. Levy, and S. W. Michnick, 2017 Exhaustive search of linear
information encoding protein-peptide recognition. PLoS Comput. Biol. 13: e1005499.

Kijima Y., D. Evans-Yamamoto, H. Toyoshima, and N. Yachie, 2023 A universal sequencing read interpreter. Sci Adv 9: eadd2793.

Konagurthu A. S., J. C. Whisstock, P. J. Stuckey, and A. M. Lesk, 2006 MUSTANG: a multiple structural alignment algorithm. Proteins 64: 559–574.

Kumar M., S. Michael, J. Alvarado-Valverde, A. Zeke, T. Lazar, *et al.*, 2024 ELM-the Eukaryotic Linear Motif resource-2024 update. Nucleic Acids Res. 52: D442–D455.

Kursula P., I. Kursula, Y. H. Song, K. Paraskevopoulos, and M. Wilmanns, 2008 YEAST SHO1 SH3 DOMAIN COMPLEXED WITH A PEPTIDE FROM PBS2

Langmead B., C. Trapnell, M. Pop, and S. L. Salzberg, 2009 Ultrafast and memory-efficient alignment of short DNA sequences to the human genome. Genome Biol. 10: R25.

Lee C. H., K. Saksela, U. A. Mirza, B. T. Chait, and J. Kuriyan, 1996 Crystal structure of the conserved core of HIV-1 Nef complexed with a Src family SH3 domain. Cell 85: 931–942.

Lemieux P., D. Bradley, A. K. Dubé, U. Dionne, and C. R. Landry, 2024 Dissection of the role of a Src homology 3 domain in the evolution of binding preference of paralogous proteins. Genetics 226. https://doi.org/10.1093/genetics/iyad175

Lemieux P., A. K. Dube, and C. R. Landry, 2025 A protein-fragment complementation assay to quantify synthetic protein scaffold efficiency. bioRxiv 2025.03.19.640584.

Li S. S.-C., 2005 Specificity and versatility of SH3 and other proline-recognition domains: structural basis and implications for cellular signal transduction. Biochem. J 390: 641–653.

Liao T.-J., H. Jang, R. Nussinov, and D. Fushman, 2020 High-affinity interactions of the nSH3/cSH3 domains of Grb2 with the C-terminal proline-rich domain of SOS1. J. Am. Chem. Soc. 142: 3401–3411.

Lim W. A., F. M. Richards, and R. O. Fox, 1994 Structural determinants of peptide-binding orientation and of sequence specificity in SH3 domains. Nature 372: 375–379.

Maeda T., M. Takekawa, and H. Saito, 1995 Activation of Yeast PBS2 MAPKK by MAPKKKs or by Binding of an SH3-Containing Osmosensor. Science 269: 554–558.

Mapes J., and I. M. Ota, 2004 Nbp2 targets the Ptc1-type 2C Ser/Thr phosphatase to the HOG MAPK pathway. EMBO J. 23: 302–311.

Marles J. A., S. Dahesh, J. Haynes, B. J. Andrews, and A. R. Davidson, 2004 Protein-Protein Interaction Affinity Plays a Crucial Role in Controlling the Sho1p-Mediated Signal Transduction Pathway in Yeast. Mol. Cell 14: 813–823.

Martin M., 2011 Cutadapt removes adapter sequences from high-throughput sequencing reads. EMBnet.journal 17: 10–12.

Masella A. P., A. K. Bartram, J. M. Truszkowski, D. G. Brown, and J. D. Neufeld, 2012 PANDAseq: paired-end assembler for illumina sequences. BMC Bioinformatics 13: 31.

McDonald M. J., 2019 Microbial Experimental Evolution - a proving ground for evolutionary
theory and a tool for discovery. EMBO Rep. 20: e46992.

Michnick S. W., E. D. Levy, C. R. Landry, J. Kowarzyk, and V. Messier, 2016 The Dihydrofolate
Reductase Protein-Fragment Complementation Assay: A Survival-Selection Assay for
Large-Scale Analysis of Protein–Protein Interactions. Cold Spring Harb. Protoc. 2016:
db.prot090027.

Miyazaki K., 2011 Chapter seventeen - MEGAWHOP Cloning: A Method of Creating Random
Mutagenesis Libraries via Megaprimer PCR of Whole Plasmids, pp. 399–406 in *Methods*
*in Enzymology*, Synthetic Biology, Part B. edited by Voigt C. Academic Press.

Monera O. D., T. J. Sereda, N. E. Zhou, C. M. Kay, and R. S. Hodges, 1995 Relationship of
sidechain hydrophobicity and alpha-helical propensity on the stability of the single-
stranded amphipathic alpha-helix. J. Pept. Sci. 1: 319–329.

Murakami Y., K. Tatebayashi, and H. Saito, 2008 Two adjacent docking sites in the yeast Hog1
mitogen-activated protein (MAP) kinase differentially interact with the Pbs2 MAP kinase
and the Ptp2 protein tyrosine phosphatase. Mol. Cell. Biol. 28: 2481–2494.

New England Biolabs, Inc., 2020 *Gibson Assembly Master Mix Instruction Manual*.

Pawson T., M. Raina, and P. Nash, 2002 Interaction domains: from simple binding events to
complex cellular behavior. FEBS Lett. 513: 2–10.

Pettersen E. F., T. D. Goddard, C. C. Huang, E. C. Meng, G. S. Couch, *et al.*, 2021 UCSF
ChimeraX: Structure visualization for researchers, educators, and developers. Protein
Sci. 30: 70–82.

Posas F., and H. Saito, 1997 Osmotic Activation of the HOG MAPK Pathway via Ste11p
MAPKKK: Scaffold Role of Pbs2p MAPKK. Science 276: 1702–1705.

R Core Team, 2023 *R: A Language and Environment for Statistical Computing*. R Foundation
for Statistical Computing, Vienna, Austria.

Rice P., I. Longden, and A. Bleasby, 2000 EMBOSS: the European Molecular Biology Open
Software Suite. Trends Genet. 16: 276–277.

Rickles R. J., M. C. Botfield, X. M. Zhou, P. A. Henry, J. S. Brugge, *et al.*, 1995 Phage display
selection of ligand residues important for Src homology 3 domain binding specificity.
Proc. Natl. Acad. Sci. U. S. A. 92: 10909–10913.

Robles J. T., H. J. Lou, G. Shi, P. L. Pan, and B. E. Turk, 2023 Linear motif specificity in
signaling through p38α and ERK2 mitogen–activated protein kinases. Proceedings of the
National Academy of Sciences 120: e2316599120.

Rochette S., G. Diss, M. Filteau, J.-B. Leducq, A. K. Dubé, *et al.*, 2015 Genome-wide protein-
protein interaction screening by protein-fragment complementation assay (PCA) in living
cells. J. Vis. Exp. https://doi.org/10.3791/52255

Rognes T., T. Flouri, B. Nichols, C. Quince, and F. Mahé, 2016 VSEARCH: a versatile open
source tool for metagenomics. PeerJ 4: e2584.

Ryan O. W., S. Poddar, and J. H. D. Cate, 2016 CRISPR-Cas9 Genome Engineering in
1088               Saccharomyces cerevisiae Cells. Cold Spring Harb. Protoc. 2016.
1089               https://doi.org/10.1101/pdb.prot086827

1090    Saito H., and F. Posas, 2012 Response to Hyperosmotic Stress. Genetics 192: 289–318.

1091    Stein A., and P. Aloy, 2008 Contextual specificity in peptide-mediated protein interactions. PLoS
1092               One 3: e2524.

1093    Stollar E. J., B. Garcia, P. A. Chong, A. Rath, H. Lin, *et al.*, 2009 Structural, functional, and
1094               bioinformatic studies demonstrate the crucial role of an extended peptide binding site for
1095               the SH3 domain of yeast Abp1p. J. Biol. Chem. 284: 26918–26927.

1096    Takaku T., K. Ogura, H. Kumeta, N. Yoshida, and F. Inagaki, 2010 Solution structure of a novel
1097               Cdc42 binding module of Bem1 and its interaction with Ste20 and Cdc42. J. Biol. Chem.
1098               285: 19346–19353.

1099    Tarassov K., V. Messier, C. R. Landry, S. Radinovic, M. M. S. Molina, *et al.*, 2008 An in Vivo
1100               Map of the Yeast Protein Interactome. Science 320: 1465–1470.

1101    Tatebayashi K., K. Yamamoto, K. Tanaka, T. Tomida, T. Maruoka, *et al.*, 2006 Adaptor
1102               functions of Cdc42, Ste50, and Sho1 in the yeast osmoregulatory HOG MAPK pathway.
1103               EMBO J. 25: 3033–3044.

1104    The pandas development team, 2023 *pandas-dev/pandas: Pandas*.

1105    Tonikian R., X. Xin, C. P. Toret, D. Gfeller, C. Landgraf, *et al.*, 2009 Bayesian Modeling of the
1106               Yeast SH3 Domain Interactome Predicts Spatiotemporal Dynamics of Endocytosis
1107               Proteins. PLoS Biol. 7: e1000218.

1108    Van Roey K., B. Uyar, R. J. Weatheritt, H. Dinkel, M. Seiler, *et al.*, 2014 Short Linear Motifs:
1109               Ubiquitous and Functionally Diverse Protein Interaction Modules Directing Cell
1110               Regulation. Chem. Rev. 114: 6733–6778.

1111    Varadi M., S. Anyango, M. Deshpande, S. Nair, C. Natassia, *et al.*, 2022 AlphaFold Protein
1112               Structure Database: massively expanding the structural coverage of protein-sequence
1113               space with high-accuracy models. Nucleic Acids Res. 50: D439–D444.

1114    Vincentelli R., K. Luck, J. Poirson, J. Polanowska, J. Abdat, *et al.*, 2015 Quantifying domain-
1115               ligand affinities and specificities by high-throughput holdup assay. Nat. Methods 12:
1116               787–793.

1117    Wang C.-H., P. Mehta, and C. J. Bashor, 2018 The strength of protein-protein interactions
1118               controls the information capacity and dynamical response of signaling networks. bioRxiv
1119               469197.

1120    Yu H., J. K. Chen, S. Feng, D. C. Dalgarno, A. W. Brauer, *et al.*, 1994 Structural basis for the
1121               binding of proline-rich peptides to SH3 domains. Cell 76: 933–945.

1122    Zarin T., and B. Lehner, 2024 A complete map of specificity encoding for a partially fuzzy
1123               protein interaction. bioRxiv 2024.04.25.591103.

Zarrinpar A., S.-H. Park, and W. A. Lim, 2003 Optimization of specificity in a cellular protein

interaction network by negative selection. Nature 426: 676–680.

Zarrinpar A., R. P. Bhattacharyya, M. P. Nittler, and W. A. Lim, 2004 Sho1 and Pbs2 Act as
Coscaffolds Linking Components in the Yeast High Osmolarity MAP Kinase Pathway.
Mol. Cell 14: 825–832.

Zhou L., T. Feng, S. Xu, F. Gao, T. T. Lam, *et al.*, 2022 Ggmsa: A visual exploration tool for
multiple sequence alignment and associated data. Brief. Bioinform. 23: bbac222.

1168
1169
1170

# Figures

1172
1173
Figure 1. The Sho1-Pbs2 interaction can be strengthened using sorbitol exposure, and the DHFR-
PCA can be used to measure the interaction strength between Sho1 and Pbs2.
a) Simplified schematic of the HOG pathway showing Sho1, Pbs2, and Hog1 interactions under
normal conditions (left) where they are largely separate, and during osmotic stress (e.g., with
sorbitol, right) where pathway induction leads to complex formation. The resulting Hog1
phosphorylation leads to nuclear translocation and gene expression modulation. b) Schematic of
the DHFR-PCA screen using a DMS *PBS2* mutant library. Single residue variants of *PBS2* were
inserted into the yeast genome at the native locus using CRISPR-Cas9 genome editing. Sho1
and Pbs2 were fused with complementary DHFR fragments. In cases where the Pbs2 variant
does not interact with Sho1, the DHFR fragments remain separated. Upon interaction between a
Pbs2 variant and Sho1, the DHFR fragments combine to reconstitute the functional DHFR, and
allow growth in the presence of methotrexate (MTX). More strongly interacting partners allow the
cell to proliferate more rapidly (Freschi *et al.* 2013).

Figure 2. <u>Mutations in Pbs2 within and around the binding motif affect binding to Sho1</u>.
a) Interaction scores (normalized log2-fold change after MTX selection) of Pbs2 mutations across
the region surrounding the SH3 binding motif (positions 71-126), measured for interaction with
Sho1 and Hog1. Wild-type Pbs2 residues are indicated above each position, with the extended
motif (positions 85-99) highlighted in blue and the canonical motif (positions 93-99) residues in
bold. Boxplots show the distribution of interaction scores for mutants at all positions (3-18
replicates per mutant; nonsense mutants excluded). b) Distribution of interaction scores for single
amino acid mutants (3-18 replicates) within the Pbs2 extended motif DMS library (positions 85-
100) when interacting with Sho1. Mutations are displayed on the y-axis, colored by type:
nonsense (stop codon), WT or silent (synonymous), and missense (amino acid change).
Missense mutants were categorized based on Mann-Whitney U test (FDR-corrected $p < 0.05$)
against WT and silent mutations: Significantly weaker (negative score), Not significantly different,
or Significantly stronger (positive score).
1202
1203

Figure 3. <u>Contacts mediated by positions outside the canonical binding motif of Pbs2 could</u>
<u>modulate binding</u>
a) Predicted structure of Sho1 in complex with Pbs2 residues 71-126, oriented to highlight the
Pbs2 helix interacting with a hydrophobic pocket on Sho1, alongside the canonical motif (positions
93-99) in its binding pocket. For Sho1, the SH3 domain spanning positions 303 to 360 is colored
in green, while the rest of the protein is in yellow. For Pbs2, the extended motif spanning positions
85 to 99 is in blue while the rest of Pbs2 is in dark red. b) Close-up view of the interaction between
Sho1 and the extended motif (positions 85-99) of Pbs2. Only residues outside the canonical motif
of Pbs2 which are in contact with Sho1 are shown, and the Sho1 residues with which they are in
contact are also shown. The coloring of the residues is as in panel a, with Sho1 residues inside
the SH3 domain in green and Sho1 residues outside the SH3 domain in yellow. c) Close-up view
of the predicted Sho1-Pbs2(I87W) complex, showing the W87 side chain and a predicted
hydrogen bond (red dotted line) with Sho1 D333. All other Sho1 residues with which the I87W
residue is predicted to form contacts are shown with their sidechains. Coloring as in a). d) Boxplots
showing interaction scores of Pbs2 mutants at selected positions (from DMS DHFR-PCA of the
extended motif), grouped by the hydrophobicity of the substituted residue. Wild-type Pbs2 residue
at each position is indicated and colored by its hydrophobicity (Hydrophobic: F, I, W, L, V, M, Y,
C, A; Neutral: T, H, E, S, Q; Hydrophilic: R, K, N, G, P, D (Monera *et al.* 1995)). Nonsense mutants
are excluded. e) Interaction scores of select Sho1 variants with either Pbs2 or Ybt1, derived from
growth rates from individual DHFR-PCA assays. Scores for individual replicates shown as points.
The median wild-type Sho1 interaction score with Pbs2 (purple) or Ybt1 (orange) is shown with
dashed lines. Stuffer indicates a Sho1 mutant where the SH3 domain was replaced by a neutral,
flexible peptide (GGSSGGGG). The scores for all Sho1 variants are shown in Figure S14 in File
S1.

**c**

| Canonical motif | | | | | | | | | | | | | | | | | | | | | |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| Standard numbering | -14 | -13 | -12 | -11 | -10 | -9 | -8 | -7 | -6 | -5 | -4 | -3 | -2 | -1 | 0 | 1 | 2 | 3 | 4 | 5 | 6 | 7 |
| Pbs2 | Q | S | P | K | H | I | Q | Q | I | V | N | K | P | L | P | P | L | P | V | A | G | S |
| Ark1 | | | | K | P | K | L | H | S | P | K | P | P | P | T | P | K | T | K | K | A |
| Ste20 | S | S | S | A | N | G | K | F | I | P | S | R | P | A | P | K | P | P | S | S | A |
| Pbs2 position | 82 | 83 | 84 | 85 | 86 | 87 | 88 | 89 | 90 | 91 | 92 | 93 | 94 | 95 | 96 | 97 | 98 | 99 | 100 | 101 | 102 | 103 |

1247

1248

<u>Figure 4. The Pbs2 motif binds specifically to Sho1, despite similarities of other SH3 domains and</u>

<u>motifs</u>

a) Interaction score of wild-type Pbs2 (green) or Pbs2-stuffer (white) with select yeast SH3-

containing proteins. All replicates shown as points. Scores are normalized relative to wild-type

Sho1 interactions, with wild-type Pbs2 median at 1 and Pbs2-stuffer median at 0. The scores for

all SH3-containing yeast proteins are found in Figure S16 in File S1. b) Root mean square

deviation (RMSD) of atom locations between matching backbone atoms in each pair of structures,

as calculated using the R package bio3d (Grant *et al.* 2006). c) Sequences of Ste20, Ark1 and

Pbs2, with standard numbering according to the motif, to compare equivalent positions (Lim *et al.*

1994). Ste20 and Ark1 positions included are those present in the structures used to generate in

panel b, with Pbs2 positions overlapping all represented positions.

1260