Residues Neighboring an SH3-Binding Motif Participate in the Interaction *In Vivo*

David F. Jordan[1,2,3,4,*], Alexandre K. Dubé[1,2,3,4,5], Ugo Dionne[2,3,4,6,7,8], David Bradley[1,2,3,4,5] and Christian R. Landry [1,2,3,4,5,*]

1. Département de Biochimie, microbiologie et bio-informatique, Université Laval, 1045 Avenue de la Médecine, Québec, QC, Canada, G1V 0A6
2. Institut de Biologie Intégrative et des Systèmes (IBIS), 1030, Avenue de la Médecine, Université Laval, Québec, QC, Canada, G1V 0A6
3. PROTEO-Regroupement Québécois de Recherche sur la Fonction, l'Ingénierie et les Applications des Protéines, Université Laval, 1030, Avenue de la Médecine Québec, QC, Canada, G1V 0A6
4. Centre de Recherche en Données Massives de l'Université Laval, Université Laval, 1065, Avenue de la Médecine, Québec, QC, Canada, G1V 0A6
5. Département de Biologie, Université Laval, 1045 Avenue de la Médecine, Québec, QC, Canada, G1V 0A6
6. Centre de Recherche du Centre Hospitalier Universitaire (CHU) de Québec-Université Laval, 11, côte du Palais, Québec, QC, Canada, G1R 2J6
7. Centre de Recherche sur le Cancer de l'Université Laval, 9, rue McMahon, Québec, QC, Canada, G1R 3S3
8. Present address: Lunenfeld-Tanenbaum Research Institute, Sinai Health, 600 University Avenue, Toronto, ON, Canada, M5G 1X5

* Correspondence: 1030, Avenue de la Médecine, Université Laval, Québec, QC, Canada, G1V 0A6 Email: david.jordan.2@ulaval.ca. 1030, Avenue de la Médecine, Université Laval, Québec, QC, Canada, G1V 0A6 Email: christian.landry@bio.ulaval.ca

Running head: Residues neighboring motif interaction
Keywords: Short Linear motif, Src Homology 3 (SH3) domain, Deep mutational scanning, Binding affinity, Structure prediction

Abstract

In signaling networks, protein-protein interactions are often mediated by modular domains that bind short linear motifs. The motifs' sequences affect many factors, among them affinity and specificity, or the ability to bind strongly and to the appropriate partners. Using Deep Mutational Scanning to create a mutant library, and protein complementation assays to measure protein-protein interactions, we determined the *in vivo* binding strength of a library of mutants of a binding motif on the MAP kinase kinase Pbs2, which binds the SH3 domain of the osmosensor protein Sho1 in *Saccharomyces cerevisiae*. These measurements were made using the full-length endogenous proteins, in their native cellular environment. We find that along with residues within the canonical motif, many mutations in the residues neighboring the motif also modulate binding strength. Interestingly, all Pbs2 mutations which increase affinity are situated outside of the Pbs2 region that interacts with the canonical SH3 binding pocket, suggesting that other surfaces on Sho1 contribute to binding. We use predicted structures to hypothesize a model of binding which involves residues neighboring the canonical Pbs2 motif binding outside of the canonical SH3 binding pocket. We compared this predicted structure with known structures of SH3 domains binding peptides through residues outside of the motif, and put forth possible mechanisms through which Pbs2 can bind specifically to Sho1. We propose that for certain SH3 domain-motif pairs, affinity and specificity are determined by a broader range of sequences than what has previously been considered, potentially allowing easier differentiation between otherwise similar partners.

Summary

Protein-protein interactions are often mediated by a binding domain on one protein and a short disordered binding motif on another protein. We measured the binding strength of a mutant library of a binding motif situated in the yeast protein Pbs2 to the SH3 domain of Sho1. Many mutations in the residues neighboring the binding motif affect binding. A protein structure prediction of the interaction partners shows that residues neighboring the motif bind residues outside the known binding pocket on the SH3 domain. The Sho1-Pbs2 interaction differs enough from other known SH3-motif pairs to allow specific binding.

Introduction

Cells possess complex and robust signaling networks that detect stimuli and trigger responses. These signaling networks are composed of a series of protein-protein interactions, which are often mediated by modular interaction domains. Many classes of protein interaction domains are shared across pathways and species, yet fulfill different roles and functions (Pawson *et al.* 2002). The different constraints placed on interaction domains by their distinct roles contribute to explaining the divergence in sequence between homologs of the same domain (Ernst *et al.* 2010; Dionne *et al.* 2022). Therefore, to understand how domain sequences have evolved and continue to evolve, the phenotypic consequences of mutations must be understood. Two of the major phenotypes for protein-protein interaction domains are affinity, that is, the strength of binding to the partner protein, and specificity, that is, the ability to bind to the appropriate partner proteins, and to not form spurious interactions with other proteins in the cellular environment (Ivarsson and Jemth 2019).

Many interaction domains bind short, intrinsically disordered stretches of their interaction partners, known as short linear motifs (SLiMs) or simply binding motifs (Gouw *et al.* 2018). Binding motifs are involved in a wide swath of cell signaling pathways, and underlie many important mechanisms in human and other cells (Kumar *et al.* 2024). Uncovering the determinants of binding can elucidate the functioning of human cells and certain diseases that affect them, such as various cancers in which motif-mediated signaling plays a part (Van Roey *et al.* 2014). Additionally, viruses also express their own binding motifs to disrupt the signaling machinery of their hosts (Davey *et al.* 2011). Understanding domain-motif binding can also allow them to be used as tools, for example by attaching therapeutic proteins to domains and motifs to hydrogels, to ensure the gradual release of the therapeutic proteins in human tissues (Delplace *et al.* 2019).

Interactions mediated by domain-motif associations are relatively weak, with dissociation constants (K_D) in the micromolar range, while domain-domain binding typically results in dissociation constants values orders of magnitude smaller, indicating much stronger binding (Van Roey *et al.* 2014). While such low affinity could simply be caused by physical limitations, such as the smaller interface, adaptive hypotheses have been put forward which suggest that an overly strong affinity can have deleterious effects on the cell, for instance by compromising specificity (Haslam and Shields 2012; Karlsson *et al.* 2016). Alternatively, this low intrinsic affinity can be compensated by other factors. For example, certain protein-protein interactions depend on the simultaneous binding of many domain-motif pairs (Liao *et al.* 2020). In other cases, motifs bind outside of the canonical binding pockets of their partner domains, although this does not always lead to increased affinity (Douangamath *et al.* 2002). The regions surrounding a domain can also have an effect on motif binding preference, as inserting domains into new protein backgrounds can change the motifs that the domain binds to (Dionne *et al.* 2021, 2022).

Binding affinity and specificity have previously been explored in the context of the interaction between binding domains and motifs, by measuring the binding of mutants *in vitro* (Yu *et al.* 1994; Zarrinpar *et al.* 2003; Tonikian *et al.* 2009; Vincentelli *et al.* 2015; Kazlauskas *et al.* 2016).

However, the interaction of proteins in the cellular environment is a more complex situation, and
many interactions detected *in vivo* are not detected *in vitro,* and vice versa (Kelil *et al.* 2017).
Many factors can modulate binding in the cell, including colocalization of partners, expression in
the same cell cycle phases, and contributions from sequences outside of the immediate binding
domain and motif, as well as other proteins that can interact with one or both of the partners
(Ivarsson and Jemth 2019; Dionne *et al.* 2022). Earlier studies have measured the effects of
limited numbers of domain and motif mutants *in vivo* (Zarrinpar *et al.* 2003; Marles *et al.* 2004).
More recently, techniques such as deep mutational scanning (DMS) have been used to study the
impact of large libraries of mutations on protein stability and function *in vivo* (Fowler and Fields
2014). Protein complementation assays (Tarassov *et al.* 2008; Michnick *et al.* 2016) have also
successfully been used to measure the *in vivo* binding strength of libraries of mutants in yeast
(Diss and Lehner 2018; Dionne *et al.* 2021; Faure *et al.* 2022; Robles *et al.* 2023; Bendel *et al.*
2024; Dibyachintan *et al.* 2025). For example, a recent study showed the effect on binding of
combining mutations in a PDZ domain and its binding motif (Zarin and Lehner 2024). These *in*
vivo techniques could be used to determine how a large library of mutations can affect domain-
motif interaction affinity.
138
SH3 domains are powerful models to study domain-motif interactions. These globular domains
are composed of around 60 residues, and bind different proline-rich motifs (Kaneko *et al.* 2008).
The model yeast *Saccharomyces cerevisiae*'s proteome contains 27 different SH3 domains
(Dionne *et al.* 2022), including one in the High Osmolarity Glycerol (HOG) signaling pathway
protein Sho1 (Saito and Posas 2012). The SH3 domain of Sho1 is known to interact with a motif
on the MAPKK Pbs2, and this interaction can be strengthened by exposing cells to osmotic stress
(Maeda *et al.* 1995; Posas and Saito 1997; Saito and Posas 2012) (Figure 1a). The Sho1-Pbs2
interaction is thus a powerful model to study domain-motif interaction strength, due to the potential
for increasing or decreasing affinity through mutations and the simple method of inducing an
increase in the interaction strength using osmotic stress.
149
To better understand the impact of mutations on affinity, we measured the binding strength of a
nearly complete library of mutations in the Pbs2 binding motif *in vivo* using a DHFR Protein-
fragment Complementation Assay (PCA) (Tarassov *et al.* 2008; Michnick *et al.* 2016). We
measured the binding strength between single residue Pbs2 motif mutants and Sho1, the
canonical partner of Pbs2. We find that all single mutations that increase the interaction strength
are found outside the canonical binding motif, and that certain residues outside the binding motif
are predicted to interact with Sho1 outside of the canonical binding pocket, and even outside of
the SH3 domain. We use this finding to propose a model of Sho1-Pbs2 binding, where residues
outside the canonical Pbs2 motif bind an additional binding pocket on Sho1. We also compare
this proposed model to other known SH3-motif interactions involving binding outside the canonical
motif.

Material and methods

<u>Growth conditions</u>

Escherichia coli cells were grown in 2YT medium with shaking at 37 °C for liquid cultures. *Saccharomyces cerevisiae* cells were grown in YPD medium, synthetic complete (SC) medium buffered to a pH of 6.0 or PCA medium (with or without 200 µg/ml methotrexate (MTX, Bioshop Canada) diluted in dimethyl sulfoxide (DMSO, Bioshop Canada), and 2.5% noble agar for solid plates), as specified. When specified, 200 µg/mL of G418 (Bioshop Canada), 100 µg/mL of nourseothricin (Nat, Jena Bioscience) or 250 µg/mL of hygromycin B (Hyg, Bioshop Canada) were used as selection agents. Also, when specified, 1 M of sorbitol (Bioshop Canada) was added to either liquid or solid PCA medium. Liquid cultures were grown at 30 °C with agitation at 250 rpm. See Table S4 in File S2 for complete composition of all growth media used.

<u>Cloning</u>

All plasmids were constructed by Gibson assembly. As template for the *PBS2* DMS library construction, a pUC19 plasmid (Addgene #50005), was used for Gibson assembly with an insert composed of a 248 base pair region of *PBS2* amplified from genomic DNA (pUC19-Pbs2). For CRISPR-Cas9 based genome editing, the pCAS plasmid (Addgene #60847) was modified as previously described (Ryan *et al.* 2016) to change the sgRNA to target either the Sho1 binding motif of Pbs2 (pCAS-Pbs2), the SH3 domain of Sho1 (pCAS-Sho1), or the stuffer sequence which was inserted into *PBS2* (pCAS-stuffer) (Dionne *et al.* 2021). The plasmids expressing fragments of Pbs2 were based on the pGD110 plasmid (Faure *et al.* 2022). Gibson cloning was used to add the Pbs2 fragments in 3' of an open reading frame containing DHFR F[3] and a linker, followed by the CYC terminator. See File S3 for details, and see Table S5 in File S2 for all primers used.

<u>Strain construction</u>

For all strains used see Table S6 in File S2, and see File S3 for all strain construction details. DHFR fragment strains were either taken from the Yeast Protein Interactome Collection (Horizon Discovery) (Tarassov *et al.* 2008), from a previously published work (Dionne *et al.* 2021), or constructed for this project using the same methods. Further deletions were done in certain DHFR fragment fused strains in order to knock out either *PBS2* or *SHO1*, using the *LEU2* cassette from pUG37 (Gueldener *et al.* 2002), through homologous recombination. This prevents the additional wild-type allele from influencing the measurements of interaction strength of the variants.

PBS2 and *SHO1* were modified by replacing either the region surrounding the binding motif (codons 71-126) or the SH3 domain and 3' sequence (codons 303-367) with a stuffer sequence (GGCGGAAGTTCTGGAGGTGGTGGT) that translates into a flexible linker sequence (GGSSGGGG), using CRISPR-Cas9 based genome editing (Biot-Pelletier and Martin 2016; Ryan *et al.* 2016). These stuffer strains were used in experiments, and also as targets into which mutant sequences were transformed using CRISPR-Cas9 based genome editing. This approach was used for the construction of the DMS libraries, as well as the reconstruction of individual mutants. For the validation growth curves of individual *PBS2* mutants, synthesized oligonucleotide sequences corresponding to the desired mutation were transformed (Integrated DNA

Technologies, see Table S7 in File S2). For individually reconstructed *SHO1* mutants, the *SHO1*
locus was deleted by the *HPHNT1* cassette from pFA6-HPHNT1 (Janke *et al.* 2004). Fusion PCR
was used to construct repair templates of mutant sequences of *SHO1*-DHFR F[1,2], and
transformed into the *SHO1* locus using CRISPR-Cas9 based genome editing (Ryan *et al.* 2016).
The inducible SH3 expression strains for the growth curves DHFR-PCA with the *PBS2* fragments
were constructed for this project following a previously published protocol (Lemieux *et al.* 2024),
using a previously constructed strain, AKD0678, which contains the GEM expression system,
where expression of the *GAL1* promoter is driven by a β-estradiol inducible artificial transcription
factor (Aranda-Díaz *et al.* 2017), and where the coding sequence of *GAL1* is replaced by the
stuffer sequence with as DHFR F[1,2] fusion.
219
<u>Construction of DMS libraries</u>
The initial DMS library covering the surrounding region of Pbs2 was constructed as previously
reported (Dionne *et al.* 2021). 56 codons (168 nucleotides) were targeted, which correspond to
codons 71 to 126 of the *YJL128C*/*PBS2* ORF. A degenerate oligonucleotide was designed for
each codon to be mutated, with the three nucleotides of the mutated codon replaced by an NNN
codon (see Table S5 in File S2 for all primers used). This series of 56 oligonucleotides was used
to amplify pUC19-Pbs2, in a two-step PCR procedure (Miyazaki 2011). The mutated but unligated
plasmids were transformed into chemocompetent *E. coli* cells. Transformant colonies were
resuspended in liquid media, and mutant plasmids were extracted using a miniprep plasmid
extraction kit (Presto Mini Plasmid Kit, Geneaid) to obtain a plasmid library containing all the DMS
mutants. Prepared libraries were sequenced (see DNA sequencing section), to verify that the
desired codon diversity was present. Once diversity was confirmed, the libraries were amplified
from the plasmids, and transformed into the Pbs2-stuffer-DHFR F[3] yeast strain using the
CRISPR-Cas9 strategy described in the strain construction section. Mutants for all codons were
mixed together in a master pool. Genomic DNA was extracted from the master pool using
phenol/chloroform DNA extraction (Amberg *et al.* 2005), and as before, the library was sequenced
to verify diversity. At this point, three codons had unsatisfactory diversity. A new template was
created for codons 103 and 119 using fusion PCR. For codon 88, a new template was ordered
as an oligonucleotide of 503 base pairs (Twist Bioscience), around codon 88, which is replaced
by an NNN codon.
240
A second DMS library was constructed for the extended motif of *PBS2*, comprising codons 85 to
100. An oligonucleotide pool (Integrated Data Technologies, see Table S8 in File S2) was
synthesized containing all possible NNK codons (K indicating either a G or T nucleotide) for the
16 codon region of interest. This pool was integrated into 2 different yeast strains: Sho1-DHFR
F[1,2]/Pbs2-stuffer-DHFR F[3] and Hog1-DHFR F[1,2]/Pbs2-stuffer-DHFR F[3], by replacing the
Pbs2 stuffer using the CRISPR-Cas9 strategy described in the strain construction section. As
before, the integrated libraries were sequenced to verify diversity.
248
The selected Pbs2 mutants for the validation competition assay were constructed in Sho1-DHFR
F[1,2]/Pbs2-stuffer-DHFR F[3], Hog1-DHFR F[1,2]/Pbs2-stuffer-DHFR F[3], and Pbs2-stuffer with
no DHFR fragments, using the CRISPR-Cas9 strategy described above. The mutant templates
were constructed through fusion PCR (see primers used in Table S5 in File S2). Most mutations

were constructed in all three DHFR strains, but a certain number could only be built in one or two
of the strains. The successfully constructed mutants are listed in Table S6 in File S2.
255

<u>DHFR-PCA competition assays</u>
The PCA selection followed a previously published protocol (Dubé *et al.* 2022), and all details can
be found in File S3. Three liquid cultures were started, each from 100 µL of the master pool
containing all mutated codon positions, in SC complete medium. These cultures were mated with
either Sho1-DHFR F[1,2] or Hog1-DHFR F[1,2], and diploids were selected using Nat and Hyg.
A volume equivalent to 5 OD was spun down and the supernatant was removed, to form a cellular
pellet, which was stored at -80 °C. This is the initial time point used for sequencing. The selected
diploid cultures were diluted to 0.1 OD/mL in 15 mL of liquid PCA medium with either 200 µg/mL
of methotrexate with 1 M of sorbitol, the same concentration of methotrexate without sorbitol, or
a control without methotrexate but with 1 M sorbitol. This PCA selection was grown at 30 °C for
96 hours, except for the control cultures which were saturated after 24 hours. After this first growth
cycle, the cultures were diluted 0.1 OD/mL into 15 mL of fresh liquid PCA medium, and a 5 OD
pellet was spun down and stored at -80 °C. The second PCA cycle lasted the same amount of
time, and at the end, as many 5 OD pellets as possible were prepared from each culture. Genomic
DNA was extracted from the pellets of the second PCA selection culture. The Pbs2 locus was
amplified and sequenced, as described in the DNA sequencing section.
272

273   The liquid PCA selection for the haploid double DHFR fused strains was done following the same
274   protocol as the first DHFR-PCA selection, only ignoring the mating step. However, a fourth control
275   condition of PCA medium without methotrexate and without sorbitol was added. Three replicate
276   pools were done for each growth condition. Frozen pellets were prepared as in the first DHFR-
277   PCA selection.
278

The individually reconstructed validation strains were pooled by combining 100 µL of saturated
overnight culture of each mutant in YPD. 6 pools were made for each condition, three pools each
from two individual reconstructed colonies of the mutant. The same conditions were used as the
second liquid PCA selection. For the cell proliferation measurements, SC medium with and
without 1 M of sorbitol was used. Two cycles of selection were done, with 96 hour cycles for the
conditions with methotrexate and 24 hour cycles for the conditions without methotrexate, and the
SC medium. Frozen pellets were prepared as for the previous liquid PCA selections.
286

<u>DNA sequencing and analysis</u>
For full details of DNA sequencing methods and analysis of DNA sequencing results, see File S3.
DNA sequencing was done in generally the same manner for all experiments, with minor changes
in certain cases. DNA was extracted from the frozen pellets using a standard phenol/chloroform
genomic DNA extraction protocol (Amberg *et al.* 2005). The extracted DNA was amplified and
barcoded using a Row-Column DNA barcoding strategy as previously described (Dubé *et al.*
2022). These barcoded pools were purified on magnetic beads, and sent for sequencing on
Illumina NGS machines. Unique combinations of barcodes were used to identify the DNA
extracted from each pellet in each replicate of each condition. After preliminary analysis of the
sequencing of the DNA from the pellets of the DHFR-PCA on the extended motif DMS library

(positions 85-100), it was determined that more reads would be needed. Additional sequencing
libraries were prepared fresh from the original DNA extractions. Instead of using a row-column
barcoding approach, the *PBS2* region was amplified using primers allowing the addition of
Illumina Nextera barcodes (Illumina). These allow automated demultiplexing by the Illumina
MiSeq instrument. All other steps were as described above.
302
The variant frequency in DMS libraries during construction and in the DHFR-PCA and proliferation
screens was evaluated using custom scripts based on already published work (Després *et al.*
2022). Briefly, to identify variants, reads were merged and demultiplexed, then identical reads
were grouped using vsearch (Rognes *et al.* 2016), and aligned to the wild-type *PBS2* sequence
using the needle function from the EMBOSS software (Rice *et al.* 2000). From this, the frequency
of each variant was obtained. From the frequency of each variant, the interaction scores and
selection coefficients of each variant were calculated using custom R 4.3.1 (R Core Team 2023)
scripts. The interaction score was calculated as the log-2-fold-change of the frequency of each
variant at the end of the second cycle of methotrexate selection and the frequency of each variant
after the selection for diploids (so immediately preceding the methotrexate selection). These log-
2-fold-change scores were rescaled for each library, so that the median of nonsense codons was
equivalent to -1 and the value of wild-type variants was 0. Variants were then filtered to keep only
those that had 20 or more reads detected after diploid selection, in order to remove bias caused
by low read counts leading to inflated scores. A unique score was calculated for each variant by
taking the median score of all codons coding for the same amino acid variant in all replicates of
the same condition. Scores were only kept for amino acid variants that had at least 3 replicates
from any combination of codons not filtered out. For the competition assay of cell proliferation, a
selection coefficient was calculated. We statistically verified which nonsense and silent mutants
were abnormal by comparing every nonsense or silent mutation to all other nonsense or silent
mutations (Two-sided Mann-Whitney U test, $p < 0.05$). The mutants which were significantly
different were removed from the interaction score dataset. Similarly, we verified which mutants
had defects in the control condition by comparing each missense mutant to the nonsense mutants
(Two-sided Mann-Whitney U test with false discovery rate corrected p-value > 0.05). The
missense mutants which were not significantly different from the nonsense mutants were also
removed from the interaction score dataset. We also removed from the interaction score dataset
the mutants for which an abundance change was detected through gain or loss of interaction
strength between Pbs2 and Hog1, as explained in the results section (Two-sided Mann-Whitney
U test with false discovery rate corrected p-value < 0.05).
331
<u>DHFR-PCA growth curves</u>
Growth curves were measured for individually reconstructed *PBS2* mutants in DHFR-PCA
medium, to validate Pbs2 interaction strength measurements. The DHFR F[3] fused strains with
the *PBS2* mutations were mated with a Sho1-DHFR F[1,2]/*pbs2::LEU2* strain. Exponential phase
precultures of diploids were diluted to 0.1 OD/mL in 80 µL of PCA medium, in a 384 well plate, in
4 replicates, at 30 °C. Cultures were grown either with and without 200 µg/mL of methotrexate,
and with and without 1 M sorbitol. The optical density of each well was measured every 15 minutes
in a Tecan Spark plate reader (Tecan). The growth rates for *SHO1* mutants were measured in
the same way, except that the mutant strains were mated with either Pbs2-DHFR

F[3]/*sho1Δ::LEU2* or Ybt1-DHFR F[3]/*sho1Δ::LEU2*. The latter strain is used to measure Sho1
stability, as Ybt1 interacts constitutively with Sho1. In cases where the mutation destabilizes
Sho1, the Ybt1 interaction will be affected, analogously to how Pbs2 destabilizing mutations affect
the Pbs2-Hog1 interaction. No sorbitol was added, and the optical density was measured in 200
µL in a 96 well plate using an Agilent Biotek Epoch 2 plate reader (Agilent). The plasmids
containing the *PBS2* fragments with the DHFR F[3] were transformed into 3 strains containing a
DHFR F[1,2] at the *GAL1* locus: Sho1-SH3-DHFR F[1,2], Abp1-SH3-DHFR F[1,2] Nbp2-SH3-
DHFR F[1,2] as well as a YFP-DHFR F[1,2] and a control wild-type strain. Three individual
colonies were taken from each transformation as independent replicates, and grown to
exponential phase. The expression of the DHFR F[1,2] fused fragments was induced using 20
nM of estradiol, which drives expression of the *GAL1* promoter by the GEM artificial transcription
factor, and should result in the same expression level of each construct (Aranda-Díaz *et al.* 2017).
The growth curves were measured in the same manner as the *SHO1* mutants.
354
Growth curves data was analyzed using a custom script written in R (R Core Team 2023). The
maximal growth rate was calculated using a custom function, by measuring growth over 5 time
points, and taking the 98th percentile to avoid outliers. *PBS2* mutant P94R was disregarded, as
growth of all replicates in the control condition in the absence of methotrexate was much lower
than all other samples, suggesting a problem with strain construction. For the *SHO1* mutants, an
interaction score was calculated by normalizing scores by the median score of wild-type Sho1
with the same partner. For the *PBS*2 fragments, an interaction score was calculated by
normalizing the growth rate using the growth rate of the strain measuring the interaction of the
same SH3 domain with a DHFR F[3] fused to no Pbs2 fragment.
364
365
<u>Structure prediction and protein-protein contact analysis</u>
Structure predictions were done for wild-type Pbs2 and Pbs2 mutants in complex with Sho1 using
AlphaFold 2.3.2 and the AlphaFold-Multimer implementation (Jumper *et al.* 2021; Evans *et al.*
2022). Default options were used, and only the top scoring model for each mutant was relaxed.
For Sho1, the sequence of the entire protein was used for prediction (Uniprot entry P40073). For
Pbs2, residues 71 to 126 were used for the prediction (Uniprot entry P08018). Predicted
structures were visualized using ChimeraX 1.8 (Pettersen *et al.* 2021).
373
Contacts were analyzed for 4 structures. Sho1-Pbs2 was generated by AlphaFold-Multimer, while
Abp1-Ark (PDB accession 2RPN), Nbp2-Ste20 (PDB accession 2LCS), and Bem1-2-Ste20 (PDB
accession 2RQW) were all previously measured NMR structure ensembles, composed of 20
individual structures. All 20 structures were used for determining the contacts, although only the
designated representative structure was used for visualization. The structures were imported into
ChimeraX 1.8 (Pettersen *et al.* 2021), and the ChimeraX "Contacts" tool was used to detect all
pairs of atoms from different proteins with a Van der Waals radius overlap ≥ -0.40 Å. This was
done for all atoms in the motif-bearing protein preceding position -3 in the standard motif
numbering. To identify equivalent positions in the four SH3 domains which form contacts, the
structures of the 4 domains were aligned using the MUSTANG tool (Konagurthu *et al.* 2006), and
a standard numbering was determined based on this alignment (See Table S9 in File S2). The

contacts between atoms were filtered to simply indicate all contacts between residues, and are
summarized in Table S10 in File S2, using the aforementioned standard numbering. The minimum
distance between residues or proteins was calculated using the bio3d package (Grant *et al.* 2006).
388

<u>Sequence and structural alignments</u>
The protein sequences of 24 yeast SH3-containing sequences were obtained from the
alliancemine server (Bult and Sternberg 2023). The sequence of the SH3 domains plus 25
residues on each side was extracted from the complete protein sequences using the SH3 location
information from Interpro (Paysan-Lafosse *et al.* 2023), and a custom R script. When an SH3
domain was within less than 25 residues of the N- or C-terminus of a protein, the sequence up to
the terminus was used. The sequences of these SH3 domains and adjoining regions were aligned
using mafft 7.526, using the L-INS_i (local alignment) algorithm (Katoh and Standley 2013). For
the structural alignment, we used previously predicted structures from the AlphaFold Protein
Structure Database (Varadi *et al.* 2022). Using the R packages AlphaMissenseR (Morgan and
Nguyen) and bio3d (Grant *et al.* 2006), we downloaded the predicted structures of the 24 same
yeast SH3-containing proteins from the AlphaFold Protein Structure Database, and trimmed the
pdb files to only contains residues composing the SH3 domains as well as 25 residues on each
side of the domain. To align the predicted sequences, we used the MUSTANG 3.2.4 tool
(Konagurthu *et al.* 2006).
404

<u>DHFR-PCA on solid media against SH3 domain-containing proteins</u>
The DHFR-PCA on solid media was based on previous work (Tarassov *et al.* 2008; Rochette *et
al.* 2015). All colony manipulation was done using a robotic pin tool platform (BM5-SC1, S&P
Robotics Inc.). The strains containing DHFR F[1,2] fused SH3-containing proteins were taken
from the Yeast Protein Interactome Collection (Horizon Discovery) or built for this project.
Colonies were cherry picked from the 96-well plates, and arrayed onto YPD+Nat plates for the
DHFR F[1,2] strains or YPD+Hyg plates for the DHFR F[3] strains, in a 384 colony array. The
DHFR F[1,2] strains were organized into a randomized pattern to avoid any effects caused by
neighboring colonies. The 384 colony arrays were condensed into 1536 colony arrays on
YPD+Nat or YPD+Hyg, with the randomized DHFR F[1,2] pattern repeated four times. The outer
two rows and columns of the 1536 arrays were composed of LSM8-DHFR F[1,2] and CDC39-
DHFR F[3], which grow well on PCA medium and serve to avoid any measurement bias from
being on the edge of the array, for the interactions of interest. The 1536 colony arrays were mated
together by pinning the repeated DHFR F[1,2] array onto YPD plates, and then pinning one of the
DHFR F[3] arrays onto each plate. In this collection of arrays, each interaction between an SH3
containing protein and a Pbs2 variant was measured in 5 or 6 replicates. After 48 hours of mating
at 30 °C, the mated colonies were pinned onto YPD+Nat+Hyg plates to select for diploid cells.
Two growth cycles of 48 hours at 30 °C on YPD+Nat+Hyg were done. After diploid selection, the
plates were photographed using an EOS Rebel T5i camera (Canon), to verify that all colonies
grew correctly. The mated arrays were then pinned onto solid PCA medium plates, either with or
without methotrexate, and with 1 M of sorbitol. The arrays were grown for two cycles of 48 hours
at 30 °C in a custom growth and imaging platform (S&P Robotics Inc.), which incubated the plates
and took a picture of each plate every 2 hours.

These plate pictures were analyzed to measure the size of every colony. First, the ImageMagick
command line tool (ImageMagick Studio LLC 2023) was used to crop the images of the selection
plates as well as change them to grayscale and invert the colors. Colony sizes were then
quantified using the Python package Pyphe (Kamrad *et al.* 2020). The colony sizes were analyzed
and visualized using a custom script written in R 4.3.1 (R Core Team 2023), based on a previous
analysis (Dionne *et al.* 2021). Colony sizes were filtered, and colonies which had not grown at the
end of the diploid selection were removed from the PCA results. Colony sizes were log2
transformed and then normalized to avoid bias from a position in the array or from the plate they
grew on. Scores were rescaled, so that the median of the scores of the Sho1 and wild-type Pbs2
interaction was equivalent to 1 and the median of scores of the Sho1 and Pbs2-stuffer interaction
was equivalent to 0.

<u>Large Language Model use</u>
Google's large language model Gemini Flash 2.5 was used after manuscript drafting to suggest
changes to grammar and syntax. This was done to increase the clarity and conciseness of the
text. Suggestions from the model were selectively applied, based on the authors' judgment. The
model was prompted by providing the text, paragraph by paragraph, and asking "Here is part of
a scientific article I am writing. Can you give me suggestions on how to make the text more clear
and more compact?".

449 Results

<u>Scan of the region surrounding the Pbs2 motif reveals that few mutations modify binding to Sho1</u>

The interaction between Sho1 and Pbs2 is modulated by an SH3 domain on Sho1 and the
canonical binding motif on Pbs2, situated in positions 93-99, and which has the sequence
KPLPPLP (Maeda *et al.* 1995). Previous computational work has suggested that the region
surrounding binding motifs could play an important role in modulating binding, including in SH3
interactions and in particular in the Sho1-Pbs2 interaction (Stein and Aloy 2008; Kelil *et al.* 2016).
To determine which Pbs2 residues play a role in binding, we undertook a Deep Mutational Scan
(DMS) on 56 codons in positions 71-126, comprising the binding motif and its surrounding region.
We generated a DNA library containing all possible single codon mutations within this region and
used CRISPR-Cas9 to insert it into the *PBS2* locus of *S. cerevisiae*, replacing the wild-type
sequence (Dionne *et al.* 2021). The *PBS2* gene was fused with a Dihydrofolate Reductase
Protein-Fragment 3 (DHFR F[3]), and these cells were mated with cells containing a DHFR F[1,2]
fusion with *SHO1*, creating diploid strains with complementary DHFR fragments on both
interaction partners. Sho1-Pbs2 interaction reconstitutes a functional DHFR, which enables cell
division in the presence of methotrexate (MTX) (Figure 1b). Growth in the presence of MTX in a
strain with both DHFR fragments thus serves as a proxy for the amount of Sho1-Pbs2 complexes
forming, which depends on both the binding affinity and the local abundance of the interacting
proteins (Tarassov *et al.* 2008; Freschi *et al.* 2013). Since the DHFR fusions are C-terminal,
nonsense mutations in the library result in the absence of DHFR fragment expression.

471
To measure the interaction strength of the various mutants in the DMS library, the mutants were
pooled and DHFR-PCA competition assays were done in media containing both MTX and 1 M of
sorbitol, in order to induce the HOG pathway (Ferrigno *et al.* 1998). The interaction score for each
variant was calculated by normalizing the log2-fold change in variant frequencies before and after
selection in MTX, determined by targeted sequencing of the *PBS2* locus. Within each replicate,
scores were normalized such that wild-type Pbs2 had a score of 0, and the median of nonsense
mutants had a score of -1. Mutants which possess a higher interaction strength than wild-type
Pbs2 therefore have a positive interaction score, while mutants which possess a lower interaction
score than wild-type Pbs2 have a negative interaction score.
481
While many mutations within the canonical motif altered interaction strength, most mutations in
the surrounding region (positions 71-126) did not affect Sho1 binding, except for a short section
neighboring the canonical motif itself (Figure 2a, Figure S1 and Figure S2 in File S1, Table S1 in
File S2). This finding supports the critical role of the motif and validates computational predictions
that extended the binding interface to include nearby residues (Stein and Aloy 2008; Kelil *et al.*
2016). Since the effect on binding was strongly position dependent, we categorized different
sections of Pbs2 as follows: the canonical binding motif is the conserved type I binding motif
sequence situated in positions 93 to 99, the extended motif is composed of the canonical motif
and the neighboring residues which have a strong impact on binding, from positions 85 to 99.
491
DHFR-PCA signal reflects both binding affinity and protein abundance. To distinguish how
mutations can impact these, we assessed the effect of Pbs2 mutants on the Hog1-Pbs2
interaction, which is independent of the SH3-binding motif (Murakami *et al.* 2008), thus serving
as a control for Pbs2 local abundance. Therefore, mutations solely affecting Sho1 binding should
not impact the Hog1-Pbs2 DHFR-PCA, whereas mutations affecting Pbs2 stability or local
abundance will influence both the Sho1-Pbs2 and Hog1-Pbs2 assays.
498
We found that most mutations in the surrounding region of the Pbs2 motif had little effect on Hog1
binding (Figure 2a). Furthermore, mutants that had a negative effect on Hog1-Pbs2 binding also
had a negative effect on Sho1-Pbs2 binding, suggesting that the reduction in signal results from
a loss of local abundance of Pbs2 (Figure S3 in File S1). However, most mutations that affected
Sho1 binding did not impact the Hog1 interaction, and therefore did not change the local
abundance of Pbs2.
505
506 <u>Extended motif DMS library identifies mutations modulating interaction strength</u>
507
Based on the initial results, we focused our subsequent analysis on the extended Pbs2 motif
(codons 85-99), including position 100 to measure potential effects immediately downstream of
the canonical motif. The initial DHFR-PCA in a diploid strain led to incomplete measurement of
Sho1-Pbs2 interaction, as some Pbs2 mutants presumably interacted with the Sho1 copy not
fused with DHFR F[1,2]. Consequently, we built a new haploid strain containing both DHFR F[1,2]
on Sho1 and DHFR F[3] on Pbs2. We also constructed a Hog1-DHFR F[1,2]/Pbs2-DHFR F[3]
haploid strain to control for Pbs2 local abundance. These new strains capture all Pbs2 binding to

Sho1 or Hog1, therefore allowing for more sensitive measurements. Using the same CRISPR-
Cas9 approach as before, we introduced a second Pbs2 DMS library, comprising all possible
single mutations in Pbs2 positions 85 to 100, into the two haploid DHFR strains. Mutant interaction
strengths were measured via DHFR-PCA as previously described (Figures S4 and S5 in File S1).
We then excluded 12 Pbs2 mutants showing significantly altered Hog1 interaction scores
compared to wild-type and synonymous Pbs2 mutants (two-sided Mann-Whitney U test, FDR-
corrected p-value < 0.05; Figure S6 in File S1). This filtering step eliminated mutants affecting
interaction through changes in Pbs2 abundance or stability, which influence both Sho1 and Hog1
interactions.
524
The impacts on the Sho1-Pbs2 interaction were varied (Figure 2b, Figure S7 in File S1, Table S2
in File S2). Only a small fraction of mutants exhibited interaction scores comparable to nonsense
mutants, indicating that single mutations rarely abolish Sho1-Pbs2 binding completely.
Comparison of missense mutant interaction scores to wild-type and synonymous variants
revealed that many mutants displayed significantly stronger or weaker interactions (two-sided
Mann-Whitney U test, FDR-corrected p < 0.05). Specifically, 18 mutants (6.5% of missense)
showed significantly increased interaction, while 173 mutants (62.7% of missense) showed
significantly decreased interaction compared to wild-type Pbs2. The impact of mutants that
reduced the interaction scores was generally greater.
534
Given that osmotic stress enhances the Sho1-Pbs2 interaction (Maeda *et al.* 1995; Posas and
Saito 1997; Saito and Posas 2012), we compared interaction strengths of Pbs2 mutants with and
without 1 M sorbitol (Figure S8 in File S1). A strong correlation was observed between interaction
scores under both conditions (Spearman's rho 0.98, p-value < 2.2×10^{-16}). As expected due to
HOG pathway induction, the effects on binding were stronger in the presence of sorbitol. Under
osmotic stress, strongly interacting Pbs2 mutants are even more often in contact with Sho1,
increasing the number of reconstituted DHFR complexes, and further outcompeting weakly
interacting mutants. Nevertheless, the high correlation suggests that Pbs2 mutants exhibit similar
relative binding changes regardless of osmotic stress, and the fundamental impact of mutations
is independent of HOG pathway activation.
545
To validate the pooled measurements, we individually reconstructed 24 mutants spanning the
range of measured interaction scores in Pbs2-DHFR F[3] strains and mated them with the Sho1-
DHFR F[1,2] strain. Growth rates of these diploids were then measured in DHFR-PCA medium.
As with the competition assay, the growth rate is a proxy for interaction strength, which reflects
affinity in the absence of a change in abundance. The growth curve results correlate strongly with
the interaction scores (Spearman's rho 0.86, p-value 2.4×10^{-6} with 1 M sorbitol, spearman's rho
of 0.8, p-value 4.3×10^{-6} without sorbitol), confirming the competition assay results (Figure S9a in
File S1). To further validate the pooled data, we performed a smaller-scale competition DHFR-
PCA, using 67 individually reconstructed mutants with diverse interaction scores (Figure S10 in
File S1, Table S3 in File S2).This validation assay showed a strong correlation with the DMS
library measurements (Spearman's rho 0.98, p-value < 2.2×10^{-16} with 1 M sorbitol, spearman's
rho of 0.98, p-value < 2.2×10^{-16} without sorbitol), thus validating the DMS library results (Figure
S9b in File S1). To assess the direct impact of mutations on cell proliferation, we constructed

strains with the same 67 mutations but lacking DHFR fragments. Cell growth and division thus
only depend on the effect of the mutations, and not interaction strength. We then performed a
pooled competitive growth assay in SC medium with 1 M sorbitol to measure the proliferation of
each mutant. With the exception of nonsense mutations, no significant effect on cell proliferation
was observed (Welch's t-test false discovery rate corrected p-value < 0.05) (Figure S11 in File
S1, Table S3 in File S2). Notably, the strongest interacting mutant, I87W, shows a detectable,
though not statistically significant, deleterious effect on cell proliferation.
566
567   <u>Structure prediction suggests a secondary contact between extended motif and Sho1</u>
568
While the crystal structure of the Sho1 SH3 domain bound to a 9-residue Pbs2 segment shows
the canonical motif (positions 93-99) occupying the entire canonical binding pocket (Kursula *et al.*
2008), our data reveals that many mutations outside this motif strongly affect binding as well. We
hypothesized that residues in the extended Pbs2 motif (positions 85-99) might interact with Sho1
outside its canonical SH3 domain pocket. This is supported by NMR structures and studies of
other SH3 domains showing that residues outside the canonical motif can modulate interaction
strength and specificity (Rickles *et al.* 1995; Feng *et al.* 1995; Stollar *et al.* 2009; Takaku *et al.*
2010; Gorelik and Davidson 2012). Given the absence of experimental structures for full-length
Sho1 or Pbs2, we used AlphaFold-Multimer predictions (Jumper *et al.* 2021; Evans *et al.* 2022)
to explore potential interactions beyond the canonical binding site (Figure S12 in File S1).
Modeling the entire Sho1 protein along with the surrounding region of the Pbs2 motif (positions
71-126), revealed potential interactions outside of the canonical binding pocket of Sho1, mediated
by residues neighboring the canonical binding motif of Pbs2 (Figure 3a). The predicted structure
shows a hydrophobic pocket formed by the Sho1 SH3 domain and a nearby loop, interacting with
an alpha helix formed by Pbs2 residues 84-92 (Figure 3b). Interestingly, all but one of the mutants
in the extended motif which significantly increase interaction strength are located within this
predicted Pbs2 helix. Although the model also predicted proximity of some C-terminal Pbs2
residues to Sho1, mutations in these positions had minimal impact on binding (Figure 2a), leading
us to focus on residues within and near the extended motif.
588
To understand how specific mutations alter binding, we predicted the structures of the five Pbs2
variants exhibiting the strongest Sho1 interaction, using the same method as for wild-type Pbs2
(Figure S12 in File S1). In particular, the strongest interacting mutant, I87W, is predicted to
position its tryptophan side chain in close proximity (1.77 Å) to the non-canonical hydrophobic
pocket, specifically within a sub-pocket formed by both SH3 and non-SH3 Sho1 residues.
Furthermore, it is predicted to form a hydrogen bond with D333 of Sho1 (Figure 3c). These two
factors potentially explain its strong interaction strength relative to all other Pbs2 variants.
Similarly, the strongly interacting mutants H86W, V91L, and V91M are all predicted to place
hydrophobic side chains within the non-canonical hydrophobic pocket, potentially contributing to
stronger interactions. For the remaining mutant, Q89D, the hydrophilic aspartate side chain is
predicted to face away from the hydrophobic pocket, thus not disrupting these interactions.
However, the mechanism of its increased binding strength remains unclear (Figure S13 in File
S1).
602

We hypothesized that replacing hydrophobic Pbs2 residues predicted to contact the hydrophobic
non-canonical pocket with non-hydrophobic ones would disrupt this interaction and weaken
binding. Consistent with this, analysis of mutations within the predicted alpha helix shows that
replacing the initially hydrophobic residues at positions 87, 90, and 91 with non-hydrophobic
residues reduces binding strength. Generally, substituting these positions with other hydrophobic
residues had a less detrimental effect, and for V91, many hydrophobic substitutions even
increased interaction strength (Figure 3d). These results support the prediction that the side
chains of residues at positions 87, 90, and 91 are located within the non-canonical hydrophobic
binding pocket. This finding helps explain the strong interactions strength of Pbs2 mutant I87W,
by showing the important role of hydrophobic interactions in position 87.

To further support the presence of contacts between Pbs2 and surfaces outside the Sho1
canonical binding pocket, we introduced a series of Sho1 mutations in residues predicted to be
in proximity to Pbs2, in strains containing the DHFR fragments. We then assessed the effect of
the mutations on interaction strength using growth assays. To control for Sho1 stability and
abundance, similar to our approach with Hog1 for Pbs2 mutants, we also measured the interaction
of the Sho1 mutants with Ybt1, a protein previously found to interact with Sho1 independently
from the SH3 domain (Dionne *et al.* 2021). We compared mutant growth rates to wild-type
controls, looking for Sho1 variants that diminished Pbs2 binding but not Ybt1 binding (One-sided
Student's t-test, FDR-corrected p-value < 0.05, Figure S14 in File S1). We identified four Sho1
positions where mutations specifically weakened the Pbs2 interaction. Three of these positions,
319, 333 and 350, are located in the SH3 domain, although outside the canonical binding site.
The fourth position, 287, is situated in a neighboring loop. All four are predicted to be within 5 Å
of the Pbs2 extended motif, with the G287 backbone notably predicted to form a hydrogen bond
with residue Q88, located in the predicted alpha helix. Together, our structural predictions and
interaction measurements indicate that residues outside the canonical Pbs2 motif could directly
interact with residues both within and without the Sho1 SH3 domain, thereby modulating binding
affinity.

<u>Comparison of Sho1-Pbs2 binding to other SH3-motif pairs</u>

Given the proposed role of the extended Pbs2 motif and the additional Sho1 binding pocket,
composed of the SH3 domain and a nearby loop, in modulating interaction strength, we
investigated if these features also contribute to the specificity of Pbs2 binding to Sho1. We
reasoned that if this additional pocket were unique among yeast SH3 domains, it could enhance
binding specificity, as the Pbs2 extended motif might be specifically adapted to this feature. First,
we performed a multiple sequence alignment of yeast SH3 domains, including 25 flanking
residues on each side, which encompasses all Sho1 residues predicted to be within 5 Å of Pbs2.
Apart from key conserved residues within the SH3 domain itself, this analysis revealed no
significant conservation of residues forming the additional Sho1 pocket (Figure S15a in File S1).
To assess potential structural conservation despite sequence divergence, we conducted a
multiple structural alignment using MUSTANG (Konagurthu *et al.* 2006) on AlphaFold-predicted
structures (Varadi *et al.* 2022) of yeast SH3 domains, with 25 neighboring residues. This analysis
showed structural conservation only within the SH3 domains, with no significant structural

conservation in the flanking regions, including those forming the additional Sho1 binding pocket
(Figure S15b in File S1). Therefore, we conclude that the additional binding pocket on Sho1
represents a unique structural feature among yeast SH3 domains.
650
Despite the lack of conserved structure of non-SH3 sequences, there are reports of three other
yeast SH3 domains binding extended motifs: Abp1, the second SH3 domain of Bem1 (referred to
as Bem1-2) and Nbp2 (Stollar *et al.* 2009; Takaku *et al.* 2010; Gorelik and Davidson 2012). We
investigated whether these SH3 domains interact with the extended Pbs2 motif *in vivo* using a
DHFR-PCA assay that measures colony growth on solid medium (Tarassov *et al.* 2008).The SH3-
containing proteins were fused with DHFR F[1,2] and Pbs2 with DHFR F[3]. As a control, we used
a Pbs2 variant where the extended motif (positions 85-99) and surrounding region (codons 71-
126, the same positions as the initial DMS library) were replaced by a neutral, flexible "stuffer"
peptide (GGSSGGGG). The interaction strength of the SH3 domains was also measured against
this Pbs2-stuffer variant.
661
Abp1 and Nbp2 interacted with Pbs2 at a similar strength to Sho1, while Bem1-2 showed no
detectable interaction (Figure 4a). However, Abp1 and Nbp2 also interacted with Pbs2-stuffer at
comparable levels to wild-type Pbs2, indicating that their binding is independent of the Sho1-
binding motif. Nbp2 is known to bind a second, distal motif on Pbs2 through its SH3 domain
(Mapes and Ota 2004), while Abp1 likely binds Pbs2 through an alternative, unknown site.
Screening all other known yeast SH3-containing proteins revealed significantly weaker
interactions with both Pbs2 variants compared to wild-type Sho1 (one-sided Mann-Whitney U test,
FDR-corrected p-value < 0.05; Figure S16 in File S1). Furthermore, wild-type Pbs2 showed a
stronger interaction with a Sho1 variant where the SH3 domain was replaced by the
aforementioned stuffer (Sho1-stuffer), than Pbs2-stuffer with Sho1-stuffer, confirming the role of
the non-SH3 portion of Sho1 in binding the extended motif of Pbs2 (Figure 4a). Given that Sho1-
stuffer abundance is similar to wild-type Sho1 (Dionne et al. 2021), these results demonstrate that
the extended Pbs2 motif specifically binds to Sho1 through both its SH3 and non-SH3 regions.
675
To further test whether the Pbs2 motif and surrounding region can bind other SH3 domains, we
used DHFR-PCA to measure the interaction of Pbs2 fragments of different lengths with the SH3
domains of Abp1, Nbp2 and Sho1. We expressed the SH3 domains at the same level to remove
bias resulting from the different cellular abundances of Abp1, Nbp2 and Sho1. We also measured
the interaction of the same Pbs2 fragments against a constitutively expressed YFP-DHFR F[1,2]
to control for proper expression of the Pbs2 fragments (Lemieux *et al.* 2025). The canonical Pbs2
motif (positions 93-99) showed a significantly stronger interaction (One-sided Student's t-test,
FDR-corrected p-value < 0.05) with the Sho1 SH3 domain compared to the Nbp2 SH3 domain
(Figure S17 in File S1). This preference for the Sho1 SH3 domain was maintained and the
interaction strength increased with two longer Pbs2 fragments: positions 85-100 (the region of the
second DMS screen) and positions 69-124 (corresponding roughly to the Pbs2-stuffer
replacement), which also showed significantly stronger binding to the Sho1 SH3 domain
compared to the Abp1 and Nbp2 SH3 domains. While interaction scores with the non-Sho1 SH3
domains increased with Pbs2 fragment length, they were always weaker than the Sho1

interaction. This confirms the role of the extended motif and residues even more distant from the
canonical site in maintaining binding specificity to the Sho1 SH3 domain.
692
To understand the basis of Pbs2 binding specificity to Sho1, we compared the predicted Sho1-
Pbs2 structure with NMR structures of Abp1-Ark1, Bem1-2-Ste20, and Nbp2-Ste20, all involving
peptides of similar length to the Pbs2 extended motif. The structures of the four SH3 domains and
their bound peptides showed a conserved binding mode, with each peptide occupying the
canonical SH3 domain pocket and extending towards the region corresponding to the additional
Sho1 binding pocket (Figure 4b, Figure S18 in File S1). However, Abp1 and Nbp2 appear to
interact with their respective extended motifs solely through the SH3 domain surface. In contrast,
Bem1-2 utilizes a second, non-SH3 interface that forms a pocket with its SH3 domain,
accommodating the extended motif, similar to Sho1. This Bem1-2 interface, identified as the
Cdc42-interacting (CI) region (Takaku *et al.* 2010), exhibits no structural or sequence similarity to
the corresponding region in Sho1. Furthermore, unlike Sho1, this additional binding region in
Bem1-2 is located C-terminal to its SH3 domain.
705
Beyond the conserved canonical motif positions, the extended motifs of Pbs2, Ark1, and Ste20
show limited sequence conservation. These extended motifs can be aligned using a previously
developed standard numbering system where the first conserved proline is assigned position 0,
preceding residues are numbered negatively, and following residues positively (Figure 4c) (Lim
et al. 1994). The binding specificity exhibited by Pbs2 may depend on the possible contacts
between the extended motif and the different SH3 domains, as well as the non-SH3 surfaces in
the cases of Sho1 and Bem1-2. By examining the interactions of these extended motifs with their
respective partners, particularly focusing on residues within the extended motif but outside the
canonical motif (positions 85-92), we aimed to identify determinants of Pbs2 binding specificity.
715
Three scenarios can differentiate Pbs2-Sho1 binding from the binding of other extended motifs to
their partners: either contacts are present in Pbs2-Sho1 but absent in other pairs, contacts are
absent in Pbs2-Sho1 but present in other pairs, or equivalent residues in extended motifs contact
different residues of their interaction partner. We find examples of all three scenarios. For
example, Pbs2 is predicted to contact Sho1 through a histidine residue in position -10, while Ark1,
for example, does not have contacts so far from the canonical motif (Figure S19a in File S1). Pbs2
is also missing certain interactions that are present in other pairs. For instance, in both Ste20 and
Ark1, the residue at position -7 is crucial for binding, as in both cases, mutation of the residue to
an alanine reduced binding affinity by at least an order magnitude, and this mutation had the
largest impact of all alanine substitutions outside the canonical motif (Stollar *et al.* 2009; Gorelik
and Davidson 2012). In contrast, position -7 in Pbs2 is not predicted to interact with Sho1, though
mutation to a hydrophilic residue increased interaction strength (Figure S19b in File S1, Table S2
in File S2). Finally, the same position in different extended motifs can form distinct interactions
with their partners. In position -6, the isoleucine in Pbs2 is predicted to contact Y319, W338 and
I350 on the SH3 domain. In contrast, the histidine in position -6 of Ark1 forms a salt bridge with
an aspartic acid on a loop of the Abp1 SH3 domain (Figure S19c in File S1). The structural
alignment of SH3 domains indicates that Sho1 has a gap at the position of the aspartic acid that
forms the salt bridge (Figure S15b in File S1, position 112). Even beyond these limited examples,

the four extended motifs show differences in how they bind their partners. This suggests that the
non-conserved regions within extended motifs enable a much greater degree of binding
differentiation than the conserved canonical motif alone.
737

Discussion

739
Binding motifs face the double challenge of binding their partner with appropriate strength while
avoiding spurious interactions with non-partner proteins. To understand this balance in the Sho1-
Pbs2 interaction, we used deep mutational scanning of the Pbs2 motif and its flanking regions to
quantify the impact of single residue mutations on Sho1 binding. We further predicted the Sho1-
Pbs2 structure and compared it to existing structures of other SH3 domain-peptide complexes.
Based on these comparisons, we propose several mechanisms by which the Pbs2 extended motif
achieves specific binding to Sho1 over other SH3 domains.
747
One important consideration in domain-peptide interactions is the potential trade-off between
interaction affinity and specificity. Intuitively, a domain that can strongly bind its target motif might
also exhibit increased unwanted interactions with similar motifs within the same family, due to
their physical and chemical resemblance. This idea is supported by previous studies on SH2 and
PDZ domains, which, like SH3 domains, bind short motifs. These studies have suggested that
mutants exhibiting higher affinity for their target tend to display lower specificity, implying a
potential incompatibility between high affinity and high specificity (Ernst *et al.* 2010; Haslam and
Shields 2012; Kaneko *et al.* 2012; Karlsson *et al.* 2016). However, a computational study of
binding motifs, including SH3 binding motifs, proposed that increased affinity can correlate with
increased specificity (Kelil *et al.* 2016). Other works have emphasized the importance of the
residues surrounding the canonical binding motif in determining both affinity and specificity (Li
2005; Ivarsson and Jemth 2019). In fact, one study computed that nearly 30% of binding energy
from the binding motif was contributed by residues outside the canonical motif, and suggested
that the residues outside the canonical motif mostly play a role in determining specificity (Stein
and Aloy 2008). Notably, our deep mutational scanning revealed that all mutations which
significantly increase the interaction strength between Sho1 and Pbs2 are situated outside of the
canonical motif. This suggests that the wild-type canonical motif might already be optimized for
maximal binding strength. Therefore, we propose a model where the canonical motif residues in
positions 93 to 99 are optimized and thus structurally constrained for affinity, while the neighboring
residues are less constrained and able to increase affinity and specificity when mutated.
768
The enhanced binding strength from mutations outside the Pbs2 canonical motif could arise from
interactions with surfaces beyond the SH3 domain's canonical pocket, potentially involving non-
SH3 surfaces on Sho1. Non-binding pocket interactions have previously been observed in
multiple SH3-mediated interactions (Lee *et al.* 1996; Dalgarno *et al.* 1997; Li 2005; Stollar *et al.*
2009; Takaku *et al.* 2010; Gorelik and Davidson 2012; Gaussmann *et al.* 2024), and we found
that mutations outside the canonical Sho1 SH3 binding pocket can disrupt binding, suggesting
that the mutated residues play a role in the interaction. Our AlphaFold-Multimer predictions
support this, revealing a hydrophobic pocket formed by the Sho1 SH3 domain and a non-SH3

loop, which residues of the Pbs2 extended motif can occupy. While the disordered nature of the
Pbs2 flanking region suggests a dynamic interaction, these models provide a plausible
mechanism for non-canonical motif-mediated binding. By removing the need for affinity to be
dependent only on canonical binding motifs, the residues that surround motifs could bind to
protein surfaces which are less conserved among SH3 domains, and therefore easier to
differentiate. The presence of less conserved components, such as extended motifs or non-
canonical binding pockets could thus contribute to specificity among domains of the same family.
The Sho1 hydrophobic pocket, formed by both SH3 and non-SH3 elements, exemplifies a unique
structural feature that might differentiate Sho1, enabling Pbs2 to achieve higher affinity without
necessarily increasing binding to other SH3 domains. This binding model, involving a larger
number of residues in tuning affinity and specificity, could potentially uncouple affinity and
specificity, avoiding a trade-off between the two. While affinity and specificity might be intrinsically
coupled within the highly conserved canonical binding interface, the less constrained surrounding
residues offer a greater capacity for independent variation of these traits.
791
SH3 domain-mediated interactions are known to be context-dependent, influenced not solely by
the domain itself but also by the surrounding protein environment. This context includes the non-
SH3 regions of the protein, interactions with other proteins, and cellular localization (Dionne *et al.*
2022). Consequently, an SH3 domain's interaction profile is not solely determined by its amino
acid sequence. Indeed, studies show that relocating an SH3 domain to a different protein or
expressing it in isolation significantly alters its interaction profile (Dionne *et al.* 2021; Lemieux *et*
al. 2024; Dibyachintan *et al.* 2025). The observed effects of Pbs2 mutations on binding strength
may be mediated not only through direct SH3 domain interactions but also within the broader
Sho1 protein context. The interaction of the Pbs2 extended motif with non-SH3 Sho1 residues
likely contributes to this context-dependent modulation. Other causes could include
stereochemical adjustments of SH3 binding loops or changes affecting Sho1's interactions with
other binding partners. Other proteins involved in the osmotic stress response, namely Ste11 and
Ste50, have been shown to interact with the Sho1 SH3 domain outside the canonical binding
pocket (Zarrinpar *et al.* 2004; Tatebayashi *et al.* 2006). Mutations in the Pbs2 peptide could
potentially influence the stability of these interactions as well.
807
Previous studies have investigated the binding of Pbs2 motif variants (Zarrinpar *et al.* 2003), and
our findings corroborate some of their observations. For instance, we also find that mutations near
the canonical Pbs2 motif can either increase or decrease binding strength. This study also
reported reduced fitness for a promiscuous Pbs2 double mutant (Zarrinpar *et al.* 2003). In
contrast, our analysis revealed no substantial fitness loss for single mutants, except for the I87W
variant, which exhibited a noticeable, though not statistically significant, decrease in binding
strength (Figure S11 in File S1). Interestingly, I87W is also the mutant showing the greatest
increase in interaction strength, potentially explaining its slight fitness cost. It has previously been
suggested that affinity increasing mutants may have deleterious effects on cell homeostasis, by
reducing the dynamicity of the pathway, and impeding dissociation of the partners (Wang *et al.*
2018). Consistent with this, prior work on the Sho1-Pbs2 interaction demonstrated that a Sho1
SH3 domain mutation increasing affinity led to a minor reduction in cell growth (Marles *et al.*

2004), indicating that the wild-type affinity may be already optimized, and that stronger binding
may disrupt signaling.
822
In conclusion, we show that mutations in positions neighboring the canonical binding motif of Pbs2
can both increase and decrease binding strength with Sho1. We propose that these non-canonical
Pbs2 residues can enhance binding affinity through interactions outside the conserved SH3
binding pocket. We hypothesize that Sho1 is similar to certain other yeast SH3 domains which
bind longer motifs through additional binding pockets (Stollar *et al.* 2009; Takaku *et al.* 2010;
Gorelik and Davidson 2012), and propose mechanisms through which an extended binding motif
could better differentiate binding to different SH3 domains. Moving forward, studies of SH3-motif
interactions should encompass a larger range of residues, considering both the SH3 domain
surface beyond the canonical pocket and the regions surrounding the motif. Our findings, along
with previous research, highlight that *in vitro* studies using isolated domains and minimal peptides
may not fully represent the complexity of domain-motif interactions within their native protein
context. By understanding and harnessing the potential of extended motif binding, researchers
may be better able to understand the impact of motifs in cell signaling and disease, and better
able to use motif binding in technical and medical applications.
837

Data availability

Strains and plasmids are available upon request. All analysis scripts and visualization scripts are
available on Github at the following address: https://github.com/Landrylab/Jordan_et_al_2024.
Supplementary File 1 contains Figures S1-S19. Supplementary File 2 contains Tables S1-S10.
Supplementary File 3 contains the Supplementary methods. All data, including Supplementary
Files 1-3, demultiplexed sequencing data, results from the DHFR-PCA experiments, growth curve
data, predicted structures, protein alignments, and all files and information necessary to run the
scripts are available in the following Dryad repository: https://doi.org/10.5061/dryad.79cnp5j3z.
846

Acknowledgments

848 The authors thank Philippe C. Després, Moïra Dion and Dan Evans-Yamamoto for help with
849 sequencing data analysis. We also thank all members of the LandryLab team for thoughtful
850 comments on the experiments and on figure conception. All molecular graphics were performed
851 using UCSF ChimeraX 1.8. The protein structure predictions were enabled by computing
852 resources provided by the Digital Research Alliance of Canada.
853

Funding

This work was supported by Canadian Institutes of Health Research Foundation grant number
387697 and Human Frontier Science Program research grant RGP34/2018 to C.R.L. C.R.L. holds
the Canada Research Chair in Cellular Systems and Synthetic Biology. D.J. was supported by
the Canada Graduate Scholarship - Master's program from NSERC, a graduate scholarship from
PROTEO, a graduate scholarship from the NSERC CREATE program EvoFunPath, a
Postgraduate Scholarship - Doctoral from NSERC and a Doctoral scholarship from FRQS. D.B.

was funded by the EMBO Long-Term Fellowship (LTF) (ALTF 1069-2019). U.D. was funded by
an NSERC graduate fellowship.
863

864 Author contributions

865 D.J., A.K.D., U.G. , D.B., and C.R.L. conceived and designed the study. C.R.L. supervised the
866 work and obtained funding. D.J. and A.K.D. built the strains and performed the experiments. D.J.
867 performed the computational analyses. D.J. wrote the first draft of the manuscript. D.J., A.K.D.,
868 U.G. , D.B., and C.R.L. contributed to revising the final version of the manuscript.
869

870 Literature cited

871
Amberg D., D. C. Burke, and J. N. Strathern, 2005 *Methods in yeast genetics: a Cold Spring Harbor*
Laboratory course manual. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y.

Aranda-Díaz A., K. Mace, I. Zuleta, P. Harrigan, and H. El-Samad, 2017 Robust synthetic circuits for two-
dimensional control of gene expression in yeast. ACS Synth. Biol. 6: 545–554.

Bendel A. M., A. J. Faure, D. Klein, K. Shimada, R. Lyautey, *et al.*, 2024 The genetic architecture of
protein interaction affinity and specificity. Nat. Commun. 15: 8868.

Biot-Pelletier D., and V. J. J. Martin, 2016 Seamless site-directed mutagenesis of the Saccharomyces
cerevisiae genome using CRISPR-Cas9. J. Biol. Eng. 10: 6.

Bult C. J., and P. W. Sternberg, 2023 The alliance of genome resources: transforming comparative
genomics. Mamm. Genome 34: 531–544.

Dalgarno D. C., M. C. Botfield, and R. J. Rickles, 1997 SH3 domains and drug design: Ligands, structure,
and biological function. Pept. Sci. 43: 383–400.

Davey N. E., G. Travé, and T. J. Gibson, 2011 How viruses hijack cell regulation. Trends Biochem. Sci.
36: 159–169.

Delplace V., A. Ortin-Martinez, E. L. S. Tsai, A. N. Amin, V. Wallace, *et al.*, 2019 Controlled release
strategy designed for intravitreal protein delivery to the retina. J. Control. Release 293: 10–20.

Després P. C., A. F. Cisneros, E. M. M. Alexander, R. Sonigara, C. Gagné-Thivierge, *et al.*, 2022
Asymmetrical dose responses shape the evolutionary trade-off between antifungal resistance and
nutrient use. Nat Ecol Evol 6: 1501–1515.

Dibyachintan S., A. K. Dubé, D. Bradley, P. Lemieux, U. Dionne, *et al.*, 2025 Cryptic genetic variation
shapes the fate of gene duplicates in a protein interaction network. Nat. Commun. 16: 1530.

Dionne U., É. Bourgault, A. K. Dubé, D. Bradley, F. J. M. Chartier, *et al.*, 2021 Protein context shapes the
specificity of SH3 domain-mediated interactions in vivo. Nat. Commun. 12: 1597.

Dionne U., L. J. Percival, F. J. M. Chartier, C. R. Landry, and N. Bisson, 2022 SRC homology 3 domains:
multifaceted binding modules. Trends Biochem. Sci. 47: 772–784.

Diss G., and B. Lehner, 2018 The genetic landscape of a physical interaction. Elife 7.
https://doi.org/10.7554/eLife.32472

Douangamath A., F. V. Filipp, A. T. J. Klein, P. Barnett, P. Zou, *et al.*, 2002 Topography for Independent
Binding of α-Helical and PPII-Helical Ligands to a Peroxisomal SH3 Domain. Mol. Cell 10: 1007–
1017.

Dubé A. K., R. Dandage, S. Dibyachintan, U. Dionne, P. C. Després, *et al.*, 2022 Deep Mutational
Scanning of Protein–Protein Interactions Between Partners Expressed from Their Endogenous
Loci In Vivo, pp. 237–259 in *Yeast Functional Genomics: Methods and Protocols*, edited by
Devaux F. Springer US, New York, NY.

Ernst A., D. Gfeller, Z. Kan, S. Seshagiri, P. M. Kim, *et al.*, 2010 Coevolution of PDZ domain-ligand
interactions analyzed by high-throughput phage display and deep sequencing. Mol. Biosyst. 6:
1782–1790.

Evans R., M. O'Neill, A. Pritzel, N. Antropova, A. Senior, *et al.*, 2022 Protein complex prediction with
AlphaFold-Multimer. bioRxiv 2021.10.04.463034.

Faure A. J., J. Domingo, J. M. Schmiedel, C. Hidalgo-Carcedo, G. Diss, *et al.*, 2022 Mapping the
energetic and allosteric landscapes of protein binding domains. Nature 604: 175–183.

Feng S., C. Kasahara, R. J. Rickles, and S. L. Schreiber, 1995 Specific interactions outside the proline-
rich core of two classes of Src homology 3 ligands. Proc. Natl. Acad. Sci. U. S. A. 92: 12408–
12415.

Ferrigno P., F. Posas, D. Koepp, H. Saito, and P. A. Silver, 1998 Regulated nucleo/cytoplasmic exchange
of HOG1 MAPK requires the importin beta homologs NMD5 and XPO1. EMBO J. 17: 5606–5614.

Fowler D. M., and S. Fields, 2014 Deep mutational scanning: a new style of protein science. Nat.
Methods 11: 801–807.

Freschi L., F. Torres-Quiroz, A. K. Dubé, and C. R. Landry, 2013 qPCA: a scalable assay to measure the
perturbation of protein-protein interactions in living cells. Mol. Biosyst. 9: 36–43.

Gaussmann S., R. Peschel, J. Ott, K. M. Zak, J. Sastre, *et al.*, 2024 Modulation of peroxisomal import by
the PEX13 SH3 domain and a proximal FxxxF binding motif. Nat. Commun. 15: 3317.

Gorelik M., and A. R. Davidson, 2012 Distinct peptide binding specificities of Src homology 3 (SH3)
protein domains can be determined by modulation of local energetics across the binding
interface. J. Biol. Chem. 287: 9168–9177.

Gouw M., S. Michael, H. Sámano-Sánchez, M. Kumar, A. Zeke, *et al.*, 2018 The eukaryotic linear motif
resource – 2018 update. Nucleic Acids Res. 46: D428–D434.

Grant B. J., A. P. C. Rodrigues, K. M. ElSawy, J. A. McCammon, and L. S. D. Caves, 2006 Bio3d: an R
package for the comparative analysis of protein structures. Bioinformatics 22: 2695–2696.

Gueldener U., J. Heinisch, G. J. Koehler, D. Voss, and J. H. Hegemann, 2002 A second set of loxP
marker cassettes for Cre-mediated multiple gene knockouts in budding yeast. Nucleic Acids Res.
30: e23.

Haslam N. J., and D. C. Shields, 2012 Peptide-Binding Domains: Are Limp Handshakes Safest? Sci.
Signal. 5: e40-pe40.

Illumina, Nextera DNA Indexes. Illumina Support Docs.

ImageMagick Studio LLC, 2023 *ImageMagick*.

Ivarsson Y., and P. Jemth, 2019 Affinity and specificity of motif-based protein-protein interactions. Curr.
Opin. Struct. Biol. 54: 26–33.

Janke C., M. M. Magiera, N. Rathfelder, C. Taxis, S. Reber, *et al.*, 2004 A versatile toolbox for PCR-
based tagging of yeast genes: new fluorescent proteins, more markers and promoter substitution
cassettes. Yeast 21: 947–962.

Jumper J., R. Evans, A. Pritzel, T. Green, M. Figurnov, *et al.*, 2021 Highly accurate protein structure
prediction with AlphaFold. Nature 596: 583–589.

Kamrad S., M. Rodríguez-López, C. Cotobal, C. Correia-Melo, M. Ralser, *et al.*, 2020 Pyphe, a python
toolbox for assessing microbial growth and cell viability in high-throughput colony screens. Elife 9.
https://doi.org/10.7554/eLife.55160

Kaneko T., L. Li, and S. S.-C. Li, 2008 The SH3 domain--a family of versatile peptide- and protein-
recognition module. Front. Biosci. 13: 4938–4952.

Kaneko T., H. Huang, X. Cao, X. Li, C. Li, *et al.*, 2012 Superbinder SH2 Domains Act as Antagonists of
Cell Signaling. Sci. Signal. 5: ra68–ra68.

Karlsson O. A., G. N. Sundell, E. Andersson, Y. Ivarsson, and P. Jemth, 2016 Improved affinity at the
cost of decreased specificity: a recurring theme in PDZ-peptide interactions. Sci. Rep. 6: 34269.

Katoh K., and D. M. Standley, 2013 MAFFT multiple sequence alignment software version 7:
improvements in performance and usability. Mol. Biol. Evol. 30: 772–780.

Kazlauskas A., C. Schmotz, T. Kesti, J. Hepojoki, I. Kleino, *et al.*, 2016 Large-Scale Screening of
Preferred Interactions of Human Src Homology-3 (SH3) Domains Using Native Target Proteins as
Affinity Ligands. Mol. Cell. Proteomics 15: 3270–3281.

Kelil A., E. D. Levy, and S. W. Michnick, 2016 Evolution of domain–peptide interactions to coadapt
specificity and affinity to functional diversity. Proc. Natl. Acad. Sci. U. S. A. 113: E3862–E3871.

Kelil A., B. Dubreuil, E. D. Levy, and S. W. Michnick, 2017 Exhaustive search of linear information
encoding protein-peptide recognition. PLoS Comput. Biol. 13: e1005499.

Konagurthu A. S., J. C. Whisstock, P. J. Stuckey, and A. M. Lesk, 2006 MUSTANG: a multiple structural
alignment algorithm. Proteins 64: 559–574.

Kumar M., S. Michael, J. Alvarado-Valverde, A. Zeke, T. Lazar, *et al.*, 2024 ELM-the Eukaryotic Linear
Motif resource-2024 update. Nucleic Acids Res. 52: D442–D455.

Kursula P., I. Kursula, Y. H. Song, K. Paraskevopoulos, and M. Wilmanns, 2008 YEAST SHO1 SH3
DOMAIN COMPLEXED WITH A PEPTIDE FROM PBS2

Lee C. H., K. Saksela, U. A. Mirza, B. T. Chait, and J. Kuriyan, 1996 Crystal structure of the conserved
core of HIV-1 Nef complexed with a Src family SH3 domain. Cell 85: 931–942.

Lemieux P., D. Bradley, A. K. Dubé, U. Dionne, and C. R. Landry, 2024 Dissection of the role of a Src
homology 3 domain in the evolution of binding preference of paralogous proteins. Genetics 226.
https://doi.org/10.1093/genetics/iyad175

Lemieux P., A. K. Dube, and C. R. Landry, 2025 A protein-fragment complementation assay to quantify
synthetic protein scaffold efficiency. bioRxiv 2025.03.19.640584.

Li S. S.-C., 2005 Specificity and versatility of SH3 and other proline-recognition domains: structural basis
and implications for cellular signal transduction. Biochem. J 390: 641–653.

Liao T.-J., H. Jang, R. Nussinov, and D. Fushman, 2020 High-affinity interactions of the nSH3/cSH3
domains of Grb2 with the C-terminal proline-rich domain of SOS1. J. Am. Chem. Soc. 142: 3401–
3411.

Lim W. A., F. M. Richards, and R. O. Fox, 1994 Structural determinants of peptide-binding orientation and
of sequence specificity in SH3 domains. Nature 372: 375–379.

Maeda T., M. Takekawa, and H. Saito, 1995 Activation of Yeast PBS2 MAPKK by MAPKKKs or by
Binding of an SH3-Containing Osmosensor. Science 269: 554–558.

Mapes J., and I. M. Ota, 2004 Nbp2 targets the Ptc1-type 2C Ser/Thr phosphatase to the HOG MAPK
pathway. EMBO J. 23: 302–311.

Marles J. A., S. Dahesh, J. Haynes, B. J. Andrews, and A. R. Davidson, 2004 Protein-Protein Interaction
Affinity Plays a Crucial Role in Controlling the Sho1p-Mediated Signal Transduction Pathway in
Yeast. Mol. Cell 14: 813–823.

Michnick S. W., E. D. Levy, C. R. Landry, J. Kowarzyk, and V. Messier, 2016 The Dihydrofolate
Reductase Protein-Fragment Complementation Assay: A Survival-Selection Assay for Large-
Scale Analysis of Protein–Protein Interactions. Cold Spring Harb. Protoc. 2016: db.prot090027.

Miyazaki K., 2011 Chapter seventeen - MEGAWHOP Cloning: A Method of Creating Random
Mutagenesis Libraries via Megaprimer PCR of Whole Plasmids, pp. 399–406 in *Methods in
Enzymology*, Synthetic Biology, Part B. edited by Voigt C. Academic Press.

Monera O. D., T. J. Sereda, N. E. Zhou, C. M. Kay, and R. S. Hodges, 1995 Relationship of sidechain
hydrophobicity and alpha-helical propensity on the stability of the single-stranded amphipathic
alpha-helix. J. Pept. Sci. 1: 319–329.

Morgan M., and T. Nguyen, *AlphaMissenseR*.

Murakami Y., K. Tatebayashi, and H. Saito, 2008 Two adjacent docking sites in the yeast Hog1 mitogen-
activated protein (MAP) kinase differentially interact with the Pbs2 MAP kinase kinase and the
Ptp2 protein tyrosine phosphatase. Mol. Cell. Biol. 28: 2481–2494.

Pawson T., M. Raina, and P. Nash, 2002 Interaction domains: from simple binding events to complex
cellular behavior. FEBS Lett. 513: 2–10.

Paysan-Lafosse T., M. Blum, S. Chuguransky, T. Grego, B. L. Pinto, *et al.*, 2023 InterPro in 2022. Nucleic Acids Res. 51: D418–D427.

Pettersen E. F., T. D. Goddard, C. C. Huang, E. C. Meng, G. S. Couch, *et al.*, 2021 UCSF ChimeraX: Structure visualization for researchers, educators, and developers. Protein Sci. 30: 70–82.

Posas F., and H. Saito, 1997 Osmotic Activation of the HOG MAPK Pathway via Ste11p MAPKKK: Scaffold Role of Pbs2p MAPKK. Science 276: 1702–1705.

R Core Team, 2023 *R: A Language and Environment for Statistical Computing.* R Foundation for Statistical Computing, Vienna, Austria.

Rice P., I. Longden, and A. Bleasby, 2000 EMBOSS: the European Molecular Biology Open Software Suite. Trends Genet. 16: 276–277.

Rickles R. J., M. C. Botfield, X. M. Zhou, P. A. Henry, J. S. Brugge, *et al.*, 1995 Phage display selection of ligand residues important for Src homology 3 domain binding specificity. Proc. Natl. Acad. Sci. U. S. A. 92: 10909–10913.

Robles J. T., H. J. Lou, G. Shi, P. L. Pan, and B. E. Turk, 2023 Linear motif specificity in signaling through p38α and ERK2 mitogen–activated protein kinases. Proceedings of the National Academy of Sciences 120: e2316599120.

Rochette S., G. Diss, M. Filteau, J.-B. Leducq, A. K. Dubé, *et al.*, 2015 Genome-wide protein-protein interaction screening by protein-fragment complementation assay (PCA) in living cells. J. Vis. Exp. https://doi.org/10.3791/52255

Rognes T., T. Flouri, B. Nichols, C. Quince, and F. Mahé, 2016 VSEARCH: a versatile open source tool for metagenomics. PeerJ 4: e2584.

Ryan O. W., S. Poddar, and J. H. D. Cate, 2016 CRISPR-Cas9 Genome Engineering in Saccharomyces cerevisiae Cells. Cold Spring Harb. Protoc. 2016. https://doi.org/10.1101/pdb.prot086827

Saito H., and F. Posas, 2012 Response to Hyperosmotic Stress. Genetics 192: 289–318.

Stein A., and P. Aloy, 2008 Contextual specificity in peptide-mediated protein interactions. PLoS One 3: e2524.

Stollar E. J., B. Garcia, P. A. Chong, A. Rath, H. Lin, *et al.*, 2009 Structural, functional, and bioinformatic studies demonstrate the crucial role of an extended peptide binding site for the SH3 domain of yeast Abp1p. J. Biol. Chem. 284: 26918–26927.

Takaku T., K. Ogura, H. Kumeta, N. Yoshida, and F. Inagaki, 2010 Solution structure of a novel Cdc42 binding module of Bem1 and its interaction with Ste20 and Cdc42. J. Biol. Chem. 285: 19346–19353.

Tarassov K., V. Messier, C. R. Landry, S. Radinovic, M. M. S. Molina, *et al.*, 2008 An in Vivo Map of the Yeast Protein Interactome. Science 320: 1465–1470.

Tatebayashi K., K. Yamamoto, K. Tanaka, T. Tomida, T. Maruoka, *et al.*, 2006 Adaptor functions of Cdc42, Ste50, and Sho1 in the yeast osmoregulatory HOG MAPK pathway. EMBO J. 25: 3033–

3044.

Tonikian R., X. Xin, C. P. Toret, D. Gfeller, C. Landgraf, *et al.*, 2009 Bayesian Modeling of the Yeast SH3
Domain Interactome Predicts Spatiotemporal Dynamics of Endocytosis Proteins. PLoS Biol. 7:
e1000218.

Van Roey K., B. Uyar, R. J. Weatheritt, H. Dinkel, M. Seiler, *et al.*, 2014 Short Linear Motifs: Ubiquitous
and Functionally Diverse Protein Interaction Modules Directing Cell Regulation. Chem. Rev. 114:
6733–6778.

Varadi M., S. Anyango, M. Deshpande, S. Nair, C. Natassia, *et al.*, 2022 AlphaFold Protein Structure
Database: massively expanding the structural coverage of protein-sequence space with high-
accuracy models. Nucleic Acids Res. 50: D439–D444.

Vincentelli R., K. Luck, J. Poirson, J. Polanowska, J. Abdat, *et al.*, 2015 Quantifying domain-ligand
affinities and specificities by high-throughput holdup assay. Nat. Methods 12: 787–793.

Wang C.-H., P. Mehta, and C. J. Bashor, 2018 The strength of protein-protein interactions controls the
information capacity and dynamical response of signaling networks. bioRxiv 469197.

Yu H., J. K. Chen, S. Feng, D. C. Dalgarno, A. W. Brauer, *et al.*, 1994 Structural basis for the binding of
proline-rich peptides to SH3 domains. Cell 76: 933–945.

Zarin T., and B. Lehner, 2024 A complete map of specificity encoding for a partially fuzzy protein
interaction. bioRxiv 2024.04.25.591103.

Zarrinpar A., S.-H. Park, and W. A. Lim, 2003 Optimization of specificity in a cellular protein interaction
network by negative selection. Nature 426: 676–680.

Zarrinpar A., R. P. Bhattacharyya, M. P. Nittler, and W. A. Lim, 2004 Sho1 and Pbs2 Act as Coscaffolds
Linking Components in the Yeast High Osmolarity MAP Kinase Pathway. Mol. Cell 14: 825–832.

1072

1073

Figures

Figure 1. <u>The Sho1-Pbs2 interaction can be strengthened using sorbitol exposure, and the DHFR-PCA can be used to measure the interaction strength between Sho1 and Pbs2.</u>
a) Simplified schematic of the HOG pathway showing Sho1, Pbs2, and Hog1 interactions under normal conditions (left) where they are largely separate, and during osmotic stress (e.g., with sorbitol, right) where pathway induction leads to complex formation. The resulting Hog1 phosphorylation leads to nuclear translocation and gene expression modulation. b) Schematic of the DHFR-PCA screen using a DMS PBS2 mutant library. Single residue variants of PBS2 were inserted into the yeast genome at the native locus using CRISPR-Cas9 genome editing. Sho1 and Pbs2 were fused with complementary DHFR fragments. In cases where the Pbs2 variant does not interact with Sho1, the DHFR fragments remain separated. Upon interaction between a Pbs2 variant and Sho1, the DHFR fragments combine to reconstitute the functional DHFR, and allow growth in the presence of methotrexate (MTX). More strongly interacting partners allow the cell to proliferate more rapidly (Freschi *et al.* 2013).

1092

Figure 2. <u>Mutations in Pbs2 within and around the binding motif affect binding to Sho1</u>.

a) Interaction scores (normalized log2-fold change after MTX selection) of Pbs2 mutations across the region surrounding the SH3 binding motif (positions 70-126), measured for interaction with Sho1 and Hog1. Wild-type Pbs2 residues are indicated above each position, with the extended motif (positions 85-99) highlighted in blue and the canonical motif (positions 93-99) residues in bold. Boxplots show the distribution of interaction scores for mutants at all positions (3-18 replicates per mutant; nonsense mutants excluded). b) Distribution of interaction scores for single amino acid mutants (3-18 replicates) within the Pbs2 extended motif DMS library (positions 85-100) when interacting with Sho1. Mutations are displayed on the y-axis, colored by type: nonsense (stop codon), WT or silent (synonymous), and missense (amino acid change). Missense mutants were categorized based on Mann-Whitney U test (FDR-corrected p < 0.05) against WT and silent mutations: Significantly weaker (negative score), Not significantly different, or Significantly stronger (positive score).

1106

1107

Figure 3. <u>Contacts mediated by positions outside the canonical binding motif of Pbs2 could</u>
<u>modulate binding</u>
a) Predicted structure of Sho1 in complex with Pbs2 residues 71-126, oriented to highlight the
Pbs2 helix interacting with a hydrophobic pocket on Sho1, alongside the canonical motif (positions
93-99) in its binding pocket. For Sho1, the SH3 domain spanning positions 303 to 360 is colored
in green, while the rest of the protein is in yellow. For Pbs2, the extended motif spanning positions
85 to 99 is in blue while the rest of Pbs2 is in red. b) Close-up view of the interaction between
Sho1 and the extended motif (positions 85-99) of Pbs2. Only residues outside the canonical motif
of Pbs2 which are in contact with Sho1 are shown, and the Sho1 residues with which they are in
contact are also shown. The coloring of the residues is as in panel a, with Sho1 residues inside
the SH3 domain in green and Sho1 residues outside the SH3 domain in yellow. c) Close-up view
of the predicted Sho1-Pbs2(I87W) complex, showing the W87 side chain and a predicted
hydrogen bond (red dotted line) with Sho1 D333. All other Sho1 residues with which the I87W
residue is predicted to form contacts are shown with their sidechains. Coloring as in a). d) Boxplots
showing interaction scores of Pbs2 mutants at selected positions (from DMS DHFR-PCA of the
extended motif), grouped by the hydrophobicity of the substituted residue. Wild-type Pbs2 residue
at each position is indicated and colored by its hydrophobicity (Hydrophobic: F, I, W, L, V, M, Y,
C, A; Neutral: T, H, E, S, Q; Hydrophilic: R, K, N, G, P, D (Monera *et al.* 1995)). Nonsense mutants
are excluded.

Figure 4. The Pbs2 motif binds specifically to Sho1, despite similarities of other SH3 domains and motifs

a) Interaction score of wild-type Pbs2 (green) or Pbs2-stuffer (white) with select yeast SH3-containing proteins. All replicates shown as points. Scores are normalized relative to wild-type Sho1 interactions, with wild-type Pbs2 median at 1 and Pbs2-stuffer median at 0. b) Root mean square deviation (RMSD, in Å) of atom locations between matching backbone atoms in each pair of structures, as calculated using the R package bio3d (Grant *et al.* 2006). c) Sequences of Ste20, Ark1 and Pbs2, with standard numbering according to the motif, to compare equivalent positions (Lim *et al.* 1994). Ste20 and Ark1 positions included are those present in the structures used to generate in panel b, with Pbs2 positions overlapping all represented positions.