Residues Neighboring an SH3-Binding Motif Participate in the Interaction *In Vivo*

David F. Jordan[1,2,3,4,*], Alexandre K. Dubé[1,2,3,4,5], Ugo Dionne[2,3,4,6,7,8], David Bradley[1,2,3,4,5] and Christian R. Landry [1,2,3,4,5,*]

1. Département de Biochimie, microbiologie et bio-informatique, Université Laval, 1045 Avenue de la Médecine, Québec, QC, Canada, G1V 0A6
2. Institut de Biologie Intégrative et des Systèmes (IBIS), 1030, Avenue de la Médecine, Université Laval, Québec, QC, Canada, G1V 0A6
3. PROTEO-Regroupement Québécois de Recherche sur la Fonction, l'Ingénierie et les Applications des Protéines, Université Laval, 1030, Avenue de la Médecine Québec, QC, Canada, G1V 0A6
4. Centre de Recherche en Données Massives de l'Université Laval, Université Laval, 1065, Avenue de la Médecine, Québec, QC, Canada, G1V 0A6
5. Département de Biologie, Université Laval, 1045 Avenue de la Médecine, Québec, QC, Canada, G1V 0A6
6. Centre de Recherche du Centre Hospitalier Universitaire (CHU) de Québec-Université Laval, 11, côte du Palais, Québec, QC, Canada, G1R 2J6
7. Centre de Recherche sur le Cancer de l'Université Laval, 9, rue McMahon, Québec, QC, Canada, G1R 3S3
8. Present address: Lunenfeld-Tanenbaum Research Institute, Sinai Health, 600 University Avenue, Toronto, ON, Canada, M5G 1X5

* Correspondance: 1030, Avenue de la Médecine, Université Laval, Québec, QC, Canada, G1V 0A6 Email: david.jordan.2@ulaval.ca. 1030, Avenue de la Médecine, Université Laval, Québec, QC, Canada, G1V 0A6 Email: christian.landry@bio.ulaval.ca

Running head: Residues neighboring motif participate in interaction
Keywords: Short Linear motif, Src Homology 3 (SH3) domain, Deep mutational scanning, Binding affinity, Structure prediction

Abstract

In signaling networks, protein-protein interactions are often mediated by modular domains that
bind short linear motifs. The motifs' sequences affect many factors, among them affinity and
specificity, or the ability to bind strongly and to bind the appropriate partners. Using Deep
Mutational Scanning to create a mutant library, and protein complementation assays to measure
protein-protein interactions, we determined the *in vivo* binding strength of a library of mutants of
a binding motif on the MAP kinase kinase Pbs2, which binds the SH3 domain of the osmosensor
protein Sho1 in *Saccharomyces cerevisiae*. These measurements were made using the full-length
endogenous proteins, in their native cellular environment. We find that along with residues within
the canonical motif, many mutations in the residues neighboring the motif also modulate binding
strength. Interestingly, all Pbs2 mutations which increase affinity are situated outside of the Pbs2
region that interacts with the canonical SH3 binding pocket, suggesting that other surfaces on
Sho1 contribute to binding. We use predicted structures to hypothesize a model of binding which
involves residues neighboring the canonical Pbs2 motif binding outside of the canonical SH3
binding pocket. We compared this predicted structure with known structures of SH3 domains
binding peptides through residues outside of the motif, and put forth possible mechanisms through
which Pbs2 can bind specifically to Sho1. We propose that for certain SH3 domain-motif pairs,
affinity and specificity are determined by a broader range of sequences than what has previously
been considered, potentially allowing easier differentiation between otherwise similar partners.

Summary

Protein-protein interactions are often mediated by a binding domain on one protein and a short
disordered binding motif on another protein. We measured the binding strength of a mutant library
of a binding motif situated in the yeast protein Pbs2 to the SH3 domain of Sho1. Many mutations
in the residues neighboring the binding motif affect binding. A protein structure prediction of the
interaction partners shows that residues neighboring the motif bind residues outside the known
binding pocket on the SH3 domain. The Sho1-Pbs2 interaction differs enough from other known
SH3-motif pairs to allow specific binding.

65
66

68
69

Introduction

Cells possess complex and robust signaling networks that detect stimuli and trigger responses. These signaling networks are composed of a series of protein-protein interactions, which are often mediated by modular interaction domains. Many classes of protein interaction domains are shared across pathways and species, yet fulfill different roles and functions (Pawson *et al.* 2002). The different constraints placed on interaction domains by their distinct roles contribute to explaining the divergence in sequence between homologs of the same domain (Ernst *et al.* 2010; Dionne *et al.* 2022). Therefore, to understand how domain sequences have evolved and continue to evolve, the phenotypic consequences of mutations must be understood. Two of the major phenotypes for protein-protein interaction domains are affinity, that is, the strength of binding to the partner protein, and specificity, that is, the ability to bind to the appropriate partner proteins, and to not form spurious interactions with other proteins in the cellular environment (Ivarsson and Jemth 2019).

Many interaction domains bind short intrinsically disordered stretches of their interaction partners, also known as short linear motifs (SLiMs) or simply binding motifs (Gouw *et al.* 2018). Binding motifs are involved in a wide swath of cell signaling pathways, and underlie many important mechanisms in human and other cells (Kumar *et al.* 2024). Uncovering the determinants of binding can elucidate the functioning of human cells and certain diseases that affect them, such as various cancers in which motif-mediated signaling plays a part (Van Roey *et al.* 2014). Additionally, viruses also use their own binding motifs to disrupt the signaling machinery of their hosts (Davey *et al.* 2011). Understanding domain-motif binding can also allow them to be used as tools, for example by attaching therapeutic proteins to domains and motifs to hydrogels, to ensure the gradual release of the therapeutic proteins in human tissues (Delplace *et al.* 2019).

Interactions mediated by domain-motif associations are relatively weak, with dissociation constants (K_D) in the micromolar range, while domain-domain binding typically results in dissociation constants values orders of magnitude smaller, indicating much stronger binding (Van Roey *et al.* 2014). While such low affinity could simply be caused by physical limitations, such as the smaller interface, adaptive hypotheses have been put forward which suggest that an overly strong affinity can have deleterious effects on the cell, for instance by compromising specificity (Haslam and Shields 2012; Karlsson *et al.* 2016). Alternatively, this low intrinsic affinity can be compensated by other factors. For example, certain protein-protein interactions depend on the simultaneous binding of many domain-motif pairs (Liao *et al.* 2020). In other cases, motifs bind outside of the canonical binding pockets of their partner domains, although this does not always lead to increased affinity (Douangamath *et al.* 2002). The regions surrounding a domain can also have an effect on motif binding preference, as inserting domains into new protein backgrounds can change the motifs that this domain binds to (Dionne *et al.* 2021, 2022).

Binding affinity and specificity have previously been explored in the context of the interaction between binding domains and motifs, by measuring the binding of mutants *in vitro* (Yu *et al.* 1994; Zarrinpar *et al.* 2003; Tonikian *et al.* 2009; Vincentelli *et al.* 2015; Kazlauskas *et al.* 2016). However, the interaction of proteins in the cellular environment is a more complex situation, and

many interactions detected *in vivo* are not detected *in vitro,* and vice versa (Kelil *et al.* 2017).
Many factors can modulate binding in the cell, including colocalization of partners, expression in
the same cell cycle phases, and contributions from sequences outside of the immediate binding
domain and motif, as well as other proteins that can interact with one or both of the partners
(Ivarsson and Jemth 2019; Dionne *et al.* 2022). Earlier studies have measured the effects of
limited numbers of domain and motif mutants *in vivo* (Zarrinpar *et al.* 2003; Marles *et al.* 2004).
More recently, techniques such as deep mutational scanning (DMS) have been used to study the
impact of large libraries of mutations on protein stability and function *in vivo* (Fowler and Fields
2014). Methods such as protein complementation assays (Tarassov *et al.* 2008; Michnick *et al.*
2016) have also successfully been used to measure the *in vivo* binding strength of libraries of
mutants in yeast (Diss and Lehner 2018; Dionne *et al.* 2021; Faure *et al.* 2022; Robles *et al.* 2023;
Dibyachintan *et al.* 2024; Bendel *et al.* 2024). For example, a recent study showed the effect on
binding of combining mutations in a PDZ domain and its binding motif (Zarin and Lehner 2024).
These *in vivo* techniques could be used to determine how a large library of mutations can affect
domain-motif interaction affinity.
128
One powerful model to study domain-motif interactions is SH3 domains. These globular domains
are composed of around 60 residues, and bind different classes of polyproline motifs defined by
the canonical forms R/KXXPXXP for class I and PXXPXR/K for class II (Kaneko *et al.* 2008). The
model yeast *Saccharomyces cerevisiae*'s proteome contains 27 different SH3 domains (Dionne
et al. 2022), including one in the High Osmolarity Glycerol (HOG) signaling pathway protein Sho1
(Saito and Posas 2012). The SH3 domain of Sho1 is known to interact with a motif on the MAPKK
Pbs2, and this interaction can be strengthened by exposing cells to osmotic stress (Maeda *et al.*
1995; Posas and Saito 1997; Saito and Posas 2012) (Figure 1a). The Sho1-Pbs2 interaction is
thus a powerful model to study domain-motif interaction strength, due to the potential for
increasing or decreasing affinity through mutations and the simple method of inducing an increase
in the interaction strength using osmotic stress.
140
To better understand the impact of mutations on affinity, we measured the binding strength of a
nearly complete library of mutations in the Pbs2 binding motif *in vivo* using a DHFR Protein-
fragment Complementation Assay (PCA) (Tarassov *et al.* 2008; Michnick *et al.* 2016). We
measured the binding strength between single residue Pbs2 motif mutants and Sho1, the
canonical partner of Pbs2. We find that all single mutations that increase the interaction strength
are found outside the canonical binding motif, and that certain residues outside the binding motif
are predicted to interact with Sho1 outside of the canonical binding pocket, and even outside of
the SH3 domain. We use this finding to propose a model of Sho1-Pbs2 binding, where residues
outside the canonical Pbs2 motif bind a second binding pocket on Sho1. We also compare this
proposed model to other known SH3-motif interactions involving binding outside the canonical
motif.

Figure 1. <u>The Sho1-Pbs2 interaction can be strengthened using sorbitol exposure, and the DHFR-PCA can be used to measure the interaction strength between Sho1 and Pbs2.</u>

a) Simplified schematic of the HOG pathway, and the interactions between Sho1, Pbs2 and Hog1. In normal conditions (left), Sho1, Pbs2 and Hog1 are mostly separated. In the presence of osmotic stress such as sorbitol (right), signaling through the pathway is induced, the proteins assemble into a complex, and Hog1 is phosphorylated, which causes it to modulate gene expression via translocation to the nucleus. b) Schematic of the DHFR-PCA screen with a DMS mutant library. Single residue variants of *PBS2* were inserted into the yeast genome at the native locus using CRISPR-Cas9 genome editing. Sho1 and Pbs2 were tagged with complementary DHFR fragments. In cases where the Pbs2 variant does not interact with Sho1, the DHFR fragments remain separated. Upon interaction between a Pbs2 variant and Sho1, the DHFR fragments combine to reconstitute the functional DHFR, and allow growth in the presence of methotrexate (MTX). More strongly interacting partners allow the cell to proliferate more rapidly (Freschi *et al.* 2013).

Results

<u>Scan of the region surrounding the Pbs2 motif reveals that few mutations modify binding to Sho1</u>
The interaction between Sho1 and Pbs2 is modulated by an SH3 domain on Sho1 and the canonical binding motif on Pbs2, which is a type I motif with the sequence KPLPPLP (Maeda *et al.* 1995). Previous computational work has suggested that the region surrounding binding motifs could play an important role in modulating binding, including in SH3 interactions and in particular in the Sho1-Pbs2 interaction (Stein and Aloy 2008; Kelil *et al.* 2016). To determine which Pbs2 residues play a role in binding, we undertook a Deep Mutational Scan (DMS) on a region of 56 codons comprising the binding motif and its surrounding region. A DNA library of every possible single codon mutant was created for the 56 codon stretch, and inserted into the *S. cerevisiae* genome at the PBS2 locus using a CRISPR Cas-9 based approach, replacing the wild-type sequence (Dionne *et al.* 2021). The *PBS2* gene was fused with a Dihydrofolate Reductase Protein-Fragment 3 (DHFR F[3]), and these cells were mated with cells containing a DHFR F[1,2] fusion with *SHO1*, in order to obtain diploid strains containing complementary DHFR F fusions on both interaction partners. An interaction between Sho1 and Pbs2 brings the two DHFR fragments into contact, forming a functional DHFR which allows cell division even in the presence of the drug methotrexate (MTX) (Figure 1b). Growth in the presence of MTX in a strain with both DHFR fragments is therefore a proxy for the amount of Sho1-Pbs2 complexes forming, which depends on both the binding affinity and the local abundance of the interacting proteins (Tarassov *et al.* 2008; Freschi *et al.* 2013). As the DHFR fusions are located at the C-terminus of proteins, nonsense mutations introduced in the mutant library lead to no DHFR fragment expression.

To measure the interaction strength of the DMS library, the mutants were pooled and DHFR-PCA competition assays were done in media containing both MTX and 1 M of sorbitol, in order to induce the HOG pathway (Ferrigno *et al.* 1998). The resulting interaction score was calculated as the normalization of log-2-fold change of variant frequencies before and after selection for the interaction using MTX, as detected by targeted sequencing of the *PBS2* locus. Interaction scores were normalized in each replicate, so that scores of 0 are equivalent to wild-type Pbs2, while interaction scores of -1 correspond to the median of all nonsense mutants. Mutants which possess a higher interaction strength than wild-type Pbs2 therefore have a positive interaction score, while mutants which possess a lower interaction score than wild-type Pbs2 have a negative interaction score.

Many mutations in the canonical motif change the interaction strength. However, we found that most mutations in the region surrounding the motif did not affect binding of Sho1, apart from a short section neighboring the canonical motif itself (Figure 2a, Figure S1, Figure S2, Table S1). This confirms the strong role of the motif in the interaction and validates the computational predictions that extended the role of binding to include neighboring residues (Stein and Aloy 2008; Kelil *et al.* 2016). Since the effect on binding was strongly position dependent, we categorized different sections of Pbs2 as follows: the canonical binding motif is the conserved type I binding motif sequence situated in positions 93 to 99, the extended motif is composed of the canonical motif and the neighboring residues which have a strong impact on binding from positions 85 to 99 (Figure 2b).

DHFR-PCA reports on the amount of protein complex formed, which depends on affinity and local
protein abundance. To differentiate between these two parameters, we measured the effect of
mutations on a Pbs2 interaction to a partner that does not depend on the SH3 binding motif. Hog1
binds Pbs2 distally to the SH3 binding motif and independently of Sho1 binding (Murakami *et al.*
2008) (Figure 1a). So, measuring Hog1-Pbs2 binding allows us to control for Pbs2 local
abundance. Pbs2 mutations that only affect Sho1 binding should not have a detectable effect in
the Hog1-Pbs2 DHFR-PCA, while Pbs2 mutations that affect Pbs2 stability or local abundance
will have an effect in the Sho1-Pbs2 as well as the Hog1-Pbs2 DHFR-PCA.

We found that most mutations in the surrounding region of the Pbs2 motif had little effect on Hog1
binding (Figure 2a). Furthermore, mutants that had a negative effect on Hog1-Pbs2 binding also
had a negative effect on Sho1-Pbs2 binding, suggesting that the reduction in signal results from
a loss of local abundance of Pbs2 (Figure S3). However, most mutations that affected Sho1
binding did not impact the Hog1 interaction, and therefore did not change the local abundance of
Pbs2. Strikingly, only mutations in the residues comprising the extended motif had Sho1-
interaction specific effects, while mutations outside the extended motif had either little effect or
destabilizing effects.

237
238
239

241 Figure 2. <u>Mutations in Pbs2 within and around the binding motif affect binding to Sho1</u>.
242 a) Effect of Pbs2 mutations in the surrounding region of the SH3 binding motif (interaction score,
243 normalized log2-fold change before and after selection using MTX), with either Sho1 or Hog1.

The wild-type residues of Pbs2 are indicated for each position at the top of the figure. The positions comprising the extended motif are highlighted in blue, while the positions in the canonical binding motif have the wild-type residue indicated in bold. Boxplots show the distribution effects for each residue mutated to, and each residue is measured in between 3 and 18 replicates. Nonsense mutants omitted. b) A predicted structure (AlphaFold2-Multimer) of the SH3 domain of Sho1 (green) in complex with the Pbs2 region surrounding the motif (burgundy). The extended motif in positions 85 to 99 is colored in light blue. c) Distribution of interaction scores between Sho1 and Pbs2 single mutants in the extended motif DMS library (positions 85 to 100). Mutations are colored and placed on the y-axis by the nature of the mutation, either a change to a stop codon (nonsense), a change to the wild-type codon or a synonymous codon (WT or silent) or a change to a codon coding for a different residue (missense). The missense mutants were compared to the WT and silent mutations, using a Mann-Whitney U test with false discovery rate correction. They were grouped into 3 categories: Significantly weaker (p-value < 0.05 and negative interaction score), Not significantly different (p-value > 0.05) or Significantly stronger (p-value < 0.05 and positive interaction score).

Extended motif DMS library identifies mutations modulating interaction strength

In light of these results, we decided to focus solely on the extended motif of Pbs2, comprising codons 85 to 99, and adding position 100 to verify no effects were taking place in the residue immediately following the canonical motif. The DHFR-PCA on the initial DMS library of the entire region surrounding the motif took place in a diploid strain. Consequently, a large part of Pbs2 binding occurred with the Sho1 copy which did not possess the DHFR F[1,2] fusion, and these interactions did not contribute to the DHFR-PCA signal. Since this limited our ability to precisely measure the Sho1-Pbs2 interaction, we built a new haploid strain containing both DHFR F[1,2] on Sho1 and DHFR F[3] on Pbs2. We also constructed a Hog1-DHFR F[1,2]/Pbs2-DHFR F[3] strain to once more measure the local abundance of Pbs2 mutants. These new strains capture all Pbs2 binding to Sho1 or Hog1, therefore allowing for more sensitive measurements. We inserted a second Pbs2 DMS library, comprising all possible single mutations in Pbs2 positions 85 to 100, into the two haploid DHFR strains, using the same CRISPR-Cas9 approach as the first library. The interaction strength of the variants was measured using the DHFR-PCA as previously (Figure S4, Figure S5). At this point, we filtered out the 12 mutants which had an interaction score with Hog1 significantly different from wild type and synonymous Pbs2 mutants (Two-sided Mann-Whitney U test with false discovery rate corrected p-value < 0.05) from our analysis (Figure S6). As before, the goal was to disregard mutants which modulated the interaction through change of local abundance or stability, which would affect both the Sho1 and Hog1 interaction.

The impacts on the Sho1-Pbs2 interaction were varied (Figure 2c, Figure S7, Table S2). Few mutants had interaction scores as low as the nonsense mutants, suggesting that few single mutations are able to completely prevent Sho1-Pbs2 binding. By comparing missense mutant interaction scores to the wild type and synonymous mutants scores, we found that many missense mutants had significantly stronger or weaker scores than the wild-type Pbs2 sequence (Two sided Mann-Whitney U test with false discovery rate corrected p-value < 0.05). 18 mutants (6.5% of all missense mutants) interacted significantly more strongly with Sho1 than wild-type Pbs2, while

173 mutants (62.7% of all missense mutants) interacted significantly more weakly than wild-type Pbs2. The impact of mutants that reduced the interaction scores was generally greater.

As the Sho1-Pbs2 interaction is strengthened under osmotic stress (Maeda *et al.* 1995; Posas and Saito 1997; Saito and Posas 2012), we also compared the interaction strength between these two proteins in the presence and absence of 1 M of sorbitol (Figure S8). We found that interaction scores in the presence and absence of sorbitol correlate strongly (Spearman's rho 0.98, p-value $< 2.2X10^{-16}$). The effects on binding were stronger in the presence of sorbitol, as a result of the induction of the HOG signaling pathway. When sorbitol is present, strongly interacting Pbs2 mutants are even more often in contact with Sho1, increasing the number of reconstituted DHFR complexes, while weakly interacting mutants are even further outcompeted by the strongly interacting mutants. Still, the high correlation in interaction scores between these two conditions indicates that Pbs2 mutants behave similarly in the presence and absence of osmotic stress, and that the general strengthening or weakening effect of mutations does not depend on the activation of the signaling pathway.

To validate the pooled measurements, we individually reconstructed 24 mutants spanning the range of measured interaction scores in Pbs2-DHFR F[3] strains which were mated with the Sho1-DHFR F[1,2] strain. We then measured the growth rate of these mated strains in growth assays in DHFR-PCA medium. As with the competition assay, the growth rate is a proxy for interaction strength, which reflects affinity in the absence of a change in abundance. The growth curve results correlate strongly with the interaction scores (Spearman's rho 0.88, p-value $7.3x10^{-5}$ for interactions in presence of 1 M sorbitol, spearman's rho of 0.73, p-value $2.9x10^{-6}$ for interactions in absence of sorbitol), thus upholding the results of the competition assay (Figure S9a). To validate further that the effects on interaction strength were not the result of noise or the result of another source, we also performed a smaller scale competition assay DHFR-PCA, individually reconstructing 67 mutants with varying interaction scores (Figure S10, Table S3). The measurements of this validation assay correlated strongly with the assay on the DMS library of the extended motif (Spearman's rho 0.98, $p < 2.2X10^{-16}$ for interactions in presence of 1 M sorbitol, spearman's rho of 0.98, $p < 2.2X10^{-16}$ for interactions in absence of sorbitol), thus validating the results of the measurement of the DMS library (Figure S9b). We also measured the effect of mutations on cell proliferation by constructing strains with the same 67 mutations, but with no DHFR tags. Cell growth and division should thus only depend on the effect of the mutations, and not interaction strength. We undertook a pooled competitive growth assay in SC synthetic media with 1 M of sorbitol, measuring how much each mutant proliferated. We found that all mutations except nonsense mutations had no significant effect on cell proliferation (Welch's t-test false discovery rate corrected p-value < 0.05) (Figure S11, Table S3). Interestingly, mutant I87W, which is the strongest interacting mutant, seems to have a detectable, though not statistically significant, deleterious effect on cell proliferation.

Structure prediction suggests a secondary contact between extended motif and Sho1

A crystal structure of the SH3 domain of Sho1 in complex with a 9 residue-long segment of Pbs2 reveals that the canonical motif occupies the entirety of the canonical binding pocket of Sho1 (Kursula *et al.* 2008). However, many mutations outside the canonical motif have a strong effect on binding. These residues must affect the interaction of Pbs2 with Sho1, but it is unclear how. One possibility is that certain residues in the extended motif are interacting with Sho1 outside of the canonical binding pocket of the SH3 domain. This would not be unprecedented, as NMR structures of different SH3 domains interacting with long peptides reveal interactions involving residues outside of the canonical binding pocket, which affected the strength and specificity of the interaction (Rickles *et al.* 1995; Feng *et al.* 1995; Stollar *et al.* 2009; Takaku *et al.* 2010; Gorelik and Davidson 2012). No experimental structure has been determined for the entirety of Sho1 or Pbs2, so we used structure predictions from AlphaFold-Multimer (Jumper *et al.* 2021; Evans *et al.* 2022) to determine if binding outside the canonical binding pocket was likely to occur (Figure S12). By modeling the entire Sho1 protein along with the surrounding region of the Pbs2 motif (positions 71-126), we found potential interactions outside of the canonical binding pocket of Sho1, mediated by residues neighboring the canonical binding motif of Pbs2 (Figure 3a). In the structural model, a hydrophobic pocket is formed between the SH3 domain and a loop of Sho1, which interacts with an alpha helix composed of Pbs2 positions 84 to 92 (Figure 3b). Interestingly, all but one of the mutants in the extended motif which significantly increase interaction strength are found in the positions comprising the helix. Certain Pbs2 residues towards the C-terminus relative to the canonical motif are also predicted to be in proximity to Sho1, with 2 residues within 2 Å of Sho1, but as mutations in these positions did not have substantial effects on binding (Figure 2a), we decided to concentrate on residues in and near the extended motif.

To help explain how different mutations could modify binding, we predicted the structures of the 5 Pbs2 variants possessing the strongest interaction strength with Sho1, using the same method as with wild-type Pbs2 (Figure S12). In particular, I87W, which is the strongest interacting mutant measured, positions its side-chain in close contact (1.77 Å) with the non-canonical hydrophobic pocket, and particularly in a sub-pocket formed of both SH3 and non-SH3 residues. Additionally, it is predicted to form a hydrogen bond with the aspartic acid side chain in position 333 of the Sho1 sequence (Figure 3c). These two factors potentially explain its strong interaction strength relative to all other Pbs2 variants. Of the other strengthening variants tested, H86W, V91L and V91M all place a hydrophobic side chain in the non-canonical hydrophobic pocket, potentially strengthening the interaction. For the last mutant tested, Q89D, the hydrophilic side-chain faces away from the hydrophobic pocket, thus not disrupting the hydrophobic interaction, although it is unclear how the interaction is strengthened (Figure S13).

We further reasoned that substituting a non-hydrophobic residue in one of the Pbs2 positions which is predicted to contact the non-canonical hydrophobic pocket could interrupt this interaction, and weaken the affinity. Indeed, when classifying the mutations in the alpha helix by their hydrophobicity (Monera *et al.* 1995), we find that replacing the initially hydrophobic residues in positions I87, I90, and V91 with a non-hydrophobic residue reduces the binding strength. In

general, a mutation to another hydrophobic residue in these positions has a weaker negative impact on binding, and in the case of V91, many hydrophobic substitutions increase the interaction strength (Figure 3d). Based on the results of the hydrophobic or non-hydrophobic substitutions, we reasoned that these three positions situate their side chains in the non-canonical hydrophobic binding pocket. This finding helps explain the strong interactions strength of Pbs2 mutant I87W, by showing the important role of hydrophobic interactions in position 87. The predicted models of interactions between Sho1 and different Pbs2 variants, along with our interaction score measurements suggest that residues outside the canonical binding motif of Pbs2 can interact directly with Sho1, and thus they have the capacity to modulate affinity.

Figure 3. <u>Contacts mediated by positions outside the canonical binding motif of Pbs2 could modulate binding</u>
a) Structure of Sho1 in complex with residues 71 to 126 of Pbs2 as predicted by AlphaFold2-Multimer (Jumper *et al.* 2021; Evans *et al.* 2022). The structure is positioned to show the helix on Pbs2 interacting with a hydrophobic pocket on Sho1, while the canonical motif interacts with the canonical binding pocket. The structure is colored by the different regions of Sho1 and Pbs2, as indicated on the panel. The orientation of Pbs2 is shown by annotations indicating which end is nearer the N-terminus and which end is nearer the C-terminus. b) Hydrophobicity map of the predicted structure of Sho1, with high molecular lipophilicity potential (MLP) indicating a hydrophobic surface, and low MLP indicating a hydrophilic surface. MLP calculated using

Chimera X 1.8. Positioning of both proteins, and coloring of Pbs2 as in panel a). c) Close-up of the predicted structure of Sho1 with residues 71 to 126 of Pbs2 with mutation I87W. The side-chain of W87 is shown, and a hydrogen bond (red dotted line) is predicted with D333 of Sho1. Coloring of the proteins is as described in panel a). The orientation of Pbs2 is shown by annotations indicating which end is nearer the N-terminus (N) and which end is nearer the C-terminus (C). d) Boxplot of interaction scores of Pbs2 mutants for selected positions from the DHFR-PCA of the DMS library of the extended motif, based on the hydrophobicity of the substituted residue. The wild-type residue of Pbs2 for every position is indicated and colored according to its hydrophobicity. Nonsense mutants omitted. Hydrophobicity for the different amino acids was obtained from (Monera *et al.* 1995), and F, I, W, L, V, M, Y, C, and A were classified as hydrophobic, T, H, E, S and Q were classified as neutral and R, K, N, G, P and D were classified as hydrophilic.

Comparison of Sho1-Pbs2 binding to other SH3-motif pairs

As we propose a role in modulating interaction strength for the extended motif and the additional binding pocket of Sho1, we wondered if these structures could play a part in modulating the binding specificity of Pbs2 to Sho1. If this additional pocket were a unique structure among yeast SH3 domains, that could increase the specificity of binding, as the Pbs2 extended motif could be uniquely tuned to bind this unique feature. First, we generated a multiple sequence alignment of yeast SH3 domain sequences with 25 adjacent residues on each side, to verify the conservation of the amino acids forming the additional binding pocket. This window includes all residues predicted to be in proximity to Pbs2 in Sho1. Apart from a few key residues in the SH3 domain, we found no significant conservation of the positions which in Sho1 are predicted to be in proximity (< 5 Å) to Pbs2 positions situated in the extended motif, but outside of the canonical motif (Figure S14a). We reasoned that the structure of the additional binding pocket might be conserved, even if the particular residues were not, so we used the MUSTANG (Konagurthu *et al.* 2006) tool to carry out a multiple structural alignment on structure predictions of yeast SH3 domains with 25 neighboring residues, obtained from the AlphaFold Protein Structure Database (Varadi *et al.* 2022). We found that only the positions in the SH3 domains were structurally conserved, with no significant structural conservation for the positions outside the SH3 domains (Figure S14b), including those forming the additional binding pocket. We therefore conclude that the additional binding pocket on Sho1 is a unique structural feature.

Even without a conserved structure of non-SH3 sequences, there are reports of three other yeast SH3 domains binding extended motifs: Abp1, the second SH3 domain of Bem1 (referred to as Bem1-2) and Nbp2 (Stollar *et al.* 2009; Takaku *et al.* 2010; Gorelik and Davidson 2012). We tested whether there was an *in vivo* interaction between these SH3 domains tuned to bind extended motifs, and the extended motif of Pbs2. To do so, we once again used a DHFR-PCA assay, though this time measuring colony growth on solid medium as is routinely done for this (Tarassov *et al.* 2008), with the SH3-containing proteins tagged with DHFR F[1,2] and Pbs2 tagged with DHFR F[3]. We used two variants of Pbs2: a wild-type, and a control where the

445 extended motif and the region surrounding it (positions 71 to 126, the same region covered in the
446 initial DMS library) were replaced by a neutral, flexible stuffer sequence.
447
448
We found that Abp1 and Nbp2 interact with Pbs2 with approximately the same strength as Sho1
does, while Bem1-2 does not detectably interact with Pbs2 (Figure 4a). However, Abp1 and Nbp2
also interact with the stuffed Pbs2 variant at the same level as the wild-type variant, indicating
that the interaction is not mediated by the Sho1-binding motif. Nbp2 is known to bind a second,
distal motif on Pbs2 through its SH3 domain (Mapes and Ota 2004), while Abp1 must bind Pbs2
through an unknown mechanism involving a different site than the one bound by Sho1. We
additionally measured the interaction with all other known yeast SH3-containing proteins, and
found that apart from Abp1 and Nbp2, all proteins interacted significantly less strongly with either
Pbs2 variant than Sho1 with wild-type Pbs2 (One sided Mann-Whitney U test with false discovery
rate corrected p-value < 0.05) (Figure S15). Additionally, measuring the interaction of Pbs2 with
a Sho1 variant where the SH3 domain was replaced by the neutral, flexible stuffer sequence
showed an interaction strength above the level of stuffed Pbs2 with the same Sho1 variant,
confirming the role of the non-SH3 portion of Sho1 in binding the extended motif of Pbs2 (Figure
S15, Sho1_stuffed). Together, these findings show that the extended motif of Pbs2 specifically
binds to Sho1 both inside and outside the SH3 domain, and doesn't interact with other SH3-
containing proteins *in vivo*.
465
To determine whether we could explain the specificity of Pbs2 binding, we compared the predicted
structure of the Sho1-Pbs2 interaction to NMR structures of the Abp1-Ark1, Bem1-2-Ste20, and
Nbp2-Ste20 interactions, all of which include long peptides with lengths comparable to the
extended motif of Pbs2. The structures of the four SH3 domains and peptides are very similar,
with all peptides binding in the canonical binding pocket of their respective SH3 domains, and
extending towards the space corresponding to what we identified as the additional binding pocket
on Sho1 (Figure 4b, Figure S16). However, Abp1 and Nbp2 only bind the extended motif through
the SH3 surface, while Bem1-2 has a second, non-SH3 interface that forms a pocket with the
SH3 domain into which the extended motif fits, much like Sho1. However, this surface has no
structural or sequence similarity with Sho1 and has been previously identified as the Cdc42
interacting (CI) region (Takaku *et al.* 2010). Contrary to Sho1, this additional binding region is
located C-terminal to the SH3 domain.
478
There is little sequence conservation between the extended motifs of Pbs2, Ark1, and Ste20,
apart from the conserved positions of the canonical motifs. These extended motifs can be aligned
using a previously developed standard numbering, where the first conserved proline is designated
0, residues preceding the motif are numbered with decreasing negative numbers, and residues
following the motif are numbered with increasing numbers (Lim *et al.* 1994) (Figure 4c). The
binding specificity exhibited by Pbs2 may depend on the possible contacts with the different SH3
domains, and the non-SH3 surfaces in the cases of Sho1 and Bem1-2. Examining the different
motifs and their contacts with their respective interaction partners, we can compare Pbs2 to the
two other extended motifs, and in particular the positions which are in the extended motif but
outside the canonical binding motif, to establish determinants of specificity.

Foremost, Pbs2 binds using residues more distal from the canonical motif than either Ste20 or
Ark1. Contacts are predicted for Q(-14) as well as H(-10) on the predicted Sho1-Pbs2 structure,
which is not the case for any other interaction. In Pbs2, I(-9), which can have a large impact on
binding when mutated, contacts the hydrophobic residues L290 and Y293 on the non-SH3 surface
of Sho1. The equivalent position on Ste20, G(-9) contacts the polar residues Q240 and K243 in
the non-SH3 surface of Bem1, which would be unfavorable for the isoleucine in Pbs2. In Ark1
binding to Abp1, the residue at position (-9) contacts the SH3 domain (Figure 5a).

Position (-7) is important for binding in both Ste20 (F(-7)) and Ark1(L(-7)). When binding to their
respective partners, including both Bem1-2 and Nbp2 for Ste20, mutation of position (-7) to an
alanine reduced the strength of binding by at least an order of magnitude. Sequentially mutating
all positions in the extended motif to an alanine showed that position (-7) had the most impact of
any position outside the canonical motif in all cases (Stollar *et al.* 2009; Gorelik and Davidson
2012). In Pbs2, position (-7) is occupied by a polar glutamine, as opposed to the hydrophobic
residues found in the same position in the other two peptides. Also, in the predicted Sho1-Pbs2
structure, the side chain of Q(-7) is pointed away from Sho1 and is not predicted to have any
contacts, while in the other structures, the equivalent side chains are pointed towards the SH3
domains (Figure S17a).

Position (-6) also marks a difference, specifically between Pbs2 and Ark1. In Pbs2, I(-6) contacts
Y42 on the SH3 domain. In Ark1, H(-6) forms a salt bridge with an aspartic acid situated on a
different loop of the Abp1 SH3 domain (Figure 5b). The structural alignment of SH3 domains
indicates that Sho1 has a gap at the position of the salt bridge forming aspartic acid (Figure S14b,
position 112). The last major difference is in position (-4), where the residues in both Ste20 and
Ark1 make contact with an SH3 domain, while N(-4) in Pbs2 is not predicted to have any contacts.
As with position (-7), the polar side-chain is pointed away from the SH3 domain (Figure S17b).

Overall, we can see that the Pbs2 extended motif is very different from the other extended motifs
known to bind SH3 domains in yeast. The additional contacts in positions (-14) and (-10), the
differing contacts in positions (-9) and (-6), and the missing contact in position (-4) and especially
the important missing contact in position (-7) may preclude Pbs2 from binding to SH3 domains
other than Sho1. If this is truly the case, the presence of a non-conserved portion in the extended
motif allows much more differentiation in binding than solely the canonical motif.

c

Standard numbering	-14	-13	-12	-11	-10	-9	-8	-7	-6	-5	-4	-3	-2	-1	0	1	2	3	4	5	6	7
Pbs2	Q	S	P	K	H	I	Q	Q	I	V	N	K	P	L	P	P	L	P	V	A	G	S
Ark1				K	P	K	L	H	S	P	K	P	P	P	T	P	K	T	K	K	A	
Ste20	S	S	S	A	N	G	K	F	I	P	S	R	P	A	P	K	P	P	S	S	A	
Pbs2 position	82	83	84	85	86	87	88	89	90	91	92	93	94	95	96	97	98	99	100	101	102	103

Figure 4. The Pbs2 motif binds specifically to Sho1, despite similarities of other SH3 domains and
motifs
a) Interaction score of wild-type Pbs2 (green) or Pbs2 where positions 76 to 126 have been
replaced with a flexible stuffer sequence (Stuffed, white) with select yeast SH3-containing
proteins. All replicates shown. Scores were normalized, with 1 being the median of Sho1 - wild-
type Pbs2 interactions and 0 being the median of Sho1 - stuffed Pbs2 interactions. b) Root mean
square deviation of atom locations between matching backbone atoms in each pair of structures,
as calculated using the R package bio3d (Grant *et al.* 2006). c) Sequences of Ste20, Ark1 and
Pbs2, with standard numbering according to the motif, to compare equivalent positions (Lim *et al.*
1994). For Ste20 and Ark1, the positions included are those present in the structures in panel b,
with Pbs2 positions chosen to overlap all represented positions.

    Figure 5. Differing contacts between SH3-extended motif pairs
a) Visualization of the side-chains and contacts of position (-9) in the Sho1-Pbs2 predicted
structure, Abp1-Ark1 structure (PDB accession 2RPN), Nbp2-Ste20 structure (PDB accession
2LCS) and Bem1-2-Ste20 (PDB accession 2RQW). Sho1-Pbs2 predicted structure colored as in
figure 3a, with the SH3 domain of Sho1 in green, the non-SH3 portions of Sho1 in orange, the
Pbs2 extended motif in light blue and the rest of Pbs2 in burgundy. All four structures were
superimposed using the Chimera X tool "Matchmaker" (Pettersen *et al.* 2021) to minimize root
mean square deviance of backbone atom distances of the SH3 domains with Sho1, then pictures
were captured separately. Position (-9) and its contacts are kept opaque while the rest of the
structures are semi-transparent. b) Visualization of the side-chains and contacts of position (-6)
of the same structures as in panel a). All four structures were superimposed using the Chimera X
tool "Matchmaker" (Pettersen *et al.* 2021) to minimize root mean square deviance of backbone
atom distances of the SH3 domains with Sho1, then pictures were captured separately. Position
(-6) and its contacts are kept opaque while the rest of the structures are semi-transparent.

Discussion

Binding motifs face the double challenge of binding their partner with appropriate strength while
avoiding spurious interactions with non-partner proteins. Using deep mutational scanning libraries
of the Pbs2 binding motif and its surrounding sequences, we measured the impact of single
residue mutations on interaction strength with Sho1. We also predicted the structure of the Sho1-
Pbs2 interaction, and compared it to known structures of SH3-mediated interactions. We use
these comparisons to propose a number of ways in which the Pbs2 extended motif may bind
specifically to Sho1, and not other SH3 domains.

One important consideration in domain-peptide interactions is the potential trade-off between
interaction affinity and specificity. Previous studies on SH2 and PDZ domains, both domains that
bind short motifs similarly to SH3 domains, have put forth that mutants which bind with higher
affinity also possess lower specificity (Ernst *et al.* 2010; Haslam and Shields 2012; Kaneko *et al.*
2012; Karlsson *et al.* 2016), suggesting that high affinity and high specificity may be incompatible.
This represents an intuitive idea: a protein capable of having strong interactions with a certain
motif will also have more frequent and stronger spurious interactions with motifs of the same
family, which are physically and chemically similar. On the other hand, a computational assay
studying binding motifs, including SH3 binding motifs, found that an increase in affinity can lead

to an increase in specificity (Kelil *et al.* 2016). Other works have emphasized the importance of
the residues surrounding the canonical binding motif — in this case defined as K/RXXPXXP,
corresponding to positions 93 to 99 of Pbs2 — in determining both affinity and specificity (Li 2005;
Ivarsson and Jemth 2019). In fact, another study computed that nearly 30% of binding energy
from the binding motif was contributed by residues outside the canonical motif, and suggests that
the residues outside the canonical motif mostly play a role in determining specificity (Stein and
Aloy 2008). In our experimental work, we found that all mutations which significantly increase the
interaction strength between Sho1 and Pbs2 are situated outside of the canonical motif.
Consequently, the wild-type canonical motif could be already optimized for the strongest possible
interaction strength. This suggests different roles for the canonical motif and its neighboring
residues in determining affinity and specificity, with canonical motif residues optimized and
therefore more constrained, while the neighboring residues are less constrained and able to
increase affinity and specificity when mutated.
593
One interesting possibility that may explain the higher binding strength conferred by mutations to
certain residues outside the Pbs2 canonical binding motif is that the mutated residues may
interact with either residues outside the canonical binding pocket in the SH3 domain or even with
non-SH3 surfaces on Sho1. Non-binding pocket interactions have previously been observed in
multiple SH3-mediated interactions (Lee *et al.* 1996; Dalgarno *et al.* 1997; Li 2005; Stollar *et al.*
2009; Takaku *et al.* 2010; Gorelik and Davidson 2012; Gaussmann *et al.* 2024). Using predicted
protein complex structures generated by AlphaFold-Multimer, we were able to hypothesize that a
hydrophobic pocket is formed between the SH3 domain and a non-SH3 loop of Sho1, which
residues of the extended motif can fill. The presence of such a pocket can explain the increased
affinity of many Pbs2 mutants in the extended motif, as predicted structures of these models show
that substituted hydrophobic side chains can insert themselves into the predicted pocket. Perhaps
the disordered region encompassing the Sho1-binding motif can place itself in such a way to
maximize the hydrophobic interaction between any mutated side chain and the hydrophobic
pocket. These predicted structures may represent only one possible form of the Sho1-Pbs2
interaction, which may in reality be dynamic due to the disordered nature of the region surrounding
the Pbs2 motif. Nevertheless, they offer a potential example of binding mediated by positions
outside the canonical motif. By removing the need for affinity to be dependent only on canonical
binding motifs, the residues surrounding motifs could bind to protein surfaces which are less
conserved among SH3 domains, and thus easier to differentiate. The presence of unique
components, such as extended motifs or extended binding surfaces could therefore contribute to
specificity among domains of the same family. The hydrophobic pocket formed by both SH3 and
non-SH3 parts of Sho1 may represent an example of a unique structural feature which may help
differentiate Sho1 from other SH3 domains, and allow increased affinity to develop in Pbs2 without
increasing the affinity to other SH3 containing proteins. This proposed model of binding also
involves a larger number of residues in tuning affinity and specificity, potentially uncoupling affinity
and specificity and therefore avoiding a trade-off between these two phenotypes. Affinity and
specificity may only be linked for highly conserved residues and structures, such as the canonical
binding motif or the canonical SH3 binding pocket, but for less constrained residues, may develop
independently.
623

Studies on interaction domains have shown that SH3 domain mediated interactions are context
specific, meaning that the properties of the interactions depend not only on the domains
themselves, but also on the protein context. This context can be composed of many factors,
including the non-SH3 sequence of the protein, other proteins interacting with the SH3 containing
protein, and cellular localization (Dionne *et al.* 2022). SH3 mediated interactions therefore do not
simply depend on the SH3 sequence. For example, when inserting an SH3 domain into a different
protein or expressing an SH3 domain without its host protein, the SH3 domain's interaction profile
substantially changes (Dionne *et al.* 2021; Lemieux *et al.* 2024; Dibyachintan *et al.* 2024). Thus,
Pbs2 mutations may increase or decrease binding strength not only with the Sho1 SH3 domain,
but also within the Sho1 context. It is not entirely clear how Pbs2 mutations modulate binding
within the Sho1 context, but the binding of the Pbs2 extended motif to non-SH3 residues could
explain a large part of this phenomenon. Other causes could include stereochemical adjustments
of SH3 binding loops or changes which contribute to stabilizing or destabilizing interactions with
other Sho1 binding partners. There are observations of other proteins, namely Ste11 and Ste50,
interacting with the Sho1 SH3 domain outside of the canonical binding pocket (Zarrinpar *et al.*
2004; Tatebayashi *et al.* 2006). Changes to the Pbs2 peptide may affect the stability of the
interaction of these proteins, which are also involved in the osmotic stress response pathway.
641
Previous work has already focused on the binding of Pbs2 motif variants (Zarrinpar *et al.* 2003).
We replicate some of the findings of this previous work, for example, showing that mutations near
the canonical Pbs2 motif can cause both gain and loss of binding strength. This study also finds
that a promiscuous double mutant of the Pbs2 motif has lower fitness (Zarrinpar *et al.* 2003). We
find that, apart from nonsense variants, no mutants confer a significant loss of fitness. However,
Pbs2 mutant I87W has a noticeable, though not statistically significant, negative effect on cell
fitness (Figure S11). I87W is also the mutant which most increases the interaction strength, which
may be a cause of the slightly reduced fitness. Mutations which increase affinity have been
suggested to have deleterious effects, by reducing the dynamicity of the pathway, and impeding
dissociation of the partners (Wang *et al.* 2018). For the Sho1-Pbs2 interaction, previous work has
found that a mutation in the Sho1 SH3 domain that increased affinity also led to a small reduction
in cell growth (Marles *et al.* 2004), indicating that the wild-type affinity may be already optimized,
and that stronger binding may disrupt signaling.
655
In conclusion, we show that mutations in positions neighboring the canonical binding motif of Pbs2
can both increase and decrease binding strength with Sho1. We propose a role for residues
outside the canonical Pbs2 binding motif in increasing binding affinity independently of binding in
the canonical SH3 binding pocket. We hypothesize that Sho1 is similar to certain other yeast SH3
domains which bind longer motifs through additional binding pockets (Stollar *et al.* 2009; Takaku
et al. 2010; Gorelik and Davidson 2012), and propose mechanisms through which an extended
binding motif could better differentiate binding to different SH3 domains. Going forward, studies
of SH3-motif binding should consider a wider range of residues, both on the SH3 domain and
surrounding the motif. Our work, along with previous work by other groups, suggests that *in vitro*
work using isolated domains and shortened peptides does not capture the full picture of domain-
motif interaction. By understanding and harnessing the potential of extended motif binding,

researchers may be better able to understand the impact of motifs in cell signaling and disease,
and better able to use motif binding in technical and medical applications.
669
670

Methods
671

672
<u>Growth conditions</u>
Escherichia coli cells were grown in 2YT medium (1% yeast extract, 1.6% tryptone, 0.2% glucose,
0.5% NaCl, and 2% agar for solid plates) with shaking at 37 °C for liquid cultures. When specified,
100 µg/mL of ampicillin (Amp, Bioshop Canada) were added as a selection agent.
Saccharomyces cerevisiae cells were grown in YPD medium (1% yeast extract, 2% tryptone, 2%
glucose, and 2% agar for solid plates), synthetic complete medium buffered to a pH of 6.0 (0.174%
yeast nitrogen base without amino acids, without ammonium sulfate, 2% glucose, 0.134% amino
acid dropout complete, 0.1% monosodium glutamate, with 1% succinic acid and 0.6% NaOH as
buffer) or PCA medium (0.67% yeast nitrogen base without amino acids and without ammonium
sulfate, 2% glucose, 10% liquid drop-out without adenine, methionine, and lysine, and 200 µg/ml
methotrexate (MTX, Bioshop Canada) diluted in dimethyl sulfoxide (DMSO, Bioshop Canada),
and 2.5% noble agar for solid plates), as specified. When specified, 200 µg/mL of G418 (Bioshop
Canada), 100 µg/mL of nourseothricin (Nat, Jena Bioscience) or 250 µg/mL of hygromycin B (Hyg,
Bioshop Canada) were used as selection agents. Also, when specified, 1 M of sorbitol (Bioshop
Canada) was added to either liquid or solid PCA medium. Yeast cultures were grown at 30 °C
with agitation at 250 rpm. See Table S4 for details of all growth media used.
689
<u>Cloning</u>
All plasmids were constructed by Gibson assembly. As template for the *PBS2* DMS library
construction (pUC19-Pbs2, see below), a pUC19 plasmid backbone was amplified (Addgene
#50005, primers 1F and 1R, all primers detailed in Table S5), then digested for 1 hour at 37 °C
using SacI (New England Biolabs). This linearized plasmid was used for Gibson assembly with
an insert composed of a 248 base pair region of *PBS2* amplified from genomic DNA, using primers
that add homology arms (Primers 2F and 2R). For the CRISPR-Cas9 based insertion of
sequences, the pCAS plasmid (Addgene #60847) was modified as previously described (Ryan *et*
al. 2016) to change the sgRNA to target either the Sho1 binding motif of Pbs2 (pCAS-Pbs2,
primers 3F and 3R), the SH3 domain of Sho1 (pCAS-Sho1, primers 4F and 4R), or the stuffer
sequence which was inserted into *PBS2* (pCAS-stuffer, primer 5F and 5R) (Dionne *et al.* 2021).
Proper plasmid construction was verified using Sanger sequencing (Primers 6F and 6R) (Messing
1983).
703
704
705 <u>Strain construction</u>
706
All strains used are summarized in Table S6. Yeast strains were constructed with either the F[1,2]
or the F[3] fragments of an engineered murine DHFR protein fused to the C-terminus of different

proteins, to measure protein interaction strength as previously described (Tarassov *et al.* 2008;
Michnick *et al.* 2016). The transformed sequence contains the DHFR F[1,2] fusion sequence as
well as a nourseothricin resistance cassette, or the DHFR F[3] fusion sequence with a hygromycin
B resistance cassette. This allows selection of the strains with DHFR fragment fusions. The DHFR
F[1,2] tagged strains were derived from MATa strain BY4741 (*his3Δ leu2Δ met15Δ ura3Δ*), while
the DHFR F[3] strains were derived from MATα strain BY4742 (*his3Δ leu2Δ lys2Δ ura3Δ*). The
Hog1-DHFR F[1,2] strain and the Pbs2-DHFR F[3] strain used in the paper were taken from the
Yeast Protein Interactome Collection (Horizon Discovery) (Tarassov *et al.* 2008). The Sho1-DHFR
F[1,2] strain additionally had a 1X FLAG tag at the end of the fragment construct, and its
construction was described in a previous publication (Dionne *et al.* 2021). Briefly, the DHFR
F[1,2], the Nat cassette, a flexible linker and the FLAG tag sequence were amplified from the
pAG25-DHFR F[1,2]-linker-FLAG plasmid (Dionne *et al.* 2021) using primers with homology arms
matching the regions either side of the *SHO1* termination codon. The cassette was transformed
into BY4741 cells and integrated by homologous recombination. Further deletions were done in
the DHFR F[1,2] tagged strains in order to knock out *PBS2*, using the *LEU2* cassette from pUG37
(Gueldener *et al.* 2002), through homologous recombination (Primers 7F and 7R). Proper deletion
was verified by PCR (Primers 8 to 11) and sequencing (Gueldener *et al.* 2002). After the mating
of the DHFR F[1,2] MATa strains with the Pbs2-DHFR F[3] MATα, only one copy of *PBS2* is
present. This prevents the additional wild-type allele from influencing the measurements of
interaction strength of the variants.
729
The Pbs2 mutants of interest for the DHFR-PCA growth curves were constructed from the Pbs2-
DHFR F[3] fusion strain. First, the region surrounding the Sho1 binding motif of Pbs2 (codons 71
to 126) was replaced with a stuffer sequence (GGCGGAAGTTCTGGAGGTGGTGGT) that
translates into a flexible linker sequence (GGSSGGGG), by co-transformation of the pCAS-Pbs2
plasmid, and of a repair template consisting of the stuffer sequence flanked by homology arms
matching the sequences around codons 71 to 126 of *PBS2* (produced by amplifying the stuffer
sequence with primers 12F and 12R)*,* following a previously published protocol (Ryan *et al.* 2016).
This produced the Pbs2-stuffed-DHFR F[3] strain, which was used for the construction of the
individually reconstructed mutants, for the construction of the DMS libraries (see below), and also
as a control to measure Pbs2 binding to SH3-containing yeast proteins. The individually
reconstructed mutants contain all mutants used in a previous paper exploring Pbs2 mutants
(Zarrinpar *et al.* 2003), including a P94A+P97A double mutant. The mutants were individually
constructed by co-transforming the Pbs2-stuffed-DHFR F[3] strain with the pCAS-stuffer plasmid
targeting the aforementioned stuffer, and synthesized oligonucleotide sequences corresponding
to the desired mutation (Integrated DNA Technologies) (Table S7). The stuffed strain as well as
all reconstructed strains were verified by PCR and Sanger sequencing (Primers 13F and 13R).
The pCAS plasmids were purged by growth in liquid YPD without G418, and loss of the plasmid
was verified by lack of growth on YPD+G418 plates.
748
For the extended motif DMS library and the individually reconstructed mutants used in the
validation competition assays, three additional haploid strains were built, into which Pbs2 DHFR
mutations were inserted. The first two strains were Hog1-DHFR F[1,2]/Pbs2-stuffed-DHFR F[3]
and Sho1-DHFR F[1,2]/Pbs2-stuffed-DHFR F[3]. These were built by adding the DHFR F[1,2] tag

to either *HOG1* or *SHO1* in the MATα Pbs2-stuffed-DHFR F[3] strain previously constructed. The
DHFR F[1,2] moieties were amplified from plasmid pAG25-DHFR F[1,2]-linker-FLAG plasmid
using primers with homology arms to either the *HOG1* or *SHO1* locus (Primers 14 F to 15R)
(Dionne *et al.* 2021). Proper tagging was verified using PCR and Sanger sequencing (Primers 16
to 18) (Freschi *et al.* 2013; Dionne *et al.* 2021). The third strain was Pbs2-stuffed, which was used
to measure the proliferation of mutants without the effect of the DHFR fragment. This was
constructed in the haploid strain BY4742, using the same strategy to insert the stuffer into *PBS2*
as detailed above.

The selected Pbs2 mutants for the validation assay were constructed in these three same strains
by transformation with repair templates containing the mutations of interest, formed through fusion
PCR. Briefly, forward and reverse primers were designed for each mutant to be constructed,
which contained the mutated sequence instead of the wild-type sequence (Primers 19F to 110R).
These primers were used for separate PCR amplification of the *PBS2* sequence along with either
common forward or reverse primers (Primers 111F and 111R), to create two overlapping
fragments. These were combined in a second PCR using only the two common primers, which
fused the overlapping sequences together, to create a single fragment of 338 base pairs, which
was identical to *PBS2* except for the mutated codon of interest. These fragments were
transformed into DHFR F[1,2]/Pbs2-stuffed-DHFR F[3] strains using the CRISPR-Cas9 strategy
described above. Mutants were confirmed by PCR and sequencing (Primers 112F and 112R).
Most mutations were constructed in all three DHFR strains, but a certain number could only be
built in one or two of the strains. The successfully constructed mutants are listed in Table S6.

For the DHFR-PCA screen on solid media, the previously constructed wild-type Pbs2-DHFR F[3]
and Pbs2-stuffed-DHFR F[3] were used. The SH3-containing proteins DHFR F[1,2] strains were
obtained from the Yeast Protein Interactome Collection (Horizon Discovery). A version of Sho1
was also built where the entire SH3 domain was replaced by the same flexible stuffer used in the
Pbs2-stuffed strain. As with Pbs2, the stuffer was amplified using primers adding homology to the
sequences either side of the Sho1 SH3 domain locus (Primers 113F and 113R). The stuffer was
inserted into the Sho1 sequence by cotransformation of the pCas-Sho1 plasmid and the repair
template of the stuffer with homology arms. Proper stuffing was verified by PCR and Sanger
sequencing (Primers 114F and 114R).

<u>Construction of the surrounding region DMS library</u>

The initial DMS library covering the surrounding region of Pbs2 was constructed as previously
reported (Dionne *et al.* 2021). 56 codons (168 nucleotides) were targeted, which correspond to
codons 71 to 126 of the *YJL128C*/*PBS2* ORF.

A degenerate oligonucleotide was designed for each codon to be mutated, with the three
nucleotides of the mutated codon replaced by an NNN codon (Primers 115 to 170). This series of
56 oligonucleotides was used to amplify pUC19-Pbs2, in a two-step PCR procedure (Miyazaki
2011). At all steps, the different codons were kept in separate reactions and the mutants for each
codon were created in separate reactions. In the first amplification, a megaprimer was created

which includes the *PBS2* coding sequence using the degenerate oligonucleotide and a common
reverse primer (Primer 171R). The megaprimer thus carries the DMS library for one codon. In the
second PCR step, this megaprimer was used to amplify the entire plasmid. The PCR products
were digested with DpnI (New England Biolabs) at 37°C for 1 hour to remove the original
methylated template, and keep only the mutated plasmids amplified by the megaprimer. The
remaining mutated amplified but unligated plasmids were transformed into chemocompetent *E.*
coli cells strain MC1061, and plated on solid 2YT+Amp. The amplified plasmids were ligated and
thus recircularized in the cells. All transformations resulted in hundreds of colonies and these
colonies were suspended in 5 mL of 2YT. Their plasmids were extracted using a miniprep plasmid
extraction kit (Presto Mini Plasmid Kit, Geneaid) to obtain a plasmid library containing all the DMS
mutants. Prepared libraries were sequenced using 300-PE MiSeq technology (see DNA
sequencing section), to verify that the desired codon diversity was present. Once diversity was
confirmed, the libraries were amplified from the plasmids (Primers 172F and 172R), and
transformed into the Pbs2-stuffed-DHFR F[3] yeast strain using the CRISPR-Cas9 strategy
described in the strain construction section.
812
Transformed cells were plated on solid YPD+G418+Hyg for selection, and then resuspended in
liquid YPD. The optical density of each suspension (each codon) was measured and all
suspensions were pooled into a masterpool, with 5 optical density units (OD) added for each
codon. Frozen glycerol stocks were prepared for the individual suspensions of each codon, and
multiple stocks were prepared for the masterpool. Genomic DNA was extracted from the
masterpool using phenol/chloroform DNA extraction (Amberg *et al.* 2005), and as before, the
library was sequenced to verify diversity. At this point, three codons had unsatisfactory diversity.
A new template was created for codons 103 and 119 using fusion PCR, creating a sequence of
967 base pairs around the mutated codon, using the original degenerate primers as well as new
reverse degenerate primers to create an NNN codon at either codon 103 or 119 (Primers 173 to
175). For codon 88, a new template was ordered as an oligonucleotide of 503 base pairs (Twist
Bioscience), around codon 88, which is replaced by an NNN codon. These templates were
transformed into Pbs2-stuffed-DHFR F[3] as described above, and diversity at the desired codon
was verified with Sanger sequencing. The three supplementary cultures were pooled into the
masterpool, with an OD corresponding to 1/56th of the OD of the master pool.
828
829 <u>DHFR-PCA competition assay for surrounding region DMS library</u>
830
The PCA selection followed a previously published protocol (Dubé *et al.* 2022). Three liquid
cultures were started, each from 100 µL of the masterpool containing all mutated codon positions,
in SC complete medium. These cultures were incubated at 30 °C for 16 hours. 300 µL of this
preculture was mixed with 150 µL of either Sho1-DHFR F[1,2] or Hog1-DHFR F[1,2], and 1.2 mL
of fresh liquid YPD. These mixes of strains were left to mate for 8 hours at 30 °C with agitation.
The mated diploid cells were then selected in two successive cycles of diploid selection in liquid
YPD+Nat+Hyg at 30 °C with agitation. The first cycle consisted of a 5/12 dilution of the mating
culture in 3 mL of liquid YPD for 16 hours, and the second cycle consisted of a 1/10 dilution in 3
mL of liquid SC media for 24 hours. A volume equivalent to 5 OD was spun down and the
supernatant was removed, to form a cellular pellet, which was stored at -80 °C. This is the initial

time point used for sequencing. The selected diploid cultures were then diluted to 0.1 OD/mL in
15 mL of liquid PCA medium with either 200 µg/mL of methotrexate with 1 M of sorbitol, the same
concentration of methotrexate without sorbitol, or a control without methotrexate but with 1 M
sorbitol. This PCA selection was grown without light for 96 hours, except for the control cultures
which were saturated after 24 hours. After this first growth cycle, the cultures were diluted 0.1
OD/mL into 15 mL of fresh liquid PCA medium, and a 5 OD pellet was spun down and stored at -
80 °C. The second PCA cycle lasted the same amount of time, and at the end, as many 5 OD
pellets as possible were prepared from each culture. Genomic DNA was extracted from the pellets
of the second diploid selection cultures and the second PCA selection culture. The Pbs2 locus
was amplified and sequenced, as described in the DNA sequencing section.

Construction of the extended motif DMS library
A second DMS library was constructed of the extended motif of *PBS2*, comprising codons 85 to
100. An oligonucleotide pool (Integrated Data Technologies, Table S8) was synthesized
containing all possible NNK codons (K indicating either a G or T nucleotide) for the 16 codon
region of interest, with homology arms for a total length of 248 base pairs. This pool was integrated
into 2 different yeast strains: Sho1-DHFR F[1,2]/Pbs2-stuffed-DHFR F[3] and Hog1-DHFR
F[1,2]/Pbs2-stuffed-DHFR F[3], to allow respectively for PCA measurements of Sho1-Pbs2
interaction in a haploid strain, and PCA measurements of Hog1-Pbs2 interaction in a haploid
strain.

The mutant pool was integrated into these strains by replacing the Pbs2 stuffer using the CRISPR-
Cas9 strategy described in the strain construction section. As before, the integrated libraries were
sequenced to verify diversity. Transformed yeast libraries were stored at -80 °C.

DHFR-PCA competition assay for extended motif DMS library

The liquid PCA selection for the haploid double DHFR tagged strains was done following the
same protocol as the first DHFR-PCA selection, only ignoring the mating step. However, a fourth
control condition of PCA medium without methotrexate and without sorbitol was added. Three
replicate pools were done for each growth condition. Frozen pellets were prepared as in the first
DHFR-PCA selection.

Validation DHFR-PCA competition assay of individually reconstructed mutants

The individually reconstructed validation strains were pooled by combining 100 µL of saturated
overnight culture of each mutant in YPD. 6 pools were made for each condition, three pools each
from two individual reconstructed colonies of the mutant. The same conditions were used as the
second liquid PCA selection: PCA medium both with and without 1 M of sorbitol and with and
without methotrexate for the DHFR tagged strains. For the cell proliferation measurements, SC
medium with and without 1 M of sorbitol was used. Two cycles of selection were done, with 96-
hour cycles for the conditions with methotrexate and 24 hour cycles for the conditions without
methotrexate, and the SC medium. Frozen pellets were prepared as for the previous liquid PCA
selections.

885
886 <u>DNA sequencing</u>
887
DNA sequencing was done in generally the same manner for all experiments, with minor changes
in certain cases. DNA was extracted from the frozen pellets using a standard phenol/chloroform
genomic DNA extraction protocol (Amberg *et al.* 2005). The extracted DNA was amplified and
barcoded using a Row-Column DNA barcoding strategy as previously described (Dubé *et al.*
2022). From the extracted genomic DNA, the *PBS2* region of interest was amplified using primers
containing 3' overhangs which allow a second amplification to add barcodes (Primers 176F and
176R). PCR products were diluted 1/2500, and used as a template to add row and column
barcode primers onto the sequence, with a unique combination of 5' and 3' barcodes for each
replicate in each condition. The row-column barcode primers were previously described (Dubé *et*
al. 2022). PCR products were migrated on an agarose gel, and DNA concentrations were
estimated based on band size using the Image Lab software (BioRad Laboratories). Samples
were then pooled with different volumes to obtain the same quantity of DNA from each sample.
The pools were purified using magnetic beads, measured using a NanoDrop (ThermoFisher), and
diluted to 0.1 ng/µL. The diluted pools were then amplified with primers adding a second set of
barcodes, called plate barcodes (Dubé *et al.* 2022). These double-barcoded pools were purified
on magnetic beads, and sent for sequencing on Illumina NGS machines. Unique combinations of
barcodes were used to identify the DNA extracted from each pellet in each replicate of each
condition. The sequencing of the first DHFR-PCA, on the larger region surrounding the Pbs2 motif
was done using Illumina NovaSeq paired-end 250 base pair technology (CHUL sequencing
platform, Quebec, Canada), while all other sequencing was done using Illumina MiSeq paired-
end 300 bp technology (IBIS Genomic Analysis Platform, Quebec, Canada).
909
After preliminary analysis of the sequencing of the DNA from the pellets of the DHFR-PCA on the
extended motif DMS library, it was determined that more reads would be needed. Additional
sequencing libraries were prepared fresh from the original DNA extractions. Instead of using a
row-column barcoding approach, the *PBS2* region was amplified using primers (Primers 177F
and 177R) with homology arms allowing the addition of Illumina Nextera barcodes (Illumina).
These allow automated demultiplexing by the Illumina MiSeq instrument. All other steps were as
described above.
917
918
<u>Analysis of DNA sequencing results</u>
The variant frequency in DMS libraries during construction and in the DHFR-PCA and proliferation
screens was evaluated using custom scripts based on already published work (Després *et al.*
2022). Python libraries pandas 1.5.2 (The pandas development team 2023), matplotlib 3.6.2
(Hunter 2007) and numpy 1.24.1 (Harris *et al.* 2020) were used for data manipulation and
visualization. Quality was assessed using FastQC 0.12.1 (Andrews 2010). Reads were trimmed
using Trimmomatic 0.39 (Bolger *et al.* 2014), then demultiplexed using bowtie 1.3.1 for the
sequencing of the surrounding region DMS library (Langmead *et al.* 2009), Interstellar 1.0
following the RCP-PCR configuration for the first sequencing run of the extended motif DMS
library (Kijima *et al.* 2023) and cutadapt 4.7 for the validation competition assay sequencing

(Martin 2011). The second sequencing run for the extended motif DMS library was barcoded using Illumina Nextera primers, and was thus automatically demultiplexed by the sequencing instrument. Forward and reverse reads were then merged using the PANDAseq software (Masella *et al.* 2012). Next, identical reads were grouped using vsearch (Rognes *et al.* 2016), and aligned to the wild-type *PBS2* sequence using the needle function from the EMBOSS software (Rice *et al.* 2000). From this, the frequency of each variant was obtained.

From the frequency of each variant, the interaction scores and selection coefficients of each variant were calculated using custom R 4.3.1 (R Core Team 2023) scripts. The R package collection tidyverse (Wickham *et al.* 2019), as well as the packages rstatix 0.7.0 (Kassambara 2021), ggpubr 0.4.0 (Kassambara 2020) and bioseq 0.1.4 (Keck 2020) were used for data analysis. R packages ggExtra 0.10.1 (Attali and Baker 2023), ggrepel 0.9.4 (Slowikowski 2023), GGally 2.2.0 (Schloerke *et al.* 2023) and cowplot 1.1.1 (Wilke 2020) were used for data visualization. Briefly, variant counts were imported, identified by the mutation type (wild type, nonsense, missense or silent) then transformed into frequencies by dividing each variant count by the total number of counts present in that sequencing library. The interaction score was calculated as the log-2-fold-change of the frequency of each variant at the end of the second cycle of methotrexate selection and the frequency of each variant after the selection for diploids (so immediately preceding the methotrexate selection). These log-2-fold-change scores were rescaled for each library, so that the median of nonsense codons was equivalent to -1 and the value of wild-type variants was 0. Variants were then filtered to keep only those that had 20 or more reads detected after diploid selection, in order to remove bias caused by low read counts leading to inflated scores. Next, a unique score was calculated for each variant by taking the median score of all codons coding for the same amino acid variant in all replicates of the same condition. Scores were only kept for amino acid variants that had at least 3 replicates from any combination of codons not filtered out. For the competition assay of cell proliferation, the number of generations elapsed during the selection for each replicate was calculated, as the log-2-fold-change the optical density of the cultures at the end and at the beginning of the growth cycles. The selection coefficient for each variant was calculated as follows:

$$s = \frac{ln\left(\frac{variant\ frequency\ after\ PCA\ selection}{variant\ frequency\ before\ PCA\ selection}\right) - ln\left(\frac{wt\ frequency\ after\ PCA\ selection}{wt\ frequency\ before\ PCA\ selection}\right)}{number\ of\ generations}$$

The frequency of the wild type used is the frequency of the wild type in the same replicate as the variant. The same filtering criteria were used for the proliferation assay as for the DHFR-PCA.

When measuring the interaction scores of the preliminary and extended motif DMS libraries, we also included control conditions without methotrexate, the selective agent allowing measurement of DHFR fragment reconstitution. The measurements in this condition should not give any information about interactions between Pbs2 and Sho1, but can inform us about any other defects in the mutant strains. We calculated DMSO scores for growth in the control condition in the same way as the interaction scores. We noticed that some nonsense mutants had growth scores

comparable to wild-type Pbs2, and that some silent mutants had growth scores comparable to nonsense mutants. These abnormal scores could indicate defects in the strains, possibly resulting from the Cas9 mediated mutation process. We statistically verified which nonsense and silent mutants were abnormal by comparing every nonsense or silent mutation to all other nonsense or silent mutations (Two-sided Mann-Whitney U test, $p < 0.05$). The mutants which were significantly different were removed from the interaction score dataset. Similarly, we wanted to verify that no missense mutants had any defects caused by other sources than the interaction. We statistically verified which mutants had defects in the control condition by comparing each missense mutant to the nonsense mutants (Two-sided Mann-Whitney U test with false discovery rate corrected p-value > 0.05). The missense mutants which were not significantly different from the nonsense mutants were also removed from the interaction score dataset.

We also removed from the interaction score dataset the mutants for which an abundance change was detected through gain or loss of interaction strength between Pbs2 and Hog1, as explained in the results section. Pbs2 missense mutants that had an interaction strength with Hog1 which was significantly different from wild-type Pbs2 and silent mutations (Two-sided Mann-Whitney U test with false discovery rate corrected p-value < 0.05), either in the presence or absence of sorbitol, were removed from the datasets.

DHFR-PCA growth curves of individually reconstructed mutants

Growth curves were measured for individually reconstructed Pbs2 mutants in DHFR-PCA medium, to validate Pbs2 interaction strength measurements. The DHFR F[3] tagged strains with the mutations were mated with a Sho1-DHFR F[1,2]/*pbs2::LEU2* strain, in order to produce diploids with complementary DHFR fusions. To mate these strains, 50 µL of precultures of each strain were mixed into 900 µL of YPD and incubated at 30 °C overnight without agitation. 2 µL of each mated culture was spotted on solid YPD+Nat+Hyg, to select for diploids. Exponential phase precultures of diploids were diluted to 0.1 OD/mL in 80 µL of PCA medium, in a 384 well plate, in 4 replicates, at 30 °C. Cultures were grown either with and without 200 µg/mL of methotrexate, and with and without 1 M sorbitol. The optical density of each well was measured every 15 minutes in a Tecan Spark plate reader (Tecan). Growth curves were analyzed using a custom script written in R (R Core Team 2023), using the tidyverse package collection (Wickham *et al.* 2019) and the R package ggpubr 0.6.0 (Kassambara 2020). A function was written to find the maximal instantaneous growth rate for each well. Mutant P94R was disregarded, as growth of all replicates in the control condition in the absence of methotrexate was much lower than all other samples, suggesting a problem with strain construction.

Structure prediction

Structure predictions were done for Pbs2 mutants in complex with Sho1 using AlphaFold 2.3.2 and the AlphaFold-Multimer implementation (Jumper *et al.* 2021; Evans *et al.* 2022). Default options were used, and only the top scoring model for each mutant was relaxed. For Sho1, the

1015   sequence of the entire protein was used for prediction (Uniprot entry P40073). For Pbs2, residues
1016   71 to 126 were used for the prediction (Uniprot entry P08018). Predicted structures were
1017   visualized using ChimeraX 1.8 (Pettersen *et al.* 2021).
1018
<u>Sequence and structural alignments</u>
The protein sequences of 24 yeast SH3-containing sequences were obtained from the
alliancemine server (Bult and Sternberg 2023). The sequence of the SH3 domains plus 25
residues on each side was extracted from the complete protein sequences using the SH3 location
information from Interpro (Paysan-Lafosse *et al.* 2023), and a custom R script. When an SH3
domain was within less than 25 residues of the N- or C-terminus of a protein, the sequence up to
the terminus was used. The sequences of these SH3 domains and adjoining regions were aligned
using mafft 7.526, using the L-INS_i (local alignment) algorithm with the command line :
"/usr/bin/mafft" --localpair --maxiterate 16 --inputorder "<input_file>" > "<output_file>" (Katoh and
Standley 2013). For the structural alignment, we used previously predicted structures from the
AlphaFold Protein Structure Database (Varadi *et al.* 2022). Using the R packages
AlphaMissenseR (Morgan and Nguyen) and bio3d (Grant *et al.* 2006), we downloaded the
predicted structures of the 24 same yeast SH3-containing proteins from the AlphaFold Protein
Structure Database, and trimmed the pdb files to only contains residues composing the SH3
domains as well as 25 residues on each side of the domain. To align the predicted sequences,
we used the MUSTANG 3.2.4 tool, with the command line ./bin/mustang-3.2.3 -f ./path_to_files.txt
-F fasta -o SH3_flanking -s off (Konagurthu *et al.* 2006). The ./path_to_files.txt file contains the
paths of all trimmed pdb files.
1037
1038   <u>DHFR-PCA on solid media against SH3 domain-containing proteins</u>
1039
The DHFR-PCA on solid media was based on previous work (Tarassov *et al.* 2008; Rochette *et
al.* 2015). All colony manipulation was done using a robotic pin tool platform (BM5-SC1, S&P
Robotics Inc.). The strains containing DHFR F[1,2]-tagged SH3-containing proteins and Pbs2-
DHFR F[3] were taken from the Yeast Protein Interactome Collection (Horizon Discovery). The
Pbs2-stuffed-DHFR F[3] strain is the same used for DMS library construction, and is described in
the strain construction section. Cultures were started from frozen stocks in YPD with
nourseothricin for DHFR F[1,2] strains and hygromycin B for DHFR F[3] strains, in deep welled
96-well plates.
1048
Colonies were cherry picked from the 96-well plates, and arrayed onto YPD+Nat plates for the
DHFR F[1,2] strains or YPD+Hyg plates for the DHFR F[3] strains, in a 384 colony array. The
DHFR F[1,2] strains were organized into a randomized pattern to avoid any effects caused by
neighboring colonies. Each 384 colony plate of DHFR F[3] strains contained only copies of the
same variant. The 384 colony arrays were condensed into 1536 colony arrays on YPD+Nat or
YPD+Hyg, with the randomized DHFR F[1,2] pattern repeated four times. The outer two rows and
columns of the 1536 arrays were composed of LSM8-DHFR F[1,2] and CDC39-DHFR F[3], which
grow well on PCA medium and serve to avoid any measurement bias from being on the edge of
the array for the interactions of interest. The 1536 colony arrays were mated together by pinning
the repeated DHFR F[1,2] array onto YPD plates, and then pinning one of the DHFR F[3] arrays

onto each plate. In this collection of arrays, each interaction between an SH3 containing protein and a Pbs2 variant was measured in 5 or 6 replicates. After 48 hours of mating at 30 °C, the mated colonies were pinned onto YPD+Nat+Hyg plates to select for diploid cells. Two growth cycles of 48 hours at 30 °C on YPD+Nat+Hyg were done. After diploid selection, the plates were photographed using an EOS Rebel T5i camera (Canon), to verify that all colonies grew correctly. The mated arrays were then pinned onto solid PCA medium plates, either with or without methotrexate, and with 1 M of sorbitol. The arrays were grown for two cycles of 48 hours at 30 °C in a custom growth and imaging platform (S&P Robotics Inc.), which incubated the plates and took a picture of each plate every 2 hours.

Analysis of DHFR-PCA of Pbs2 mutants binding to SH3 domain-containing proteins

First, the ImageMagick command line tool (ImageMagick Studio LLC 2023) was used to crop the images of the selection plates as well as change them to grayscale and invert the colors, using the commands: convert -colorspace LinearGray, convert -crop 4200X2800+530+320 and convert -negate. Colony sizes were then quantified using the Python package Pyphe (Kamrad *et al.* 2020), using the command pyphe-quantify timecourse --grid auto_1536 --t 1 --d 3 --s 0.05. The colony sizes were analyzed and visualized using a custom script written in R 4.3.1 (R Core Team 2023), based on a previous analysis (Dionne *et al.* 2021), and using R packages from the tidyverse collection (Wickham *et al.* 2019). Colony sizes were filtered, and colonies which had not grown at the end of the diploid selection were removed from the PCA results. Colony sizes were log2 transformed and then normalized to avoid bias from a position in the array or from the plate they grew on by subtracting the log2-transformed median colony size for each row, column and 1536 array, including the border colonies. Scores were rescaled, so that the median of the scores of the Sho1 and wild-type Pbs2 interaction was equivalent to 1 and the median of scores of the Sho1 and stuffed Pbs2 interaction was equivalent to 0.

Analysis of contacts in SH3-motif structures

Contacts were analyzed for 4 structures. Sho1-Pbs2 was generated by AlphaFold-Multimer, as described above. Abp1-Ark (PDB accession 2RPN), Nbp2-Ste20 (PDB accession 2LCS), and Bem1-2-Ste20 (PDB accession 2RQW) were all previously measured NMR structure ensembles, composed of the 20 individual structures. All 20 structures were used for determining the contacts, although only the designated representative structure was used for visualization. The structures were imported into ChimeraX 1.8 (Pettersen *et al.* 2021), and the ChimeraX "Contacts" tool was used to detect all pairs of atoms from different proteins with a Van der Waals radius overlap ≥ -0.40 Å. This was done for all atoms in the motif-bearing protein preceding position -3 in the standard motif numbering. So, in Pbs2, this was done for all atoms in positions 71 to 92 (SNQSEQDKGSSQSPKHIQQIVN), for Ark1 this was done for positions -10 to -4 (KPKLHSP), and for Ste20 this was done for positions -14 to -4 (SSSANGKFIPS). To identify equivalent positions in the four SH3 domains which form contacts, the structures of the 4 domains were aligned using the MUSTANG tool (Konagurthu *et al.* 2006), and a standard numbering was determined based on this alignment (Table S9). The contacts between atoms were filtered to simply indicate all contacts between residues, and are summarized in Table S10, using the

1103 aforementioned standard numbering. The minimum distance between residues or proteins was
1104 calculated in a custom R script, using the bio3d package (Grant *et al.* 2006).

Data availability

Strains and plasmids are available upon request. All analysis scripts and visualization scripts are
available on Github at the following address: https://github.com/Landrylab/Jordan_et_al_2024. All
data, including demultiplexed sequencing data, results from the DHFR-PCA experiments, growth
curve data, predicted structures, protein alignments, and all files and information necessary to run
the scripts are available in the following Dryad repository:
https://doi.org/10.5061/dryad.79cnp5j3z.
1112
1113

Acknowledgments

1115 The authors thank Philippe C. Després, Moïra Dion and Dan Evans-Yamamoto for help with
1116 sequencing data analysis. We also thank all members of the LandryLab team for thoughtful
1117 comments on the experiments and on figure conception. All molecular graphics were performed
1118 using UCSF ChimeraX 1.8. The protein structure predictions were enabled by computing
1119 resources provided by the Digital Research Alliance of Canada.
1120

Funding

This work was supported by Canadian Institutes of Health Research Foundation grant number
387697 and Human Frontier Science Program research grant RGP34/2018 to CRL. C.R.L. holds
the Canada Research Chair in Cellular Systems and Synthetic Biology. D.J. was supported by
the Canada Graduate Scholarship - Master's program from NSERC, a graduate scholarship from
PROTEO, a graduate scholarship from the NSERC CREATE program EvoFunPath, a
Postgraduate Scholarship - Doctoral from NSERC and a Doctoral scholarship from FRQS. D.B.
was funded by the EMBO Long-Term Fellowship (LTF) (ALTF 1069-2019). U.D. was
funded by an NSERC graduate fellowship.
1130
1131

Author contributions

1133 All authors were involved in the design of the study. C.R.L. supervised this work and obtained
1134 funding. D.J. and A.K.D. performed the experiments. D.J. performed the computational analyses.

1135 D.J. wrote the first draft of the manuscript. All authors contributed to revising and writing the final
1136 version of the manuscript.

1137

1138

1139

Supplementary Figures

1140

1141

Figure S1.

Sho1-Pbs2 interaction scores for Pbs2 mutants in the DHFR-PCA on the DMS library of the region
surrounding the Pbs2 motif. Scores are indicated for each mutant according to their position in
Pbs2 and the residue that it was mutated to. The wild-type residue for each position is indicated
by an asterisk (*). Mutants which significantly affect the Hog1-Pbs2 interaction are marked with
the letter H (Two-sided Mann-Whitney U test with false discovery rate corrected p-value < 0.05).
Mutants which were removed from the dataset for having too few reads in the initial timepoint are
marked in black. Each measurement is the median of all codons in each replicate coding for the
same residue, with 3 to 18 replicates for each mutant.

1151

1152

Figure S2.

Correlation between replicates in the DHFR-PCA experiments on DMS libraries surrounding

region of the Pbs2 binding motif a) Scatterplot between the 3 replicates of the Sho1-Pbs2 DHFR-

PCA interaction scores for mutants in the DMS library of the region surrounding the Pbs2 motif.

Each point represents one Pbs2 codon variant in one condition (with or without sorbitol, with MTX

or with only DMSO), as indicated by the color in the corresponding panel on the top right. Pearson

correlations also indicated in the top right panels. In the panels on the diagonal, the distribution

of scores for a given condition and replicate is shown. The replicate number is indicated in the
gray rectangle. b) Scatterplot between the 3 replicates of the Hog1-Pbs2 DHFR-PCA interaction
scores for mutants in the DMS library of the region surrounding the Pbs2 motif. Each point
represents one Pbs2 codon variant in one condition (with or without sorbitol, with MTX or with
only DMSO), as indicated by the color in the corresponding panel on the top right. Pearson
correlations also indicated in the top right panels. In the panels on the diagonal, the distribution
of scores for a given condition and replicate is shown. The replicate number is indicated in the
gray rectangle.
1168

1169
1170
1171
Figure S3.
Scatterplot of interaction scores for missense and silent mutants in the DMS library of the region
surrounding the motif, for the interaction of Pbs2 with either Sho1 or Hog1. Points are colored
according to whether the mutants are situated inside or outside the extended motif. The dotted
line represents the diagonal. The gray rectangles on the left and bottom of the panel represent
the 97.5 percentile of scores for the nonsense variants of Pbs2, which do not express the DHFR
F[3] fragment. As such, they represent the limit of detectable signal in the assay. The points which
have scores comparable to the nonsense mutations, and therefore no detectable interaction, were
placed at the limit of the gray rectangle to indicate that no interaction was detected. In the margins,
density plots show the distribution of scores for mutants situated either inside or outside the
extended motif. Each point represents the median of all codons in each replicate coding for the
same mutation, with 3 to 18 replicates for each mutant.
1184

1185

Figure S4.

<u>Correlation between replicates in the DHFR-PCA experiments on DMS libraries of the extended</u>

<u>Pbs2 binding motif</u> a) Scatterplot between the 6 replicates of the Sho1-Pbs2 DHFR-PCA

interaction scores for mutants in the DMS library of the extended Pbs2 motif. Each point

represents one Pbs2 codon variant in one condition (with or without sorbitol, with MTX or with

only DMSO), as indicated by the color in the corresponding panel on the top right. Pearson

correlations also indicated in the top right panels. In the panels on the diagonal, the distribution

of scores for a given condition and replicate is shown. The replicate number is indicated in the

gray rectangle. b) Scatterplot between the 6 replicates of the Hog1-Pbs2 DHFR-PCA interaction
scores for mutants in the DMS library of the extended Pbs2 motif. Each point represents one Pbs2
codon variant in one condition (with or without sorbitol, with MTX or with only DMSO), as indicated
by the color in the corresponding panel on the top right. Pearson correlations also indicated in the
top right panels. In the panels on the diagonal, the distribution of scores for a given condition and
replicate is shown. The replicate number is indicated in the gray rectangle.
1200
1201

1202
Figure S5.
Scatterplot of interaction scores of Pbs2 mutants in the initial DHFR-PCA screen on the DMS
library of the surrounding region of the motif, and the subsequent DHFR-PCA screen on the DMS
library of only the extended motif (positions 85 to 100). Scores shown were measured in the
presence of methotrexate and 1 M of sorbitol. Each point represents the median of all codons in
each replicate coding for the same mutation, with 3 to 18 replicates for each mutant. Spearman's
rho 0.91, p < $2.2X10^{-16}$.
1210
1211
1212
1213

1214
1215
Figure S6.
Scatterplot of interaction scores of the Pbs2 extended motif DMS library mutants with Sho1 and
Hog1, in the presence of 1 M of sorbitol. Purple missense points represent Pbs2 mutants which
have an interaction score with Hog1 which is significantly different to the combination of silent
mutants and wild-type Pbs2 (Two-sided Mann-Whitney U test with false discovery rate corrected
p-value < 0.05), while blue missense points represent mutants for which the Hog1 interaction
score is not significantly different to the combination of silent mutants and wild-type Pbs2. The
combination of silent mutations and wild type represents all mutants expressing the wild-type
protein sequence. Since the Hog1 interaction is used to measure abundance and stability of the
Pbs2 mutants, the significantly different mutants are considered to have an effect on the
abundance or the stability of Pbs2 and were not kept for the remainder of the analysis. The only
exceptions are the nonsense mutants, which were kept because they provide a useful reference.
Each point represents the median of all codons in each replicate coding for the same mutation,
with 3 to 18 replicates for each mutant.

1235
Figure S7.
Sho1-Pbs2 interaction scores for Pbs2 mutants in the DHFR-PCA on the DMS library of the
extended Pbs2 motif. Scores are indicated for each mutant according to their position in Pbs2
and the residue that it was mutated to. The wild-type residue for each position is indicated by an
asterisk (*). Mutants which significantly affect the Hog1-Pbs2 interaction are marked with the letter
H (Two-sided Mann-Whitney U test with false discovery rate corrected p-value < 0.05). Mutants
which were removed from the dataset for having too few reads in the initial timepoint are marked
in black. Each measurement represents the median of all codons in each replicate coding for the
same residue, with 3 to 18 replicates for each mutant.

1257
1258
Figure S8.
Scatterplot of the Sho1 interaction scores for all mutants in the Pbs2 extended motif DMS library,
in the presence and absence of 1 M of sorbitol. Measurements in DHFR-PCA media with
methotrexate. The correlation has a spearman's rho of 0.98, and a p-value $< 2.2 \times 10^{-16}$. The
measurements for the interaction in 1 M sorbitol have a greater range due to the greater activation
of the HOG pathway. Each point represents the median of all codons in each replicate coding for
the same mutation, with 3 to 18 replicates for each mutant.
1266
1267
1268

1269

1270
1271
Figure S9.
<u>Validation of pooled DHFR-PCA effects by individually reconstructed mutants</u>
a) Scatterplot of Sho1-Pbs2 interaction scores as measured in pooled competition DHFR-PCA of
the DMS libraries (x-axis) and growth rate of the same mutations individually reconstructed, in
DHFR-PCA growth curves (y-axis). Results from the DMS library of the extended motif (left) and
the DMS library of the surrounding region (right), in the presence (orange) or absence (purple) of
1 M of sorbitol. The spearman correlations between the two methods of measuring the interaction
strength are indicated on the plots. The values for the DMS library interaction scores represent
the median of all codons in each replicate coding for the same mutation, with 3 to 18 replicates
for each mutant. The values for the growth rate represent the median of 3 replicates. b) Scatterplot
of Sho1-Pbs2 interaction scores as measured in pooled competition DHFR-PCA of the DMS
libraries (x-axis) and measured in pooled competition DHFR-PCA of individually reconstructed
mutants (y-axis). Results from the DMS library of the extended motif (left) and the DMS library of
the surrounding region (right), in the presence (orange) or absence (purple) of 1 M of sorbitol.
The spearman correlations between the two methods of measuring the interaction strength are
indicated on the plots. The values for the DMS library interaction scores represent the median of
all codons in each replicate coding for the same mutation, with 3 to 18 replicates for each mutant.
The values for the reconstructed mutants represent the median of 3 to 6 replicates.
1290

1291

1292
Figure S10.
Correlation between replicates in the DHFR-PCA experiments on DMS libraries of the extended
Pbs2 binding motif
a) Scatterplot between the 6 replicates of the Sho1-Pbs2 DHFR-PCA interaction scores for
mutants in the Pbs2 validation DMS library. Each point represents one Pbs2 codon variant in one
condition (with or without sorbitol, with MTX or with only DMSO), as indicated by the color in the
corresponding panel on the top right. Pearson correlations also indicated in the top right panels.

In the panels on the diagonal, the distribution of scores for a given condition and replicate is
shown. The replicate number is indicated in the gray rectangle. b) Scatterplot between the 6
replicates of the Hog1-Pbs2 DHFR-PCA interaction scores for mutants in the Pbs2 validation
DMS library. Each point represents one Pbs2 codon variant in one condition (with or without
sorbitol, with MTX or with only DMSO), as indicated by the color in the corresponding panel on
the top right. Pearson correlations also indicated in the top right panels. In the panels on the
diagonal, the distribution of scores for a given condition and replicate is shown. The replicate
number is indicated in the gray rectangle. c) Scatterplot between the 6 replicates of selection
coefficients for mutants in the Pbs2 validation DMS library measuring cell proliferation. Each point
represents one Pbs2 codon variant in one condition (with or without sorbitol), as indicated by the
color in the corresponding panel on the top right. Pearson correlations also indicated in the top
right panels. In the panels on the diagonal, the distribution of scores for a given condition and
replicate is shown. The replicate number is indicated in the gray rectangle.
1313
1314

1315

Figure S11.
Impact on growth rate measured as selection coefficients for individually reconstructed Pbs2
mutants in SC synthetic media with 1 M sorbitol. The strains contain no DHFR fragments on any
proteins. Boxplots show the distribution of 6 replicates per mutant. The selection coefficient is
normalized by the growth of the wild-type sequence, making the wild-type selection coefficient
zero. Only the 3 nonsense mutants grew significantly less than wild-type Pbs2 and the silent
mutants (Welch's t-test with false discovery rate correction p-value < 0.05), although I87W
(highlighted in red on axis), which is the mutant with the highest Sho1-Pbs2 interaction score
shows a marked decrease in selection coefficient (p-value = 0.066).
1325

1326
Figure S12.
<u>Confidence metrics for the seven structures predicted using AlphaFold2-Multimer.</u> Sho1 SH3-
Pbs2 WT is the structure in figure 2b, Sho1-Pbs2 WT is the structure in figure 3a and 3b, Sho1-
Pbs2 I87W is the structure in figure 3c, and all others are the structures with the same Pbs2
mutants in figure S13. a) Predicted aligned error (PAE) values for the 7 predicted structures,
output from AlphaFold-Multimer. b) pLDDT value for each position in the 7 predicted structures,
output from AlphaFold-Multimer. Positions corresponding to the SH3 domain of Sho1 are
indicated by a green background, and positions corresponding to the extended motif of Pbs2 are
indicated by a blue background. Since AlphaFold-Multimer combines the sequences of both
proteins into a single model, a red dashed line was used to mark the end of the Sho1 portion of
the model, and the beginning of the Pbs2 portion of the model.

1338
Figure S13.
<u>Close-up details of AlphaFold-Multimer predicted complexes of Sho1 with Pbs2 mutants</u> a) H86W,
b) V91L, c) V91M and d) Q89D. The predicted structures are colored as in panel a of figure 3,
with the SH3 domain of Sho1 in green, the non-SH3 portions of Sho1 in orange, the extended
motif of Pbs2 in light blue and the rest of Pbs2 in burgundy. The side chain of the mutated residue
is shown with oxygen atoms in red, nitrogen atoms in dark blue, sulfur atoms in yellow, hydrogen
atoms in white and carbon atoms the same color as the extended motif, while all other side chains
in Pbs2 are hidden. The orientation of Pbs2 is shown by annotations indicating which end is nearer
the N-terminus (N) and which end is nearer the C-terminus (C).

Figure S14.

<u>Sequence and structural alignments of yeast SH3 domains and adjacent sequences</u> a) Multiple
sequence alignment of yeast SH3 domains with 25 residues on each side. Residues which in the
predicted Sho1-Pbs2 structure are within 5 Å of Pbs2 residues situated in the extended motif but
not the canonical motif are surrounded by a black rectangle. Beginning and end of the SH3
domain of Sho1 marked above the alignment. b) Multiple structural alignment of predicted yeast
SH3 domains with 25 residues on each side, from MUSTANG. Residues which in the predicted
Sho1-Pbs2 structure are within 5 Å of Pbs2 residues situated in the extended motif but not the
canonical motif are surrounded by a black rectangle. Beginning and end of the SH3 domain of
Sho1 marked above the alignment.

1359

1360
1361
Figure S15.

Interaction score of wild-type Pbs2 (green) or Pbs2 where positions 76 to 126 have been replaced
with a flexible stuffer sequence (Stuffed, white) with all yeast SH3-containing proteins. Controls
grown on DMSO and not MTX, and therefore not measuring the interaction, are shown in the top
panel. All replicates shown as points. Scores normalized with 1 being the median of Sho1 - wild
type Pbs2 interactions and 0 being the median of Sho1 - stuffed Pbs2 interactions. A control
where the SH3 domain of Sho1 was replaced by the same neutral, flexible stuffer sequence used
for Pbs2-stuffed was also included, with the name Sho1_stuffed.

1370

1371
Figure S16.
Predicted structure of the Sho1-Pbs2 interaction superimposed with other SH3-motif pairs. The
Sho1-Pbs2 structure is semi-transparent and colored as in figure 3a, with the SH3 domain of
Sho1 in green, the non-SH3 portions of Sho1 in orange, the Pbs2 extended motif in light blue and
the rest of Pbs2 in burgundy. Sho1-Pbs2 structure was superimposed with the a) Abp1-Ark
structure (Abp1 - light green, Ark1 - pink), b) Nbp2-Ste20 structure (Nbp2 - orange-pink, Ste20 -
light pink), and c) Bem1-2-Ste20 structure (Bem1 - yellow, Ste20 - light pink). Structures were
superimposed using the Chimera X tool "Matchmaker" (Pettersen *et al.* 2021) to minimize root
mean square deviance of backbone atom distances of the SH3 domains with Sho1.
1381

1382

Figure S17.
Additional differing contacts between SH3-extended motif pairs
a) Visualization of the side-chains and contacts of position (-7) in the Sho1-Pbs2 predicted
structure , Abp1-Ark1 structure (PDB accession 2RPN), Nbp2-Ste20 structure (PDB accession
2LCS) and Bem1-Ste20 (PDB accession 2RQW). Sho1-Pbs2 predicted structure colored as in
figure 3a, with the SH3 domain of Sho1 in green, the non-SH3 portions of Sho1 in orange, the
Pbs2 extended motif in light blue and the rest of Pbs2 in burgundy. All four structures were
superimposed using the Chimera X tool "Matchmaker" (Pettersen *et al.* 2021) to minimize root
mean square deviance of backbone atom distances of the SH3 domains with Sho1, then pictures
were captured separately. b) Visualization of the side-chains and contacts of position (-4) of the
same structures as in panel a). All four structures were superimposed using the Chimera X tool
"Matchmaker" (Pettersen *et al.* 2021) to minimize root mean square deviance of backbone atom
distances of the SH3 domains with Sho1, then pictures were captured separately.
1396
1397

Literature cited
1398

1399

Amberg D., D. C. Burke, and J. N. Strathern, 2005 *Methods in yeast genetics: a Cold Spring*

Harbor Laboratory course manual. Cold Spring Harbor Laboratory Press, Cold Spring

Harbor, N.Y.

Andrews S., 2010 FastQC: a quality control tool for high throughput sequence data

Attali D., and C. Baker, 2023 *ggExtra: Add Marginal Histograms to "ggplot2", and More "ggplot2"*

Enhancements.

Bendel A. M., A. J. Faure, D. Klein, K. Shimada, R. Lyautey, *et al.*, 2024 The genetic

architecture of protein interaction affinity and specificity. Nat. Commun. 15: 8868.

Bolger A. M., M. Lohse, and B. Usadel, 2014 Trimmomatic: a flexible trimmer for Illumina

sequence data. Bioinformatics 30: 2114–2120.

Bult C. J., and P. W. Sternberg, 2023 The alliance of genome resources: transforming

comparative genomics. Mamm. Genome 34: 531–544.

Dalgarno D. C., M. C. Botfield, and R. J. Rickles, 1997 SH3 domains and drug design: Ligands,

structure, and biological function. Pept. Sci. 43: 383–400.

Davey N. E., G. Travé, and T. J. Gibson, 2011 How viruses hijack cell regulation. Trends

Biochem. Sci. 36: 159–169.

Delplace V., A. Ortin-Martinez, E. L. S. Tsai, A. N. Amin, V. Wallace, *et al.*, 2019 Controlled

release strategy designed for intravitreal protein delivery to the retina. J. Control. Release

293: 10–20.

Després P. C., A. F. Cisneros, E. M. M. Alexander, R. Sonigara, C. Gagné-Thivierge, *et al.*,

2022 Asymmetrical dose responses shape the evolutionary trade-off between antifungal

resistance and nutrient use. Nat Ecol Evol 6: 1501–1515.

Dibyachintan S., A. K. Dube, D. Bradley, P. Lemieux, U. Dionne, *et al.*, 2024 Cryptic genetic

variation shapes the fate of gene duplicates in a protein interaction network. bioRxiv

2024.02.23.581840.

Dionne U., É. Bourgault, A. K. Dubé, D. Bradley, F. J. M. Chartier, *et al.*, 2021 Protein context

shapes the specificity of SH3 domain-mediated interactions in vivo. Nat. Commun. 12:

1597.

Dionne U., L. J. Percival, F. J. M. Chartier, C. R. Landry, and N. Bisson, 2022 SRC homology 3

domains: multifaceted binding modules. Trends Biochem. Sci. 47: 772–784.

Diss G., and B. Lehner, 2018 The genetic landscape of a physical interaction. Elife 7.

https://doi.org/10.7554/eLife.32472

Douangamath A., F. V. Filipp, A. T. J. Klein, P. Barnett, P. Zou, *et al.*, 2002 Topography for

Independent Binding of α-Helical and PPII-Helical Ligands to a Peroxisomal SH3 Domain.

1434   Mol. Cell 10: 1007–1017.

1435 Dubé A. K., R. Dandage, S. Dibyachintan, U. Dionne, P. C. Després, *et al.*, 2022 Deep

1436   Mutational Scanning of Protein–Protein Interactions Between Partners Expressed from

1437   Their Endogenous Loci In Vivo, pp. 237–259 in *Yeast Functional Genomics: Methods and*

1438   *Protocols*, edited by Devaux F. Springer US, New York, NY.

1439 Ernst A., D. Gfeller, Z. Kan, S. Seshagiri, P. M. Kim, *et al.*, 2010 Coevolution of PDZ domain-

1440   ligand interactions analyzed by high-throughput phage display and deep sequencing. Mol.

1441   Biosyst. 6: 1782–1790.

1442 Evans R., M. O'Neill, A. Pritzel, N. Antropova, A. Senior, *et al.*, 2022 Protein complex prediction

1443   with AlphaFold-Multimer. bioRxiv 2021.10.04.463034.

1444 Faure A. J., J. Domingo, J. M. Schmiedel, C. Hidalgo-Carcedo, G. Diss, *et al.*, 2022 Mapping

1445   the energetic and allosteric landscapes of protein binding domains. Nature 604: 175–183.

1446 Feng S., C. Kasahara, R. J. Rickles, and S. L. Schreiber, 1995 Specific interactions outside the

1447   proline-rich core of two classes of Src homology 3 ligands. Proc. Natl. Acad. Sci. U. S. A.

1448   92: 12408–12415.

1449 Ferrigno P., F. Posas, D. Koepp, H. Saito, and P. A. Silver, 1998 Regulated nucleo/cytoplasmic

1450   exchange of HOG1 MAPK requires the importin beta homologs NMD5 and XPO1. EMBO J.

1451   17: 5606–5614.

1452 Fowler D. M., and S. Fields, 2014 Deep mutational scanning: a new style of protein science.

1453   Nat. Methods 11: 801–807.

1454 Freschi L., F. Torres-Quiroz, A. K. Dubé, and C. R. Landry, 2013 qPCA: a scalable assay to

1455   measure the perturbation of protein-protein interactions in living cells. Mol. Biosyst. 9: 36–

1456 43.

1457 Gaussmann S., R. Peschel, J. Ott, K. M. Zak, J. Sastre, *et al.*, 2024 Modulation of peroxisomal
1458  import by the PEX13 SH3 domain and a proximal FxxxF binding motif. Nat. Commun. 15:
1459  3317.

1460 Gorelik M., and A. R. Davidson, 2012 Distinct peptide binding specificities of Src homology 3
1461  (SH3) protein domains can be determined by modulation of local energetics across the
1462  binding interface. J. Biol. Chem. 287: 9168–9177.

1463 Gouw M., S. Michael, H. Sámano-Sánchez, M. Kumar, A. Zeke, *et al.*, 2018 The eukaryotic
1464  linear motif resource – 2018 update. Nucleic Acids Res. 46: D428–D434.

1465 Grant B. J., A. P. C. Rodrigues, K. M. ElSawy, J. A. McCammon, and L. S. D. Caves, 2006
1466  Bio3d: an R package for the comparative analysis of protein structures. Bioinformatics 22:
1467  2695–2696.

1468 Gueldener U., J. Heinisch, G. J. Koehler, D. Voss, and J. H. Hegemann, 2002 A second set of
1469  loxP marker cassettes for Cre-mediated multiple gene knockouts in budding yeast. Nucleic
1470  Acids Res. 30: e23.

1471 Harris C. R., K. J. Millman, S. J. van der Walt, R. Gommers, P. Virtanen, *et al.*, 2020 Array
1472  programming with NumPy. Nature 585: 357–362.

1473 Haslam N. J., and D. C. Shields, 2012 Peptide-Binding Domains: Are Limp Handshakes Safest?
1474  Sci. Signal. 5: e40–pe40.

1475 Hunter J. D., 2007 Matplotlib: A 2D Graphics Environment. Comput. Sci. Eng. 9: 90–95.

1476 Illumina, Nextera DNA Indexes. Illumina Support Docs.

ImageMagick Studio LLC, 2023 *ImageMagick*.

Ivarsson Y., and P. Jemth, 2019 Affinity and specificity of motif-based protein-protein

interactions. Curr. Opin. Struct. Biol. 54: 26–33.

Jumper J., R. Evans, A. Pritzel, T. Green, M. Figurnov, *et al.*, 2021 Highly accurate protein

structure prediction with AlphaFold. Nature 596: 583–589.

Kamrad S., M. Rodríguez-López, C. Cotobal, C. Correia-Melo, M. Ralser, *et al.*, 2020 Pyphe, a

python toolbox for assessing microbial growth and cell viability in high-throughput colony

screens. Elife 9. https://doi.org/10.7554/eLife.55160

Kaneko T., L. Li, and S. S.-C. Li, 2008 The SH3 domain--a family of versatile peptide- and

protein-recognition module. Front. Biosci. 13: 4938–4952.

Kaneko T., H. Huang, X. Cao, X. Li, C. Li, *et al.*, 2012 Superbinder SH2 Domains Act as

Antagonists of Cell Signaling. Sci. Signal. 5: ra68–ra68.

Karlsson O. A., G. N. Sundell, E. Andersson, Y. Ivarsson, and P. Jemth, 2016 Improved affinity

at the cost of decreased specificity: a recurring theme in PDZ-peptide interactions. Sci.

Rep. 6: 34269.

Kassambara A., 2020 ggpubr: "ggplot2" Based Publication Ready Plots

Kassambara A., 2021 rstatix: Pipe-Friendly Framework for Basic Statistical Tests

Katoh K., and D. M. Standley, 2013 MAFFT multiple sequence alignment software version 7:

improvements in performance and usability. Mol. Biol. Evol. 30: 772–780.

Kazlauskas A., C. Schmotz, T. Kesti, J. Hepojoki, I. Kleino, *et al.*, 2016 Large-Scale Screening

of Preferred Interactions of Human Src Homology-3 (SH3) Domains Using Native Target

Proteins as Affinity Ligands. Mol. Cell. Proteomics 15: 3270–3281.

Keck F., 2020 Handling biological sequences in R with the bioseq package. Methods Ecol. Evol.

11: 1728–1732.

Kelil A., E. D. Levy, and S. W. Michnick, 2016 Evolution of domain–peptide interactions to

coadapt specificity and affinity to functional diversity. Proc. Natl. Acad. Sci. U. S. A. 113:

E3862–E3871.

Kelil A., B. Dubreuil, E. D. Levy, and S. W. Michnick, 2017 Exhaustive search of linear

information encoding protein-peptide recognition. PLoS Comput. Biol. 13: e1005499.

Kijima Y., D. Evans-Yamamoto, H. Toyoshima, and N. Yachie, 2023 A universal sequencing

read interpreter. Sci Adv 9: eadd2793.

Konagurthu A. S., J. C. Whisstock, P. J. Stuckey, and A. M. Lesk, 2006 MUSTANG: a multiple

structural alignment algorithm. Proteins 64: 559–574.

Kumar M., S. Michael, J. Alvarado-Valverde, A. Zeke, T. Lazar, *et al.*, 2024 ELM-the Eukaryotic

Linear Motif resource-2024 update. Nucleic Acids Res. 52: D442–D455.

Kursula P., I. Kursula, Y. H. Song, K. Paraskevopoulos, and M. Wilmanns, 2008 YEAST SHO1

SH3 DOMAIN COMPLEXED WITH A PEPTIDE FROM PBS2

Langmead B., C. Trapnell, M. Pop, and S. L. Salzberg, 2009 Ultrafast and memory-efficient

alignment of short DNA sequences to the human genome. Genome Biol. 10: R25.

Lee C. H., K. Saksela, U. A. Mirza, B. T. Chait, and J. Kuriyan, 1996 Crystal structure of the

conserved core of HIV-1 Nef complexed with a Src family SH3 domain. Cell 85: 931–942.

Lemieux P., D. Bradley, A. K. Dubé, U. Dionne, and C. R. Landry, 2024 Dissection of the role of

a Src homology 3 domain in the evolution of binding preference of paralogous proteins.

Genetics 226. https://doi.org/10.1093/genetics/iyad175

Li S. S.-C., 2005 Specificity and versatility of SH3 and other proline-recognition domains:

structural basis and implications for cellular signal transduction. Biochem. J 390: 641–653.

Liao T.-J., H. Jang, R. Nussinov, and D. Fushman, 2020 High-affinity interactions of the

nSH3/cSH3 domains of Grb2 with the C-terminal proline-rich domain of SOS1. J. Am.

Chem. Soc. 142: 3401–3411.

Lim W. A., F. M. Richards, and R. O. Fox, 1994 Structural determinants of peptide-binding

orientation and of sequence specificity in SH3 domains. Nature 372: 375–379.

Maeda T., M. Takekawa, and H. Saito, 1995 Activation of Yeast PBS2 MAPKK by MAPKKKs or

by Binding of an SH3-Containing Osmosensor. Science 269: 554–558.

Mapes J., and I. M. Ota, 2004 Nbp2 targets the Ptc1-type 2C Ser/Thr phosphatase to the HOG

MAPK pathway. EMBO J. 23: 302–311.

Marles J. A., S. Dahesh, J. Haynes, B. J. Andrews, and A. R. Davidson, 2004 Protein-Protein

Interaction Affinity Plays a Crucial Role in Controlling the Sho1p-Mediated Signal

Transduction Pathway in Yeast. Mol. Cell 14: 813–823.

Martin M., 2011 Cutadapt removes adapter sequences from high-throughput sequencing reads.

EMBnet.journal 17: 10–12.

Masella A. P., A. K. Bartram, J. M. Truszkowski, D. G. Brown, and J. D. Neufeld, 2012

PANDAseq: paired-end assembler for illumina sequences. BMC Bioinformatics 13: 31.

Messing J., 1983 [2] New M13 vectors for cloning, pp. 20–78 in *Methods in Enzymology*,

Academic Press.

Michnick S. W., E. D. Levy, C. R. Landry, J. Kowarzyk, and V. Messier, 2016 The Dihydrofolate

Reductase Protein-Fragment Complementation Assay: A Survival-Selection Assay for

Large-Scale Analysis of Protein–Protein Interactions. Cold Spring Harb. Protoc. 2016:

db.prot090027.

Miyazaki K., 2011 Chapter seventeen - MEGAWHOP Cloning: A Method of Creating Random

Mutagenesis Libraries via Megaprimer PCR of Whole Plasmids, pp. 399–406 in *Methods in*

Enzymology, Synthetic Biology, Part B. edited by Voigt C. Academic Press.

Monera O. D., T. J. Sereda, N. E. Zhou, C. M. Kay, and R. S. Hodges, 1995 Relationship of

sidechain hydrophobicity and alpha-helical propensity on the stability of the single-stranded

amphipathic alpha-helix. J. Pept. Sci. 1: 319–329.

Morgan M., and T. Nguyen, *AlphaMissenseR*.

Murakami Y., K. Tatebayashi, and H. Saito, 2008 Two adjacent docking sites in the yeast Hog1

mitogen-activated protein (MAP) kinase differentially interact with the Pbs2 MAP kinase

kinase and the Ptp2 protein tyrosine phosphatase. Mol. Cell. Biol. 28: 2481–2494.

Pawson T., M. Raina, and P. Nash, 2002 Interaction domains: from simple binding events to

complex cellular behavior. FEBS Lett. 513: 2–10.

Paysan-Lafosse T., M. Blum, S. Chuguransky, T. Grego, B. L. Pinto, *et al.*, 2023 InterPro in

2022. Nucleic Acids Res. 51: D418–D427.

Pettersen E. F., T. D. Goddard, C. C. Huang, E. C. Meng, G. S. Couch, *et al.*, 2021 UCSF

ChimeraX: Structure visualization for researchers, educators, and developers. Protein Sci.

30: 70–82.

Posas F., and H. Saito, 1997 Osmotic Activation of the HOG MAPK Pathway via Ste11p

MAPKKK: Scaffold Role of Pbs2p MAPKK. Science 276: 1702–1705.

R Core Team, 2023 *R: A Language and Environment for Statistical Computing.* R Foundation
for Statistical Computing, Vienna, Austria.

Rice P., I. Longden, and A. Bleasby, 2000 EMBOSS: the European Molecular Biology Open
Software Suite. Trends Genet. 16: 276–277.

Rickles R. J., M. C. Botfield, X. M. Zhou, P. A. Henry, J. S. Brugge, *et al.*, 1995 Phage display
selection of ligand residues important for Src homology 3 domain binding specificity. Proc.
Natl. Acad. Sci. U. S. A. 92: 10909–10913.

Robles J. T., H. J. Lou, G. Shi, P. L. Pan, and B. E. Turk, 2023 Linear motif specificity in
signaling through p38α and ERK2 mitogen–activated protein kinases. Proceedings of the
National Academy of Sciences 120: e2316599120.

Rochette S., G. Diss, M. Filteau, J.-B. Leducq, A. K. Dubé, *et al.*, 2015 Genome-wide protein-
protein interaction screening by protein-fragment complementation assay (PCA) in living
cells. J. Vis. Exp. https://doi.org/10.3791/52255

Rognes T., T. Flouri, B. Nichols, C. Quince, and F. Mahé, 2016 VSEARCH: a versatile open
source tool for metagenomics. PeerJ 4: e2584.

Ryan O. W., S. Poddar, and J. H. D. Cate, 2016 CRISPR-Cas9 Genome Engineering in
Saccharomyces cerevisiae Cells. Cold Spring Harb. Protoc. 2016.
https://doi.org/10.1101/pdb.prot086827

Saito H., and F. Posas, 2012 Response to Hyperosmotic Stress. Genetics 192: 289–318.

Schloerke B., D. Cook, J. Larmarange, F. Briatte, M. Marbach, *et al.*, 2023 *GGally: Extension to*
"ggplot2."

Slowikowski K., 2023 *ggrepel: Automatically Position Non-Overlapping Text Labels with
"ggplot2."*

Stein A., and P. Aloy, 2008 Contextual specificity in peptide-mediated protein interactions. PLoS
One 3: e2524.

Stollar E. J., B. Garcia, P. A. Chong, A. Rath, H. Lin, *et al.*, 2009 Structural, functional, and
bioinformatic studies demonstrate the crucial role of an extended peptide binding site for
the SH3 domain of yeast Abp1p. J. Biol. Chem. 284: 26918–26927.

Takaku T., K. Ogura, H. Kumeta, N. Yoshida, and F. Inagaki, 2010 Solution structure of a novel
Cdc42 binding module of Bem1 and its interaction with Ste20 and Cdc42. J. Biol. Chem.
285: 19346–19353.

Tarassov K., V. Messier, C. R. Landry, S. Radinovic, M. M. S. Molina, *et al.*, 2008 An in Vivo
Map of the Yeast Protein Interactome. Science 320: 1465–1470.

Tatebayashi K., K. Yamamoto, K. Tanaka, T. Tomida, T. Maruoka, *et al.*, 2006 Adaptor
functions of Cdc42, Ste50, and Sho1 in the yeast osmoregulatory HOG MAPK pathway.
EMBO J. 25: 3033–3044.

The pandas development team, 2023 *pandas-dev/pandas: Pandas.*

Tonikian R., X. Xin, C. P. Toret, D. Gfeller, C. Landgraf, *et al.*, 2009 Bayesian Modeling of the
Yeast SH3 Domain Interactome Predicts Spatiotemporal Dynamics of Endocytosis
Proteins. PLoS Biol. 7: e1000218.

Van Roey K., B. Uyar, R. J. Weatheritt, H. Dinkel, M. Seiler, *et al.*, 2014 Short Linear Motifs:
Ubiquitous and Functionally Diverse Protein Interaction Modules Directing Cell Regulation.
Chem. Rev. 114: 6733–6778.

Varadi M., S. Anyango, M. Deshpande, S. Nair, C. Natassia, *et al.*, 2022 AlphaFold Protein

Structure Database: massively expanding the structural coverage of protein-sequence

space with high-accuracy models. Nucleic Acids Res. 50: D439–D444.

Vincentelli R., K. Luck, J. Poirson, J. Polanowska, J. Abdat, *et al.*, 2015 Quantifying domain-

ligand affinities and specificities by high-throughput holdup assay. Nat. Methods 12: 787–

793.

Wang C.-H., P. Mehta, and C. J. Bashor, 2018 The strength of protein-protein interactions

controls the information capacity and dynamical response of signaling networks. bioRxiv

469197.

Wickham H., M. Averick, J. Bryan, W. Chang, L. D. McGowan, *et al.*, 2019 Welcome to the

tidyverse. Journal of Open Source Software 4: 1686.

Wilke C. O., 2020 cowplot: Streamlined Plot Theme and Plot Annotations for "ggplot2"

Yu H., J. K. Chen, S. Feng, D. C. Dalgarno, A. W. Brauer, *et al.*, 1994 Structural basis for the

binding of proline-rich peptides to SH3 domains. Cell 76: 933–945.

Zarin T., and B. Lehner, 2024 A complete map of specificity encoding for a partially fuzzy

protein interaction. bioRxiv 2024.04.25.591103.

Zarrinpar A., S.-H. Park, and W. A. Lim, 2003 Optimization of specificity in a cellular protein

interaction network by negative selection. Nature 426: 676–680.

Zarrinpar A., R. P. Bhattacharyya, M. P. Nittler, and W. A. Lim, 2004 Sho1 and Pbs2 Act as

Coscaffolds Linking Components in the Yeast High Osmolarity MAP Kinase Pathway. Mol.

Cell 14: 825–832.

1628