SARS-CoV-2 Infection of Salivary Glands Compromises Oral Antifungal Innate

Immunity and Predisposes to Oral Candidiasis

Areej A. Alfaifi[1,2,3], Tristan W. Wang[1], Paola Perez [4], Ahmed S. Sultan[1,5], Timothy F.

Meiller[1,5], Peter Rock[6], David E. Kleiner[7], Daniel S. Chertow[8,9], Stephen M. Hewitt[7],

Billel Gasmi[7], Sydney Stein[8], Sabrina Ramelli[8], Daniel Martin[10], Blake M. Warner*[4],

Mary Ann Jabra-Rizk*[1,11]

[1] Department of Oncology and Diagnostic Sciences, School of Dentistry, University of

Maryland Baltimore, Maryland, United States of America

[2] Department of Restorative and Prosthetic Dental Sciences, College of Dentistry, King

Saud bin Abdulaziz University for Health Sciences, Riyadh, Saudi Arabia

[3] King Abdullah International Medical Research Center (KAIMRC), Riyadh, Saudi Arabia

[4] Salivary Disorders Unit, National Institute of Dental and Craniofacial Research, National

Institutes of Health, Bethesda, Maryland, United States of America

[5] University of Maryland Greenebaum Cancer Center, University of Maryland Baltimore,

Maryland, United States of America

[6] Department of Anesthesia, School of Medicine, University of Maryland Baltimore,

Maryland, United States of America

[7] Laboratory of Pathology, Center for Cancer Research, National Cancer Institute,

National Institutes of Health, Bethesda, Maryland, United States of America

[8] Critical Care Medicine Department, Clinical Center, National Institutes of Health,

Bethesda, MD, USA

[9] Laboratory of Virology, National Institute of Allergy and Infectious Diseases, National

Institutes of Health, Hamilton, MT, United States of America, USA

[10] Genomics and Computational Biology Core, National Institute of Dental and

Craniofacial Research, National Institutes of Health, Bethesda, Maryland, United States

of America

[11] Department of Microbiology and Immunology, School of Medicine, University of

Maryland, Baltimore, Maryland, United States of America

* Corresponding authors

Mary Ann Jabra-Rizk, PhD

E-mail address: mrizk@umaryland.edu

Blake M. Warner, DDS, PhD, MPH

E-mail address: blake.warner@nih.gov

Short title: SARS-CoV-2 Infection of Salivary Glands Predisposes to Oral Candidiasis

ABSTRACT

Saliva contains antimicrobial peptides considered integral components of host innate

immunity, and crucial for protection against colonizing microbial species. Most notable is

histatin-5 which is exclusively produced in salivary glands with uniquely potent antifungal

activity against the opportunistic pathogen *Candida albicans*. Recently, SARS-CoV-2 was

shown to replicate in salivary gland acinar cells eliciting local immune cell activation. In

this study, we performed mechanistic and clinical studies to investigate the implications

of SARS-CoV-2 infection on salivary histatin-5 production and *Candida* colonization. Bulk

RNA-sequencing of parotid salivary glands from COVID-19 autopsies demonstrated

statistically significant decreased expression of histatin genes. *In situ* hybridization,

coupled with immunofluorescence for co-localization of SARS-CoV-2 spike and histatin

in salivary gland cells, showed that histatin was absent or minimally present in acinar cells

with replicating viruses. To investigate the clinical implications of these findings, salivary

histatin-5 levels and oral *Candida* burden in saliva samples from three independent

cohorts of mild and severe COVID-19 patients and matched healthy controls were

evaluated. Results revealed significantly reduced histatin-5 in SARS-CoV-2 infected

subjects, concomitant with enhanced prevalence of *C. albicans*. Analysis of prospectively

recovered samples indicated that the decrease in histatin-5 is likely reversible in mild-

moderate disease as concentrations tended to increase during the post-acute phase.

Importantly, salivary cytokine profiling demonstrated correlations between activation of

the Th17 inflammatory pathway, changes in histatin-5 concentrations, and subsequent

clearance of *C. albicans* in a heavily colonized subject. The importance of salivary

histatin-5 in controlling the proliferation of *C. albicans* was demonstrated using an *ex vivo*

assay where *C. albicans* was able to proliferate in COVID-19 saliva with low histatin-5,

but not with high histatin-5. Taken together, the findings from this study provide direct

evidence implicating SARS-CoV-2 infection of salivary glands with compromised oral

innate immunity, and potential predisposition to oral candidiasis.

AUTHOR SUMMARY

Saliva contains antimicrobial peptides part of host innate immunity crucial for protection against colonizing microbial species. Most notable is the antifungal peptide histatin-5 produced in salivary glands cells. SARS-CoV-2 was shown to replicate in salivary gland cells causing tissue inflammation. In this study, we showed decreased expression of histatin genes in salivary glands from COVID-19 autopsies, and co-localization studies of SARS-CoV-2 spike and histatin revealed absence or minimal presence of histatin in acinar cells with replicating virus. To investigate the clinical implications of these findings, we conducted studies using saliva samples from subjects with mild to severe COVID-19, matched with healthy controls. Results revealed significantly reduced histatin-5 in SARS-CoV-2 infected subjects with enhanced prevalence of *C. albicans*. Prospective analysis indicated the decrease in histatin-5 is reversible in mild-moderate disease, and salivary cytokine profiling demonstrated activation of the Th17 inflammatory pathway. The importance of salivary histatin-5 in controlling the proliferation of *C. albicans* was demonstrated using an *ex vivo* assay where *C. albicans* was able to proliferate in saliva with low histatin-5, but not with high histatin-5. Collectively, the findings provide direct evidence implicating SARS-CoV-2 infection of salivary glands with compromised oral innate immunity and predisposition to oral candidiasis.

INTRODUCTION

The oral cavity remains an underappreciated site for SARS-CoV-2 infection despite the evident myriad of oral conditions observed in COVID-19 patients, and the presence of the virus in saliva (1-6). The homeostasis of the oral cavity is maintained by saliva, an

extracellular fluid produced by salivary glands possessing a wealth of protective

properties (7). Specifically, saliva is enriched with antimicrobial peptides considered to be

part of the host Th17-type adaptive immune response that play a vital role in innate

immune defenses against microbial species (7-10). Most notable are histatins, a family of

peptides exclusively produced in the acinar cells of the salivary glands and secreted into

saliva (11, 12). There are two genes for histatins, *HTN1* and *HTN3* which encode

histatins-1 and 3, respectively (13). Histatin-5, a proteolytic product of histatin-3, is the

most abundant and unique as it exhibits potent antifungal activity against the fungal

pathogen *Candida albicans* (*C. albicans*) (12, 14-18). Although *C. albicans* is a

commensal colonizer of the oral cavity, any changes in the host microenvironment

favoring its proliferation allows this opportunistic species to transition into a pathogen

causing oral candidiasis (thrush) (19, 20). As a commensal colonizer of the oral mucosa,

the host immune response to *Candida* is oriented toward a more tolerogenic state and,

therefore, local innate immune defenses and specifically histatin-5 play a central role in

maintaining *Candida* in its commensal state preventing development of oral candidiasis

(8, 21, 22).

Recently, the salivary glands were shown to be a potential target for SARS-CoV-2

infection as several studies demonstrated the concomitant expression of

ACE2/transmembrane serine proteases 2 (TMPRSS2) in salivary glands epithelial cells

(23-27). In fact, virus entry into salivary glands cells was found to be higher, as compared

with entry into lung cells (25). Most notable are findings from a landmark study by Huang

and Perez *et al.* (2021) (28) which comprehensively demonstrated that salivary gland

acinar and ductal cells are robust sites for SARS-CoV-2 infection and replication, eliciting

local immune cell activation. Additionally, architectural distortion, atrophy, fibrosis, and

ductal rupture were also revealed, establishing that the minor and major salivary glands

are susceptible sites for infection, replication and local immune cell activation. Moreover,

the authors demonstrated that saliva from acutely infected patients could infect and

replicate in Vero cells, *ex vivo*, indicating that the source of virus in saliva is likely derived

from infected cells in the salivary glands (28). In fact, clinical observations are in support

of SARS-CoV-2 mediated damage to the salivary glands as COVID-19 patients frequently

present with gustatory dysfunction and xerostomia, and cases of inflamed salivary glands

have been reported in this population (26, 29).

Unlike other antimicrobial peptides, histatins are exclusively expressed and secreted

by acinar cells of salivary glands; saliva containing histatins is secreted from the acini into

the lumen which transits the ductal network into the oral cavity where they can exert their

antimicrobial activities. Patients infected with SARS-CoV-2 have the propensity to

develop superimposed infection including oral candidiasis. While SARS-CoV-2 infects

and replicates in the acinar and ductal cells of the salivary glands, it is not known if the

risk of oral candidiasis is due to affects on the acinar expression of anti-candidal proteins,

or due to viral and immune-mediated destruction of the glands. To that end, we performed

mechanistic and clinical studies to investigate the implications of SARS-CoV-2 infection

of salivary gland tissue on oral innate immune defenses, salivary histatin-5 production,

and predisposition to opportunistic infections. Using *in situ* hybridization and

immunofluorescence, we performed co-localization studies to assess histatin production

in SARS-CoV-2 infected salivary gland acinar cells. To provide mechanistic insights, the

expression of histatin genes was comparatively evaluated in infected and uninfected

salivary gland tissue. The clinical implications of findings from mechanistic studies were

demonstrated using various cohorts of SARS-CoV-2 infected subjects to evaluate salivary

histatin-5 levels and oral *Candida* colonization. Collectively, the novel findings from this

study establish the oral cavity as a robust site for SARS-CoV-2 infection warranting

reassessment of the risks for oral opportunistic infections in COVID-19 patients.

RESULTS

SARS-CoV-2 infection of the parotid glands affects the expression of histatins. To

determine the effect of SARS-CoV-2 infection on the expression of anti-candidal proteins

in the salivary glands, RNA sequencing (RNAseq) and confirmatory hybrid *in situ*

hybridization and immunofluorescence microscopy, was used. First, RNAseq of SARS-

CoV-2 infected parotid glands showed significantly reduced expression of histatin genes

(*HTN1,* >100-fold lower, p=0.0022*; HTN3,* >100-fold lower, p=0.0038; Fig. 1a)*.* Next, we

confirmed the loss of histatin expression and the co-localization of SARS-CoV-2 and

histatin protein expression in acinar cells. Hybrid *in situ* hybridization and

immunofluorescence microscopy co-localization studies of SARS-CoV-2 and histatins,

respectively, demonstrated significant (p<0.001) reduction of histatin protein expression

in virus infected acinar cells in parotid gland tissues obtained from deceased COVID-19

patients. As previously reported, the detection of SARS-CoV-2 is non-uniform across the

acini of the glands (28). Comparing infected vs non-infected acinar cells, histatin protein

expression intensity was inversely proportional to viral count (Fig.1b-f) indicating that

direct viral infection of the glands suppresses histatin mRNA and protein expression.

SARS-infected patients secrete significantly less histatins potentiating oral candidiasis. Clinical studies were conducted using two COVID-19 patient cohorts (hospitalized [UMD] and outpatient [NIDCR]) with ranging disease severity, and matched healthy controls. In the hospitalized cohort, a histatin-5 specific immunoassay to measure salivary histatin-5 concentrations was used and significant (p=0.0303) differences in values between the COVID-19 patients and matched healthy controls were seen. Average concentrations for the control group was 21.3 µg/ml (11.1 µg/ml-34.5 µg/ml), and 18.4 µg/ml (1.4 µg/ml-53.5 µg/ml) for the COVID-19 group. Strikingly, 14 of the COVID-19 patients had concentrations below 10 µg/ml (Fig. 2a), and although unclear why, 3 subjects had concentrations higher than any of the controls. The difference in concentrations between the two groups was also seen in age- and race-matched comparison between each COVID-19 patient and their matched control subject (Fig. 2b). *Candida* colonization was assessed by culturing samples; where no *Candida* was recovered from any of the healthy controls, 45% (9/20) of COVID-19 patients sampled were positive for *C. albicans*, some heavily colonized (Fig. 2a).

To corroborate these findings, a separate outpatient cohort collected by the National Institute of Dental and Craniofacial Research (NIDCR) (28) of serially sampled acutely infected patients was analyzed for histatin-5 expression. Compared to healthy control samples collected prior to the pandemic (n=9), the acute phase COVID-19 whole unstimulated saliva (n=23) had significantly lower histatin-5 concentrations (3.63 vs 10.41 µg/ml, *p*<0.0001; Fig. 2c). *Candida* colonization was assessed as above; no *Candida* was recovered from the healthy controls, yet 17% (4/23) of COVID-19 patients sampled were positive for *C. albicans.* Despite high variability in the expression of histatin-5 across the

cohort, prospective analysis of available serially-collected samples (n=8; 3-15 days from

symptom onset, 6-months, and 1-year from initial infection) from these COVID-19

subjects indicated a slight, but statistically significant trend in the restoration histatin-5

concentration from the post-acute to the chronic phase (R^2=0.30, p=0.0046; Fig. 2d).

However, some patients exhibited persistence of low salivary histatin concentration for up

to 1 year after recovery in the chronic phase. No statistical differences (p values > 2.0) in

histatin-5 concentrations were seen between subjects based on disease severity or other

variables (gender, race, age).

Significant changes in histatin-5 concentrations over the course of COVID-19

disease in prospectively sampled subjects. A total of 5 subjects with mild-moderate

disease were prospectively sampled and salivary histatin-5 was measured to monitor

changes in concentrations. Two of the subjects were longitudinally sampled up to 39 and

80 days, respectively (Fig. 3). For these subjects, saliva and histatin-5 concentrations

prior to COVID-19 infection were available. The baseline histatin-5 concentration

measured prior to COVID-19 disease for *Subject #1* was 15.7 µg/ml. However, the

concentration in the first sample recovered during the acute phase of the disease was 5.6

µg/ml (64.33% drop) and 8.8 µg/ml for the second sample recovered 2 days later. The

concentration gradually began to increase during the post-acute phase with a spike noted

on Day 15 (25.3 µg/ml) returning to baseline level in the last sample analyzed (Fig. 3a).

For *Subject #2*, the histatin-5 concentration measured prior to COVID-19 disease was

32.8 µg/ml; however, the concentration in the first sample recovered during the post-acute

phase of the disease was 15.65 µg/ml (52.29% drop) which gradually increased over

subsequent days with a spike noted on Day 45 (43.6 µg/ml) before returning to pre-

COVID-19 levels on last day sampled (Fig. 3c). Significantly, *C. albicans* was recovered

from the initial 6 samples recovered from *Subject #2*, but not from the last 3 samples that

followed the restoration of histatin-5 to pre-COVID-19 level (Fig. 3c). Three additional

subjects (Subjects 3, 4, and 5) were also prospectively sampled (Fig. 4), however, for this

group, pre-COVID-19 saliva was not available. For these subjects, the histatin-5

concentrations in the first samples recovered during the acute phase of the disease were

2.4, 4.1 and 5.6 µg/ml, respectively which increased during the post-acute phase to 10.9,

13.2, and 12.5 µg/ml, respectively in the last samples tested (Fig. 4). No *Candida* was

recovered from any of the samples from these 3 subjects.

Activation of Th17 inflammatory pathway concomitant with changes in histatin-5

concentrations. Salivary cytokine profiling was performed on samples recovered from

the 5 prospectively sampled subjects. Comparative analysis of samples from *Subject #1*

demonstrated a notable increase in the Th17 associated inflammatory cytokines;

compared to baseline pre-COVID-19 sample (IL-17A, IL-17F, IL-6, IL-10, TNF-α, IFN-**γ**,

IL-22, GMCSF and IL-23); concentrations gradually increased with a spike on Day 15,

decreasing over subsequent days and returning to baseline level in the last sample

measured (Day 39) (Fig. 3b). Although IL-1β followed the same trend, the spike in

concentration was seen in the first sample recovered during the acute phase of the

disease. Cytokine levels in *Subject #2* similarly demonstrated an increase in the Th17

associated inflammatory cytokines; compared to baseline pre-COVID-19 sample,

concentrations of all cytokines (IL-17A, IL-17F, IL-6, IL-10, TNF-α, IFN-**γ**, IL-22, GMCSF,

IL-23 and IL-1β) spiked on Day 38, gradually decreasing and returning to baseline level

on Day 45. IL-17E was not detectable in all samples analyzed for *Subject #1* and *Subject*

#2 (Fig. 3b, 3d). For the remaining 3 subjects, increase in Th17-associated inflammatory

cytokines was similarly observed during the post-acute phase for *Subject #3* (Fig. 4). Only

the first and last samples recovered from these 3 subjects were subjected for cytokine

analysis.

***C. albicans* proliferates in saliva from COVID-19 positive subjects with low histatin-**

5. Saliva recovered from subjects (n=3) during acute COVID-19 infection (low histatin-5)

and post-acute recovery (high histatin-5) phases were separately pooled and histatin-5

concentrations determined. Samples were subsequently tested *in vitro* for ability to control

C. albicans proliferation (Fig. 5a). Histatin-5 concentration in the pooled saliva sample

obtained during acute COVID-19 was 0.5 µg/ml (low), and in the pooled sample recovered

during post-recovery concentration was 11.5 µg/ml (high) (Fig. 5b). Based on *C. albicans*

growth (CFUs) following *in vitro* inoculation of saliva with $1x10^4$ cells/ml *C. albicans*, an

average of $2.55x10^4$ cells/ml *C. albicans* was recovered from the low histatin-5 sample

and $1.29x10^4$ cells/ml from the high histatin-5 sample (Fig. 5c).

DISCUSSION

To date, little is known about the physiological mechanisms of oral manifestations in

COVID-19 disease and the impact of SARS-CoV-2 infection on salivary gland function.

Clinical observations are in support of SARS-CoV-2 mediated damage to the salivary

glands, as COVID-19 patients frequently present with gustatory dysfunction

and xerostomia, and clinical cases of inflamed salivary glands have been reported in this

patient population (26).

Oral candidiasis is the most common opportunistic infection, particularly in

immunocompromised individuals, although immunocompromization due to COVID-19 is

ardently debated (1). However, the onset of oral candidiasis early in HIV disease indicates

that immunity to *Candida* is site-specific and involves secondary local innate defenses

(21, 22). In fact, innate immunity not only represents the first line of defense providing the

initial host response to pathogens, but also activates the adaptive immunity (8). These

systems exhibit coordinated regulation and response to establish and maintain tissue

homeostasis (8). Host-produced salivary antimicrobial peptides are considered to play a

vital role in innate immune defenses against microbial species, specifically histatins which

are localized in the serous acinar cells and secreted into saliva (7, 12, 14).

The recent demonstration of SARS-CoV-2 replication in salivary glands and

destruction of gland integrity including at sites where histatins are produced. These

findings indicate that histatin production may be compromised in infected individuals. This

hypothesis was validated using transcriptional analysis where the expression of both

histatin genes (*HTN1* and *HTN3)* were shown to be downregulated in SARS-CoV-2

infected salivary gland tissue (Fig. 1a). Importantly, SARS-CoV-2 and histatin co-

localization studies indicated an inversely proportional association where little or no

histatin signal was detected in virus infected acinar cells compared to uninfected cells

(Fig. 1b-f). The clinical implications of these findings were subsequently revealed using

various cohorts of SARS-CoV-2 infected subjects. Overall, analysis of saliva samples

demonstrated significant reduction in histatin-5 levels in infected subjects compared to

healthy controls (Fig. 2). However, despite an association between infection and reduced

histatin expression, the exact temporal relationship could not be perfectly gleaned. To

better understand the temporal relationship between infection and histatin expression, we

included a panel of prospective samples from moderately symptomatic COVID-19

subjects, which allowed us to glean some insights (Fig. 3, 4). Specifically, analysis

indicated that the decrease in histatin-5 during COVID-19 is likely reversible as

concentrations tended to increase during the post-acute phase of the disease. However,

it is important to iterate that co-localization studies exhibited an inversely proportional

association between histatin and viral counts in infected acinar cells (Fig. 1c-f), and some

patients experienced long-term deficits in salivary histatin-5 levels (42 days to 1 year).

Therefore, in severe COVID-19, a robust salivary infection and the ensuing adaptive

immune response may irreparably damage the salivary gland parenchyma. Thus, due to

pathologically lower histatin-5 secretion (and likely reduced saliva secretion overall),

affected individuals may remain predisposed to both recurrent and recalcitrant to

antifungal therapy oral candidiasis as part of the Long COVID syndrome (30)

(https://www.covid.gov/be-informed/longcovid/about#term). It is important to note here

that there are no known set "*normal*" values for salivary histatin-5 concentrations and

what is considered physiological is dependent on many factors including: host factors

(e.g, age, sex), time of collection, the collection method, and post-processing. Moreover,

histatin-5 is expression is secreted primarily by serous acinar cells; these cells

predominate in the parotid glands but are half the acinar cells in the submandibular

glands, and almost entirely absent from the sublingual glands. The relative contributions

of each gland to the total saliva collected is also host specific. Therefore, it is expected to

see a wide range of histatin-5 concentrations among the healthy control subjects.

However, based on our experience using our assay, prior clinical studies, and evaluations

of anti-candidal activity, we have set arbitrary concentrations as a general guidelines for

what is physiological (31).

With limited available clinical information on the hospitalized cohort population

(including presence of oral lesions), we recognize that we cannot fully account for

confounding factors that may influence histatin-5. However, our study focuses on the

presence of SARS-CoV-2 virus rather than disease severity; importantly, on average we

observed lower histatin-5 levels for infected subjects in all cohorts studied (hospitalized,

outpatient and individual case). In our analysis, we did not find strong association

between histatin-5 concentrations and *Candida* colonization regardless of disease status

or demographics. One intriguing observation in the samples from the hospitalized cohort

is the exceptionally high histatin-5 concentrations seen for 3 of the subjects with values

higher than that of any of the healthy controls (Fig. 2a). Although unclear as to why, it is

important to note that in the two prospectively sampled subjects with known pre-COVID-

19 histatin-5 values, a spike in histatin-5 and inflammatory cytokines was noted over the

course of the disease progression prior to return to baseline levels (Fig. 3). Therefore, we

speculate that these three hospitalized subjects may have been sampled during these

transient spikes. Interestingly, all 3 subjects with high histatin-5 levels were also positive

for *C. albicans* (Fig. 2a), and it is tempting to also speculate that these high values may

be part of the host immune response to *Candida* presence.

Importantly, our findings demonstrated that changes in histatin-5 concentrations were

concomitant with activation of the Th17 inflammatory pathway. Antimicrobial peptides,

including histatins, are part of the host Th17-type adaptive immune response (8).

Interleukin-17 (IL-17) is a pro-inflammatory cytokine that regulates multiple immune

events (32). In the tissues of the oral cavity, including the salivary glands, the IL-17/Th17

signaling pathway is essential for host protection against *C. albicans* infection (32, 33).

Therefore, it was not surprising that cytokine profiling of saliva samples from prospectively

sampled subjects displayed a trend for the Th17 associated cytokines similar to that for

histatin-5 (Fig. 3). Interestingly, in the subject heavily colonized with *C. albicans*, the spike

in cytokine concentrations coincided with increase in salivary histatin-5, and more

significantly, with the clearance of *C. albicans* (Fig. 3c). The activation of the Th17 immune

response in COVID-19 disease is expected as immunophenotyping and microscopic

analysis of SARS-CoV-2 infected salivary gland tissues demonstrated chronic, focal

lymphocytic sialadenitis with a T-lymphocytic infiltrates (28, 29). As IL-17 antifungal

activity involves regulation of the expression of antimicrobial peptides, including histatins

(8, 32, 33), the gradual increase in histatin-5 concomitant with clearance of *C. albicans*

may also be mediated by activation of Th17 response. Therefore, based on these

collective findings, we presume that decreased histatin-5 salivary levels due to SARS-

CoV-2 infection of salivary gland serous acini permits proliferation of colonizing *C.*

albicans leading to the development of oral candidiasis.

The importance of salivary histatin-5 in controlling the proliferation of *C. albicans* was

shown using an *ex vivo* anti-candidal assay where saliva with low histatin-5 was able to

allow *C. albicans* proliferation *in vitro* (Fig. 5). These findings provide experimental

evidence establishing the importance of salivary histatin-5 in preventing *Candida*

proliferation by maintaining colonizing *Candida* in the commensal state. Further, in

addition to anti-candidal activity, given the established anti-inflammatory and wound

healing properties of histatins (34-36), it is possible that compromised levels of histatins

(and other antimicrobial proteins) may contribute to other immune- and dysbiosis-

mediated mucosal conditions observed in COVID-19 patients such as burning mouth,

blisters and non-specific ulcerations, and aphthous-like lesions, all of which have been

associated with COVID-19 (5). With the current lack of knowledge on the implications of

SARS-CoV-2 infection on oral health, the findings from this study provide mechanistic

insights underscoring the oral cavities' diverse susceptibilities to SARS-CoV-2 infection,

prompting a reassessment of oral opportunistic infection risks and their potential long-

term impacts on oral health.

METHODS

Co-localization of SARS-CoV-2 and histatin in salivary gland tissue using *in situ*

hybridization and immunofluorescence, respectively. To investigate the expression

of histatin during SARS-CoV-2 infection, we employed the RNAscope Multiplex

Fluorescent v2 assay with antigen co-detection, adhering to the manufacturer guidelines.

Studies were performed on parotid FFPE (formalin-fixed, paraffin-embedded) biopsies

from uninfected donors (n=4) and from COVID-19 subjects (n=8) at autopsy with high and

low copy virus number. Uninfected donors were procured from the Human Cooperative

Tissue Network, and COVID-19 autopsy tissues were procured by the National Institutes

of Health COVID-19 Autopsy Consortium, as described previously (28, 37). Initially, slides

were dried in an air oven at 60°C for 30 min and subsequently rehydrated, and target

retrieval was performed using 1X Co-Detection Target Retrieval for 20 min at 99°C. Slides

were then rinsed briefly in water and washed in PBS-T (1X PBS, 0.1% Tween-20, pH 7.2),

and a hydrophobic barrier was established using an ImmEdge pen. Primary antibodies

against histatin-3 (Biorbyt; diluted 1:300 in Co-detection Antibody Diluent from ACD) and

AQP5 (SantaCruz; diluted 1:300 in Co-detection Antibody Diluent from ACD) were applied

and slides were incubated overnight at 4°C. Following washing in PBS-T, tissues and

antibodies were fixed in 10% Neutral Buffered Formalin (NBF) for 30 min at room

temperature, followed by another wash in PBS-T. Tissue sections were then treated with

RNAscope Protease Plus for 30 min at 40°C and immediately followed by probe

hybridization using either a positive control probe (320861), negative control probe

(320871), or the V-nCoV2019-S probe (848561). The V-nCoV2019-S probe is antisense

and specifically targets the spike protein sequence of viral RNA, providing insights into

infection and viral load within the tissue. Following hybridization, signals were amplified

and visualized using fluorescence Tyramine signal amplification. The primary antibodies

were detected using secondary antibodies conjugated to AF488 (histatin-3), and AF568

(AQP5; aquaporin-5) and counterstaining was performed with DAPI. Images were

captured using a Zeiss Axio Scan Z1 slide scanner light microscope equipped with a

40x/0.95 N.A. objective, and image files were imported into Visiopharm software version

2017.2. We first segmented the cells using the AI app from Visiopharm, once the cells

were segmented, using the APP Author function we designed an algorithm that detects

the dots of ISH and counts the number of dots per cell, and quantify the mean intensity

of histatin-3 per cell. The data was exported and analyzed in Prism 10. Control

experiments for *in situ* hybridization and immunofluorescence studies were performed

prior to sample analysis (Supp. Fig. 1).

Bulk RNAseq analysis of salivary gland tissue. Total RNA was extracted from

RNAlater (Invitrogen)-preserved parotid gland at autopsy from uninfected donors (n=3)

and COVID-19 subjects (n=8), using the RNeasy Mini (Qiagen) according to

manufacturer protocols. Following standard bulk RNAseq using Illumina platform,

normalized counts per million (CPM) were plotted for *HTN1* and *HTN3* and expression

levels of both genes were compared.

Study subjects. *Hospitalized cohort.* Adult COVID-19 patients (n=26) hospitalized at the

University of Maryland Medical Center and a matched control (age, race and gender)

group of healthy volunteers (n=26) were enrolled in the study. The University of Maryland

Baltimore Institutional Review Board approved this study, no patient identifiers were used

and informed consent was obtained from all subjects. Inclusion criteria for control subjects

included: healthy adults over 18 years of age with no history of oral candidiasis, any

predisposing factors or antifungal therapy. As the premise of the study is to investigate

the presence of the virus in the oral cavity regardless or disease severity, the only

inclusion criteria for the COVID-19 cohort was COVID-19 positivity. Of the 26 subjects, 11

were COVID-19 positive but asymptomatic and were hospitalized for other medical

reasons; the remaining 15 were symptomatic with severe disease, admitted to the

intensive care unit and/or receiving oxygen supplementation. The hospitalized COVID-19

population included 14 males and 12 females; 15 Caucasians and 11 African American

with age range of 29-76. *Outpatient cohort.* This cohort included 33 subjects with mild-

moderate disease; a total of 68 samples were prospectively collected from these subjects

at NIDCR (previously described) (28). *Prospective cases.* In addition to the cohorts, 5

otherwise healthy outpatient subjects (4 females, one male; age range 30-58) were

prospectively sampled at different timepoints during the course of their COVID-19

disease. A total of 29 samples were collected and analyzed for these subjects.

Clinical Samples. Saliva samples were collected from participating subjects using the

Salivette collection system as we previously performed (31, 38). Saliva was recovered,

aliquoted and immediately stored at -80°C with protease inhibitors. In addition, oral swabs

of the oral mucosal tissue were recovered for culturing to evaluate *Candida* recovery. Due

to inaccessibility to some patients the result of medical status, samples for fungal culturing

were only obtained from 20 of the 26 hospitalized patients. For the outpatient cohort and

their control subjects, whole unstimulated saliva samples were collected and processed

as above.

Evaluation of salivary *Candida* colonization. Oral swabs from all sampled subjects

were immediately cultured on fungal Yeast Peptone Dextrose (YPD) agar media (Difco

Laboratories) and incubated at 35°C for 24-48 h for fungal growth evaluation. The

chromogenic medium CHROMagar Candida was used for *Candida* speciation.

Measurement of histatin-5 (Hst-5) salivary levels using ELISA. ELISA was performed

as we previously described (31). A high purity Hst-5 peptide was synthesized by

GenScript and Hst-5 specific rabbit polyclonal antibody was produced by Lampire

Biological Laboratories. For measurement of Hst-5 levels, a standard curve was

performed with each assay using Hst-5 peptide concentrations ranging from 0.5-500

µg/ml. Wells of high-binding 96-well plates were coated with 100 µl of each Hst-5

concentration or 1/100 dilution of saliva. Following overnight incubation at 4°C, wells were

blocked with 0.1% dry milk in PBS for 1 h and anti-Hst-5 antibody (1/1000) (100 µl) was

added for 1 h at 37°C. Following washing, HRP-labeled goat anti-rabbit secondary

antibody (1/3000) (Abcam) was added, and plates incubated for 1 h at 37°C. Following

washing, 100 µl of ABTS Peroxidase Substrate (KPL, Inc.) was added and plates

incubated for 20 min until color develops. The reaction was stopped by the addition of 50

µl of Stop Solution (KPL, Inc.) and optical density (OD) was measured at 405nm using a

microtiter plate reader. A standard curve was plotted with each run; samples were tested

in triplicate on two separate occasions and the average Hst-5 concentration calculated in

µg/ml. As there are no set normal Hst-5 salivary concentrations, based on our experience

with clinical studies on Hst-5 salivary levels using our immunoassay, arbitrary

concentrations of approximately 9-10 µg/ml was considered the cutoff whereby lower

concentrations are considered in the low range (31).

Salivary cytokines profiling. Saliva samples from outpatient subjects were analyzed at

the University of Maryland Cytokine Core using the Luminex Multianalyte System. Each

sample was measured in triplicate and results expressed in Pg/ml. Not enough saliva was

available from other cohorts for cytokine analysis.

Ex vivo **salivary** *C. albicans* **inhibition assay.** In order to assess the anti-candidal

potency of saliva with respect to Hst-5 concentration, saliva samples were recovered from

3 subjects during COVID-19 and after recovery, then pooled to generate a COVID-19

positive and a COVID-19 negative sample. Hst-5 concentration in the pooled samples

was pre-determined by ELISA and samples were filter-sterilized and comparatively tested

for efficacy in inhibiting *C. albicans* proliferation *in vitro*. For these assays, cultures of the

standard *C. albicans* SC5314 strain were grown in YPD broth (Difco Laboratories)

overnight at 30°C with shaking and cells were washed and resuspended in sterile PBS

(1mM). *C. albicans* cells were added to each of the saliva samples (100 µl) at final cell

density of 1×10^4 cells/ml in the wells of a 96-well microtiter plate and plates were

incubated for 1 h at 37°C with shaking. Aliquots from reactions were diluted with PBS and

plated on YPD agar and incubated for 24–48 h at 35°C. The number of single colonies

on each plate was counted and the level of *C. albicans* proliferation was determined

based on CFU counts (cells/ml).

Statistical analysis. Statistical analysis was performed using GraphPad Prism 10.2.1. A

Wilcoxon rank-sum test was used to compare histatin-5 levels between COVID-19

patients and healthy control subjects and a t-test was used to evaluate associations

between histatin-5 values and patient characteristics (disease severity, *Candida*

colonization, demographics). For *in vitro* assays, histatin-5 levels in pooled saliva of

COVID-19 positive and negative subjects and *Candida* CFU counts were determined

using the unpaired two-sample t-tests. Significance of *HTN1* and *HTN3* expression levels

in parotid glands was determined using ANOVA analysis with Bonferroni correction for

multiple measurement. Figures were constructed using GraphPad Prism 10.2.1 and R

statistical programming software.

ACKNOWLEDGMENT. We would like to acknowledge the contributions of LaToya Stubbs

and Kendra Petrick who enrolled subjects and organized sampling of hospitalized cohort

at the University of Maryland Medical Center. We would like to thank the NIH COVID-19

Autopsy Consortium (PIs-Chertow & Kleiner) for coordination and access to tissues from

fatal COVID-19 cases. We give our special thanks to the members of the NIDCR

Sjögren's Disease Clinical Research Team for their coordination of patients and collection

of research data and tissues. Biorender.com was used to create elements of figures.

FUNDING. This work was funded by the National Institutes of Health (NIH), National

Institute of Dental and Craniofacial Research (NIDCR), Division of Extramural Research

(R21-DE031888, PI-Rizk-Jabra, Co-I-Warner). This project was also funded by the

University of Maryland Baltimore, Institute for Clinical & Translational Research (ICTR)

(1UL-1TR00309, PIs-M.A.J-R & P.R). Funding was also provided through the Intramural

Research Programs of the NIH/NIDCR (Z01-DE000704, PI-Warner), NIH Clinical Center,

and the National Institute of Allergy and Infectious Diseases (ZIA CL090070-03, PI-

Chertow). This research was supported in part by the NIDCR Genomics and

Computational Biology Core: ZIC DC000086.

AUTHOR CONTRIBUTIONS. M.A.J-R., B.M.W., A.S., T.F.M., P.R. conceived and

designed this research, M.A.J-R., B.M.W. provided funding; A.A.A., T.W.W., P.P., B.M.W.

performed the experiments; D.E.K, D.S.C., S.M.H., B.G., S.S., S.R., collected and

processed tissues and clinical data for COVID-19 autopsies; D.E.K, S.M.H., B.G.,

performed the autopsies; D.S.C. and D.E.K., led and oversaw the NIH COVID-19 Autopsy

Consortium; M.A.J-R., A.A.A., T.W.W., P.P., B.M.W., A.S., T.F.M., analyzed data; M.A.J-

R., A.A.A., T.W.W., B.M.W. wrote the paper; B.M.W. led and oversaw the execution of NIH

clinical and molecular studies supporting this work; M.A.J-R. oversaw the entire study.

REFERENCES

1.　　Amorim Dos Santos J, Normando AGC, Carvalho da Silva RL, De Paula RM,

Cembranel AC, Santos-Silva AR, et al. Oral mucosal lesions in a COVID-19 patient: New

signs or secondary manifestations? Int J Infect Dis. 2020;97:326-8.

2.　　Salehi M, Ahmadikia K, Badali H, Khodavaisy S. Opportunistic fungal Infections in

the epidemic area of COVID-19: A cinical and diagnostic perspective from Iran.

Mycopathologia. 2020 185(4):607-11.

3.   Salehi M, Ahmadikia K, Mahmoudi S, Kalantari S, Jamalimoghadamsiahkali S,

Izadi A, et al. Oropharyngeal candidiasis in hospitalised COVID-19 patients from Iran:

Species identification and antifungal susceptibility pattern. Mycoses. 2020; 63(8):771-8.

4.   Riad A, Gad A, Ockova B, Klugar M. Oral candidiasis in non-severe COVID-19

patients: Call for antibiotic stewardship. Oral Surg. 2020;15(3): 465-466.

5.   Amorim Dos Santos J, Normando AGC, Carvalho da Silva RL, Acevedo AC, De

Luca Canto G, Sugaya N, et al. Oral manifestations in patients with COVID-19: A living

systematic review. Dent Res. 2021;100(2):141-54.

6.   Brandini DA, Takamiya AS, Thakkar P, Schaller S, Rahat R, Naqvi AR. Covid-19

and oral diseases: Crosstalk, synergy or association? Rev Med Virol. 2021; 31(6): e2226.

7.   Vila T, Rizk AM, Sultan AS, Jabra-Rizk MA. The power of saliva: Antimicrobial and

beyond. Plos Pathog. 2019;15(11).

8.   Salvatori O, Puri S, Tati S, Edgerton M. Innate immunity and saliva in *Candida*

albicans-mediated oral diseases. J Dent Res. 2016;95(4):365-71.

9.   Sultan AS, Kong EF, Rizk AM, Jabra-Rizk MA. The oral microbiome: A lesson in

co-existence. PLoS Pathog. 2018;14(1):e1006719.

10.   Peters BM, Shirtliff ME, Jabra-Rizk MA. Antimicrobial peptides: Primeval

molecules or future drugs? PLoS Pathog. 2010;6(10):e1001067.

11.   Oppenheim FG, Xu T, McMillian FM, Levitz SM, Diamond RD, Offner GD, et al.

Histatins, a novel family of histidine-rich proteins in human parotid secretion. J Biol Chem.

1988;263(16):7472-7.

12.   Ahmad M, Piludu M, Oppenheim FG, Helmerhorst EJ, Hand AR.

Immunocytochemical localization of histatins in human salivary glands. J Histochem

Cytochem. 2004;52(3):361-70.

13.   Sabatini L, Azen E. Two coding change mutations in the HIS2(2) allele characterize

the salivary histatin 3-2 protein variant. Human Mutation. 1994;4(1):12-9.

14.   Edgerton M, Koshlukova SE, Araujo MWB, Patel RC, Dong J, Bruenn J. Salivary

histatin 5 and human neutrophil defensin 1 kill *Candida albicans* via shared pathways.

Animicrob Agents Chemother. 2000;44(12):3310-6.

15.   Edgerton M, Koshlukova S, Lo T, Chrzan B, Straubinger R, Raj P. Candidacidal

activity of salivary histatins. J Biol Chem. 1998;272(32):20438-47.

16.   Helmerhorst EJ, Troxler RF, Oppenheim FG. The human salivary peptide histatin

5 exerts its antifungal activity through the formation of reactive oxygen species. Proc Nat

Acad Sci. 2001;98(25):14637-42.

17.   Nikawa H, Jin C, Fukushima H, Makihira S, Hamada T. Antifungal activity of

histatin-5 against non-albicans *Candida* species. Oral Microbiol Immunol. 2001;16:250-

2.

18.   Situ H, Bobek L. In vitro assessment of antifungal therapeutic potential of salivary

histatin-5, two variants of histatin-5, and salivary mucin (MUC7) domain 1. Animicrob

Agents Chemother 2000;44(6):1485-93.

19.   Jabra-Rizk MA, Kong E, Tsui C, Nguyen M, Clancy C, Fidel P, et al. *Candida*

albicans pathogenesis: Fitting within the host-microbe damage response framework.

Infect Immun. 2016;84(10):2724-39.

20.   Vila T, Sultan AS, Montelongo-Jauregui D, Jabra-Rizk MA. Oral candidiasis: A

disease of opportunity. J Fungi (Basel). 2020;6(1):pii: E15.

21.   Fidel PJ. Immunity to *Candida*. Oral Dis. 2002;8(Suppl 2):69-75.

22.   Fidel PL, Jr. *Candida*-host interactions in HIV disease: Relationships in

oropharyngeal candidiasis. Adv Dent Res. 2006;19:80-4.

23.   Song J, Li Y, Huang X, Chen Z, Li Y, Liu C, et al. Systematic analysis of ACE2

and TMPRSS2 expression in salivary glands reveals underlying transmission mechanism

caused by SARS-CoV-2. J Med Virol. 2020 92(11):2556-66.

24.   da Silva Pedrosa M, Sipert CR, Nogueira FN. Are the salivary glands the key

players in spreading COVID-19 asymptomatic infection in dental practice? J Med Virol.

2021;93(1):204-5.

25.   Pascolo L, Zupin L, Melato M, Tricarico PM, Crovella S. TMPRSS2 and ACE2

coexpression in SARS-CoV-2 salivary glands infection. J Dent Res. 2020;99(10):1120-1.

26.   Corchuelo J, Ulloa FC. Oral manifestations in a patient with a history of

asymptomatic COVID-19: Case report. Int J Infect Dis. 2020;100:154-7.

27.   Tsuchiya H. Oral symptoms associated with COVID-19 and their pathogenic

mechanisms: A literature review. Dent J (Basel). 2021;9(3):32.

28.   Huang N, Pérez P, Kato T, al. e. SARS-CoV-2 infection of the oral cavity and saliva.

Nat Med 2021;27(5):892-903.

29.   Shen Y, Voigt A, Goranova L, Abed M, Kleiner DE, Maldonado JO, et al. Evidence

of a Sjögren's disease-like phenotype following COVID-19 in mice and humans. JCI

Insight. 2023;8(24):e166540.

30.   Pisano M, Romano A, Di Palo MP, Baroni A, Serpico R, Contaldo M. Oral

candidiasis in adult and pediatric patients with COVID-19. Biomedicines.

2023;11(13):846.

31.   Khan SA, Fidel Jr P, Al Thunayyan A, Meiller T, Jabra-Rizk MA. Impaired histatin-

5 level and salivary antimicrobial activity against *C. albicans* in HIV-infected individuals. J

AIDS Clin Res. 2013;4(2):1-6.

32.   Gaffen SL, Moutsopoulos NM. Regulation of host-microbe interactions at oral

mucosal barriers by type 17 immunity. Sci Immunol. 2020;5(43):eaau4594.

33.   Pavlova A, Sharafutdinov I. Recognition of *Candida albicans* and role of innate

type 17 immunity in oral candidiasis. Microorganisms. 2020;8(9):1340.

34.   Kong E, Tsui C, Boyce H, Ibrahim A, Hoag S, Karlsson A, et al. Development and

in vivo evaluation of a novel histatin-5 bioadhesive hydrogel formulation against oral

candidiasis. Antimicrob Agents Chemother. 2015;60(2):881-9.

35.   Sultan AS, Rizk AM, Vila T, Ji Y, Masri R, Jabra-Rizk MA. Digital design of a

universal rat intraoral device for therapeutic evaluation of topical formulation against

Candida-associated denture stomatitis Infect Immun. 2019 87(12):e00617-19.

36.   Sultan AS, Vila T, Hefni E, Karlsson AJ, Jabra-Rizk MA. Evaluation of the antifungal

and wound healing properties of a novel peptide-based bioadhesive hydrogel formulation.

Antimicrob Agents Chemother. 2019;63(10):e00888-19.

37.   Stein SR, Ramelli SC, Grazioli A, Chung JY, Singh M, Yinda CK, et al. SARS-CoV-

2 infection and persistence in the human body and brain at autopsy. Nature Biotech.

2022;612(7941):758-63.

38.   Alfaifi A, Sultan A, Montelongo-Jauregui D, Meiller TF, Jabra-Rizk MA. Long-term

post-COVID-19 associated oral inflammatory sequelae. Front Cell Infect Microbiol.

2022;12:831744.

FIGURE LEGEND

Figure 1. Impact of SARS-CoV-2 salivary gland infection on histatin production,

gene expression and salivary levels. (a) Parotid expression of histatin genes (*HTN1*

and *HTN3*) from deceased COVID-19 subjects (n=8) and healthy control subjects (n=3)

using bulk RNA sequencing. **(b-f)** Co-localization studies using *in situ* hybridization for

SARS-CoV-2 (*white dots* and *arrows*) and immunofluorescence for histatin-3 (*green*) with

quantitative correlation between the intensity of histatin and SARS-CoV-2 counts within

infected acinar cells of parotid tissue from **(b)** healthy subject (*white asterisk,* non-specific

signal) and **(c, e)** deceased COVID-19 patients. Acini (dotted circles) identified based on

expression of AQP5 (*red,* apical membrane); *inset* (*orange box*) shows the presence of

the viral genome in acinar cells of the parotid glands. **(d, f)** Per acinar cell expression of

histatin and the per cell viral count are inversely proportional as shown in two

representative COVID-19 cases (P7, P19).

Figure 2. Salivary histatin-5 (Hst-5) levels and fungal colonization in prospectively

sampled hospitalized and outpatient COVID-19 cohorts. (a) Boxplot with Hst-5

concentrations and *Candida* recovery from 26 saliva samples from hospitalized COVID-

19 patients and matched healthy controls. **(b)** Waterfall plot depicting 26 pairs of COVID-

19 and healthy subjects matched for race and age. Bar height represents differences

among pairs in Hst-5 levels (µg/ml) between healthy controls and COVID-19 patients. **(c)**

Bar plot depicts Hst-5 concentrations and *Candida* recovery of 9 healthy subjects and 23

COVID-19 outpatients' saliva. Red marked dots indicate cultured saliva contained

candidal outgrowth **(d)** Linear regression analysis of serially sampled COVID-19

outpatients' saliva (n=8) shows time-dependent restoration of Hst-5 concentration from

the post-acute phase to the chronic phase, based on longitudinally collected samples

from acute phase (3-15 days) to chronic phase (6-12 months). Blue line, linear fit; light

blue shading, confidence interval of the linear fit.

Figure 3. Prospective time course longitudinal analysis of saliva samples for Hst-5

from two subjects with moderate COVID-19. Changes in Hst-5 and cytokine

concentrations in two longitudinally sampled subjects prior to and during the acute and

post-acute phases of COVID-19 disease. Timeline of sampling during COVID-19 infection

up to 49 and 85 days post-COVID-19 infection for subjects 1 and 2 **(a and c,**

respectively). Measurement of Hst-5 concentrations and fungal culturing of samples

from the two subjects prior to and during the acute and post-acute phases of COVID-19

disease. Line graphs in bottom rows depict similar trends for Hst-5 and Th17 associated

cytokine levels for subjects 1 and 2 **(b and d, respectively)**. Upon culturing, *C. albicans*

was recovered from the initial 6 samples from subject 2 **(c, d)**.

Figure 4. Prospective sampling and analysis of saliva from three subjects with

moderate COVID-19. Timeline of sampling and measurement of salivary Hst-5 and

cytokine concentrations in samples recovered during the acute phase of the disease and

upon recovery. Table presents cytokine values measured in the first and last samples

recovered from the three subjects and the percent change in levels between the samples.

Figure 5. *Ex vivo* proliferation assay using pooled saliva. (a) Workflow for histatin-5

measurement and proliferation assay using pooled saliva samples from 3 subjects under

a COVID-19 infected state (PC+) and recovered state (PC-). **(b)** Bar plots depict Hst-5

concentrations (μg/ml) from pooled saliva samples; **(c)** recovered *C. albicans* (cells/ml)

following 1 h incubation in pooled saliva samples seeded with 1×10^4 cells/ml of *C.*

albicans.

SUPPLEMENTAL FIGURES

Supp Figure 1. *In situ* hybridization and immunofluorescence controls. (a)

Immunofluorescence (IF) and *in situ* hybridization (ISH) controls (see also **Fig. 1**).

Positive and negative controls for IF, antibody anti-Hst-3 in control parotid shows strong

IF (green) in acini structures and no signal is observed in ducts. The isotype control shows

no signal. **(b)** Probe anti-human *PPIB* shows strong signal in all the cells (white); negative

control probe shows sparse, rare positive signal in the tissue.

a

b

Pre-COVID-19 infection Acute phase (COVID-19 test positive) ● *C. albicans* negative

c

d

Pre-COVID-19 infection Acute phase (COVID-19 test positive) ● *C. albicans* positive ● *C. albicans* negative

Cytokine	Day 0	Day 35	% Change
IL-1ß (pg/ml)	122.56	150.51	22.80
IL-6 (pg/ml)	0.57	0.57	0
IFN- γ (pg/ml)	2.31	2.24	-2.92
IL-10 (pg/ml)	0.94	0.80	-14.82
IL-17A (pg/ml)	5.26	2.17	-58.78
TNF-α (pg/ml)	6.31	6.63	5.14
IL-22 (ng/ml)	ND	ND	N/A
IL-23 (ng/ml)	0.41	0.13	-69.46
IL-17F (ng/ml)	0.002	0.001	-69.64
GMCSF (ng/ml)	0.011	0.005	-52.69
IL-17E (ng/ml)	ND	ND	N/A

Cytokine	Day 0	Day 35	% Change
IL-1ß (pg/ml)	218.31	2991.20	1270.15
IL-6 (pg/ml)	2.02	13.80	584.19
IFN- γ (pg/ml)	4.67	4.06	-12.96
IL-10 (pg/ml)	1.28	3.53	175.77
IL-17A (pg/ml)	6.34	19.49	207.40
TNF-α (pg/ml)	7.91	38.98	392.69
IL-22 (ng/ml)	0.050	0.135	170.62
IL-23 (ng/ml)	0.60	1.54	157.01
IL-17F (ng/ml)	0.012	0.017	42.62
GMCSF (ng/ml)	0.016	0.052	221.11
IL-17E (ng/ml)	ND	ND	N/A

Cytokine	Day 0	Day 12	% Change
IL-1ß (pg/ml)	61.71	44.32	-28.18
IL-6 (pg/ml)	0.047	0.57	1103.07
IFN- γ (pg/ml)	1.35	2.72	100.48
IL-10 (pg/ml)	0.66	0.80	21.07
IL-17A (pg/ml)	ND	ND	N/A
TNF-α (pg/ml)	9.49	9.64	1.62
IL-22 (ng/ml)	ND	ND	N/A
IL-23 (ng/ml)	ND	0.08	N/A
IL-17F (ng/ml)	ND	ND	N/A
GMCSF (ng/ml)	ND	0.003	N/A
IL-17E (ng/ml)	ND	ND	N/A

Acute phase (COVID-19 test positive)

a

b

c