Microplastics found in high concentrations around Kantamanto,

the world's largest secondhand textile market in Accra, Ghana: A citizen science study

Dimitri D. Deheyn[1,♦,†], Joe Ayesu[2,♦], The Or Foundation[2,*] and Branson Skinner[†]

[1]Marine Biology Research Division, Scripps Institution of Oceanography, UC San Diego, 9500

Gilman Drive, La Jolla, CA 92093, USA.

[2]The Or Foundation, Unit 2C, 603 Southern Liberia Rd, Accra, Ghana

♦ Contributed equally to the work.

* List of personnel of The Or Foundation and citizen scientists who provided valuable help: Patrick

Abesiyine Anyebuno, Frank Aboyinga, Emmanuel Akyea, Brown Addoquaye, Bright Anokye,

Jacob Ayesu, Bright Ayikpa, Priscilla Danso, Abena Essoun, Emmanuel Ewudzei, Al-Hassan

Fattawu, Clémence Faure, Musah Iddrisu, Suleman Issifu, Maxwell Klutse, Freeheart Kordah,

Somed Mohammed, Nutifafa Mensah, Katia Osei, Sammy Oteng, Peter Yeboah, and Liz Ricketts

and Megan Schuknecht.

[†] Corresponding author: ddeheyn@ucsd.edu; lizandbranson@theor.org

Microplastics found in high concentrations around Kantamanto, the world's largest second-hand textile market in Accra, Ghana: A citizen science study

Dimitri D. Deheyn[1,♦,†], Joe Ayesu[2,♦], The Or Foundation[2,*] and Branson Skinner[†]

[1]Marine Biology Research Division, Scripps Institution of Oceanography, UC San Diego, 9500 Gilman Drive, La Jolla, CA 92093, USA.

[2]The Or Foundation, Unit 2C, 603 Southern Liberia Rd, Accra, Ghana

♦ Contributed equally to the work.

* List of personnel of The Or Foundation and citizen scientists who provided valuable help: Patrick Abesiyine Anyebuno, Frank Aboyinga, Emmanuel Akyea, Brown Addoquaye, Bright Anokye, Jacob Ayesu, Bright Ayikpa, Priscilla Danso, Abena Essoun, Emmanuel Ewudzei, Al-Hassan Fattawu, Clémence Faure, Musah Iddrisu, Suleman Issifu, Maxwell Klutse, Freeheart Kordah, Somed Mohammed, Nutifafa Mensah, Katia Osei, Sammy Oteng, Peter Yeboah, and Liz Ricketts and Megan Schuknecht.

† Corresponding author: ddeheyn@ucsd.edu, lizandbranson@theor.org

Authors biography

Dimitri Deheyn, PhD

Dimitri is a Research Scientist at the Marine Biology Research Division of Scripps Institution of Oceanography, University of California San Diego, where he conducts cross-disciplinary research on biomimicry. In this frame, he studies alternative materials to emerging pollutants, which includes plastics, textile microfibers and associated additives. Some key steps of his studies include assessing plastic and fiber biodegradation (from both natural and synthetic fibers, untreated and treated with additives). His research has a global perspective, across a diversity of ecosystems, around the world. Dimitri grew up in Africa, and earned his PhD in marine sciences and ecotoxicology from the Free-Thinking University in Brussels, Belgium where he started his academic career.

Joe Ayesu

Joe is an integral member of the Accra Arts Centre beachfront community, where he has a shop selling works from across West Africa's arts and crafts industry, including many of his own creations. Living and working in this location for decades he has observed Accra's waste management challenges firsthand, particularly those related to textiles. Joe majored in science in his secondary school days and studied the fundamentals of medical laboratory technology at Accra Technical University. He manages The Or Foundation's team of citizen scientists, mobilizing dozens of people for beach monitoring and cleanup efforts, sample collection and imaging.

The Or Foundation

Headquartered in Accra, Ghana, The Or Foundation is the leading not-for-profit implementing Globally Accountable Extended Producer Responsibility for textiles through a voluntary program in service of Accra's Kantamanto Market community, the largest secondhand clothing market in the world. Working at the intersection of environmental justice, education and fashion innovation since 2011, the mission of the NGO is to identify and manifest alternatives to the dominant model of fashion – alternatives that bring forth ecological prosperity, as opposed to destruction, and that inspire citizens to form a relationship with fashion that extends beyond their role as consumer.

Branson Skinner

Branson was studying in Ghana when he was confronted with the impact of textile waste on the environment and communities in the country. This led him to co-found The Or Foundation, a Ghana-based non-profit working at the intersection of environmental justice, education and fashion development. At The Or Foundation, Branson has combined his diverse background as an educator, farmer, urban planner, and musician to work closely with the local community in Accra and lead the organization's work implementing innovative solutions and science-backed research initiatives to address fashion's global waste crisis. Branson studied Globalization of Public Health and Societies at NYU Gallatin, where he was a Reynolds Scholar for Social Entrepreneurship. He also holds a Masters of Community Planning degree from the University of Cincinnati.

Abstract:

Located in Accra, Ghana, Kantamanto is the world's largest secondhand resale and upcycling

market for clothing and textiles, receiving tons of garments every week exported under HS Code

6309 by countries in the Global North. We looked at the abundance of microplastics and

microfibers from textiles and other sources in this dynamic environment. Our primary interest was

assessing how microfibers in a textile-rich environment contribute to overall microplastic

pollution. Continuous monitoring of air quality using PurpleAir sensors showed large

concentrations of airborne microparticles around the market, which were shown to drop with rain

events. Water collected from rain contained dozens to hundreds of microfibers and microplastics

(roughly in a 1:10 ratio), depending on the intensity of the rain and on whether the rain was

preceded by a significant period of dry weather. Microfiber and microplastic counts from airborne

samples showed concentrations about >20x to >100x greater, respectively, than those reported

from other metropolises around the world. A comparison of concentrations from water in the

adjacent Korle Lagoon was similarly striking, with up to 45x more microfibers, and up to 200x

more microplastics than what has previously been reported for lagoon waters worldwide. Beyond

the microscale, the issue of whole garments as waste was also demonstrated, with a strong

similarity found between the global brand garment tags from fashion items discarded in the market

and those that were washed up on nearby beaches, highlighting the need for more holistic

regulatory frameworks to address textile waste.

Keywords: Microplastic epifluorescence; Textile microfibers; Airborne dispersion; Rain flush;

Lagoon water contamination; Fast Fashion Waste; Sampling methods for citizen science

INTRODUCTION

Significant work has been undertaken to establish the environmental footprint of textile manufacturing operations (Madhav et al. 2018; Niinimäki et al. 2020; Grazzini et al. 2021; Farhana et al. 2022; Lim et al. 2022) and to characterize pollution resulting from garment use and maintenance at the primary consumer level (Carr 2017; Carney Almroth et al. 2018; Liu et al. 2019; Nowack et al. 2021; Liu et al. 2023). In contrast, efforts remain largely limited with regards to assessing the environmental impact of garments once they are discarded for secondhand reuse or for end-of-life in a landfill or incinerator. In general, fashion waste originating worldwide ultimately ends up in the Global South, particularly in African countries (Manieson and Ferrero-Regis 2021; UNComtrade 2023; OEC 2024), where the socio-environmental impacts of the waste remain poorly addressed.

Accra (Ghana) is the recipient of tons of fast fashion garments weekly and is home to Kantamanto Market, the largest secondhand textile trade market in the world. The market is adjacent to Korle Lagoon, a major water body found in the city's center with an outlet that connects to the Gulf of Guinea. The market is connected to the lagoon via a network of small drainage systems and outflow pipes feeding a tributary canal that leads to the lagoon. The area is mostly pedestrian, and the market has over 10,000 stalls, all exposed to some degree of open-air circulation though most are protected under a partial roof. About 15,000,000 garments exported from Global North countries enter the market every week, primarily on Thursdays, in the form of 55 kg bales that the market retailers purchase from importers (The Or Foundation 2022). The retailers work to recommodify the individual items through resale, repair and remanufacturing during the week. As for the undesired or damaged items that go unsold (the majority of which are pre-sorted for

reusability prior to arrival in Ghana), they are eventually cast aside, accumulating as waste around

the market, with piles of them being burned during the night to avoid excessive build-up, as more

space is continuously needed. As such, with Accra's waste infrastructure unable to handle and

safely process the continuously increasing volume of low-quality garments (Niinimäki et al. 2020),

the resulting textile waste can easily enter the surrounding ecosystems. This can take place through

airborne processes, by burning textiles with microparticles flying into the air, and/or though

waterborne processes by dumping garments regarded as waste (full or in pieces) in uncontrolled

sites of the lagoon shoreline that are adjacent to the market.

Over the years, the market has grown to become a critical economic center for the region, where

an estimated 30,000 people converge every day to work as market traders, upcyclers or to provide

auxiliary support. Considering customers, the number of people buzzing through the market goes

up to around 60,000 total daily. This large number of people combined with the constant traffic of

large volumes of textiles across the market makes Kantamanto a place with potential exposure to

microparticles, both microplastics (from packaging material, bags or film wraps) and microfibers

(from clothing). Indeed, although the washing and drying of textiles are obvious sources of shed

microfibers, another important source is related to wearing them and hanging them outside where

they are exposed to various environmental conditions (including high temperatures, humidity,

wind and UV) that facilitate their shedding (Carney Almroth et al. 2018; Belzagui et al. 2019; De

Falco et al. 2020; Lim et al. 2022; Zhang et al. 2022). Thus, this study offers new evidence on the

release of textile microfibers (viz. fiber fragments, with a length greater than the diameter) in a

dense urban setting and additional paths through which textile waste can enter the surrounding

ecosystems.

The dominant textile materials in Kantamanto are made of cotton but also plastic material,

such as polyester, nylon and elastane (and often in blends of two or more of these), particularly

given the global proliferation of cheaply made (and quicker to break down) fossil-based fast

fashion and the quick turnover between consumption and then disposal as secondhand textiles

(Niinimäki et al. 2020). The area is also the recipient of single use plastics like plastic bags, plastic

bottles and food containers that litter the market and surroundings.

Knowing all of the above, we hypothesized that the air and water would contain high

concentrations of microplastics, and plastic microfibers as well as cotton microfibers. It is also fair

to assume that anyone living or working in Kantamanto, or passing through the market, would be

exposed to air pollution, and potentially water pollution, in relation to the ambient load of

microplastics and microfibers (both synthetic and natural). We also hypothesized that whole

garments found as waste on beaches outside the lagoon would originate from the textile dumpsite

of the market, which overflows into the lagoon. As such, the garments would show similarities (in

terms of brand assemblage and polymer constituent as identifier) to the garments discarded in the

market.

Within this context, the research presented here involved a large contingent of citizen scientists

who worked to establish whether Kantamanto Market and its main dumpsite nearby generate

environmental changes on the surroundings, specifically with regard to the load of microparticles

in the air and in lagoon waters. Our effort was cross-disciplinary, performing the best possible

environmental sampling with tools and supplies readily available for citizen scientists to monitor

the environment in which they live, to address their concerns in terms of environmental and public

health, as well as for strategies to develop more sustainable socio-eco-environmental services. This

study is the first to establish the close relationship between fast fashion waste and its subsequent

release of microplastics and microfibers in the environment, both in the air and water. Our results

add to the evidence emphasizing the significant environmental burden disproportionately borne by

the communities that have become the de facto stewards of global waste (Clapp 2025).

METHODS

The study area was the Kantamanto secondhand textile market in Accra, Ghana, the nearby Korle

Lagoon and the contiguous beachfront (Fig. 1), which we investigated in May 2023 for

microplastic and specifically microfiber concentrations, using waterborne samples as well as

airborne samples. Considering the dependence of our analyses on environmental conditions, we

used an ATMOS41 ZL6 weather station (https://metergroup.com/products/atmos-41/) placed at

the headquarters of The Or Foundation, which is located on the outskirts of the market, to track

weather data. The weather station recorded a series of parameters every 15 min., the most

important ones used in this study being the time of day when episodes of rain occurred and the

amount of rainfall recorded in mm. For microplastic counts, we needed to collect actual volumes

of rainwater (expressed in mL), which we did using a Stratus Rain Gauge

(https://www.scientificsales.com/6330-Stratus-Rain-Gauge-p/6330.htm) placed next to the

ATMOS Station (Online Resource 6). Although the ATMOS Station collected data continuously,

the rain gauge collected rainfall water only during time periods that were possible and safe for the

citizen scientists to place the gauge in and out of its operating location on the roof. As such, the

volumes of rainwater analyzed did not correlate exactly with the precision rainfall measurements

of the ATMOS Station due to the difference in the respective range of operating periods. Our study

does not intend to be a weather study and for this reason we referred to the amount of rainwater

collected in mL, which is the most accurate for our purpose, while disclosing mm of precipitation

for reference. Following collection, rainwater was then analyzed for concentration of microplastics

and microfibers as described below, based on volumes of mL collected.

Sample collection for microplastics and microfibers has become (since 2020) a routine

monitoring activity of The Or Foundation (Ghana), which was put together and optimized over the

years with the Deheyn lab at Scripps Oceanography (California, USA). The work in the field, and

in the lab, is always done with what is readily available locally in a non-academic infrastructure,

keeping in mind the possibility of contamination (Chen et al. 2020; Prata et al. 2020; Song et al.

2021; Ma et al. 2024), which is addressed with various precautions and controls. In the field,

collectors wear bright yellow safety jackets and bright blue/yellow gloves (the microfibers of

which are easy to detect under an imaging microscope in brightfield illumination), and samples

are collected standing downstream from the actual collection process. Samples are brought back

in closed and cleaned containers to The Or Foundation headquarters. There they are processed in

an enclosed laboratory with no windows and no direct exposure to outside air, thus being relatively

sealed off from the surrounding activities of the local area. The samples are processed using

conventional guidelines against contamination, only using tap water passed through a ceramic clay

filter (pore size < 1μm) and limiting direct exposure to air flow (samples are protected by non-

sticky parchment paper and aluminum foil when needed). Negative (blank) controls of all the

processing steps are performed on a regular basis (on average once a month or when something is

changed in the laboratory, e.g. water containers) in order to identify possible airborne and

waterborne sources of contamination. This includes running the processing steps of sample

preparation as usual but without the sample. Such blanks are also run for each solution used, for

each step (e.g., sitting in an oven, fridge or under the microscope), considering the same volumes

and times as for regular sample processing. The concentrations of microparticles found in this

range of controls are considered "negative" because they are not associated with the sample and

are removed from the data presented. For this paper we focused on the month of May 2023,

because this was a month of heavy rain during which all of our sensors worked without interruption

(viz. there was no power outage from the city grid).

On-line monitoring of airborne microparticles. A key air quality monitoring sensor is the

PurpleAir, a small "accessible-to-everyone" instrument that detects concentrations of particulate

matter in air at a given time point (Ko et al. 2020), which has been used widely around the world

by people (technically citizen scientists) interested in knowing the quality of the air they breathe

(note that the PurpleAir does not collect or trap the airborne microparticles). The monitoring is

continuous and the data (often made publicly available in real-time) characterizes the air quality

in a few categories (www.PurpleAir.com), which are in line with the US EPA Clean Air Act

quality standards (Barkjohn et al. 2021; Jaffe et al. 2023).

In this study, PurpleAir PA-II-Flex air quality sensors were used to monitor concentrations

of PM2.5 (smaller than 2.5 microns) and PM10.0 (smaller than 10 microns) particles expressed as

mass concentration ($\mu g/m^3$) in the air, which is the conventional unit to report air quality under the

standards of the US EPA Clean Air Act. This mass concentration output, however, originates from

raw numbers of particles that are transformed into mass considering specific algorithms (see

www.PurpleAir.com) and assumptions about the particles (mostly that they are pollen, dust, and/or

black carbon from improper fuel combustion). This mass concentration could therefore be biased

under conditions with heavy concentrations of airborne microplastics (which we assumed would

be the case around Kantamanto Market, considering its substantial volume of synthetic clothing).

As such, we also used the raw data of these sensors, using the PM2.5 and PM10.0 particle counts

directly, which are expressed in dL by the sensor (and expressed in L here). These particle count

concentrations can therefore be more reliable to compare to the data obtained from the Coriolis®

air sampler (see section below).

One PurpleAir sensor was located next to the ATMOS41 weather station, in the area near

the market corresponding to the upper deck (4[th] floor, about 12 m above street level) of The Or

Foundation headquarters (Fig. 1). Another sensor was positioned on a metal pole at either 5 m or

10 m height and moved around for measurements in different locations, including in the heart of

the market. Measurements were also taken 5 km away from the market, in Osu, a residential area

of Accra. Data from the PurpleAir were processed according to the protocol recommended by the

vendor. (The raw data, which was collected every 2 min, was organized in a spreadsheet and

curated for any missing data, such as for when the instrument was saturated e.g., or for when power

supply instability would make it miss some measurements). Although PurpleAir sensors measure

the concentration of particles within their range size, they do not identify what the particles are in

terms of their chemical makeup. As such, they do not identify whether particles have a natural

basis (dust, pollen vs black carbon) or are purely synthetic, like microplastics (e.g., such as tire

shred particles or microfibers from textile fabrics). For this reason, airborne particles had to be

collected for infrared fingerprint analysis to identify their chemical nature.

Airborne sample collection. A Coriolis Compact portable air sampler instrument (Bertin

Technologies) was used to collect airborne microparticles between 0.5 and 10 μm from the same

locations as the PurpleAir sensors. (Note that the manufacturer has validated the optimal efficiency

of the Coriolis for this size range of microparticles, which does not exclude that larger particles

can also be collected, since there is no physical size exclusion mechanism in place but rather

pelleting of particles from generated cyclonic air flow conditions in the instrument;

https://www.bertin-technologies.com/products-range/air-samplers/). Because the instrument

collects the microparticles, it cannot be run 24/7 like the PurpleAir, which is used for monitoring

the microparticles in the air without collecting them and knowing what they are. Thus, the

PurpleAir and the Coriolis instruments appear to complete complementary functions, which we

took advantage of in this study. Here, the Coriolis was run for 2 hrs each time at each location,

with a vacuum rate of 50 L/min as indicated by the manufacturer specifications. The particles that

were collected in the 10 mL cone receiver of the instrument were retrieved according to the

following process (given that MilliQ water is not readily available): Tap water filtered through

ceramic clay filters with pore size <1 µm was added to the cone, shaken using a Vortex, and poured

into a 50 mL Falcon tube. The process was repeated five times to make sure all the particles were

transferred.

Waterborne sample collection from the lagoon. Water was collected from 7 sites (named OKS8

through OKS18) along the lagoon, in order to have samples upstream, running next to, and

downstream from the dumpsite that receives significant quantities of textile waste leaving

Kantamanto Market (Fig. 1). A permanent cement beam structure (aka surface barrier) crosses the

width of the lagoon between OKS20 and OKS26. This surface barrier is about 40 m long and about

1.5 m tall, and the watermark reaches about half-way up the height of the barrier. As such, the

surface barrier stops floating waste from moving further down the lagoon and causes items to build

up in massive amounts upstream, to the point where the water surface can barely be seen in OKS20.

In contrast, OKS26 (located just downstream from the surface barrier) has much less floating waste

since it receives the water that flows under the surface barrier, from about 0.75 m below the water

surface. The surface barrier thus does not prevent heavier waste (sub-surface, midwater or

tumbling on the seafloor) from flowing under the structure, downstream in the lagoon and into the

ocean.

Considering the poor quality of the water in the lagoon, which is due in part to raw sewage

input (Nixon et al. 2007; Koppelaar et al. 2018; Owusu et al. 2023), surface water collection was

performed using an extended pole. The pole was equipped with a dedicated 5 L bucket made of a

clean, bright blue plastic easily identifiable under microscope imaging (for blank background).

This way, the water collection was safe for the operator. Water was collected with the bucket, the

bucket was rinsed and emptied on the ground, and water was collected again 2 separate times at

the same site. Each time two 50 mL Falcon tubes (N=4 total) were rinsed three times with lagoon

water from the bucket, shaken vigorously. emptied, and rinsed/emptied again two times, and then

filled with water from the bucket for the fourth time (for analysis), ensuring that the tubes were

submerged sub-surface during collection (to limit any possibility of airborne contamination). The

tubes were capped, labelled and transported in a cooler back to the laboratory.

The water collection method put in place in this study, based on collecting 50 mL Falcon

tubes of liquid, is unique in allowing samples to be collected from almost anywhere, and by almost

anyone. This is powerful for gathering a large number of samples (and replicates) from places that

are difficult to access in time or space. This methodology therefore leads to numbers of

microplastics and microfibers for every 50mL of sample collected, which is the concentration that

will be expressed throughout the study (counts/50mL), since expressing the data per L (as is

common) could induce a biased extrapolation. Our data can be compared to microplastic

concentrations found in the literature, yet keeping in mind the difference in methodology and the

difference in sample size and processing. Indeed, most studies focusing on microplastics use the

Manta Net to trawl large volumes of water, thus results are expressed in unitary m^3 or L, and only

available from coastal areas where boats can operate. The Manta Net usually has a mesh size of

100-500 μm, thus collecting larger microplastics, rather than the smaller microplastics (<100 μm),

as we do in this study. This difference in methodology relative to coastal baseline studies on

microplastics and the difference in size range of microplastics collected, needs to be kept at the

forefront when reading this paper, with the difference in units a reminder of these fundamental

differences.

Samples processing for microplastic and microfiber counts. Samples were brought back to the

laboratory where they were processed for microplastic and microfiber counts following a routine

protocol used in the Deheyn Lab at Scripps Oceanography, (UC San Diego, USA) and

implemented at The Or Foundation (see Online Resources). This involved filtering the samples on

glass fiber filters (0.6 μm mesh size to retain very small microplastics) and imaging the dried filters

under a stereomicroscope. The imaging was done in bright field as well as epifluorescence (ex.

360-380 nm, em. >415 nm Long Pass filter; see Online Resource 1-2 for details). Microplastics

and microfibers were counted by at least 3 individuals (and up to 6 individuals) using both the

bright field image and the epifluorescence images, since they showed different items (Online

Resources 1-2).

The epifluorescence property of materials of different colors was addressed in the Deheyn

Lab, using reference fabric material provided by industry (Lenzing AG and Archroma). We

showed that synthetic white polymers (e.g., polyester, elastane, nylon) displayed vivid blue/cyan

blue fluorescence, while their colored counterparts showed no visible fluorescence under the same

settings. When mixed with plant debris or phytoplankton, the blue/cyan blue fluorescence was

visible together with the red fluorescence of chlorophyll, which we expected was the most

representative of what we would find on our filters. As for the nature-based polymers (e.g., cotton,

lyocell), they had no visible fluorescence, whether white or colored, under the settings used for

viewing their synthetic equivalent (Online Resource 1).

The intent was not to perform an exhaustive study on the effect of color dyes on the

fluorescence but to demonstrate using key polymers and representative colors that our combination

of bright field and epifluorescence counting was comprehensive enough to sort microparticles into

four categories: (1) Colored microparticles from a variety of sources, including textiles, degrading

plastic items, or even just silt grains (bright field counts); (2) Microparticles from natural debris

rich in chlorophyll, whether decaying plants, filamentous algae or cyanobacteria (red fluorescence

counts); (3) Synthetic microplastics (blue fluorescence dots, flakes or particles); and (4) Synthetic

microfibers (blue fluorescence fibers) (Online Resource 2). This combination of bright field and

epifluorescence imaging thus allowed us to broadly cover different situations of the life of plastic.

For example, an unweathered piece of colored plastic will be seen in bright field but not necessarily

in fluorescence, since dyes could affect the innate synthetic polymer fluorescence by shifting its

intensity and/or color (Grieve et al. 2005; Biermann 2007). In contrast, weathered pieces of plastic,

invisible in bright field, will emit bright blue fluorescence. As for red fluorescent items, they will

likely represent a rich content in chlorophyll, which could be seen in bright field as well since they

display a green color, unless the red fluorescence originates from a synthetic dye, which would

likely be seen in red in bright field (Online Resource 2).

The synthetic microfiber counts, which represented our main interest, are discussed in the Results

section. (The counts from the other analyses, which are from bright field, microparticles red

fluorescence, and synthetic blue, fluorescent microparticles are available as Online Resources.

They show the same trends as the synthetic microfibers but in different ranges. Nonetheless,

similar conclusions can be drawn from this data.)

A subset of samples (35 items) was processed for polymer chemical identification using Quantum

Cascade Laser InfraRed (QCL-IR) spectral imaging, using the Spero® IR system (DLS

DayLightSolutions Inc.) (Primpke et al. 2020), which is available at the Deheyn Lab (see Online

Resource for detailed methodology). The infrared hyperspectral cubes (950 to 1800 cm^{-1})

generated from each sample were compared to those of 23 reference standards for plastics

(PolymerKit 1.0; Hawaii Pacific University) and 7 raw textile fibers that were provided by industry

collaborators at Lenzing AG and Archroma. The infrared fingerprint spectral data for three

common polymers standards were overlapped over the spectral data collected from some

microparticles to illustrate our polymer assignment (see Online Resource 3).

Textile waste by brand tag. We assessed the relationship between the textile waste as whole

garments collected directly from retailers in Kantamanto Market and the macro textile waste

pollution found in the lagoon and at the beaches flanking the opening of the lagoon. For this, we

compared the brand tags from garments found at the market (172 garments) with those found at

the beach (128 garments) (see Online Resource for details). We expected the assembly of brands

from the beach textiles to reflect the ones from the market, since we presumed that the discarded

garments somehow (through the lagoon flow) made their way to the ocean, where the tides then

pushed them back on the local beaches. We then correlated the two sets of tags using a Pearson

correlation coefficient to assess their potential relationship, viz. the market, via the dumpsite, being

the source of the textile waste found on the beach. In addition, we recorded the polymer textile

fiber composition of the garments (relative content of cotton and polyester in each garment, for

example) as indicated in the product tag. In that case, we expected their polymer composition to

reflect the one of microfibers found in the airways and waterways.

Statistical analysis. This study aims to provide a snapshot of the concentration of microplastics

and microfibers in the dynamic environment of the world's largest second-hand textile market. A

diversity of variables is considered, from textile-related merchandising activities to textile burning,

in addition to weather variables and to the schedule of the crowd coming in and out of the market.

Our analyses therefore mainly rely on descriptive statistics using box plot displays, with data

curing performed in Excel (Microsoft Corp) and figures made in Delta Graph (Red Rock

Software). Although our intentions were not to test the particular level of significance of variables,

we performed the research with as many replicates and independent controls as possible, in order

to provide a robust assessment of effects and trends. The only "experimental" assessment we did,

which deserved statistical assessment, was when we compared the assembly of brand tags from

the market textile waste (which we manually pulled out of the market) versus the one from the

beach textile waste (which we dug out from the sand). We therefore ranked the brands from the

most abundant to the least abundant based on the number of tags found for each brand and used

Excel to run a Spearman ρ correlation coefficient analysis considering a significance level of

$\alpha<0.05$.

RESULTS

The results showed that the clothing from global sources sold into Kantamanto Market and

discarded from the market as waste is a source of microparticles, but that the release and

distribution of these microparticles can show large discrepancies, based mostly on market activities

and weather (rain), which are variables both in time and space. The synthetic microfiber counts

we report here always showed high values compared to their blank controls equivalent for

background contamination. In general, our various controls accounted for <2 % of counts

compared to those from processed samples (Online Resource 4-8).

Of the microparticles collected via air sampling (Coriolis analysis) and counted by microscope

imaging, we show here the data for the synthetic microfibers and microplastics identified in

epifluorescence, while the other imaging data is shown in Online Resource 5-8. Overall, all the

imaging data followed the same trend but in different ranges of counts. In general, bright field

counts were always the lowest and below one hundred, followed by red fluorescence counts and

the blue fluorescence microfiber counts (both about 10x greater than the bright field counts) and

blue fluorescence microparticle counts (about 10x greater than the blue fluorescence microfiber

counts). Broadly, this indicates our samples contained a sustained level of recently released

microparticles: a large amount of natural microparticles and synthetic microfibers and a much

greater amount of microplastics (Online Resource 5-8).

PurpleAir data were up to 5x greater within the market. PurpleAir sensors allowed us to collect

data over a continuous 24 hr period and to measure both mass concentration and raw counts of

particles (N=720 data points each). The PurpleAir sensors across the different areas of town and

at the different height (5 m, 10 m) all showed similar mass concentration data in terms of the

relative abundance of PM10 and PM2.5 particles (as expected since PM10 data encompass PM2.5

data), which systematically showed PM10 mass concentration data up to 125% greater than that

of PM2.5 (Online Resource 4A). The most striking difference across sites was regarding the

variability of the data, which was more variable in the market than off market, and which thus

translated into a greater difference between the means and the medians for each of these sites

(Online Resource 4A). Over the course of May 29, 2023, which was a day with no rain (last rain

event was about 40 hrs prior) PM2.5 in the air at a given timepoint ranged from 3 to 356 $\mu g/m^3$,

while those of PM10 ranged from 4 to 372 $\mu g/m^3$. The mass concentration was systematically

greater for the sensors placed in the market, especially for the one at 10 m height (PM10 up to 372

$\mu g/m^3$), which was slightly greater than the one at 5 m height (PM10 up to 332 $\mu g/m^3$). The mass

concentration decreased when moving away from the market, whether at its outskirts at The Or

Foundation headquarters at 12 m (up to 103 $\mu g/m^3$) or in the residential area control at 2 m (with

up to 94 $\mu g/m^3$) (Online Resource 4A). The corresponding data with respect to microparticle count

concentration (raw data generated by the PurpleAir) showed similar characteristics, with greater

count concentration and greater variability for sites within the market (Online Resource 4B). This

was especially true for data of PM2.5 particles, which was up to 3,467 counts/L in the market and

up to 912 counts/L off the market. The particle count concentration for PM10 were lower than for

PM2.5, as expected, reaching 172 counts/L in the market and up to 226 counts/L off the market,

indicating that larger microparticles can also be found off market (Online Resource 4B), but

whether they originate from the market remains to be determined (see assessment from the section

on the Coriolis data below).

The trend in the data, showing greater values in the market relative to the outskirts and

residential areas, and, within the market, higher concentrations at 10 m relative to 5 m, was

observed throughout our study, yet was highly variable from day to day and within a given day.

This was dependent on the activities at the market (most active from 05:00 until 18:00 every day

but starting as early as 02:00 for the vendors) with possible burning of waste in locations

immediately surrounding the market occurring overnight. Vehicle traffic on the market periphery

and foot traffic across its web of stalls could also affect the PurpleAir data, but also, and most

importantly perhaps, the weather, especially in relation to wind and rain. Indeed, data collected

during days when rain occurred decreased sharply as soon as rain started before increasing again

gradually after the rain stopped (Fig. 2). Microplastic particles can make up a portion of the overall

PM10 and PM2.5 data, and therefore are a component of the air we breathe, with potential impacts

on human health. It was therefore critical for us to look at the actual content of microplastics and

microfibers for both airborne and rainwater samples.

Airborne microparticles were dominated by synthetic textile microfibers. Data from the Coriolis

air sampler showed relatively high concentrations of microfibers across all sites of investigation

within the market, ranging from about 20 at HQ to about 180 microfibers per 6,000 L in the market

at 10 m height, equivalent to 0.004 to 0.029 microfibers/L (Fig. 3). In general, we found that the

microfiber concentrations were greater for the samples analyzed at 10 m height compared to the 5

m analysis (even within a ventilated warehouse); this was particularly evident for the samples

taken in the open-air area of the market (Fig. 3). Measurements in the market were about 2x to 5x

greater than the ones made 1 km away, at the outskirts of the market (Fig. 3), which correlated

with the data found with the PurpleAir. Greater concentrations of microfibers were also observed

in bright field conditions, indicating that these microfibers were not weathered and thus likely

originating from the area where the air sample itself was collected (see BF MFs; Online Resource

5). We also found greater concentrations of microplastic particles at 10 m in the market compared

to the same location at 5 m. We assumed that such greater concentrations of microplastic particles

at 10 m were mostly originating from textiles as well considering that their values were also high

in the warehouse where textiles are shredded for upcycling activity, while lower for HQ where

textile work is not performed. (see UV MPs, Online Resource 5). These microplastic particles were

in much greater concentrations than their synthetic microfiber equivalent, which correlated well

with the trend observed for the PurpleAir data where PM2.5 count concentrations were greater

than the ones for PM10.

Rainwater and lagoon samples contained large concentrations of synthetic textile microfibers.

Rainfall was highly variable and characteristic of a tropical weather regime, thus more often with

heavy "showers" scattered in space and time across Accra. We recorded rainfall from 1.5 mm/day

to 80 mm/day. Rainwater samples that were collected from the rainfall ranged from a few mL to a

couple hundred mL. Analyses showed that the rainwater contained synthetic microfibers (Table 1;

Online Resource 6-7) within a range of concentrations (counts/50 mL) that varied with the duration

of the rain event, with greater concentrations of synthetic microfibers (up to 300/50 mL) found in

rainfall from shorter rain events. Indeed, in situations where we collected over 40 mL of rainwater,

which equate to about 19.8 mm of rainfall under weather standards, the concentration of synthetic

microfibers decreased sharply (Fig. 4A). In addition, the concentration of synthetic microfibers

found in rainwater changed based on when the last rain event occurred, and as such, the greater

the extent of dry weather between two rain events, the greater the concentration of synthetic

microfibers found in the rainwater. This was especially observed after 3 days of dry weather (Fig.

4B).

Water samples from Korle Lagoon contained large concentrations of synthetic microfibers,

ranging from 2 to 80 counts/50 mL (Table 1; Online Resource 8-9), which were greater when

closer to the dumpsite and market compared to the locations that were further away (Fig. 5). In

addition, and as expected, the concentrations of synthetic microfibers peaked at OKS20 (between

60 and 80 counts/50 mL, where surface waste is stopped from flowing downstream by the surface

barrier crossing the lagoon (Fig. 5). The data was not as clear for the concentration of microfibers

identified in bright field, suggesting that the microfibers in the water were mostly weathered and

not originating directly from airborne microfibers recently generated in the market (see BF MFs,

Online Resource 8). As for microplastic particles, they too increased in concentration in the lagoon

areas facing the market (up to 550 particles/50 mL), up to where the surface barrier crosses the

lagoon.

Chemical composition of waterborne microparticles reflects that of garments from the market.

Infrared fingerprint analyses (see Online Resource 3 for matching of some spectra with standards)

showed that waterborne microparticles represented a variety of polymers, the most dominant ones

being cotton (29%), polypropylene (26%), polyester (14%), nylon (11%), and polyethylene (9%).

Garments in the market reflect this pattern of polymer ratio as well (Fig. 6), with 40% of the

garments made primarily of cotton, 13% of polyester and 10% of nylon. As for polypropylene and

polyethylene, these polymers are mostly used in plastic bags (used as single-use water containers

and takeaway bags) and single-use water bottles, respectively, which are also found spread all over

the market and lagoon where they are being weathered and breaking down.

Textile waste showed similar brand assemblages between beach and market. Collected tags

showed that the 10 most common brands found in the textile waste were Marks & Spencer,

Associated British Foods (Primark), Asda (George), Sainsbury's (Tu), Next, Boohoo Group, Tesco

(F&F), H&M Group, New Look Group, and GAP, representing up to 63% and 77% of the items

found at the beach and in the market, respectively (Fig. 7). There was a 59% correlation of brands

occurring between the two sites of waste collection, based on the tags analyzed, which was

statistically significant ($P < 0.05$; Pearson ρ correlation coefficient).

DISCUSSION

Our research around Kantamanto Market showed the extensive presence of microplastics and

synthetic microfibers, in both the surrounding air and water, with concentrations seemingly

representative of the varying attributes of the market activity. The data indicate that the

concentrations of airborne microparticles we found varied depending on the day of the week and

activities taking place on those days, with concentrations notably being higher after every

Thursday, when bales of globally sourced secondhand clothing are delivered to the market. The

concentrations were also higher during early morning on Saturdays, a busy market day when bales

are opened and garments are displayed for sale in Kantamanto Market (Skinner 2019). Our study

focused on microfibers, especially their synthetic version, because those are directly related to the

secondhand textile activities in the market, yet keeping in mind that such microfibers could also

have other origins (from outside the market or even a far-away location), while not dismissing the

microplastic components, which can come from a diversity of sources (from tire wear to

microplastics from burning waste, including textiles).

Our study showed, at least during the daytime when analyses were conducted, that

microfibers that are released can float around and be carried to higher elevations (we tested at 10

m height), likely due to warm upward convection currents caused by heat generated by the hot

surfaces heated by the sun, and to which the high number of people in the market can also

contribute (Belzagui et al. 2019; De Falco et al. 2020; Periyasamy and Tehrani-Bagha 2022). Such

upward hot air convection currents are known to occur in nature, where they help with plant seed

dispersal for example (Inatsu et al. 2021; Mun et al. 2024) although they can also be very important

in urban areas where they are considered by city managers for airflow improvements (Xie et al.

2006; Xue and Li 2017; Di Bernardino et al. 2018; Muniz-Gäal et al. 2020). Here, we evidenced

that the microfibers could be carried at least as high as 10 m into the air, although one cannot

exclude that some of them could originate from other sources nearby as well. From there, they

could potentially spread around with winds (we indeed found unweathered microfibers in

neighborhoods around the market; see Online Resource 5) and fall back onto the ground at night,

either when heat-driven convection currents stop or when they become entrained down onto the

ground by rain events (Miao et al. 2023; Liu et al. 2025). Indeed, rainwater was polluted with

microfibers (and microplastics), which was especially true for rain events occurring after 3-4 day

periods of no rain. In such a case, we showed that the first flush of the rain (within the first hour

or so, varying based on the rain intensity) could bring down microfibers in high concentrations.

Practically, from a public health perspective, this indicates that people who collect rainwater for

drinking usage should skip the first flush of the rain event, especially if it has not rained in several

days, or should deploy filtration technology if any is available (like the clay filters commonly

found across Africa).

Our data thus indicates that microplastics and microfibers can enter Korle Lagoon waters

via two routes: One concentrated around the piles of discarded garments that shed fragments upon

mechanical abrasion and weathering in the environment; and the other one more diffuse and

associated with aerial deposition of airborne microplastics and microfibers, that can also be

precipitated during rain events, and reach the lagoon via runoff and surface water drain outflows.

Ultimately, using Kantamanto as a case study representing the communities that receive,

sort and process global fashion waste, we show that the high volume and low quality of garments

arriving in these communities and the lack of infrastructure and capital to safely and efficiently

deal with this waste, results in significant microfiber pollution. This pollution has the capacity to

impact local air quality as well as local aquatic ecosystems and their associated services, thus

potentially contributing to chronic public health issues (Manisalidis et al. 2020; Sridharan et al.

2021; Zhao et al. 2023; Song et al. 2024; Vasse and Melgert 2024).

Global textiles as a source of airborne microparticles. Using the PurpleAir sensors, we

demonstrated that the mass concentrations of PM2.5 and PM10 particles were high and varied

extensively around and near the market, decreasing when going further away from it. This

indicated that the market was a source for airborne microparticles.

PM2.5 airborne microparticle mass concentrations in different neighborhoods in Accra

have been reported to often be above the healthy air quality guidelines recommended by the World

Health Organization, which is set at 10 μg/m^3 (Zhou et al. 2011; Ofosu et al. 2012), while rarely

surpassing 100 μg/m^3 (Alli et al. 2021). Our study showed PM2.5 mass concentrations within the

market can often reach levels up to between 125 and 175 μg/m^3 (Online Resource 4A), sometimes

peaking at much higher concentration levels around 1,300 μg/m^3 (Fig. 2), which thus can be 10x-

20x greater than for various areas in Accra. Because our study took place in the core of a textile

market, and knowing that textiles shed microfibers (and/or fiber fragments that could be shorter

and more like "elongated" microplastics) that can be <10 μm in size, it is fair to assume that our

PM2.5 as well as PM10 mass concentrations could also include microfiber and/or shorter fiber

fragments. This would be in addition to other microparticles typically found in cities (related for

example to energy combustion, industry and road traffic) but also pollen and sand dust, often

carried by sub-Saharan winds to Accra (Amegah et al. 2017).

Aside from the abnormally high airborne microparticle mass concentrations in

Kantamanto, another peculiarity was that the PM10 mass concentrations were only about 1.0-1.2x

greater than those of PM2.5. This was unlike other studies in Accra that showed PM10 values

being 2.0-2.5x greater than PM2.5 (Arku et al. 2008; Dionisio et al. 2010). Because our PM10

mass concentrations were relatively elevated, the lower PM2.5/PM10 ratio in our study indicates

that the source of PM2.5 is likely greater in the market environment, relative to its surroundings.

PM2.5 values tend to be associated with byproducts of combustion such as those from fuel-

powered engines (Arku et al. 2008; Dionisio et al. 2010). However, because the interior of the

market (where measurements were made) is largely closed to vehicular traffic, the increased

PM2.5 count concentration cannot be from on-site fuel-powered engines but likely stems from

incomplete fire combustion of plastic material. Burning plastic waste is indeed a common practice

in Kantamanto, which is done without wood, and not for the need of cooking or getting warmer,

but for the need to discard waste and gain space. Such frequent (almost every night) and

uncontrolled incineration of waste material is also done using textile waste, which is mostly made

of plastic that, when burned, releases large numbers of PM2.5 microparticles (Bardales Cruz et al.

2022; Islam et al. 2022). Although another source of PM2.5 could originate from the individual

use of cooking stoves across the market (Zhou et al. 2011), the importance of this practice is

definitely less than the textile waste burning. The latter is likely the most prevalent indeed

considering the many small fluorescent particles that were observed under epifluorescence in our

study, which makes them synthetic (plastic), since black carbon microparticles from engine

combustion and incineration practices with wood (both important sources of PM2.5 and PM10

microparticles in general) have not been previously reported to show native epifluorescence

properties. In fact, additional staining methods are needed to image black carbon microparticles in

fluorescence (Chen et al. 2019). The native fluorescence of microparticles in our study then

indicates their synthetic (plastic) nature.

As expected, air quality in Accra shows daily variation relative to people's activities,

whether from cooking or driving to/from work (Alli et al. 2021), and/or from textile market

activities and the burning of garments, as indicated by this study. We also showed here that

airborne PM2.5 and PM10 microparticle concentrations decreased sharply in events of rain

indicating that rainfall can be a major process to help microparticles, including microplastics and

microfibers, transition from the airborne phase to the waterborne phase. This indicates that rain

can be a major process by which airborne microfibers enter the waterways, in addition to wind

driven processes that can deposit them onto water surfaces. Such an effect of rain bringing down

airborne microplastics was recently described in other industrial settings (Do et al. 2023).

Global textile waste as a source of microparticles for waterways. We evidenced that Korle Lagoon

is a receptacle of textile microfibers from runoff. Indeed, water samples collected from the lagoon

showed high concentrations of microfibers (and other microplastics as well), with a spatial

distribution along the lagoon and chemical makeup indicative of an origin associated with market

activities. In particular, runoff following rain events and/or weathering fragmentation of garments

discarded directly in the environment are likely a sustained source of microparticles for the lagoon.

This is in addition to the microfibers probably reaching the water surface from airborne deposition.

A majority of garments in the market contain polyester and/or cotton, which was also reflected in

the waterborne microparticles, especially in waterways next to, and downstream from, the market.

Our study thus highlights the extent to which the increase in textile production, and therefore textile

waste, contributes to the release of microplastics and microfibers in the environment, either

atmospheric or aquatic. This information is critical because microplastics and plastic microfibers

are not biodegradable in a biologically relevant timeframe, making them accumulate in the

environment where their long residence time makes them available to enter the food web and

accumulate in organisms, including humans (Sharma et al. 2022; Thompson et al. 2024; Zhao et

al. 2024). This, in turn, can affect the overall ecosystem services (Santana et al. 2017, Olmo-

Gilabert et al. 2024).

Concentrations of microfibers and microplastics were greater than in other parts of the world.

The concentrations of microplastics, including microfibers, found in our study, whether in the air

or in the water, were usually by far the highest ever reported in an environment (e.g., Sazli et al.

2023). This does not come as a surprise considering that methodologies used can lead to

underestimation of microplastics, both from airborne and waterborne sources (Xu et al. 2020). In

addition, a direct comparison across studies is challenging because the methodology for sample

collection and/or for microparticle counting usually varies from study to study, based on local

environmental limitations, local expertise, the distinction (or not) between microplastics and

microfibers, but also on the scientific question driving a particular study. Zhao et al. (2023)

provided a relevant range of comparison for levels of airborne microplastics in megacities, with

concentrations of microplastics possibly ranging broadly from <1 to >1,000 per m^3, depending on

environmental factors (e.g., winds, wildfires, smog conditions) and methods of collection (e.g.,

dry or wet passive atmospheric fallout vs. active suspended particle measurement). However,

under normal circumstances and using suspended particle measurement as a reference (thus not

brought down by rain or snow), others have measured relatively low concentrations, e.g. around 4

microplastics per m^3 in Shanghai (Sridharan et al. 2021), 1.5 per m^3 in Paris, and 0.6 m^3 in the

French Alps (Xu et al. 2024), often with no quantified distinction between microfiber and

microplastic particles, except for Xu and colleagues who identified that 10% to 25% of the

microplastics were microfibers. In our study, the Coriolis data showed about 100 to 700

microplastics per m^3, thus about 100x what was found in Shanghai, and about 3 to 30 microfibers

per m^3 (considering only UV synthetic MFs, and not the BF MFs; see Online Resource 5), which

is about 20x greater than in Shanghai and about 125x the numbers from the French Alps.

A similar analysis can be done with waterborne microfibers and microplastics.

Microplastics in Kantamanto can originate from degrading plastic bags and disposable individual

drinking water pouches (the main source of drinking water around the market), used extensively

and discarded on the ground, with limited direct contribution from tire wear as otherwise found

elsewhere in the city (Arku et al. 2008; Zhou et al. 2011; Knight et al. 2020; Amegah et al. 2022;

Safo-Adu et al. 2024). As for microfibers their main source would be shedding from clothing (Carr

2017; Carney Almroth et al. 2018; Liu et al. 2019; Nowack et al. 2021; Liu et al. 2023).

Concentrations of microfibers have been reported at around 2-30 counts per L in coastal waters

(Suaria et al. 2020; Tang et al. 2023), which does not include lagoons as these have been much

less investigated around the world for their content in microfibers. The same can be said for

microplastics, although in this case one comprehensive study gathered data from the literature

since 2000 (Garcés-Ordóñez et al. 2022), showing concentrations of 20 to 90 counts per L on

average for lagoons around the world, with about 40% of these numbers attributed to microfibers,

thus making the count to about 12-54 microplastics per L and thus about 8-36 microfibers per L.

This is in line with studies reporting up to 30 counts of microplastics per L in harbors and with

0.5-2.0 counts per L in coastal and more open ocean waters (Suaria et al. 2020; Tsang et al. 2020).

The difference in concentrations relative to what we found in Korle Lagoon waters (between 100

to 1,600 microfibers per L and between 1,000 to 11,000 microplastics per L; Online Resource 8)

is staggering and likely due to the source being the abundance of textiles in the market as well as

in the dumpsite (as waste) along the lagoon. Such high concentrations (up to 25,000 counts per L)

were previously observed in lagoons in the Florida Keys (USA) (Badylak et al. 2021) and were

considered the result of debris (mostly polystyrene pellets) coming from recent hurricane activities

and accumulation in specific areas of lagoons due to winds and currents. This was not the case for

the data collected in Korle Lagoon, which shows a slow but steady flow to the ocean.

Another factor possibly contributing to the difference in concentrations of microplastics

found across studies could also originate from the differences in methods used to collect water

samples and to analyze the water samples. Indeed, a common way to collect microplastics involves

a Manta net, which is not easily used in a shallow lagoon (Royer and Deheyn 2019); this method

has also been documented to underestimate the numbers of microfibers that can pass through the

mesh due to their elongated shape, and because the mesh size is usually larger, around 500 μm

(Barrows et al. 2018; Du et al. 2022; Concato et al. 2023). Hence, the technique to collect set

volumes of water for subsequent filtration and counts of microfibers (and microplastics) retained

on filters, is likely more representative of the actual waterborne microfiber concentrations.

Textile waste from global sources passing through Kantamanto Market contaminates the coastal

ecosystem. Beaches in Accra have masses of entangled garments that arise on and under the sand.

Composed of dozens of garments, these masses develop into long tentacles (bundles as much as

30 cm in diameter and meters long) that go deep into the sand, and their extraction, through beach

cleanups organized weekly by The Or Foundation, represent a monumental effort. It is unclear

how long it takes for garments to form tentacles like these with the back and forth of the waves

and tides, and for them to be buried deep into the beach. However, the local coastal community,

including The Or Foundation's Beach Monitoring team, which counts the tentacles every week,

has observed them for several years (The Or Foundation 2022). Taking tags from garments on the

beach and comparing them to the tags from the textile waste in the market demonstrated that

garments found at the beach originate as part of the global secondhand clothing trade passing

through Kantamanto.

The tags showed that the 10 most abundant brands found at the beach and as market waste

correlate well with one another. While it is possible that some of the garments are trashed directly

at the beach after purchase and use for a little while, it is unlikely that such "isolated"

circumstances would lead to the formation of the long tentacles that have been observed.

Brands of the garments most commonly found as waste on the beach and as waste within

Kantamanto Market were from the Global North, with companies headquartered and distributed

principally within the USA, the EU and the UK. Most of these companies are well known to be

involved in the fast fashion economy (Boohoo, Gap, George, H&M, New Look, Next, Primark),

although not always recognized as such by consumers (as is the case for Marks & Spencer, for

example). Thus, within the fast fashion economy, whereby consumers are influenced to rapidly

change wardrobes at relatively affordable prices (The Economist 2015), garments can be

considered environmental pollutants of emerging concern when discarded as waste at their end of

life due to the rapid turnover and high volumes of garments in circulation.

A simple collection and processing technique to enable citizen science. In this study, we worked

together with the local community to access sites for collecting samples, to engage community

leaders to implement new waste management practices, and to empower citizens to help with

sample collection. As such, we used a simple and straightforward technique that easily could be

performed by any citizen, with only brief training needed. The technique is effective in collecting

and quantifying microfibers, but also microplastics. The technique relies on small amounts of

water (50 mL) that can easily be sampled by scientists and citizen scientists alike using

standardized tubes, making this attractive for large sampling collection involving the community.

The use of water volumes greater than 50 mL was tested as well (including 1,000 mL, 500 mL,

200 mL; data not shown) but shown to be more challenging due simply to the larger volume to

handle (with increased risks of clogging). It was also more challenging for counting the

microfibers, due to excess debris on the filters, or due to difficulty counting because of the

excessive number of microfibers overlapping each other in the field of view. It must be noted that

in some instances here, the 50 mL volume had to be analyzed 10 mL at a time, using several filters,

to facilitate accurate counting and then added together at the end to represent the one sample (see

Online Resources for more details). The proposed technique thus provides flexibility when dealing

with water masses that are heavily charged with silt, plankton and/or debris, which can clog the

filtration process early on (before reaching the complete 50 mL volume). The proposed technique

(which by default uses triplicates of 50 mL volumes) thus complements the most common one

used for microplastics in particular, using Manta Net towing/trawling, which is bulky, requires a

boat, and is thus difficult for citizens to access.

649
650
CONCLUSION

This citizen science-led study provides a comprehensive assessment of microplastic and microfiber

concentrations in the air, rainwater and lagoon water around Kantamanto Market and its known

dumpsites of global secondhand textile waste. The global textiles and textile waste passing through

the market appear to be a significant source of microfibers (and microplastics), with concentrations

within and around the market sometimes more than 100x greater than in other metropolises around

the world. We demonstrated the dynamic distribution of microparticles following their generation

in the market, with particles in the air precipitating back down with rain events, and possibly with

wind and cooler temperatures. The effects of global textile waste passing through the market and

entering the surrounding environment go beyond the microscopic particles. Indeed, we showed a

strong correlation between garments considered waste at the market and the ones found on the

surrounding beaches, by comparing brand composition of the garments. Overall, we emphasize

the fact that fast fashion contributes to poor water and air quality, which is likely the case across

other densely populated cities worldwide considering that the processes we highlighted in

Kantamanto are not site specific, and could take place in other secondhand clothing markets. As

such, the study highlights the global need for new policy regulations and programs to ensure that

cleanup programs and proper waste management efforts are sufficiently funded along the entire

life cycle of the garments, including for existing and future waste, repurposing and recycling

streams.

GLOSSARY

Fast fashion - An approach to the design, creation, and marketing of clothing fashions that emphasizes making fashion trends quickly and cheaply available to consumers. Fast fashion is characterized by overproduction and the use of cheap (often exclusively synthetic) materials and fabrication techniques.

Fiber fragment - A microscopic fragment that arises from the shedding or degradation of fiber threads from either natural or plastic (synthetic) textile materials. Fiber fragments are characterized by a diameter usually around 8-12 μm and a length that can be greater than the diameter, in which case it would be known as "microfiber." Shorter fiber fragments (with diameter shorter than their length) would then be reported as microplastics.

Kantamanto Market - The world's largest secondhand clothing and upcyling market, located in Accra, Ghana, and featuring over 10,000 vendor stalls.

Microfiber - A microscopic fragment that arises from the shedding or disintegration (physical break off) of both natural and plastic textile thread materials. Also known as a "fiber fragment."

Microparticle - Any microscopic particle, regardless of shape, whether synthetic or originating directly from nature (mineral dust or small organic debris).

Microplastic - A microscopic fragment that arises from the disintegration of plastic materials, or that is made small to begin with, such as microplastic beads from cosmetics. As used here, the

term "microplastic" is distinct from "microfiber", although microplastics could be short

microfibers (see fiber fragment definition above) As such, in this study, microplastic counts

represent something distinct from microfiber counts.

Natural microfiber - A microscopic fiber fragment shed from a natural textile material, such as

cotton, wool, or other plant-based sources.

Plastic (or synthetic) microfiber - A type of microplastic with a "fiber shape". Plastic microfibers

typically arise from the shedding or disintegration of textiles, although they can also originate from

tire wear, carpets or upholstery.

706
707
708
709

 FIGURE CAPTIONS

Figure 1. Map (ESRI) of the areas of investigation around Kantamanto Market, indicating the

locations for the PurpleAir sensors and the Coriolis air sampler (green and blue stars, respectively),

and the water sampling sites in the Korle Lagoon (orange circles). Location of the surface barrier

is also visible (green arrow). Main directions of surface water runoff and surface wastewater are

also indicated (yellow arrows). HQ: Headquarters of The Or Foundation; WH: Warehouse, large,

enclosed space immediately adjacent to the market where textile waste is prepared for reuse; OSU:

Residential neighborhood in central Accra, about 5 km from the market.

Figure 2. PM2.5 (grey line) and PM10 (black line) particles mass concentration ($\mu g/m^3$) from the

PurpleAir data recorded every 2 min during a 24 hr period starting at midnight, for 3 different days

with distinct intensity of rain events (date, duration of rain and volume of rainwater collected are

indicated). Period of rain (gray shade) and volume of rainfall was provided by ATMOS41 weather

station. Period of market activity is indicated on each panel. Of note: Thursdays are when new

bales of garments are delivered to the market to be opened; the garments inside are sorted on

Wednesdays and Saturdays.

Figure 3. Airborne synthetic microfiber concentrations (box plots; N=9) across the various sites

of investigation showing that the values were greater in the market and, within the market, greater

at 10 m compared to 5 m. Data were collected with the Coriolis air sampler for 2 hrs each time,

thus representing concentrations of "counts/6,000 L of air". HQ: Headquarters of The Or

Foundation; WH: Warehouse; KM: Kantamanto Market.

Figure 4. Rainwater microparticle count concentrations (box plots; N=9) collected with the Stratus

Rain Gauge (with rain event volumes validated by an ATMOS41 weather station) set on the roof

of The Or Foundation headquarters. **A.** Concentrations of synthetic microfibers from rainwater

samples that correspond to different rain events, thus each with their own volumes (on the X axis).

Data counts on the Y axis are expressed per set volume of 50 mL. The concentration of synthetic

microfibers was high for short rain events (up to 2 hrs long) and decreased steadily as rainfall got

heavier and/or lasted longer. *These samples were collected the same day as the 185 mL rain

event, but just a few hours earlier. This shows that the concentration of synthetic microfibers in

volume of rainwater depends on precedent episodes of rain. **B.** Concentrations of synthetic

microfibers for rain events expressed in relation to the time (number of days) since the last rain

event. Data show that the concentration of synthetic microfibers increases greatly in the rainwater

after 4 days with no rain. Data counts are expressed per set volume of 50 mL.

Figure 5. Concentrations (counts/50 mL) of synthetic microfiber (box plots; N=9) from Korle

Lagoon water showing that the site directly flanking the market in terms of runoff (OKS13) has

the greatest concentrations compare to upstream/downstream of the market, this until the flow of

the lagoon water is slowed down by a surface barrier (OKS20), where floating debris and

microparticles accumulated. All data counts are expressed per set volume of 50 mL of water

collected.

Figure 6. Pie chart representation of the percent occurrence of polymer types found in the

microparticles collected in the lagoon water adjacent to the landfill (OKS 12.5, OKS20) (A, from

InfraRed fingerprint analysis of microplastics) and those found in the waste pile of garments at

Kantamanto Market (B, from material composition tags of garments).

Figure 7. Pie chart representation of the percent occurrence of name tags for each brand found in

the waste pile at Kantamanto Market (top) and found entangled and buried in the sand at the nearby

beach (bottom). More than 60% of the tags originated from just 10 brands (Marks & Spencer,

Associated British Foods, Asda, Sainsbury's, Next, Boohoo Group, Tesco, H&M Group, New

Look Group, and GAP).

Table 1. Minimum (Min), Maximum (Max), Mean and Median values for the counts of synthetic

microfibers and microplastics (both found with native blue fluorescence) from samples of air (in

6,000 L), rainwater (in variable volumes), and Korle Lagoon water (in 50 mL set volume sampling

standard of our methodology). For all counts, N=9, except for the airborne sampling counts (N=3).

Airborne microparticle samples originated from 6,000 L of processed air, and were thus expressed

in concentration of "counts/6,000 L", whereas lagoon water samples were expressed in "counts/50

mL". Data on concentrations for unweathered microparticles (bright field microscope imaging)

and natural microparticles (red fluorescence) can be found in Online Resource 9.

<u>Abbreviations.</u> HQ: Headquarters of The Or Foundation (roof) at two different times (a,b); WH:

Warehouse space adjacent to the market at two different heights (5 m, 10 m); KM Kantamanto

Market (center) at two different heights (5 m, 10 m).

		Synthetic Microfibers					Microplastics			
		Min	**Max**	**Mean**	**Median**		**Min**	**Max**	**Mean**	**Median**
Airborne particles (counts/6,000 L)	**HQa**	8	22	14	11		311	485	414	446
	HQb	7	15	11	10		252	290	275	282
	WH – 5m	12	35	21	16		1034	1700	1454	1629
	WH – 10m	27	48	37	35		1348	1855	1636	1704
	KM – 5m	52	74	60	54		1170	1400	1290	1300
	KM – 10m	99	144	124	128		1195	2930	1790	1244
Rainwater (HQ)	**16 mL**	92	96	94	94		946	1020	971	947
	20 mL	92	101	95	92		1358	1900	1600	1541
	160 mL	23	32	28	30		385	663	486	411
	190 mL	14	18	16	15		421	661	515	462
	180 mL	25	43	35	36		3149	3601	3331	3243
	200 mL	14	16	15	14		1365	1484	1427	1433
	185 mL	100	112	107	108		972	1264	1082	1009
	70 mL	62	71	67	69		738	1092	949	1017
	40 mL	94	126	108	105		1181	1479	1370	1449
	20 mL	9	18	12	10		273	285	279	279
Korle lagoon sites (counts/50 mL)	**OKS18**	3	11	7	6		125	253	187	174
	OKS14	4	10	7	7		251	489	328	269
	OKS13	5	46	24	22		293	600	443	435
	OKS12.5	9	18	13	12		208	532	350	395
	OKS20	53	81	68	68		246	485	357	364
	OKS26	7	15	11	12		74	284	194	182
	OKS8	6	19	13	13		42	267	170	165

779 **Author's contribution.** DDD and BS conceptualized the project. JA performed the field and
780 laboratory work in Ghana, under DDD guidance. DDD performed analysis in California. DDD
781 wrote the first draft of the paper, which was then reviewed and finalized by BS and JA, and several
782 members of The Or Foundation.

Acknowledgments
This work benefited from the work of some volunteers at Scripps Oceanography, including Layla
Nazif and Avery Dougherty, to whom we are thankful for their time. In Ghana the work has been
greatly facilitated by the Accra Metropolitan Assembly, the Kantamanto Used Clothing Sellers
Association, the Kantamanto Hard Workers Association, and the ADAMOG group in Old Fadama.
Funding for the project was provided to BS from The Or Foundation, leveraging financial support
from the SHEIN Global EPR Fund; The Biomimicry Institute; Soliii Earth; and general public
contributions.

REFERENCES

Alli, A. S., S. N. Clark, A. Hughes, J. Nimo, J. Bedford-Moses, S. Baah, J. Wang, J. Vallarino, et al. 2021. Spatial-temporal patterns of ambient fine particulate matter (PM2.5) and black carbon (BC) pollution in Accra. *Environmental Research Letters* 16. doi:10.1088/1748-9326/ac074a.

Amegah, A. K., and S. Agyei-Mensah. 2017. Urban air pollution in Sub-Saharan Africa: Time for action. *Environmental Pollution* 220: 738–743. doi:10.1016/j.envpol.2016.09.042

Amegah, A. K., G. Dakuu, P. Mudu, and J. J. K. Jaakkola. 2022. Particulate matter pollution at traffic hotspots of Accra, Ghana: levels, exposure experiences of street traders, and associated respiratory and cardiovascular symptoms. *Journal of Exposure Science and Environmental Epidemiology* 32: 333–342. doi:10.1038/s41370-021-00357-x.

Arku, R. E., J. Vallarino, K. L. Dionisio, R. Willis, H. Choi, J. G. Wilson, C. Hemphill, S. Agyei-Mensah, et al. 2008. Characterizing air pollution in two low-income neighborhoods in Accra, Ghana. *Science of the Total Environment* 402: 217–231. doi:10.1016/j.scitotenv.2008.04.042.

Badylak, S., E. Phlips, C. Batich, M. Jackson, and A. Wachnicka. 2021. Polystyrene microplastic contamination versus microplankton abundances in two lagoons of the Florida Keys. *Scientific Reports* 11: 1–10. doi:10.1038/s41598-021-85388-y.

Bardales Cruz, M., E. Saikawa, M. Hengstermann, A. Ramirez, J. P. McCracken, and L. M. Thompson. 2022. Plastic waste generation and emissions from the domestic open burning of plastic waste in Guatemala. Environmental Science: Atmospheres 3. *Royal Society of Chemistry*: 156–167. doi:10.1039/d2ea00082b.

Barkjohn, K. K., B. Gantt, and A. L. Clements. 2021. Development and application of a United States-wide correction for PM2.5 data collected with the PurpleAir sensor. *Atmospheric*

Measurement Techniques 14: 4617–4637. doi:10.5194/amt-14-4617-2021.

Barrows, A. P. W., S. E. Cathey, and C. W. Petersen. 2018. Marine environment microfiber

contamination: Global patterns and the diversity of microparticle origins. *Environmental*

Pollution 237: 275–284. doi:10.1016/j.envpol.2018.02.062.

Belzagui, F., M. Crespi, A. Álvarez, C. Gutiérrez-Bouzán, and M. Vilaseca. 2019. Microplastics'

emissions: Microfibers' detachment from textile garments. *Environmental Pollution* 248:

1028–1035. doi:10.1016/j.envpol.2019.02.059.

Biermann, T. W. 2007. Blocks of colour IV: The evidential value of blue and red cotton fibres.

Science and Justice 47: 68–87. doi:10.1016/j.scijus.2006.08.001.

Carney Almroth, B. M., L. Åström, S. Roslund, H. Petersson, M. Johansson, and N. K. Persson.

2018. Quantifying shedding of synthetic fibers from textiles; a source of microplastics

released into the environment. *Environmental Science and Pollution Research* 25: 1191–

1199. doi:10.1007/s11356-017-0528-7.

Carr, S. A. 2017. Sources and dispersive modes of micro-fibers in the environment. *Integrated*

Environmental Assessment and Management 13: 466–469. doi:10.1002/ieam.1916.

Chen, G., Z. Fu, H. Yang, and J. Wang. 2020. An overview of analytical methods for detecting

microplastics in the atmosphere. *TrAC - Trends in Analytical Chemistry* 130: 115981.

doi:10.1016/j.trac.2020.115981.

Chen, L. W. A., M. Zhang, T. Liu, K. Fortier, J. C. Chow, F. Alonzo, R. Kolberg, J. Cao, et al.

2019. Evaluation of epifluorescence methods for quantifying bioaerosols in fine and coarse

particulate air pollution. *Atmospheric Environment* 213: 620–628.

doi:10.1016/j.atmosenv.2019.05.051.

Clapp, A. 2025. *Waste wars: the wild afterlife of your trash.* New York, NY. Little, Brown and

Company. 400 p.

Concato, M., C. Panti, M. Baini, M. Galli, D. Giani, and M. C. Fossi. 2023. Detection of

anthropogenic fibres in marine organisms: Knowledge gaps and methodological issues.

Marine Pollution Bulletin 191: 114949. doi:10.1016/j.marpolbul.2023.114949.

De Falco, F., M. Cocca, M. Avella, and R. C. Thompson. 2020. Microfiber Release to Water, Via

Laundering, and to Air, via Everyday Use: A Comparison between Polyester Clothing with

Differing Textile Parameters. *Environmental Science and Technology* 54: 3288–3296.

doi:10.1021/acs.est.9b06892.

Di Bernardino, A., P. Monti, G. Leuzzi, and G. Querzoli. 2018. Pollutant fluxes in two-

dimensional street canyons. *Urban Climate* 24: 80–93. doi:10.1016/j.uclim.2018.02.002.

Dionisio, K. L., R. E. Arku, A. F. Hughes, O. Jose Vallarin, H. Carmichael, J. D. Spengler, S.

Agyei-Mensah, and M. Ezzati. 2010. Air Pollution in Accra Neighborhoods: Spatial,

Socioeconomic, and Temporal Patterns. *Environmental Science and Technology* 44: 2270–

2276. doi:10.1021/es903276s.

Do, T., Y. Park, B. Lim, S. Kim, M. Y. Chae, and C. H. Chun. 2023. Effect of the first-flush

phenomenon on the quantification of microplastics in rainwater. *Marine Pollution Bulletin*

187. Elsevier Ltd: 114559. doi:10.1016/j.marpolbul.2022.114559.

Du, R., X. Sun, H. Lin, and Z. Pan. 2022. Assessment of manta trawling and two newly-developed

surface water microplastic monitoring techniques in the open sea. *Science of the Total*

Environment 842: 156803. doi:10.1016/j.scitotenv.2022.156803.

Farhana, K., K. Kadirgama, A. S. F. Mahamude, and M. T. Mica. 2022. Energy consumption,

environmental impact, and implementation of renewable energy resources in global textile

industries: an overview towards circularity and sustainability. *Materials Circular Economy*

4. doi:10.1007/s42824-022-00059-1.

Garcés-Ordóñez, O., J. F. Saldarriaga-Vélez, L. F. Espinosa-Díaz, M. Canals, A. Sánchez-Vidal,

and M. Thiel. 2022. A systematic review on microplastic pollution in water, sediments, and

organisms from 50 coastal lagoons across the globe. *Environmental Pollution* 315.

doi:10.1016/j.envpol.2022.120366.

Grazzini, L., D. Acuti, and G. Aiello. 2021. Solving the puzzle of sustainable fashion consumption:

The role of consumers' implicit attitudes and perceived warmth. *Journal of Cleaner*

Production 287: 125579. doi:10.1016/j.jclepro.2020.125579.

Grieve, M. C., T. W. Biermann, and K. Schaub. 2005. The individuality of fibres used to provide

forensic evidence - Not all blue polyesters are the same. *Science and Justice - Journal of the*

Forensic Science Society 45: 13–28. doi:10.1016/S1355-0306(05)71616-6.

Inatsu, M., R. Yoshida, S. Karino, S. Takeuchi, and S. Kobayashi. 2021. A high-resolution

prediction system for birch pollen in Sapporo. *Agricultural and Forest Meteorology* 297:

108229. doi:10.1016/j.agrformet.2020.108229.

Islam, M. R., J. Welker, A. Salam, and E. A. Stone. 2022. Plastic Burning Impacts on Atmospheric

Fine Particulate Matter at Urban and Rural Sites in the USA and Bangladesh. *ACS*

Environmental Au 2: 409–417. doi:10.1021/acsenvironau.1c00054.

Jaffe, D. A., C. Miller, K. Thompson, B. Finley, M. Nelson, J. Ouimette, and E. Andrews. 2023.

An evaluation of the U.S. EPA's correction equation for PurpleAir sensor data in smoke, dust,

and wintertime urban pollution events. *Atmospheric Measurement Techniques* 16: 1311–

1322. doi:10.5194/amt-16-1311-2023.

Knight, L. J., F. N. F. Parker-Jurd, M. Al-Sid-Cheikh, and R. C. Thompson. 2020. Tyre wear

particles: an abundant yet widely unreported microplastic? *Environmental Science and*

Pollution Research 27: 18345–18354. doi:10.1007/s11356-020-08187-4.

Ko, K., S. Cho, and R. R. Rao. 2020. Performance evaluation of low-cost purpleair sensors in

ambient air. *Proceedings - 2020 IEEE 7th International Conference on Data Science and*

Advanced Analytics, DSAA 2020: 563–568. doi:10.1109/DSAA49011.2020.00071.

Koppelaar, R. H. E. M., M. N. Sule, Z. Kis, F. K. Mensah, X. Wang, C. Triantafyllidis, K. H. van

Dam, and N. Shah. 2018. Framework for WASH sector data improvements in data-poor

environments, applied to Accra, Ghana. *Water (Switzerland)* 10: 1–24.

doi:10.3390/W10091278.

Lim, J., J. Choi, A. Won, M. Kim, S. Kim, and C. Yun. 2022. Cause of microfibers found in the

domestic washing process of clothing; focusing on the manufacturing, wearing, and washing

processes. *Fashion and Textiles* 9. doi:10.1186/s40691-022-00306-8.

Liu, J., B. Zhu, L. An, J. Ding, and Y. Xu. 2023. Atmospheric microfibers dominated by natural

and regenerated cellulosic fibers: Explanations from the textile engineering perspective.

Environmental Pollution 317. doi:10.1016/j.envpol.2022.120771.

Liu, J., Y. Yang, J. Ding, B. Zhu, and W. Gao. 2019. Microfibers: a preliminary discussion on

their definition and sources. *Environmental Science and Pollution Research*.

doi:10.1007/s11356-019-06265-w.

Liu, Y., Z. Nie, Y. Meng, G. Liu, Y. Chen, and G. Chai. 2025. Influence of meteorological

conditions on atmospheric microplastic transport and deposition. *Environmental Research*

265. doi:10.1016/j.envres.2024.120460

Ma, H., L. Chao, H. Wan, and Q. Zhu. 2024. Microplastic Pollution in Water Systems:

Characteristics and Control Methods. *Diversity* 16: 1–14. doi:10.3390/d16010070.

Madhav, S., A. Ahamad, P. Singh, and P. K. Mishra. 2018. A review of textile industry: Wet

processing, environmental impacts, and effluent treatment methods. *Environmental Quality*
Management 27: 31–41. doi:10.1002/tqem.21538

Manieson, L. A., and T. Ferrero-Regis. 2021. Castoff from London, Pearls in Kantamanto? A
Critique of Second-hand Clothing. 4th PLATE 2021 Virtual Conference: 1–6.

Manisalidis, I., E. Stavropoulou, A. Stavropoulos, and E. Bezirtzoglou. 2020. Environmental and
Health Impacts of Air Pollution: A Review. *Frontiers in Public Health* 8: 1–13.
doi:10.3389/fpubh.2020.00014.

Miao, C., S. Yu, Y. Zhang, Y. Hu, X. He, and W. Chen. 2023. Assessing outdoor air quality
vertically in an urban street canyon and its response to microclimatic factors. *Journal of*
Environmental Sciences (China) 124: 923–932. doi:10.1016/j.jes.2022.02.021.

Mun, D. S., G. Kang, M. Yang, and J. J. Kim. 2024. How trees' drag and cooling effects influence
airflow and temperature distributions around a street canyon. *Building and Environment* 264:
111913. doi:10.1016/j.buildenv.2024.111913.

Muniz-Gäal, L. P., C. C. Pezzuto, M. F. H. de Carvalho, and L. T. M. Mota. 2020. Urban geometry
and the microclimate of street canyons in tropical climate. *Building and Environment* 169.
doi:10.1016/j.buildenv.2019.106547

Niinimäki, K., G. Peters, H. Dahlbo, P. Perry, T. Rissanen, and A. Gwilt. 2020. The environmental
price of fast fashion. *Nature Reviews – Earth and Environment* 1: 189–200.
doi:10.1038/s43017-020-0039-9.

Nixon, S. W., B. A. Buckley, S. L. Granger, M. Entsua-Mensah, O. Ansa-Asare, M. J. White, R.
A. McKinney, and E. Mensah. 2007. Anthropogenic enrichment and nutrients in some
tropical Lagoons of Ghana, West Africa. *Ecological Applications* 17: 144–164.
doi:10.1890/05-0684.1.

Nowack, B., Y. Cai, D. M. Mitrano, and R. Hufenus. 2021. Formation of fiber fragments during
abrasion of polyester textiles. *Environmental Science and Technology* 55: 8001–8009.
doi:10.1021/acs.est.1c00650.

OEC. 2024. Observatory of Economic Complexity. https://oec.world/en/profile/hs/textiles and
https://oec.world/en/profile/hs/textiles?multihierarchySelector1736-value=af&multihierarchySelector1736-
type=Exporter+Continent

Ofosu, F. G., P. K. Hopke, I. J. K. Aboh, and S. A. Bamford. 2012. Characterization of fine
particulate sources at Ashaiman in Greater Accra, Ghana. *Atmospheric Pollution Research* 3:
301–310. doi:10.5094/APR.2012.033.

Olmo-Gilabert, R., V. Fagiano, C. Alomar, B. Rios-Fuster, M. Compa, and S. Deudero. 2024.
Plastic webs, the new food: Dynamics of microplastics in a Mediterranean food web, key
species as pollution sources and receptors. *Science of the Total Environment* 918: 170719.
doi:10.1016/j.scitotenv.2024.170719.

Owusu, E., E. R. Blankson, R. G. Awunyo, Y. Adams, V. Oduro, B. S. Kpakpo, and F. Gbogbo.
2023. Trace metal load of two urban wetlands with varied catchment activities in Ghana.
Environmental Monitoring and Assessment 195: 1–12. doi:10.1007/s10661-023-11684-x.

Periyasamy, A. P., and A. Tehrani-Bagha. 2022. A review on microplastic emission from textile
materials and its reduction techniques. *Polymer Degradation and Stability* 199: 109901.
doi:10.1016/j.polymdegradstab.2022.109901

Prata, J. C., J. L. Castro, J. P. da Costa, A. C. Duarte, T. Rocha-Santos, and M. Cerqueira. 2020.
The importance of contamination control in airborne fibers and microplastic sampling:
Experiences from indoor and outdoor air sampling in Aveiro, Portugal. *Marine Pollution*
Bulletin 159: 111522. doi:10.1016/j.marpolbul.2020.111522.

Primpke, S., M. Godejohann, and G. Gerdts. 2020. Rapid Identification and Quantification of

Microplastics in the Environment by Quantum Cascade Laser-Based Hyperspectral Infrared Chemical Imaging. *Environmental Science and Technology* 54: 15893–15903. doi:10.1021/acs.est.0c05722.

Royer, S.-J., and D. D. Deheyn. 2019. The technological challenges of dealing with plastics in the environment. *Marine Technology Society Journal* 53. doi:10.4031/MTSJ.53.5.3.

Safo-Adu, G., F. Attiogbe, F. G. Ofosu, I. Emahi, O. Gyampo, and H. Ahiamadjie. 2024. Assessment of Chemical Composition and Health Implications Associated with PM10 Exposure: A Comparative Study of an Urban Road Intersection and a Wood-Burning Vicinity. *Environmental Forensics* 0: 1–15. doi:10.1080/15275922.2024.2431321

Santana, M. F. M., F. T. Moreira, and A. Turra. 2017. Trophic transference of microplastics under a low exposure scenario: Insights on the likelihood of particle cascading along marine food-webs. *Marine Pollution Bulletin* 121: 154–159. doi:10.1016/j.marpolbul.2017.05.061.

Sazli, D., D. Nassouhi, M. B. Ergönül, and S. Atasağun. 2023. A Comprehensive Review on Microplastic Pollution in Aquatic Ecosystems and Their Effects on Aquatic Biota. *Aquatic Sciences and Engineering* 38: 12–46. doi:10.26650/ASE20221186783.

Sharma, S., B. Sharma, and S. Dey Sadhu. 2022. Microplastic profusion in food and drinking water: are microplastics becoming a macroproblem? *Environmental Science: Processes and Impacts* 24: 992–1009. doi:10.1039/d1em00553g.

Skinner, B. 2019. Fashioning Waste: Considering the Global and Local Impacts of the Secondhand Clothing Trade in Accra, Ghana and Charting an Inclusive Path Forward. *Thesis dissertation.* University of Cincinnati. Cincinnati, Ohio.163 p.

Song, S., F. van Dijk, G. F. Vasse, Q. Liu, I. F. Gosselink, E. Weltjens, A. H. V. Remels, M. H. de Jager, et al. 2024. Inhalable Textile Microplastic Fibers Impair Airway Epithelial

Differentiation. *American Journal of Respiratory and Critical Care Medicine* 209: 427–443.

doi:10.1164/rccm.202211-2099OC

Song, Z., K. Liu, X. Wang, N. Wei, C. Zong, C. Li, C. Jiang, Y. He, et al. 2021. To what extent

are we really free from airborne microplastics? *Science of the Total Environment* 754:

142118. doi:10.1016/j.scitotenv.2020.142118.

Sridharan, S., M. Kumar, L. Singh, N. S. Bolan, and M. Saha. 2021. Microplastics as an emerging

source of particulate air pollution: A critical review. *Journal of Hazardous Materials* 418:

126245. doi:10.1016/j.jhazmat.2021.126245.

Suaria, G., A. Achtypi, V. Perold, J. R. Lee, A. Pierucci, T. G. Bornman, S. Aliani, and P. G. Ryan.

2020. Microfibers in oceanic surface waters: A global characterization. *Science Advances* 6:

1–8. doi:10.1126/sciadv.aay8493.

Tang, L., J. C. Feng, C. Li, J. Liang, S. Zhang, and Z. Yang. 2023. Global occurrence, drivers, and

environmental risks of microplastics in marine environments. *Journal of Environmental*

Management 329: 116961. doi:10.1016/j.jenvman.2022.116961.

The Economist 2015. Faster, cheaper fashion - https://www.economist.com/business/2015/09/05/faster-

cheaper-fashion. *Business*, September 5.

The Or Foundation 2022. *Waste makes visible our separation from nature: A "waste landscape"*

report. https://theordev2.s3.amazonaws.com/2023-01/Waste%20Landscape%20Report%20-

%20Compressed.pdf

Thompson, R. C., W. Courtene-Jones, J. Boucher, S. Pahl, K. Raubenheimer, and A. A. Koelmans.

2024. Twenty years of microplastic pollution research-what have we learned? *Science* 386:

eadl2746. doi:10.1126/science.adl2746

Tsang, Y. Y., C. W. Mak, C. Liebich, S. W. Lam, E. T. P. Sze, and K. M. Chan. 2020. Spatial and

temporal variations of coastal microplastic pollution in Hong Kong. *Marine Pollution*

Bulletin 161: 111765. doi:10.1016/j.marpolbul.2020.111765.

UNComtrade. 2023. United Nations Commodity Trade Statistics Database. Retrieved from

http://comtrade.un.org/db.

Vasse, G. F., and B. N. Melgert. 2024. Microplastic and plastic pollution: impact on respiratory

disease and health. *European Respiratory Review* 33. doi:10.1183/16000617.0226-2023.

Xie, X., C. H. Liu, D. Y. C. Leung, and M. K. H. Leung. 2006. Characteristics of air exchange in

a street canyon with ground heating. *Atmospheric Environment* 40: 6396–6409.

doi:10.1016/j.atmosenv.2006.05.050

Xu, C., B. Zhang, C. Gu, C. Shen, S. Yin, M. Aamir, and F. Li. 2020. Are we underestimating the

sources of microplastic pollution in terrestrial environment? *Journal of Hazardous Materials*

400: 123228. doi:10.1016/j.jhazmat.2020.123228.

Xu, X., T. Li, J. Zhen, Y. Jiang, X. Nie, Y. Wang, X. Z. Yuan, H. Mao, et al. 2024. Characterization

of Microplastics in Clouds over Eastern China. *Environmental Science and Technology*

Letters 11: 16–22. doi:10.1021/acs.estlett.3c00729.

Xue, F., and X. Li. 2017. The impact of roadside trees on traffic released PM10 in urban street

canyon: Aerodynamic and deposition effects. *Sustainable Cities and Society* 30: 195–204.

doi:10.1016/j.scs.2017.02.001

Zhang, Q., F. Du, W. Liang, Q. Chen, J. Meng, and H. Shi. 2022. Microfiber fallout during dining

and potential human intake. *Journal of Hazardous Materials* 430: 128477.

doi:10.1016/j.jhazmat.2022.128477.

Zhao, B., P. Rehati, Z. Yang, Z. Cai, C. Guo, and Y. Li. 2024. The potential toxicity of

microplastics on human health. Science of the Total Environment 912: 168946.

doi:10.1016/j.scitotenv.2023.168946

Zhao, X., Y. Zhou, C. Liang, J. Song, S. Yu, G. Liao, P. Zou, K. H. D. Tang, et al. 2023. Airborne

microplastics: Occurrence, sources, fate, risks and mitigation. *Science of the Total*

Environment 858: 159943. doi:10.1016/j.scitotenv.2022.159943.

Zhou, Z., K. L. Dionisio, R. E. Arku, A. Quaye, A. F. Hughes, J. Vallarino, J. D. Spengler, A. Hill,

et al. 2011. Household and community poverty, biomass use, and air pollution in Accra,

Ghana. *Proceedings of the National Academy of Sciences of the United States of America*

108: 11028–11033. doi:10.1073/pnas.1019183108.

1036

1037

Figure 1

OKS18

HQ

Old Fadama:
Informal
Settlement,
Textile Waste,
Dumpsite

Kantamanto market
shops & trade stalls

Other activities: reuse
and repurpose of textile

OKS14

OKS13

WH

OKS12.5

OSU

Residential area (5 km)

OKS20

OKS26

OKS8

500m

N

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7

Market Waste Known Conglomerate %

- Ralph Lauren Corp. 1.2%
- Hanesbrands Inc. 1.7%
- Authentic Brands Group 1.7%
- GAP Inc. 1.7%
- Target Corporation 1.7%
- Frasers Group 1.7%
- Shein Inc 2.3%
- Inditex 2.3%
- Gildan 2.3%
- Asos plc 2.3%
- New Look Group 5.2%
- H&M Group 6.4%
- Tesco plc 6.4%
- Boohoo Group plc 6.4%
- Mark & Spencer plc 13.4%
- Associated British Foods plc 11.6%
- Asda 8.7%
- Sainsbury's 8.7%
- Next plc 8.7%

Beach Waste Known Conglomerate %

- Bestseller 1.6%
- Adidas Group 1.6%
- Liz Clairborne 1.6%
- Target Corporation 1.6%
- C&A 2.3%
- Puma SE 2.3%
- Nike, Inc 2.3%
- Carter's 2.3%
- The Walt Disney Company 2.3%
- Cornerstone Apparel, Inc. 2.3%
- Frasers Group 2.3%
- Tesco plc 2.3%
- Sainsbury's 2.3%
- Boohoo Group plc 2.3%
- Asda 3.1%
- Gildan 3.9%
- Associated British Foods plc 3.9%
- Mark & Spencer plc 14.8%
- Next plc 12.5%
- GAP Inc. 12.5%
- H&M Group 7.8%
- Asos plc 4.7%